

# SMALL COMBINATION OF SLICES, DENTABILITY AND STABILITY RESULTS OF SMALL DIAMETER PROPERTIES IN BANACH SPACES

SUDESHNA BASU<sup>1</sup>, SUSMITA SEAL<sup>2</sup>

ABSTRACT. In this work we study three different versions of small diameter properties of the unit ball in a Banach space and its dual. The related concepts for all closed bounded convex sets of a Banach space was initiated and developed in [B3], [BR], [EW], [GM] was extensively studied in the context of dentability, huskability, Radon Nikodym Property and Krein Milman Property in [GGMS]. We introduce the the Ball Huskable Property (*BHP*), namely, the unit ball has relatively weakly open subsets of arbitrarily small diameter. We compare this property to two related properties, *BSCSP* namely, the unit ball has convex combination of slices of arbitrarily small diameter and *BDP* namely, the closed unit ball has slices of arbitrarily small diameter. We show *BDP* implies *BHP* which in turn implies *BSCSP* and none of the implications can be reversed. We prove similar results for the  $w^*$ -versions. We prove that all these properties are stable under  $l_p$  sum for  $1 \leq p \leq \infty$ ,  $c_0$  sum and Lebesgue Bochner spaces. Finally, we explore the stability of these with properties in the light of three space property. We show that *BHP* is a three space property provided  $X/Y$  is finite dimensional and same is true for *BSCSP* when  $X$  has *BSCSP* and  $X/Y$  is strongly regular ([GGMS]).

## 1. INTRODUCTION

Let  $X$  be a *real* nontrivial Banach space and  $X^*$  its dual. We will denote by  $B_X$ ,  $S_X$  and  $B_X(x, r)$  the closed unit ball, the unit sphere and the closed ball of radius  $r > 0$  and center  $x$ . We refer to the monograph [B1] for notions of convexity theory that we will be using here.

- Definition 1.**
- (i) We say  $A \subseteq B_{X^*}$  is a norming set for  $X$  if  $\|x\| = \sup\{|x^*(x)| : x^* \in A\}$ , for all  $x \in X$ . A closed subspace  $F \subseteq X^*$  is a norming subspace if  $B_F$  is a norming set for  $X$ .
  - (ii) Let  $x^* \in X^*$ ,  $\alpha > 0$  and bounded set  $C \subseteq X$ . Then the set  $S(C, x^*, \alpha) = \{x \in C : x^*(x) > \sup x^*(C) - \alpha\}$  is called the slice determined by  $x^*$  and  $\alpha$ . For any slice we assume without loss of generality that  $\|x^*\| = 1$ , because if  $x^* \neq 0$  then  $S(C, x^*, \alpha) = S(C, \frac{x^*}{\|x^*\|}, \frac{\alpha}{\|x^*\|})$ . One can analogously define  $w^*$  slices in  $X^*$  by choosing the functional from the predual.
  - (iii) A point  $x$  in a bounded set  $C \subseteq X$  is called a denting point point of  $C$ , if for every  $\varepsilon > 0$ , there exists a slice  $S$  of  $C$ , such that  $x \in S$  and  $dia(S) < \varepsilon$ . One can analogously define  $w^*$ -denting point of a bounded set in  $X^*$  by considering  $w^*$ -slices.

---

2020 *Mathematics Subject Classification.* 46B20, 46B28.

*Key words and phrases.* Slices, Huskable, Denting, Dentable, Small Combination of Slices.

(iv) A point  $x$  in a bounded convex set  $C \subseteq X$  is called a small combination of slices (SCS) point of  $C$ , if for every  $\varepsilon > 0$ , there exists a convex combination of slices,  $S = \sum_{i=1}^n \lambda_i S_i$  of  $C$  (where  $0 \leq \lambda_i \leq 1$  and  $\sum_{i=1}^n \lambda_i = 1$ ) such that  $x \in S$  and  $dia(S) < \varepsilon$ . One can analogously define  $w^*$ -SCS point of a bounded set in  $X^*$  by considering  $w^*$ -slices.

We recall the following two definitions from [BR] and [B2].

**Definition 1.1.** A Banach space  $X$  has

- (i) *Ball Dentable Property (BDP)* if  $B_X$  has a slice of arbitrarily small diameter.
- (ii) *Ball Small Combination of Slice Property (BSCSP)* if  $B_X$  has a convex combination of slices of arbitrarily small diameter.

We now define,

**Definition 1.2.** A Banach space  $X$  has *Ball Huskable Property (BHP)* if  $B_X$  has a relatively weakly open subset of arbitrarily small diameter.

**Remark 1.3.** Analogously we can define  $w^*$ -BSCSP,  $w^*$ -BHP and  $w^*$ -BDP in a dual space by considering  $w^*$ -SCS,  $w^*$  open sets and  $w^*$ -slices of  $B_{X^*}$  respectively.

Observe that for a Banach space, *BDP* always implies *BHP*, in fact, any slice of the unit ball is relatively weakly open. Also *BHP* implies *BSCSP*, by Bourgain's Lemma (see[B3], [GGMS]), which says that every non-empty relatively weakly open subset of  $B_X$  contains a finite convex combination of slices. Similar observations are true for  $w^*$ -versions. Since every  $w^*$ -slice ( $w^*$ -open set) of  $B_{X^*}$  is also a slice (weakly open set) of  $B_{X^*}$ , so we have the following diagram :

$$\begin{array}{ccccc}
 BDP & \implies & BHP & \implies & BSCSP \\
 \Uparrow & & \Uparrow & & \Uparrow \\
 w^*BDP & \implies & w^*BHP & \implies & w^*BSCSP
 \end{array}$$

In general, none of the reverse implications of the diagram hold, which we will discuss later.

SCS points were first introduced in [GGMS] as a "slice generalization" of denting points as well as the notion PC (i.e. points for which the identity mapping on the unit ball, from weak topology to norm topology is continuous) and subsequently analyzed in detail in [R] and [S]. It is well known that  $X$  has Radon Nikodym Property (*RNP*) if and only if every closed bounded and convex subset of  $X$  has slices with arbitrarily small diameter.  $X$  has the Point of Continuity Property (*PCP*) if every closed bounded and convex subset of  $X$  has relatively weakly open subsets with arbitrarily small diameter.  $X$  is said to be Strongly Regular (*SR*) if every closed, convex and bounded subset of  $X$  has convex combination of slices with arbitrarily small diameter. For more details, see [B3], [GGMS] and

[GMS]. It is clear then that  $RNP$  implies  $PCP$  and  $PCP$  implies  $SR$ . It is also well known that none of these implications can be reversed. Clearly,  $RNP$  implies  $BDP$ ,  $PCP$  implies  $BHP$  and  $SR$  implies  $BSCSP$ . The diagram below gives a clear picture.

$$\begin{array}{ccccc}
 RNP & \implies & PCP & \implies & SR \\
 \Downarrow & & \Downarrow & & \Downarrow \\
 BDP & \implies & BHP & \implies & BSCSP
 \end{array}$$

It was proved in [GGMS] that  $X$  is strongly regular if and only if every nonempty bounded convex set  $K$  in  $X$  is contained in norm closure of  $SCS(K)$  i.e.  $SCS$  points of  $K$ . Later it was proved in [S] that a Banach space has  $RNP$  if and only if it is  $SR$  and has the Krein Milman Property (KMP), i.e. every closed bounded convex subset  $K$  of  $X$  is the norm-closed convex hull of its extreme points. All the three properties discussed in this paper in a way, are "localised"(to the closed unit ball) versions of the three geometric properties  $RNP$ ,  $PCP$  and  $SR$ .

In this work we introduce Ball Huskable Property ( $BHP$ ) and explore its relation with  $BSCSP$  and  $BDP$ . We observe that  $BDP$  implies  $BHP$  which in turn implies  $BSCSP$  and none of the implications can be reversed. We prove certain stability results for  $BDP$ ,  $BHP$  and  $BSCSP$ . We further explore these properties in the context of three space property. The spaces that we will be considering have been well studied in literature. A large class of function spaces like the Bloch spaces, Lorentz and Orlicz spaces, spaces of vector valued functions and spaces of compact operators are examples of the spaces we will be considering, for details, see [HWW]

## 2. STABILITY RESULTS

The following result will be useful in our discussion.

**Proposition 2.1.** A Banach space  $X$  has  $BDP$  (resp.  $BHP$ ,  $BSCSP$ ) if and only if  $X^{**}$  has  $w^*$ - $BDP$  (resp.  $w^*$ - $BHP$ ,  $w^*$ - $BSCSP$ ).

*Proof.* Suppose  $X$  has  $BDP$ . Let  $\varepsilon > 0$ . Then there exists a slice  $S(B_X, x^*, \alpha)$  of  $B_X$  with diameter less than  $\frac{\varepsilon}{2}$ .

Claim:  $S(B_X, x^*, \alpha)$  is  $w^*$  dense in the  $w^*$ -slice  $S(B_{X^{**}}, x^*, \alpha)$  of  $B_{X^{**}}$ .

Indeed, fix  $x^{**} \in S(B_{X^{**}}, x^*, \alpha)$ . By Goldstine's Theorem, there is a net  $(x_\beta)$  in  $B_X$  which converges to  $x^{**}$  in the  $w^*$ -topology. Since,

$$\lim_{\beta} x^*(x_\beta) = x^{**}(x^*) > 1 - \alpha$$

So, there exists  $\beta_0$  such that  $(x_\beta) \in S(B_X, x^*, \alpha)$  for all  $\beta \geq \beta_0$ . Hence the claim.

Now let  $x^{**}, \tilde{x}^{**} \in S(B_{X^{**}}, x^*, \alpha)$ . Then there exist net  $(x_\beta)$  and  $(\tilde{x}_\beta)$  in  $S(B_X, x^*, \alpha)$  such that  $(x_\beta - \tilde{x}_\beta)$  converges to  $x^{**} - \tilde{x}^{**}$  in the  $w^*$ -topology. So,

$$\|x^{**} - \tilde{x}^{**}\| \leq \liminf_{\beta} \|x_\beta - \tilde{x}_\beta\| \leq \frac{\varepsilon}{2} < \varepsilon.$$

Thus  $\text{dia } S(B_{X^{**}}, x^*, \alpha) < \varepsilon$ . Hence  $X^{**}$  has  $w^*$ -BDP.

Conversely, if  $X^{**}$  has  $w^*$ -BDP, it immediately follows that  $X$  has BDP.

The proofs for BHP and BSCSP follow similarly.  $\square$

We immediately have,

**Corollary 2.2.** If  $X$  has BDP (resp. BHP, BSCSP) then  $X^{**}$  has BDP (resp. BHP, BSCSP)

The following lemma will be useful.

**Lemma 2.3.** Let  $Z = X \oplus_p Y$ ,  $1 \leq p < \infty$ , For any  $\varepsilon > 0$  and for any slice there exists a slice  $S(B_Z, z^*, \mu)$  of  $B_Z$  such that  $S(B_Z, z^*, \mu) \subset S(B_X, x^*, \alpha) \times \varepsilon B_Y$ . Similarly, there exists a slice  $S(B_Z, z^*, \gamma)$  of  $B_Z$  such that  $S(B_Z, z^*, \gamma) \subset \varepsilon B_X \times S(B_Y, y^*, \alpha)$ .

*Proof.* Let  $S(B_X, x^*, \alpha)$  be any slice of  $B_X$ . Also let  $\varepsilon > 0$ . Put  $z^* = (x^*, 0) \in S_{Z^*}$ . Choose  $0 < \mu < \alpha$  such that  $[1 - (1 - \mu)^p]^{1/p} < \varepsilon$ . Now, consider a slice of  $B_Z$  as,  $S(B_Z, z^*, \mu) = \{z \in B_Z : z^*(z) > 1 - \mu\} = \{z \in B_Z : x^*(x) > 1 - \mu\}$ . Then  $S(B_Z, z^*, \mu) \subset S(B_X, x^*, \alpha) \times \varepsilon B_Y$ . Indeed, let  $z \in S(B_Z, z^*, \mu)$ . Then,

$$1 \geq \|z\|^p = \|x\|^p + \|y\|^p > (1 - \mu)^p + \|y\|^p$$

Thus,  $\|y\|^p < 1 - (1 - \mu)^p$  and so  $\|y\| < [1 - (1 - \mu)^p]^{1/p} < \varepsilon$ . Also since,  $0 < \mu < \alpha$ , it follows that,  $x \in S(B_X, x^*, \alpha)$ . Hence,  $z = (x, y) \in S(B_X, x^*, \alpha) \times \varepsilon B_Y$ . The other proof is similar.  $\square$

**Proposition 2.4.** Let  $X$  and  $Y$  be two Banach spaces and  $Z = X \oplus_p Y$ ,  $1 \leq p < \infty$ . Then  $Z$  has BDP if and only if  $X$  or  $Y$  has BDP.

*Proof.* Suppose  $Z$  has BDP. We prove by contradiction. If possible, let  $X$  and  $Y$  do not have BDP. Then there exists  $\varepsilon > 0$  such that every slice of  $B_X$  and  $B_Y$  has diameter greater than  $\varepsilon$ . Since  $Z$  has BDP, there exists a slice  $S(B_Z, z^*, \alpha)$  of  $B_Z$  with diameter less than  $\varepsilon$ .

Case 1 :  $x^* = 0$  or  $y^* = 0$

Without loss of generality, let  $y^* = 0$ . Then  $x^* \in S_{X^*}$ . Then  $S(B_X, x^*, \alpha) \times \{0\} \subset S(B_Z, z^*, \alpha)$ . Thus,

$$\text{dia} S(B_X, x^*, \alpha) = \text{dia}(S(B_X, x^*, \alpha) \times \{0\}) \leq \text{dia} S(B_Z, z^*, \alpha) < \varepsilon,$$

a contradiction.

Case-2 :  $x^* \neq 0$  and  $y^* \neq 0$ . Choose  $z_0 = (x_0, y_0) \in S(B_Z, z^*, \frac{\alpha}{4})$  with  $\|z_0\| = 1$ . Now,  $S(B_X, \frac{x^*}{\|x^*\|}, \frac{\alpha}{2})$  and  $S(B_Y, \frac{y^*}{\|y^*\|}, \frac{\alpha}{2})$  are slices of  $B_X$  and  $B_Y$  respectively. Hence,  $diaS(B_X, \frac{x^*}{\|x^*\|}, \frac{\alpha}{2}) > \varepsilon$  and  $diaS(B_Y, \frac{y^*}{\|y^*\|}, \frac{\alpha}{2}) > \varepsilon$ . There exists  $x, \tilde{x} \in S(B_X, \frac{x^*}{\|x^*\|}, \frac{\alpha}{2})$  and  $y, \tilde{y} \in S(B_Y, \frac{y^*}{\|y^*\|}, \frac{\alpha}{2})$  such that  $\|x - \tilde{x}\| > \varepsilon$  and  $\|y - \tilde{y}\| > \varepsilon$ .

Let  $z = (\|x_0\|x, \|y_0\|y)$  and  $\tilde{z} = (\|x_0\|\tilde{x}, \|y_0\|\tilde{y})$ . Clearly,  $z, \tilde{z} \in S(B_Z, z^*, \alpha)$ . Also,  $\|z - \tilde{z}\|^p = \|x_0\|^p \|x - \tilde{x}\|^p + \|y_0\|^p \|y - \tilde{y}\|^p > \varepsilon^p (\|x_0\|^p + \|y_0\|^p) = \varepsilon^p$  which implies  $\|z - \tilde{z}\| > \varepsilon$ , a contradiction.

Hence either  $X$  or  $Y$  has  $BDP$ .

Conversely, assume that either  $X$  or  $Y$  has  $BDP$ . Without loss of generality let  $X$  have  $BDP$ . Let  $\varepsilon > 0$ . Then there exists a slice  $S(B_X, x^*, \alpha)$  of  $B_X$  with diameter  $< \varepsilon$ , where  $x^* \in S_{X^*}$  and  $\alpha > 0$ . From Lemma 2.3, there exists a slice  $S(B_Z, z^*, \mu)$  of  $B_Z$  such that  $S(B_Z, z^*, \mu) \subset S(B_X, x^*, \alpha) \times \varepsilon B_Y$ . Consequently,

$$dia(B_Z, z^*, \mu) \leq diaS(B_X, x^*, \alpha) + dia(\varepsilon B_Y) < \varepsilon + 2\varepsilon = 3\varepsilon$$

□

**Corollary 2.5.** Let  $X = \bigoplus_p X_i$ . If  $X_i$  has  $BDP$  for some  $i$ , then  $X$  has  $BDP$ .

We quote the following Lemma from [L1].

**Lemma 2.6.** [L1] Let  $Z = X \oplus_\infty Y$  then for every slice  $S(B_Z, z^*, \alpha)$  of  $B_Z$  there exists a slice  $S(B_X, x^*, \mu_1)$  of  $B_X$ , a slice  $S(B_Y, y^*, \mu_2)$  of  $B_Y$ ,  $x_0 \in B_X$  and  $y_0 \in B_Y$  such that  $S(B_X, x^*, \mu_1) \times \{y_0\} \subset S(B_Z, z^*, \alpha)$  and  $\{x_0\} \times S(B_Y, y^*, \mu_2) \subset S(B_Z, z^*, \alpha)$ .

**Proposition 2.7.**  $Z = X \oplus_\infty Y$  has  $BDP$  if and only if both  $X$  and  $Y$  have  $BDP$ .

*Proof.* First suppose that  $Z$  has  $BDP$ . Let  $0 < \varepsilon < 2$ . Then there exists a slice  $S(B_Z, z^*, \alpha)$  of  $B_Z$ , such that  $dia(S(B_Z, z^*, \alpha)) < \varepsilon$ , where  $z^* = (x^*, y^*) \in S_{Z^*}$  and  $\alpha > 0$ .

Claim :  $x^* \neq 0$  and  $y^* \neq 0$ .

If not, let  $x^* = 0$ . Then  $\|y^*\| = 1$ . Choose any fixed  $y_0 \in S(B_Y, y^*, \alpha)$ . Then  $B_X \times \{y_0\} \subset S(B_Z, z^*, \alpha)$ . So,  $2 = dia(B_X \times \{y_0\}) \leq diaS(B_Z, z^*, \alpha) < \varepsilon$ , a contradiction. Hence the claim. Now from Lemma 2.6, there exists a slice  $S(B_X, x^*, \mu)$  of  $B_X$  and  $y_1 \in B_Y$  such that  $S(B_X, x^*, \mu) \times \{y_1\} \subset S(B_Z, z^*, \alpha)$ . Consequently,  $diaS(B_X, x^*, \mu) = dia(S(B_X, x^*, \mu) \times \{y_1\}) \leq diaS(B_Z, z^*, \alpha) < \varepsilon$ . Thus,  $X$  has  $BDP$ . Similarly,  $Y$  has  $BDP$ .

Conversely let both  $X$  and  $Y$  have  $BDP$  and  $\varepsilon > 0$ . So, there exist slices  $S(B_X, x^*, \alpha_1)$  and  $S(B_Y, y^*, \alpha_2)$  of  $B_X$  and  $B_Y$  respectively such that  $diaS(B_X, x^*, \alpha_1) < \varepsilon$  and  $diaS(B_Y, y^*, \alpha_2) < \varepsilon$ . Choose  $0 < \gamma < \min\{\alpha_1, \alpha_2\}$ . Consider slice  $S(B_Z, z^*, \gamma)$  of  $B_Z$  such that  $z^* = (\frac{x^*}{2}, \frac{y^*}{2})$ . Then  $\|z^*\| = 1$ . Then  $S(B_Z, z^*, \gamma) \subset S(B_X, x^*, \alpha_1) \oplus_\infty S(B_Y, y^*, \alpha_2)$ . Indeed, let  $z = (x, y) \in S(B_Z, z^*, \gamma)$ . Then

$$z^*(z) = \frac{x^*}{2}(x) + \frac{y^*}{2}(y) > 1 - \gamma$$

$$\begin{aligned} \Rightarrow 1 + y^*(y) &\geq x^*(x) + y^*(y) > 2 - 2\gamma \\ \Rightarrow y^*(y) &> 1 - 2\gamma > 1 - \alpha_2 \end{aligned}$$

Thus  $y \in S(B_Y, y^*, \alpha_2)$  and similarly  $x \in S(B_X, x^*, \alpha_1)$ . Finally,  $\text{dia}S(B_Z, z^*, \gamma) \leq \varepsilon$  as both  $S(B_X, x^*, \alpha_1)$  and  $S(B_Y, y^*, \alpha_2)$  are of diameter  $< \varepsilon$ .  $\square$

**Proposition 2.8.** Let  $X$  and  $Y$  be two Banach spaces and  $Z = X \oplus_p Y$ ,  $1 \leq p < \infty$ . Then  $Z$  has *BHP* if and only if  $X$  or  $Y$  has *BHP*.

*Proof.* Suppose  $Z$  has *BHP*. If possible let  $X$  and  $Y$  fail *BHP*. Then there exists  $\varepsilon > 0$  such that every relatively weakly open subset of  $B_X$  and  $B_Y$  has diameter greater than  $\varepsilon$ . Now since  $Z$  has *BHP* so there exists a relatively weakly open subset  $W$  of  $B_Z$  with diameter less than  $\varepsilon$ . Fix  $z_0 = (x_0, y_0) \in W \cap S_Z$ . Then there exists a basic weakly open subset,  $W_0 = \{z \in B_Z : |z_i^*(z - z_0)| < 1, i = 1, 2, \dots, n\} \subset W$  where  $z_i^* = (x_i^*, y_i^*)$ ,  $i = 1, 2, \dots, n$ . We consider two cases.

Case 1 :  $x_0 = 0$  or  $y_0 = 0$

without loss of generality let  $y_0 = 0$ . Thus  $x_0 \in S_X$ . Then

$U = \{x \in B_X : |x_i^*(x - x_0)| < 1; i = 1, 2, \dots, n\}$  is nonempty relatively weakly open subset of  $B_X$ . By our assumption  $\text{dia}(U) > \varepsilon$ . So there exists  $x, \tilde{x} \in U$  such that  $\|x - \tilde{x}\| > \varepsilon$ . Now,  $z = (x, 0)$  and  $\tilde{z} = (\tilde{x}, 0)$  are in  $W_0$  and  $\|z - \tilde{z}\| = \|x - \tilde{x}\| > \varepsilon$ , a contradiction.

Case 2 :  $x_0 \neq 0$  and  $y_0 \neq 0$

Consider,  $U = \{x \in B_X : |x_i^*(x - \frac{x_0}{\|x_0\|})| < \frac{1}{2\|x_0\|}; i = 1, 2, \dots, n\}$

and  $V = \{y \in B_Y : |y_i^*(y - \frac{y_0}{\|y_0\|})| < \frac{1}{2\|y_0\|}; i = 1, 2, \dots, n\}$

Then  $U$  and  $V$  are nonempty relatively weakly open subsets of  $B_X$  and  $B_Y$  respectively and so  $\text{dia}(U) > \varepsilon$  and  $\text{dia}(V) > \varepsilon$ . Hence, there exists  $x, \tilde{x} \in U$  and  $y, \tilde{y} \in V$  such that  $\|x - \tilde{x}\| > \varepsilon$  and  $\|y - \tilde{y}\| > \varepsilon$ .

Thus  $z = (\|x_0\|x, \|y_0\|y)$  and  $\tilde{z} = (\|x_0\|\tilde{x}, \|y_0\|\tilde{y})$  are in  $W$ .

Indeed,  $\|z\|^p = \|x_0\|^p \|x\|^p + \|y_0\|^p \|y\|^p \leq \|x_0\|^p + \|y_0\|^p = 1$ , and

$\forall i = 1, 2, \dots, n$  we have,

$$\begin{aligned} |z_i^*(z - z_0)| &= |x_i^*(\|x_0\|x - x_0) + y_i^*(\|y_0\|y - y_0)| \leq \|x_0\| |x_i^*(x - \frac{x_0}{\|x_0\|})| + \|y_0\| |y_i^*(y - \frac{y_0}{\|y_0\|})| \\ &< \|x_0\| \frac{1}{2\|x_0\|} + \|y_0\| \frac{1}{2\|y_0\|} = 1 \end{aligned}$$

Similarly for  $\tilde{z}$ .

Finally,  $\|z - \tilde{z}\|^p = \|x_0\|^p \|x - \tilde{x}\|^p + \|y_0\|^p \|y - \tilde{y}\|^p > \varepsilon^p (\|x_0\|^p + \|y_0\|^p) = \varepsilon^p$

and so  $\|z - \tilde{z}\| > \varepsilon$ , a contradiction.

Hence either  $X$  or  $Y$  has *BHP*.

Conversely assume that either  $X$  or  $Y$  has *BHP*. Without loss of generality, suppose  $X$  has *BHP*. Let  $\varepsilon > 0$ . Then there exists a relatively weakly open set  $W$  of  $B_X$  with diameter  $< \varepsilon$ . Since slices of  $B_X$

forms a subbase for relatively weakly open subset of  $B_X$ , so there exists slices  $S(B_X, x_i^*, \alpha_i)$ ,  $i=1,2,\dots,n$  of  $B_X$  such that  $\bigcap_{i=1}^n S(B_X, x_i^*, \alpha_i) \subset W$ . Now from Lemma 2.3 for each  $i$ , we get a slice  $S(B_Z, z_i^*, \mu_i)$  of  $B_Z$  such that  $S(B_Z, z_i^*, \mu_i) \subset S(B_X, x_i^*, \alpha_i) \times \varepsilon B_Y$ . Choose  $\mu < \min\{\mu_1, \mu_2, \dots, \mu_n\}$ .

Thus,  $\bigcap_{i=1}^n S(B_Z, z_i^*, \mu) \subset \bigcap_{i=1}^n S(B_X, x_i^*, \alpha_i) \times \varepsilon B_Y \subset W \times \varepsilon B_Y$ .

Hence  $T = \bigcap_{i=1}^n S(B_Z, z_i^*, \mu)$  is a relatively weakly open subset of  $B_Z$  with diameter less than  $3\varepsilon$ . Consequently  $Z$  has *BHP*. □

**Corollary 2.9.** Let  $X = \bigoplus_p X_i$ . If  $X_i$  has *BHP* for some  $i$ , then  $X$  has *BHP*.

The following Lemma from [ALN] will be useful.

**Lemma 2.10.** [ALN] Let  $X$  and  $Y$  be Banach spaces and  $W$  be a nonempty relatively weakly open subset in unit ball of  $Z = X \oplus_\infty Y$ . Then  $U$  and  $V$  can be chosen to be relatively weakly open subsets of  $B_X$  and  $B_Y$  respectively such that  $U \times V \subset W$ .

**Proposition 2.11.**  $Z = X \oplus_\infty Y$  has *BHP* if and only if both  $X$  and  $Y$  have *BHP*.

*Proof.* First suppose that  $Z$  has *BHP*. Also let  $\varepsilon > 0$ . Then  $B_Z$  has a relatively weakly open subset  $W_0$  with diameter less than  $< \varepsilon$ . Then by Lemma 2.10, there exists a relatively weakly open subset  $U_0$  in  $B_X$  and  $V_0$  in  $B_Y$  such that  $U_0 \times V_0 \subset W_0$ . Fix  $u_0 \in U_0$  and  $v_0 \in V_0$ . Then,  $\text{dia}(U_0) = \text{dia}(U_0 \times \{v_0\}) \leq \text{dia}(U_0 \times V_0) \leq \text{dia} W_0 < \varepsilon$ .

Similarly for  $V_0$ . Consequently, both  $X$  and  $Y$  have *BHP*.

Conversely suppose  $X$  and  $Y$  have *BHP* and  $\varepsilon > 0$ . So, there exists relatively weakly open subset  $U$  and  $V$  of  $B_X$  and  $B_Y$  respectively such that  $\text{dia}(U) < \varepsilon$  and  $\text{dia}(V) < \varepsilon$ . Since slices of  $B_X$  forms a subbase for relatively weakly open subset of  $B_X$ , so there exists slices  $S(B_X, x_i^*, \alpha_i)$ ,  $i=1,2,\dots,n$  of  $B_X$  such that  $\bigcap_{i=1}^n S(B_X, x_i^*, \alpha_i) \subset U$  and similarly there exists slices  $S(B_Y, y_j^*, \beta_j)$ ,  $j=1,2,\dots,m$  of  $B_Y$  such that  $\bigcap_{j=1}^m S(B_Y, y_j^*, \beta_j) \subset V$ . Without loss of generality let  $n \geq m$ . Then proceeding same way as in Proposition 2.7 we get slices of  $B_Z$ ,  $S(B_Z, z_i^*, \gamma_i) \quad \forall i = 1, \dots, n$

where  $\gamma_i < \min\{\alpha_i, \beta_i\}$  if  $i = 1, \dots, m$  and  $\gamma_i < \min\{\alpha_i, \beta_m\}$  if  $i = m+1, \dots, n$  such that

$$S(B_Z, z_i^*, \gamma_i) \subset S(B_X, x_i^*, \alpha_i) \oplus_\infty S(B_Y, y_i^*, \beta_i) \quad \forall i = 1, \dots, m$$

and

$$S(B_Z, z_i^*, \gamma_i) \subset S(B_X, x_i^*, \alpha_i) \oplus_\infty S(B_Y, y_i^*, \beta_i) \quad \forall i = m+1, \dots, n$$

Thus,  $\bigcap_{i=1}^n S(B_Z, z_i^*, \gamma_i) \subset \bigcap_{i=1}^n S(B_X, x_i^*, \alpha_i) \oplus_\infty \bigcap_{i=1}^m S(B_Y, y_i^*, \beta_i) \subset U \oplus_\infty V$ .

Hence,  $\text{dia}(\bigcap_{i=1}^n S(B_Z, z_i^*, \gamma_i)) < \varepsilon$  □

**Proposition 2.12.** Let  $X$  and  $Y$  be two Banach spaces and  $Z = X \oplus_p Y$ ,  $1 \leq p < \infty$ .  $Z$  has *BSCSP* if  $X$  or  $Y$  has *BSCSP*.

*Proof.* Without loss of generality, let  $X$  have *BSCSP*. Let  $\varepsilon > 0$ . Then there exists a convex combination of slices  $\sum_{i=1}^n \lambda_i S(B_X, x_i^*, \alpha_i)$ ,  $\lambda_i \geq 0$ ,  $\sum_{i=1}^n \lambda_i = 1$  of  $B_X$  with diameter less than  $\varepsilon$ . By Lemma 2.3, for each  $i$ , there exists a slice  $S(B_Z, z_i^*, \mu_i)$  of  $B_Z$  such that  $S(B_Z, z_i^*, \mu_i) \subset S(B_X, x_i^*, \alpha_i) \times \varepsilon B_Y$ . Thus,  $\sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \mu_i) \subset \sum_{i=1}^n \lambda_i [S(B_X, x_i^*, \alpha_i) \times \varepsilon B_Y]$ . Hence,  $\text{dia}(\sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \mu_i)) < 3\varepsilon$ . Consequently  $Z$  has *BSCSP*. □

**Corollary 2.13.** Let  $X = \oplus_p X_i$ . If  $X_i$  has *BSCSP* for some  $i$ , then  $X$  has *BSCSP*.

**Proposition 2.14.** If  $Z = X \oplus_1 Y$  has *BSCSP*, then either  $X$  or  $Y$  has *BSCSP*.

*Proof.* If possible, let  $X$  and  $Y$  fail *BSCSP*. Then there exists  $\varepsilon > 0$  such that every convex combination of slices of  $B_X$  and  $B_Y$  have diameter greater than  $\varepsilon$ . Now since  $Z$  has *BSCSP*, so there exists convex combination of slices  $\sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \alpha_i)$  of  $B_Z$  with diameter less than  $\varepsilon$ . Observe,  $1 = \|z_i^*\| = \max\{\|x_i^*\|, \|y_i^*\|\}$ ,  $i = 1, 2, \dots, n$ . We consider two disjoint subsets  $I$  and  $J$  where,  $I = \{i : \|x_i^*\| = 1\}$  and  $J = \{j : \|y_j^*\| = 1\}$ . Now,  $S(B_X, x_i^*, \alpha_i) \times \{0\} \subset S(B_Z, z_i^*, \alpha_i) \quad \forall i \in I$  and  $\{0\} \times S(B_Y, y_j^*, \alpha_j) \subset S(B_Z, z_j^*, \alpha_j) \quad \forall j \in J$ .

Let  $\lambda_I = \sum_{i \in I} \lambda_i$  and  $\lambda_J = \sum_{j \in J} \lambda_j$

Case-1 :  $\lambda_I = 0$  or  $\lambda_J = 0$

Without loss of generality, let  $\lambda_I = 0$

Then  $\lambda_J = 1$  and so  $\sum_{j \in J} \lambda_j S(B_Y, y_j^*, \alpha_j)$  is a convex combination of slices of  $B_Y$ , hence  $\text{dia}(\sum_{j \in J} \lambda_j S(B_Y, y_j^*, \alpha_j)) > \varepsilon$ . So, there exist  $y, \tilde{y} \in \sum_{j \in J} \lambda_j S(B_Y, y_j^*, \alpha_j)$  such that  $\|y - \tilde{y}\| > \varepsilon$ . Hence,  $\|(0, y) - (0, \tilde{y})\| > \varepsilon$ , which is a contradiction.

Case-2 :  $\lambda_I \neq 0$  or  $\lambda_J \neq 0$

So we have,  $\sum_{i \in I} \frac{\lambda_i}{\lambda_I} S(B_X, x_i^*, \alpha_i) \times \{0\} \subset \sum_{i \in I} \frac{\lambda_i}{\lambda_I} S(B_Z, z_i^*, \alpha_i)$  and  $\{0\} \times \sum_{j \in J} \frac{\lambda_j}{\lambda_J} S(B_Y, y_j^*, \alpha_j) \subset \sum_{j \in J} \frac{\lambda_j}{\lambda_J} S(B_Z, z_j^*, \alpha_j)$ . Again,  $\text{dia}(\sum_{i \in I} \frac{\lambda_i}{\lambda_I} S(B_X, x_i^*, \alpha_i)) > \varepsilon$  and  $\text{dia}(\sum_{j \in J} \frac{\lambda_j}{\lambda_J} S(B_Y, y_j^*, \alpha_j)) > \varepsilon$ . So, there exist  $x, \tilde{x} \in \sum_{i \in I} \frac{\lambda_i}{\lambda_I} S(B_X, x_i^*, \alpha_i)$  and  $y, \tilde{y} \in \sum_{j \in J} \frac{\lambda_j}{\lambda_J} S(B_Y, y_j^*, \alpha_j)$  such that  $\|x - \tilde{x}\| > \varepsilon$  and  $\|y - \tilde{y}\| > \varepsilon$

Observe,

$$\begin{aligned} (\lambda_I x, \lambda_J y) &= (\lambda_I x, 0) + (0, \lambda_J y) \in \sum_{i \in I} \lambda_i S(B_X, x_i^*, \alpha_i) \times \{0\} + \{0\} \times \sum_{j \in J} \lambda_j S(B_Y, y_j^*, \alpha_j) \\ &\subset \sum_{i \in I} \lambda_i S(B_Z, z_i^*, \alpha_i) + \sum_{j \in J} \lambda_j S(B_Z, z_j^*, \alpha_j) \\ &= \sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \alpha_i) \end{aligned}$$

Similarly,  $(\lambda_I \tilde{x}, \lambda_J \tilde{y}) \in \sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \alpha_i)$

Also,  $\|(\lambda_I x, \lambda_J y) - (\lambda_I \tilde{x}, \lambda_J \tilde{y})\| = \|(\lambda_I(x - \tilde{x}), \lambda_J(y - \tilde{y}))\| = \lambda_I \|x - \tilde{x}\| + \lambda_J \|y - \tilde{y}\| > \varepsilon(\lambda_I + \lambda_J) = \varepsilon$ , a contradiction. Hence, either  $X$  or  $Y$  has *BSCSP*.



□

**Proposition 2.15.** If  $Z = X \oplus_{\infty} Y$  has *BSCSP*, then both  $X$  and  $Y$  have *BSCSP*.

*Proof.* Suppose  $Z$  has *BSCSP*. So, for any  $\varepsilon > 0$ , there exists a convex combination of slices  $\sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \alpha_i), \lambda_i \geq 0, \sum_{i=1}^n \lambda_i = 1$  of  $B_Z$  such that  $dia(\sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \alpha_i)) < \varepsilon$ . By Lemma 2.6, for each  $i$ , there exists a slice  $S(B_X, x_i^*, \mu_i)$  of  $B_X$  and  $y_i \in B_Y$  such that  $S(B_X, x_i^*, \mu_i) \times \{y_i\} \subset S(B_Z, z_i^*, \alpha_i)$ . Let  $y_0 = \sum_{i=1}^n \lambda_i y_i$ .  $[\sum_{i=1}^n \lambda_i S(B_X, x_i^*, \mu_i)] \times \{y_0\} = \sum_{i=1}^n \lambda_i [S(B_X, x_i^*, \mu_i) \times \{y_i\}] \subset \sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \alpha_i)$ . Hence,  $dia(\sum_{i=1}^n \lambda_i S(B_X, x_i^*, \mu_i)) = dia([\sum_{i=1}^n \lambda_i S(B_X, x_i^*, \mu_i)] \times \{y_0\}) \leq dia(\sum_{i=1}^n \lambda_i S(B_Z, z_i^*, \alpha_i)) \leq \varepsilon$ . So,  $X$  has *BSCSP*. Similarly for  $Y$ . □

Similar results are true for  $w^*$ -versions. We omit the proofs which are similar.

**Proposition 2.16.** Let  $X$  and  $Y$  be Banach spaces and  $Z = X \oplus_p Y, 1 < p \leq \infty$ . Then

- (i)  $Z^*$  has  $w^*$ *BHP* ( $w^*$ *BDP*) if and only if  $X^*$  or  $Y^*$  has  $w^*$ *BHP* ( $w^*$ *BDP*).
- (ii)  $X^*$  or  $Y^*$  has  $w^*$ *BSCSP* implies  $Z^*$  has  $w^*$ *BSCSP*.

**Proposition 2.17.** Let  $X$  and  $Y$  be Banach spaces and  $Z = X \oplus_1 Y$ , Then

$Z^*$  has  $w^*$ *BHP* ( resp.  $w^*$ *BDP*,  $w^*$ *BSCSP* ) implies both  $X^*$  and  $Y^*$  has  $w^*$ *BHP* ( resp.  $w^*$ *BDP*,  $w^*$ *BSCSP*).

**Proposition 2.18.** Let  $X = \oplus_{c_0(\Gamma)} X_i$ . If  $X$  has *BDP* (resp. *BHP*, *BSCSP*), then each of  $X_i$  has *BDP* (resp. *BHP*, *BSCSP*).

*Proof.* Fix  $i$ . Note that  $X = \oplus_{c_0(\Gamma)} X_j = X_i \oplus_{\infty} (\oplus_{c_0(\Gamma \setminus \{i\})} X_j)$ . The rest follows from Propositions 2.7, 2.11 and 2.15. □

**Remark 2.19.** The converse of the above Proposition may not be true. Indeed,  $\mathbb{R}$  has *BDP* (resp. *BHP*, *BSCSP*) but  $c_0$  does not have *BDP* (resp. *BHP*, *BSCSP*).

We now show that none of the implications in the following diagram can be reversed.

$$\begin{array}{ccccc}
 BDP & \implies & BHP & \implies & BSCSP \\
 \uparrow & & \uparrow & & \uparrow \\
 w^*BDP & \implies & w^*BHP & \implies & w^*BSCSP
 \end{array}$$

**Example 2.20.** (i)  $BHP \not\Rightarrow BDP$ .

It was proved in [BGLPRZ1] that, if a Banach space  $X$  contains an isomorphic copy of  $c_0$  then it can be equivalently renormed, so that every slice of unit ball of  $X$  has diameter two but it has a relatively weakly open subset of arbitrarily small diameter. Hence a Banach space

containing an isomorphic copy of  $c_0$  can be equivalently renormed so that it has *BHP* but not *BDP*.

(ii) *BSCSP*  $\not\Rightarrow$  *BHP*.

It was proved in [BGLPRZ2] that , if a Banach space  $X$  contains an isomorphic copy of  $c_0$  then it can be equivalently renormed, so that every relatively weakly open subset of unit ball of  $X$  has diameter two but it has convex combination of slices of arbitrarily small diameter. Hence a Banach space containing an isomorphic copy of  $c_0$  can be equivalently renormed so that it has *BSCSP* but not *BHP*.

(iii)  $w^*$ *BHP*  $\not\Rightarrow$   $w^*$ *BDP*.

If we consider bidual of the space in (i), then from Proposition 2.1, we get a space with  $w^*$ *BHP* but not  $w^*$ *BDP* .

(iv)  $w^*$ *BSCSP*  $\not\Rightarrow$   $w^*$ *BHP*.

If we consider bidual of the space in (ii), then from Proposition 2.1, we get a space with  $w^*$ *BSCSP* but not  $w^*$ *BHP*.

(v) *BDP*  $\not\Rightarrow$   $w^*$ *BDP*, *BHP*  $\not\Rightarrow$   $w^*$ *BHP*, *BSCSP*  $\not\Rightarrow$   $w^*$ *BSCSP*

Let  $X = C[0, 1]$ . It is well known that  $X^* = L_1[0, 1] \oplus_1 Z$ , for some subspace  $Z$  of  $X^*$  with *RNP* and hence  $Z$  has *BDP*. Since  $Z$  has *BDP*, it follows from Proposition 2.4,  $X^*$  has *BDP* and hence it has *BHP* and *BSCSP*. But every convex combination of  $w^*$  slices of  $B_{X^*}$  has diameter two ( see [BGLPRZ3] for details). Hence,  $C[0, 1]^*$  does not have  $w^*$ -*BSCSP* and consequently it does not have  $w^*$ -*BDP* ,  $w^*$ -*BHP*.

**Remark 2.21.** In recent times,the study of diameter two properties has been a very active area of research in the geometry of Banach spaces( see [BGLPRZ1],[BGLPRZ2],[BGLPRZ3],[ALN],[L1]). As is evident from the examples above that a Banach space may have one version of small diameter property and simultaneously may have a version of diameter two property. More investigations about the geometric implications of these observations will be an interesting topic of research in future.

### 3. LEBESGUE BOCHNER FUNCTION SPACES

Let  $(\Omega, \mathcal{A}, \mu)$  be a finite measure space. Then for  $1 \leq p < \infty$ ,  $L^p(\mu, X)$  denote the Banach space of all Lebesgue-Bochner function of  $p$ -integrable  $X$  valued functions defined on  $\Omega$  with norm  $\|f\|_p = \left( \int_{\Omega} \|f(t)\|^p d\mu(t) \right)^{1/p}$  ( see [DU] for details).

**Proposition 3.1.** Let  $(\Omega, \mathcal{A}, \mu)$  be a finite measure space.Let  $1 \leq p < \infty$ . Then  $L^p(\mu, X)$  has *BHP* implies  $X$  has *BHP*.

*Proof.* Suppose  $X$  does not have *BHP*. Then there exists  $\varepsilon > 0$  such that every relatively weakly open subset of  $B_X$  has diameter  $> \varepsilon$ . Since  $L^p(\mu, X)$  has *BHP*, there exists  $V$ , a weakly open subset of  $L^p(\mu, X)$  such that  $V \cap B_{L^p(\mu, X)} \neq \emptyset$  and  $\text{dia}(V \cap B_{L^p(\mu, X)}) < \varepsilon$ . If  $X$  is finite dimensional then it has *RNP* and so it has *BHP*. So we consider the case when  $X$  is infinite dimensional. Then  $L^p(\mu, X)$  is also infinite dimensional and so  $V \cap S_{L^p(\mu, X)} \neq \emptyset$ . Choose  $f_0 \in V \cap S_{L^p(\mu, X)}$ . Since simple functions are dense in  $L^p(\mu, X)$  so let  $f_0 = \sum_{i=1}^n x_i \chi_{A_i}$  where  $A_i (i = 1, 2, \dots, n)$  are disjoint sets in  $\mathcal{A}$  and  $x_i (\neq 0) \in X \forall i = 1, 2, \dots, n$ . Since weak topology on  $L^p(\mu, X)$  is linear so  $(x, y) \mapsto x + y$  from  $L^p(\mu, X) \times L^p(\mu, X) \rightarrow L^p(\mu, X)$  is weak continuous. Thus there exist weakly open subsets  $V_i$  containing  $x_i \chi_{A_i}$  in  $L^p(\mu, X)$  such that

$$(3.1) \quad \sum_{i=1}^n V_i \subset V$$

Since the map  $x \mapsto x \chi_{A_i}$  from  $X$  to  $L^p(\mu, X)$  is linear and norm continuous so also weak continuous for each  $i$ . Thus for each  $i$ , there exists weakly open subset  $W_i \subset X$  containing  $x_i$  such that

$$(3.2) \quad x \chi_{A_i} \in V_i \quad \forall x \in W_i$$

Note that  $\frac{W_i}{\|x_i\|}$  is a weakly open set in  $X$  and  $\frac{W_i}{\|x_i\|} \cap B_X \neq \emptyset$  as  $\frac{x_i}{\|x_i\|} \in \frac{W_i}{\|x_i\|} \cap B_X$ . Then  $\text{diam}(\frac{W_i}{\|x_i\|} \cap B_X) > \varepsilon$  implies  $\text{diam}(W_i \cap \|x_i\| B_X) > \varepsilon \|x_i\|$ . Then for each  $i$ , there exist  $y_i, z_i \in W_i \cap \|x_i\| B_X$  such that  $\|y_i - z_i\| > \varepsilon \|x_i\|$ . Consider  $g = \sum_{i=1}^n y_i \chi_{A_i}$  and  $h = \sum_{i=1}^n z_i \chi_{A_i}$ . From 3.1 and 3.2 we get  $g, h \in V$ . Also  $g, h \in B_{L^p(\mu, X)}$ . Indeed,  $\|g\|_p^p = \sum_{i=1}^n \|y_i\|^p \mu(A_i) \leq \sum_{i=1}^n \|x_i\|^p \mu(A_i) = 1$ .

We argue similarly for  $h$ . Finally,

$\text{dia}(V \cap B_{L^p(\mu, X)}) \geq \|g - h\|_p = \left( \sum_{i=1}^n \|y_i - z_i\|^p \mu(A_i) \right)^{1/p} > \left( \sum_{i=1}^n \varepsilon^p \|x_i\|^p \mu(A_i) \right)^{1/p} = \varepsilon$ , which is a contradiction.  $\square$

**Proposition 3.2.** Let  $(\Omega, \mathcal{A}, \mu)$  be a finite measure space. Let  $1 \leq p < \infty$ . Then  $L^p(\mu, X)$  has *BDP* implies  $X$  has *BDP*.

*Proof.* Suppose  $X$  does not have *BDP*. Then there exists  $\varepsilon > 0$  such that any slice of  $B_X$  has diameter  $> \varepsilon$ . Since  $L^p(\mu, X)$  has *BDP* so there exists a slice  $S(B_{L^p(\mu, X)}, f^*, \alpha)$  of  $B_{L^p(\mu, X)}$  with diameter  $< \varepsilon$ . Choose  $f \in S(B_{L^p(\mu, X)}, f^*, \alpha)$  with  $\|f\|_p = 1$ . Since simple functions are dense in  $L^p(\mu, X)$  space, so without loss of generality we can assume that  $f = \sum_{i=1}^n x_i \chi_{A_i}$  where  $A_i (i = 1, 2, \dots, n)$  are disjoint sets in  $\mathcal{A}$  and  $x_i (\neq 0) \in X \forall i = 1, 2, \dots, n$ . For each  $i = 1, \dots, n$  we define  $x_i^*(x) = f^*(x \chi_{A_i})$ . Then clearly  $x_i^*$  is linear and bounded. Now for each  $i$ , consider the slice  $S(\|x_i\| B_X, x_i^*, \|x_i^*\| \|x_i\| - x_i^*(x_i) + \alpha_i) = \{y \in \|x_i\| B_X : x_i^*(y) > x_i^*(x_i) - \alpha_i\}$  where  $\alpha_i > 0$  are such that  $\sum_{i=1}^n \alpha_i \leq f^*(f) - (1 - \alpha)$ . Since any slice of  $B_X$  has diameter  $> \varepsilon$  so  $\text{dia}(S(\|x_i\| B_X, x_i^*, \|x_i^*\| \|x_i\| - x_i^*(x_i) + \alpha_i)) > \|x_i\| \varepsilon$ . Thus for each  $i$ , there exist  $y_i, z_i \in S(\|x_i\| B_X, x_i^*, \|x_i^*\| \|x_i\| - x_i^*(x_i) + \alpha_i)$  such that  $\|y_i - z_i\| > \|x_i\| \varepsilon$ . Define

$$g = \sum_{i=1}^n y_i \chi_{A_i}, h = \sum_{i=1}^n z_i \chi_{A_i}.$$

$$\|g\|_p^p = \sum_{i=1}^n \|y_i\|^p \mu(A_i) \leq \sum_{i=1}^n \|x_i\|^p \mu(A_i) = 1$$

Similarly for  $h$ . Thus  $g, h \in B_{L^p(\mu, X)}$ . Also

$$\begin{aligned} f^*(g) &= f^*\left(\sum_{i=1}^n y_i \chi_{A_i}\right) = \sum_{i=1}^n f^*(y_i \chi_{A_i}) = \sum_{i=1}^n x_i^*(y_i) \\ &> \sum_{i=1}^n (x_i^*(x_i) - \alpha_i) \\ &= \sum_{i=1}^n x_i^*(x_i) - \sum_{i=1}^n \alpha_i \\ &= \sum_{i=1}^n f^*(x_i \chi_{A_i}) - \sum_{i=1}^n \alpha_i \\ &= f^*(f) - \sum_{i=1}^n \alpha_i \geq 1 - \alpha \end{aligned}$$

Thus  $g \in S(B_{L^p(\mu, X)}, f^*, \alpha)$ . Similarly  $h \in S(B_{L^p(\mu, X)}, f^*, \alpha)$ .

Hence

$$\text{dia}(S(B_{L^p(\mu, X)}, f^*, \alpha)) \geq \|g - h\|_p = \left( \sum_{i=1}^n \|y_i - z_i\|^p \mu(A_i) \right)^{1/p} > \left( \sum_{i=1}^n \varepsilon^p \|x_i\|^p \mu(A_i) \right)^{1/p} = \varepsilon$$

which is a contradiction.  $\square$

**Remark 3.3.** We do not know whether the converse of the Propositions 3.1, 3.2 are true or not in general. However, converse of Proposition 3.1, fails for  $p = 1$ . Indeed,  $\mathbb{R}$  has *BHP* but  $L^1(\mu, \mathbb{R})$  where  $\mu$  is Lebesgue measure on  $[0, 1]$  (which is simply denoted by  $L^1[0, 1]$ ) does not have *BHP*.

#### 4. THREE SPACE PROPERTY

We quote the following result from [GGMS]

**Proposition 4.1.** [GGMS]  $X$  is strongly regular if and only if every closed convex bounded subset  $D$  of  $X$  is the norm closure of its SCS points.

**Proposition 4.2.** If  $X/Y$  is strongly regular and  $Y$  has *BSCSP*, then  $X$  has *BSCSP*.

*Proof.* Suppose  $X$  does not have *BSCSP*. Then there exists  $\varepsilon > 0$  such that diameter of any convex combination of slices of  $B_X$  is greater than  $\varepsilon$ . Since  $Y$  has *BSCSP*, for  $\varepsilon > 0$ , there exists slices  $S(B_Y, y_i^*, \delta)$  and  $0 \leq \lambda_i \leq \sum_{i=1}^n \lambda_i = 1$  such that  $\text{dia}(\sum_{i=1}^n \lambda_i S(B_Y, y_i^*, \delta)) < \frac{\varepsilon}{2}$ . Choose  $\tilde{\varepsilon}, \delta_0 > 0$  such that  $\tilde{\varepsilon} + 2\delta_0 < \delta$ , and  $0 < \delta_0 < \frac{\varepsilon}{8}$ . By Hahn Banach theorem, we can extend  $y_i^*$  to a norm preserving extension  $x_i^*$  for all  $i = 1, 2, \dots, n$ . Let  $P : X \rightarrow X/Y$  be the map such that  $P(x) = x + Y$ . Now,  $A_i = P(S(B_X, x_i^*, \tilde{\varepsilon}))$  is a convex subset of  $B_{X/Y}$ , since  $\|P\| \leq 1$  and  $A_i$  also contains zero. Also, by Proposition 4.1,  $\bar{A}_i = \overline{(\text{SCS}(\bar{A}_i))}$ . Thus  $\forall i = 1, 2, \dots, n$ , there exists SCS

point  $a_i$  of  $\bar{A}_i$  such that  $\|a_i\| < \delta_0$ . Hence  $a_i = \sum_{j=1}^{n_i} \gamma_j^i a_j^i \in \sum_{j=1}^{n_i} \gamma_j^i (S(B_{X/Y}, (a_j^i)^*, \eta_j^i) \cap \bar{A}_i)$  with  $\text{dia}(\sum_{j=1}^{n_i} \gamma_j^i (S(B_{X/Y}, (a_j^i)^*, \eta_j^i) \cap \bar{A}_i)) < \delta_0$  for all  $i = 1, 2, \dots, n$  where  $\sum_{j=1}^{n_i} \gamma_j^i = 1$  with  $\gamma_j^i > 0 \forall j = 1, \dots, n_i$ ,  $(a_j^i)^* \in S_{(X/Y)^*}$  and  $\eta_j^i > 0$  for all  $i = 1, 2, \dots, n$ . Now,  $S(B_{X/Y}, (a_j^i)^*, \eta_j^i) \cap \bar{A}_i \neq \emptyset$  implies  $S(B_{X/Y}, (a_j^i)^*, \eta_j^i) \cap A_i \neq \emptyset$ . Consequently,  $S(B_X, P^*(a_j^i)^*, \eta_j^i) \cap S(B_X, x_i^*, \tilde{\varepsilon}) \neq \emptyset$ . Indeed, let  $P(z) \in S(B_{X/Y}, (a_j^i)^*, \eta_j^i)$  for some  $z \in S(B_X, x_i^*, \tilde{\varepsilon})$ . hence  $P^*(a_j^i)^*(z) = (a_j^i)^*(P(z)) > 1 - \eta_j^i \geq \sup_{w \in B_X} P^*(a_j^i)^*(w) - \eta_j^i$ . Now,  $D = \sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i ((S(B_X, P^*(a_j^i)^*, \eta_j^i) \cap S(B_X, x_i^*, \tilde{\varepsilon})))$  is a convex combination of nonempty relatively weakly open subset of  $B_X$ . By Bourgain's lemma,  $D$  contains a convex combination of slices of  $B_X$  and since diameter of any convex combination of slices of  $B_X$  is greater than  $\varepsilon$ , so  $\text{dia} \sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i ((S(B_X, P^*(a_j^i)^*, \eta_j^i) \cap S(B_X, x_i^*, \tilde{\varepsilon}))) > \varepsilon$ . Then there exists  $x_j^i, z_j^i \in (S(B_X, P^*(a_j^i)^*, \eta_j^i) \cap S(B_X, x_i^*, \tilde{\varepsilon}))$  such that  $\|\sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i x_j^i - \sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i z_j^i\| > \varepsilon$ . Note that,  $\sum_{j=1}^{n_i} \gamma_j^i x_j^i \in \sum_{j=1}^{n_i} \gamma_j^i ((S(B_X, P^*(a_j^i)^*, \eta_j^i) \cap S(B_X, x_i^*, \tilde{\varepsilon}))) \Rightarrow P(\sum_{j=1}^{n_i} \gamma_j^i x_j^i) \in \sum_{j=1}^{n_i} \gamma_j^i (S(B_{X/Y}, (a_j^i)^*, \eta_j^i) \cap A_i)$ . Since  $\text{diam}(\sum_{j=1}^{n_i} \gamma_j^i (S(B_{X/Y}, (a_j^i)^*, \eta_j^i) \cap A_i)) < \delta_0, \forall i = 1, 2, \dots, n$ , we have ,

$$\|P(\sum_{j=1}^{n_i} \gamma_j^i x_j^i) - \sum_{j=1}^{n_i} \gamma_j^i a_j^i\| < \delta_0. \Rightarrow \|P(\sum_{j=1}^{n_i} \gamma_j^i x_j^i)\| < \delta_0 + \|\sum_{j=1}^{n_i} \gamma_j^i a_j^i\| < 2\delta_0 \quad \forall i = 1, 2, \dots, n$$

Thus,  $d(\sum_{j=1}^{n_i} \gamma_j^i x_j^i, Y) < 2\delta_0 \Rightarrow$  for each  $i = 1, 2, \dots, n$  there exists  $v_i \in B_Y$  such that  $\|v_i - \sum_{j=1}^{n_i} \gamma_j^i x_j^i\| < 2\delta_0$ . Similarly , for each  $i = 1, 2, \dots, n$  there exists  $w_i \in B_Y$  such that  $\|w_i - \sum_{j=1}^{n_i} \gamma_j^i z_j^i\| < 2\delta_0$ . Now ,

$$y_i^*(v_i) = y_i^*(\sum_{j=1}^{n_i} \gamma_j^i x_j^i) + y_i^*(v_i - \sum_{j=1}^{n_i} \gamma_j^i x_j^i) > 1 - \tilde{\varepsilon} - 2\delta_0$$

Similarly ,  $y_i^*(w_i) > 1 - \tilde{\varepsilon} - 2\delta_0$ . Thus  $v_i, w_i \in S(B_Y, y_i^*, \tilde{\varepsilon} + 2\delta_0)$

$$\text{dia}(\sum_{i=1}^n \lambda_i S(B_Y, y_i^*, \tilde{\varepsilon} + 2\delta_0)) \geq \|\sum_{i=1}^n \lambda_i v_i - \sum_{i=1}^n \lambda_i w_i\|$$

$$\begin{aligned}
 &\geq \|\sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i x_j^i - \sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i z_j^i\| - \|\sum_{i=1}^n \lambda_i v_i - \sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i x_j^i\| \\
 &\quad - \|\sum_{i=1}^n \lambda_i w_i - \sum_{i=1}^n \lambda_i \sum_{j=1}^{n_i} \gamma_j^i z_j^i\| \\
 &> \varepsilon - \sum_{i=1}^n \lambda_i \|v_i - \sum_{j=1}^{n_i} \gamma_j^i x_j^i\| - \sum_{i=1}^n \lambda_i \|w_i - \sum_{j=1}^{n_i} \gamma_j^i z_j^i\| \\
 &> \varepsilon - 2\delta_0 - 2\delta_0 \\
 &= \varepsilon - 4\delta_0
 \end{aligned}$$

Since ,  $\sum_{i=1}^n \lambda_i S(B_Y, y_i^*, \tilde{\varepsilon} + 2\delta_0) \subset \sum_{i=1}^n \lambda_i S(B_Y, y_i^*, \delta)$  so

$$\text{dia}(\sum_{i=1}^n \lambda_i S(B_Y, y_i^*, \delta)) \geq \text{dia}(\sum_{i=1}^n \lambda_i S(B_Y, y_i^*, \tilde{\varepsilon} + 2\delta_0)) > \varepsilon - 4\delta_0 > \varepsilon - \frac{\varepsilon}{2} = \frac{\varepsilon}{2},$$

a contradiction. □

**Proposition 4.3.** Let  $Y$  be a closed subspace of  $X$  such that  $X/Y$  is finite dimensional and  $Y$  has *BHP*, then  $X$  has *BHP*.

*Proof.* Suppose  $X$  does not have *BHP*. Then there exists  $\varepsilon > 0$  such that diameter of any relatively weakly open set in  $B_X$  is  $> \varepsilon$ . Since  $Y$  has *BHP*, so let  $W = \{y \in B_Y : |y_i^*(y - y_0)| < \varepsilon_i \ \forall i = 1, 2, \dots, n\}$  where  $y_0 \in B_Y$  is relatively weakly open set in  $B_Y$  with diameter  $< \frac{\varepsilon}{2}$ . Now for each  $i = 1, 2, \dots, n$ , we can find  $\tilde{\varepsilon}_i > 0$  and  $\frac{\varepsilon}{4} > \delta_0 > 0$  such that  $\tilde{\varepsilon}_i + \delta_0 \|y_i^*\| < \varepsilon_i$ . By Hahn Banach theorem, we can extend  $y_i^*$  to a norm preserving extension  $x_i^*$  for all  $i = 1, 2, \dots, n$ . Define  $U = \{x \in B_X : |x_i^*(x - y_0)| < \tilde{\varepsilon}_i \ \forall i = 1, 2, \dots, n\}$ . Clearly  $U \neq \emptyset$  and  $U$  is relatively weakly open in  $B_X$ . Let  $P : X \rightarrow X/Y$  be the map such that  $P(x) = x + Y$ . Then clearly  $P$  is onto and linear. Also  $P$  is open map by Open Mapping Theorem. Thus  $P(U)$  is a norm open set in  $X/Y$  and  $y_0 \in U \cap Y$ . Thus  $P(U)$  is a norm open set containing zero. So, there exists  $0 < \delta < \frac{\delta_0}{2}$  such that  $B(0, \delta) \subset P(U)$ . Put,  $B = P^{-1}(B(0, \delta)) \cap U \subset B_X$ . Now using the fact that norm norm continuous implies weak weak continuous and  $X/Y$  is finite dimensional we can conclude that  $B$  is relatively weakly open in  $B_X$ . Then  $dia(B) > \varepsilon$ . Thus there exists  $v_1, v_2 \in B$  such that  $\|v_1 - v_2\| > \varepsilon$ . Now,  $\|P(v_1)\| < \delta \Rightarrow d(v_1, Y) < \delta \Rightarrow \exists u_1 \in Y$  such that  $\|u_1 - v_1\| < \delta$ . Similarly for  $v_2$  there exists  $u_2 \in Y$  such that  $\|u_2 - v_2\| < \delta$ . Without loss of generality we can assume that  $u_1, u_2 \in B_Y$ . Otherwise we will choose  $\frac{u_1}{\|u_1\|}$  and  $\frac{u_2}{\|u_2\|}$ . Now  $\forall i = 1, 2, \dots, n$ ,  $|y_i^*(u_1 - y_0)| \leq |y_i^*(u_1 - v_1)| + |y_i^*(v_1 - y_0)| \leq \|y_i^*\| 2\delta + \tilde{\varepsilon}_i < \|y_i^*\| \delta_0 + \tilde{\varepsilon}_i < \varepsilon_i$

Thus,  $u_1 \in W$ . Similarly  $u_2 \in W$  and

$$\|u_1 - u_2\| \geq \|v_1 - v_2\| + \|u_1 - v_1\| + \|u_2 - v_2\| > \varepsilon - 4\delta > \varepsilon - 2\delta_0 > \varepsilon - \frac{\varepsilon}{2} = \frac{\varepsilon}{2}.$$

Thus,  $dia(W) > \frac{\varepsilon}{2}$ , a contradiction.  $\square$

**Acknowledgement.** *The second author's research is funded by the National Board for Higher Mathematics (NBHM), Department of Atomic Energy (DAE), Government of India, Ref No: 0203/11/2019-R&D-II/9249.*

#### REFERENCES

- [ALN] T. Abrahamsen, V. Lima, O. Nygaard; *Remarks on diameter 2 properties*, J.Conv. Anal. **20** (1) 439-452 (2013).
- [B1] R. D. Bourgin; *Geometric aspects of convex sets with the Radon-Nikodym property*, Lecture Notes in Mathematics Springer-Verlag Berlin **993** (1983).
- [B2] S. Basu; *On Ball dentable property in Banach Spaces*, Math. Analysis and its Applications in Modeling (IC-MAAM 2018) Springer Proceedings in Mathematics and Statistics **302**, 145-149 (2020).
- [B3] J. Bourgain; *La propriété de Radon-Nikodym*, Publ. Mathé. de l'Université Pierre et Marie Curie **36** (1979).
- [BGLPRZ1] J. Becerra Guerrero, G. López Pérez, A. Rueda Zoca; *Big Slices versus big relatively weakly open subsets in Banach spaces*, J.Math. Anal. Appl. **428** (2) 855-865 (2015).
- [BGLPRZ2] J. Becerra Guerrero, G. López Pérez, A. Rueda Zoca; *Extreme differences between weakly open subsets and convex combination of slices in Banach spaces*, Adv. Math. **269** 56-70 (2015).
- [BGLPRZ3] J. Becerra Guerrero, G. López Pérez, A. Rueda Zoca; *Octahedral norms and convex combination of slices in Banach spaces*, J.Funct.Anal. **266** (4) 2424-2435 (2014).

- [BR] S. Basu, T. S. S. R. K. Rao; *On Small Combination of slices in Banach Spaces*, Extracta Mathematica **31** 1-10 (2016).
- [DU] J. Diestel, J. J. Uhl; *Vector Measures*, Amer. Math. Soc. **15** (1977).
- [EW] G. À Edgar, R. Wheeler; *Topological properties of Banach spaces*, Pacific J. Math., **115** 317-350 (1984).
- [GGMS] N. Ghoussoub, G. Godefroy, B. Maurey, W. Schachermayer; *Some topological and geometrical structures in Banach spaces*, Mem. Amer. Math. Soc. **70** 378 (1987).
- [GM] N. Ghoussoub, B. Maurey;  *$G_\delta$  Embeddings in Hilbert Space*, J. Funct. Anal. **61** (1) 72-97 (1985).
- [GMS] N. Ghoussoub, B. Maurey, W. Schachermayer; *Geometrical implications of certain infinite-dimensional decomposition*, Trans. Amer. Math. Soc. **317**, 541–584 (1990).
- [HWW] P. Harmand, D. Werner and W. Werner; *M-ideals in Banach spaces and Banach algebras*, Lecture Notes in Mathematics, Springer-Verlag, Berlin. **1547** (1993).
- [L1] J. Langemets; *Geometrical structure in diameter 2 Banach spaces*, Dissertationes Mathematicae Universitatis Tartuensis **99** (2015).
- [R] H. P. Rosenthal; *On the structure of non-dentable closed bounded convex sets*, Adv. in Math. **70** 1-58 (1988).
- [S] W. Schachermayer; *The Radon Nikodym Property and the Krein-Milman Property are equivalent for strongly regular sets*, Trans. Amer. Math. Soc. **303** (2) 673-687 (1987).

<sup>1</sup> SUDESHNA BASU, DEPARTMENT OF MATHEMATICS, GEORGE WASHINGTON UNIVERSITY, WASHINGTON DC 20052 USA AND, DEPARTMENT OF MATHEMATICS, RAM KRISHNA MISSION VIVEKANANDA EDUCATION AND RESEARCH INSTITUTE, BELUR MATH, HOWRAH 711202 WEST BENGAL, INDIA

*Email address:* sudeshname1ody@gmail.com

<sup>2</sup> SUSMITA SEAL, DEPARTMENT OF MATHEMATICS, , RAM KRISHNA MISSION VIVEKANANDA EDUCATION AND RESEARCH INSTITUTE, BELUR MATH, HOWRAH 711202, WEST BENGAL, INDIA

*Email address:* susmitaseal1996@gmail.com