# A CAMPANATO REGULARITY THEORY FOR MULTI-VALUED FUNCTIONS WITH APPLICATIONS TO MINIMAL SURFACE REGULARITY THEORY

#### PAUL MINTER

ABSTRACT. The regularity theory of the Campanato space  $\mathcal{L}_k^{(q,\lambda)}(\Omega)$  has been successfully used to understand the regularity of solutions to certain geometric variational problems. Here we extend this theory to multi-valued functions, adapting for the most part Campanato's original ideas in [Cam64]. We then give an application within the regularity theory of stationary integral varifolds. More precisely, we prove a regularity theorem for certain blow-up classes of functions, which typically arise when studying blow-ups of sequences of stationary integral varifolds converging to higher multiplicity planes or unions of half-planes. In such a setting, based in part on ideas in [Wic14] and [BKW21], we are able to deduce a boundary regularity theory for multi-valued harmonic functions, which is the first of its kind. In conjunction with [Min21], the results here establish a regularity theorem for stable codimension one integral varifolds near classical cones of density  $\frac{5}{2}$ .

## Contents

1.	Preliminaries	2
2.	Multi-Valued Campanato Regularity Theory	5
3.	Applications to Minimal Surface Theory	23
References		35

Multi-valued functions were first introduced by F. Almgren [AJ00] to study the branching behaviour of area-minimising currents. In recent years multi-valued functions have successfully been used to describe the structure of large classes of stationary integral varifolds near multiplicity two planes, and in particular near multiplicity two branch points (see [Wic08], [Wic21], [BKW21]). In order to understand the structure of stationary integral varifolds close to higher multiplicity planes or higher multiplicity non-flat cones (see [Min21]), we need to develop suitable regularity results for special classes of multi-valued functions; for the applications we have in mind, such classes are known as a proper blow-up class (see [Wic14] for a key example). The functions within a proper blow-up class are generated by taking scaling limits of approximate graphical representations of sequences of certain stationary integral varifolds converging to a fixed stationary cone. The functions within a given blow-up class are typically multi-valued, in most cases of interest defined on either an open ball of half-ball in a plane, and inherit certain integral estimates from the stationary assumption on the varifolds. When it is possible to use these estimates to deduce  $C^{1,\alpha}$  regularity of the functions within a blowup class is a key problem in geometric measure theory, and is the motivation for our work here. In conjunction with [Min21], this work establishes a  $C^{1,\alpha}$  regularity theory for stable codimension one integral varifolds which are close to a stationary classical cone of density  $\frac{5}{2}$ ; this is the first instance of a regularity theorem in a non-flat setting of higher multiplicity when branch points may be present.

The present paper is divided into two parts. In Sections 1 and 2 we define Campanato spaces for multi-valued functions, and develop a regularity theory for these multi-valued Campanato spaces which mirrors that seen in Campanato's original single-valued work [Cam64]; other than some technical changes, Campanato's main ideas extend readily to this setting. Such a result we anticipate to have more applications than just to minimal surface regularity theory, which is why we choose to present it separately; for our purposes the results serve as a black box to apply to the minimal surface setting. We also give some adaptations of the Campanato regularity theorem suited to the minimal surface setting. In Section 3 we then define the notion of a proper blow-up class over a half-plane. Using the results from Section 2, and adapting ideas in [Wic14] and [BKW21], we deduce that the functions within such a class are in fact  $C^{1,\alpha}$ , with  $C^{1,\alpha}$  extensions up to the boundary of the half-plane, for some  $\alpha>0$  depending only on the class. We note that in particular, as the functions in these proper blow-up classes are multi-valued  $C^{1,\alpha}$  harmonic in the interior, these results give the first instance of a  $C^{1,\alpha}$  boundary regularity theory for multi-valued harmonic functions.

## 1. Preliminaries

The standard references for multi-valued functions are [AJ00], [DL10], and [DLS13]; the reader wishing to attain a broader background on multi-valued functions is recommended to consult these, as we shall only recall the notions we need.

For  $T \in \mathbb{R}^m$  we write  $[\![T]\!]$  to denote the Dirac mass centred at T. For  $Q \in \{1, 2, ...\}$  we write  $\mathcal{A}_Q(\mathbb{R}^m)$  for the space of unordered Q-tuples, i.e.

$$\mathcal{A}_Q(\mathbb{R}^m) := \left\{ \sum_{i=1}^Q \llbracket T_i \rrbracket : T_i \in \mathbb{R}^m \text{ for } i = 1, \dots, Q \right\}.$$

We equip  $\mathcal{A}_Q(\mathbb{R}^m)$  with a metric  $\mathcal{G}$  defined by

$$\mathcal{G}\left(\sum_{i} \llbracket R_{i} \rrbracket , \sum_{i} \llbracket T_{i} \rrbracket \right) := \min_{\sigma \in S_{Q}} \left(\sum_{i} |R_{i} - T_{\sigma(i)}|^{2}\right)^{1/2}$$

where  $S_Q$  is the group of permutations of  $\{1,\ldots,Q\}$ . It is easy to check that  $(\mathcal{A}_Q(\mathbb{R}^m),\mathcal{G})$  is a complete metric space. For  $T\in\mathcal{A}_Q(\mathbb{R}^m)$  we also write  $|T|:=\mathcal{G}(T,Q[0])$ . We stress that  $\mathcal{A}_Q(\mathbb{R}^m)$  is *not* a vector space, as there is no natural notion of addition for unordered Q-tuples.

In the same way as in [Cam64], we work in certain domains of  $\mathbb{R}^n$  which obey a certain mass condition:

**Definition 1.1.** Let A > 0. We say a bounded domain  $\Omega \subset \mathbb{R}^n$  is A-weighted if

$$\mathcal{H}^n(\Omega \cap B_\rho(x)) \ge A\rho^n$$

for all  $x \in \overline{\Omega}$  and  $\rho \in (0, \operatorname{diam}(\Omega)]$ , where  $\mathcal{H}^n$  denotes the *n*-dimensional Hausdorff measure and  $\operatorname{diam}(\Omega) := \sup_{x,y \in \Omega} |x-y|$  is the diameter of  $\Omega$ .

An A-weighted domain always take up a fixed proportion of every ball centred at points on  $\overline{\Omega}$ . Key examples of A-weighted domains (for some fixed A > 0) include the open ball  $B_1(0) \subset \mathbb{R}^n$  and half-ball  $B_1(0) \cap \{x^1 > 0\} \subset \mathbb{R}^n$ . We note that A-weighted domains are referred to as type(A) domains in [RSS13] and type(I) domains in [Cam64].

Fix  $\Omega \subset \mathbb{R}^n$  a domain. A Q-valued function is a function  $u: \Omega \to \mathcal{A}_Q(\mathbb{R}^m)$ ; we write  $u(x) = \sum_i \llbracket u_i(x) \rrbracket$  for some  $u_i(x) \in \mathbb{R}^m$  which are unique up to permutations. In the case m = 1, we can define single-valued functions  $\tilde{u}_i: \Omega \to \mathbb{R}$  with  $\tilde{u}_1 \geq \tilde{u}_2 \geq \cdots \geq \tilde{u}_Q$  and  $u(x) = \sum_i \llbracket \tilde{u}_i(x) \rrbracket$  for all x. Even though  $\mathcal{A}_Q(\mathbb{R}^m)$  is not a vector space, we shall abuse notation and write for a single-valued function  $f: \Omega \to \mathbb{R}^m$  and Q-valued function  $g: \Omega \to \mathcal{A}_Q(\mathbb{R}^m)$  the function f + g to mean the Q-valued function given by  $x \mapsto \sum_i \llbracket f(x) + g_i(x) \rrbracket$ .

For  $\alpha \in (0,1]$ , we define the space of Q-valued  $\alpha$ -Hölder continuous functions  $C^{0,\alpha}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  in the usual way for functions between metric spaces. Similarly we define  $C^{0,\alpha}(\overline{\Omega}; \mathcal{A}_Q(\mathbb{R}^m))$  to be those functions in  $C^{0,\alpha}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  which have  $\alpha$ -Hölder continuous extensions to  $\overline{\Omega}$ . For each  $p \in [1,\infty)$  we define  $L^p(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  for the Q-valued functions which have  $\|u\|_p := \|\mathcal{G}(u,Q[0])\|_{L^p(\Omega)} < \infty$ .

**Definition 1.2.** We say that a Q-valued function  $u: \Omega \to \mathcal{A}_Q(\mathbb{R}^m)$  is differentiable at  $x_0 \in \Omega$  if there exist Q matrices  $A_i \in \mathbb{R}^{m \times n}$ ,  $i = 1, \ldots, Q$ , which satisfy

$$\lim_{h\to 0} \frac{\mathcal{G}(u(x_0+h), L(h))}{|h|} \to 0$$

where  $L(h) := \sum_i \llbracket u_i(x_0) + A_i(h) \rrbracket$ ; we then write  $Du_i(x_0) := A_i$ , and  $Du(x_0)$  is the Q-valued function given by  $\sum_i \llbracket A_i \rrbracket : \mathbb{R}^n \to \mathcal{A}_Q(\mathbb{R}^m)$ .

We can then define spaces such as  $C^k(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ ,  $C^{k,\alpha}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ ,  $k = 1, 2, \ldots$ , in the natural ways, and we denote the corresponding "norms" and "semi-norms" by  $|u|_{k,\alpha;\Omega}$ ,  $[u]_{k,\alpha;\Omega}$ , respectively. Note that as there is no vector space structure, these are not norms in the usual sense, however we shall still refer to them as norms and semi-norms.

One result which we need later is the following version of the Lebesgue differentiation theorem for multi-valued functions.

**Theorem 1.1** (Multi-Valued Lebesgue Differentiation Theorem). Suppose  $\Omega \subset \mathbb{R}^n$  is a domain,  $q \in [1, \infty)$ , and  $u \in L^q(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ . Then for  $\mathcal{H}^n$ -a.e.  $x_0 \in \Omega$  we have

$$\lim_{\rho \to 0} \frac{1}{\omega_n \rho^n} \int_{B_{\rho}(x_0)} \mathcal{G}(u(x), u(x_0))^q \, \mathrm{d}x = 0$$

where here  $\omega_n := \mathcal{H}^n(B_1(0))$  is the volume of the n-dimensional unit ball in  $\mathbb{R}^n$ .

*Proof.* From [AJ00] or [DL10, Theorem 2.1] we know that there exists N = N(m, Q) and a bi-Lipschitz function  $\xi : \mathcal{A}_Q(\mathbb{R}^m) \to \xi(\mathcal{A}_Q(\mathbb{R}^m)) \subset \mathbb{R}^N$  such that  $\text{Lip}(\xi) \leq 1$  and  $\text{Lip}(\xi^{-1}) \leq C(m, Q)$ . Now consider the function  $\tilde{u} := \xi \circ u : \Omega \to \mathbb{R}^N$ . Then since  $u \in L^q(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ 

and  $\xi$  is Lipschitz we see that  $\tilde{u} \in L^q_{loc}(\Omega)$ ; indeed, fixing some  $p \in \Omega$  we have for all  $x \in \Omega$ ,

$$|\tilde{u}(x)| \le |\tilde{u}(p)| + \mathcal{G}(u(x), u(p)) \le |\tilde{u}(p)| + |u(x)| + |u(p)|.$$

Hence applying the usual Lebesgue differentiation theorem to  $\tilde{u}$  we see that for  $\mathcal{H}^n$ -a.e.  $x_0 \in \Omega$ ,

$$\lim_{\rho \to 0} \frac{1}{\omega_n \rho^n} \int_{B_{\rho}(x_0)} |\tilde{u}(x) - \tilde{u}(x_0)|^q \, dx = 0.$$

Hence for  $\mathcal{H}^n$ -a.e.  $x_0 \in \Omega$ ,

$$\frac{1}{\omega_{n}\rho^{n}} \int_{B_{\rho}(x_{0})} \mathcal{G}(u(x), u(x_{0}))^{q} = \frac{1}{\omega_{n}\rho^{n}} \int_{B_{\rho}(x_{0})} \mathcal{G}(\xi^{-1}(\tilde{u}(x)), \xi^{-1}(\tilde{u}(x_{0})))^{q} dx 
\leq \frac{C(m, Q)^{q}}{\omega_{n}\rho^{n}} \int_{B_{\rho}(x_{0})} |\tilde{u}(x) - \tilde{u}(x_{0})|^{q} dx 
\to 0$$

and so we are done.

**Definition 1.3.** We say that  $P: \mathbb{R}^n \to \mathcal{A}_Q(\mathbb{R}^m)$  is a Q-valued polynomial if there exist Q functions  $p_i: \mathbb{R}^n \to \mathbb{R}^m$  such that for each  $i = 1, \ldots, Q$  and  $j = 1, \ldots, m, p_i^j: \mathbb{R}^n \to \mathbb{R}$  is a polynomial function, and for all  $x \in \mathbb{R}^n$ :

$$p(x) = \sum_{i} \llbracket p_i(x) \rrbracket.$$

We define the degree of p by  $deg(p) := \max_{i,j} deg(p_i^j)$ .

For each k = 0, 1, ..., we write  $\mathcal{P}_k$  for the set of all Q-valued polynomials  $p : \mathbb{R}^n \to \mathbb{R}^m$  with  $\deg(p) \leq k$ .

**Definition 1.4.** Let  $q \in [1, \infty)$ ,  $\lambda \in (0, \infty)$ ,  $k \in \{0, 1, 2, \dots\}$ , and  $\Omega \subset \mathbb{R}^n$  be a domain. We then define the *Q-valued Campanato space*  $\mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  to be the set of functions  $u \in L^q(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  which obey

$$|||u|||_{k,q,\lambda;\Omega} := \sup_{\substack{x_0 \in \overline{\Omega} \\ \rho \in (0,\operatorname{diam}(\Omega)]}} \left[ \rho^{-\lambda} \inf_{P \in \mathcal{P}_k} \int_{\Omega \cap B_{\rho}(x_0)} \mathcal{G}(u(x),P(x))^q \, dx \right]^{1/q} < \infty.$$

Remark. We will always suppress the domain dependence on our norms and semi-norms when the domain is clear from context, and again we still refer to them as norms and semi-norms despite  $\mathcal{A}_Q(\mathbb{R}^m)$  not being a vector space.

Of course  $||u||_{k,q,\lambda} = 0$  whenever  $u \in \mathcal{P}_k$ , and so  $|\cdot|_{k,q,\lambda}$  is only a semi-norm. To make  $\mathcal{L}_k^{q,\lambda}(\Omega;\mathcal{A}_Q(\mathbb{R}^m))$  into a norm we instead work with

$$||u||_{k,q,\lambda}^q := ||u||_q^q + ||u||_{k,q,\lambda}^q.$$

#### 2. Multi-Valued Campanato Regularity Theory

The purpose of this section is to develop the regularity theory of the space  $\mathcal{L}_k^{q,\lambda}(\Omega;\mathcal{A}_Q(\mathbb{R}^m))$ , following the ideas seen in [Cam64]. At the end of this section we also provide extensions relevant to geometric problems, such as the study of stationary integral varifolds. The main general regularity result of this section is the following theorem:

**Theorem 2.1** (Regularity of  $\mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ ). Suppose  $\Omega \subset \mathbb{R}^n$  is a convex A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ . Suppose that  $\lambda$  obeys  $n + \ell q < \lambda < n + (\ell+1)q$  for some  $\ell \in \{0,1,\ldots,k\}$ . Then  $u \in C^{\ell,\alpha}(\overline{\Omega}; \mathcal{A}_Q(\mathbb{R}^m))$  with the estimate

$$[u]_{\ell,\alpha} \le C ||u||_{\ell,q,\lambda}$$

where  $C = C(n, m, k, \ell, q, Q, A, \lambda)$ . Moreover  $\mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  is continuously isomorphic to  $\mathcal{L}_\ell^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ .

We fix  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  throughout. First note that for each  $x_0 \in \overline{\Omega}$  and  $\rho \in (0, \operatorname{diam}(\Omega)]$ , we can find a Q-valued polynomial  $\tilde{P} \in \mathcal{P}_k$  such that

$$\inf_{P \in \mathcal{P}_k} \int_{\Omega \cap B_o(x_0)} \mathcal{G}(u, P)^q = \int_{\Omega \cap B_o(x_0)} \mathcal{G}(u, \tilde{P})^q.$$

Indeed, each  $P \in \mathcal{P}_k$  is determined by polynomials  $p^{i,j} : \mathbb{R}^n \to \mathbb{R}$ , which are in turn determined by coefficients  $a = (a_p^{i,j})_{i,j,p}$  where  $i = 1, \ldots, Q, \ j = 1, \ldots, m$ , and  $p \in \mathbb{N}^n$  is a multi-index with  $|p| \leq k$ . Then the function  $f : \mathbb{R}^M \to \mathbb{R}, \ M = M(n, m, k, Q)$ , sending

$$f: ((a_p^{i,j})_{i,j,p}) \longmapsto \int_{\Omega \cap B_{\varrho}(x_0)} \mathcal{G}(u,P)^q$$

where P is the polynomial generated by  $a=(a_p^{i,j})_{i,j,p}$  as above, is a continuous function. Therefore to see the infimum is attained, we just need to show that  $f(a) \to \infty$  as  $|a| \to \infty$ . But the triangle inequality gives

$$f(a) := \int_{\Omega \cap B_{\rho}(x_0)} \mathcal{G}(u, P)^q \ge \left| \|u\|_{L^q(\Omega \cap B_{\rho}(x_0))} - \|P\|_{L^q(\Omega \cap B_{\rho}(x_0))} \right|^q$$

which clearly  $\to \infty$  as  $|a| \to \infty$ . We write  $P_{x_0,\rho} := \tilde{P}$  for a choice of Q-valued polynomial  $\tilde{P}$  attaining the infimum.

We will write  $a_p^{i,j}(x_0,\rho) := D^p P^{i,j}(x_0,\rho)$  so that

$$P_{x_0,\rho}^{i,j}(x) = \sum_{|p| \le k} \frac{a_p^{i,j}(x_0,\rho)}{p!} (x-x_0)^p.$$

We first prove a crucial integral estimate for comparing two Q-valued polynomials. Whilst the estimate is similar to that seen in [Cam64, Lemma 2.I], we need to be more careful due to the lack of any vector space structure. Indeed, when Q > 1,  $\mathcal{A}_Q(\mathbb{R}^{n \times m}) \not\cong \mathcal{A}_Q(\mathbb{R}^n) \times \mathcal{A}_Q(\mathbb{R}^m)$ , since in the former space the  $(n \times m)$ -tuples need to be close to one another for a given

permutation, whilst in the later we are able to use different permutations. For us, this would correspond to being able to permute the constants in the Q-valued polynomials independently of the linear coefficients, which will not be allowed.

**Lemma 2.1.** Let  $q \in [1, \infty)$  and  $E \subset B_{\rho}(x_0)$  be a measurable subset with  $\mathcal{H}^n(E) \geq A\rho^n$  for some A > 0. Then for any pair of Q-valued polynomials  $F, G \in \mathcal{P}_k$ , given by

$$F^{i,j}(x) := \sum_{|p| \le k} \frac{a_p^{i,j}}{p!} (x - x_0)^p \quad and \quad G^{i,j}(x) := \sum_{|p| \le k} \frac{b_p^{i,j}}{p!} (x - x_0)^p$$

we have, defining  $a_{\rho} \in \mathcal{A}_{Q}(\mathbb{R}^{M})$  and  $b_{\rho} \in \mathcal{A}_{Q}(\mathbb{R}^{M})$ , M = M(n, m, k), by  $a_{\rho}^{i} := (\rho^{|p|} a_{p}^{i,j})_{j,p}$  and  $b_{\rho}^{i} := (\rho^{|p|} b_{p}^{i,j})_{j,p}$ ,

$$\mathcal{G}(a_{\rho}, b_{\rho})^q \le \frac{C_1}{\rho^n} \int_E \mathcal{G}(F(x), G(x))^q \, \mathrm{d}x$$

for some  $C_1 = C_1(n, m, k, q, Q, A)$ , which is in particular independent of F, G, E.

*Proof.* First note that if we set  $\tilde{F}(x) := F(x_0 + \rho x)$  and  $\tilde{G}(x) := F(x_0 + \rho x)$ , then we see

$$\tilde{F}^{i,j}(x) = \sum_{|p| \le k} \frac{(\rho^{|p|} a_p^{i,j})}{p!} x^p$$
 and  $\tilde{G}^{i,j}(x) = \sum_{|p| \le k} \frac{(\rho^{|p|} b_p^{i,j})}{p!} x^p$ 

and so we see that it suffices to prove the case  $x_0 = 0$  and  $\rho = 1$ . So we restrict to this case.

So suppose we have two Q-valued polynomials F, G, and write

$$F = \sum_{i=1}^{Q} \llbracket F^i \rrbracket$$
 and  $G = \sum_{i=1}^{Q} \llbracket G^i \rrbracket$ 

where  $F^i = (F^{i,j})_{j=1}^m$ ,  $G^i = (G^{i,j})_{j=1}^m$ , and

$$F^{i,j}(x) = \sum_{|p| \le k} a_p^{i,j} x^p$$
 and  $G^{i,j}(x) = \sum_{|p| \le k} b_p^{i,j} x^p$ .

Note that we have combined with the coefficients the p! factors for notational simplicity, and these can simply be removed at the end. Define  $a, b \in \mathcal{A}_Q(\mathbb{R}^M)$ , where M = M(n, m, k), where  $a^i = (a_p^{i,j})_{j,p}$  and  $b^i = (b_p^{i,j})_{j,p}$ , i.e.,  $a^i$  determines all the polynomials in  $F^i$ , etc. Now define:

$$\alpha_{F,G} := \mathcal{G}(a,b) \equiv \inf_{\sigma} \left( \sum_{i=1}^{Q} \sum_{j=1}^{m} \sum_{|p| \le k} |a_p^{i,j} - b_p^{\sigma(i),j}|^2 \right)^{1/2}.$$

Let  $\mathcal{F}$  denote all pairs (F,G) of Q-valued polynomials in  $\mathcal{P}_k$  which have  $\alpha_{F,G}=1$ . Let  $\mathcal{E}$  be the class of subsets  $E\subset B_1(0)$  which have  $\mathcal{H}^n(E)\geq A$ . Then define:

$$\gamma := \inf_{\substack{(F,G) \in \mathcal{F} \\ F \subseteq \mathcal{E}}} \int_E \mathcal{G}(F,G)^q.$$

We claim that  $\gamma \geq C$  for some constant C = C(n, m, k, q, Q, A). Indeed, fix  $(F, G) \in \mathcal{F}$ , and using the same notation as just defined, for each  $\sigma \in S_Q$ ,  $i = 1, \ldots, Q$ ,  $j = 1, \ldots, m$ , and  $|p| \leq k$ , set

$$c_{i,j,p}^{\sigma}:=a_p^{i,j}-b_p^{\sigma(i),j} \quad \text{ and } \quad P_{i,j}^{\sigma}(x):=\sum_{|p|\leq k}c_{i,j,p}^{\sigma}x^p.$$

Then we have for each  $x \in B_1(0)$ ,

$$\mathcal{G}(F(x), G(x))^2 := \inf_{\sigma} \sum_{i=1}^{Q} |F^i(x) - G^i(x)|^2 = \inf_{\sigma} \sum_{i=1}^{Q} \sum_{j=1}^{m} \left| \sum_{|p| \le k} (a_p^{i,j} - b_p^{\sigma(i),j}) x^p \right|^2$$

i.e.,

$$\mathcal{G}(F(x), G(x))^2 = \inf_{\sigma} \sum_{i,j} |P_{i,j}^{\sigma}(x)|^2.$$

Now define for each  $\sigma \in S_Q$ ,  $E_{\sigma} \subset E$  to be the set of points  $x \in E$  for which this infimum equals  $\sum_{i,j} |P_{i,j}^{\sigma}(x)|^2$ . Clearly  $E = \bigcup_{\sigma} E_{\sigma}$ , and so we must be able to find  $\sigma' \in S_Q$  for which  $\mathcal{H}^n(E_{\sigma'}) \geq A/|S_Q| = A/(Q!)$ . Now note that

$$\int_E \mathcal{G}(F,Q)^q \geq \int_{E_{\sigma'}} \mathcal{G}(F,G)^q = \int_{E_{\sigma'}} \sum_{i,j} |P_{i,j}^{\sigma'}|^2.$$

Now fix each  $\sigma \in S_Q$  we have by definition of  $\alpha_{F,G}$  and since  $(F,G) \in \mathcal{F}$ ,

$$\sum_{i=1}^{Q} \sum_{j=1}^{m} \sum_{|p| \le k} |c_{i,j,p}^{\sigma}|^2 \ge \alpha_{F,G}^2 = 1$$

and hence applying this with  $\sigma = \sigma'$ , we must be able to find some  $i' \in \{1, ..., Q\}, j' \in \{1, ..., m\}$ , and  $|p'| \leq k$  for which

$$|c_{i',j',p'}^{\sigma'}| \ge \frac{1}{mQN(n,k)}.$$

Now applying [Cam64, Lemma 2.I] with A/(Q!) in place of A, F in place of E, where  $F \subset E_{\sigma'}$  has  $\mathcal{H}^n(F) = A/(Q!)$ , p = p', we have:

$$\int_{E\sigma'} \sum_{i,j} |P_{i,j}^{\sigma'}|^q \ge \int_F |P_{i',j'}^{\sigma'}|^q \ge \tilde{C} |D^{p'} P_{i',j'}^{\sigma'}(0)|^q = \tilde{C} |(p_{\sigma'}!) \cdot c_{i',j',p'}^{\sigma'}|^q \ge C_*$$

where  $\tilde{C} = \tilde{C}(n, k, q, A/(Q!))$  and  $C_* = C_*(n, m, k, q, Q, A)$ . Hence we see that

$$\int_{E} \mathcal{G}(F, Q)^{q} \ge C$$

and so as  $(F,G) \in \mathcal{F}$  and  $E \in \mathcal{E}$  were arbitrary, we see  $\gamma \geq C_*$ , as desired.

Now given any arbitrary pair of distinct Q-valued functions (F, G), note that  $(F/\alpha_{F,G}, G/\alpha_{F,G}) \in \mathcal{F}$ , and so for any  $E \subset A$  with  $\mathcal{H}^n(E) \geq A$  we have

$$\int_{E} \mathcal{G}(F,G)^{q} \ge C\alpha_{F,G}^{q}$$

which completes the proof.

Remark. In particular we see from Lemma 2.1 that, if we write  $a^i_{(r)}:=(a^{i,j}_p)_{j=1,\dots,m,|p|\leq r}$  and  $b^i_{(r)}:=(b^{i,j}_p)_{j=1,\dots,m,|p|\leq r}$ , then,

$$\mathcal{G}(a_{(r)}, b_{(r)})^q \equiv \inf_{\sigma} \left( \sum_{i=1}^{Q} \sum_{j=1}^{m} \sum_{|p| \le r} |a_p^{i,j} - b_p^{\sigma(i),j}|^2 \right)^{q/2} \le \frac{C_1}{\min\{1, \rho\}^{n+rq}} \int_E \mathcal{G}(F, G)^q \, dx.$$

**Lemma 2.2.** Let  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ . Then for each  $x_0 \in \overline{\Omega}$ ,  $\rho \in (0, \operatorname{diam}(\Omega)]$ , and  $\ell \in \{0, 1, 2, \dots\}$ , we have for some  $C = C(q, \lambda)$ ,

$$\int_{\Omega \cap B_{2^{-\ell-1}\rho}(x_0)} \mathcal{G}(P_{x_0,2^{-\ell}\rho}, P_{x_0,2^{-\ell-1}\rho})^q \le C2^{-\ell\lambda}\rho^{\lambda} \cdot |||u|||_{k,q,\lambda}^q.$$

*Proof.* We simply have by the triangle inequality for  $\mathcal{G}$ ,

$$\begin{split} \int_{\Omega \cap B_{2^{-\ell-1}\rho}(x_0)} & \mathcal{G}(P_{x_0,2^{-\ell}\rho},P_{x_0,2^{-\ell-1}\rho})^q \\ & \leq 2^q \int_{\Omega \cap B_{2^{-\ell}\rho}(x_0)} & \mathcal{G}(P_{x_0,2^{-\ell}\rho},u)^q + 2^q \int_{\Omega \cap B_{2^{-\ell-1}\rho}(x_0)} & \mathcal{G}(P_{x_0,2^{-\ell-1}\rho},u)^q \\ & \leq 2^q \cdot (2^{-\ell}\rho)^\lambda \|\|u\|_{k,q,\lambda}^q + 2^q \cdot (2^{-\ell-1}\rho)^\lambda \|\|u\|_{k,q,\lambda}^q \\ & = \left[2^q + 2^{q-\lambda}\right] \cdot 2^{-\ell\lambda}\rho^\lambda \|\|u\|_{k,q,\lambda}^q. \end{split}$$

The next lemma will be the first step towards a Hölder estimate for the k'th order derivatives and differentiability properties for lower order derivatives.

**Lemma 2.3.** Suppose  $\Omega \subset \mathbb{R}^n$  is an A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ . Then for each pair of points  $x_0, y_0 \in \overline{\Omega}$  with  $\rho := |x_0 - y_0| \leq \frac{1}{2} \mathrm{diam}(\Omega)$ , we have

$$\mathcal{G}(a_k(x_0, 2\rho), a_k(y_0, 2\rho))^q \le C_1 2^{q+1+\lambda} |||u|||_{k,q,\lambda}^q \cdot \rho^{\lambda - n - kq}$$

where  $C_1$  is as in Lemma 2.1, and  $a_k^i(x_0, 2\rho) = (a_p^i(x_0, 2\rho))_{|p|=k}$ , etc.

*Proof.* Fix  $x_0, y_0 \in \overline{\Omega}$  as in the statement and set  $\rho := |x_0 - y_0|$ . Then by the triangle inequality for  $\mathcal{G}$  and simple inclusions we have

$$\int_{\Omega \cap B_{\rho}(x_{0})} \mathcal{G}(P_{x_{0},2\rho}, P_{y_{0},2\rho})^{q} \leq 2^{q} \int_{\Omega \cap B_{2\rho}(x_{0})} \mathcal{G}(P_{x_{0},2\rho}, u)^{q} + 2^{q} \int_{\Omega \cap B_{2\rho}(y_{0})} \mathcal{G}(P_{y_{0},2\rho}, u)^{q} \\
\leq 2^{q} \cdot (2\rho)^{\lambda} ||u||_{k,q,\lambda}^{q} + 2^{q} \cdot (2\rho)^{\lambda} ||u||_{k,q,\lambda}^{q} \\
= 2^{q+\lambda+1} ||u||_{k,q,\lambda}^{q} \cdot \rho^{\lambda}.$$

Moreover from Lemma 2.1, since  $\Omega$  is A-weighted and because the k'th derivative of any polynomial of degree k is constant (so in particular does not depend on the point we evaluate at, as Lemma 2.1 requires both polynomials to be centred at the same point) we have

$$\mathcal{G}(a_k(x_0, 2\rho), a_k(y_0, 2\rho))^q \le \frac{C_1}{\rho^{n+kq}} \int_{\Omega \cap B_\rho(x_0)} \mathcal{G}(P_{x_0, 2\rho}, P_{y_0, 2\rho})^q$$

since we can just look at the order k terms. Combining these inequalities we get the result.  $\Box$ 

**Lemma 2.4.** Suppose  $\Omega \subset \mathbb{R}^n$  is an A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ . Then  $\exists C = C(n, m, k, q, \lambda, Q, A)$  such that, for all  $x \in \overline{\Omega}$ ,  $\rho \in (0, \min\{1, \operatorname{diam}(\Omega)\}]$ ,  $i = 0, 1, 2, \ldots$ , and  $r \leq k$  we have

$$\mathcal{G}(a_{(r)}(x_0, \rho), a_{(r)}(x_0, 2^{-i}\rho)) \le C \|\|u\|\|_{k,q,\lambda} \sum_{j=0}^{i-1} 2^{j\left(\frac{n+rq-\lambda}{q}\right)} \cdot \rho^{\frac{\lambda-n-rq}{q}}.$$

*Proof.* For any such  $x_0, \rho, i, r$ , we have

$$\begin{split} \mathcal{G}(a_{(r)}(x_{0},\rho),a_{(r)}(x_{0},2^{-i}\rho)) &\leq \sum_{j=0}^{i-1} \mathcal{G}(a_{(r)}(x_{0},2^{-j}\rho),a_{(r)}(x_{0},2^{-j-1}\rho)) \\ &\leq \sum_{j=0}^{i-1} \left( \frac{C_{1}}{(2^{-j-1}\rho)^{n+rq}} \int_{\Omega \cap B_{2^{-j-1}\rho}(x_{0})} \mathcal{G}(P_{x_{0},2^{-j}\rho},P_{x_{0},2^{-j-1}})^{q} \right)^{1/q} \\ &\leq \sum_{j=0}^{i-1} \left( C_{1}2^{(j+1)(n+rq)}\rho^{-n-rq} \cdot C \cdot 2^{-j\lambda}\rho^{\lambda} |||u|||_{k,q,\lambda}^{q} \right)^{1/q} \\ &\leq C |||u|||_{k,q,\lambda} \sum_{j=0}^{i-1} 2^{j\left(\frac{n+rq-\lambda}{q}\right)} \cdot \rho^{\frac{\lambda-n-rq}{q}} \end{split}$$

as desired, where the second inequality is from Lemma 2.1 and the third inequality is from Lemma 2.2.  $\Box$ 

**Lemma 2.5.** Suppose  $\Omega \subset \mathbb{R}^n$  is an A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ , with  $n+rq < \lambda \leq n+(r+1)q$  for some  $r \in \{0,1,\ldots,k\}$ . Then for every  $|p| \leq r$ , there exists a function  $v_p : \overline{\Omega} \to \mathcal{A}_Q(\mathbb{R}^m)$  such that for every  $x_0 \in \overline{\Omega}$  and  $\rho \in (0, \min\{1, \operatorname{diam}(\Omega)\}]$  we have

$$\mathcal{G}(a_{(r)}(x_0, \rho), v_{(r)}(x_0)) \le C \|u\|_{k,q,\lambda} \cdot \rho^{\frac{\lambda - n - rq}{q}}$$

where  $C = C(n, m, k, q, \lambda, Q, A)$ . In particular  $a_{(r)}(\cdot, \rho) \to v_{(r)}(\cdot)$  uniformly as  $\rho \to 0$ .

*Proof.* Fix  $x_0, \rho, r$  as in the statement of the lemma. For  $i, j \in \{0, 1, 2, ...\}$  sufficiently large with j > i so that  $2^{-i}\rho < 1$ , apply Lemma 2.4 with  $2^{-i}\rho$  in place of  $\rho$  and j - i in place of i to get

$$\mathcal{G}(a_{(r)}(x_0, 2^{-j}\rho), a_{(r)}(x_0, 2^{-i}\rho)) \le C \|\|u\|_{k, q, \lambda} \sum_{h=i}^{j-1} 2^{h\left(\frac{n+rq-\lambda}{q}\right)} \cdot \rho^{\frac{\lambda-n-rq}{q}}.$$

Thus as  $\lambda > n + rq$ , we see that  $\sum_{h=0}^{\infty} 2^{h\left(\frac{n+rq-\lambda}{q}\right)} < \infty$ , and thus the above shows that  $(a_{(r)}(x_0, 2^{-j}\rho))_j$  is a Cauchy sequence. Since  $(\mathcal{A}_Q(\mathbb{R}^M), \mathcal{G})$  is a metric space for every  $M \geq 1$ , this sequence therefore converges.

We claim that this limit is independent of the choice of  $\rho$ , and so only depends on  $x_0$ . Indeed, for any  $0 < \rho_1 \le \rho_2 \le \text{diam}(\Omega)$  we have for all sufficiently large i, by Lemma 2.1,

$$\begin{split} &\mathcal{G}(a_{(r)}(x_0, 2^{-i}\rho_1), a_{(r)}(x_0, 2^{-i}\rho_2))^q \\ &\leq \frac{C_1}{(2^{-i}\rho_1)^{n+rq}} \int_{\Omega \cap B_{2^{-i}\rho_1}(x_0)} \mathcal{G}(P_{x_0, 2^{-i}\rho_1}, P_{x_0, 2^{-i}\rho_2})^q \\ &\leq \frac{C_1 \cdot 2^{i(n+rq)} \cdot 2^q}{\rho_1^{n+rq}} \int_{\Omega \cap B_{2^{-i}\rho_1}(x_0)} \mathcal{G}(P_{x_0, 2^{-i}\rho_1}, u)^q + \frac{C_1 \cdot 2^{i(n+rq)} \cdot 2^q}{\rho_1^{n+rq}} \int_{\Omega \cap B_{2^{-i}\rho_2}(x_0)} \mathcal{G}(P_{x_0, 2^{-i}\rho_2}, u)^q \\ &\leq C_1 \cdot 2^q \cdot \|\|u\|_{k, q, \lambda}^q \cdot 2^{i(n+rq)} \rho^{-(n+rq)} \left( (2^{-i}\rho_1)^{\lambda} + (2^{-i}\rho_2)^{\lambda} \right) \\ &= C \|\|u\|_{k, q, \lambda}^q 2^{-i(\lambda - n - rq)} \cdot \left( \frac{\rho_1^{\lambda} + \rho_2^{\lambda}}{\rho_1^{n+rq}} \right) \end{split}$$

and so we see that, as  $i \to \infty$  the right-hand side of this inequality  $\to 0$ , showing that the limit is independent of  $\rho$ .

Now define for each  $x_0 \in \overline{\Omega}$  and  $|p| \le r$ ,

$$v_p(x_0) := \lim_{i \to \infty} a_p(x_0, 2^{-i}\rho)$$

for any choice of  $\rho \in (0, \operatorname{diam}(\Omega)]$ . Thus  $v_p : \overline{\Omega} \to \mathcal{A}_Q(\mathbb{R}^m)$ . But now note that for  $\rho < 1$ , by Lemma 2.4 we have for every i sufficiently large and  $\rho < 1$ ,

$$\mathcal{G}(a_{(r)}(x_0, \rho), a_{(r)}(x_0, 2^{-i}\rho)) \le C \|u\|_{k,q,\lambda} \rho^{\frac{\lambda - n - rq}{q}}$$

which follows by bounding the convergent sum in j by a constant depending only on  $n, r, q, \lambda$ . Thus taking  $i \to \infty$  we see

$$\mathcal{G}(a_{(r)}(x_0, \rho), v_{(r)}(x_0)) \le C \|u\|_{k,q,\lambda} \cdot \rho^{\frac{\lambda - n - rq}{q}}$$

and this is true independently of  $x_0$ . In particular this shows that  $\lim_{\rho\to 0} a_{(r)}(\cdot,\rho) = v_{(r)}(\cdot)$  uniformly.

We now study the regularity properties of the functions  $v_p$ . Ultimately we will show that  $D^p u = v_p$ , and thus properties for the  $v_p$  will provide the necessary conclusions for u. The first step is to show that the "top"  $v_p$ , i.e. those with |p| = k, are  $\alpha$ -Hölder continuous in  $\overline{\Omega}$  for appropriate  $\lambda$ . Note that from Lemma 2.1,  $v_p$  is independent of the choice of polynomials  $P_{x_0,\rho}$  attaining the infimum.

**Lemma 2.6.** Suppose  $\Omega \subset \mathbb{R}^n$  is a convex A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  for some  $\lambda > n + kq$ . Then the function  $v_k = (v_p)_{|p|=k}$  is Hölder continuous in  $\overline{\Omega}$ , with the estimate

$$\mathcal{G}(v_k(x), v_k(y)) \le C \|\|u\|_{k,q,\lambda} \cdot |x-y|^{\frac{\lambda - n - kq}{q}}$$

for some  $C = C(n, m, k, q, Q, \lambda, A)$ .

*Proof.* Suppose  $x, y \in \overline{\Omega}$  are such that  $\rho := |x - y| \le \frac{1}{2} \operatorname{diam}(\Omega)$ . Then,

$$\begin{split} \mathcal{G}(v_{k}(x), v_{k}(y)) &\leq \mathcal{G}(v_{k}(x), a_{k}(x, 2\rho)) + \mathcal{G}(a_{k}(x, 2\rho), a_{k}(y, 2\rho)) + \mathcal{G}(v_{k}(y), a_{k}(y, 2\rho)) \\ &\leq C \|u\|_{k, q, \lambda} \cdot (2\rho)^{\frac{\lambda - n - kq}{q}} + \tilde{C} \|u\|_{k, q, \lambda} \cdot \rho^{\frac{\lambda - n - kq}{q}} + C \|u\|_{k, q, \lambda} \cdot (2\rho)^{\frac{\lambda - n - kq}{q}} \\ &= C \|u\|_{k, q, \lambda} \cdot |x - y|^{\frac{\lambda - n - kq}{q}} \end{split}$$

where in the second inequality we have used Lemma 2.5 for the first and third terms, and Lemma 2.3 for the second term (which is where we need the condition on  $\rho$  and the fact that we can only deal with the |p| = k case).

If  $\rho > \frac{1}{2} \operatorname{diam}(\Omega)$ , then since  $\operatorname{diam}(\Omega) < \infty$  and  $\Omega$  is convex, the midpoint z := (x+y)/2 lies in  $\Omega$  and obeys  $|x-z|, |z-y| < \frac{1}{2} \operatorname{diam}(\Omega)$ . Hence applying the above with x, z and z, y, we get

$$\begin{split} \mathcal{G}(v_k(x), v_k(y)) &\leq \mathcal{G}(v_k(x), v_k(z)) + \mathcal{G}(v_k(z), v_k(y)) \\ &\leq C \|\|u\|_{k,q,\lambda} \left( |x-z|^{\frac{\lambda - n - kq}{q}} + |z-y|^{\frac{\lambda - n - kq}{q}} \right) \\ &= \tilde{C} \|\|u\|_{k,q,\lambda} |x-y|^{\frac{\lambda - n - kq}{q}} \end{split}$$

for some  $\tilde{C}$  independent of x, y. Hence we are done.

**Lemma 2.7.** Suppose  $\Omega \subset \mathbb{R}^n$  is a convex A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ , where  $k \in \{1, 2, ...\}$  and  $\lambda > n + kq$ . Then for any  $|p| \leq k - 1$  the function  $v_p$  is differentiable, and moreover for each  $i \in \{1, ..., Q\}$ ,  $j \in \{1, ..., n\}$ ,

$$D_j v_p^i = v_{p+e_j}^i$$

where  $e_j \in \mathbb{R}^n$  denotes the j'th standard basis vector.

*Proof.* We work by downwards induction on |p|. Fix a multi-index p with  $|p| \leq k-1$  and  $x_0 \in \overline{\Omega}$ . We know from Lemma 2.6 that  $v_p$ , where |p| = k, is Hölder continuous on  $\overline{\Omega}$ , and so in our induction we may assume without loss of generality that  $v_{p+\ell e_j}$  is continuous on  $\overline{\Omega}$  for  $\ell = 1, 2, \ldots, k - |p|$ .

Set  $L^i(\rho) := v^i_p(x_0) + \rho v^i_{p+e_j}(x_0)$ : note this is a well-defined Q-valued (affine) function, since we know that  $a_{(k)}(\cdot,\rho)$  converges to  $v_{(k)}$  uniformly by Lemma 2.5, and so for each i we have  $v^i_{(k)} = (v^i_p)_{|p| \le k}$  is a function  $\Omega \to \mathbb{R}^N$  for some N = N(n,m,k), and so given an i we have a well-defined way of choosing  $v^i_p$  and  $v^i_{p+e_j}$ . We will show that  $L(\rho)$  provides the suitable linear approximation to  $v_p$  at  $x_0$  for the definition of differentiability.

Firstly, since the  $v^i$  provide a natural ordering to the unordered tuples, for each  $x_0, \rho$  we can reorder  $a(x, \rho)$  to assume without loss of generality that

$$\mathcal{G}(a(x,\rho),v(x))^2 := \inf_{\sigma} \sum_{i} |v^i(x) - a^{\sigma(i)}(x,\rho)|^2 = \sum_{i} |v^i(x) - a^i(x,\rho)|^2$$

i.e., the infimum is attained at  $\sigma = id$ .

Now note that for any  $\sigma_1 \in S_Q$  and any  $\rho$  with  $|\rho|$  sufficiently small:

$$\rho^{-1}\mathcal{G}(v_{p}(x_{0}+\rho e_{j}),L(\rho)) \leq \underbrace{\rho^{-1}\mathcal{G}(v_{p}(x_{0}+\rho e_{j}),a_{p}(x_{0}+\rho e_{j},2|\rho|))}_{(1)} + \underbrace{\rho^{-1}\mathcal{G}(a_{p}(x_{0}+\rho e_{j},2|\rho|),a_{p}(x_{0},2|\rho|)+\rho v_{p+e_{j}}^{\sigma_{1}(\cdot)}(x_{0}))}_{(2)} + \underbrace{\rho^{-1}\mathcal{G}(a_{p}(x_{0},2|\rho|)+\rho v_{p+e_{j}}^{\sigma_{1}(\cdot)}(x_{0}),L(\rho))}_{(3)}.$$

Let us look at each term individually. For (1) we simply have from Lemma 2.5, as  $|p| \le k-1$ ,

$$(1) \le \rho^{-1} \mathcal{G}(v_{(k-1)}(x_0 + \rho e_j), a_{(k-1)}(x_0 + \rho e_j, 2|\rho|)) \le C \|u\|_{k,q,\lambda} \cdot \rho^{\frac{\lambda - n - (k-1+1)q}{q}}$$

and so (1)  $\rightarrow$  0 as  $\rho \rightarrow 0$ , since  $\lambda > n + kq$ . For (2), note that for any  $\sigma \in S_Q$ ,

$$(2)^{2} \leq \rho^{-2} \sum_{i} \left| a_{p}^{\sigma(i)}(x_{0} + \rho e_{j}, 2|\rho|) - a_{p}^{i}(x_{0}, 2|\rho|) - \rho v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0}) \right|^{2}$$

$$= \rho^{-2} \sum_{i} \left| D^{p} P_{x_{0} + \rho e_{j}, 2|\rho|}^{\sigma(i)}(x)|_{x=x_{0} + \rho e_{j}} - D^{p} P_{x_{0}, 2|\rho|}^{i}(x)|_{x=x_{0}} - \rho v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0}) \right|^{2}.$$

Now we can readily check that

$$D^{p}P_{x_{0}+\rho e_{j},2|\rho|}^{\sigma(i)}(x_{0}+\rho e_{j}) - D^{p}P_{x_{0}+\rho e_{j},2|\rho|}^{\sigma(i)}(x_{0}) = -\sum_{\ell=1}^{|k|-p} \frac{(-1)^{\ell}}{\ell!} a_{p+\ell e_{j}}^{\sigma(i)}(x_{0}+\rho e_{j},2|\rho|) \rho^{\ell}$$

and thus substituting this in and using the triangle inequality we get:

$$(2)^{2} \leq 2\rho^{-2} \sum_{i} \left| D^{p} [P_{x_{0} + \rho e_{j}, 2|\rho|}^{\sigma(i)} - P_{x_{0}, 2|\rho|}^{i}] |_{x=x_{0}} \right|^{2} + 2 \sum_{i} \left| v_{p+e_{j}}^{\sigma(i)}(x_{0}) + \sum_{\ell=1}^{k-|p|} \frac{(-1)^{\ell}}{\ell!} a_{p+\ell e_{j}}^{\sigma(i)}(x_{0} + \rho e_{j}, 2|\rho|) \rho^{\ell-1} \right|^{2}.$$

Now choose  $\sigma = \sigma_{\rho}$  such that the first sum here equals  $\mathcal{G}(D^{p}P_{x_{0}+\rho e_{j},2|\rho|}(x_{0}), D^{p}P_{x_{0},2|\rho|}(x_{0}))^{2}$ . Since  $|x_{0} + \rho e_{j} - x_{0}| = \rho$ , we can apply Lemma 2.1 and argue as in the proof of Lemma 2.3 to see that this first term  $\to 0$  as  $\rho \to 0$  (again using the fact that  $|p| \le k - 1$  so that we can absorb the  $\rho^{-2}$  factor and still get the decay). Moreover note that, since

$$D^{p}P_{x_{0}+\rho e_{j},2|\rho|}^{\sigma_{\rho}(i)}(x_{0}) - D^{p}P_{x_{0},2|\rho|}^{i}(x_{0}) = a_{p}^{\sigma_{\rho}(i)}(x_{0}+\rho e_{j},2|\rho|) - a_{p}^{i}(x_{0},2|\rho|) + \sum_{\ell=1}^{k-|p|} \frac{(-1)^{\ell}}{\ell!} a_{p+\ell e_{j}}^{\sigma_{\rho}(i)}(x_{0}+\rho e_{j},2|\rho|) \rho^{\ell}$$

we see that necessarily:

$$|v_p^{\sigma_\rho(i)}(x_0 + \rho e_j) - v_p^i(x_0)| \to 0.$$

Hence we see that

$$(2)^{2} \leq o(1) + 4 \sum_{i} \left| v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0}) - a_{p+e_{j}}^{\sigma_{\rho}(i)}(x_{0} + \rho e_{j}, 2|\rho|) \right|^{2}$$

$$\leq o(1) + 8 \sum_{i} \left| v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0}) - v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0} + \rho e_{j}) \right|^{2} + 8 \sum_{i} \left| v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0} + \rho e_{j}) - a_{p+e_{j}}^{\sigma_{\rho}(i)}(x_{0} + \rho e_{j}, 2|\rho|) \right|^{2}$$

and so if we choose  $\sigma_1 = \sigma_{\rho}$ , we know that this  $\to 0$ , since then by construction the second sum equals  $\mathcal{G}(v_{p+e_j}(x_0 + \rho e_j), a_{p+e_j}(x_0 + \rho e_j, 2|\rho|))^2$  which  $\to 0$ , and the first sum  $\to 0$  by the assumed continuity of  $v_{p+e_j}$ . Note that  $\sigma_1$  was arbitrary, so we can choose it dependent on  $\rho$ .

Finally for (3) we have by any  $\sigma \in S_Q$ ,

$$(3)^{2} \leq \rho^{-2} \sum_{i} \left| a_{p}^{i}(x_{0}, 2|\rho|) + \rho v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0}) - v_{p}^{\sigma(i)}(x_{0}) - \rho v_{p+e_{j}}^{\sigma(i)}(x_{0}) \right|^{2}$$

$$(\dagger) \qquad \leq 2\rho^{-2} \sum_{i} \left| a_{p}^{i}(x_{0}, 2|\rho|) - v_{p}^{\sigma(i)}(x_{0}) \right|^{2} + 2\sum_{i} |v_{p+e_{j}}^{\sigma_{1}(i)}(x_{0}) - v_{p+e_{j}}^{\sigma(i)}(x_{0})|^{2}.$$

In particular, we can re-run this argument to see that  $v_p$  is continuous at  $x_0$  in the  $e_j$  direction: indeed, if instead we just had  $L^i \equiv v_p^i(x_0)$ , and considered  $\mathcal{G}(v_p(x_0 + \rho e_j), L)$ , from the above we would have shown that, for each j,

$$\mathcal{G}(v_p(x_0 + \rho e_j), L) \le o(1) + \sum_i |a_p^i(x_0, 2|\rho|) - v_p^{\sigma(i)}(x_0)|^2$$

and so just taking  $\sigma = \mathrm{id}$ , we see this  $\to 0$ . Similarly we can show  $v_p$  is continuous at every point.

So now choosing  $\sigma = \sigma_1 (= \sigma_\rho)$  in (†), the second sum vanishes. So we have

$$(3)^{2} \le \rho^{-2} \sum_{i} \left| a_{p}^{i}(x_{0}, 2|\rho|) - v_{p}^{\sigma_{\rho}(i)}(x_{0}) \right|^{2}.$$

Now, for any sequence  $\rho_t \to 0$ , we can find a subsequence  $\rho_{t'}$  for such  $\sigma_{\rho_{t'}} \equiv \sigma'$  is constant. Hence since we know  $v_p$  is continuous, from  $(\star)$  we see that  $v_p^{\sigma'(i)}(x_0) = v_p^i(x_0)$  for all i. In particular we get along this subsequence,

$$(3)^{2} \leq \rho_{t'}^{-2} \sum_{i} |a_{p}^{i}(x_{0}, 2|\rho_{t'}|) - v_{p}^{i}(x_{0})|^{2} \leq \rho_{t'}^{-2} \mathcal{G}(a_{(k-1)}(x_{0}, 2|\rho_{t'}|), v_{(k-1)}(x_{0}))^{2}$$

which  $\to 0$  by Lemma 2.5. Thus we see that every sequence  $\rho_t \to 0$  has a further subsequence  $(\rho_{t'})_{t'}$  such that

$$\rho_{t'}^{-1}\mathcal{G}(v_p(x_0 + \rho_{t'}e_j), L(\rho_{t'})) \to 0$$

which, by elementary analysis, implies that we have  $\lim_{\rho\to 0} \rho^{-1} \mathcal{G}(v_p(x_0+\rho e_j), L(\rho)) = 0$ , i.e.  $v_p$  is differentiable at  $x_0$  with derivative given by  $D_j v_p^i = v_{p+e_j}^i$ , as required.

Combining everything so far, we have now shown:

Corollary 2.1. Suppose that  $\Omega \subset \mathbb{R}^n$  is a convex A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$  with  $\lambda > n + kq$ . Then  $v_{(0)} \in C^{k,\alpha}(\overline{\Omega}; \mathcal{A}_Q(\mathbb{R}^m))$ , where  $\alpha = \frac{\lambda - n - kq}{q}$ , and moreover for all  $|p| \leq k$  we have

$$D^p v_{(0)} = v_p$$

where  $v_{(0)} = v_{(0,0,\dots,0)}$ .

Remark. If for each  $x_0, \rho$  the polynomial  $P_{x_0,\rho}$  was of degree at most r < k, then we would have  $a_p(x_0, \rho) \equiv 0$  for all |p| > r, and so  $v_p \equiv 0$  for all |p| > r. In particular,  $D^{r+1}v_{(0)} \equiv 0$ , and so  $v_{(0)}$  is a polynomial of degree at most r.

Remark. If  $\lambda > n + (k+1)q$ , then  $\alpha > 1$ , and so  $D^k v_{(0)}$  is Hölder continuous with exponent > 1, implying that  $D^k v_{(0)} \equiv \text{constant}$ . Hence in this case  $v_{(0)}$  is a polynomial of degree at most k.

Next we prove a special case of Theorem 2.1, which we will use to prove the general result.

**Theorem 2.2.** Suppose  $\Omega$  is a convex A-weighted domain and  $u \in \mathcal{L}_k^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ . Suppose that  $\lambda > n + kq$ . Then  $u \in C^{k,\alpha}(\overline{\Omega}; \mathcal{A}_Q(\mathbb{R}^m))$ , with the estimate

$$[u]_{k,\alpha} \leq C \|u\|_{k,q,\lambda}$$

where  $C = C(n, m, k, q, Q, A, \lambda)$ .

*Proof.* Fix  $x_0 \in \Omega$  and let  $\rho > 0$ . Then by the triangle inequality and integrating over  $\Omega \cap B_{\rho}(x_0)$ , since  $\Omega$  is A-weighted we get for some constant C = C(q),

$$\mathcal{G}(a_{(0)}(x,\rho), u(x_0))^q \leq \underbrace{\frac{C(q)}{A\rho^n} \int_{\Omega \cap B_{\rho}(x_0)} \mathcal{G}(P_{x_0,\rho}(x), a_{(0)}(x_0,\rho))^q \, dx}_{(1)} + \underbrace{\frac{C(q)}{A\rho^n} \int_{\Omega \cap B_{\rho}(x_0)} \mathcal{G}(P_{x_0,\rho}(x), u(x))^q \, dx}_{(2)} + \underbrace{\frac{C(q)}{A\rho^n} \int_{\Omega \cap B_{\rho}(x_0)} \mathcal{G}(u(x), u(x_0))^q \, dx}_{(3)}.$$

For (1) notice that for all  $\rho < 1$ ,

$$\mathcal{G}(P_{x_0,\rho}(x), a_{(0)}(x_0, \rho))^2 \le \sum_{i} |P_{x_0,\rho}^i(x) - a_{(0)}^i(x_0, \rho)|^2 = \sum_{i} \left| \sum_{|p| \ne 0} \frac{a_p^i(x_0, \rho)}{p!} \cdot (x - x_0)^p \right|^2 \le K\rho^2$$

for some constant independent of  $\rho$ , since  $a_p^i$  converge uniformly and so are bounded. Hence we see that  $(1) \to 0$  as  $\rho \to 0$ . For (2) simply notice that  $(2) \le \frac{C(q)}{A\rho^n} \cdot \rho^{\lambda} \cdot |||u|||_{k,q,\lambda}^q \to 0$  since  $\lambda > n + kq$ . For the final term, for  $\rho$  sufficiently small we have  $B_{\rho}(x_0) \subset \Omega$ , and so (3)  $\to 0$  follows for  $\mathcal{H}^n$ -a.e.  $x \in \Omega$  from the Lebesgue differentiation theorem, Theorem 1.1.

Hence we see that for  $\mathcal{H}^n$ -a.e.  $x_0 \in \mathcal{H}^n$ ,  $\lim_{\rho \to 0} \mathcal{G}(a_{(0)}(x,\rho),u(x_0))^q = 0$  i.e., we have  $v_{(0)} = u$   $\mathcal{H}^n$ -a.e. in  $\Omega$ . Since  $v_{(0)} \in C^{k,\alpha}(\overline{\Omega};\mathcal{A}_Q(\mathbb{R}^m))$  from Corollary 2.1, this completes the proof.  $\square$ 

Finally, using Theorem 2.2 we are able to prove the full Campanato regularity theorem, Theorem 2.1.

Proof of Theorem 2.1. Clearly since we are taking an infimum over a larger set,  $\mathcal{L}^{q,\lambda}_{\ell}(\Omega;\mathcal{A}_Q(\mathbb{R}^m)) \subset \mathcal{L}^{q,\lambda}_k(\Omega;\mathcal{A}_Q(\mathbb{R}^m))$  and  $||u||_{k,q,\lambda} \leq ||u||_{\ell,q,\lambda}$ . Now take  $u \in \mathcal{L}^{q,\lambda}_k(\Omega;\mathcal{A}_Q(\mathbb{R}^m))$ , where  $\lambda,\ell,k$  obey the stated relations. Fix a multi-index p with  $\ell < |p| \leq k$ . Fixing  $\rho_0$ , we get from the argument in Lemma 2.4 that for any  $i \geq 1$  and  $x_0 \in \overline{\Omega}$ , since  $n + |p|q \geq \lambda$ ,

$$\begin{split} \mathcal{G}(a_p(x_0, \rho_0), a_p(x_0, 2^{-i}\rho_0)) &\leq C \|u\|_{k,q,\lambda} \sum_{j=0}^{i-1} 2^{j\left(\frac{n+|p|q-\lambda}{q}\right)} \cdot \rho_0^{\frac{\lambda-n-|p|q}{q}} \\ &= C \|u\|_{k,q,\lambda} \cdot \rho_0^{\frac{\lambda-n-|p|q}{q}} \cdot \frac{2^{i\left(\frac{n+|p|q-\lambda}{q}\right)} - 1}{2^{\frac{n+|p|q-\lambda}{q}} - 1} \\ &\leq \tilde{C} \|u\|_{k,q,\lambda} \cdot \left(\frac{\rho_0}{2^i}\right)^{\frac{\lambda-n-|p|q}{q}} \end{split}$$

where  $\tilde{C}$  can depend on |p|, although this will not matter. Now for any  $\rho < \rho_0$  we can find  $i \in \{0, 1, ...\}$  with  $\rho/\rho_0 \in [2^{-(i+1)}, 2^{-i})$ , and so by the above inequality with this i we have

$$\mathcal{G}(a_p(x_0, \rho_0), a_p(x_0, 2^{-i}\rho_0)) \le \tilde{C} \|u\|_{k,q,\lambda} \cdot \rho^{\frac{\lambda - n - |p|q}{q}}.$$

Hence we have from Lemma 2.1 that, estimating as in the proof of Lemma 2.5,

$$\mathcal{G}(a_p(x_0, \rho), a_p(x_0, 2^{-i}\rho_0))^q \le \frac{C_1}{\rho^{n+|p|q}} \int_{\Omega \cap B_\rho(x_0)} \mathcal{G}(P_{x_0, \rho}, P_{x_0, 2^{-i}\rho_0})^q \le C \|u\|_{k, q, \lambda}^q \cdot \left(\frac{\rho^{\lambda} + (2^{-i}\rho_0)^{\lambda}}{\rho^{n+|p|q}}\right)^{\frac{1}{2}} d\mu$$

i.e.

$$\mathcal{G}(a_p(x_0, \rho), a_p(x_0, 2^{-i}\rho_0)) \le C \|u\|_{k,q,\lambda} \rho^{\frac{\lambda - n - |p|q}{q}}.$$

Combining the last inequalities we see that for any  $\rho \leq \rho_0$ ,

$$|a_p(x_0, \rho)| \le \bar{C} ||u||_{k,q,\lambda} \cdot \rho^{\frac{\lambda - n - |p|q}{q}} + |a_p(x_0, \rho_0)|.$$

Now applying Lemma 2.1 with  $F = P_{x_0,\rho_0}$  and  $G \equiv 0$ , we have

$$|a_{p}(x_{0}, \rho_{0})|^{q} \leq \frac{C_{1}}{\rho_{0}^{n+|p|q}} \int_{\Omega \cap B_{\rho_{0}}(x_{0})} |P_{x_{0},\rho_{0}}|^{q}$$

$$\leq \frac{\tilde{C}}{\rho_{0}^{n+|p|q}} \left[ \int_{\Omega \cap B_{\rho_{0}}(x_{0})} \mathcal{G}(P_{x_{0},\rho_{0}}, u)^{q} + \int_{\Omega} |u|^{q} \right]$$

$$\leq \frac{\tilde{C}}{\rho_{0}^{n+|p|q}} \left[ |||u|||_{k,q,\lambda}^{q} \cdot \rho_{0}^{\lambda} + ||u||_{L^{q}(\Omega)}^{q} \right]$$

$$= \bar{C} ||u||_{k,q,\lambda}^{q}$$

where  $\overline{C}$  depends on p and  $\rho_0$ . Note that this is  $||u||_{k,q,\lambda}$  and not  $|||u||_{k,q,\lambda}$ . Hence we see that for any  $\rho < 1$ ,  $x_0 \in \overline{\Omega}$  and  $|p| > \ell$ , we have since  $\lambda < n + |p|q$ ,

$$|a_p(x_0, \rho_0)| \le C||u||_{k,q,\lambda} \cdot \rho^{\frac{\lambda - n - |p|q}{q}}.$$

Thus for all such  $\rho, x_0, p$  we see

$$|a_p(x_0, \rho)| \le C||u||_{k,q,\lambda} \cdot \rho^{\frac{\lambda - n - |p|q}{q}}$$

and so, for any  $x_0 \in \overline{\Omega}$  and  $\rho < 1$ ,

$$\inf_{P \in \mathcal{P}_{\ell}} \int_{\Omega \cap B_{\rho}(x_{0})} \mathcal{G}(u, P)^{q} \leq \int_{\Omega \cap B_{\rho}(x_{0})} \mathcal{G}\left(u, \sum_{|p| \leq \ell} \frac{a_{p}(x_{0}, \rho)}{p!} (x - x_{0})^{p}\right)^{q} \\
\leq 2^{q} \int_{\Omega \cap B_{\rho}(x_{0})} \mathcal{G}(u, P_{x_{0}, \rho})^{q} + 2^{q} \int_{\Omega \cap B_{\rho}(x_{0})} \mathcal{G}\left(P_{x_{0}, \rho}, \sum_{|p| \leq \ell} \frac{a_{p}(x_{0}, \rho)}{p!} \cdot (x - x_{0})^{p}\right)^{q} \\
\leq 2^{q} \|u\|_{k, q, \lambda}^{q} \cdot \rho^{\lambda} + 2^{q} \int_{\Omega \cap B_{\rho}(x_{0})} \left(\sum_{i} \left|\sum_{\ell < |p| \leq k} \frac{a_{p}(x_{0}, \rho)}{p!} \cdot (x - x_{0})^{p}\right|^{2}\right)^{q/2} \\
\leq 2^{q} \|u\|_{k, q, \lambda}^{q} \cdot \rho^{\lambda} + C \|u\|_{k, q, \lambda}^{q} \cdot \rho^{\lambda} \\
\leq \tilde{C} \|u\|_{k, q, \lambda}^{q} \rho^{\lambda}.$$

Now when  $\rho \geq 1$ , this bound is immediate because we know that

$$\inf_{P\in\mathcal{P}_{\ell}}\int_{\Omega\cap B_{\rho}(x_0)}\mathcal{G}(u,P)^q\leq \int_{\Omega\cap B_{\rho}(x_0)}\mathcal{G}(u,Q[0])^q\leq \|u\|_q^q\leq \|u\|_{k,q,\lambda}^q\rho^{\lambda}.$$

So hence we see that

$$|||u||_{\ell,q,\lambda}^q \le C||u||_{k,q,\lambda}^q$$

i.e. we see that  $u \in \mathcal{L}_{\ell}^{q,\lambda}(\Omega; \mathcal{A}_Q(\mathbb{R}^m))$ . Hence as  $\lambda > n + \ell q$  we can apply Theorem 2.2 to get the result.

- 2.1. Modifications relevant to minimal surface regularity theory. We now give two additional lemmas which are relevant for the regularity theory of stationary integral varifolds, which we shall be considering in Section 3. The reason behind needing these modifications is that often we will have certain integral expressions over balls which decay with the radius at a certain rate, but only for radii such that the ball does not intersect some bad set. One key example is a boundary point, where we may only have estimates in the interior and so cannot say anything about the boundary regularity unless we have some control of the integrals at the boundary. Another key example for us is interior branch points of multi-valued harmonic functions. Not having a uniform lower bound on the radii which we can apply the estimates to means we cannot immediately deduce that the functions lie in a certain Campanato space, and so cannot deduce any regularity near the bad set. However if the points on the bad set also have certain integral expressions which decay, and there is a uniform lower bound on the radii we can apply this to, we are able to deduce regularity up to the bad set so long as we can compare the integral expressions as the good points to those at the bad points. This is exactly the set up for the first case, Lemma 2.8, which we then use to prove a more general version, where there are multiple "layers" of bad points: these can be thought of as "very bad" points, "bad points", and "good points", and so forth.
- **Lemma 2.8.** Suppose  $\Omega \subset \mathbb{R}^n$  is a convex domain and  $u: \Omega \to \mathcal{A}_Q(\mathbb{R}^m)$ . Let  $\Gamma \subset \overline{\Omega}$  be a closed subset and let  $\Omega' \subset \Omega'' \subset \overline{\Omega}$  be open such that  $\overline{\Omega'} \subset \Omega''$  and  $\overline{\Omega''} \subset \Omega$  are compact. Suppose also  $\Omega'$  is a convex A-weighted domain. Suppose that there are numbers  $k \in \{0, 1, \ldots\}$ ,  $q \in [1, \infty)$ ,  $\beta_1, \beta_2, \beta \in (0, \infty)$ ,  $\mu \in (0, 1)$ , and  $\epsilon \in (0, 1/4)$  such that we have the following:
  - (I) For each  $x_0 \in \Gamma \cap \overline{\Omega''}$ , there exists  $P_{x_0} \in \mathcal{P}_k$  with  $\sup_{\Omega} |P_{x_0}| \leq \beta$  such that for all  $0 < \sigma \leq \rho/2 \leq \epsilon/2$ ,

$$\sigma^{-n-kq} \int_{B_{\sigma}(x_0) \cap \Omega} \mathcal{G}(u(x), P_{x_0}(x))^q \, dx \leq \beta_1 \left(\frac{\sigma}{\rho}\right)^{q\mu} \cdot \rho^{-n-kq} \int_{B_{\rho}(x_0) \cap \Omega} \mathcal{G}(u(x), P_{x_0}(x))^q \, dx;$$

(II) For each  $x_0 \in \overline{\Omega''} \setminus \Gamma$ , there exists  $P_{x_0} \in \mathcal{P}_k$  such that for every  $y \in \Gamma \cap \overline{\Omega''}$  and every  $0 < \sigma \le \rho/2 < \frac{1}{2} \min\{1/4, \operatorname{dist}(x_0, \Gamma)\},$ 

$$\sigma^{-n-kq} \int_{B_{\sigma}(x_0) \cap \Omega} \mathcal{G}(u(x), P_{x_0}(x))^q \, dx \le \beta_2 \left(\frac{\sigma}{\rho}\right)^{q\mu} \cdot \rho^{-n-kq} \int_{B_{\rho}(x_0) \cap \Omega} \mathcal{G}(u(x), P_y(x))^q \, dx.$$

Then  $u \in C^{k,\lambda}(\overline{\Omega'}; \mathcal{A}_Q(\mathbb{R}^m))$  for some  $\lambda = \lambda(n, m, k, q, Q, A, \beta_1, \beta_2, \epsilon, \mu, \operatorname{dist}(\Omega', \Omega''))$  with the estimate

$$||u||_{k,\lambda;\Omega'} \le C \left(\beta^q + \int_{\Omega''} |u|^q\right)^{1/q}$$

where  $C = C(n, m, k, q, Q, A, \beta_1, \beta_2, \epsilon, \operatorname{dist}(\Omega', \Omega'')).$ 

Remark. Unlike in [Wic14, Lemma 4.3], in (II) we only require the integral on the right-hand side to include those  $P \in \mathcal{P}_k$  which are of the form  $P = P_y$  for some  $y \in \Gamma \cap \overline{\Omega''}$ . Whilst this does not modify the proof, we need this weaker hypothesis later on to choose certain  $\Gamma$ .

*Proof.* Consider an arbitrary point  $z \in \overline{\Omega'}$  and  $\rho \in (0, \epsilon)$ . Fix  $\gamma < \frac{1}{2} \min\{1/4, \operatorname{dist}(\Omega', \Omega'')\}$  to be chosen later with the desired dependencies. Now if there is a point  $y \in \Gamma \cap \overline{B_{\gamma\rho}(z)}$ , then

we have  $y \in \Omega''$  and so from property (II),

$$(\gamma \rho)^{-n-kq} \int_{B_{\gamma \rho}(z) \cap \Omega} \mathcal{G}(u, P_y)^q \leq 2^{n+kq} (\gamma \rho + |z - y|)^{-n-kq} \int_{B_{\gamma \rho + |z - y|}(y) \cap \Omega} \mathcal{G}(u, P_y)^q$$

$$\leq 2^{n+kq} \beta_1 \left( \frac{\gamma \rho + |z - y|}{\rho - |z - y|} \right)^{q\mu} \cdot (\rho - |z - y|)^{-n-kq} \int_{B_{\rho - |z - y|}(y) \cap \Omega} \mathcal{G}(u, P_y)^q$$

$$\leq 4^{n+kq} \beta_1 \left( \frac{2\gamma}{1 - \gamma} \right)^{q\mu} \cdot \rho^{-n-kq} \int_{B_{\rho}(y) \cap \Omega} \mathcal{G}(u, P_y)^q.$$

So choosing  $\tilde{\gamma} = \tilde{\gamma}(n, k, q, \beta_1, \mu)$  such that  $4^{n+kq}\beta_1 \left(\frac{2\tilde{\gamma}}{1-\tilde{\gamma}}\right)^{q\mu} < \frac{1}{4}$ , we see that if  $\gamma = \frac{1}{2}\min\{\tilde{\gamma}, \epsilon\}$  we have

$$(2.1) \qquad (\gamma \rho)^{-n-kq} \int_{B_{\gamma \rho}(z) \cap \Omega} \mathcal{G}(u, P_y)^q \le \frac{1}{4} \cdot \rho^{-n-kq} \int_{B_{\rho}(y) \cap \Omega} \mathcal{G}(u, P_y)^q$$

for any  $z \in \overline{\Omega'}$  and  $\rho \in (0, \epsilon)$ , provided  $y \in \Gamma \cap \overline{B_{\gamma\rho}(z)}$ . If on the other hand  $\Gamma \cap \overline{B_{\gamma\rho}(z)} = \emptyset$ , then by (II) we have for any  $P \in \mathcal{P}_k$  of the form  $P = P_y$  for some  $y \in \Gamma \cap \overline{\Omega''}$  that, for any  $\sigma \in (0, 1/2]$ ,

$$(2.2) \qquad (\sigma \gamma \rho)^{-n-kq} \int_{B_{\sigma \gamma \rho}(z) \cap \Omega} \mathcal{G}(u, P_z)^q \leq \beta_2 \sigma^{q\mu} \cdot (\gamma \rho)^{-n-kq} \int_{B_{\gamma \rho}(z) \cap \Omega} \mathcal{G}(u, P)^q.$$

Now fix any  $z \in \overline{\Omega'} \setminus \Gamma$  with  $\operatorname{dist}(z,\Gamma) \leq \gamma$ . Let  $j_* \in \{1,2,\dots\}$  be minimal such that  $\Gamma \cap \overline{B_{\gamma^{j_*+1}}(z)} = \emptyset$ . Then we can apply (2.1) with  $\rho = \gamma, \gamma^2, \dots, \gamma^{j_*-1}$  to get that for each  $j = 2, \dots, j_*$  (if  $y_* \in \Gamma \cap \overline{\Omega''}$  is chosen such that  $|z - y_*| = \operatorname{dist}(z,\Gamma)$ ),

$$(2.3) \qquad (\gamma^{j})^{-n-kq} \int_{B_{\gamma^{j}}(z)\cap\Omega} \mathcal{G}(u, P_{y_{*}})^{q} \leq \frac{1}{4} \cdot (\gamma^{j-1})^{-n-kq} \int_{B_{\gamma^{j-1}}(y)\cap\Omega} \mathcal{G}(u, P_{y_{*}})^{q}$$

and applying (2.2) with  $\rho = \gamma^{j*}$  that, for each  $\sigma \in (0, 1/2]$  and  $P = P_y$  for  $y \in \Gamma \cap \overline{\Omega''}$ ,

$$(2.4) \quad (\sigma \gamma^{j_*+1})^{-n-kq} \int_{B_{\sigma \gamma^{j_*+1}}(z) \cap \Omega} \mathcal{G}(u, P_z)^q \leq \beta_2 \sigma^{q\mu} \cdot (\gamma^{j_*+1})^{-n-kq} \int_{B_{\gamma^{j_*+1}}(z) \cap \Omega} \mathcal{G}(u, P)^q.$$

Set  $P_* := P_{y_*}$ . Take  $\sigma = 1/2$ ,  $P = P_*$  in (2.4) and  $j = j_*$  in (2.3) to get (after iterating (2.3)):

$$\begin{split} &\left(\frac{1}{2}\gamma^{j_*+1}\right)^{-n-kq} \int_{B_{\frac{1}{2}\gamma^{j_*+1}}(z)\cap\Omega} \mathcal{G}(P_*,P_z)^q \\ &\leq 2^q \left(\frac{1}{2}\gamma^{j_*+1}\right)^{-n-kq} \int_{B_{\frac{1}{2}\gamma^{j_*+1}}(z)\cap\Omega} \mathcal{G}(u,P_z)^q + 2^q \left(\gamma/2\right)^{-n-kq} \cdot \left(\gamma^{j_*}\right)^{-n-kq} \int_{B_{\gamma^{j_*}}(z)\cap\Omega} \mathcal{G}(u,P_*)^q \\ &\leq C\beta_2(1/2)^{q\mu} (\gamma^{j_*+1})^{-n-kq} \int_{B_{\gamma^{j_*+1}}(z)\cap\Omega} \mathcal{G}(u,P_*)^q + C \cdot 4^{-(j_*-1)} \cdot \gamma^{-n-kq} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u,P_*)^q \\ &\leq C4^{-(j_*-1)} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u,P_*)^q \end{split}$$

where  $C = C(n, k, q, \gamma, \mu, \beta_2)$ . In the last inequality we have again used (2.3). Now since these are Q-valued polynomials of degree at most k, for each  $j = 1, 2, \ldots, j_*$  we have by substituting  $\tilde{x} = f(x) := \frac{1}{2} \gamma^{j^* + 1 - j} (x - z) + z$  in the integral (and noting that by convexity of  $\Omega$  we have  $f(\Omega) \subset \Omega$ ):

$$(\gamma^{j})^{-n-kq} \int_{B_{\gamma^{j}}(z)\cap\Omega} \mathcal{G}(P_{*}(x), P_{z}(x))^{q} dx$$

$$= (\gamma^{j})^{-n-kq} \int_{B_{\frac{1}{2}\gamma^{j_{*}+1}}(z)\cap\Omega} \mathcal{G}(P_{*}(f^{-1}(\tilde{x})), P_{z}(f^{-1}(\tilde{x})))^{q} \cdot \left(\frac{1}{2}\gamma^{j_{*}+1-j}\right)^{-n} d\tilde{x}$$

$$\leq (\gamma^{j})^{-n-kq} \cdot \left(\frac{1}{2}\gamma^{j_{*}+1-j}\right)^{-n-kq} \int_{B_{\frac{1}{2}\gamma^{j_{*}+1}}(z)\cap\Omega} \mathcal{G}(P_{*}(\tilde{x}), P_{z}(\tilde{x}))^{q} d\tilde{x}$$

$$\leq C4^{-(j_{*}-1)} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u, P_{*})^{q}$$

$$\leq C4^{-j} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u, P_{*})^{q}$$

for some  $C = C(n, k, q, \beta_1, \beta_2, \mu, \epsilon, \operatorname{dist}(\Omega', \Omega''))$  independent of j, i.e. for each  $j = 1, 2, \dots, j_*$  we have

(2.5) 
$$(\gamma^{j})^{-n-kq} \int_{B_{\gamma^{j}}(z)\cap\Omega} \mathcal{G}(P_{*}, P_{z})^{q} \leq C4^{-j} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u, P_{*})^{q}.$$

Using (2.5) with (2.3) we see that for each  $j = 1, ..., j_*$ ,

$$(2.6) \qquad (\gamma^j)^{-n-kq} \int_{B_{\gamma^j}(z)\cap\Omega} \mathcal{G}(u, P_z)^q \le C4^{-(j-1)} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u, P_*)^q.$$

Now let  $\rho \in (0, \gamma/2]$ . Then if we have  $\rho \leq \frac{1}{2}\gamma^{j_*+1}$ , then we can write  $\rho = \sigma\gamma^{j_*+1}$  for some  $\sigma \in (0, 1/2]$ , and then by (2.4) we have (choosing  $P = P_*$  as estimating as in the derivation of (2.5)):

$$\rho^{-n-kq} \int_{B_{\rho}(z)\cap\Omega} \mathcal{G}(u, P_z)^q \le C\sigma^{q\mu} 4^{-(j_*-1)} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u, P_*)^q$$
$$\le C(\sigma\gamma^{j_*})^{q\mu'} \int_{B_{\gamma}(y)\cap\Omega} \mathcal{G}(u, P_*)^q$$

where  $\mu' = \min\{\mu, q^{-1} \log_{\gamma}(1/4)\}$ : recall  $\gamma < 1/4$  which ensures  $\mu' \in (0, 1)$ ; thus we get for such  $\rho$ ,

(2.7) 
$$\rho^{-n-kq} \int_{B_{\Omega}(z) \cap \Omega} \mathcal{G}(u, P_z)^q \le C \rho^{q\mu} \int_{B_{\Omega}(u) \cap \Omega} \mathcal{G}(u, P_*)^q.$$

On the other hand, if  $\rho \in (0, \gamma/2]$  has  $\rho > \frac{1}{2}\gamma^{j_*+1}$ , then we can find  $j \in \{1, 2, \dots, j_*\}$  for which  $\gamma^{j+1} < 2\rho \le \gamma^j$ , and so by (2.6) we have

(2.8) 
$$\rho^{-n-kq} \int_{B_{\rho}(z) \cap \Omega} \mathcal{G}(u, P_z)^q \le C \rho^{q\lambda} \int_{B_{\gamma}(y) \cap \Omega} \mathcal{G}(u, P_*)^q$$

where  $\lambda = \log_{\gamma}(1/4) \in (0,1)$ . Combining (2.7) and (2.8) we see that for any  $z \in \overline{\Omega'} \backslash \Gamma$  with  $\operatorname{dist}(z,\Gamma) \leq \gamma$ , we have for all  $\rho \in (0,\gamma/2]$ , if  $\tilde{\mu} := \min\{\lambda,\mu'\}$ ,

(2.9) 
$$\rho^{-n-kq} \int_{B_{\rho}(z) \cap \Omega} \mathcal{G}(u, P_z)^q \le C \rho^{q\tilde{\mu}} \int_{B_{\gamma}(y) \cap \Omega} \mathcal{G}(u, P_*)^q.$$

In particular since  $\sup_{\Omega} |P_*| \leq \beta$  we have that

$$\rho^{-(n+q(k+\tilde{\mu}))} \int_{B_{\rho}(z)\cap\Omega} \mathcal{G}(u, P_z)^q \le C \left(\beta^q + \int_{\Omega''} |u|^q\right)$$

where now C also depends on m,Q. For  $z \in \overline{\Omega'} \setminus \Gamma$  with  $\operatorname{dist}(z,\Gamma) > \gamma$  the same inequality follows immediately from (II), for all  $\rho \in (0,\gamma/2]$ . If  $z \in \overline{\Omega'} \cap \Gamma$ , again the same inequality follows from (I). In all of this modifications to  $\tilde{\mu}$ , C, may need to be made but they have the same dependencies. Hence we see that  $u \in \mathcal{L}_k^{q,\tilde{\lambda}}(\Omega'; \mathcal{A}_Q(\mathbb{R}^m))$  where  $\tilde{\lambda} = n + q(k + \tilde{\mu})$ , and so we can apply Theorem 2.1 to conclude that  $u \in C^{k,\tilde{\lambda}}(\Omega'; \mathcal{A}_Q(\mathbb{R}^m))$ , with the desired estimate.

The following more general version of Lemma 2.8 will be necessary when there are two different types of "bad" points, which in our case will be boundary points and interior branch points for multi-valued harmonic functions.

**Lemma 2.9.** Suppose  $\Omega \subset \mathbb{R}^n$  is a convex domain and we have  $(Q_i)_{i=1}^N$ ,  $(m_i)_{i=1}^N \subset \mathbb{Z}_{\geq 1}$  and  $u_i: \Omega \to \mathcal{A}_{Q_i}(\mathbb{R}^{m_i})$  for  $i=1,\ldots,N$ . Suppose we have  $\Omega' \subset \subset \Omega'' \subset \subset \overline{\Omega}$ , with  $\Omega'$  a convex A-weighted domain. Suppose also that  $\Gamma \subset \overline{\Omega}$  is a closed, non-empty subset, and we have  $\Gamma_1,\ldots,\Gamma_N\subset \overline{\Omega}\setminus\Gamma$  such that  $\overline{\Gamma}_i\subset\Gamma\cup\Gamma_i$ . Suppose that there are numbers  $k\in\{0,1,2,\ldots\}$ ,  $q\in[1,\infty)$ ,  $\beta,\beta_0\in(0,\infty)$ ,  $(\beta_i)_{i=1}^N$ ,  $(\tilde{\beta}_i)_{i=1}^N\subset(0,\infty)^N$ ,  $\mu\in(0,1)$ ,  $\epsilon\in(0,1/4)$  and subsets  $\mathfrak{P}_0,\mathfrak{P}_i,\mathfrak{P}\subset\mathcal{P}_k$  with  $\sup_{\Omega}|P|\leq\beta$  for all  $P\in\mathfrak{P}_0$ , such that we have the following:

(I) For each  $x_0 \in \Gamma \cap \overline{\Omega''}$ , there exists  $P_{x_0}^1, \ldots, P_{x_0}^N \in \mathfrak{P}_0$  such that for all  $0 < \sigma \le \rho/2 \le \epsilon/2$  we have:

$$\sigma^{-n-kq} \int_{B_{\sigma}x_0 \cap \Omega} \sum_{i=1}^{N} \mathcal{G}(u_i, P_{x_0}^i)^q \le \beta_0 \left(\frac{\sigma}{\rho}\right)^{q\mu} \rho^{-n-kq} \int_{B_{\rho}(x_0) \cap \Omega} \sum_{i=1}^{N} \mathcal{G}(u_i, P_{x_0}^i)^q;$$

(II) For each  $i \in \{1, ..., N\}$ , we have for each  $x_1 \in \overline{\Omega''} \cap \Gamma_i \backslash \Gamma$  there exists  $P_{x_1}^i \in \mathfrak{P}_i$  such that for every  $P \in \mathfrak{P}_0$  and every  $0 < \sigma \le \rho/2 \le \frac{1}{2} \min\{1/4, \operatorname{dist}(x_1, \Gamma)\}$  we have:

$$\sigma^{-n-kq} \int_{B_{\sigma}(x_1) \cap \Omega} \mathcal{G}(u_i, P_{x_1}^i)^q \leq \beta_i \left(\frac{\sigma}{\rho}\right)^{q\mu} \rho^{-n-kq} \int_{B_{\sigma}(x_1) \cap \Omega} \mathcal{G}(u_i, P)^q;$$

(III) For each  $i \in \{1, ..., N\}$ , we have for each  $x_2 \in \overline{\Omega''} \setminus (\Gamma \cup \Gamma_i)$  there exists  $P_{x_2}^i \in \mathfrak{P}$  such that for every  $P \in \mathfrak{P}_0 \cup \mathfrak{P}_i$  and every  $0 < \sigma \le \rho/2 \le \frac{1}{2} \min\{1/4, \operatorname{dist}(x_2, \Gamma \cup \Gamma_i)\}$  we have:

$$\sigma^{-n-kq} \int_{B_{\sigma}(x_2) \cap \Omega} \mathcal{G}(u_i, P_{x_2}^i)^q \leq \tilde{\beta}_i \left(\frac{\sigma}{\rho}\right)^{q\mu} \rho^{-n-kq} \int_{B_{\rho}(x_2) \cap \Omega} \mathcal{G}(u_i, P)^q.$$

Then  $u_i \in C^{k,\lambda}(\overline{\Omega'}; \mathcal{A}_{Q_i}(\mathbb{R}^{m_i}))$  for each i, with the estimate

$$\|u_i\|_{k,\lambda;\overline{\Omega'}} \le C \left(\beta^q + \int_{\Omega''} \sum_{i=1}^N |u_i|^q\right)^{1/q}$$

where  $\lambda, C$  both depend on  $n, (m_i)_i, k, q, (Q_i)_i, N, A, \beta_0, (\beta_i)_i, (\tilde{\beta}_i)_i, \epsilon, \operatorname{dist}(\Omega', \Omega''))$ , and  $\lambda$  can also depend on  $\mu$ .

Remark. The proof will follow in the same way as Lemma 2.8: essentially whenever a point  $z \in \overline{\Omega'} \setminus (\Gamma \cup \Gamma_i)$  is either closer to  $\Gamma$  than  $\Gamma_i$ , or the distances have the same order, we can complete the proof in a similar manner to Lemma 2.8 using (I) and (III). The only issue comes when  $\Gamma_i$  is much closer in order to z than  $\Gamma$ , in which case we need to use (II) with (III). Then we can use (I) and (II) in the same manner as Lemma 2.8 to complete the proof.

Remark. In our applications, we will have  $\Omega = B_1(0) \cap H \subset \mathbb{R}^n$ , where  $H := \{x^1 > 0\}$  is a half-plane, and  $\Omega' = B_{1/8}(0) \cap H$ .  $\Gamma$  will be a subset of  $\partial H$  of a priori "bad" points, but which we do have good integral estimates for uniformly lower bounded radii. The  $\Gamma_i$  will then be the set of branch points of a multi-valued harmonic function: this will include some boundary points which are branch points of an odd reflection of the function, which will exist so long as we have  $C^1$  regularity on a neighbourhood of the boundary a priori. The condition on  $\overline{\Gamma}_i$  is to ensure that  $\Gamma \cup \Gamma_i$  is a closed set for each i, and so if  $z \in \overline{\Omega'} \setminus (\Gamma \cup \Gamma_i)$ , then  $\mathrm{dist}(z, \Gamma \cup \Gamma_i) > 0$ : this prevents us having a point on  $\partial H$  in the complement of  $\Gamma \cup \Gamma_i$  which is a limit point of interior branch points.

*Proof.* Note first that for each  $z \in \Gamma_i \backslash \Gamma$  we may repeat the argument seen in the proof of Lemma 2.8 (with only minor modifications to deal with the new form of (I)) to find that we can find  $\tilde{\mu}$  such that for all  $\rho > 0$ ,

(2.10) 
$$\rho^{-n-kq} \int_{B_{\rho}(z)\cap\Omega} \mathcal{G}(u_i, P_z^i)^q \le C \rho^{q\tilde{\mu}} \left(\beta^q + \int_{\Omega''} \sum_i |u_i|^q\right)$$

where  $\tilde{\mu}$ , C have all the allowed dependencies.

Now fix  $i \in \{1, ..., N\}$ . Suppose  $z \in \overline{\Omega'} \setminus (\Gamma \cup \Gamma_i)$ . Let  $\gamma < \frac{1}{2} \min\{1/4, \operatorname{dist}(\Omega', \Omega'')\}$  to be chosen later and let  $\rho \in (0, \epsilon)$ . Note that if  $\overline{B_{\gamma\rho}(z)} \cap \Gamma \neq \emptyset$ , then in the same way as Lemma 2.8, choosing  $y_{\Gamma} \in \Gamma$  with  $\operatorname{dist}(z, \Gamma) = |z - y_{\Gamma}|$ , we have for suitable  $\gamma$ ,

$$(\gamma \rho)^{-n-kq} \int_{B_{\gamma \rho}(z) \cap \Omega} \sum_{j} \mathcal{G}(u_j, P_{y_{\Gamma}}^j)^q \leq \frac{1}{4} \cdot \rho^{-n-kq} \int_{B_{\rho}(y_{\Gamma}) \cap \Omega} \sum_{j} \mathcal{G}(u_j, P_{y_{\Gamma}}^j)^q.$$

However, when  $B_{\gamma\rho}(z) \cap \Gamma = \emptyset$ , we cannot proceed as in the proof of Lemma 2.8, since we could have  $\mathcal{B}_{\gamma\rho}(z) \cap \Gamma_i \neq \emptyset$ , and so (III) is not applicable in the form we want, i.e. with a uniform lower bound on the radii we can apply it to.

Assume now dist $(z, \Gamma) < \gamma$ : if this is not true then (i) if dist $(z, \Gamma_i) < \gamma/4$ , we can follow the proof of Lemma 2.8 just with  $\Gamma_i$  in place of  $\Gamma$  to get the result at z, or (ii) If dist $(z, \Gamma_i) \ge \frac{\gamma/4}{4}$ , then we can just apply (III) to get the result at z. So choose  $j_{\Gamma}$  minimal such that  $B_{\gamma^{j_{\Gamma}+1}}(z) \cap \Gamma = \emptyset$ . Then in the same way as Lemma 2.8, for each  $j = 2, \ldots, j_{\Gamma}$  we have (choosing  $y_{\Gamma}$  as above and setting  $P_{\Gamma} := P_{y_{\Gamma}}$ )

$$(2.11) \qquad (\gamma^j)^{-n-kq} \int_{B_{\gamma^j}(z)\cap\Omega} \sum_{\ell} \mathcal{G}(u_\ell, P_\Gamma^\ell)^q \leq \frac{1}{4} \cdot (\gamma^{j-1})^{-n-kq} \int_{B_{\gamma^{j-1}}(y_\Gamma)\cap\Omega} \sum_{\ell} \mathcal{G}(u_\ell, P_\Gamma^\ell)^q.$$

Now, if  $\operatorname{dist}(z,\Gamma_i) \geq \gamma^{j_\Gamma+M}$ ,  $M \geq 3$  to be chosen (dependent only on the allowed parameters), then we can still follow the proof of Lemma 2.8 to get the result (as we get the same estimates on  $P_i^z$  and  $P_\Gamma^i$ , and we have a large enough region around z to apply (III)). So assume  $\operatorname{dist}(z,\Gamma_i) < \gamma^{j_\Gamma+M}$ , and let  $y_* \in \Gamma_i$  be such that  $|z-y_*| = \operatorname{dist}(z,\Gamma_i)$ ; note that since  $z \notin \Gamma_i \cup \Gamma = \overline{\Gamma} \cup \overline{\Gamma_i}$ , we have  $\operatorname{dist}(z,\Gamma_i) > 0$ . So let  $j_* \in \{0,1,2,\dots\}$  be minimal such that  $\overline{B_{\gamma^{j_\Gamma+M+j_*+1}}(z)} \cap \Gamma_i = \emptyset$ .

We know that  $\operatorname{dist}(y_i, \Gamma) > \gamma^{j_{\Gamma}+1} - \gamma^{j_{\Gamma}+M} \ge \gamma^{j_{\Gamma}+1} (1-\gamma^2)$ , meaning that we can apply (II) for all  $0 < \sigma \le \rho/2 \le \frac{1}{2} \gamma^{j_{\Gamma}+1} (1-\gamma^2)$ . Thus, since  $B_{\rho}(z) \subset B_{\rho+\gamma^{j_{\Gamma}+M}}(y_*)$ , for  $j=2,\ldots,M+j_*-1$ , we have from (II), in the same way as (2.1), for suitable  $\gamma$ ,

$$(2.12) (\gamma^{j_{\Gamma}+j+1})^{-n-kq} \int_{B_{\gamma^{j_{\Gamma}+j+1}}(z)\cap\Omega} \mathcal{G}(u_i, P_{y_*}^i)^q \leq \frac{1}{4} \cdot (\gamma^{j_{\Gamma}+j})^{-n-kq} \int_{B_{\gamma^{j_{\Gamma}+j}}(y_*)\cap\Omega} \mathcal{G}(u_i, P_{y_*}^i)^q.$$

Now since  $\overline{B_{\gamma^{j_{\Gamma}+M+j_{*}+1}}(z)} \cap \Gamma_{i} = \emptyset$ , we can apply (III) to get that, for any  $\sigma \in (0,1/2]$ ,

$$(\sigma\gamma^{j_{\Gamma}+M+j_{*}+1})^{-n-kq}\int_{B_{\sigma\gamma^{j_{\Gamma}+M+j_{*}-1}}(z)\cap\Omega}\mathcal{G}(u_{i},P_{z}^{i})^{q}\leq \tilde{\beta}_{i}\sigma^{q\mu}(\gamma^{j_{\Gamma}+M+j_{*}+1})^{-n-kq}\int_{B_{\gamma^{j_{\Gamma}+M+1+j_{*}}}(z)\cap\Omega}\mathcal{G}(u,P)^{q}.$$

Now following the proof of Lemma 2.8, we get that for each  $j = 2, ..., M + j_* - 1$ ,

$$(\gamma^{j_{\Gamma}+j+1})^{-n-kq} \int_{B_{\gamma^{j_{\Gamma}+j+1}}(z) \cap \Omega} \mathcal{G}(u_i, P_z^i)^q \leq C4^{-j} (\gamma^{j_{\Gamma}+2})^{-n-kq} \int_{B_{\gamma^{j_{\Gamma}+2}}(y_*) \cap \Omega} \mathcal{G}(u_i, P_{y_*}^i)^q.$$

Hence following the proof of Lemma 2.8 again, we see that for any  $\rho < \frac{1}{2}\gamma^{j_{\Gamma}+M+j_*+1}$  we have for some  $\mu' \in (0,1)$ ,

$$\rho^{-n-kq} \int_{B_{\rho}(z) \cap \Omega} \mathcal{G}(u_i, P_z^i)^q \le C \rho^{q\mu'} \cdot (\gamma^{j_{\Gamma}})^{-n-kq-q\mu'} \int_{B_{-j_{\Gamma}}(y_*) \cap \Omega} \mathcal{G}(u^i, P_{y_*}^i)^q.$$

Also, if  $\gamma^{j_{\Gamma}+j+1} < 2\rho \le \gamma^{j_{\Gamma}+j}$  holds for some  $j=3,\ldots,M+j_*-1$ , then again we have the same inequality, e.g. if M=4, we see that for any  $\rho < \frac{1}{2}\gamma^{j_{\Gamma}+3}$  we have

$$\rho^{-n-kq-q\mu'} \int_{B_{\rho}(z)\cap\Omega} \mathcal{G}(u_i, P_z^i)^q \le C(\gamma^{j_{\Gamma}})^{-n-kq-q\mu'} \int_{B_{\gamma^{j_{\Gamma}}}(y_*)\cap\Omega} \mathcal{G}(u^i, P_{y_*}^i)^q$$

from which the desired bound (for a suitable exponent) follows from (2.10).

Now suppose  $\rho \ge \frac{1}{2} \gamma^{j_{\Gamma}+3}$ . Then we simply have from inclusions, since  $|z-y_*| \le \gamma^{j_{\Gamma}+4} < \rho$ ,

$$\rho^{-n-kq-q\mu'} \int_{B_{\rho}(z)} \mathcal{G}(u_i, P_{y_*}^i)^q \leq 2^{n+kq+q\mu'} \left(\rho + |z-y_*|\right)^{-n-kq-q\mu'} \int_{B_{\rho+|z-y_*|}(y_*)\cap\Omega} \mathcal{G}(u_i, P_{y_*}^i)^q$$

which then applying (2.10) tells us that, for every  $\rho > 0$ ,

$$\rho^{-n-kq-q\mu'} \inf_{P \in \mathcal{P}_k} \int_{B_{\rho}(z)} \mathcal{G}(u_i, P)^q \le C \left( \beta^q + \int_{\Omega''} \sum_i |u_i|^q \right)$$

and so since  $z \in \overline{\Omega'} \setminus (\Gamma \cup \Gamma_i)$  as arbitrary, and since this inequality has already been established for  $z \in \overline{\Omega'} \cap (\Gamma \cup \Gamma_i)$ , we see that  $u_i \in \mathcal{L}_k^{\lambda,q}(\Omega'; \mathcal{A}_{Q_i}(\mathbb{R}^{m_i}))$ , as so applying Theorem 2.1 we are done.

## 3. Applications to Minimal Surface Theory

In this section we apply the results of Section 2 to classes of functions known as *proper blow-up classes*; such classes arise naturally in the regularity theory of stationary integral varifolds. For illustrations of this relation as it appears here, we recommend the reader to consult the sources [Wic14], [BK17], [Min21], [BKW21]; in particular, the general ideas originated in [Sim93] for the multiplicity one setting, and [Wic14] for the higher multiplicity setting.

Throughout this section we will just need the results of Section 2 when  $\Omega = B_1(0) \cap H \subset \mathbb{R}^{n+k}$ , where  $H := \{x^1 > 0, x^{n+1} = \cdots = x^{n+k} = 0\} \subset \mathbb{R}^n$  is a half-plane. This situation arises when considering blow-ups of sequences of stationary integral varifolds converging to either a higher multiplicity stationary non-flat classical cone, or converge to a higher multiplicity plane yet have smaller order excess relative to a sequence of classical cones (see [Min21] for an explanation of this terminology, and the sources above).

Remark. In codimension > 1 it is not possible to guarantee that the blow-up functions generated are  $C^{1,\alpha}$ , but only that they are generalised  $C^1$  regular (see [BKW21]). This is due to the presence of stationary integral varifolds which are only Lipschitz regular. However, as we shall see in this section, if they always are  $C^{1,\alpha}$  in the interior with suitable  $C^{1,\alpha}$  regularity properties, we will be able to deduce  $C^{1,\alpha}$  boundary regularity.

Remark. We do not discuss the case when  $\Omega = B_1(0) \subset \mathbb{R}^n$ , which arises in the study of interior branch point singularities and sequences of stationary integral varifolds converging to a higher multiplicity plane. The reason for this is that the blow-up class takes a different form, and the interior regularity is not known. In some cases this difficulty has been bypassed: the interested reader should consult [Wic14], [BKW21], and [Wic21]. In our setting we are assuming suitable interior regularity properties, which will be true when working inductively on the multiplicity, provided we understand the regularity of the varifolds near lower multiplicity planes.

3.1. Initial Properties of Multi-Valued Harmonic Functions. We first need to define a stronger notion of a multi-valued harmonic function; for the basic notions and properties of two-valued harmonic functions, which we mimic, see [SW16]. Fix  $U \subset \mathbb{R}^n$  open and a  $u: U \to \mathcal{A}_q(\mathbb{R}^k)$  a q-valued  $C^1$  function.

**Definition 3.1.** The branch set of u, denoted  $\mathcal{B}_u$ , is the set of points  $x \in U$  such that there is no  $\rho > 0$  with the property that on  $B_{\rho}(x) \cap U$  we have  $u|_{B_{\rho}(x)} = \sum_{i=1}^{q} \llbracket u_i \rrbracket$ , for some single-valued  $C^1$  functions  $u_i : B_{\rho}(x) \cap U \to \mathbb{R}$ .

**Definition 3.2.** We say u is harmonic if for each  $x \in U \setminus \mathcal{B}_u$ , there is a  $\rho > 0$  with the property that  $u|_{B_{\rho}(x)} = \sum_{i=1}^{q} \llbracket u_i \rrbracket$ , where each  $u_i : B_{\rho}(x) \to \mathbb{R}^k$  is a single-valued  $C^1$  harmonic function.

**Definition 3.3.** We shall say a q-valued function  $u: \Omega \to \mathcal{A}_q(\mathbb{R}^k)$  is a good q-valued harmonic function if  $u \in C^{1,1/q}(\Omega; \mathcal{A}_q(\mathbb{R}^k))$  and if  $\mathcal{B}_u$  denotes the branch set of u, then  $\dim_{\mathcal{H}}(\mathcal{B}_u) \leq n-2$  and moreover at every branch point  $x \in \mathcal{B}_u$  we have for every  $0 < \sigma \leq \rho < \operatorname{dist}(x, \partial\Omega)$ ,

$$\sigma^{-n} \int_{B_{\sigma}(x)} |u_s|^2 \le \left(\frac{\sigma}{\rho}\right)^{2(1+1/q)} \rho^{-n} \int_{B_{\rho}(x)} |u_s|^2$$

where  $u_s := u - u_a$  is the symmetric part of u and  $u_a := q^{-1} \sum_{i=1}^q u_i$  its average-part.

Remark. In the case q=2 we already know from [SW16] that any two-valued  $C^{1,\alpha}$  harmonic function,  $\alpha \in (0,1)$ , is a good two-valued harmonic function in the above sense. For  $q \geq 3$  it is natural to conjecture that the same claim holds, and indeed the fact of whether q-valued  $C^{1,\alpha}$  harmonic functions are always good in the above sense is a direction of work we intend to pursue.

Remark. If u is a good q-valued harmonic function, then since  $\dim_{\mathcal{H}}(\mathcal{B}_u) \leq n-2$  we automatically have that  $u_a$  is a smooth single-valued harmonic function on  $\Omega$ .

**Lemma 3.1** (Unique Continuation for Good q-Valued Harmonic Functions). Suppose  $U \subset \mathbb{R}^n$  is open and that  $u_1, u_2 \in C^{1,\alpha}(U; \mathcal{A}_q(\mathbb{R}^k))$  are good q-valued harmonic functions. Then if  $u_1, u_2$  agree on an open subset of U, then we must have  $u_1 \equiv u_2$  on U.

Proof. By definition of a good multi-valued harmonic function, we know that  $B := \mathcal{B}_{u_1} \cup \mathcal{B}_{u_2}$  has  $\dim_{\mathcal{H}}(B) \leq n-2$  and is closed in U. In particular  $U \setminus B$  is open and connected, and thus is path connected. It suffices to show that  $u_1 \equiv u_2$  on  $U \setminus B$ , since both are continuous and  $\overline{U \setminus B} = U$ . So suppose there is some  $x \in U \setminus B$  for which  $u_1(x) \neq u_2(x)$ . We can find a path  $\gamma : [0,1] \to U \setminus B$  connecting a point  $x_0$  to x, where  $x_0$  lies in the open set on which  $u_1 \equiv u_2$ ; clearly we can choose  $x_0 \in U \setminus B$ . Moreover since the image of  $\gamma$  is compact and thus closed, we can find some r > 0 for which  $B_r(\gamma([0,1])) \subset U \setminus B$ . Restricting to  $B_r(\gamma([0,1]))$ , we can thus reduce to the case where on U we have  $\mathcal{B}_{u_1} \cup \mathcal{B}_{u_2} = \emptyset$ .

So therefore without loss of generality we can assume  $u_1, u_2 : U \to \mathcal{A}_q(\mathbb{R}^k)$  have  $u_1 \equiv u_2$  on some open set, and  $\mathcal{B}_{u_1}, \mathcal{B}_{u_2} = \emptyset$ . But now the result follows immediately from the unique continuation principle of single-valued harmonic functions.

Finally we will also need the following classification of  $C^1$  homogeneous of degree one q-valued harmonic functions. This is known in the case q = 2 from [SW16], and it is natural to

conjecture it holds for all q; as mentioned above, we intend to pursue these questions in aa later work. For now, it will be an assumption we make.

Classification Hypothesis. Every q-valued symmetric harmonic function  $u : \mathbb{R}^N \setminus \{0\} \to \mathcal{A}_q(\mathbb{R}^k)$  which is  $C^1$  and homogeneous of degree one is necessarily linear.

Note that there is no assumption that the q-valued harmonic function is good in the above sense. We do however remark that we can prove this under an inductive-like assumption, and in particular can prove this in the case q = 3, using a rectifiability result in [KW13].

**Lemma 3.2.** Fix  $q \in \{3, 4, ...\}$ . Suppose we know that for Q = 2, 3, ..., q-1, every Q-valued harmonic  $\tilde{u}$  function on an open set in  $\mathbb{R}^n$  has that  $\mathcal{B}_u$  is countably (n-2)-rectifiable. Then every q-valued symmetric harmonic function  $u : \mathbb{R}^n \to \mathcal{A}_q(\mathbb{R}^k)$  which is  $C^1$  on  $\mathbb{R}^n \setminus \{0\}$  and homogeneous of degree one is necessarily linear.

Proof. We may assume without loss of generality that k=1 just by working with the components, as that is enough to prove the result here. Set  $\tilde{\mathcal{K}}_u:=\{x\in\mathbb{R}^n:u(x)=q[0]\}$  and  $Du(x)=q[0]\}\subset\mathcal{K}_u$ . On  $\mathbb{R}^n\backslash\tilde{\mathcal{K}}_u$  it can be seen that we can locally decompose u, i.e., in some neighbourhood of each point  $x\in\mathbb{R}^n\backslash\tilde{\mathcal{K}}_u$ , into two multi-valued harmonic functions, both with < q values. In particular, away from  $\tilde{\mathcal{K}}_u$  the set of points where u does not decompose into a sum of harmonic functions is by assumption (n-2)-rectifiable. In particular it has vanishing 2-capacity, and so for each  $\epsilon>0$  we may find a cut-off function  $\eta_\epsilon\in C_c^1(\mathbb{R}^n)$  such that  $\eta_\epsilon\equiv 1$  on an  $\epsilon$ -neighbourhood of the multiplicity < q branch set,  $0\leq \eta_\epsilon\leq 1$ ,  $\eta_\epsilon\to 0$   $\mathcal{H}^n$ -a.e. as  $\epsilon\to 0$ , and  $\int_{\mathbb{R}^n}|D\eta_\epsilon|^2\leq \epsilon^2$ .

Now following the proof of [SW16, Lemma 2.5], for any  $\delta > 0$  choose a smooth function  $\gamma_{\delta} : \mathbb{R} \to \mathbb{R}$  which is an odd function, is convex for  $t \geq 0$ , has  $\gamma_{\delta} \equiv 0$  on some neighbourhood of 0, and  $\gamma'_{\delta} \equiv 1$  for all  $t \geq \delta$ . Then notice that, if we write  $f_{\delta} := \gamma_{\delta}(D_{j}u_{i})$ , then  $f_{\delta}$  has the property that it has compact support on  $S^{n-1} \setminus \tilde{\mathcal{K}}_{u}$ . In particular, by the same argument as in [SW16, Lemma 2.5] we have:

$$\int_{S^{n-1}} \eta_{\epsilon} \cdot |\nabla_{S^{n-1}} \gamma_{\delta}(D_j u_i)|^2 \le -\int_{S^{n-1}} \eta_{\epsilon} \cdot [\Delta_{S^{n-1}}(D_j u_i)] \gamma_{\delta}(D_j u_i) + o(1) = o(1)$$

where here o(1) means a term which  $\to 0$  as  $\epsilon \to 0$ , and the last equality comes from the fact that Du is homogeneous of degree one. Thus taking  $\epsilon \to 0$ , followed by  $\delta \to 0$ , we see for each i, j, that  $D_j u_i$  is constant, and thus u is linear.

Remark. In our desired application, we will only have a  $C^1$  multi-valued harmonic function — we will not know whether it is a good multi-valued harmonic function or not, and we won't have any more regularity information.

3.2. **Proper Blow-Up Classes.** Throughout this section we fix  $N \in \{1, 2, ..., \}$  and integers  $(q_i)_{i=1}^N \subset \{1, 2, ...\}$ . Recall that  $\Omega = B_1(0) \cap H$ .

**Definition 3.4.** Fix integers  $q_1, \ldots, q_N \in \{1, 2, \ldots\}$ , and write  $q = (q_1, \ldots, q_N)$ . We say that  $\mathfrak{B}_q(\Omega)$  is a proper (half-plane) blow-up class if it obeys the following properties:

(31) Each element  $v = (v^1, \dots, v^N) \in \mathfrak{B}_q(\Omega)$  has  $v^i \in L^2(\Omega; \mathcal{A}_{q_i}(\mathbb{R}^k)) \cap W^{1,2}_{loc}(\Omega; \mathcal{A}_{q_i}(\mathbb{R}^k))$  for each  $i = 1, \dots, N$ ;

- (32) (Interior Regularity). If  $v \in \mathfrak{B}_q(\Omega)$  then  $v^i : \Omega \to \mathcal{A}_{q_i}(\mathbb{R}^k)$  is a good  $q_i$ -valued harmonic function for each  $i = 1, \ldots, N$ ;
- (**33**) (Boundary Estimates). If  $v \in \mathfrak{B}_q(\Omega)$  and  $z \in B_1(0) \cap \partial H$ , then for each  $\rho \in (0, \frac{3}{8}(1 |z|)]$  we have:

$$\int_{B_{\rho/2}(z)\cap\Omega} \sum_{i=1}^{N} \frac{|v^i(x) - \kappa^i(z)|^2}{|x - z|^{n+3/2}} \, \mathrm{d}x \le C\rho^{-n-3/2} \int_{B_{\rho}(z)\cap\Omega} \sum_{i=1}^{N} |v^i(x) - \kappa^i(z)|^2 \, \mathrm{d}x$$

where  $\kappa: B_1(0) \cap \partial H :\to (\mathbb{R}^k)^N$  is is a smooth, single-valued function, which obeys:

$$\sup_{B_{5/16}(0)\cap\partial H} |\kappa|^2 \le C \int_{B_{1/2}(0)\cap\Omega} \sum_i |v^i|^2$$

where  $C = C(\mathfrak{B}_q(\Omega));$ 

(34) (Hardt–Simon Inequality). For  $v \in \mathfrak{B}_q(\Omega)$  and  $z \in B_1(0) \cap \partial H$ , for each  $\rho \in (0, \frac{3}{8}(1 - |z|)]$  we have:

$$\int_{B_{\rho/2}(z)\cap\Omega}\sum_{i=1}^NR_z^{2-n}\left|\frac{\partial}{\partial R_z}\left(\frac{v^i-v_a^i(z)}{R_z}\right)\right|^2\leq C\rho^{-n-2}\int_{B_{\rho}(z)\cap\Omega}\sum_{i=1}^N|v^i-\ell_{v^i,z}|^2$$

where  $R_z(x) := |x - z|$  and  $\ell_{v^i,z}(x) := v_a^i(z) + (x - z) \cdot Dv_a^i(z)$  is the first-order linear approximation to the average-part  $v_a^i$  at z;

- ( $\mathfrak{B}5$ ) (Closure Properties). If  $v \in \mathfrak{B}_q(\Omega)$ , then:
  - $\begin{array}{ll} (\mathfrak{B}5\mathrm{I}) \ v_{z,\sigma}(\cdot) := \|v(z+\sigma(\cdot))\|_{L^2(\Omega)}^{-1} v(z+\sigma(\cdot)) \in \mathfrak{B}_q(\Omega) \ \text{for each} \ z \in B_1(0) \cap \partial H \ \text{and each} \\ \sigma \in (0, \frac{3}{8}(1-|z|)], \ \text{whenever} \ v \not\equiv 0 \ \text{in} \ B_\sigma(z) \cap \Omega; \end{array}$
  - $(\mathfrak{B}5II) \ \|v-\ell_v\|_{L^2(\Omega)}^{-1}(v-\ell_v) \in \mathfrak{B}_q(\Omega) \text{ whenever } v-\ell_v \not\equiv 0 \text{ in } \Omega, \text{ where } \ell_v = (\ell_{v^1,0},\dots,\ell_{v^N,0});$
- ( $\mathfrak{B}6$ ) (Compactness Property). If  $(v_m)_m \subset \mathfrak{B}_q(\Omega)$ , then there is a subsequence  $(m') \subset (m)$  and a function  $v \in \mathfrak{B}_q(\Omega)$  such that  $v_{m'} \to v$  strongly in  $L^2_{\text{loc}}(B_1(0) \cap \overline{H})$  and weakly in  $W^{1,2}_{\text{loc}}(\Omega)$ ;
- ( $\mathfrak{B}$ 7) ( $\epsilon$ -Regularity Property). There exists  $\alpha = \alpha(\mathfrak{B}_q(\Omega))$  and  $c = c(\mathfrak{B}_q(\Omega))$  such that the following is true: if  $\sigma \in (0,1]$ , then there exists  $\epsilon = \epsilon(\mathfrak{B}_q(\Omega), \sigma)$  such that whenever  $v \in \mathfrak{B}_q(\Omega)$  has  $v_a(0) = 0$ ,  $Dv_a(0) = 0$ ,  $||v||_{L^2(\Omega)} = 1$ , and  $v_i^* : H \to \mathcal{A}_{q_i}(\mathbb{R}^k)$  is such that for each i,  $(v_i^*)_a = 0$ ,  $v_i^*$  is a union of  $q_i$  half-planes with boundaries meeting along  $\partial H$ , and  $\int_{B_1(0)\cap H} |v^*|^2 \geq c$ , then if:

$$\int_{\Omega} \mathcal{G}(v, v^*)^2 < \epsilon$$

then  $v|_{B_{\sigma}(0)\cap\Omega} \in C^{1,\alpha}(B_{\sigma}(0)\cap\overline{H}).$ 

Remark. We expect such a class of functions to arise when performing either (i) a coarse blow-up procedure relative to a non-flat stationary classical cone, or (ii) a fine blow-up procedure, relative to a higher multiplicity non-flat classical cone (see [Min21]). In each case, the multiplicity on each half-hyperplane will be strictly smaller than the multiplicity at the origin of the cone, so one would expect an inductive approach to give rise to the above properties of  $\mathfrak{B}_q(\Omega)$ , as long as we have first understood the structure of the relevant varifolds near hyperplanes of strictly smaller multiplicity. In this setting, we have therefore inductively assumed we understand the interior regularity, and so the boundary regularity becomes the main problem. Slight modifications need to be made for the fine blow-up process however – see Section 3.4.

It is also possible that such classes of functions will arise when studying the boundary regularity of certain classes of stationary integral varifolds, and so the results of this section may have applications there.

We will prove the following theorem:

**Theorem 3.1.** For each  $q=(q_1,\ldots,q_N)$  as above, we have that  $\mathfrak{B}_q(\Omega)\subset C^{1,\mu}(B_{1/8}(0)\cap\overline{H})$  for some  $\mu=\mu(\mathfrak{B}_q(\Omega))$ . Moreover for each  $v\in\mathfrak{B}_q(\Omega)$  we have  $\|v\|_{1,\mu;B_{1/8}(0)\cap\overline{H}}\leq C\int_{B_{1/2}(0)\cap H}|v|^2$ .

Remark. With this regularity at the boundary, as the symmetric part  $v_s^i$  of each component has zero boundary values, we are able to perform an odd reflection across the boundary. This gives rise to a  $C^{1,\mu}$   $q_i$ -valued harmonic function on all of  $\mathbb{R}^n$ . Therefore if we know that the interior branch set of such functions is countably (n-2)-rectifiable (which we do in the  $q_i=2$  case, see [SW16], [KW13]) we can then deduce that the boundary branch set is in fact also countably (n-2)-rectifiable.

The proof will consist of three parts. Firstly we are able to use  $(\mathfrak{B}3)$  with Lemma 2.9 to deduce initial  $C^{0,\alpha}(B_{1/8}(0)\cap \overline{H})$  regularity; the exponent  $\lambda=n+3/2$  present in  $(\mathfrak{B}3)$  is however not enough to push this to  $C^1$  regularity. We will be able to use the larger exponent present in the Hardt–Simon inequality,  $(\mathfrak{B}4)$ , to push this regularity to  $C^{1,\alpha}$ . Indeed, we will first classify the homogeneous degree one elements of  $\mathfrak{B}_q(\Omega)$  by showing that they are indeed  $C^1(B_1(0)\cap \overline{H})$ , which is a sufficiently high regularity to perform a reflection across  $\partial H$  and apply our Classification Hypothesis. Once we have achieved this classification we will be able to combine it with  $(\mathfrak{B}4)$  in order to deduce that there is sufficiently high decay in the integral estimates to apply Lemma 2.9 in order to deduce the desired regularity.

Before starting the proof, we first make the following observation: there exists  $\epsilon = \epsilon(\mathfrak{B}_q(\Omega)) \in (0,1)$  such that if  $v \in \mathfrak{B}_q(\Omega)$  has  $v_a(0) = 0$  and  $Dv_a(0) = 0$ , then

(3.1) 
$$\epsilon \int_{\Omega} |v|^2 < \int_{\Omega \setminus B_{1/2}(0)} |v|^2.$$

Indeed if this were not true, for each  $\ell=1,2,\ldots$ , we could find  $v_\ell\in\mathfrak{B}_q(\Omega)$  with  $(v_\ell)_a(0)=0$  and  $D(v_\ell)_a(0)=0$  for which  $\frac{1}{\ell}\int_{\Omega}|v_\ell|^2>\int_{\Omega\backslash B_{1/2}}|v_\ell|^2$ . Setting  $w_\ell:=v_\ell/\|v_\ell\|_{L^2(\Omega)}$  (note  $w_\ell\in\mathfrak{B}_q(\Omega)$  by ( $\mathfrak{B}$ 5II), since  $(v_\ell)_a(0)=0$  and  $D(v_\ell)_a(0)=0$ ), we can then use ( $\mathfrak{B}$ 6) to pass to a subsequence such that  $w_\ell\to w\in\mathfrak{B}_q(\Omega)$ , where the convergence is strong in  $L^2_{\mathrm{loc}}(B_1(0)\cap\overline{H})$ . In particular since  $\|w_\ell\|_{L^2(\Omega)}=1$  we have  $\int_{B_{1/2}\cap H}|w_\ell|^2>1-1/\ell$ , and thus  $\int_{B_{1/2}\cap H}|w|^2=1$ .

But then we also see that  $\int_K |w|^2 = 0$  for each compact  $K \subset \Omega \backslash B_{1/2}(0)$ , and so  $w \equiv 0$  on  $\Omega \backslash B_{1/2}(0)$ . But then from ( $\mathfrak{B}2$ ), by the unique continuation property of good Q-valued harmonic functions (Lemma 3.1), since the 0 function is a good Q-valued harmonic function for each i, we would therefore have that  $w \equiv 0$  on  $\Omega$ , which contradicts  $\int_{B_{1/2} \cap H} |w|^2 = 1$ , proving (3.1).

3.3. **Proof of Theorem 3.1.** Fix  $q = (q_1, \ldots, q_N)$  throughout this section.

**Lemma 3.3** (Initial Boundary Regularity Estimate). There exists  $\alpha = \alpha(\mathfrak{B}_q(\Omega))$  for which  $\mathfrak{B}_q(\Omega) \subset C^{0,\alpha}(B_{1/8}(0) \cap \overline{H})$ , and moreover for each  $v \in \mathfrak{B}_q(\Omega)$  we have

$$||v||_{0,\alpha;B_{1/8}(0)\cap\overline{H}}^2 \le C \int_{B_{1/2}(0)\cap H} |v|^2$$

where  $C = C(\mathfrak{B}_q(\Omega))$  and  $|v|^2 = \sum_{i=1}^N |v^i|^2$ .

*Proof.* From ( $\mathfrak{B}3$ ) we get that, for each  $z \in B_{1/8}(0) \cap \partial H$  and any  $0 < \sigma \le \rho/2 \le 1/32$ , if we set  $v^i(z) := q_i[\![\kappa^i(z)]\!]$ :

$$\sigma^{-n-3/2} \int_{B_{\sigma}(z) \cap \Omega} \sum_{i} \mathcal{G}(v^{i}, v^{i}(z))^{2} \leq C \rho^{-n-3/2} \int_{B_{\rho}(z) \cap \Omega} \sum_{i} \mathcal{G}(v^{i}, v^{i}(z))^{2}.$$

Set  $\Gamma := B_{1/8}(0) \cap \partial H$  and  $\Gamma_i := \mathcal{B}_{v^i}$ , the branch set of  $v^i$ . Fix  $z \in B_{1/8}(0) \cap H$ . Then if  $z \in \Gamma_i$ , since for any Q-valued function w we have  $|w|^2 = |w_s|^2 + Q|w_a|^2$ , and since for any good Q-valued function we know  $w_a$  is harmonic, we thus get for all  $0 < \sigma \le \rho/2 < \frac{1}{2} \mathrm{dist}(z, \Gamma)$  that for any constant  $b \in \mathbb{R}$ :

$$\sigma^{-n} \int_{B_{\sigma}(z) \cap \Omega} \mathcal{G}(w, w_a(z))^2 = \sigma^{-n} \int_{B_{\sigma}(z)} |w - w_a(z)|^2$$

$$= \sigma^{-n} \int_{B_{\sigma}(z)} Q|w_a - w_a(z)|^2 + |w_s|^2$$

$$\leq C \left(\frac{\sigma}{\rho}\right)^2 \int_{B_{\rho}(z)} Q|w_a - b|^2 + \left(\frac{\sigma}{\rho}\right)^{2(1+1/q)} \int_{B_{\rho}(z)} |w_s|^2$$

$$= C \left(\frac{\sigma}{\rho}\right)^2 \int_{B_{\rho}(z) \cap \Omega} \mathcal{G}(w, b)^2.$$

Moreover if  $z \in B_{1/8}(0) \cap H \setminus (\Gamma_i \cup \Gamma)$ , then on some open set disjoint from  $\Gamma \cup \Gamma_i$  we can express  $v^i$  as q single-valued harmonic functions, for which a similar result holds. Thus we see that we can apply Lemma 2.9, along with the assumed bounds on the boundary values from  $(\mathfrak{B}3)$ , to get  $v^i \in C^{0,\alpha}(B_{1/8}(0) \cap \overline{H}, \mathcal{A}_{q_i}(\mathbb{R}))$  for each i, and

$$||v||_{0,\alpha;B_{1/8}(0)\cap\overline{H}}^2 \le C \int_{B_{1/2}(0)\cap H} |v|^2$$

as desired.  $\Box$ 

Remark. From Lemma 3.3 we see that the average-part of each  $v^i$  is a  $C^{0,\alpha}(B_{1/8}(0) \cap \overline{H}) \cap C^{2,\alpha}(B_{1/8}(0) \cap H)$  harmonic function, which from ( $\mathfrak{B}3$ ) has smooth boundary data. As a consequence of classical elliptic boundary regularity results (see [GT15] or [Mor66]) we have that  $v_a^i \in C^{1,\alpha}(B_{1/8}(0) \cap \overline{H})$  for each i.

**Lemma 3.4** (Classification of Homogeneous Degree One Elements). Suppose that  $v \in \mathfrak{B}_q(\Omega)$  is homogeneous of degree one on  $\Omega \backslash B_{1/4}$ , i.e.  $\frac{\partial v}{\partial R} = 0$ , where R = |x|. Then v is a linear function on  $\overline{\Omega}$ , and moreover the symmetric part  $v_a^i$  takes the form  $x \mapsto \sum_{j=1}^q \llbracket a_j^i x^1 \rrbracket$  for each i, where  $a_j^i \in \mathbb{R}^k$  have the property that  $\sum_j a_j^i = 0$ .

*Proof.* The proof will combine the argument of [Wic14, Proposition 4.2] and its adaptation in [BKW21], with a reflection principle in order to reduce the proof to the classification hypothesis.

Fix  $v \in \mathfrak{B}_q(\Omega)$  which is homogeneous of degree one on  $\Omega \backslash B_{1/4}(0)$ . First, note that the homogeneous degree one extension of  $v^i$  to all of  $\Omega$  is also a good  $q_i$ -valued harmonic function: hence by the unique continuation property, we have that v is homogeneous of degree one on all of  $B_1 \cap H$ . From the above remark, we know that for each i  $v_a^i$  is a  $C^{1,\alpha}(B_{1/8}(0) \cap \overline{H})$  homogeneous degree one harmonic function. In particular,  $Dv_a^i$  is homogeneous of degree zero and in  $C^{0,\alpha}(B_{1/8}(0) \cap \overline{H})$ , and so is constant along rays and continuous at the origin, which implies  $Dv_a^i$  must be constant. Hence  $v_a^i$  is linear for each i. So if  $v \equiv v_a$ , we are done.

So suppose  $v \not\equiv v_a$ . Applying ( $\mathfrak{B}5II$ ), since we know  $\ell_v^i = v_a^i$ , we may reduce to the setting where  $v_a \equiv 0$  and  $\|v\|_{L^2(\Omega)} = 1$ . In particular from ( $\mathfrak{B}3$ ) we must have  $v^i|_{B_1(0)\cap\partial H} \equiv q_i[0]$ . Also since  $v \in C^{0,\alpha}(B_{1/8}(0)\cap\overline{H})$  from Lemma 3.3 and v is homogeneous of degree one, we see that by extending v to all of  $\overline{H}$  by a homogeneous degree one extension, we can assume without loss of generality that  $v \in C^{0,\alpha}(\overline{H})$ , and  $v^i$  is a good  $q_i$ -valued harmonic function on H.

Now define for each i:

$$T_i(v) := \{ z \in \overline{H} : v^i(z+x) = v^i(x) \text{ for all } x \in \overline{H} \}$$

i.e.,  $T_i(v)$  is the set of points in  $\overline{H}$  for which  $v^i$  is translation invariant. Since  $v^i|_{\partial H} = q[0]$ , we must have  $T_i(v) \subset \partial H$  if  $v^i \not\equiv 0$ ; by assumption this is true for some i. Since v is homogeneous of degree one and continuous on  $\overline{H}$ , it is straightforward to check that  $T_i(v)$  must be a subspace of  $\partial H$  whenever  $v^i \not\equiv 0$ , and so in particular in this case  $\dim(T_i(v)) \leq n-1$ .

Set  $d_i(v) := \dim(T_i(v))$ , with  $d(v) = (d_i(v))_i$ . For  $d = (d_1, \ldots, d_N)$ , write  $\mathcal{H}_d$  for the set of  $v \in \mathfrak{B}_q(\Omega)$  for which  $d_i(v) = d_i$  for each i. First note that if  $d_i(v) \geq n-2$  for some i, then  $v^i$  is translation invariant along a subspace of  $\partial H$  of dimension at least n-2, meaning that it is independent of at least (n-2)-coordinates. Therefore in suitable coordinates we can write  $v^i(x^1,\ldots,x^n) = \tilde{v}(x^1,x^2)$ , where  $\tilde{v}$  is a good  $q_i$ -valued symmetric harmonic function on  $H_2 := \{(x^1,x^2) \in \mathbb{R}^2 : x^1 > 0\}$  which is  $C^{0,\alpha}$  on  $\overline{H_2}$ . Moreover from the definition of a good q-valued harmonic we know  $\dim_{\mathcal{H}}(\mathcal{B}_{v^i}) \leq n-2$ , which implies  $\mathcal{B}_{\tilde{v}} \cap H_2 = \emptyset$ , since otherwise from the homogeneity and translation invariance we would have  $\dim_{\mathcal{H}}(\mathcal{B}_{v^i}) \geq n-1$ . But then we would be able to write, on all of H since it is simply connected,  $\tilde{v} = \sum_{j=1}^{q_i} \llbracket \phi_j \rrbracket$ , where each  $\phi_j : H \to \mathbb{R}$  is a smooth harmonic function. Then since  $\phi_j|_{\partial H_2} = 0$  for each i, again by the boundary regularity theory of harmonic functions, this would imply  $\phi_j \in C^{\infty}(\overline{H_2})$ ,

implying as before that  $D\phi_j$  is constant and so  $\phi_j$  is linear for each j. The zero boundary values then imply that we must have  $\phi_j(x^1, x^2) = a_j x^1$  for some constant  $a_j \in \mathbb{R}$ , i.e. we have  $v^i = \sum_{j=1}^{q_i} \llbracket a_j x^1 \rrbracket$ , and so in particular  $d_i(v) = n-1$  (provided  $v^i \not\equiv 0$ ). Thus we see that if  $d_i(v) \geq n-2$  for each i, then  $v \in \mathcal{H}_{(n-1,n-1,\dots,n-1)}$  and v takes the form as claimed.

Now suppose that there is some  $d \neq (n-1, n-1, \ldots, n-1)$  for which  $\mathcal{H}_d \neq \emptyset$ . Without loss of generality we can choose d such that among all such  $d' \neq (n-1, n-1, \ldots, n-1)$  with  $\mathcal{H}_{d'} \neq \emptyset$ , we have  $\sum_i d_i$  maximal. We will derive a contradiction from this, meaning that the only  $\mathcal{H}_d$  which is non-empty is  $\mathcal{H}_{(n-1,n-1,\ldots,n-1)}$ , which proves the result.

For such a d let  $v \in \mathcal{H}_d$ . By the above we know that we can find some  $i_*$  with  $d_{i_*}(v) < n-2$ . Let K be any compact subset of  $\overline{H} \backslash T_{i_*}(v)$ . We claim that there exists  $\epsilon = \epsilon(v, K, \mathfrak{B}_q(\Omega)) \in (0, 1)$  such that at each  $z \in K \cap \partial H$  for which v is not  $C^{1,\alpha}$  on any neighbourhood of z in  $\overline{H}$  (here  $\alpha$  is as in  $(\mathfrak{B}7)$ , so is independent of v, K), and for each  $\rho \in (0, \epsilon]$ , we have the reverse Hardt-Simon inequality:

(3.2) 
$$\int_{H \cap B_{\rho}(z)} \sum_{i} R_{z}^{2-n} \left( \frac{\partial (v^{i}/R_{z})}{\partial R_{z}} \right)^{2} \ge \epsilon \rho^{-n-2} \int_{H \cap B_{\rho}(z)} \sum_{i} |v^{i}|^{2}.$$

We shall prove this by contradiction. Clearly if  $v \equiv 0$  on  $H \cap B_{\rho}(z)$ , then there is nothing to prove, so suppose  $v \not\equiv 0$  on any such  $H \cap B_{\rho}(z)$ . If this were false, then we could find points  $z, (z_{\ell})_{\ell}$  in  $K \cap \partial H$ ,  $(\epsilon_{\ell})_{\ell}$ , and radii  $(\rho_{\ell})_{\ell}$  with  $z_{\ell} \to z$ ,  $\epsilon_{\ell} \downarrow 0$ ,  $\rho_{\ell} \downarrow 0$ , with  $v \not\equiv 0$  in  $B_{\rho_{\ell}}(z_{\ell})$  for each  $\ell$ , v not  $C^{1,\alpha}$  on  $B_{\rho_{\ell}}(z_{\ell}) \cap \overline{H}$ , and

$$\int_{H \cap B_{\rho_{\ell}}(z_{\ell})} \sum_{i} R_{z_{\ell}}^{2-n} \left( \frac{\partial (v^{i}/R_{z_{\ell}})}{\partial R_{z_{\ell}}} \right)^{2} < \epsilon_{\ell} \rho_{\ell}^{-n-2} \int_{H \cap B_{\rho_{\ell}}(z_{\ell})} \sum_{i} |v^{i}|^{2}.$$

For each  $\ell$  set  $w_{\ell}(\cdot) := \|v(z_{\ell} + \rho_{\ell}(\cdot))\|_{L^{2}(\Omega)}^{-1} v(z_{\ell} + \rho_{\ell}(\cdot))$ . We know from ( $\mathfrak{B}5I$ ) that  $w_{\ell} \in \mathfrak{B}_{q}(\Omega)$  for each  $\ell$ . Thus by ( $\mathfrak{B}6$ ) we can pass to a subsequence to ensure that  $w_{\ell} \to w_{*} \in \mathfrak{B}_{q}(\Omega)$ , where the convergence is strongly in  $L^{2}_{\text{loc}}(B_{1}(0) \cap \overline{H})$  and weakly in  $W^{1,2}_{\text{loc}}(\Omega)$ . In particular the above inequality gives for each  $\ell$ ,

$$\int_{\Omega} \sum_{i} R^{2-n} \left( \frac{\partial (w_{\ell}^{i}/R)}{\partial R} \right)^{2} < \epsilon_{\ell}.$$

It follows from this and the local weak convergence in  $W^{1,2}$  that  $w_*$  is homogeneous of degree one on each set of the form  $\{x \in \Omega : \operatorname{dist}(x, \partial\Omega) \geq \epsilon\}$ , for  $\epsilon > 0$ , and thus  $w_*$  is homogeneous of degree one on  $\Omega$ . Moreover from Lemma 3.3 we know  $w_*$  is continuous on  $B_1(0) \cap \overline{H}$ , and so we see that  $w_*$  has a continuous homogeneous degree one extension to all of  $\overline{H}$ , which from the unique continuation property Lemma 3.1 and ( $\mathfrak{B}2$ ) we know is a good multi-valued harmonic function in H.

We now claim that  $w_* \not\equiv 0$ . To see this, for any Q-valued  $C^{1,\alpha}$  function f defined on any interval  $I = [a, b] \subset \mathbb{R}$  we have (see e.g. [DL10, Equation (1.2)]):

$$\mathcal{G}(f(a), f(b)) \le \int_a^b |Df|.$$

Now fix  $\omega \in S^{n-1} \cap H$  and  $s, r \in (1/4, 1)$  with s < r. Applying this inequality with  $f(t) := w_{\ell}^{i}(t\omega)/t$  and I = [s, r] for each i, we get

$$\mathcal{G}\left(r^{-1}w_{\ell}(r\omega), s^{-1}w_{\ell}(s\omega)\right) \leq \int_{s}^{r} \left| \frac{\partial(w_{\ell}(R\omega)/R)}{\partial R} \right| dR \leq \int_{1/4}^{1} \left| \frac{\partial(w_{\ell}(R\omega)/R)}{\partial R} \right| dR$$

which implies that

$$|w_{\ell}(r\omega)|^2 \le c \left( |w_{\ell}(s\omega)|^2 + \int_{1/4}^1 \left[ \frac{\partial (w_{\ell}(R\omega)/R)}{\partial R} \right]^2 \right)$$

where c = c(n), as  $r, s \ge 1/4$ . Integrating this now over  $\omega \in S^{n-1} \cap H$  we get

$$\int_{S^{n-1}\cap H} |w_{\ell}(r\omega)|^2 d\omega \le c \left( \int_{S^{n-1}\cap H} |w_{\ell}(s\omega)|^2 d\omega + \int_{\Omega \setminus B_{1/4}(0)} \left[ \frac{\partial (w_{\ell}/R)}{\partial R} \right]^2 \right).$$

Now multiply both sides by  $r^{n-1}$  and integrate over  $r \in (1/2, 1)$ , and then multiply the resulting inequality by  $s^{n-1}$  and integrate over  $s \in (1/4, 1/2)$  (this ensures s < r holds) to get

$$\int_{\Omega \setminus B_{1/2}(0)} |w_{\ell}|^2 \le c \left( \int_{\Omega \cap B_{1/2}(0) \setminus B_{1/4}(0)} |w_{\ell}|^2 + \int_{\Omega \setminus B_{1/4}(0)} R^{2-n} \left[ \frac{\partial (w_{\ell}/R)}{\partial R} \right]^2 \right)$$

(again, using the fact that  $R \in (1/4, 1)$  in the last integral). Then by (3.1) we get, for all  $\ell$ ,

$$\epsilon \le c \left( \int_{\Omega \cap B_{1/2}(0) \setminus B_{1/4}(0)} |w_{\ell}|^2 + \epsilon_{\ell} \right)$$

for some  $\epsilon$  independent of  $\ell$ . Thus from the strong  $L^2$  convergence in  $B_{1/2}(0) \cap \overline{H}$ , we see that  $\epsilon \leq c \int_{\Omega \setminus B_{1/4}(0)} |w_*|^2$ , and so we must have  $w_* \not\equiv 0$ .

From the definition of  $w_\ell$  is straightforward to see that for each i we have  $T_i(v) \subset T_i(w_*)$ . We now claim that  $z \in T_i(w_*)$  for each i. Indeed for each  $\ell$  and i write  $w_\ell^i = \sum_j \llbracket f_\ell^{i,j} \rrbracket$ , where the  $f_\ell^{i,j}$  are single-valued functions on  $\Omega$  with  $f_\ell^{i,q_i} \leq f_\ell^{i,q_i-1} \leq f_\ell^{i,1}$  for all  $\ell$  and i (this is the case when k=1: otherwise we just work component-wise). For notational simplicity we shall just fix i,j and write  $f_\ell^{i,j} \equiv f_\ell$ . Then by homogeneity, for each  $y \in \overline{H}$  and  $\sigma > 0$  we have:

$$\sigma^{-n} \int_{B_{\sigma}(y)\cap H} f_{\ell}(x+z) \, \mathrm{d}x = \delta_{\ell}^{-1} \sigma^{-n} \int_{B_{\sigma}(y)\cap H} v_{j}^{i}(z_{\ell} + \rho_{\ell}(x+z)) \, \mathrm{d}x$$

$$= (1+\rho_{\ell}) \delta_{\ell}^{-1} \sigma^{-n} \int_{B_{\sigma}(y)\cap H} v_{j}^{i}(z_{\ell} + (1+\rho_{\ell})^{-1} \rho_{\ell}(z-z_{\ell}) + (1+\rho_{\ell})^{-1} \rho_{\ell}x) \, \mathrm{d}x$$

$$= (1+\rho_{\ell})^{n+1} \delta_{\ell}^{-1} \sigma^{-n} \int_{B_{(1+\rho_{\ell})^{-1}\sigma}((1+\rho_{\ell})^{-1}(z-z_{\ell}+y))\cap H} v_{j}^{i}(z_{\ell} + \rho_{\ell}x) \, \mathrm{d}x$$

$$= (1+\rho_{\ell}) \cdot \left[ (1+\rho_{\ell})^{-1} \sigma \right]^{-n} \int_{B_{(1+\rho_{\ell})^{-1}\sigma}((1+\rho_{\ell})^{-1}(z-z_{\ell}+y))\cap H} f_{\ell}(x) \, \mathrm{d}x$$

where we have written  $\delta_{\ell} := \|v_{z_{\ell},\rho_{\ell}}\|_{L^{2}(\Omega)}$  for notational simplicity.

Now letting  $\ell \to \infty$ , using the strong convergence in  $L^2_{\text{loc}}(\overline{H})$ , and then letting  $\sigma \downarrow 0$  we can apply the Lebesgue differentiation theorem for single-valued functions to get that for  $\mathcal{H}^n$ -a.e.  $y \in \overline{H}$  we have  $w_*(y+z) = w_*(y)$ . Then from the continuity of  $w_*$  provided by Lemma 3.3 we get that this is true for every  $y \in \overline{H}$ , and so as i, j were arbitrary,  $z \in T_i(w_*)$  for each i.

In particular we see that  $z \in T_{i_*}(w_*)$ , and so  $d_{i_*}(w_*) > d_{i_*}(v)$ . Thus we have  $\sum_i d_i(w_*) > \sum_i d_i(v)$ , with  $w_* \not\equiv 0$ , and so recalling that d we chosen to be maximal with  $\mathcal{H}_d \neq \emptyset$  and  $d \neq (n-1,n-1,\ldots,n-1)$ , the only way we can avoid a contradiction is if  $w_* \in \mathcal{H}_{(n-1,n-1,\ldots,n-1)}$ , i.e. graph $(w_*)$  is a union of half-hyperplanes meeting along a common boundary of  $\partial H$ . But then since  $w_* \not\equiv 0$  and  $(w_*)_a \equiv 0$ , there must be some i with  $w_*^i$  consisting of at least two distinct half-hyperplanes. Thus we can apply  $(\mathfrak{B}7)$  to see that for all  $\ell$  sufficiently large, we must have that  $w_\ell$  is  $C^{1,\alpha}$  on  $B_1(0) \cap \overline{H}$ , i.e. v is  $C^{1,\alpha}$  on  $B_{\rho_\ell}(z_\ell) \cap \overline{H}$ , which gives a contradiction. So (3.2) must hold.

Combining (3.2) with ( $\mathfrak{B}4$ ) we see that for any  $z \in K \cap \partial H$  at which v is not  $C^{1,\alpha}$  on any neighbourhood of z in  $\overline{H}$ , and each  $\rho \in (0, \epsilon]$  we have

$$\int_{B_{\rho}(z)\cap H} \sum_{i} R_{z}^{2-n} \left( \frac{\partial (v^{i}/R_{z})}{\partial R_{z}} \right)^{2} \geq \frac{\epsilon}{C} \int_{B_{\rho/2}(z)\cap H} \sum_{i} R_{z}^{2-n} \left( \frac{\partial (v^{i}/R_{z})}{\partial R_{z}} \right)^{2}.$$

Set  $\theta := \epsilon/C$ , which depends only on  $v, K, \mathfrak{B}_q(\Omega)$ . Iterating this inequality for a fixed z with  $2^{-\ell}\rho$  in place of  $\rho$ , for  $\ell=1,2,\ldots$ , and then interpolating between these geometric scales in the usual way (see the proof of Lemma 2.8) we see that for any  $z \in K \cap \partial H$  at which v is not  $C^{1,\alpha}$  on a neighbourhood of z in  $\overline{H}$ , and any  $0 < \sigma \le \rho/2 \le \epsilon/2$ , there are constants  $\beta = \beta(v, K, \mathfrak{B}_q(\Omega)) \in (0, \infty)$  and  $\mu = \mu(v, K, \mathfrak{B}_q(\Omega)) \in (0, 1)$  independent of z for which

$$\int_{B_{\sigma}(z)\cap H} \sum_{i} R_{z}^{2-n} \left( \frac{\partial (v^{i}/R_{z})}{\partial R_{z}} \right)^{2} \leq \beta \left( \frac{\sigma}{\rho} \right)^{2\mu} \int_{B_{\rho}(z)\cap H} \sum_{i} R_{z}^{2-n} \left( \frac{\partial (v^{i}/R_{z})}{\partial R_{z}} \right)^{2}.$$

Combining this with ( $\mathfrak{B}4$ ) and (3.2) we see that for every such  $z \in K \cap \partial H$  at which v is not  $C^{1,\alpha}$  on a neighbourhood of z in  $\overline{H}$ , we have for every  $0 < \sigma \le \rho/2 \le \epsilon/4$ :

$$\sigma^{-n-2} \int_{B_{\sigma}(z) \cap H} \sum_{i} |v^{i}|^{2} \leq C \left(\frac{\sigma}{\rho}\right)^{2\mu} \rho^{-n-2} \int_{B_{\rho}(z) \cap H} \sum_{i} |v^{i}|^{2}$$

for some  $C=C(v,K,\mathfrak{B}_q(\Omega))\in (0,\infty)$  independent of z.

Now set  $\Gamma$  to be the set of  $z \in K \cap \partial H$  at which v is not  $C^{1,\alpha}$  in a neighbourhood of z in  $\overline{H}$ ; we know this is a closed subset of  $\overline{H}$  by definition. Now, at  $z \in K \cap \partial H$  at which v is  $C^{1,\alpha}$  in a neighbourhood of z in  $\overline{H}$ , since  $v|_{\partial H} \equiv 0$ , we are able to apply an odd reflection across  $\partial H$  to get a  $C^{1,\alpha}$  multi-valued harmonic function on an open ball (see below for a more precise argument of this). In particular, the branch set of this reflection must also have a closed, codimension at most two branch set. So define  $\Gamma_i$  to be the union of the interior branch set (in H) of  $v^i$ , along with any branch points on  $\partial H$  which occur in such a reflection process at points on the boundary when v is  $C^{1,\alpha}$  on a neighbourhood of  $\overline{H}$ . It is simple to then check that  $\overline{\Gamma}_i \subset \Gamma \cup \Gamma_i$ , as away from  $\partial H$  the interior branch set is closed, and so if the closure of

 $\Gamma_i$  includes a point on  $\partial H$ , either this is in  $\Gamma$  or if not then it must be a "boundary branch point" in the above sense, as the branch set of the reflection is still closed and coincides with the interior branch set on H. We can therefore apply Lemma 2.9 to this setting (remember that those points in  $\partial H \setminus (\Gamma \cup \Gamma_i)$  have the usual harmonic function decomposition locally), for convex A-weighted K, to get that v is  $C^{1,\tilde{\alpha}}$  on K for some  $\tilde{\alpha} = \tilde{\alpha}(v, K, \mathfrak{B}_q(\Omega))$ . So in particular, since K was arbitrary, we are able to deduce that  $v \in C^1(\overline{H} \setminus T_{i_*}(v))$ ; we aren't able to deduce  $C^{1,\alpha'}$  regularity for some fixed  $\alpha'$  since we could have  $\tilde{\alpha} \to 0$  for different K.

Now consider  $v^{i_*}$ : this is translation invariant along  $T_{i_*}(v)$ , and so after we quotient out by this subspace and rotate appropriately, we see that  $v^{i_*}$  is determined entirely by a good  $q_{i_*}$ -valued harmonic function f defined on  $H_N:=\{(x^1,\ldots,x^N):\in\mathbb{R}^N:x^1>0\}\subset\mathbb{R}^N$  for some  $N\geq 3$   $(N:=n-d_{i_*}(v))$ . We then know that f is  $C^1(\overline{H_N}\setminus\{0\})$ , and has  $f|_{\partial H_N}=q_{i_*}[\![0]\!]$ .

We now define the odd reflection  $F: \mathbb{R}^N \to \mathcal{A}_{q_{i_*}}(\mathbb{R}^k)$  of f by

$$F(x) := \begin{cases} f(x^1, x^2, \dots, x^N) & \text{if } x^1 \ge 0; \\ -f(-x^1, x^2, \dots, x^N) & \text{if } x^1 < 0. \end{cases}$$

Now since we know that f is  $C^1(\overline{H}\setminus\{0\})$ , we see that  $F \in C^1(\mathbb{R}^N\setminus\{0\})$ . Moreover F is a  $q_{i_*}$ -valued harmonic function: this is clear away from  $\{x^1=0\}$ , and on  $\{x^1=0\}$  if we have any point away from  $\mathcal{K}_F := \{x: F^i(x) = F^j(x) \text{ and } DF^i(x) = DF^j(x) \text{ for some } i \neq j\}$ , then locally about this point f will be given by  $q_{i_*}$  single-valued harmonic functions on a neighbourhood of this point in  $\overline{H_N}$ , and so the usual reflection principle for single-valued harmonic functions tells us that F is given by  $q_{i_*}$  single-valued harmonic functions on a neighbourhood.

Thus F is a homogeneous degree one  $C^1$   $q_{i_*}$ -harmonic function on  $\mathbb{R}^N \setminus \{0\}$ . Note that we do not know whether F is a  $good\ q_{i_*}$  harmonic function, as we do not know anything about the size of the branch set on  $\{x^1=0\}$ . But then from the Classification Hypothesis, we can conclude that F must be linear. However since  $F|_{\{x^1=0\}}=q_{i_*}[0]$ , we must have that F takes the form  $x\mapsto \sum_j [a_jx^1]$  for some  $a_i$ , which in turn implies that  $v^{i_*}(x)=\sum_j [a_jx^1]$ . But then this would mean that  $d_{i_*}(v)=n-1$ , providing the contradiction.

Remark. It is worth noting that the crucial difference between the multi-valued case and the single-valued case is the ability to reflect. The reflection principle for single-valued harmonic functions requires no assumption on the boundary regularity of the derivative, whilst in the multi-valued setting we first need to establish  $C^1$  regularity at the boundary before we can reflect to get a  $C^1$  function on the whole plane.

With the characterisation provided by Lemma 3.4, we can now prove Theorem 3.1.

*Proof of Theorem 3.1.* We will prove another reverse Hardt–Simon inequality, similar to that seen in the proof of Lemma 3.4.

We claim that there is  $\epsilon = \epsilon(\mathfrak{B}_q(\Omega))$  such that the following holds: if  $v \in \mathfrak{B}_q(\Omega)$  is such that v is not  $C^{1,\alpha}$  on a neighbourhood of 0 in  $\overline{H}$ , where  $\alpha$  is as in  $(\mathfrak{B}7)$ , then we have for all

 $\rho \in (0, 1/32]$ :

$$\int_{\Omega \cap B_{\rho}(z)} \sum_{i} R_{z}^{2-n} \left( \frac{\partial ((v^{i} - v^{i}(z))/R_{z})}{\partial R_{z}} \right)^{2} \ge \epsilon \rho^{-n-2} \int_{\Omega \cap B_{\rho}(z)} \sum_{i} |v^{i} - \ell_{v^{i},z}|^{2}.$$

Indeed, setting  $w := \|v(z + \rho(\cdot))\|_{L^2(\Omega)}^{-1} v(z + \rho(\cdot))$  and then  $\tilde{w} := \|w - \ell_w\|_{L^2(\Omega)}^{-1} (w - \ell_w)$ , which both belong to  $\mathfrak{B}_q(\Omega)$  by ( $\mathfrak{B}5I$ ) and ( $\mathfrak{B}5II$ ) respectively, it suffices to consider the case  $\rho = 1$ , z = 0, and  $v \in \mathfrak{B}_q(\Omega)$  with  $v_a(0) = 0$ ,  $Dv_a(0) = 0$ , i.e. it suffices to prove for any such v,

$$\int_{\Omega} \sum_{i} R^{2-n} \left( \frac{\partial (v^{i}/R)}{\partial R} \right)^{2} \ge \epsilon \int_{\Omega} \sum_{i} |v^{i}|^{2}.$$

Indeed if this was not true, we could find sequences  $(v_{\ell})_{\ell}$  of such v with

$$\int_{\Omega} \sum_{i} R^{2-n} \left( \frac{\partial (v_{\ell}^{i}/R)}{\partial R} \right)^{2} < \frac{1}{\ell} \int_{\Omega} \sum_{i} |v_{\ell}^{i}|^{2}.$$

Setting  $w_{\ell} := \|v_{\ell}\|_{L^2(\Omega)}^{-1} v_{\ell}$ , as  $w_{\ell} \in \mathfrak{B}_q(\Omega)$  by ( $\mathfrak{B}5II$ ), we may pass to a subsequence by ( $\mathfrak{B}6$ ) to find a limit  $w_{\ell} \to w_* \in \mathfrak{B}_q(\Omega)$ , which arguing in exactly the same way as Lemma 3.4, must be homogeneous of degree one, and so by Lemma 3.4 must be linear. Moreover in the same way as Lemma 3.4, using (3.1) we may prove that  $w_* \not\equiv 0$ . However as  $(w_*)_a(0) = 0$  and  $D(w_*)_a(0) = 0$ , we must have that  $(w_*)_a \equiv 0$ , and so  $w_* \not\equiv 0$  implies that we can apply ( $\mathfrak{B}7$ ) to get that the  $w_{\ell}$  must be  $C^{1,\alpha}$  on a neighbourhood of 0 for all  $\ell$  sufficiently large, providing the contradiction.

Having established this reverse Hardt–Simon inequality, we may proceed in exactly the same way as Lemma 3.4, using Lemma 2.9 to deduce the desired regularity, thus completing the proof.

3.4. Modifications to Fine Blow-Up Classes. We wish to now make some remarks relevant to the setting in [Min21]. When performing a fine blow-up procedure, the class of functions which arise are determined by numerous multiplicity parameters, but also a closeness constant, M > 1 (see [Min21, Section 8]). Ideally we would like the class of fine blow-up functions to be a proper blow-up class, however this is no longer true because the class is not closed under the closure operations ( $\mathfrak{B}5$ ): indeed, the closeness parameter can change. Crucially, it turns out that instead of remaining in the same class, the operations in ( $\mathfrak{B}5$ ), in particular rescaling, instead cause the function to lie in another fine blow-up class, of a fixed larger closeness parameter. As all other properties other than ( $\mathfrak{B}5$ ) can still be established, we can still use the same proof as in Section 3.3 to conclude the desired regularity. Thus we make the following notion and observation:

**Definition 3.5.** Fix integers  $q_1, \ldots, q_N \in \{1, 2, \ldots, \}$  and write  $q = (q_1, \ldots, q_N)$ . We say that  $(\mathfrak{B}_{q;M}^F(\Omega))_{M \in (0,\infty)}$  is a proper (fine) blow-up family if for each  $M \in (0,\infty)$ ,  $\mathfrak{B}_{q;M}^F(\Omega)$  obeys properties  $(\mathfrak{B}1), (\mathfrak{B}2), (\mathfrak{B}3), (\mathfrak{B}4), (\mathfrak{B}6), (\mathfrak{B}7)$  of a proper blow-up class with constants  $C, \alpha$  independent of M, and there is a fixed constant  $M_0 = M_0 > 1$  independent of M such that if  $v \in \mathfrak{B}_{q;M}^F(\Omega)$  and  $v_*$  is one of the functions as in  $(\mathfrak{B}5)$  generated from v, then  $v_* \in \mathfrak{B}_{q;M_0}^F(\Omega)$ .

Then following the proof in Section 3.3, we still have:

**Theorem 3.2.** For each  $q = (q_1, \ldots, q_N)$ ,  $M \in (0, \infty)$ , we have that  $\mathfrak{B}_{q;M}^F(\Omega) \subset C^{1,\mu}(B_{1/8}(0) \cap \overline{H})$ , for some  $\mu \in (0,1)$  independent of M, and moreover given appropriate regularity theorems on multi-valued harmonic functions, the boundary branch set is countably (n-2)-rectifiable.

## ACKNOWLEDGEMENTS

This work was supported by the UK Engineering and Physical Sciences Research Council (EPSRC) grant EP/L016516/1 for the University of Cambridge Centre for Doctoral Training, the Cambridge Centre for Analysis.

#### References

- [AJ00] Frederick Almgren Jr. Almgren's big regularity paper: Q-valued functions minimizing dirichlet's integral and the regularity of area-minimizing rectifiable currents up to codimension 2, 2000.
- [BK17] Spencer Becker-Kahn. Transverse singularities of minimal two-valued graphs in arbitrary codimension. *Journal of Differential Geometry*, 107(2):241–325, 2017.
- [BKW21] Spencer Becker-Kahn and Neshan Wickramasekera. A regularity theorem for stationary integral varifolds near multiplicity 2 planes. 2021. Manuscript in preparation.
- [Cam64] Sergio Campanato. Proprieta di una famiglia di spazi funzionali. Annali della Scuola Normale Superiore di Pisa-Classe di Scienze, 18(1):137–160, 1964.
- [DL10] Camillo De Lellis. Almgren's q-valued functions revisited. In Proceedings of the International Congress of Mathematicians 2010 (ICM 2010) (In 4 Volumes) Vol. I: Plenary Lectures and Ceremonies Vols. II–IV: Invited Lectures, pages 1910–1933. World Scientific, 2010.
- [DLS13] Camillo De Lellis and Emanuele Spadaro. Multiple valued functions and integral currents. arXiv preprint arXiv:1306.1188, 2013.
- [GT15] David Gilbarg and Neil S Trudinger. *Elliptic partial differential equations of second order*, volume 224. springer, 2015.
- [KW13] Brian Krummel and Neshan Wickramasekera. Fine properties of branch point singularities: two-valued harmonic functions. arXiv preprint arXiv:1311.0923, 2013.
- [Min21] Paul Minter. The structure of stable codimension one integral varifolds near classical cones of density  $\frac{5}{2}$ . arXiv preprint arXiv:2108.02614, 2021.
- [Mor66] Charles B Morrey. Multiple integrals in the calculus of variations. Springer, 1966.
- [RSS13] Humberto Rafeiro, Natasha Samko, and Stefan Samko. Morrey-campanato spaces: an overview. Operator theory, pseudo-differential equations, and mathematical physics, pages 293–323, 2013.
- [Sim93] Leon Simon. Cylindrical tangent cones and the singular set of minimal submanifolds. *Journal of Differential Geometry*, 38(3):585–652, 1993.
- [SW16] Leon Simon and Neshan Wickramasekera. A frequency function and singular set bounds for branched minimal immersions. Communications on Pure and Applied Mathematics, 69(7):1213–1258, 2016.
- [Wic08] Neshan Wickramasekera. A regularity and compactness theory for immersed stable minimal hypersurfaces of multiplicity at most 2. *Journal of Differential Geometry*, 80(1):79–173, 2008.
- [Wic14] Neshan Wickramasekera. A general regularity theory for stable codimension 1 integral varifolds. Annals of Mathematics, pages 843–1007, 2014.
- [Wic21] Neshan Wickramasekera. 2021. Personal communication.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE

Email address: pdtwm2@cam.ac.uk