

# On the relation between Gegenbauer polynomials and the Ferrers function of the first kind

Howard S. Cohl<sup>†</sup> and Roberto S. Costas-Santos<sup>§</sup>

<sup>†</sup> *Applied and Computational Mathematics Division, National Institute of Standards and Technology, Mission Viejo, CA 92694, USA*

URL: <http://www.nist.gov/itl/math/msg/howard-s-cohl.cfm>

E-mail: [howard.cohl@nist.gov](mailto:howard.cohl@nist.gov)

<sup>§</sup> *Dpto. de Física y Matemáticas, Universidad de Alcalá, c.p. 28871, Alcalá de Henares, Spain*

URL: <http://www.rscosan.com>

E-mail: [rscosa@gmail.com](mailto:rscosa@gmail.com)

Received August 10, 2021 in final form ????. Published online ????

doi:10.3842/JOURNAL.202\*.\*\*\*

**Abstract.** Using the direct relation between the Gegenbauer polynomials and the Ferrers function of the first kind, we compute interrelations between certain Jacobi polynomials, Meixner polynomials, and the Ferrers function of the first kind. We then compute Rodrigues-type and orthogonality relations for Ferrers functions of the first and second kinds. In the remainder of the paper using the relation between Gegenbauer polynomials and the Ferrers function of the first kind we derive connection and linearization relations, some definite integral and series expansions, some asymptotic expansions of Mehler-Heine type, Christoffel-Darboux summation formulas, and infinite series closure relations (Dirac delta distribution).

*Key words:* Ferrers functions; Gegenbauer polynomials; orthogonal polynomials

*2010 Mathematics Subject Classification:* 33C05; 33C15; 33C45; 42C10

## 1 Introduction

The generalized hypergeometric function [8, Chapter 16] is defined by the infinite series [8, (16.2.1)]

$${}_rF_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; z \right) := \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r)_k}{(b_1, \dots, b_s)_k} \frac{z^k}{k!}, \quad (1)$$

where  $b_i \notin -\mathbb{N}_0$ , for  $i = 1, \dots, s$ ; and elsewhere by analytic continuation. One interesting limit which we will use below is [12, (1.4.4)]

$$\lim_{\lambda \rightarrow \infty} {}_rF_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_{s-1}, \lambda b_s \end{matrix}; \lambda z \right) = {}_rF_{s-1} \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_{s-1} \end{matrix}; \frac{z}{b_s} \right). \quad (2)$$

The Pochhammer symbol for  $a \in \mathbb{C}$ ,  $n \in \mathbb{N}_0$  is given by [8, (5.2.4-5)]

$$(a)_n := (a)(a+1) \cdots (a+n-1).$$

The gamma function [8, Chapter 5] is related to the Pochhammer symbol, namely for  $a \in \mathbb{C} \setminus -\mathbb{N}_0$ , one has

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)},$$

which allows one to extend the definition to non-positive integer values of  $n$ . We will also use the common notational product convention, e.g.,

$$(a_1, \dots, a_r)_k := (a_1)_k (a_2)_k \cdots (a_r)_k.$$

The special case of  $r = 0$ ,  $s = 1$  in (1) is connected with the the Bessel function of the first kind [8, (10.16.9)]

$$J_\lambda(z) := \frac{1}{\Gamma(\lambda+1)} {}_0F_1\left(\begin{matrix} - \\ \lambda+1 \end{matrix}; -\frac{z^2}{4}\right). \quad (3)$$

Furthermore, the Hermite polynomials, related to the  $r = 2$ ,  $s = 0$  case, can be defined by [12, (9.15.1)]

$$H_n(x) := (2x)^n {}_2F_0\left(\begin{matrix} -\frac{n}{2}, \frac{1-n}{2} \\ - \end{matrix}; -\frac{1}{x^2}\right). \quad (4)$$

The Hermite polynomials are examples of generalized hypergeometric functions which are terminating, that is the infinite series (1) truncates at a finite value of  $k$ . In fact, all of the polynomials discussed in this paper will be defined in terms of a terminating generalized hypergeometric series.

The special case of  $r = 2$ ,  $s = 1$  is referred to as the Gauss hypergeometric series [8, Chapter 15]. Many of the functions used in this paper are defined in terms of the Gauss hypergeometric function. Therefore in this manuscript, we will require properties for this function. A useful asymptotic expansion for Gauss hypergeometric series is given as follows. [8, (15.12.2)], since  $a$  or  $b \in -\mathbb{N}_0$ , for example, if  $a = -m$ , then

$${}_2F_1\left(\begin{matrix} a, b \\ c \end{matrix}; z\right) = \sum_{k=0}^{m-1} \frac{(a, b)_k}{(c)_k} \frac{z^k}{k!} + \mathcal{O}\left(\frac{1}{c^m}\right), \quad (5)$$

as  $c \rightarrow \infty$ .

The Jacobi polynomial is defined as [8, (18.5.7)]

$$P_n^{(\alpha, \beta)}(x) := \frac{(\alpha+1)_n}{n!} {}_2F_1\left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; \frac{1-x}{2}\right), \quad (6)$$

and the Gegenbauer function is defined as [8, (15.9.15)]

$$C_\alpha^\lambda(z) := \frac{\Gamma(2\lambda+\alpha)}{\Gamma(2\lambda)\Gamma(\alpha+1)} {}_2F_1\left(\begin{matrix} -\alpha, 2\lambda+\alpha \\ \lambda+\frac{1}{2} \end{matrix}; \frac{1-z}{2}\right), \quad (7)$$

and the classical orthogonal Gegenbauer (ultraspherical) polynomial is given when  $\alpha \in \mathbb{N}_0$ , which makes the Gauss hypergeometric function terminating. Note that the Gegenbauer polynomial can be given in terms of the symmetric Jacobi polynomial as [8, (18.7.1)]

$$C_n^\lambda(x) = \frac{(2\lambda)_n}{(\lambda+\frac{1}{2})_n} P_n^{(\lambda-\frac{1}{2}, \lambda-\frac{1}{2})}(x). \quad (8)$$

Furthermore, there is also [8, (18.7.15)]

$$C_{2n}^\lambda(x) = \frac{(\lambda)_n}{(\frac{1}{2})_n} P_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(2x^2-1), \quad (9)$$

and [8, (18.7.16)]

$$C_{2n+1}^\lambda(x) = \frac{(\lambda)_{n+1}}{\left(\frac{1}{2}\right)_{n+1}} x P_n^{(\lambda-\frac{1}{2}, \frac{1}{2})}(2x^2 - 1). \quad (10)$$

Ferrers functions and Legendre functions are given in terms of Gauss hypergeometric functions which satisfy both linear and quadratic transformations. There are many such transformations and therefore there are many hypergeometric representations for the Gauss hypergeometric function. The Ferrers function of the first kind (associated Legendre function of the first kind on-the-cut)  $P_\nu^{-\mu} : (-1, 1) \rightarrow \mathbb{C}$  can for instance be defined as [8, (14.3.1)]

$$P_\nu^{-\mu}(x) = \frac{1}{\Gamma(\mu+1)} \left(\frac{1-x}{1+x}\right)^{\frac{\mu}{2}} {}_2F_1\left(\begin{matrix} -\nu, \nu+1 \\ \mu+1 \end{matrix}; \frac{1-x}{2}\right). \quad (11)$$

The Ferrers function of the second kind  $Q_\nu^\mu : (-1, 1) \rightarrow \mathbb{C}$  can also be defined in terms of Gauss hypergeometric representations. See for instance the recent publication [5] where all Gauss hypergeometric representations of the Ferrers functions of the second kind are given. The Ferrers function of the second kind (associated Legendre function of the second kind on-the-cut) for instance, is given in [8, (14.3.2)]

$$\begin{aligned} Q_\nu^\mu(x) := & \frac{\pi}{2 \sin(\pi\mu)} \left( \frac{\cos(\pi\mu)}{\Gamma(1-\mu)} \left(\frac{1+x}{1-x}\right)^{\frac{\mu}{2}} {}_2F_1\left(\begin{matrix} -\nu, \nu+1 \\ 1-\mu \end{matrix}; \frac{1-x}{2}\right) \right. \\ & \left. - \frac{\Gamma(\nu+\mu+1)}{\Gamma(\nu-\mu+1)} \left(\frac{1-x}{1+x}\right)^{\frac{\mu}{2}} \frac{1}{\Gamma(1+\mu)} {}_2F_1\left(\begin{matrix} -\nu, \nu+1 \\ 1+\mu \end{matrix}; \frac{1-x}{2}\right) \right), \quad (12) \end{aligned}$$

where  $\mu \notin \mathbb{Z}$ . However,  $Q_\nu^\mu$  can be analytically continued for  $\mu \in \mathbb{Z}$  which is demonstrated by [8, (14.3.12)]. Note that in this paper we will often indicate that the domain of the Ferrers function of the first and second kinds is  $(-1, 1)$ . But it should be emphasized that, in general and for specific formulas, this region can be analytically continued to a much larger region in the complex plane. It will often be convenient to express the Ferrers function of the first kind in terms of a Gauss hypergeometric series in a different way. One powerful such series representation is given as follows.

**Theorem 1.1.** *Let  $\nu, \mu \in \mathbb{C}$ ,  $x \in (-1, 1)$ . Then*

$$P_\nu^{-\mu}(x) = \frac{x^{\nu-\mu}(1-x^2)^{\frac{\mu}{2}}}{2^\mu \Gamma(\mu+1)} {}_2F_1\left(\begin{matrix} \frac{\mu-\nu}{2}, \frac{\mu-\nu+1}{2} \\ \mu+1 \end{matrix}; 1 - \frac{1}{x^2}\right). \quad (13)$$

**Proof.** Applying the connection relation [8, (14.9.1)]

$$P_\nu^{-\mu}(x) = -\frac{2}{\pi} \csc(\pi\mu) \frac{\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)} Q_\nu^\mu(x) + \frac{2}{\pi} \cot(\pi\mu) Q_\nu^{-\mu}(x),$$

to the Gauss hypergeometric representation of the Ferrers function of the second kind [3, (49)]

$$\begin{aligned} Q_\nu^\mu(x) = & \frac{2^{\mu-1} \cos(\pi\mu)}{(1-x^2)^{\frac{\mu}{2}}} \Gamma(\mu) x^{\nu+\mu} {}_2F_1\left(\begin{matrix} \frac{-\nu-\mu}{2}, \frac{-\nu-\mu+1}{2} \\ 1-\mu \end{matrix}; 1 - \frac{1}{x^2}\right) \\ & + \frac{\Gamma(\nu+\mu+1)\Gamma(-\mu)}{2^{\mu+1}\Gamma(\nu-\mu+1)} (1-x^2)^{\frac{\mu}{2}} x^{\nu-\mu} {}_2F_1\left(\begin{matrix} \frac{\mu-\nu}{2}, \frac{\mu-\nu+1}{2} \\ \mu+1 \end{matrix}; 1 - \frac{1}{x^2}\right), \end{aligned}$$

and after some simplification, one completes the proof. ■

One special case which we will encounter frequently below is as follows.

**Corollary 1.2.** *Let  $n \in \mathbb{N}_0$ ,  $x \in (-1, 1)$ . Then*

$$\mathbf{P}_{n+\lambda}^{-\lambda}(x) = \frac{x^n(1-x^2)^{\frac{\lambda}{2}}}{2^\lambda \Gamma(\lambda+1)} {}_2F_1\left(\begin{matrix} -\frac{n}{2}, \frac{1-n}{2} \\ 1+\lambda \end{matrix}; 1 - \frac{1}{x^2}\right) = \frac{n!}{(2\lambda+1)_n} \mathbf{P}_\lambda^{-\lambda}(x) C_n^{\lambda+\frac{1}{2}}(x). \quad (14)$$

**Proof.** Letting  $\mu \mapsto \lambda$ , and  $\nu \mapsto \lambda + n$  in (13) proves the first relation. The second identity holds due to the definition of the Gegenbauer function (7) for  $\alpha \in \mathbb{N}_0$  and taking into account [8, (14.5.18)]

$$\mathbf{P}_\lambda^{-\lambda}(x) = \frac{(1-x^2)^{\frac{\lambda}{2}}}{2^\lambda \Gamma(\lambda+1)}.$$

This completes the proof. ■

Some other special cases which are interesting and will prove to be useful include [8, (14.5.11)]

$$\mathbf{P}_\nu^{\frac{1}{2}}(\cos \theta) = \sqrt{\frac{2}{\pi \sin \theta}} \cos\left(\left(\nu + \frac{1}{2}\right)\theta\right), \quad (15)$$

$$\mathbf{P}_\nu^{-\frac{1}{2}}(\cos \theta) = \sqrt{\frac{2}{\pi \sin \theta}} \frac{\sin\left(\left(\nu + \frac{1}{2}\right)\theta\right)}{\nu + \frac{1}{2}}. \quad (16)$$

Using the connection relation [8, (14.9.1)], the Ferrers function of the first kind can be expressed in terms of the Ferrers function of the second kind, namely

$$\mathbf{P}_{k+n-\frac{1}{2}}^{\frac{1}{2}-n}(x) = \frac{2(-1)^n k!}{\pi(2n+k-1)!} \mathbf{Q}_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x). \quad (17)$$

The Ferrers function of the second kind has an interesting trigonometric special case for  $\mu = \frac{1}{2}$ , namely [8, (14.5.13)]

$$\mathbf{Q}_\nu^{\frac{1}{2}}(\cos \theta) = -\sqrt{\frac{\pi}{2 \sin \theta}} \sin\left(\left(\nu + \frac{1}{2}\right)\theta\right), \quad (18)$$

and for  $\mu = -\frac{1}{2}$ , namely [8, (14.5.14)]

$$\mathbf{Q}_\nu^{-\frac{1}{2}}(\cos \theta) = \sqrt{\frac{\pi}{2 \sin \theta}} \frac{\cos\left(\left(\nu + \frac{1}{2}\right)\theta\right)}{\nu + \frac{1}{2}}. \quad (19)$$

The Ferrers function of the first kind is related to the Gegenbauer function [8, (14.3.21)]

$$\begin{aligned} \mathbf{P}_\nu^{-\mu}(x) &= \frac{\Gamma(2\mu+1)\Gamma(\nu-\mu+1)}{2^\mu \Gamma(\nu+\mu+1)\Gamma(\mu+1)} (1-x^2)^{\frac{\mu}{2}} C_{\nu-\mu}^{\mu+\frac{1}{2}}(x) \\ &= \frac{\Gamma(2\mu+1)\Gamma(\nu-\mu+1)}{\Gamma(\nu+\mu+1)} \mathbf{P}_\mu^{-\mu}(x) C_{\nu-\mu}^{\mu+\frac{1}{2}}(x), \end{aligned} \quad (20)$$

where  $2\mu+1, \nu-\mu+1 \notin -\mathbb{N}_0$ . Equivalently

$$C_\lambda^\mu(x) = \frac{\sqrt{\pi} \Gamma(2\mu+1)}{2^{\mu-\frac{1}{2}} \Gamma(\mu) \Gamma(\lambda+1)} \frac{\mathbf{P}_{\lambda+\mu-\frac{1}{2}}^{\frac{1}{2}-\mu}(x)}{(1-x^2)^{\frac{\mu}{2}-\frac{1}{4}}} = \frac{\Gamma(2\mu+1)}{\Gamma(2\mu)\Gamma(\lambda+1)} \frac{\mathbf{P}_{\lambda+\mu-\frac{1}{2}}^{\frac{1}{2}-\mu}(x)}{\mathbf{P}_{\mu-\frac{1}{2}}^{\frac{1}{2}-\mu}(x)}, \quad (21)$$

or with  $\lambda = n \in \mathbb{N}_0$ , then

$$C_n^\mu(x) = \frac{\sqrt{\pi} \Gamma(2\mu + n)}{2^{\mu - \frac{1}{2}} \Gamma(\mu) n!} \frac{P_{n+\mu-\frac{1}{2}}^{\frac{1}{2}-\mu}(x)}{(1-x^2)^{\frac{\mu-1}{4}}}. \quad (22)$$

In order to compute Sobolev orthogonality which arises from Gegenbauer polynomials in terms of Ferrers functions we will require several particular interrelations between Gegenbauer polynomials in terms of Ferrers functions.

**Theorem 1.3.** *Let  $x \in (-1, 1)$ ,  $n \in \mathbb{N}_0$ ,  $N \in \mathbb{N}$  such that  $n \leq 2N - 1$ . Then*

$$C_n^{\frac{1}{2}-N}(x) = \frac{2^{N-1}}{\sqrt{\pi}} \Gamma(N + \frac{1}{2}) (1-x^2)^{\frac{N}{2}} \left( P_{n-N}^{-N}(x) + (-1)^n P_{n-N}^{-N}(-x) \right) \quad (23)$$

$$= \frac{2^N}{\sqrt{\pi}} \Gamma(N + \frac{1}{2}) (1-x^2)^{\frac{N}{2}} \left( P_{n-N}^{-N}(x) + \frac{(-1)^{n+N}}{n!(2N-n-1)!} Q_{n-N}^N(x) \right). \quad (24)$$

**Proof.** The connection relation [8, (14.9.7)] produces

$$P_{n+\lambda-\frac{1}{2}}^{\frac{1}{2}-\lambda}(x) = \frac{n!}{2 \sin(\pi\lambda) \Gamma(2\lambda + n)} \left( P_{n+\lambda-\frac{1}{2}}^{\lambda-\frac{1}{2}}(x) + (-1)^n P_{n+\lambda-\frac{1}{2}}^{\lambda-\frac{1}{2}}(-x) \right). \quad (25)$$

Applying (25) to (22) and substituting  $\lambda = \frac{1}{2} - N$  produces (23). The application of the connection relation [8, (14.9.9)] to the Ferrers function of the first kind with argument  $-x$  in (23) with simplification produces (24). This completes the proof. ■

**Theorem 1.4.** *Let  $x \in (-1, 1)$ ,  $n \in \mathbb{N}_0$ ,  $N \in \mathbb{N}$  such that  $n \geq 2N$ . Then,*

$$C_n^{\frac{1}{2}-N}(x) = \frac{2^N}{\sqrt{\pi}} \Gamma(N + \frac{1}{2}) \frac{(n-2N)!}{n!} (1-x^2)^{\frac{N}{2}} P_{n-N}^N(x). \quad (26)$$

**Proof.** Starting with (22) and substituting  $\lambda = \frac{1}{2} - N$  completes the proof. ■

Here we give interrelations between certain Jacobi polynomials and the Ferrers function of the first kind from Gegenbauer polynomials using a quadratic transformation.

**Corollary 1.5.** *Let  $n \in \mathbb{N}_0$ ,  $\lambda \in \mathbb{C}$ . Then*

$$P_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(2x^2-1) = \frac{2^\lambda \Gamma(\lambda+n+1)}{n!(1-x^2)^{\frac{\lambda}{2}}} P_{2n+\lambda}^{-\lambda}(x). \quad (27)$$

**Proof.** Start with (9) then inserting (22), and setting  $\lambda \mapsto \lambda + \frac{1}{2}$  and after simplification completes the proof. ■

**Corollary 1.6.** *Let  $n \in \mathbb{N}_0$ ,  $\lambda \in \mathbb{C}$ . Then*

$$P_n^{(\lambda, \frac{1}{2})}(2x^2 - 1) = \frac{2^\lambda \Gamma(\lambda + n + 1)}{n! x (1 - x^2)^{\frac{\lambda}{2}}} P_{2n+\lambda+1}^{-\lambda}(x). \quad (28)$$

**Proof.** Start with (10) then inserting (22), and setting  $\lambda \mapsto \lambda + \frac{1}{2}$ , and after simplification completes the proof.  $\blacksquare$

Below, the following related functions will be used. The associated Legendre function of the first kind  $P_\nu^\mu : \mathbb{C} \setminus (-\infty, 1] \rightarrow \mathbb{C}$  is defined as [8, (14.3.6)]

$$P_\nu^{-\mu}(z) = \frac{(z^2 - 1)^{\frac{\mu}{2}}}{2^\mu \Gamma(\mu + 1)} {}_2F_1\left(\begin{matrix} \nu + \mu + 1, -\nu + \mu \\ 1 + \mu \end{matrix}; \frac{1 - z}{2}\right). \quad (29)$$

Note that the Legendre polynomials are given by [8, (14.7.1), (18.7.9)]

$$P_n(x) = P_n^0(x) = P_n^0(x) = C_n^{\frac{1}{2}}(x). \quad (30)$$

**Remark 1.7.** *The Chebyshev polynomials of the first kind [12, (9.8.35)] can be given by*

$$T_n(\cos \theta) := \cos(n\theta) = \sqrt{\frac{\pi \sin \theta}{2}} P_{n-\frac{1}{2}}^{\frac{1}{2}}(\cos \theta) = n \sqrt{\frac{2 \sin \theta}{\pi}} Q_{n-\frac{1}{2}}^{-\frac{1}{2}}(\cos \theta), \quad (31)$$

and the Chebyshev polynomials of the second kind [12, (9.8.36)] can be given by

$$U_n(\cos \theta) := \frac{\sin((n+1)\theta)}{\sin \theta} = (n+1) \sqrt{\frac{\pi}{2 \sin \theta}} P_{n+\frac{1}{2}}^{-\frac{1}{2}}(\cos \theta) = -\sqrt{\frac{2}{\pi \sin \theta}} Q_{n+\frac{1}{2}}^{\frac{1}{2}}(\cos \theta), \quad (32)$$

where we have used (15), (16), (18), (19). See Remark 2.8 below with a concrete orthogonality example where the prefactor  $n$  cancels for the Chebyshev polynomials of the first kind, even for  $n = 0$ .

The associated Legendre function of the second kind  $Q_\nu^\mu : \mathbb{C} \setminus (-\infty, 1] \rightarrow \mathbb{C}$  can be defined in terms of the Gauss hypergeometric function as [8, (14.3.10) and Section 14.21]

$$Q_\nu^\mu(z) := \frac{\sqrt{\pi}(z^2 - 1)^{\mu/2}}{2^{\nu+1} \Gamma(\nu + \frac{3}{2}) z^{\nu+\mu+1}} {}_2F_1\left(\begin{matrix} \frac{\nu+\mu+1}{2}, \frac{\nu+\mu+2}{2} \\ \nu + \frac{3}{2} \end{matrix}; \frac{1}{z^2}\right), \quad (33)$$

for  $|z| > 1$  and, by analytic continuation of the Gauss hypergeometric function, elsewhere on  $z \in \mathbb{C} \setminus (-\infty, 1]$ . The normalized notation  $Q_\nu^\mu(z)$  is due to Olver [14, p. 178] and is defined in terms of the more commonly appearing Hobson notation for the associated Legendre function of the second kind  $Q_\nu^\mu(z)$  as [8, (14.3.10)]

$$Q_\nu^\mu(z) =: e^{i\pi\mu} \Gamma(\nu + \mu + 1) Q_\nu^\mu(z).$$

The Meixner polynomials are defined by [12, (9.10.1)]

$$M_n(x; \beta, c) := {}_2F_1\left(\begin{matrix} -n, -x \\ \beta \end{matrix}; 1 - \frac{1}{c}\right),$$

where  $n \in \mathbb{N}_0$ ,  $c \in \mathbb{C} \setminus \{0, 1\}$ . Specialized Meixner polynomials are related to the Ferrers function of the first kind.

**Corollary 1.8.** *Let  $n \in \mathbb{N}_0$ ,  $\lambda \in \mathbb{C}$ ,  $x \in (-1, 1)$ . Then*

$$P_{n+\lambda}^{-\lambda}(x) = \frac{(1-x^2)^{\frac{\lambda}{2}}}{2^\lambda \Gamma(\lambda+1)} M_n \left( -2\lambda - n - 1; \lambda + 1, \frac{2}{1+x} \right). \quad (34)$$

**Proof.** Start with the relation between the Meixner polynomials and the Jacobi polynomials [12, p. 236]

$$M_n(x; \beta, c) = \frac{n!}{(\beta)_n} P_n^{(\beta-1, -\beta-n-x)} \left( \frac{2-c}{c} \right),$$

and let  $x = 1 - n - 2\beta$ . This converts the Jacobi polynomial to symmetric form and then using (8), one obtains

$$C_n^\lambda(\omega) = \frac{(2\lambda)_n}{n!} M_n \left( -2\lambda - n; \lambda + \frac{1}{2}, \frac{2}{1+\omega} \right).$$

Finally, using (22), completes the proof. ■

## 1.1 Rodrigues-type relations

From the Gegenbauer polynomial Rodrigues-type relations, we can derive Rodrigues-type relations for the Ferrers function of the first and second kinds.

**Corollary 1.9.** *Let  $x \in (-1, 1)$ ,  $\mu \in \mathbb{C}$ ,  $n \in \mathbb{N}_0$ . Then,*

$$P_{n+\mu}^{-\mu}(x) = \frac{(-1)^n}{2^{\mu+n} \Gamma(\mu+n+1) (1-x^2)^{\frac{\mu}{2}}} \left( \frac{d}{dx} \right)^n \left[ (1-x^2)^{\mu+n} \right]. \quad (35)$$

**Proof.** Start with (20), setting  $\nu \rightarrow \mu + n$  and by using the Rodrigues-type formula for the Gegenbauer polynomials [8, (18.5.5)]

$$C_n^\mu(x) = \frac{(-1)^n (2\mu)_n}{(\mu + \frac{1}{2})_n 2^n n! (1-x^2)^{\mu-\frac{1}{2}}} \left( \frac{d}{dx} \right)^n \left[ (1-x^2)^{\mu+n-\frac{1}{2}} \right],$$

and after straightforward calculation completes the proof. ■

Similarly we have a Rodrigues-type relation for the Ferrers function of the second kind.

**Corollary 1.10.** *Let  $x \in (-1, 1)$ ,  $\mu \in \mathbb{C}$ ,  $n \in \mathbb{N}_0$ . Then,*

$$Q_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x) = \frac{\pi (-1)^{n+k} (2n+k-1)!}{2^{k+n+\frac{1}{2}} k! \Gamma(k+n+\frac{1}{2}) (1-x^2)^{\frac{n}{2}-\frac{1}{4}}} \left( \frac{d}{dx} \right)^k \left[ (1-x^2)^{k+n-\frac{1}{2}} \right]. \quad (36)$$

**Proof.** Start with (35), let  $n \mapsto k$ ,  $\mu \mapsto n - \frac{1}{2}$  then use (17), and after simplification, this completes the proof. ■

## 2 Orthogonality relations

In this section, we derive several new forms of orthogonality for Ferrers functions of the first and second kinds which are equivalent to the orthogonality of Gegenbauer (ultraspherical) polynomials. Due to this equivalence, it is surprising that these orthogonality relations have not been noticed previously. We will use these orthogonality conditions to verify our derived eigenfunctions for the Laplace-Beltrami operator on the  $d$ -dimensional  $R$ -radius hypersphere  $\mathbb{S}_R^d$ . For detailed information about the special functions we use below, namely the gamma function  $\Gamma : \mathbb{C} \setminus -\mathbb{N}_0 \rightarrow \mathbb{C}$ , Ferrers functions  $P_\nu^\mu$ ,  $Q_\nu^\mu$ , and the Gegenbauer polynomials  $C_n^\mu$ , and their properties, see [8, Chapters 5, 14, 18].

### 2.1 Continuous orthogonality from Gegenbauer polynomials

**Theorem 2.1.** *Let  $\mu \in \mathbb{C}$ ,  $k, k' \in \mathbb{N}$ ,  $\mu + k + \frac{1}{2} \neq 0$ ,  $\Re\mu > -1$ . Then,*

$$\int_{-1}^1 P_{k+\mu}^{-\mu}(x) P_{k'+\mu}^{-\mu}(x) dx = \frac{k!}{\Gamma(2\mu + k + 1)(\mu + k + \frac{1}{2})} \delta_{k,k'}. \quad (37)$$

**Proof.** Combining [8, (14.3.21)] (20) and orthogonality for the Gegenbauer polynomials [8, Sections 18.2-3]

$$\int_{-1}^1 C_n^\mu(x) C_{n'}^\mu(x) (1-x^2)^{\mu-\frac{1}{2}} dx = \frac{\pi \Gamma(2\mu + n)}{2^{2\mu-1} (\mu + n) n! \Gamma^2(\mu)} \delta_{n,n'}, \quad (38)$$

where  $n, n' \in \mathbb{N}_0$ ,  $\mu \in (-\frac{1}{2}, \infty) \setminus \{0\}$ , with  $\nu = k + \mu$  (resp.  $\nu = k' + \mu$ ) produces (37). The restriction on  $\Re\mu$  comes from ensuring that the singularities at  $x = \pm 1$  are integrable. This completes the proof. ■

Now specializing the above orthogonality relation using  $\mu$  as either an integer or an odd-half-integer produces other orthogonality relations as corollaries.

**Corollary 2.2.** *Let  $n, k, k' \in \mathbb{N}_0$ . Then*

$$\int_{-1}^1 P_{k+n}^n(x) P_{k'+n}^n(x) dx = \frac{(2n+k)!}{k!(n+k+\frac{1}{2})} \delta_{k,k'}. \quad (39)$$

**Proof.** Let  $\mu = n \in \mathbb{N}_0$  in (37), and the connection relation [8, (14.9.3)] expresses the Ferrers functions of the first kind with negative integer order in terms of Ferrers functions of the first kind with positive integer order. ■

Similarly we can derive orthogonality with non-positive integer order.

**Corollary 2.3.** *Let  $n, k, k' \in \mathbb{N}_0$ . Then*

$$\int_{-1}^1 P_{k+n}^{-n}(x) P_{k'+n}^{-n}(x) dx = \frac{k!}{(2n+k)!(n+k+\frac{1}{2})} \delta_{k,k'}. \quad (40)$$

**Proof.** Setting  $\mu = n \in \mathbb{N}_0$  in Theorem 2.1 the result follows. ■

**Corollary 2.4.** *Let  $n, k, k' \in \mathbb{N}_0$ . Then*

$$\int_{-1}^1 P_{k+n}^n(x) P_{k'+n}^{-n}(x) dx = \frac{(-1)^n}{n+k+\frac{1}{2}} \delta_{k,k'}. \quad (41)$$

**Proof.** Using the connection relation [8, (14.9.3)] once with (39) completes the proof. ■

**Remark 2.5.** *The orthogonality relations (37), (39) generalize [8, (14.17.6)],*

$$\int_{-1}^1 P_k^l(x) P_{k'}^l(x) dx = \frac{(k+l)!}{(k-l)!(k+\frac{1}{2})} \delta_{k,k'}, \quad (42)$$

*by setting  $n \mapsto n+l$ , and taking  $n=0$  in (39). Similarly, the orthogonality relation (41) generalizes [8, (14.17.7)],*

$$\int_{-1}^1 P_k^l(x) P_{k'}^{-l}(x) dx = \frac{(-1)^l}{k+\frac{1}{2}} \delta_{k,k'}, \quad (43)$$

*by setting  $n=0$ .*

One also has the following orthogonality relation for the Ferrers function of the second kind.

**Corollary 2.6.** *Let  $n, k, k' \in \mathbb{N}_0$ . Then*

$$\int_{-1}^1 Q_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x) Q_{k'+n-\frac{1}{2}}^{n-\frac{1}{2}}(x) dx = \frac{\pi^2(2n+k-1)!}{4(n+k)k!} \delta_{k,k'}. \quad (44)$$

**Proof.** Let  $\mu = n - \frac{1}{2}$ ,  $n \in \mathbb{N}_0$  in (37), and the connection relation [8, (14.9.1)] expresses the Ferrers functions of the first kind with negative integer order in terms of Ferrers functions of the second kind with positive integer order. ■

**Corollary 2.7.** *Let  $l, k, k' \in \mathbb{Z}$ ,  $l+k+1 \geq 0$ ,  $k+1 \neq 0$ ,  $l \geq -1$ . Then*

$$\int_{-1}^1 Q_{k+\frac{1}{2}}^{l+\frac{1}{2}}(x) Q_{k'+\frac{1}{2}}^{l+\frac{1}{2}}(x) dx = \frac{\pi^2(k+l+1)!}{4(k-l)!(k+1)} \delta_{k,k'}. \quad (45)$$

**Proof.** Specializing (44) with  $n=0$  completes the proof. ■

**Remark 2.8.** *For  $l \in \{-1, 0\}$ , (45) reduces to orthogonality for trigonometric functions, or Chebyshev polynomials of the first and second kinds. For  $l = -1$ , (45) reduces to orthogonality for Chebyshev polynomials of the first kind (31), namely [12, (9.8.37)]*

$$\int_0^\pi \cos(k\theta) \cos(k'\theta) d\theta = \int_{-1}^1 T_k(x) T_{k'}(x) \frac{dx}{\sqrt{1-x^2}} = \frac{\pi}{\epsilon_k} \delta_{k,k'}, \quad (46)$$

*where  $\epsilon_k = 2 - \delta_{k,0}$  is the Neumann factor. For  $l = 0$ , (45) reduces to*

$$\int_0^\pi \sin(k\theta) \sin(k'\theta) d\theta = \delta_{k,k'} \begin{cases} 0, & \text{if } k = 0, \\ \frac{\pi}{2}, & \text{if } k \geq 1, \end{cases} \quad (47)$$

*or for  $k \geq 0$ , [12, (9.8.38)]*

$$\int_{-1}^1 \sin((k+1)\theta) \sin((k'+1)\theta) d(\cos \theta) = \int_{-1}^1 U_k(x) U_{k'}(x) \sqrt{1-x^2} dx = \frac{\pi}{2} \delta_{k,k'}. \quad (48)$$

## 2.2 Sobolev orthogonality from Gegenbauer and Meixner polynomials

**Theorem 2.9.** *Let  $N, k, k' \in \mathbb{N}_0$ ,  $x \in (-1, 1)$ . Then the Ferrers function of the first kind fulfills the property of orthogonality*

$$\mathcal{B}(\mathbf{P}_k^N(x), \mathbf{P}_{k'}^N(x)) = \delta_{k,k'} \begin{cases} h_k^I, & \text{if } k < 2N, \\ h_k^{II}, & \text{if } k \geq 2N, \end{cases} \quad (49)$$

where

$$h_k^I = \frac{(\frac{1}{2}-N, 1-2N)_k}{k!(\frac{3}{2}-N)_k} = \frac{(-1)^k(2N-1)(2N-1)!}{k!(2N-2k-1)(2N-k-1)!}, \quad (50)$$

$$h_k^{II} = \frac{2k!(2N-1)!!^2}{(1+2k-2N)(k-2N)!} = \frac{2^{2N+1}k! \left(\frac{1}{2}\right)_N^2}{(1+2k-2N)(k-2N)!}, \quad (51)$$

$\mathcal{B}$  is the bilinear form defined as follows [15, Theorem 4.3]

$$\mathcal{B}(f, g) := \left( (1-x^2)^{\frac{N}{2}} f, (1-x^2)^{\frac{N}{2}} g \right)_1 + \int_{-1}^1 ((1-x^2)^{\frac{N}{2}} f)^{(2N)}(x) ((1-x^2)^{\frac{N}{2}} g)^{(2N)}(x) (1-x^2)^N dx, \quad (52)$$

where

$$(f, g)_1 := \sum_{k=0}^{N-1} \binom{N-1}{k} \frac{2^{k-1}}{(2-2N)_k} \left[ \left( \frac{d^k}{dx^k} f(x)g(x) \right)_{x=1} + (-1)^k \left( \frac{d^k}{dx^k} f(x)g(x) \right)_{x=-1} \right]. \quad (53)$$

**Proof.** The proof follows by using the identity (14), the recurrence relations for the Gegenbauer polynomials [12, (9.8.21)], and the property of orthogonality of the Gegenbauer polynomials. If  $k < 2N$ , then

$$\begin{aligned} h_k^I &= \left( \mathbf{P}_k^N(x)(1-x^2)^{\frac{N}{2}}, \mathbf{P}_k^N(x)(1-x^2)^{\frac{N}{2}} \right)_1 = \left( \frac{\sqrt{\pi}k!}{2^N \Gamma(N + \frac{1}{2})(k-2N)!} \right)^2 \left( C_k^{\frac{1}{2}-N}(x), C_k^{\frac{1}{2}-N}(x) \right)_1 \\ &= \left( \frac{\sqrt{\pi}k!}{2^N \Gamma(N + \frac{1}{2})(k-2N)!} \right)^2 \left( C_k^{\frac{1}{2}-N}(x), \frac{x(2k-1-2N)}{k} C_{k-1}^{\frac{1}{2}-N}(x) - \frac{k-1-2N}{k} C_{k-2}^{\frac{1}{2}-N}(x) \right)_1 \\ &= \frac{(2k-1-2N)}{k} \left( \frac{\sqrt{\pi}k!}{2^N \Gamma(N + \frac{1}{2})(k-2N)!} \right)^2 \left( x C_k^{\frac{1}{2}-N}(x), C_{k-1}^{\frac{1}{2}-N}(x) \right)_1 \\ &= \frac{(2k-1-2N)}{k} \frac{k-2N}{2k+1-2N} \left( \frac{\sqrt{\pi}k!}{2^N \Gamma(N + \frac{1}{2})(k-2N)!} \right)^2 \left( C_{k-1}^{\frac{1}{2}-N}(x), C_{k-1}^{\frac{1}{2}-N}(x) \right)_1. \end{aligned}$$

After some straightforward calculations we get the desired value. If  $k \geq 2N$ , due to the following factorization identity:

$$C_k^{\frac{1}{2}-N}(x) = (x^2-1)^N C_{k-2N}^{\frac{1}{2}+N}(x),$$

using (20) and after some algebraic manipulations one demonstrates

$$\left( \mathbf{P}_k^N(x)(1-x^2)^{\frac{N}{2}}, \mathbf{P}_k^N(x)(1-x^2)^{\frac{N}{2}} \right)_1 = 0.$$

Moreover, taking into account the identity [3, Remark 12]

$$\frac{d^n}{dx^n} \frac{P_\nu^{-\mu}(x)}{(1-x^2)^{\frac{\mu}{2}}} = (-1)^n (\nu + \mu + 1)_n (\mu - \nu)_n \frac{P_\nu^{-\mu-n}(x)}{(1-x^2)^{\frac{\mu+n}{2}}}, \quad (54)$$

then

$$\begin{aligned} h_k^{II} &= \int_{-1}^1 ((1-x^2)^{\frac{N}{2}} P_k^N(x))^{(2N)}(x) ((1-x^2)^{\frac{N}{2}} P_k^N(x))^{(2N)}(x) (1-x^2)^N dx \\ &= (k-N+1)_{2N}^2 (-N-k)_{2N}^2 \int_{-1}^1 p_k^{-N}(x) p_k^{-N}(x) (1-x^2)^{-N} dx \\ &= \frac{(k-N+1)_{2N}^2 (-N-k)_{2N}^2}{(2N+1)_{k-N}^2} \int_{-1}^1 C_{k-N}^{N+1/2}(x) C_{k-N}^{N+1/2}(x) dx. \end{aligned}$$

After some straightforward calculations we obtain the desired value. Hence the result follows.  $\blacksquare$

More details about the Gegenbauer/ultraspherical polynomials for non-classical parameters, their recurrence relations, and the Sobolev-type orthogonality can be found in [15].

### 3 Properties which follow from orthogonality and completeness

#### 3.1 Connection and linearization relations

Here we find the connection and linearization relations which arise from Gegenbauer polynomials for the Ferrers function of the first kind.

**Corollary 3.1.** *Let  $n \in \mathbb{N}_0$ ,  $\nu, \mu \in \mathbb{C}$ . Then*

$$\begin{aligned} P_{n+\nu}^{-\nu}(x) &= \frac{2^{\nu-\mu} n! (1-x^2)^{\frac{\nu-\mu}{2}}}{\Gamma(2\nu+2+n)} \\ &\times \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (\mu+n-2k+\frac{1}{2}) \frac{(\nu-\mu)_k \Gamma(\nu+\frac{1}{2}+n-k) \Gamma(2\mu+n-2k+1)}{k!(n-2k)! \Gamma(\mu+\frac{3}{2}+n-k)} P_{n-2k+\mu}^{-\mu}(x). \end{aligned} \quad (55)$$

**Proof.** The connection relation for Gegenbauer polynomials is given by [8, (18.18.16)]

$$C_n^\nu(x) = \frac{1}{\mu} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (\mu+n-2k) \frac{(\nu-\mu)_k (\nu)_{n-k}}{k! (\mu+1)_{n-k}} C_{n-2k}^\mu(x). \quad (56)$$

First replace the Gegenbauer polynomials in (56) using (22) then setting  $\lambda \mapsto \lambda + \frac{1}{2}$  completes the proof.  $\blacksquare$

**Corollary 3.2.** *Let  $n, m \in \mathbb{N}_0$ ,  $\lambda \in \mathbb{C}$ . Then*

$$\begin{aligned} P_{n+\lambda}^{-\lambda}(x) P_{m+\lambda}^{-\lambda}(x) &= \frac{2^\lambda \Gamma(\lambda + \frac{1}{2}) m! n! (1-x^2)^{\frac{\lambda}{2}}}{\sqrt{\pi} \Gamma(2\lambda+m+1) \Gamma(2\lambda+n+1)} \sum_{k=0}^m B_{k,m,n}^{\lambda+\frac{1}{2}} \frac{\Gamma(2\lambda+n+m-2k+1)}{(n+m-2k)!} P_{n+m-2k+\lambda}^{-\lambda}(x). \end{aligned} \quad (57)$$

**Proof.** The Gegenbauer polynomials have a linearization formula given by [8, (18.18.22)]

$$C_m^\lambda(x)C_n^\lambda(x) = \sum_{k=0}^m B_{k,m,n}^\lambda C_{m+n-2k}^\lambda(x), \quad (58)$$

where  $m, n \in \mathbb{N}_0$ ,  $n \geq m$ , and

$$B_{k,m,n}^\lambda := \frac{(m+n+\lambda-2k)(m+n-2k)!(\lambda)_k(\lambda)_{m-k}(\lambda)_{n-k}(2\lambda)_{m+n-k}}{(m+n+\lambda-k)k!(m-k)!(n-k)!(\lambda)_{m+n-k}(2\lambda)_{m+n-2k}}.$$

First replace the Gegenbauer polynomials in (58) using (22) then setting  $\lambda \mapsto \lambda + \frac{1}{2}$  completes the proof.  $\blacksquare$

### 3.2 Some definite integrals and sums

Along the lines of [7], one can use these orthogonality relations to compute some new definite integrals. Furthermore, using the derived closure relations one can convert certain definite integrals into infinite series expressions. In this subsection, we give some carefully chosen examples of these procedures.

The following result follows from a sample generating function for the Gegenbauer polynomials (60) and then re-expressing as a definite integral using orthogonality for the Ferrers function of the first kind.

**Corollary 3.3.** *Let  $|x|, |t| < 1$ ,  $\alpha, \gamma \in \mathbb{C}$ ,  $k, \ell \in \mathbb{N}_0$ ,  $\alpha + k + \frac{1}{2} \neq 0$ ,  $\Re\alpha > -\ell - 1$ . Then*

$$\begin{aligned} \int_{-1}^1 (1+t^2-2xt)^{\frac{\alpha-\gamma+\ell}{2}} P_{\ell+\alpha-\gamma}^{-\ell-\alpha} \left( \frac{1-xt}{\sqrt{1+t^2-2xt}} \right) P_{k+\alpha}^{-\ell-\alpha}(x) dx \\ = \frac{(-1)^\ell (\gamma-\ell)_k t^{\alpha+k}}{(1-\gamma)_\ell \Gamma(2\alpha+\ell+k+1)(\alpha+k+\frac{1}{2})}. \end{aligned} \quad (59)$$

*Special care must be taken when  $\gamma \in \mathbb{Z}$ .*

**Proof.** Start with the generating function  $\lambda, \gamma \in \mathbb{C}$ , [8, (9.8.33)]

$$(1-xt)^{-\gamma} {}_2F_1 \left( \begin{matrix} \frac{\gamma}{2}, \frac{\gamma+1}{2} \\ \lambda + \frac{1}{2} \end{matrix}; \frac{t^2(x^2-1)}{(1-xt)^2} \right) = \sum_{n=0}^{\infty} \frac{(\gamma)_n}{(2\lambda)_n} C_n^\lambda(x) t^n, \quad (60)$$

and making the replacement  $n \mapsto k - \ell$ ,  $k \geq \ell \geq 0$  converts the sum over all  $k \geq \ell \in \mathbb{N}_0$ . Then convert the Gegenbauer polynomial to a Ferrers function of the first kind using (20). Multiply the resulting sum by  $(1-x^2)^{\frac{\alpha+\ell}{2}} P_{k'+\alpha}^{-\ell-\alpha}(x)$  and integrate both sides of the equation over  $(-1, 1)$  using the orthogonality relation (37) produces a definite integral of the Gauss hypergeometric function multiplied by a Ferrers function of the first kind. One can then convert the Gauss hypergeometric function to a Ferrers function of the first kind using (14) which completes the proof.  $\blacksquare$

The following result follows from an expansion over Gegenbauer polynomials [2, Corollary 1] which generalizes many classical expansions including the generating function for Legendre polynomials, Heine's formula and Heine's reciprocal square root Fourier cosine series expansion [4, Figure 1].

**Corollary 3.4.** *Let  $\nu, \mu, x \in \mathbb{C}$  such that if  $z \in \mathbb{C} \setminus (-\infty, 1]$  lies on any ellipse with foci at  $\pm 1$  then  $x$  can lie on any point interior to that ellipse. Then*

$$\frac{(1-x^2)^{\frac{\mu}{2}}(z^2-1)^{\frac{\nu-\mu-1}{2}}}{(z-x)^\nu} = \sum_{n=0}^{\infty} \frac{(2\mu+2n+1)(\nu)_n \Gamma(2\mu+n+1)}{n!} P_{n+\mu}^{-\mu}(x) Q_{n+\mu}^{\nu-\mu-1}(z). \quad (61)$$

**Proof.** Start with [2, Corollary 1]

$$\frac{1}{(z-x)^\nu} = \frac{2^{\mu+\frac{1}{2}}\Gamma(\mu)}{\sqrt{\pi}(z^2-1)^{\frac{\nu-\mu}{2}-\frac{1}{4}}} \sum_{n=0}^{\infty} (n+\mu)(\nu)_n Q_{n+\mu-\frac{1}{2}}^{\nu-\mu-\frac{1}{2}}(z) C_n^\mu(x),$$

and convert the Gegenbauer polynomial to a Ferrers function of the first kind using (22), then set  $\mu \mapsto \mu + \frac{1}{2}$ . This completes the proof. ■

**Corollary 3.5.** *Let  $n, l \in \mathbb{N}_0$ ,  $n \geq l$ ,  $\nu, \mu \in \mathbb{C}$ , such that  $z \in \mathbb{C} \setminus (-\infty, 1]$ . Then*

$$\int_{-1}^1 \frac{P_{n+\mu}^{-\mu}(x)}{(z-x)^\nu} (1-x^2)^{\frac{\mu}{2}} dx = \frac{2(\nu)_n Q_{n+\mu}^{\nu-\mu-1}(z)}{(z^2-1)^{\frac{\nu-\mu-1}{2}}}. \quad (62)$$

**Proof.** Starting with (61), multiplying both sides of the equation by  $P_{n+\mu}^{-\mu}(x)$  and integrating over  $(-1, 1)$  using the orthogonality relation (37) completes the proof. ■

The following result follows from an integral derived in [1] by Askey and Razban for the Jacobi polynomials, but then we specialize to Gegenbauer polynomials and then re-express using the Ferrers function of the first kind.

**Corollary 3.6.** *Let  $n \in \mathbb{N}_0$ ,  $\lambda, \gamma \in \mathbb{C}$ . Then*

$$\int_{-1}^1 (1-x)^{\lambda-\gamma} (1+x)^{\lambda} P_{n+\lambda}^{-\lambda}(x) dx = \frac{2^{\lambda-\gamma+1} \Gamma(\lambda-\gamma+1) \Gamma(\gamma+n)}{\Gamma(\gamma) \Gamma(2\lambda-\gamma+n+2)}. \quad (63)$$

**Proof.** Start with [1] (see also [9, (16.4.2)])

$$\int_{-1}^1 P_n^{(\alpha, \beta)}(x) (1-x)^{\alpha-\gamma} (1+x)^\beta dx = \frac{2^{\alpha+\beta-\gamma+1} \Gamma(\alpha-\gamma+1) \Gamma(\beta+n+1) \Gamma(\gamma+n)}{n! \Gamma(\gamma) \Gamma(\alpha+\beta-\gamma+n+2)},$$

then replace  $\alpha, \beta \mapsto \lambda - \frac{1}{2}$  and using (8), then (22) produces the following integral over Gegenbauer polynomials

$$\int_{-1}^1 (1-x)^{\lambda-\gamma-\frac{1}{2}} (1+x)^{\lambda-\frac{1}{2}} C_n^\lambda(x) dx = \frac{\sqrt{\pi} \Gamma(\lambda-\gamma+\frac{1}{2}) \Gamma(2\lambda+n) \Gamma(\gamma+n)}{2^{\gamma-1} n! \Gamma(\lambda) \Gamma(\gamma) \Gamma(2\lambda-\gamma+n+1)}, \quad (64)$$

using (22) and replacing  $\lambda \mapsto \lambda + \frac{1}{2}$  completes the proof. ■

**Remark 3.7.** *The above result is equivalent to the following expansion*

$$\sum_{n=0}^{\infty} \frac{(\lambda+n)(\gamma)_n}{(2\lambda-\gamma+1)_n} C_n^\lambda(x) = \frac{\sqrt{\pi}\Gamma(2\lambda-\gamma+1)(1-x)^{-\gamma}}{2^{2\lambda-\gamma}\gamma(\lambda)\Gamma(\lambda-\gamma+\frac{1}{2})}, \quad (65)$$

which we recently proved in [4, Corollary 5.12].

### 3.3 Asymptotic expansions and Mehler-Heine relations

The asymptotic expansion of the Ferrers function of the first kind which is related to the limit transition [8, (18.6.4)]

$$\lim_{\lambda \rightarrow \infty} \frac{C_n^\lambda(x)}{C_n^\lambda(1)} = n! \lim_{\lambda \rightarrow \infty} \frac{C_n^\lambda(x)}{(2\lambda)_n} = x^n, \quad (66)$$

since [8, Table 18.6.1]  $C_n^\lambda(1) = \frac{(2\lambda)_n}{n!}$ , is given as follows.

**Corollary 3.8.** *Let  $n \in \mathbb{N}_0$ ,  $x \in (-1, 1)$ . Then*

$$\frac{2^\lambda \Gamma(\lambda+1)}{(1-x^2)^{\frac{\lambda}{2}}} \mathbf{P}_{n+\lambda}^{-\lambda}(x) = x^n \left( 1 - \frac{n(n-1)(1-x^2)}{4x^2\lambda} + \frac{n(n-1)(n-2)(n-3)(1-x^2)^2}{32x^4\lambda^2} + \mathcal{O}\left(\frac{1}{\lambda^3}\right) \right), \quad (67)$$

as  $\lambda \rightarrow \infty$ .

**Proof.** Start with (66) then apply (22) with the replacement  $\lambda \mapsto \lambda + \frac{1}{2}$ . This gives the zeroth order term. In order to compute an arbitrary asymptotic expansion near  $\lambda = \infty$ , use (14), expanding the resulting terminating expansion using a Laurent series near  $\lambda = \infty$  with (5). This completes the proof. ■

**Theorem 3.9.** *Let  $n \in \mathbb{N}_0$ ,  $x \in (-1, 1)$ . Then*

$$\begin{aligned} & \frac{\sqrt{\pi}\Gamma(2\lambda+n+1)}{2^\lambda(\lambda+\frac{1}{2})^{\frac{n}{2}}\Gamma(\lambda+\frac{1}{2})(1-\frac{x^2}{\lambda})^{\frac{\lambda}{2}}} \mathbf{P}_{n+\lambda}^{-\lambda} \left( x \left( \lambda + \frac{1}{2} \right)^{-\frac{1}{2}} \right) \\ &= (2x)^n {}_2F_1 \left( \begin{matrix} -\frac{n}{2}, \frac{1-n}{2} \\ \lambda+1 \end{matrix}; -\frac{\lambda}{x^2} \right) \left( 1 + \frac{n(n+1)}{4\lambda} + \frac{n(n-1)(n+1)(3n+2)}{96\lambda^2} + \mathcal{O}\left(\frac{1}{\lambda^3}\right) \right), \end{aligned} \quad (68)$$

as  $\lambda \rightarrow \infty$ .

**Proof.** Using (14) one obtains

$$\begin{aligned} & \frac{\sqrt{\pi}\Gamma(2\lambda+n+1)}{2^\lambda(\lambda+\frac{1}{2})^{\frac{n}{2}}\Gamma(\lambda+\frac{1}{2})(1-\frac{x^2}{\lambda})^{\frac{\lambda}{2}}} \mathbf{P}_{n+\lambda}^{-\lambda} \left( x \left( \lambda + \frac{1}{2} \right)^{-\frac{1}{2}} \right) \\ &= \frac{\Gamma(2\lambda+n+1)x^n}{\Gamma(2\lambda)(\lambda+\frac{1}{2})^n} {}_2F_1 \left( \begin{matrix} -\frac{n}{2}, \frac{1-n}{2} \\ \lambda+1 \end{matrix}; 1 - \frac{\lambda+\frac{1}{2}}{x^2} \right). \end{aligned} \quad (69)$$

Then by performing an asymptotic expansion up to second order in  $1/\lambda$  using [8, (5.11.12)] and expanding the resulting terminating expansion using a Laurent series near  $\lambda = \infty$ , produces the result. ■

**Corollary 3.10.** *Let  $n \in \mathbb{N}_0$ ,  $x \in (-1, 1)$ . Then*

$$\frac{2^\lambda \Gamma(\lambda + 1) (2\lambda + 1)_n}{(\lambda + \frac{1}{2})^{\frac{n}{2}} (1 - x^2)^{\frac{\lambda}{2}}} P_{n+\lambda}^{-\lambda} \left( x \left( \lambda + \frac{1}{2} \right)^{-\frac{1}{2}} \right) \sim H_n(x), \quad (70)$$

as  $\lambda \rightarrow \infty$ , where  $H_n(x)$  are the Hermite polynomials.

**Proof.** Start with [8, (18.7.24)]

$$\lim_{\lambda \rightarrow \infty} \frac{1}{\lambda^{\frac{n}{2}}} C_n^\lambda \left( \frac{x}{\sqrt{\lambda}} \right) = \frac{1}{n!} H_n(x),$$

then inserting (22) produces

$$\lim_{\lambda \rightarrow \infty} \frac{\sqrt{\pi} \Gamma(2\lambda + n)}{2^{\lambda - \frac{1}{2}} \lambda^{\frac{n}{2}} \Gamma(\lambda) (1 - \frac{x^2}{\lambda})^{\frac{\lambda}{2} - \frac{1}{4}}} P_{n+\lambda - \frac{1}{2}}^{\frac{1}{2} - \lambda} \left( \frac{x}{\sqrt{\lambda}} \right) = H_n(x).$$

Setting  $\lambda \mapsto \lambda + \frac{1}{2}$  produces

$$\frac{\sqrt{\pi} \Gamma(2\lambda + n + 1)}{2^\lambda (\lambda + \frac{1}{2})^{\frac{n}{2}} \Gamma(\lambda + \frac{1}{2}) \left( 1 - \frac{x^2}{\lambda + \frac{1}{2}} \right)^{\frac{\lambda}{2}}} P_{n+\lambda}^{-\lambda} \left( x \left( \lambda + \frac{1}{2} \right)^{-\frac{1}{2}} \right) \sim H_n(x),$$

which completes the proof. ■

**Remark 3.11.** *Corollary 3.10 clearly matches up to Theorem 3.9 by using (2) and (4).*

Now we examine this relevant asymptotic expansion in connection with the Mehler-Heine formula for Gegenbauer polynomials in terms of the Ferrers function of the first kind.

**Theorem 3.12.** *Let  $z, \lambda \in \mathbb{C}$ ,  $1 \ll n \in \mathbb{N}_0$ . Then*

$$n^\lambda P_{n+\lambda}^{-\lambda} \left( 1 - \frac{z^2}{2n^2} \right) = J_\lambda(z) \left( 1 + \frac{2\lambda + 1}{n} + \mathcal{O} \left( \frac{1}{n^2} \right) \right), \quad (71)$$

as  $n \rightarrow \infty$ .

**Proof.** The Mehler-Heine formula for Jacobi polynomials [8, (18.11.5)]

$$\lim_{n \rightarrow \infty} \frac{1}{n^\alpha} P_n^{(\alpha, \beta)} \left( 1 - \frac{z}{2n^2} \right) = \left( \frac{2}{z} \right)^\alpha J_\alpha(z), \quad (72)$$

can be used to obtain the Mehler-Heine formula for Gegenbauer polynomials using (8), namely

$$\lim_{n \rightarrow \infty} \frac{n!}{\Gamma(2\lambda + n + 1)} C_n^{\lambda + \frac{1}{2}} \left( 1 - \frac{z^2}{2n^2} \right) = \frac{\sqrt{\pi} J_\lambda(z)}{\Gamma(\lambda + \frac{1}{2})(2z)^\lambda}. \quad (73)$$

Now evaluate the left-hand side of (73) in terms of the Ferrers function of the first kind with (22), which produces

$$C_n^{\lambda+\frac{1}{2}}\left(1-\frac{z^2}{2n^2}\right) = \frac{\sqrt{\pi}\Gamma(2\lambda+n+1)}{2^\lambda\Gamma(\lambda+\frac{1}{2})n!} \left(1-\left(1-\frac{z^2}{2n^2}\right)^2\right)^{-\frac{\lambda}{2}} P_{n+\lambda}^{-\lambda}\left(1-\frac{z^2}{2n^2}\right), \quad (74)$$

then simplification using [8, (5.11.12)]

$$\frac{\Gamma(z+a)}{\Gamma(z+b)} \sim z^{a-b},$$

as  $z \rightarrow \infty$ , obtains the prefactor. Then use (11) and expand the numerator Pochhammer symbols in the  ${}_2F_1$  in descending powers of  $n$ , namely

$$(-n-\lambda)_k(n+\lambda+1)_k = (-1)^k n^{2k} \left(1 + \frac{2\lambda+1}{n} + \mathcal{O}\left(\frac{1}{n^2}\right)\right),$$

and using the definition of the Bessel function of the first kind (3), completes the proof.  $\blacksquare$

### 3.4 Christoffel-Darboux formulas

The Christoffel-Darboux formula for orthogonal Ferrers functions is given in the following theorem which gives the Christoffel-Darboux formula and its confluent form for the Ferrers function of the first kind.

**Theorem 3.13.** *Let  $x, x' \in (-1, 1)$ ,  $\mu \in \mathbb{C}$ ,  $n \in \mathbb{N}_0$ . Then, the Christoffel-Darboux formula and its confluent form for the Ferrers function of the first kind are given by*

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{(2\mu+2k+1)\Gamma(2\mu+k+1)}{k!} P_{k+\mu}^{-\mu}(x) P_{k+\mu}^{-\mu}(x') \\ &= \frac{\Gamma(2\mu+n+1)}{(n-1)!(x-x')} \left( P_{n+\mu}^{-\mu}(x) P_{n+\mu-1}^{-\mu}(x') - P_{n+\mu-1}^{-\mu}(x) P_{n+\mu}^{-\mu}(x') \right), \end{aligned} \quad (75)$$

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{(2\mu+2k+1)\Gamma(2\mu+k+1)}{k!} \left( P_{k+\mu}^{-\mu}(x) \right)^2 \\ &= \frac{\Gamma(2\mu+n+1)}{(n-1)!(1-x^2)} \left( (n+2\mu) \left( P_{n+\mu}^{-\mu}(x) \right)^2 + n \left( P_{n+\mu-1}^{-\mu}(x) \right)^2 \right. \\ & \quad \left. - 2x(n+\mu) P_{n+\mu-1}^{-\mu}(x) P_{n+\mu}^{-\mu}(x) \right). \end{aligned} \quad (76)$$

**Proof.** Start with the Christoffel-Darboux summation formula for the Gegenbauer polynomials [17, (3.10)]

$$\sum_{k=0}^{n-1} \frac{k!(k+\lambda)}{\Gamma(2\lambda+k)} C_k^\lambda(x) C_k^\lambda(x') = \frac{n! \left( C_n^\lambda(x) C_{n-1}^\lambda(x') - C_{n-1}^\lambda(x) C_n^\lambda(x') \right)}{2\Gamma(2\lambda-1+n)(x-x')}. \quad (77)$$

The confluent form of the Christoffel-Darboux summation formula follows from the first formula using [8, (18.2.13)] and recurrence relations for the Gegenbauer polynomials [12, (9.8.21)], [10, (8.933.4)], and is given by

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{k!(k+\lambda)}{\Gamma(2\lambda+k)} (C_k^\lambda(x))^2 \\ &= \frac{n! \left( n(C_n^\lambda(x))^2 + (2\lambda+n-1)(C_{n-1}^\lambda(x))^2 - x(2\lambda+2n-1)C_n^\lambda(x)C_{n-1}^\lambda(x) \right)}{2(1-x^2)\Gamma(2\lambda-1+n)}. \end{aligned} \quad (78)$$

The conversion to the Ferrers functions of the first kind is accomplished using (22). This completes the proof.  $\blacksquare$

Using (17), the above Christoffel-Darboux formulas for the Ferrers function of the first kind can be expressed in terms of the Ferrers function of the second kind.

**Corollary 3.14.** *Let  $n \in \mathbb{N}_0$ ,  $x, x' \in (-1, 1)$ . Then*

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{(n+k)k!}{(2n+k-1)!} \mathbf{Q}_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x) \mathbf{Q}_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x') \\ &= \frac{n!}{2(3n-2)!(x-x')} \left( \mathbf{Q}_{2n-\frac{1}{2}}^{n-\frac{1}{2}}(x) \mathbf{Q}_{2n-\frac{3}{2}}^{n-\frac{1}{2}}(x') - \mathbf{Q}_{2n-\frac{3}{2}}^{n-\frac{1}{2}}(x) \mathbf{Q}_{2n-\frac{1}{2}}^{n-\frac{1}{2}}(x') \right), \end{aligned} \quad (79)$$

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{(n+k)k!}{(2n+k-1)!} \left( \mathbf{Q}_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x) \right)^2 \\ &= \frac{n!}{2(3n-2)!(1-x^2)} \left( n \left( \mathbf{Q}_{2n-\frac{1}{2}}^{n-\frac{1}{2}}(x) \right)^2 + (3n-1) \left( \mathbf{Q}_{2n-\frac{3}{2}}^{n-\frac{1}{2}}(x) \right)^2 \right. \\ & \quad \left. - x(4n-1) \mathbf{Q}_{2n-\frac{3}{2}}^{n-\frac{1}{2}}(x) \mathbf{Q}_{2n-\frac{1}{2}}^{n-\frac{1}{2}}(x) \right). \end{aligned} \quad (80)$$

**Proof.** Let  $\mu = n - \frac{1}{2}$  in (75), (76), respectively using (17) completes the proof.  $\blacksquare$

### 3.5 Poisson kernels

The Poisson kernel  $K_t(x, y)$  of an orthogonal polynomial  $p_n(x; \mathbf{a})$ , where  $\mathbf{a}$  is a set of parameters that the orthogonal polynomial depends upon is given by

$$K_t(x, y; \mathbf{a}) = \sum_{n=0}^{\infty} \frac{t^n}{h_n} p_n(x; \mathbf{a}) p_n(y; \mathbf{a}), \quad (81)$$

where  $h_n$  (98) is the norm of the polynomial. In this subsection we start with the Poisson kernel of Gegenbauer polynomials and some companion identities to derive addition theorems for the Ferrers function of the first kind.

**Theorem 3.15.** *Let  $t, \lambda \in \mathbb{C}$ ,  $|t| < 1$ ,  $x, y \in (-1, 1)$ ,  $x = \cos \theta$ ,  $y = \cos \theta'$ . Then*

$$\sum_{n=0}^{\infty} \frac{(2\lambda+1)_n t^n}{n!} \mathbf{P}_{n+\lambda}^{-\lambda}(x) \mathbf{P}_{n+\lambda}^{-\lambda}(y) = \frac{Q_{\lambda-\frac{1}{2}}(x)}{\pi \Gamma(2\lambda+1) t^{\lambda+\frac{1}{2}} \sqrt{\sin \theta \sin \theta'}}, \quad (82)$$

where

$$\chi := \frac{1 + t^2 - 2t \cos \theta \cos \theta'}{2 \sin \theta \sin \theta'}. \quad (83)$$

**Proof.** Start with the Poisson kernel for the Gegenbauer polynomials [13, (19)]

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{n! t^n}{(2\lambda)_n} C_n^\lambda(x) C_n^\lambda(y) &= \frac{1}{(1 + t^2 - 2t \cos \theta \cos \phi)^\lambda} {}_2F_1 \left( \begin{matrix} \frac{\lambda}{2}, \frac{\lambda+1}{2} \\ \lambda + \frac{1}{2} \end{matrix}; \frac{1}{\chi^2} \right) \\ &= \frac{\Gamma(\lambda + \frac{1}{2}) Q_{\lambda-1}(\chi)}{\sqrt{\pi} \Gamma(\lambda) t^\lambda (1-x^2)^{\frac{\lambda}{2}} (1-y^2)^{\frac{\lambda}{2}}}, \end{aligned} \quad (84)$$

then replacing  $\lambda \mapsto \lambda + \frac{1}{2}$ , and using (22) completes the proof.  $\blacksquare$

**Remark 3.16.** Note that for  $\lambda = m$ , and setting  $n \mapsto n + m$ , Theorem 3.15 specializes to the second addition theorem for spherical harmonics presented in [6, (2.4)]

$$\sum_{n=|m|}^{\infty} t^n \frac{(n-m)!}{(n+m)!} P_n^m(\cos \theta) P_n^m(\cos \theta') = \frac{Q_{m-\frac{1}{2}}(\chi)}{\pi \sqrt{t \sin \theta \sin \theta'}}, \quad (85)$$

after letting  $t = r_{<}/r_{>}$ , where  $r_{\leq} := \min_{\max} \{r, r'\}$  with  $r, r' \in [0, \infty)$ .

**Theorem 3.17.** Let  $t, \lambda \in \mathbb{C}$ ,  $|t| < 1$ ,  $x, y \in (-1, 1)$ ,  $x = \cos \theta$ ,  $y = \cos \theta'$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda + n + \frac{1}{2})(2\lambda + 1)_n t^n}{n!} P_{n+\lambda}^{-\lambda}(x) P_{n+\lambda}^{-\lambda}(y) \\ = - \frac{1-t^2}{2\pi \Gamma(2\lambda + 1) t^{\lambda+\frac{3}{2}} (1-x^2)^{\frac{3}{4}} (1-y^2)^{\frac{3}{4}}} \frac{Q_{\lambda-\frac{1}{2}}^1(\chi)}{\sqrt{\chi^2 - 1}}. \end{aligned} \quad (86)$$

**Proof.** Start with the companion Poisson kernel for Gegenbauer polynomials [13, (18)]

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\lambda + n}{\lambda} \frac{n! t^n}{(2\lambda)_n} C_n^\lambda(x) C_n^\lambda(y) &= \frac{1-t^2}{(1 + t^2 - 2t \cos \theta \cos \phi)^{\lambda+1}} {}_2F_1 \left( \begin{matrix} \frac{\lambda+1}{2}, \frac{\lambda+2}{2} \\ \lambda + \frac{1}{2} \end{matrix}; \frac{1}{\chi^2} \right) \\ &= - \frac{\Gamma(\lambda + \frac{1}{2})(1-t^2)}{\sqrt{\pi} \Gamma(\lambda + 1) t^{\lambda+1} (1-x^2)^{\frac{\lambda+1}{2}} (1-y^2)^{\frac{\lambda+1}{2}}} \frac{Q_{\lambda-1}^1(\chi)}{\sqrt{\chi^2 - 1}}, \end{aligned} \quad (87)$$

then replacing  $\lambda \mapsto \lambda + \frac{1}{2}$ , and using (22) completes the proof.  $\blacksquare$

**Theorem 3.18.** Let  $t, \lambda \in \mathbb{C}$ ,  $|t| < 1$ ,  $x, y \in (-1, 1)$ ,  $x = \cos \theta$ ,  $y = \cos \theta'$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda + n + \frac{1}{2})(\lambda + n + \frac{3}{2})(2\lambda + 1)_n t^n}{n!} P_{n+\lambda}^{-\lambda}(x) P_{n+\lambda}^{-\lambda}(y) \\ = \frac{1}{\pi t^{\lambda-\frac{1}{2}} \Gamma(2\lambda + 1) (1-x^2)^{\frac{3}{4}} (1-y^2)^{\frac{3}{4}}} \left( \frac{Q_{\lambda-\frac{1}{2}}^1(\chi)}{\sqrt{\chi^2 - 1}} + \frac{(1-t^2)^2 Q_{\lambda-\frac{1}{2}}^2(\chi)}{4t^3 \sqrt{1-x^2} \sqrt{1-y^2} (\chi^2 - 1)} \right). \end{aligned} \quad (88)$$

**Proof.** Start with Theorem 3.17, and put all powers of  $t$  on the left-hand side of the equation and differentiate with respect to  $t$  using

$$\frac{d}{dt} \left( \frac{t^{\lambda+n+\frac{3}{2}}}{1-t^2} \right) = \frac{2t^{\lambda+n+\frac{5}{2}}}{(1-t^2)^2} + \frac{(\lambda+n+\frac{3}{2})t^{\lambda+n+\frac{1}{2}}}{1-t^2},$$

$$\frac{\partial \chi}{\partial t} = \frac{-(1-t^2)}{2t^2\sqrt{1-x^2}\sqrt{1-y^2}},$$

and the following derivative formula for the associated Legendre function of the second kind (cf. [3, Remark 4])

$$\frac{d}{d\chi} \frac{Q_{\lambda-\frac{1}{2}}^m(\chi)}{(1-\chi^2)^{\frac{m}{2}}} = \frac{Q_{\lambda-\frac{1}{2}}^{m+1}(\chi)}{(\chi^2-1)^{\frac{m+1}{2}}},$$

for  $m \in \mathbb{N}_0$ . Then applying Theorem 3.17 again and the recurrence relation

$$Q_{\lambda-\frac{1}{2}}^{\mu+1}(\chi) = \frac{1}{\sqrt{\chi^2-1}} \left( -(\lambda+\mu+\frac{1}{2})\chi Q_{\lambda-\frac{1}{2}}^{\mu}(\chi) + (\lambda-\mu+\frac{1}{2})Q_{\lambda+\frac{1}{2}}^{\mu}(\chi) \right),$$

completes the proof. ■

**Corollary 3.19.** *Let  $t, \lambda \in \mathbb{C}$ ,  $|t| < 1$ ,  $x, y \in (-1, 1)$ ,  $x = \cos \theta$ ,  $y = \cos \theta'$ . Then*

$$\begin{aligned} & \sum_{n=0}^{\infty} (\lambda+n)(\lambda+n+1) \frac{n!t^n}{(2\lambda)_n} C_n^{\lambda}(x) C_n^{\lambda}(y) \\ &= \frac{\Gamma(\lambda+\frac{1}{2})}{\sqrt{\pi}t^{\lambda-1}\Gamma(\lambda)(1-x^2)^{\frac{\lambda+1}{2}}(1-y^2)^{\frac{\lambda+1}{2}}} \left( \frac{Q_{\lambda-1}^1(\chi)}{\sqrt{\chi^2-1}} + \frac{(1-t^2)^2 Q_{\lambda-1}^2(\chi)}{4t^3\sqrt{1-x^2}\sqrt{1-y^2}(\chi^2-1)} \right). \end{aligned} \quad (89)$$

**Proof.** Using (22) in Theorem 3.18, completes the proof. ■

**Corollary 3.20.** *Let  $t, \lambda \in \mathbb{C}$ ,  $|t| < 1$ ,  $x, y \in (-1, 1)$ ,  $x = \cos \theta$ ,  $y = \cos \theta'$ . Then, one has the following algebraic generating functions for the Gegenbauer polynomials:*

$$\sum_{n=0}^{\infty} t^n C_n^{\lambda}(x) = \frac{1}{(1+t^2-2tx)^{\lambda}}, \quad (90)$$

$$\sum_{n=0}^{\infty} (\lambda+n)t^n C_n^{\lambda}(x) = \frac{\lambda(1-t^2)}{(1+t^2-2tx)^{\lambda+1}}, \quad (91)$$

$$\sum_{n=0}^{\infty} (\lambda+n)(\lambda+n+1)t^n C_n^{\lambda}(x) = \frac{\lambda(\lambda+1)(1-t^2)^2}{(1+t^2-2tx)^{\lambda+2}} - \frac{2\lambda t^2}{(1+t^2-2tx)^{\lambda+1}}. \quad (92)$$

**Proof.** Consider the limit as  $y \rightarrow 1^-$  as follows. Let  $y = 1 - \frac{1}{2}\epsilon^2$  with  $\epsilon > 0$ , such that  $\epsilon \ll 1$ . Then  $\sqrt{1 - y^2} \sim \epsilon$  and

$$\chi \sim \frac{1 + t^2 - 2tx}{2t\sqrt{1 - x^2}\epsilon},$$

as  $\epsilon \rightarrow 0^+$ . Note that [8, Table 18.6.1]  $C_n^\lambda(1) = (2\lambda)_n/n!$ . Applying this and the above estimates in (84), (87) and Corollary 3.19, then taking the limit as  $\epsilon \rightarrow 0^+$ , completes the proof. ■

**Corollary 3.21.** *Let  $t, \lambda \in \mathbb{C}$ ,  $|t| < 1$ ,  $x, y \in (-1, 1)$ ,  $x = \cos \theta$ ,  $y = \cos \theta'$ . Then*

$$\sum_{n=0}^{\infty} \frac{t^n (2\lambda + 1)_n}{n!} \mathbf{P}_{n+\lambda}^{-\lambda}(x) = \frac{(1 - x^2)^{\frac{\lambda}{2}}}{2^\lambda \Gamma(\lambda + 1) (1 + t^2 - 2tx)^{\lambda + \frac{1}{2}}}, \quad (93)$$

$$\sum_{n=0}^{\infty} (\lambda + n + \frac{1}{2}) \frac{t^n (2\lambda + 1)_n}{n!} \mathbf{P}_{n+\lambda}^{-\lambda}(x) = \frac{(\lambda + \frac{1}{2})(1 - t^2)(1 - x^2)^{\frac{\lambda}{2}}}{2^\lambda \Gamma(\lambda + 1) (1 + t^2 - 2tx)^{\lambda + \frac{3}{2}}}, \quad (94)$$

$$\begin{aligned} \sum_{n=0}^{\infty} (\lambda + n + \frac{1}{2})(\lambda + n + \frac{3}{2}) \frac{t^n (2\lambda + 1)_n}{n!} \mathbf{P}_{n+\lambda}^{-\lambda}(x) \\ = \frac{(1 - x^2)^{\frac{\lambda}{2}} (\lambda + \frac{1}{2})}{2^\lambda \Gamma(\lambda + 1)} \left( \frac{(\lambda + \frac{3}{2})(1 - t^2)^2}{(1 + t^2 - 2tx)^{\lambda + \frac{5}{2}}} - \frac{2t^2}{(1 + t^2 - 2tx)^{\lambda + \frac{3}{2}}} \right). \end{aligned} \quad (95)$$

**Proof.** Starting with Corollary 3.20 and replacing the Gegenbauer polynomials using (22) completes the proof. ■

### 3.6 Closure relations

Closure relations are infinite series representations of the Dirac delta distribution. Let  $n, m \in \mathbb{N}_0$ , the closure relations are obtained from a complete set of orthogonal functions  $\psi_n(x)$  on  $(a, b)$  with a weight function  $w(x)$ , with respect to an inner product defined as

$$\langle f, g \rangle := \int_a^b f(x)g(x)w(x) dx, \quad (96)$$

so that the moments  $(m_n)$  are bounded, where

$$m_n := \langle x^n, 1 \rangle. \quad (97)$$

Then, the integral orthogonality relation is given by

$$\langle \psi_n, \psi_m \rangle = \int_a^b \psi_n(x)\psi_m(x)w(x) dx = h_n \delta_{n,m}. \quad (98)$$

**Proposition 3.22.** *Let  $x, x' \in [a, b]$  and the Poisson kernel is non-negative. Then, one has the following closure relation:*

$$\sum_{n=0}^{\infty} \frac{w(x)}{h_n} \psi_n(x)\psi_n(x') = \delta(x - x'), \quad (99)$$

where both the left- and right-hand sides should be treated as distributions.

**Proof. (formal)** One may expand a function  $f \in L^2(a, b)$  in the infinite series of the orthogonal functions as

$$f(x) = \sum_{n=0}^{\infty} a_n \psi_n(x). \quad (100)$$

Now multiplying both sides of this equation by  $w(x)\psi_m(x)$  and integrating over the interval  $(a, b)$  produces

$$a_n = \frac{1}{h_n} \int_a^b f(x) \psi_n(x) w(x) dx.$$

Multiplying the above expression by  $\psi_n(x')$  and summing over all non-negative integers produces

$$f(x') = \int_a^b f(x) \left( \sum_{n=0}^{\infty} \frac{w(x)}{h_n} \psi_n(x) \psi_n(x') \right) dx = \int_a^b f(x) \delta(x - x') dx,$$

where we have used the sifting property of the Dirac delta distribution [8, (1.16.11)]. This completes the proof. See [11, Theorem 2.1] for a different proof. ■

The following is the closure relation for Ferrers function of the first kind.

**Theorem 3.23.** *Let  $x, x' \in (-1, 1)$ ,  $\mu \in \mathbb{C}$ ,  $2\mu + 1 \notin -\mathbb{N}_0$ . Then*

$$\sum_{n=0}^{\infty} \frac{(\mu + n + \frac{1}{2}) \Gamma(2\mu + n + 1)}{n!} P_{n+\mu}^{-\mu}(x) P_{n+\mu}^{-\mu}(x') = \delta(x - x'). \quad (101)$$

**Proof.** Using the orthogonality relation for Gegenbauer polynomials (38) in order to obtain  $h_n$  with (99) produces

$$\sum_{n=0}^{\infty} \frac{(\mu + n) n!}{\Gamma(2\mu + n)} C_n^{\mu}(x) C_n^{\mu}(x') = \frac{\pi \delta(x - x')}{(1 - x^2)^{\mu - \frac{1}{2}} 2^{2\mu - 1} \Gamma^2(\mu)}. \quad (102)$$

Applying (22) twice completes the proof. ■

**Remark 3.24.** *Note that in the limit as  $n \rightarrow \infty$  the Christoffel-Darboux formula (75) becomes the closure relation (101). Furthermore, the Poisson kernel can be obtained from the Christoffel-Darboux sum with each coefficient multiplied by some  $t^k$ , with  $|t| < 1$ . This interrelation as well as the interrelation with universality for orthogonal polynomials are quite intriguing (see the excellent review article by Barry Simon entitled “The Christoffel-Darboux Kernel” [16, pp. 295–335]).*

**Corollary 3.25.** *Let  $x \in (-1, 1)$ ,  $\lambda, \gamma \in \mathbb{C}$  such that  $2\lambda - \gamma + 2 \notin -\mathbb{N}_0$ . Then*

$$\sum_{n=0}^{\infty} \frac{(\lambda + n + \frac{1}{2})(\gamma)_n (2\lambda + 1)_n}{n! (2\lambda - \gamma + 2)_n} P_{n+\lambda}^{-\lambda}(x) = \frac{\Gamma(2\lambda - \gamma + 2)(1 - x)^{\frac{\lambda}{2} - \gamma} (1 + x)^{\frac{\lambda}{2}}}{2^{\lambda - \gamma + 1} \Gamma(2\lambda + 1) \Gamma(\lambda - \gamma + 1)}. \quad (103)$$

**Proof.** Applying the closure relation (101) to (63), multiplying by the necessary factors and integrating over  $(-1, 1)$  completes the proof. ■

**Remark 3.26.** *It is interesting to see that using orthogonality for Jacobi polynomials [8, Table 18.3.1], the closure relation for Jacobi polynomials is given by*

$$\sum_{n=0}^{\infty} \frac{(2n + \alpha + \beta + 1)\Gamma(\alpha + \beta + n + 1)n!}{\Gamma(\alpha + n + 1)\Gamma(\beta + n + 1)} P_n^{(\alpha, \beta)}(x)P_n^{(\alpha, \beta)}(x') = \frac{2^{\alpha+\beta+1}\delta(x - x')}{(1 - x)^\alpha(1 + x)^\beta}. \quad (104)$$

**Remark 3.27.** *It should be noted that if one takes  $\mu = 0$  in (101) then one obtains the closure relation for Legendre polynomials (30) [8, (1.17.22)]*

$$\sum_{n=0}^{\infty} (n + \frac{1}{2})P_n(x)P_n(x') = \delta(x - x'). \quad (105)$$

On the other hand, using  $\mu = \pm 1/2$  in (101) produces the following closure relations for the Chebyshev polynomials of the first and second kinds [8, Table 18.3.1], namely:

$$\sum_{n=0}^{\infty} T_n(x)T_n(x') = \frac{\pi}{2}(1 - x^2)^{\frac{1}{4}}(1 - x'^2)^{\frac{1}{4}}\delta(x - x'), \quad (106)$$

$$\sum_{n=0}^{\infty} U_n(x)U_n(x') = \frac{\pi\delta(x - x')}{2(1 - x^2)^{\frac{1}{4}}(1 - x'^2)^{\frac{1}{4}}}. \quad (107)$$

Using (44) we can obtain a closure relation for the Ferrers function of the second kind.

**Corollary 3.28.** *Let  $n \in \mathbb{N}_0$ ,  $x, x' \in (-1, 1)$ . Then*

$$\sum_{k=0}^{\infty} \frac{(n + k)k!}{(2n + k - 1)!} Q_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x)Q_{k+n-\frac{1}{2}}^{n-\frac{1}{2}}(x') = \frac{\pi^2}{4}\delta(x - x'). \quad (108)$$

**Proof.** From (44) we obtain  $h_n = \pi^2(2n + k - 1)!/(4(n + k)k!)$ , then using (99) with  $w(x) = 1$  completes the proof. ■

The  $n = 0, 1$  specializations of (108) correspond with the closure relation for the Chebyshev polynomials of the first and second kinds (106), (107).

**Corollary 3.29.** *Let  $x, x' \in (-1, 1)$ ,  $x = \cos \theta$ ,  $x' = \cos \theta'$ . Then*

$$\sum_{k=0}^{\infty} k^2 Q_{k-\frac{1}{2}}^{-\frac{1}{2}}(x)Q_{k-\frac{1}{2}}^{-\frac{1}{2}}(x') = \frac{\pi}{2\sqrt{\sin \theta \sin \theta'}} \sum_{k=0}^{\infty} \cos(k\theta) \cos(k\theta') = \frac{\pi^2}{4}\delta(x - x'), \quad (109)$$

$$\sum_{k=0}^{\infty} Q_{k+\frac{1}{2}}^{\frac{1}{2}}(x)Q_{k+\frac{1}{2}}^{\frac{1}{2}}(x') = \frac{\pi}{2\sqrt{\sin \theta \sin \theta'}} \sum_{k=1}^{\infty} \sin(k\theta) \sin(k\theta') = \frac{\pi^2}{4}\delta(x - x'). \quad (110)$$

**Proof.** Start with (108) with  $n = 0, 1$ , and using (18), (19), completes the proof. ■

## Acknowledgements

Much thanks to Adri Olde Daalhuis and Nico Temme for valuable discussions. The research of R.S.C.-S. was funded by Agencia Estatal de Investigación of Spain, grant number PGC-2018-096504-B-C33.

## References

- [1] R. Askey and B. Razban. An integral for Jacobi polynomials. *Simon Stevin*, 46:165–169, 1972.
- [2] H. S. Cohl. Fourier, Gegenbauer and Jacobi expansions for a power-law fundamental solution of the polyharmonic equation and polyspherical addition theorems. *Symmetry, Integrability and Geometry: Methods and Applications*, 9(042):26, 2013.
- [3] H. S. Cohl and R. S. Costas-Santos. Multi-integral representations for associated Legendre and Ferrers functions. *Symmetry*, 12(1598):22, 2020.
- [4] H. S. Cohl, R. S. Costas-Santos, and T. V. Wakhare. On a generalization of the Rogers generating function. *Journal of Mathematical Analysis and Applications*, 475(2):1019–1043, 2019.
- [5] H. S. Cohl, J. Park, and H. Volkmer. Gauss hypergeometric representations of the ferrers function of the second kind. *SIGMA. Symmetry, Integrability and Geometry. Methods and Applications*, 17:Paper No. 053, 33, 2021.
- [6] H. S. Cohl, A. R. P. Rau, J. E. Tohline, D. A. Browne, J. E. Cazes, and E. I. Barnes. Useful alternative to the multipole expansion of  $1/r$  potentials. *Physical Review A: Atomic and Molecular Physics and Dynamics*, 64(5):052509, Oct 2001.
- [7] H. S. Cohl and H. Volkmer. Definite integrals using orthogonality and integral transforms. *Symmetry, Integrability and Geometry: Methods and Applications, Special Issue on Super-integrability, Exact Solvability, and Special Functions*, 8, 2012.
- [8] *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov>, Release 1.1.2 of 2021-06-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [9] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Tables of Integral Transforms. Vol. II*. McGraw-Hill Book Company, Inc., New York-Toronto-London, 1954.
- [10] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products*. Elsevier/Academic Press, Amsterdam, seventh edition, 2007.
- [11] M. E. H. Ismail, R. Zhang, and K. Zhou.  $q$ -Fractional Askey-Wilson Integrals and Related Semigroups of Operators. *in preparation*, 2021.
- [12] R. Koekoek, P. A. Lesky, and R. F. Swarttouw. *Hypergeometric orthogonal polynomials and their  $q$ -analogues*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2010. With a foreword by Tom H. Koornwinder.
- [13] R. S. Maier. Algebraic generating functions for Gegenbauer polynomials. In *Frontiers in orthogonal polynomials and  $q$ -series*, volume 1 of *Contemp. Math. Appl. Monogr. Expo. Lect. Notes*, pages 425–444. World Sci. Publ., Hackensack, NJ, 2018.

- 
- [14] F. W. J. Olver. *Asymptotics and Special Functions*. AKP Classics. A K Peters Ltd., Wellesley, MA, 1997. Reprint of the 1974 original [Academic Press, New York].
- [15] J. F. Sánchez-Lara. On the Sobolev orthogonality of classical orthogonal polynomials for non standard parameters. *Rocky Mountain J. Math.*, 47(1):267–288, 2017.
- [16] B. Simon. The Christoffel-Darboux kernel. In *Perspectives in partial differential equations, harmonic analysis and applications, A Volume in Honor of Vladimir G. Maz'ya's 70th Birthday*, volume 79, pages 295–335, Providence, RI, USA, 2008. American Mathematical Society.
- [17] R. Szmytkowski. A note on parameter derivatives of classical orthogonal polynomials. *arXiv:0901.2639*, 2018.