

A NON-COMMUTATIVE NULLSTELLENSATZ

ZHENGHENG BAO AND ZINOVY REICHSTEIN

ABSTRACT. Let K be a field and D be a finite-dimensional central division algebra over K . We prove a variant of the Nullstellensatz for 2-sided ideals in the ring of polynomial maps $D^n \rightarrow D$. In the case where $D = K$ is commutative, our main result reduces to the K -Nullstellensatz of Laksov and Adkins-Gianni-Tognoli. In the case, where $K = \mathbb{R}$ is the field of real numbers and D is the algebra of Hamilton quaternions, it reduces to the quaternionic Nullstellensatz recently proved by Alon and Paran.

1. INTRODUCTION

Let K be a field and $P_{K,n} = K[x_1, \dots, x_n]$ be the commutative polynomial ring in n variables over K . For any ideal $J \subset P_{K,n}$ we will denote the set of K -points cut out by J in the affine space \mathbb{A}^n by

$$\mathcal{Z}(J) = \{(a_1, \dots, a_n) \in K^n \mid f(a_1, \dots, a_n) = 0 \ \forall f \in J\}.$$

Conversely, for any set X of K -points in \mathbb{A}^n , we will denote the ideal of polynomials vanishing at every point of X by $\mathcal{I}(X) \subset P_{K,n}$. If K is an algebraically closed field, then the celebrated Hilbert Nullstellensatz asserts that

$$(1) \quad \mathcal{I}(\mathcal{Z}(J)) = \text{Rad}(J),$$

where $\text{Rad}(J) = \{f \in P_{K,n} \mid f^m \in J \text{ for some integer } m \geq 1\}$ is the radical of J .

If K is not algebraically closed, then $\mathcal{I}(\mathcal{Z}(J))$ is larger than $\text{Rad}(J)$ in general. However, several variants of the Nullstellensatz are known to hold in this setting. In particular, if $K = \mathbb{R}$ is the field of real numbers, then for every ideal $J \subset \mathbb{R}[x_1, \dots, x_n]$, we have

$$(2) \quad \mathcal{I}(\mathcal{Z}(J)) = \text{Rad}_{\mathbb{R}}(J),$$

where

$$\begin{aligned} \text{Rad}_{\mathbb{R}}(J) = \{f \in P_{K,n} \mid \exists m, r \geq 1 \text{ and } f_1, \dots, f_{m-1} \in P_{K,n} \\ \text{such that } f_1^2 + \dots + f_{m-1}^2 + f^{2r} \in J\}. \end{aligned}$$

This assertion is known as the ‘‘real Nullstellensatz’’; see [PD, Corollary 4.2.5]. The ideal $\text{Rad}_{\mathbb{R}}(J)$ is called the real radical of J . For the history of the real Nullstellensatz and further references, we refer the reader to [PD, Section 4.7].

2020 *Mathematics Subject Classification.* 16S36, 14A25.

Key words and phrases. The Nullstellensatz, real Nullstellensatz, central division algebra.

Zhengheng Bao was partially supported by a Work Learn International Undergraduate Research Award (WLIURA) at the University of British Columbia.

Zinovy Reichstein was partially supported by National Sciences and Engineering Research Council of Canada Discovery grant 253424-2017.

There is also a version of the Nullstellensatz for a general field K . To state it, recall that a homogeneous polynomial $p \in K[z_1, \dots, z_m]$ is called anisotropic if $p(b_1, \dots, b_m) = 0$ for some $b_1, \dots, b_m \in K$ is only possible when $b_1 = \dots = b_m = 0$. Similarly we will say that a homogeneous polynomial $p \in K[z_1, \dots, z_m]$ is quasi-anisotropic if $p(b_1, \dots, b_m) = 0$ for some $b_1, \dots, b_m \in K$ is only possible when $b_m = 0$. Then

$$(3) \quad \mathcal{I}(\mathcal{Z}(J)) = \text{Rad}_K(J),$$

where

$$\begin{aligned} \text{Rad}_K(J) = \{f \in P_{K,n} \mid \exists m \geq 1, \text{ a homogeneous quasi-anisotropic } p \in K[z_1, \dots, z_m] \\ \text{and } f_1, \dots, f_{m-1} \in P_{K,n} \text{ such that } p(f_1, \dots, f_{m-1}, f) \in J\}. \end{aligned}$$

This identity, known as the K -Nullstellensatz, is due to Laksov [La, p. 324]. In weaker form it appeared earlier in the work of Adkins-Gianni-Tognoli [AGT]; see Section 4. The ideal $\text{Rad}_K(J) \subset K[x_1, \dots, x_n]$ is called the K -radical of J . Laksov [La] showed that (3) can fail if “quasi-anisotropic” is replaced by “anisotropic” in the definition of $\text{Rad}_K(J)$.

The purpose of this note is to explore a non-commutative version of the Nullstellensatz, following up on a recent paper by Alon and Paran [AP]. In the sequel K will denote an infinite field and D a finite-dimensional central division algebra of degree s over K . The role of the polynomial ring $P_{K,n}$ will be played by the ring $P_{D,n}$ of polynomial functions $f: D^n \rightarrow D$. Here we refer to a function $f(x_1, \dots, x_n): D^n \rightarrow D$ as a polynomial function if

- (1) f is expressible as a sum of finitely many “monomial” functions of the form

$$(x_1, \dots, x_n) \mapsto a_1 x_{i_1} a_2 x_{i_2} \cdot \dots \cdot a_r x_{i_r} a_{r+1},$$

for some $a_1, \dots, a_{r+1} \in D$.

Alternatively, if b_1, \dots, b_{s^2} is a K -basis for D , then we can identify D with K^{s^2} and D^n with K^{ns^2} . Writing $x_i = y_{i1}b_1 + \dots + y_{is^2}b_{s^2}$ for each $i = 1, \dots, n$ and expanding $f(x_1, \dots, x_n)$, we obtain

$$f(x_1, \dots, x_n) = f_1(y_{ij})b_1 + \dots + f_{s^2}(y_{ij})b_{s^2},$$

where each f_i is a function $K^{ns^2} \rightarrow K$. We can now define $f(x_1, \dots, x_n): D^n \rightarrow D$ to be a polynomial function if

- (2) each $f_k(y_{ij})$ is a polynomial in the variables y_{ij} ($i = 1, \dots, n$ and $j = 1, \dots, s^2$) with coefficients in K .

It is easy to see that (2) does not depend on the choice of K -basis b_1, \dots, b_{s^2} in D . Moreover, (1) and (2) are equivalent; see Wilczyński [Wi, Theorem 4.1] or Alon-Paran [AP, Theorem 5].

As in the commutative case, for any two-sided ideal $J \subset P_{D,n}$, we can define the zero locus of J in D^n as

$$\mathcal{Z}(J) = \{(d_1, \dots, d_n) \in D^n \mid f(d_1, \dots, d_n) = 0 \forall f \in J\}.$$

Similarly for any subset $X \subset D^n$, we can define the two-sided ideal

$$\mathcal{I}(X) = \{f \in P_{D,n} \mid f(d_1, \dots, d_n) = 0 \forall (d_1, \dots, d_n) \in X\}$$

of $P_{D,n}$.

In the case, where $K = \mathbb{R}$ is the field of real numbers, and $D = \mathbb{H}$, the algebra of Hamilton quaternions, Alon and Paran proved the following “quaternionic” version of the Nullstellensatz:

$$(4) \quad \mathcal{I}(\mathcal{Z}(J)) = \{f \in P_{\mathbb{H},n} \mid \exists m, r \geq 1 \text{ and } f_1, \dots, f_{m-1} \in P_{\mathbb{H},n} \\ \text{such that } N_{\mathbb{H}}(f_1) + \dots + N_{\mathbb{H}}(f_{m-1}) + N_{\mathbb{H}}(f)^r \in J\};$$

see [AP, Theorem 1]. Here $N_{\mathbb{H}}(f)$ denotes the composition of $f: \mathbb{H}^n \rightarrow \mathbb{H}$ with the norm function $N_{\mathbb{H}}: \mathbb{H} \rightarrow \mathbb{R} \hookrightarrow \mathbb{H}$. Their proof is based on the real Nullstellensatz (2).

Our goal in this note is to extend (4) to two-sided ideals J in $P_{D,n}$, for an arbitrary infinite field K and arbitrary central division K -algebra D . To state our main result, we need to introduce some additional notations.

We will denote the center of $P_{D,n}$ by $C_{D,n}$. It is easy to see that $C_{D,n}$ is the ring of polynomial maps $D^n \rightarrow K$, where we identify K with the center of D . In other words, $C_{D,n}$ is the commutative polynomial ring $K[y_{ij}]$ in ns^2 variables, where $i = 1, \dots, n$ and $j = 1, \dots, s^2$. Note that in the case, where $D = K$ is a field (i.e., $s = 1$), $P_{D,n} = C_{D,n} = P_{K,n} = K[x_1, \dots, x_n]$.

There are two natural candidates for the radical of a 2-sided ideal $J \subset P_{D,n}$ suggested by (4), $\text{Rad}_D(J)$ and $\text{Rad}'_D(J)$ defined below. Here $J_c = J \cap C_{D,n}$.

$$\begin{aligned} \text{Rad}_D(J) &= \{f \in P_{D,n} \mid N_D(f) \in \text{Rad}_K(J_c)\} \\ &= \{f \in P_{D,n} \mid \exists m \geq 1, \text{ a homogeneous quasi-anisotropic } p \in K[z_1, \dots, z_m] \\ &\quad \text{and } f_1, \dots, f_{m-1} \in C_{D,n} \text{ such that } p(f_1, \dots, f_{m-1}, N_D(f)) \in J\}. \end{aligned}$$

and

$$\begin{aligned} \text{Rad}'_D(J) &= \{f \in P_{D,n} \mid \exists m \geq 1, \text{ a homogeneous quasi-anisotropic } p \in K[z_1, \dots, z_m] \\ &\quad \text{and } f_1, \dots, f_{m-1} \in P_{D,n} \text{ such that } p(N_D(f_1), \dots, N_D(f_{m-1}), N_D(f)) \in J\}. \end{aligned}$$

Once again, by the norm $N_D(f)$ of $f \in P_{D,n}$ we mean the composition of $f: D^n \rightarrow D$ with the norm map $N_D: D \rightarrow K$, where we identify K with the center of D . We are now ready to state the main result of this paper.

Theorem 1. *Let K be an infinite field, D be a finite-dimensional central division algebra of degree s over K , and J be a 2-sided ideal in $P_{D,n}$. Then*

- (a) $\mathcal{I}(\mathcal{Z}(J)) = \text{Rad}_D(J)$.
- (b) *Moreover, if $\text{char}(K)$ does not divide $s!$, then $\text{Rad}_D(J) = \text{Rad}'_D(J)$.*

In particular, Theorem 1 tells us that $\text{Rad}_D(J)$ is always a 2-sided ideal in $P_{D,n}$, which is not obvious from the definition. Similarly Theorem 1(b) tells us that $\text{Rad}'_D(J)$ is a 2-sided ideal of $P_{D,n}$, provided that $\text{char}(K)$ does not divide $s!$.

When $D = K$ (i.e., $s = 1$), then $P_{D,n} = C_{D,n} = K[x_1, \dots, x_n]$ is the usual commutative polynomial ring, $N_D(f) = f$, and Theorem 1 reduces to the K -Nullstellensatz (3). Note however that we will not give a new proof in this case. We will assume the K -Nullstellensatz and deduce Theorem 1 from it.

The assumption that K is infinite is made to ensure that formal polynomials are in bijective correspondence with polynomial maps $D^n \rightarrow D$. This assumption is harmless: if K is finite, then by Wedderburn’s Little Theorem, the only finite-dimensional central

division algebra over K is $D = K$ itself. This places us back in the setting of the commutative K -Nullstellensatz (3). Thus we are not missing anything by assuming that K is infinite.

The main weakness of the K -Nullstellensatz (3) is that for most fields K the set of quasi-anisotropic polynomials is difficult to describe explicitly. Theorem 1 tells us that, as in the quaternionic case (4), no additional complications arise in passing from K to D ; see also Proposition 4.

2. PROOF OF THEOREM 1(A)

Let J be a 2-sided ideal of $P_{D,n}$. Set $J_c = J \cap C_{D,n}$, as we did in the Introduction. By [GW, Proposition 17.5], J is generated by J_c as an ideal. In particular,

$$(5) \quad Z_D(J) = Z_K(J_c).$$

Here $Z_D(J)$ is a subset of D^n and $Z_K(J_c)$ is a subset of K^{ns^2} , and we are using a K -basis b_1, \dots, b_{s^2} to identify D^n with K^{ns^2} .

Suppose $f \in \text{Rad}_D(J)$, i.e., $N_D(f) \in \text{Rad}_K(J_c)$. Then there exists a positive integer m , a homogeneous quasi-anisotropic polynomial $p \in K[z_1, \dots, z_m]$ and $f_1, \dots, f_{m-1} \in J_c$ such that $p(f_1, \dots, f_{m-1}, N_D(f)) \in J_c$. Hence, for any $a \in \mathcal{Z}(J_c) = \mathcal{Z}(J)$, we have $p(f_1(a), \dots, f_{m-1}(a), N_D(f)(a)) = 0$. Since p is quasi-anisotropic, this implies $N_D(f)(a) = 0$. Since D is a central division algebra over K , this is only possible if $f(a) = 0$. We conclude that $f(a) = 0$ for any $a \in \mathcal{Z}(J)$. So $f \in \mathcal{I}(\mathcal{Z}(J))$. This shows that $\text{Rad}_D(J) \subset \mathcal{I}(\mathcal{Z}(J))$.

To prove the opposite inclusion, let $f \in \mathcal{I}(\mathcal{Z}(J))$. Write $f = f_1 b_1 + \dots + f_{s^2} b_{s^2}$, where each f_i lies in $C_{D,n}$. To say that f vanishes at every point of $\mathcal{Z}(J)$ is equivalent to saying that each f_i vanishes at every point of $\mathcal{Z}(J)$. In view of (5), the latter is equivalent to each f_i vanishing at every point of $\mathcal{Z}(J_c)$. In other words, $f_i \in \mathcal{I}(\mathcal{Z}(J_c))$ for every $i = 1, \dots, s^2$. By the K -Nullstellensatz (3), $\mathcal{I}(\mathcal{Z}(J_c)) = \text{Rad}_K(J_c)$, so each f_i lies in $\text{Rad}_K(J_c)$. Since $N_D(f)$ is a homogeneous polynomial of degree s in f_1, \dots, f_{s^2} with coefficients in K , we conclude $N_D(f) \in \text{Rad}_K(J_c)$. By the definition of $\text{Rad}_D(J)$, this is equivalent to $f \in \text{Rad}_D(J)$, as desired. \square

Example 2. The quaternionic Nullstellensatz (4) is readily recovered from Theorem 1(a). Indeed, let $K = \mathbb{R}$ be the field of real numbers and $D = \mathbb{H}$ be the algebra of Hamilton quaternions. Following Along and Paran [AP], we define the quaternionic radical of an ideal $J \subset P_{\mathbb{H},n}$ as

$$\text{Rad}_{AP}(J) = \{f \in P_{\mathbb{H},n} \mid \exists m, k \geq 1, \text{ and } f_1, \dots, f_{m-1} \in P_{\mathbb{H},n} \text{ such that} \\ N_{\mathbb{H}}(f_1) + \dots + N_{\mathbb{H}}(f_{m-1}) + N_{\mathbb{H}}(f)^k \in J\}.$$

To recover the quaternionic Nullstellensatz (4) from Theorem 1, it suffices to show that

$$\text{Rad}_{\mathbb{H}}(J) \subset \text{Rad}_{AP}(J) \subset \mathcal{I}(\mathcal{Z}(J)).$$

To prove that $\text{Rad}_{AP}(J) \subset \mathcal{I}(\mathcal{Z}(J))$, suppose that $f \in \text{Rad}_{AP}(J)$. Then there exists $m, k \geq 1$, and $f_1, \dots, f_{m-1} \in P_{\mathbb{H},n}$, such that $N_{\mathbb{H}}(f_1) + \dots + N_{\mathbb{H}}(f_{m-1}) + N_{\mathbb{H}}(f)^k \in J$. So for any $a \in \mathcal{Z}(J)$, we have $(N_{\mathbb{H}}(f_1) + \dots + N_{\mathbb{H}}(f_{m-1}) + N_{\mathbb{H}}(f)^k)(a) = 0$. Since $N_{\mathbb{H}}(b)$ is

non-negative for any $b \in \mathbb{H}$, this implies $f(a) = 0$. In other words, f vanishes at every point of $\mathcal{Z}(J)$, i.e., $f \in \mathcal{I}(\mathcal{Z}(J))$, as desired.

To prove that $\text{Rad}_{\mathbb{H}}(J) \subset \text{Rad}_{AP}(J)$, suppose that $f \in \text{Rad}_{\mathbb{H}}(J)$. Then by our definition of $\text{Rad}_{\mathbb{H}}(J)$, we have $N_{\mathbb{H}}(f) \in \text{Rad}_{\mathbb{R}}(J_c)$. By the Real Nullstellensatz (2), there exists $f_1, \dots, f_{m-1} \in C_{D,n}$ and $m \geq 1$, such that $f_1^2 + \dots + f_{m-1}^2 + N_{\mathbb{H}}(f)^{2k} \in J_c$ for some $k \geq 1$. Since $f_i \in C_{D,n}$, $f_i^2 = N_{\mathbb{H}}(f_i)$. In other words, $N_{\mathbb{H}}(f_1) + \dots + N_{\mathbb{H}}(f_{m-1}) + N_{\mathbb{H}}(f)^{2k} \in J$. Thus $f \in \text{Rad}_{AP}(J)$, as desired. \square

3. PROOF OF THEOREM 1(B)

It is clear from the definition that $\text{Rad}'_D(J) \subset \text{Rad}_D(J)$. Our proof of the opposite inclusion will rely on the following elementary proposition.

Proposition 3. *Let K be a field and $s \geq 1$ be an integer. If $\text{char}(K)$ does not divide $s!$, then the s -powers span the polynomial ring $K[y_1, \dots, y_m]$ as a K -vector space. In other words, any polynomial $f \in K[y_1, \dots, y_m]$, can be written as a K -linear combination of polynomials of the form g^s , where $g \in K[y_1, \dots, y_m]$.*

Proof. We claim that the following identity holds in any commutative ring R :

$$(6) \quad t_1 \dots t_s = \frac{1}{s!} \sum_{I \subset \{1, \dots, s\}} (-1)^{s-|I|} \left(\sum_{i \in I} t_i \right)^s.$$

Here $t_i \in R$ and $|I|$ denotes the number of elements in a subset I of $\{1, \dots, s\}$. Once (6) is established, we can substitute $R = K[y_1, \dots, y_m]$, $t_1 = f$ and $t_2 = \dots = t_s = 1$ to obtain a desired expression for f as a linear combination of s -powers in $K[y_1, \dots, y_m]$.

It thus remains to prove (6). This identity is a special case of the ‘‘polarization formula’’, relating a homogeneous polynomial of degree s to its associated s -linear form; see [SE]. For the sake of completeness, we include a short self-contained proof of (6)¹.

The right hand side of (6) is a homogeneous polynomial of degree s . To determine the coefficient of a monomial M in t_1, \dots, t_s of degree s , let us consider two cases.

Case 1: One of the variables t_1, \dots, t_s does not occur in M . For notational simplicity, let us say this variable is t_s . To calculate the coefficient of M on the right hand side of (6), we may set t_s equal to 0. The index sets $I \subset \{1, \dots, s\}$ not containing s are in bijective correspondence with index sets $I' = I \cup \{s\}$ containing s . Since the size of I' is one greater than the size of I the terms $(\sum_{i \in I} t_i)^s$ and $(\sum_{i \in I'} t_i)^s$ appear with opposite signs in the outer sum. When we substitute $t_s = 0$, these terms will cancel, and the right hand side will sum up to 0. This shows that the coefficient of M on the right hand side of (6) is 0.

Case 2. The only monomial M of degree s not covered by Case 1 is $M = t_1 \dots t_s$. This monomial can only come from the $(t_1 + \dots + t_s)^s$ term on the right hand side of (6), and it comes with coefficient $s!$.

This completes the proof of (6) and thus of Proposition 3. \square

¹A slightly modified version of this same argument can be used to prove the general polarization formula.

We are now ready to finish the proof of Theorem 1(b) by showing that

$$\text{Rad}_D(J) \subset \text{Rad}'_D(J).$$

Let $f \in \text{Rad}_D(J)$. Our goal is to show that $f \in \text{Rad}'_D(J)$. By the definition of $\text{Rad}_D(J)$, there exists a homogeneous quasi-anisotropic $p \in K[z_1, \dots, z_m]$ and $f_1, \dots, f_{m-1} \in C_{D,n}$, such that $p(f_1, \dots, f_{m-1}, N_D(f)) \in J$.

As we explained in the Introduction, after choosing a K -basis b_1, \dots, b_{s^2} in D , we can identify $C_{D,n}$ with the polynomial ring $K[y_{ij}]$ in ns^2 variables. By Proposition 3, there exist finitely many polynomials $g_1, \dots, g_l \in C_{D,n}$, such that each f_i can be written as

$$f_i = c_{i1}g_1^s + \dots + c_{il}g_l^s$$

for some $c_{ij} \in K$. (The proof of Proposition 3 shows that we can take each c_{ij} to be either 0 or $\pm \frac{1}{s!}$, but this will not matter in the sequel.) Let

$$q(w_1, \dots, w_l, w_{l+1}) = p(z_1, \dots, z_{m-1}, y_m),$$

where $z_i = c_{i1}w_1 + \dots + c_{il}w_l$ for each $i = 1, \dots, m-1$ and $z_m = w_{l+1}$. This way we obtain a new homogeneous polynomial $q(w_1, \dots, w_l, w_{l+1})$ in $l+1$ variables. Since p is quasi-anisotropic, it is clear from the definition that q is also quasi-anisotropic and

$$q(g_1^s, \dots, g_l^s, N_D(f)) = p(f_1, \dots, f_{m-1}, N_D(f)) \in J.$$

Since $g_i \in C_{D,n}$, we have $N_D(g_i) = g_i^s$ and thus

$$q(N_D(g_1), \dots, N_D(g_l), N_D(f)) = q(g_1^s, \dots, g_l^s, N_D(f)) \in J.$$

This shows that $f \in \text{Rad}'_D(J)$, as desired. \square

4. D -RADICAL IDEALS

An ideal J in $P_{K,n} = K[x_1, \dots, x_n]$ is called K -radical if $J = \mathcal{Z}(\mathcal{I}(J))$. The following description of K -radical ideals is due to Adkins, Gianni and Tognoli [AGT]: J is K -radical if and only $\text{Rad}_K(J) = J$. This assertion is a precursor to (and, in turn, an easy consequence of) the K -Nullstellensatz (3), due to Laksov.

Now let D be a finite-dimensional division algebra over K . We will say that a two-sided ideal $J \subset P_{D,n}$ is a D -radical if $J = \mathcal{I}(\mathcal{Z}(J))$. In view of Theorem 1 this is equivalent to $J = \text{Rad}_D(J)$. We conclude this paper with the following observation.

Proposition 4. *A two-sided ideal $J \subset P_{D,n}$ is D -radical if and only if $J_c = J \cap C_{D,n}$ is K -radical.*

Proof. Recall that $\mathcal{Z}(J) = \mathcal{Z}(J_c)$ for any 2-sided ideal $J \subset P_{D,n}$; see (5).

First suppose $J \subset P_{D,n}$ is D -radical. Clearly $J_c \subset \mathcal{I}(\mathcal{Z}(J_c))$. To prove the opposite inclusion, assume that $f \in C_{D,n}$ lies in $\mathcal{I}(\mathcal{Z}(J_c))$. Since J is D -radical, we conclude that f lies in J and hence, in $J \cap C_{D,n} = J_c$, as desired.

Conversely, suppose J_c is K -radical. Once again, $J \subset \mathcal{I}(\mathcal{Z}(J))$ for any two-sided ideal $J \subset P_{D,n}$, so we only need to prove the opposite inclusion. Assume that $f \in \mathcal{I}(\mathcal{Z}(J)) \subset P_{D,n}$. Choose a K -basis b_1, \dots, b_{s^2} of D and write $f = f_1b_1 + \dots + f_{s^2}b_{s^2}$, where each f_i lies in $C_{D,n}$. Since f vanishes on $\mathcal{Z}(J) = \mathcal{Z}(J_c)$, so does each f_i . Since J_c is K -radical, we conclude that each f_i lies in J_c and hence, $f = f_1b_1 + \dots + f_{s^2}b_{s^2}$ lies in J . \square

REFERENCES

- [AGT] W. A. Adkins, P. Gianni and A. Tognoli, A Nullstellensatz for an algebraically non-closed field, *Boll. Un. Mat. Ital. B* (5) **15** (1978), no. 1, 338–343. MR0498701
- [AP] G. Alon and E. Paran, A quaternionic Nullstellensatz, *J. Pure Appl. Algebra* **225** (2021), no. 4, 106572, 9 pp. MR4158764
- [GW] K. R. Goodearl and R. B. Warfield, Jr., *An introduction to noncommutative Noetherian rings*, second edition, London Mathematical Society Student Texts, 61, Cambridge University Press, Cambridge, 2004. MR2080008
- [La] D. Laksov, Radicals and Hilbert Nullstellensatz for not necessarily algebraically closed fields, *Enseign. Math. (2)* **33** (1987), no. 3-4, 323–338. MR0925995
- [PD] A. Prestel and C. N. Delzell, *Positive polynomials*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2001. MR1829790
- [SE] Polarization formula,
<https://math.stackexchange.com/questions/481167/polarization-formula>.
- [Wi] D. M. Wilczyński, On the fundamental theorem of algebra for polynomial equations over real composition algebras, *J. Pure Appl. Algebra* **218** (2014), no. 7, 1195–1205. MR3168491

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC, CANADA,
V6T 1Z2

Email address: `zhengheng.bao@alumni.ubc.ca`

Email address: `reichst@math.ubc.ca`