SECOND ORDER ESTIMATES FOR FULLY NONLINEAR ELLIPTIC EQUATIONS WITH GRADIENT TERMS ON HERMITIAN MANIFOLDS

BO GUAN AND XIAOLAN NIE

ABSTRACT. We derive a priori second order estimates for fully nonlinear elliptic equations which depend on the gradients of solutions in critical ways on Hermitian manifolds. The global estimates we obtained apply to an equation arising from a conjecture by Gauduchon which extends the Calabi conjecture; this was one of the original motivations to this work. We were also motivated by the fact that there had been increasing interests in fully nonlinear pde's from complex geometry in recent years, and aimed to develop general methods to cover as wide a class of equations as possible.

Mathematical Subject Classification (MSC2020): 35J15, 35J60, 58J05, 53C21, 53C55.

Keywords: Fully nonlinear elliptic equations, Hermitian manifolds; *a priori* estimates; concave functions and the associated tangent cones at infinity.

1. INTRODUCTION

Let (M^n, ω) be a compact Hermitian manifold of complex dimension $n \ge 2$, and f a symmetric function of n variables. We consider fully nonlinear elliptic equations of the form

(1.1)
$$f(\lambda(\sqrt{-1}\partial\bar{\partial}u + \chi)) = \psi \text{ on } M$$

where χ is a real (1,1) form which may depend on u and its gradient, and $\lambda(X) = (\lambda_1, \dots, \lambda_n)$ denotes the eigenvalues of a (1,1) form X with respect to the metric ω .

The function f is assumed to be defined in a symmetric open and convex cone $\Gamma \subset \mathbb{R}^n$ with vertex at the origin,

(1.2)
$$\Gamma_n = \{\lambda \in \mathbb{R}^n : \lambda_i > 0\} \subset \Gamma,$$

and satisfies the following structure conditions

(1.3)
$$f_i \equiv f_{\lambda_i} = \frac{\partial f}{\partial \lambda_i} > 0 \text{ in } \Gamma, \ 1 \le i \le n$$

Research of the first author was supported in part by NSF grants. The second author was partially supported by NSFC (Grant No. 11801516).

BO GUAN AND XIAOLAN NIE

and

(1.4)
$$f$$
 is a concave function in Γ .

These conditions, first introduced by Caffarelli-Nirenberg-Spruck [6], have become standard in the theory of fully nonlinear PDEs. According to [6], equation (1.1) is elliptic by (1.3) for a function $u \in C^2(M)$ with $\chi_u = \chi + \sqrt{-1}\partial \bar{\partial} u \in \Gamma$; we call such functions *admissible*.

We assume in addition

(1.5)
$$\sup_{\partial \Gamma} f < \psi < \sup_{\Gamma} f$$

where

$$\sup_{\partial \Gamma} f \equiv \sup_{\lambda_0 \in \partial \Gamma} \limsup_{\lambda \to \lambda_0} f(\lambda)$$

in order for equation (1.1) to remain non-degenerate.

Equation (1.1) covers some of the important elliptic equations in complex geometry. In particular, it includes the complex Monge-Ampère equation which has received extensive study from different aspects, going back at least to the work of Aubin [1] and Yau [74] on compact Kähler manifolds in proof of Calabi conjectures, Caffarelli-Kohn-Nirenberg-Spruck [5] on the Dirichlet problem in \mathbb{C}^n , and Bedford-Taylor [2, 3] on weak solutions and pluriportential theory; see e.g. [56] for an excellent survey and references. In recent years, there have been increasing interests from complex geometry in more general fully nonlinear elliptic and parabolic equations, including those of form (1.1) in which χ and ψ may depend on u and/or its gradient. Exciting successes have been achieved in the study of some of these equations, such as the Fu-Yau equation [23, 24] in which $\chi = \chi(z, u)$ and its extensions [51, 53, 55, 9], and an equation treated by Székelyhidi-Tosatti-Weinkove [68] in connection to a conjecture of Gauduchon which will be discussed in more details in Section 5. In |54| and subsequent papers, Phong-Picard-Zhang introduced new geometric flows; see also [50] and references therein for equations from geometry and physics. In this paper we wish to treat equations in the general form (1.1) for $\chi = \chi(\cdot, \partial u, \partial u)$.

Typical examples of function f satisfying conditions (1.3)-(1.4) include $f = \sigma_k^{\frac{1}{k}}$ and more generally $f = (\sigma_k/\sigma_l)^{\frac{1}{k-l}}, 0 \le l < k \le n \ (\sigma_0 = 1)$ defined on the Garding cone

$$\Gamma_k = \{\lambda \in \mathbb{R}^n : \sigma_j(\lambda) > 0, \text{ for } 1 \le j \le k\},\$$

 $\mathbf{2}$

as well as $f = \sigma_k / \sigma_{k-1}$ on Γ_{k-1} for $1 < k \le n$, where

$$\sigma_k(\lambda) = \sum_{1 \le i_1 < \dots < i_k \le n} \lambda_{i_1} \cdots \lambda_{i_k}$$

is the k-th elementary symmetric function. Another important example is

$$f(\lambda) = \sum \tan^{-1} \lambda_i$$

which, deeply rooted in calibration geometry of Harvey-Lawson [37] and theory of special Lagrangian manifolds in the real case, corresponds to the deformed Hermitian-Yang-Mill equation; see [45] as well as e.g. [43, 10, 13, 14, 7, 36, 41, 47] for recent related results.

So far most of attentions to equation (1.1) from a more PDE point of view have been on the case where χ is independent of u and its gradient. In [46] Li treated the Dirichlet problem in \mathbb{C}^n . For $f = \sigma_k^{\frac{1}{k}}$ and $\chi = \omega$ on a compact Kähler manifold, equation (1.1) was first studied by Hou-Ma-Wu [40] who derived second order estimates of the form

(1.6)
$$\max_{M} |\partial \bar{\partial} u| \le C(1 + \max_{M} |\nabla u|^2),$$

followed by Dinew-Kolodziej [16] who used a blow-up argument and Liouville type theorem to derive a gradient bound from (1.6). In [61], Song-Weinkove introduced a necessary and sufficient cone condition to solve equation (1.1) for $f = \sigma_n/\sigma_{n-1}$, known as J-equation, in answer to a question raised by Donaldson [17] in the Kähler case where χ is another Kähler metric on M and ψ is an invariant constant given by

$$\psi = \frac{\int_M \chi^n}{\int_M \chi^{n-1} \wedge \omega};$$

see also [72, 73]. Their results were extended to $f = (\sigma_n/\sigma_l)^{\frac{1}{n-l}}$, $1 \leq l < n$ by Fang-Lai-Ma [19] and by Sun [64, 65] who treated general ψ . From a point of view in analogue to Yau-Tian-Donaldson stability [75, 69, 18], Lejmi-Székelyhidi [44] proposed as a conjecture an integral condition for J-equation and solved it when n = 2. In higher dimensions, the conjecture was proved for toric manifolds by Collins-Székelyhidi [12] who also treated Krylov type equations, extending results of [19] and Zheng [78], and by Song [60] in the general case; see also Chen [7] and Datar-Pingali [15].

For equation (1.1) on compact Hermitian manifolds, Zhang [77] and Sun [66] solved the cases $f = \sigma_k^{\frac{1}{k}}$ and $f = (\sigma_k/\sigma_l)^{\frac{1}{k-l}}$ respectively, and Székelyhidi [67] gave a thorough treatment for general f and $\psi = \psi(z)$, under the assumption in addition to (1.3)-(1.5),

(1.7)
$$\sum f_i \lambda_i \ge 0, \quad \sum f_i \ge c_0 > 0.$$

A crucial ingredient in solving equation (1.1) is to establish a priori bounds for second derivatives of admissible solutions, on which we shall focus in this paper. Our main results will be stated in Section 2; see Theorems 2.2, 2.9 and 2.14. We obtain in Theorems 2.2 an interior estimates for second derivatives under an assumption on χ (see (2.5)) analogous to Ma-Trudinger-Wang condition [49] in optimal transport theory. Theorems 2.9 and 2.14 concern global second order estimates in which the primary assumptions are in terms of the tangent cone at infinity introduced in [27] for the level hypersurface $\{f = \psi\}$ which is smooth and convex by conditions (1.3) and (1.4). In the case $\chi = \chi(z)$, i.e. when it is independent of ∂u and $\bar{\partial} u$, Theorem 2.9 recovers the second order estimates of Székelyhidi [67] without assumption (1.7) for type I cones (as defined in Caffarelli-Nirenberg-Spruck [6]); for details see Section 2.

In what follows we describe an application of Theorem 2.9 which was one of the original motivations to our work in the current article.

Let ω_0 be another Hermitian metric on M^n . We consider the Monge-Ampère formtype equation

(1.8)
$$\det \Phi_u = e^{h+b} \det(\omega^{n-1}) \text{ in } M$$

with

(1.9)
$$\Phi_u = \omega_0^{n-1} + \sqrt{-1}\partial\bar{\partial}u \wedge \omega^{n-2} + c \Re \left\{\sqrt{-1}\partial u \wedge \bar{\partial}\omega^{n-2}\right\} > 0$$

where h is a given function, and b, c are constant.

Equation (1.8) is related to the general notion of plurisubharmonic functions of Harvey-Lawson [38, 39]. It was studied by Fu-Wang-Wu [21, 22], Tosatti-Weinkove [70, 71] for c = 0, and by Székelyhidi-Tosatti-Weinkove [68] for c = 1, proving a conjecture of Gauduchon [25] (see also Conjecture 1.3 in [71]). As a consequence of Theorem 2.9 we obtain

Corollary 1.1. Let $u \in C^4(M)$ be a solution of equation (1.8) satisfying (1.9). Then (1.6) holds for some constant C depending on $|u|_{C^0(M)}$ and known data.

It was proved by Tosatti-Weinkove [71] that the solvability of equation (1.8) reduces to establishing (1.6); see also Székelyhidi [67]. In their joint paper [68], these authors carried out such an estimate. Among other examples of f satisfying (1.3)-(1.4) are $f = \log \rho_k$, $1 \le k \le n$, where

$$\rho_k(\lambda) := \prod_{1 \le i_1 < \dots < i_k \le n} (\lambda_{i_1} + \dots + \lambda_{i_k})$$

defined in the cone

$$\mathcal{P}_k := \{ \lambda \in \mathbb{R}^n : \lambda_{i_1} + \dots + \lambda_{i_k} > 0, \ \forall \ 1 \le i_1 < \dots < i_k \le n \}$$

In particular, $\rho_1 = \sigma_n$ and $\rho_n = \sigma_1$.

We note that in the literature of nonlinear PDE, a domain $\Omega \in \mathbb{R}^n$ or its boundary $\partial \Omega$ is often called k-convex or k-mean convex if $\kappa \in \Gamma_k$ where $\kappa = (\kappa_1, \ldots, \kappa_{n-1})$ denotes the principal curvatures of $\partial \Omega$, while in areas of geometry/geometric analysis such as mean curvature flows, that a hypersurface Σ in \mathbb{R}^n is k-convex sometimes means that its principal curvatures $\kappa \in \mathcal{P}_k$. The latter concept was first introduced in 1986 by Sha [59]. In [42] and [4], Brandle, Huisken and Sinestrari studied regularity of mean curvature flow of two-convex hypersurfaces in \mathbb{R}^n .

Using the Hodge star operator, Tosatti-Weinkove [70] were able to rewrite equation (1.8) in the form of (1.1) for $f = \log \rho_{n-1}$. We shall derive Corollary 1.1 by verifying the conditions in Theorem 2.9 for this equation.

In [52], Phong-Picard-Zhang considered equation (1.1) in which $\psi = \psi(z, u, \nabla u)$ for $f = \sigma_k^{\frac{1}{k}}$ and $\chi = \chi(z, u) \ge c_0 \omega > 0$ on a compact Kähler manifold (M^n, ω) , and derived second order estimates which allow arbitrary dependence of ψ on ∇u , in the spirit of Guan-Ren-Wang [34]. We obtain similar results under assumption (2.5); see Remarks 2.3 and ??.

The Dirichlet problem for equation (1.1) on Hermitian manifolds was treated for $f = (\sigma_n/\sigma_l)^{\frac{1}{n-l}}, 1 \leq l < n$ by Li, Sun and the first author [31, 33] where gradient estimates were derived directly using the maximum principle method, and more recently by Collins-Picard [11] for $f = \sigma_k^{\frac{1}{k}}, 1 < k < n$ and Feng-Ge-Zheng [20] for Hessian quotient equations. See also [32, 58, 76] where equations with gradient dependence were studied. For general f it seems still open, while the corresponding Dirichlet problem on Riemannian manifolds was studied in [28] for $\chi = \chi(z), \psi = \psi(z)$ and [29, 30] in the more general case $\chi = \chi(z, u, \nabla u), \psi = \psi(z, u, \nabla u)$.

The article is organized as follows. Section 2 contains the statements of our main theorems. In Section 3 we prove a key inequality needed in the proof of Theorem 2.14 for f, extending a result in [27]. Section 4 is devoted to second derivative estimates, completing the proof of the main results while in Section 5 we prove Corollary 1.1.

BO GUAN AND XIAOLAN NIE

Part of research described in this paper was done while the second author was a Ross Assistant Professor in Department of Mathematics at The Ohio State University. More recently we were able to refine some of the calculations, so hopefully it was more accessible to the reader and especially those new to the field. We also up-dated the references, adding papers appeared more recently in the field. We wish to thank Gabor Székelyhidi, Valentino Tosatti and Ben Weinkove for communications on their results in [68].

2. NOTATIONS AND MAIN RESULTS

Throughout the paper we write in local coordinates (z_1, \ldots, z_n)

$$\omega = \sqrt{-1}g_{i\bar{j}}\,dz_i \wedge d\bar{z}_j.$$

Let $\{g^{i\bar{j}}\} = \{g_{i\bar{j}}\}^{-1}$ denote the inverse matrix of $\{g_{i\bar{j}}\} > 0$.

For fixed $z \in M$, $h \in \mathbb{R}$ and (1,0) form ζ , we use the notation

$$\chi(z,\zeta,\bar{\zeta},h) = \sqrt{-1}\chi_{i\bar{j}}(z,\zeta,\bar{\zeta},h)dz_i \wedge d\bar{z}_j$$

or simply $\chi = \sqrt{-1}\chi_{i\bar{j}}dz_i \wedge d\bar{z}_j$, and

$$\chi_{\xi\bar{\eta}} = \sum \chi_{i\bar{j}}\xi_i\bar{\eta}_j$$

for

$$\xi = \sum \xi_i \frac{\partial}{\partial z_i}, \ \eta = \sum \eta_i \frac{\partial}{\partial z_i}.$$

For a function $u \in C^2(M)$, we write $\chi[u] = \chi(\cdot, \nabla u, u)$ or $\chi[u] = \chi(\cdot, \partial u, \bar{\partial} u, u)$ to indicate the dependence of χ on u and its gradient, and

$$\mathfrak{g}_{i\bar{j}} = \chi_{i\bar{j}}[u] + \nabla_{\bar{j}}\nabla_i u = \chi_{i\bar{j}}[u] + \partial_{\bar{j}}\partial_i u$$

 \mathbf{SO}

$$\chi_u := \chi[u] + \sqrt{-1} \partial \bar{\partial} u = \sqrt{-1} \mathfrak{g}_{i\bar{j}} \, dz_i \wedge d\bar{z}_j.$$

In the current paper we shall consider the case $\psi = \psi(z)$, $\chi[u] = \chi(z, \partial u, \bar{\partial} u)$, and use the following expressions to distinguish different derivatives

$$\chi_{i\bar{j},k} = \frac{\partial \chi_{i\bar{j}}}{\partial z_k}, \ \chi_{i\bar{j}k} = \chi_{i\bar{j},k} - \Gamma^p_{ki}\chi_{p\bar{j}}, \ \chi_{i\bar{j}k\bar{l}} = \chi_{i\bar{j}k,\bar{l}} - \overline{\Gamma^q_{lj}}\chi_{i\bar{q}k},$$
$$\chi_{i\bar{j},\zeta_\alpha} = \frac{\partial \chi_{i\bar{j}}}{\partial \zeta_\alpha}, \ \chi_{i\bar{j},\zeta_\alpha k} = \frac{\partial^2 \chi_{i\bar{j}}}{\partial z_k \partial \zeta_\alpha}, \ \chi_{i\bar{j}k,\zeta_\alpha} = \frac{\partial \chi_{i\bar{j}k}}{\partial \zeta_\alpha},$$

as well as

(2.1)
$$\partial_k \chi_{i\bar{j}}[u] = \chi_{i\bar{j},k}[u] + \chi_{i\bar{j},\zeta_\alpha}[u]\partial_k\partial_\alpha u + \chi_{i\bar{j},\bar{\zeta}_\alpha}[u]\partial_k\partial_{\bar{\alpha}} u$$

 $\mathbf{6}$

and

$$(2.2) \quad \nabla_k \chi_{i\bar{j}}[u] = \partial_k \chi_{i\bar{j}}[u] - \Gamma^p_{ki} \chi_{p\bar{j}}[u] = \chi_{i\bar{j}k}[u] + \chi_{i\bar{j},\zeta_\alpha}[u]\partial_k \partial_\alpha u + \chi_{i\bar{j},\bar{\zeta}_\alpha}[u]\partial_k \partial_{\bar{\alpha}} u$$

etc; we shall drop [u] in the expressions when no confusions would arise. Similarly,

(2.3)
$$\nabla_{\bar{l}}\chi_{i\bar{j}k} = \chi_{i\bar{j}k\bar{l}} + \chi_{i\bar{j}k,\zeta_{\alpha}}\partial_{\bar{l}}\partial_{\alpha}u + \chi_{i\bar{j}k,\bar{\zeta}_{\alpha}}\partial_{\bar{l}}\partial_{\bar{\alpha}}u$$

and

$$(2.4) \qquad \nabla_{\bar{l}}\chi_{i\bar{j},\zeta_{\alpha}} = \partial_{\bar{l}}\chi_{i\bar{j},\zeta_{\alpha}} + \overline{\Gamma}_{lj}^{q}\chi_{i\bar{q},\zeta_{\alpha}} \\ = \chi_{i\bar{j},\zeta_{\alpha}\bar{l}} + \chi_{i\bar{j},\zeta_{\alpha}\zeta_{\beta}}\partial_{\bar{l}}\partial_{\beta}u + \chi_{i\bar{j},\zeta_{\alpha}\zeta_{\bar{\beta}}}\partial_{\bar{l}}\partial_{\bar{\beta}}u + \overline{\Gamma}_{lj}^{q}\chi_{i\bar{q},\zeta_{\alpha}} \\ = \chi_{i\bar{j},\bar{l}\zeta_{\alpha}} + \chi_{i\bar{j},\zeta_{\alpha}\zeta_{\beta}}\partial_{\bar{l}}\partial_{\beta}u + \chi_{i\bar{j},\zeta_{\alpha}\zeta_{\bar{\beta}}}\partial_{\bar{l}}\partial_{\bar{\beta}}u + \overline{\Gamma}_{lj}^{q}\chi_{i\bar{q},\zeta_{\alpha}} \\ = \chi_{i\bar{j}\bar{l},\zeta_{\alpha}} + \chi_{i\bar{j},\zeta_{\alpha}\zeta_{\beta}}\partial_{\bar{l}}\partial_{\beta}u + \chi_{i\bar{j},\zeta_{\alpha}\zeta_{\bar{\beta}}}\partial_{\bar{l}}\partial_{\bar{\beta}}u.$$

Example 2.1. We explain the above notation by a simple example. Let

$$\chi[u] = \partial u \wedge \bar{\partial} u = \partial_i u \bar{\partial}_j u \, dz_i \wedge d\bar{z}_j.$$

Then

$$\chi_{i\bar{j},k} = 0, \quad \chi_{i\bar{j}k} = -\Gamma^p_{ki}\partial_p u\bar{\partial}_j u, \quad \chi_{i\bar{j},\zeta_\alpha} = \delta_{i\alpha}\bar{\partial}_j u, \quad \chi_{i\bar{j},\bar{\zeta}_\alpha} = \delta_{j\alpha}\partial_i u,$$

and

$$\nabla_k \chi_{i\bar{j}}[u] = -\Gamma^p_{ki} \partial_p u \bar{\partial}_j u + \partial_k \partial_i u \bar{\partial}_j u + \partial_i u \partial_k \bar{\partial}_j u, = \nabla_k \nabla_i u \nabla_{\bar{j}} u + \partial_i u \partial_k \bar{\partial}_j u.$$

In our first result we establish an interior estimate for second derivatives, which requires the additional assumption that there exists $c_0 > 0$ such that

(2.5)
$$\sum \chi_{i\bar{j},\zeta_k\bar{\zeta}_l}(z,\cdot,\cdot)\xi_i\xi_{\bar{j}}\eta_k\bar{\eta}_l \le -c_0|\xi|^2|\eta|^2, \ \forall \xi,\eta \in T_z^{1,0}M, \ \omega(\xi,\bar{\eta}) = 0.$$

This is an analogue of assumption (A3) of Ma-Trudinger-Wang [49].

Theorem 2.2. Let $u \in C^{4,\alpha}(B_R)$ be an admissible solution of equation (1.1) in a geodesic ball $B_R \subset M$ of radius R, where $0 < \alpha < 1$. Assume $\psi = \psi(z)$ and that (1.3)-(1.5), (2.5) holds. Then u satisfies the interior a priori estimate

$$(2.6) |\nabla^2 u|_{C^{\alpha}(B_{R/2})} \le C$$

where C depends on c_0 , R^{-1} , $|\nabla u|_{C^1(\overline{B}_{3R/4})}$ and

$$c_1 = \sup_{\Gamma} f - \sup_{B_R} \psi > 0.$$

Remark 2.3. Theorem 2.2 still holds for $\psi = \psi(z, u, \nabla u)$ provided in addition that either $\psi(z, u, \nabla u)$ is convex in ∇u or

(2.7)
$$\lim_{|\lambda| \to +\infty} \sum f_i = +\infty, \ \forall \ \lambda \in \Gamma.$$

This holds for $f = \sigma_k^{\frac{1}{k}}$ on $\Gamma_k, k > 1$

It is well known that in general there are no interior second order estimates for fully nonlinear elliptic equations. In particular, Theorem 2.2 fails without condition (2.5), even in the case $\chi = \chi(z)$.

Turning to global second derivative estimates where assumption (2.5) is dropped, we first recall some notions from [27]. For $\sigma \in (\sup_{\partial \Gamma} f, \sup_{\Gamma} f)$ define

$$\Gamma^{\sigma} = \{\lambda \in \Gamma : f(\lambda) > \sigma\}.$$

By (1.3) and (1.4), $\partial\Gamma^{\sigma} = \{\lambda \in \Gamma : f(\lambda) = \sigma\}$ is a smooth and convex complete hypersurface in Γ . For $\lambda \in \partial\Gamma^{\sigma}$ let $\nu_{\lambda} = Df(\lambda)/|Df(\lambda)|$ denote the unit normal vector to $\partial\Gamma^{\sigma}$ at λ .

Definition 2.4 ([27]). For $\mu \in \mathbb{R}^n$ let

$$S^{\sigma}_{\mu} = \{\lambda \in \partial \Gamma^{\sigma} : \nu_{\lambda} \cdot (\mu - \lambda) \le 0\}.$$

The tangent cone at infinity to Γ^{σ} is defined as

$$\mathcal{C}_{\sigma}^{+} = \{ \mu \in \mathbb{R}^{n} : S_{\mu}^{\sigma} \text{ is compact} \}.$$

Clearly, \mathcal{C}^+_{σ} is a symmetric convex cone, and $\Gamma^{\sigma} \subset \mathcal{C}^+_{\sigma}$. The following results were proved in [27].

Theorem 2.5 ([27]). a) C^+_{σ} is open. b) Let $\mu \in C^+_{\sigma}$. There exist $\varepsilon, R > 0$ such that (2.8) $f_i(\lambda)(\mu_i - \lambda_i) \ge \varepsilon \sum f_i(\lambda) + \varepsilon, \quad \forall \ \lambda \in \partial \Gamma^{\sigma} \setminus B_R(0).$

We next introduce a new quantity which will also play a key role in a forthcoming paper [26].

Definition 2.6. The rank of C_{σ}^+ is defined to be

 $\min\{r(\nu) : \nu \text{ is the unit normal vector of a supporting plane to } C_{\sigma}^+\}.$

where for a unit vector $\nu \in \overline{\Gamma}_n$, $r(\nu)$ denotes the number of non-zero components of ν . For convenience we define the rank of \mathbb{R}^n to be n.

Remark 2.7. For $f = \sigma_k^{\frac{1}{k}}$ defined on Γ_k , the rank of \mathcal{C}_{σ}^+ is n - k + 1. This follows from an inequality of Lin-Trudinger [48].

It is also easy to see

Lemma 2.8. For $f = \log \rho_k$ defined on \mathcal{P}_k , $1 \le k \le n$, the rank of \mathcal{C}_{σ}^+ is k.

Our second main result may be stated as follows.

Theorem 2.9. Let $u \in C^4(M)$ be an admissible solution of (1.1) with $\psi \in C^2(M)$ and let

$$r_0 = \min \{ \operatorname{rank} of \mathcal{C}^+_{\psi(z)} : z \in M \}.$$

Assume in addition to (1.3)-(1.5) that

(2.9)
$$\chi_{\xi\bar{\xi}}(\cdot,p)$$
 is a concave function in $p \in T_z^*(M), \ \forall \ \xi \in T_z^{1,0}M$

where $T_z^*(M)$ denotes the real cotangent space of M at z, and that at any point on M where in local coordinates $g_{i\bar{j}} = \delta_{ij}$ and $\mathfrak{g}_{i\bar{j}} = \delta_{ij}\lambda_i$ with $\lambda_1 \geq \cdots \geq \lambda_n$,

(2.10)
$$\left|\sum f_i \chi_{i\bar{i}\bar{1},\zeta_{\alpha}}\right| + \sum f_i |\chi_{i\bar{1},\zeta_{\alpha}}|^2 \le C\lambda_1 f_{\alpha}, \quad \forall \alpha \le n - r_0.$$

Then the following estimate holds

(2.11)
$$\max_{M} |\partial \bar{\partial} u| \le C_1 e^{C_2(u - \inf_M u)}$$

where C_1 depends on $|\nabla u|_{C^0(M)}$ and C_2 is a uniform constant, provided that there exists a function $\underline{u} \in C^2(M)$ satisfying

(2.12)
$$\lambda(\chi_{\underline{u}}(z)) \in \mathcal{C}^+_{\psi(z)} \quad \forall \ z \in M$$

Remark 2.10. For $\chi = \chi(z)$ we obtain (2.11) under conditions (1.3)-(1.5) and (2.12).

Remark 2.11. In [67], Székelyhidi introduced the notation of C-subsolution which has been used widely. A function $\underline{u} \in C^2(M)$ is called a C-subsolution if

(2.13)
$$(\lambda(\chi_{\underline{u}}(z)) + \Gamma_n) \cap \partial \Gamma^{\psi(z)} \text{ is compact}, \quad \forall z \in M.$$

Among other interesting results, he derived the second order estimate assuming (1.3)-(1.5), (1.7) and the existence of a C-subsolution.

Remark 2.12. It was shown in [28] that if Γ is a type I cone, then conditions (2.12) and (2.13) are equivalent. A cone Γ is of type I if each positive λ_i -axis belongs to the boundary of Γ ; see Caffarelli-Nirenberg-Spruck [6] for definition. For k > 1, Γ_k is a type I cone while Γ_1 is a type II cone, meaning not of type I.

Remark 2.13. For $f = (\sigma_n/\sigma_l)^{\frac{1}{n-l}}$, $1 \leq l < n$, conditions (2.12) and (2.13) are all equivalent to the cone condition of Song-Weinkove [61] and Fang-Lai-Ma [19]; see e.g. [67].

We now introduce a larger cone containing C_{σ}^+ . Note that the unit normal vector of any supporting hyperplane to C_{σ}^+ belongs to $\overline{\Gamma_n}$. We define \widetilde{C}_{σ}^+ to be the region in \mathbb{R}^n bounded by those supporting hyperplanes to C_{σ}^+ with unit normal vector in $\partial \Gamma_n$; so $\widetilde{C}_{\sigma}^+ = \mathbb{R}^n$ if there are no such supporting planes.

so $\widetilde{\mathcal{C}}_{\sigma}^{+} = \mathbb{R}^{n}$ if there are no such supporting planes. Clearly, the rank of $\widetilde{\mathcal{C}}_{\sigma}^{+}$ is equal to that of \mathcal{C}_{σ}^{+} . Moreover, $\mu + \Gamma_{n} \subset \widetilde{\mathcal{C}}_{\sigma}^{+}$ for $\mu \in \widetilde{\mathcal{C}}_{\sigma}^{+}$ and $\widetilde{\mathcal{C}}_{\sigma}^{+} \subset \widetilde{\mathcal{C}}_{\rho}^{+}$ if $\sigma \geq \rho$.

We have the following extension of Theorem 2.9.

Theorem 2.14. Suppose that f satisfies the additional assumption

(2.14)
$$\sum f_i \ge c_0 > 0 \quad in \ \{\lambda \in \Gamma : \inf_M \psi \le f(\lambda) \le \sup_M \psi\}.$$

Theorem 2.9 then still holds with assumption (2.12) replaced by

(2.15)
$$\lambda(\chi_{\underline{u}}(z)) \in \widetilde{\mathcal{C}}_{\psi(z)}^+ \quad \forall \ z \in M.$$

3. The cone $\widetilde{\mathcal{C}}_{\sigma}^+$ and extension of Theorem 2.5

A crucial tool in the proof of Theorem 2.9 is Theorem 2.5. To prove Theorem 2.14, we need the following extension to $\widetilde{\mathcal{C}}_{\sigma}^+$ in a slightly weaker form.

Theorem 3.1. Let $\mu \in \widetilde{\mathcal{C}}_{\sigma}^+$. There exist $\delta, \varepsilon > 0$ such that for any $\lambda \in \partial \Gamma^{\sigma}$, either (3.1) $f_k(\lambda) \ge \delta \sum f_i(\lambda), \quad \forall k$

or

(3.2)
$$\sum f_i(\lambda)(\mu_i - \lambda_i) \ge \varepsilon \sum f_i(\lambda).$$

Proof. We prove by contradiction. Assume Theorem 3.1 is false. There exists $\tilde{\mu} \in \widetilde{C}_{\sigma}^+$ and $\lambda_k \in \partial \Gamma^{\sigma}$ for each positive integer k such that

(3.3)
$$\sum f_i(\lambda_k)(\tilde{\mu}_i - (\lambda_k)_i) \le \frac{1}{k\sqrt{n}} \sum f_j(\lambda_k) \le \frac{1}{k} |\nabla f(\lambda_k)|,$$

(3.4)
$$\min_{i} \{f_i(\lambda_k)\} \le \frac{1}{k\sqrt{n}} \sum f_j(\lambda_k) \le \frac{1}{k} |\nabla f(\lambda_k)|$$

and therefore

(3.5)
$$\operatorname{dist}(\nu_{\lambda_k}, \partial \Gamma_n) = \min_i \frac{f_i(l_k)}{|\nabla f(l_k)|} \le \frac{1}{k}.$$

By the concavity of f, for any $\mu \in \Gamma^{\sigma}$ we have

(3.6)
$$0 < f(\mu) - f(\lambda_k) \le \sum f_i(\lambda_k)(\mu_i - (\lambda_k)_i).$$

Thus $\nu_{\lambda_k} \cdot \lambda_k \leq |\mu|$ (and is therefore uniformly bounded above).

On the other hand, from (3.3) we see that

$$\nu_{\lambda_k} \cdot \lambda_k \ge \nu_{\lambda_k} \cdot \tilde{\mu} - \frac{1}{k} \ge -|\tilde{\mu}| - 1.$$

This shows that $\{\nu_{\lambda_k} \cdot \lambda_k\}$ is bounded from below.

Consequently, passing to a subsequence we may assume

(3.7)
$$\lim_{k \to \infty} \nu_{\lambda_k} = \nu$$

(3.8)
$$\lim_{k \to \infty} \nu_{\lambda_k} \cdot \lambda_k = c$$

We have $\nu \in \partial \Gamma_n$ by (3.5), and it follows from (3.6) and (3.8) that Γ^{σ} is contained in the half space

$$H_{\nu}^+ = \{ \mu \in \mathbb{R}^n : \nu \cdot \mu > c \}.$$

As $\nu \in \partial \Gamma_n$, this implies $\mathcal{C}^+_{\sigma} \subset H^+_{\nu}$ and therefore $\widetilde{\mathcal{C}}^+_{\sigma} \subset H^+_{\nu}$. Moreover, it follows from (3.3) that $\nu \cdot \tilde{\mu} = c$ showing $\tilde{\mu} \in \partial H^+_{\nu}$. Consequently, $\tilde{\mu} \in \partial \widetilde{\mathcal{C}}^+_{\sigma}$ which is a contradiction.

It would be desirable to improve (3.2) to (2.8); whether this is possible, however, is not clear to us at the moment. For fixed $\mu \in \widetilde{C}_{\sigma}^+$ let $\varepsilon > 0$ satisfy (3.2) in Theorem 3.1 and denote

$$A = \{\lambda \in \partial \Gamma^{\sigma} : (3.2) \text{ holds} \}$$

For $\lambda \in A$ let

$$t_{\lambda} = \min\{t \ge 0 : t\lambda + (1-t)\mu \in \overline{\Gamma^{\sigma}}\}.$$

We see that $t_{\lambda} < 1$, for otherwise (3.2) would not hold. As in [27] by the concavity of f,

$$\sum f_i(\lambda)(\mu_i - \lambda_i) \ge \sup_{t_\lambda \le t \le 1} f(t\lambda + (1-t)\mu) - \sigma > 0$$

for $\lambda \in A$, as otherwise $t\lambda + (1-t)\mu \in \partial \Gamma^{\sigma}$ for all $t_{\lambda} \leq t \leq 1$ which contradicts (3.2). Clearly for r large,

$$h_{\mu}(r) := \min_{\lambda \in A \cap \partial B_{r}(0)} \sup_{t_{\lambda} \le t \le 1} f(t\lambda + (1-t)\mu) - \sigma > 0$$

since $A \cap \partial B_r(0)$ is compact. We end this section with the following question: Is $h_{\mu}(r)$ nondecreasing in r? A positive answer to this question would give an improvement of Theorem 3.1 which enables us to drop assumption (2.14) in Theorem 2.14.

BO GUAN AND XIAOLAN NIE

4. The second order estimates

In this section we derive the second order estimates in Theorems 2.2 and 2.9. Throughout the section, we use ∇ to denote the Chern connection of (M, ω) and let $u \in C^4(M)$ be an admissible solution of equation (1.1).

In local coordinates $z = (z_1, \ldots, z_n)$, equation (1.1) can be written in the form

(4.1)
$$F(\mathfrak{g}_{i\bar{j}}) = \psi$$

where the function F is defined by $F(X) = f(\lambda(X))$ for a real (1, 1) form X on M. As usual we denote

$$F^{i\bar{j}} = \frac{\partial F}{\partial \mathfrak{g}_{i\bar{j}}}, \ F^{i\bar{j},k\bar{l}} = \frac{\partial^2 F}{\partial \mathfrak{g}_{k\bar{l}}\partial \mathfrak{g}_{i\bar{j}}}$$

We use an idea of Tosatti-Weinkove [70] and consider the quantity which is given in local coordinates

(4.2)
$$A := \sup_{z \in M} \max_{\xi \in T_z^{1,0}M} e^{(1+\gamma)\phi} \mathfrak{g}_{p\bar{q}} \xi_p \bar{\xi}_q (g^{k\bar{l}} \mathfrak{g}_{l\bar{l}} \mathfrak{g}_{k\bar{j}} \xi_i \bar{\xi}_j)^{\frac{\gamma}{2}} / |\xi|^{2+\gamma}$$

where ϕ is a function depending on u and $|\nabla u|$, and $\gamma > 0$ is a small constant to be determined; one may as well follow the approach of Szekelyhidi [67]. Assume that Ais achieved at an interior point $z_0 \in M$ for some $\xi \in T_{z_0}^{1,0}M$, $|\xi| = 1$. We choose local coordinates around z_0 such that $g_{i\bar{j}} = \delta_{ij}$ and that $\mathfrak{g}_{i\bar{j}}$ is diagonal at z_0 with

$$\mathfrak{g}_{1\bar{1}} \geq \mathfrak{g}_{2\bar{2}} \geq \cdots \geq \mathfrak{g}_{n\bar{n}}.$$

We shall assume $\mathfrak{g}_{1\overline{1}} \geq 1$; otherwise we are done.

As pointed out in [70], an important fact is that when γ is chosen sufficient small we have $\xi = \partial_1$ and $A = e^{(1+\gamma)\phi} \mathfrak{g}_{1\bar{1}}^{1+\gamma}$ at z_0 . Indeed, this is obvious if $\mathfrak{g}_{n\bar{n}} \geq -\mathfrak{g}_{1\bar{1}}$. Suppose $\mathfrak{g}_{n\bar{n}} < -\mathfrak{g}_{1\bar{1}}$. Then $n \geq 3$ and $(n-1)\mathfrak{g}_{1\bar{1}} \geq -\mathfrak{g}_{n\bar{n}}$ since

$$\sum \mathfrak{g}_{k\bar{k}} \ge 0.$$

Clearly, $\xi_i = 0$ for 1 < i < n and $\xi_1^2 + \xi_n^2 = 1$. It follows that

$$Ae^{-(1+\gamma)\phi} \le \mathfrak{g}_{1\bar{1}}\xi_1^2(\mathfrak{g}_{1\bar{1}}^2\xi_1^2 + \mathfrak{g}_{n\bar{n}}^2\xi_n^2)^{\frac{\gamma}{2}} \le \xi_1^2(\xi_1^2 + (n-1)^2\xi_n^2)^{\frac{\gamma}{2}}\mathfrak{g}_{1\bar{1}}^{1+\gamma} \le \mathfrak{g}_{1\bar{1}}^{1+\gamma}$$

provided that $\gamma \leq \frac{2}{n(n-2)}$. This shows that $\xi_n = 0$.

Let $W = g_{1\bar{1}}^{-1} g^{k\bar{l}} \mathfrak{g}_{1\bar{l}} \mathfrak{g}_{k\bar{1}}$. The function $e^{(1+\gamma)\phi} g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{1}} W^{\frac{\gamma}{2}}$ which is locally well defined attains a maximum $A = (e^{\phi} \mathfrak{g}_{1\bar{1}})^{1+\gamma}$ at z_0 . It follows that at z_0

(4.3)
$$\frac{\frac{\partial_i(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}})}{\mathfrak{g}_{1\bar{1}}} + \frac{\gamma\partial_i W}{2W} + (1+\gamma)\partial_i\phi = 0,}{\frac{\bar{\partial}_i(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}})}{\mathfrak{g}_{1\bar{1}}} + \frac{\gamma\bar{\partial}_i W}{2W} + (1+\gamma)\bar{\partial}_i\phi = 0}$$

for each $1 \leq i \leq n$, and

(4.4)
$$0 \geq \frac{1}{\mathfrak{g}_{1\bar{1}}} F^{i\bar{i}} \bar{\partial}_i \partial_i (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{1}}) - \frac{1}{\mathfrak{g}_{1\bar{1}}^2} F^{i\bar{i}} \partial_i (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{1}}) \bar{\partial}_i (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{1}}) \\ + \frac{\gamma}{2W} F^{i\bar{i}} \bar{\partial}_i \partial_i W - \frac{\gamma}{2W^2} F^{i\bar{i}} \partial_i W \bar{\partial}_i W + (1+\gamma) F^{i\bar{i}} \bar{\partial}_i \partial_i \phi.$$

Recall that in local coordinates the Christoffel symbols Γ^k_{ij} are defined by

$$\nabla_{\frac{\partial}{\partial z_i}} \frac{\partial}{\partial z_j} = \Gamma^k_{ij} \frac{\partial}{\partial z_k}$$

 \mathbf{SO}

$$\Gamma^{k}_{ij} = g^{k\bar{l}} \frac{\partial g_{j\bar{l}}}{\partial z_i} = g^{k\bar{l}} \partial_i g_{j\bar{l}}$$

and

$$\partial_i g_{j\bar{k}} = g_{l\bar{k}} \Gamma^l_{ij}, \ \bar{\partial}_i g_{j\bar{k}} = g_{j\bar{l}} \overline{\Gamma^l_{ik}}.$$

The torsion and curvature tensors are given by

(4.5)
$$T_{ij}^k = \Gamma_{ij}^k - \Gamma_{ji}^k$$

and, respectively,

(4.6)
$$R_{i\bar{j}k\bar{l}} = -g_{m\bar{l}}\frac{\partial\Gamma^m_{ik}}{\partial\bar{z}_j} = -\frac{\partial g_{k\bar{l}}}{\partial z_i\partial\bar{z}_j} + g^{p\bar{q}}\frac{\partial g_{k\bar{q}}}{\partial z_i}\frac{\partial g_{p\bar{l}}}{\partial\bar{z}_j}.$$

By a lemma of Streets-Tian [63], we may assume $T_{ij}^k = 2\Gamma_{ij}^k$ at z_0 .

We record the following basic formulas

$$\partial_i g^{k\bar{l}} = -g^{k\bar{q}} g^{p\bar{l}} \partial_i g_{p\bar{q}} = -g^{p\bar{l}} \Gamma^k_{ip}, \ \bar{\partial}_j g^{k\bar{l}} = \overline{\partial_j g^{l\bar{k}}} = -g^{k\bar{q}} \overline{\Gamma^l_{jq}}$$
$$\bar{\partial}_j \partial_i g^{k\bar{l}} = -g^{p\bar{l}} \overline{\partial}_j \Gamma^k_{ip} + g^{p\bar{q}} \Gamma^k_{ip} \overline{\Gamma^l_{jq}}$$

and

$$\partial_{i}\mathfrak{g}_{k\bar{l}} = \nabla_{i}\mathfrak{g}_{k\bar{l}} + \Gamma^{m}_{ik}\mathfrak{g}_{m\bar{l}}, \quad \bar{\partial}_{j}\mathfrak{g}_{k\bar{l}} = \nabla_{\bar{j}}\mathfrak{g}_{k\bar{l}} + \overline{\Gamma^{m}_{jl}}\mathfrak{g}_{k\bar{m}},$$
$$\bar{\partial}_{j}\partial_{i}\mathfrak{g}_{k\bar{l}} = \nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{k\bar{l}} + \overline{\Gamma^{q}_{jl}}\nabla_{i}\mathfrak{g}_{k\bar{q}} + \Gamma^{m}_{ik}\nabla_{\bar{j}}\mathfrak{g}_{m\bar{l}} + \Gamma^{m}_{ik}\overline{\Gamma^{q}_{jl}}\mathfrak{g}_{m\bar{q}} + \bar{\partial}_{j}\Gamma^{m}_{ik}\mathfrak{g}_{m\bar{l}}.$$

Therefore, for any indices i, j, k, l, r, s,

$$(4.7) \qquad \begin{aligned} \partial_i (g^{k\bar{l}} \mathfrak{g}_{r\bar{s}}) &= g^{k\bar{l}} \partial_i \mathfrak{g}_{r\bar{s}} + \partial_i g^{k\bar{l}} \mathfrak{g}_{r\bar{s}} \\ &= g^{k\bar{l}} \nabla_i \mathfrak{g}_{r\bar{s}} + g^{k\bar{l}} \Gamma^m_{ir} \mathfrak{g}_{m\bar{s}} - g^{p\bar{l}} \Gamma^k_{ip} \mathfrak{g}_{r\bar{s}}, \\ \bar{\partial}_j (g^{k\bar{l}} \mathfrak{g}_{r\bar{s}}) &= g^{k\bar{l}} \bar{\partial}_j \mathfrak{g}_{r\bar{s}} + \bar{\partial}_j g^{k\bar{l}} \mathfrak{g}_{r\bar{s}} = g^{k\bar{l}} \nabla_{\bar{j}} \mathfrak{g}_{r\bar{s}} + g^{k\bar{l}} \overline{\Gamma^m_{js}} \mathfrak{g}_{r\bar{m}} - g^{k\bar{m}} \overline{\Gamma^l_{jm}} \mathfrak{g}_{r\bar{s}} \end{aligned}$$

and

$$(4.8) \qquad \begin{split} \bar{\partial}_{j}\partial_{i}(g^{k\bar{l}}\mathfrak{g}_{r\bar{s}}) &= g^{k\bar{l}}(\nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{r\bar{s}} + \overline{\Gamma_{js}^{q}}\nabla_{i}\mathfrak{g}_{r\bar{q}} + \Gamma_{ir}^{m}\nabla_{\bar{j}}\mathfrak{g}_{m\bar{s}} + \Gamma_{ir}^{m}\overline{\Gamma_{js}^{q}}\mathfrak{g}_{m\bar{q}} + \bar{\partial}_{j}\Gamma_{ir}^{m}\mathfrak{g}_{m\bar{s}}) \\ &- g^{k\bar{q}}\overline{\Gamma_{jq}^{l}}(\nabla_{i}\mathfrak{g}_{r\bar{s}} + \Gamma_{ir}^{m}\mathfrak{g}_{m\bar{s}}) - g^{p\bar{l}}\Gamma_{ip}^{k}(\nabla_{\bar{j}}\mathfrak{g}_{r\bar{s}} + \overline{\Gamma_{js}^{q}}\mathfrak{g}_{r\bar{q}}) \\ &- (g^{p\bar{l}}\bar{\partial}_{j}\Gamma_{ip}^{k} - g^{p\bar{q}}\Gamma_{ip}^{k}\overline{\Gamma_{jq}^{l}})\mathfrak{g}_{r\bar{s}}. \end{split}$$

Let r = k and sum over k,

(4.9)
$$\partial_i(g^{k\bar{l}}\mathfrak{g}_{k\bar{s}}) = g^{k\bar{l}}\nabla_i\mathfrak{g}_{k\bar{s}},$$

(4.10)
$$\bar{\partial}_j(g^{k\bar{l}}\mathfrak{g}_{k\bar{s}}) = g^{k\bar{l}}\nabla_{\bar{j}}\mathfrak{g}_{k\bar{s}} + g^{k\bar{l}}\overline{\Gamma_{js}^m}\mathfrak{g}_{k\bar{m}} - g^{k\bar{m}}\overline{\Gamma_{jm}^l}\mathfrak{g}_{k\bar{s}},$$

(4.11)
$$\bar{\partial}_{j}\partial_{i}(g^{k\bar{l}}\mathfrak{g}_{k\bar{s}}) = g^{k\bar{l}}\nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{k\bar{s}} + g^{k\bar{l}}\overline{\Gamma_{js}^{t}}\nabla_{i}\mathfrak{g}_{k\bar{t}} - g^{k\bar{t}}\overline{\Gamma_{jt}^{l}}\nabla_{i}\mathfrak{g}_{k\bar{s}}$$

which also follows from

$$\begin{split} \bar{\partial}_{j}\partial_{i}(g^{k\bar{l}}\mathfrak{g}_{k\bar{s}}) &= \bar{\partial}_{j}(g^{k\bar{l}}\nabla_{i}\mathfrak{g}_{k\bar{s}}) \\ &= \bar{\partial}_{j}g^{k\bar{l}}\nabla_{i}\mathfrak{g}_{k\bar{s}} + g^{k\bar{l}}\bar{\partial}_{j}\nabla_{i}\mathfrak{g}_{k\bar{s}} \\ &= g^{k\bar{l}}\nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{k\bar{s}} + g^{k\bar{l}}\overline{\Gamma_{js}^{t}}\nabla_{i}\mathfrak{g}_{k\bar{t}} - g^{k\bar{t}}\overline{\Gamma_{jt}^{l}}\nabla_{i}\mathfrak{g}_{k\bar{s}}. \end{split}$$

We shall also need

$$\begin{split} \partial_i (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{l}}) &= g_{1\bar{1}}^{-1} \partial_i \mathfrak{g}_{1\bar{l}} - g_{1\bar{1}}^{-2} \mathfrak{g}_{1\bar{l}} \partial_i g_{1\bar{1}} \\ &= g_{1\bar{1}}^{-1} \nabla_i \mathfrak{g}_{1\bar{l}} + g_{1\bar{1}}^{-2} \Gamma_{i1}^m (g_{1\bar{1}} \mathfrak{g}_{m\bar{l}} - g_{m\bar{1}} \mathfrak{g}_{1\bar{l}}), \\ \partial_{\bar{j}} (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{l}}) &= g_{1\bar{1}}^{-1} \partial_{\bar{j}} \mathfrak{g}_{1\bar{l}} - g_{1\bar{1}}^{-2} \mathfrak{g}_{1\bar{l}} \partial_{\bar{j}} g_{1\bar{1}} \\ &= g_{1\bar{1}}^{-1} (\nabla_{\bar{j}} \mathfrak{g}_{1\bar{l}} + \overline{\Gamma_{jl}^m} \mathfrak{g}_{1\bar{m}}) - g_{1\bar{1}}^{-2} \overline{\Gamma_{j1}^m} g_{1\bar{m}} \mathfrak{g}_{1\bar{l}} \end{split}$$

and

$$\begin{split} \bar{\partial}_{j}\partial_{i}(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}}) &= g_{1\bar{1}}^{-1}\nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{1\bar{1}} + g_{1\bar{1}}^{-2}(g_{1\bar{1}}\overline{\Gamma_{j1}^{m}}\nabla_{i}\mathfrak{g}_{1\bar{m}} - g_{1\bar{m}}\overline{\Gamma_{j1}^{m}}\nabla_{i}\mathfrak{g}_{1\bar{1}}) \\ &+ g_{1\bar{1}}^{-2}\Gamma_{i1}^{m}(g_{1\bar{1}}\nabla_{\bar{j}}\mathfrak{g}_{m\bar{1}} - g_{m\bar{1}}\nabla_{\bar{j}}\mathfrak{g}_{1\bar{1}}) \\ &+ \bar{\partial}_{j}(g_{1\bar{1}}^{-2}\Gamma_{i1}^{m})(g_{1\bar{1}}\mathfrak{g}_{m\bar{1}} - g_{m\bar{1}}\mathfrak{g}_{1\bar{1}}) + g_{1\bar{1}}^{-2}\Gamma_{i1}^{m}(\bar{\partial}_{j}g_{1\bar{1}}\mathfrak{g}_{m\bar{1}} - \bar{\partial}_{j}g_{m\bar{1}}\mathfrak{g}_{1\bar{1}}). \end{split}$$

At z_0 where $g_{i\bar{j}} = \delta_{ij}$, $T_{ij}^k = 2\Gamma_{ij}^k$, and $\mathfrak{g}_{i\bar{j}}$ is diagonal, we have $\partial (a^{k\bar{l}}\mathfrak{q}_{i\bar{j}}) = \nabla_i \mathfrak{q}_{l\bar{j}}$

$$\partial_i(g^{\kappa l}\mathfrak{g}_{k\bar{s}}) = \nabla_i\mathfrak{g}_{l\bar{s}},$$
$$\bar{\partial}_j(g^{k\bar{l}}\mathfrak{g}_{k\bar{s}}) = \nabla_{\bar{j}}\mathfrak{g}_{l\bar{s}} + \overline{\Gamma_{js}^l}(\mathfrak{g}_{l\bar{l}} - \mathfrak{g}_{s\bar{s}}),$$
$$\bar{\partial}_j\partial_i(g^{k\bar{l}}\mathfrak{g}_{k\bar{s}}) = \nabla_{\bar{j}}\nabla_i\mathfrak{g}_{l\bar{s}} + \overline{\Gamma_{js}^t}\nabla_i\mathfrak{g}_{l\bar{t}} - \overline{\Gamma_{jt}^l}\nabla_i\mathfrak{g}_{t\bar{s}},$$

$$\begin{aligned} \partial_i (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{l}}) &= \nabla_i \mathfrak{g}_{1\bar{l}} + (\Gamma_{i1}^l \mathfrak{g}_{l\bar{l}} - \Gamma_{i1}^1 \mathfrak{g}_{1\bar{l}}), \\ \partial_{\bar{j}} (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{l}}) &= \nabla_{\bar{j}} \mathfrak{g}_{1\bar{l}} + \overline{\Gamma_{j1}^1} \mathfrak{g}_{1\bar{1}} - \overline{\Gamma_{j1}^1} \mathfrak{g}_{1\bar{l}}. \end{aligned}$$

In particular,

$$\partial_i(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}}) = \nabla_i\mathfrak{g}_{1\bar{1}}, \ \bar{\partial}_j(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}}) = \nabla_j\mathfrak{g}_{1\bar{1}},$$

and

$$\begin{split} \bar{\partial}_{j}\partial_{i}(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}}) &= \nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{1\bar{1}} + (\overline{\Gamma_{j1}^{m}}\nabla_{i}\mathfrak{g}_{1\bar{m}} - \overline{\Gamma_{j1}^{1}}\nabla_{i}\mathfrak{g}_{1\bar{1}}) \\ &+ (\Gamma_{i1}^{m}\nabla_{\bar{j}}\mathfrak{g}_{m\bar{1}} - \Gamma_{i1}^{1}\nabla_{\bar{j}}\mathfrak{g}_{1\bar{1}}) + (\Gamma_{i1}^{1}\overline{\Gamma_{j1}^{1}} - \Gamma_{i1}^{m}\overline{\Gamma_{j1}^{m}})\mathfrak{g}_{1\bar{1}}. \end{split}$$

It follows that

(4.12)
$$\partial_i W = g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{l}} \partial_i (g^{k\bar{l}} \mathfrak{g}_{k\bar{1}}) + g^{k\bar{l}} \mathfrak{g}_{k\bar{1}} \partial_i (g_{1\bar{1}}^{-1} \mathfrak{g}_{1\bar{l}}) = 2\mathfrak{g}_{1\bar{1}} \nabla_i \mathfrak{g}_{1\bar{1}},$$

and

$$\begin{split} \bar{\partial}_{j}\partial_{i}W &= g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{l}}\bar{\partial}_{j}\partial_{i}(g^{k\bar{l}}\mathfrak{g}_{k\bar{1}}) + \bar{\partial}_{j}(g^{k\bar{l}}\mathfrak{g}_{k\bar{1}})\partial_{i}(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{l}}) \\ &\quad + \bar{\partial}_{j}(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{l}})\partial_{i}(g^{k\bar{l}}\mathfrak{g}_{k\bar{1}}) + g^{k\bar{l}}\mathfrak{g}_{k\bar{1}}\bar{\partial}_{j}\partial_{i}(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{l}}) \\ &\quad = \mathfrak{g}_{1\bar{1}}(\nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{1\bar{1}} + \overline{\Gamma_{j1}^{t}}\nabla_{i}\mathfrak{g}_{1\bar{t}} - \overline{\Gamma_{j1}^{t}}\nabla_{i}\mathfrak{g}_{t\bar{1}}) \\ &\quad + (\nabla_{\bar{j}}\mathfrak{g}_{l\bar{1}} + \overline{\Gamma_{j1}^{l}}\mathfrak{g}_{l\bar{l}} - \overline{\Gamma_{j1}^{l}}\mathfrak{g}_{1\bar{1}})(\nabla_{i}\mathfrak{g}_{1\bar{l}} + \Gamma_{i1}^{l}\mathfrak{g}_{l\bar{l}} - \Gamma_{i1}^{1}\mathfrak{g}_{1\bar{l}}) \\ &\quad + \nabla_{i}\mathfrak{g}_{l\bar{1}}(\nabla_{\bar{j}}\mathfrak{g}_{1\bar{l}} + \overline{\Gamma_{j1}^{l}}\mathfrak{g}_{1\bar{1}} - \overline{\Gamma_{j1}^{1}}\mathfrak{g}_{1\bar{l}}) \\ &\quad + \mathfrak{g}_{1\bar{1}}[\nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{1\bar{1}} + (\overline{\Gamma_{j1}^{m}}\nabla_{i}\mathfrak{g}_{1\bar{m}} - \overline{\Gamma_{j1}^{1}}\nabla_{i}\mathfrak{g}_{1\bar{1}}) \\ &\quad + (\Gamma_{i1}^{m}\nabla_{\bar{j}}\mathfrak{g}_{m\bar{1}} - \Gamma_{i1}^{1}\nabla_{\bar{j}}\mathfrak{g}_{1\bar{1}}) + (\Gamma_{i1}^{1}\overline{\Gamma_{j1}^{1}} - \Gamma_{i1}^{m}\overline{\Gamma_{j1}^{m}})\mathfrak{g}_{1\bar{1}}]. \end{split}$$

After some cancellations, this can be rewritten as

$$(4.13)$$

$$\bar{\partial}_{j}\partial_{i}W = 2\mathfrak{g}_{1\bar{1}}\nabla_{\bar{j}}\nabla_{i}\mathfrak{g}_{1\bar{1}} + 2\nabla_{i}\mathfrak{g}_{1\bar{1}}\nabla_{\bar{j}}\mathfrak{g}_{1\bar{1}} + \sum_{l>1}\nabla_{i}\mathfrak{g}_{l\bar{1}}\nabla_{\bar{j}}\mathfrak{g}_{1\bar{l}}$$

$$+ \sum_{l>1} (\nabla_{i}\mathfrak{g}_{1\bar{l}} + \Gamma_{i1}^{l}\mathfrak{g}_{l\bar{l}})(\nabla_{\bar{j}}\mathfrak{g}_{l\bar{1}} + \overline{\Gamma_{j1}^{l}}\mathfrak{g}_{l\bar{l}})$$

$$+ \mathfrak{g}_{1\bar{1}}\sum_{l>1} (\overline{\Gamma_{j1}^{l}}\nabla_{i}\mathfrak{g}_{1\bar{l}} + \Gamma_{i1}^{l}\nabla_{\bar{j}}\mathfrak{g}_{l\bar{1}})$$

$$- \mathfrak{g}_{1\bar{1}}\sum_{l>1} \Gamma_{i1}^{m}\overline{\Gamma_{j1}^{m}}(\mathfrak{g}_{1\bar{1}} + \mathfrak{g}_{l\bar{l}}).$$

Finally, we obtain

(4.14)
$$F^{i\bar{i}}\partial_i W \bar{\partial}_i W = 4\mathfrak{g}_{1\bar{1}}^2 F^{i\bar{i}} \nabla_i \mathfrak{g}_{1\bar{1}} \nabla_{\bar{i}} \mathfrak{g}_{1\bar{1}},$$

(4.15)
$$F^{i\bar{i}}\partial_i(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}})\bar{\partial}_i(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}}) = F^{i\bar{i}}\nabla_i\mathfrak{g}_{1\bar{1}}\nabla_{\bar{i}}\mathfrak{g}_{1\bar{1}}$$

and, by Cauchy-Schwarz inequality,

$$(4.16) F^{i\bar{i}}\bar{\partial}_{i}\partial_{i}W \geq 2\mathfrak{g}_{1\bar{1}}F^{i\bar{i}}\nabla_{\bar{i}}\mathfrak{g}_{1\bar{1}} + 2F^{i\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{1}}\nabla_{\bar{i}}\mathfrak{g}_{1\bar{1}} + \sum_{l>1}F^{i\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{l}}\nabla_{\bar{i}}\mathfrak{g}_{l\bar{1}} + \frac{1}{2}\sum_{l>1}F^{i\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{l}}\nabla_{\bar{i}}\mathfrak{g}_{l\bar{1}} - C\mathfrak{g}_{1\bar{1}}^{2}\sum F^{i\bar{i}},$$

$$(4.17) \qquad F^{i\bar{i}}\bar{\partial}_{i}\partial_{i}(g_{1\bar{1}}^{-1}\mathfrak{g}_{1\bar{1}}) \geq F^{i\bar{i}}\nabla_{\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{1}} - \frac{\gamma}{8\mathfrak{g}_{l\bar{1}}}\sum_{l>1}F^{i\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{l}}\nabla_{\bar{i}}\mathfrak{g}_{l\bar{1}} - C\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}}.$$

In summary, we can rewrite (4.3) as

(4.18)
$$\nabla_i \mathfrak{g}_{1\bar{1}} + \mathfrak{g}_{1\bar{1}} \partial_i \phi = 0, \ \nabla_{\bar{i}} \mathfrak{g}_{1\bar{1}} + \mathfrak{g}_{1\bar{1}} \bar{\partial}_i \phi = 0,$$

and, plugging (4.14)-(4.17) into (4.4), we derive

$$(4.19) \qquad \begin{array}{l} 0 \geq \frac{1}{\mathfrak{g}_{1\bar{1}}} F^{i\bar{i}} \nabla_{\bar{i}} \nabla_{i} \mathfrak{g}_{1\bar{1}} - \frac{1}{\mathfrak{g}_{1\bar{1}}^{2}} F^{i\bar{i}} \nabla_{i} \mathfrak{g}_{1\bar{1}} \nabla_{\bar{i}} \mathfrak{g}_{1\bar{1}} + F^{i\bar{i}} \overline{\partial}_{i} \partial_{i} \phi \\ + \frac{\gamma}{\mathfrak{g}_{1\bar{1}}^{2}} \sum_{l>1} F^{i\bar{i}} \nabla_{i} \mathfrak{g}_{1\bar{l}} \nabla_{\bar{i}} \mathfrak{g}_{l\bar{1}} + \frac{\gamma}{16\mathfrak{g}_{1\bar{1}}^{2}} \sum_{l>1} F^{i\bar{i}} \nabla_{i} \mathfrak{g}_{1\bar{l}} \nabla_{\bar{i}} \mathfrak{g}_{l\bar{1}} - C \sum F^{i\bar{i}} \end{array}$$

Next, differentiate equation (4.1) twice to obtain (at z_0),

(4.20)
$$F^{ii}\nabla_k \mathfrak{g}_{i\bar{i}} = \nabla_k \psi,$$

(4.21)
$$F^{i\bar{i}}\nabla_{\bar{1}}\nabla_{1}\mathfrak{g}_{i\bar{i}} + F^{i\bar{j},k\bar{l}}\nabla_{1}\mathfrak{g}_{i\bar{j}}\nabla_{\bar{1}}\mathfrak{g}_{k\bar{l}} = \nabla_{\bar{1}}\nabla_{1}\psi.$$

Recall the formulas for communication of covariant derivatives

$$(4.22) \qquad \begin{cases} u_{i\bar{j}k} - u_{k\bar{j}i} = T^{l}_{ik} u_{l\bar{j}}, \quad u_{i\bar{j}k} - u_{ik\bar{j}} = -g^{l\bar{m}} R_{k\bar{j}i\bar{m}} u_{l}, \\ u_{i\bar{j}k\bar{l}} - u_{i\bar{j}\bar{l}k} = g^{p\bar{q}} R_{k\bar{l}i\bar{q}} u_{p\bar{j}} - g^{p\bar{q}} R_{p\bar{l}k\bar{j}} u_{i\bar{q}}, \\ u_{i\bar{j}k\bar{l}} - u_{k\bar{l}i\bar{j}} = g^{p\bar{q}} (R_{k\bar{l}i\bar{q}} u_{p\bar{j}} - R_{i\bar{j}k\bar{q}} u_{p\bar{l}}) + T^{p}_{ik} u_{p\bar{j}\bar{l}} + \overline{T^{q}_{jl}} u_{i\bar{q}k} - T^{p}_{ik} \overline{T^{q}_{jl}} u_{p\bar{q}} \\ where for simplicity, a_{\bar{l}} = \nabla \nabla \alpha = \partial \partial \alpha. \end{cases}$$

where for simplicity, $u_{i\bar{j}} = \nabla_{\bar{j}} \nabla_i u = \partial_{\bar{j}} \partial_i u$,

$$u_{i\bar{j}k} = \nabla_k u_{i\bar{j}} = \partial_k u_{i\bar{j}} - \Gamma^l_{ki} u_{l\bar{j}},$$

and

$$u_{i\bar{j}k\bar{l}} = \nabla_{\bar{l}} u_{i\bar{j}k} = \partial_{\bar{l}} u_{i\bar{j}k} - \overline{\Gamma_{lj}^m} u_{i\bar{m}k}.$$

Therefore at z_0 , by (4.22),

(4.23)
$$\nabla_{\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{1}} - \nabla_{\bar{1}}\nabla_{1}\mathfrak{g}_{i\bar{i}} = R_{i\bar{i}1\bar{1}}\mathfrak{g}_{1\bar{1}} - R_{1\bar{1}i\bar{i}}\mathfrak{g}_{i\bar{i}} - T_{i1}^{l}\nabla_{\bar{i}}\mathfrak{g}_{l\bar{1}} - \overline{T_{i1}^{l}}\nabla_{i}\mathfrak{g}_{1\bar{l}} - T_{i1}^{l}\nabla_{i}\mathfrak{g}_{1\bar{l}} - T_{i1}^{l}\nabla_{i}\mathfrak{g}_{1\bar{l}}$$

where

$$H_{i\bar{i}} = \nabla_{\bar{i}} \nabla_{i} \chi_{1\bar{1}} - \nabla_{\bar{1}} \nabla_{1} \chi_{i\bar{i}} - 2 \Re \mathfrak{e} \{ T_{i1}^{l} \nabla_{\bar{i}} \chi_{l\bar{1}} \} + R_{i\bar{i}1\bar{l}} \chi_{l\bar{1}} - R_{1\bar{1}i\bar{l}} \chi_{l\bar{i}} - T_{i1}^{j} \overline{T_{i1}^{l}} \chi_{j\bar{l}}$$

It follows from Schwarz inequality that

(4.24)
$$F^{i\bar{i}}\nabla_{\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{1}} \geq F^{i\bar{i}}\nabla_{1}\mathfrak{g}_{1\bar{i}} - \frac{\gamma}{16\mathfrak{g}_{1\bar{1}}}F^{i\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{l}}\nabla_{\bar{i}}\mathfrak{g}_{l\bar{1}} - C\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}} + F^{i\bar{i}}H_{i\bar{i}}.$$

Combining (4.19), (4.21) and (4.24), by Schwarz inequality we derive

$$(4.25) \qquad \qquad \mathfrak{g}_{1\bar{1}}F^{i\bar{i}}\bar{\partial}_{i}\partial_{i}\phi \leq -\nabla_{\bar{1}}\nabla_{1}\psi - E - F^{i\bar{i}}H_{i\bar{i}} + C\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}} - \frac{\gamma}{32\mathfrak{g}_{1\bar{1}}}\sum_{l>1}F^{i\bar{i}}(\nabla_{i}\mathfrak{g}_{l\bar{1}}\nabla_{\bar{i}}\mathfrak{g}_{1\bar{l}} + \nabla_{i}\mathfrak{g}_{1\bar{l}}\nabla_{\bar{i}}\mathfrak{g}_{l\bar{1}})$$

where

$$E = -F^{i\bar{j},k\bar{l}}\nabla_{1}\mathfrak{g}_{i\bar{j}}\nabla_{\bar{1}}\mathfrak{g}_{k\bar{l}} - \frac{1+\gamma}{\mathfrak{g}_{1\bar{1}}}F^{i\bar{i}}\nabla_{i}\mathfrak{g}_{1\bar{1}}\nabla_{\bar{i}}\mathfrak{g}_{1\bar{1}}$$

In the rest of this section we shall not need the last nonpositive term on the right hand side of (4.25) and therefore drop it.

4.1. The term E. To estimate the term E, set

$$\begin{split} J &= \{i : |\mathfrak{g}_{i\overline{i}}| \geq \gamma \mathfrak{g}_{1\overline{1}}\}, \\ K &= \{i : |\mathfrak{g}_{i\overline{i}}| < \gamma \mathfrak{g}_{1\overline{1}}, \ \gamma F^{i\overline{i}} > F^{1\overline{1}}\}, \\ L &= \{i : |\mathfrak{g}_{i\overline{i}}| < \gamma \mathfrak{g}_{1\overline{1}}, \ \gamma F^{i\overline{i}} \leq F^{1\overline{1}}\} \end{split}$$

where $\gamma > 0$ is same as in (4.2). By an inequality due to Caffarelli-Nirenberg-Spruck, Andrews and Gerhardt (see e.g. [62]) we have

$$-F^{i\overline{j},k\overline{l}}\nabla_{1}\mathfrak{g}_{i\overline{j}}\nabla_{\overline{1}}\mathfrak{g}_{k\overline{l}} \geq \sum_{i\neq j}\frac{F^{i\overline{i}}-F^{j\overline{j}}}{\mathfrak{g}_{j\overline{j}}-\mathfrak{g}_{i\overline{i}}}|\nabla_{1}\mathfrak{g}_{i\overline{j}}|^{2} \geq \sum_{i\geq 2}\frac{F^{i\overline{i}}-F^{1\overline{1}}}{\mathfrak{g}_{1\overline{1}}-\mathfrak{g}_{i\overline{i}}}|\nabla_{1}\mathfrak{g}_{i\overline{1}}|^{2}.$$

By the first formula in (4.22),

$$\nabla_1 \mathfrak{g}_{i\bar{1}} = \nabla_i \mathfrak{g}_{1\bar{1}} + T_{i1}^1 \mathfrak{g}_{1\bar{1}} + \kappa_i$$

where

$$\begin{aligned} \kappa_i &= \nabla_1 \chi_{i\bar{1}} - \nabla_i \chi_{1\bar{1}} - T_{i1}^l \chi_{l\bar{1}} \\ &= \chi_{i\bar{1}1} + \chi_{i\bar{1},\zeta_\alpha} \partial_1 \partial_\alpha u - \chi_{1\bar{1}i} - \chi_{1\bar{1},\zeta_\alpha} \partial_i \partial_\alpha u - T_{i1}^l \chi_{l\bar{1}}. \end{aligned}$$

This yields

$$(4.26) \qquad -F^{i\bar{j},k\bar{l}}\nabla_{1}\mathfrak{g}_{i\bar{j}}\nabla_{\bar{1}}\mathfrak{g}_{k\bar{l}} \geq \frac{1-\gamma}{(1+\gamma)\mathfrak{g}_{1\bar{1}}}\sum_{i\in K}F^{i\bar{i}}|\nabla_{i}\mathfrak{g}_{1\bar{1}}+T^{1}_{i1}\mathfrak{g}_{1\bar{1}}+\kappa_{i}|^{2}$$
$$\geq \frac{1-2\gamma}{(1+\gamma)\mathfrak{g}_{1\bar{1}}}\sum_{i\in K}F^{i\bar{i}}|\nabla_{i}\mathfrak{g}_{1\bar{1}}|^{2}-C\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}}-\frac{C}{\mathfrak{g}_{1\bar{1}}}\sum F^{i\bar{i}}|\nabla_{i}\mathfrak{g}_{1\bar{1}}|^{2}+|\partial_{i}\partial_{\alpha}u|^{2}).$$

Therefore,

(4.27)
$$E \geq -4\gamma \mathfrak{g}_{1\bar{1}} \sum_{i \in K} F^{i\bar{i}} |\nabla_i \phi|^2 - (1+\gamma) \mathfrak{g}_{1\bar{1}} \sum_{J \cup L} F^{i\bar{i}} |\nabla_i \phi|^2 - \frac{C}{\mathfrak{g}_{1\bar{1}}} \sum_{i,\alpha} F^{i\bar{i}} (|\chi_{i\bar{1},\zeta_\alpha} \nabla_1 \nabla_\alpha u|^2 + |\nabla_i \nabla_\alpha u|^2) - C \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}}.$$

4.2. The term $H = F^{i\bar{i}}H_{i\bar{i}}$. We need to handle the first three terms in $H_{i\bar{i}}$ carefully. First, by (2.2),

(4.28)
$$F^{i\bar{i}}\mathfrak{Re}\{T^l_{i1}\nabla_{\bar{i}}\chi_{l\bar{1}}\} \le C\sum F^{i\bar{i}}|\nabla_i\nabla_\alpha u| + C\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}},$$

Next, a straightforward calculation using (2.2)-(2.4) shows

$$\begin{aligned}
\nabla_{\bar{l}}\nabla_{k}\chi_{i\bar{j}} &= \nabla_{\bar{l}}(\chi_{i\bar{j}k} + \chi_{i\bar{j},\zeta_{\alpha}}\partial_{k}\partial_{\alpha}u + \chi_{i\bar{j},\bar{\zeta}_{\alpha}}\partial_{k}\partial_{\bar{\alpha}}u) \\
&= \chi_{i\bar{j}k\bar{l}} + \chi_{i\bar{j}k,\zeta_{\alpha}}\partial_{\bar{l}}\partial_{\alpha}u + \chi_{i\bar{j}k,\bar{\zeta}_{\alpha}}\partial_{\bar{l}}\partial_{\bar{\alpha}}u \\
&+ (\chi_{i\bar{j}\bar{l},\zeta_{\alpha}} + \chi_{i\bar{j},\zeta_{\alpha}\zeta_{\beta}}\partial_{\bar{l}}\partial_{\beta}u + \chi_{i\bar{j},\zeta_{\alpha}\bar{\zeta}_{\beta}}\partial_{\bar{l}}\partial_{\bar{\beta}}u)\partial_{k}\partial_{\alpha}u \\
&+ (\chi_{i\bar{j}\bar{l},\bar{\zeta}_{\alpha}} + \chi_{i\bar{j},\bar{\zeta}_{\alpha}\zeta_{\beta}}\partial_{\bar{l}}\partial_{\beta}u + \chi_{i\bar{j},\bar{\zeta}_{\alpha}\bar{\zeta}_{\beta}}\partial_{\bar{l}}\partial_{\bar{\beta}}u)\partial_{k}\partial_{\bar{\alpha}}u \\
\end{aligned}$$

$$(4.29) \qquad \qquad + \chi_{i\bar{j},\zeta_{\alpha}}\nabla_{\bar{l}}(\nabla_{k}\nabla_{\alpha}u + \Gamma^{m}_{k\alpha}\nabla_{m}u) + \chi_{i\bar{j},\bar{\zeta}_{\alpha}}\nabla_{\bar{l}}\nabla_{k}\nabla_{\bar{\alpha}}u \\
&= \chi_{i\bar{j}k\bar{l}} + \chi_{i\bar{j}k,\zeta_{\alpha}}\partial_{\bar{l}}\partial_{\alpha}u + \chi_{i\bar{j}k,\bar{\zeta}_{\alpha}}\partial_{\bar{l}}\partial_{\bar{\alpha}}u + \chi_{i\bar{j}\bar{l},\bar{\zeta}_{\alpha}}\partial_{k}\partial_{\alpha}u + \chi_{i\bar{j}\bar{l},\bar{\zeta}_{\alpha}}\partial_{k}\partial_{\bar{\alpha}}u \\
&+ \chi_{i\bar{j},\bar{\zeta}_{\alpha}\zeta_{\beta}}\partial_{k}u_{\alpha}\partial_{\bar{l}}\partial_{\beta}u + \chi_{i\bar{j},\zeta_{\alpha}\bar{\zeta}_{\beta}}\partial_{k}\partial_{\alpha}u\partial_{\bar{l}}\partial_{\beta}u \\
&+ \chi_{i\bar{j},\bar{\zeta}_{\alpha}\bar{\zeta}_{\beta}}\partial_{k}\partial_{\bar{\alpha}}u\partial_{\bar{l}}\partial_{\bar{\beta}}u + \chi_{i\bar{j},\bar{\zeta}_{\alpha}}\nabla_{\bar{l}}\nabla_{k}\nabla_{\alpha}u \\
&+ \chi_{i\bar{j},\bar{\zeta}_{\alpha}}(\nabla_{\bar{l}}\Gamma^{m}_{k\alpha}\nabla_{m}u + \Gamma^{m}_{k\alpha}\nabla_{\bar{l}}\nabla_{m}u).
\end{aligned}$$

By (2.2),

$$\nabla_k \nabla_{\bar{i}} \nabla_i u = \nabla_k \mathfrak{g}_{i\bar{i}} - \nabla_k \chi_{i\bar{i}} = \nabla_k \mathfrak{g}_{i\bar{i}} - \chi_{i\bar{i}k} - \chi_{i\bar{i},\zeta_\alpha} \partial_k \partial_\alpha u - \chi_{i\bar{i},\bar{\zeta}_\alpha} \partial_k \partial_{\bar{\alpha}} u,$$

$$\nabla_{\bar{k}} \nabla_{\bar{i}} \nabla_i u = \nabla_{\bar{k}} \mathfrak{g}_{i\bar{i}} - \nabla_{\bar{k}} \chi_{i\bar{i}} = \nabla_{\bar{k}} \mathfrak{g}_{i\bar{i}} - \chi_{i\bar{i}\bar{k}} - \chi_{i\bar{i},\zeta_\alpha} \partial_{\bar{k}} \partial_\alpha u - \chi_{i\bar{i},\bar{\zeta}_\alpha} \partial_{\bar{k}} \partial_{\bar{\alpha}} u.$$

18

Therefore by (4.18), (4.20) and (4.22),

$$\begin{aligned} F^{i\bar{i}} \nabla_{\bar{i}} \nabla_{i} \chi_{1\bar{1}} &\geq 2F^{i\bar{i}} \Re \mathfrak{e} \{ \chi_{1\bar{1},\zeta_{\alpha}} \nabla_{\alpha} \nabla_{\bar{i}} \nabla_{i} u \} - C \sum_{i,\alpha} F^{i\bar{i}} |\partial_{i} \partial_{\alpha} u|^{2} - CF^{i\bar{i}} (\mathfrak{g}_{i\bar{i}}^{2} + 1) \\ &\geq 2 \Re \mathfrak{e} \{ \chi_{1\bar{1},\zeta_{\alpha}} \nabla_{\alpha} \psi \} - 2F^{i\bar{i}} \Re \mathfrak{e} \{ \chi_{1\bar{1},\zeta_{\alpha}} \chi_{i\bar{i},\zeta_{\beta}} \partial_{\alpha} \partial_{\beta} u \} \\ &- C \sum_{i,\alpha} F^{i\bar{i}} |\nabla_{i} \nabla_{\alpha} u|^{2} - CF^{i\bar{i}} (\mathfrak{g}_{i\bar{i}}^{2} + 1), \end{aligned}$$

$$(4.31) \qquad F^{i\bar{j}} \nabla_{\bar{1}} \nabla_{1} \chi_{i\bar{i}} \leq 2F^{i\bar{i}} \Re \mathfrak{e} \{ \chi_{i\bar{i},\zeta_{\alpha}} \nabla_{\alpha} \nabla_{\bar{1}} \nabla_{1} u + \chi_{i\bar{i}\bar{1},\zeta_{\alpha}} \partial_{1} \partial_{\alpha} u \} + C \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}} + Q$$
$$= -2\mathfrak{g}_{1\bar{1}} F^{i\bar{i}} \Re \mathfrak{e} \{ \chi_{i\bar{i},\zeta_{\alpha}} \nabla_{\alpha} \phi \} - 2F^{i\bar{i}} \Re \mathfrak{e} \{ \chi_{i\bar{i},\zeta_{\alpha}} \chi_{1\bar{1},\zeta_{\beta}} \partial_{\beta} \partial_{\alpha} u \}$$
$$+ 2F^{i\bar{i}} \Re \mathfrak{e} \{ \chi_{i\bar{i}\bar{1},\zeta_{\alpha}} \partial_{1} \partial_{\alpha} u \} + C \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}} + Q$$

where

(4.32)
$$Q = F^{ii}\chi_{i\bar{i},\zeta_{\alpha}\bar{\zeta}_{\beta}}(\partial_{1}\partial_{\alpha}u\partial_{\bar{1}}\partial_{\bar{\beta}}u + \partial_{\bar{1}}\partial_{\alpha}u\partial_{1}\partial_{\bar{\beta}}u) + 2F^{i\bar{i}}\mathfrak{Re}\{\chi_{i\bar{i},\zeta_{\alpha}\zeta_{\beta}}\partial_{1}\partial_{\alpha}u\partial_{\bar{1}}\partial_{\beta}u\}.$$

_

It follows that

$$(4.33) F^{i\overline{i}}H_{i\overline{i}} \ge 2\mathfrak{Re}\{\chi_{1\overline{1},\zeta_{\alpha}}\nabla_{\alpha}\psi\} + 2\mathfrak{g}_{1\overline{1}}F^{i\overline{i}}\mathfrak{Re}\{\chi_{i\overline{i},\zeta_{\alpha}}\nabla_{\alpha}\phi\} - 2F^{i\overline{i}}\mathfrak{Re}\{\chi_{i\overline{i}\overline{1},\zeta_{\alpha}}\partial_{1}\partial_{\alpha}u\} - CF^{i\overline{i}}\mathfrak{g}_{i\overline{i}}^{2} - Q - C\sum_{i,k}F^{i\overline{i}}|\nabla_{i}\nabla_{k}u|^{2} - C\mathfrak{g}_{1\overline{1}}\sum F^{i\overline{i}}.$$

4.3. The function ϕ . We choose $\phi = \eta - \log(1 - b|\nabla u|^2)$ where η is a function to be determined, and b is a positive constant satisfying $2b|\nabla u|^2 \leq 1$. Write

$$h = \log(1 - b|\nabla u|^2).$$

By straightforward calculations,

(4.34)
$$-b^{-1}e^{h}\partial_{i}h = \nabla_{k}u\nabla_{i}\nabla_{\bar{k}}u + \nabla_{\bar{k}}u\nabla_{i}\nabla_{k}u, -b^{-1}e^{h}\partial_{\bar{i}}h = \nabla_{k}u\nabla_{\bar{i}}\nabla_{\bar{k}}u + \nabla_{\bar{k}}u\nabla_{\bar{i}}\nabla_{k}u$$

and

$$(4.35) -b^{-1}e^{h}(\partial_{\bar{i}}h\partial_{i}h + \partial_{\bar{i}}\partial_{i}h) = \nabla_{\bar{k}}u\nabla_{\bar{i}}\nabla_{i}\nabla_{k}u + \nabla_{k}u\nabla_{\bar{i}}\nabla_{i}\nabla_{\bar{k}}u + \sum_{k}(\nabla_{i}\nabla_{\bar{k}}u\nabla_{k}\nabla_{\bar{i}}u + \nabla_{i}\nabla_{k}u\nabla_{\bar{i}}\nabla_{\bar{k}}u) = \nabla_{\bar{k}}u\nabla_{k}\nabla_{\bar{i}}\nabla_{i}u + \nabla_{k}u\nabla_{\bar{k}}\nabla_{\bar{i}}\nabla_{i}u + 2\mathfrak{Re}\{T^{l}_{ik}\nabla_{\bar{i}}\nabla_{l}u)\} + \sum_{k}(|\nabla_{i}\nabla_{\bar{k}}u|^{2} + |\nabla_{i}\nabla_{k}u|^{2}).$$

Therefore, by (2.2),

Let \mathcal{L} be the linear operator given by

$$\mathcal{L}v = F^{i\bar{i}}\partial_{\bar{i}}\partial_{i}v + 2F^{i\bar{i}}\mathfrak{Re}\{\chi_{i\bar{i},\zeta_{\alpha}}\partial_{\alpha}v\}.$$

By (4.36) we derive

(4.37)
$$\mathcal{L}\phi \ge \mathcal{L}\eta + \frac{b}{2}F^{i\bar{i}}\mathfrak{g}_{i\bar{i}}^2 + b\sum_{i,k}F^{i\bar{i}}|\nabla_i\nabla_k u|^2 - Cb\sum F^{i\bar{i}} - Cb.$$

Note that

(4.38)
$$|\nabla_i \phi|^2 \le 2|\nabla_i \eta|^2 + Cb^2 \mathfrak{g}_{i\bar{i}}^2 + Cb^2 \sum_k |\nabla_i \nabla_k|^2 + Cb^2$$

where C depends on $|\nabla u|_{C^0(\bar{M})}$.

Finally, combining (4.25), (4.27), (4.33), (4.37) and (4.38) we obtain

$$(4.39)\qquad\qquad\qquad \mathfrak{g}_{1\bar{1}}\mathcal{L}\eta \leq A_1 + A_2 + A_3 + 2bA_4$$

if b and $\mathfrak{g}_{1\bar{1}}$ are sufficiently small and large, respectively, where

$$(4.40) \qquad \begin{aligned} A_{1} &= -\nabla_{\bar{1}} \nabla_{1} \psi - 2 \Re \mathfrak{e} \{ \chi_{1\bar{1},\zeta_{\alpha}} \nabla_{\alpha} \psi \} + Cb \mathfrak{g}_{1\bar{1}} | \Re \mathfrak{e} \{ \nabla_{\bar{k}} u \nabla_{k} \psi \} | \\ A_{2} &= 8 \gamma \mathfrak{g}_{1\bar{1}} F^{i\bar{i}} | \nabla_{i} \eta |^{2} + 2 \mathfrak{g}_{1\bar{1}} \sum_{J \cup L} F^{i\bar{i}} | \nabla_{i} \eta |^{2} + C \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}} \\ A_{3} &= 2 F^{i\bar{i}} \Re \mathfrak{e} \{ \chi_{i\bar{i}\bar{1},\zeta_{\alpha}} \nabla_{1} \nabla_{\alpha} u \} + \frac{C}{\mathfrak{g}_{1\bar{1}}} \sum_{i,\alpha} F^{i\bar{i}} | \chi_{i\bar{1},\zeta_{\alpha}} \nabla_{1} \nabla_{\alpha} u |^{2} + Q \\ A_{4} &= - \frac{\mathfrak{g}_{1\bar{1}}}{8} F^{i\bar{i}} \mathfrak{g}_{i\bar{i}}^{2} - \frac{\mathfrak{g}_{1\bar{1}}}{4} \sum_{i,k} F^{i\bar{i}} | \nabla_{i} \nabla_{k} u |^{2}. \end{aligned}$$

4.4. Proof of Theorems 2.2. By Cauchy-Schwarz inequality we obtain

Lemma 4.1. Under assumption (2.5),

(4.41)
$$Q \leq -\frac{c_0}{8} \left(\mathfrak{g}_{1\bar{1}}^2 + \sum_k |\nabla_1 \nabla_k u|^2 \right) \sum F^{i\bar{i}} + C \sum F^{i\bar{i}} |\nabla_1 \nabla_i u|^2 + C \sum F^{i\bar{i}}.$$

Remark 4.2. If $\chi = \chi(z, u)$ and is independent of ∂u , $\overline{\partial} u$, then

$$F^{i\bar{i}}H_{i\bar{i}} \ge -C\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}}.$$

20

By Lemma 4.1 and (4.39) we derive under assumption (2.5),

(4.42)
$$\mathfrak{g}_{1\bar{1}}\mathcal{L}\eta \leq 4\mathfrak{g}_{1\bar{1}}F^{i\bar{i}}|\nabla_i\eta|^2 + A_1 + bA_4 - \frac{c_0}{32}\left(\mathfrak{g}_{1\bar{1}}^2 + \sum_k |\nabla_1\nabla_k u|^2\right)\sum F^{i\bar{i}}$$

if b and $\mathfrak{g}_{1\overline{1}}$ are sufficiently small and large, respectively.

Next, by the concavity of f and Schwarz inequality,

$$\begin{split} \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}} &= \sum F^{i\bar{i}}(\mathfrak{g}_{1\bar{1}} - \mathfrak{g}_{i\bar{i}}) + \sum F^{i\bar{i}}\mathfrak{g}_{i\bar{i}}\\ &\geq f(\mathfrak{g}_{1\bar{1}}\mathbf{1}) - \psi - \frac{1}{\mathfrak{g}_{1\bar{1}}} \sum F^{i\bar{i}}\mathfrak{g}_{i\bar{i}}^2 - \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}}. \end{split}$$

It follows that when $\mathfrak{g}_{1\overline{1}}$ is sufficiently large,

(4.43)
$$\mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}} \ge c_1 - \frac{1}{2\mathfrak{g}_{1\bar{1}}} \sum F^{i\bar{i}} \mathfrak{g}_{i\bar{i}}^2$$

where

$$c_1 = \frac{1}{2} \left(\sup_{\Gamma} f - \sup_{M} \psi \right) > 0.$$

Following Guan-Wang [35] we choose $\eta = \log \rho$ where ρ is a smooth function with compact support in $B_R \subset M$ satisfying

(4.44)
$$0 \le \rho \le 1, \ \rho|_{B_{\frac{3R}{4}}} \equiv 1, \ |\nabla\rho| \le C_R \sqrt{\rho}, \ |\nabla^2\rho| \le C_R.$$

Clearly,

(4.45)
$$\mathcal{L}\eta = \frac{1}{\rho}\mathcal{L}\rho - \frac{1}{\rho^2}F^{i\overline{i}}|\nabla_i\rho|^2 \ge -\frac{C}{\rho}\sum F^{i\overline{i}}.$$

By (4.42), (4.43) and (4.45),

(4.46)
$$(c_0 \rho \mathfrak{g}_{1\bar{1}} - C) \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}} + (b \mathfrak{g}_{1\bar{1}} - C) \rho F^{i\bar{i}} \mathfrak{g}_{i\bar{i}}^2 + (c_0 c_1 - Cb) \rho \mathfrak{g}_{1\bar{1}} \le C$$

We now fix b sufficiently small to derive a bound $\rho \mathfrak{g}_{1\bar{1}} \leq C$. This yields an interior estimates for $|\partial \bar{\partial} u|$, and a bound for the Hölder norm of $|\partial \bar{\partial} u|$ follows from Evans-Krylov Theorem. The proof of Theorem 2.2 is complete.

4.5. Proof of Theorems 2.9 and 2.14. As in [27] we take

$$\eta = A(\underline{u} - u + 1 + \sup(u - \underline{u}))$$

and denote $\underline{\mathfrak{g}}_{i\overline{j}} = \sqrt{-1}\partial_{\overline{j}}\partial_i\underline{u} + \chi_{i\overline{j}}[\underline{u}]$. Then

(4.47)
$$A_2 \le C(\gamma A^2 + 1)\mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}} + CA^2 \mathfrak{g}_{1\bar{1}} \sum_J F^{i\bar{i}} + CA^2 \mathfrak{g}_{1\bar{1}} F^{1\bar{1}}.$$

By the concavity of $\chi_{i\bar{i}}$ we have

$$\chi_{i\bar{i},\zeta_{\alpha}}(\underline{u}_{\alpha}-u_{\alpha})+\chi_{i\bar{i},\zeta_{\bar{a}}}(\underline{u}_{\bar{a}}-u_{\bar{a}})\geq\chi_{i\bar{i}}(z,\nabla\underline{u})-\chi_{i\bar{i}}(z,\nabla u).$$

Therefore,

(4.48)
$$\mathcal{L}(\underline{u}-u) \ge F^{i\overline{i}}(\underline{u}_{i\overline{i}}-u_{i\overline{i}}) + F^{i\overline{i}}(\chi_{i\overline{i}}[\underline{u}]) - \chi_{i\overline{i}}[u]) = F^{i\overline{i}}(\underline{\mathfrak{g}}_{i\overline{i}}-\mathfrak{g}_{i\overline{i}}).$$

Lemma 4.3. There exist uniform constants θ , N > 0 such that either

(4.49)
$$\mathcal{L}\eta \ge \theta \Big(\sum F^{i\bar{i}} + 1\Big)$$

or

(4.50)
$$F^{1\bar{1}} \ge \theta \sum F^{i\bar{i}}$$

provided that $\mathfrak{g}_{1\overline{1}} \geq N$.

Proof. Let $\mu = (\mu_1, \ldots, \mu_n)$ and $\lambda = (\lambda_1, \ldots, \lambda_n)$ be the eigenvalues of $\sqrt{-1}\partial \bar{\partial}\underline{u} + \chi[\underline{u}]$ and $\sqrt{-1}\partial \bar{\partial}u + \chi[u]$, respectively. Suppose (4.50) does not hold. By a lemma in [6] and Theorem 3.1 we have

(4.51)
$$F^{i\bar{i}}(\underline{\mathfrak{g}}_{i\bar{i}} - \mathfrak{g}_{i\bar{i}}) \ge \sum f_i(\mu_i - \lambda_i) \ge \theta \Big(\sum F^{i\bar{i}} + 1\Big).$$

This proves (4.49).

Suppose $\mathfrak{g}_{1\bar{1}} \geq N$ and (4.50) holds. Then

$$A_2 + A_3 \le C \left(A^2 \mathfrak{g}_{1\bar{1}} + \mathfrak{g}_{1\bar{1}}^2 + \sum |\nabla_1 \nabla_k u|^2 \right) \sum F^{i\bar{i}}$$

and

$$A_{4} \leq -\frac{\mathfrak{g}_{1\bar{1}}}{8}F^{1\bar{1}}\mathfrak{g}_{1\bar{1}}^{2} - \frac{\mathfrak{g}_{1\bar{1}}}{4}\sum F^{1\bar{1}}|\nabla_{1}\nabla_{k}u|^{2}$$
$$\leq -\frac{\theta\mathfrak{g}_{1\bar{1}}}{8}\left(\mathfrak{g}_{1\bar{1}}^{2} + \sum |\nabla_{1}\nabla_{k}u|^{2}\right)\sum F^{i\bar{i}}.$$

Moreover, by (4.48)

$$\mathcal{L}\eta \ge -C\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}}.$$

Combining these inequalities (4.43), from (4.39) we derive a bound for $\mathfrak{g}_{1\overline{1}}$.

Suppose now that (4.49) holds. To control the term A_3 in (4.39) we need condition (2.10) which is equivalent to $Q \leq 0$ so that

(4.52)
$$A_3 \le (C + \epsilon \mathfrak{g}_{1\bar{1}}) \sum F^{i\bar{i}} |\nabla_1 \nabla_i u|^2 + C \mathfrak{g}_{1\bar{1}} \sum F^{i\bar{i}}.$$

We make use of Lemma 4.3 to derive from (4.39)

(4.53)
$$0 \ge (b\mathfrak{g}_{1\bar{1}}^2 - CA^2)F^{1\bar{1}} + \sum_J F^{i\bar{i}}(b\mathfrak{g}_{i\bar{i}}^2 - CA^2) + \frac{b}{16}F^{1\bar{1}}\mathfrak{g}_{1\bar{1}}^2 + (b - \epsilon - C\mathfrak{g}_{1\bar{1}}^{-1})F^{i\bar{i}}|\nabla_1\nabla_i u|^2 + (\theta A - C\gamma A^2 - C)\sum F^{i\bar{i}}$$

22

Finally, fixing A sufficiently large and γ , ϵ sufficiently small, we obtain a bound $\mathfrak{g}_{1\bar{1}} \leq CA/\sqrt{b}$, completing the proof of Theorem 2.9. Clearly the proof also applies to Theorem 2.14 with slight modifications.

4.6. **Proof of Remark 2.3.** We now refine the above arguments to prove the claims in Remark 2.3 when $\psi = \psi(z, u, \nabla u)$. By straightforward calculations,

$$\mathfrak{Re}\{\nabla_{\bar{k}}u\nabla_{k}\psi\}=\mathfrak{Re}\{\psi_{\zeta_{\alpha}}\nabla_{\alpha}(|\nabla u|^{2}\})+O(1)$$

and therefore

$$2be^{-h}\mathfrak{Re}\{\nabla_{\bar{k}}u\nabla_{k}\psi\}=-2\mathfrak{Re}\{\psi_{\zeta_{\alpha}}\nabla_{\alpha}h\}+O(1).$$

Let $\widetilde{\mathcal{L}}$ be defined by

$$\widetilde{\mathcal{L}}v = \mathcal{L}v - 2\mathfrak{Re}\{\psi_{\zeta_{\alpha}}\nabla_{\alpha}v\} = F^{i\overline{i}}\partial_{\overline{i}}\partial_{\overline{i}}v + 2\mathfrak{Re}\{(F^{i\overline{i}}\chi_{i\overline{i},\zeta_{\alpha}} - \psi_{\zeta_{\alpha}})\nabla_{\alpha}v\}.$$

From (4.36) we see that

(4.54)
$$-\widetilde{\mathcal{L}}h \ge \frac{b}{2}F^{i\bar{i}}\mathfrak{g}_{i\bar{i}}^2 + b\sum_{i,k}F^{i\bar{i}}|\nabla_i\nabla_k u|^2 - Cb\sum F^{i\bar{i}} - Cb.$$

Next, we calculate

$$(4.55) \qquad \begin{aligned} -\nabla_{\bar{1}}\nabla_{1}\psi - 2\mathfrak{Re}\{\chi_{1\bar{1},\zeta_{\alpha}}\nabla_{\alpha}\psi\} \\ \leq -2\mathfrak{Re}\{\psi_{\zeta_{\alpha}}\nabla_{\alpha}\mathfrak{g}_{1\bar{1}}\} + C\sum |\nabla_{1}\nabla_{k}u| + C\mathfrak{g}_{1\bar{1}} + R\\ \leq 2\mathfrak{g}_{1\bar{1}}\mathfrak{Re}\{\psi_{\zeta_{\alpha}}\nabla_{\alpha}\phi\} + C\sum |\nabla_{1}\nabla_{k}u| + C\mathfrak{g}_{1\bar{1}} - R\end{aligned}$$

where

$$(4.56) R = \psi_{\zeta_{\alpha}\bar{\zeta}_{\beta}}(\partial_{1}\partial_{\alpha}u\partial_{\bar{1}}\partial_{\bar{\beta}}u + \partial_{\bar{1}}\partial_{\alpha}u\partial_{1}\partial_{\bar{\beta}}u) + 2\Re \mathfrak{e}\{\psi_{\zeta_{\alpha}\zeta_{\beta}}\partial_{1}\partial_{\alpha}u\partial_{\bar{1}}\partial_{\beta}u\}.$$

So in place of (4.37) we derive

$$(4.57) \qquad \qquad \mathfrak{g}_{1\bar{1}}\widetilde{\mathcal{L}}\phi \geq \mathfrak{g}_{1\bar{1}}\widetilde{\mathcal{L}}\eta + \frac{b\mathfrak{g}_{1\bar{1}}}{2}F^{i\bar{i}}\mathfrak{g}_{i\bar{i}}^2 + b\mathfrak{g}_{1\bar{1}}\sum_{i,k}F^{i\bar{i}}|\nabla_i\nabla_k u|^2 - Cb\mathfrak{g}_{1\bar{1}}\sum F^{i\bar{i}} - C\sum |\nabla_1\nabla_k u| - C\mathfrak{g}_{1\bar{1}} + R.$$

Consequently,

(4.58)
$$\mathfrak{g}_{1\bar{1}}\widetilde{\mathcal{L}}\eta \leq 4\mathfrak{g}_{1\bar{1}}F^{i\bar{i}}|\nabla_i\eta|^2 + A_3 + bA_4 + R + C\sum |\nabla_1\nabla_k u| + C\mathfrak{g}_{1\bar{1}}$$

provided that $\mathfrak{g}_{1\overline{1}}$ is sufficiently large.

We have $R \ge 0$ if ψ is convex in ∇u ; otherwise

$$R \ge -C\sum |\nabla_1 \nabla_k u|^2 - C\mathfrak{g}_{1\bar{1}}^2$$

and can therefore be controlled by

$$\left(\sum |\nabla_1 \nabla_k u|^2 + \mathfrak{g}_{1\bar{1}}^2\right) \sum F^{i\bar{i}}$$

under assumption (2.7). In either case, the rest of proof is same as in Subsection 4.4.

5. The equation from Gauduchon conjecture

Let Ω be a closed real (1,1) form on a compact Hermitian manifold (M^n, ω) with $[\Omega] = c_1^{BC}(M)$ in the Bott-Chern cohomology group $H^{1,1}_{BC}(M, \mathbb{R})$. Gauduchon [25] conjectured that there exists a Gauduchon metric $\hat{\omega}$ on M with Chern-Ricc curvature $\operatorname{Ric}_{\hat{\omega}} = \Omega$.

This is a natural extension of the Calabi conjecture for Kähler manifolds solved by Yau [74]. It was discovered by Popovici [57] and Tosatti-Weinkove [71] independently that Gauduchon conjecture reduces to solving the form-type Monge-Ampère equation (1.8)-(1.9) with c = 1 and $\sup_M u = 0$, where ω_0 is a Gauduchon metric; see [71] for details.

When n = 2, Equation (1.8) is the standard complex Monge-Ampère equation so Gauduchon conjecture follows affirmatively from results of Cherrier [8].

Using the Hodge star operator *, Tosatti-Weinkove [71] converted equation (1.8) into a Monge-Ampère type equation for a (1, 1)-form. Recall that

$$\Phi_u = \omega_0^{n-1} + \sqrt{-1}\partial\bar{\partial}u \wedge \omega^{n-2} + c \Re\left(\sqrt{-1}\partial u \wedge \bar{\partial}\omega^{n-2}\right) > 0.$$

Define

$$\tilde{\omega} = \frac{1}{(n-1)!} * \Phi_u.$$

As in [71],

$$\tilde{\omega} = \frac{1}{(n-1)!} * \Phi_u = \frac{\tilde{\chi} + (\Delta u)\omega - \sqrt{-1}\partial\bar{\partial}u}{n-1} > 0$$

where

$$\tilde{\chi} = \frac{1}{(n-2)!} * (\omega_0^{n-1} + c \operatorname{\mathfrak{Re}}\{\sqrt{-1}\partial u \wedge \bar{\partial}\omega^{n-2}\}),$$

and equation (1.8) becomes

(5.1)
$$\tilde{\omega}^n = e^{h+b}\omega^n.$$

It was shown by Tosatti-Weinkove [71] that the classical solvability of equation (5.1) reduces to the second order estimate

(5.2)
$$\Delta u \le C \Big(1 + \sup_{M} |\nabla u|^2 \Big);$$

see Conjecture 1.5 in [71]. Later on, Szekelyhidi-Tosatti-Weinkove [68] derived such an estimate and consequently proved Gauduchon conjecture.

Let

$$\chi = \frac{\mathrm{tr}_{\omega}\tilde{\chi}}{n-1}\omega - \tilde{\chi}$$

so $\tilde{\chi} = (tr_{\omega}\chi)\omega - \chi$. Equation (5.1) can therefore be rewritten in the form

(5.3)
$$\log \rho_{n-1}(\lambda(\sqrt{-1}\partial\bar{\partial}u + \chi)) = \psi(z).$$

In the rest of this section we verify that Theorem 2.9 applies to equation (5.3).

First of all, note that $\chi[u]$ is linear in ∇u so (2.9) is satisfied. We may take $\underline{u} = 0$ as $\chi_0 > 0$ and $\widetilde{\mathcal{C}}_{\sigma}^+ = \mathcal{P}_{n-1}$ for $f = \log \rho_{n-1}$. It remains to verify (2.10) for $\alpha = 1$ since $r_0 = n - 1$ by Lemma 2.8.

In local coordinates computing at a point where $g_{k\bar{l}} = \delta_{kl}$ and $T^k_{ij} = 2\Gamma^k_{ij}$, we have

(5.4)
$$\sqrt{-1}\partial u \wedge \bar{\partial}\omega^{n-2} = -(n-2)\partial_p u \partial_{\bar{q}} g_{k\bar{l}} dz_p \wedge d\bar{z}_q \wedge dz_k \wedge d\bar{z}_l \wedge \omega^{n-3}$$
$$= \frac{n-2}{2} \Big(\sum_i \sum_{p,l \neq i} \partial_p u \overline{T_{pl}^l} \mu_{i\bar{i}} - \sum_{i \neq j} \sum_{l \neq j} (\partial_j u \overline{T_{il}^l} + \partial_l u \overline{T_{li}^j}) \mu_{i\bar{j}} \Big)$$

where $\mu_{i\bar{j}} = s_{ij}dz_1 \wedge \cdots \wedge \widehat{dz_i} \wedge d\bar{z_i} \wedge \cdots \wedge dz_j \wedge \widehat{dz_j} \wedge \cdots \wedge dz_n \wedge d\bar{z_n}$ and

$$s_{ij} = \begin{cases} (\sqrt{-1})^{n-1}, & i \le j, \\ -(\sqrt{-1})^{n-1}, & i > j. \end{cases}$$

Similarly,

(5.5)
$$-\sqrt{-1}\bar{\partial}u \wedge \partial\omega^{n-2} = -(n-2)\partial_{\bar{q}}u\partial_{p}g_{k\bar{l}}dz_{p} \wedge d\bar{z}_{q} \wedge dz_{k} \wedge d\bar{z}_{l} \wedge \omega^{n-3}$$
$$= \frac{n-2}{2} \Big(\sum_{i}\sum_{p,l\neq i}\partial_{\bar{p}}uT^{l}_{pl}\mu_{i\bar{i}} - \sum_{i\neq j}\sum_{l\neq i}(\partial_{\bar{i}}uT^{l}_{jl} + \partial_{\bar{l}}uT^{i}_{lj})\mu_{i\bar{j}}\Big).$$

It follows that (see [71])

$$\frac{1}{(n-2)!} * \mathfrak{Re}\{\sqrt{-1}\partial u \wedge \bar{\partial}\omega^{n-2}\} = \sqrt{-1}\widetilde{E}_{i\bar{j}}dz_i \wedge d\bar{z}_j$$

where

(5.6)
$$\widetilde{E}_{i\bar{i}} = \frac{1}{2} \sum_{p,l \neq i} (\partial_p u \overline{T_{pl}^l} + \partial_{\bar{p}} u T_{pl}^l),$$

and

$$\widetilde{E}_{i\overline{j}} = -\frac{1}{2} \Big(\sum_{l \neq i} (\partial_i u \overline{T_{jl}^l} + \partial_l u \overline{T_{lj}^i}) + \sum_{l \neq j} (\partial_{\overline{j}} u T_{il}^l + \partial_{\overline{l}} u T_{li}^j) \Big), \quad i \neq j.$$

Clearly $\widetilde{E}_{i\bar{j}}$ does not contain $\partial_j u$. Therefore, as $\tilde{\chi}_{i\bar{j}\bar{k}} = \tilde{\chi}_{i\bar{j},\bar{k}} - \overline{\Gamma_{kj}^l} \tilde{\chi}_{i\bar{l}}$ and $\Gamma_{kk}^l = 0$,

(5.7)
$$\tilde{\chi}_{i\bar{j},\zeta_j} = c \,\widetilde{E}_{i\bar{j},\zeta_j} = 0, \quad \tilde{\chi}_{i\bar{i}\bar{i},\zeta_i} = \tilde{\chi}_{i\bar{i},\zeta_i\bar{i}} = c \,\widetilde{E}_{i\bar{i},\zeta_i\bar{i}} = 0, \quad \forall \, i,j.$$

Consider

$$F(\sqrt{-1}\partial\bar{\partial}u + \chi) = \log \rho_{n-1}(\lambda(\sqrt{-1}\partial\bar{\partial}u + \chi)).$$

Denote $\lambda_i = \mathfrak{g}_{i\overline{i}}$ and $\eta_j = \lambda_1 + \cdots + \widehat{\lambda}_j + \cdots + \lambda_n$ at p. Then

(5.8)
$$F^{i\overline{i}} = \sum_{j \neq i} \frac{1}{\eta_j}.$$

As $\lambda_1 \geq \cdots \geq \lambda_n$, we have $F^{1\bar{1}} \leq \cdots \leq F^{n\bar{n}}$ and $F^{h\bar{h}} \geq \frac{1}{2} \geq \frac{1}{2} = \frac{1}{2} F^{h\bar{h}}$

$$F^{k\bar{k}} \ge \frac{1}{\eta_1} \ge \frac{1}{n-1} F^{k\bar{k}}, \quad \forall k \ge 2.$$

Next,

(5.9)
$$\sum F^{i\overline{i}}\chi_{i\overline{i}\overline{1},\zeta_{1}} = \sum_{i}\sum_{j\neq i}\frac{\chi_{i\overline{i}\overline{1},\zeta_{1}}}{\eta_{j}} = \sum_{j}\sum_{i\neq j}\frac{\chi_{i\overline{i}\overline{1},\zeta_{1}}}{\eta_{j}}$$
$$= \sum_{j}\frac{\tilde{\chi}_{j\overline{j}\overline{1},\zeta_{1}}}{\eta_{j}} = \sum_{j\neq 1}\frac{\tilde{\chi}_{j\overline{j}\overline{1},\zeta_{1}}}{\eta_{j}}$$

since $\tilde{\chi}_{1\bar{1}\bar{1},\zeta_1} = c \widetilde{E}_{1\bar{1}\bar{1},\zeta_1} = 0$ by (5.7). Finally, from (5.8) and (5.9) we derive

(5.10)
$$|\sum F^{i\bar{i}}\chi_{i\bar{i}\bar{1},\zeta_1}| \le C\sum_{j\neq 1}\frac{1}{\eta_j} = CF^{1\bar{1}}.$$

Thus (2.10) is verified.

References

- T. Aubin, Équations du type Monge-Ampère sur les variétés kählériennes compactes, (French) Bull. Sci. Math. (2) 102 (1978), 63–95.
- [2] E. Bedford and B. A. Taylor, The Dirichlet problem for a complex Monge-Ampère equation, Invent. Math. 37 (1976), 1–44.
- [3] E. Bedford and B. A. Taylor, A new capacity for plurisubharmonic functions, Acta Math. 149 (1982) 1–40.
- [4] S. Brendle and G. Huisken, A fully nonlinear flow for two-convex hypersurfaces in Riemannian manifolds, Invent. Math. 210 (2017), 559–613.
- [5] L. A. Caffarelli, J. J. Kohn, L. Nirenberg and J. Spruck, The Dirichlet problem for nonlinear second-order elliptic equations II. Complex Monge-Ampère and uniformly elliptic equations, Comm. Pure Applied Math. 38 (1985), 209–252.
- [6] L. A. Caffarelli, L. Nirenberg and J. Spruck, The Dirichlet problem for nonlinear second-order elliptic equations III: Functions of eigenvalues of the Hessians, Acta Math. 155 (1985), 261– 301.
- [7] G. Chen, The J-equation and the supercritical deformed Hermitian-Yang-Mills equation, Invent. Math. 225 (2021), 529?602.

- [8] P. Cherrier, Equations de Monge-Ampère sur les variétés hermitiennes compactes, Bull. Sci. Math. 111 (1987), 343–385.
- [9] J.-C. Chu, L.-D. Huang and X.-H. Zhu, The Fu-Yau equation in higher dimensions, Peking Math. J. 2 (2019), 71–97.
- [10] T. Collins, A. Jacob and S.-T. Yau, (1,1) forms with specified Lagrangian phase: a priori estimates and algebraic obstructions, Cambridge J. Math. 8 (2020), 407–452.
- [11] T. Collins and S. Picard, The Dirichlet problem for k-Hessian equation on a complex manifold, arXiv:1909.00447.
- [12] T. Collins and G. Székelyhidi, Convergence of the J-flow on toric manifolds, J. Differential Geom. 107 (2017), 47–81.
- [13] T. Collins, D. Xie and S.-T. Yau, The deformed Hermitian-Yang-Mills equation in geometry and physics, Geometry and Physics. Vol. I, 69-90, Oxford Univ. Press, Oxford, 2018.
- [14] T. Collins and S.-T. Yau, Moment maps, nonlinear PDE and stability in mirror symmetry, I: geodesics, Ann. PDE 7 (2021), no. 1, Paper No. 11, 73 pp.
- [15] V. V. Datar and V. P. Pingali, A numerical criterion for generalized Monge-Ampère equations on projective manifolds, arXiv:2006.01530.
- [16] S. Dinew and S. Kolodziej, Liouville and Calabi-Yau type theorems for complex Hessian equations, Amer. J. Math. 139 (2017), 403–415.
- [17] S. K. Donaldson, Moment maps and diffeomorphisms, Asian J. Math. 3 (1999), 1–16.
- [18] S. K. Donaldson, Scalar curvature and stability of toric varieties, J. Differential Geom. 62 (2002), 289–349.
- [19] H. Fang, M.-J. Lai and X.-N. Ma, On a class of fully nonlinear flows in Kähler geometry, J. Reine Angew. Math. 653 (2011), 189–220.
- [20] K. Feng, H.-B. Ge and T. Zheng, The Dirichlet problem of fully nonlinear elliptic equations on Hermitian manifolds, arXiv:1905.02412.
- [21] J.-X. Fu, Z.-Z. Wang and D.-M. Wu, Form-type Calabi-Yau equations, Math. Res. Lett. 17 (2010), 887–903.
- [22] J.-X. Fu, Z.-Z. Wang and D.-M. Wu, Form-type equations on Kähler manifolds of nonnegative orthogonal bisectional curvature, Calc. Var. PDE 52 (2015), 327–344.
- [23] J.-X. Fu and S.-T. Yau, A Monge-Ampère-type equation motivated by string theory, Comm. Anal. Geom. 15 (2007), 29–75.
- [24] J.-X. Fu and S.-T. Yau, The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampère equation, J. Differential Geom. 78 (2008), 369–428.
- [25] P. Gauduchon, La 1-forme de torsion dúne variété hermitienne compacte Math. Ann. 267 (1984), 495–518.
- [26] M. George, B. Guan and C.-H. Qiu, On a class of fully nonlinear elliptic equations on Hermitian manifolds, preprint.
- [27] B. Guan, Second order estimates and regularity for fully nonlinear elliptic equations on Riemannian manifolds, Duke Math. J. 163 (2014) 1491–1524.
- [28] B. Guan, The Dirichlet problem for fully nonlinear elliptic equations on Riemannian manifolds, arXiv:1403.2133.
- [29] B. Guan and H.-M. Jiao, Second order estimates for Hessian type fully nonlinear elliptic equations on Riemannian manifolds, Calc. Var. PDE 54 (2015), 2693–2712.
- [30] B. Guan and H.-M. Jiao, The Dirichlet problem for Hessian type elliptic equations on Riemannian manifolds, Discrete Contin. Dyn. Syst. 36 (2016), 701–714.
- [31] B. Guan and Q. Li, The Dirichlet problem for a Monge-Ampère type equation on Hermitian manifolds, Adv. Math. 246 (2013), 351–367.

BO GUAN AND XIAOLAN NIE

- [32] B. Guan, C.-H. Qiu and R.-R. Yuan, Fully nonlinear elliptic equations for conformal deformations of Chern-Ricci forms, Adv. Math. 343 (2019), 538–566.
- [33] B. Guan and W. Sun, On a class of fully nonlinear elliptic equations on Hermitian manifolds, Calc. Var. PDE 54 (2015), 901–916.
- [34] P.-F. Guan, C.-Y. Ren and Z.-Z. Wang, Global C²-estimates for convex solutions of curvature equations, Comm. Pure Appl. Math. 68 (2015), 1287–1325.
- [35] P.-F. Guan and G.-F. Wang, Local estimates for a class of fully nonlinear equations arising from conformal geometry, Int. Math. Res. Not. 2003, no.26, 1413–1432.
- [36] X.-L. Han and X.-S. Jin, A rigidity theorem for the deformed Hermitian-Yang-Mills equation, Calc. Var. PDE 60 (2021), no. 1, Paper No. 13, 16 pp.
- [37] R. Harvey and H. B. Lawson Jr., *Calibrated geometries*, Acta Math. 148 (1982), 47–157.
- [38] R. Harvey and H. B. Lawson Jr., *Plurisubharmonicity in a general geometric context*, Geometry and Analysis. No. 1, 363–402, Adv. Lect. Math. (ALM), 17, Int. Press, Somerville, MA, 2011.
- [39] R. Harvey and H. B. Lawson Jr., Geometric plurisubharmonicity and convexity: an introduction, Adv. Math. 230 (2012), 2428–2456.
- [40] Z.-L. Hou, X.-N. Ma and D.-M. Wu, A second order estimate for complex Hessian equations on a compact Kähler manifold, Math. Res. Lett. 17 (2010), 547–561.
- [41] L. Huang, J. Zhang and X. Zhang, The deformed Hermitian-Yang-Mills equation on almost Hermitian manifolds, arXiv:2011.14091.
- [42] G. Huisken and C. Sinestrari, Mean curvature flow with surgeries of two-convex hypersurfaces Invent. Math. 175 (2009), 137–221.
- [43] A. Jacob and S.-T. Yau, A special Lagrangian type equation for holomorphic line bundles, Math. Ann. 369 (2017), 869–898.
- [44] M. Lejmi and G. Székelyhidi, The J-flow and stability, Adv. Math. 274 (2015), 404–431.
- [45] N.-C. Leung, S.-T. Yau and E. Zaslow, From special Lagrangian to Hermitian-Yang-Mills via Fourier-Mukai transform, Adv. Theor. Math. Phys. 4 (2000), 1319–1341.
- [46] S.-Y. Li, On the Dirichlet problems for symmetric function equations of the eigenvalues of the complex Hessian, Asian J. Math. 8 (2004), 87–106.
- [47] C.-M. Lin, Deformed Hermitian-Yang-Mills equation on compact Hermitian manifolds, arXiv 2012.00487.
- [48] M. Lin. and N. Trudinger, On some inequalities for elementary symmetric functions, Bull. Austral. Math. Soc. 50 (1994), 317–326.
- [49] X.-N. Ma, N. Trudinger and X.-J. Wang, Regularity of potential functions of the optimal transportation problem, Arch. Ration. Mech. Anal. 177 (2005), 151–183.
- [50] D. H. Phong, Geometric partial differential equations from unified string theories, arXiv: 1906.03693.
- [51] D. H. Phong, S. Picard and X.-W. Zhang, On estimates for the Fu-Yau generalization of a Strominger system, J. Reine Angew. Math. (2016), 1–32.
- [52] D. H. Phong, S. Picard and X.-W. Zhang, A second order estimate for general complex Hessian equations, Anal. PDE 9 (2016), 1693–1709.
- [53] D. H. Phong, S. Picard and X.-W. Zhang, The Fu-Yau equation with negative slope parameter, Invent. Math. 209 (2017), 541–576.
- [54] D. H. Phong, S. Picard, X.-W. Zhang, Geometric flows and Strominger systems, Math. Z. 288 (2018), 101–113.
- [55] D. H. Phong, S. Picard and X.-W. Zhang, Fu-Yau Hessian equations, J. Differential Geom. 118 (2021), 147–187.
- [56] D. H. Phong, J. Song and J. Sturm, Complex Monge Ampère Equations, Surveys in Differential Geometry, 17 (2012), 327–411.

- [57] D. Popovici, Aeppli cohomology classes associated with Gauduchon metrics on compact complex manifolds, Bull. Soc. Math. France 143 (2015), 763–800.
- [58] C.-H. Qiu and R.-R. Yuan, On the Dirichlet problem for fully nonlinear elliptic equations on annuli of metric cones, Discrete Contin. Dyn. Syst. 37 (2017), 5707–5730.
- [59] J.-P. Sha, p-convex Riemannian manifolds, Invent. Math. 83 (1986), 437–447.
- [60] J. Song, Nakai-Moishezon criterions for complex Hessian equations, arXiv:2012.07956.
- [61] J. Song and B. Weinkove, On the convergence and singularities of the J-flow with applications to the Mabuchi energy, Comm. Pure Appl. Math. 61 (2008), 210–229.
- [62] J. Spruck, Geometric aspects of the theory of fully nonlinear elliptic equations, Global theory of minimal surfaces, 283–309, Clay Math. Proc., 2, Amer. Math. Soc., Providence, RI, 2005.
- [63] J. Streets and G. Tian, Hermitian curvature flow, J. Eur. Math. Soc. 13 (2011), 601–634.
- [64] W. Sun, On a class of fully nonlinear elliptic equations on closed Hermitian manifolds, J. Geom. Anal. 26 (2016), 2459–2473.
- [65] W. Sun, On a class of fully nonlinear elliptic equations on closed Hermitian manifolds II: L[∞] estimate, Comm. Pure Appl. Math. 70 (2017), 172–199.
- [66] W. Sun, On uniform estimate of complex elliptic equations on closed Hermitian manifolds, Commun. Pure Appl. Anal. 16 (2017), 1553–1570.
- [67] G. Székelyhidi, Fully non-linear elliptic equations on compact Hermitian manifolds, J. Differential Geom. 109 (2018), 337–378.
- [68] G. Székelyhidi, V. Tosatti and B. Weinkove, Gauduchon metrics with prescribed volume form, Acta Math. 219 (2017), 181–211.
- [69] G. Tian, Kähler-Einstein metrics with positive scalar curvature, Invent. Math. 130 (1997), 1–37.
- [70] V. Tosatti, B. Weinkove, The Monge-Ampère equation for (n-1)-plurisubharmonic functions on a compact Kähler manifold, J. Amer. Math. Soc. 30 (2017), 311–346.
- [71] V. Tosatti, B. Weinkove, Hermitian metrics, (n-1, n-1) forms and Monge-Ampère equations, J. Reine Angew. Math. 755 (2019), 67–101.
- [72] B. Weinkove, Convergence of the J-flow on Kähler surfaces, Comm. Anal. Geom. 12 (2004), 949–965.
- [73] B. Weinkove, On the J-flow in higher dimensions and the lower boundedness of the Mabuchi energy, J. Differential Geom. 73 (2006), 351–358.
- [74] S.-T. Yau, On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I. Comm. Pure Appl. Math. 31 (1978), 339–411.
- [75] S.-T. Yau, Open problems in geometry, Proc. Symposia Pure Math. 54 (1993), 1–28.
- [76] R.-R. Yuan, On a class of fully nonlinear elliptic equations containing gradient terms on compact Hermitian manifolds, Canad. J. Math. 70 (2018), 943–960.
- [77] D.-K. Zhang, Hessian equations on closed Hermitian manifolds, Pacific J. Math. 291 (2017), 485–510.
- [78] K. Zheng, *I-properness of Mabuchi's K-energy* Calc. Var. PDE **54** (2015), 2807–2830.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, OH 43210, USA *Email address*: guan@math.ohio-state.edu

College of of Mathematics and Computer Science, Zhejiang Normal University, Jinhua, Zhejiang Province, 321004 China

Email address: nie@zjnu.edu.cn