

# A NOTE ON THE WEAK CONVERGENCE OF CONTINUOUSLY INTEGRABLE SEQUENCES

GANE SAMB LO AND ALADJI BABACAR NIANG

**ABSTRACT.** A uniformly continuously integrable sequence of real-valued measurable functions, defined on some probability space, is relatively compact in the  $\sigma(L^1, L^\infty)$  topology. In this paper, we link such a result to weak convergence theory of bounded measures as exposed in Billingsley (1968) and in Lo(2021) to offer a detailed and new proof using the ideas beneath the proof of prohorov's theorem where the continuous integrability replaces the uniform or asymptotic tightness.

† Gane Samb Lo.

LERSTAD, Gaston Berger University, Saint-Louis, Sénégal (main affiliation).  
 LSTA, Pierre and Marie Curie University, Paris VI, France.  
 AUST - African University of Sciences and Technology, Abuja, Nigeria  
 gane-samb.lo@edu.ugb.sn, gslo@aust.edu.ng, ganesamblo@ganesamblo.net  
 Permanent address : 1178 Evanston Dr NW T3P 0J9, Calgary, Alberta, Canada.

†† Aladji Babacar Niang

LERSTAD, Gaston Berger University, Saint-Louis, Sénégal.  
 Email: niang.aladji-babacar@ugb.edu.sn, aladjibacar93@gmail.com

**Keywords.** weak convergence in the  $\sigma(L^1, L^\infty)$  topology; weak convergence of bounded measure from a probabilistic point of view; relative compactness; tightness; continuous integrability.

**AMS 2010 Mathematics Subject Classification:** 60F05; 54C08

## 1. INTRODUCTION

The following result is proved, for example in Meyer (1966).

**Proposition 1.** *Let  $(f_n)_{n \geq 1}$  be a sequence of measurable mappings:*

$$f_n : \Omega \longrightarrow \mathbb{R},$$

where  $(\Omega, \mathcal{A}, \mathbb{P})$  is a probability space. Suppose that  $(f_n)_{n \geq 0}$  is uniformly continuously integrable (u.c.i), that is

$$\lim_{c \rightarrow +\infty} \bar{\phi}(c) := \lim_{c \rightarrow +\infty} \sup_{n \geq 1} \int_{(|f_n| > c)} |f_n| d\mathbb{P} = 0.$$

Then  $(f_n)_{n \geq 1}$  has a sub-sequence  $(f_{n_k})_{k \geq 1}$  converging to some  $f \in L^1$  in the  $\sigma(L^1, L^\infty)$  topology, i.e.,

$$\forall g \in L^\infty, \int g f_{n_k} d\mathbb{P} \rightarrow \int g f d\mathbb{P}.$$

Let us make some remarks.

**Remark 1.** Proposition 1 implies that: Any uniformly continuously integrable sequence possesses a subsequence such that the sequence of its mathematical expectations converges in  $\mathbb{R}$ , since

$$\mathbb{E}_{\mathbb{P}}(f_{n_k}) = \int 1_{\Omega} f_{n_k} d\mathbb{P} \rightarrow \int 1_{\Omega} f d\mathbb{P} = \mathbb{E}_{\mathbb{P}}(f)$$

This fact is very important in some mathematical problems. For example, it is used in the extension of the Doob decomposition of super-martingales for discrete sequences to time-continuous versions.

**Remark 2.** Proposition 1 is very important in some mathematical problems. For example, it is used in the extension of the Doob decomposition of super-martingales for discrete sequences to time-continuous versions.

**Remark 3.** From the weak convergence theory in the sense of Billingsley (1968), the problem posed here is very similar to Prohorov's theorem except we do not have a topology on the domain  $\Omega$  of the  $f_n$ 's. The idea is that we might give a proof following the lines of that of Prohorov's theorem (see Theorem in Lo et al. (2016)) in which the tightness hypothesis will be replaced by the *u.c.i* condition. The proof given below realizes that idea. Let us proceed to the proof.

## 2. PROOF

Let us divide the proof into two parts.

**Part 1.** Let us suppose that the  $f_n$ 's are non-negative. Let us set measures  $\nu_n$ :

$$d\nu_n = f_n d\mathbb{P}, \quad n \geq 1.$$

Let

$$\mathcal{C}_0 = \{g \in L^\infty, \|g\| \leq 1\}.$$

Since the  $f_n$ 's are *u.c.i*, we have for  $n \geq 1$ , for  $c > 0$ ,

$$\begin{aligned} \int |f_n| d\mathbb{P} &= \int_{(|f_n| \leq c)} |f_n| d\mathbb{P} + \int_{(|f_n| > c)} |f_n| d\mathbb{P} \\ &\leq c + \sup_{n \geq 1} \int_{(|f_n| > c)} |f_n| d\mathbb{P} \\ &= c + \bar{\phi}(c) \end{aligned}$$

and

$$\bar{\phi}(c) \rightarrow 0, \text{ as } c \rightarrow +\infty.$$

So, for some  $c_0 > 0$ , we have

$$\sup_{n \geq 1} \int |f_n| d\mathbb{P} \leq c_0 + 1 =: C.$$

Now, for  $g \in \mathcal{C}_0, \forall n \geq 1$ ,

$$\left| \int g f_n d\mathbb{P} \right| \leq \int |f_n| d\mathbb{P} \leq C.$$

Hence

$$\bar{\nu}_n = (\nu_n(g), g \in \mathcal{C}_0) \in [0, C]^{\mathcal{C}_0}.$$

Since  $[0, C]^{\mathcal{C}_0}$  as an arbitrary product of the compact set  $[0, C]$ , by Tychonov's theorem, we may extract a sub-sequence  $(\bar{\nu}_{n_k})_{k \geq 1}$  from  $(\bar{\nu}_n)_{n \geq 1}$  converging to  $\bar{\nu} = (\nu(g), g \in \mathcal{C}_0)$ , that is

$$\forall g \in \mathcal{C}_0, \nu_{n_k}(g) \rightarrow \nu(g),$$

with

$$\forall g \in \mathcal{C}_0, \nu(g) \leq C.$$

We have to show that  $\nu$  is still a measure on  $(\Omega, \mathcal{A}, \mathbb{P})$  for  $g = 1_A$ ,  $A \in \mathcal{A}$ . But in general, we have for  $(\alpha, \beta) \in \mathbb{R}^2$ :  $|\alpha| + |\beta| \leq 1$ ,  $(g, h) \in \mathcal{C}_0^2$ ,

$$\begin{aligned} \nu_{n_k}(\alpha g + \beta h) &= \alpha \nu_{n_k}(g) + \beta \nu_{n_k}(h) \\ &\rightarrow \alpha \nu(g) + \beta \nu(h) \text{ as } k \rightarrow +\infty \end{aligned}$$

and since  $\alpha g + \beta h \in \mathcal{C}_0$ , we have

$$\nu_{n_k}(\alpha g + \beta h) \rightarrow \nu(\alpha g + \beta h) \text{ as } k \rightarrow +\infty$$

and hence

$$\nu(\alpha g + \beta h) = \alpha \nu(g) + \beta \nu(h).$$

We prove similarly that for  $(g_1, \dots, g_p) \in \mathcal{C}_0^p$ ,  $p \geq 2$ , for  $(\alpha_1, \dots, \alpha_p) \in \mathbb{R}^p$ :  $|\alpha_1| + \dots + |\alpha_p| \leq 1$ ,

$$\nu_{n_k}(\alpha_1 g_1 + \dots + \alpha_p g_p) \rightarrow \alpha_1 \nu(g_1) + \dots + \alpha_p \nu(g_p).$$

Now for any pair-wise disjoint measurable sets  $(A_j)_{1 \leq j \leq p}$ ,  $1_{\sum_{j=1}^p A_j} \in \mathcal{C}_0$ , and

$$\nu_{n_k} \left( \sum_{j=1}^p A_j \right) =: \nu_{n_k} \left( 1_{\sum_{j=1}^p A_j} \right) \rightarrow \nu \left( 1_{\sum_{j=1}^p A_j} \right)$$

and

$$\nu_{n_k} \left( \sum_{j=1}^p A_j \right) = \sum_{j=1}^p \nu_{n_k}(1_{A_j}) \rightarrow \sum_{j=1}^p \nu(1_{A_j}).$$

So,

$$\nu \left( \sum_{j=1}^p A_j \right) = \sum_{j=1}^p \nu(A_j).$$

Let us prove that  $\nu$  is a measure by extending the finite additivity to a  $\sigma$ -additivity.

But since  $\nu$  is non-negative, finite and additive, so it is enough to show that  $\nu$  is continuous at  $\emptyset$ :

$$\forall A_p \downarrow \emptyset, \nu(A_p) \downarrow 0 \text{ as } p \uparrow +\infty.$$

By Theorem [Lo \(2018\)](#), an *u.c.i* sequence is also uniformly absolutely:

$$\forall \varepsilon > 0, \left( \exists \delta > 0, \forall A \in \mathcal{A} : \mathbb{P}(A) < \delta, \sup_{n \geq 1} \int_A |f_n| d\mathbb{P} < \varepsilon \right).$$

Let  $A_p \downarrow \emptyset$ ,  $A_p \in \mathcal{A}$ . So there exists  $p_0 \in \mathbb{N}$ , such that for any  $p \geq p_0$ ,  $\mathbb{P}(A_p) < \varepsilon$  and hence

$$\sup_{k \geq 1} \int_{A_p} |f_{n_k}| d\mathbb{P} < \varepsilon,$$

that is,

$$\forall p \geq p_0, \forall k \geq 1, |\nu_{n_k}(A_p)| \leq \int_{A_p} |f_{n_k}| d\mathbb{P} < \varepsilon.$$

By letting  $k \rightarrow +\infty$ , we get

$$\forall p \geq p_0, \nu(A_p) < \varepsilon.$$

We proved that

$$\nu(A_p) \rightarrow 0 \text{ as } p \rightarrow +\infty.$$

$\nu$  is non-decreasing as an additive and non-negative mapping on  $\mathcal{A}$ . So  $\nu$  is a measure on  $(\Omega, \mathcal{A}, \mathbb{P})$ , Let us show that  $\nu$  is continuous with respect to  $\mathbb{P}$ , that is

$$d\nu = f d\mathbb{P}, f \in L^1.$$

We have for  $A \in \mathcal{A}$  such that  $\mathbb{P}(A) = 0$ ,

$$0 = \int_A f_{n_k} d\mathbb{P} = \nu_{n_k}(A) \rightarrow \nu(A)$$

and then  $\nu(A) = 0$ . So  $\nu$  is continuous with respect to  $\mathbb{P}$ . By Random Nikodym theorem, there exists  $f \in L^1$  (since  $\nu$  is finite) such that

$$d\nu = f d\mathbb{P}.$$

So

$$(2.1) \quad \forall A \in \mathcal{A}, \quad \int 1_A f_{n_k} d\mathbb{P} \rightarrow \int 1_A f d\mathbb{P}.$$

Now, (2.1) is automatically holds if  $1_A$  is replaced by an elementary function on  $(\Omega, \mathcal{A}, \mathbb{P})$ . Next any element  $g$  of  $\mathcal{C}_0$  is limit of a sequence  $(g_p)$  of elementary functions with  $|g_p| \leq |g| \leq 1$ . So for  $p \geq 1, k \geq 1, \varepsilon > 0$ , we have

$$\begin{aligned} & \left| \int g f_{n_k} d\mathbb{P} - \int g f d\mathbb{P} \right| \\ & \leq \left| \int (g - g_p) f_{n_k} d\mathbb{P} \right| + \left| \int g_p f_{n_k} d\mathbb{P} - \int g_p f d\mathbb{P} \right| \\ & \leq \left| \int_{(|g-g_p| \leq \varepsilon)} (g - g_p) f_{n_k} d\mathbb{P} \right| + \left| \int_{(|g-g_p| > \varepsilon)} (g - g_p) f_{n_k} d\mathbb{P} \right| + \left| \int g_p f_{n_k} d\mathbb{P} - \int g_p f d\mathbb{P} \right| \\ & \leq \varepsilon \int f_{n_k} d\mathbb{P} + \int 1_{(|g-g_p| > \varepsilon)} f_{n_k} d\mathbb{P} + \left| \int g_p f_{n_k} d\mathbb{P} - \int g_p f d\mathbb{P} \right|. \end{aligned}$$

By letting  $p \geq 1$  and  $\varepsilon > 0$  fixed, and letting  $k \rightarrow +\infty$ , we get

$$\limsup_{k \rightarrow +\infty} \left| \int g f_{n_k} d\mathbb{P} - \int g f d\mathbb{P} \right| \leq \varepsilon \int f d\mathbb{P} + \int 1_{(|g-g_p| > \varepsilon)} f d\mathbb{P} + \left| \int g_p f d\mathbb{P} - \int g f d\mathbb{P} \right|.$$

Now, we let  $p \rightarrow +\infty$ , to get

$$\left| \int g_p f d\mathbb{P} - \int g f d\mathbb{P} \right| \rightarrow 0,$$

by the dominated convergence theorem applied to  $L^1 \ni (2f) \geq |g - g_p| \rightarrow 0$  and

$$\int 1_{(|g-g_p| > \varepsilon)} f d\mathbb{P} \rightarrow 0,$$

since  $g_p \rightarrow g$  and hence  $g_p \rightarrow_{\mathbb{P}} g$  and next  $\mathbb{P}(|g - g_p| > \varepsilon) \rightarrow 0$  and finally the integrable function  $f$  is continuous. We get for any  $\varepsilon > 0$

$$\limsup_{k \rightarrow +\infty} \left| \int g f_{n_k} d\mathbb{P} - \int g f d\mathbb{P} \right| \leq \varepsilon \int f d\mathbb{P}.$$

We conclude by letting  $\varepsilon \downarrow 0$  and get

$$\forall g \in \mathcal{C}_0, \int g f_{n_k} d\mathbb{P} \rightarrow \int g f d\mathbb{P}.$$

Finally, for any  $g \in L^\infty$ ,  $\|g\|_\infty \geq 1$ ,

$$\bar{g} = \frac{g}{\|g\|_\infty} \in \mathcal{C}_0$$

and

$$\int g f_{n_k} d\mathbb{P} = \|g\|_\infty \int \bar{g} f_{n_k} d\mathbb{P} \rightarrow \|g\|_\infty \int \bar{g} f d\mathbb{P} = \int g f d\mathbb{P}.$$

The proof of **Part 1** is complete.

**Part 2.** Here the  $f_n$ s are not necessarily non-negative. We define for  $g \in \mathcal{C}_0$

$$\begin{aligned} \nu_n(g) &= \int g f_n^+ d\mathbb{P} - \int g f_n^- d\mathbb{P} \\ &= \nu_n^+(g) - \nu_n^-(g). \end{aligned}$$

The sequences  $(f_n^+)_{n \geq 1}$  and  $(f_n^-)_{n \geq 1}$  are still *u.c.i* since for example

$$\int_{(f_n^+ > c)} f_n^+ d\mathbb{P} \leq \int_{(|f_n| > c)} f_n^+ d\mathbb{P} \leq \int_{(|f_n| > c)} |f_n| d\mathbb{P} \leq \bar{\phi}(c).$$

So for some constant  $C > 0$ , we have

$$0 \leq \nu_n^+ \leq C \text{ and } 0 \leq \nu_n^- \leq C.$$

By **Part 1**, we extract a subsequence  $(\nu_{n'}^+)_{n' \geq 1}$  from  $(\nu_n^+)_{n \geq 1}$  that weakly converges to  $\nu^+$  with  $d\nu^+ = f^+ d\mathbb{P}$ ,  $f^+ \in L^1$ . Next From  $(\nu_{n'}^-)_{n' \geq 1}$  we extract a subsequence  $(\nu_{n_k}^-)_{k \geq 1}$  and  $(\nu_{n_k}^+)_{k \geq 1}$  that weakly converges to  $\nu^-$  with  $d\nu^- = f^- d\mathbb{P}$ ,  $f^- \in L^1$ . Hence

$$\nu_{n_k}^+ \rightarrow_{\sigma(L^1, L^\infty)} \nu^+ \quad \text{and} \quad \nu_{n_k}^- \rightarrow_{\sigma(L^1, L^\infty)} \nu^-,$$

and

$$\nu_{n_k} = \nu_{n_k}^+ - \nu_{n_k}^- \rightarrow_{\sigma(L^1, L^\infty)} \nu^+ - \nu^- = \nu,$$

with  $d\nu = f d\mathbb{P}$ , with  $f = f^+ - f^- \in L^1$ . ■

#### REFERENCES

- Billingsley, P.(1968). *Convergence of Probability measures*. John Wiley, New-York.
- Kallianpur G.(1980) *Stochastic filtering Theory*. Springer+Business Medias, new York.
- Meyer I.(1966) *Probability and Potential*. Blaisdell Publishing Company, Toronto
- Lo, G.S.(2016). Weak Convergence (IA). Sequences of random vectors. SPAS Books Series. Saint-Louis, Senegal - Calgary, Canada. Doi : 10.16929/sbs/2016.0001. Arxiv : 1610.05415. ISBN : 978-2-9559183-1-9
- Lo, G.S.(2018). *Mathematical Foundation of Probability Theory*. SPAS Books Series. Saint-Louis, Senegal - Calgary, Canada. Doi : <http://dx.doi.org/10.16929/sbs/2016.0008>. Arxiv : [arxiv.org/pdf/1808.01713](http://arxiv.org/pdf/1808.01713)