

COUNTING WEIGHTED MAXIMAL CHAINS IN THE CIRCULAR BRUHAT ORDER

GOPAL GOEL, OLIVIA MCGOUGH, AND DAVID PERKINSON

ABSTRACT. The totally nonnegative Grassmannian $\text{Gr}(k, n)_{\geq 0}$ is the subset of the real Grassmannian $\text{Gr}(k, n)$ consisting of points with all nonnegative Plücker coordinates. The circular Bruhat order is a poset isomorphic to the face poset of Postnikov’s ([Pos05]) positroid cell decomposition of $\text{Gr}_{\geq 0}(k, n)$. We provide a closed formula for the sum of its weighted chains in the spirit of Stembridge ([Ste02]).

1. INTRODUCTION.

Let S_n be the symmetric group on $[n] := \{1, \dots, n\}$. An *inversion* of $\pi \in S_n$ is a pair $i, j \in [n]$ such that $i < j$ and $\pi(i) > \pi(j)$. The number of inversions of π is its *length*, denoted $\ell(\pi)$. The *Bruhat order* on S_n is a partial ordering on S_n , graded by length. Its cover relations have the form $\pi s_{ij} \lessdot \pi$ where $s_{ij} := (i, j)$ is a transposition such that $\ell(\pi) = \ell(\pi s_{ij}) + 1$. The maximal element of the Bruhat order, written in row notation, is $\pi_{\text{top}} = [n, n-1, \dots, 1]$ of length $r := \binom{n}{2}$ and the smallest element is the identity permutation $\text{id} = [1, 2, \dots, n]$ of length 0. In the Bruhat order, each maximal chain has the form $\text{id} = \pi_0 \lessdot \pi_1 \lessdot \dots \lessdot \pi_r = \pi_{\text{top}}$. Let $\alpha_1, \dots, \alpha_n$ be indeterminates. Define the *weight* of a covering $\pi s_{ij} \lessdot \pi$ with $i < j$ to be $\alpha_i + \alpha_{i+1} + \dots + \alpha_{j-1}$, and then define the weight of a maximal chain to be the product of the weights of its cover relations. In a result that extends to all Weyl groups, Stembridge ([Ste02]) shows that the sum of the weights of the maximal chains is

$$\frac{\binom{n}{2}!}{1^{n-1} 2^{n-2} \dots (n-1)!} \prod_{1 \leq i < j \leq n} (\alpha_i + \dots + \alpha_{j-1}).$$

For instance, this formula reduces to $\binom{n}{2}!$ after setting all weights $\alpha_i = 1$.

The *totally nonnegative Grassmannian* $\text{Gr}(k, n)_{\geq 0}$ is introduced in [Pos05] as the subset of points in the real Grassmannian $\text{Gr}(k, n)$ which have all nonnegative Plücker coordinates. It is related to areas as diverse as cluster algebras ([GL]), electrical networks ([Lam18]), solitons ([KW11]), scattering amplitudes in Yang-Mills theory ([AHBC⁺16]), and the mathematical theory of juggling ([KLS13]). Postnikov gave a decomposition of $\text{Gr}(k, n)_{\geq 0}$ into *positroid cells* defined by setting certain Plücker coordinates equal to zero, and he conjectured that this decomposition forms a regular CW-complex. A generalization of that conjecture due to Williams ([Wil07]) was proved by Galashin, Karp, and Lam ([GKL20]). Our object of interest is the face poset of this complex, known as the *circular Bruhat order* ([Pos05, Section 17]). Postnikov’s work provides characterizations in terms of many different combinatorial objects, e.g., decorated permutations, Grassmannian necklaces, Le-diagrams, and equivalence classes of certain plabic (planar, bi-colored) graphs. The list is extended by Knutson, Lam, and Speyer ([KLS13]) to include bounded affine permutations, bounded juggling patterns, and equivalence classes of intervals in the k -Bruhat order for S_n . In this paper, we use the language of bounded affine permutations.

Our purpose is to give a Stembridge-like formula for the circular Bruhat order. We define “circular” analogues of Stembridge’s weights (Definition 3.1) and our main result, Theorem 3.2, provides a closed formula for the sum of the weights of the maximal chains in the circular Bruhat order:

$$f(k, n)(\alpha_1 + \dots + \alpha_n)^{k(n-k)},$$

where $f(k, n)$ is the number of Young tableaux for the $k \times (n-k)$ rectangle (cf. Example 3.3). Section 2 provides background and notation. Section 3 states and proves the main result, Theorem 3.2. The proof relies on two technical lemmas whose proofs are relegated to Section 4. These proofs rely on the interpretation of

2010 *Mathematics Subject Classification.* primary 05E15, secondary 05A15.

Key words and phrases. circular Bruhat order, k -Bruhat order, positroid, totally nonnegative Grassmannian.

The second author’s work is partially supported by the Reed College Robert & Louise Rosenbaum Fellowship.

the circular Bruhat order in terms of intervals in the k -Bruhat order for S_n developed in [KLS13]. We also use a result of Bergeron and Sottile ([BS98, Corollary 1.3.1]) on cyclic shifts of k -Bruhat intervals. Their proof is a consequence of a symmetry they find for Littlewood-Richardson coefficients using geometry. It would be nice to have a purely combinatorial proof of their cyclic shift result.

Acknowledgments. We would like to thank Alex Postnikov for suggesting this project.

2. CIRCULAR BRUHAT ORDER

We recall ideas and notation introduced in [KLS13], which built on earlier work by Postnikov on the totally nonnegative Grassmannian ([Pos05]). Our reference for the affine symmetric group is [BB96]. Let \tilde{S}_n denote the group of *affine permutations* consisting of bijections $f: \mathbb{Z} \rightarrow \mathbb{Z}$ satisfying $f(i+n) = f(i) + n$ for all $i \in \mathbb{Z}$. We use the standard *window* notation $f = [f(1), f(2), \dots, f(n)]$ to represent $f \in \tilde{S}_n$. Define the averaging function on \tilde{S}_n by $\text{av}(f) = \frac{1}{n} \sum_{i=1}^n (f(i) - i)$, and for $0 \leq k \leq n$, let $\tilde{S}_n^k := \text{av}^{-1}(k)$. In particular, \tilde{S}_n^0 is the *affine symmetric group*.

The affine symmetric group is generated by its *simple reflections*:

$$s_i = \begin{cases} [0, 2, 3, \dots, n-1, n+1] & \text{if } i = 0, \\ [1, 2, \dots, i-1, i+1, i, i+2, \dots, n] & \text{if } 0 < i \leq n. \end{cases}$$

For instance,

$$[f(1), \dots, f(n)]_{s_0} = [f(0), f(2), \dots, f(n-1), f(n+1)] = [f(n) - n, f(2), \dots, f(n-1), f(1) + n].$$

Then $(\tilde{S}_n^0, \{s_0, \dots, s_{n-1}\})$ is the affine Coxeter group \tilde{A}_{n-1} and is thus a graded poset under the Bruhat order. The *reflections* for \tilde{A}_{n-1} , i.e., the conjugates of the simple reflections, are

$$(1) \quad [1, 2, \dots, i-1, j-rn, i+1, \dots, j-1, i+rn, j+1, \dots, n]$$

for $1 \leq i < j \leq n$ and $r \in \mathbb{Z}$.

The mapping $[f(1), \dots, f(n)] \mapsto [f(1)-k, \dots, f(n)-k]$ is a bijection $\tilde{S}_n^k \rightarrow \tilde{S}_n^0$, and thus the Bruhat order on \tilde{S}_n^0 induces a graded poset structure on \tilde{S}_n^k for which we now give an explicit description. A pair $(i, j) \in \mathbb{Z}^2$ is an *inversion* for $f \in \tilde{S}_n^k$ if $i < j$ and $f(j) > f(i)$. Define an equivalence relation on the set of inversions by $(i, j) \sim (i', j')$ if $i' = i + rn$ and $j' = j + rn$ for some integer r . Then the *length* of f , denoted $\ell(f)$, is the number of equivalence classes of inversions of f . (This notion of length coincides with that inherited from the Bruhat order ([BB96, Proposition 4.1]).) In general, if $f \in \tilde{S}_n^{k'}$ and $g \in \tilde{S}_n^k$, then $fg \in \tilde{S}_n^{k'+k}$. In particular, \tilde{S}_n^0 acts on \tilde{S}_n^k . If $f, g \in \tilde{S}_n^k$, then f covers g , denoted $g < f$, exactly when $g = ft$ from some reflection t from \tilde{A}_{n-1} and $\ell(f) = \ell(g) + 1$.

A permutation $f \in \tilde{S}_n$ is *bounded* if $i \leq f(i) \leq i+n$ for all $i \in \mathbb{Z}$. For each $0 \leq k \leq n$ the bounded elements of \tilde{S}_n^k are denoted

$$\text{Bound}(k, n) := \left\{ f \in \tilde{S}_n^k : i \leq f(i) \leq i+n \text{ for all } i \in \mathbb{Z} \right\}.$$

By Lemma 3.6 of [KLS13], $\text{Bound}(k, n)$ is a lower order ideal in \tilde{S}_n^k and thus forms a graded poset with rank function given by length. The *dual* of a poset $P = (P, <_P)$, is the poset $P^* = (P, <_{P^*})$ for which $a <_{P^*} b$ if and only if $b <_P a$. We now arrive at our object of study:

Definition 2.1. The *circular Bruhat order* $\text{CB}(k, n)$ is the poset $\text{Bound}(k, n)^*$.

The circular Bruhat order was originally defined in [Pos05] in terms of *decorated permutations*. These are permutations $\pi \in S_n$ for which each fixed point is assigned a color—either black or white. The *anti-excedances* of a decorated permutation π are $i \in [n]$ for which either $\pi^{-1}(i) > i$ or i is a white fixed point. Then $\text{CB}(k, n)$ was defined to be the set of decorated permutations with k anti-excedances and with a poset structure determined by *alignments* and *crossings* in *chord diagrams*. See [Pos05] for details. To go from a bounded affine permutation f to a decorated permutation π , reduce the window of f modulo n , and then color each fixed point i in the resulting permutation black if $f(i) = i$ or white if $f(i) = i + n$. We translate the notion of an anti-excedance from decorated permutations to bounded affine permutations:

Definition 2.2. The *anti-excedances* of a bounded affine permutation $f \in \text{CB}(k, n)$ are the integers $f(i) - n$ such that $i \in [n]$ and $f(i) > n$.

One may check that the elements of $\text{CB}(k, n)$ are exactly the bounded affine permutations with k anti-excedances.

To describe the poset structure on $\text{CB}(k, n)$ in detail, note that for a reflection of an element of $\text{CB}(k, n)$ to remain in $\text{CB}(k, n)$, it is necessary (but not sufficient) that the integer r in (1) be 0 or 1. Thus, for $i, j \in [n]$ with $i \neq j$, we define

$$t_{ij} = \begin{cases} [1, 2, \dots, i-1, j, i+1, \dots, j-1, i, j+1, \dots, n] & \text{if } i < j, \\ [1, 2, \dots, j-1, i-n, j+1, \dots, i-1, j+n, i+1, \dots, n] & \text{if } i > j. \end{cases}$$

The cover relations in $\text{CB}(k, n)$ are given by $g < f$ if and only if there exists t_{ij} such that $g = ft_{ij}$ and $\ell(f) = \ell(g) - 1$ (recalling that $\text{CB}(k, n)$ is the dual of $\text{Bound}(k, n)$).

By Lemma 17.6 of [Pos05], the unique maximal element of $\text{CB}(k, n)$ is

$$f_{\text{top}} := [1+k, 2+k, \dots, n+k].$$

The minimal elements are in bijection with $\binom{[n]}{k}$. Given $\lambda \in \binom{[n]}{k}$, the corresponding minimal element is

$$f_{\text{min}, \lambda}(i) = \begin{cases} i+n & \text{if } i \in \lambda, \\ i & \text{otherwise.} \end{cases}$$

We have $\ell(f_{\text{top}}) = 0$, and $\ell(f_{\text{min}, \lambda}) = k(n-k)$ for any minimal element. Thus, the rank function for $\text{CB}(k, n)$ is $\text{rk}(f) = k(n-k) - \ell(f)$. By Proposition 23.1 of [Pos05], the exponential generating function for the cardinality of $\text{CB}(k, n)$ is

$$\sum_{0 \leq k \leq n} |\text{CB}(k, n)| x^k \frac{y^n}{n!} = e^{xy} \frac{x-1}{x - e^{y(x-1)}}.$$

For the rank generating function of $\text{CB}(k, n)$, see [Wil05].

3. MAIN THEOREM

Definition 3.1. Let $\alpha_1, \dots, \alpha_n$ be indeterminates. The *weight* of a covering $ft_{ij} < f$ in $\text{CB}(k, n)$ is the sum of α_i through α_{j-1} in cyclic order:

$$\text{wt}(ft_{ij} < f) = \begin{cases} \alpha_i + \alpha_{i+1} + \dots + \alpha_{j-1} & \text{if } i < j, \\ \alpha_i + \dots + \alpha_n + \alpha_1 + \alpha_2 + \dots + \alpha_{j-1} & \text{if } i > j. \end{cases}$$

The *weight* of a saturated chain in $\text{CB}(k, n)$ is the product of the weights of its cover relations (the empty chain is assigned weight 1). For $r \in [n]$, a covering is *r-good* if α_r appears in its weight. A saturated chain in $\text{CB}(k, n)$ is *r-good* if all of its cover relations are *r-good*. For arbitrary $r \in \mathbb{Z}$, we define *r-good* covers and chains by replacing r with its representative in $[n]$ modulo n .

Our main theorem is the following.

Theorem 3.2. *The sum of the weights of the maximal chains in $\text{CB}(k, n)$ is*

$$f(k, n)(\alpha_1 + \dots + \alpha_n)^{k(n-k)}$$

where $f(k, n)$ is the number of standard Young tableaux of a $k \times (n-k)$ rectangle.

Example 3.3. Figure 1 illustrates $\text{CB}(2, 3)$ with its cover weights. The sum of the weights of its six maximal chains is


$$\alpha_1\alpha_2 + \alpha_1(\alpha_1 + \alpha_3) + \alpha_2(\alpha_1 + \alpha_2) + \alpha_2\alpha_3 + \alpha_3\alpha_1 + \alpha_3(\alpha_2 + \alpha_3) = f(2, 3)(\alpha_1 + \alpha_2 + \alpha_3)^2,$$

where $f(2, 3) = 1$ since there is only one Young tableau for the 2×1 rectangle.

The proof of Theorem 3.2 follows from two lemmas whose proofs appear in next section.

Lemma 3.4. *Let $f \in \text{CB}(k, n)$. Then the number of *r-good* downward-saturated chains in $\text{CB}(k, n)$ with maximal element f is independent of r .*

Lemma 3.5. *The number of *n-good* maximal chains in $\text{CB}(k, n)$ is $f(k, n)$.*

FIGURE 1. $CB(2,3)$ with edge weights.

Proof of Theorem 3.2. Let $\delta(f)$ be the number of r -good downward-saturated chains with maximal element $f \in CB(k, n)$. This number is independent of r by Lemma 3.4. Noting linearity in the α_i , it is straightforward to check that

$$(2) \quad \sum_{t_{ij}: ft_{ij} < f} \delta(ft_{ij}) \text{wt}(ft_{ij} < f) = \delta(f)(\alpha_1 + \cdots + \alpha_n).$$

Let $\mathcal{C}(m)$ denote the set of maximal chains C in $CB(k, n)$ such that $\max(C) = f_{\text{top}}$ and $\ell(\min(C)) = m$. We now show by induction on m that

$$\sum_{C \in \mathcal{C}(m)} \delta(\min(C)) \text{wt}(C) = \delta(f_{\text{top}})(\alpha_1 + \cdots + \alpha_n)^m$$

for $0 \leq m \leq k(n-k)$. The case $m = 0$ is a tautology since the only element of $CB(k, n)$ with length 0 is f_{top} . On the other hand, in the case $m = k(n-k)$ we are summing over maximal chains C of $CB(k, n)$. For these $\delta(\min(C)) = 1$, and Theorem 3.2 will then follow from Lemma 3.5. To proceed with induction, fix some m with $0 \leq m < k(n-k)$. Then

$$\begin{aligned} \sum_{C \in \mathcal{C}(m+1)} \delta(\min(C)) \text{wt}(C) &= \sum_{C' \in \mathcal{C}(m)} \sum_{f < \min(C')} \delta(f) \text{wt}(f < \min(C')) \text{wt}(C') \\ &= \sum_{C' \in \mathcal{C}(m)} \delta(\min(C')) (\alpha_1 + \cdots + \alpha_n) \text{wt}(C') && \text{(by (2))} \\ &= \delta(f_{\text{top}}) (\alpha_1 + \cdots + \alpha_n)^{m+1} && \text{(by induction).} \end{aligned}$$

□

4. PROOFS OF LEMMAS

4.1. **k -Bruhat order.** Our references for the k -Bruhat order are [BS98] and [KLS13].

Definition 4.1. The k -Bruhat order \leq_k on the symmetric group S_n is given by $u \leq_k v$ if

- (1) $u(i) \leq v(i)$ for $1 \leq i \leq k$;
- (2) $u(j) \geq v(j)$ for $k < j \leq n$;
- (3) $u(i) < u(j)$ implies $v(i) < v(j)$ if $1 \leq i < j \leq k$ or if $k < i < j \leq n$

The cover relations for the k -Bruhat order have the form $u \lessdot_k v$ if $u \lessdot v$ (in the ordinary Bruhat order) and $\{u(1), \dots, u(k)\} \neq \{v(1), \dots, v(k)\}$. Each interval $[u, w]_k$ in the k -Bruhat order is a graded poset of rank $\ell(w) - \ell(u)$.

Definition 4.2. A permutation $w \in S_n$ is k -Grassmannian (or just *Grassmannian* when k is clear from context) if $w(1) < \cdots < w(k)$ and $w(k+1) < \cdots < w(n)$. These are in bijection with $\lambda \in \binom{[n]}{k}$ by letting w_λ be the unique k -Grassmannian permutation such that $\{w(1), \dots, w(k)\} = \lambda$.

Denote the positions of the anti-excedances of $f \in \text{CB}(k, n)$ by

$$\Lambda(f) = \{i \in [n] : f(i) - n \text{ is an anti-excedance of } f\} = \{i \in [n] : f(i) > n\}.$$

Then associate a k -Grassmannian permutation to f by

$$w_f = w_{\Lambda(f)}.$$

Fixing $\lambda \in \binom{[n]}{k}$, we define two posets. The first is the principal order ideal in the k -Bruhat order generated by w_λ :

$$S_{n,\lambda} = \{u \in S_n : u \leq_k w_\lambda\}.$$

The second is

$$\text{CB}(k, n)_\lambda = \{f \in \text{CB}(k, n) : \Lambda(f) = \lambda\}$$

with partial order \leq_γ defined by its cover relations: $g \leq_\gamma f$ if g is covered by f in $\text{CB}(k, n)$ and the covering $g \leq f$ is n -good (γ is a mnemonic for “good”). Note that a covering $ft_{ij} \leq f$ in $\text{CB}(k, n)$ is n -good if and only if $i > j$, in which case $\Lambda(ft_{ij}) = \Lambda(f)$.

Embed S_n in \tilde{S}_n via $u \mapsto [u(1), \dots, u(n)]$ and define the *translation element* $t_k = [1 + n, 2 + n, \dots, k + n, k + 1, k + 2, \dots, n] \in \tilde{S}_n^k$. Taking our lead from [KLS13], for each $u \in S_{n,\lambda}$ define $f_u = f_{u,\lambda} = ut_k w_\lambda^{-1}$. Therefore,

$$f_u(w_\lambda(i)) = \begin{cases} u(i) + n & \text{if } 1 \leq i \leq k \\ u(i) & \text{if } k < i \leq n. \end{cases}$$

Since w_λ is k -Grassmannian and $u \leq_k w_\lambda$, we have $1 \leq u(i) \leq w_\lambda(i) \leq n$ for $1 \leq i \leq k$, and $w_\lambda(i) \leq u(i) \leq n$ for $k < i \leq n$. Therefore, $i \leq f(i) \leq i + n$ for all i . Further, $\Lambda(f_u) = \lambda$. Hence, $f_u \in \text{CB}(k, n)_\lambda$.

For each $f \in \text{CB}(k, n)_\lambda$, define $u_f = u_{f,\lambda} = fw_\lambda t_k^{-1}$ so that

$$u_f(i) = \begin{cases} f(w_\lambda(i)) - n & \text{if } 1 \leq i \leq k \\ f(w_\lambda(i)) & \text{if } k < i \leq n. \end{cases}$$

Since $\lambda = \Lambda(f)$, it follows that $1 \leq u_f(i) \leq n$ for $i \in [n]$. To see that $u \leq_k w_\lambda$, first note that $w_\lambda(i) \leq f(w_\lambda(i)) \leq w_\lambda(i) + n$ for all i since $f \in \text{CB}(k, n)$. Properties (1) and (2) of Definition 4.1 then follow. Property 2 holds since w_λ is a k -Grassmannian element and, therefore, $w_\lambda(i)$ is increasing for $1 \leq i \leq k$ and for $k < i \leq n$.

Example 4.3. Let $\lambda = \{2, 4, 5\} \in \binom{[5]}{3}$ and $f = [3, 6, 5, 9, 7]$. Then $w_\lambda = [2, 4, 5, 1, 3]$, and $f \in \text{CB}(3, 5)_\lambda$ since its anti-excedances appear in positions 2, 4, and 5. We have $u_f = [1, 4, 2, 3, 5]$, which is formed by first listing the anti-excedances of f , reduced modulo 5, as they appear in order by position in f , i.e., $1 = 6 - 5$, $4 = 9 - 5$, and $2 = 7 - 5$, and then listing the non-anti-excedances, 3 and 5. Reversing this process yields $f_{u_f} = f$.

Proposition 4.4. ¹ Let $\lambda \in \binom{[n]}{k}$. Then the mapping

$$\begin{aligned} (S_{n,\lambda}, \leq_k) &\rightarrow (\text{CB}(k, n)_\lambda, \leq_\gamma) \\ u &\mapsto f_u \end{aligned}$$

is an anti-isomorphism of posets with inverse $f \mapsto u_f$.

Proof. It is clear that $u \mapsto f_u$ and $f \mapsto u_f$ are inverses. We must show they reverse cover relations. Let $u, v \in S_{n,\lambda}$ with corresponding $f := f_u$ and $g := f_v$ in $\text{CB}(k, n)$. The condition that $u \leq_k v$ is equivalent to:

1. There exists $p \leq k < q$ such that $v = us_{p,q}$ where $s_{p,q} = (p, q)$ is the transposition swapping p and q , and
2. $\ell(v) = \ell(u) + 1$, i.e.,
 - (i) $u(p) < u(q)$, and
 - (ii) there is no integer r such that $p < r < q$ and $u(p) < u(r) < u(q)$.

On the other hand, the condition that $g \leq_\gamma f$ is equivalent to:

- 1*. There exists $i < j$ such that $f(i)$ is a non-anti-excedance, $f(j)$ is an anti-excedance, $g = ft_{ji}$, and

¹For a closely related result, see [KLS13, Theorem 3.16].

- 2*. $\ell(g) = \ell(f) + 1$, i.e.,
- (i) $f(j) < f(i) + n$, and
 - (ii) there is no integer a such that $j < a < i + n$ and $f(j) < f(a) < f(i) + n$.

To show equivalence of these two sets of conditions, first suppose that $u \leq_k v \leq_k w_\lambda$. We will show that $g \leq_\gamma f$. Take $p \leq k < q$ as in condition 1, and let $i := w_\lambda(q)$ and $j := w_\lambda(p)$. It follows from the k -Bruhat order that $i < j$:


$$i = w_\lambda(q) \leq v(q) = u(p) \leq v(p) \leq w_\lambda(p) = j.$$

We have that $f(r) = g(r)$ for $r \in [n] \setminus \{i, j\}$, and

$$\begin{aligned} g(i) &= g(w_\lambda(q)) = v(q) = u(p) = f(w_\lambda(p)) - n = f(j) - n \\ g(j) &= g(w_\lambda(p)) = v(p) + n = u(q) + n = f(w_\lambda(q)) + n = f(i) + n. \end{aligned}$$

Therefore, condition 1* holds, and 2*(i) follows from 2(i).

Condition 2*(ii) says that the graph of f has no points inside a certain box:



To verify condition 2*(ii) holds, it helps to divide the sequence of integers $j, j+1, \dots, i+n$ into two parts: $X := \{a \in \mathbb{Z} : j < a \leq n\}$, and $Y := \{a \in \mathbb{Z} : n < a < i+n\}$. If $a \in X$ and $f(a)$ is not an anti-excedance, then $f(a) < n < f(j)$ and, hence, condition 2*(ii) is not violated. Similarly, if $a \in Y$ and $f(a-n)$ is an anti-excedance, then $f(i+n) \leq 2n < f(a-n) + n = f(a)$ and, hence, 2*(ii) is again not violated.

It remains to check anti-excedances whose positions are in X and non-anti-excedances whose positions are between 1 and i (i.e., are in $-n + Y$). Take $a \in X$ and suppose that $f(a)$ is an anti-excedance. Since $a > j = w_\lambda(p)$, there exists r with $p < r \leq k < q$ such that $f(a) = u(r) + n$. By condition 2(ii), $u(r)$ is not between $u(p)$ and $u(q)$, which implies that $f(a)$ is not between $f(i) + n = u(q) + n$ and $f(j) = u(p) + n$ in accordance with 2*(ii). Now, instead, take $a \in Y$ and suppose that $f(a-n)$ is not an anti-excedance. Since $a-n < i = w_\lambda(q)$, there exists r with $p \leq k < r < q$ such that $f(a-n) = u(r)$. As above, $u(r)$ is not between $u(p)$ and $u(q)$, and this implies that $f(a) = u(r) + n$ is not between $f(i) + n = u(q) + n$ and $f(j) = u(p) + n$.

We have shown that the mapping $u \mapsto f_u$ reverses cover relations. The proof that its inverse $f \mapsto u_f$ reverses cover relations is similar. \square

Corollary 4.5. *Let $f \in \text{CB}(k, n)$ with corresponding Grassmannian permutation w_f . Then the n -good downward-saturated chains in $\text{CB}(k, n)$ with maximal element f are in bijection with the maximal chains in the k -Bruhat interval $[u_f, w_f]_k$.*

Proof. Let $\lambda = \Lambda(f)$. Since an n -good covering preserves anti-excedance positions, downward-saturated n -good chains in $\text{CB}(k, n)$ with maximal element f are exactly downward saturated chains in $\text{CB}(k, n)_\lambda$ with maximal element f . By Proposition 4.4 these are in bijection with upward saturated chains in $S_{n, \lambda}$ with minimal element u_f . Since $S_{n, \lambda}$ has unique maximal element $w_\lambda = w_f$, these upward-saturated chains are exactly the maximal chains in $[u_f, w_f]_k$. \square

Example 4.6. Let $f = [2, 5, 4, 7] \in \text{CB}(2, 4)$. Then $w_f = [2, 4, 1, 3] \in S_n$. We have $u_f = [1, 3, 2, 4]$ and $f_{w_f} = [1, 6, 3, 8]$. As seen in Figure 2, the interval $[u_f, w_f]_2$ has two maximal chains:

$$[1, 3, 2, 4] \leq_2 [1, 4, 2, 3] \leq_2 [2, 4, 1, 3] \quad \text{and} \quad [1, 3, 2, 4] \leq_2 [2, 3, 1, 4] \leq_2 [2, 4, 1, 3].$$

Under the isomorphism of Proposition 4.4, these correspond to the two n -good downward-saturated chains in $\text{CB}(2, 4)$ with maximal element f :

$$[1, 6, 3, 8] < [2, 5, 3, 8] < [2, 5, 4, 7] \quad \text{and} \quad [1, 6, 3, 8] < [1, 6, 4, 7] < [2, 5, 4, 7].$$




FIGURE 2. The posets $S_{4,\lambda}$ and $CB(2,4)_\lambda$ in the case $\lambda = \{2,4\} \in \binom{[4]}{2}$. They are anti-isomorphic in accordance with Proposition 4.4.

4.2. **Cyclic shifts.** Define the *cyclic shift* of $f \in CB(k,n)$ by

$$\chi(f) := [f(0) + 1, f(1) + 1, \dots, f(n-1) + 1].$$

The following properties of χ are immediate: (i) $i \leq \chi(f)(i) \leq i+n$ for all $i \in [n]$, (ii) $\chi(ft_{ij}) = \chi(f)t_{i+1,j+1}$ for all reflections $t_{i,j}$ (with indices taken modulo n), and (iii) $(i,j) \in \mathbb{Z}^2$ represents an inversion for f if and only if $(i+1,j+1)$ represents an inversion for $\chi(f)$. Therefore, χ is an automorphism of the graded poset $CB(k,n)$ and induces a faithful action of the cyclic group of order n on $CB(k,n)$. The following is an immediate implication of property (ii).

Proposition 4.7. *A saturated chain C in $CB(k,n)$ is r -good if and only if $\chi(C)$ is $(r+1)$ -good.*

The following result of Bergeron and Sottile is an important step in the proof of Lemma 3.4.

Theorem 4.8 ([BS98, Corollary 1.3.1]). *Let $u \leq_k v$ and $x \leq_k y$ in the k -Bruhat order on S_n , and suppose that $cvu^{-1}c^{-1} = yx^{-1}$ where $c = [2, 3, \dots, n, 1] = (1, 2, \dots, n)$. Then the intervals $[u, v]_k$ and $[x, y]_k$ have the same number of maximal chains.*

4.3. Proofs of lemmas.

Proof of Lemma 3.4. For general $g \in CB(k,n)$, let $\delta_r(g)$ denote the number of r -good downward-saturated chains in $CB(k,n)$ with maximal element g . Fix $f \in CB(k,n)$. We first show that Theorem 4.8 applies to the pair of intervals $[u_f, w_f]_k$ and $[u_{\chi(f)}, w_{\chi(f)}]_k$ by checking that $cu_f w_f^{-1} = u_{\chi(f)} w_{\chi(f)}^{-1} c$ where $c = [2, 3, \dots, n, 1]$. Let $i \in [n]$. Working modulo n ,

$$cu_f w_f^{-1}(i) = cu_f t_k w_f^{-1}(i) = f(i) + 1 = \chi(f)(i+1) = u_{\chi(f)} t_k w_{\chi(f)}^{-1} c(i) = u_{\chi(f)} w_{\chi(f)}^{-1} c(i),$$

and the conclusion follows. Therefore, the number of maximal chains in $[u_f, w_f]_k$ is the same as the number of maximal chains in $[u_{\chi(f)}, w_{\chi(f)}]_k$ and, by induction, as the number of maximal chains in $[u_{\chi^r(f)}, w_{\chi^r(f)}]_k$ for all $r \in [n]$. Applying Corollary 4.5 and Proposition 4.7,

$$\delta_n(f) = \delta_n(\chi^r(f)) = \delta_r(f)$$

for all $r \in [n]$. □

Proof of Lemma 3.5. Let $w_{\max} = w_{\{n-k+1, n-k+2, \dots, n\}}$ and $\text{id} = [1, 2, \dots, n]$. Then $w \leq_k w_{\max}$ for all k -Grassmannian permutations w and $\text{id} \leq_k u$ for all $u \in S_n$. Thus, by Corollary 4.5, maximal n -good chains in $CB(k,n)$ are in correspondence with maximal chains in $[\text{id}, w_{\max}]_k$, an interval of rank $k(n-k)$. This interval is exactly the set of all k -Grassmannian elements of S_n .

Let $L(k, n-k)$ denote the poset of Young diagrams fitting inside a $k \times (n-k)$ rectangle, ordered by containment, as usual. There is a well-known correspondence between $\binom{[n]}{k}$ and $L(k, n-k)$: given $\lambda \in \binom{[n]}{k}$ with $\lambda = \{\lambda_1 < \dots < \lambda_k\}$, let $Y_{p(\lambda)}$ be the Young diagram corresponding to the partition $p(\lambda) = \{p_1 > \dots > p_k\}$ where $p_i := (n-k) - \lambda_i + i$. In English notation, $Y_{p(\lambda)}$ is the diagram determined by the left-down walk in \mathbb{Z}^2 from $(k, n-k)$ to $(0,0)$ whose λ_i -th step is its i -th *vertical* step. This correspondence yields an anti-isomorphism $w_\lambda \mapsto Y_{p(\lambda)}$ from the interval $[\text{id}, w_{\max}]_k$ with its k -Bruhat order and $L(k, n-k)$. The

result now follows since Young tableaux for the $k \times (n - k)$ rectangle are in bijection with maximal chains in $L(k, n - k)$. \square

REFERENCES

- [AHBC⁺16] Nima Arkani-Hamed, Jacob Bourjaily, Freddy Cachazo, Alexander Goncharov, Alexander Postnikov, and Jaroslav Trnka. *Grassmannian geometry of scattering amplitudes*. Cambridge University Press, Cambridge, 2016.
- [BB96] Anders Björner and Francesco Brenti. Affine permutations of type A . *Electron. J. Combin.*, 3(2):Research Paper 18, approx. 35, 1996. The Foata Festschrift.
- [BS98] Nantel Bergeron and Frank Sottile. Schubert polynomials, the Bruhat order, and the geometry of flag manifolds. *Duke Math. J.*, 95(2):373–423, 1998.
- [GKL20] Pavel Galashin, Steven N. Karp, and Thomas Lam. Regularity theorem for totally nonnegative flag varieties. *Sém. Lothar. Combin.*, 84B:Art. 31, 12, 2020.
- [GL] Pavel Galashin and Thomas Lam. Positroid varieties and cluster algebras. To appear in *Ann. Sci. Éc. Norm. Supér.*
- [KLS13] Allen Knutson, Thomas Lam, and David E. Speyer. Positroid varieties: juggling and geometry. *Compos. Math.*, 149(10):1710–1752, 2013.
- [KW11] Yuji Kodama and Lauren K. Williams. KP solitons, total positivity, and cluster algebras. *Proc. Natl. Acad. Sci. USA*, 108(22):8984–8989, 2011.
- [Lam18] Thomas Lam. Electroid varieties and a compactification of the space of electrical networks. *Adv. Math.*, 338:549–600, 2018.
- [Pos05] Alexander Postnikov. Total positivity, Grassmannians, and networks. Preprint (2005) <http://www-math.mit.edu/~apost/papers/tpgrass.pdf>, 2005.
- [Ste02] John R. Stembridge. A weighted enumeration of maximal chains in the Bruhat order. *J. Algebraic Combin.*, 15(3):291–301, 2002.
- [Wil05] Lauren K. Williams. Enumeration of totally positive Grassmann cells. *Adv. Math.*, 190(2):319–342, 2005.
- [Wil07] Lauren K. Williams. Shelling totally nonnegative flag varieties. *J. Reine Angew. Math.*, 609:1–21, 2007.

M.I.T., CAMBRIDGE, MA
Email address: `gopalkg@mit.edu`

REED COLLEGE, PORTLAND, OR
Email address: `mcgougho@reed.edu`

REED COLLEGE, PORTLAND, OR
Email address: `davidp@reed.edu`