

**THE p -PRIMARY SUBGROUPS OF THE COHOMOLOGY OF
 BPU_n IN DIMENSIONS LESS THAN $2p + 5$**

XING GU, YU ZHANG, ZHILEI ZHANG, AND LINAN ZHONG*

ABSTRACT. Let PU_n denote the projective unitary group of rank n and BPU_n be its classifying space. For an odd prime p , we extend previous results to a complete description of $H^s(BPU_n; \mathbb{Z})_{(p)}$ for $s < 2p + 5$ by showing that the p -primary subgroups of $H^s(BPU_n; \mathbb{Z})$ are trivial for $s = 2p + 3$ and $s = 2p + 4$.

1. INTRODUCTION

Let U_n denote the group of $n \times n$ unitary matrices. The unit circle S^1 can be viewed as the normal subgroup of scalar matrices of U_n . We let PU_n denote the quotient group of U_n by S^1 , and BPU_n be the classifying space of PU_n . In this paper we consider $H^*(BPU_n; \mathbb{Z})$, the ordinary cohomology of BPU_n with coefficients in \mathbb{Z} .

A review of the literature. The ordinary and generalized cohomology of BPU_n for special n has been the subject of various works such as Kono-Mimura [15], Kameko-Yagita [14], Kono-Yagita [16], Toda [19], and Vavpetič-Viruel [21]. Vezzosi [22] and Vistoli [23] studied the Chow ring of the classifying space (in the sense of Totaro [20]) of $BPGL_3(\mathbb{C})$ and $BPGL_p(\mathbb{C})$ for p an odd prime, respectively. Much of their results applies to the ordinary cohomology of BPU_p .

None of the works above dealt with $H^*(BPU_n; \mathbb{Z})$ for n not a prime number. The first named author considered $H^*(BPU_n; \mathbb{Z})$, as well as the Chow ring of $BPGL_n(\mathbb{C})$ for an arbitrary n in [10], [12] and [13]. In particular, in [12], the first named author determined the ring structure of $H^*(BPU_n; \mathbb{Z})$ in dimensions less than or equal to 10.

Other related works include Duan [6], in which the integral cohomology of PU_n is fully determined, and Crowley-Gu [5], in which the image of the canonical map $H^*(BPU_n; \mathbb{Z}) \rightarrow H^*(BU_n; \mathbb{Z})$ is studied.

The cohomology of BPU_n plays significant roles in the study of the topological period-index problem ([1], [2], [9] and [11]), and in the study of anomalies in particle physics ([4], [8]).

Notations. Throughout the rest of this paper, $H^*(-)$ denotes $H^*(-; \mathbb{Z})$. For an abelian group A and a prime number p , let $A_{(p)}$ be the localization of A at p , and let ${}_pA$ denotes the p -primary subgroup of A , i.e., the subgroup of A of all torsion elements with torsion order a power of p . In particular, we have a canonical isomorphism ${}_pH^*(-) \cong {}_p[H^*(-)_{(p)}]$, and we will not distinguish the two throughout this paper. Tensor products of $\mathbb{Z}_{(p)}$ -modules are always taken over $\mathbb{Z}_{(p)}$.

2020 *Mathematics Subject Classification.* 55T10, 55R35, 55R40.

* Corresponding author.

The main theorem and some remarks. We review a basic fact on the cohomology of BPU_n . Consider the short exact sequence of Lie groups

$$1 \rightarrow \mathbb{Z}/n \rightarrow SU_n \rightarrow PSU_n \simeq PU_n \rightarrow 1,$$

which induces a fiber sequence of their classifying spaces

$$(1.1) \quad B(\mathbb{Z}/n) \rightarrow BSU_n \rightarrow BPU_n$$

When $p \nmid n$, the space $B(\mathbb{Z}/n)$ is p -locally contractible and we have

$$(1.2) \quad H^*(BPU_n; \mathbb{Z}_{(p)}) \cong H^*(BSU_n; \mathbb{Z}_{(p)})$$

Since $\mathbb{Z}_{(p)}$ is a flat \mathbb{Z} -module, and in particular, $H^*(-; \mathbb{Z}_{(p)}) \cong H^*(-)_{(p)}$, we have an isomorphism of $\mathbb{Z}_{(p)}$ -algebras

$$(1.3) \quad H^*(BPU_n)_{(p)} \cong H^*(BSU_n)_{(p)} = \mathbb{Z}_{(p)}[c_2, c_3, \dots, c_n],$$

which shows $H^*(BPU_n)_{(p)}$ is torsion-free for $p \nmid n$. In other words, we have the following

Proposition 1.1. *Suppose $x \in H^*(BPU_n)$ is a torsion class. Then there exists some $i \geq 0$ such that $n^i x = 0$.*

Therefore, to determine the graded abelian group structure of $H^s(BPU_n)$, it suffices to consider the p -primary subgroup ${}_p H^s(BPU_n)$ for $p \mid n$.

Remark 1.2. In the case of Chow rings, Vezzosi [22] proved the stronger result that all torsion classes in the Chow ring of $BPGL_n(\mathbb{C})$ are n -torsion.

To state the main theorem, recall that, as shown in [12], the integral cohomology group $H^3(BPU_n)$ is generated by a class denoted by x_1 . In addition, P^i will denote the i th Steenrod reduced power operation, and

$$\delta : H^*(-, \mathbb{Z}/p) \rightarrow H^{*+1}(-)$$

will denote the connecting homomorphism. Finally, a bar over an integral cohomology class will denote the mod p reduction of this class. For instance, \bar{x}_1 denotes the mod p reduction of x_1 , which is in $H^3(BPU_n; \mathbb{Z}/p)$.

Theorem 1. *Let $p > 2$ be a prime number, and $n = p^r m$ for a positive integer m co-prime to p . Then the p -primary subgroup of $H^s(BPU_n)$ in dimensions less than $2p + 5$ is as follows:*

(1) *For $r > 0$, we have*

$${}_p H^s(BPU_n) \cong \begin{cases} \mathbb{Z}/p^r, & s = 3, \\ \mathbb{Z}/p, & s = 2p + 2, \\ 0, & s < 2p + 5, \quad s \neq 3, 2p + 2. \end{cases}$$

The group ${}_p H^{2p+2}(BPU_n)$ is generated by $\delta P^1(\bar{x}_1)$.

(2) *For $r = 0$, we simply have ${}_p H^s(BPU_n) = 0$ for all $s \geq 0$.*

Remark 1.3. Note ${}_p H^s(BPU_n) \cong {}_p H^s(BPU_n)_{(p)}$. By the discussion preceding Remark 1.2, Theorem 1 completely determines $H^s(BPU_n; \mathbb{Z}_{(p)})$ for $0 \leq s < 2p + 5$.

For $s \leq 3$, the groups ${}_p H^s(BPU_n)$ are well known and are part of Theorem 1.1 of [12]. For $3 < s < 2p + 2$, they are given in Theorem 1.2 of [12]. Therefore, what remains to show is

$$(1.4) \quad {}_p H^{2p+2}(BPU_n) \cong \mathbb{Z}/p, \quad {}_p H^{2p+3}(BPU_n) = {}_p H^{2p+4}(BPU_n) = 0.$$

Remark 1.4. For $p = 2$, it was shown by the first named author [12] that the 2-torsion subgroup of $H^s(BPU_n)$ in dimension $s = 2p + 3 = 7$ is $\mathbb{Z}/2$ if $n \equiv 2 \pmod{4}$, and is 0 otherwise. In particular, Theorem 1 does not generalize to the case $p = 2$.

Remark 1.5. For $p = 3$, (1.4) follows immediately from the computation in [12] of $H^s(BPU_n)$ in dimensions 8, 9 and 10.

Organization of the paper. In Section 2, we discuss some preliminary results of the Serre spectral sequence ${}^U E$ associated to the fiber sequence $U : BU_n \rightarrow BPU_n \rightarrow K(\mathbb{Z}, 3)$. This will be our main tool for computing the p -primary subgroup ${}_p H^s(BPU_n)$. We will also show that (1.4) can be deduced from Theorem 1.2 of [12] and Proposition 2.10, which says that certain chain complex \mathcal{M} constructed from the differentials in ${}^U E$ is exact.

In Section 3, we prove Proposition 2.10. The proof is based on the explicit computation of some relevant differentials in ${}^U E$. This section finishes our proof of Theorem 1.

Acknowledgments. The authors would like to thank Ben Williams for various helpful editorial suggestions. The first named author would like to thank the hospitality of Professor Jie Wu, as well as the Center for Topology and Geometry based Technology (CTGT) and the School of Mathematical Sciences at Hebei Normal University, and acknowledge the supports from the National Natural Science Foundation of China (No. 21113062) and from the High-level Scientific Research Foundation of Hebei Province (No. 13113093). The second named author and the third named author would like to thank Xiangjun Wang for helpful discussions. The second named author and the third named author were supported by the National Natural Science Foundation of China (No. 11871284). The fourth named author was supported by the National Natural Science Foundation of China (No. 12001474; 11761072). All authors contribute equally.

2. THE SPECTRAL SEQUENCES

The Serre spectral sequence ${}^U E$. We follow the strategy employed in [12] to compute the cohomology of BPU_n . The short exact sequence of Lie groups

$$1 \rightarrow S^1 \rightarrow U_n \rightarrow PU_n \rightarrow 1$$

induces a fiber sequence of their classifying spaces

$$BS^1 \rightarrow BU_n \rightarrow BPU_n.$$

Notice that BS^1 is of the homotopy type of the Eilenberg-Mac Lane space $K(\mathbb{Z}, 2)$ and indeed we obtain another fiber sequence

$$(2.1) \quad U : BU_n \rightarrow BPU_n \xrightarrow{X} K(\mathbb{Z}, 3).$$

Remark 2.1. In general, it is not always possible to obtain a fiber sequence of the form $F \rightarrow E \rightarrow B$ from a fiber sequence $\Omega B \rightarrow F \rightarrow E$. See Ganea [7] for more.

We will use the Serre spectral sequence associated to the last fiber sequence to compute the cohomology of BPU_n . For notational convenience, we denote this spectral sequence by ${}^U E$. The E_2 page of ${}^U E$ has the form

$${}^U E_2^{s,t} = H^s(K(\mathbb{Z}, 3); H^t(BU_n)) \implies H^{s+t}(BPU_n).$$

In principle, Cartan and Serre [3] determined the cohomology of $K(A, n)$ for all finitely generated abelian groups A . Also see Tamanoi [18] for a nice treatment.

We summarize the p -local cohomology of $K(\mathbb{Z}, 3)$ in low dimensions as follows.

Proposition 2.2. *Let $p > 2$ be a prime. In degrees up to $2p + 5$, we have*

$$(2.2) \quad H^s(K(\mathbb{Z}, 3))_{(p)} = \begin{cases} \mathbb{Z}_{(p)}, & s = 0, 3, \\ \mathbb{Z}/p, & s = 2p + 2, 2p + 5, \\ 0, & s < 2p + 5, s \neq 0, 3, 2p + 2. \end{cases}$$

where $x_1, y_{p,0}, x_1 y_{p,0}$ are generators on degree $3, 2p + 2, 2p + 5$ respectively. In addition, we have $y_{p,0} = \delta P^1(\bar{x}_1)$.

Here we use the same notations for the generators as in [12]. Sometimes we abuse notations and let $x_1, y_{p,0}$ denote $\chi^*(x_1), \chi^*(y_{p,0})$, where $\chi : BPU_n \rightarrow K(\mathbb{Z}, 3)$ is defined in (2.1). For instance, we have

Proposition 2.3 (Theorem 1.2, [12]). *Let p be a prime. In $H^{2p+2}(BPU_n)$, we have $y_{p,0} \neq 0$ of order p when $p \mid n$, and $y_{p,0} = 0$ otherwise. Furthermore, the p -torsion subgroup of $H^k(BPU_n)$ is 0 for $3 < k < 2p + 2$.*

Also recall

$$(2.3) \quad H^*(BU_n) = \mathbb{Z}[c_1, c_2, \dots, c_n], \quad |c_i| = 2i.$$

In particular, $H^*(BU_n)$ is torsion-free. We have

$${}^U E_2^{s,t} \cong H^s(K(\mathbb{Z}, 3)) \otimes H^t(BU_n).$$

The auxiliary fiber sequences and spectral sequences. To determine some of the differentials in ${}^U E$, we consider two more fiber sequences.

Let T^n be the maximal torus of U^n with the inclusion denoted by

$$\psi : T^n \rightarrow U_n.$$

Passing to quotients over S^1 , we have another inclusion of maximal torus

$$\psi' : PT^n \rightarrow PU_n.$$

The quotient map $T^n \rightarrow PT^n$ fits in an exact sequence of Lie groups

$$1 \rightarrow S^1 \rightarrow T^n \rightarrow PT^n \rightarrow 1,$$

which induces a fiber sequence

$$T : BT^n \rightarrow BPT^n \rightarrow K(\mathbb{Z}, 3).$$

Notice that we have

$$(2.4) \quad H^*(BT^n) = \mathbb{Z}[v_1, v_2, \dots, v_n], \quad |v_i| = 2.$$

The next fiber sequence is simply the path fibration for the space $K(\mathbb{Z}, 3)$

$$K : K(\mathbb{Z}, 2) \rightarrow * \rightarrow K(\mathbb{Z}, 3)$$

where $*$ denotes a contractible space. We denote their associated Serre spectral sequences as ${}^T E$ and ${}^K E$ respectively.

We denote the corresponding differentials of ${}^U E$, ${}^T E$, and ${}^K E$ by ${}^U d_*^{*,*}$, ${}^T d_*^{*,*}$, and ${}^K d_*^{*,*}$, respectively, if there are risks of ambiguity. Otherwise, we simply denote the differentials by $d_*^{*,*}$.

These fiber sequences fit into the following homotopy commutative diagram:

$$(2.5) \quad \begin{array}{ccccccc} K : & & K(\mathbb{Z}, 2) & \longrightarrow & * & \longrightarrow & K(\mathbb{Z}, 3) \\ & & \downarrow B\varphi & & \downarrow & & \downarrow = \\ T : & & BT^n & \longrightarrow & BPT^n & \longrightarrow & K(\mathbb{Z}, 3) \\ & & \downarrow B\psi & & \downarrow B\psi' & & \downarrow = \\ U : & & BU_n & \longrightarrow & BPU_n & \longrightarrow & K(\mathbb{Z}, 3) \end{array}$$

Here, the map $B\varphi : K(\mathbb{Z}, 2) \simeq BS^1 \rightarrow BT^n$ is the de-looping of the diagonal map $S^1 \rightarrow T^n$. The induced homomorphism between cohomology rings is as follows:

$$B\varphi^* : H^*(BT^n) = \mathbb{Z}[v_1, v_2, \dots, v_n] \rightarrow H^*(BS^1) = \mathbb{Z}[v], \quad v_i \mapsto v.$$

The map $B\psi : BT^n \rightarrow BU_n$ induces the injective ring homomorphism

$$B\psi^* : H^*(BU_n) = \mathbb{Z}[c_1, \dots, c_n] \rightarrow H^*(BT^n) = \mathbb{Z}[v_1, \dots, v_n], \\ c_i \mapsto \sigma_i(v_1, \dots, v_n),$$

where $\sigma_j(t_1, t_2, \dots, t_n)$ be the j th elementary symmetric polynomial in variables t_1, t_2, \dots, t_n :

$$(2.6) \quad \begin{aligned} \sigma_0(t_1, t_2, \dots, t_n) &= 1, \\ \sigma_1(t_1, t_2, \dots, t_n) &= t_1 + t_2 + \dots + t_n, \\ \sigma_2(t_1, t_2, \dots, t_n) &= \sum_{i < j} t_i t_j, \\ &\vdots \\ \sigma_p(t_1, t_2, \dots, t_n) &= t_1 t_2 \cdots t_n. \end{aligned}$$

We will use the associated maps of spectral sequences to compute the differentials in $^U E$. This is possible because we have a good understanding of the corresponding differentials in $^T E$ and $^K E$. In particular, we have the following results.

Proposition 2.4 ([12], Corollary 2.16). *The higher differentials of $^K E_*^{*,*}$ satisfy*

$$\begin{aligned} d_3(v) &= x_1, \\ d_{2p-1}(x_1 v^{lp^e-1}) &= v^{lp^e-1-(p-1)} y_{p,0}, \quad e > 0, \quad \gcd(l, p) = 1, \\ d_r(x_1) &= d_r(y_{p,0}) = 0, \quad \text{for all } r, \end{aligned}$$

and the Leibniz rule.

Remark 2.5. Proposition 2.4 is a special case of Corollary 2.16, [12]. Here, we take the opportunity to correct a typo in the original Corollary 2.16, [12], where the condition $k \geq e$ should be replaced by $e > k$.

Proposition 2.6 ([12], Proposition 3.2). *The differential $^T d_r^{*,*}$, is partially determined as follows:*

$$(2.7) \quad ^T d_r^{*,2t}(v_i^t \xi) = (B\pi_i)^*(^K d_r^{*,2t}(v^t \xi)),$$

where $\xi \in ^T E_r^{*,0}$, a quotient group of $H^*(K(\mathbb{Z}, 3))$, and $\pi_i : T^n \rightarrow S^1$ is the projection of the i th diagonal entry. In plain words, $^T d_r^{*,2t}(v_i^t \xi)$ is simply $^K d_r^{*,2t}(v^t \xi)$ with v replaced by v_i .

Remark 2.7. Here we correct another typo in the original Proposition 3.2 in [12], in which “ $\xi \in {}^T E_r^{0,*}$ ” should be replaced by “ $\xi \in {}^T E_r^{*,0}$ ”.

Proposition 2.8 ([12], Proposition 3.3). (1) *The differential ${}^T d_3^{0,t}$ is given by the “formal divergence”*

$$\nabla = \sum_{i=1}^n (\partial/\partial v_i) : H^t(BT^n; R) \rightarrow H^{t-2}(BT^n; R),$$

in such a way that ${}^T d_3^{0,} = \nabla(-) \cdot x_1$. For any ground ring $R = \mathbb{Z}$ or \mathbb{Z}/m for any integer m .*

(2) *The spectral sequence degenerates at ${}^T E_4^{0,*}$. Indeed, we have ${}^T E_\infty^{0,*} = {}^T E_4^{0,*} = \text{Ker } {}^T d_3^{0,*} = \mathbb{Z}[v_1 - v_n, \dots, v_{n-1} - v_n]$.*

Corollary 2.9 ([12], Corollary 3.4).

$${}^U d_3^{0,*}(c_k) = \nabla(c_k)x_1 = (n - k + 1)c_{k-1}x_1.$$

Computations in the spectral sequence ${}^U E$. In order to study

$${}_p H^*(BPU_n) \cong {}_p [H^*(BPU_n)_{(p)}],$$

it suffices to look at the p -localized spectral sequence, where the E_2 page becomes

$$(2.8) \quad ({}^U E_2^{s,t})_{(p)} = H^s(K(\mathbb{Z}, 3))_{(p)} \otimes H^t(BU_n) = H^s(K(\mathbb{Z}, 3)) \otimes H^t(BU_n)_{(p)}.$$

By abuse of notation, for the rest of this paper, we let ${}^U E$, ${}^T E$ and ${}^K E$ denote the corresponding p -localized Serre spectral sequences.

By Proposition 2.2 and (2.3), in the range $s \leq 2p + 5$, the only cases in which ${}^U E_2^{s,t}$ could be nonzero are when $s = 0, 3, 2p + 2, 2p + 5$ and t is even.

To simplify the notations, we let

$$M^0 = {}^U E_2^{0,2p+2}, \quad M^1 = {}^U E_2^{3,2p}, \quad M^2 = {}^U E_2^{2p+2,2}, \quad M^3 = {}^U E_2^{2p+5,0}.$$

Inspection of degrees shows that ${}^U E_*^{3,2p}$ can receive only the d_3 differential and support the d_{2p-1} differential. Similarly, ${}^U E_*^{2p+2,2}$ can receive only the d_{2p-1} differential and support the d_3 differential. In addition, all d_2 's are trivial and therefore we have ${}^U E_2^{*,*} = {}^U E_3^{*,*}$.

We let δ^0 be the map

$$\delta^0 : M^0 = {}^U E_3^{0,2p+2} \xrightarrow{d_3} {}^U E_3^{3,2p} = M^1.$$

We let δ^1 be the composition

$$\delta^1 : M^1 = {}^U E_3^{3,2p} \rightarrow {}^U E_3^{3,2p} / \text{Im } d_3 = {}^U E_{2p-1}^{3,2p} \xrightarrow{d_{2p-1}} {}^U E_{2p-1}^{2p+2,2} = \text{Ker } d_3 \subset M^2.$$

We let δ^2 be the map

$$\delta^2 : M^2 = {}^U E_3^{2p+2,2} \xrightarrow{d_3} {}^U E_3^{2p+5,0} = M^3.$$

One immediately sees that

$$M^0 \xrightarrow{\delta^0} M^1 \xrightarrow{\delta^1} M^2 \xrightarrow{\delta^2} M^3$$

is a chain complex of $\mathbb{Z}_{(p)}$ -modules, which we denote by \mathcal{M} . We will show later that Theorem 1 is a consequence of the following

Proposition 2.10. *Let $p \geq 3$ be a prime number such that $p \mid n$. The chain complex \mathcal{M} defined above is exact.*

Proof of Theorem 1 assuming Proposition 2.10. Let $n = p^r m$. For $r = 0$, the theorem follows from Proposition 1.1. In the rest of the proof we assume $r > 0$. First, we prove

$${}_p H^{2p+2}(BPU_n) \cong \mathbb{Z}/p.$$

By Proposition 2.3, $y_{p,0} \in {}^U E_2^{2p+2,0}$ survives to a nonzero element in $H^{2p+2}(BPU_n)$ of order p . Therefore, we have

$${}^U E_\infty^{2p+2,0} = {}^U E_2^{2p+2,0} \cong \mathbb{Z}/p.$$

Since the only nontrivial entries in ${}^U E_2^{*,*}$ of total degree $2p+2$ are ${}^U E_2^{2p+2,0}$ and ${}^U E_2^{0,2p+2}$, we have a short exact sequence of $\mathbb{Z}_{(p)}$ -modules

$$0 \rightarrow {}^U E_\infty^{2p+2,0} \rightarrow H^{2p+2}(BPU_n)_{(p)} \rightarrow {}^U E_\infty^{0,2p+2} \rightarrow 0.$$

Since ${}^U E_\infty^{0,2p+2} \subset {}^U E_2^{0,2p+2}$ is a free $\mathbb{Z}_{(p)}$ -module, the above short exact sequence splits and we have

$$H^{2p+2}(BPU_n)_{(p)} \cong {}^U E_\infty^{2p+2,0} \oplus {}^U E_\infty^{0,2p+2},$$

from which we deduce

$${}_p H^{2p+2}(BPU_n) \cong {}^U E_\infty^{2p+2,0} \cong \mathbb{Z}/p.$$

Since the row $E_\infty^{*,0}$ is the image of χ^* , the above implies

$$(2.9) \quad {}_p H^{2p+2}(BPU_n) = \chi^*(H^{2p+2}(K(\mathbb{Z}, 3))).$$

From (2.9) and Proposition 2.2, it follows that ${}_p H^{2p+2}(BPU_n)$ is generated by $\delta P^1(\bar{x}_1)$.

Next, we prove

$${}_p H^{2p+3}(BPU_n) = H^{2p+3}(BPU_n)_{(p)} = 0.$$

The exactness of \mathcal{M} at M^1 implies ${}^U E_\infty^{3,2p} = 0$. On the other hand, ${}^U E_2^{3,2p}$ is the only nontrivial entry in ${}^U E_2^{*,*}$ of total degree $2p+3$. Hence, we have

$${}_p H^{2p+3}(BPU_n) \subset H^{2p+3}(BPU_n)_{(p)} = {}^U E_\infty^{3,2p} = 0.$$

Finally, we prove

$${}_p H^{2p+4}(BPU_n) = 0.$$

The exactness of \mathcal{M} at M^2 implies ${}^U E_\infty^{2p+2,2} = 0$. Since ${}^U E_2^{0,2p+4}$ and ${}^U E_2^{2p+2,2}$ are the only nontrivial entries in ${}^U E_2^{*,*}$ of total degree $2p+4$, we have

$$H^{2p+4}(BPU_n)_{(p)} \cong {}^U E_\infty^{0,2p+4},$$

which is torsion-free. In particular, we have ${}_p H^{2p+4}(BPU_n) = 0$. □

The proof of Proposition 2.10 occupies Section 3.

3. THE PROOF OF PROPOSITION 2.10

From (2.8), we can write out the $\mathbb{Z}_{(p)}$ -modules M^0, M^1, M^2, M^3 more explicitly:

$$M^0 = H^0(K(\mathbb{Z}, 3)) \otimes H^{2p+2}(BU_n)_{(p)} \cong H^{2p+2}(BU_n)_{(p)}$$

is the free $\mathbb{Z}_{(p)}$ -module generated by monomials in c_1, \dots, c_{p+1} in dimension $2p+2$, and

$$M^1 = H^3(K(\mathbb{Z}, 3)) \otimes H^{2p}(BU_n)_{(p)} \cong H^{2p}(BU_n)_{(p)}$$

is the free $\mathbb{Z}_{(p)}$ -module generated by elements of the form cx_1 where c is a monomial in c_1, \dots, c_p in dimension $2p$. Furthermore, we have

$$M^2 = H^{2p+2}(K(\mathbb{Z}, 3)) \otimes H^2(BU_n)_{(p)} = \mathbb{Z}_{(p)}\{c_1 y_{p,0}\}/p \cong \mathbb{Z}/p$$

and

$$M^3 = H^{2p+5}(K(\mathbb{Z}, 3)) \otimes H^0(BU_n)_{(p)} = \mathbb{Z}_{(p)}\{x_1 y_{p,0}\}/p \cong \mathbb{Z}/p.$$

The exactness of \mathcal{M} at M^2 .

Lemma 3.1. *In the spectral sequence ${}^T E$, we have*

$$(3.1) \quad \begin{cases} v_n^k x_1 \in \text{Im } {}^T d_3, & 0 \leq k \leq p-2 \text{ or } k = p, \\ {}^T d_{2p-1}^{3,*}(v_n^{p-1} x_1) = y_{p,0}. \end{cases}$$

Proof. When $p \nmid k+1$, the first formula in Proposition 2.4 together with Proposition 2.6 imply that

$$v_n^k x_1 = \frac{1}{k+1} {}^T d_3(v_n^{k+1})$$

is in the image of ${}^T d_3$. This completes the proof for the case $0 \leq k \leq p-2$ or $k = p$.

The remaining case is proved by applying the second formula in Proposition 2.4, taking $e = l = 1$, and then Proposition 2.6. \square

Lemma 3.2. *The map $\delta^1 : M^1 \rightarrow M^2 \cong \mathbb{Z}/p$ is surjective.*

Proof. Recall the morphism of fiber sequences Ψ introduced in (2.5), and the induced morphism $\Psi^* : {}^U E \rightarrow {}^T E$ of spectral sequences.

For $1 \leq i \leq n$, let $v'_i = v_i - v_n$. It follows from (2) of Proposition 2.8 that the v'_i 's are permanent cycles. To determine the value of δ^1 at $c_p x_1 \in M^1$, we have

$$(3.2) \quad \begin{aligned} & \Psi^* \delta^1(c_p x_1) \\ &= \Psi^* {}^U d_{2p-1}^{3,2p}(c_p x_1) = {}^T d_{2p-1}^{3,2p} \Psi^*(c_p x_1) \\ &= {}^T d_{2p-1}^{3,2p} \left(\sum_{n \geq i_1 > i_2 > \dots > i_p \geq 1} v_{i_1} v_{i_2} \dots v_{i_p} x_1 \right) \\ &= {}^T d_{2p-1}^{3,2p} \left(\sum_{n \geq i_1 > i_2 > \dots > i_p \geq 1} (v'_{i_1} + v_n)(v'_{i_2} + v_n) \dots (v'_{i_p} + v_n) x_1 \right) \\ &= {}^T d_{2p-1}^{3,2p} \left(\sum_{n \geq i_1 > i_2 > \dots > i_p \geq 1} \sum_{j=0}^p \sigma_j(v'_{i_1}, \dots, v'_{i_p}) v_n^{p-j} x_1 \right). \end{aligned}$$

where $\Psi^* : {}^U E \rightarrow {}^T E$ is the morphism of spectral sequences induced by the inclusions of maximal tori $T^n \rightarrow U_n$ and $PT^n \rightarrow PU_n$, as introduced in (2.5), and σ_i the elementary symmetric polynomials in p variables, as in (2.6).

By Lemma 3.1, we simplify (3.2) and obtain

$$(3.3) \quad \Psi^* \delta^1(c_p x_1) = {}^T d_{2p-1} \left(\sum_{n \geq i_1 > i_2 > \dots > i_p \geq 1} \sigma_1(v'_{i_1}, \dots, v'_{i_p}) v_n^{p-1} x_1 \right).$$

To proceed, we evaluate the expression

$$\sum_{n \geq i_1 > i_2 > \dots > i_p \geq 1} \sigma_1(t_{i_1}, \dots, t_{i_p})$$

for variables t_i , $1 \leq i \leq n$. Since it is multi-linear and symmetric in the variables t_1, \dots, t_n , we have

$$\sum_{n \geq i_1 > i_2 > \dots > i_p \geq 1} \sigma_1(t_{i_1}, \dots, t_{i_p}) = \lambda \sum_{i=1}^n t_i$$

for some $\lambda \in \mathbb{Z}$. Taking the substitution $t_1 = \dots = t_n = 1$ and comparing both sides of the above, we obtain

$$\lambda = \frac{p}{n} \binom{n}{p} = \binom{n-1}{p-1} \not\equiv 0 \pmod{p}$$

and

$$(3.4) \quad \sum_{n \geq i_1 > i_2 > \dots > i_p \geq 1} \sigma_1(t_{i_1}, \dots, t_{i_p}) = \binom{n-1}{p-1} \sum_{i=1}^n t_i.$$

Consider the following commutative diagram:

$$\begin{array}{ccc} M^1 = U E_2^{3,2p} & \xrightarrow{\Psi^*} & T E_2^{3,2p} \\ \downarrow & & \downarrow \\ U E_{2p-1}^{3,2p} & \xrightarrow{\Psi^*} & T E_{2p-1}^{3,2p} \\ \downarrow U d_{2p-1} & & \downarrow T d_{2p-1} \\ U E_{2p-1}^{2p+2,2} & \xrightarrow{\Psi^*} & T E_{2p-1}^{2p+2,2} \\ \downarrow & & \downarrow \\ M^2 = U E_2^{2p+2,2} & \xrightarrow{\Psi^*} & T E_2^{2p+2,2} \end{array}$$

where the composition of the left vertical maps is δ^1 and we resume the computation of $\Psi^* \delta^1(c_p x_1)$ started in (3.3):

$$\begin{aligned} & \Psi^* \delta^1(c_p x_1) \\ &= {}^T d_{2p-1} \left(\binom{n-1}{p-1} \sum_{i=1}^n v'_i v_n^{p-1} x_1 \right) \quad (\text{by (3.4)}) \\ (3.5) \quad &= \binom{n-1}{p-1} \sum_{i=1}^n v'_i y_{p,0} \quad (\text{since } v'_i \text{'s are permanent cocycles}) \\ &= \binom{n-1}{p-1} \sum_{i=1}^n v_i y_{p,0} \quad (\text{since } y_{p,0} \text{ is } p\text{-torsion}) \\ &= \Psi^* \left(\binom{n-1}{p-1} c_1 y_{p,0} \right). \end{aligned}$$

By the injectivity of

$$\Psi^* : M^2 = U E_2^{2p+2,2} \rightarrow T E_2^{2p+2,2}$$

together with (3.5), we have

$$\delta^1(c_p x_1) = \binom{n-1}{p-1} c_1 y_{p,0} \neq 0$$

and we conclude. \square

Lemma 3.3. *The chain complex \mathcal{M} is exact at M^2 .*

Proof. By Lemma 3.2, and the fact that \mathcal{M} is a chain complex, we have $\delta^2 = 0$ and the lemma follows.

Alternatively, one may compute $\delta^2 = d_3^{2p+2,2}$ directly with Corollary 2.9 and obtain the same result. \square

The exactness of \mathcal{M} at M^1 . Recall that the $\mathbb{Z}_{(p)}$ -module M^1 is freely generated by elements of the form $c x_1$ for

$$c \in S' := \{c_1^{i_1} c_2^{i_2} \cdots c_p^{i_p} \mid i_k \geq 0, \sum_k k i_k = p\}.$$

Indeed, S' is simply the set of monomials in c_1, c_2, \dots, c_n in $H^{2p}(BU_n)$. We define a total ordering \mathfrak{D} on monomials in c_1, c_2, \dots, c_n as follows. We assert

$$c_1^{i_1} c_2^{i_2} \cdots c_p^{i_p} > c_1^{j_1} c_2^{j_2} \cdots c_p^{j_p}$$

if and only if

- (1) there is at least one k such that $i_k \neq j_k$, and
- (2) for the smallest such k , we have $i_k > j_k$.

Let $S := S' - \{c_p\}$. Then \mathfrak{D} defines total orderings on S , S' and $S'x_1$ as well. To compare $c x_1, c' x_1 \in S'x_1$, we assert $c x_1 > c' x_1$ if and only if $c > c'$.

Let L be the $\mathbb{Z}_{(p)}$ -submodule of $H^{2p}(BU_n)_{(p)}$ spanned by S . We define a $\mathbb{Z}_{(p)}$ -linear map

$$\tau : L \rightarrow M^0 = H^{2p+2}(BU_n)_{(p)}$$

as follows. Each element in S is of the form $c_1^{i_1} c_2^{i_2} \cdots c_k^{i_k}$ such that $k < p$ and $i_k > 0$, and we define

$$\tau(c_1^{i_1} c_2^{i_2} \cdots c_k^{i_k}) := (c_1^{i_1} c_2^{i_2} \cdots c_{k-1}^{i_{k-1}})(c_k^{i_k-1} c_{k+1}).$$

Lemma 3.4. *Let $\bar{\tau} : L/pL \rightarrow M^0/pM^0$ and $\bar{\delta}^0 : M^0/pM^0 \rightarrow M^1/pM^1$ denote the mod p reductions of τ and δ^0 , respectively. Then the image of the composition*

$$L/pL \xrightarrow{\bar{\tau}} M^0/pM^0 \xrightarrow{\bar{\delta}^0} M^1/pM^1$$

is Lx_1/pLx_1 . In particular, we have

$$(3.6) \quad \text{Im } \delta^0 \tau \subset W := Lx_1 + (pc_p x_1) \subset M^1.$$

Proof. Consider the $\mathbb{Z}_{(p)}$ -basis $S, S'x_1$ for L and M^1 , respectively, both in the descending order with respect to the ordering \mathfrak{D} . Notice that $c_p x_1$ is the smallest element in S' . We study the $(N+1) \times N$ matrix A of the map

$$\delta^0 \tau : L \rightarrow M^1$$

with respect to these basis, where N is the cardinality of S .

Consider an arbitrary element

$$c := c_1^{i_1} \cdots c_k^{i_k} \in S$$

with $k < p$ and $i_k > 0$. By Corollary 2.9 and the Leibniz's formula, we have

$$\begin{aligned} \delta^0 \tau(c) &= \delta^0(c_1^{i_1} \cdots c_{k-1}^{i_{k-1}} c_k^{i_k-1} c_{k+1}) \\ &= \begin{cases} (n-k)cx_1 + ni_1 c_1^{i_1-1} c_2^{i_2} \cdots c_k^{i_k-1} c_{k+1} x_1 + (\text{higher order terms}), & i_1 > 0, \\ (n-k)cx_1 + (\text{higher order terms}), & i_1 = 0. \end{cases} \end{aligned}$$

In both cases, we have

$$\delta^0 \tau(c) \equiv (n-k)cx_1 + (\text{higher order terms}) \pmod{p}.$$

Therefore, the matrix A satisfies

$$A \equiv \begin{pmatrix} \lambda_1 & * & \cdots & * \\ 0 & \lambda_2 & * & \vdots \\ 0 & 0 & \ddots & * \\ 0 & \cdots & 0 & \lambda_N \\ 0 & 0 & \cdots & 0 \end{pmatrix} \pmod{p},$$

where the λ_i 's are of the form $n-k$ for $k < p$, which are invertible in $\mathbb{Z}_{(p)}$, and we have verified that the image of the composition

$$L/pL \xrightarrow{\bar{\tau}} M^0/pM^0 \xrightarrow{\bar{\delta}^0} M^1/pM^1$$

is Lx_1/pLx_1 . The equation (3.6) follows from the above and the fact

$$M^1 = Lx_1 + (c_p x_1).$$

□

Lemma 3.5. *Consider the $\mathbb{Z}_{(p)}$ -submodule $V = \tau(L) + (c_1 c_p - c_{p+1})$ of M^0 . We have $\delta^0(V) \subset W$ where*

$$W := Lx_1 + (pc_p x_1) \subset M^1$$

is the $\mathbb{Z}_{(p)}$ -submodule of M^1 defined in Lemma 3.4.

Proof. By Lemma 3.4 we have $\delta^0(\tau(L)) \subset W$. On the other hand, we have

$$(3.7) \quad \delta^0(c_1 c_p - c_{p+1}) = (n-p+1)c_1 c_{p-1} x_1 + pc_p x_1 \in W,$$

and we conclude. □

Lemma 3.6. *The chain complex \mathcal{M} is exact at M^1 .*

Proof. By Lemma 3.5, the restriction of δ^0 to V has image in W . Therefore, we write $\delta_V^0 := \delta^0|_V : V \rightarrow W$ and consider its mod p reduction

$$\bar{\delta}_V^0 : V/pV \rightarrow W/pW = Lx_1/pLx_1 + (pc_p x_1)/(p^2 c_p x_1).$$

By Lemma 3.4, we have $Lx_1/pLx_1 \subset \text{Im } \bar{\delta}_V^0$.

By $Lx_1/pLx_1 \subset \text{Im } \bar{\delta}_V^0$ and (3.7), we have $[pc_p x_1] \in \text{Im } \bar{\delta}_V^0$, where $[pc_p x_1]$ is the class in W/pW represented by $pc_p x_1$. Therefore, $\bar{\delta}_V^0 : V/pV \rightarrow W/pW$ is surjective. By Nakayama's lemma in commutative algebra (Theorem 2.2, Chapter 1, [17]), $\delta_V^0 : V \rightarrow W$ is surjective.

Therefore, we have

$$(3.8) \quad \text{Im } \delta^0 \supset \text{Im } \delta_V^0 = W = Lx_1 + (pc_p x_1).$$

On the other hand, we have $\text{Ker } \delta^1 \supset \text{Im } \delta^0$, and therefore $\text{Ker } \delta^1 \supset W$. Now, by Lemma 3.2, we have

$$\mathbb{Z}/p \cong M^1/(L + (pc_p x_1)) = M^1/W \rightarrow M^1/\text{Ker } \delta^1 \cong \mathbb{Z}/p,$$

where the arrow is the tautological quotient map, which is surjective. Therefore, the above composition is a bijection. It follows that we have

$$(3.9) \quad W = \text{Ker } \delta^1 \supset \text{Im } \delta^0,$$

and the lemma follows from (3.8) and (3.9). \square

Lemma 3.6 and Lemma 3.3 complete the proof of Proposition 2.10.

REFERENCES

- [1] Benjamin Antieau and Ben Williams. The period-index problem for twisted topological K-theory. *Geometry & Topology*, 18(2):1115–1148, 2014.
- [2] Benjamin Antieau and Ben Williams. The topological period–index problem over 6-complexes. *Journal of Topology*, 7(3):617–640, 2014.
- [3] Henri Cartan and Jean-Pierre Serre. Séminaire Henri Cartan vol. 7. pages 283–288, 1954–1955.
- [4] Clay Cordova, Daniel Freed, Ho Tat Lam, and Nathan Seiberg. Anomalies in the space of coupling constants and their dynamical applications II. *SciPost Physics Proceedings*, 8(1), 2020.
- [5] Diarmuid Crowley and Xing Gu. On $H^*(BPU_n; \mathbb{Z})$ and Weyl group invariants. *arXiv preprint arXiv:2103.03523*, 2021.
- [6] Haibao Duan. The cohomology and K-theory of the projective unitary groups $PU(n)$. *arXiv preprint arXiv:1710.09222*, 2017.
- [7] Tudor Ganea. Induced fibrations and cofibrations. *Transactions of the American Mathematical Society*, 127(3):442–459, 1967.
- [8] Iñaki García-Etxebarria and Miguel Montero. Dai-Freed anomalies in particle physics. *Journal of High Energy Physics*, 2019(8):3, 2019.
- [9] Xing Gu. The topological period–index problem over 8-complexes, I. *Journal of Topology*, 12(4):1368–1395, 2019.
- [10] Xing Gu. An almost polynomial subring of the Chow ring and cohomology of $BPGL_n$. *arXiv preprint arXiv:2012.00199*, 2020.
- [11] Xing Gu. The topological period-index problem over 8-complexes, II. *Proceedings of the American Mathematical Society*, 148:4541–4545, 2020.
- [12] Xing Gu. On the cohomology of the classifying spaces of projective unitary groups. *Journal of Topology and Analysis*, 13(02):535–573, 2021.
- [13] Xing Gu. Some torsion classes in the Chow ring and cohomology of $BPGL_n$. *Journal of the London Mathematical Society*, 103(1):127–160, 2021.
- [14] Masaki Kameko and Nobuaki Yagita. The Brown–Peterson cohomology of the classifying spaces of the projective unitary groups $PU(p)$ and exceptional Lie groups. *Transactions of the American Mathematical Society*, 360(5):2265–2284, 2008.
- [15] Akira Kono and Mamoru Mimura. On the cohomology of the classifying spaces of $PSU(4n+2)$ and $PO(4n+2)$. *Publications of the Research Institute for Mathematical Sciences*, 10(3):691–720, 1975.
- [16] Akira Kono and Nobuaki Yagita. Brown–Peterson and ordinary cohomology theories of classifying spaces for compact Lie groups. *Transactions of the American Mathematical Society*, 339(2):781–798, 1993.
- [17] Hideyuki Matsumura. *Commutative ring theory*. Cambridge university press, 1989.
- [18] Hirotaka Tamanoi. Q -subalgebras, Milnor basis, and cohomology of Eilenberg–MacLane spaces. *Journal of Pure and Applied Algebra*, 137(2):153–198, 1999.
- [19] Hiroshi Toda et al. Cohomology of classifying spaces. In *Homotopy theory and related topics*, pages 75–108. Mathematical Society of Japan, 1987.
- [20] Burt Totaro. The Chow ring of a classifying space. In *Proceedings of symposia in pure mathematics*, volume 67, pages 249–284. Providence, RI; American Mathematical Society; 1998, 1999.

- [21] Aleš Vavpetič and Antonio Viruel. On the mod p cohomology of $BPU(p)$. *Transactions of the American Mathematical Society*, pages 4517–4532, 2005.
- [22] Gabriele Vezzosi. On the Chow ring of the classifying stack of $PGL_{3,\mathbb{C}}$. *Journal für die reine und angewandte Mathematik*, 2000(523):1–54, 2000.
- [23] Angelo Vistoli. On the cohomology and the Chow ring of the classifying space of PGL_p . *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2007(610):181–227, 2007.

CENTER FOR TOPOLOGY AND GEOMETRY BASED TECHNOLOGY, SCHOOL OF MATHEMATICAL SCIENCES, HEBEI NORMAL UNIVERSITY

Email address: `gux2006@outlook.com`

DEPARTMENT OF MATHEMATICS, NANKAI UNIVERSITY

Email address: `zhang.4841@osu.edu`

DEPARTMENT OF MATHEMATICS, NANKAI UNIVERSITY

Email address: `15829207515@163.com`

DEPARTMENT OF MATHEMATICS, YANBIAN UNIVERSITY

Email address: `lnzhong@ybu.edu.cn`