

A PROOF OF A CONJECTURE ON RAMSEY NUMBERS $B(2, 2, 3)$

YASER ROWSHAN¹ AND MOSTAFA GHOLAMI¹

ABSTRACT. The bipartite Ramsey number $B(n_1, n_2, \dots, n_t)$ is the least positive integer b such that, any coloring of the edges of $K_{b,b}$ with t colors will result in a monochromatic copy of K_{n_i, n_i} in the i -th color, for some i , $1 \leq i \leq t$. In this paper we obtain the exact values of bipartite Ramsey numbers $B(2, 2, 3)$. In particular, we prove the conjecture of Radziszowski et al. about $B(2, 2, 3)$ which was introduced in 2015. In fact we prove that $B(2, 2, 3) = 17$.

1. INTRODUCTION

The bipartite Ramsey number $B(n_1, n_2, \dots, n_t)$ is the least positive integer b such that, any coloring of the edges of $K_{b,b}$ with t colors will result in a monochromatic copy of K_{n_i, n_i} in the i -th color, for some i , $1 \leq i \leq t$. The existence of such a positive integer is guaranteed by a result of Erdős and Rado [6].

The Zarankiewicz number $z(K_{m,n}, t)$ is defined as the maximum number of edges in any subgraph G of the complete bipartite graph $K_{m,n}$, such that G does not contain $K_{t,t}$ as a subgraph. Zarankiewicz numbers and related extremal graphs have been studied by numerous authors, including Kóvari [12], Reiman [14] and Goddard, Henning, and Oellermann in [9].

The study of bipartite Ramsey numbers was initiated by Beineke and Schwenk in 1976 [1], and continued by others, in particular Exoo [7], Hattingh and Henning [11]. The following exact values have been established. $B(2, 5) = 17$ [4], $B(2, 2, 2, 2) = 19$ [5], $B(2, 2, 2) = 11$ [7]. In the smallest open case for 5 colors, it is known that $26 \leq B(2, 2, 2, 2, 2) \leq 28$ [5]. One can refer to [3, 5, 8, 10, 12, 13, 15] and its references for further studies. Radziszowski et al. in [5] showed that $17 \leq B(2, 2, 3) \leq 18$ and in the same source has made the following conjecture:

Conjecture 1. ([5]) $B(2, 2, 3) = 17$.

We intend to compute the exact values of the multicolor bipartite Ramsey numbers $B(2, 2, 3)$. Actually, we prove the following results:

Theorem 1.1. $B(2, 2, 3) = 17$.

In this paper, we only concerned with undirected, simple and finite graphs. We follow [2] for terminology and notations not defined here. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. The degree of a vertex $v \in V(G)$ is denoted by $\deg_G(v)$, or simply by $\deg(v)$. The neighborhood $N_G(v)$ of a vertex v is the set of all vertices of G adjacent to v and satisfies $|N_G(v)| = \deg_G(v)$. The minimum and maximum degrees of vertices of G are denoted by $\delta(G)$ and $\Delta(G)$, respectively. Also, the complete bipartite graph with bipartition (X, Y) , where $|X| = m$ and $|Y| = n$ is denoted by $K_{m,n}$. We use $[X, Y]$ to denote the set of edges between a bipartition (X, Y) of G . The degree of a vertex $v \in X \subseteq V(G)$ is denoted by $D_G(X) = (d_1, d_2, \dots, d_n)$ where $|X| = n$. The complement of a graph G , denoted by \overline{G} , is the graph with the same vertices as G and contains those edges which are not in G . G is n -colorable to (G_1, G_2, \dots, G_n) if there exist a n -edge decomposition of G , say (H_1, H_2, \dots, H_n) where $G_i \not\subseteq H_i$ for each $i = 1, 2, \dots, n$.

2010 *Mathematics Subject Classification.* 05C35, 05C55.

Key words and phrases. Ramsey numbers, Bipartite Ramsey numbers, Zarankiewicz number.

2. PROOF OF THE MAIN THEOREM

To prove our main results, namely Theorem 1.1, we need to establish some preliminary results. We start with the following proposition:

Proposition 2.1. (*[3, 4]*) *The following results about Zarankiewicz number are true:*

- $z(K_{17,17}, 2) = 74.$
- $z(K_{16,17}, 2) = 71.$
- $z(K_{17,17}, 3) = 141.$
- $z(K_{16,17}, 3) = 133.$
- $z(K_{13,17}, 3) = 110.$
- $z(K_{12,17}, 3) = 103.$
- $z(K_{11,17}, 3) = 96.$

Proof. By using the bounds in Table 3 and Table 4 of [5] and Table C.3 of [3] the proposition holds. ■

Theorem 2.2. *[5]* $17 \leq B(2, 2, 3) \leq 18.$

Proof. The lower bound witness is found in Table 2 of [5]. The upper bound is implied by using the bounds in Table 3 and Table 4 of [5]. We know that $z(K_{18,18}, 2) = 81$, $z(K_{18,18}, 3) \leq 156$, and $2 \times 81 + 156 = 318 < 324 = |E(K_{18,18})|$. ■

Suppose that (G^r, G^b, G^g) be 3-edges coloring of $K_{17,17}$ where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$, in the following theorem, we specify some properties of the subgraph with color g . The properties regarding $\Delta(G^g)$, $\delta(G^g)$, $E(G^g)$ and degree sequence of vertices X, Y in induced graph with color g .

Theorem 2.3. *Assume that (G^r, G^b, G^g) be 3-edges coloring of $K_{17,17}$ where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$. Hence we have:*

- (a) $|E(G^g)| = 141,$
- (b) $\Delta(G^g) = 9$ and $\delta(G^g) = 8,$
- (c) $D_{G^g}(X) = D_{G^g}(Y) = (9, 9, 9, 9, 9, 8, 8, \dots, 8).$

Proof. Assume that $X = \{x_1, x_2, \dots, x_{17}\}, Y = \{y_1, y_2, \dots, y_{17}\}$ be the partition set of $K = K_{17,17}$ and (G^r, G^b, G^g) be 3-edges coloring of K where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$. Since $|E(K)| = 289$, if $|E(G^g)| \leq 140$ then $|E(\overline{G^g})| \geq 149$ that is, either $|E(G^r)| \geq 75$ or $|E(G^b)| \geq 75$. In any case by Proposition 2.1, either $K_{2,2} \subseteq G^r$ or $K_{2,2} \subseteq G^b$, a contradiction. Hence assume that $|E(G^g)| \geq 141$. If $|E(G^g)| \geq 142$ then by Proposition 2.1, $K_{3,3} \subseteq G^g$, a contradiction again that is, we have $|E(G^g)| = 141$ and the part (a) is true.

To prove the part (b), by part (a) since $|E(G^g)| = 141$, we can check that $\Delta(G^g) \geq 9$. Assume that there exist a vertex of $V(K)$ say x such that, $|N_{G^g}(x)| \geq 10$ that is, $\Delta(G^g) \geq 10$. Consider x , set $G_1^g = G^g \setminus \{x\}$, hence by part (a) we have $|E(G_1^g)| \leq 141 - 10 = 131$ therefore, since $|E(K_{16,17})| = 272$ we have $|E(\overline{G_1^g})| \geq 141$ that is, either $|E(G_1^r)| \geq 71$ or $|E(G_1^b)| \geq 71$. In any case by Proposition 2.1 either $K_{2,2} \subseteq G_1^r \subseteq G^r$ or $K_{2,2} \subseteq G_1^b \subseteq G^b$, a contradiction. So, we have $\Delta(G^g) = 9$. To prove $\delta(G^g) = 8$, assume that $M = \{x \in X, |N_{G^g}(x)| = 9\}$ and $N = \{x \in X, |N_{G^g}(x)| = 8\}$, by part (a) one can say that $|M| \geq 5$, if $|M| = 6$ then, we have $\delta(G^g) \leq 7$ that is, there exist a vertex of X say x such that, $|N_{G^g}(x)| \leq 7$ therefore, $|N| \leq 10$. If $|N| = 10$ then, we have $|E(G^g[M \cup N, Y])| = 134$, so by Proposition 2.1 we have $K_{3,3} \subseteq G^g$, a contradiction. Now assume that $|N| \leq 9$ therefore, $|E(G^g)| \leq (6 \times 9) + (9 \times 8) + (2 \times 7) = 140$, a contradiction again. For $|M| = 7$, if $|N| \geq 6$, we have $|E(G^g[M \cup N', Y])| = 111$ where $N' \subseteq N$ and $|N'| = 6$ therefore, by Proposition 2.1 we have $K_{3,3} \subseteq G^g$, a contradiction, hence assume

that $|N| \leq 5$ therefore, we have $|E(G^g)| \leq (7 \times 9) + (5 \times 8) + (5 \times 7) = 138$, a contradiction again. For $|M| = 8$, if $|N| \geq 5$ then, $|E(G^g[M \cup N', Y])| = 112$ where $N' \subseteq N$ and $|N'| = 5$ therefore, by Proposition 2.1 we have $K_{3,3} \subseteq G^g$, a contradiction, so assume that $|N| \leq 4$ that is, $|E(G^g)| \leq (8 \times 9) + (4 \times 8) + (5 \times 7) = 139$, a contradiction again. For $|M| = 9$ if $|N| \geq 3$, then $|E(G^g[M \cup N', Y])| = 105$ where $N' \subseteq N$ and $|N'| = 3$, so by Proposition 2.1 we have $K_{3,3} \subseteq G^g$, a contradiction, so $|N| \leq 2$ that is, $|E(G^g)| \leq (9 \times 9) + (2 \times 8) + (6 \times 7) = 139$ which is a contradiction again. For $|M| = 10$, if $|N| \geq 1$ then, $|E(G^g[M \cup N', Y])| = 98$ where $N' \subseteq N$ and $|N'| = 1$ therefore, by Proposition 2.1 $K_{3,3} \subseteq G^g$, a contradiction. Thus assume that $|N| = 0$ so, $|E(G^g)| \leq (10 \times 9) + (7 \times 7) = 139$, a contradiction again. Therefore, $|M| = 5$ and $|N| = 12$ that is, $\delta(G^g) = 8$ and the part (b) is true.

Now by parts (a) and (b) it is easy to say that $D_{G^g}(X) = D_{G^g}(Y) = (9, 9, 9, 9, 9, 8, 8, \dots, 8)$ that is, the part (c) is true and the proof is complete. \blacksquare

Suppose that (G^r, G^b, G^g) be 3-edge coloring of $K_{17,17}$ where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$. In the following theorem we discuss about the maximum number of common neighbors of $G^g(x)$ and $G^g(x')$ for $x, x' \in X$.

Theorem 2.4. *Assume that (G^r, G^b, G^g) be 3-edge coloring of $K_{17,17}$ where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$. Let $|N_{G^g}(x)| = 9$ and $N_{G^g}(x) = Y_1$, the following results are true:*

- (a) *For each $x \in X \setminus \{x_1\}$, we have $|N_{G^g}(x) \cap Y_1| \leq 5$,*
- (b) *Assume that $n = \sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1|$, then $72 \leq n \leq 73$.*

Proof. Assume that $X = \{x_1, x_2, \dots, x_{17}\}$, $Y = \{y_1, y_2, \dots, y_{17}\}$ be the partition set of $K = K_{17,17}$ and (G^r, G^b, G^g) be 3-edges coloring of K where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$. Without loss of generality (W.l.g) assume that $x = x_1$ and $Y_1 = \{y_1, \dots, y_9\}$. To prove the part (a), by contrary assume that there exist a vertex of $X \setminus \{x_1\}$ say x such that, $|N_{G^g}(x) \cap Y_1| \geq 6$. W.l.g assume that $x = x_2$ and $Y_2 = \{y_1, y_2, \dots, y_6\} \subseteq N_{G^g}(x_2)$. Since $K_{3,3} \not\subseteq G^g$, for each $x \in X \setminus \{x_1, x_2\}$, we have $|N_{G^g}(x) \cap Y_2| \leq 2$ that is, $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_2| \leq 6 + 6 + (15 \times 2) \leq 42$. Now, since $|E(G^g[X, Y_2])| \leq 42$, one can check that there exist at least one vertex of Y_2 say y such that, $|N_{G^g}(y)| \leq 7$, a contradiction to part (c) of Theorem 2.3. Hence $|N_{G^g}(x) \cap Y_1| \leq 5$ for each $x \in X \setminus \{x_1\}$ that is, the part (a) is true.

To prove the part (b), if $n \leq 71$ then, by part (c) of Theorem 2.3, one can check that there exist at least one vertex of Y_1 say y such that, $|N_{G^g}(y)| \leq 7$, a contradiction. Therefore, $n \geq 72$. Assume that $n \geq 74$ and let $D_{G^g}(Y_1) = (d_1, d_2, \dots, d_9)$. Since $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1| \geq 74$, there exist at least two vertices of Y_1 say y', y'' such that, $|N_{G^g}(y')| = |N_{G^g}(y'')| = 9$. Since $n \geq 74$ and $|X \setminus \{x_1\}| = 16$, there exist at least one vertex of $X \setminus \{x_1\}$ say x' such that, $|N_{G^g}(x') \cap Y_1| = 5$. W.l.g assume that $x' = x_2$ and $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$. Now we note the following claims:

Claim 2.5. *For each $x \in X \setminus \{x_1, x_2\}$ we have $|N_{G^g}(x) \cap Y_2| = 2$ and $D_{G^g}(Y_2) = (8, 8, 8, 8, 8)$.*

Proof of the Claim. Since $K_{3,3} \not\subseteq G^g$ for each $x \in X \setminus \{x_1, x_2\}$ thus, $|N_{G^g}(x) \cap Y_2| \leq 2$ that is, $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_2| \leq 5 + 5 + (15 \times 2) \leq 40$. Now, since $|E(G^g[X, Y_2])| \leq 40$ and $|Y_2| = 5$, if there exist a vertex of X_1 say x' such that, $|N_{G^g}(x) \cap Y_2| \leq 1$ then, $|E(G^g[X, Y_2])| \leq 39$ therefore, one can check that there exist at least one vertex of Y_2 say y such that, $|N_{G^g}(y)| \leq 7$, a contradiction to part (c) of Theorem 2.3. So $|N_{G^g}(x) \cap Y_2| = 2$ and $\sum_{y \in Y_2} |N_{G^g}(y)| = 40$ therefor, by part (c) of

Theorem 2.3 $D_{G^g}(Y_2) = (8, 8, 8, 8, 8)$ and the proof of the claim is complete.

Claim 2.6. $D_{G^g}(X_1) = (5, 4, 4, \dots, 4)$ where $X_1 = X \setminus \{x_1\}$ in other word, $|N_{G^g}(x_i) \cap Y_1| = 4$ for each $i \in \{3, 4, \dots, 17\}$.

Proof of the Claim. By contrary, assume that there exist a vertex of $X \setminus \{x_1, x_2\}$ say x such that, $|N_{G^g}(x) \cap Y_1| = 5$. W.l.g assume that $x = x_3$ and $N_{G^g}(x_3) \cap Y_1 = Y_3$, now by Claim 2.5 we have $|N_{G^g}(x_3) \cap Y_2| = 2$. W.l.g assume that $Y_3 = \{y_1, y_2, y_6, y_7, y_8\}$ therefore, by Claim 2.5, $D_{G^g}(Y_3) = (8, 8, 8, 8, 8)$ that is, $|N_{G^g}(y)| = 8$ for each $y \in Y_1 \setminus \{y_9\}$. Since $\Delta(G^g) = 9$, we can check that $n = \sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1| = \sum_{i=1}^{i=9} |N_{G^g}(y_i)| \leq (8 \times 8) + 9 = 73$, a contradiction. So $D_{G^g}(X_1) = (5, 4, 4, \dots, 4)$ and the proof of the claim is complete.

Assume that $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$, by Claim 2.5 $D_{G^g}(Y_2) = (8, 8, 8, 8, 8)$. Since there exist at least two vertices of Y_1 say y', y'' such that, $|N_{G^g}(y')| = |N_{G^g}(y'')| = 9$ thus, $y', y'' \in \{y_6, y_7, y_8, y_9\}$. W.l.g we may assume that $y' = y_6$ and $N_{G^g}(y_6) = X_2 = \{x_1, x_3, \dots, x_{10}\}$. By Claim 2.6 $|N_{G^g}(x) \cap Y_1| = 4$ and $|N_{G^g}(x) \cap Y_2| = 2$ for each $x \in X_2 \setminus \{x_1\}$ that is, $|N_{G^g}(x) \cap \{y_7, y_8, y_9\}| = 1$ for each $x \in X_2 \setminus \{x_1\}$. Since $|X_2 \setminus \{x_1\}| = 8$ and $|N_{G^g}(x) \cap \{y_7, y_8, y_9\}| = 1$, by the pigeon-hole principle we can check that there exist a vertex of $\{y_7, y_8, y_9\}$ say y such that, $|N_{G^g}(y) \cap X_2 \setminus \{x_1\}| \geq 3$. W.l.g we may assume that $y = y_7$ and $\{x_3, x_4, x_5\} \subseteq N_{G^g}(y_7) \cap X_2 \setminus \{x_1\}$. Since $|Y_2| = 5$ and $|N_{G^g}(x_i) \cap Y_2| = 2$ for $i = 3, 4, 5$, we can say that there exist $i, i' \in \{3, 4, 5\}$ such that, $|N_{G^g}(x_i) \cap N_{G^g}(x_{i'}) \cap Y_2| \neq 0$, w.l.g assume that $i = 3, i' = 4$ and $y_1 \in N_{G^g}(x_3) \cap N_{G^g}(x_4) \cap Y_2$. Therefore, $K_{3,3} \subseteq G^g[\{x_1, x_3, x_4\}, \{y_1, y_6, y_7\}]$, a contradiction. So, $n \leq 73$ and the proof of the theorem is complete. ■

In the following theorem we prove that in any 3-edge coloring of $K_{17,17}$ say (G^r, G^b, G^g) where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$, if there exist a vertex of $V(K)$ say x such that, $|N_{G^g}(x)| = 9$ and $\sum_{x_i \in X \setminus \{x\}} |N_{G^g}(x_i) \cap N_{G^g}(x)| = 64$ then, $K_{3,3} \subseteq G^g$.

Theorem 2.7. Assume that (G^r, G^b, G^g) be 3-edge coloring of $K = K_{17,17}$ such that, $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$. Assume that there exist a vertex of $V(K)$ say x such that, $|N_{G^g}(x)| = 9$. If $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1| = 73$ where $Y_1 = N_{G^g}(x)$ then, $K_{3,3} \subseteq G^g$.

Proof. By contrary, assume that $K_{3,3} \not\subseteq G^g$. Therefore, by Theorem 2.3 and Theorem 2.4 we have the following results:

- (a) $|E(G^g)| = 141$,
- (b) $\Delta(G^g) = 9$ and $\delta(G^g) = 8$,
- (c) $D_{G^g}(X) = D_{G^g}(Y) = (9, 9, 9, 9, 9, 8, 8, \dots, 8)$,
- (d) For each $x' \in X \setminus \{x\}$ we have $|N_{G^g}(x) \cap N_{G^g}(x')| \leq 5$,
- (e) If $A = \{x \in X, |N_{G^g}(x)| = 9\}$, then $|A| = 5$ and $72 \leq \sum_{y \in N_{G^g}(x)} |N_{G^g}(y)| \leq 73$, for each $x \in A$.

Assume that $X = \{x_1, x_2, \dots, x_{17}\}$, $Y = \{y_1, y_2, \dots, y_{17}\}$ be the partition set of $K = K_{17,17}$ and (G^r, G^b, G^g) be 3-edges coloring of K where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$. W.l.g assume that $x = x_1$, $Y_1 = \{y_1, y_2, \dots, y_9\}$ and $n = \sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1| = 73$. Since $n = 73$ by (c) we can say that $D_{G^g}(Y_1) = (d_1, d_2, \dots, d_9) = (9, 8, 8, \dots, 8)$ that is, there exist a vertex of Y_1 say y such that, $|N_{G^g}(y)| = 9$. By (d) we have $|N_{G^g}(x_1) \cap N_{G^g}(x)| \leq 5$ for each $x \in X \setminus \{x_1\}$. Set $C = \{x \in X, |N_{G^g}(x) \cap N_{G^g}(x_1)| = 5\}$. Now by argument similar to proof of the Claim 2.5 we have the following claim:

Claim 2.8. Assume that $x \in C$ and $N_{G^g}(x) \cap Y_1 = Y'$ then, for each $x' \in X \setminus \{x_1, x\}$ we have $|N_{G^g}(x') \cap Y'| = 2$ and $D_{G^g}(Y') = (8, 8, 8, 8, 8)$.

Now we have the following claim about $|C|$:

Claim 2.9. $|C| \leq 2$.

Proof of the Claim. By contrary, assume that $|C| \geq 3$. W.l.g assume that $\{x_2, x_3, x_4\} \subseteq C$ and $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$. By Claim 2.8 $|N_{G^g}(x) \cap Y_2| = 2$ for each $x \in X \setminus \{x_1, x_2\}$. W.l.g assume that $N_{G^g}(x_3) \cap Y_1 = Y_3 = \{y_1, y_2, y_6, y_7, y_8\}$. Since $x_4 \in C$ and $|N_{G^g}(x_4) \cap Y_i| = 2$ for $i = 2, 3$, we have $y_9 \in N_{G^g}(x_4) \cap Y_1$. Hence, for each $y \in Y_1$ there is at least one $i \in \{2, 3, 4\}$ such that $y \in N_{G^g}(x_i)$ therefore, by Claim 2.8 $D_{G^g}(Y_1) = (8, 8, 8, 8, 8, 8, 8, 8, 8)$, which is in contrast to $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1| = \sum_{i=1}^{i=9} |N_{G^g}(y_i)| = 73$ so, $|C| \leq 2$.

Now by considering $|C|$ there are three cases as follow:

Case 1: $|C| = 0$. Since $n = 73$, $|Y_1| = 9$ and $|C| = 0$ so, $D_{G^g}(X \setminus \{x_1\}) = (4, 4, \dots, 4, 4)$, $D_{G^g}(Y_1) = (9, 8, 8, 8, 8, 8, 8, 8, 8)$, $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y'| = 68$ and $D_{G^g}(Y') = (9, 9, 9, 9, 8, 8, 8, 8)$ where $Y' = Y \setminus Y_1$. Set $B = \{y \in Y', |N_{G^g}(y)| = 9\}$ so, $|B| = 4$. Now we are ready to prove the following claim:

Claim 2.10. There exist a vertex of $A \setminus \{x_1\}$ say x such that:

$$\sum_{y \in N_{G^g}(x)} |N_{G^g}(y)| \geq 74$$

Where $A = \{x \in X, |N_{G^g}(x)| = 9\}$.

Proof of the Claim . Since $D_{G^g}(X_1) = (4, 4, \dots, 4, 4)$ and $D_{G^g}(Y_1) = (9, 8, 8, 8, 8, 8, 8, 8, 8)$ for each $x \in A \setminus \{x_1\}$, one can say that:

$$\sum_{y \in N_{G^g}(x) \cap Y_1} |N_{G^g}(y)| \geq 32$$

As $|Y'| = 8$, $|B| = 4$ and $|N_{G^g}(x_i) \cap Y'| = 5$ for each $x \in A \setminus \{x_1\}$, there exist at least one vertex of $A \setminus \{x_1\}$ say x such that, $|N_{G^g}(x) \cap B| \geq 2$ otherwise, $K_{3,3} \subseteq G^g[A, Y' \setminus B]$, a contradiction. Hence w.l.g assume that $x_2 \in A$ where $|N_{G^g}(x_2) \cap B| \geq 2$, so we have:

$$\sum_{y \in N_{G^g}(x_2) \cap Y'} |N_{G^g}(y)| \geq 42$$

That is, we have:

$$\sum_{y \in N_{G^g}(x_2)} |N_{G^g}(y)| = \sum_{y \in N_{G^g}(x_2) \cap Y'} |N_{G^g}(y)| + \sum_{y \in N_{G^g}(x_2) \cap Y_1} |N_{G^g}(y)| \geq 42 + 32 = 74$$

Now by considering x_2 and $N_{G^g}(x_2)$ and by (e) (or part (b) of Theorem 2.4) $K_{3,3} \subseteq G^g$, a contradiction again.

Case 2: $|C| = 1$. W.l.g assume that $C = \{x_2\}$, $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$. By Claim 2.8 $|N_{G^g}(x_2) \cap N_{G^g}(x) \cap Y_1| = 2$ for each $x \in X \setminus \{x_1, x_2\}$ and $|N_{G^g}(y_i)| = 8$ for each $i \in \{1, 2, \dots, 5\}$. Since there exist a vertex of Y_1 say y such that, $|N_{G^g}(y)| = 9$, w.l.g we may assume that $y = y_6$ and $N_{G^g}(y_6) = \{x_1, x_3, x_4, \dots, x_{10}\}$. Since $n = 73$ and $|C| = 1$ thus, $D_{G^g}(X_1) = (5, 4, 4, \dots, 4, 3)$ that is, there exist at least seven vertices of $N_{G^g}(y_6) \setminus \{x_1\}$ say $X_3 = \{x_3, x_4, \dots, x_9\}$ such that,

$|N_{G^g}(x) \cap Y_1| = 4$ for each $x \in X_3$. Since $|X_3| = 7$, $|Y_2| = 5$, $|N_{G^g}(x) \cap Y_1| = 4$ and $|N_{G^g}(x_i) \cap Y_2| = 2$ for each $x \in X_3$ thus, $|N_{G^g}(x) \cap \{y_7, y_8, y_9\}| = 1$ for each $x \in X_3$. Therefore, by the pigeon-hole principle we can say that there exist a vertex of $\{y_7, y_8, y_9\}$ say y' such that, $|N_{G^g}(y') \cap X_3| \geq 3$. W.l.g assume that $y' = y_7$ and $\{x_3, x_4, x_5\} \subseteq N_{G^g}(y_7)$. Therefore, since $|Y_2| = 5$ one can check that there exist $i, i' \in \{3, 4, 5\}$ such that, $|N_{G^g}(x_i) \cap N_{G^g}(x_{i'}) \cap Y_2| \neq 0$. W.l.g assume that $i = 3, i' = 4$ and $y_1 \in N_{G^g}(x_3) \cap N_{G^g}(x_4) \cap Y_2$. Therefore, $K_{3,3} \subseteq G^g[\{x_1, x_3, x_4\}, \{y_1, y_6, y_7\}]$ which is a contradiction.

Case 3: $|C| = 2$. W.l.g assume that $C = \{x_2, x_3\}$, $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$. By Claim 2.8 we have $|N_{G^g}(x_2) \cap N_{G^g}(x_3) \cap Y_1| = 2$. So, w.l.g we may assume that $N_{G^g}(x_3) \cap Y_1 = Y_3 = \{y_1, y_2, y_6, y_7, y_8\}$. Now by Claim 2.8 we have $|N_{G^g}(y_i)| = 8$ for each $i \in \{1, 2, \dots, 8\}$. Since there is a vertex of Y_1 say y such that, $|N_{G^g}(y)| = 9$, we have $y = y_9$. W.l.g we may assume that $N_{G^g}(y_9) = X_2 = \{x_1, x_4, x_5, \dots, x_{11}\}$. Since $n = 73$ and $|C| = 2$ thus, $D_{G^g}(X_1) = (5, 5, 4, 4, \dots, 4, 3, 3)$ that is, there exist two vertices of X say x, x' such that, $|N_{G^g}(x) \cap Y_1| = 3$. If $|N_{G^g}(y_9) \cap \{x, x'\}| \leq 1$ then, there exist at least seven vertices of $N_{G^g}(y_9) \setminus \{x_1\}$ such that, $|N_{G^g}(x) \cap Y_1| = 4$, in this case the proof is same as Case 1. Hence, assume that $x, x' \in N_{G^g}(y_9)$. Since $|N_{G^g}(x) \cap Y_2| = |N_{G^g}(x') \cap Y_2| = 2$, one can check that $|N_{G^g}(x) \cap \{y_6, y_7, y_8\}| = |N_{G^g}(x') \cap \{y_6, y_7, y_8\}| = 0$. Assume that $X_i = N_{G^g}(y_i)$ for $i = 6, 7, 8$. Since $|X_i| = 8$ and $x, x' \notin X_i$ then, for each $x \in X_i \setminus \{x_1\}$ we have $|N_{G^g}(x) \cap Y_1| = 4$. Therefore, by considering $X_i \setminus \{x_1\}$ and y_i for each $i \in \{6, 7, 8\}$, the proof is same as Case 1 and $K_{3,3} \subseteq G^g$, a contradiction again.

Therefore, by Cases 1, 2, 3 the assumption dose not hold that is, $K_{3,3} \subseteq G^g$ and the proof is complete. \blacksquare

In the following theorem we prove that in any 3-edge coloring of $K_{17,17}$ say (G^r, G^b, G^g) where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$, if there exist a vertex of $V(K)$ say x such that, $|N_{G^g}(x)| = 9$ and $\sum_{x_i \in X \setminus \{x\}} |N_{G^g}(x_i) \cap N_{G^g}(x)| = 63$ then, $K_{3,3} \subseteq G^g$.

Theorem 2.11. *Assume that (G^r, G^b, G^g) be 3-edge coloring of $K = K_{17,17}$ where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$. Assume that there exist a vertex of $V(K)$ say x such that, $|N_{G^g}(x)| = 9$. If $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1| = 72$ where $Y_1 = N_{G^g}(x)$ then, $K_{3,3} \subseteq G^g$.*

Proof. By contrary, assume that $K_{3,3} \not\subseteq G^g$. Therefore, by Theorems 2.3 and 2.4 the following results are true:

- (a) $|E(G^g)| = 141$,
- (b) $\Delta(G^g) = 9$ and $\delta(G^g) = 8$,
- (c) $D_{G^g}(X) = D_{G^g}(Y) = (9, 9, 9, 9, 9, 8, 8, \dots, 8)$,
- (d) For each $x \in X \setminus \{x_1\}$, we have $|N_{G^g}(x) \cap Y_1| \leq 5$,
- (e) If $A = \{x \in X, |N_{G^g}(x)| = 9\}$ then, $|A| = 5$ and $72 \leq \sum_{y \in N_{G^g}(x)} |N_{G^g}(y)| \leq 73$, for each

$x \in A$.

Assume that $X = \{x_1, x_2, \dots, x_{17}\}$, $Y = \{y_1, y_2, \dots, y_{17}\}$ be the partition set of $K = K_{17,17}$ and (G^r, G^b, G^g) be 3-edges coloring of K where $K_{2,2} \not\subseteq G^r$, $K_{2,2} \not\subseteq G^b$ and $K_{3,3} \not\subseteq G^g$. W.l.g assume that $x = x_1$, $Y_1 = \{y_1, y_2, \dots, y_9\}$ and $n = \sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y_1| = 72$. Since $n = 73$ by (c) we can say that $D_{G^g}(Y_1) = (d_1, d_2, \dots, d_9) = (8, 8, 8, \dots, 8)$. Set $C = \{x \in X, |N_{G^g}(x) \cap N_{G^g}(x_1)| = 5\}$. Define D and E as follow:

$$D = \{x \in X \setminus \{x_1\}, \text{ such that } |N_{G^g}(x) \cap Y_1| = 5\}$$

$$E = \{x \in X \setminus \{x_1\}, \text{ such that } |N_{G^g}(x) \cap Y_1| = 3\}$$

Now we have the following claim:

Claim 2.12. $|D| \leq 3$ and $|E| \leq 4$.

Proof of the Claim. By contrary, assume that $|D| \geq 4$. W.l.g suppose that $\{x_2, x_3, x_4, x_5\} \subseteq D$, $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$. Now, by Claim 2.8 we have $|N_{G^g}(x) \cap Y_2| = 2$ for each $x \in X \setminus \{x_1, x_2\}$. W.l.g we may assume that $N_{G^g}(x_3) \cap Y_1 = Y_3 = \{y_1, y_2, y_6, y_7, y_8\}$. Consider $N_{G^g}(x_i) \cap Y_1 (i = 4, 5)$. Since $|N_{G^g}(x_i) \cap Y_j| = 2 (i = 4, 5, j = 2, 3)$ and $x_i \in A$ we can check that $|N_{G^g}(x_i) \cap \{y_3, y_4, y_5\}| = 2$, $|N_{G^g}(x_i) \cap \{y_6, y_7, y_8\}| = 2$ and $y_9 \in N_{G^g}(x_i)$ for $i = 4, 5$ otherwise, if there exist a vertex of $\{x_4, x_5\}$ say x such that, $|N_{G^g}(x_i) \cap \{y_1, y_2\}| \neq 2$ then, one can say that $K_{3,3} \subseteq G^g[\{x_1, x_i, x\}, Y_1]$ for some $i \in \{1, 2\}$, a contradiction. Therefore, since $|\{y_3, y_4, y_5\}| = |\{y_6, y_7, y_8\}| = 3$ and $x_4, x_5 \in A$, by the pigeon-hole principle $|N_{G^g}(x_4) \cap N_{G^g}(x_5) \cap \{y_3, y_4, y_5\}| \geq 1$ and $|N_{G^g}(x_4) \cap N_{G^g}(x_5) \cap \{y_6, y_7, y_8\}| \geq 1$. W.l.g we may assume that $y_3, y_6 \in N_{G^g}(x_4) \cap N_{G^g}(x_5)$ therefore, since $y_9 \in N_{G^g}(x_4) \cap N_{G^g}(x_5)$, we have $K_{3,3} \subseteq G^g[\{x_1, x_4, x_5\}, \{y_3, y_6, y_9\}]$, a contradiction. Therefore, $|D| \leq 3$. Now, as $\sum_{i=2}^{i=17} |N_{G^g}(x_i) \cap Y_1| = 63$ and $|D| \leq 3$, we can say that $|E| \leq 4$ and the proof of claim is complete.

Now, by considering $|D|$, there are three cases as follow:

Case 1: $|D| = 0$. Since $n = 72$ and $|D| = 0$ thus? $D_{G^g}(X \setminus \{x_1\}) = (4, 4, \dots, 4, 3)$ and $D_{G^g}(Y_1) = (8, 8, 8, 8, 8, 8, 8, 8, 8, 8)$, $\sum_{i=1}^{i=17} |N_{G^g}(x_i) \cap Y'| = 69$ and $D_{G^g}(Y') = (9, 9, 9, 9, 9, 8, 8, 8, 8)$ where $Y' = Y \setminus Y_1$. Set $B = \{y \in Y', |N_{G^g}(y)| = 9\}$ Hence, $|B| = 5$. Now, we have the following claim:

Claim 2.13. *There exist a vertex of $A \setminus \{x_1\}$ say x such that:*

$$\sum_{y \in N_{G^g}(x)} |N_{G^g}(y)| \geq 75$$

Where $A = \{x \in X, |N_{G^g}(x)| = 9\}$.

Proof of the Claim. Since $D_{G^g}(X_1) = (4, 4, \dots, 4, 3)$ and $D_{G^g}(Y_1) = (8, 8, 8, 8, 8, 8, 8, 8, 8, 8)$ so, for at least three vertices of $A \setminus \{x_1\}$ one can say that:

$$\sum_{y \in N_{G^g}(x) \cap Y_1} |N_{G^g}(y)| \geq 32$$

Therefore, since $|N_{G^g}(x_i) \cap Y'| = 5$ for each $x \in A \setminus \{x_1\}$ and $D_{G^g}(Y') = (9, 9, 9, 9, 9, 8, 8, 8, 8)$, there exist at least one vertex of $A \setminus \{x_1\}$ say x such that, $|N_{G^g}(x) \cap B| \geq 3$ otherwise, $K_{3,3} \subseteq G^g[A, Y' \setminus B]$, a contradiction. Hence, w.l.g assume that $x_2 \in A$ and $|N_{G^g}(x_2) \cap B| \geq 3$ therefore, we have:

$$\sum_{y \in N_{G^g}(x) \cap Y''} |N_{G^g}(y)| \geq 3 \times 9 + 2 \times 8 = 43$$

That is, we have:

$$\sum_{y \in N_{G^g}(x_2)} |N_{G^g}(y)| = \sum_{y \in N_{G^g}(x_2) \cap Y'} |N_{G^g}(y)| + \sum_{y \in N_{G^g}(x_2) \cap Y_1} |N_{G^g}(y)| \geq 43 + 32 = 75$$

Now by considering x_2 and $N_{G^g}(x_2)$ and by (e) (or by part (b) of Theorem 2.4), $K_{3,3} \subseteq G^g$, a contradiction again.

Case 2: $|D| = 1$ (for the case that $(|D| = 2)$ the proof is same). W.l.g assume that $D = \{x_2\}$, $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$. Since $n = 72$, $|D| = 1$ and $|N_{G^g}(x) \cap Y_1| \leq 5$ we can say that $|E| = 2$. As $|N_{G^g}(x) \cap Y_2| = 2$ for each $x \in X \setminus \{x_1, x_2\}$ and $|E| = 2$, we can check that

there exist a vertex of $\{y_6, y_7, y_8, y_9\}$ say y such that, for each vertex of $N_{G^g}(y) \cap X \setminus \{x_1\}$ say x , $|N_{G^g}(x) \cap Y_1| = 4$. W.l.g we may assume that $y = y_6$, $N_{G^g}(y_6) \cap X \setminus \{x_1\} = \{x_3, x_4, \dots, x_9\}$. Since $|N_{G^g}(y_6) \cap X \setminus \{x_1\}| = 7$ and $|N_{G^g}(x) \cap Y_2| = 2$ for each $x \in N_{G^g}(y_6) \cap X \setminus \{x_1\}$ so, $|N_{G^g}(x) \cap \{y_7, y_8, y_9\}| = 1$. Therefore, by the pigeon-hole principle we can check that there exist a vertex of $\{y_7, y_8, y_9\}$ say y' such that, $|N_{G^g}(y_6) \cap N_{G^g}(y') \cap X \setminus \{x_1\}| \geq 3$. W.l.g assume that $y' = y_7$ and $\{x_3, x_4, x_5\} \subseteq N_{G^g}(y_6) \cap N_{G^g}(y_7) \cap X \setminus \{x_1\}$. Therefore, since $|Y_2| = 5$ and $|N_{G^g}(x) \cap Y_2| = 2$, there exist at least two vertices of $\{x_3, x_4, x_5\}$ say x', x'' such that, $|N_{G^g}(x') \cap N_{G^g}(x'') \cap Y_2| \neq 0$. W.l.g assume that $x' = x_3, x'' = x_4$ and $y_1 \in N_{G^g}(x_3) \cap N_{G^g}(x_4)$ therefore, we have $K_{3,3} \subseteq G^g[\{x_1, x_3, x_4\}, \{y_1, y_6, y_7\}]$, a contradiction.

Case 3: $|D| = 3$. W.l.g assume that $D = \{x_2, x_3, x_4\}$, $N_{G^g}(x_2) \cap Y_1 = Y_2 = \{y_1, \dots, y_5\}$. By Claim 2.8 we have $|N_{G^g}(x_2) \cap N_{G^g}(x_3) \cap Y_1| = 2$. W.l.g we may assume that $N_{G^g}(x_3) \cap Y_1 = Y_3 = \{y_1, y_2, y_6, y_7, y_8\}$. Since $x_4 \in D$ and $|N_{G^g}(x_4) \cap Y_i| = 2$ for $i = 2, 3$ thus, $y_9 \in N_{G^g}(x_4)$. If $|N_{G^g}(x_4) \cap \{y_1, y_2\}| \neq 0$, as $|N_{G^g}(x_2) \cap N_{G^g}(x_4) \cap Y_1| = 2$ and $x_4 \in D$, one can check that $|N_{G^g}(x_4) \cap \{y_6, y_7, y_8\}| = 2$ that is, $K_{3,3} \subseteq G^g[\{x_1, x_3, x_4\}, Y_1]$, a contradiction. Hence, $|N_{G^g}(x_4) \cap \{y_1, y_2\}| = 0$ therefore, $|N_{G^g}(x_4) \cap \{y_3, y_4, y_5\}| = 2$ and $|N_{G^g}(x_4) \cap \{y_6, y_7, y_8\}| = 2$. W.l.g we may assume that $N_{G^g}(x_4) \cap Y_1 = Y_4 = \{y_3, y_4, y_6, y_7, y_9\}$. Since $|D| = 3$ we can say that $|E| = 4$. W.l.g assume that $E = \{x_5, x_6, x_7, x_8\}$. Now, we have the following claim:

Claim 2.14. $|N_{G^g}(y_9) \cap E| = 0$.

Proof of the Claim. By contrary, assume that $|N_{G^g}(y_9) \cap E| \neq 0$. Assume that $x_5 \in N_{G^g}(y_9) \cap E$ that is, $x_5 y_9 \in E(G^g)$. Since $x_5 \in E$ and $\{x_2, x_3, x_4\} = D$, by Claim 2.8 we have $|N_{G^g}(x_5) \cap N_{G^g}(x_i)| = |N_{G^g}(x_5) \cap Y_i| = 2$ for $i = 2, 3, 4$. Consider $N_{G^g}(x_5) \cap Y_2$, assume that $N_{G^g}(x_5) \cap Y_2 = \{y', y''\}$, if $\{y', y''\} = \{y_1, y_2\}$ then, $|N_{G^g}(x_5) \cap Y_4| = 1$, a contradiction. Therefore, we may assume that $|\{y', y''\} \cap \{y_1, y_2\}| \leq 1$. If $|\{y', y''\} \cap \{y_1, y_2\}| = 0$ then, $|N_{G^g}(x_5) \cap Y_3| = 0$ and if $|\{y', y''\} \cap \{y_1, y_2\}| = 1$ then, $|N_{G^g}(x_5) \cap Y_3| \leq 1$, in any case there exist a vertex of D say x' such that, $|N_{G^g}(x_5) \cap N_{G^g}(x')| = 1$, a contradiction. So, the assumption dose not hold and the claim is true.

Therefore, by Claim 2.14, since $|N_{G^g}(y_9) \cap D| = 0$, we can say that for any vertex of $N_{G^g}(y_9) \cap X \setminus \{x_1\}$ say x , $|N_{G^g}(x) \cap Y_1| \geq 4$ therefore, by considering Y_2 and y_9 , as $|N_{G^g}(y_9) \cap X \setminus \{x_1\}| = 7$ and $|N_{G^g}(x) \cap Y_1| \geq 4$ for each $x \in N_{G^g}(y_9) \cap X \setminus \{x_1\}$ the proof is same as Case 1, a contradiction. Therefore, by Cases 1, 2, 3 the assumption dose not hold that is, $K_{3,3} \subseteq G^g$ and the proof is complete. ■

Now, combining Theorems 2.3, 2.4 2.7 and 2.11, yield the proof of the Theorem 1.1.

REFERENCES

- [1] BEINERE, L., LW, B., AJ, S.: On a bipartite form of the ramsey problem. (1976)
- [2] Bondy, J.A., Murty, U.S.R., et al.: Graph theory with applications, vol. 290. Macmillan London (1976)
- [3] Collins, A.F.: Bipartite Ramsey Numbers and Zarankiewicz Numbers. Rochester Institute of Technology (2015)
- [4] Collins, A.F., Riasanovsky, A.W., Wallace, J.C.: Zarankiewicz numbers and bipartite ramsey numbers. Journal of Algorithms and Computation **47**, 63–78 (2016)
- [5] Dybizbański, J., Dzido, T., Radziszowski, S.: On some zarankiewicz numbers and bipartite ramsey numbers for quadrilateral. ARS COMBINATORIA **119**, 275–287 (2015)
- [6] Erdős, P., Rado, R.: A partition calculus in set theory. Bulletin of the American Mathematical Society **62**(5), 427–489 (1956)
- [7] Exoo, G.: A bipartite ramsey number. Graphs and Combinatorics **7**, 395–396 (1991)
- [8] Gholami, M., Rowshan, Y.: The bipartite ramsey numbers $br(c_8, c_{2n})$ (2021)
- [9] Goddard, W., Henning, M.A., Oellermann, O.R.: Bipartite ramsey numbers and zarankiewicz numbers. Discrete Mathematics **219**(1-3), 85–95 (2000)

- [10] Hatala, I., Héger, T., Mattheus, S.: New values for the bipartite ramsey number of the four-cycle versus stars. *Discrete Mathematics* **344**(5), 112320 (2021)
- [11] Hattingh, J.H., Henning, M.A.: Star-path bipartite ramsey numbers. *Discrete Mathematics* **185**(1-3), 255–258 (1998)
- [12] Kóvari T., S.V.T.P.: On a problem of k. zarankiewicz. *Colloquium Mathematicae* **3**(1), 50–57 (1954). URL <http://eudml.org/doc/210011>
- [13] Raeisi, G.: Star-path and star-stripe bipartite ramsey numbers in multicoloring. *Transactions on Combinatorics* **4**(3), 37–42 (2015)
- [14] Reiman, I.: Über ein problem von k. zarankiewicz. *Acta Mathematica Academiae Scientiarum Hungarica* **9**, 269–273 (1958)
- [15] Rowshan, Y., Gholami, M., Shateyi, S.: The size, multipartite ramsey numbers for nk_2 versus path–path and cycle. *Mathematics* **9**(7), 764 (2021)

¹DEPARTMENT OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDIES IN BASIC SCIENCES (IASBS), ZANJAN 66731-45137, IRAN

Email address: `y.rowshan@iasbs.ac.ir`

Email address: `gholami.m@iasbs.ac.ir`