

Gromov-Hausdorff distance with boundary and its application to Gromov hyperbolic spaces and uniform spaces

Hyogo Shibahara*

*Department of Mathematical Sciences, University of Cincinnati,
P.O. Box 210025, Cincinnati, OH 45221-0025, U.S.A.*

Abstract

In this paper we introduce a notion of the Gromov-Hausdorff distance with boundary, denoted by d_{GHB} , to construct a framework of convergence of noncomplete metric spaces. We show that a class of bounded A -uniform spaces with diameter bounded from below is a complete metric space with respect to d_{GHB} . As an application we show the stability of Gromov hyperbolicity, roughly starlike property, uniformization, quasihyperbolization, and boundary of Gromov hyperbolic spaces under appropriate notions of convergence and assumptions.

Keywords and phrases : Gromov-Hausdorff distance, Gromov hyperbolic space, uniform domain, uniform space, uniformization, quasihyperbolic metric.

Mathematics Subject Classification (2020) : Primary: 53C23; Secondary: 51F30

1 Introduction

The notion of uniform domains was first introduced in [MS]. Since then, they have played a key role in deriving many important results in the context of quasiconformal mappings, such as extendability of Sobolev functions. Although a uniform domain was classically considered to be in the Euclidean plane, recent studies on uniform domains in metric measure spaces with doubling measure supporting p -Poincaré inequality have been also conducted, see, for example, [A], [BHK], [BBS], [BSh], [GeO],[Mar], [V1]). Also, in [BHK], uniform domains were generalized as noncomplete locally compact metric spaces that satisfy the criteria of Definition 2.19, which we call uniform spaces. Moreover, in [BHK] it has been shown that there is a connection between proper geodesic roughly starlike Gromov hyperbolic spaces and uniform spaces and that if a roughly starlike Gromov hyperbolic space

*Email: shibahho@mail.uc.edu

(X, d) is uniformized with a sufficiently small parameter $\epsilon > 0$, then the resulting space, denoted by (X^ϵ, d_ϵ) , is a uniform space. Conversely, a uniform space Ω with the quasihyperbolic metric k is a Gromov hyperbolic space. Definitions of these terms are introduced in the next section. The above connection can be seen as the generalization of the correspondence between the Poincaré disk model and the Euclidean disk. Quite recently Butler obtained another uniformization result, which is motivated by the correspondence between the upper half plane and the hyperbolic space, see [Bu].

The Gromov-Hausdorff distance was originally defined in [Ed]. It has been widely used in Riemannian geometry and analysis on metric spaces. Since every uniform space Ω is a noncomplete locally compact metric space with additional properties as in Definition 2.19, we need to consider the boundary $\partial\Omega := \overline{\Omega} \setminus \Omega$, in addition to the metric space Ω itself. There are a few studies on convergence of Riemannian manifolds and metric spaces with boundary, see [PS] and [P]. In [PS], a new notion of limit spaces called glued limit spaces was introduced for a sequence of noncomplete precompact metric spaces through δ -inner regions. It was observed there that in some situations there is no Gromov-Hausdorff limit but a glued limit space still exists. Another feature is that this limit is realized as a subset of the Gromov-Hausdorff limit if, of course, the Gromov-Hausdorff limit exists. Glued limit spaces, however, do not take care of the issue of the limit of the sequence of boundaries very well in general even if there is a Gromov-Hausdorff limit for a sequence of metric spaces, when we take the completion of a glued limit space. On the other hand, as we mentioned above, uniform spaces have to be treated together with their boundaries. It is therefore natural to ask how we could extend the notion of the Gromov-Hausdorff distance, taking boundary into consideration. One possible way to do so is a simple modification of the Gromov-Hausdorff distance. For metric spaces X and Y , we consider

$$d_{GHB}(X, Y) := \inf d_H^Z(\varphi(X), \psi(Y)) + d_H^Z(\varphi(\partial X), \psi(\partial Y))$$

where d_H^Z is the Hausdorff distance in Z and infimum is taken over all metric spaces Z and isometric embeddings $\varphi : \overline{X} \rightarrow Z$ and $\psi : \overline{Y} \rightarrow Z$. Note that $\partial X := \overline{X} \setminus X$ where \overline{X} is a metric completion of X . This definition allows us to distinguish between a metric space and its completion while the Gromov-Hausdorff distance does not see the difference between them. We adopt d_{GHB} as our definition of the Gromov-Hausdorff distance with boundary and we will show basic properties of d_{GHB} for the space of all noncomplete precompact locally compact metric spaces.

In [HRS] various types of convergence of a sequence of sets with quasihyperbolic distances, such as the Gromov-Hausdorff convergence and the Carathéodory convergence, were investigated. Since a notion of Gromov-Hausdorff distance with boundary has not been established, the paper [HRS] deals with a sequence of sets in a same metric space. So one could ask how the results in [HRS] would be extended for a sequence of noncomplete metric spaces, see Theorem 1.4 for a related result.

The goal of this paper is to examine the metric d_{GHB} and properties that a limit space

has from the geometric function theoretic viewpoint. Hence, the following questions are addressed in this paper:

1. Is d_{GHB} a metric? Is the class of all bounded uniform spaces complete with respect to this metric?
2. Are Gromov hyperbolicity and roughly starlike property stable under the pointed Gromov-Hausdorff convergence?
3. Are uniformized spaces and their boundaries stable under the pointed Gromov-Hausdorff convergence?
4. Is quasihyperbolization stable under the Gromov-Hausdorff convergence with boundary?

In this paper, we provide positive answers for the first three questions. Here we list the theorems.

Theorem 1.1. *Let $A > 0$ and $R > 0$. Let \mathcal{M} be the class of all noncomplete precompact locally compact metric spaces and $\mathcal{U}(A, R)$ be the class of all bounded A -uniform spaces with the diameter bounded from below by $R > 0$. Then (\mathcal{M}, d_{GHB}) is a (noncomplete) metric space and $(\mathcal{U}(A, R), d_{GHB})$ is a complete metric space.*

We remark that there is one direct application to analysis on metric spaces. The fact that $\mathcal{U}(A, R)$ is complete with respect to d_{GHB} together with [BSh, Theorem 4.4], [Kei, Theorem 3] and [Ch, Chapter 9] gives us a stability of metric measure spaces with doubling measure supporting p -Poincaré inequality, which is new for a sequence of noncomplete metric measure spaces, see Remark 4.9.

Theorem 1.2. *Let $(X_n, d_n, p_n)_n$ be a sequence of pointed proper geodesic δ -Gromov hyperbolic spaces which is pointed Gromov-Hausdorff convergent to a complete metric space (X, d, p) . Then, X is a proper geodesic δ -Gromov hyperbolic space. Moreover, if the spaces $(X_n, d_n, p_n)_n$ are further assumed to be M -roughly starlike, then there exists $\tilde{M} := \tilde{M}(M, \delta)$ such that X is \tilde{M} -roughly starlike.*

Theorem 1.3. *Let $(X_n, d_n, p_n)_n$ be a sequence of pointed proper geodesic M -roughly starlike δ -Gromov hyperbolic spaces which is pointed Gromov-Hausdorff convergent to a complete metric space (X, d, p) . For each $0 < \epsilon \leq \epsilon_0(\delta)$, let $(X_n^\epsilon, d_{n,\epsilon})_n$ the sequence of uniformized spaces. Then $(X_n^\epsilon, d_{n,\epsilon})_n$ Gromov-Hausdorff converges to (X^ϵ, d_ϵ) with boundary.*

We remark that the stability of the Gromov hyperbolicity holds true, under the assumption that the limit exists for a given sequence. This assumption is valid if one considers a sequence of intrinsic A -uniform spaces endowed with quasihyperbolic metric where diameter of spaces and doubling constants with respect to original metrics are uniformly bounded, see Theorem 1.4 and Theorem 4.7. On the other hand, in general a pointed

Gromov-Hausdorff limit may not exist for some sequence of Gromov hyperbolic spaces.

It is clear that for given noncomplete precompact locally compact metric spaces X and Y , we have

$$d_{GH}(\overline{X}, \overline{Y}) + d_{GH}(\partial X, \partial Y) \leq d_{GHB}(X, Y). \quad (1)$$

Hence Theorem 1.3 also states that $(\overline{X}_n^\epsilon, d_{n,\epsilon})_n$ and $(\partial_{d_{n,\epsilon}} X_n^\epsilon, d_{n,\epsilon})_n$ Gromov-Hausdorff converge to $(\overline{X}^\epsilon, d_\epsilon)$ and $(\partial_{d_\epsilon} X^\epsilon, d_\epsilon)$, respectively. On the other hand, the converse inequality of (1) does not hold, see Remark 3.10 where we provide a simple example which also suggests that the sum in the left-hand side of (1) does not induce a metric. Also, by [BHK, Proposition 4.13], for a given proper geodesic M -roughly starlike δ -Gromov hyperbolic space (X, d, p) , the canonical bijective identification between the metric boundary of X^ϵ with respect to uniformization metric d_ϵ , $(\partial_{d_\epsilon} X^\epsilon, d_\epsilon)$, and the Gromov boundary of X with the visual metric, $(\partial_G X, d_{p,\epsilon})$ are bilipschitz equivalent. Hence, Theorem 1.3 can be seen as follows : If we change the visual metric $d_{p,\epsilon}$ of Gromov boundary to the metric d_ϵ , which is bilipschitz equivalent to $d_{p,\epsilon}$, then the sequence of Gromov boundaries $(\partial_G X_n, d_{n,\epsilon})$ converges to $(\partial_G X, d_\epsilon)$.

On the other hand, regarding the fourth question, in general the desired conclusion is false. Instead, we prove that if a sequence of uniform spaces $(\Omega_n, d_n)_n$ are all length spaces, then the quasihyperbolization is stable. To make the meaning precise, we state the last theorem.

Theorem 1.4. *Let $(\Omega_n, d_n)_n \subseteq \mathcal{U}(A, R)$ and $(\Omega, d) \in \mathcal{M}$. Suppose $(\Omega_n, d_n)_n$ are all length spaces and $(\Omega_n, d_n)_n$ converges to (Ω, d) with respect to the Gromov-Hausdorff distance with boundary. Then $(\Omega_n, k_n, p_n)_n$ is pointed Gromov-Hausdorff convergent to (Ω, k, p) for some $p_n \in \Omega_n$ and $p \in \Omega$, where k_n and k are quasihyperbolic metrics defined on Ω_n and Ω , respectively.*

The rest of this paper consists of six parts. In Section 2, we review basic concepts and propositions needed to state our results. In Section 3, we define the Gromov-Hausdorff distance with boundary d_{GHB} on \mathcal{M} and prove the first assertion of Theorem 1.1. We also explore some basic properties of d_{GHB} that the Gromov-Hausdorff distance also has. In Section 4, we prove the second assertion of Theorem 1.1. A compactness theorem for $\mathcal{U}(A, R)$ is also provided. In Section 5 we prove Theorem 1.2. In section 6, we prove Theorem 1.3. In Section 7, we give a proof of Theorem 1.4.

Acknowledgement

The author would like to thank Nageswari Shanmugalingam and Jeff Lindquist for countless helpful discussions and for carefully reading the manuscript. He also would like to appreciate his former advisor Atsushi Katsuda who brought him to this research area and gave some important ideas, remarks on negatively curved spaces and an alternative proof of Theorem 1.3, which were invaluable. The author's research was partially supported by the grant DMS #1800161 from NSF (U.S.A.).

2 Preliminaries

We fix some notations. Let (X, d) be a metric space. For $x \in X$ and $r > 0$, we set $B_d(x, r) := \{y \in X \mid d(x, y) < r\}$ and $\bar{B}_d(x, r) := \{y \in X \mid d(x, y) \leq r\}$. The length of a curve γ with respect to d is denoted by $l_d(\gamma)$. The minimum and the maximum of n real numbers $(a_i)_{i=1}^n$ is denoted by $a_1 \wedge \cdots \wedge a_n$ and $a_1 \vee \cdots \vee a_n$, respectively. We first review the Gromov-Hausdorff distance and the pointed Gromov-Hausdorff convergence.

Definition 2.1. (ϵ -neighborhood) Let $\epsilon \geq 0$ be given and X be a metric space. An ϵ -neighborhood of a given set $A \subseteq X$, denoted by A_ϵ , is defined by

$$A_\epsilon := \{x \in X \mid \text{There exists } y \in A \text{ such that } d(x, y) \leq \epsilon\}.$$

Note that if $\epsilon = 0$, then $A_0 = A$ for every set $A \subseteq X$.

Definition 2.2. (Hausdorff distance) Let $A, B \subseteq X$ be given. The Hausdorff distance, denoted by $d_H(A, B)$, is defined by

$$d_H(A, B) := \inf\{r \geq 0 \mid A \subseteq B_r \text{ and } B \subseteq A_r\}.$$

When we want to emphasize which space or metric is used, we sometimes write d_H^X or d_H^d .

Using the Hausdorff distance, we define the Gromov-Hausdorff distance.

Definition 2.3. (Gromov-Hausdorff distance) Let X, Y be bounded metric spaces. The Gromov-Hausdorff distance, denoted by $d_{GH}(X, Y)$, is defined by

$$d_{GH}(X, Y) := \inf d_H^Z(\varphi(X), \psi(Y))$$

where infimum is taken over all metric spaces Z and isometric embeddings $\varphi : X \rightarrow Z$ and $\psi : Y \rightarrow Z$.

We remark that $d_{GH}(X, \bar{X}) = 0$ where \bar{X} is the completion of X . It is also well-known that for compact metric spaces X and Y , $d_{GH}(X, Y) = 0$ if and only if there exists an isometry from X to Y . Moreover, the Gromov-Hausdorff distance d_{GH} is a metric on the set of all isometry classes of compact metric spaces, see [HKST] and [BBI].

Definition 2.4. Let (X, d) be a metric space. Define

$$Net_\epsilon(X) := \inf \{|M| \mid M \text{ is an } \epsilon\text{-net in } X\}$$

and

$$Sep_\epsilon(X) := \sup \{|M| \mid M \text{ is } \epsilon\text{-separated in } X\}.$$

where $|\cdot|$ is the cardinality of a set. Here we say that M is an ϵ -net in X if $X \subseteq M_\epsilon$ and M is ϵ -separated in X if $d(x, y) \geq \epsilon$ for every $x, y \in M$ with $x \neq y$.

The following two useful propositions are due to Gromov, see [Gr, p.64-65].

Proposition 2.5. *Let \mathcal{C} be a subset of the set of all compact metric spaces. Then, TFAE :*

1. *There exist $D > 0$ and $N : (0, \infty) \rightarrow \mathbb{N}$ such that for every $X \in \mathcal{C}$, $\text{diam}(X) \leq D$ and for every $\epsilon > 0$, $\text{Net}_\epsilon(X) \leq N(\epsilon)$.*
2. *There exist $D > 0$ and $N : (0, \infty) \rightarrow \mathbb{N}$ such that for every $X \in \mathcal{C}$, $\text{diam}(X) \leq D$ and for every $\epsilon > 0$, $\text{Sep}_\epsilon(X) \leq N(\epsilon)$.*
3. *\mathcal{C} is totally bounded with respect to d_{GH} .*

Proposition 2.6. *Let \mathcal{C} be a subset of the set of all compact metric spaces. If \mathcal{C} satisfies any of the conditions in Proposition 2.5, then there is a compact set $K \subseteq l^\infty$ such that every $X \in \mathcal{C}$ admits an isometric embedding into K .*

Next, we define the pointed Gromov-Hausdorff convergence. Although some equivalent notions can be found in many papers, we follow the pointed Gromov-Hausdorff convergence in [BBI, Definition 8.1.1] and [HKST, Definition 11.3.1]. One can also check [H] for other equivalent conditions and detailed explanation on the pointed Gromov-Hausdorff topology.

Definition 2.7. (Pointed Gromov-Hausdorff convergence) We say that $(X_n, d_n, p_n)_n$ is *pointed Gromov-Hausdorff convergent* to (X, d, p) if for every $r > 0$ and $0 < \epsilon < r$, there exists $N \in \mathbb{N}$ such that for each $n \geq N$, there exists $f_n^\epsilon : B_{d_n}(p_n, r) \rightarrow X$ such that

1. $f_n^\epsilon(p_n) = p$,
2. $|d(f_n^\epsilon(x), f_n^\epsilon(y)) - d_n(x, y)| < \epsilon$ for all $x, y \in B_{d_n}(p_n, r)$,
3. $B_d(p, r - \epsilon) \subseteq f_n^\epsilon(B_{d_n}(p_n, r))_\epsilon$.

Remark 2.8. A metric space (X, d) is said to be a length space if for every pair of points $x, y \in X$, the infimum of the lengths of curves from x to y coincides with $d(x, y)$. In the above definition, if the pointed metric spaces $(X_n, d_n, p_n)_n$ are all length spaces, then we can further assume that $f_n^\epsilon(B_{d_n}(p_n, r)) \subseteq B_d(p, r)$ for every $r > 0$ and $0 < \epsilon < r$. We leave it to the reader to verify this.

We set the terminology to state the compactness theorem for pointed metric spaces. A metric space (X, d) is said to be proper if every bounded closed subset is compact. If each pair of points in (X, d) is joined by a curve whose length equals the distance between them, we say that (X, d) is a geodesic metric space.

Definition 2.9. A sequence of pointed metric spaces $(X_n, d_n, p_n)_n$ is said to be *pointed uniformly totally bounded* if there is a function $N : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ such that, for each $0 < \epsilon < R$ and $n \in \mathbb{N}$, the closed ball $\bar{B}_{d_n}(p_n, R)$ in X_n contains an ϵ -net of cardinality at most $N(\epsilon, R)$.

The compactness theorem for pointed proper metric spaces first appeared in [Gr, p.64]. We state the compactness theorem under the pointed Gromov-Hausdorff convergence based on [HKST, Theorem 11.3.16].

Proposition 2.10. *Let $(X_n, d_n, p_n)_n$ be a sequence of pointed proper metric spaces. Suppose $(X_n, d_n, p_n)_n$ is pointed uniformly totally bounded. Then there exists a subsequence $(X_{n_k}, d_{n_k}, p_{n_k})_k$ which is pointed Gromov-Hausdorff convergent to a complete metric space (X, d, p) .*

Remark 2.11. If pointed metric spaces $(X_n, d_n, p_n)_n$ are all assumed to be proper and geodesic, then the limit pointed complete metric space (X, d, p) is also proper and geodesic by the Hopf-Rinow theorem, see [HKST, Proposition 11.3.12 and Proposition 11.3.14] and [BBI, Theorem 7.5.1] for further discussion on the pointed Gromov-Hausdorff convergence.

Definition 2.12 (Gromov product). Let X be a metric space. The *Gromov product* of two points $x, y \in X$ with respect to $m \in X$ is defined by

$$(x|y)_m := \frac{1}{2}(d(m, x) + d(m, y) - d(x, y)).$$

In order to use uniformization results from [BHK], we assume δ -Gromov hyperbolic spaces to be geodesic even though it is not required in the following four-point condition. Note that there are equivalent definitions of δ -Gromov hyperbolicity, such as δ -thinness of a geodesic triangle. See [Bo], [BH], [GH], and [V2] for more discussion about Gromov hyperbolic spaces.

Definition 2.13 (Gromov hyperbolic space). Let (X, d) be an unbounded proper geodesic metric space. We say that X is a δ -Gromov hyperbolic space if for all $x, y, z, m \in X$,

$$(x|z)_m \geq (x|y)_m \wedge (y|z)_m - \delta.$$

Definition 2.14. (M -Roughly starlike property) We say that a pointed metric space (X, d, p) is M -roughly starlike if for every $x \in X$, there exists a geodesic ray $\gamma : [0, \infty) \rightarrow X$ with $\gamma(0) = p$ such that $\text{dist}(x, \gamma) := \inf_{t \in [0, \infty)} d(x, \gamma(t)) \leq M$.

Definition 2.15. (Quasiisometric embedding/path) Let $\mu \geq 1$ and $C > 0$ be constants. We say that a map $f : X \rightarrow Y$ is a (μ, C) -quasi-isometric embedding if for every pair of points $x, y \in X$,

$$\mu^{-1}d(x, y) - C \leq d(f(x), f(y)) \leq \mu d(x, y) + C$$

holds. In particular, if X is an interval $[a, b]$, then $f : [a, b] \rightarrow Y$ satisfying the above condition is called a (μ, C) -quasiisometric path. Also, if a map $f : X \rightarrow Y$ is $(1, \epsilon)$ -isometric embedding and $Y \subseteq (f(X))_\epsilon$, we call f an ϵ -isometry.

Remark 2.16. Given compact metric spaces X and Y , the existence of an ϵ -isometry $f : X \rightarrow Y$ implies that $d_{GH}(X, Y) \leq 3\epsilon$. Conversely, if $d_{GH}(X, Y) < \epsilon$, then there exists a 5ϵ -isometry $f : X \rightarrow Y$, see [BBI, Proposition 7.4.11] for the proof. We use this relation to prove the convergence of uniformized spaces and boundaries.

We next review the definitions of uniform spaces, uniformization, and quasihyperbolization. These definitions can also be found in [BHK].

Definition 2.17. (*A*-uniform curve) Let $A > 0$ and a metric space (Ω, d) be given. We say that a curve $\gamma : [0, 1] \rightarrow \Omega$ is an *A-uniform curve* if the curve γ satisfies

1. $l_d(\gamma) \leq Ad(x, y)$,
2. $l_d(\gamma|_{[0,t]}) \wedge l_d(\gamma|_{[t,1]}) \leq A \operatorname{dist}(\gamma(t), \partial\Omega)$ for every $t \in [0, 1]$.

Remark 2.18. This definition does not depend on the parametrization of a curve γ .

Definition 2.19. (*A*-uniform space) A locally compact, rectifiably connected noncomplete metric space (Ω, d) is called an *A-uniform space* if every pair of points in Ω can be connected by an *A-uniform curve*.

Definition 2.20. (Uniformization) Let (X, d) be a δ -Gromov hyperbolic space. Fix a base point $p \in X$ and $\epsilon > 0$. The *uniformization metric* d_ϵ is defined by

$$d_\epsilon(x, y) := \inf_{\gamma} \int_0^{l_d(\gamma)} e^{-\epsilon d(p, \gamma(t))} dt$$

where infimum is taken over all arc-length parametrized curves γ from x to y . The metric space (X, d_ϵ) is called a *uniformized space*. We denote this uniformized space (X, d_ϵ) by (X^ϵ, d_ϵ) . We note that a *Harnack type inequality*

$$e^{-\epsilon d(x, y)} \leq \frac{e^{-\epsilon d(p, x)}}{e^{-\epsilon d(p, y)}} \leq e^{\epsilon d(x, y)} \quad (2)$$

holds for every $x, y \in X$, see [BHK, Chapter 5]

Remark 2.21. As mentioned in the previous section, it has been shown in [BHK, Proposition 4.5 and Chapter 5] that there exists $\epsilon_0(\delta) > 0$ such that for each $0 < \epsilon \leq \epsilon_0(\delta)$, the uniformized space (X^ϵ, d_ϵ) is a uniform space. Throughout this paper, we fix the constant $\epsilon_0(\delta)$ and use it in the rest of this paper.

Definition 2.22. (Quasihyperbolization) Let (Ω, d) be an *A-uniform space*. The *quasihyperbolic metric* k is defined by

$$k(x, y) := \inf_{\gamma} \int_0^{l_d(\gamma)} \frac{1}{d(\gamma(t))} dt$$

where infimum is taken over all arc-length parametrized curves γ from x to y , and $d(\cdot) := \operatorname{dist}(\cdot, \partial\Omega)$.

By [BHK, Theorem 3.6], if (Ω, d) is a uniform space, then (Ω, k) is a proper geodesic δ -Gromov hyperbolic space for some $\delta = \delta(A)$. Moreover, if (Ω, d) is bounded, then (Ω, k) is M -roughly starlike for some $M = M(A)$.

In order to prove the boundary convergence in the next section, we briefly review the connection between the Gromov boundary and the metric boundary $(\partial_{d_\epsilon} X^\epsilon, d_\epsilon)$. We refer the interested reader to [BHK] for a detailed discussion and [V2] and [BH] for Gromov boundary.

Definition 2.23. (Gromov boundary) We say that for a fixed point $p \in X$, a sequence $(x_n)_n$ is a *Gromov sequence* if $(x_i|x_j)_p \rightarrow \infty$ as $i, j \rightarrow \infty$. Also we say two Gromov sequences $(x_n)_n$ and $(y_n)_n$ are equivalent if $(x_n|y_n)_p \rightarrow \infty$ as $n \rightarrow \infty$. We write $(x_n)_n \sim (y_n)_n$ if they are equivalent. Since the space X is δ -Gromov hyperbolic, this is an equivalence relation. Hence we can consider the quotient space of the set of all Gromov sequences, denoted by $\partial_G X$. We call $\partial_G X$ the *Gromov boundary*.

Let (X, d) be a δ -Gromov hyperbolic space and $p \in X$ be a fixed base point. The Gromov product is extended to $X \cup \partial_G X$. We refer the reader to [BH] and [V1] for more discussion on extension of the definition of Gromov product to $X \cup \partial_G X$.

Definition 2.24. (The metric on $\partial_G X$) Let (X, d) be a δ -Gromov hyperbolic space. Let $p \in X$ be a fixed point and $0 < \epsilon < 1/(5\delta)$. For $x, y \in \partial_G X$, define

$$\rho_{p,\epsilon}(x, y) := e^{-\epsilon(x|y)_p}. \quad (3)$$

We set the metric

$$d_{p,\epsilon}(x, y) := \inf \sum_{i=1}^n \rho_{p,\epsilon}(a_{i-1}, a_i) \quad (4)$$

where infimum is taken over all finite sequences $(a_i)_{i=0}^n \subseteq X \cup \partial_G X$ with $a_0 = x$ and $a_n = y$.

Remark 2.25. There is another characterization of the Gromov boundary using geodesic rays. We say that two geodesic rays γ and $\tilde{\gamma}$ in X with $\gamma(0) = \tilde{\gamma}(0) = p$ are equivalent if $\sup_{t \geq 0} d(\gamma(t), \tilde{\gamma}(t))$ is finite. Then we can consider a quotient space of a set of all geodesic rays emanating from $p \in X$ by the above equivalence relation, denoted by $\partial_r X$. By [BH, Lemma 3.13], if a metric space X is proper geodesic, then the map $\partial_r X \ni [\gamma] \mapsto [(\gamma(n))_n] \in \partial_G X$ is a bijective canonical identification between the Gromov boundary $\partial_G X$ and $\partial_r X$. Moreover, by [BHK, Lemma 4.10, Theorem 4.13], we know that the map $\partial_G X \ni [(\gamma(n))_n] \mapsto \lim_{n \rightarrow \infty} \gamma(n) \in \partial_{d_\epsilon} X^\epsilon$ is well-defined and is a canonical bilipschitz map for $0 < \epsilon \leq \epsilon_0(\delta)$ where $\epsilon_0(\delta)$ is a constant defined in [BHK, Chapter 5] and $\partial_{d_\epsilon} X^\epsilon$ is the metric boundary defined by $\overline{X^\epsilon} \setminus X^\epsilon$. This fact plays a key role to prove the convergence of boundary.

3 The notion of Gromov-Hausdorff distance with boundary

In this section, we define the notion of Gromov-Hausdorff distance with boundary, denoted by d_{GHB} , and examine basic properties of d_{GHB} . We omit some proofs of the propositions provided below if there is no change in their proofs in comparison to the analogs of the Gromov-Hausdorff distance found in [HKST] and [BBI].

Definition 3.1. (Hausdorff distance with boundary) Let (X, d) be a metric space and $\mathcal{M}(X)$ denote the set of all noncomplete precompact locally compact subsets of X . For $A, B \in \mathcal{M}(X)$, we set the *Hausdorff distance with boundary* by

$$d_{HB}(A, B) := d_H^X(A, B) + d_H^X(\partial A, \partial B)$$

where $\partial A := \overline{A} \setminus A$ and $\partial B := \overline{B} \setminus B$ are the boundaries of A and B respectively. One can check that this d_{HB} is a metric on $\mathcal{M}(X)$.

For a given metric space X , denote the metric boundary by $\partial X := \overline{X} \setminus X$ where \overline{X} is the completion of X . We now define the Gromov-Hausdorff distance with boundary.

Definition 3.2 (Gromov-Hausdorff metric with boundary). Let X and Y be noncomplete precompact metric spaces. Set

$$d_{GHB}(X, Y) := \inf d_H^Z(\varphi(X), \psi(Y)) + d_H^Z(\varphi(\partial X), \psi(\partial Y))$$

where infimum is taken over all metric spaces Z and isometric embeddings $\varphi : \overline{X} \rightarrow Z$ and $\psi : \overline{Y} \rightarrow Z$.

In this section, unless otherwise stated, we always assume that metric spaces are all noncomplete and precompact. We first record the following elementary yet useful remarks.

Remark 3.3. If a map f is an isometry from X to Y , then both the canonical extension $\tilde{f} : \overline{X} \rightarrow \overline{Y}$ and the restriction $\tilde{f}|_{\partial X} : \partial X \rightarrow \partial Y$ are isometries.

Remark 3.4. If there exists an isometry $f : X \rightarrow Y$, then $d_{GHB}(X, Y) = 0$.

Definition 3.5. Let X, Y be metric spaces. We say that $X \sqcup Y := (X \times \{0\}) \cup (Y \times \{1\})$ is a *disjoint union* of the sets X and Y . The canonical maps $\iota_X : X \hookrightarrow X \sqcup Y$ and $\iota_Y : Y \hookrightarrow X \sqcup Y$ are defined by

$$\iota_X(x) = (x, 0) \quad \text{and} \quad \iota_Y(y) = (y, 1)$$

for each $x \in X$ and $y \in Y$.

Definition 3.6. (Admissible metric) Let X, Y be metric spaces. We say a metric d on $X \sqcup Y$ is *admissible* if the canonical maps $\iota_X : X \hookrightarrow X \sqcup Y$ and $\iota_Y : Y \hookrightarrow X \sqcup Y$ are isometric embeddings with respect to the metric d .

We only give a sketch of the proof of the following lemma since the proof is identical with the case of the Gromov-Hausdorff distance. See [BBI, Remark 7.3.12 and Proposition 7.3.16] and [HKST, Proposition 11.1.9].

Lemma 3.7. *Let X and Y be noncomplete precompact metric spaces. Then*

$$d_{GHB}(X, Y) = \inf_d (d_H^d(\iota_{\overline{X}}(X), \iota_{\overline{Y}}(Y)) + d_H^d(\iota_{\overline{X}}(\partial X), \iota_{\overline{Y}}(\partial Y))) =: \tilde{d}_{GHB}(X, Y),$$

where the infimum is taken over all admissible metrics on $\overline{X} \sqcup \overline{Y}$ and d_H^d is the Hausdorff distance with respect to the metric d on $\overline{X} \sqcup \overline{Y}$.

Proof. Since $d_{GHB}(X, Y) \leq \tilde{d}_{GHB}(X, Y)$ is obvious, it suffices to show that $\tilde{d}_{GHB}(X, Y) \leq d_{GHB}(X, Y)$. For any $\epsilon > 0$, choose a metric space Z and the isometric embeddings $\varphi : \overline{X} \rightarrow Z$ and $\psi : \overline{Y} \rightarrow Z$ such that

$$d_H^Z(\varphi(X), \psi(Y)) + d_H^Z(\varphi(\partial X), \psi(\partial Y)) < d_{GHB}(X, Y) + \epsilon \quad (5)$$

By considering $Z' := Z \times [0, 1]$ with the product metric, $\varphi(\overline{X})' := \varphi(\overline{X}) \times \{0\}$, and $\psi(\overline{Y})' := \psi(\overline{Y}) \times \{\epsilon\}$ if needed, we may assume that $\varphi(\overline{X})$ and $\psi(\overline{Y})$ are disjoint. Define d on $\overline{X} \sqcup \overline{Y}$ by

$$d(x, y) := \begin{cases} d_Z(\varphi(x), \varphi(y)) & \text{if } x, y \in \overline{X} \\ d_Z(\varphi(x), \psi(y)) & \text{if } x \in \overline{X}, y \in \overline{Y} \\ d_Z(\psi(x), \psi(y)) & \text{if } x, y \in \overline{Y} \end{cases}$$

This defines a metric on $\overline{X} \sqcup \overline{Y}$ due to the fact that $\varphi(\overline{X})$ and $\psi(\overline{Y})$ are disjoint. This implies that

$$\tilde{d}_{GHB}(X, Y) \leq d_{GHB}(X, Y) + \epsilon.$$

This completes the proof. □

Through Lemma 3.7, triangle inequality with respect to d_{GHB} is derived. Since the proof is a simple modification of the case of d_{GH} , we omit the proof. See the argument to prove triangle inequality in case of the Gromov-Hausdorff distance in [BBI] and [HKST].

Lemma 3.8. *Let X, Y, Z be noncomplete precompact metric spaces. Then*

$$d_{GHB}(X, Z) \leq d_{GHB}(X, Y) + d_{GHB}(Y, Z).$$

Under the additional assumption that given spaces are locally compact, we get the following equivalence.

Lemma 3.9. *Let X, Y be noncomplete precompact locally compact spaces. Then X is isometric to Y if and only if $d_{GHB}(X, Y) = 0$.*

Proof. We follow the case of the Gromov-Hausdorff distance, but with a modification. By Remark 3.4, if X is isometric to Y , then $d_{GHB}(X, Y) = 0$. Now suppose $d_{GHB}(X, Y) = 0$. By Lemma 3.7, for each $k \in \mathcal{N}$, there exist metrics d_k on $\overline{X} \sqcup \overline{Y}$ such that

$$d_H^{d_k}(\iota_{\overline{X}}(X), \iota_{\overline{Y}}(Y)) + d_H^{d_k}(\iota_{\overline{X}}(\partial X), \iota_{\overline{Y}}(\partial Y)) < 1/k$$

where $d_H^{d_k}$ is the Hausdorff distance on the metric space $(\overline{X} \sqcup \overline{Y}, d_k)$. Then there exist maps (possibly not continuous) $I_k : \overline{X} \rightarrow \overline{Y}$ and $J_k : \overline{Y} \rightarrow \overline{X}$ such that for all $x \in \overline{X}$ and $y \in \overline{Y}$,

$$d_k(\iota_{\overline{X}}(x), \iota_{\overline{Y}}(I_k(x))) < 1/k \quad \text{and} \quad d_k(\iota_{\overline{Y}}(y), \iota_{\overline{X}}(J_k(y))) < 1/k. \quad (6)$$

Since \overline{X} and \overline{Y} are separable, we can take $S_X := (x_n)_n \subseteq \overline{X}$ and $S_Y := (y_n)_n \subseteq \overline{Y}$ that are dense in \overline{X} and \overline{Y} respectively. For each $x \in S_X$, $(I_k(x))_k \in \overline{Y}$. The compactness of \overline{Y} allows us to pick a subsequence $(I_{k_l}(x))_l$ such that $I_{k_l}(x) \rightarrow y_x \in \overline{Y}$ in d_Y . Doing the same argument for other elements in S_X for the subsequence $(I_{k_l}(x))_l$ and by the diagonal argument, we can choose $(I_{k_l})_l$ such that $I_{k_l}(x) \rightarrow y_x$ in d_Y for each $x \in S_X$. Define $I : S_X \rightarrow \overline{Y}$ by $S_X \ni x \mapsto y_x \in \overline{Y}$. Since d_{k_l} is admissible, for every $x, y \in S_X$ we have

$$\begin{aligned} d_Y(I(x), I(y)) &= \lim_{l \rightarrow \infty} d_Y(I_{k_l}(x), I_{k_l}(y)) \\ &= \lim_{l \rightarrow \infty} d_{k_l}(\iota_{\overline{Y}}(I_{k_l}(x)), \iota_{\overline{Y}}(I_{k_l}(y))) \\ &= d_X(x, y), \end{aligned}$$

where the last equality holds due to (6). Hence we have the canonical extension of I , denoted by $\tilde{I} : \overline{X} \rightarrow \overline{Y}$. Note that \tilde{I} is a distance preserving map. By exactly the same argument for $J_k : \overline{Y} \rightarrow \overline{X}$, we get $\tilde{J} : \overline{Y} \rightarrow \overline{X}$. Since $\tilde{J} \circ \tilde{I}$ and $\tilde{I} \circ \tilde{J}$ are distance preserving self-maps of \overline{X} and \overline{Y} respectively, both \tilde{I} and \tilde{J} are bijective isometries. We will prove that $\tilde{I}(x) \in Y$ for every $x \in X$. To prove this, suppose that there exists $x \in X$ such that $\tilde{I}(x) \in \partial Y$. Set $\alpha := \text{dist}(x, \partial X) > 0$. Note that α is positive due to the local compactness of the space X . Let $\epsilon := \alpha/4 > 0$ be given. Then there exists $x_\epsilon \in S_X$ such that

$$d_X(x_\epsilon, x) = d_Y(\tilde{I}(x_\epsilon), \tilde{I}(x)) < \epsilon.$$

Hence $\text{dist}(x_\epsilon, \partial X) \geq 3\alpha/4$ and $\text{dist}(\tilde{I}(x_\epsilon), \partial Y) \leq \alpha/4$. For this x_ϵ , there exists $N \in \mathbb{N}$ such that for each $k_l \geq N$, we have

$$d_Y(I_{k_l}(x_\epsilon), \tilde{I}(x_\epsilon)) = d_{k_l}(\iota_{\overline{Y}}(I_{k_l}(x_\epsilon)), \iota_{\overline{Y}}(\tilde{I}(x_\epsilon))) < \epsilon.$$

Also, by the construction of I_{n_k} , we know that

$$d_{k_l}(\iota_{\overline{X}}(x_\epsilon), \iota_{\overline{Y}}(I_{k_l}(x_\epsilon))) < \frac{1}{k_l}.$$

Combining these inequalities, for $k_l \geq N$, we have

$$\text{dist}_Y(\tilde{I}(x_\epsilon), \partial Y) \geq \text{dist}_Y(I_{k_l}(x_\epsilon), \partial Y) - \epsilon$$

$$\begin{aligned}
&= \text{dist}_{d_{k_l}}(\iota_{\overline{Y}}(I_{k_l}(x_\epsilon)), \iota_{\overline{Y}}(\partial Y)) - \epsilon \\
&\geq \text{dist}_{d_{k_l}}(\iota_{\overline{Y}}(I_{k_l}(x_\epsilon)), \iota_{\overline{X}}(\partial X)) - \frac{1}{k_l} - \epsilon \\
&\geq \text{dist}_{d_{k_l}}(\iota_{\overline{X}}(x_\epsilon), \iota_{\overline{X}}(\partial X)) - \frac{2}{k_l} - \epsilon \\
&\geq \text{dist}_X(x_\epsilon, \partial X) - \frac{2}{k_l} - \epsilon \\
&\geq \alpha/2 - \frac{2}{k_l}, \tag{7}
\end{aligned}$$

where $\text{dist}_{d_{n_k}}$ represents the distance between a point and a set with respect to the metric d_{n_k} . Since $\text{dist}_Y(\tilde{I}(x_\epsilon), \partial Y) \leq \alpha/4$, by letting $l \rightarrow \infty$ in the inequality (7), we get a contradiction. Hence $\tilde{I}|_X : X \rightarrow Y$ is well-defined. By the similar argument we did above, we can also prove that $\tilde{I}(x) \in \partial Y$ for every $x \in \partial X$. This implies that $\tilde{I}|_X : X \rightarrow Y$ is bijective. This completes the proof. \square

Remark 3.10. As we remarked in Section 1, it is immediate that for noncomplete precompact metric spaces X and Y , we have

$$d_{GH}(\overline{X}, \overline{Y}) + d_{GH}(\partial X, \partial Y) \leq d_{GHB}(X, Y).$$

On the other hand, the following example shows that the the converse inequality does not hold. Set

$$X := [0, 1] \times [0, 1] \setminus \{(1/2, 0), (1/2, 1)\}$$

and

$$Y := [0, 1] \times [0, 1] \setminus \{(1, 0), (1, 1)\}.$$

where the standard Euclidean metric is defined on both X and Y . Then $d_{GH}(\overline{X}, \overline{Y}) = d_{GH}(\partial X, \partial Y) = 0$ but it follows that $d_{GHB}(X, Y) > 0$ since if $d_{GHB}(X, Y) = 0$, then by Lemma 3.9 there exists an isometry between X and Y . Under this isometry, the line

$$\{(1/2, t) : 0 \leq t \leq 1\} \subseteq \overline{X}$$

should get mapped to

$$\{(1, t) : 0 \leq t \leq 1\} \subseteq \overline{Y},$$

which is impossible by looking at the connected components.

Set a class of metric spaces \mathcal{M} by

$$\mathcal{M} := \{X \mid X \text{ is a noncomplete, precompact, and locally compact metric space.}\} / \sim.$$

where the equivalence relation \sim between X and Y is defined by existence of an isometry between X and Y . Then combining Lemma 3.8 and Lemma 3.9, we get the following.

Theorem 3.11 (First assertion of Theorem 1.1). *The space \mathcal{M} is a metric space with the metric d_{GHB} .*

Remark 3.12. The assumption that metric spaces are locally compact in Theorem 1.1 is essential in order to make d_{GHB} a metric. One can see this by considering

$$X := [0, 1] \cap \mathbb{Q} \quad \text{and} \quad Y := [0, 1] \cap (\mathbb{R} \setminus \mathbb{Q}).$$

Note that $d_{GHB}(X, Y) = 0$ but there is no way to construct a bijective map between them.

Remark 3.13. Note that \mathcal{M} is not a complete metric space. Consider a sequence of metric spaces

$$X_n := (0, 1) \times (0, 1) \setminus \bigcup_{k=1}^{n-1} [1/2, 1) \times \{k/n\}.$$

It is easy to verify that X_n with the Euclidean metric are noncomplete precompact locally compact metric spaces. Moreover, $(X_n)_n$ is Cauchy in d_{GHB} . If \mathcal{M} is complete, we can find a limit X such that $\bar{X} := [0, 1] \times [0, 1]$, and $d_{GHB}(X_n, X) \rightarrow 0$ as $n \rightarrow \infty$. However, ∂X_n converges to

$$\partial X = [0, 1] \times [0, 1] \setminus (0, 1/2) \times (0, 1),$$

which is not the boundary of the metric space X .

Here, in order to take the boundary into account, we define the notion of correspondence with boundary.

Definition 3.14. (Correspondence) Let X, Y be metric spaces. A *correspondence between X and Y with boundary* is a subset $R \subseteq \bar{X} \times \bar{Y}$ such that $\pi_X(R \cap (X \times Y)) = X$, $\pi_Y(R \cap (X \times Y)) = Y$, $\pi_X(R \cap (\partial X \times \partial Y)) = \partial X$, and $\pi_Y(R \cap (\partial X \times \partial Y)) = \partial Y$ where $\pi_X : \bar{X} \times \bar{Y} \rightarrow \bar{X}$ and $\pi_Y : \bar{X} \times \bar{Y} \rightarrow \bar{Y}$ are the canonical projections.

Definition 3.15. (Distortion) Let X, Y be metric spaces and R be a correspondence between X and Y with boundary. A *distortion of R* is defined by

$$\text{dis}(R) := \sup_{(x,y),(x',y') \in R} |d_X(x, x') - d_Y(y, y')|.$$

Remark 3.16. One can check that if $\text{dis}(R) = 0$ for a given correspondence with boundary R , then it is not hard to show that there exists an isometry $f : \bar{X} \rightarrow \bar{Y}$ such that $f|_X : X \rightarrow Y$ and $f|_{\partial X} : \partial X \rightarrow \partial Y$.

Proposition 3.17. *Let X, Y be noncomplete precompact locally compact metric spaces. Then*

$$\frac{1}{2} d_{GHB}(X, Y) \leq \frac{1}{2} \inf_R \text{dis}(R) \leq d_{GHB}(X, Y),$$

where the infimum is taken over all correspondences with boundary $R \subseteq X \times Y$.

Proof. We will first prove that

$$d_{GHB}(X, Y) \geq \frac{1}{2} \inf_R \text{dis}(R).$$

For each $r > d_{GHB}(X, Y)$, there exist a metric space Z and isometric embeddings $\varphi : \overline{X} \rightarrow Z$ and $\psi : \overline{Y} \rightarrow Z$ such that

$$d_H^Z(\varphi(X), \psi(Y)) + d_H^Z(\varphi(\partial X), \psi(\partial Y)) < r,$$

where d_H^Z is the Hausdorff distance in Z . Set $s > d_H^Z(\varphi(X), \psi(Y))$ and $t > d_H^Z(\varphi(\partial X), \psi(\partial Y))$ such that $s + t, r$. Define a correspondence with boundary

$$R := \left\{ (x, y) \in \overline{X} \times \overline{Y} \left| \begin{array}{l} d_Z(\varphi(x), \psi(y)) \leq s, x \in X, \text{ and } y \in Y \\ \text{or} \\ d_Z(\varphi(x), \psi(y)) \leq t, x \in \partial X, \text{ and } y \in \partial Y. \end{array} \right. \right\}.$$

If $(x, y), (x', y') \in R \cap (X \times Y)$, then

$$\begin{aligned} |d_X(x, x') - d_Y(y, y')| &\leq |d_Z(\varphi(x), \varphi(x')) - d_Z(\varphi(x), \psi(y'))| \\ &\quad + |d_Z(\varphi(x), \psi(y')) - d_Z(\psi(y), \psi(y'))| \\ &\leq d_Z(\varphi(x'), \psi(y')) + d_Z(\varphi(x), \psi(y)) \\ &\leq 2s, \end{aligned}$$

Similarly, if $(x, y), (x', y') \in R \cap (\partial X \times \partial Y)$, then

$$|d_X(x, x') - d_Y(y, y')| \leq 2t,$$

Also, if $(x, y) \in R \cap (X \times Y)$ and $(x', y') \in R \cap (\partial X \times \partial Y)$, then

$$|d_X(x, x') - d_Y(y, y')| \leq s + t.$$

Therefore, we conclude that

$$\frac{1}{2} \inf_R \text{dis}(R) \leq s + t < r.$$

Letting $r \rightarrow d_{GHB}(X, Y)$, the claim is proved. Next we will prove

$$d_{GHB}(X, Y) \leq \inf_R \text{dis}(R).$$

Let R be a correspondence with boundary and $r := \frac{1}{2} \text{dis}(R)$. If $r = 0$, by Remark 3.16, we have an isometry $f : X \rightarrow Y$, which implies $d_{GHB}(X, Y) = 0$ by Lemma 3.7. Hence we assume $r > 0$. Define an admissible metric d on $\overline{X} \sqcup \overline{Y}$ by

$$d(\iota_{\overline{X}}(x), \iota_{\overline{Y}}(y)) := \begin{cases} d_X(x, y) & \text{if } x, y \in \overline{X} \\ \inf_{(x', y') \in R} (d_X(x, x') + d_Y(y, y')) + r & \text{if } x \in \overline{X}, y \in \overline{Y} \\ \inf_{(x', y') \in R} (d_X(y, x') + d_Y(x, y')) + r & \text{if } x \in \overline{Y}, y \in \overline{X} \\ d_Y(x, y) & \text{if } x, y \in \overline{Y}. \end{cases}$$

One can check that d satisfies triangle inequality due to the choice of r . We claim that

$$d_{\overline{X} \sqcup \overline{Y}}^H(\iota_{\overline{X}}(X), \iota_{\overline{Y}}(Y)) \leq r.$$

For each $y \in Y$, there exists $x \in X$ such that $(x, y) \in R$ since $\pi_Y(R \cap (X \times Y)) = Y$. Then

$$d(\iota_{\overline{X}}(x), \iota_{\overline{Y}}(y)) = \inf_{(x', y') \in R} (d_X(x, x') + d_Y(y, y')) + r \leq d_X(x, x) + d_Y(y, y) + r = r.$$

By the same argument, for every $x \in X$, we can find $y \in Y$ such that $d(x, y) \leq r$, which implies that $d_{\overline{X} \sqcup \overline{Y}}^H(X, Y) \leq r$. Similarly, we have

$$d_{\overline{X} \sqcup \overline{Y}}^H(\iota_{\overline{X}}(\partial X), \iota_{\overline{Y}}(\partial Y)) \leq r.$$

Hence we get

$$\frac{1}{2}d_{GHB}(X, Y) \leq \frac{1}{2}(d_{\overline{X} \sqcup \overline{Y}}^H(\iota_{\overline{X}}(X), \iota_{\overline{Y}}(Y)) + d_{\overline{X} \sqcup \overline{Y}}^H(\iota_{\overline{X}}(\partial X), \iota_{\overline{Y}}(\partial Y))) \leq r.$$

We conclude that

$$\frac{1}{2}d_{GHB}(X, Y) \leq \frac{1}{2} \inf_R \text{dis}(R).$$

□

We introduce a notion of ϵ -isometry with boundary.

Definition 3.18. (ϵ -isometry with boundary) A map $f : \overline{X} \rightarrow \overline{Y}$ is called an ϵ -isometry with boundary if

$$\text{dis}(f) := \sup_{x, x' \in \overline{X}} |d_Y(f(x), f(x')) - d_X(x, x')| \leq \epsilon,$$

and

$$Y \subseteq (f(X))_\epsilon, \quad \text{and} \quad \partial Y \subseteq (f(\partial X))_\epsilon.$$

Here is a simple generalization of the relationship between d_{GH} and ϵ -isometry. We omit the proof since there is no change in it in comparison to the case of the Gromov-Hasudorff distance. This proposition will be used to prove the convergence of uniformized spaces.

Proposition 3.19. *Let X, Y be noncomplete precompact locally compact metric spaces.*

1. *If $d_{GHB}(X, Y) < \epsilon$, then there exists a 2ϵ -isometry with boundary $f : \overline{X} \rightarrow \overline{Y}$.*
2. *If there exists an ϵ -isometry with boundary $f : \overline{X} \rightarrow \overline{Y}$, then $d_{GHB}(X, Y) \leq 3\epsilon$.*

Here we define the Gromov-Hausdorff convergence with boundary. As we will see later in Proposition 3.21, the convergence of spaces with respect to d_{GHB} is equivalent to the notion defined below.

Definition 3.20. (Gromov-Hausdorff convergence with boundary) Let $(X_n, d_n)_n, (X, d) \subseteq \mathcal{M}$. We say that $(X_n, d_n)_n$ is *Gromov-Hausdorff convergent to (X, d) with boundary* if there exist a metric space Z and isometric embeddings $\varphi_n : \overline{X}_n \rightarrow Z$ and $\varphi : \overline{X} \rightarrow Z$ such that

$$d_H^Z(\varphi_n(X_n), \varphi(X)) \rightarrow 0$$

and

$$d_H^Z(\varphi_n(\partial X_n), \varphi(\partial X)) \rightarrow 0$$

where d_H^Z is the Hausdorff distance on Z .

The following equivalence seems to be well-known in case of the Gromov-Hausdorff distance. Here we provide the proof for the reader's convenience.

Proposition 3.21. *Let $(X_n, d_n)_n, (X, d) \subseteq \mathcal{M}$. Then the following notions of convergence are equivalent :*

1. $(X_n, d_n)_n$ is *Gromov-Hausdorff convergent to (X, d) with boundary in Definition 3.20*
2. $d_{GHB}(X_n, X) \rightarrow 0$ as $n \rightarrow \infty$.

Proof. Although the proof is the same one as the case of d_{GH} , we would like to record the proof for the sake of completeness. It is enough to prove that $d_{GHB}(X_n, X) \rightarrow 0$ as $n \rightarrow \infty$ implies $(X_n, d_n)_n$ is Gromov-Hausdorff convergent to (X, d) with boundary in a sense of Definition 3.20. We would like to remark that in this proof, we identify \overline{X}_n and \overline{X} with the copies in $\overline{X}_n \sqcup \overline{X}$. Also, if $x \in \overline{X}$, then the corresponding element $\iota_{\overline{X}}(x)$ in $\overline{X}_n \sqcup \overline{X}$ by the canonical map $\iota_{\overline{X}} : \overline{X} \hookrightarrow \overline{X}_n \sqcup \overline{X}$ is still denoted by x . Same principle is applied to elements in \overline{X}_n .

Let $\epsilon_n := d_{GHB}(X_n, X) + 1/n$. Then By Lemma 3.7, there exist admissible metrics δ_n on $\overline{X}_n \sqcup \overline{X}$ such that

$$d_H^{\delta_n}(X_n, X) + d_H^{\delta_n}(\partial X_n, \partial X) \leq \epsilon_n$$

where $d_H^{\delta_n}$ is the Hausdorff distance on $(X_n \sqcup X, \delta_n)$. Set $Z := \overline{X} \sqcup \overline{X}_1 \sqcup \cdots \sqcup \overline{X}_n \sqcup \cdots := \overline{X} \times \{0\} \cup (\bigcup_{k=1}^{\infty} \overline{X}_k \times \{k\})$. We will define a metric d_Z on Z by

$$d_Z(x, y) := \begin{cases} \delta_n(x, y) & \text{if } x, y \in \overline{X} \cup \overline{X}_n \\ \inf_{w \in \overline{X}} (\delta_n(x, w) + \delta_m(w, y)) & \text{if } x \in \overline{X}_n, y \in \overline{X}_m \text{ with } n \neq m. \end{cases}$$

Then we claim that this is a metric. Suppose $\delta(x, y) = 0$. We only consider the case where $x \in \overline{X}_n$ and $y \in \overline{X}_m$. Take $(w_k)_k \subseteq \overline{X}$ such that

$$\delta_n(x, w_k) + \delta_m(w_k, y) \rightarrow 0,$$

as $k \rightarrow \infty$. This implies that $x \in \overline{X}$ and $y \in \overline{X}$, which is not possible. Since the symmetry of this metric is obvious, it suffices to show that triangle inequality holds. We check cases $x \in \overline{X}_n, y \in \overline{X}_m, z \in \overline{X}$ with $n \neq m$ and $x \in \overline{X}_n, y \in \overline{X}_m, z \in \overline{X}_l$ with n, m, l not all distinct since all other cases are straightforward.

(Case 1) $x \in \overline{X}_n, y \in \overline{X}_m$, and $z \in \overline{X}$. Since \overline{X} is compact, we can pick a point $w \in \overline{X}$ such that

$$d_Z(x, y) = \delta_n(x, w) + \delta_m(w, y).$$

Then

$$\begin{aligned} d_Z(x, y) + d_Z(y, z) &= \delta_n(x, w) + \delta_m(w, y) + \delta_m(y, z) \\ &\geq \delta_n(x, w) + \delta_m(w, z) = d_Z(x, z) \end{aligned}$$

where we used the fact that $\delta_m(w, z) = \delta_n(w, z)$ for every $w, z \in \overline{X}$.

(Case 2) $x \in \overline{X}_n, y \in \overline{X}_m$, and $z \in \overline{X}_l$. Take $w \in \overline{X}$ and $\bar{w} \in \overline{X}$ such that

$$d_Z(x, y) = \delta_n(x, w) + \delta_m(w, y)$$

and

$$d_Z(y, z) = \delta_m(y, \bar{w}) + \delta_l(\bar{w}, z).$$

Then,

$$\begin{aligned} d_Z(x, z) &\leq \delta_n(x, \bar{w}) + \delta_l(\bar{w}, z) \\ &\leq \delta_n(x, w) + \delta_n(w, \bar{w}) + \delta_l(\bar{w}, z) \\ &= \delta_n(x, w) + \delta_m(w, \bar{w}) + \delta_l(\bar{w}, z) \\ &\leq \delta_n(x, w) + \delta_m(w, y) + \delta_m(y, \bar{w}) + \delta_l(\bar{w}, z) \\ &= d_Z(x, y) + d_Z(y, z). \end{aligned}$$

Hence d_Z is a metric on Z . By taking a metric space (Z, d_Z) , $\varphi_n := \iota_{\overline{X}_n} : \overline{X}_n \hookrightarrow Z$, and $\varphi := \iota_{\overline{X}} : \overline{X} \hookrightarrow Z$, we have

$$d_H^Z(\varphi_n(X_n), \varphi(X)) + d_H^Z(\varphi_n(\partial X_n), \varphi(\partial X)) \leq \epsilon_n \rightarrow 0$$

as $n \rightarrow \infty$. This completes the proof. \square

4 An application to a class of bounded A -uniform spaces

In this section we will present an application of the Gromov-Hausdorff distance with boundary d_{GHB} to a class of bounded A -uniform spaces and prove Theorem 1.1. Here we will first prove the stability of A -uniform spaces with respect to d_{GHB} . Recall from Definition 2.19 that a noncomplete locally compact metric space (Ω, d) is called an A -uniform space if for each pair of points $x, y \in \Omega$ there exists a curve γ from x to y such that

1. $l_d(\gamma) \leq Ad(x, y)$,
2. $l_d(\gamma|_{[0,t]}) \wedge l_d(\gamma|_{[t,1]}) \leq A \operatorname{dist}(\gamma(t), \partial\Omega)$ for every $t \in [0, 1]$.

Before stating the next lemma, we would like to remark that by [BHK, Proposition 2.20], every bounded A -uniform space is precompact. Hence the isometry class of a bounded A -uniform space is in \mathcal{M} .

Lemma 4.1. *Let $(\Omega_n, d_n)_n \subseteq \mathcal{M}$ be a sequence of bounded A -uniform spaces and $(\Omega, d) \in \mathcal{M}$. Suppose $(\Omega_n, d_n)_n$ Gromov-Hausdorff converges to (Ω, d) with boundary. Then (Ω, d) is also an A -uniform space.*

Proof. First note that in this proof we do not take the arc-length parametrization for A -uniform curves γ' . We always take the parametrization set by the following procedure. First consider the arc-length parametrization $\gamma' : [0, l_d(\gamma')] \rightarrow \Omega$ and define $\alpha(t) := l_d(\gamma')t$ for every $t \in [0, 1]$. Then we set $\gamma := \gamma' \circ \alpha : [0, 1] \rightarrow \Omega$, which is the curve considered in this proof. By Proposition 3.21, we may assume that $(\Omega_n)_n$ and Ω are all subsets of the same space (Z, d) . Recall again that Ω_n and Ω need not be open in (Z, d) . Set $\epsilon_n > d_H^Z(\Omega_n, \Omega)$. Let $x, y \in \Omega$ be given. Take $x_n, y_n \in \Omega_n$ such that

$$d(x, x_n) \leq \epsilon_n \quad \text{and} \quad d(y, y_n) \leq \epsilon_n.$$

Take A -uniform curves $\gamma_n : [0, 1] \rightarrow \Omega_n$ from x_n to y_n parametrized as described above. We record the following inequality for the later use. For each $t, t' \in [0, 1]$, we get

$$\begin{aligned} d(\gamma_n(t), \gamma_n(t')) &\leq l_{d_n}(\gamma_n|[t, t']) \\ &= l_{d_n}(\gamma_n)|t - t'| \\ &\leq Ad(x_n, y_n)|t - t'| \\ &\leq A(d(x, y) + 2\epsilon_n)|t - t'|. \end{aligned} \tag{8}$$

Let $t_1 \in [0, 1] \cap \mathbb{Q}$ and $n \in \mathbb{N}$ be given. Then there exists $y_n(t_1) \in \Omega$ such that $d(y_n(t_1), \gamma_n(t_1)) \leq \epsilon_n$. Since we know that $\overline{\Omega}$ is compact, we can take a subsequence $(y_{n_k}(t))_k$ convergent to some $y_t \in \overline{\Omega}$. By a standard diagonalization argument we may assume that for every $t \in [0, 1] \cap \mathbb{Q}$, there exists $y_t \in \overline{\Omega}$ such that $d(y_t, \gamma_{n_k}(t)) \rightarrow 0$ as $k \rightarrow \infty$. Note that, by triangle inequality and (8),

$$\begin{aligned} d(y_t, y_{t'}) &\leq d(y_t, y_{n_k}(t)) + d(y_{n_k}(t), \gamma_{n_k}(t)) + d(\gamma_{n_k}(t), \gamma_{n_k}(t')) \\ &\quad + d(\gamma_{n_k}(t'), y_{n_k}(t')) + d(y_{n_k}(t'), y_{t'}) \\ &\leq 2\epsilon_{n_k} + d(y_t, y_{n_k}(t)) + d(\gamma_{n_k}(t), \gamma_{n_k}(t')) + d(y_{n_k}(t'), y_{t'}) \\ &\leq 2\epsilon_{n_k} + d(y_t, y_{n_k}(t)) + A(d(x, y) + 2\epsilon_n)|t - t'| + d(y_{n_k}(t'), y_{t'}) \end{aligned}$$

for every $t, t' \in [0, 1] \cap \mathbb{Q}$. By taking the limit as $k \rightarrow \infty$ in the above inequality, we get

$$d(y_t, y_{t'}) \leq Ad(x, y)|t - t'|,$$

which implies that the map $t \in [0, 1] \cap \mathbb{Q} \mapsto y_t \in \overline{\Omega}$ is $Ad(x, y)$ -lipschitz. Hence, by canonical extension to the interval $[0, 1]$, we have an $Ad(x, y)$ -lipschitz map $\gamma : [0, 1] \rightarrow \overline{\Omega}$ such that $\gamma(t) = y_t$ for every $t \in [0, 1] \cap \mathbb{Q}$. We also note that $\gamma_{n_k}(t) \rightarrow \gamma(t)$ as $k \rightarrow \infty$ for every $t \in [0, 1]$ since

$$\begin{aligned} \limsup_{k \rightarrow \infty} d(\gamma_{n_k}(t), \gamma(t)) &\leq \limsup_{k \rightarrow \infty} d(\gamma_{n_k}(t), \gamma_{n_k}(t_j)) + \limsup_{k \rightarrow \infty} d(\gamma_{n_k}(t_j), \gamma(t_j)) + d(\gamma(t_j), \gamma(t)) \\ &\leq 2Ad(x, y)|t - t_j| \end{aligned}$$

for $t_j \in [0, 1] \cap \mathbb{Q}$. Letting $t_j \rightarrow t$ shows that $\gamma_{n_k}(t) \rightarrow \gamma(t)$. We claim that the curve γ is A -uniform. Since γ is $Ad(x, y)$ -lipschitz, we know that $l_d(\gamma) \leq Ad(x, y)$. Next we will prove that for each $t \in [0, 1]$,

$$l_d(\gamma|_{[0,t]}) \wedge l_d(\gamma|_{[t,1]}) \leq A \operatorname{dist}(\gamma(t), \partial\Omega).$$

Since γ_{n_k} are all A -uniform curves, we know that

$$\begin{aligned} l_d(\gamma_{n_k})(t \wedge (1 - t)) &= l_{d_{n_k}}(\gamma_{n_k}|_{[0,t]}) \wedge l_{d_{n_k}}(\gamma_{n_k}|_{[t,1]}) \leq A \operatorname{dist}(\gamma_{n_k}(t), \partial\Omega_{n_k}) \\ &\leq A \operatorname{dist}(\gamma_{n_k}(t), \partial\Omega) + A\tilde{\epsilon}_{n_k} \end{aligned} \quad (9)$$

where $\tilde{\epsilon}_{n_k} := d_H^Z(\partial\Omega_{n_k}, \partial\Omega)$. Since length does not increase under uniform limits of curves, we get

$$\limsup_{k \rightarrow \infty} l_{d_{n_k}}(\gamma_{n_k}|_{[0,t]}) \wedge l_{d_{n_k}}(\gamma_{n_k}|_{[t,1]}) \leq A \operatorname{dist}(\gamma(t), \partial\Omega). \quad (10)$$

Since the curve γ is a uniform limit of $(\gamma_{n_k})_k$, we can get

$$l_d(\gamma|_{[0,t]}) \wedge l_d(\gamma|_{[t,1]}) \leq \limsup_{k \rightarrow \infty} l_{d_{n_k}}(\gamma_{n_k}|_{[0,t]}) \wedge l_{d_{n_k}}(\gamma_{n_k}|_{[t,1]}) \quad (11)$$

for each $t \in [0, 1]$. By inequalities (10) and (11), we conclude that

$$l_d(\gamma|_{[0,t]}) \wedge l_d(\gamma|_{[t,1]}) \leq A \operatorname{dist}(\gamma(t), \partial\Omega).$$

This tells us that even though initially the curve γ was defined in $\overline{\Omega}$, indeed γ is in Ω from the last inequality. This completes the proof. \square

Definition 4.2. Let (X, d) be a metric space and set

$$\mathcal{U}_X(A, R) := \{ \Omega \subseteq X \mid \Omega \text{ is a bounded } A\text{-uniform space and } \operatorname{diam}(\Omega) \geq R \}.$$

Remark 4.3. Since our focus is uniform spaces, we do not require $\Omega \subseteq X$ to be open in X .

Proposition 4.4. *Let (X, d) be a compact metric space and $A \geq 1$ and $R > 0$ be fixed. Then $(\mathcal{U}_X(A, R), d_{HB})$ is a compact metric space.*

Proof. First we prove that $\mathcal{U}_X(A, R)$ is complete with respect to d_{HB} . Let $(\Omega_n)_n$ be a Cauchy sequence in $(\mathcal{U}_X(A, R), d_{HB})$. Note that $\overline{\Omega}_n$ and $\partial\Omega_n$ are all compact. Since $(\overline{\Omega}_n)_n$ and $(\partial\Omega_n)_n$ are Cauchy sequences with respect to the Hausdorff distance d_H^X , there exist compact sets $S, T \subseteq X$ such that $d_H^X(\overline{\Omega}_n, S) \rightarrow 0$, $d_H^X(\partial\Omega_n, T) \rightarrow 0$ as $n \rightarrow \infty$ by [BBI, Theorem 7.3.8]. Note that $T \subseteq S$. Set $\Omega := S \setminus T$. We now prove that Ω is a nonempty set. Suppose $\Omega = \emptyset$. Then we have $S = T$. For each $n \in \mathbb{N}$, we can take a pair of points $x_n, y_n \in \Omega_n$ with $d(x_n, y_n) \geq R/2$ and an arc-length parametrized A -uniform curve $\gamma_n : [0, l_d(\gamma_n)] \rightarrow \Omega_n$ from x_n to y_n . Then, by A -uniformity, we get

$$\frac{R}{4} \leq \text{Adist}(\gamma_n(R/4), \partial\Omega_n) \rightarrow 0$$

as $n \rightarrow \infty$, which is a contradiction. Hence Ω is nonempty. We next claim that Ω is a dense subset of S . Suppose that there exists $x \in T$ such that $B_d(x, r) \cap S \subseteq T$ for some $r > 0$. We may assume that $r < R/2$. Set $\epsilon := \frac{r}{16A}$. Then there exists $N_\epsilon \in \mathbb{N}$ such that for $n \geq N_\epsilon$, we get $d_H^X(\Omega_n, S) + d_H^X(\partial\Omega_n, T) < \epsilon$. We claim that for every point $z_n \in B_d(x, r/2) \cap \Omega_n$, $\text{dist}(z_n, T) \leq \epsilon$. In fact, we can take $z' \in S$ such that $d(z', z_n) \leq \epsilon$. Also,

$$d(x, z') \leq d(x, z_n) + d(z_n, z') \leq r/2 + \epsilon < r.$$

Hence $z' \in B_d(x, r) \cap S \subseteq T$, which implies that

$$\text{dist}(z_n, T) \leq d(z_n, z') \leq \epsilon.$$

Take $x_n \in \Omega_n$ satisfying $d(x_n, x) < \epsilon$. Since $\text{diam}(\Omega_n) \geq R$ and $r < R/2$, we have $\Omega_n \neq B_d(x_n, r) \cap \Omega_n$. Hence we can pick $y_n \in \Omega_n$ such that $d(x_n, y_n) \geq r$. Let $\gamma_n : [0, l_d(\gamma_n)] \rightarrow \Omega_n$ be an A -uniform curve in Ω_n from x_n to y_n . Set $t := r/4$. Noting that $\gamma_n(t) \in B_d(x, r/2) \cap \Omega_n$ and the above claim, we get

$$t = l_d(\gamma_n|_{[0,t]}) \wedge l_d(\gamma_n|_{[t, l_d(\gamma_n)]}) \leq \text{Adist}(\gamma_n(t), \partial\Omega_n) \leq A(\text{dist}(\gamma_n(t), T) + \epsilon) \leq 2A\epsilon \leq \frac{r}{8},$$

which is a contradiction. Hence Ω is dense in S and noncomplete. Local compactness of Ω follows directly from the fact that $d(x, T) > 0$ for each $x \in \Omega$ and S and T are compact. From Lemma 4.1, we conclude that Ω is an A -uniform space. Therefore $\mathcal{U}_X(A, R)$ is complete. Regarding the compactness of $\mathcal{U}_X(A, R)$, since X is compact, for every sequence $(\Omega_n)_n$ from $\mathcal{U}_X(A, R)$ we can take out a subsequence $(\Omega_{n_k})_k$ such that both $(\overline{\Omega_{n_k}})_k$ and $(\partial\Omega_{n_k})_k$ are Cauchy in d_H , which means $(\Omega_{n_k})_k$ is Cauchy in d_{HB} . Since $\mathcal{U}_X(A, R)$ is complete, there exists a limit $\Omega \in \mathcal{U}_X(A, R)$ such that $d_{HB}(\Omega_{n_k}, \Omega) \rightarrow 0$ as $k \rightarrow \infty$. \square

Definition 4.5. Let $A \geq 1$ and $R > 0$. Define

$$\mathcal{U}(A, R) := \{[\Omega] \in \mathcal{M} \mid \Omega \text{ is a bounded } A\text{-uniform space and } \text{diam}(\Omega) \geq R\}$$

where $[\Omega]$ is an equivalence class of Ω .

Theorem 4.6 (Second assertion of Theorem 1.1). *The metric space $(\mathcal{U}(A, R), d_{GHB})$ is a complete metric space.*

Proof. Let $(\Omega_n)_n$ be a Cauchy sequence of uniform spaces with respect to d_{GHB} . Note that $(\overline{\Omega_n})_n$ is Cauchy with respect to d_{GH} and hence it is totally bounded. By Proposition 2.6 there exists a compact set $K \subseteq l^\infty$ such that every $\overline{\Omega_n}$ admits an isometric embedding into K . Since $\mathcal{U}_K(A, R)$ is compact by Proposition 4.4, there exists a subsequence $(\Omega_{n_k})_k$ and $\Omega \in \mathcal{U}_K(A, R)$ such that $\Omega_{n_k} \rightarrow \Omega$ in d_{HB} as $k \rightarrow \infty$, which implies $d_{GHB}(\Omega_{n_k}, \Omega) \rightarrow 0$ as $k \rightarrow \infty$. Since the original sequence is Cauchy, $d_{GHB}(\Omega_n, \Omega) \rightarrow 0$ as $n \rightarrow \infty$. \square

Theorem 4.7. *(Compactness theorem for uniform spaces) Let $(\Omega_n)_n \subseteq \mathcal{U}(A, R)$. If one of the conditions in Proposition 2.5 holds for $(\overline{\Omega_n})_n$, then there exists a subsequence $(\Omega_{n_k})_k$ such that $\Omega_{n_k} \rightarrow \Omega \in \mathcal{U}(A, R)$ as $k \rightarrow \infty$ with respect to d_{GHB} .*

Proof. Since $(\overline{\Omega_n})_n$ is totally bounded with respect to d_{GH} , by Proposition 2.6 there exists a compact set $K \subseteq l^\infty$ such that every $\overline{\Omega_n}$ admits an isometric embedding into K . By Proposition 4.4, there exists $(\Omega_{n_k})_k$ such that $\Omega_{n_k} \rightarrow \Omega \in \mathcal{U}_K(A, R)$ with respect to d_{HB} , which implies that $\Omega_{n_k} \rightarrow \Omega$ with respect to d_{GHB} . \square

Remark 4.8. A type of compactness theorem for a sequence of noncomplete precompact metric spaces has also been proved in [PS, Theorem 4.1]. It was proven there that for a sequence of noncomplete precompact metric spaces $(X_n)_n$ converging to a limit space X and a given $(\delta_i)_i$ monotonically decreasing to 0, there exists a subsequence $(X_{n_k})_k$ such that δ_i -inner regions converge to the set inside the limit space X for all $i \in \mathbb{N}$. Note that a δ -inner region is the set of all points δ -away from the boundary. Our result, however, states that under the additional assumption that metric spaces are A -uniform, we can find a limit space such that the metric spaces and their boundaries converge to the limit and its metric boundary, respectively.

Remark 4.9. There is a remark on stability of noncomplete PI spaces. Here we say that a metric measure space (X, d, μ) is a PI space if μ is a doubling measure and X supports a Poincaré inequality. We refer readers to [HK] for the exact definition of PI spaces. We say that $(X_n, d_n, \mu_n)_n \subseteq \mathcal{M}$ measured Gromov-Hausdorff converges to $(X, d, \mu) \in \mathcal{M}$ with boundary if there exist a metric space (Z, d_Z) and isometric embeddings $\iota_n : \overline{X_n} \hookrightarrow Z$, $\iota : \overline{X} \hookrightarrow Z$ such that

- $d_H^Z(\iota_n(X_n), \iota(X)) + d_H^Z(\iota(\partial X_n), \iota(\partial X)) \rightarrow 0$ as $n \rightarrow \infty$,
- $\iota_{n,*}\mu_n \rightharpoonup \iota_*\mu$ (weak* convergence in $C_b(Z)^*$),

where $C_b(Z)$ is the set of all continuous functions with bounded support and $\iota_{n,*}\mu_n, \iota_*\mu$ are pushforward measures of μ_n and μ , respectively. For a given sequence of bounded PI uniform spaces $(X_n, d_n, \mu_n)_n$ measured Gromov-Hausdorff converging to $(X, d, \mu) \in \mathcal{M}$ with boundary, we claim that (X, d, μ) is a PI space. In fact, we have already shown,

by Theorem 1.1, that X is a uniform space. Additionally, the sequence $(\overline{X}_n, d_n, \mu_n)$ is a PI space by [HKST, Lemma 8.2.3]. Moreover, by [Ch, Chapter 9] and [Kei, Theorem 3], (\overline{X}, d, μ) supports a Poincaré inequality. Since the limit space X is a uniform domain in the PI space \overline{X} , we conclude that X supports a Poincaré inequality by [BSh, Theorem 4.4].

5 Stability of M -roughly starlike δ -Gromov hyperbolic spaces

In this section we prove the stability of a sequence of M -roughly starlike δ -Gromov hyperbolic spaces, under the assumption that a limit exists. Recall again that curves are parametrized by arclength with respect to d . We first prove that δ -Gromov hyperbolicity is stable under the pointed Gromov-Hausdorff convergence.

Proposition 5.1. *Let $(X_n, d_n, p_n)_n$ be a sequence of pointed δ -Gromov hyperbolic spaces. Suppose that $(X_n, d_n, p_n)_n$ is pointed Gromov-Hausdorff convergent to (X, d, p) as $n \rightarrow \infty$. Then X is a δ -Gromov hyperbolic space.*

Proof. Let $\epsilon > 0$ be fixed. Pick arbitrary four points $x, y, z, m \in X$. Then let $R > 0$ so that all of these four points are in $B_d(p, R - \epsilon)$. From Property (3) in Definition 2.7, we know that for sufficiently large $n \in \mathbb{N}$, there exist $m_n, x_n, y_n, z_n \in X_n$ such that

$$d(f_n^\epsilon(m_n), m) < \epsilon, \quad d(f_n^\epsilon(x_n), x) < \epsilon, \quad d(f_n^\epsilon(y_n), y) < \epsilon, \quad d(f_n^\epsilon(z_n), z) < \epsilon.$$

Since X_n is a δ -Gromov hyperbolic space, we have

$$(x_n|z_n)_{m_n} \geq (x_n|y_n)_{m_n} \wedge (y_n|z_n)_{m_n} - \delta.$$

From Property (2) in Definition 2.7, we notice that

$$(x_n|z_n)_{m_n} \leq (f_n^\epsilon(x_n)|f_n^\epsilon(z_n))_{f_n^\epsilon(m_n)} + \frac{3}{2}\epsilon.$$

Similarly, we can get

$$(x_n|y_n)_{m_n} \wedge (y_n|z_n)_{m_n} \geq (f_n^\epsilon(x_n)|f_n^\epsilon(y_n))_{f_n^\epsilon(m_n)} \wedge (f_n^\epsilon(y_n)|f_n^\epsilon(z_n))_{f_n^\epsilon(m_n)} - \frac{3}{2}\epsilon$$

Combining all of these inequalities, we have

$$\begin{aligned} (f_n^\epsilon(x_n)|f_n^\epsilon(z_n))_{f_n^\epsilon(m_n)} + \frac{3}{2}\epsilon &\geq (f_n^\epsilon(x_n)|f_n^\epsilon(y_n))_{f_n^\epsilon(m_n)} \wedge (f_n^\epsilon(y_n)|f_n^\epsilon(z_n))_{f_n^\epsilon(m_n)} \\ &\quad - \frac{3}{2}\epsilon - \delta. \end{aligned}$$

Also, from direct calculation, we have

$$(f_n^\epsilon(x_n)|f_n^\epsilon(z_n))_{f_n^\epsilon(m_n)} \leq (x|z)_m + 3\epsilon/2,$$

and

$$(f_n^\epsilon(x_n)|f_n^\epsilon(y_n))_{f_n^\epsilon(m_n)} \wedge (f_n^\epsilon(y_n)|f_n^\epsilon(z_n))_{f_n^\epsilon(m_n)} \geq (x|y)_m \wedge (y|z)_m - 3\epsilon/2.$$

We conclude that

$$(x|z)_m + 3\epsilon \geq (x|y)_m \wedge (y|z)_m - 3\epsilon - \delta.$$

Letting ϵ tend to 0 completes the proof. \square

We next prove that the roughly starlike property is stable.

Proposition 5.2. *Let $(X_n, d_n, p_n)_n$ be a sequence of pointed proper geodesic M -roughly starlike δ -Gromov hyperbolic spaces which is pointed Gromov-Hausdorff convergent to some metric space (X, d, p) . Then, there exists $\tilde{M} := \tilde{M}(M, \delta)$ such that (X, d, p) is \tilde{M} -roughly starlike.*

Proof. For each $k \in \mathbb{N}$, set $\epsilon := 1/k$ and $r := k$. Then there exists $n_k \in \mathbb{N}$ such that $f_{n_k}^{1/k} : B(p_{n_k}, k) \rightarrow X$ with the properties in Definition 2.7. For each $x \in X$, take $k_x \in \mathbb{N}$ such that $k_x \geq d(p, x) + 1$. Note that we can find $x_{n_k} \in B_{d_{n_k}}(p_{n_k}, k)$ such that $d(f_{n_k}^{1/k}(x_{n_k}), x) \leq 1/k$ for each $k \geq k_x$. Since X_{n_k} is M -roughly starlike and $x_{n_k} \in X_{n_k}$, there is a geodesic ray $\gamma_{n_k} \subseteq X_{n_k}$ such that $\gamma_{n_k}(0) = p_{n_k}$ and $\text{dist}(x_{n_k}, \gamma_{n_k}) \leq M$. Note that since γ_{n_k} is a geodesic ray, $\gamma_{n_k}([0, k-1])$ is in $B_{d_{n_k}}(p_{n_k}, k)$ and $f_{n_k}^{1/k} \circ \gamma_{n_k}|_{[0, k-1]}$ is a $(1, 1/k)$ -quasi-isometric path (see Definition ??). Since the limit space X is geodesic, we have a geodesic curve $\tilde{\gamma}_{n_k}$ emanating from p to $f_{n_k}^{1/k}(\gamma_{n_k}(k-1))$. Note that the limit space is proper geodesic δ -Gromov hyperbolic by Proposition 5.1. Therefore from the geodesic stability theorem [V2, Theorem 3.7] we get

$$d_H(\tilde{\gamma}_{n_k}, f_{n_k}^{1/k} \circ \gamma_{n_k}|_{[0, k-1]}) \leq M(\delta) \quad (12)$$

for some constant $M(\delta)$ depending only on δ . Pick $t_{n_k} \in [0, \infty)$ such that $d_{n_k}(x_{n_k}, \gamma_{n_k}(t_{n_k})) \leq M$. We claim that $(t_{n_k})_k$ has a uniform bound from above. In fact, we have

$$d_{n_k}(p_{n_k}, x_{n_k}) \leq d(p, f_{n_k}^{1/k}(x_{n_k})) + 1/k \leq d(p, x) + d(x, f_{n_k}^{1/k}(x_{n_k})) + 1/k < C(x) \quad (13)$$

where $C(x) = d(p, x) + 2$. Therefore $x_{n_k} \in B_{d_{n_k}}(p_{n_k}, C(x))$ and we conclude that

$$t_{n_k} = d_{n_k}(p_{n_k}, \gamma_{n_k}(t_{n_k})) \leq d_{n_k}(p_{n_k}, x_{n_k}) + d_{n_k}(x_{n_k}, \gamma_{n_k}(t_{n_k})) \leq C(x) + M.$$

Therefore, for k large enough, $\gamma_{n_k}(t_{n_k}) \in B_{d_{n_k}}(p_{n_k}, k)$, $t_{n_k} \leq k-1$, and we get

$$d(f_{n_k}^{1/k}(x_{n_k}), f_{n_k}^{1/k}(\gamma_{n_k}(t_{n_k}))) \leq d_{n_k}(x_{n_k}, \gamma_{n_k}(t_{n_k})) + 1/k \leq M + 1/k. \quad (14)$$

From the stability consequence (12), there exists $s_{n_k} \in [0, l_d(\tilde{\gamma}_{n_k})]$ such that

$$d(\tilde{\gamma}_{n_k}(s_{n_k}), f_{n_k}^{1/k}(\gamma_{n_k}(t_{n_k}))) \leq M(\delta). \quad (15)$$

Combining (14) and (15), we have

$$d(f_{n_k}^{1/k}(x_{n_k}), \tilde{\gamma}_{n_k}) \leq d(f_{n_k}^{1/k}(x_{n_k}), \tilde{\gamma}_{n_k}(s_{n_k})) \leq M + M(\delta) + 1/k.$$

Moreover, since $d(x, f_{n_k}^{1/k}(x_{n_k})) < 1/k$, note that

$$d(x, \tilde{\gamma}_{n_k}) \leq d(x, \tilde{\gamma}_{n_k}(s_{n_k})) \leq M + M(\delta) + 2/k. \quad (16)$$

By an Arzela-Ascoli type argument, there exists a geodesic ray γ and a further subsequence of $\tilde{\gamma}_{n_k}$ (still denoted by $\tilde{\gamma}_{n_k}$) such that $\tilde{\gamma}_{n_k} \rightarrow \gamma$ locally uniformly as $k \rightarrow \infty$. From the estimates (13) and (15), we have

$$\begin{aligned} s_{n_k} = d(p, \tilde{\gamma}_{n_k}(s_{n_k})) &\leq d(p, f_{n_k}^{1/k}(\gamma(t_{n_k}))) + d(f_{n_k}^{1/k}(\gamma(t_{n_k})), \tilde{\gamma}_{n_k}(s_{n_k})) \\ &\leq C(x) + M + 2/k + M(\delta). \end{aligned}$$

Hence $(s_{n_k})_k$ is bounded. By taking a subsequence again if needed, we may assume that s_{n_k} is convergent to some $s \in \mathbb{R}$. Finally, from the uniform convergence of $\tilde{\gamma}_{n_k}|_{[0, l]}$ where $l > s$ and the estimate (16), we have

$$d(x, \gamma(s)) = \lim_{k \rightarrow \infty} d(x, \tilde{\gamma}_{n_k}(s_{n_k})) \leq M + M(\delta).$$

□

Proof of Theorem 1.2. Combine Remark 2.11, Proposition 5.1, and Proposition 5.2. □

6 Stability of uniformized spaces, and boundaries

In this section, we prove Theorem 1.3. Recall again that we fix the constant $\epsilon_0(\delta) > 0$ from Remark 2.21. Also, unless otherwise stated, all curves in a metric space (X, d) are assumed to be parametrized by arclength with respect to d .

In order to prove Theorem 1.3, we need the following lemmas.

Lemma 6.1. *Let (X, d, p) be a pointed proper geodesic δ -Gromov hyperbolic space and $0 < \epsilon \leq \epsilon_0(\delta) := \frac{1}{14L}$ be a fixed constant where*

$$L := 6(1 + 24\delta)(2 + 40\delta) + 48\delta + 3.$$

Then for each $R > 0$ and every pair of points x and y in $\bar{B}_d(p, R)$, there exists a curve γ that is geodesic with respect to d_ϵ such that $\gamma \subseteq \bar{B}_d(p, 3R + L)$.

Remark 6.2. Notice that whether the uniformized space (X^ϵ, d_ϵ) is geodesic or not is nontrivial since we do not have access to the Hopf-Rinow theorem. On the other hand by Definition 2.20, it is obvious that (X^ϵ, d_ϵ) is a length space. What this proposition tells us is that for $0 < \epsilon \leq \epsilon_0(\delta)$, (X^ϵ, d_ϵ) is a geodesic metric space with geodesic curves that do not wander too far.

Proof. Given $x, y \in \bar{B}_d(p, R)$, we first construct a geodesic curve with respect to d_ϵ . Take a curve $\tilde{\gamma}_k$ from x to y satisfying

$$l_{d_\epsilon}(\tilde{\gamma}_k) \leq d_\epsilon(x, y) + \frac{1}{k}$$

for each $k \in \mathbb{N}$. By [BHK, Lemma 5.7], we can take another curve $\hat{\gamma}_k$ with the same endpoints as $\tilde{\gamma}_k$ with $l_{d_\epsilon}(\hat{\gamma}_k) \leq l_{d_\epsilon}(\tilde{\gamma}_k)$ and

$$l_d(\hat{\gamma}_k|_{(s,t)}) \leq 3d(\hat{\gamma}_k(s), \hat{\gamma}_k(t)) + 1$$

whenever $0 \leq s \leq t \leq l_d(\hat{\gamma}_k)$ are such that $d(\hat{\gamma}_k(s), \hat{\gamma}_k(t)) \leq \frac{1}{12\epsilon}$. Since the curves $\hat{\gamma}_k$ satisfy the assumptions of [BHK, Lemma 5.21] by the choice of $\epsilon_0(\delta)$, the curve $\hat{\gamma}_k$ belongs to the L -neighborhood of each geodesic $[x, y]$ with respect to the original metric d . Since $[x, y] \subseteq \bar{B}_d(p, 3R)$, we conclude that $\hat{\gamma}_k \subseteq ([x, y])_L \subseteq \bar{B}_d(p, 3R + L)$. Since the diameter of the uniformized space X^ϵ is at most $\frac{2}{\epsilon}$, we get

$$2 + \frac{2}{\epsilon} \geq 1 + 1/k + l_d(\tilde{\gamma}_k) \geq \int_0^{l_d(\hat{\gamma}_k)} e^{-\epsilon d(p, \hat{\gamma}_k(t))} dt \geq e^{-\epsilon(3R+L)} l_d(\hat{\gamma}_k).$$

Therefore,

$$l_d(\hat{\gamma}_k) \leq \left(2 + \frac{2}{\epsilon}\right) e^{\epsilon(3R+L)} =: M \quad (17)$$

for every $k \in \mathbb{N}$. Set $\alpha_k(t) := l_d(\hat{\gamma}_k)t$ for each $t \in [0, 1]$ and consider $\gamma_k := \hat{\gamma}_k \circ \alpha_k : [0, 1] \rightarrow X$. Then we have, for each $t, \tilde{t} \in [0, 1]$ with $t \leq \tilde{t}$,

$$d(\gamma_k(t), \gamma_k(\tilde{t})) \leq l_d(\gamma_k|_{[t, \tilde{t}]}) \leq l_d(\hat{\gamma}_k|_{[\alpha_k(t), \alpha_k(\tilde{t})]}) \leq l_d(\hat{\gamma}_k)|\tilde{t} - t| \leq M|\tilde{t} - t|,$$

which implies that $(\gamma_k)_k \subseteq C([0, 1] : X)$ is equicontinuous. Hence by the Arzela-Ascoli theorem, we have a curve $\gamma : [0, 1] \rightarrow X$ and a subsequence γ_k (still denoted by γ_k) such that $\gamma_k \rightarrow \gamma$ uniformly with respect to the metric d . Since the identity map $Id : (X, d) \rightarrow (X, d_\epsilon)$ is homeomorphic by [BHK, Appendix], for every partition $t_1 = 0 < \dots < t_N = 1$,

$$\sum_{i=1}^{N-1} d_\epsilon(\gamma(t_i), \gamma(t_{i+1})) \leq \lim_{k \rightarrow \infty} \sum_{i=1}^{N-1} d_\epsilon(\gamma_k(t_i), \gamma_k(t_{i+1})) \leq \lim_{k \rightarrow \infty} l_{d_\epsilon}(\gamma_k) \leq d_\epsilon(x, y),$$

which implies that the curve γ is geodesic with respect to d_ϵ which stays in $[x, y]_L$. This completes the proof. \square

Lemma 6.3. *Let (X, d, p) be a pointed proper geodesic δ -Gromov hyperbolic space and $R \geq 1$, $0 < \epsilon \leq \epsilon_0(\delta)$, and $\delta' > 0$ be given. Set $T := 4 + L$ where L is the constant in Lemma 6.1. For every pair of points $x, y \in \bar{B}_d(p, R)$ with $d(x, y) \geq \delta'/2$, let $\gamma \subseteq B_d(p, TR)$ be a geodesic curve from x to y with respect to d_ϵ taken from Lemma 6.1. Then we can take subcurves $(\gamma_i)_{i=1}^N$ of γ such that*

$$\gamma = \sum_{i=1}^N \gamma_i, \quad \delta'/2 \leq l_d(\gamma_i) < \delta' \quad \text{and} \quad N \leq \frac{4}{\epsilon\delta'} e^{\epsilon TR}.$$

Proof. By Lemma 6.1, for every pair of points $x, y \in \bar{B}_d(p, R)$, we can always find a geodesic curve γ with respect to d_ϵ such that $\gamma \subseteq B_d(p, TR)$. Since the diameter of the uniformized space X^ϵ is at most $\frac{2}{\epsilon}$, we get

$$\frac{2}{\epsilon} \geq \int_0^{l_d(\gamma)} e^{-\epsilon d(p, \gamma_m(t))} dt \geq e^{-\epsilon TR} l_d(\gamma).$$

Therefore, we have

$$l_d(\gamma) \leq \frac{2e^{\epsilon TR}}{\epsilon}. \quad (18)$$

Now split this curve γ into N subcurves $(\gamma_i)_{i=1}^N$ so that $\delta'/2 \leq t_i := l(\gamma_i) < \delta'$ for $1 \leq i \leq N$. Since

$$\delta'/2 \times N \leq \sum_{i=1}^N l_d(\gamma_i) = l_d(\gamma) \leq \frac{2e^{\epsilon TR}}{\epsilon},$$

we have $N \leq \frac{4}{\epsilon \delta'} e^{\epsilon TR}$. \square

Remark 6.4. In order to prove the next lemma, we note that for each $\epsilon > 0$, the function $t \mapsto e^{-\epsilon t}$ is uniformly continuous on $[0, \infty)$, i.e., for any $\tilde{\epsilon} > 0$, there exists $0 < \delta' \leq \tilde{\epsilon} \wedge 1$ such that

$$|e^{-\epsilon t} - e^{-\epsilon t'}| < \tilde{\epsilon}$$

holds for every $t, t' \in [0, \infty)$ with $|t - t'| \leq 4\delta'$. This choice of δ' plays an important role in proving the convergence of uniformized spaces.

Lemma 6.5. *Let (X, d, p) be a pointed proper geodesic δ -Gromov hyperbolic space, $R \geq 1$ and $0 < \epsilon \leq \epsilon_0(\delta)$. Let $T = 4 + L$ be the constant from Lemma 6.3. For any $\tilde{\epsilon} > 0$, let $\delta' > 0$ be as in Remark 6.4 corresponding to $\tilde{\epsilon} > 0$. Then for every $x, y \in \bar{B}_d(p, R)$ with $d(x, y) \geq \delta'/2$, we have*

$$d_\epsilon(x, y) = \int_0^{l_d(\gamma)} e^{-\epsilon d(p, \gamma(t))} dt \geq \sum_{i=1}^N e^{-\epsilon d(p, \gamma_i(t_i))} l_d(\gamma_i) - \frac{2\tilde{\epsilon}}{\epsilon} e^{\epsilon TR}$$

where $\gamma \subseteq B_d(p, TR)$ and subcurves $(\gamma_i)_{i=1}^N$ of γ are as in the statement of Lemma 6.3 corresponding to the constant δ' and $t_i := l_d(\gamma_i)$ for $i = 1, \dots, N$.

Proof. By Lemma 6.3, there exist a geodesic curve $\gamma \subseteq B_d(p, TR)$ from x to y with respect to d_ϵ and subcurves $(\gamma_i)_{i=1}^N$ of γ such that

$$\delta'/2 \leq l_d(\gamma_i) < \delta' \quad \text{and} \quad N \leq \frac{4}{\epsilon \delta'} e^{\epsilon TR}.$$

Then we have, by (18) and the uniform continuity of $e^{-\epsilon t}$,

$$\int_0^{l_d(\gamma)} e^{-\epsilon d(p, \gamma(t))} dt = \sum_{i=1}^N \int_0^{l_d(\gamma_i)} e^{-\epsilon d(p, \gamma_i(t))} dt$$

$$\begin{aligned}
&\geq \sum_{i=1}^N \int_0^{l_d(\gamma_i)} (e^{-\epsilon d(p, \gamma_i(t_i))} - \tilde{\epsilon}) dt \\
&\geq \sum_{i=1}^N e^{-\epsilon d(p, \gamma_i(t_i))} l_d(\gamma_i) - \tilde{\epsilon} l_d(\gamma) \\
&\geq \sum_{i=1}^N e^{-\epsilon d(p, \gamma_i(t_i))} l_d(\gamma_i) - \frac{2\tilde{\epsilon}}{\epsilon} e^{\epsilon TR}.
\end{aligned}$$

This completes the proof. \square

Lemma 6.6. *Let $(X_n, d_n, p_n)_n$ and (X, d, p) be pointed proper geodesic δ -Gromov hyperbolic spaces. Suppose that $(X_n, d_n, p_n)_n$ is pointed Gromov-Hausdorff convergent to (X, d, p) . Fix $0 < \epsilon \leq \epsilon_0(\delta)$ and let $T := 3 + L$ be the constant from Lemma 6.3. Let $R \geq 1$ be given. For any $\tilde{\epsilon} > 0$, let $\delta' > 0$ be the constant given in Remark 6.4. Then there exists $N := N_{T, R, \tilde{\epsilon}} \in \mathbb{N}$ such that for $n \geq N$, we have*

$$|d_{n, \epsilon}(x_n, y_n) - d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n))| \leq S(\epsilon, T, R)\tilde{\epsilon}$$

for every pair of points $x_n, y_n \in \bar{B}_{d_n}(p_n, R)$ where $f_n^{\delta'^2} : B_{d_n}(p_n, TR) \rightarrow X$ is a map as in Definition 2.7 and $S(\epsilon, T, R)\tilde{\epsilon} \rightarrow 0$ as $\tilde{\epsilon} \rightarrow 0$.

Proof. Let $\tilde{\epsilon} > 0$ and $R \geq 1$ be given. We also fix the constant $\delta' > 0$ in Remark 6.4. Since $(X_n, d_n, p_n)_n$ pointed Gromov-Hausdorff converges to (X, d, p) , there exists $N := N_{T, R, \tilde{\epsilon}} \in \mathbb{N}$ such that for $n \geq N$, there exists a map $f_n^{\delta'^2} : B_{d_n}(p_n, TR) \rightarrow X$ satisfying the properties in Definition 2.7. We may assume that $f_n^{\delta'^2}(\bar{B}_{d_n}(p_n, R)) \subseteq \bar{B}_d(p, R)$ since X_n are all geodesic, see Remark 2.8. For given $x_n, y_n \in \bar{B}_{d_n}(p_n, R)$, first note that if $d_n(x_n, y_n) < \delta'/2$, then it is always true that

$$|d_{n, \epsilon}(x_n, y_n) - d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n))| \leq 3\tilde{\epsilon}$$

since $d_\epsilon(x, y) \leq d(x, y)$ for all $x, y \in X$ and $d_{n, \epsilon}(x_n, y_n) \leq d_n(x_n, y_n)$ for all $x_n, y_n \in X_n$. Hence we may assume that $d_n(x_n, y_n) \geq \delta'/2$. By Lemma 6.5, we can find a geodesic curve $\gamma_n \subseteq B_{d_n}(p_n, TR)$ with respect to $d_{n, \epsilon}$, and subcurves $(\gamma_{n, i})_{i=1}^N$ of γ_n such that

$$d_{n, \epsilon}(x_n, y_n) = \int_0^{l_{d_n}(\gamma_n)} e^{-\epsilon d_n(p_n, \gamma_n(t))} dt \geq \sum_{i=1}^N e^{-\epsilon d_n(p_n, \gamma_{n, i}(t_i))} l_{d_n}(\gamma_{n, i}) - \frac{2\tilde{\epsilon}}{\epsilon} e^{\epsilon TR}, \quad (19)$$

where $t_i := l_{d_n}(\gamma_{n, i})$ for $i = 1, \dots, N$. Note that by the property of the map $f_n^{\delta'^2}$ in Definition 2.7, for $(f_n^{\delta'^2}(\gamma_{n, i}(t_i)))_{i=1}^N$,

$$|d_n(\gamma_{n, i}(t_i), \gamma_{n, i+1}(t_{i+1})) - d(f_n^{\delta'^2}(\gamma_{n, i}(t_i)), f_n^{\delta'^2}(\gamma_{n, i+1}(t_{i+1})))| < \delta'^2$$

and

$$|d(p, f_n^{\delta'^2}(\gamma_{n,i}(t_i))) - d_n(p_n, \gamma_{n,i}(t_i))| \leq \delta'^2$$

for $1 \leq i \leq N$. Let $\tilde{\gamma}_1$ be a geodesic curve from $f_n^{\delta'^2}(x_n)$ to $f_n^{\delta'^2}(\gamma_{n,1}(t_1))$ and $\tilde{\gamma}_i$ be a geodesic curve from $f_n^{\delta'^2}(\gamma_{n,i-1}(t_{i-1}))$ to $f_n^{\delta'^2}(\gamma_{n,i}(t_i))$ for $2 \leq i \leq N$ with respect to d . Then for $2 \leq i \leq N$, we get

$$t'_i := l_d(\tilde{\gamma}_i) = d(f_n^{\delta'^2}(\gamma_{n,i-1}(t_{i-1})), f_n^{\delta'^2}(\gamma_{n,i}(t_i))) \leq d_n(\gamma_{n,i-1}(t_{i-1}), \gamma_{n,i}(t_i)) + \delta'^2 \leq 2\delta'.$$

It is also clear that $t'_i := l_d(\tilde{\gamma}_1) \leq 2\delta'$. Hence, from (19) and the uniform continuity of e^{-ct} , we have

$$\begin{aligned} \sum_{i=1}^N e^{-cd(\tilde{\gamma}_i(t'_i), p)} l_d(\tilde{\gamma}_i) &\leq \sum_{i=1}^N (e^{-cd_n(p_n, \gamma_{n,i}(t_i))} + \tilde{\epsilon}) l_d(\tilde{\gamma}_i) \\ &\leq \sum_{i=1}^N (e^{-cd_n(p_n, \gamma_{n,i}(t_i))} + \tilde{\epsilon}) (l_{d_n}(\gamma_{n,i}) + \delta'^2) \\ &\leq \sum_{i=1}^N e^{-cd_n(p_n, \gamma_{n,i}(t_i))} l_{d_n}(\gamma_{n,i}) + \sum_{i=1}^N e^{-cd_n(p_n, \gamma_{n,i}(t_i))} \delta'^2 \\ &\quad + \tilde{\epsilon} l_{d_n}(\gamma_n) + N\tilde{\epsilon}\delta'^2 \\ &\leq \int_0^{l_{d_n}(\gamma_n)} e^{-cd_n(p_n, \gamma_n(t))} dt + \frac{2\tilde{\epsilon}}{\epsilon} e^{\epsilon TR} + N\delta'^2 \\ &\quad + \tilde{\epsilon} l_{d_n}(\gamma_n) + N\tilde{\epsilon}\delta'^2 \\ &\leq \int_0^{l_{d_n}(\gamma_n)} e^{-cd_n(p_n, \gamma_n(t))} dt + \frac{2\tilde{\epsilon}}{\epsilon} e^{\epsilon TR} + \left(\frac{4}{\epsilon\delta'} e^{\epsilon TR}\right) \delta'^2 \\ &\quad + \frac{\tilde{\epsilon}}{\epsilon} e^{\epsilon TR} + \left(\frac{4}{\epsilon\delta'} e^{\epsilon TR}\right) \tilde{\epsilon}\delta'^2 \\ &= \int_0^{l_{d_n}(\gamma_n)} e^{-cd_n(p_n, \gamma_n(t))} dt + C(\epsilon, T, R)\tilde{\epsilon} \end{aligned} \tag{20}$$

where

$$C(\epsilon, T, R) := \frac{11}{\epsilon} e^{\epsilon TR}.$$

Also, noting that for each $t \in [0, l(\tilde{\gamma}_i)]$ ($1 \leq i \leq N$),

$$d(\tilde{\gamma}_i(t), \tilde{\gamma}_i(t_i)) \leq l_d(\tilde{\gamma}_i) \leq 2\delta',$$

we get

$$\sum_{i=1}^N e^{-cd(p, \tilde{\gamma}_i(t'_i))} l_d(\tilde{\gamma}_i) \geq \sum_{i=1}^N \int_0^{l_d(\tilde{\gamma}_i)} (e^{-cd(p, \tilde{\gamma}_i(t))} - \tilde{\epsilon}) dt$$

$$\begin{aligned}
&\geq \sum_{i=1}^N \int_0^{l_d(\tilde{\gamma}_i)} e^{-\epsilon d(p, \tilde{\gamma}_i(t))} dt - (N\tilde{\epsilon})(2\delta') \\
&\geq d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n)) - 2\tilde{\epsilon}\delta' \left(\frac{4}{\epsilon\delta'} e^{\epsilon TR} \right) \\
&\geq d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n)) - 2\tilde{\epsilon} \left(\frac{4}{\epsilon} e^{\epsilon TR} \right) \\
&\geq d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n)) - K(\epsilon, T, R)\tilde{\epsilon} \tag{21}
\end{aligned}$$

where $K(\epsilon, T, R) := \frac{8}{\epsilon} e^{\epsilon TR}$ and we used the estimate for N in Lemma 6.3. Combining the inequalities (20) and (21), we get

$$d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n)) \leq d_{n,\epsilon}(x_n, y_n) + K(\epsilon, T, R)\tilde{\epsilon} + C(\epsilon, T, R)\tilde{\epsilon}.$$

Hence we conclude that

$$d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n)) - d_{n,\epsilon}(x_n, y_n) \leq K(\epsilon, T, R)\tilde{\epsilon} + C(\epsilon, T, R)\tilde{\epsilon}.$$

Next by picking x_n and y_n from $\bar{B}_{d_n}(p_n, R)$ and a geodesic curve γ between $f_n^{\delta'^2}(x_n)$ and $f_n^{\delta'^2}(y_n)$ with respect to the uniformization metric d_ϵ , and doing the same argument for γ instead of γ_n , we also have,

$$d_{n,\epsilon}(x_n, y_n) - d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n)) \leq \tilde{K}(\epsilon, T, R)\tilde{\epsilon} + \tilde{C}(\epsilon, T, R)\tilde{\epsilon}.$$

where

$$\tilde{C}(\epsilon, T, R)\tilde{\epsilon} := \frac{33}{\epsilon} e^{\epsilon TR}$$

and

$$\tilde{K}(\epsilon, T, R)\tilde{\epsilon} := \frac{24}{\epsilon} e^{\epsilon TR}.$$

Therefore, we conclude that

$$|d_{n,\epsilon}(x_n, y_n) - d_\epsilon(f_n^{\delta'^2}(x_n), f_n^{\delta'^2}(y_n))| \leq S(\epsilon, T, R)\tilde{\epsilon}$$

for every $x_n, y_n \in \bar{B}_{d_n}(p_n, R)$ where

$$S(\epsilon, T, R) := 2(K(\epsilon, T, R) \vee \tilde{K}(\epsilon, T, R) \vee C(\epsilon, T, R) \vee \tilde{C}(\epsilon, T, R)).$$

□

Proposition 6.7. *Let $(X_n, d_n, p_n)_n$ and (X, d, p) be pointed proper geodesic δ -Gromov hyperbolic spaces. Suppose that $(X_n, d_n, p_n)_n$ is pointed Gromov-Hausdorff convergent to (X, d, p) and $R \geq 1$ and $0 < \epsilon \leq \epsilon_0(\delta)$ are given. Then*

$$\lim_{n \rightarrow \infty} d_{GH}((\bar{B}_{d_n}(p_n, R), d_{n,\epsilon}), (\bar{B}_d(p, R), d_\epsilon)) = 0.$$

Proof. Set $T := 4 + L$ as in Lemma 6.3. For any $\tilde{\epsilon} > 0$, by Lemma 6.5 and Lemma 6.6, there exist $\delta' > 0$ and $N := N_{T,R,\tilde{\epsilon}} \in \mathbb{N}$ such that for $n \geq N$, we have

$$|d_{n,\epsilon}(x_n, y_n) - d_\epsilon(f_n^{\delta'}(x_n), f_n^{\delta'}(y_n))| \leq S(\epsilon, T, R)\tilde{\epsilon}$$

for every pair of points $x_n, y_n \in \bar{B}_{d_n}(p_n, R)$ where $f_n^{\delta'} : B_{d_n}(p_n, TR) \rightarrow X$ is the map in Definition 2.7 and $S(\epsilon, T, R)\tilde{\epsilon} \rightarrow 0$ as $\tilde{\epsilon} \rightarrow 0$. Recall again that $0 < \delta' \leq \tilde{\epsilon}$ by Remark 6.4. We may assume again that $f_n^{\delta'}(\bar{B}_{d_n}(p_n, R)) \subseteq \bar{B}_d(p, R)$. From Property (3) in Definition 2.7, we know that, with respect to the metric d ,

$$\bar{B}_d(p, R - \delta'^2) \subseteq (f_n^{\delta'}(\bar{B}_{d_n}(p_n, R)))_{\delta'^2}$$

which tells us that

$$\bar{B}_d(p, R) \subseteq (f_n^{\delta'}(\bar{B}_{d_n}(p_n, R)))_{2\delta'^2}$$

since the space is geodesic. This above inclusion holds with respect to d_ϵ as well, so that we get

$$\bar{B}_d(p, R) \subseteq (f_n^{\delta'}(\bar{B}_{d_n}(p_n, R)))_{2\delta'^2}$$

with respect to d_ϵ . Hence, the map $f_n^{\delta'} : \bar{B}_{d_n}(p_n, R) \rightarrow \bar{B}_d(p, R)$ is $M(\epsilon, T, R)\tilde{\epsilon}$ -isometry where $M(\epsilon, T, R) := S(\epsilon, T, R) \vee 2$. Hence we conclude that

$$\limsup_{n \rightarrow \infty} d_{GH}((\bar{B}_d(p_n, R), d_{n,\epsilon}), (\bar{B}_d(p, R), d_\epsilon)) \leq M(\epsilon, T, R)\tilde{\epsilon}.$$

Since $\tilde{\epsilon}$ is arbitrary, the proof is complete. \square

Remark 6.8. From the proof of this proposition, we know that for fixed $R \geq 1$ and $\tilde{\epsilon} > 0$, there exists $N \in \mathbb{N}$ such that for each $n \geq N$, there exists a map $\phi_n : \bar{B}_{d_n}(p_n, R) \rightarrow \bar{B}_d(p, R)$ which is an $\tilde{\epsilon}$ -isometry with respect to both the uniformization metric and the original metric such that $\phi_n(p_n) = p$.

Theorem 6.9 (Theorem 1.3). *Let $(X_n, d_n, p_n)_n$ be pointed proper geodesic M -roughly star-like δ -Gromov hyperbolic spaces. Suppose $(X_n, d_n, p_n)_n$ is pointed Gromov-Hausdorff convergent to (X, d, p) . Then for a fixed $0 < \epsilon \leq \epsilon_0(\delta)$, the sequence of uniformized spaces $(X_n^\epsilon, d_{n,\epsilon})_n$ Gromov-Hausdorff converges to (X^ϵ, d_ϵ) with boundary.*

Proof. Let $\tilde{\epsilon} > 0$ be given. Choose $R > 0$ such that

$$\frac{e^{-\epsilon(R-2\tilde{\epsilon}-M)}}{\epsilon} \leq \tilde{\epsilon}. \quad (22)$$

By Remark 6.8, there exists $N \in \mathbb{N}$ such that for each $n \geq N$, there exists an $\tilde{\epsilon}$ -isometry $\phi_n : \bar{B}_{d_n}(p_n, R) \rightarrow \bar{B}_d(p, R)$ with respect to both the uniformization metric and the original metric such that $\phi_n(p_n) = p$. Let $x_n \in \bar{X}_n^\epsilon \setminus \bar{B}_{d_n}(p_n, R)$. Then there exists a geodesic curve

(or ray) γ_n from p_n to x_n with respect to d_n , see Remark 2.25. Set $\tilde{x}_n := \gamma_n(R) \in \bar{B}_{d_n}(p_n, R)$. We define a map $\Phi_n : \bar{X}_n^\epsilon \rightarrow \bar{X}^\epsilon$ by

$$\Phi_n(x_n) := \begin{cases} \phi_n(x_n) & \text{if } x_n \in \bar{B}(p_n, R) \\ \phi_n(\tilde{x}_n) & \text{if } x_n \in \bar{X}_n^\epsilon \setminus \bar{B}_{d_n}(p_n, R). \end{cases}$$

By Proposition 3.19, it is sufficient to prove that the map Φ_n is a $5\tilde{\epsilon}$ -isometry with boundary. Note that for every $x_n \in \bar{X}_n^\epsilon \setminus \bar{B}_{d_n}(p_n, R)$,

$$d_{n,\epsilon}(\tilde{x}_n, x_n) \leq \int_R^{l_{d_n}(\gamma)} e^{-\epsilon d(p_n, \gamma_n(t))} dt \leq \int_R^\infty e^{-\epsilon t} dt \leq \tilde{\epsilon} \quad (23)$$

where γ_n is a geodesic curve(or ray) from p_n to x_n with respect to the metric d . Hence, for every pair of points $x_n, x'_n \in \bar{X}_n^\epsilon$, we have

$$\begin{aligned} |d_\epsilon(\Phi_n(x_n), \Phi_n(x'_n)) - d_{n,\epsilon}(x_n, x'_n)| &\leq |d_\epsilon(\Phi_n(x_n), \Phi_n(x'_n)) - d_{n,\epsilon}(\tilde{x}_n, \tilde{x}'_n)| \\ &\quad + |d_{n,\epsilon}(\tilde{x}_n, \tilde{x}'_n) - d_{n,\epsilon}(x_n, x'_n)| \\ &\leq \tilde{\epsilon} + |d_{n,\epsilon}(\tilde{x}_n, \tilde{x}'_n) - d_{n,\epsilon}(\tilde{x}_n, x'_n)| \\ &\quad + |d_{n,\epsilon}(\tilde{x}_n, x'_n) - d_{n,\epsilon}(x_n, x'_n)| \\ &\leq 3\tilde{\epsilon}. \end{aligned}$$

where we set $\tilde{x}_n := x_n$ if $x_n \in \bar{B}_{d_n}(p_n, R)$ and used (23). Therefore we obtain

$$\text{dis}(\Phi_n) \leq 3\tilde{\epsilon}$$

with respect to the uniformization metric. We next prove that $X^\epsilon \subseteq (\Phi_n(X_n^\epsilon))_{2\tilde{\epsilon}}$. Let $x \in X^\epsilon$. We may assume that $x \in X^\epsilon \setminus \bar{B}_d(p, R)$ and take a geodesic curve γ with respect to d from p to x , setting $\tilde{x} := \gamma(R)$. By the same argument as in (23), we know that $d_\epsilon(x, \tilde{x}) \leq \tilde{\epsilon}$. Since $\phi_n : \bar{B}_{d_n}(p_n, R) \rightarrow \bar{B}_d(p, R)$ is $\tilde{\epsilon}$ -isometry with respect to the uniformization metric, there exists $x_n \in \bar{B}_d(p, R)$ such that $d_\epsilon(\phi_n(x_n), \tilde{x}) \leq \tilde{\epsilon}$. Hence we obtain

$$d_\epsilon(x, \Phi_n(x_n)) = d_\epsilon(x, \phi_n(x_n)) \leq d_\epsilon(x, \tilde{x}) + d_\epsilon(\tilde{x}, \phi_n(x_n)) \leq 2\tilde{\epsilon}.$$

We next claim that $\partial_{d_\epsilon} X^\epsilon \subseteq (\Phi(\partial_{d_{n,\epsilon}} X_n^\epsilon))_{5\tilde{\epsilon}}$. By Remark 2.25, for any $x \in \partial_{d_\epsilon} X^\epsilon$, we can take a geodesic ray $\gamma : [0, \infty) \rightarrow X$ with respect to the metric d such that $d_\epsilon(x, \gamma(k)) \rightarrow 0$ as $k \rightarrow \infty$. Set $\tilde{x} := \gamma(R)$. As in (23), we know that

$$d_\epsilon(x, \tilde{x}) \leq \int_R^\infty e^{-\epsilon t} dt \leq \tilde{\epsilon}. \quad (24)$$

Since $\phi_n : \bar{B}_{d_n}(p_n, R) \rightarrow \bar{B}_d(p, R)$ is an $\tilde{\epsilon}$ -isometry with respect to the uniformization metric, there exists $x_n \in \bar{B}_{d_n}(p_n, R)$ such that $d(\phi_n(x_n), \tilde{x}) \leq \tilde{\epsilon}$. The M -roughly starlike property of X_n allows us to take a geodesic ray $\gamma_n : [0, \infty) \rightarrow X_n$ emanating from p_n with

respect to d_n such that $\text{dist}(x_n, \gamma_n) \leq M$. Pick $z_n \in \gamma_n$ such that $d(x_n, z_n) = \text{dist}(x_n, \gamma_n)$. Then since

$$d_n(p_n, x_n) \geq d(p, \phi_n(x_n)) - \tilde{\epsilon} \geq d(p, \tilde{x}) - d(\tilde{x}, \phi_n(x_n)) - \tilde{\epsilon} \geq R - 2\tilde{\epsilon},$$

we get

$$d_{n,\epsilon}(x_n, z_n) \leq \int_0^{l_d(\beta)} e^{-\epsilon d(p, \beta(t))} dt \leq e^{-\epsilon(R-2\tilde{\epsilon})} \int_0^{l_d(\beta)} e^{\epsilon t} dt \leq \tilde{\epsilon} \quad (25)$$

where β is a geodesic curve from x_n to z_n and we employed the Harnack inequality (2). Also, noting that

$$d_n(p_n, z_n) \geq d_n(p_n, x_n) - d_n(x_n, z_n) \geq d(p, \phi_n(x_n)) - \tilde{\epsilon} - M \geq R - 2\tilde{\epsilon} - M,$$

we have

$$d_{n,\epsilon}(z_n, \gamma_n(R)) \leq \int_{R-2\tilde{\epsilon}-M}^{\infty} e^{-\epsilon t} dt \leq \tilde{\epsilon}. \quad (26)$$

Set $y := \Phi(a_n) := \Phi_n(\gamma_n(R)) = \phi_n(\gamma_n(R))$ where $a_n \in \partial_{d_{n,\epsilon}} X_n^\epsilon$ is the unique point such that $\gamma_n(k) \rightarrow a_n$ as $k \rightarrow \infty$. Then, combining (24), (25), and (26), we have

$$\begin{aligned} d_\epsilon(x, y) &\leq d_\epsilon(x, \tilde{x}) + d_\epsilon(\tilde{x}, \phi_n(x_n)) + d_\epsilon(\phi_n(x_n), y) \\ &\leq d_\epsilon(x, \tilde{x}) + d(\tilde{x}, \phi_n(x_n)) + d_\epsilon(\phi_n(x_n), y) \\ &\leq 3\tilde{\epsilon} + d_{n,\epsilon}(x_n, \gamma_n(R)) \\ &\leq 3\tilde{\epsilon} + d_{n,\epsilon}(x_n, z_n) + d_{n,\epsilon}(z_n, \gamma_n(R)) \\ &\leq 5\tilde{\epsilon}. \end{aligned}$$

Therefore we obtain

$$\partial_{d_\epsilon} X^\epsilon \subseteq (\Phi(\partial_{d_{n,\epsilon}} X_n^\epsilon))_{5\tilde{\epsilon}}.$$

This completes the proof. \square

7 Stability of quasihyperbolization

The rest of the paper is devoted to the proof of Theorem 1.4. The proof is based on the approach used already in the section on the convergence of uniformized spaces. Recall that for a metric space (Ω, d) , $d(\cdot) := d(\cdot, \partial\Omega)$ and all curves are parametrized by arc-length with respect to d unless otherwise stated.

Proposition 7.1. *Let (Ω, d) be an A -uniform space. Then for each $R > 0$ and $p \in \Omega$,*

$$B_k(p, R) \subseteq \Omega \setminus (\partial\Omega)_c$$

where $c := \frac{d(p)/2}{e^R - 1} \wedge d(p)/2$, $B_k(p, R)$ is a ball with radius R with respect to the quasihyperbolic metric k , and $(\partial\Omega)_c$ is the c -neighborhood of the boundary $\partial\Omega$ with respect to d .

Proof. For each $x \in B_k(p, R)$, if $d(p, x) < d(p)/2$, then $x \in \Omega \setminus (\partial\Omega)_c$. Hence we assume that $d(p, x) \geq d(p)/2$. Take a curve $\gamma : [0, l_d(\gamma)] \rightarrow \Omega$ from x to p that is geodesic with respect to the quasihyperbolic metric k and the closest point $y \in \partial\Omega$ from x . Then we have

$$d(\gamma(t)) \leq d(\gamma(t), y) \leq d(\gamma(t), x) + d(x, y) = d(\gamma(t), x) + d(x) = t + d(x).$$

Hence we get

$$R > \int_0^{l_d(\gamma)} \frac{1}{d(\gamma(t))} dt \geq \int_0^{l_d(\gamma)} \frac{1}{d(x) + t} dt \geq \log\left(\frac{d(x) + l_d(\gamma)}{d(x)}\right) \geq \log\left(\frac{d(x) + d(p)/2}{d(x)}\right).$$

We conclude that

$$e^R > \frac{d(x) + d(p)/2}{d(x)},$$

which implies $d(p)/2 < d(x)(e^R - 1)$. Therefore $x \notin (\partial\Omega)_c$. \square

Lemma 7.2. *Let (Ω, d) be an A -uniform space and $\delta' > 0$ be given. For every pair of points $x, y \in \Omega$ with $d(x, y) \geq \delta'/2$, let γ be a geodesic curve from x to y with respect to the quasihyperbolic metric k . Then we can take subcurves $(\gamma_i)_{i=1}^N$ such that*

$$\gamma = \sum_{i=1}^N \gamma_i, \quad \delta'/2 \leq l_d(\gamma_i) < \delta', \quad \text{and} \quad N \leq \frac{2B \operatorname{diam}(\Omega)}{\delta'}.$$

where the constant B only depends on the constant A .

Proof. Let $x, y \in \Omega$ and γ be a geodesic curve between x and y with respect to the quasihyperbolic metric k . Note that, by [BHK, Theorem 2.10], γ is a $B := B(A)$ -uniform curve, in particular,

$$l_d(\gamma) \leq Bd(x, y) \leq B \operatorname{diam}(\Omega). \quad (27)$$

Split this curve γ into N subcurves $(\gamma_i)_{i=1}^N$ so that $\delta'/2 \leq t_i := l_d(\gamma_i) < \delta'$ for $1 \leq i \leq N$. Then we have

$$\frac{\delta'}{2} \times N \leq \sum_{i=1}^N l_d(\gamma_i) \leq B \operatorname{diam}(\Omega),$$

which tells us that $N \leq \frac{2B \operatorname{diam}(\Omega)}{\delta'}$. \square

Remark 7.3. Let $\alpha > 0$ be given. In order to show the next lemma, we note that for any $\epsilon > 0$, the function $t \mapsto 1/t$ is uniformly continuous on $[\alpha, \infty)$, i.e., for any $\epsilon > 0$, there exists $0 < \delta' \leq \epsilon \wedge 1$ such that

$$\left| \frac{1}{t} - \frac{1}{t'} \right| \leq \epsilon \quad (28)$$

holds for every $t, t' \in [0, \infty)$ with $|t - t'| \leq 4\delta'$.

Lemma 7.4. *Let (Ω, d) be an A -uniform space and $p \in \Omega$ and $R > 0$ be given. Let $0 < \alpha < c$ be a positive constant where c is the constant given for $B_k(p, 3R)$ in Lemma 7.1. Then for any $\epsilon > 0$, let $\delta' > 0$ be a constant from Remark 7.3 for the interval $[\alpha, \infty)$. Then for every pair of points $x, y \in B_k(p, R)$ with $d(x, y) \geq \delta'/2$, we have*

$$\int_0^{l_d(\gamma)} \frac{1}{d(\gamma(t))} dt \geq \sum_{i=1}^N \frac{1}{d(\gamma_i(t_i))} l_d(\gamma_i) - B \operatorname{diam}(\Omega) \epsilon$$

where γ is a geodesic curve from x to y with respect to the quasihyperbolic metric k , $(\gamma_i)_{i=1}^N$ are the subcurves of γ taken from Lemma 7.2, and $t_i := l_d(\gamma_i)$.

Proof. For any $\epsilon > 0$, choose $\delta' > 0$ as in Remark 7.3. Let $x, y \in B_k(p, R)$ with $d(x, y) \geq \delta'/2$ and $\gamma \subseteq B_k(p, 3R)$ be a geodesic curve from x to y with respect to k . Then, we can take subcurves $(\gamma_i)_{i=1}^N$ as in Lemma 7.2. Since $d(\gamma(t)) \geq c > \alpha$ by Lemma 7.1, from Remark 7.3 we have

$$\begin{aligned} \int_0^{l_d(\gamma)} \frac{1}{d(\gamma(t))} dt &= \sum_{i=1}^N \int_0^{l_d(\gamma_i)} \frac{1}{d(\gamma_i(t))} dt \\ &\geq \sum_{i=1}^N \int_0^{l_d(\gamma_i)} \left(\frac{1}{d(\gamma_i(t_i))} - \epsilon \right) dt \\ &\geq \sum_{i=1}^N \frac{1}{d(\gamma_i(t_i))} l_d(\gamma_i) - \epsilon l_d(\gamma) \\ &\geq \sum_{i=1}^N \frac{1}{d(\gamma_i(t_i))} l_d(\gamma_i) - B \operatorname{diam}(\Omega) \epsilon. \end{aligned} \tag{29}$$

where we used (27) for the last inequality. This completes the proof. \square

Proposition 7.5. *Let $(\Omega_n, d_n)_n \subseteq \mathcal{U}(A, R)$ and $(\Omega, d) \in \mathcal{M}$. Suppose that $(\Omega_n, d_n)_n$ are all length spaces and $d_{GHB}(\Omega_n, \Omega) \rightarrow 0$ as $n \rightarrow \infty$. Then there exist $p_n \in \Omega_n$ and $p \in \Omega$ such that for each $R > 0$ and $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \geq N$, we have a map $I_n : B_{k_n}(p_n, R) \rightarrow \Omega$ with $I(p_n) = p$ such that*

$$|k_n(x_n, y_n) - k(I_n(x_n), I_n(y_n))| \leq S \epsilon$$

for some $S > 0$ which only depends on the constants $A > 0$, $R > 0$ and $p \in \Omega$.

Proof. By Proposition 3.21, we may assume that $(\Omega_n)_n$ and Ω are all in a same metric space (Z, d_Z) and $d_H^Z(\Omega_n, \Omega) + d_H^Z(\partial\Omega_n, \partial\Omega) \rightarrow 0$ as $n \rightarrow \infty$. Pick any $p \in \Omega$ and take $p_n \in \Omega_n$ such that $d_Z(p_n, p) \rightarrow 0$ as $n \rightarrow \infty$. Let $R > 0$ and $\epsilon > 0$ be fixed. Then by Lemma 7.1, we know that

$$B_k(p, 3R) \subseteq \Omega \setminus (\partial\Omega)_c$$

and

$$B_{k_n}(p_n, 3R) \subseteq \Omega_n \setminus (\partial\Omega_n)_{c_n}$$

where $c = \frac{d(p)/2}{e^{3R}-1} \wedge d(p)/2$ and $c_n = \frac{d_n(p_n)/2}{e^{3R}-1} \wedge d_n(p_n)/2$. Note that $c_n \rightarrow c$ as $n \rightarrow \infty$. Set $\alpha := \frac{c}{2} \wedge \frac{\tilde{c}}{2} > 0$ where $\tilde{c} > 0$ is a constant given later. For this $\alpha > 0$, let us fix the constant $\delta' > 0$ in Remark 7.3. We may assume that $\delta' \leq \frac{c}{20} \wedge \frac{\tilde{c}}{20}$ for the later estimates. Then, there exists $N \in \mathbb{N}$ such that for each $n \geq N$, we have

$$d_Z(p_n, p) \leq \delta'^2, \quad \frac{19c}{20} \leq c_n \leq \frac{21c}{20}, \quad \text{and} \quad d_H^Z(\Omega_n, \Omega) + d_H^Z(\partial\Omega_n, \partial\Omega) \leq \delta'^2 =: \tilde{\epsilon}. \quad (30)$$

Let $n \geq N$ be fixed. Since for each $x_n \in \Omega_n$ we can find $x \in \Omega$ such that $d_Z(x_n, x) \leq \tilde{\epsilon}$, define a map $I_n : B_{k_n}(p_n, R) \rightarrow \Omega$ by $B_{k_n}(p_n, R) \ni x_n \mapsto x \in \Omega$ with $I_n(p_n) := p$. Let $x_n, y_n \in B_{k_n}(p_n, R)$ be given. We may assume that $d(x_n, y_n) \geq \delta'/2$. Take a geodesic curve γ_n from x_n to y_n with respect to k_n . By Lemma 7.2, we have subcurves $(\gamma_{n,i})_{i=1}^N$ of γ_n such that

$$\gamma_n = \sum_{i=1}^N \gamma_{n,i}, \quad \delta'/2 \leq l_{d_n}(\gamma_{n,i}) < \delta' \quad \text{and} \quad N \leq \frac{2BD}{\delta'}. \quad (31)$$

where $D := \sup_n \text{diam}(\Omega_n)$. Also note that $d_n(\gamma_{n,i}(t)) \geq c_n \geq \frac{c}{2}$ for each $t \in [0, l_{d_n}(\gamma_{n,i})]$ by the choice of c_n in (30) and $\gamma_n \subseteq B_{k_n}(p_n, 3R)$. Then, by Lemma 7.4, we have

$$\int_0^{l_{d_n}(\gamma_n)} \frac{1}{d_n(\gamma_n(t))} dt \geq \sum_{i=1}^N \frac{1}{d_n(\gamma_{n,i}(t_i))} l_{d_n}(\gamma_{n,i}) - BD\epsilon \quad (32)$$

where $t_i := l_{d_n}(\gamma_{n,i})$. By the definition of the map $I_n : B_{k_n}(p_n, R) \rightarrow \Omega$, for $(I_n(\gamma_n(t_i)))_{i=1}^N$ we have

$$|d_n(\gamma_{n,i}(t_i), \gamma_{n,i+1}(t_{i+1})) - d(I_n(\gamma_{n,i}(t_i)), I_n(\gamma_{n,i+1}(t_{i+1})))| < 2\tilde{\epsilon}. \quad (33)$$

Since (Ω, d) is a length space, we can pick a curve $\tilde{\gamma}_1$ from $I_n(x_n)$ to $I_n(\gamma_{n,1}(t_1))$ whose length is less than $d(I_n(x_n), I_n(\gamma_{n,1}(t_1))) + \tilde{\epsilon}$ and $\tilde{\gamma}_i$ from $I_n(\gamma_{n,i-1}(t_{i-1}))$ to $I_n(\gamma_{n,i}(t_i))$ whose length is less than $d(I_n(\gamma_{n,i-1}(t_{i-1})), I_n(\gamma_{n,i}(t_i))) + \tilde{\epsilon}$ for $2 \leq i \leq N$. Let $t'_i := l_d(\tilde{\gamma}_i)$. Then from (33) we get

$$t'_i := l_d(\tilde{\gamma}_i) \leq d_n(\gamma_{n,i-1}(t_{i-1}), \gamma_{n,i}(t_i)) + 3\tilde{\epsilon} \leq l_{d_n}(\gamma_{n,i}) + 3\tilde{\epsilon} \leq 4\delta'$$

for $1 \leq i \leq N$, $t_1 \leq 4\delta'$, and

$$\begin{aligned} |d(\tilde{\gamma}_i(t'_i)) - d_n(\gamma_{n,i}(t_i))| &\leq |d(I_n(\gamma_{n,i}(t_i))) - d(\gamma_{n,i}(t_i))| \\ &\quad + |d(\gamma_{n,i}(t_i)) - d_n(\gamma_{n,i}(t_i))| \\ &\leq 2\tilde{\epsilon} \leq 2\delta'. \end{aligned}$$

for $1 \leq i \leq N$. Also, in order to employ (28), we also note that

$$d(\tilde{\gamma}_i(t'_i)) \geq d(\gamma_{n,i}(t_i)) - \tilde{\epsilon} \geq d_n(\gamma_{n,i}(t_i)) - 2\tilde{\epsilon} \geq \frac{19c}{20} - \frac{2c}{20} = \frac{17c}{20}. \quad (34)$$

Hence, from (27), (28), (32), (34), and the estimate

$$\sum_{i=1}^N \frac{1}{d_n(\gamma_{n,i}(t_i))} \tilde{\epsilon} \leq \frac{6N\tilde{\epsilon}}{c} \leq \frac{12BD}{c\delta'} \tilde{\epsilon} \leq \frac{12BD}{c} \epsilon,$$

we have

$$\begin{aligned} \sum_{i=1}^N \frac{1}{d(\tilde{\gamma}_i(t'_i))} l_d(\tilde{\gamma}_i) &\leq \sum_{i=1}^N \left(\frac{1}{d_n(\gamma_{n,i}(t_i))} + \epsilon \right) (l_{d_n}(\gamma_{n,i}) + 3\tilde{\epsilon}) \\ &\leq \sum_{i=1}^N \frac{1}{d_n(\gamma_{n,i}(t_i))} l_{d_n}(\gamma_{n,i}) + 3 \sum_{i=1}^N \frac{1}{d_n(\gamma_{n,i}(t_i))} \tilde{\epsilon} \\ &\quad + \epsilon l_{d_n}(\gamma_n) + 3\tilde{\epsilon} \epsilon \\ &\leq \int_0^{l_{d_n}(\gamma_n)} \frac{1}{d_n(\gamma_n(t))} dt + BD\epsilon + \frac{36BD}{c} \epsilon \\ &\quad + BD\epsilon + 3\tilde{\epsilon} \epsilon \\ &= \int_0^{l_{d_n}(\gamma_n)} \frac{1}{d_n(\gamma_n(t))} dt + C(B, D, c)\epsilon \end{aligned} \tag{35}$$

where

$$C(B, D, c) := 2BD + \frac{36BD}{c} + 3.$$

Since we have the estimate

$$d(\tilde{\gamma}_i(t), \tilde{\gamma}_i(t'_i)) \leq l_d(\tilde{\gamma}_i) \leq d_n(\gamma_{n,i}(t_i), \gamma_{n,i+1}(t_{i+1})) + 3\tilde{\epsilon} \leq l_{d_n}(\gamma_{n,i+1}) + 3\tilde{\epsilon} \leq 4\delta,$$

for each $t \in [0, l_d(\tilde{\gamma}_i)]$ ($1 \leq i \leq N$), one can see, together with (34), that

$$d(\tilde{\gamma}_i(t)) \geq d(\tilde{\gamma}_i(t'_i)) - 4\delta \geq \frac{17c}{20} - \frac{4c}{100} \geq \frac{13c}{20} \geq \frac{c}{2}.$$

Hence by (28) and (34), we have

$$\begin{aligned} \sum_{i=1}^N \frac{1}{d(\tilde{\gamma}_i(t'_i))} l_d(\tilde{\gamma}_i) &\geq \sum_{i=1}^N \int_0^{l_d(\tilde{\gamma}_i)} \left(\frac{1}{d(\tilde{\gamma}_i(t))} - \epsilon \right) dt \\ &\geq \sum_{i=1}^N \int_0^{l_d(\tilde{\gamma}_i)} \frac{1}{d(\tilde{\gamma}_i(t))} dt - (N\epsilon)(4\delta) \\ &\geq k(I_n(x_n), I_n(y_n)) - 8BD\epsilon \\ &\geq k(I_n(x_n), I_n(y_n)) - K(B, D)\epsilon \end{aligned} \tag{36}$$

where $K(B, D) := 8BD$. Therefore, combining the inequalities (35) and (36), we conclude that

$$k(I_n(x_n), I_n(y_n)) \leq k_n(x_n, y_n) + K(B, D)\epsilon + C(B, D, c)\epsilon$$

Hence

$$k(I_n(x_n), I_n(y_n)) - k_n(x_n, y_n) \leq K(B, D)\epsilon + C(B, D, c)\epsilon.$$

Next, by following almost the same proof as above, we will prove that for every pair points $x_n, y_n \in B_{k_n}(p_n, R)$

$$k_n(x_n, y_n) - k(I_n(x_n), I_n(y_n)) \leq \tilde{K}(B, D)\epsilon + \tilde{C}(\tilde{c}, B, D)\epsilon$$

for some $\tilde{K}(B, D)$ and $\tilde{C}(\tilde{c}, B, D)$. Again pick x_n and y_n from $B_{k_n}(p_n, R)$ and a geodesic curve γ between $I_n(x_n)$ and $I_n(y_n)$ with respect to the quasihyperbolic metric k . We first claim that $I_n(x_n)$ and $I_n(y_n)$ are in the large enough ball $B_k(p, M)$ for some $M > 0$. Since we have the estimate

$$d(I_n(x_n)) \geq d(x_n) - \tilde{\epsilon} \geq d_n(x_n) - 2\tilde{\epsilon} \geq c_n - 2\tilde{\epsilon} \geq \frac{19c}{20} - \frac{2c}{20} \geq \frac{17c}{20}$$

for all $n \geq N$, by [BHK, Lemma 2.13], we have

$$\begin{aligned} k(p, I_n(x_n)) &\leq 4A^2 \log\left(1 + \frac{d(p, I_n(x_n))}{d(p) \wedge d(I_n(x_n))}\right) \\ &\leq 4A^2 \log\left(1 + \frac{20D}{17c}\right) =: M(A, c, D) =: M. \end{aligned}$$

Hence a geodesic curve γ connecting $I_n(x_n)$ and $I_n(y_n)$ stays in $B_k(p, 3M)$. By Lemma 7.1, we find \tilde{c} such that

$$B_k(p, 3M) \subseteq \Omega \setminus (\partial\Omega)_{\tilde{c}}$$

where $\tilde{c} = \frac{d(p)/2}{e^{3M}-1} \wedge d(p)/2$. From here by following the same argument we did above we have

$$k_n(x_n, y_n) - k(I_n(x_n), I_n(y_n)) \leq \tilde{K}(B, D)\epsilon + \tilde{C}(\tilde{c}, B, D)\epsilon$$

where

$$\tilde{C}(\tilde{c}, B, D) := 2BD + \frac{36BD}{\tilde{c}} + 3 \quad \text{and} \quad \tilde{K}(B, D) := 8BD.$$

Therefore, we conclude that

$$|k_n(x_n, y_n) - k(I_n(x_n), I_n(y_n))| \leq S\epsilon$$

where

$$S := S(B, D, \tilde{c}, c) := 2(\tilde{K}(B, D) \vee \tilde{C}(\tilde{c}, B, D) \vee K(B, D) \vee C(B, D, c)).$$

□

Theorem 7.6 (Theorem 1.4). *Let $(\Omega_n, d_n)_n \subseteq \mathcal{U}(A, R)$ and $(\Omega, d) \in \mathcal{M}$. Suppose $(\Omega_n, d_n)_n$ are all length spaces and $d_{GH}(\Omega_n, \Omega) \rightarrow 0$ as $n \rightarrow \infty$. Then $(\Omega_n, p_n, k_n)_n$ is pointed Gromov-Hausdorff convergent to (Ω, p, k) for some $p_n \in \Omega_n$ and $p \in \Omega$ where k_n and k are quasihyperbolic metrics defined on Ω_n and Ω , respectively.*

Proof. By Proposition 7.5, for each $R > 0$ and $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \geq N$, we have a map $I_n : B_{k_n}(p_n, R) \rightarrow \Omega$ with $I(p_n) = p$ such that

$$|k_n(x_n, y_n) - k(I_n(x_n), I_n(y_n))| \leq S\epsilon$$

for some S which only depends on the constants A and $R > 0$ and a point $p \in \Omega$. It suffices to prove that I_n satisfies Property (3) in Definition 2.7. Recall that by the choice of $N \in \mathbb{N}$ in the proof of Proposition 7.5, for every $n \geq N$, we have

$$d_Z(p_n, p) \leq \delta'^2, \quad \frac{19c}{20} \leq c_n \leq \frac{21c}{20}, \quad \text{and} \quad d_H^Z(\Omega_n, \Omega) + d_H^Z(\partial\Omega_n, \partial\Omega) \leq \delta'^2 =: \tilde{\epsilon}.$$

where $0 < \delta' \leq \frac{c}{20} \wedge \frac{\tilde{c}}{20}$ and c_n , c , and \tilde{c} are the constants in Proposition 7.5 for Ω_n and Ω , respectively. Define a map $J : B_k(p, R) \rightarrow X_n$ by $x \in B_k(p, R) \mapsto x_n \in X_n$ such that $J(p) = p_n$ and $d_z(x_n, x) \leq \tilde{\epsilon}$. Then by exactly the same argument (think of X as X_n and X_n as X in the proof of Proposition 7.5), we have

$$|k(p, x) - k_n(p_n, J(x))| \leq S\epsilon,$$

for every $x \in B_k(p, R)$. This implies that

$$k_n(p_n, J(x)) \leq k(p, x) + S\epsilon < R$$

for every $x \in B_k(p, R - S\epsilon)$. Hence $J(x) \in B_{k_n}(p_n, R)$. Also, since

$$d_n(I_n(J(x)), x) \leq d_Z(I_n(J(x)), J(x)) + d_Z(J(x), x) \leq 2\tilde{\epsilon},$$

by taking a curve γ from $I_n(J(x))$ to x with the length $d(I_n(J(x)), x) + \tilde{\epsilon} \leq 3\tilde{\epsilon}$, we have

$$\begin{aligned} k(I_n(J(x)), x) &\leq \int_0^{l_d(\gamma)} \frac{1}{d(\gamma(t))} dt \\ &\leq \left(\frac{1}{d(x)} + \epsilon\right) l_d(\gamma) \\ &\leq \left(\frac{1}{c} + \epsilon\right) 3\tilde{\epsilon}. \end{aligned}$$

This implies that

$$B_k(p, R - S\epsilon) \subseteq (I_n(B_{k_n}(p_n, R)))_{3(\frac{1}{c} + \epsilon)\tilde{\epsilon}}.$$

Taking the maximum $S\epsilon \vee 3(\frac{1}{c} + \epsilon)\tilde{\epsilon}$, we complete the proof. \square

We close this paper by giving the following two corollaries. We would like to point out that by [BHK, Proposition 4.37], it is known that a proper geodesic roughly starlike Gromov hyperbolic space (X, d) is bilipschitz equivalent to the space (X^ϵ, k) where X^ϵ is the uniformized space and k is the quasihyperbolic metric with respect to d_ϵ . It is not straightforward, however, that the pointed Gromov-Hausdorff convergence $(X_n, d_n, p_n) \rightarrow (X, d, p)$ implies the pointed Gromov-Hausdorff convergence $(X_n^\epsilon, k_n, p_n) \rightarrow (X^\epsilon, k, p)$.

Corollary 7.7. *Let $(X_n, d_n, p_n)_n$ be a sequence of pointed proper geodesic M -roughly starlike δ -Gromov hyperbolic spaces convergent to a pointed metric space (X, d, p) . Then, for a fixed $0 < \epsilon \leq \epsilon_0(\delta)$, the sequence of uniformized spaces with quasihyperbolic metrics $(X_n^\epsilon, k_n, p_n)_n$ pointed Gromov-Hausdorff converges to (X^ϵ, k, p) .*

Corollary 7.8. *Let $(\Omega_n, d_n)_n$ be a sequence of bounded A -uniform length metric spaces convergent to a metric space (Ω, d) with respect to d_{GHB} . Then there exist $p_n \in \Omega_n$ and $p \in \Omega$ such that the sequence $(\Omega_n, k_{n,\epsilon})$ converges to (Ω, k_ϵ) with respect to d_{GHB} for a fixed $0 < \epsilon \leq \epsilon_0(\delta)$ where $(\Omega_n, k_{n,\epsilon})$ and (Ω, k_ϵ) are the uniformized spaces of (Ω_n, p_n, k_n) and (Ω, p, k) , respectively.*

References

- [A] H. Aikawa, *Potential-theoretic characterizations of nonsmooth domains*. Bull. London Math. Soc. **36** (2004), no. 4, 469–482.
- [BBS] A. Björn, J. Björn, and N. Shanmugalingam, *Bounded geometry and p -harmonic functions under uniformization and hyperbolization*, arXiv:1908.04644.
- [BSh] J. Björn and N. Shanmugalingam, *Poincaré inequalities, uniform domains and extension properties for Newton-Sobolev functions in metric spaces*. J. Math. Anal. Appl. **332** (2007), no. 1, 190–208.
- [Bo] M. Bonk, *Quasi-geodesic segments and Gromov hyperbolicity*. Geom. Dedicata **62** (1996), no. 3, 281–298.
- [BHK] M. Bonk, J. Heinonen, and P. Koskela, *Uniformizing Gromov hyperbolic spaces*. Astérisque **270** (2001), viii+99 pp.
- [BH] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*. Grundlehren der Mathematischen Wissenschaften **319**, Springer-Verlag, Berlin, 1999. xxii+643 pp.
- [BBI] D. Burago, Y. Burago, and S. Ivanov, *A course in metric geometry*, volume 33 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2001.

- [Bu] C. Butler, *Uniformizing Gromov hyperbolic spaces with Busemann functions*, arXiv:2007.11143.
- [Ch] J. Cheeger, *Differentiability of Lipschitz Functions on Metric Measure Spaces*, *Geom. funct. anal.* **9**, 428–517 (1999).
- [Ed] D.A. Edwards, *The structure of superspace*. Studies in Topology, Academic Press, 1975, 121-133.
- [GeO] F.W. Gehring and B.G. Osgood, *Uniform domains and the quasihyperbolic metric*. *J. Analyse Math.* **36** (1979), 50–74.
- [GH] E. Ghys and P. de la Harpe, *Le bord d'un espace hyperbolique*. Sur les groupes hyperboliques d'après Mikhael Gromov (Bern, 1988), 117–134, *Progr. Math.* **83**, Birkhäuser Boston, Boston, MA, 1990.
- [Gr] M. Gromov, *Groups of polynomial growth and expanding maps*, *Inst. Hautes Etudes Sci. Publ. Math.* (1981), no. 53, 53-73.
- [HK] P. Hajlasz and P. Koskela, *Sobolev met Poincaré.*, *Mem. Amer. Math. Soc.* **145**, no. 688 (2000): x+101
- [HKST] J. Heinonen, P. Koskela, N. Shanmugalingam, and J. Tyson, *Sobolev Spaces on Metric Measure Spaces: An Approach Based on Upper Gradients*, Cambridge University Press (2015).
- [H] D. Herron, *Gromov-Hausdorff distance for pointed metric spaces*, *J. Anal.* **24** (2016), 1-38.
- [HRS] D. Herron, A. Richard, and M. Snipes, *Chordal Hausdorff convergence and quasi-hyperbolic distance*, *Anal. Geom. Metr. Spaces* **8**(2020), 60-91
- [Kei] S. Keith, *Modulus and the Poincaré inequality on metric measure spaces*. *Mathematische Zeitschrift*, **245** (2003), no. 2, 255–292.
- [Mar] O. Martio, *Definitions for uniform domains*. *Ann. Acad. Sci. Fenn. Ser. A I Math.* **5** (1980), no. 1, 197–205.
- [MS] O. Martio and J. Sarvas, *Injectivity theorems in plane and space*. *Ann. Acad. Sci. Fenn. Ser. A I Math.* **4** (1979), no. 2, 383–401.
- [P] R. Perales, *Convergence of manifolds and metric spaces with boundary*, *Journal of Topology and Analysis* (2018), 1-40.
- [PS] R. Perales and C. Sormani, *Sequences of open Riemannian manifolds with boundary*, *Pac. J. Math.* **270**(2), 423-471 (2014).

- [V1] J. Väisälä, *Uniform domains*, Tohoku Math. J. 40(1) (1988), 101-118.
- [V2] J. Väisälä, *Gromov hyperbolic spaces*, Expo. Math. 23 (2005), no.3, 187-231.