

Normal subgroups of iterated wreath products of symmetric groups and alternating with symmetric groups

Ruslan V. Skuratovskii

Abstract

Normal subgroups and their properties for finite and infinite iterated wreath products $S_{n_1} \wr \dots \wr S_{n_m}$, $n, m \in \mathbb{N}$ and $A_n \wr S_n$ are founded. The special classes of normal subgroups and their orders are investigated. Special classes of normal subgroups are investigated and their generators are found and presented in the form of Kaloujnine tables. Further, the monolith (i.e. the unique minimal normal subgroup) of these wreath products is investigated.

Inverse limit of wreath product of permutation groups is found.

Key words: normal subgroups, iterated wreath product of symmetric groups, normal subgroups of wreath product of symmetric group and alternating group.

MSC: 20D10, 20F05, 20B05, 20B25, 20B22, 20B07, 20E08, 20E28, 20B35.

1 Introduction

This work is the continuation of the previous researches of the author [1, 29]. Also this research clarifies our previous results from [1]. One of the modern branches of group theory researches is a study of isometry of geometry structures as graph trees and of combinatorial-topological structures [34]. In many cases it is related with studying of groups of automorphisms and normal structures of these groups. Isometry group of Baire space of type (n_1, n_2, \dots) is isomorphic to spherically homogeneous tree with the same type. The results were presented on conferences [31, 32].

The modern direction of group theory is the so-called problems of studying groups of isometries of combinatorial - topological structures that are traditional for geometric theory groups. This is often closely related to the study of groups of automorphisms of different types of trees. Thus, the group of isometries of the Baire space type (n_1, n_2, \dots) is isomorphic to spherically homogeneous rooted tree with the same type.

One of the reasons for the interest in groups acting on trees is that various interesting examples of groups are naturally depicted as groups acting in particular on root trees,

for example, infinite finitely generated periodic groups. In addition, groups of automorphisms (isometries) of root trees are still being actively studied due to the fact that they are quite well arranged and immerse in them numerous finitely approximate groups containing a number of subgroups with different properties. Thus, the periodic groups of V.I. Sushchansky and R. I. Grigorchuk are embedded in the groups of automorphisms of such trees. Thus, it is natural to consider infinitely iterated wreath products of finite groups of permutations. In particular, the main purpose of this work is to describe all normal subgroups of the group of automorphisms of a finite (restricted) regular rooted tree. The paper develops tools for the study of the normal structure, based on the specifics of the wreath product of symmetric groups.

A form of commutators of wreath product $A \wr B$ was briefly considered in [5]. We consider the form of commutator presentation [5] in form of wreath recursion [7, 12] and as Kaloujnine tableaux, its properties as a normal subgroup is studied here. We impose our results about of normal subgroups of permutational wreath product [1].

In this paper we continue the research of subgroups of wreath product which was started in [1, 12, 19, 21, 22] in particular in [12] a commutator subgroup as a normal subgroup of wreath product with non-effective action was considered by us. Despite this, it was an open question whether the subgroup of $\wr_{i=1}^k \mathcal{S}_{n_i}$ is normal. obtain an exact commutator width. We generalize the passive group of this wreath product $S_n \wr S_n$ to any permutation subgroup B instead of only considering the wreath product of symmetric groups.

2 Preliminaries

Let G be a group. The intersection of all non-trivial normal subgroups $Mon(G)$ of G is called the monolith of a group G . If $Mon(G) \neq \langle 1 \rangle$, then the group G is called monolithic, and, in this case, $Mon(G)$ is the least non-trivial normal subgroup of G .

Definition 1. The unique minimal normal subgroup is called the monolith.

According to [11, 15] by tableaux we call all possible infinite tuple of the form:

$$a = [a_k]_{k \in \mathbb{N}}, \quad \text{where } a_k \in Fun^{(k-1)}(M, G_k), \quad k \in \mathbb{N}. \quad (1)$$

A group of isometries of the generalized Baire space E [6, 9, 10] over a family of sets M_k , $k \in Z$, we denote by I according to [6, 10, 15]. We suppose that for every $k \in N$ set M_k is finite.

A group of isometries I_{Z_p} of metric space of integer ring of p -adic numbers is isomorphic as a permutation group to the iterated wreath product of permutation groups of degree p .

Definition 2. We will say that the subset $U < G$ is determined by its k -coordinate sets $[U]_k$, $k \in N$, if this subset consists of the most possible tables $a \in I_{Z_p}$ such that for arbitrary $k \in N$ holds $[a]_k \in [U]_k$.

Definition 3. Let A be some set of the tableaux of form (1), then we will call by the k -projection of the set A the set $[A]_k = \{a|_k : a \in A\}$. We call the set A as k -coordinate set if $[A]_k = A$ holds.

Definition 4. We denote by $A|_k$ a set of all functions a_k such that $[\varepsilon, \dots, \varepsilon, a_k, \varepsilon, \dots] \in [A]_k$.

Recall that such tableau action on element $a = [a_k]_{k \in \mathbb{N}}$ on element $m = (m_k)_{k \in \mathbb{N}} \in M$ is defined by the rule on element:

$$m^a = (m_k^{a_k((k-1)m)})_{k \in \mathbb{N}}.$$

the iterated wreath product of permutation groups of degree p .

A group of isometries of the generalized Baire E [10, 15] [8, 7] over a family of sets M_k , $k \in N$, (then will be generalized to $k \in Z$), we will denote as I according to [8, 7]. We will consider that for every $k \in N$ set M_k is finite.

The set G of all such tableaux of the form (1) with such action which equipped by such multiplication rule forms a group. The neutral element is the table of form $e = [\dots, \varepsilon, \varepsilon, \dots]$.

Definition 5. We will say that the subset $U < G$ is determined by its k -coordinate sets $[U]_k$, $k \in N$, if this subset consists of the most possible tables $a \in I$ such that for arbitrary $k \in N$ holds $[a]_k \in [U]_k$. If U is a splitting subgroup [15] of the group I , which is defined in [15], then k -coordinate subset $[U]_k$ are subgroups of the group U , and $[U_k] = {}^{(k)}U \cap U^{(k)}$. Besides, family $[U]_k$, where $k \in N$ will generate a uniformly dense subgroup U' of a group U . The splitting normal subgroups of the group I were also studied in [10].

Let G be a group acting (from the right) by permutations on a set X and let H be an arbitrary group. Then the (permutational) wreath product $H \wr G$ is the semidirect product $H^X \rtimes G$, where G acts on the direct power H^X by the respective permutations of

the direct factors. The group C_p or (C_p, X) is equipped with a natural action by the left shift on $X = \{1, \dots, p\}$, $p \in \mathbb{N}$. It is well known that the wreath product of permutation groups is associative.

The multiplication rule of automorphisms g, h which are presented in the form of wreath recursion [7] $g = (g_{(1)}, g_{(2)}, \dots, g_{(d)})\sigma_g$, $h = (h_{(1)}, h_{(2)}, \dots, h_{(d)})\sigma_h$, is given by the formula:

$$g \cdot h = (g_{(1)}h_{(\sigma_g(1))}, g_{(2)}h_{(\sigma_g(2))}, \dots, g_{(d)}h_{(\sigma_g(d))})\sigma_g\sigma_h.$$

We define σ as $(1, 2, \dots, p)$, where p is clear from context.

The set X^* is naturally the vertex set of a regular rooted tree, i.e. a connected graph without cycles and a designated vertex v_1 called the root, in which two words are connected by an edge if and only if they are of form v and vx , where $v \in X^*$, $x \in X$. The set $X^n \subset X^*$ is called the n -th level of the tree X^* and $X^1 = \{v_1\}$. Denote the i -th the vertex of X^j by v_{ji} . Note that the unique vertex $v_{k,i}$ corresponding to each word v in alphabet X . For every automorphism $g \in \text{Aut}X^*$ and every word $v \in X^*$ we define the section (state) $g_{(v)} \in \text{Aut}X^*$ of g at v by the rule: $g_{(v)}(x) = y$ for $x, y \in X^*$ if and only if $g(vx) = g(v)y$. The subtree of X^* induced by the set of vertices $\cup_{i=1}^k X^i$ is denoted by $X^{[k]}$. The restriction of the action of an automorphism $g \in \text{Aut}X^*$ to the subtree $X^{[l]}$ is denoted by $g_{(v)}|_{X^{[l]}}$. The restriction $g_{(v_{ij})}|_{X^{[1]}}$ is called the vertex permutation (v.p.) of g at the vertex v_{ij} and denoted by g_{ij} . We refer to the endomorphism $\alpha|_v$ restriction of g in a vertex v [7]. For example, if $|X| = 2$ then we just have to distinguish active vertices, i.e. the vertices for which $\alpha|_v$ is non-trivial.

We label every vertex of X^l , $0 \leq l < k$ by sign 0 or 1 depending on the action of v.p. on it. value of α in v_{ki} as $s_{ki}(\alpha)$ we put that $s_{ki}(\alpha) = 1$ if $\alpha|_{v_{ki}}$ is non-trivial, The resulting vertex-labeled regular tree is an element of $\text{Aut}X^{[k]}$.

We remind a definition of level subgroup $G(l)$ presented by us in [27].

Definition 6. We define the subgroup of l -th level ($G(l)$) as a subgroup generated by all possible vertex permutations of this level.

Let us fix some notations. From here and throughout whole paper we will use left action in wreath product.

For convenience we will determine and present the conjugation of a by b by the expression

$$a^b = bab^{-1}.$$

3 Normal subgroups of wreath product of symmetric groups

3.1 Normal subgroups in $S_n \wr S_m$

We will notice at once, that further statements and expositions will be true for $n \geq 3$ even in case if degrees of two our symmetric groups from wreath product are different. Elements of our wreath product $S_n \wr S_n, n \geq 5$, will present in the form of tableaux, where $[a]_1$ – the active element of the table and $a \in S_n, [a_1, a_2, \dots, a_n]_2$ are the passive elements of the table, $a_i \in S_n$, viz.:

$$[a_1]_1, [a_1, a_2, \dots, a_n]_2.$$

The rule of multiplication of elements is described in book of Meldrum [5]. Let us give the following definitions.

Let $k(\pi)$ be the number of cycles in decomposition of permutation π of degree n . The number $n - k(\pi)$ is denoted by $dec(\pi)$, and is called a *decrement* [16] of permutation π .

As well known [16] the minimal number of transpositions in factorization of a permutation π on transpositions is happen to be equal to $dec(\pi)$. We set $dec(e) = 0$. Therefore the decrement of n -cycle is $n - 1$. Therefore the decrement of n -cycle is $n - 1$. If $\pi_1, \pi_2 \in S_n$, then the following formula holds:

$$dec(\pi_1 \cdot \pi_2) = dec(\pi_1) + dec(\pi_2) - 2m, m \in \mathbf{N}, \quad (2)$$

where m is number of joint simplifying transpositions in π_1 and π_2 . Note, that a decrement of permutation's product $\pi_1 \pi_2$ can be lesser then decrement of each permutation π_1, π_2 .

For example, independent transpositions from the Moore generating system [4] for A_n can be used for factorization of g in product of transpositions.

Definition 7. The set of elements from $S_n \wr S_n, n \geq 3$ presented by the tables [11] form $[e]_1, [a_1, a_2, \dots, a_n]_2$, satisfying the following condition

$$\sum_{i=1}^n dec([a_i]_2) = 2k, k \in \mathbf{N}, \quad (3)$$

we will call set of type $\tilde{A}_n^{(2)}$ and denote this set by $e \wr \tilde{A}_n$.

For brevity of notation this subgroup be also denoted by $\tilde{A}_n^{(2)}$. It follows directly from the definition that the set of these elements supplemented by the operation of multiplication in the subdirect product, coincides with the group $E \times \underbrace{(S_n \boxtimes S_n \boxtimes S_n \dots \boxtimes S_n)}_n$,

where subdirect product satisfies to condition (3).

In other words such subset of element g from $S_n \wr S_n, n \geq 3$ belongs to type $\widetilde{A}^{(2)}$ if g corresponds to automorphism of $X^{[2]}$ having on zero level trivial permutation and vertex permutations from $g \in \text{Aut}X^{[2]}$ on the first level form an even permutation.

It follows directly from the definition that the set of these elements supplemented by the operation of multiplication in the subdirect product, coincides with the group $E \times \underbrace{(S_n \boxtimes S_n \boxtimes S_n \dots \boxtimes S_n)}_n$, where subdirect product satisfies to condition (3).

Proposition 1. *The set of elements of type $\widetilde{A}_n^{(2)}$ forms subgroup $e \wr \widetilde{A}_n \triangleleft S_n \wr S_n$.*

Proof. Let us check that $e \wr \widetilde{A}_n$ is closed with respect to multiplication and existing of inverse element.

(1B, a) Let $\tilde{a}, \tilde{b}, \tilde{c}$ are elements of $e \wr \widetilde{A}_n$ presented by tableaux: $[e]_1, [a_1, a_2, \dots, a_n]_2$; $[e]_1, [b_1, b_2, \dots, b_n]_2$; $[e]_1, [c_1, c_2, \dots, c_n]_2$. Therefore inequality of their products $(\tilde{a} \cdot \tilde{b}) \cdot \tilde{c} = \tilde{a} \cdot (\tilde{b} \cdot \tilde{c})$ is based on the associativity of permutations multiplication.

(1B, b) The neutral element is $[e]_1, [e, e, \dots, e]_2$.

(1B, c) For every element of the form $[e]_1, [a_1, a_2, \dots, a_n]_2$ there is an inverse element $\tilde{a}^{-1} \in e \wr \widetilde{A}_n$ having form: $[e]_1, [a_1^{-1}, a_2^{-1}, \dots, a_n^{-1}]_2$.

Therein decrement permanency $\sum_{i=1}^n \text{dec}([a_i]_2) = \sum_{i=1}^n \text{dec}([a_i^{-1}]_2) = 2k, k \in \mathbb{N}, 0 \leq i \leq n$ holds. Therefore $e \wr \widetilde{A}_n$ is a subgroup in $S_n \wr S_n$.

For proving a normality of $e \wr \widetilde{A}_n$ in $S_n \wr S_n$ we consider a conjugation by an arbitrary element $s \in S_n \wr S_n$ and show that $s \cdot \tilde{a} \cdot s^{-1}$ belongs to $e \wr \widetilde{A}_n$. For this we take into account that for $s^{-1} \in S_n \wr S_n$ invariance of decrements with respect to taking the inverse element $\sum_{i=1}^n \text{dec}([s_i]_2) = \sum_{i=1}^n \text{dec}([s_i^{-1}]_2) = k$ holds. Let us denote the tableaux representation of $s \cdot \tilde{a} \cdot s^{-1}$ by $[b_1]_1, [b_1, b_2, \dots, b_n]_2$ by virtue of definition of $e \wr \widetilde{A}_n$ an element $\tilde{a} \in e \wr \widetilde{A}_n$ satisfies the following property of parity $\sum_{i=1}^n \text{dec}([a_i]_2) = 2k, k \in \mathbb{N}$. Then taking into account the multiplication rule in the wreath product of groups and (2) we have $[b_1]_1 = [s_1 \cdot e \cdot s_1^{-1}]_1 = [e]_1$, and also $\sum_{i=1}^n \text{dec}([b_i]_2) = \sum_{i=1}^n \text{dec}([s_i]_2) + \sum_{i=1}^n \text{dec}([a_i]_2) - 2a + \sum_{i=1}^n \text{dec}([s_i^{-1}]_2) - 2b = k + 2m - 2a + k - 2b = 2(k + m - a - b)$. Hence the sum is even that satisfies the condition (3).

Thus, $s \cdot \tilde{a} \cdot s^{-1} \in e \wr \widetilde{A}_n$. This accomplished the proof. \square

Now we can recursively construct easiest and elegant subgroup $E \wr \widetilde{A}_n^{(2)}$ of $S_n \wr S_n \wr S_n$.

Definition 8. *The subgroup $E \wr \widetilde{A}_n^{(2)}$ be denoted by $\widetilde{A}_n^{(3)}$.*

Furthermore we will prove that $E \wr \widetilde{A}_n^{(2)} \triangleleft S_n \wr S_n \wr S_n$. The order of $E \wr \widetilde{A}_n^{(2)}$ is $(n!)^{3n} : 2^3$.

Definition 9. The set of elements from $S_n \wr S_n \wr S_n, n \geq 3$ presented by the tables [11] form:

$[e]_1, [e, e, \dots, e]_2, [a_1, a_2, \dots, a_n]_3$, satisfying the following condition

$$\sum_{i=1}^n dec([a_i]_3) = 2k, k \in \mathbb{N}, \quad (4)$$

be denoted by $\tilde{A}_{n^2}^{(3)}$.

Proposition 2. *The set of elements of type $\tilde{A}_{n^2}^{(3)}$ forms a subgroup in $S_n \wr S_n \wr S_n$. Moreover $\tilde{A}_{n^2}^{(3)} \triangleleft S_n \wr S_n \wr S_n$.*

Remark 3. *We note that $\tilde{A}_n^{(3)} < \tilde{A}_{n^2}^{(3)}$. The order of $\tilde{A}_{n^2}^{(3)}$ is $(n!)^{n^2} : 2$. Furthermore $\tilde{A}_n^{(3)} \triangleleft S_n \wr S_n \wr S_n$.*

Remind that the condition on commutator subgroup of permutation wreath product base was obtained by us in [12], and commutator subgroup is also normal.

The condition (4) accords with the condition setting the subdirect product in a commutator subgroup of wreath product base [12].

Definition 10. *A subgroup in $S_n \wr S_n$ is called \tilde{T}_n if it consists of:*

1. *elements of $E \wr A_n$,*
2. *elements with the tableau [11] presentation $[e]_1, [\pi_1, \dots, \pi_n]_2$, that $\pi_i \in S_n \setminus A_n$.*

One easy can validates a correctness of this definition, i.e. that the set of such elements form a subgroup and its normality. This subgroup has structure

$$\tilde{T}_n \simeq \underbrace{(A_n \times A_n \times \dots \times A_n)}_n \rtimes C_2 \simeq \underbrace{S_n \boxplus S_n \dots \boxplus S_n}_n,$$

where the operation \boxplus of a subdirect product is subject of item 1) and 2).

Definition 11. *A subgroup in $S_n \wr S_n \wr S_n$ is of the type $\tilde{T}_{n^2}^{(3)}$ if it consists of:*

1. *elements of the form $E \wr E \wr A_n$,*
2. *elements with the tableau [9] presentation $[e]_1, [e \dots, e]_2, [\pi_1 \dots, \pi_n, \pi_{n+1} \dots, \pi_{n^2}]_3$, wherein $\forall i = 1, \dots, n: \pi_i \in S_n \setminus A_n$.*

We define recursively the subgroup $\tilde{T}_n^{(3)}$ having n different intervals of elements with the same parity permutations on X^2 .

Definition 12. The subgroup of $S_n \wr S_n \wr S_n$ having structure $E \wr \widetilde{T}_n^{(3)}$ is denoted by $\widetilde{T}_n^{(3)}$. The following isomorphism $\widetilde{T}_n^{(3)} \simeq \underbrace{S_n \boxplus S_n \dots \boxplus S_n}_n \times \underbrace{S_n \boxplus S_n \dots \boxplus S_n}_n \times \dots \times \underbrace{S_n \boxplus S_n \dots \boxplus S_n}_n$, where a tuple $S_n \boxplus S_n \dots \boxplus S_n$ repeats n times, holds. The operation \boxplus is determined in Definition 10.

The operation \boxplus accords with the properties described in item 1 and 2 of Definition (10), also \boxplus is determined by automorphism in $\widetilde{T}_n \simeq \underbrace{(A_n \times A_n \times \dots \times A_n)}_n \rtimes C_2$ in this case.

Remark 4. Note that in $\widetilde{T}_n^{(3)}$ vertex permutation of tableau third part $[e]_1, [e \dots, e]_2, [\pi_1 \dots, \pi_n; \pi_{n+1} \dots, \pi_{2n}]_3$ satisfy the condition that either all $\pi_{i_j} \in S_n \setminus A_n$ or all $[\pi_{i_j}]_3 \in A_n$ for $1 \leq i_1 \leq n, n+1 \leq i_2 < 2n, \dots, n^2 - n + 1 \leq i_n \leq n^2$.

Let $\bigwedge_{i=1}^k S_{n_i}$ be denoted by W_k .

Remark 5. The set of of tableau [11] with vertex permutation in of tableau [11] 3-rd part $[e]_1, [e \dots, e]_2, [e \dots, e]_3, [\pi_1 \dots, \pi_n; \pi_{n+1}, \dots, \pi_{n^3}]_4$, satisfying the condition that either all $\pi_{i_j} \in S_n \setminus A_n$ or all $[\pi_{i_j}]_4 \in A_n$ for $1 \leq i_1 \leq n^2, n^2 + 1 \leq i_2 < 2n^2, \dots, (n-1)n^2 + 1 < i_n \leq n^3$ be denoted by $\widetilde{T}_{n^2}^{(4)}$. Furthermore $\widetilde{T}_{n^2}^{(4)}$ forms normal subgroup in $St_W(4)$.

The **interesting problem** is to describe all possible sets of generators of $e \wr \widetilde{A}_n$ with respect of conjugation in $S_n \wr S_n$. Let us denote the element of $e \wr \widetilde{A}_n$ having tableaux presentation $[e]_1, [(12), (12), e, \dots, e]_2$ by \widetilde{a} .

Proposition 6. The normal closure of \widetilde{a} in $S_n \wr S_n$ is $e \wr \widetilde{A}_n$.

Proof. In order to prove this it is sufficient to show that $N(\widetilde{a})$ is closed in $e \wr \widetilde{A}_n$ with respect to multiplication and conjugation.

1) this element of $N(\widetilde{a})$ is closed with respect of conjugation by elements of $S_n \wr S_n$ in the subgroup $e \wr \widetilde{A}_n$ i.e. $\forall s \in S_n \wr S_n, s \cdot a \cdot s^{-1} \in N(\widetilde{a})$.

2) In Proposition 1 is shown that multiplication of two arbitrary elements $\forall a, b \in N(\widetilde{a})$ satisfies the closeness property $a \cdot b \in N(\widetilde{a})$. But it remains to show that all elements of $e \wr \widetilde{A}_n$ can be generated by elements from $N(\widetilde{a})$.

To describe form of such a, b we denote an element of $e \wr \widetilde{A}_n$ having form of the tableaux $[e]_1, [\pi_1, \pi_2, \dots, \pi_n]_2$, where $\pi_1, \pi_2, \dots, \pi_n \in S_n$ and $\sum_{i=0}^n dec([\pi_i]_2) = 2k, k \in \mathbb{N}$ by $a_{\pi_1, \pi_2, \dots, \pi_n}$. In particular $\widetilde{a} = a_{(12), (12), e, \dots, e}$.

We make the constructive prove showing that every element satisfying condition (3) belongs to $(\widetilde{a})^{S_n \wr S_n}$.

(a) Let $\pi_1 = (i_1 i_2), \pi_2 = (j_1 j_2), \pi_3 = \dots = \pi_n = e$. Then $a_{\pi_1, \pi_2, \dots, \pi_n} = s \cdot \tilde{a} \cdot s^{-1} \in N(\tilde{a})$, where s has tableaux of form: $[e]_1, [((i_1 i_2)(2i_2)), ((j_1 j_2)(2j_2)), e, e, \dots, e]_2$.

(b) Let $\pi_1 = (i_1 i_2)(i_3 i_4), \pi_2 = \dots = \pi_n = e$. Then an element $a_{\pi_1, \pi_2, \dots, \pi_n}$ can be expressed as product of elements of type (a) where $\pi_2 = (1, 2)$: $b_{(i_1 i_2), (i_1 i_2), e, \dots, e} = a_{(i_1 i_2), (1, 2), \dots, \pi_n} \cdot a_{(i_3 i_4), (1, 2), \dots, \pi_n} \in N(\tilde{a})$. So as A_n can be generated by pairs of transpositions then $A_n \subseteq N(\tilde{a})$.

Analogous case is $\pi_2 = (j_1 j_2)(j_3 j_4), \pi_1 = \pi_3 = \dots = \pi_n = e$ and the similar reasoning as above are applicable.

(c) If $\pi_1 = (i_1 i_2) \dots (i_{2x-1} i_{2x}), \pi_2 = (j_1 j_2) \dots (j_{2y-1} j_{2y}), \pi_3 = \dots = \pi_n = e$. Without losing the generality, we assume $x \geq y$. Since $dec(\pi_1 \pi_2) = 2k$, then $(x - y) \equiv 0 \pmod{2}$, since: $a_{\pi_1, \pi_2, \dots, \pi_n} = a_{(i_1 i_2), (j_1 j_2), e, \dots, e} \cdot \dots \cdot a_{(i_{2x-1} i_{2x}), (j_{2x-1} j_{2x}), e, \dots, e} \cdot a_{(i_{2x+1} i_{2x+2})(j_{2x+3} j_{2x+4}), e, \dots, e} \cdot \dots \cdot a_{(i_{2y-4} i_{2y-3})(j_{2y-1} j_{2y}), e, \dots, e} \in N(\tilde{a})$. Hence such elements are the product of the elements which are described above in the items (a) and (b).

(d) If $\pi_1 = (i_1 i_2) \dots (i_{4x-1} i_{4x}), \pi_2 = \dots = \pi_n = e$. Then we can write $a_{\pi_1, \pi_2, \dots, \pi_n}$ as product of elements from item (b): $b_{(i_1 i_2), (i_3 i_4), e, \dots, e} \cdot \dots \cdot b_{(i_{4x-4} i_{4x-3})(j_{4x-1} j_{4x}), e, \dots, e} = a_{\pi_1, \pi_2, \dots, \pi_n} \in N(\tilde{a})$. Thus arbitrary even permutation π_1 of first coordinate of table can be expressed.

(e) If $\pi_i \neq e, \pi_j \neq e, dec(\pi_i \pi_j) = 2z, z \in \mathbb{N}, \pi_k = e, k \neq i, j$. Then $a_{\pi_2, \dots, \pi_n} = s \cdot a_{\pi_1, \pi_2, e, \dots, e} \cdot s^{-1} \in N(\tilde{a})$, where s has tableau presentation: $[(1i)(2j)]_1, [e, \dots, e]_2$. So such elements are conjugated to the elements described in (c).

(f) If $\pi_i \neq e, dec(\pi_i) = 2z, z \in \mathbb{N}, \pi_k = e, k \neq i$. Then $a_{\pi_1, \pi_2, \dots, \pi_n} = s \cdot a_{\pi_1, e, \dots, e} \cdot s^{-1} \in N(\tilde{a})$, where s has presentation: $[(1i)]_1, [e, \dots, e]_2$. So such elements are conjugated to the elements described in (d).

(g) If $\pi_i \neq e, i = \overline{1, n}$ and the order of permutation is even i.e. $\sum_{i=1}^n dec([\pi_i]) = 2z, z \in \mathbb{N}$ then $a_{\pi_1, \pi_2, \dots, \pi_n}$ is a product of the elements described in (e), (f).

So the group closure is $e \wr \widetilde{A}_n$.

Thus, the normality of $E \wr \widetilde{A}_n$ is proved in new way. \square

The generalization of \tilde{a} which is $[e]_1, [e, e, \dots, e, \pi_i, e, \dots, \pi_j, \dots, e]_2 = a_{e, \dots, e, (i_1 i_2), e, \dots, e, (j_1 j_2), e, \dots, e}$, where $\pi_i = (i_1 i_2), \pi_j = (j_1 j_2) \in S_n$ are transpositions on i -th and j -th coordinate, we denote by \tilde{b} .

Remark 7. The normal closure of \tilde{b} is $e \wr \widetilde{A}_n$.

Proof. To established the proof it is sufficient to show $\tilde{a} \in N(\tilde{b})$, since the inclusion $\tilde{b} \in N(\tilde{a})$ follows from the previous statement.

For this goal we express $a_{e, \dots, e, (12), e, \dots, e, (12), e, \dots, e} = a_{e, \dots, e, (i_1 i_2), e, \dots, e, (j_1 j_2), e, \dots, e}$.

$$\tilde{b} \cdot (a_{e, \dots, e, (i_1 1 i_2 2), e, \dots, e, (j_1 1 j_2 2), e, \dots, e})^{-1} \in N(\tilde{b}).$$

Suppose an expressed element has non trivial permutations on coordinates i and j . Then

$$\tilde{a} = s \cdot a_{e, \dots, e, (12), e, \dots, e, (12), e, \dots, e} \cdot s^{-1} \in N(\tilde{b}),$$

where s has tableau form as $[(1i)(2j)]_1, [e \dots, e]_2$. □

Let us denote by \tilde{c} an element of $e \wr \widetilde{A}_n$ having the tableaux presentation $[e]_1, [(123), e, \dots, e]_2$.

Proposition 8. *The normal closure of \tilde{c} is $E \wr A_n$.*

Proof. We use for the proving notifications of previous statements.

In order to prove normality of the $e \wr A_n$ we show that it is normal closure of \tilde{c} in $S_n \wr S_n$.

(a) If $\pi_1 \in A_n, \pi_2 = \dots = \pi_n = e$ then such element \tilde{c} be denoted by $c_{\pi_1, e, \dots, e}$.

Therefore the element $a_{\pi_1, \pi_2, \dots, \pi_n} \in N(\tilde{c})$ can be obtained by using of transformation of first and second type from Proposition 6 (conjugations and multiplications) of an element \tilde{c} .

Since $A_n \triangleleft S_n$ and A_n is simple, then a normal closure its elements coincides with hole A_n . Hence an element $c_{\pi_1, e, \dots, e} \in N(\tilde{c})$, where π_i is 3-cycle, can be expressed from \tilde{c} by using transformations from prof of previous statements.

(b) If $\pi_i \neq e, \pi_i \in A_n$, but for another $k \neq i$ we set $\pi_k = e$, then

$$c_{e, \dots, \pi_i, \dots, e} = s \cdot c_{\pi_1, e, \dots, e} \cdot s^{-1} \in N(\tilde{c}),$$

where s is presented by the tableau: $[(1i)]_1, [e \dots, e]_2$. Therefore, $c_{e, \dots, \pi_i, \dots, e}$ are conjugated with elements from item (a).

(c) If $\pi_i \in A_n$, for every $i = \overline{1, n}$. Then $c_{\pi_1, \pi_2, \dots, \pi_n}$ can be expressed as a product of elements described in the item (b).

Thus, all elements of $e \wr A_n$ are constructed in the items (a)-(c), that is what accomplished the proof. □

Corollary 9. *The subgroup $E \wr A_n$ is normal in $S_n \wr S_n$.*

Proof. Since we proved that the $E \wr A_n$ is normal closure of \tilde{c} in $S_n \wr S_n$ then $E \wr A_n$ is normal. □

Corollary 10. *The normal rank [17] of $E \wr \widetilde{A}_n$ is 2.*

Proof. If we take normal closure only of \tilde{c} then we generate only subgroup $E \wr A_n$. But if we restrict generating set of normal closure on \tilde{a} then we obtain in normal closure whole $E \wr \tilde{A}_n$ because of an element $\tilde{a} = a_{(12),(12),e,\dots,e} \in E \wr \tilde{A}_n$. \square

Corollary 11. *Previous statement remains to be correct for the normal closure $N(\tilde{b})$ of $\tilde{b} = a_{e,\dots,e,(i_1 i_2 i_3),e,\dots,e}$, $i_1, i_2, i_3 \in \{1, 2, \dots, n\}$.*

Proof. To prove the statement it is sufficient to establish $\tilde{a} \in N(\tilde{b})$, because of the belonging $\tilde{b} \in N(\tilde{a})$ follows from the $\tilde{a} \in N(\tilde{b})$.

Let us prove it. So we have

$$a_{e,\dots,e,(123),e,\dots,e} = a_{e,\dots,e,(i_1 i_2 i_3),e,\dots,e} \cdot \tilde{b} \cdot (a_{e,\dots,e,(i_1 i_2 i_3),e,\dots,e})^{-1} \in N(\tilde{b}).$$

Without losing the generality we can set non trivial permutations in obtained an element is on coordinate j . Therefore using the conjugation by the tableau $[(1j)]_1, [e, \dots, e]_2$ we obtain the mentioned above element \tilde{c} : $\tilde{c} = s \cdot a_{e,\dots,e,(123),e,\dots,e} \cdot s^{-1} \in N(\tilde{b})$. Thus, the required element \tilde{c} was expressed in $N(\tilde{b})$. \square

Proposition 12. *The subgroup $A_n \wr A_n \wr A_n$ of $S_n \wr S_n \wr S_n$ has normalizer*

$$N_{S_n \wr S_n \wr S_n}(A_n \wr A_n \wr A_n) \simeq S_n \wr A_n \wr A_n.$$

Remark 13. *As previously [33] we denote l -th level subgroup by $AutX^{[k]}(l)$ i.e. such subgroup of $AutX^{[k]}$ that contains all v.p. from X^l , $l < k - 1$. In other words it contains all v.p. from $Stab_{AutX^{[k]}(l)}$ and do not contains v. p. from $Stab_{AutX^{[k]}(l+1)}$, $l < k - 1$. In our paper about sylow 2-subgroups we introduced a concept of $G_k(k-1)$ subgroup of $AutX^{[k]}$ that consists of v.p. which are located on X^{k-1} and isomorphic to W_{k-1} [33]. Now we note that $G_k(k-1)$ contains $\tilde{A}_{n^k}^{(k)}$.*

Definition 13. An element g of iterated wreath product $\prod_{i=1}^{\infty} S_{n_i}$ is said to be a finite if, starting from some level, all vertex permutations of the level are trivial.

We define *fixator* of a permutation $g \in S_n$ as the number of elements of $\{1, \dots, n\}$, which are fixed under action of g , and denote a fixator by $Fix(g)$.

Definition 14. Tableau u has **depth** m if all $[u]_i = \varepsilon$ provided $i = 1, 2, \dots, m - 1$; and $[u]_m \neq \varepsilon$.

Definition 15. We denote by $W^{(k)}$ the subgroup of depth $\geq k$ and ${}^{(k)}W$ is subgroup of tableaux $[u]_{\alpha} = \varepsilon$ if $\alpha > k$.

A mapping $g|_v$ is an automorphism of tree X^* and $g|_v$ calls *restriction* of g on word v or the *state* g in word v . This mapping $g|_v$ defined by the following rule $g|_v : X^* \rightarrow X^*$ iff $g(vx) = uy$, where $|x| = |y|$ and $g(vx) = uy$.

Automorphism g is called *finite* if it exists $n \geq 1$ such that $g|_v = e$ for all $v \in X^*$, $|v| \geq n$. The set of all finite automorphisms generates the group $Aut_f X^*$.

Let N normal subgroup in $Aut_f X^*$ possessing the even product of vertex permutation inside each subgroup $RiSt_W(v_k)$ [7]. If $H \triangleleft \prod_{i=1}^{\infty} S_{n_i}$ of depth k and $H \triangleright N$, where $k \in N$, $n_i \geq 5$ and m can tends to ∞ , then the subgroup H contains proper normal subgroup $P \in Aut_f X^*$ which have even permutations in all vertexes of k -th level states. Such permutation can be generated as left based commutators of level permutation of $\prod_{i=1}^{\infty} S_{n_i}$ with depth no less than k . Now we are going to prove it.

Since $\prod_{i=1}^m S_{n_i} \simeq Aut X^{[m]}$, then we make our proof in terms of states of automorphisms of rooted spherically homogeneous threes. Let $H \triangleleft Aut X^*$ be a proper normal subgroup with depth k . For convenience of notation we denote $Aut X^*$ by G .

We call level of $Aut X^*$ as active if it has at least one non-trivial permutation. A maximal active level exists if all tableau are locally bounded [9]. Denote by $Aut_f X^*$ the group of all finite automorphism of spherically homogeneous rooted tree.

Proposition 14. *Let $H \triangleleft Aut_f X^*$ with depth k then H contains k -th level subgroup $P = H(k)$ having all even vertex permutations $p_{ki} \in A_n$ on X^k and trivial permutations in vertices of rest of levels. Furthermore P is normal in $Aut_f X^*$ provided k is last active level of $Aut_f X^*$.*

Furthermore states of $h \in H$ on X^k possess an even product of vertex permutations belonging to subtree $v_{ki} X^$ for each level $j : j > k$. i.e.*

$$\prod_{j>k, t=(i-1)n^{j-k}}^{in^{j-k}} h_{jt} \in A_n,$$

for each $j > k$.

If subgroup H of depth k contains normal subgroups with increasing depth $j \geq k$, then H contains all level subgroups P_j having all even vertex permutations $p_{ji} \in A_n$, where $k \geq j \geq m$.

If depth of H is 0, then the normality of H yields the possibility to express in v_0 an arbitrary element of A_{n_1} .

Let v be a vertex of the k -th level of $Aut X^{[m]}$ with the non-trivial vertex permutation σ , we set, without loosing the generality, that $\sigma = (1, 2, \dots, l)$, $\sigma \in S_{n_k}$. In general

case the permutation σ can consists of some cycles but for the proof it is sufficient to consider one cycle. Therefore a state of automorphism in v can be presented in the form of the following tableau $x = (b_1, b_2, \dots, b_{l-1}, b_l)\sigma$, $b_i \in Aut_v X^{[m-k]}$ or equivalently $b_i \in \mathcal{L}_{i=k+1}^{m-1} S_{n_i}$, $k \leq m$, $l = n_k$. Let $y = (e, e, \dots, e, e)\sigma_1$, where $\sigma_1 = (1, 2, \dots, l-2)$, then $z = yxy^{-1} = (b_2, b_3, \dots, b_{l-2}, b_1, b_{l-1}, b_l)\sigma_1\sigma_1^{-1}$, and $zx^{-1} = yxy^{-1}x^{-1} = [y, x] = zx^{-1} = (a_1, \dots, a_{l-2}, e, e)[\sigma, \sigma_1]$, where $a_1 = b_2b_1^{-1}$, $a_2 = b_3b_2^{-1}$, ... , because of $x^{-1} = (b_l^{-1}, b_1^{-1}, b_2^{-1}, \dots, b_{l-1}^{-1})\sigma^{-1}$.

Since H is normal then $\sigma_1\sigma\sigma_1^{-1} \in H$ and in view of $H \triangleright N$ the we can choose σ from A_n therefore $\sigma_1\sigma\sigma_1^{-1}$ is even permutation again.

Now we see that zx^{-1} expressed by us is the even permutation, moreover $zx^{-1} = [y, x]$. Choosing degree $l-i$, $2 \leq i < l$ of σ_1 from y we can express in $v \in X^k$ each even permutation in form $[y, x]$ with arbitrary degree no grater then n . Alternatively we can put $\sigma_1 = (i, j)$ $\sigma = (i, k)$ then their commutator $[(i, j), (i, k)] = (i, k, j)$ to generate even permutation in v . After we take a normal closure of this automorphisms which contain \tilde{a} and \tilde{b} by the same method as in Proposition (6) and Remark (7). But degree of σ_1 and σ are equal in this case introducing some restrictions on the construction of permutations on $k+1$ level.

The product of all vertex permutations of $k+1$ level is even that satisfy the condition (10).

Then we can spread this method on the rest of vertexes of X^k using conjugating by elements of $Aut_f X^*$ for taking normal closure of this automorphisms which contain \tilde{a} and \tilde{b} by the same method as in Proposition (6) and Remark (7) to generate even permutations on the rest of vertexes of X^k .

If we have nested in H proper normal subgroup H_{k+1} of depth $k+1$ then we can apply recursively this method in each active vertex $v_{k+1,i}$ of X^{k+1} to obtain even permutation satisfying on first level of subtree $v_{k+1,i}X^*$ the condition (3). After this generating such elements with even permutation on X^k we can takes normal closure of this automorphisms using the same method as in Proposition (6) and Remark (7) to generate even permutations on rest of vertexes of X^{k+1} .

Thus, the subgroup $N \triangleleft H$ such that $N < Aut_f X^*$ having depth k -th and possessing the property of even product of level vertexes permutations (4) for each vertex subgroup $rist(v_{ki})$ is constructed.

For constructing in H an arbitrary even permutation on a first level of a state g in v we have to variate not only a degree of σ_1 but also a fixator of σ_1 . Thus, we can obtain arbitrary even permutation in v_{ki} . Vertex permutations on the next levels can be as even as odd but they form even level permutation.

It remains to express from $[y, x]$ states of k -th level with trivial permutation on a next $k + i$, $i > 0$ levels. Consider the element of $AutX^*$ stabilizing elements in vertexes $1, \dots, l-2$ i.e. the element of form $g = (e, e, e, e, e)(l-1, l)$. In general case, such g can be chosen from $(1, \dots, l) \setminus Fix_H(y)$ with respect to the restriction of action in v is such that $g|_v = (e, e, e, e, e)(l-1, l)$. Due to H is normal in G the inclusion $gyxy^{-1}x^{-1}g^{-1} \in H$ holds. We chose $y \in St_G(v_1) \cap St_G(v_2) \cap \dots \cap St_G(v_{l-2})$ and g from complementary vertex stabilizer intersection of y i.e. $g \in St_G(v_{l-1}) \cap St_G(v_l)$.

Since automorphism in H is finite then using recursive procedure described above we can construct even vertex permutation in vertexes of all levels in finitely many steps.

Now we conjugate $gyxy^{-1}x^{-1}$ by such $g = (e, e, e, e, e)(l-1, l)$, which permutes only trivial elements in tuple $(a_1, \dots, a_{l-2}, e, e)\sigma$ fixing all non trivial elements, this the tuple presents the element $gyxy^{-1}x^{-1}$ in the form of wreath recursion. Thus, obtained by us state in v_{ki} arbitrary even permutation in root trivial permutation on the rest levels.

The obtained element $gyxy^{-1}x^{-1}g^{-1} = (a_1, a_2, \dots, a_{l-2}, e, e)\pi$ we multiply by the inverse element to commutator $[y, x]$: $[x, y]gyxy^{-1}x^{-1}g^{-1} = (zx^{-1})^{-1}gyxy^{-1}y^{-1}g^{-1} = (zx^{-1})^{-1}gzx^{-1}g^{-1} = [x, y]g[y, x]g^{-1} = (e, e, e, e, e)(\tau)$, where $\tau = [[\sigma_1, \sigma], \pi]$, σ_1 is a permutation from the first level of the state of portrait automorphism y , σ is a root permutation of state of automorphism of x in v and π is root permutation of state of $gyxy^{-1}x^{-1}g^{-1}$ in v . In terms of commutators $(zx^{-1})^{-1}gzx^{-1}g^{-1}$ can be presented as $[zx^{-1}, g] = [[y, x], g]$. Thus, we obtain the automorphism $\rho = [[y, x], g]$ having trivial states in vertexes of each $k + j$ levels, where $1 \leq j \leq m - k$, and a permutation ρ on the k -th level in v_{ki} . Being a left based commutator ρ is an even permutation from S_{n_k} , therefore $\rho \in A_{n_k}$. Hence, such a vertex permutation can be generated as left based commutators of 3 automorphisms from $AutX^{[m]}$.

To complete the proof it is necessary to show each element with depth k corresponds to an element with the same depth l and having all but one vertex permutations on X^k are trivial. We emphasize that non - trivial vertex permutation on X^l in v_{lm} is even because of it expressed as a commutator $[[x, y], c]$. It remains to prove that in rest of vertices of l -th level we can obtain trivial permutations. For this goal we can use the same procedure as for constructing in vertex

In the similar way elements of arbitrary depth i having an even permutation on i -th level and trivial permutations on other levels can be constructed.

Corollary 15. *If the group $Aut_f X^*$ of all finite automorphism of spherically homogeneous rooted tree has all normal subgroups with each depth $0 \leq k \leq m$, then containing all level subgroups $H(l)$, $l \leq m$, where all vertex permutation are even.*

In Proposition 14 we express tuple of arbitrary even vertex permutation on k -th level. Now we can spread this method of constructing on rest of level subgroups $H_{Aut_f X^*}(l)$, $2 \leq l \leq m$ with even vertex permutations.

Consider first level consider the first level X^0 of the tree X^* separately. The even permutation in v_0 . In the case of existing in H automorphism h_0 having non-trivial permutation only in v_0 we compute h_0^2 which is even permutation.

In opposite case of non-existing such h_0 of heith 1 we make the same computation $[xz^{-1}, g] = [[y, x], g]$ to form on X^1 tuple of even vertex permutations. Note that obtained automorphism $[[y, x], g]$ have even permutation in v_0 and trivial permutation on X^1 with the same property described above in Proposition 14.

Having automorphisms with constructed above level subgroups elements of levels $0 \leq l \leq m$ with even permutation we can multiply them to express arbitrary $h \in \bigwedge_{i=1}^m A_{n_i} \triangleleft Aut_f X^*$.

If we consider $S_3 \wr S_2$, then we find 7 proper normal subgroups: $[[\text{"1"}, 1], [\text{"A}_4\text{"}, 1], [\text{"C}_2\text{"}, 1], [\text{"C}_2 \times \text{A}_4\text{"}, 1], [\text{"C}_2 \times \text{C}_2\text{"}, 1], [\text{"C}_2 \times \text{C}_2 \times \text{C}_2\text{"}, 1], [\text{"C}_2 \times \text{S}_4\text{"}, 1], [\text{"S}_4\text{"}, 2]]$.

Remark 16. *The following isomorphism $S_3 \wr S_2 \simeq C_2 \times S_4$ is true.*

Proof. The proof follows from direct computations. □

For instance we analysis all normal subgroups of $S_3 \wr S_3$. Computations in GAP lead us to following results and conclusions:

The group $S_3 \wr S_3$ has 10 normal subgroups, where $G_0 = E$ and $G_1 = S_3 \wr S_3 = \langle (1, 2, 3), (1, 2), (4, 5, 6), (4, 5), (7, 8, 9), (7, 8), (1, 4, 7)(2, 5, 8)(3, 6, 9), (1, 4)(2, 5)(3, 6) \rangle$. The list of proper normal subgroups with respect of their generators is the following:

$$G_2 = \langle (7, 8, 9), (4, 5, 6), (1, 2, 3) \rangle,$$

$$G_3 = \langle (7, 8, 9), (4, 5, 6), (1, 2, 3), (2, 3)(8, 9), (2, 3)(5, 6) \rangle,$$

$$G_4 = \langle (7, 8, 9), (4, 5, 6), (1, 2, 3), (1, 4, 7)(2, 5, 8)(3, 6, 9), (2, 3)(5, 6) \rangle,$$

$$G_5 = \langle (7, 8, 9), (4, 5, 6), (1, 2, 3), (1, 4)(2, 6)(3, 5)(8, 9), (1, 4, 7)(2, 5, 8)(3, 6, 9), (2, 3)(5, 6) \rangle,$$

$$G_6 = \langle (7, 8, 9), (4, 5, 6), (1, 2, 3), (1, 4)(2, 5)(3, 6), (1, 4, 7)(2, 5, 8)(3, 6, 9), (2, 3)(5, 6) \rangle,$$

$$G_7 = \langle (2, 3)(5, 6)(8, 9), (7, 8, 9), (4, 5, 6), (1, 2, 3) \rangle,$$

$$G_8 = \langle (8, 9), (5, 6), (2, 3), (7, 8, 9), (4, 5, 6), (1, 2, 3) \rangle,$$

$$G_9 = \langle (1, 4, 7)(2, 5, 8)(3, 6, 9), (8, 9), (5, 6), (2, 3), (7, 8, 9), (4, 5, 6), (1, 2, 3) \rangle.$$

In total we have 8 proper normal subgroups in $S_3 \wr S_3$.

By analyzing their structures in terms of a semidirect product and determination them as the subgroup of wreath product with respect of their orders we obtain:

1) $S_3 \wr \widetilde{A}_3 \simeq (((C_3 \times C_3 \times C_3) \times (C_2 \times C_2)) \times C_3) \times C_2, 3]$ here are 3 non equal isomorphic subgroups, also we compute $ord(S_3 \wr \widetilde{A}_3) = ((6^3) : 2) \cdot 6 = 3^4 \cdot 2^3 = 108 \cdot 6 = 648$.

2) $\langle (1, 2, 3), (1, 2), (4, 5, 6), (4, 5), (7, 8, 9), (7, 8), (1, 4, 7)(2, 5, 8)(3, 6, 9), (1, 4)(2, 5)(3, 6) \rangle = S_3 \wr S_3$.

All subgroups, except G_7 , from this list can be easily identified and presented with new indexing.

We denote $S_3 \wr S_3$ by H_1 and we compute that $[H_1 = (((C_3 \times C_3 \times C_3) \rtimes (C_2 \times C_2)) \rtimes C_3) \rtimes C_2 \rtimes C_2, 1]$. Order of H_1 is 1296.

We find the structures of these 8 proper normal subgroups of $S_3 \wr S_3$:

$$H_2 = [(((C_3 \times C_3 \times C_3) \rtimes (C_2 \times C_2)) \rtimes C_3) \rtimes C_2, 3],$$

$$H_3 = [((C_3 \times C_3 \times C_3) \rtimes (C_2 \times C_2)) \rtimes C_3, 1],$$

$$H_4 = [(C_3 \times C_3 \times C_3) \rtimes (C_2 \times C_2), 1],$$

$$H_5 = [(C_3 \times C_3 \times C_3) \rtimes C_2, 1],$$

$$H_6 = [C_3 \times C_3 \times C_3, 1],$$

$$H_7 = [S_3 \times S_3 \times S_3, 1].$$

Thus, $S_3 \wr S_3$ contains 8 proper normal subgroups (one of them, numbered by 2, determined by the same products but which has non isomorphic three different structures). Furthermore form of their generators we can deduce the splitability of this groups.

By analyzing their structures in terms of a semidirect product and determination them as the subgroup of wreath product with respect of their generators we obtain:

1) Consider the three isomorphic subgroups corresponding to G_7 having GAP structure description $H_2 = [“(C_3 \times C_3 \times C_3) : (C_2 \times C_2) : C_3 : C_2”, 3]$ here are 3 non equal isomorphic subgroups of order 648. Algebraic structure of them is the following:

$$H_2 \simeq (((C_3 \times C_3 \times C_3) \rtimes (C_2 \times C_2)) \rtimes C_3) \rtimes C_2,$$

these subgroups have order 648 and identifiers according to GAP system of this small group: id =703, 704 and 705 respectively.

At the same time, we compute $ord(S_3 \wr \tilde{A}_3) = ((6^3) : 2) \cdot 6 = 3^4 \cdot 2^3 = 108 \cdot 6 = 648$.

We will denote it by H_{703} , H_{704} and H_{705} correspondingly to their "id" in GAP. To classify these groups we compute its commutator subgroups and centers. The subgroup H'_{703} and H'_{704} have order 324 and GAP identifier is 160, this order is exactly half of 648. This means that commutator of these subgroups is the same therefore H_{703} and H_{704} **are isomorphic but they are embedded** in $S_3 \wr S_3$ in **different ways** as different copies. The center of this subgroups are trivial subgroup. The commutator subgroup of

$$S_3 \wr \tilde{A}_3 \simeq H_{703} \simeq H_{704}$$

is exactly $A_3 \wr \tilde{A}_3$, so active subgroup has order in 2 times less then order of S_3 . Thus, we can conclude that $H_{703} \simeq S_3 \wr \tilde{A}_3$ as well as $H_{704} \simeq S_3 \wr \tilde{A}_3$. The center of $S_3 \wr \tilde{A}_3$ is the trivial subgroup too that confirms isomorphism $H_{703} \simeq H_{704} \simeq S_3 \wr \tilde{A}_3$.

In H_{705} the homomorphism from top group of semidirect product in $\text{Aut} \tilde{A}_3$ has another structure of the kernel, $\text{ord}(H'_{705}) = 108$ which in 3 time less than order of H'_{703} .

Also we compute $\text{ord}(S_3 \wr \tilde{A}_3) = ((6^3) : 2) \cdot 6 = 3^4 \cdot 2^3 = 108 \cdot 6 = 648$.

On the other side the normal subgroup $A_3 \wr S_3$ has the commutator subgroup $e \wr \tilde{A}_3$ with the same order 108. Thus, the **third copy** of H_2 from the GAP list is

$$A_3 \wr S_3 = H_{705}.$$

But their commutators subgroup have different orders 324 (for id 703 and 704) and 108 for H_{705} , also commutator subgroups H'_{703} , H'_{704} have Id 160 in GAP and Id of H'_{705} is 40. Furthermore second commutators H''_{703} and H''_{704} has order 108 and Id=40, and order of exceptional subgroup H''_{705} is 27 and its Id =5.

2) $\langle (1, 2, 3), (1, 2), (4, 5, 6), (4, 5), (7, 8, 9), (7, 8), (1, 4, 7)(2, 5, 8)(3, 6, 9), (1, 4)(2, 5)(3, 6) \rangle = S_3 \wr S_3$.

3) $T_3 = (C_3 \times C_3 \times C_3) \rtimes C_2$ taking into account that $C_3 \simeq A_3$ we see that in general case when $G = S_n \wr S_n$ this subgroup has structure $\tilde{T}_n \simeq \underbrace{(A_n \times A_n \times \cdots \times A_n)}_n \rtimes C_2$.

4) $((C_3 \times C_3 \times C_3) \rtimes (C_2 \times C_2)) \rtimes C_3 = A_3 \wr \tilde{A}_3$.

Remark 17. If $G = S_n \wr S_m$ then the subgroup of type \tilde{T}_m has the structure $T_{n,m} \simeq \underbrace{(A_m \times A_m \times \cdots \times A_m)}_n \rtimes C_2$.

The parity of product of permutations tuple (g_1, g_2, \dots, g_n) , where $g_i \in A_m$, on $\delta \in C_2$ depends on homomorphic image of $\delta \in C_2$ in $\text{Aut}(\prod_{i=1}^n A_m^{(i)})$ as an element $((g_1, g_2, \dots, g_n), \delta)$ of the semidirect product $\prod_{i=1}^n A_m^{(i)} \rtimes C_2$.

We recall that a center of the group $(\mathcal{A}, X) \wr \mathcal{B}$ is direct product of normal closure of center of diagonal of \mathcal{B}^n i.e. $Z(\Delta(\mathcal{B}^n))$, trivial an element, and intersection of $(\mathcal{K}) \times E$ with $Z(\mathcal{A})$ [12], where \mathcal{K} is a kernel of action \mathcal{A} on X .

Note that derived by us subgroup \tilde{T}_m is **generalization of diagonal subgroup**.

But the following generalization of the diagonal of base of wreath product is a non-trivial normal subgroup. A base group of $N \triangleleft A \wr B$ consists of equivalence classes of an algebraic subgroup B , where conjugation by elements $A \wr B$ of acts imprimitive. These

generalization of diagonal subgroups $\Delta Z(S_m^n)$ [12] of base of wreath product are \widetilde{T}_m , $T_{n,m}$ and $E \wr \widetilde{A}_m$.

In this generalization a passive subgroup of $N \triangleleft A \wr B$ consists of equivalence classes of an algebraic subgroup B , where conjugation by elements $A \wr B$ acts imprimitive.

Remark 18. Taking into account formulated above properties of $S_n \wr S_m$ we conclude that $E \times \Delta Z(S_m^n)$ [12] is normal, where $Z(\Delta S_m^n)$ denotes the center of diagonal subgroup of $S_m^n = \prod_1^n S_m$. This statement can be generalized on $S_n \wr G$.

Proof. The intersection of kernel of action of active subgroup and its center is trivial because center of S_n is trivial and kernel of action (S_n, N_n) , where $N_n = \{1, \dots, n\}$ is trivial because of action is faithful. Also in this case of the center of diagonal of base of wreath product $S_n \wr S_m$ i.e $\Delta Z(S_m^n)$ [12] is trivial too. Therefore usual diagonal subgroup of base subgroup S_m^n from $S_n \wr S_m$ does not form normal subgroup. Thus, center of $S_n \wr S_m$ is trivial. \square

The cycle-type of a permutation of S_n is the partition of n obtained from the lengths of the cycles that appear in its decomposition into product of disjoint cycles.

In this paper we will denote the cycle-type of a permutation ω by $ct(\omega)$.

We recall that group is *monolithic* if the intersection of all nontrivial normal subgroups is nontrivial; the *monolith* is this intersection.

Theorem 19. Proper normal subgroups in $S_n \wr S_m$ (where action is left), where $n, m \geq 3$ with $n, m \neq 4$ are of the following types:

1. subgroups that act only on the second level (stabilizing the first level [27, 30]) are

$$E \wr \widetilde{A}_m, \widetilde{T}_m, E \wr S_m, E \wr A_n,$$

2. subgroups that act on both levels are $A_n \wr \widetilde{A}_m$, $S_n \wr \widetilde{A}_m$, $A_n \wr S_m$,

wherein the subgroup $S_n \wr \widetilde{A}_m \simeq S_n \triangleleft (\underbrace{S_m \boxtimes S_m \boxtimes S_m \dots \boxtimes S_m}_n)$ endowed with the subdirect product satisfying to condition (3), moreover $S_n \wr \widetilde{A}_m$ has two isomorphic copies, embedded into $S_n \wr S_m$ in different ways.

In total there are 8 proper normal subgroups in $S_n \wr S_m$.

Proof. We recall that in case of a trivial action G on X a wreath product $(G, X) \wr H$ degrades in $G \times H^X$. If $G = E$ we have trivial action on X , therefore the subgroup $E \times S_m^n$ appears.

First of all, note that all our normal subgroups contain subgroup $e \wr A_m$. If we prove the normality of $e \wr \widetilde{A}_m$ then last two facts would imply that $S_n \wr \widetilde{S}_m$ is *monolithic* group and its *monolith* is $e \wr A_m$. To prove the normality of $e \wr A_m$ as well as $A_n \wr S_m$ and $S_n \wr \widetilde{A}_m$ it is sufficient to validate the embedednes of the normal closure $S_n \wr S_m$ in of its generators to the correspondent initial subgroup $e \wr \widetilde{A}_m$ as well as $A_n \wr S_m$ and $S_n \wr \widetilde{A}_m$ for $n, m \geq 3, n, m \neq 4$.

Generators of each of subgroups will be written in a similar way with the generators of normal subgroups $S_3 \wr S_3$. The proof of split ups of any these subgroups will follow from the form of their generators elements.

1. a) $e \wr A_m$: Its generating elements have the following form: $a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$, where $i_1, i_2, i_3 \in \{1, 2, \dots, m\}$, as well as $\tilde{a} = [e]_1, [(12), (12), e, e \dots, e]_2$. The normal closure of \tilde{a} is found in 6. Its coincide with $e \wr A_m$. Moreover, A_n can be generated by all 3-cycles, and all 3-cycles can be generated by conjugating $(i_1 i_2 i_3)$ by even substitutions, such substitutions are in S_m . Proposition 8 and Corollary 11 show us that the conjugated generator $a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$ by elements of $S_n \wr S_m$ pertains to $e \wr A_m$ too.

Normality of the subgroup. Consider an arbitrary generator a , without losing the generality a can be in of the form: $a = a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$, $i_1, i_2, i_3 \in \{1, 2, \dots, m\} \in A_m$. Then for any element $b \in S_n \wr S_m$, that has tableau presentation in the form: $[\pi_0]_1, [\pi_1, \dots, \pi_n]_2, \pi_0 \in S_n, \pi_1, \dots, \pi_n \in S_m$ we have: $[b \cdot a \cdot b^{-1}]_1 = [\pi_0 \cdot e \cdot \pi_0^{-1}] = [e]_1$; $[b \cdot a \cdot b^{-1}]_2 = [b(x) \cdot a(x^{\pi_0}) \cdot b^{-1}(x^{\pi_0 \cdot \pi_0^{-1}})]_2 =$
 $= [\pi_1 \cdot e \cdot \pi_1^{-1}, \dots, \pi(x^{\pi_0}) \cdot (i_1 i_2 i_3) \cdot \pi(x^{\pi_0})^{-1}, \dots, \pi_n \cdot e \cdot \pi_n^{-1}]_2 =$
 $= [e, \dots, e, \pi(x^{\pi_0}) \cdot (i_1 i_2 i_3) \cdot \pi(x^{\pi_0})^{-1}, e, \dots, e]_2$.

Obviously, the mentioned above element pertains to $e \wr A_m$ because of $\pi(x^{\pi_0}) \cdot (i_1 i_2 i_3) \cdot \pi(x^{\pi_0})^{-1}$ is an even substitution.

b) *Generators of $e \wr \widetilde{A}_m$* are the following: $a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$, where $i_1, i_2, i_3 \in \{1, 2, \dots, m\}$ and $a_{(i_1 i_2), e, \dots, e, (i_1 i_2), e, \dots, e}$, $i_1, i_2 \in \{1, 2, \dots, m\}$. The normal closure of these generators is found in Proposition 6. This closure $N_{S_n \wr S_n}(\tilde{a})$ equals to $e \wr \widetilde{A}_m$.

d) Consider $E \wr S_m$. Recall that two element generating set of S_m has form $\tau = (1, 2)$, $\lambda = (1, 2, \dots, m)$. Without losing the generality we can consider generators for S_3 . Taking into consideration the mentioned above generating set the generators of $e \wr S_m$ have the form: $a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$, where $i_1, i_2, i_3 \in \{1, 2, \dots, m\}$ and $a_{e, \dots, e, (i_1 i_2), e, \dots, e}$, $i_1, i_2, i_3 \in \{1, 2, \dots, m\}$.

Let us verify the *normality of this subgroup*. It is obviously, because on the first levels in result of the conjugation of generators by permutation of S_n always be identical substitution, and on the second level be any substitution from S_m which was required to

prove in the theorem.

2. Here we prove that the groups mentioned in item 2 are normal with using the fact of normality in S_m of subgroups appearing on second level. Such groups are normal in S_m .

a) $A_n \wr \widetilde{A}_m$: Generators are the same as for $e \wr \widetilde{A}_m$ that are $a_{e, \dots, e, (i_1, i_2, i_3), e, \dots, e}$, and one new generator of wreath product $A_n \wr \widetilde{A}_m$ with the following tableau presentation:

$$[(i_1, i_2, i_3)]_1, [e, \dots, e]_2, i_1, i_2, i_3 \in \{1, 2, \dots, n\}. \quad (5)$$

Taking into account that A_n is generated by all 3-cycles and the fact that all 3-cycles can be generated by conjugating of (i_1, i_2, i_3) by even substitutions π we use for conjugation elements of form $[\pi]_1, [e, \dots, e]_2, i_1, i_2, i_3 \in \{1, 2, \dots, n\}$. As a result we see that normal closure of the new generator (5) is $A_n \wr E$. And normal closure of all generators is $A_n \wr \widetilde{A}_m$.

b) *Generating elements of $S_n \wr \widetilde{A}_m$* have the same presentation as for $A_n \wr \widetilde{A}_m$ except one additional generator presented by the next tableau:

$$[(i_1 i_2)]_1, [e, \dots, e]_2, i_1, i_2, \in \{1, 2, \dots, n\}.$$

The last generator completes generation set of active group to generating set of S_n , its normal closure pertain to $S_n \wr E$, the normal closure of $(i_1 i_2)$ is considered above in item c) for generators of \widetilde{T}_m . Therefore this completed generating set generates $S_n \wr \widetilde{A}_m$.

c) *The generating set of $A_n \wr S_m$* are the same as for described above $e \wr S_m$ except one additional generator in the normal closure which has tableau presentation:

$$[(i_1 i_2 i_3)]_1, [e, \dots, e]_2, i_1, i_2, i_3 \in \{1, 2, \dots, n\}.$$

As it is mentioned above A_n is generated by all 3-cycles and the fact that all 3-cycles can be generated by conjugating of $(i_1 i_2 i_3)$ by even permutations. Therefore normal closure of additional generator $[(i_1 i_2 i_3)]_1, [e, \dots, e]_2, i_1, i_2, i_3 \in \{1, 2, \dots, n\}$. \square

Proposition 20. *The normal closure of \widetilde{T}_m in $S_n \wr S_m$ is $\widetilde{T}_m^{S_n \wr S_m} = \widetilde{T}_m$.*

Proof. Consider the generators of \widetilde{T}_m . Since for $n \geq 3$, the group A_n is generated by 3-cycles of the form $(12i)$ then elements of first type can be generated by following elements: $a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$, where $i_1, i_2, i_3 \in \{1, 2, \dots, m\}$. Generators for the elements of second type: $a_{(i_1 i_2), \dots, (i_3 i_4), \dots, (i_5 i_6), i_1, \dots, i_6 \in \{1, 2, \dots, m\}}$ [14].

Normal closure of the first type generators in $S_n \wr S_m$ is $e \wr \widetilde{A}_m$, that is studied above in the item a) of Theorem 19.

It remains to prove the same for the generators of the second type. For this goal we consider the generators of the form: $a_{(i_1 i_2), \dots, (i_1 i_2), \dots, (i_1 i_2)} \in \widetilde{T}_m$. Then for any element $b \in S_n \wr S_m$, having tableau presentation $[\pi_0]_1, [\pi_1, \dots, \pi_n]_2, \pi_0 \in S_n, \pi_1, \dots, \pi_n \in S_m$ we have:

$$\begin{aligned} [b \cdot a \cdot b^{-1}]_1 &= [\pi_0 \cdot e \cdot \pi_0^{-1}] = [e]_1; \\ [b \cdot a \cdot b^{-1}]_2 &= [b(x) \cdot a(x^{\pi_0}) \cdot b^{-1}(x^{\pi_0 \cdot \pi_0^{-1}})]_2 = \\ &= [\pi_1 \cdot (i_1 i_2) \cdot \pi_1^{-1}, \dots, \pi_n \cdot (i_1 i_2) \cdot \pi_n^{-1}]_2 = \\ &= [\pi_k(x^{\pi_0}) \cdot (i_1 i_2) \cdot \pi_k(x^{\pi_0})^{-1}]_2. \end{aligned} \tag{6}$$

It is obvious that the obtained element $[\pi_k(x^{\pi_0}) \cdot (i_1 i_2) \cdot \pi_k(x^{\pi_0})^{-1}]_2$ is an odd permutation, therefore $b \cdot a \cdot b^{-1}$ belongs to the second type. This accomplished the prove for whole \widetilde{T}_m . \square

Corollary 21. *The subgroup \widetilde{T}_m is normal in $S_n \wr S_m$.*

Proof. In view of Proposition 20 we can conclude that parity of decrement of each coordinate in \widetilde{T}_m is invariant under conjugation in $S_n \wr S_m$ so \widetilde{T}_m is normal. \square

The example of $S_5 \wr S_5$ confirms last statement because of it gives us 8 proper normal subgroups too.

The lattice of invariant subgroups for $S_n \wr S_n$ is presented on Fig. 1.

Remark 22. *We have to note that $A_n \wr A_m$ is not normal as in $S_n \wr S_m$ as well as in $S_n \wr A_m$ because of wreath product having specific conjugation as a result of action.*

Theorem 23. *Proper normal subgroups in $A_n \wr S_m$, $n, m \geq 3$, $n, m \neq 4$ are the following 2 types:*

1. *the subgroups the of first level stabilizer ($St_{S_n \wr S_m}(1)$), which are the following: $E \wr \widetilde{A}_m, \widetilde{T}_m, E \wr S_m$;*

2. *subgroups, acting on both levels: $A_n \wr \widetilde{A}_m$*

Thus we have 5 in $A_n \wr S_m$ and 8 proper normal subgroups in $S_n \wr S_m$.

Proof. The proofs for all subgroups except one is similar to proofs in Theorem 19. The proof of normality for $A_n \wr \widetilde{A}_m$ is such:

$A_n \wr \widetilde{A}_m$: Its generators are the same as for $e \wr \widetilde{A}_m$ that are $a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$, and one new generator of wreath product $A_n \wr \widetilde{A}_m$ with the following tableau presentation:

$$[(i_1 i_2 i_3)]_1, [e, \dots, e]_2, i_1, i_2, i_3 \in \{1, 2, \dots, n\}.$$




Figure 1: Lattice of invariant subgroups $S_n \wr S_n$ for the case $n \equiv 1 \pmod{2}$

Taking in account that A_n is generated by all 3-cycles and the fact that all 3-cycles can be generated by conjugating of $(i_1 i_2 i_3)$ by even substitutions π we use for conjugation elements of form $[\pi]_1, [e, \dots, e]_2, \pi \in \{1, 2, \dots, n\}$. As a result we see that normal closure of the last kind of generators is $A_n \wr E$. And normal closure of all generators is $A_n \wr \tilde{A}_n$. \square

Remark 24. Thus we have 8 proper normal subgroups in $S_n \wr S_m$ and 5 in $A_n \wr S_m$. The list of new proper subgroups in $E \wr S_m, S_n \wr \tilde{A}_m, A_n \wr S_m$. Thus, we have 3 new proper normal subgroups.

Theorem 25. There are exactly 5 proper normal subgroups in the wreath product $W = A_n \wr S_n$. The list of these subgroups is:

1. $E \wr A_n$, this is the subgroup of the stabilizer of the first level $St_W(1)$ [27, 30],
2. subgroup \tilde{T}_n , that is the subgroup of $St_W(1)$,




Figure 2: Lattice of invariant subgroups $S_n \wr S_n$ for the case $n \equiv 0 \pmod{2}$

3. $St_{W'}(1)$, this is a subgroup of the first type in terms of Theorem 23 having the structure $E \wr \tilde{A}_n$,
4. W' , this is a subgroup of the second type and W' has the structure $A_n \wr \tilde{A}_n$,
5. $St_W(1)$, this is a proper normal subgroup of W . In this case as well as in case $W = S_n \wr S_n$ the normal subgroup $St_W(1)$ has structure the same as $E \wr S_n$.

Proof. The proofs for all subgroups except one is similar to proofs in Theorem 19. The proof of normality for $A_n \wr \tilde{A}_m$ is such:

$A_n \wr \tilde{A}_m$: Its generators are the same as for $e \wr \tilde{A}_m$ that are $a_{e, \dots, e, (i_1 i_2 i_3), e, \dots, e}$, and one new generator of wreath product $A_n \wr \tilde{A}_m$ with the following tableau presentation:

$$[(i_1 i_2 i_3)]_1, [e, \dots, e]_2, i_1, i_2, i_3 \in \{1, 2, \dots, n\}.$$

Taking in account that A_n is generated by all 3-cycles and the fact that all 3-cycles can be generated by conjugating of $(i_1 i_2 i_3)$ by even substitutions π we use for conjugation elements of form $[\pi]_1, [e, \dots, e]_2, \pi \in \{1, 2, \dots, n\}$. As a result we see that normal closure of the last kind of generators is $A_n \wr E$. And normal closure of all generators is $A_n \wr \widetilde{A}_m$. \square

Remark 26. Note that subgroup $S_n \wr \widetilde{A}_m$ and $A_n \wr S_m$ is the new normal proper subgroup of $S_n \wr S_m$ relatively to the normal subgroups of $A_n \wr S_m$ which has 5 proper subgroups.

Remark 27. A normalizer of $A_n \wr A_m$ in $S_n \wr S_m$ is the following:

$$N_{S_n \wr S_m}(A_n \wr A_m) = S_n \wr A_m.$$

Also we have to remark that $A_n \wr A_m \triangleleft S_n \wr A_m$.

Let K_4 be Klein subgroup.

The group $W_4 = S_4 \wr S_4$ has the same 11 normal subgroups as $S_n \wr S_n$, for $n = 3$ or $n > 4$, and **3 additional** subgroups: $E \wr K_4, K_4 \wr \widetilde{A}_4, K_4 \wr S_4$ provided by the condition $n = 4$.

Lemma 28. All normal subgroups N_i of $S_n \wr S_m$ are presented in one of the following form:

1. $E \wr N_i$, where $N_i \leq \prod_{i=1}^n S_m^{(i)}$ i.e. N_i is a normal subgroup in the base of $S_n \wr S_m$ and invariant relatively to conjugation in wreath product [14, 27, 29].
2. $A_j \wr N_i$, where N_i is form 1) and $A_i \leq S_n$ subgroups which possesses the quotient subgroup of the form $A_j \wr E$ such that $E \wr N_i$ can be extended by A_j to a subgroup in $S_n \wr S_m$, i.e. for which extension problem for $E \wr N_i$ by A_j to $A_j \wr N_i$ is solvable.

Remark 29. Subgroups from 1) can be partitioned into 2 classes:

- a) subgroups which possesses the normal subgroup of form $E \wr N_i$ which can be extended by A_j to $S_n \wr S_m$. I.e. for which extension problem for A_j by $E \wr N_i$ to $A_j \wr N_i$ is solvable.
- b) Subgroups from 1) which has not such extension.

Let the set of normal subgroup of $S_n \wr S_n$ be denoted by $N(S_n \wr S_n)$, wherein a subgroup with number i from $N(S_n \wr S_n)$ be denoted by $N_i(S_n \wr S_n)$.

To obtain embedding theorem of $N_i(S_n \wr S_n)$ in $S_n \wr S_n \wr S_n$ we rely on Theorem 1 and Lemma 1 taking into account that $N_i(S_n \wr S_n)'$ is subgroup of $St_{S_n \wr S_n \wr S_n}(1)$ described in Theorem 19.

The set of normal subgroup of $S_n \wr S_n$ is denoted by $N(S_n \wr S_n)$. Subgroup with number i from $N(S_n \wr S_n)$ is denoted by $N_i(S_n \wr S_n)$.

Theorem 30. *The full list of normal subgroups of $S_n \wr S_n \wr S_n$ consists of 50 normal subgroups. These subgroups are the following:*

- 1 **Type** T_{023} contains: $E \wr \tilde{A}_n \wr H$, $\widetilde{T}_n \wr H$, where $H \in \{\tilde{A}_n, \tilde{A}_{n^2}, S_n\}$. There are 6 subgroups.
- 2 **The second type of subgroups is subclass in T_{023} with new base of wreath product subgroup \tilde{A}_{n^2} :** $E \wr S_n \wr \tilde{A}_{n^2}$, $E \wr N_i(S_n \wr S_n)$. Therefore this class has 12 new subgroups. Thus, the total number of normal subgroups in **Type** T_{023} is 18.
- 3 **Type** T_{003} : $A_{00(n^2)}^{(3)} = E \wr E \wr \tilde{A}_{n^2}, \widetilde{T}_{n^2}, \widetilde{T}_n^{(3)}$. Hence, here are 3 new subgroups.
- 4 **Type** T_{123} : $N_i(S_n \wr S_n) \wr S_n$, $N_i(S_n \wr S_n) \wr \tilde{A}_n$ and $N_i(S_n \wr S_n) \wr \tilde{A}_{n^2}$. Thus, there are 29 new normal subgroups in T_{123} , taking into account repetition.

To obtain the embedding theorem of $N_i(S_n \wr S_n)$ in $S_n \wr S_n \wr S_n$ we rely on Theorem 19 and Lemma 25.

Definition 16. The set of elements from $\wr_{i=1}^k S_{n_i}, n \geq 3$ with depth j satisfying the following condition

$$\sum_{i=1}^{n^j} \text{dec}([a_i]_k) = 2t, t \in \mathbb{N}, [a_i]_j = e, \text{ whenever } j = \overline{0, k-1} \quad (7)$$

is called $\tilde{A}_{n^k}^{(k)}$.

Definition 17. The set of elements from $\wr_{i=1}^k S_{n_i}, n \geq 3$ with depth j having tableaux presentation which satisfy the following conditions

$$\sum_{i=1}^{n^j} \text{dec}([a_i]_j) = 2z, z \in \mathbb{N}, j = \overline{m, k}; [a_i]_j = e, \text{ whenever } j = \overline{0, m-1} \quad (8)$$

is called $\tilde{A}_{n^k}^{(m,k)}$.

Definition 18. The set of elements from $\wr_{i=1}^{\infty} S_{n_i}, n \geq 3$ with depth j satisfying for each $j \geq m$ the following condition

$$\sum_{i=1}^{n^j} \text{dec}([a_i]_j) = 2t, t \in \mathbb{N}, j \geq k, [a_i]_j = e, \text{ whenever } j = \overline{0, k-1} \quad (9)$$

is called $\widetilde{A}_{n^j}^{\geq k}$.

As in our previous work [27] we denote by $G_k(l)$ a level subgroup of G_k , which consists of the tuples of v.p. from X^l , $l < k - 1$ of any $\alpha \in G_k$. Note that $G_k(l)$ is in bijective correspondence (and isomorphism) with l -coordinate subgroup $[U]_l$ [10].

Statement 31. In $\widetilde{A}_{n^m}^{\geq l}$ the following $m - l$ equalities are true:

$$\sum_{i=l}^m \text{dec}(g_{ij}) = 2k, k \in \mathbb{N}, 0 < j \leq n^i, \quad (10)$$

Remind that the condition on commutator subgroup of permutation wreath product base was obtained by us in [12], and commutator subgroup is also normal. The condition (4) accords with the condition setting the subdirect product in a commutator subgroup of wreath product base [12].

We call *permutation of l -th level* $\pi(X^l)$ a permutation of automorphism $g \in \text{Aut}X$ formed by action of all vertex permutations of l -th level on vertexes of $l + 1$ level. Such permutations are elements of level subgroup defined in our paper [27].

Proposition 32. *The set of elements of type $\widetilde{A}_{n^2}^{(3)}$ forms the subgroup in $S_n \wr S_n \wr S_n$, furthermore $\widetilde{A}_{n^2}^{(3)} \triangleleft S_n \wr S_n \wr S_n$.*

Proof. For convenience we will identify $\widetilde{A}_{n^2}^{(3)}$ with its isomorphic image in $\text{Aut}(X^{[k]})$ and automorphisms from $\text{Aut}(X^{[k]})$ identify with elements of $\widetilde{A}_{n^2}^{(3)}$.

The subgroup of 3-rd level of $\widetilde{A}_{n^2}^{(3)}$ has even product of vertex permutation in view of condition (4).

Consider an arbitrary automorphisms $a \in \text{Aut}(X^{[3]})$ and $b \in \widetilde{A}_{n^2}^{(3)}$. We specify our attention on $a^{-1}ba$.

Let H be a group acting (from the left) by a permutation π upon a set X and the composition will occur from the right to left, and let G be an arbitrary group. Then the (permutational) wreath product $H \wr G$ is the semi-direct product $H \ltimes G^X$, where H acts on the direct power G^X by the respective permutations of the direct factors.

$$ba = \sigma(b_1, \dots, b_d)\pi(a_1, \dots, a_d) = \sigma\pi(b_{\pi(1)}a_{(1)}, \dots, b_{\pi(d)}a_{(d)})$$

An inverse an element has form $a^{-1} = \pi^{-1}(a_{\pi^{-1}(1)}^{-1}, \dots, a_{\pi^{-1}(d)}^{-1})$. Let $b = (b_1, \dots, b_n)\sigma \in \widetilde{A}_{n^2}^{(3)}$ so we need to verify invariance of property 4 under conjugation. Consider a conjugation in $\text{Aut}X^{\{3\}} \simeq S_n \wr S_n \wr S_n$

$$\begin{aligned} \pi^{-1} \left(a_{\pi^{-1}(1)}^{-1}, \dots, a_{\pi^{-1}(d)}^{-1} \right) \sigma(b_1, \dots, b_d) \pi(a_1, \dots, a_d) = \\ \pi^{-1} \sigma \pi \left(a_{\pi^{-1}\sigma\pi(1)}^{-1} b_{\pi(1)} a_{(1)}, \dots, a_{\pi^{-1}\sigma\pi(d)}^{-1} b_{\pi(d)} a_{(d)} \right), \end{aligned} \quad (11)$$

where right-to-left composition of permutations is used. Taking into account the invariance of decrements of $a \in \text{Aut}(X^{[3]})$ with respect to taking the inverse element and the fact that acting of π and σ only rearranges states of automorphisms a^{-1} in right part of (11) over the vertices of the X^2 in tuple of elements $(a_{\pi^{-1}\sigma\pi(1)}^{-1} b_{\pi(1)} a_{(1)}, \dots, a_{\pi^{-1}\sigma\pi(d)}^{-1} b_{\pi(d)} a_{(d)})$ respectively to the initial element $\pi^{-1}(a_{\pi^{-1}(1)}^{-1}, \dots, a_{\pi^{-1}(n)}^{-1})$ we conclude that $\sum_{j=1}^n \text{dec}(a_{\pi^{-1}\sigma\pi(j)}^{-1}) = \sum_{j=1}^n \text{dec}(a_j)$. By the same reasoning we obtain $\sum_{j=1}^n \text{dec}(b_{\pi\sigma\pi(j)}^{-1}) = \sum_{j=1}^n \text{dec}(b_j)$.

Thus, conjugation $(b_1, \dots, b_n)\sigma$ by $(a_1, \dots, a_n)\pi$ [7, 8, 27] do not change parity of its sum of vertex permutations b_{2i} in $v_{2i} \in X^2$, $1 \leq i \leq n^3$ decrements. Therefore the characteristic element property (4) of $\tilde{A}_{n^2}^{(3)}$ holds after conjugation too.

The generalization of proof for such extension of this group as $\tilde{A}_{n^2}^{\geq l}$ is the following.

Consider an arbitrary automorphisms $h \in \text{Aut}(X^{[k]})$ and $g \in \tilde{A}_{n^2}^{(k)}$. We specify our attention on $h^{-1}gh$.

Let $\pi(X^k)$ be *level permutation* of the l -th level of an conjugating automorphism $h \in \text{Aut}(X^{[k]})$ and $\tau \in A_{n^j}^{(3)}$. As it was mentioned and proved above a $\text{dec}(\pi) = \text{dec}(\pi^{-1})$. Therefore permutation of the l -th level of automorphism $\pi^{-1}\sigma\pi$ is even again. Thus, $\tilde{A}_{n^j}^{(k)} \triangleleft \text{Aut}(X^{[k]}) \simeq \wr_{i=1}^k S_n$.

Corollary 33. *The set of elements $\tilde{A}_{n^2}^{\geq 3}$ form a normal subgroup in $\wr_{i=1}^m S_n$.*

The permutations of l -th level subgroup of $\tilde{A}_{n^2}^{\geq 3}$ possess even product of vertex permutation in view of definition 18 Let π be level permutation of i -th level, $j \geq l$, of an conjugating automorphism $h \in S_n \wr S_n \wr S_n$ and $g \in \tilde{A}_{n^2}^{\geq 3}$. As it was mentioned and proved above $\text{dec}(\pi) = \text{dec}(\pi^{-1})$, so they have the same parity. Therefore permutation of j -th level of automorphism $\pi^{-1}\tau\pi$ is even again.

Let us verify that the similar condition to (11) holds for rest of levels after j of $\tilde{A}_{n^2}^{\geq 3}$.

Let $H = \tilde{A}_{n^{i-1}}^{(l)}$ be a group acting (from the left) by a permutation π upon a set X , wherein $|X| = n$, and the composition will occur from the right to left, and let G be an wreath product of symmetric groups $\wr_{i=l}^{\infty} \tilde{S}_n$ satisfying the condition (10). Then the (permutational) wreath product $H \wr G$ is the semi-direct product $H \ltimes G^X$, where H acts on the direct power G^X by the respective permutations of the direct factors.

$$ba = \sigma(b_1, \dots, b_d)\pi(a_1, \dots, a_d) = \sigma\pi(b_{\pi(1)}a_{(1)}, \dots, b_{\pi(d)}a_{(d)})$$

An inverse element has form $a^{-1} = \pi^{-1}(a_{\pi^{-1}(1)}^{-1}, \dots, a_{\pi^{-1}(d)}^{-1})$. Let $b = (b_1, \dots, b_n)\sigma \in \tilde{A}_{n^2}^{(3)}$ so we need to verify invariance of property 4 under conjugation. Consider a conjugation in $AutX^{\{3\}} \simeq S_n \wr S_n \wr S_n$

In more detail result of conjugation of $l+1$ level states b_j of automorphism $b \in AutX^{[k]}$ which belongs to subgroup $Aut(v_{lj}X^{k-l})$ of automorphisms $b \in \tilde{A}_{n^2}^{\geq 3}$ has form

$$\begin{aligned} \pi^{-1} \left(a_{\pi^{-1}(1)}^{-1}, \dots, a_{\pi^{-1}(n)}^{-1} \right) \sigma(b_1, \dots, b_n) \pi(a_1, \dots, a_n) = \\ \pi^{-1} \sigma \pi \left(a_{\pi^{-1}\sigma\pi(1)}^{-1} b_{\pi(1)} a_{(1)}, \dots, a_{\pi^{-1}\sigma\pi(n)}^{-1} b_{\pi(n)} a_{(n)} \right), \end{aligned} \quad (12)$$

where right-to-left composition of permutations is used. As can be seen from the formula, the conjugating elements $h_{(i)}$ are only rearranged by action of σ , but in the aggregate the sum of their decrements remained unchanged. And the sum of decrements of conjugating elements $\sum_{i=l}^m dec(h_{\sigma(i)}^{-1}) = \sum_{i=l}^m dec(h_{(i)}), i \in \mathbb{N}, 1 < i \leq n^3$, remained invariant, because of $dec(h_{(i)}) = dec(h_{(i)}^{-1})$.

Taking into account the invariance of decrements of $a \in Aut(X^{[k]})$ with respect to taking the inverse element and the fact that acting of π and σ only rearranges states of automorphisms a^{-1} in right part of (12) over the vertices of the $v_{lj}X^{k-m}$ in the tuple of elements $(a_{\pi^{-1}\sigma\pi(1)}^{-1} b_{\pi(1)} a_{(1)}, \dots, a_{\pi^{-1}\sigma\pi(d)}^{-1} b_{\pi(d)} a_{(d)})$ respectively to the initial element $(a_{\pi^{-1}(1)}^{-1}, \dots, a_{\pi^{-1}(n)}^{-1})$ we conclude that $\sum_{j=1}^n dec(a_{\pi^{-1}\sigma\pi(j)}^{-1}) = \sum_{j=1}^n dec(a_j)$. By the same reasoning we obtain $\sum_{j=1}^n dec(b_{\pi\sigma\pi(j)}^{-1}) = \sum_{j=1}^n dec(b_j)$.

Thus, conjugation (b_1, \dots, b_n) by $(a_1, \dots, a_n)\pi$ [7, 8, 27] do not change parity of its decrements sum of vertex permutations $b_{l+1,i}$ in $v_{l,i}X^{l+1}$ for each $1 \leq i \leq n^l$. Therefore the characteristic element property (4) of $\tilde{A}_{n^2}^{(3)}$ holds after conjugation too.

Consequently a decrements sum of all vertexes permutation from X^{l+1} has the same parity $\sum_{j=1}^{n^{l+1}} dec(a_{\pi^{-1}\sigma\pi(j)}^{-1} b_{\pi(i)} a_{(i)}) \equiv \sum_{j=1}^{n^{l+1}} dec(b_j) \pmod{2}$.

$g = (g_{(1)}, g_{(2)}, \dots, g_{(d)})\pi_g$, $h = (h_{(1)}, h_{(2)}, \dots, h_{(d)})\sigma_h$, is given by the formula:

$$g \cdot h = (g_{(1)}h_{(\pi_g(1))}, g_{(2)}h_{(\pi_g(2))}, \dots, g_{(d)}h_{(\pi_g(d))})\pi_g\sigma_h.$$

We define σ as permutation in $v_{li} \sigma \in A_n$, where, in general case, n is degree of l -th

level of spherically homogeneous tree.

$$g = (g_{(1)}, g_{(2)}, \dots, g_{(d)})\pi_g, \quad h = (h_{(1)}, h_{(2)}, \dots, h_{(d)})\sigma_h, \quad \text{is given by the formula: } \quad \square$$

Theorem 34. *Let $H \triangleleft \text{Aut}_f X^*$ with depth k then H contains k -th level subgroup P having all even vertex permutations $p_{ki} \in A_n$ on X^k and trivial permutations in vertices of rest of levels. Furthermore P is normal in W provided k is last active level of $\text{Aut}_f X^*$.*

Let H_i be a subgroup of $St_W(k)$, then there is k -th level subgroup $H_i(k)$ of H_i isomorphic to $\tilde{A}_{n^{k-i}}^{(k)}$, $0 \leq i < k$.

In particular for the case $i = k - 1$, $H_1 \simeq \tilde{A}_n^{(k)}$ and determined by the following congruences

$$\sum_{i=1}^n \text{dec}(\pi_{ki}) \equiv \sum_{i=n+1}^{2n} \text{dec}(\pi_{ki}) \equiv \dots \equiv \sum_{i=ln+1}^{(l+1)n} \text{dec}(\pi_{ki}) \equiv \dots \equiv \sum_{n^{k-n+1}}^{n^k} (\pi_{ki}) \equiv 0 \pmod{2}.$$

Note that there are n^{k-1} such sums.

The subgroup P is partial case where $i = 0$ and H_i degenerates in extrimal subgroup $H_0 \simeq \tilde{A}_{n^0}^{(k)}$. The congruences takes form: $\text{dec}(\pi_{ki}) \equiv 0 \pmod{2}$, $1 \leq i \leq n^k$.

If $i = k - 2$, then subgroup H_{k-2} is isomorphic to $\tilde{A}_{n^2}^{(k)}$ which is determined by characteristic equations:

$$\sum_{i=1}^{n^2} \text{dec}(\pi_{ki}) \equiv \sum_{i=n^2+1}^{2n^2} \text{dec}(\pi_{ki}) \equiv \dots \equiv \sum_{i=ln^2+1}^{(l+1)n^2} \text{dec}(\pi_{ki}) \equiv \dots \equiv \sum_{n^{k-n^2+1}}^{n^k} (\pi_{ki}) \equiv 0 \pmod{2}.$$

there are n^{k-2} such sums. In general for $W^{(i)}$ there are n^i such sums.

In the case H_i this subgroup takes form $\tilde{A}_{n^{k-i}}^{(k)}$ which is determined by the following n^i characteristic equations:

$$\sum_{i=1}^{n^{k-i}} \text{dec}(\pi_{ki}) \equiv \sum_{i=n^{k-i}+1}^{2n^{k-i}} \text{dec}(\pi_{ki}) \equiv \dots \equiv \sum_{i=(l)n^i+1}^{i=(l+1)n^i} \text{dec}(\pi_{ki}) \equiv \dots \equiv \sum_{n^{k-n^i+1}}^{n^k} (\pi_{ki}) \equiv 0 \pmod{2}.$$

In total there are n^{i+1} sums of decrements and so on.

In the case $i = 1$ this subgroup takes form $\tilde{A}_{n^{k-1}}^{(k)}$ which is determined by characteristic equations:

$$\sum_{i=1}^{n^{k-1}} \text{dec}(\pi_{ki}) \equiv \sum_{i=n^{k-1}+1}^{2n^{k-1}} \text{dec}(\pi_{ki}) \equiv 0 \pmod{2}$$

Process of jointing of sums is continuing and ends on the condition $i = 0$ with the sum

$$\sum_{i=1}^{n^k} dec(\pi_{ki}) \equiv 0 \pmod{2}$$

which is characterize the level subgroup of normal subgroup $\tilde{A}_{n^k}^{(k)}$.

Proof. To construct the described in conditions of this Theorem subgroup P , we set the following restrictions on root permutations of automorphisms x, y, c having form $x = \sigma_x(h_1, \dots, h_n)$, $y = \sigma_y(g_1, \dots, g_n)$ and introduce auxiliary notation Let $\sigma_\pi \in S_n$, then $Supp_{S_n}(\sigma_\pi)$ be a subset of $N_n = \{1, 2, \dots, n\}$ such that $\sigma_\pi(k) \neq k$, $\sigma_\pi \in S_n$, $k \in N_n$ also $Fix_{S_n}(\sigma_\pi) = N_n \setminus Supp_{S_n}(\sigma_\pi)$. We choose σ_x, σ_y and σ_c with such properties: 1) $Supp_{A_n}(\sigma_y) \subseteq Supp_{A_n}(\sigma_x)$. 2) $Supp_{A_n}(\sigma_c) \subseteq Fix_{A_n}(\sigma_y)$, wherein $|Fix(\sigma_y)| > 1$, hence $Supp_{A_n}(\sigma_y) \cap Supp_{A_n}(\sigma_c) = \emptyset$. 3) We also demand $Supp_{A_n}(\sigma_x) \cap Supp_{A_n}(\sigma_c) \neq \emptyset$. As a direct conclusion from 1) and 3) we have that $Supp[\sigma_y, \sigma_x] \cap Supp[\sigma_c] \neq \emptyset$. Note that 2) yields commutating of σ_y and σ_c . Let σ_x and σ_y are the permutations in v_0 (or in general case it is arbitrary vertex $v_{ki} \in X^*$). As we researched with Y.A. Drozd, conjugation of automorphism x of rooted homogeneous tree X^* by automorphism $y \in AutX^*$ permutes only vertices on which y acts non-trivially and fixes only those vertices from X^1 where action of σ_y from $y = \sigma_y(g_1, \dots, g_n)$ is trivial. Therefore, some of the vertices of the first level will remain rigid (invariant) if $Fix(\sigma_y) \neq \emptyset$, furthermore in this part of vertices we reduce the permutations using multiplication in a form similar to the commutator: $(\sigma_y \sigma_x \sigma_y^{-1}) \cdot \sigma_x^{-1}$. Thus we will simplify those permutations located in the vertices of X^1 indexed by numbers belonging to $Fix(\sigma_y)$. Then in order to make permutations in the remaining vertices (namely in $X^1 \setminus v_{1i} : i \in Fix(\sigma_y)$) trivial, we multiply again on automorphism $c[y, x]^{-1} c^{-1}$ conjugated to inverse to obtained above automorphism $(yxy^{-1}) \cdot x^{-1} = [y, x]$. Note, that we conjugate it with such an element c that acts trivially precisely on those vertices of X^1 , the permutations in which have to be simplified. Thus we express $[y, x] c[y, x]^{-1} c^{-1}$. In this automorphism all vertices from X^1 already contain trivial permutations as a result of the provided above reduction.

To meet the condition $Supp[\sigma_y, \sigma_x] \cap Supp[\sigma_c] \neq \emptyset$ it is sufficiently to present the common elements in the non-invariant elements of permutation $[\sigma_y, \sigma_x]$ and $Supp[\sigma_c]$. For this purpose we choose I such that $y(i) = j$, $i \notin Fix(y)$, provided the condition $x(j) = l \in Fix(y)$ that is possible due to 1), and $x^{-1}(l) = k \in Supp(c)$ that is possible because of 3). Since $l \in Fix(y)$ then $y^{-1}(l) = l$. Now we can spread our method of constructing of even permutation in the root of $T(X^*)$ on vertices of X^k . Let σ_x and σ_y are the permutations in v_{ki} from automorphisms x and y correspondently. We can exactly

implement the described above method to automorphisms: $y = \sigma_y(g_{k+1,1}, \dots, g_{k+1,n})$, $x = \sigma_x(h_{k+1,1}, \dots, h_{k+1,n})$. Thus we express $[y, x]c[y, x]^{-1}c^{-1}$. In this automorphism all vertices from X^{k+1} that belong to the subtree $v_{ki}X^*$ are trivial result of the reduction, but the root of the tree already contains non-trivial even permutations as a commutator. Also, all subtrees having root in these vertices, namely $v_{k1}X^*$, $v_{k2}X^*$, ..., $v_{kn}X^*$ contain only trivial permutations in their vertices. Note that we have computed a double left-based commutator because of $\alpha = [y, x]c[y, x]^{-1}c^{-1} = [[y, x], c]$. As a result of simplification we obtain automorphism $\alpha = \sigma_\alpha(\alpha_{k+1,1}, \dots, \alpha_{k+1,n})$ having all states of form $\alpha_{k+1,i}$ that are trivial. This automorphism α contains a non-trivial permutation only at the root of the tree. The root permutation σ_α of α is even since α is a commutator. Let us show that σ_α can be expressed as an arbitrary permutation from A_n .

In order to generate the whole A_n , it is sufficient to show that we can generate all 3-cycles. Let us show that such elements $\sigma_x, \sigma_y, \sigma_c$ can always be found for $n > 3$. Now, following the similar construction from the proof of the Galois theorem, we assume that the cyclic type of x consists of one cycle $(1, 2, 3, \dots, k)$. For instance, let $k = 5$. Then similarly to Galois theorem proof we compute $[\sigma_y, \sigma_x] = (3, 4, 5)(1, 2, 3, 4, 5)(3, 5, 4)(5, 4, 3, 2, 1) = (2, 4, 5)$. The element σ_c acting only on $v_{2i} : i \in \text{Fix}(\sigma_y)$ for simplification of permutations in subset of vertices from dual set $v_{k+1i} \in \text{Supp}_{A_n}(\sigma_y)$ under and vertices on consecutive lower levels under $v_{k+1i} \in \text{Supp}_{A_n}(\sigma_y)$, at the same time commuting with $[\sigma_x, \sigma_y]$. Now according our construction we set that $\sigma_c = (1, 2)$. Let \mathbb{B}^{TM} s verify that our conditions are satisfied:

$$\begin{cases} \text{Supp}\{1, 2\} \cap \text{Supp}\{3, 4, 5\} = \emptyset \\ \text{Supp}\{1, 2\} \cap \text{Supp}\{2, 5, 4\} = 2 \neq \emptyset. \end{cases}$$

It remains to compute left-based composition of commutators: $[[\sigma_y, \sigma_x], \sigma_c] = [(1, 2), (2, 5, 4)] = (1, 2)(2, 5, 4) \times (1, 2)(2, 4, 5) = (4, 1, 2)$ as well as $[[\sigma_x, \sigma_y], \sigma_c] = [(2, 5, 4), (1, 2)] = (2, 5, 4)(1, 2) \times (2, 4, 5)(1, 2) = (2, 1, 4)$.

Varying the choice of the elements for $\text{Supp}(\sigma_c)$, taking into account the constraint $\text{Supp}_{A_n}(c) \subseteq \text{Fix}_{A_n}(y)$, we can get different 3-cycles after calculating $[[\sigma_y, \sigma_x], \sigma_c]$. \square

Definition 19. The set of Kaloujnine tables with vertex permutation in tableau 2-nd part $[e]_0, [e \dots, e]_1, [e \dots, e]_2, [\pi_1 \dots, \pi_n; \pi_{n+1}, \dots, \pi_{n^3}]_2$, satisfying the condition that either all $\pi_{i_j} \in S_n \setminus A_n$ or all $[\pi_{i_j}]_3 \in A_n$ for $1 \leq i_1 \leq n^2, n^2 + 1 \leq i_2 < 2n^2, \dots, (n-1)n^2 + 1 < i_n \leq n^3$ be denoted by $\tilde{T}_{n^2}^{(3)}$.

Proposition 35. Furthermore $\tilde{T}_{n^2}^{(3)}$ forms normal subgroup in $St_W(3)$ as well as in $S_n \wr S_n \wr S_n$.

The *monolith* of $S_n \wr S_m$ is $e \wr A_m$.

Corollary 36. If H has depth k and its k -th level subgroup $H(k) \simeq \widetilde{A}_{n^j}^k, 1 < j < k$ then H is subgroup of normal subgroups denoted by $\widetilde{A}^{(k)} \triangleleft W_m = \wr_{i=1}^m S_{n_i}$, where $\widetilde{A}^{(k)} \simeq \widetilde{A}_{n^k} \wr H_{k+1} \wr H_{k+2} \wr \dots \wr H_{m-1} \wr H_m \triangleleft \wr_{i=1}^m S_{n_i}$ and $H_j \in \{S_n, \widetilde{A}_{n^j}\}$ and $m \geq j > k$. Furthermore the subgroup $\widetilde{T}^{(k)}$ is subgroup from saturation set [18] of normal subgroup of $\wr_{i=1}^m S_{n_i}$ and forms an finite proper normal subgroup.

Corollary 37. If H has depth k and its k -th level subgroup $H(k) \simeq \widetilde{T}_{n^j}^k, 1 < j < k$ then H is subgroup of normal subgroups denoted by $\widetilde{T}^{(k)} \triangleleft W_m = \wr_{i=1}^m S_{n_i}$, where $\widetilde{T}^{(k)} \simeq \widetilde{T}_{n^k} \wr H_{k+1} \wr H_{k+2} \wr \dots \wr H_{m-1} \wr H_m \triangleleft \wr_{i=1}^m S_{n_i}$ and $H_j \in \{S_n, \widetilde{A}_{n^j}\}$ and $m \geq j > k$. Furthermore the subgroup $\widetilde{T}^{(k)}$ is subgroup from saturation set [18] of normal subgroup of $\wr_{i=1}^m S_{n_i}$ and forms an finite proper normal subgroup.

Remark 38. Subgroups from 1) can be partitioned into 2 classes: a) subgroups which possesses the quotient subgroup of form $A_j \wr E$ such that A_j can be extended by N_i to $S_n \wr S_m$. I.e. for which extension problem for A_j by $E \wr N_i$ to $A_j \wr N_j$ is solvable.

Subgroups from 1) which has not such extension.

Theorem 39. The set of normal subgroups of $\wr_{i=1}^\infty S_{n_i} \widetilde{A}_{n^i}^{(\geq k)}, k > i$ is saturated by $\widetilde{A}_{n^i}^{(\geq k)}, k > i$.

Proof. In view of Remark we have embedding each $\widetilde{A}_{n^i}^{(k)}, k > i$ is normal in $\widetilde{A}_{n^k}^{(k)}$. \square

Theorem 40. The full list of normal subgroups of $S_n \wr S_n \wr S_n$ consists of 50 normal subgroups. These subgroups are the following:

- 1 **Type** T_{023} contains: $E \wr \widetilde{A}_n \wr H, \widetilde{T}_n \wr H$, where $H \in \{\widetilde{A}_n, \widetilde{A}_{n^2}, S_n\}$. There are 6 subgroups.
- 2 **Additional class** of subgroups in T_{023} with new base of wreath product subgroup \widetilde{A}_{n^2} : $E \wr S_n \wr \widetilde{A}_{n^2}, E \wr N_i(S_n \wr S_n)$. Therefore this class has 12 new subgroups. Thus, the total number of normal subgroups in **Type** T_{023} is 18.
- 3 **Type** T_{003} : $A_{00(n^2)}^{(3)} = E \wr E \wr \widetilde{A}_{n^2}, \widetilde{T}_{n^2}^{(3)}$. Hence, here are 2 new subgroups.
- 4 **Type** T_{123} : $N_i(S_n \wr S_n) \wr S_n, N_i(S_n \wr S_n) \wr \widetilde{A}_n$ and $N_i(S_n \wr S_n) \wr \widetilde{A}_{n^2}$. Thus, there 30 new subgroups in T_{123} .

Futhermore all of these groups are splitable.

Thus, we have find the complete list of 50 normal subgroups in $S_n \wr S_n \wr S_n$ there are 2 of them in non-proper subgroups.

The proof is based on Theorem 23 stating that there are 10 normal subgroups in $S_n \wr S_n$.

3.2 Normal subgroups in $\prod_{i=1}^{\infty} S_{n_i}$

Consider the set N_i consisting of elements of S_n with even product of all level elements (from any level). This set forms the subgroup N_i of $\prod_{i=1}^{\infty} S_{n_i}$. Here the index i means minimal depth of element on N_i and calls the depth of the subgroup N_i . Thus, the i -th level has a subdivision of the tree \mathbf{T} at which acts $\prod_{i=1}^j S_{n_i}$ on $\prod_{i=1}^j n_i$ subtrees, at each level of which the product of vertex permutations is even.

One easy can verify that H_j is normal subgroup in $\prod_{i=1}^{\infty} S_{n_i}$.

Statement. The following subgroup embedding is valid:

$$H_1 \supset H_2 \supset H_3 \supset \dots$$

Proof. Let $1 \leq m \leq k$ then at the i -th level, where $m \leq i \leq k$, in the elements of N_k are identical substitutions e . Therefore, their product at the m -th level is an even permutations, which is a characteristic property of N_m , so N_k belongs to N_m .

Note that is, N_m is a group. The definition of $\prod_{i=1}^{\infty} S_{n_i}$ implies that the product of its elements from the k -th level forms as the product of permutations of elements the k -th level. It means that elements k -th level satisfy property (9) too.

The presence of the inverse element in such a set N_m can be proved by contradiction.

Definition 20. We call by k -coordinate tableau such tableau having $[u]_l = \varepsilon$ when $l \neq k$. The subgroup of all l -coordinate tableaux is denoted by $[W]_l$.

Theorem 41. The bases B_i forms topology of wreath product by determining of open neighborhoods system of neutral element $t = [\varepsilon, \varepsilon, \dots, \varepsilon, \dots]$.

Definition 21. Tableau u has depth m if all $[u]_i = \varepsilon$ provided $i = 1, 2, \dots, m-1$; and $[u]_m \neq \varepsilon$. We set that depth of e is equal to ∞ .

Statement. Each subgroup H_i is a normal subgroup in $\prod_{i=1}^{\infty} S_{n_i}$. Furthermore H_i form the i -th base of wreath product $\prod_{i=1}^{\infty} S_{n_i}$.

Obviously, because the conjugation at each level can only add an even number of odd permutations. (Moreover, the conjugation does not change the cyclic type of substitution). Each substitution of the i -th level conjugates only one substitution of the same level, therefore the rule (2) is applicable.

In general case conjugation by arbitrary element does not change parity of product of elements of H_i .

Theorem 42. Let H be a normal subgroup of $W = \prod_{i=1}^{\infty} S_{n_i}$. Let us assume that $H_1 < H$.

Then $\prod_{i=1}^{\infty} S_{n_i}/H$ is isomorphic to some subgroup of the group $\prod_1^{\infty} \mathbb{Z}_2$.

Proof. Recall that we have defined subgroup of l -th level $W(l)$ in [27] of the group W .

Define a homomorphism from $W(l)$ onto \mathbb{Z}_2 , whose kernel is $N_1(l)$. Another words kernel of this homomorphism is subgroup $N_1(l)$ of l -th level of group $N_1(l)$. If products of transpositions of l -th level that belongs to $N_1(l)$ is even then the image of $W(l)$ is non trivial element 1 from \mathbb{Z}_2 , if products of its transpositions is odd permutation then its image is 0.

$N_1(l)$. Another words kernel of this homomorphism is subgroup $N_1(l)$ of j -th level of group N_1 .

$H_j(l)$. Another words kernel of this homomorphism is subgroup $H_j(l)$ of j -th level of group H_j .

The proof will be constructed by adding elements to the group $N < H$. Since, N contains arbitrary element that having an even product of level permutations, we need to add those elements that are characterized only by product of permutations on each level. We are interested only in existence of odd permutation on each level, because by conjugation it is possible to place permutations of arbitrarily level in arbitrarily order. Let us consider two arbitrarily elements $a, b \in \prod_{i=1}^{\infty} S_{n_i}$. Let us also consider $l - th$ level of each of these elements. Product of permutations in $a^{-1}ba$ of $l - th$ level is even if and only if their product as elements of a group S_{n_l} forms even permutation of S_{n_l} . In general case, we have to notice that conjugation by an arbitrary element cannot change the product of permutations on each level each sub-tree, that corresponds to H . Indeed, by the definition in $\prod_{i=1}^{\infty} S_{n_i}$ each permutation $i + 1$ -th level conjugate one permutation of $i + 1$ -th level of initial element. Therefore, all H are normal subgroups in $\prod_{i=1}^{\infty} S_{n_i}$.

Let's consider bijection $\phi : \prod_{i=1}^{\infty} S_{n_i} \rightarrow \prod_1^{\infty} \mathbb{Z}_2$. In other words, ϕ gives to each element from $\prod_{i=1}^{\infty} S_{n_i}$ infinite sequence of zeros and ones. This bijection is homomorphism of the group. If we assume that all elements of the group $\prod_{i=1}^{\infty} S_{n_i}$ are equivalent if and only if parity of products of elements in all levels coincides. Since, each representative of the class will correspond to only one element from $\prod_1^{\infty} \mathbb{Z}_2$, we get statement of the theorem. \square

Theorem 43. *Normal subgroups consisting of finite elements, form a lattice that is isomorphic to a lattice of subgroups of infinite iterate in one side Cartesian product of cyclic groups of order 2.*

Proof. Proof is obvious from the previous *Theorem*. \square

Theorem 44. *Each N_i is a normal subgroup of $\prod_{i=1}^{\infty} S_{n_i}$.*

Proof. We need to prove that conjugation of elements from N_i with elements from $\prod_{i=1}^{\infty} S_{n_i}$ belongs to N_i . Consider N_1 . In this case everything is obvious: on each level conjugation can add only even number of odd permutations, that will not change the parity of the product of permutations on each level. In general case, we have to notice that conjugation by an arbitrary element cannot change the product of permutations on each level each sub-tree, that corresponds to N_i . Indeed, by the definition in $\prod_{i=1}^{\infty} S_{n_i}$ each permutation $i + 1$ -th level conjugate one permutation of $i + 1$ -th level of initial element. Therefore, all N_i are normal subgroups in $\prod_{i=1}^{\infty} S_{n_i}$.

Let $(\prod_{i=1}^{\infty} S_{n_i})_{l-1}$ is a subgroup of $\prod_{i=1}^{\infty} S_{n_i}$, where identical permutations are on the first $l - 1$ levels. The following theorem holds:

Consider some matrix space M over \mathbb{Z}_2 , each matrix has infinite number of rows. We assign to each depth l of some element \mathbb{Z}_2 the number $\prod_{i=1}^{l-1} n_i$.

Definition Two elements of matrix space M , that is characterized by a integer l and group $\prod_{i=1}^{\infty} S_{n_i}$, is ω -equivalent, if one of these elements can be obtained from another by the following transformations:

1. aloud to permute first n_{l-1} columns, second n_{l-1} columns, etc.

2. aloud to permute first $n_l - 2$ groups of columns, that consist of n_{l-1} columns, second n_{l-2} groups of columns , etc.
3. aloud to permute first n groups of columns. □

4 Inverse limit of wreath product of permutation groups

Definition. **Baire space** is defined as the Cartesian product of an **infinitely countable set** of copies of a set of natural numbers and endowed with a **product topology** (where each copy of the set of natural numbers is assigned a *discrete topology*).

Baire space is often presented by a tree of **finite sequences of natural numbers**.

Let $W_1 = S_X$ symmetric group of permutations on X , then we can determine group recursively $W_{n+1} = (W_n, X^n) \wr S_X$ as wreath product of W_n acting of X^n with permutation group acting on X . The base of wreath product consists of functions $Fun(X^n, S_X)$. But taking into account that S_X acts on X we can consider functions $Fun(X^n, X) =: B_n$. This subgroup B is kernel of natural epimorphism:

$$\rho_n : W_{n+1} \rightarrow W_n.$$

Or in general case $\rho_{k,n} : W_k \rightarrow W_n$. In such a way we can construct inverse system of groups:

$$I = \langle \rho_n, W_{n+1} \rangle$$

In general case we can consider different alphabets X_i and $W_k = \wr_{i=1}^k S_{n_i}$ then $\langle \rho_n, W_{n+1} \rangle = \left\langle \rho_n, \wr_{i=1}^{n+1} S_{n_i} \right\rangle$ This projective limit we will denote by

$$PW_\infty(X) := \varprojlim W_n.$$

We define canonical projection as the following homomorphism from projective limit to objects of inverse system: $\pi_n : PW_\infty(X) \rightarrow W_n(X)$. Provided this projection, the group $PW_\infty(X)$ acts on Baire space over alphabet X .

The group $PW_\infty(X)$ is the projective limit of I because of:

1) there exist set of mappings $\pi_n : PW_\infty(X) \rightarrow W_n(X^n)$, $\pi_k : PW_\infty(X) \rightarrow W_k(X^k)$ and epimorphism $\rho_{n,k} : W_k(X^k) \rightarrow W_n(X^n)$, $\forall n \leq k$ such that $\pi_n = \rho_{n,k} \pi_k$, where π_i is canonical projection on i -th component of algebraic system of groups $W_i(X)$. The mapping $\rho_n : W_{n+1}(X) \rightarrow W_n(X)$ is epimorphism with the kernel $\ker(\rho_n) = Fun(X^n, S_X)$.

2) Also the following composition of epimorphism is epimorphism again: $\rho_{n,m} =$

$\rho_{n,k}\rho_{k,m}, \forall n \leq k \leq m.$

We consider the bases B_i as open neighborhoods system of neutral element e of $W = \varprojlim_{i=1}^{\infty} S_{n_i}$ that is a normal subgroup in $\varprojlim_{i=1}^{\infty} S_{n_i}$. Furthermore B_i form the i -th base of wreath product $\varprojlim_{i=1}^{\infty} S_{n_i}$.

For this goal firstly open neighborhoods B_i have to satisfy to conditions of there must be consistency over multiplication (or an action on an open set) $gB_i = B_j$ for $\forall i \in \mathbb{N}$.

Secondly for an arbitrary open neighborhoods B_i of e there must exists an inverse element B_i^{-1} in the group W .

It is obviously that these conditions are satisfied for bases B_i of W .

Theorem 45. *Theorem. An isometry group of Baire space is isomorphic to $PW_{\infty}(X)$.*

We call by k -coordinate tableau such tableau having $[u]_k = \varepsilon$ when $k \neq l$. The subgroup of all l -coordinate tableaux is denoted by $[W]_l$.

Tableau u has depth m if all $[u]_i = \varepsilon$ provided $i = 1, 2, \dots, m$; and $[u]_i \neq \varepsilon$.

Remark 46. *The subgroup $[W]_l$ is isomorphic to μ_l -th direct power of subgroup S_l from wreath product $W_l = S_{X_1} \wr S_{X_2} \wr \dots \wr S_{X_l} \wr S_{X_{l+1}}$, where $\mu_{l-1} = |X_1 \times X_2 \times \dots \times X_{l-1}|$.*

Conclusion. Further, the monolith (i.e. the unique minimal normal subgroup) of these wreath products has been investigated. The monolith exists when $n \equiv 0 \pmod{2}$.

References

- [1] *Skuratovskii R.V.*, Invariant structures of wreath product of symmetric groups. Naukovy Chasopus of Science hour writing of the National Pedagogical University named after M.P. Dragomanova. (in ukrainian) Series 01. Physics and Mathematics. — 2009. Issue 10. — P. 163 B– 178. [Source [http : //enpuir.npu.edu.ua : 8080/bitstream/handle/123456789/13876/skuratokskyyi163 – 178.pdf](http://enpuir.npu.edu.ua:8080/bitstream/handle/123456789/13876/skuratokskyyi163-178.pdf)]
- [2] *Robert Carmichael* Introduction to the theory of Groups of Finite Order. Dover, 1956, 431. ISBN 0-486-60300-8.
- [3] *R. D. Carmichael* Tactical Configurations of Rank Two. American Journal of Mathematics, Vol. 53, No. 1 (Jan., 1931), pp. 217-240.
- [4] *Aviv Rotbart* Generator sets for the alternating group. Seminaire Lotharingien de Combinatoire 65 (2011), Article B65b.

- [5] *J.D.P. Meldrum*, Wreath Products of Groups and Semigroups. Pitman Monographs and Surveys in Pure and Applied Mathematic. 1st Edition. Jun (1995). 425 p.
- [6] Leonov U. G., On the representation of groups approximated by finite p -groups. // *Ukr. Math. Journ.*, 2011.—*ГЗБ'*. 63, *Г,Б№* 11.—*Р.* 15018-1511.
- [7] *Nekrashevych V.*, Self-similar groups. International University Bremen. American Mathematical Society. Monographs. Volume 117. 230 p.
- [8] *Lavrenyuk Y.*, On the finite state automorphism group of a rooted tree Algebra and Discrete Mathematics Number 1. (2002). pp. 79-87
- [9] Vitaly Ivanovich Sushchansky. Normal structure of the isometric metric group spaces of p -adic integers. *Algebraic structures and their application. Kiev*, Visn of KNU, 1988. pages 113–121.
- [10] *Sushchansky V.I.* ℓ -Wreath products and isometries of generalized Baire metrics. // *Translated from Ukrainskii Matematicheskii Zhurnal*,.Vol. 43, No. 7 and 8, Issue – 1991.— pp. 1031-1038. *N 8.—ГЗБ'*. 1013–1038.
- [11] Kaloujnine L. A. Sur les p -group de Sylow. // *C. R. Acad. Sci. Paris.* — 1945. — **221**. — P. 222–224.
- [12] R. V. Skuratovskii, A. Williams "Irreducible bases and subgroups of a wreath product in applying to diffeomorphism groups acting on the Mobius band", **2021**. *Rendiconti del Circolo Matematico di Palermo Series 2*, 70(2), 721-739. <https://doi.org/10.1007/s12215-020-00514-5>.
- [13] *Guralnick R.*, Commutators and wreath products // *Contemporary Mathematics*. Volume 524, 2010.
- [14] *Drozd, Y.A., Skuratovskii R.V.*, Generators and relations for wreath products of groups. *Ukr. Math. J.* (2008), 60, pp. 1168–1171.
- [15] *Sushchansky V.I.* Normal structure of the isometry group of the metric space of p -adic integers. Algebraic structures and their application. Kyiv: KSU, 1988. 113-121 pp.
- [16] *Sachkov, V.N.*, Combinatorial methods in discrete Mathematics. Encyclopedia of mathematics and its applications 55. Cambridge Press. 2008. P. 305.
- [17] *Dashkova O. Yu.* On groups of finite normal rank. Algebra Discrete Math. 2002. 1, No. 1. P. 64-68.

- [18] *A.K. Shlepin*, On periodic groups saturated with finite simple subgroups, Groups. Group rings: abstract. report intl. Symp., Lvov, 1996.
- [19] *Skuratovskii R.*, Generators and relations for sylows p -subgroup of group S_n . *Naukovi Visti KPI*. 4 (2013), pp. 94–105. (in Ukrainian)
- [20] *Skuratovskii R. V.*, Minimal generating systems and properties of $Syl_2 A_{2^k}$ and $Syl_2 A_n$. X International Algebraic Conference in Odessa dedicated to the 70th anniversary of Yu. A. Drozd. (2015), pp. 104.
- [21] *Skuratovskii R. V.*, Structure and minimal generating sets of Sylow 2-subgroups of alternating groups. Source: <https://arxiv.org/abs/1702.05784v2>
- [22] *Skuratovskii R.*, Commutators subgroups of sylow subgroups of alternating and symmetric groups their minimal generating sets. The XII International Algebraic Conference in Ukraine (2019) Vinnytsia, p. 75.
- [23] *Dmitruk U., Suschansky V.*, Structure of 2-sylow subgroup of alternating group and normalizers of symmetric and alternating group. *UMJ*. (1981), N. 3, pp. 304-312.
- [24] *Ward D.*, Topics in Finite Groups: Homology Groups, Pi-product Graphs, Wreath Products and Cuspidal Characters. Manchester Institute for Mathematical Sciences School of Mathematics. July (2015) P. 253.
- [25] *Birkhoff G.* Theory of lattice. M. Nauka 1984, P. 564.
- [26] *Coxeter G.S., Moser U.O.* Generating elements and determining relations of discrete groups. M. Nauka., 1980, 240p.
- [27] *Ruslan V. Skuratovskii.* On commutator subgroups of Sylow 2-subgroups of the alternating group, and the commutator width in wreath products. / Ruslan V. Skuratovskii // European Journal of Mathematics. – 2021. – vol. 7, no. 1. – P. 353-373.
- [28] *Sushchansky V.I., Sikora V.S.* Operations on the groups of permutations. Chernivtsi: Ruta, 2003.14.
- [29] *Ruslan V. Skuratovskii.* On commutator subgroups of Sylow 2-subgroups of the alternating group, and the commutator width in wreath products. / Ruslan V. Skuratovskii // European Journal of Mathematics. – 2021. – vol. 7, no. 1. – P. 353-373. <https://doi.org/10.1007/s40879-020-00418-9>
- [30] *R. Grigorchuk, V. Nekrashevich, V. Sushchanskii*, Automata, Dynamical Systems, and Groups, *Trudy mat. inst. imeny Steklova*. BN°231. (2000) 134–214.

- [31] Skuratovskii R.V. Normal subgroups of iterated wreath products of symmetric alternating groups // Modern problems in mathematics and its applications International (53rd National) Youth School-Conference, 2022, Yekaterinburg, 2022 from January 31 to February 4 section Group theory, pp. 1-2. <https://sopromat.imm.uran.ru>.
- [32] Skuratovskii R.V., Aled Williams. Normal subgroups of iterated wreath products of symmetric groups. // International Algebraic Conference, dedicated to the 90th anniversary of the birth of A.I. Starostin. Book of Abstracts. 05 October 2021 - 09 October 2021. pp. 108-110. Source: [https://www.imm.uran.ru/rus/Library/Lists/New entries to the library/Attachments/641/contKonfStar.pdf](https://www.imm.uran.ru/rus/Library/Lists/New%20entries%20to%20the%20library/Attachments/641/contKonfStar.pdf)
- [33] Ruslan Skuratovskii. Involutive irreducible generating sets and structure of sylow 2-subgroups of alternating groups. ROMAI J., v.13, no.1(2017), 117 - 139. ROMAI J., v.13, no.1 (2017), 117-139.
- [34] I. V. Kuznetsova, Yu. Yu. Soroka. The first numbers of Betty orbits of Morse functions on surfaces // Ukr. math. Journal, 2021, vol. 73, No. 2, pp. 179-200.
- [35] P. J. Morandi, Group Extensions and H^3 Archived 2018-05-17 at the Wayback Machine. From his collection of short mathematical notes.