The signless Laplacian spectral radius of graphs without intersecting odd cycles *

Ming-Zhu Chen, A-Ming Liu, School of Science, Hainan University, Haikou 570228, P. R. China,

Xiao-Dong Zhang[†] School of Mathematical Sciences, MOE-LSC, SHL-MAC Shanghai Jiao Tong University, Shanghai 200240, P. R. China

Abstract

Let F_{a_1,\ldots,a_k} be a graph consisting of k cycles of odd length $2a_1 + 1, \ldots, 2a_k + 1$, respectively which intersect in exactly a common vertex, where $k \ge 1$ and $a_1 \ge a_2 \ge \cdots \ge a_k \ge 1$. In this paper, we present a sharp upper bound for the signless Laplacian spectral radius of all F_{a_1,\ldots,a_k} -free graphs and characterize all extremal graphs which attain the bound. The stability methods and structure of graphs associated with the eigenvalue are adapted for the proof.

AMS Classification: 05C50, 05C35

Key words: Brualdi-Solheid-Turán type problem; signless Laplacian spectral radius; intersecting odd cycles free; extremal graph.

1 Introduction

Let G be an undirected simple graph with vertex set $V(G) = \{v_1, \ldots, v_n\}$ and edge set E(G), where e(G) is the number of edges of G. For $v \in V(G)$, the *neighborhood* $N_G(v)$ of v is $\{u : uv \in E(G)\}$ and the *degree* $d_G(v)$ of v is $|N_G(v)|$. We write N(v) and d(v) for $N_G(v)$ and $d_G(v)$ respectively if there is no ambiguity. Denote by $\Delta(G)$ and $\delta(G)$ the maximum and minimum degree of G, respectively. Denote by P_n and C_n the path and the cycle of order n, respectively. For $A, B \subseteq V(G)$, e(A) denotes the number of the edges of G with both endvertices in A and e(A, B) denotes the number of the edges of G with one endvertex in A and the other in B. For two vertex disjoint graphs G and H, we denote by $G \bigcup H$ and $G \nabla H$ the union of G and H, and the join of G and H, i.e., joining every vertex of G to every vertex of H, respectively. Denote by kG the union of k disjoint copies of G. Denote by $S_{n,t} = K_t \bigcup \overline{K}_{n-t}$, i.e., the join of a complete graph K_t and the independent set I of size n-t. Denote by $L_{r,t} = K_1 \nabla r K_t$, i.e., the graph consists of r complete graph K_{t+1} which intersect in exactly a common vertex. For graph notation and terminology undefined here, we refer the readers to [1].

^{*}This work is supported by the National Natural Science Foundation of China (Nos. 11971311, 12026230), and the Hainan Provincial Natural Science Foundation of China (Nos. 120RC453, 120MS002).

 $[\]label{eq:mail:mzchen@hainanu.edu.cn, amliu@hainanu.edu.cn, xiaodong@sjtu.edu.cn.$

[†]Corresponding author: Xiao-Dong Zhang (Email: xiaodong@sjtu.edu.cn),

We say a graph G is *H*-free if it does not contain *H* as a subgraph. The Turán number of a graph *H* is the maximum number of edges that is in an *H*-free graph of order *n*, and is denoted by ex(n, H). An *H*-free graph of order *n* with ex(n, H) edges is called an *extremal graph* for *H* and denote Ex(n, H) by the set of all extremal graphs of order *n* for *H*. To determine ex(n, H) and characterize those graphs in Ex(n, H) is a fundamental problem (called Turán-type problem) in extremal graph theory. It will be interesting to look for some nice graphs *H* such that ex(n, H) and Ex(n, H) will be characterized. The graph $F_{a_1,...,a_k}$ which is a graph consisting of *k* cycles of odd length $2a_1 + 1, \ldots, 2a_k + 1$, respectively which intersect in exactly a common vertex may be of interest, where $k \ge 1$ and $a_1 \ge a_2 \ge \cdots \ge a_k \ge 1$. If k = 1, then $F_{a_1,...,a_k}$ is an odd cycle C_{2a_1+1} . Simonovits [22] proved that $ex(n, C_{2a_1+1}) = \lfloor \frac{n^2}{4} \rfloor$ and $Ex(n, C_{2a_1+1})$ is a balance complete bipartite graph. If $k \ge 2$ and $a_1 = \cdots = a_k = 1$, then $F_{a_1,...,a_k}$ is denoted by $F^{(k)}$ which is called the *friendship graph*. In 1995, Erdős, Füredi, Gould, Gunderson [11] significantly extended Mantel's result and proved the following interesting result.

Theorem 1.1. [11] Let $k \ge 1$ and $n \ge 50k^2$. Then

$$ex(n, F^{(k)}) = \left\lfloor \frac{n^2}{4} \right\rfloor + \begin{cases} k^2 - k, & \text{if } k \text{ is odd,} \\ k^2 - \frac{3}{2}k, & \text{if } k \text{ is even.} \end{cases}$$

Furthermore, if k is odd, then $Ex(n, F^{(k)})$ consists of graphs which are constructed by taking a complete bipartite graph with two partite of size $\lceil \frac{n}{2} \rceil$ and $\lfloor \frac{n}{2} \rfloor$ and embedding two vertex disjoint copies of K_k in one side. If k is even, then $Ex(n, F^{(k)})$ consists of graphs which are constructed by taking a complete bipartite graph with two partite of size $\lceil \frac{n}{2} \rceil$ and $\lfloor \frac{n}{2} \rfloor$ and embedding a graph with 2k - 1 vertices, $k^2 - \frac{3}{2}k$ edges with maximum degree k - 1 in one partite.

If $k \geq 2$ and $a_1 \geq \cdots \geq a_s \geq 2$, $a_{s+1} = \cdots = a_k = 1$ with $1 \leq s \leq k$, then F_{a_1, \cdots, a_k} is denoted by $H_{k,s}$, i.e., $H_{k,s}$ is the graph consisting of k odd cycles and k - s triangles which intersect in exactly a common vertex. In 2018, Hou, Qiu and Liu [13], and Yuan [23] independently proved the following result.

Theorem 1.2. [13, 23] Let $k \ge 2$ and $1 \le s \le k$. Then

$$ex(n, H_{k,s}) = \left\lfloor \frac{n^2}{4} \right\rfloor + (k-1)^2$$

for sufficient large n. Moreover, $Ex(n, H_{k,s})$ consists of a balance complete bipartite graph with a complete bipartite graph $K_{k-1,k-1}$ embedded in one partite if $(k, s) \neq (4, 1)$; a balance complete bipartite graph with a complete bipartite graph $K_{3,3}$ or $3K_3$ embedded in one partite if (k, s) = (4, 1)

In spectral extremal graph theory, there is an analogy between the Turán type problem and the Brualdi-Solheid-Turán type problem which are proposed by Nikiforov [17]. The Brualdi-Solheid-Turán type problem is how to determine the maximum spectral radius of an *H*-free graph of order *n* and characterize those graphs which attain the maximum spectral radius. The Brualdi-Solheid-Turán type problem of the adjacent spectral spectral radius has been studied for various kinds of *H* such as the complete graph [16], the complete bipartite graph [18], the cycles or paths of specified length [17], the linear forest [3], and star forest [4] and so on. In addition, the Brualdi-Solheid-Turán type problem of the signless Laplacian spectral radius has also been investigated extensively in the literature. For more details, readers may be referred to [8, 12, 19–21, 24, 25]. It is of interest to consider this problem for F_{a_1,\dots,a_k} . The adjacency matrix of G is the $n \times n$ matrix $A(G) = (a_{ij})$, where $a_{ij} = 1$ if v_i is adjacent to v_j , and 0 otherwise. Moreover, the matrix Q(G) = D(G) + A(G) is known as the signless Laplacian matrix of G, where D(G) is the degree diagonal matrix of G. The largest eigenvalues of A(G) is called the spectral radius of G. The largest eigenvalues of Q(G) is called the signless Laplacian spectral radius of G and denoted by q(G).

In fact, if k = 1 and $a_1 = 1$, Nikiforov [16] determined the maximum spectral radius of all F_1 -free graphs of order n and proved that the only balance bipartite graphs has the maximum spectral radius; while He, Jin and Zhang [12] obtained a sharp bound for the signless Laplacian spectral radius of all F_1 -free graphs of order n and proved that the corresponding extremal graphs are all complete bipartite graphs $K_{r,s}$ with r+s=n. Later, if k = 1 and $a_1 \ge 2$, Nikiforov [17] and Yuan [24] determined the maximum spectral radius and the signless spectral radius among F_{a_1} -free graphs of order n, respectively. Recently, Cioabă, Feng, Tait and Zhang [5] studied the Brualdi-Solheid-Turán type problem for $F^{(k)}$ free graphs and determined the corresponding spectral extremal graphs, which can be viewed as a spectral analogue of Theorem 1.1.

Theorem 1.3. [5] Let G be an $F^{(k)}$ -free graph of order n with $k \ge 2$. If G has the maximum spectral radius, then

$$G \in Ex(n, F^{(k)})$$

for sufficiently large n.

Inspired by the work of Cioabă, Feng, Tait, and Zhang [5], Zhao, Huang, and Guo [25] focused on the maximum signless Laplacian spectral radius of all $F^{(k)}$ -free graphs of order n and proved the following result.

Theorem 1.4. [25] Let G be an $F^{(k)}$ -free graph of order n. If $k \ge 2$ and $n \ge 3k^2 - k - 2$, then

$$q(G) \le q(S_{n,k})$$

with equality if and only if $G = S_{n,k}$.

Recently, Li and Peng [14] proved the spectral result of all $H_{k,s}$ -free graphs of order n, which can be viewed as a spectral analogue of Theorem 1.2.

Theorem 1.5. [14] Let $k \ge 2$ and $1 \le s \le k$. If G is an $H_{k,s}$ -free graph of order n with the maximum spectral radius, then

$$G \in Ex(n, H_{k,s})$$

for sufficiently large n.

Furthermore, they proposed the following conjecture on the signless Laplacian spectral radius of F_{a_1,\dots,a_k} -free graphs of order n with $k \ge 2$ and $a_1 = \dots = a_k \ge 2$.

Conjecture 1.6. [14] Let G be an F_{a_1,\dots,a_k} -free graph of order n. If $k \ge 2$ and $a_1 = \dots = a_k = a \ge 2$, then

$$q(G) \le q(S_{n,ka})$$

for sufficiently large n with equality if and only if $G = S_{n,ka}$.

Inspired by the above known results and the conjecture on the signless Laplacian spectral radius of F_{a_1,\dots,a_k} -free graphs with $k \geq 2$ and $a_1 = \dots = a_k \geq 2$, we studied the maximum signless Laplacian spectral radius of all F_{a_1,\dots,a_k} -free graphs. Combined with known results, we present the main result of this paper as follows.

Theorem 1.7. Let G be an F_{a_1,\ldots,a_k} -free graph of order $n \ge 8t^2 - 12t + 9$ with $t = \sum_{i=1}^k a_i$. (1) [12] If k = 1 and $a_1 = 1$, then $q(G) \leq q(S_{n,t})$ with equality if and only if G is a complete bipartite graphs $K_{r,s}$ with r + s = n.

(2) [24] If k = 1, $a_1 \ge 2$, and $n \ge 110t^2$, then $q(G) \le q(S_{n,t})$ with equality if and only if $G = S_{n,t}$.

(3) [25] If $k \ge 2$ and $a_1 = \cdots = a_k = 1$, then $q(G) \le q(S_{n,t})$ with equality if and only if $G = S_{n,t}$.

(4) If $k \ge 2$ and $a_1 \ge 2$, then $q(G) \le q(S_{n,t})$ with equality if and only if $G = S_{n,t}$.

Remark. It is worth mentioning that the extremal graphs in Theorem 1.7 (4) are not the same as those of Theorems 1.2 and 1.5. The extremal graphs in Theorems 1.2 and 1.5 are only depend on the number of intersecting triangles and the number of all intersecting odd cycles; while the extremal graph in Theorem 1.7 (4) is not only depend on the number of intersecting odd cycles, but also the lengths of all intersecting odd cycles.

The rest of this paper is organized as follows. In Section 2, some known lemmas are presented. In Section 3, we give the proof of Theorem 1.7.

2 Some Lemmas

Lemma 2.1. [10] Let $k \geq 3$. If G is P_k -free graph of order n, then $e(G) \leq \frac{(k-2)n}{2}$ with equality if and only if G is union of disjoint copies of K_{k-1} .

Lemma 2.2. [10] Let $k \ge 2$. If G is graph of order n with no cycle longer than k, then $e(G) \leq \frac{k(n-1)}{2}$ with equality if and only if $G = L_{r,k-1}$ with n = r(k-1) + 1.

Lemma 2.3. [9] If G is a graph with $\delta(G) \geq 2$, then G contains a cycle of length at least $\delta(G) + 1.$

We also need the stability result on the disjoint paths.

Lemma 2.4. [2] Let $H = \bigcup_{i=1}^{k} P_{2a_i}$ with $k \ge 2$, $a_1 \ge \cdots \ge a_k \ge 1$, and $t = \sum_{i=1}^{k} a_i$. If $\delta(G) \ge t-1$ and G is an H-free connected graph of order $n \ge 2t$. then one of the following holds:

(1)
$$G \subseteq S_{n,t-1}$$

(2) $F = 2P_{2a_1}$ and $G = L_{r,t-1}$, where n = r(t-1) + 1.

The structure of graphs and the spectral radius of graphs will be stated as follows.

Lemma 2.5. [20] Let $t \ge 2$ and $n > 5t^2$. Then (1) $q(S_{n,t}) > n + 2t - 2 - \frac{2(t^2 - t)}{n + 2t - 3} > n + 2t - 3.$ (2) If G is a graph of order n with $q(G) \ge q(S_{n,t})$, then $e(G) \ge tn - t^2 + 1.$

Lemma 2.6. If n = r(t-1) + 2 with $r \ge 1$ and $t \ge 3$, then $q(K_1 \nabla L_{r,t-1}) < n + 2t - 3$.

Proof. Let $G = K_1 \nabla L_{r,t-1}$. Note that the nonincreasing degree sequence (d_1, \ldots, d_n) of G is (n-1, n-1, t, ..., t). By [7, Theorem 4.2], we have

$$q(G) = \frac{d_1 + 2d_3 - 1 + \sqrt{(2d_3 - d_1 + 1)^2 + 8\sum_{i=1}^2 (d_i - d_3)}}{2}$$

= $\frac{n + 2t - 2 + \sqrt{(n - 2t - 2)^2 + 16(n - 2t - 2) + 16t + 16}}{2}$
 $\leq \frac{n + 2t - 2 + (n - 2t - 2) + \frac{3t + 7}{2}}{2}$
 $< n + 2t - 3.$

This completes the proof.

3 Proof of Theorem 1.7

In order to prove Theorem 1.7, we only to prove the assertion holds for F_{a_1,\dots,a_k} with $k \ge 2$ and $a_1 \ge 2$. Hence we always assume that F_{a_1,\dots,a_k} is the graph with $k \ge 2$ and $a_1 \ge 2$ in this section. Firstly, we present several technical lemmas.

Lemma 3.1. Let G be an F_{a_1,\ldots,a_k} -free graph of order $n \ge 8t^2 - 12t + 9$ with $t = \sum_{i=1}^k a_i$. If $q(G) \ge q(S_{n,t})$, then $\Delta(G) = n - 1$.

Proof. By the upper bound for the signless Laplacian spectral radius from Merris [15], there exists a vertex u such that

$$q(G) \le \max_{v \in V(G)} \left\{ d(v) + \frac{1}{d(v)} \sum_{z \in N(v)} d(z) \right\} = d(u) + \frac{1}{d(u)} \sum_{z \in N(u)} d(z).$$

Let A = N(u) and $B = V(G) \setminus (N(u) \bigcup \{u\})$. Then |A| + |B| + 1 = n and

$$q(S_{n,t}) \le q(G) \le d(u) + \frac{1}{d(u)} \sum_{z \in N(u)} d(z) = |A| + 1 + \frac{2e(A) + e(A, B)}{|A|}.$$
(1)

Next we show that d(u) = n - 1. Assume for a contradiction that d(u) < n - 1. We prove the following claims.

Claim 1. No vertex in B is adjacent to every vertex in A.

Suppose that there exists a vertex $v \in B$ which is adjacent to every vertex in A. Since G is $F_{a_1,...,a_k}$ -free, we have G[A] is $P_{2a_1-1} \bigcup (\bigcup_{i=2}^k P_{2a_i})$ -free. Otherwise we can add the path P_{2a_1-1} in G[A] to a path of order $2a_1$ by deleting the first edge e of P_{2a_1-1} with $a_1 \geq 2$ and adding two edges from u to the endvertices of e and thus we obtain an $F_{a_1,...,a_k}$, which is a contradiction. Hence G[A] is P_{2t-1} -free, where $t = \sum_{i=1}^k a_i$. By Lemma 2.1, we have $2e(A) \leq (2t-3)|A|$. By (1) and $e(A, B) \leq |A||B|$, we have

$$q(S_{n,t}) \leq |A| + 1 + \frac{2e(A) + e(A, B)}{|A|}$$

$$\leq |A| + 1 + \frac{(2t-3)|A| + |A||B|}{|A|}$$

$$= n + 2t - 3 < q(S_{n,t}),$$

where the last inequality is from Lemma 2.5 (1) with $8t^2 - 12t + 9 \ge 5t^2$, which is a contraction. This proves Claim 1.

Claim 2. $|B| \le 2t^2 - 2t$.

By Claim 1, we have $e(A, B) \leq (|A| - 1)|B|$. In addition, since G is $F_{a_1,...,a_k}$ -free, we have G[A] is $(\bigcup_{i=1}^k P_{2a_i})$ -free, which implies that G[A] is P_{2t} -free. By Lemma 2.1, we have

$$2e(A) \le (2t-2)|A|.$$

Together with (1), we have

$$q(S_{n,t}) \leq |A| + 1 + \frac{2e(A) + e(A, B)}{|A|}$$

$$\leq |A| + 1 + \frac{(2t-2)|A| + (|A|-1)|B|}{|A|}$$

$$= n + 2t - 2 - \frac{|B|}{n - 1 - |B|}$$

$$< n+2t-2-\frac{|B|}{n+2t-3}$$

Hence $|B| \leq 2t^2 - 2t$ follows from Lemma 2.5 (1). This proves Claim 2.

Let A' be the set of all vertices in A which are adjacent to every vertex in B. **Claim 3.** $|A'| \ge |A| - 2t^2 + 2t$. Since $2e(A) \le (2t-2)|A|$ and

$$e(A,B) \le |A'||B| + (|A| - |A'|)(|B| - 1) = |A||B| - |A| + |A'|,$$

we have

$$q(S_{n,t}) \leq |A| + 1 + \frac{2e(A) + e(A, B)}{|A|}$$

$$\leq |A| + 1 + \frac{(2t-2)|A| + |A||B| - |A| + |A'|}{|A|}$$

$$= n + 2t - 2 - \frac{|A| - |A'|}{|A|}$$

$$\leq n + 2t - 2 - \frac{|A| - |A'|}{n + 2t - 3}.$$

Hence $|A'| \ge |A| - 2t^2 + 2t$ follows from Lemma 2.5 (1). This proves Claim 3.

Let G_1 be the union of all components of G[A] each of which contains at least a vertex in A', and let G_2 be the union of the remaining components of G[A]. Set $n_1 = |V(G_1)|$ and $n_2 = |V(G_2)|$. Note that G_2 is also $(\bigcup_{i=1}^k P_{2a_i})$ -free which implies that G_2 is P_{2t} -free. By Lemma 2.1 again, $e(G_2) \leq (t-1)n_2$.

Claim 4. G_1 contains no cycle longer than 2t - 3.

Assume for a contradiction that G contains a cycle C_p with $p \ge 2t-2$. Since G is F_{a_1,\ldots,a_k} free graph, G[A] must be $\bigcup_{i=1}^k P_{2a_i}$ -free, which implies $p \le 2t-1$. Hence $2t-2 \le p \le 2t-1$. By the definition of G_1 , there must be a vertex $z \in A'$ that either belongs to C_p or can be joined to a vertex of C_p by a path Q contained in G_1 . By Claims 2-3,

$$|A'| \ge |A| - 2t^2 + 2t = n - 1 - |B| - 2t^2 + 2t \ge n - 4t^2 + 4t - 1 \ge 2t > 2.$$

We consider the following two cases.

Case 1. $z \in V(C_p)$.

There exists a vertex $v \in A' \setminus V(C_p)$ by $|A'| \geq 2t > p$. Choose a vertex $w \in B$ and construct a path P of order 2t whose first three successive vertices are v, w, z respectively and all vertices of P except w are in A. As a result, $G[A \bigcup \{w\}]$ contains $\bigcup_{i=1}^{k} P_{2a_i}$ as a subgraph, where $\bigcup_{i=1}^{k} V(P_{2a_i}) = V(P)$ and the first three successive vertices of P_{2a_1} are v, w, z respectively. Thus G contains F_{a_1,\ldots,a_k} as a subgraph, which is a contradiction.

Case 2. z is joined to a vertex of C_p by a path Q contained in G_1 .

Choose a vertex $w \in B$. If there exists a vertex $v \in A' \setminus (V(C_p) \bigcup V(Q))$, then we can get a path P of order 2t whose first three successive vertices are v, w, z respectively and all vertices P except w are in A. Otherwise, we can also construct a path P of order 2t whose first three successive vertices are z, w, v respectively and all vertices of P except w are in A. Then $G[A \bigcup \{w\}]$ contains $\bigcup_{i=1}^{k} P_{2a_i}$ as a subgraph, implying that G contains F_{a_1,\ldots,a_k} as a subgraph. This is a contradiction. Hence Claim 4 holds.

By Claim 4 and Lemma 2.2, we have

$$2e(G_1) \le (2t-3)n_1.$$

Then

$$2e(A) = 2e(G_1) + 2e(G_2)$$

$$\leq (2t-3)n_1 + (2t-2)n_2 = (2t-3)(n_1+n_2) + n_2$$

$$\leq (2t-3)|A| + |A| - |A'| \leq (2t-3)|A| + 2t^2 - 2t.$$

By (1),

$$q(S_{n,t}) \leq |A| + 1 + \frac{2e(A) + e(A, B)}{|A|}$$

$$\leq |A| + 1 + \frac{(2t-3)|A| + 2t^2 - 2t + |A||B|}{|A|}$$

$$= n + 2t - 3 + \frac{2t^2 - 2t}{|A|}.$$

Hence it follow from Lemma 2.5(1) that

$$n+2t-2-\frac{2t^2-2t}{|A|} \le n+2t-2-\frac{2t^2-2t}{n+2t-3} \le n+2t-2-1+\frac{2t^2-2t}{|A|}$$

which implies that $|A| \leq 4t^2 - 4t$. Hence by Claim 3,

$$n = 1 + |A| + |B| \le 1 + 4t^2 - 4t + 2t^2 - 2t < 8t^2 - 12t + 9,$$

which is a contradiction. Hence d(u) = n - 1 and it completes the proof.

Lemma 3.2. Let G be a $\bigcup_{i=1}^{k} P_{2a_i}$ -free graph of order $n > t^2 - t - 1$ with $k \ge 2$, $a_1 \ge \cdots \ge a_k \ge 1$, and $t = \sum_{i=1}^{k} a_i$. If $e(G) \ge (t-1)n - (t^2 - t - 1)$, then there exists an induced subgraph H of G such that $|V(H)| \ge n - (t^2 - t - 1)$, $\delta(H) \ge t - 1$, and $d_H(v) \le t - 2$ for every vertex $v \in V(G) \setminus V(H)$.

Proof. If $\delta(G) \geq t-1$, then let H = G. Otherwise we delete a vertex v_1 with the minimum degree $\delta(G) \leq t-2$ and obtain $G_1 = G - v_1$. If $\delta(G_1) \geq t-1$, let $H = G_1$. Otherwise we may continue this way until $\delta(G_{r-1}) \leq t-2$ and G_r obtained from G_{r-1} by deleting a vertex with the minimum degree and $\delta(G_r) \geq t-1$. We show that $r \leq t^2 - t - 1$.

Since G_r is $\bigcup_{i=1}^k P_{2a_i}$ -free, we have G_r is P_{2t} -free. By $|V(G_r)| = n - r$ and Lemma 2.1,

$$(t-1)(n-r) \ge e(G_r) \ge e(G) - r(t-2) \ge (t-1)n - (t^2 - t - 1) - r(t-2),$$

which implies $r \leq t^2 - t - 1$. Hence $|V(H)| \geq n - (t^2 - t - 1)$. From the procedure constructing $\{G_i\}_{i=1}^r, d_H(v) \leq t - 2$ for every vertex $v \in V(G) \setminus V(H)$.

Lemma 3.3. Let G_1 be a $\bigcup_{i=1}^k P_{2a_i}$ -free graph of order n_1 with $t = \sum_{i=1}^k a_i \ge 3$ and G_2 be a graph of order n_2 with $1 \le n_2 \le t^2 + t - 2$. If $G = K_1 \nabla(G_1 \bigcup G_2)$ is a graph of order n with $n = n_1 + n_2 + 1 \ge 8t^2 - 12t + 9$, then $q(G) < q(S_{n,t})$.

Proof. Without loss of generality, we assume that $G_2 = K_{n_2}$. Since G_1 is $\bigcup_{i=1}^k P_{2a_i}$ -free, we have G_1 is P_{2t} -free. By Lemma 2.1,

$$e(G_1) \le (t-1)(n-n_2-1)$$

Thus,

$$e(G - V(G_2)) = e(G_1) + n - n_2 - 1 \le t(n - n_2 - 1).$$

By [6, Theorem 3.1],

$$q(G - V(G_2)) \leq \frac{2e(G - V(G_2))}{|V(G - V(G_2))| - 1} + |V(G - V(G_2))| - 2$$

$$\leq \frac{2t(n - n_2 - 1)}{n - n_2 - 1} + n - n_2 - 2$$

$$\leq n + 2t - n_2 - 2.$$

Let $\mathbf{x} = (x_1, \ldots, x_n)^T$ be a unit positive eigenvector of Q(G) corresponding to the signless Laplacian spectral radius q(G). By symmetry, all components of x corresponding to the vertices of G_2 are the same, say a. By the eigenequations of Q(G) on u with the maximum degree and a vertex of G_2 , respectively, we have

$$(q(G) - n + 1)x_u = \sum_{v \in V(G) \setminus \{u\}} x_v \le \sqrt{(n-1)\sum_{v \in V(G) \setminus \{u\}} x_v^2} = \sqrt{(n-1)(1-x_u^2)},$$

$$(q(G) - n_2)a = (n_2 - 1)a + x_u.$$

Note that $q(G) \ge q(S_{n,t}) > n + 2t - 3$ and $n_2 \le t^2 + t - 2$, we have

$$x_u^2 \le \frac{n-1}{(q(G)-n+1)^2+n-1} \le \frac{n-1}{n-1+4(t-1)^2} < 1 - \frac{4(t-1)^2}{n+4t^2},$$

and

$$a = \frac{x_u}{q(G) - 2n_2 + 1} \le \frac{x_u}{n + 2t - 2n_2 - 2} \le \frac{x_u}{n - 2t^2 + 2}.$$

Then

$$\begin{split} q(G) &= \sum_{ij \in E(G)} (x_i + x_j)^2 = \sum_{ij \in E(G) \setminus E(G_2)} (x_i + x_j)^2 + n_2(a + x_u)^2 + \sum_{ij \in E(G_2)} (x_i + x_j)^2 \\ &< q(G - V(G_2)) + n_2(a + x_u)^2 + 2n_2(n_2 - 1)a^2 \\ &\leq n + 2t - n_2 - 2 + n_2 \left(1 + \frac{1}{(n - 2t^2 + 2)^2} + \frac{2}{n - 2t^2 + 2} + \frac{2(n_2 - 1)}{(n - 2t^2 + 2)^2}\right) x_u^2 \\ &< n + 2t - n_2 - 2 + n_2 \left(1 + \frac{3}{n - 2t^2 + 2}\right) \left(1 - \frac{4(t - 1)^2}{n + 4t^2}\right) \\ &< n + 2t - 2 - \left(\frac{4(t - 1)^2}{n + 4t^2} - \frac{3}{n - 2t^2 + 2}\right) \\ &< n + 2t - 2 - \frac{2t(t - 1)}{n - 2t^2 + 2} \\ &< n + 2t - 2 - \frac{2t(t - 1)}{n + 2t - 3} < q(S_{n,t}). \end{split}$$

This completes the proof.

Now we are ready to prove Theorem 1.7.

Proof of Theorem 1.7. Suppose that $q(G) \ge q(S_{n,t})$. We will show that $G = S_{n,t}$. By Lemma 3.1, $\Delta(G) = n - 1$. Let $u \in V(G)$ be the vertex with the maximum degree $\Delta(G)$, i.e., d(u) = n - 1. By Lemma 2.5 (2), $e(G) \ge tn - t^2 + 1$, which implies that

$$e(G-u) = e(G) - n + 1$$

$$\geq (t-1)n - t^2 + 2$$

= $(t-1)(n-1) - (t^2 - t - 1).$

Note that G-u is $\bigcup_{i=1}^{k} P_{2a_i}$ -free, otherwise G contains F_{a_1,\ldots,a_k} as a subgraph. By Lemma 3.2 , there exists an induced subgraph H of G-u such that $\delta(H) \ge t-1$, $|V(H)| = n_1 \ge n - (t^2 - t)$, and $d_H(v) \le t-2$ for every vertex $v \in V(G) \setminus (V(H) \bigcup \{u\})$. Let $H = \bigcup_{i=1}^{s} H_i$ and $|V(H_i)| = h_i$, where H_i is a component of H for $i = 1, \ldots, s$.

Claim 1. Every component of G - u contains at most one graph of H_1, \ldots, H_s as an induced subgraph.

Note that $\delta(H_i) \geq t-1 \geq 2$ for $i = 1, \ldots, s$. By Lemma 2.3, H_i contains a cycle of length at least t for $i = 1, \ldots, s$. In fact, if there is a component of G-u containing at least two graphs H_i and H_j as an induced subgraph for $1 \leq i \neq j \leq s$, then G-u contains P_{2t+1} as a subgraph and thus G contains F_{a_1,\ldots,a_k} as a subgraph, which is a contradiction. This proves Claim 1.

By Claim 1, we let T_i be the component of G - u containing H_i as an induced subgraph, and $G - u = (\bigcup_{i=1}^{s} T_i) \bigcup T_0$, where T_0 is the union of the remaining components of G - u. Note that G - u is $\bigcup_{i=1}^{k} P_{2a_i}$ -free, we have T_i is P_{2t} -free for $i = 0, \ldots, s$.

Claim 2. $T_0 = \emptyset$.

In fact, if $T_0 \neq \emptyset$, then

$$1 \le |V(T_0)| = n - 1 - \sum_{i=1}^{s} |V(T_i)| \le n - 1 - \sum_{i=1}^{s} |V(H_i)|$$
$$= n - 1 - n_1 \le t^2 - t - 1.$$

By Lemma 3.3, $q(G) < q(S_{n,t})$, which is a contradiction. This proves Claim 2.

Claim 3. $h_i \ge 2t$ for i = 1, ..., s.

Suppose there exist an h_i such that $h_i \leq 2t - 1$. Since $\delta(H_i) \geq \delta(H) \geq t - 1$, we have $h_i \geq t$. Then

$$t \le h_i \le |V(T_i)| \le h_i + |V(G - u) \setminus V(H)| \le 2t - 1 + t^2 - t - 1 = t^2 + t - 2.$$

By Lemma 3.3, $q(G) < q(S_{n,t})$, which is a contradiction. This proves Claim 3.

By the definition of $L_{r,t}$, we have the following claim directly.

Claim 4. For any fixed $1 \le i \le s$, if $H_i = L_{r_i,t-1}$ with $h_i = r_i(t-1) + 1$, then H_i contains P_{2t-1} as a subgraph.

Claim 5. For any fixed $1 \le i \le s$, if H_i is a subgraph of $S_{h_i,t-1}$, then H_i contains P_{2t-1} as a subgraph. Moreover, T_i is subgraph of $S_{|V(T_i)|,t-1}$.

If H_i is a subgraph of $S_{h_i,t-1}$, then there exists $I_i \subseteq V(H_i)$ of size $h_i - t + 1$ such that I_i induces an independent set of H_i . Since $\delta(H_i) \ge t-1$, every vertex in I_i is adjacent to every vertex in $V(H_i) \setminus I_i$. Then H_i contains a path P of order 2t-1 with both endvertices in I_i as a subgraph. If $(V(T_i) \setminus V(H_i)) \bigcup I_i$ induces at least an edge, then we can get a path of order 2t from P. Hence G contains F_{a_1,\ldots,a_k} as a subgraph, which is a contradiction. This implies that $(V(T_i) \setminus V(H_i)) \bigcup I_i$ is an independent set and thus T_i is subgraph of $S_{|V(T_i)|,t-1}$. This proves Claim 5.

Note that $q(G) \ge q(S_{n,t})$. Since H_i is $\bigcup_{i=1}^k P_{2a_i}$ -free and $\delta(H_i) \ge \delta(H) \ge t-1$, by Lemmas 2.4, H_i is a subgraph of $S_{h_i,t-1}$ or $H_i = L_{r_i,t-1}$ with $h_i = r_i(t-1) + 1$ and $k = 2, a_1 = a_2$ for $i = 1, \ldots, s$. If $s \ge 2$, then by Claims 3 and 4, G - u contains $2P_{2t-1}$ as a subgraph and thus G contains F_{a_1,\ldots,a_k} as a subgraph, which is a contradiction. So s = 1. This implies that

$$G-u=T_1$$

and H_1 is an induced graph of T_1 , where H_1 is a subgraph of $S_{h_1,t-1}$ or $H_1 = L_{r_1,t-1}$ with $h_1 = r_1(t-1) + 1$ and $k = 2, a_1 = a_2$.

First suppose that H_1 is a subgraph of $S_{h_1,t-1}$. By Claim 5, T_1 is a subgraph of $S_{|V(T_1)|,t-1}$, i.e., G is a subgraph of $S_{n,t-1}$. If G is a proper subgraph of $S_{n,t}$, then it follows from Perron-Fronbenius theorem that $q(G) < q(S_{n,t})$, which is also a contradiction. Hence $G = S_{n,t}$.

Next suppose that $H_1 = L_{r_1,t-1}$ with $h_1 = r_1(t-1) + 1$ and $k = 2, a_1 = a_2$. If $H_1 = T_1$, then $G = K_1 \nabla L_{r_1,t-1}$ with $n = r_1(t-1) + 2$. By Lemma 2.6, $q(G) < n + 2t - 3 < q(S_{n,t})$, which is a contradiction. Thus H_1 is a proper subgraph of T_1 . Let $H' = T_1 - V(H_1)$ and $|V(H')| = n_2$. Since $d_{H_1}(v) \le t - 2$ for every vertex $v \in V(H')$, we have

$$e(V(H'), V(H_1)) \le (t-2)n_2 \le (t-2)(t^2-t-1).$$

Then

$$\begin{split} e(H') &= e(T_1) - e(H_1) - e(V(H'), V(H_1)) \\ &> (t-1)n - t^2 + 2 - \frac{t(n_1-1)}{2} - (t-2)(t^2 - t - 1) \\ &= (t-1)n - t^2 + 2 - \frac{t(n-n_2-2)}{2} - (t-2)(t^2 - t - 1) \\ &= \frac{(t-2)n - (2t^3 - 4t^2)}{2} + \frac{tn_2}{2} > \frac{tn_2}{2}. \end{split}$$

By Lemma 2.1, H' contains P_{t+2} as a subgraph. Together with Claim 4, T_1 contains $P_{2t-1} \bigcup P_{t+2}$ as a subgraph. This implies that G contains F_{a_1,a_2} as a subgraph, which is a contradiction. This completes the proof.

References

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory, Springer, New York, 2007.
- [2] M.-Z. Chen, X.-D. Zhang, Erdős–Gallai stability theorem for linear forests, Discrete Math. 342 (2019) 904–916.
- [3] M.-Z. Chen, A.-M. Liu, X.-D. Zhang, Spectral extremal results with forbidding linear forests, Graphs Combin. 35 (2019) 335–351.
- [4] M.-Z. Chen, A-M. Liu, X.-D. Zhang. On the spectral radius of graphs without a star forest. Discrete Math. 344 (2021), 112269.
- [5] S. Cioabă, L. Feng, M. Tait, X.-D. Zhang, The maximum spectral radius of graphs without friendship subgraphs, Electron. J. Combin. 27 (4) (2020) P4.22.
- [6] K. Das, Maximizing the sum of the squares of the degrees of a graph, Discrete Math. 285 (2004) 57–66.
- [7] X. Duan, B. Zhou, Sharp bounds on the spectral radius of a nonnegative matrix, Linear Algebra Appl. 439 (2013) 2961–2970.
- [8] M.A.A. de Freitas, V. Nikiforov, L. Patuzzi, Maxima of the Q-index: graphs with no K_{s,t}, Linear Algebra Appl. 496 (2016) 381–391.
- [9] G.A. Dirac, Some theorems on abstract graphs, Proc. London Math. Soc. 2 (1952) 69-81.

- [10] P. Erdős, T. Gallai, On maximal paths and circuits of graphs. Acta Math. Acad. Sci. Hungar. 10 (1959) 337–356.
- [11] P. Erdős, Z. Füredi, R.J. Gould, D.S. Gunderson, Extremal graphs for intersecting triangles, J. Combin. Theory. Ser. B, 64 (1995) 89–100.
- [12] B. He, Y.-L. Jin, X.-D. Zhang, Sharp bounds for the signless Laplacian spectral radius in terms of clique number, Linear Algebra Appl. 438 (2013) 3851–3861.
- [13] X. Hou, Y. Qiu, B. Liu, Turán number and decomposition number of intersecting odd cycles, Discrete Math. 341 (2018) 126–137.
- [14] Y. Li, Y. Peng, The spectral radius of graphs with no intersecting odd cycles, arXiv:2106.00587v1.
- [15] R. Merris, A note on Laplacian graph eigenvalues, Linear Algebra Appl. 285 (1998) 33–35.
- [16] V. Nikiforov, Bounds on graph eigenvalues II, Linear Algebra Appl. 427 (2007) 183–189.
- [17] V. Nikiforov, The spectral radius of graphs without paths and cycles of specified length, Linear Algebra Appl. 432 (2010) 2243–2256.
- [18] V. Nikiforov, A contribution to the Zarankiewicz problem, Linear Algebra Appl. 432 (2010) 1405–1411.
- [19] V. Nikiforov, Some new results in extremal graph theory, Surveys in combinatorics 2011, 141–181, London Math. Soc. Lecture Note Ser., 392, Cambridge Univ. Press, Cambridge, 2011.
- [20] V. Nikiforov, X. Yuan, Maxima of the Q-index: graphs without long paths, Electron. J. Linear Algebra 27 (2014) 504–514.
- [21] V. Nikiforov, X. Yuan, Maxima of the Q-index: forbidden even cycles, Linear Algebra Appl. 471 (2015) 636–653.
- [22] M. Simonovits, A method for solving extremal problems in graph theory, stability problems, Theory of Graphs (Proc. Colloq., Tihany, 1966) (1968) 279-319.
- [23] L.-T. Yuan, Extremal graphs for the k-flower, J. Graph Theory 89 (2018) 26–39.
- [24] X. Yuan, Maxima of the Q-index: forbidden odd cycles, Linear Algebra Appl. 458 (2014) 207–216.
- [25] Y. Zhao, X. Huang, H. Guo, The signless Laplacian spectral radius of graphs with no intersecting triangles, Linear Algebra Appl. 618 (2021) 12–21.