

# Distributionally Robust Resource Planning Under Binomial Demand Intakes

Ben Black<sup>\*†</sup>, Russell Ainslie<sup>‡</sup>, Trivikram Dokka<sup>§</sup>, Christopher Kirkbride<sup>¶</sup>

January 7, 2023

## Abstract

In this paper, we consider a distributionally robust resource planning model inspired by a real-world service industry problem in which there is a mixture of known demand and uncertain future demand. Prior to having full knowledge of the demand, we must decide upon how many jobs we will complete on each day of the plan. Any jobs that are not completed by the end of their due date incur a cost and become due the following day. We present a distributionally robust optimisation (DRO) model where we treat the number of uncertain jobs due on each day as a binomial random variable with unknown parameters. We make use of theoretical properties of the binomial distribution to present a simple and fast algorithm for the two-day model. We present theoretical results regarding the near-optimality of this algorithm, namely that it finds the worst-case distribution for a subset of the solution space. For the multi-day model, we extend this algorithm and present two others, one based on the cutting surface algorithm commonly seen in the DRO literature, and one based on a probability-based reduction of the uncertainty set for the uncertain parameters. We test these algorithms on a number of different problem instances to establish their performance.

**Keywords:** Uncertainty modelling, distributionally robust optimisation, heuristics, resource planning.

## 1 Introduction

In this paper, we consider a resource planning problem motivated by a real-world service company. This real problem consists of optimising the use of a large workforce of service engineers, in the face of a mixture of known and uncertain jobs.

---

<sup>\*</sup>STOR-i Centre for Doctoral Training, Lancaster University, United Kingdom. Email: b.black1@lancaster.ac.uk

<sup>†</sup>Corresponding author.

<sup>‡</sup>Applied Research, BT Technology, Adastral Park, Ipswich, United Kingdom. Email: russell.ainslie@bt.com

<sup>§</sup>Management Section, Queens Management School, Queens University Belfast, United Kingdom. Email: T.Dokka@qub.ac.uk

<sup>¶</sup>Department of Management Science, Lancaster University Management School, United Kingdom. Email: c.kirkbride@lancaster.ac.uk.

## 1.1 Problem Setting

The planning process for a service company is subject to three stages, named *the three stages of planning*. Each serves a different purpose, covers a different time horizon, and creates results that feed into the next. The three stages are *strategic*, *tactical*, and *operational* planning, and the way that they interact is shown in Figure 1.



Figure 1: The three stages of planning

Strategic planning covers a period of multiple years, and concerns long term decisions such as how many employees to be hired and in which skills they should be trained. Tactical planning concerns a period of weeks or months, and involves aggregate decisions such as deciding upon the capacity needed in each period, or how many jobs can and cannot be completed in each period. Operational planning concerns short-term decisions such as scheduling the day-to-day activities of the workforce at the individual level. We focus on the tactical planning stage in this paper. The decisions that we make are at the aggregate level, i.e. we do not plan the specific activities of every worker but we instead aggregate their availability into a daily capacity value. Since it is typically not possible to move capacity between days, planners manipulate demand to make the best use of what they have.

The aggregate capacity gives the number of jobs that our workforce can complete, for each day in a period of length  $L$ . At the time of making the plan, each day also has a number of jobs that are due on said day and are known about in advance. As well as this, there are an uncertain number of jobs that will arrive in the future to be due on each day, referred to as *intake* jobs. By default, we use all available capacity on each day to complete jobs on that day. Jobs that are in the system at the time of planning can be completed before their due date, and this is referred to as *pulling forward*. If any are still incomplete by the end of their due date then they will not leave the system but incur a cost, and become due on the following day. This is referred to as *rollover*. Pulling forward is employed in order to complete jobs that would otherwise roll over. With pulling forward as our decision variable, our baseline model minimises the total rollover cost over all days in the plan. The output of the model gives the day on which the jobs should be completed in order to achieve the minimal rollover cost.

More formally, the problem can be defined as follows. Given a planning horizon of  $L$  days, on each day  $\tau \in [L] = \{1, \dots, L\}$ , there are  $D_\tau$  jobs due on  $\tau$  that are known about in advance, and there is an overall processing capacity of  $c_\tau$ . The set of known jobs is referred to as the *workstack*. Apart from the known jobs, on each day an unknown number of intake jobs arrive in the system. The number of intake jobs that will be due on day  $\tau$  is  $I_\tau$ . Any job not completed by its due date, say  $\tau$ , will roll over to the next day and incur a cost of  $a_\tau$ . Any job can incur multiple rollover costs. All jobs are assumed to require one unit of capacity to complete. The task is to efficiently manage the overall capacity to avoid unnecessary rollover cost by pulling

some jobs forward to complete them before their due date. Since intake jobs are not known about *a priori*, we assume that only workstack jobs can be pulled forward. In the deterministic, full-information case, that is, when  $I_\tau$  is known *a priori*, the problem reduces to simple MIP that can be solved easily using off-the-shelf solvers.

The most important quality of our models is that they account for the uncertainty in the intakes, that is, we treat  $I_\tau$ ,  $\tau \in [L]$ , as unknown. In practice, the number of additional jobs is predicted using a forecasting model, and this value is often taken as the truth. In this paper, we treat the intakes as random variables and directly model the uncertainty in both their values and their distributions. More specifically, we model intakes as binomial random variables where  $I_\tau$  is parameterised by the number of trials,  $i_\tau^{\max}$ , and success probability  $p_\tau$ . Furthermore, we assume that the intake random variables for any two days in the plan are independent of one another. In many practical applications, the forecasting model used will give a range of potential values for  $I_\tau$  alongside a point estimate that falls somewhere in this range. This means that we are given  $i_\tau^{\max}$  prior to building the model. Therefore,  $p_\tau$  is the only unknown parameter for each  $\tau$ . Our decision to use the binomial distribution can be justified by the following three reasons:

1. The number of intake jobs due on each day is a discrete quantity and any two jobs arriving on the same day arrive independently of one another.
2. There is a fixed and finite set of values that each intake random variable can take. Other discrete distributions such as the Poisson distribution are unbounded, and hence not fitting for these random variables.
3. Apart from naturality, it gives a concise way of modelling the uncertainty. We can represent each distribution uniquely by one choice of  $p$ , which is a vector of length  $L$ . Using a non-parametric approach would mean having to analyse the entire distribution, which is a larger vector that has one entry for every realisation of intake.

We emphasize that the binomial assumption is in contrast with much of the DRO literature, in which assumptions are often non-parametric. The reason for this is that parametric distributions, such as the binomial, often lead to intractability. However, in the context of our problem we show that it is possible to derive algorithms which are both tractable and close to optimal.

Similar problems have been studied in the literature, although these papers either do not treat demand as uncertain or do not treat the distribution of demand as uncertain. Ainslie et al. (2015) presented a linear programming model that considers a multi-skilled workforce, and allocates the available capacity to the different skills in order to maximise completions. In this paper, demand was considered deterministic. Ross (2016) used two-stage stochastic programming models for a similar problem, for which time series models and copulas were used to generate demand scenarios. In stochastic programming, however, the distribution of the uncertain parameters is assumed known, and this is very unlikely to be the case in reality. These models also do not account for due dates, and instead model demand as the number of jobs that exist on a given day. Modelling the due dates of jobs not only allows for an understanding of which jobs can be

completed on which days but also a better representation of which jobs will roll over.

The framework that we use to model our problem is Distributionally Robust Optimisation (DRO). This framework combines elements from robust optimisation and stochastic programming, and avoids most of their pitfalls. It is similar to robust optimisation in that we seek a worst-case solution, but we do not assume that the worst-case realisation always occurs. Hence, DRO is less conservative than robust optimisation. It is also similar to stochastic programming due to its minimisation of expected costs, but we do not assume that the distribution of unknown parameters is known. This leads to a much more realistic modelling framework, that can incorporate the outputs of multiple experts or forecasting models via a set of distributions. Solutions to DRO models are therefore more robust than stochastic programming solutions and also less conservative than robust optimisation solutions.

## 1.2 Modelling Framework

DRO is a field concerning optimisation under uncertainty, in the values of some of the model's parameters and also in the probability distribution of said parameters. In order to introduce DRO, it is key to understand robust optimisation (RO) first. In RO, we have a model where some of the parameters, say  $i = (i_1, \dots, i_L)$ , are unknown but it is known or assumed to be known that they must lie in some *uncertainty set*  $\mathcal{I}$ . If the vector of decision variables is denoted  $y$ , then an RO model is one that aims to minimise the worst-case cost, over all possible realisations of the uncertain parameters. A general RO model is shown in expression (1).

$$\min_{y \in \mathcal{Y}} \max_{i \in \mathcal{I}} h(y, i) \quad (1)$$

As described by Gorissen et al. (2015), RO models can easily be reformulated to linear programs, although the robustness results a model with significantly more constraints than its deterministic counterpart. RO models are known to be over-conservative, since the worst-case values of the parameters are extremely unlikely to occur. Hence, in DRO, instead of treating the unknown parameters as constant but unknown, we consider them as random variables that behave according to some probability distribution. We do not know the distribution of said parameters, but we assume it is contained in some *ambiguity set*  $\mathcal{U}$ . Hence, let us represent the uncertain parameters as a vector  $I = (I_1, \dots, I_L)$  of random variables, which has realisations  $i \in \mathcal{I}$ . The objective is now to find the minimal worst-case *expected* cost:

$$\min_{y \in \mathcal{Y}} \max_{q \in \mathcal{U}} \mathbb{E}_q[h(y, I)]. \quad (2)$$

The ambiguity set  $\mathcal{U}$  can be defined in a number of ways, although the most common is to use all distributions that can be considered *close* to some nominal distribution using some measure such as a  $\phi$ -divergence (Bayraksan and Love, 2015). This allows for the requirement  $q \in \mathcal{U}$  to be represented as a constraint of the inner problem where the distribution is treated as a decision variable, and this inner problem can then be dualised and solved. This method applies when the set  $\mathcal{U}$  is continuous, and is not so easily used when it is discrete.

In this paper, we consider the random parameters to be binomial random variables and the set of distributions to be defined by a discrete set of binomial success probabilities. We derive a

number of algorithms for approximately solving the resulting DRO model and illustrate their performance by comparing to optimal solutions.

### 1.3 Our Contributions

We consider a DRO model for a resource planning problem with an unknown number of intake jobs on each day. Using the problem structure, we model intakes as binomial random variables and study the resulting DRO model. Due to the uncertainty in the number of intake jobs, the problem is considerably harder from a computational point of view. Our contributions in the paper include the following:

1. An almost-optimal algorithm, *Squash and Search (S&S)*, to solve the two-day problem. The algorithm uses properties of the problem itself and of the binomial distribution to efficiently search within the feasible space of the uni-dimensional pull-forward decision range in this case, and also to find the worst-case distribution for each pull-forward solution. See Section 3.5 for more details.
2. A non-trivial extension of S&S, to the multi-day problem, detailed in Section 4.3.
3. Two solver-based algorithms: a Cutting Surface (CS) algorithm and an Approximate Objective (AO) algorithm (see Sections 4.4.1 and 4.4.2). Our CS algorithm, while not exact, considerably simplifies the main bottleneck step of other CS algorithms: the distribution separation problem. This makes it much more scalable.
4. Extensive computational experiments on a variety of constructed instances which show the efficacy of our methods. See Section 6 for these results.

## 2 Literature Review

In this section we review relevant literature relating to our problem and problems of a similar nature. In Section 2.1, we review the workforce planning literature and highlight the methodologies used there. In Section 2.2, we summarise the recent DRO literature and discuss how our research differs from it.

### 2.1 Workforce Planning

This research is motivated by a workforce planning problem, a service industry one in particular. Mathematical and stochastic programming have been very common in workforce planning since very early in operational research's history. One such application of mathematical programming in workforce scheduling comes from Holt et al. (1955), who formulated a model that decided upon the optimal staffing of a workforce and their production rate over a number of periods. This model is a non-linear program (NLP) that minimises expected cost of staffing, layoff and overtime subject to inventory constraints. At the time, however, most practical applications used cost functions that were assumed linear. Hence, this model was subsequently extended by Hanssmann and Hess (1960), who formulated the same problem as a linear program (LP).

This was one of the first general models for workforce planning that can be found in the literature, and laid the foundation for later extensions. For example, Fetter (1961) extended the model of Hanssmann and Hess (1960) by including probabilistic demand and studying such uncertain changes to the model as those in ownership of the company, in cost of work hours, etc. Since these papers, it has generally been the case that LP has been appropriate for planning problems concerning minimising cost as a single objective.

Uncertain elements have also been considered, with the most common way to represent uncertainty being through stochastic programming models. Early examples used two- and three-stage stochastic programs to account for demand uncertainty. For example, J. Abernathy et al. (1973) studied a three-stage stochastic model that both plans and schedules a workforce, motivated by the nurse-scheduling problem. Most models making use of stochastic programming are, however, two-stage models, since strategic planning is normally not formulated as an optimisation problem but tactical and operational planning are. An early two-stage model came from Martel and Price (1978), who studied a model for recruiting and promoting a military organisation. Through showing that their model could be reformulated as an LP, these authors showed that modelling uncertainty can be computationally efficient as well as realistic.

With increased computing power and algorithmic developments, more recent research has aimed at representing demand uncertainty in planning models. Zhu and Sherali (2009) consider a 2-stage workforce planning model under both uncertain and fluctuating demand, where in the first stage workforce allocation is carried out and then in the second stage the demand is smoothed among the workers. They use a Bender’s decomposition algorithm to solve their model. Bastian et al. (2020) use both stochastic programming and robust optimisation for planning the cyber branch of the US army. Ross (2016) used a two-stage workforce planning model that incorporated multi-skilled workers and how they are cross-trained. Demand uncertainties were modelled using bivariate copula sampling. More generally, robust optimisation is less common in workforce planning, due to its conservative nature, but it is more commonly used in planning for disasters. Sun et al. (2021) use a robust model in allocating geographical resource in disaster response planning. The authors acknowledged that they were making a trade-off between robustness and optimality.

Modelling the uncertainty in demand can make models incredibly difficult to solve, especially when they are based on workforces as large as those used in the service industry. A common approach in workforce planning is to use a forecast of demand and assume some level of trust in the forecast, then treat the plan as more of a prediction of what will happen. Ainslie et al. (2015) consider the problem of planning the activity of a multi-skilled workforce by matching skill hours to demand hours. Combinatorially this model is already very large, making the modelling of uncertainty difficult. Ainslie et al. (2018) use a bi-level optimisation model under assumed demand in order to set capacity levels and pull “planning levers” such as overtime and contractor hours to meet more demand. The bi-level model was solved using a genetic algorithm due to the size of the problem. One recent workforce planning paper that does consider uncertain demands was given by Angalakudati et al. (2014). In their work, they consider planning the activities of

the workforce of a gas utility company, who complete a mixture of known and unknown jobs. The unknown jobs are emergency jobs that are only known about when they arrive, and must be dealt with immediately. These jobs are treated as random, as in our paper, but in this problem the workforce do not control when jobs are completed but instead manipulate capacity to reduce the overtime required to complete all jobs.

This recent work is among the growing evidence that the focus of workforce planning papers has shifted from simple LPs to large, complex and often stochastic models that aim to utilise a large, multi-skilled workforce under uncertain demand. Typically the larger and more complex models will not encompass demand uncertainty, due to the longer solve times and lack of well-known solution methods. Our algorithms are designed to be used on complex workforce planning models but to also be scalable so that problem size does not have an effect on them.

## 2.2 Distributionally Robust Optimisation

DRO is a combination of ideas from stochastic programming and robust optimisation, nicely summarised in a recent review by Rahimian and Mehrotra (2019). Typically, DRO models lead to semi-infinite formulations due to variables that encode probability distributions. The algorithmic research on dealing with such formulations is still in its infancy. One recently proposed method is the cutting surface (CS) algorithm (Mehrotra and Papp, 2014). Using similar ideas we propose a version of CS to solve our problem. The CS algorithm has some similarities with the L-shaped decomposition algorithm for stochastic programs (Laporte and Louveaux, 1993). The L-shaped algorithm is a branch and cut algorithm that iteratively adds scenarios to the set in a two-stage program by solving an optimisation model to choose which to add. The CS algorithm behaves similarly, but it adds a distribution to the ambiguity set at each iteration. The CS algorithm was presented for general semi-infinite convex optimization problems (SCP), not necessarily DRO models, by Kortanek and No (1993) and Hettich and Kortanek (1993).

The SCP algorithm was adapted for a DRO model relating to portfolio selection by Pflug and Wozabal (2007), who used it as a heuristic without claiming optimality. The algorithm was again used in DRO by Mehrotra and Papp (2014), this time rebranded as the CS algorithm, who drew insights about the algorithm's relation to risk aversion and stochastic programming. They also showed that if it converges, then it converges to an  $\varepsilon$ -optimal solution. Luo and Mehrotra (2019) later used it for a class of DRO problems where the set of distributions is defined using a Wasserstein distance. For these problems, the algorithm was shown to be  $\varepsilon$ -optimal. CS was also used for identifying scenarios in DRO models by Rahimian et al. (2019). Bansal et al. (2018) compares CS with a new version of the L-shaped decomposition algorithm for DRO models and found that, more than optimality, computing times were the cause of difficulty. CS can face run-time issues because of its need to solve a distribution separation problem at every iteration, an issue we avoid in this paper by applying a heuristic procedure instead of solving the distribution separation problem to optimality.

The CS algorithm does not make any assumptions about the method used to define the ambiguity

set and does not make use of dualisation of the inner problem. More commonly in the literature, the models are solved using the so-called Dual method. This method consists of dualising the inner model and optimising this, which can lead to optimal primal solutions under some conditions. See, for example, Delage and Ye (2010) for a model under moment uncertainty as well as distributional uncertainty, or Ben-Tal et al. (2013) for cases where ambiguity sets are defined using  $\phi$ -divergences. Using  $\phi$ -divergences, the choice of distribution can be represented by decision variables with constraints defined by the divergence. The inner model can often then be dualised and made easier to solve.

Iterative methods using dualisation and  $\phi$ -divergences are also common, due to the difficulty of solving the dual directly. For example, Liu et al. (2017) studied DRO models solved using discretisation and they used a so-called Primal Dual Hybrid Gradient method. Other iterative methods include that of Lam and Ghosh (2013), who worked with the problem of estimating the value of a performance metric. Their algorithm is designed for DRO models where the set of distributions is defined by a  $\chi^2$ -divergence in particular. Ghosh et al. (2020) also apply a gradient descent to the outer problem and use sample average approximation for the inner problem. They work with a general  $\phi$ -divergence, and use duality results for the inner problem. A tutorial on using  $\phi$ -divergences for DRO has been written by Bayraksan and Love (2015), who link the various types of divergence to different attitudes to risk and discuss ways to select a divergence measure.

Our methodology consists of two differing conditions when compared with most papers in the DRO literature. The first and most important one is that we make an additional assumption about the family of distributions in which our uncertain distribution lies. This allows us to incorporate some prior knowledge about the behaviour of the uncertain parameters and also gives us some theoretical properties that can be used to develop algorithms. It also allows us to restrict our search from a space with the same dimension as the number of realisations of the uncertain parameters to a much smaller one, with the same dimension as the vector of parameters of our multivariate distribution. It does, however, mean that our ambiguity set cannot easily be defined by constraints. The second is that we consider a discretisation of the set of distributions, which adds another level of computational complexity to the models.

### 3 A Two-day Planning Model

In this section, we detail a two-day DRO model for resource planning under uncertain demand. This model forms the basis for the remainder of the paper and gives the justification for the algorithms we create for the full model. We first detail the notation in Section 3.1 and the model in Section 3.2. Following this, we detail our reformulation of the model and the ambiguity sets used. We then present our near-optimal solution algorithm, Squash and Search (S&S), in Section 3.5.



### 3.1 Notation

Basic notation for capacity, intakes and workstacks was defined in Section 1.1. We further assume that the intakes  $I_1$  and  $I_2$  are independent of one another. Each  $I_\tau$  can be considered as a random variable, with realisations  $i_\tau \in \mathcal{I}_\tau$ , where  $\mathcal{I}_\tau = \{0, 1, \dots, i_\tau^{\max}\}$ . The number of jobs to pull forward from day 2 to day 1 is denoted  $y$ , and we can only pull forward jobs that are known in the system and due on day 2. The rollover on day  $\tau$  is the number of jobs due on  $\tau$  that will be incomplete at the end of  $\tau$ , is denoted by  $R_\tau$ , and is a random variable. The vector  $R = (R_1, R_2)$  has realisations  $R^i = (R_1^i, R_2^i)$ , corresponding to intake realisation  $i$  for each  $i \in \mathcal{I}$ . We define the realisations in this way (i.e. both values indexed by  $i$ ) because  $R_2^i$  depends on both  $i_1$  and  $i_2$ , and so it cannot be defined based only on  $i_2$ . The day 1 rollover can be defined based only on  $i_1$ , but for consistency and interpretability we keep our notation the same for both days. The cost of leaving one job incomplete at the end of day  $\tau$  that was due on  $\tau$  is  $a_\tau$ . Given this, a deterministic version of our problem is to choose the pulling forward variable  $y$  so as to minimise the total rollover cost. When considering a stochastic problem, we will use the acronym PMF to mean probability mass function (for discrete random variables), PDF to mean probability density function (for continuous random variables) and CDF to mean cumulative distribution functions.

### 3.2 Distributionally Robust Model

The model we describe here is *distributionally robust* (DR). We assume that there is a discrete set  $\mathcal{U}$  of bivariate distributions  $q$  over the set of possible intakes  $\mathcal{I} = \mathcal{I}_1 \times \mathcal{I}_2$ . By independence of the intakes, each distribution can be written as  $q(i) = q_1(i_1) \times q_2(i_2)$  for all  $i \in \mathcal{I}$ . To characterise the worst-case distribution, we define it to be the one with the worst-case *expected rollover* cost. Hence, our DR model is given by:

$$\min_{y, R} \max_{q \in \mathcal{U}} \sum_{\tau=1}^2 a_\tau \mathbb{E}_q(R_\tau) \quad (3)$$

s.t.

$$R_1^i \geq i_1 - (c_1 - D_1 - y) \quad \forall i \in \mathcal{I}, \quad (4)$$

$$R_2^i \geq R_1^i + i_2 - (c_2 - D_2 + y) \quad \forall i \in \mathcal{I}, \quad (5)$$

$$y \leq \min\{c_1 - D_1, D_2\}, \quad (6)$$

$$y \in \mathbb{N}, \quad (7)$$

$$R_1^i, R_2^i \geq 0 \quad \forall i \in \mathcal{I}. \quad (8)$$

Constraint (4) calculates the day 1 rollover by subtracting from the intake  $i_1$  the remaining capacity after completing the workstack,  $D_1$ , and all jobs pulled forward to day 1,  $y$ . Constraint (5) defines the day 2 rollover by taking the day 2 intake  $i_2$  and adding  $R_1^i$  to get the number of additional jobs requiring completion, and subtracting the remaining capacity after completing the workstack  $D_2$  after reducing its size by  $y$ .

### 3.3 Tractable Reformulation

Due to the non-linearity of most PMFs and PDFs with respect to the distribution's parameters, we cannot solve this model treating them as decision variables. Hence, we instead consider the following (equivalent) model when solving to optimality. The model is the same as the above one, but with objective:

$$\min_{y, R, t} t \quad (9)$$

and an additional constraint:

$$t \geq \sum_{\tau=1}^2 a_{\tau} \mathbb{E}_q(R_{\tau}) \quad \forall q \in \mathcal{U}. \quad (10)$$

At optimality,  $t = \max_{q \in \mathcal{U}} \sum_{\tau=1}^2 a_{\tau} \mathbb{E}_q(R_{\tau})$  is guaranteed and we can extract the optimal distribution from this.

### 3.4 Binomial Intakes

The intakes are two independent and discrete random variables. Jobs arrive independently of one another. There is a fixed, bounded set of intake jobs that can arrive. Hence, we treat the intakes as binomial random variables, where we think of  $i_{\tau}^{\max}$  as the number of trials. In other words, we assume that:

$$I_{\tau} \sim \text{Bin}(i_{\tau}^{\max}, p_{\tau}).$$

Since the number of trials  $i_{\tau}^{\max}$  is fixed and equal to  $\max(\mathcal{I}_{\tau})$ , finding the worst-case distribution reduces to finding the worst-case success probabilities  $p = (p_1, p_2)$ . Let  $\mathcal{U}^p$  be a discrete set of probability vectors  $p$ , where each  $p$  uniquely defines a bivariate binomial distribution. Specifically, we consider ambiguity sets of the form:

$$\mathcal{U}_{\alpha}^p = \{p \in \mathcal{U}^p : \mathbb{P}(I_1 + I_2 > (c_1 - D_1) + (c_2 - D_2); p) \leq \alpha\}. \quad (11)$$

In our analysis, we aim to ensure that  $I_1 + I_2 \leq (c_1 - D_1) + (c_2 - D_2)$  because if this is not true then some jobs will always roll over, regardless of the decision that we make. However, since  $\alpha = 0$  would theoretically leave no distributions if  $i_1^{\max} + i_2^{\max} \geq (c_1 - D_1) + (c_2 - D_2)$ , we take  $\alpha \approx 0$ . In fact, we take  $\alpha$  to be so small that any distribution in  $\mathcal{U}_{\alpha}^p$  gives a probability of total intake exceeding total residual capacity that is treated as zero by any modern software environment such as Python. This allows some distributions to be left in the set, whilst still ensuring that any such distribution will not allow the intake to go above the remaining capacity. In a slight abuse of notation, we refer to this set as  $\mathcal{U}_0^p$  and treat  $\mathbb{P}(I_1 + I_2 > (c_1 - D_1) + (c_2 - D_2); p)$  as 0 for all  $p$  in the set. This is reasonable because, in any computations done on these distributions, this probability will be treated as zero. Equation (11) ensures that overly conservative possibilities such as  $p = (1, 1)$  are excluded, hence leads to a more realistic setting where there is a trade-off between allocating probability to day 1 and to day 2.

### 3.5 S&S: A Fast Solution Algorithm

We now propose an algorithm for solving the DR model with an uncertainty set of the form given in (11). The algorithm is based on the following principles:

1. We will not pull forward less than the minimum spare capacity on day 1. Pulling forward less than this would always result in more unused capacity on day 1, and can never reduce the total rollover. Hence, there is no reason to do so.
2. We will not pull forward more than the maximum spare capacity on day 1, as any jobs on top of this will roll over into day 2.
3. We will not pull forward more than the maximum potential rollover on day 2, as doing so would mean that we are pulling forward jobs do not need to be pulled forward.
4. Given the uncertainty set in (11), the expected rollover cost can be maximised for a fixed  $y$  using the following steps. Firstly, allocate maximal probability to one of the two days. Then, allocate the other day the maximal probability that it can be given, conditional on the probability already allocated.

Furthermore, we consider cases where  $D_2 > \max\{c_1 - D_1 - i_1^{\max}, 0\}$  and  $\max\{c_1 - D_1 - i_1^{\max}, 0\} < \max\{i_2^{\max} - (c_2 - D_2), 0\}$ . This is because if either of these are not true, then the model has a trivial optimal solution. In fact, if the first inequality does not hold then all  $D_2$  jobs can be moved to day 1 without causing day 1 rollover. Since no more can be pulled forward and any less would not decrease day 2 rollover, this is always optimal. If the second inequality does not hold, then all incomplete jobs on day 2 can be moved to day 1 without causing day 1 rollover. Again, this is always optimal. See Appendix A for a formal presentation of these results.

The algorithm will be referred to as Squash and Search (S&S), because it simply reduces the search space for  $y$  and  $p$  by eliminating values that will not be chosen, then searches what remains. The algorithm is given below.

1. Find the two sets:

$$c_{\text{spare}} = \{\max\{c_1 - D_1 - i_1, 0\} \mid i_1 \in \mathcal{I}_1\} \quad (12)$$

$$d_{\text{inc}} = \{\max\{i_2 - (c_2 - D_2), 0\} \mid i_2 \in \mathcal{I}_2\}. \quad (13)$$

Here,  $c_{\text{spare}}$  is the set of possible realisations of residual capacity on day 1 and  $d_{\text{inc}}$  is that of the number of incomplete jobs on day 2, both before pulling forward is done.

2. Find the range  $\mathcal{Y}$  for  $y$ :

$$y_{\min} = \min(c_{\text{spare}}), \quad (14)$$

$$y_{\max} = \min\{\max(c_{\text{spare}}), \max(d_{\text{inc}})\}, \quad (15)$$

$$\mathcal{Y} = \{y_{\min}, \dots, y_{\max}\}. \quad (16)$$

3. Find  $p_{\tau}^{\max} = \max_{p \in \mathcal{U}} p_{\tau}$  for each  $\tau$  and find

$$\mathcal{P}_{\max} = \{p \in \mathcal{U}^p : p_{\tau} = p_{\tau}^{\max} \text{ for some } \tau \in \{1, 2\}\},$$

$$\mathcal{U}_{\max}^p = \mathcal{P}_{\max} \cap \operatorname{argmax}_p \left\{ \sum_{\tau=1}^2 p_{\tau} \right\}.$$

4. For  $y \in \mathcal{Y} = \{y_{\min}, \dots, y_{\max}\}$ :
  - (a) For  $p \in \mathcal{U}_{\max}^p$  calculate  $C_p = \sum_{\tau=1}^2 a_{\tau} \mathbb{E}_p(R_{\tau})$ .
  - (b) Choose  $p_y$  such that  $C_{p_y} = \max_{p \in \mathcal{U}_{\max}^p} (C_p)$ .
5. Choose  $y^*$  to minimise maximum cost:

$$y^* = \underset{y \in \mathcal{Y}}{\operatorname{argmin}} C_{p_y}. \quad (17)$$

To clarify step 3, we are first finding the set of success probabilities where at least one day is given the largest probability it can be given,  $\mathcal{P}_{\max}$ . In order to extract the most extreme distributions from this set, we look at those distributions among those in the set where the second probability is highest. This is represented here by taking the sum over the two days. Note that  $\mathcal{U}^p$  is used here because the algorithm will work with any ambiguity set for  $p$ .

### 3.6 Theoretical Results on the Near-optimality of S&S

To recap, the reasoning behind the  $y$  choices in S&S is that an optimal  $y$  must lie in the range  $\mathcal{Y}$ , otherwise we are cannot be reducing rollover. The reasoning behind the set  $\mathcal{U}_{\max}^p$  is that a higher success probability should lead to higher expected rollover. In this section, we present some formal statements of these principles and we give the proofs for them in Appendix B. Lemma 1 formally establishes that the optimal  $y$  must lie in the set  $\mathcal{Y}$  searched by S&S. Lemma 2 is a theoretical statement of our justification for using  $\mathcal{U}_{\max}^p$ , which states that the expected rollover is increasing in the success probabilities. Despite our belief that this holds for all  $y \in \mathcal{Y}$ , we have proven the statement for a subset of  $\mathcal{Y}$ . We also have Theorem 1, which establishes optimality of S&S when selecting  $p$  for each fixed  $y$  in a subset of  $\mathcal{Y}$ .

**Lemma 1.** *The optimal  $y$  decision for the two-day DRO model falls in the set:*

$$\mathcal{Y} = \{\min(c_{\text{spare}}), \dots, \min\{\max(c_{\text{spare}}), \max(d_{\text{inc}})\}\}. \quad (18)$$

**Lemma 2.** 1.  $\mathbb{E}_p(R_1)$  is increasing in  $p_1$  for any integer  $y$  solution such that:

$$y \geq (c_1 - D_1) - \left( i_1^{\max} - \frac{\sqrt{i_1^{\max}}}{2} \right)$$

2. For any fixed  $p_1$ ,  $\mathbb{E}_p(R_2)$  is increasing in  $p_2$  for any integer  $y$  solution such that:

$$y \leq (D_2 - c_2) + \left( i_2^{\max} - \frac{\sqrt{i_2^{\max}}}{2} \right)$$

3.  $\mathbb{E}_p(R_1)$  is independent of  $p_2$  and  $\mathbb{E}_p(R_2)$  is independent of  $p_1$ .

**Theorem 1.** *For any  $y$  solution such that:*

$$y \in \tilde{\mathcal{Y}} = \left\{ (c_1 - D_1) - i_1^{\max} + \frac{\sqrt{i_1^{\max}}}{2}, \dots, (D_2 - c_2) + i_2^{\max} - \frac{\sqrt{i_2^{\max}}}{2} \right\},$$

*S&S finds the worst-case distribution and hence the worst-case expected cost.*

Since the set  $\tilde{\mathcal{Y}}$  given in Theorem 1 is entirely contained in  $\mathcal{Y}$ , S&S finds the worst-case distribution for each  $y$  in the interval. Therefore, if the optimal solution lies in this interval then we have proven that S&S finds the minimum worst-case expected cost, but not necessarily that it correctly identifies this solution as optimal. This would follow if both of the expected rollovers were increasing for  $y \in \mathcal{Y} \setminus \tilde{\mathcal{Y}}$ . The result in Lemma 2 proves that, for any  $y \in \mathcal{Y} \setminus \tilde{\mathcal{Y}}$ , at least one of the expected rollovers are increasing, but not that both are. This guarantees that the optimal  $p$  is in  $\mathcal{P}_{\max}$  but not that it is in  $\mathcal{U}_{\max}^p$ . Note that the range of  $y$  that is not included Lemma 1 is indeed very narrow, even more so when  $i_{\tau}^{\max}$  are large.

In order to test S&S, we ran it on 23,000 problem instances based on different input parameters. We used a variety of workstacks, capacities and rollover costs. The uncertainty set was always  $\mathcal{U}_{\alpha}^p$  where  $\alpha \approx 0$ , so that the total intake does not exceed the total remaining capacity. The DRO model was solved using the CPLEX Python API and the S&S algorithm, and the results compared. We found that in every single case, S&S reached an optimal solution in an average time of 0.15 seconds. CPLEX took 1.74 seconds on average to find an optimal solution. In fact, the optimality gap tolerance for CPLEX had to be lowered in order for CPLEX to arrive at solutions of the same quality as S&S found. These results suggest that this algorithm is not only optimal but also efficient.

## 4 An $L$ -day Planning Model

In this section, we detail our extension of the two-day planning model to the general  $L$ -day case. This adds complexity to the pulling forward decisions, since there are more pairs of days between which pulling forward is possible, and also to the rollover, as a result of the more complex pulling forward decisions. Specifically, we assume that a job can be pulled forward a maximum of  $K$  days, where  $K$  is some positive integer. We make this assumption because, in practice, jobs cannot be pulled forward more than a few days due to such things as inconvenience to the customer and disruption to the resources' schedules.

### 4.1 Distributionally Robust Model

We now consider the  $L$ -day DRO model, which is a direct extension of the model from Section 3.2. We assume that we have a discrete set  $\mathcal{U}$  of  $L$ -dimensional distributions  $q$  over the set of intakes  $\mathcal{I} = \mathcal{I}_1 \times \mathcal{I}_2 \times \dots \times \mathcal{I}_L$ , where  $\mathcal{I}_{\tau} = \{0, 1, \dots, i_{\tau}^{\max}\}$  as before. We continue to assume that the intake random variables for different days are independent. We now have a set of decision variables for pulling forward, due to the larger number of days between which pulling forward is possible. We define  $y_{\tau_1, \tau_2}$  as the number of jobs due on day  $\tau_1 \in \{2, \dots, L\}$  to complete on  $\tau_2 \in \{\tau_1 - K, \dots, \tau_1 - 1\}$ . Tables of notation for this section are shown in Appendix F. Our objective is, again, to minimise the worst-case expected rollover cost, but now the sum is taken over  $L$  days. Our constraints need to be changed to account for the new length of the plan. The model is given by:

$$\min_{y, R} \max_{q \in \mathcal{U}} \sum_{\tau=1}^L a_{\tau} \mathbb{E}_q(R_{\tau}). \quad (19)$$

s.t.

$$\sum_{\tau_2=\max\{\tau_1-K,1\}}^{\tau_1-1} y_{\tau_1,\tau_2} \leq \max\{D_{\tau_1} - c_{\tau_1}, 0\} \quad \forall \tau_1 = 2, \dots, L, \quad (20)$$

$$\sum_{\tau_1=\tau_2+1}^{\min\{\tau_2+K,L\}} y_{\tau_1,\tau_2} \leq \max\{c_{\tau_2} - D_{\tau_2}, 0\} \quad \forall \tau_2 = 1, \dots, L-1, \quad (21)$$

$$R_1^i \geq i_1 + \sum_{\tau_1=2}^{\min\{1+K,L\}} y_{\tau_1,1} - (c_1 - D_1) \quad \forall i \in \mathcal{I}. \quad (22)$$

$$R_\tau^i \geq R_{\tau-1}^i + i_\tau + \sum_{\tau_1=\tau+1}^{\min\{\tau+K,L\}} y_{\tau_1,\tau} - \left( c_\tau - D_\tau + \sum_{\tau_2=\max\{\tau-K,1\}}^{\tau-1} y_{\tau,\tau_2} \right) \quad \forall \tau = 2, \dots, L-1 \quad \forall i \in \mathcal{I}, \quad (23)$$

$$R_L^i \geq R_{L-1}^i + i_L - \left( c_L - D_L + \sum_{\tau_2=\max\{L-K,1\}}^{L-1} y_{\tau,\tau_2} \right) \quad \forall i \in \mathcal{I}, \quad (24)$$

$$y_{\tau_1,\tau_2} \in \mathbb{N}_0 \quad \forall \tau_1, \tau_2, \quad (25)$$

$$R_\tau^i \geq 0 \quad \forall \tau = 1, \dots, L \quad \forall i \in \mathcal{I}. \quad (26)$$

The general idea in calculating rollover in the  $L$ -day model is as follows. For a given day  $\tau$ , we first compute the number of jobs to be done on day  $\tau$ . To do this, we take the rollover from day  $\tau - 1$  and day  $\tau$ 's intake as a baseline number of jobs, and then we add the number of jobs pulled forward to day  $\tau$ , i.e.  $\sum_{\tau_1=\tau+1}^{\min\{\tau+K,L\}} y_{\tau_1,\tau}$ . We secondly compute the capacity that can be used to complete these jobs. This is done by taking the capacity  $c_\tau$  and subtracting the capacity required to complete those workstack jobs that are not pulled forward from day  $\tau$ , i.e.  $D_\tau - \sum_{\tau_2=\max\{\tau-K,1\}}^{\tau-1} y_{\tau,\tau_2}$ .

Constraints (20) and (21) provide upper bounds on the pulling forward totals, ensuring no jobs are pulled forward if they cannot be completed on the day to which they are moved, or if they could be completed on the day that they are being moved from. Constraint (22) reflects that jobs cannot be pulled forward from day 1 and hence we only subtract those jobs pulled forward to day 1 from its remaining capacity and do not reduce rollover by pulling forward from it. Similarly, constraint (24) says that jobs cannot be pulled forward to the final day of the plan and hence we only pull forward from this day and not to it. For every other day, constraint (23) captures that we can pull forward to *and* from said day and hence add and subtract jobs from its capacity to calculate the rollover.

To solve this model, we can reformulate it in the same way as we did the two-day model in Section 3.3, i.e. by defining a dummy decision variable  $t$ , adding a constraint  $t \geq \sum_{\tau=1}^L a_\tau \mathbb{E}_q(R_\tau) \quad \forall q \in \mathcal{U}$ , and minimising  $t$ . This is the formulation that we use when solving using a solver.

## 4.2 Binomial Intakes and Ambiguity Sets

Now let us, again, assume that  $I_\tau \sim \text{Bin}(i_\tau^{\max}, p_\tau)$ , where  $i_\tau^{\max} = \max(\mathcal{I}_\tau)$ . Suppose that we have a discrete uncertainty set  $\mathcal{U}$  of binomial distributions  $q$ , from which we can extract a set

$\mathcal{U}^p$  of success probability vectors  $p$ . As before, in the model (19)-(26), we replace  $q \in \mathcal{U}$  with  $p \in \mathcal{U}^p$ . In the  $L$ -day model, we consider ambiguity sets based on the normalised  $\ell^2$  distance from a forecasted or nominal one,  $p^f$ . We start by discretising  $[0, 1]^L$  using the formula in (27), where  $n_{\text{probs}}$  is chosen by the planner, and details the fineness of the discretisation.

$$\mathcal{U}_{\text{base}}^p = \left\{ \frac{j}{n_{\text{probs}}} \mid j = 0, \dots, n_{\text{probs}} \right\}^L \quad (27)$$

Secondly, we assume that we have some given point forecast,  $i^f$ , of the intake, alongside the range  $\mathcal{I}$ . From this, we obtain a forecasted probability vector  $p^f = i^f / i^{\text{max}}$  which results from assuming that  $i^f$  is the mean of the forecasted distribution. Given the forecast  $p^f$ , we consider only  $p \in \mathcal{U}_{\text{base}}^p$  that can be considered *close* to  $p^f$ . In these experiments, we use an ambiguity set of the form:

$$\mathcal{U}^p = \left\{ p \in \mathcal{U}_{\text{base}}^p \mid \frac{\|p^f - p\|_2}{\|p^f\|_2} \leq \gamma \right\}. \quad (28)$$

This set contains all  $p$  with an  $\ell^2$  distance from  $p^f$  that is no more than  $100\gamma\%$  of the  $\ell^2$  norm of  $p^f$ . Using a normalised distance like this allows us to use the same formulation for different  $L$  without needing to adjust the parameters, and  $\gamma$  represents our trust in the forecast.

Another common way of constructing  $\mathcal{U}^p$  is to consider a set of joint distributions within a prescribed distance from a nominal joint distribution constructed from forecasts, i.e. apply the distance measure to the distributions and not their parameters. For example, constructing a set using the  $\chi^2$  distance between joint distributions ensures a probabilistic guarantee by the way of the  $\chi^2$  goodness of fit hypothesis test, according to the schema proposed in Bertsimas et al. (2018). However, due to the univariate distributions being binomial with only one unknown parameter each, the same can be achieved by our construction in (28) with much less computational burden.

### 4.3 An Extension of S&S via Basic Heuristics

Here we detail the extension of S&S based on using a set of policies to *squash* the search space for  $y$ . In the two-day model, we used knowledge of the problem mechanics to construct a set  $\mathcal{Y}$  in which we were sure the optimal solution would lie. We did this by using the fact that pulling forward too much caused rollover and not pulling forward enough was wasteful of resources. Each value in this set acted as a candidate solution, for which we had to find the worst-case distribution in order to determine whether it was optimal.

The brute-force approach used in constructing the range for  $y$  is no longer possible, since our decision variable is too large in dimension for it to be computationally feasible. Hence, we instead generate candidate solutions using a selection of common-sense heuristics that emulate how a planner may pull forward. We employ the same logic as before when constructing each candidate solution, e.g. not pulling forward jobs if they will cause rollover. However, our set of candidate solutions will now be smaller in cardinality due to the computational effort involved in computing its elements. We are no longer certain that the set of candidate solutions will contain the optimal  $y$ , and we hence label the extended S&S as a heuristic.

### 4.3.1 Top Down and Bottom Up

The first heuristic, Top Down (TD), is based on a greedy approach where we start from day 1 and pull forward as much as we can from the following days, prioritising the earliest, and then move to day 2 and do the same, etc. Note that TD uses an adjusted upper bound on pulling forward that accounts for the forecasted jobs. This is to ensure it doesn't pull forward too much and cause rollover. The algorithm works as follows:

1. Initialise  $y_{\tau_1, \tau_2} = 0$  for all  $\tau_1, \tau_2$ .
2. Set  $\tau = 1$ .
3. Define  $y_{\tau}^{\max} = \left\lfloor \max \left\{ c_{\tau} - D_{\tau} - i_{\tau}^f, 0 \right\} \right\rfloor$  and set  $k = 1$ .
4. While  $y_{\tau}^{\max} > 0$  and  $k \leq K$  and  $\tau + k \leq L$ :
  - (a) Set  $y_{\tau+k, \tau} = \min \left\{ \max \{ D_{\tau+k} - c_{\tau+k}, 0 \} - \sum_{l=\max\{\tau+k-K, 1\}}^{\tau+k-1} y_{\tau+k, l}, y_{\tau}^{\max} \right\}$ .
  - (b)  $y_{\tau}^{\max} = y_{\tau}^{\max} - y_{\tau+k, \tau}$ .
  - (c)  $k = k + 1$ .
5. If  $\tau < L$  set  $\tau = \tau + 1$  and go to 3.
6. Return  $y$ .

When choosing the value of  $y_{\tau+k, \tau}$ , we calculate the number of incomplete jobs that have not been moved and compare it with the maximal amount of jobs that can be moved to  $\tau$ . The minimum of these two numbers is what we pull forward, to ensure none of the jobs are rolled over or did not need to be pulled forward. Bottom Up (BU) is the same as TD apart from that we start at the final day and work backwards, pulling forward from as far in the future as we can, i.e. prioritising later days. BU is shown in Appendix C.

### 4.3.2 Pull Half Forward

This is an algorithm that attempts to distribute the workload more evenly across the plan than the previous two heuristics. We first perform a load balancing step to smooth demand, where we split the days into pairs and attempt to pull forward jobs to make it so that each pair has approximately the same number of incomplete jobs. After doing this, we pull forward half of the incomplete jobs on day 2 of the pair to day 1 of the pair.

1. Initialise  $y_{\tau_1, \tau_2} = 0$  for all pairs of days.
2. Split the days  $1, \dots, L$  into  $\frac{L}{2}$  pairs  $\mathcal{L}_P = \{(1, 2), \dots, (L-1, L)\}$  if  $L$  is even. Else create a set of the form  $\mathcal{L}_P = \{(1, 2), \dots, (L-2, L-1)\}$  and do not consider  $L$ .
3. Perform load balancing:
  - (a) Calculate incomplete jobs for each day  $\tau$  as  $r_{\tau} = \max\{D_{\tau} - c_{\tau}, 0\}$ .
  - (b) Calculate total incomplete jobs for each pair as  $r_{(\tau_1, \tau_2)} = r_{\tau_1} + r_{\tau_2}$ .



(c) Calculate (rounded) average incomplete jobs per pair as  $\bar{r}_P = \left\lfloor \frac{1}{|\mathcal{L}_P|} \sum_{(\tau_1, \tau_2) \in \mathcal{L}_P} r_{(\tau_1, \tau_2)} \right\rfloor$ .

(d) For  $(\tau_1, \tau_2) \in \mathcal{L}_P$  such that  $r_{(\tau_1, \tau_2)} < \bar{r}_P$ :

i. Set  $y^{\max} = \max\{c_{\tau_2} - D_{\tau_2} - \sum_{k=1}^K y_{\tau_2+k, \tau_2}, 0\}$ .

ii. If  $r_{(\tau_2+1, \tau_2+2)} > \bar{r}_P$  and  $r_{\tau_2+1} > 0$ , pull forward:

$$y_{\tau_2+1, \tau_2} = \min\{r_{\tau_2+1}, \bar{r}_P - r_{(\tau_1, \tau_2)}, y_{\max}\}.$$

iii. Update  $r$  via

$$r_{\tau_2} = \max\{D_{\tau_2} + y_{\tau_2+1, \tau_2} - c_{\tau_2}, 0\},$$

$$r_{\tau_2+1} = \max\{D_{\tau_2+1} - y_{\tau_2+1, \tau_2} - c_{\tau_2+1}, 0\},$$

and redefine  $r_{(\tau_1, \tau_2)}$  and  $r_{(\tau_2+1, \tau_2+2)}$ .

iv. If we still have  $r_{(\tau_1, \tau_2)} < \bar{r}_P$  and  $r_{(\tau_2+1, \tau_2+2)} > \bar{r}_P$  and  $r_{\tau_2+2} > 0$ , pull forward:

$$y_{\tau_2+2, \tau_2} = \min\{r_{\tau_2+2}, \bar{r}_P - r_{(\tau_1, \tau_2)}, y_{\max} - y_{\tau_2+1, \tau_2}\}.$$

v. Do update steps as in iii.

4. For each pair  $(\tau_1, \tau_2) \in \mathcal{L}_P$ , set

$$y_{\tau_2, \tau_1} = \min \left\{ \left\lfloor \frac{1}{2} \left( \max\{D_{\tau_2} - c_{\tau_2}, 0\} - \sum_{k=1}^K y_{\tau_2, \tau_2-k} \right) \right\rfloor, \left( \max\{c_{\tau_1} - D_{\tau_1}, 0\} + \sum_{k=1}^K y_{\tau_1, \tau_1-k} \right) \right\}.$$

The load balancing step in 3. moves from pairs with a number of incomplete jobs above the average per pair,  $\bar{r}_P$ , to pairs below the average. The maximum it can pull forward to the pair that is below average is the amount of spare capacity it has on the second day in the pair after the pulling forward that has already been done. This ensures the jobs stay in that pair and do not roll over into the next pair.

### 4.3.3 Extended S&S Algorithm

We now detail the extension of S&S using these basic heuristics. The idea behind S&S is to construct a set  $\mathcal{Y}$  of potential  $y$  solutions, and a set  $\mathcal{P}$  of potential  $p$  solutions, and then test every combination of the two sets. In order to keep the algorithm fast, these sets cannot grow large. In the case of  $y$ , the number of potential choices is very large, although the time in computing the solutions is not. For  $p$ , however, if we were to test every distribution then the computational effort involved would be very large. To see this, consider an example with  $|\mathcal{U}^p| = 3883$  distributions and  $|\mathcal{I}| = 20000$  potential intakes. Then for each of 3883 distributions we will need to compute a product of  $L$  binomial PMF values, for each of 20000 intakes. This means computing  $L \times 3883 \times 20000 = L \times (77.66 \times 10^6)$  PMF values.

We have already discussed the use of heuristics to construct  $\mathcal{Y}$ , but it may be beneficial to explain further the choice of  $\mathcal{P}$ . The model needs to find the  $p$ -vector which leads to the highest

total expected rollover cost for any fixed  $y$  solution. Since each  $p_\tau$  is a success probability corresponding to the distribution of intake, higher  $p_\tau$  leads to increased expected intake on day  $\tau$ . Since capacity is fixed on day  $\tau$ , higher  $I_\tau$  leads to higher (or the same) rollover. Following this logic it is clear that the distributions with the highest expected rollover will be those with the highest values for  $p_\tau$  for at least one  $\tau$ . If this is not the case, more expected rollover could be caused on some day  $\tau$  by simply increasing  $p_\tau$ . Furthermore, the expected rollovers for the days other than  $\tau$  should also be increasing in their success probabilities if  $p_\tau$  is fixed, and so they should be the highest that they can be, given that this is  $p_\tau$  has been maximised. This leads to our set  $\mathcal{P}^{\text{ext}}$  described in step 2 of S&S.

Let us define the vector  $S = (c, D, a, L, K)$  to be the inputs to the  $L$ -day model. Then, we describe our extension of S&S as follows.

1. Construct a set  $\mathcal{X}$  of algorithms  $x$  that take input  $S = (c, D, a, L, K)$  to create a solution  $y$  to our  $L$ -day model. We write the solution given by algorithm  $x \in \mathcal{X}$  as  $x(S)$ .
2. Find  $p_\tau^{\text{max}} = \max_{p \in \mathcal{U}^p} p_\tau$  for each  $\tau = 1, \dots, L$  and find

$$\mathcal{P}_\tau^{\text{max}} = \{p \in \mathcal{U}^p : p_\tau = p_\tau^{\text{max}}\} \text{ for } \tau = 1, \dots, L,$$

$$\mathcal{P}_\tau^{\text{ext}} = \operatorname{argmax}_{p \in \mathcal{P}_\tau^{\text{max}}} \left\{ \sum_{k=1}^L p_k \right\}, \quad \mathcal{P}^{\text{ext}} = \bigcup_{\tau=1}^L \mathcal{P}_\tau^{\text{ext}}.$$

3. Construct the set of candidate solutions for  $y$  given by applying all algorithms:

$$\mathcal{Y} = \{x(S) : x \in \mathcal{X}\}.$$

4. For  $y \in \mathcal{Y}$ :

- (a) For  $p \in \mathcal{P}^{\text{ext}}$  calculate  $C_p = \sum_{\tau=1}^L a_\tau \mathbb{E}_p(R_\tau)$  given  $y$ .
- (b) Choose  $p_y$  such that  $C_{p_y} = \max_{p \in \mathcal{P}^{\text{ext}}} (C_p)$ .

5. Choose  $y^*$  such that

$$y^* = \operatorname{argmin}_{y \in \mathcal{Y}} C_{p_y}. \tag{29}$$

6. Return values  $y^*, x^*, p^*$ .

The set  $\mathcal{P}^{\text{ext}}$  is designed to contain only the most extreme probability vectors. The set  $\mathcal{P}_\tau^{\text{max}}$  contains all probability vectors  $p$  such that at  $p_\tau$  is at its maximum. Among these vectors, the other values are not important. Hence,  $\mathcal{P}_1^{\text{max}}$  might contain, for  $L = 2$ , the vectors  $(0.9, 0.1)$ ,  $(0.9, 0.2)$  and  $(0.9, 0.3)$ . We are suggesting here that the most extreme vector will be the worst, and so given that  $p_1 = 0.9$  we want to extract the most extreme, i.e. the one where the other probabilities are highest. In this case, we would select  $(0.9, 0.3)$ . These vectors will have the highest sum, meaning we take those elements of  $\mathcal{P}_1^{\text{max}}$  with the largest sum. Then  $\mathcal{P}^{\text{ext}}$  is the union of the extreme sets for each day. Note that typically  $\mathcal{P}^{\text{ext}}$  will contain distributions where only one probability is at its highest.

## 4.4 Solver-based Heuristics

From initial experiments, S&S performs well in selecting  $p$  but not so well in selecting  $y$ . Hence, we now detail two algorithms that make use of a solver to choose  $y$ . This will improve our ability to optimise the pulling forward decisions without sacrificing our ability to identify the worst-case distribution for a fixed  $y$ .

### 4.4.1 Cutting Surface Algorithm

In this section, we describe our adaptation of the cutting surface (CS) algorithm as detailed in the literature review, which has been commonly used in the DRO literature. The algorithm has a number of different forms, but the one that we base our adaptation on is that of the review paper by Rahimian and Mehrotra (2019). The general idea of the algorithm is as follows. In order to deal with the large number of constraints implied by the ambiguity set, the algorithm uses the following steps. We start with a singleton set containing one distribution, and solve the outer problem with this small ambiguity set. Then, for the generated solution, we find the worst-case distribution. We add this distribution to the ambiguity set and then repeat the above. These steps are repeated until optimality criteria are met.

In more detail, suppose we have some initial subset  $\mathcal{U}^0$  of our set of distributions  $\mathcal{U}$  and we solve the full model but only considering distributions in  $\mathcal{U}^0$ , and get an optimal decision  $y^1$ . Then, we find the worst-case distribution,  $p^1 \in \mathcal{U}$ , for the fixed solution  $y^1$ , and add it to our set to create a new subset  $\mathcal{U}^1 = \mathcal{U}^0 \cup \{p^1\}$ . We then solve the full model with subset  $\mathcal{U}^1$ , and repeat. We stop iterating when we have reached  $\varepsilon$ -optimality, i.e. if the solution from the full problem at iteration  $k$ ,  $y^k$ , gives an expected cost of within  $\varepsilon/2$  of the worst-case expected cost for  $y^k$  over all distributions. The algorithm returns an  $\varepsilon$ -optimal solution to the DRO model in a finite number of iterations.

The issue with CS is that, even if  $y$  is fixed at  $y^k$ , finding the worst-case distribution  $p^k$  can be a very difficult task. To find this distribution, we would need to first compute the values of every PMF in the set  $\mathcal{U}$  and then solve a large LP to maximise the expected rollover cost over all distributions in this set. Since computing the PMF values is computationally intensive in itself, this would take almost as long as solving the full model without  $y$  being fixed. As well as this, if we were to treat our inner model as an LP with  $p$  as a decision variable, then we would have a model of the form:

$$\max_{p \in \mathcal{U}^p} \sum_{\tau=1}^L a_{\tau} \sum_{i \in \mathcal{I}} R_{\tau}^i \left[ \prod_{l=1}^L \binom{i_l^{\max}}{i_l} p_l^{i_l} (1-p_l)^{i_l^{\max}-i_l} \right] \quad (30)$$

which is highly non-linear in  $p_l$  for each  $l$ , and also non-concave in general. Instead of following either of these approaches, we use a hybrid S&S-CS algorithm where we solve the distribution separation problem using the  $p$ -squash step of S&S. By this, we mean that we apply Step 4. of the S&S algorithm from Section 4.3.3 and only consider distributions in the set  $\mathcal{P}^{\text{ext}}$  as defined in Step 2. of this algorithm. This allows us to heuristically generate a distribution to add to our set in significantly less time than if we were to solve to optimality. The use of the  $p$ -squash step

from S&S is motivated by the strong performance of S&S in finding the worst case  $p$  for a given  $y$ . The algorithm, which we will refer to from now on simply as CS, is given below.

1. Initialise  $\mathcal{U}^{p,0} = \{p^0\}$ , where  $p^0 = p^f$  for example.
2. For  $k = 0, \dots, k^{\max}$ :
  - (a) Solve the full model to optimality using the set  $\mathcal{U}^{p,k}$  to generate solution  $(y^k, t^k)$  where  $t^k$  is worst-case expected cost of  $y^k$  over the set  $\mathcal{U}^{p,k}$  passed to the model.
  - (b) Solve distribution separation problem  $\max_{p \in \mathcal{U}^p} \sum_{\tau=1}^L a_\tau \mathbb{E}_p(R_\tau \mid y = y^k)$  to get solution  $p^k$  (via  $p$ -squash of S&S):
    - i. Calculate  $\mathcal{P}^{\text{ext}}$  as in S&S step 2.
    - ii. For  $p \in \mathcal{P}^{\text{ext}}$  calculate  $C_p = \sum_{\tau=1}^L a_\tau \mathbb{E}_p(R_\tau \mid y = y^k)$ .
    - iii. Choose  $p^k$  such that  $C_{p^k} = \max_{p \in \mathcal{P}^{\text{ext}}}(C_p)$ .
  - (c) If  $C_{p^k} \leq t^k + \frac{\varepsilon}{2}$  or  $p^k \in \mathcal{U}^{p,k}$  then stop and return solution  $(y^k, p^k)$ .
  - (d) Else,  $\mathcal{U}^{p,k+1} = \mathcal{U}^{p,k} \cup \{p^k\}$  and  $k = k + 1$ .

Note that we use  $\mathcal{U}^{p,k}$  instead of  $\mathcal{U}^k$  as we apply the algorithm to the set of  $p$ -vectors, not the set of distributions. The logic behind 2(c), where we check if  $p^k \in \mathcal{U}^{p,k}$ , is that calculation differences might cause  $t^k$  and  $C_{p^k}$  to slightly differ even if  $p^k$  really is the worst case distribution for  $y^k$ . Solvers use some simplifications in their calculations that the function used in 2(b) will not use. If this is the case and we add  $p^k$  to the set and run the model again, then the set of distributions will be unchanged and so we will simply be running the same model again. Then the same  $p^k$  will be generated and nothing will change. By adding this extra stopping criterion we catch these cases. The new criterion cannot cause early stopping, since the only time it is triggered is when the other stopping criteria should also have been triggered.

We could also take the outputs  $y^k$  and  $p$  from the outer problem and calculate  $t^k$  using our own function for the objective value. However, this will simply cause more computations than are necessary, and has been verified to give the same solutions in the same number of iterations.

#### 4.4.2 Approximate-objective Algorithm

The final algorithm we describe is the Approximate-objective algorithm (AO). When solving the model to optimality, we are required to compute the distributions  $q \in \mathcal{U}$  based on our probabilities  $p$ . For each intake  $i$  we can easily compute:

$$q_{\max}(i) = \max_{q \in \mathcal{U}} q(i) = \max_{p \in \mathcal{U}^p} \mathbb{P}(I = i \mid p), \quad (31)$$

and then we can consider a new set of intakes in the model defined by:

$$\tilde{\mathcal{I}} = \{i \in \mathcal{I} : q_{\max}(i) > \beta\} \quad (32)$$

where  $\beta$  is our minimum accepted probability. By tuning  $\beta$ , we are removing intakes from our set that are very unlikely. When solving the model, we are approximating the expected

value by removing some small terms, but since the intakes removed have low probability, this approximation should be strong. For this paper, we use  $\beta = 10^{-3}$  as our initial testing showed that this value led to good improvements in computation times.

## 5 Design of Computational Experiments

This section details our experiments evaluating the performance of the algorithms described in Sections 4.3 and 4.4. In this section, we discuss how the parameters for the experiments will be chosen to ensure that they are representative of typical real-life scenarios.

To discuss experimental design, we need to define which parameters of the model will be varied and the values that they will take. The vector of inputs to the model for a fixed set  $\mathcal{I}$  of intakes and  $\mathcal{U}$  of distributions is  $S = (c, D, a, L, K)$ , where  $c$  is the capacity,  $D$  the workstack,  $a$  the rollover cost,  $L$  the length of the plan in days and  $K$  the maximum number of days a job can be pulled forward.

### 5.1 Parameter Hierarchy

It is helpful to consider a hierarchy of parameter choices, which is defined by the following ordering:

1.  $(L, K)$  defines the difficulty of the problem in terms of the MIP itself.
2.  $c$  and  $D$  define the set of solutions that are possible for a given model with fixed  $L$  and  $K$ , and need to be constructed for each combination of these parameters to ensure we have a varied range of instances when it comes to pulling forward opportunities:
  - (a) Pulling forward is not possible. In this case, the optimal solution should be somewhat easy to find, and we would expect all algorithms to be optimal.
  - (b) Pulling forward opportunities are small. In this case, we might only have a couple of  $y$  variables that can be non-zero, and all algorithms might be expected to find these pairs and choose them optimally.
  - (c) Pulling forward opportunities in abundance. This is where we will expect large deviations between heuristic and optimal solutions, since it is less likely our heuristics will find the optimal  $y$ .

The conditions under which these possibilities occur are defined in Section 5.2.

3.  $\mathcal{I}$  and  $\mathcal{U}$  define how the uncertainty is encoded in the model, depending on the planner's attitude to risk. If  $|\mathcal{I}|$  or  $|\mathcal{U}|$  are large, solving to optimality will be very slow, and we would like to use a heuristic that is not significantly affected.  $|\mathcal{I}|$  is defined by  $i^{\max}$  and  $|\mathcal{U}|$  is defined by two parameters. The initial discretisation of the interval  $[0, 1]$  in which each  $p_\tau$  lies is defined by  $n_{\text{probs}}$ . The maximum distance from the forecasted distribution that  $p \in \mathcal{U}^p$  can be is defined by the second parameter  $\gamma$ . A larger value of  $\gamma$  indicates a more risk-averse planner.

4.  $a$  will be left as the ones vector for these experiments as it has not been seen to have an effect on solutions.

We choose  $L = 5$  due to it being the number of days in a typical working week. We take the maximum pulling forward window length to be  $K = 2$ , since pulling forward is not enacted until the operational planning phase, where the planning horizon is very short. These choices are partly motivated by usual practices, and also partly by the fact that we aim to test our heuristics against optimal solutions, and for larger  $L$  or  $K$  the model becomes very difficult to solve to optimality.

Note that the optimality tolerance for CS,  $\varepsilon$ , will be set as 0.01 and it will be ran for a maximum of  $k_{\max} = 10$  iterations. From initial testing, 10 iterations was easily enough to reach  $\varepsilon$ -optimality, and even with  $\varepsilon = 0.01$ , CS returned optimal solutions. In fact, it takes CS typically only 1 or 2 iterations to return a solution.

## 5.2 Capacity and Workstacks

The factors affecting the potential solutions of a model the most are  $c$  and  $D$ , due to the fact that they define rollover and pulling forward opportunity. In this section, we detail the capacities  $c$  and workstacks  $D$  used in our experiments, which are constructed with the aim of ensuring that 2(a)-(c) from Section 5.1 are all represented by at least one  $(c, D)$  pair. We assume for this section that the previous parameters in the hierarchy, i.e.  $L, K$  are given. We define the situations 2(a)-(c) mathematically below.

Our choices for  $c$  and  $D$  are based on differing opportunities for pulling forward. We define the set of pairs of days under consideration for pulling forward as:

$$\mathcal{F} = \{(\tau_1, \tau_2) \mid \tau_1 \in \{2, \dots, L\}, \tau_2 \in \{\tau_1 - K, \dots, \tau_1 - 1\}\} \quad (33)$$

and the set of pairs such that the corresponding  $y$  can feasibly be positive given  $c$  and  $D$  as:

$$\mathcal{F}^+(c, D) = \{(\tau_1, \tau_2) \in \mathcal{F} \mid c_{\tau_1} < D_{\tau_1}, c_{\tau_2} > D_{\tau_2}\}. \quad (34)$$

This is the set of all pairs of days such that the second is within pulling forward range of the first, the second has spare capacity and the first has incomplete jobs. Case 2(a) corresponds to the cases where  $\mathcal{F}^+(c, D) = \emptyset$ . This can easily be constructed by ensuring, for example, that  $c_\tau > D_\tau \forall \tau \in [L]$ . In our experiments, for each  $(L, K)$  we will have  $c$  and  $D$  such that that the set of combinations of the two has  $(c, D)$  such that  $|\mathcal{F}^+(c, D)| = m$  for every possible  $m$ . This ensures we will have a good variety of potential solutions to our model. We ensure this by creating one value for  $c$  and then constructing multiple  $D$  vectors for that  $c$ . We can ensure that  $|\mathcal{F}^+(c, D)| = 1$  by, for example, setting  $D_1 = c_1 - 1$ ,  $D_2 = c_2 + 1$ , then  $D_\tau < c_\tau$  for all  $\tau > 2$ . This can be done for  $m \geq 2$  in a similar way.

The main effect that  $c$  and  $D$  has on decision making is defining the constraints on  $y$ , meaning their only significant quality is how much pulling forward they do or do not allow. Using this set of values for  $c$  and  $D$  we will be able to see how well our algorithms detect and make use of opportunities for pulling forward.

### 5.3 Uncertainty/Ambiguity Sets

As a reminder, the term “uncertainty set” refers to  $\mathcal{I}$  and “ambiguity set” refers to  $\mathcal{U}^p$ . We now detail the parameters used to construct these sets in our instances.

#### 5.3.1 Uncertainty Set

We noted in Section 1.1 that, in practice, as a result of most forecasting models giving upper and lower confidence bounds, we would be given a set  $\mathcal{I}$  and would then extract  $i^{\max}$  from this set. However, in these experiments it is more convenient to define  $i^{\max}$  and then use this to construct  $\mathcal{I}$ . Since there is a one-to-one mapping between the two, the two methods achieve the same result. In the two-day model, we considered distributions that constrained the probability that the total intake exceeded the total residual capacity. This was to establish realism through not allowing high intakes to occur on all days simultaneously. However, this calculation is very computationally intensive in the  $L$ -day problem, since it is the sum of a large number of individual multivariate PMF values, which are products of univariate PMF values. For this reason, we do not require the aforementioned constraint any more, but we do enable a similar restriction on intakes using constraints on  $i^{\max}$ . Specifically, we will consider  $i^{\max}$  satisfying:

$$\sum_{\tau=1}^L i_{\tau}^{\max} \leq \sum_{\tau=1}^L \max(c_{\tau} - D_{\tau}, 0). \quad (35)$$

This is a relaxed version of constraint (11), where we required the intakes not to exceed the total remaining capacity. This is reasonable because if the total number of jobs arriving in the system exceeds the RHS of (35) then some intake jobs will always remain incomplete at the end of day  $L$ , regardless of our pulling forward decision. Furthermore, we can consider varying numbers of high-intake days, through the number  $n(i^{\max})$  as defined below:

$$n(i^{\max}) = |\{\tau \in [L] : i_{\tau}^{\max} > c_{\tau} - D_{\tau}\}|, \quad (36)$$

which corresponds to the number of days with potential spikes in demand. Depending on  $c$  and  $D$ , it can range between 0 and  $L - 1$ , but for these experiments we consider  $n(i^{\max}) \in \{1, \lfloor \frac{L}{2} \rfloor, L - 1\}$  for sufficient coverage of cases. The case of 1 day with high intake corresponds to a one-day spike caused by an event such as a major weather event. The case of  $\lfloor \frac{L}{2} \rfloor$  days with high intake could correspond to an extended spike lasting for multiple consecutive days, for example, a network problem causing lots of service devices to break. The final case of  $L - 1$  will have  $L - 1$  small spikes in intake, marking a period of consistently high intake.

#### 5.3.2 Ambiguity Set

The choice of ambiguity set depends on the choice of discretisation of  $[0, 1]^L$  and also the way we in which we then reduce its size. The choice of discretisation is defined by the parameter  $n_{\text{probs}}$ , and increasing this value increases the size of the ambiguity set. For these experiments, we consider  $n_{\text{probs}} \in \{5, 10, 15\}$ . In our preliminary testing we found that any value larger than 15 can lead to intractability when solving to optimality. In these experiments, we assume

$i^f = (0.75i_1^{\max}, \dots, 0.75i_L^{\max})$ . Hence, we will obtain  $p^f = (0.75, \dots, 0.75)$ , which corresponds to a safe forecast, in that we are preparing for high intakes.

The value of the forecast is not particularly relevant in itself, since our trust in the forecast is what really defines the size and properties of the ambiguity set. When defining  $\gamma$ , we take values in  $\{0.05, 0.10, 0.15\}$ . These values are not based on any statistical confidence, but are designed to represent high, medium and low confidence in the forecast. The largest ambiguity set we obtain from these parameters is of size 3883, and occurs when  $\gamma = 0.15$  and  $n_{\text{probs}} = 15$ . In our experiments, this choice led to multiple problems that did not solve to optimality within the maximum run-time of 4 hours. However, CS and S&S are still viable for this set, and our results suggest that they would be viable for even larger ambiguity sets as well.

## 6 Results

We now detail the results of our experiments that test the algorithms on problem instances in which  $L = 5$  and  $K = 2$ . The total number of instances run is 186 for this  $(L, K)$  pair. For each algorithm, the maximum time allowed for model building and solving was 4 hours. If the total time taken reached a value above this at any point then the algorithm returned the best solution found so far. In this section, we report the results for instances where all algorithms finished running. This was the case in 150 of 168 instances. We report the results for the 18 instances where opt timed out in Appendix E.

These experiments were run in parallel on a computing cluster (STORM) which has almost 500 CPU cores. We used 40 of the cores to perform the experiments, and the solver used in all cases was the Gurobi Python package, `gurobipy`. The version of `gurobipy` used was 9.0.1. The node used on STORM was the Dantzig node, which runs the Linux Ubuntu 16.04.6 operating system, Python version 2.7.12, and 48 AMD Opteron 638 CPUs.

### 6.1 Summary of Instances and Their Sizes

In Table 1, we summarise the sizes of the sets  $\mathcal{I}$  and  $\mathcal{U}^p$ , that form the basis for constraints and variables in the model. A summary of the sizes of  $\mathcal{I}$  is given in Table 1a. The table shows 7 of the 31  $i^{\max}$  values considered and the size of the resulting set  $\mathcal{I}$ . The other  $i^{\max}$  values considered were permutations of the values shown in the table, and hence led to  $|\mathcal{I}|$  values that are already listed in the table.

We can see here that our choices of  $i^{\max}$  gave instances with as many distinct intakes (and rollover vectors) as 20000, and as few as 392. The sizes of the ambiguity sets (of distributions) is given in Table 1b. We found that this had a much more significant effect on solution times than  $i^{\max}$ , and hence these sets are not as large.

In the following subsections we detail the results for those instances where all algorithms finished running in the allocated time of 4 hours. Of the 18 instances where the solver algorithm (opt) timed out, there were 5 in which AO also timed out. As mentioned earlier, the results for these



$i^{\max}$	$ \mathcal{I} $
(1, 6, 6, 1, 1)	392
(1, 3, 3, 3, 3)	512
(2, 2, 2, 6, 2)	567
(2, 2, 8, 8, 2)	2187
(5, 5, 1, 5, 5)	2592
(1, 7, 7, 7, 7)	8192
(9, 9, 1, 9, 9)	20000

(a) Example  $i^{\max}$  values and sizes of the associated uncertainty sets  $\mathcal{I}$  considered.

$n_{\text{probs}}$	$\gamma$	$ \mathcal{U}^p $
5	0.15	16
10	0.10	32
10	0.15	512
15	0.05	16
15	0.10	562
15	0.15	3883

(b) Parameters defining ambiguity sets and size of corresponding sets.

Table 1: Summary of input parameters and corresponding set sizes

instances are given in Appendix E.

## 6.2 Optimality and Time Taken

Comparing results for DRO problems is not as simple as comparing final objective values. Our optimal objective value can be written as:

$$z^* = \min_y \max_p f(y, p),$$

where  $f(y, p) = \sum_{\tau=1}^L a_{\tau} \mathbb{E}_p(R_{\tau} | y)$  is the total expected rollover cost. Suppose we have an instance where  $y^{\text{CS}} = y^{\text{opt}}$  but  $p^{\text{CS}} \neq p^{\text{opt}}$ . Then, if CS gives a lower objective value than opt, it may appear to have given a better solution to the minimisation problem. However, this means that CS did not successfully choose the worst-case  $p$  for its chosen  $y$ . This leads to a lower objective but a suboptimal solution overall. Similarly, we can say that CS is suboptimal if  $p^{\text{opt}} = p^{\text{CS}}$  but  $y^{\text{CS}} \neq y^{\text{opt}}$  and CS gave a higher objective value. Hence, both a higher and a lower objective value can suggest suboptimality for a DRO model.

Given this, we summarise the results using a number of counts. An algorithm  $x$  is said to be:

1.  **$y$ -optimal** if  $\max_{p \in \mathcal{U}^p} f(y^x, p) = z^*$ .
2.  **$p$ -optimal** for a given  $y^x$  if  $f(y^x, p^x) = \max_{p \in \mathcal{U}^p} f(y^x, p)$ .
3. **Optimal** if  $f(y^x, p^x) = z^*$ . Note that this is met if the algorithm is both  $y$ -optimal and  $p$ -optimal.

We display the number of times each algorithm was optimal,  $p$ -optimal and  $y$ -optimal in Table 2.

Table 2 shows that both CS and AO were optimal in more than 93% of instances, and  $y$ -optimal in more than 98% of instances. S&S has weaker performance in selecting the optimal  $y$ , but

Alg.	No. (%) Optimal Sol	No. (%) $p$ -Optimal Sol	No. (%) $y$ -Optimal Sol
CS	161 (95.83%)	162 (96.43%)	166 (98.81%)
S&S	59 (35.12%)	162 (96.43%)	59 (35.12%)
AO	157 (93.45%)	160 (95.24%)	165 (98.21%)

Table 2: Summary of optimality of heuristics

performs very well (just as well as CS) in selecting the worst-case  $p$  for its chosen  $y$ . Closeness to optimality of the algorithms is discussed in Section 6.3.

A summary of the computation times of each algorithm is given in Table 3. S&S takes around 1 minute on average and CS takes around 1 minute 51 seconds. To find the optimal solution, it took approximately 28 minutes on average, which is quite a significant difference. AO is somewhere in the middle, taking around 9 minutes on average.

Algorithm	Min t.t.	Avg. t.t.	Max t.t.
opt	0:00:05.25	0:28:59.01	3:41:37.82
S&S	0:00:00.34	0:00:54.59	0:09:11.77
CS	0:00:02.11	0:01:51.11	0:12:09.45
AO	0:00:02.76	0:08:40.86	1:07:21.78

Table 3: Summary of times taken

Based on the optimality counts and time taken, CS is the strongest performing heuristic. These results only give the number of instances in which each algorithm was optimal. Closeness to optimality is discussed in detail in the following section.

### 6.3 Algorithm Performance in Detail

To illustrate further how well the algorithms performed, we define the following two metrics. Note that a positive value for *either* of these metrics suggests suboptimality.

1. **Quality of  $p$  choices.** For a solution  $y^x$  that was selected by an algorithm  $x \in \{\text{CS}, \text{S\&S}, \text{AO}\}$ , we calculate the worst-case expected cost over all distributions in  $\mathcal{U}^p$  using brute force. We can then compare this cost with the expected cost obtained by the algorithm, i.e. from  $p^x$ , the  $p$  that the algorithm selected. This allows us to establish how close to worst-case the choices in  $p$  were. We refer to this difference as the  $p$ -gap, and it is defined as:

$$g_p(y^x, p^x) = \max_{p \in \mathcal{U}^p} f(y^x, p) - f(y^x, p^x).$$

2. **Quality of  $y$  choices.** For a given solution  $y^x$  from algorithm  $x$ , we compute the worst-case expected cost using brute force, as we did when finding  $g_p(y^x, p^x)$ . We can then

compare this worst-case cost with that of the optimal  $y$ , to assess how good  $y^x$  is. This is referred to as the  $y$ -gap, and is defined as:

$$g_y(y^x) = \max_{p \in \mathcal{U}^p} f(y^x, p) - z^*.$$

In Table 4 we summarise the average  $p$ -gaps and  $y$ -gaps of the three heuristics, along with the average absolute percentage gaps (APGs). The  $p$ -APG is obtained by taking the  $p$ -gap as an absolute percentage of the worst-case expected cost for the chosen solution  $y^x$ . The  $y$ -APG is obtained by taking the  $y$ -gap as an absolute percentage of the optimal solution value.

Alg.	Avg. $p$ -gap	Avg. $p$ -APG	Avg. $y$ -gap	Avg. $y$ -APG
CS	0.0065	0.0061%	0.0004	0.0003%
S&S	0.0057	0.0053%	0.8309	0.9227%
AO	0.0015	0.0079%	0.0002	0.0002%

Table 4: Summary of gaps and APGs of the heuristics

This suggests that all algorithms perform very well at choosing the worst-case  $p$  for a fixed  $y^x$ , and all have an average  $p$ -APG of less than 0.008%. AO arguably performs the best at selecting  $p$ , if we use  $p$ -gap as our metric rather than  $p$ -APG. This is due to the fact that it's the only heuristic which chooses  $p$  from the entire set of distributions. CS and AO are very good at selecting the optimal  $y$ , since they both use the solver to select it, and on average they both have a  $y$ -APG of less than 0.0004%. The  $y$  solution they chose had, on average, a worst-case expected cost that was less than 0.0005 away from the optimal objective value. S&S does not perform as well at selecting  $y$ , although it still has an average  $y$ -APG of less than 1%.

We can also study the results broken down by the size of the set of distributions. We present these results in Table 6, which is in Appendix D.1 due to space considerations. In summary, the table suggests that S&S and CS do not encounter many problems in finding the worst-case  $p$  for their chosen  $y^x$  until the set reaches its largest size, i.e. 3883. This is because the set they are using will only contain a small fraction of the total number of distributions in these cases. These two algorithms are consistent in their selection of the optimal  $y$  across all values of  $|\mathcal{U}^p|$ . AO performs consistently with respect to  $p$  and  $y$  across all sizes, with the highest average  $p$ -APG and  $y$ -APG occurring when  $|\mathcal{U}^p| = 16$ .

The algorithms that use Gurobi on the full set of distributions, i.e. opt and AO, do not scale well with  $|\mathcal{U}^p|$ . For AO, solve times average approximately 1 hour when  $|\mathcal{U}^p| = 3883$ . The optimal algorithm reaches an average solution time of nearly 1 hour when  $|\mathcal{U}^p| = 512$ . We can therefore conclude that AO scales better with  $|\mathcal{U}^p|$  than opt. S&S and CS's solutions times are not significantly affected by  $|\mathcal{U}^p|$ , taking less than 3 minutes.

Finally, we can look at the performance of algorithms by the size of the set of intakes  $\mathcal{I}$ . These results are shown in Table 8, which is given in Appendix D.3. The computation times of each heuristic algorithm are not strongly impacted by the number of intakes. They typically increase

with this number, but only by a few minutes. AO in particular is less affected by  $|\mathcal{I}|$  due to its removal of low-probability intakes. The  $p$ -APGs are also not significantly affected, and for AO and CS neither are the  $y$ -APGs. The main thing we notice here is that the performance of S&S in terms of  $y$ -APG does seem to vary with the number of intakes, although it is not a strictly increasing relationship. S&S performs best for those instances where  $|\mathcal{I}|= 20000$  and 2592, with poorer performance occurring for the smaller sets of intakes. The optimal algorithm, as expected, takes the longest for  $|\mathcal{I}|= 20000$ .

We see from Tables 6 and 8 that the longest solution times from opt occurred when  $|\mathcal{U}^p| = 562$  and when  $|\mathcal{I}| = 20000$  and 567. This is due to the fact that the instances where  $|\mathcal{U}^p| = 3883$  and  $|\mathcal{I}| > 2000$  timed out and are hence not shown in these tables.

#### 6.4 The Effect of Workstacks on Solutions

In our experiments, we used only one value of the capacity  $c$  but varied the workstacks  $D$  to give a variety of possibilities for pulling forward. This was based on the number of pairs between which pulling forward was possible, i.e.  $|\mathcal{F}^+(c, D)|$  from Section 5.2. We summarise the values of  $c - D$  and  $|\mathcal{F}^+(c, D)|$  in Table 5.

$c - D$	$ \mathcal{F}^+(c, D) $
(8, -15, -15, 8, -15)	3
(8, -15, -15, 8, 8)	2
(8, -15, 8, 8, 8)	1
(8, 8, 8, 8, 8)	0

Table 5: Summary of  $c - D$  values and corresponding number of pairs

Any more pairs than 3 is not possible for  $L = 5$  and  $K = 2$ . We did not obtain any solutions that pulled forward between more than 1 pair of days. This could be explained by the way that rollover passes through the plan, incurring a cost multiple times if it rolls over on multiple days. Days 1 and 2 are typically prioritised for rollover reduction via pulling forward, since jobs due on these days have the potential to roll over the most times. There was one instance in which we found  $y_{5,4}^{\text{opt}} > 0$ , and all the rest had only  $y_{2,1}^{\text{opt}} > 0$ . Even in this case,  $y_{2,1}^{\text{opt}} > 0$  was returned by the other algorithms and was also optimal.

For reference, the results broken down by  $|\mathcal{F}^+(c, D)|$  are shown in Table 7, in Appendix D.2. Firstly, we note that all algorithms were always  $y$ -optimal when there were no feasible pairs (since  $y = 0$  is the only feasible solution). Other than this there is no clear effect of  $|\mathcal{F}^+(c, D)|$  on  $y$ -optimality. We also note that S&S and CS perform the worst in terms of  $p$ -gap when the number of feasible pairs is largest. Outside of this there is no clear relationship between  $|\mathcal{F}^+(c, D)|$  and  $p$ -optimality. The invariance in  $y$ -APGs is likely due to the fact that making use of any more pairs than 1 is was suboptimal in all cases, and no algorithm ever did this. The number of feasible pairs does not affect the  $p$  choices, explaining the invariance in this

regard. We see that average solution times generally decrease as  $|\mathcal{F}^+(c, D)|$  increases. This effect (and the number of instances for each number of pairs) can be explained by the corresponding  $i^{\max}$  values. Differences in  $c - D$  lead to differences in the possible values for  $i^{\max}$ . This is caused by constraint (35), which leads to the effect that more days with incomplete jobs, i.e.  $\max(c_\tau - D_\tau, 0) = 0$ , leads to a lower total maximum intake. More days with  $\max(c_\tau - D_\tau, 0) = 0$  also leads to more days from which jobs can be pulled forward. Hence, instances with more feasible pairs will have a lower total maximum intake, as is reflected in Table 7. This suggests that we are experiencing the effect of  $|\mathcal{I}|$ , i.e. large  $|\mathcal{F}^+(c, D)|$  leads to lower  $|\mathcal{I}|$  and hence faster solution times.

## 6.5 Analysis of Suboptimal Solutions

There were 5 instances, out of the 168 where opt finished solving, in which CS chose the optimal  $y$  but a suboptimal  $p$  for that  $y$ . We now study the  $p$ 's given by CS and those given by opt for the same instances. Recall that CS uses the  $p$ -squash step from S&S, i.e. it uses step 4. from S&S to search for the worst-case distribution over the set defined in step 2. of S&S. The nature of this method is that it results in  $p$  solutions with one  $p_\tau$  being large and the others being smaller. In the instances where CS and S&S were suboptimal, it was in fact optimal to spread the probability more evenly across the days. For example, in one of these instances, CS returned  $p^{\text{CS}} = (0.933, 0.867, 0.8, 0.8, 0.8)$ , where day 1 has high probability. It was optimal, however, to choose  $p^{\text{opt}} = (0.867, 0.867, 0.867, 0.867, 0.8)$ , where 4 days have equal probability. This is the same pattern that we found for S&S, but not for AO. In the cases where AO was  $p$ -suboptimal, it gave  $p$  solutions which had the same spread of probability. However, it gave solutions that were permutations of the ones that opt gave. This is due to the fact that it removes some intakes from the set  $\mathcal{I}$ , thereby approximating the objective function.

We can also study the suboptimal pulling forward ( $y$ ) decisions. Since S&S was the only algorithm that gave a large number of suboptimal  $y$  solutions, we now investigate the reasons behind this behaviour. In the 108 instances where S&S was  $y$ -suboptimal, both S&S and opt pulled forward between the same single pair of days (day 2 to day 1). The only difference was in the number of jobs pulled forward,  $y_{2,1}$ . In Figure 2 we plot the values of  $y_{2,1}^{\text{opt}}$  against  $y_{2,1}^{\text{S\&S}}$ . Since the extended S&S is an ensemble heuristic, we add the algorithm that returned each solution as shown by the colour of the dots.

The plot does not highlight any specific pattern in the points overall, but it does suggest the following three points. Firstly, TD typically pulls forward too little. This is due to the upper bound that TD uses for  $\sum_{l=\max(\tau-K, 1)}^{\tau-1} y_{\tau, l}$ . This upper bound is shown in step 3. of Top Down (Section 4.3.1) as:

$$y_\tau^{\max} = \left\lfloor \max \left\{ c_\tau - D_\tau - i_\tau^f, 0 \right\} \right\rfloor, \quad (37)$$

which is lower than the upper bound given in constraint (21). This corresponds to reserving capacity for intake jobs, and hence reducing the potential for pulling forward. Secondly, PH typically pulls forward too much. As described in step 3(d)i. of PH in Section 4.3.2, the upper bound PH uses for pulling forward to  $\tau_2$  is  $\max\{c_{\tau_2} - D_{\tau_2}, 0\}$ . This upper bound does not





Figure 2: Correct vs. Incorrect  $y_{2,1}$  values, colour coded by the algorithm that gave them

account for intake at all, and the greedy nature of PH means it will always reach this upper bound if possible. Finally, BU was not responsible for any of the solutions in Figure 2. Our results show that BU was never the algorithm responsible for S&S’s solution. BU typically chose to pull forward between day 3 and 1 instead of 2 and 1 when this was possible, and this was never optimal in the instances that we considered.

Figure 3 shows the difference between the expected rollover cost found by S&S and the worst-case expected rollover cost, for each day. Positive values suggests S&S’s expected rollover was higher. Each line in the plot corresponds to the solution for a different instance. Figure 3a shows that pulling forward too little results in less rollover on day 1, due to there being more spare capacity, but more on every subsequent day. Figure 3b shows that pulling forward leads to much higher rollover on day 1, followed by less on the subsequent days.



(a) Cases where S&S pulled forward too little (b) Cases where S&S pulled forward too much

Figure 3: Worst-case expected rollover difference when S&S chose the right pair of days

## 7 Conclusions and Further Research

The general conclusions that we can make from our results for instances where all algorithms finished running are as follows. S&S is the fastest heuristic (54 sec average) but its speed comes

with a reduction in performance compared to the other algorithms. It is not very strong when selecting  $y$  (35% optimal), but it performs very well in selecting the worst-case  $p$  (96% optimal). CS, on the other hand, offers a substantial increase in performance over S&S when choosing  $y$  (98% optimal) and the same performance when choosing  $p$  (96% optimal again), with only a slight increase in time taken (1 minute 51 seconds on average). AO performs the same as CS in choosing  $y$ , and slightly worse in choosing  $p$  (95% optimal), but it takes quite a bit longer (8 min average). We can also draw some conclusions from the results for the cases where opt timed out. Since AO can also time out for the largest of problems, CS is the best algorithm for practical use. It performs very well and takes a small amount of time to generate a solution, i.e. 12 minutes at most. It has been verified that, in every instance, CS finished in 1-2 iterations of the inner loop. It always stopped when it had reached the optimal  $p$  within the set it was given. These two facts along with further testing suggest that increasing  $k^{\max}$  or reducing the optimality tolerance for CS will not improve solutions, and that the only way this can be done is by improving the set of distributions passed to it. However, this would result in larger ambiguity sets which may cause CS to lose some time efficiency.

The conclusions regarding S&S can be summarised as follows. Typically, when S&S chose the wrong  $y$ , TD and PH were capable of choosing the correct pair of days between which pulling forward could be carried out. They were not accurate in selecting the value of this variable, however, with TD typically being too low and PH typically being too high. This was due to the upper bounds given to the two algorithms for the pulling forward variable, with the bound for TD being too conservative and the bound for PH not being conservative enough. We conjecture that constructing the set of extreme distributions before the candidate  $y$  solutions might improve S&S's performance. If we were to do this, then we could use the expected intake when computing  $y$  to avoid unnecessary rollover and/or missing an opportunity for pulling forward. The bounds to use for these algorithms is a subject for future research.

It is true that the algorithms in this paper were created for our problem specifically, but their logic is generalisable to other models. The basic idea of S&S is to reduce the search space for both the outer and inner problems using some intuition about the problem under consideration. The outer problem is solved using an ensemble of basic heuristics, and this could be applied to any DRO problem. The inner model is solved using our observation that the objective is increasing in the distribution's parameters in order to construct a set of extreme distributions. Where we applied this process to the success probabilities, this can be done analogously by considering extreme PMF values instead. For example, if we were to consider general distributions instead of binomial ones, this could be done by constructing a set of distributions with the heaviest upper tails, since these give the highest probabilities to the highest intakes.

The main contribution that we have made to the existing DRO literature is a selection of algorithms resulting from incorporating information about the family in which the distribution of the uncertain parameters lies. In the literature, the distributions are usually considered to be more general and no such assumptions are made. We have shown that this additional information can be very beneficial in constructing algorithms for the DRO model, since CS and S&S both

exploit the behaviour of the binomial distribution in order to produce solutions. This has not only allowed these algorithms to perform very well, but also to produce solutions very quickly to a model with a very large number of constraints and decision variables. It has been noted in the literature that CS suffers runtime issues due to the complexity of the distribution separation problem, and we have presented a version of CS that does not suffer from this problem.

There are a number of natural extensions to our work which would be of further interest from a practical viewpoint. Firstly, this model considers a simplified version of the planning problem in which each job requires one unit of capacity to complete. This is not typically the case in real life, since different jobs will naturally take a different amount of time to complete. Adding more varied completion times would be a clear next step in improving this model. Secondly, the model considers the case where there is only one skill, and is equivalent to assuming all engineers can complete all jobs. In some scenarios this is not the case, and the model could account for this by considering separate demand values and decision variables for each skill. Finally, we have treated the capacity as fixed and aimed to optimise its use. In some cases, if not all, however, capacity can be manipulated in the planning phase using such things as ordering extra units of existing resources (overtime), hiring outside resources for a cost (contractors), and reducing the capacity on certain days (reductions). These ways to manipulate capacity will form the basis of some of our future research.

## Acknowledgements

We would like to acknowledge the support of Engineering and Physical Sciences Research Council funded (EP/L015692/1) STOR-i Centre for Doctoral Training. We would also like to thank BT for their funding, and Mathias Kern and Gilbert Owusu from BT for their support during this project.

## Bibliography

- Ainslie, R., McCall, J., Shakya, S., and Owusu, G. (2018). Tactical plan optimisation for large multi-skilled workforces using a bi-level model. *2018 IEEE Congress on Evolutionary Computation (CEC)*, pages 1–8.
- Ainslie, R. T., Shakya, S., McCall, J., and Owusu, G. (2015). Optimising skill matching in the service industry for large multi-skilled workforces. In Bramer, M. and Petridis, M., editors, *Research and Development in Intelligent Systems XXXII*, pages 231–243, Cham. Springer International Publishing.
- Angalakudati, M., Balwani, S., Calzada, J., Chatterjee, B., Perakis, G., Raad, N., and Uichanco, J. (2014). Business analytics for flexible resource allocation under random emergencies. *Management Science*, 60(6):1552–1573.
- Bansal, M., Huang, K.-L., and Mehrotra, S. (2018). Decomposition algorithms for two-stage



- distributionally robust mixed binary programs. *SIAM Journal on Optimization*, 28(3):2360–2383.
- Bastian, N. D., Lunday, B. J., Fisher, C. B., and Hall, A. O. (2020). Models and methods for workforce planning under uncertainty: Optimizing U.S. army cyber branch readiness and manning. *Omega*, 92:102171.
- Bayraksan, G. and Love, D. K. (2015). Data-driven stochastic programming using phi-divergences. In *The Operations Research Revolution*, chapter 1, pages 1–19. INFORMS.
- Ben-Tal, A., den Hertog, D., De Waegenaere, A., Melenberg, B., and Rennen, G. (2013). Robust solutions of optimization problems affected by uncertain probabilities. *Management Science*, 59(2):341–357.
- Bertsimas, D., Gupta, V., and Kallus, N. (2018). Data-driven robust optimization. *Mathematical Programming*, 167(2):235–292.
- Delage, E. and Ye, Y. (2010). Distributionally robust optimization under moment uncertainty with application to data-driven problems. *Operations Research*, 58(3):595–612.
- Fetter, R. B. (1961). A linear programming model for long range capacity planning. *Management Science*, 7(4):372–378.
- From, S. G. (2020). Some new upper and lower bounds for the Mills ratio. *Journal of Mathematical Analysis and Applications*, 486(1):123872.
- Ghosh, S., Squillante, M., and Wollega, E. (2020). Efficient stochastic gradient descent for learning with distributionally robust optimization. *arXiv preprint. arXiv:1805.08728*.
- Gorissen, B. L., İhsan Yanıkoğlu, and den Hertog, D. (2015). A practical guide to robust optimization. *Omega*, 53:124–137.
- Hanssmann, F. and Hess, S. W. (1960). A linear programming approach to production and employment scheduling. *Management Technology*, 1(1):46–51.
- Hettich, R. and Kortanek, K. O. (1993). Semi-infinite programming: Theory, methods, and applications. *SIAM Review*, 35(3):380–429.
- Holt, C. C., Modigliani, F., and Simon, H. A. (1955). A linear decision rule for production and employment scheduling. *Management Science*, 2(1):1–30.
- J. Abernathy, W., Baloff, N., Hershey, J., and Wandel, S. (1973). A three-stage manpower planning and scheduling model – a service-sector example. *Operations Research*, 21:693–711.
- Kortanek, K. O. and No, H. (1993). A central cutting plane algorithm for convex semi-infinite programming problems. *SIAM Journal on Optimization*, 3(4):901–918.

- Lam, H. and Ghosh, S. (2013). Iterative methods for robust estimation under bivariate distributional uncertainty. In *Proceedings of the 2013 Winter Simulation Conference: Simulation: Making Decisions in a Complex World*, WSC '13, page 193–204. IEEE Press.
- Laporte, G. and Louveaux, F. V. (1993). The integer L-shaped method for stochastic integer programs with complete recourse. *Operations Research Letters*, 13(3):133–142.
- Liu, Y., Yuan, X., Zeng, S., and Zhang, J. (2017). Primal–dual hybrid gradient method for distributionally robust optimization problems. *Operations Research Letters*, 45(6):625–630.
- Luo, F. and Mehrotra, S. (2019). Decomposition algorithm for distributionally robust optimization using Wasserstein metric with an application to a class of regression models. *European Journal of Operational Research*, 278(1):20–35.
- Martel, A. and Price, W. (1978). A normative model for manpower planning under risk. In *Manpower Planning and Organization Design*, pages 291–305. Springer.
- Mehrotra, S. and Papp, D. (2014). A cutting surface algorithm for semi-infinite convex programming with an application to moment robust optimization. *arXiv preprint. arXiv:1306.3437*.
- Pflug, G. and Wozabal, D. (2007). Ambiguity in portfolio selection. *Quantitative Finance*, 7(4):435–442.
- Rahimian, H., Bayraksan, G., and Homem-De-Mello, T. (2019). Identifying effective scenarios in distributionally robust stochastic programs with total variation distance. *Mathematical Programming*, 173(1–2):393–430.
- Rahimian, H. and Mehrotra, S. (2019). Distributionally robust optimization: A review. *arXiv preprint. arXiv:1908.05659*.
- Ross, E. (2016). *Cross-trained Workforce Planning Models*. PhD thesis, University of Lancaster.
- Sun, H., Wang, Y., and Xue, Y. (2021). A bi-objective robust optimization model for disaster response planning under uncertainties. *Computers & Industrial Engineering*, 155:107213.
- Zhu, X. and Sherali, H. D. (2009). Two-stage workforce planning under demand fluctuations and uncertainty. *Journal of the Operational Research Society*, 60(1):94–103.

# Appendices

## A Optimal Solutions for Trivial Cases of the Two-day Model

Define the two sets:

$$c_{\text{spare}} = \{\max\{c_1 - D_1 - i_1, 0\} \forall i_1 \in \mathcal{I}_1\}, \quad (38)$$

$$d_{\text{inc}} = \{\max\{i_2 - (c_2 - D_2), 0\} \forall i_2 \in \mathcal{I}_2\}, \quad (39)$$

as the set of potential values of the spare capacity on day 1 before pulling forward and number of incomplete jobs on day 2 before pulling forward. We have the following results.

**Theorem 2.** *In the two-day model, if  $D_2 < \min(c_{\text{spare}})$  then  $y = D_2$  is an optimal solution.*

*Proof.* Since  $D_2 \geq 0$  this means that  $\min(c_{\text{spare}}) > 0$  and hence  $c_1 - D_1 - i_1^{\max} > 0$ . Hence:

$$D_2 < \min(c_{\text{spare}}) = c_1 - D_1 - i_1^{\max} \leq c_1 - D_1. \quad (40)$$

Therefore, the constraint on  $y$  is  $y \leq \min\{D_2, c_1 - D_1\} = D_2$  and thus  $y = D_2$  is feasible and we cannot pull forward more than  $D_2$ . Recall that, for any  $i \in \mathcal{I}$ :

$$R_1^i = \max\{i_1 - (c_1 - D_1 - y), 0\} = \max\{i_1 + y - (c_1 - D_1), 0\} \quad (41)$$

and we have:

$$i_1 + y - (c_1 - D_1) \leq i_1^{\max} + D_2 - (c_1 - D_1) \quad (42)$$

$$= D_2 - (c_1 - D_1 - i_1^{\max}) \quad (43)$$

$$= D_2 - \min(c_{\text{spare}}) \quad (44)$$

$$< 0 \quad \forall i \in \mathcal{I}. \quad (45)$$

Here, (42) is because  $i_1 \leq i_1^{\max}$  and  $y \leq D_2$ , and (45) is because  $D_2 < \min(c_{\text{spare}})$ . This means that  $R_1^i = 0 \forall i \in \mathcal{I}$ , and  $\forall y \in \{0, \dots, D_2\}$ . Hence,  $\mathbb{E}_p(R_1) = 0 \forall p \in \mathcal{U}^p$  and  $\forall y \in \{0, \dots, D_2\}$ . This means that the  $y$  with the lowest worst-case expected rollover cost on day 2 is optimal. Since  $R_1 = 0$  for any  $y$ , we have that:

$$R_2^i = \max\{i_2 - (c_2 - D_2) - y, 0\}$$

and clearly  $R_2^i$  is decreasing in  $y$  for any  $i \in \mathcal{I}$ . Hence, choosing  $y$  to be its highest value, i.e.  $D_2$ , must lead to the lowest  $R_2^i$  for all  $i \in \mathcal{I}$ , and hence the lowest expected day 2 rollover cost for any distribution  $p$ . It follows that  $y = D_2$  gives the lowest worst-case expected day 2 rollover cost and is hence optimal.  $\square$

**Theorem 3.** *In the two-day model, if  $\min(c_{\text{spare}}) > \max(d_{\text{inc}})$  then  $y = \min\{\max(d_{\text{inc}}), D_2\}$  is an optimal solution.*

*Proof.* If  $\min(c_{\text{spare}}) > \max(d_{\text{inc}})$  then, since  $\max(d_{\text{inc}}) \geq 0$ , we have  $\min(c_{\text{spare}}) > 0$  and so

$$\min(c_{\text{spare}}) = c_1 - D_1 - i_1^{\max}.$$

Also let us note that, if  $y = \min\{\max(d_{\text{inc}}), D_2\}$  then  $y \leq D_2$ . Also, since  $\max(d_{\text{inc}}) < \min(c_{\text{spare}})$  we also have:

$$y \leq \max(d_{\text{inc}}) < \min(c_{\text{spare}}) = c_1 - D_1 - i_1^{\max} \leq c_1 - D_1.$$

This means that  $y \leq \min\{c_1 - D_1, D_2\}$  and so  $y$  is feasible. Now, if  $\max(d_{\text{inc}}) = 0$  then we have  $i_2^{\max} - (c_2 - D_2) \leq 0$ , and hence

$$i_2 - (c_2 - D_2) - y \leq i_2^{\max} - (c_2 - D_2) \leq 0.$$

This means  $R_2^i = 0$  for all  $i \in \mathcal{I}$  and all  $y$ . If we take  $y = 0 = \min\{\max(d_{\text{inc}}), D_2\}$ , then we also have  $R_1^i = 0$  for all  $i \in \mathcal{I}$ , since  $\min(c_{\text{spare}}) > 0$ . Thus,  $\sum_{\tau=1}^2 \mathbb{E}_p(R_\tau) = 0$  for all  $p \in \mathcal{U}^p$ , and notably for the worst-case  $p$  as well. This means  $y = \min\{\max(d_{\text{inc}}), D_2\}$  is optimal, since the worst-case expected rollover cost cannot be less than zero.

If  $\max(d_{\text{inc}}) > 0$  then  $i_2^{\max} - (c_2 - D_2) > 0$ . Consider again the choice  $y = \min\{\max(d_{\text{inc}}), D_2\}$ . Then  $y \leq \max(d_{\text{inc}}) < \min(c_{\text{spare}})$ , so we have

$$i_1 + y - (c_1 - D_1) \leq \min(c_{\text{spare}}) + i_1^{\max} - (c_1 - D_1), \quad (46)$$

$$= 0, \quad (47)$$

for all  $i \in \mathcal{I}$ . Hence,  $R_1^i = 0 \forall i \in \mathcal{I}$  and  $\mathbb{E}_p(R_1) = 0 \forall p \in \mathcal{U}^p$ . Furthermore, if  $D_2 > \max(d_{\text{inc}})$  then  $y = \max(d_{\text{inc}})$  and:

$$i_2 - (c_2 - D_2) - y \leq \max(d_{\text{inc}}) - y = 0 \forall i \in \mathcal{I}. \quad (48)$$

This means that:

$$R_2^i = \max\{i_2 - (c_2 - D_2) - y, 0\} = 0 \forall i \in \mathcal{I}.$$

Hence,  $\mathbb{E}_p(R_2) = 0 \forall p \in \mathcal{U}^p$ . Therefore, both expected rollover costs are zero in the worst case and so our choice of  $y$  is optimal. Finally, if  $D_2 < \max(d_{\text{inc}})$  then  $y = D_2$ . Hence,  $y$  cannot be increased any further, and reducing it could only cause some values of  $R_2^i$  to be higher, while all  $R_1^i$  remain at zero. Hence,  $R_2^i$  cannot be decreased further for any  $i \in \mathcal{I}$ . Therefore,  $R_2^i$  is also minimised for all  $i \in \mathcal{I}$ , and thus  $\mathbb{E}_p(R_2)$  is minimised for all  $p \in \mathcal{U}^p$ . Again, the total expected rollover cost is minimised for all distributions, and so our choice of  $y$  is optimal.  $\square$

## B Proofs of Results in Section 3.6

### B.1 Proof of Lemma 1

*Proof.* This follows from common sense; if we pull forward any less than  $\min(c_{\text{spare}})$  then one of two things must happen. Firstly, we could be missing an opportunity to pull forward a job from day 2 that will be incomplete. Hence, the solution will have more rollover and hence is not optimal. If this is not the case, then we are missing the opportunity to pull forward a job that will currently be completed on day 2. Hence, the rollover from this solution is the same as the one resulting from pulling forward  $\min(c_{\text{spare}})$ . If we pull forward any more than  $\max(c_{\text{spare}})$ , then we will cause jobs to roll over from day 1. If we pull forward more than  $\max(d_{\text{inc}})$ , then we will be pulling forward a job that would otherwise not have rolled over. This job will either roll over on day 1, causing more total rollover than pulling forward  $\max(d_{\text{inc}})$ , or not roll over on day 1, causing the same rollover as pulling forward  $\max(d_{\text{inc}})$ . Hence, pulling forward  $y \notin \mathcal{Y}$  cannot have a lower rollover cost than pulling forward  $y \in \mathcal{Y}$ , so the optimal  $y$  must lie in  $\mathcal{Y}$ .  $\square$

### B.2 Proof of Lemma 2

We now prove Lemma 2 by taking a normal approximation to the binomial distribution and taking derivatives of the resulting expected values.

#### B.2.1 Normal Approximation of Binomial Distribution

Since the binomial PMF is not “nice” to work with in respect to taking derivatives and proving properties such as monotonicity, we use a normal approximation to the binomial distribution when proving optimality of our algorithm. For large  $i_\tau^{\max}$  we can assume that  $I_\tau \sim N(i_\tau^{\max}p_\tau, i_\tau^{\max}p_\tau(1-p_\tau))$  approximately. In order to calculate an expression for the expected rollover, we define the incomplete jobs via equations (49)-(50).

$$r_1 = I_1 - (c_1 - D_1 - y) = I_1 - c_1^r, \quad (49)$$

$$r_2 = I_2 - (c_2 - D_2 + y) = I_2 - c_2^r, \quad (50)$$

so that  $R_1 = \max\{r_1, 0\}$  and  $R_2 = \max\{r_2 + R_1, 0\}$ . It is then clear that  $r_\tau$  is normal for  $\tau \in \{1, 2\}$ . Furthermore, we can write  $R_1$  as a mixture of zero with probability  $\mathbb{P}(r_1 \leq 0)$  and a *zero-truncated normal* distribution (i.e.  $r_1|r_1 > 0$ ) with probability  $\mathbb{P}(r_1 > 0)$ . Defining:

$$\mu_1 = i_1^{\max}p_1 - c_1^r, \quad (51)$$

$$\sigma_1^2 = \text{Var}(I_1), \quad (52)$$

as the mean and variance of  $r_1$ , using the formula for the expected value of a left-truncated normal distribution truncated at zero, we can find that the expected value of  $R_1$  is given by:

$$\mathbb{E}_p(R_1) = \mathbb{P}(r_1 > 0) \times \left( \mu_1 + \sigma_1 \frac{\phi(\alpha_1)}{1 - \Phi(\alpha_1)} \right) \quad (53)$$

$$= (1 - \Phi(\alpha_1))\mu_1 + \phi(\alpha_1)\sigma_1 \quad (54)$$

$$= \Phi(-\alpha_1)\mu_1 + \phi(\alpha_1)\sigma_1 \quad (55)$$

where  $\alpha_1 = -\frac{\mu_1}{\sigma_1}$ ,  $\phi$  is the PDF of the standard normal distribution and  $\Phi$  is the CDF of said distribution. By (11), we have that  $I_1 + I_2 \leq c_1^r + c_2^r$  with probability 1, or equivalently  $r_1 + r_2 \leq 0$  with probability 1. Hence, we obtain:

$$\mathbb{P}(R_2 > 0) = \mathbb{P}(\max(r_2 + R_1, 0) > 0) \quad (56)$$

$$= \mathbb{P}(r_2 + R_1 > 0) \quad (57)$$

$$= \mathbb{P}(r_2 + \max(r_1, 0) > 0) \quad (58)$$

$$= \mathbb{P}(r_1 > 0 \cap r_2 + r_1 > 0) + \mathbb{P}(r_1 \leq 0 \cap r_2 > 0) \quad (59)$$

$$= \mathbb{P}(r_1 > 0 \cap r_2 + r_1 > 0) + \mathbb{P}(r_1 \leq 0)\mathbb{P}(r_2 > 0) \quad (60)$$

$$= \mathbb{P}(r_1 \leq 0)\mathbb{P}(r_2 > 0) \quad (61)$$

$$= (1 - \mathbb{P}(r_1 > 0))\mathbb{P}(r_2 > 0) \quad (62)$$

$$= \mathbb{P}(r_2 > 0) - \mathbb{P}(r_1 > 0 \cap r_2 > 0) \quad (63)$$

$$= \mathbb{P}(r_2 > 0). \quad (64)$$

Here, equation (59) uses the law of total probability conditioning on  $r_1 > 0$  and  $r_1 \leq 0$  and (61) uses that  $r_1 + r_2 \leq 0$  with probability 1, meaning that the event  $\{r_1 > 0 \cap r_1 + r_2 > 0\}$  has probability zero. Therefore, we have:

$$\mathbb{E}_p(R_2) = \mathbb{P}(R_2 > 0) \times \mathbb{E}_p(R_2 | R_2 > 0) \quad (65)$$

$$= \mathbb{P}(r_2 > 0) \times \left( \mu_2 + \sigma_2 \frac{\phi(\alpha_2)}{1 - \Phi(\alpha_2)} \right) \quad (66)$$

$$= \Phi(-\alpha_2)\mu_2 + \phi(\alpha_2)\sigma_2, \quad (67)$$

where  $\mu_2, \sigma_2, \alpha_2$  are defined in the same way as for day 1. Note that these expressions show that  $\mathbb{E}_p(R_2)$  is independent of  $p_1$  and  $\mathbb{E}_p(R_1)$  is independent of  $p_2$ .

### B.2.2 Proof for Day 1 Expected Rollover

We now use derivatives to prove that the expected day 1 rollover is increasing in  $p_1$ . The derivative of  $\mathbb{E}_p(R_1)$  is given by:

$$\frac{d}{dp_1} \mathbb{E}_p(R_1) = \left( \Phi(-\alpha_1)i_1^{\max} - \mu_1 \frac{d\alpha_1}{dp_1} \phi(-\alpha_1) \right) + \left( \phi(\alpha_1) \frac{i_1^{\max}}{2\sigma_1} (1 - 2p_1) - \sigma_1 \alpha_1 \frac{d\alpha_1}{dp_1} \phi(\alpha_1) \right) \quad (68)$$

$$= \left( \Phi(-\alpha_1)i_1^{\max} - \mu_1 \frac{d\alpha_1}{dp_1} \phi(\alpha_1) \right) + \left( \phi(\alpha_1) \frac{i_1^{\max}}{2\sigma_1} (1 - 2p_1) + \mu_1 \frac{d\alpha_1}{dp_1} \phi(\alpha_1) \right) \quad (69)$$

$$= \Phi(-\alpha_1)i_1^{\max} + \phi(\alpha_1) \frac{i_1^{\max}}{2\sigma_1} (1 - 2p_1), \quad (70)$$

where (69) uses the symmetry of the normal distribution, i.e. that  $\phi(-\alpha_1) = \phi(\alpha_1)$ . Clearly, when  $p_1 \in [0, \frac{1}{2}]$  this derivative is non-negative. Hence, consider  $p_1 \in (\frac{1}{2}, 1]$ . Since  $1 - 2p_1 \geq -1$  when  $p_1 > \frac{1}{2}$ , we have:

$$g_1(p_1) := \frac{d}{dp_1} \mathbb{E}_p(R_1) \geq i_1^{\max} \left[ \Phi(-\alpha_1) - \frac{\phi(\alpha_1)}{2\sigma_1} \right]. \quad (71)$$

We can further split  $(\frac{1}{2}, 1]$  into two parts based on when  $\alpha_1 > 0$  and when  $\alpha_1 \leq 0$ . Since  $\alpha_1 = -\frac{\mu_1}{\sigma_1}$ ,  $\alpha_1 > 0$  when  $\mu_1 < 0$  i.e.  $p_1 < p_1^{\text{crit}} := \frac{c_1^r}{i_1^{\max}}$ . We then have  $\alpha_1 \leq 0$  when  $p_1 \geq p_1^{\text{crit}}$ .

We discuss the case where  $p_1 \geq p_1^{\text{crit}}$  first. If  $p_1 \geq p_1^{\text{crit}}$  then this must mean that  $p_1^{\text{crit}} \leq 1$  and hence  $i_1^{\text{max}} \geq c_1^r$ . By finding the derivative of  $-\alpha_1$  to be:

$$\frac{d}{dp_1} \frac{\mu_1}{\sigma_1} = \frac{i_1^{\text{max}}}{2\sigma_1^3} (p_1(i_1^{\text{max}} - 2c_1^r) + c_1^r). \quad (72)$$

we find that it is increasing in  $p_1$  when  $i_1^{\text{max}} \geq c_1^r$ , as this implies that:

$$p_1(i_1^{\text{max}} - 2c_1^r) + c_1^r \geq p_1(-c_1^r) + c_1^r = c_1^r(1 - p_1) > 0. \quad (73)$$

Since  $\Phi$  is a CDF, it is increasing in  $-\alpha_1$  and hence  $\Phi(-\alpha_1)$  is increasing in  $p_1$ . Therefore, since  $p_1 \geq p_1^{\text{crit}}$  we have:

$$\Phi(-\alpha_1) \geq \Phi(-\alpha_1 | p_1 = p_1^{\text{crit}}) = \Phi(0) = \frac{1}{2}. \quad (74)$$

We also have that:

$$\frac{\phi(\alpha_1)}{2\sigma_1} = \frac{1}{2} \times \left[ \frac{1}{\sqrt{2\pi\sigma_1^2}} \exp\left(-\frac{1}{2} \left(\frac{0 - \mu_1}{\sigma_1}\right)^2\right) \right] = \frac{1}{2} f_{r_1}(0) \leq \frac{1}{2}, \quad (75)$$

where  $f_{r_1}$  is the PDF of  $r_1 \sim N(\mu_1, \sigma_1)$ . Combining (74) and (75), we have  $g_1(p_1) \geq 0$  for  $p_1 \in [p_1^{\text{crit}}, 1]$ . The only remaining part of the proof for day 1 is the region  $p_1 \in [\frac{1}{2}, p_1^{\text{crit}})$ .

For this part of the proof we use the *Mills ratio*, defined in (76). The Mills ratio is a known function which has proven upper and lower bounds written in terms of its argument  $z$ .

$$M(z) = \frac{1 - \Phi(z)}{\phi(z)}, \quad z > 0. \quad (76)$$

Using this, we have:

$$g_1(p_1) = i_1^{\text{max}} \phi(\alpha_1) \left[ M(\alpha_1) + \frac{1 - 2p_1}{2\sigma_1} \right], \quad (77)$$

and hence the proof reduces to showing that  $M(\alpha_1) \geq -\frac{1-2p_1}{2\sigma_1}$  for  $p_1 \in [\frac{1}{2}, p_1^{\text{crit}})$ . We can prove this result under one condition. From the paper on bounds on the Mills ratio by From (2020), we have:

$$M(z) > \frac{2}{\sqrt{z^2 + 4} + z}, \quad x \geq 0. \quad (78)$$

Note that, for  $\frac{1}{2} \leq p_1 \leq p_1^{\text{crit}}$ , we have  $\mu_1 \leq 0$  and hence  $\alpha_1 = -\frac{\mu_1}{\sigma_1} \geq 0$ . Using the above, we have:

$$\frac{d}{dp_1} \mathbb{E}_p(R_1) \geq i_1^{\text{max}} \phi(\alpha_1) \left[ M(\alpha_1) - \frac{1}{2\sigma_1} \right] \quad (79)$$

$$> i_1^{\text{max}} \phi(\alpha_1) \left[ \frac{2}{\sqrt{\alpha_1^2 + 4} + \alpha_1} - \frac{1}{2\sigma_1} \right] \quad (80)$$

$$= i_1^{\text{max}} \phi(\alpha_1) \left[ \frac{2}{\sqrt{\left(\frac{\mu_1}{\sigma_1}\right)^2 + 4} - \frac{\mu_1}{\sigma_1}} - \frac{1}{2\sigma_1} \right] \quad (81)$$

$$= i_1^{\text{max}} \phi(\alpha_1) \left[ \frac{2\sigma_1}{\sqrt{\mu_1^2 + 4\sigma_1^2} - \mu_1} - \frac{1}{2\sigma_1} \right] \quad (82)$$

$$= i_1^{\max} \phi(\alpha_1) \left[ \frac{4\sigma_1^2 - \sqrt{\mu_1^2 + 4\sigma_1^2} + \mu_1}{2\sigma_1 (\sqrt{\mu_1^2 + 4\sigma_1^2} - \mu_1)} \right] \quad (83)$$

and we can write:

$$\mu_1^2 + 4\sigma_1^2 = (\mu_1 - 2\sigma_1)^2 + 4\mu_1\sigma_1 \quad (84)$$

$$< (\mu_1 - 2\sigma_1)^2 \quad (85)$$

since  $\mu_1 < 0$ . Hence:

$$\sqrt{\mu_1^2 + 4\sigma_1^2} < |\mu_1 - 2\sigma_1| = 2\sigma_1 - \mu_1. \quad (86)$$

Therefore, we get:

$$\frac{d}{dp_1} \mathbb{E}_p(R_1) \geq i_1^{\max} \phi(\alpha_1) \left[ \frac{4\sigma_1^2 - (2\sigma_1 - \mu_1) + \mu_1}{2\sigma_1 (\sqrt{\mu_1^2 + 4\sigma_1^2} - \mu_1)} \right] \quad (87)$$

$$= i_1^{\max} \phi(\alpha_1) \left[ \frac{4\sigma_1^2 - 2\sigma_1 + 2\mu_1}{2\sigma_1 (\sqrt{\mu_1^2 + 4\sigma_1^2} - \mu_1)} \right] \quad (88)$$

Hence, since the denominator is positive, we analyse the function given by:

$$h_1(p_1) = 4\sigma_1^2 - 2\sigma_1 + 2\mu_1.$$

In fact, we have:

$$h_1(p_1) = 4i_1^{\max} p_1(1 - p_1) - 2\sigma_1 + 2(i_1^{\max} p_1 - c_1^r) \quad (89)$$

$$= 4i_1^{\max} p_1 \left( \frac{3}{2} - p_1 \right) - 2\sigma_1 - 2c_1^r \quad (90)$$

$$> 4i_1^{\max} \frac{1}{2} \left( \frac{3}{2} - \frac{1}{2} \right) - 2\sigma_1 - 2c_1^r \quad (91)$$

$$= 2i_1^{\max} - 2\sigma_1 - 2c_1^r \quad (92)$$

$$> 2(i_1^{\max} - c_1^r) - \sqrt{i_1^{\max}}. \quad (93)$$

Here, (90) is due to the fact that  $p_1 \left( \frac{3}{2} - p_1 \right)$  is concave for  $p_1 \in \left[ \frac{1}{2}, p_1^{\text{crit}} \right]$  and has its minimum when  $p_1 = \frac{1}{2}$ . Also, (93) is true because  $\sigma_1 < \sqrt{i_1^{\max} \times \frac{1}{2} \times \left( 1 - \frac{1}{2} \right)} = \frac{\sqrt{i_1^{\max}}}{2}$ . Hence, we have proved that  $\mathbb{E}_p(R_1)$  is increasing in  $p_1$  for  $p_1 \in \left[ \frac{1}{2}, p_1^{\text{crit}} \right]$  if:

$$i_1^{\max} - c_1^r \geq \frac{\sqrt{i_1^{\max}}}{2}. \quad (94)$$

Using the definition for  $c_1^r$ , this corresponds to:

$$y \geq (c_1 - D_1) - \left( i_1^{\max} - \frac{\sqrt{i_1^{\max}}}{2} \right). \quad (95)$$



### B.2.3 Proof for Day 2 Expected Rollover

From Appendix B.2.1 we have:

$$\mathbb{E}_p(R_2) = \Phi(-\alpha_2)\mu_2 + \phi(\alpha_2)\sigma_2, \quad (96)$$

and we wish to prove  $\mathbb{E}_p(R_2)$  is increasing in  $p_2$  for any fixed  $p_1$ . Similarly to in Section B.2.2, we find that:

$$g_2(p_2) = \frac{d}{dp_2}\mathbb{E}_p(R_2) = \Phi(-\alpha_2)i_2^{\max} + \phi(\alpha_2)\frac{i_2^{\max}}{2\sigma_2}(1 - 2p_2) \quad (97)$$

Again, when  $p_2 \leq \frac{1}{2}$  it is clear that  $g_2(p_2)$  is increasing in  $p_2$ , and so we consider  $p_2 > \frac{1}{2}$ . We have

$$g_2(p_2) \geq \Phi(-\alpha_2)i_2^{\max} - \phi(\alpha_2)\frac{i_2^{\max}}{2\sigma_2}, \quad (98)$$

and we define the point at which  $\alpha_2 = -\frac{\mu_2}{\sigma_2} = 0$  as:

$$p_2^{\text{crit}} = \frac{c_2^r}{i_2^{\max}}, \quad (99)$$

and from our  $y$  range we have that

$$y \leq \max(d_{\text{inc}}) = \max\{\max\{i_2 - (c_2 - D_2), 0\} \mid i_2 \in \mathcal{I}_2\} = \max\{i_2^{\max} - (c_2 - D_2), 0\}.$$

If  $\max\{i_2^{\max} - (c_2 - D_2), 0\} = 0$  then  $i_2^{\max} < c_2 - D_2$  and hence no day 2 rollover is possible. This means that  $R_2 = 0$  and is hence trivially increasing since it is constant. Hence we consider the cases where:

$$0 \leq y \leq i_2^{\max} - (c_2 - D_2)$$

and therefore  $i_2^{\max} \leq c_2^r$  and so  $p_2^{\text{crit}} \geq 1$ . Hence, we always have  $p_2 \leq p_2^{\text{crit}}$  and so  $\mu_2 \leq 0$ . Given that  $p_2 \leq p_2^{\text{crit}}$ , we use the Mills ratio approach from Appendix B.2.2 to get:

$$g_2(p_2) \geq i_2^{\max}\phi(\alpha_2) \left[ M(\alpha_2) - \frac{1}{2\sigma_2} \right] \quad (100)$$

$$> i_2^{\max}\phi(\alpha_2) \left[ \frac{2}{\sqrt{\alpha_2^2 + 4} + \alpha_2} - \frac{1}{2\sigma_2} \right] \quad (101)$$

$$= i_2^{\max}\phi(\alpha_2) \left[ \frac{4\sigma_2^2 - \sqrt{\mu_2^2 + 4\sigma_2^2} + \mu_2}{2\sigma_2(\sqrt{\mu_2^2 + 4\sigma_2^2} - \mu_2)} \right] \quad (102)$$

and we have:

$$\mu_2^2 + 4\sigma_2^2 = (\mu_2 - 2\sigma_2)^2 + 4\mu_2\sigma_2 \quad (103)$$

$$\leq (\mu_2 - 2\sigma_2)^2 \quad (104)$$

since  $\mu_2 \leq 0$ . Hence:

$$\sqrt{\mu_2^2 + 4\sigma_2^2} \leq |\mu_2 - 2\sigma_2| = 2\sigma_2 - \mu_2. \quad (105)$$

This means we have:

$$\frac{d}{dp_2}\mathbb{E}_p(R_2) \geq i_2^{\max}\phi(\alpha_2) \left[ \frac{4\sigma_2^2 - (2\sigma_2 - \mu_2) + \mu_2}{2\sigma_2(\sqrt{\mu_2^2 + 4\sigma_2^2} - \mu_2)} \right] \quad (106)$$

$$= i_2^{\max} \phi(\alpha_2) \left[ \frac{4\sigma_2^2 - 2\sigma_2 + 2\mu_2}{2\sigma_2 (\sqrt{\mu_2^2 + 4\sigma_2^2} - \mu_2)} \right] \quad (107)$$

and again we study the function:

$$h_2(p_2) = 4\sigma_2^2 - 2\sigma_2 + \mu_2 \quad (108)$$

$$= 4i_2^{\max} p_2 \left( \frac{3}{2} - p_2 \right) - 2\sigma_2 - 2c_2^r \quad (109)$$

$$> 4i_2^{\max} \frac{1}{2} \left( \frac{3}{2} - \frac{1}{2} \right) - 2\sigma_2 - 2c_2^r \quad (110)$$

$$> 2(i_2^{\max} - c_2^r) - \sqrt{i_2^{\max}} \quad (111)$$

$$= 2i_2^{\max} - \sqrt{i_2^{\max}} - 2(c_2 - D_2) - 2y \quad (112)$$

$$= 2(D_2 - c_2) + 2i_2^{\max} - \sqrt{i_2^{\max}} - 2y. \quad (113)$$

Thus,  $g_2(p_2)$  is increasing in  $p_2$  if:

$$y \leq (D_2 - c_2) + \left( i_2^{\max} - \frac{1}{2} \sqrt{i_2^{\max}} \right). \quad (114)$$

### B.3 Proof of Theorem 1

*Proof.* For any  $y \in \tilde{\mathcal{Y}}$ , both expected rollovers are increasing in their respective success probabilities by Lemma 2. This means that the maximum total expected rollover cost must occur when at least one of the two probabilities is at its maximum, i.e. it occurs at some  $p \in \mathcal{P}_{\max}$ , otherwise neither is at its maximum and hence the sum of the two cannot be either. Given that one of  $p_1, p_2$  is at its max, since the other day's expected rollover is increasing in its own success probability and independent of the other, it must be optimal to set the remaining  $p_\tau$  as high as possible. Hence, the worst-case distribution lies in  $\mathcal{U}_{\max}^p$  and will therefore be found by S&S.  $\square$

## C Bottom Up Algorithm

The BU algorithm is as follows:

1. Initialise  $y_{\tau_1, \tau_2} = 0$  for all  $\tau_1, \tau_2$ .
2. Set  $\tau = L - 1$ .
3. Define  $y_{\tau}^{\max} = \lfloor \max\{c_{\tau} - D_{\tau} - i_{\tau}^f, 0\} \rfloor$  and  $k = \min\{L - \tau, K\}$ .
4. While  $y_{\tau}^{\max} > 0$  and set  $k \geq 1$  and  $\tau + k \leq L$ :
  - (a) Set  $y_{\tau+k, \tau} = \min \left\{ \max\{D_{\tau+k} - c_{\tau+k}, 0\} - \sum_{l=1}^K y_{\tau+k, \tau+k-l}, y_{\tau}^{\max} \right\}$ .
  - (b)  $y_{\tau}^{\max} = y_{\tau}^{\max} - y_{\tau+k, \tau}$ .
  - (c)  $k = k - 1$ .
5. If  $\tau > 1$  set  $\tau = \tau - 1$  and go to 3.
6. Return  $y$ .

This algorithm may perform well in cases where there are significant spikes of demand at the end of the plan, which need to be smoothed as much as possible.

## D Large Results Tables

### D.1 Results by $|\mathcal{U}^p|$

$ \mathcal{U}^p $	Count	Algorithm	Avg. $p$ -APG	Avg. $y$ -APG	Min t.t.	Avg. t.t.	Max t.t.
16	62	opt	0.0%	0.0%	0:00:05.25	0:01:46.22	0:07:23.93
		S&S	0.0%	0.9841%	0:00:00.34	0:00:55.59	0:08:21.85
		CS	0.0%	0.0%	0:00:02.29	0:01:53.8	0:11:25.06
		AO	0.0211%	0.0002%	0:00:02.76	0:00:19.72	0:01:20.18
32	31	opt	0.0%	0.0%	0:00:10.38	0:03:03.62	0:11:51.53
		S&S	0.0%	0.8742%	0:00:00.34	0:00:12.49	0:01:00.22
		CS	0.0%	0.0%	0:00:02.11	0:01:24.59	0:06:40.94
		AO	0.0%	0.0004%	0:00:05.61	0:00:30.47	0:01:39.89
512	31	opt	0.0%	0.0%	0:03:25.97	0:52:51.43	3:17:24.39
		S&S	0.0%	0.693%	0:00:02.37	0:01:51.87	0:09:11.77
		CS	0.0%	0.0%	0:00:04	0:02:45.95	0:12:09.45
		AO	0.0004%	0.0%	0:02:22.88	0:11:39.66	0:32:45.82
562	31	opt	0.0%	0.0%	0:03:55.46	0:59:29.46	3:41:37.82
		S&S	0.0%	0.8669%	0:00:01.22	0:00:57.77	0:04:40.75
		CS	0.0%	0.0%	0:00:02.92	0:02:00.37	0:08:45.78
		AO	0.0%	0.0%	0:02:16.57	0:09:47.41	0:25:24.44
3883	13	opt	0.0%	0.0%	1:12:21.91	1:30:54.48	1:47:25.8
		S&S	0.0685%	1.4263%	0:00:04.53	0:00:06.04	0:00:07.36
		CS	0.0793%	0.0044%	0:00:05.28	0:00:08.73	0:00:12.08
		AO	0.0%	0.0%	0:48:02.39	0:58:15.19	1:07:21.78

Table 6: Summary of results and times taken by size of  $\mathcal{U}^p$

Table 6 is referred to in Section 6.3.

## D.2 Results by $|\mathcal{F}^+(c, D)|$

$ \mathcal{F}^+(c, D) $	Count	Mean $ \mathcal{I} $	Algorithm	Avg. $p$ -APG	Avg. $y$ -APG	Avg. t.t.
0	25	20000.00	opt	0.0%	0.0%	1:26:56.75
			S&S	0.0%	0.0%	0:04:38.47
			CS	0.0%	0.0%	0:08:43.75
			AO	0.038%	0.0%	0:12:12.08
1	25	8192.00	opt	0.0%	0.0%	0:37:29.34
			S&S	0.0%	1.7741%	0:01:01.83
			CS	0.0%	0.0%	0:02:42.02
			AO	0.0093%	0.0%	0:07:08.66
2	40	2440.12	opt	0.0%	0.0%	0:11:05.17
			S&S	0.0%	0.3281%	0:00:11.58
			CS	0.0%	0.0%	0:00:27.98
			AO	0.0%	0.0007%	0:03:22.21
3	78	505.46	opt	0.0%	0.0%	0:16:51.48
			S&S	0.0114%	1.2504%	0:00:02.57
			CS	0.0132%	0.0007%	0:00:05.18
			AO	0.0018%	0.0%	0:10:46.12

Table 7: Results by  $|\mathcal{F}^+(c, D)|$

Table 7 is referred to in Section 6.4.

### D.3 Results by $|\mathcal{I}|$

$ \mathcal{I} $	Count	Algorithm	Avg. $p$ -APG	Avg. $y$ -APG	Min t.t.	Avg. t.t.	Max t.t.
392	18	opt	0.0%	0.0%	0:00:05.25	0:13:25.58	1:12:43.51
		S&S	0.0064%	1.066%	0:00:00.34	0:00:02.04	0:00:05.09
		CS	0.0064%	0.0032%	0:00:02.11	0:00:03.9	0:00:08.03
		AO	0.0%	0.0%	0:00:02.76	0:09:14.97	0:50:59.18
512	30	opt	0.0%	0.0%	0:00:07.81	0:16:44.74	1:35:58.35
		S&S	0.0124%	1.2578%	0:00:00.43	0:00:02.62	0:00:06.94
		CS	0.0147%	0.0%	0:00:03.07	0:00:05.27	0:00:12.08
		AO	0.0008%	0.0%	0:00:03.26	0:10:22.41	0:59:37.25
567	30	opt	0.0%	0.0%	0:00:07.77	0:19:01.75	1:47:25.8
		S&S	0.0135%	1.3536%	0:00:00.49	0:00:02.84	0:00:07.36
		CS	0.0158%	0.0%	0:00:03.36	0:00:05.85	0:00:11.93
		AO	0.0038%	0.0%	0:00:03.31	0:12:04.52	1:07:21.78
2187	15	opt	0.0%	0.0%	0:00:34.98	0:09:45.96	0:24:39
		S&S	0.0%	0.5093%	0:00:02.2	0:00:10.25	0:00:20.01
		CS	0.0%	0.0%	0:00:15.83	0:00:22.85	0:00:33.64
		AO	0.0%	0.0018%	0:00:07.57	0:03:12.07	0:08:29.45
2592	25	opt	0.0%	0.0%	0:00:43.01	0:11:52.69	0:29:57.85
		S&S	0.0%	0.2195%	0:00:02.65	0:00:12.38	0:00:24.58
		CS	0.0%	0.0%	0:00:19.11	0:00:31.06	0:00:55.29
		AO	0.0%	0.0%	0:00:08.75	0:03:28.29	0:09:50.88
8192	25	opt	0.0%	0.0%	0:02:18.52	0:37:29.34	1:34:39.83
		S&S	0.0%	1.7741%	0:00:12.73	0:01:01.83	0:02:00.29
		CS	0.0%	0.0%	0:01:30.43	0:02:42.02	0:04:45.69
		AO	0.0093%	0.0%	0:00:22.52	0:07:08.66	0:18:56.39
20000	25	opt	0.0%	0.0%	0:05:33.23	1:26:56.75	3:41:37.82
		S&S	0.0%	0.0%	0:00:44.08	0:04:38.47	0:09:11.77
		CS	0.0%	0.0%	0:04:59.26	0:08:43.75	0:12:09.45
		AO	0.038%	0.0%	0:00:46.7	0:12:12.08	0:32:45.82

Table 8: Summary of results and times taken by size of  $\mathcal{I}$

Table 8 is referred to in Section 6.3.

## E Results for Instances Where opt Timed Out

There were 18 instances in which opt timed out, and in 5 of these instances AO also timed out. CS and S&S never timed out. All of the instances where opt timed out were those with  $n_{\text{probs}} = 15$  and  $\gamma = 0.15$ , i.e. they had the largest set of distributions. The sets of intakes were always above 2000 in size in these instances, and the intake set was always of size 20000 when AO timed out. In all of the instances where the algorithms timed out, they timed out before they even started solving. Hence, no solution is available from these algorithms in these cases. However, in those instances where the heuristics finished running, we can still brute-force compute the worst-case distribution for each chosen  $y$ . The results for the 13 instances in which only opt timed out are summarised in Table 9. The results for the 5 where both opt and AO timed out are summarised in Table 10.

Algorithm	Avg. Worst-case Objective	Avg. $p$ -APG	Min t.t.	Avg. t.t.	Max t.t.
S&S	84.55	0.002997%	0:00:25.73	0:01:19.87	0:02:40.07
CS	84.33	0.003511%	0:00:28.54	0:02:01.55	0:04:30.47
AO	84.30	0.000218%	1:54:05.16	2:46:11.82	4:04:41.95

Table 9: Summary of instances where only opt timed out.

Algorithm	Avg. Worst-case Objective	Avg. $p$ -APG	Min t.t.	Avg. t.t.	Max t.t.
S&S	2.0	0.090396%	0:11:36.86	0:11:39.86	0:11:44.75
CS	2.0	0.090396%	0:11:26.1	0:11:31.59	0:11:36.38

Table 10: Summary of instances where AO and opt both timed out.

Table 9 shows that all algorithms found  $y$ -decisions with worst-case expected costs that were quite close to each other on average, and that they were all very close to choosing the worst-case  $p$  on average. AO found the best solutions on average, but in significantly more time. Table 10 reflects that, in those 5 instances, S&S and CS returned the same solutions. These tables highlight the potential time savings by using S&S or CS. Just looking at Table 9, we see that S&S and CS took between 1 and 2 minutes on average, where AO took almost 3 hours. Table 10 shows that, in cases where AO timed out, the other two heuristics returned a solution in 12 minutes on average.

## F Tables of Notation

### F.1 General Model Notation

Notation	Meaning
$L$	Number of days in a plan
$K$	Maximum number of days a job can be pulled forward
$\tau, \tau_1, \tau_2$	A day in the plan, value in $\{1, \dots, L\}$
$y_{\tau_1, \tau_2}$	Number of jobs to pull forward from day $\tau_1 \in \{2, \dots, L\}$ to $\tau_2 \in \{\max \tau_1 - K, 1, \dots, \tau_1 - 1\}$ .
$R_\tau$	Number of jobs to roll over from day $\tau$ to $\tau + 1$ .
$a_\tau$	Cost of a job rolling over from day $\tau$ to $\tau + 1$ .
$c_\tau$	Number of hours of capacity available on day $\tau$ .
$D_\tau$	Number of jobs currently due on day $\tau$ .
$\mathbb{N}_0$	Set of non-negative integers.
$I_\tau$	Random variable representing number of jobs arriving between the time of planning and day $\tau$ that will be due on day $\tau$ (intake).
$i_\tau$	Realisation of $I_\tau$ .
$R^i$	Realisation of $R = (R_1, \dots, R_L)$ corresponding to realisation $i$ of $I$ .
$\mathcal{I}_\tau$	Set of all possible realisations of $I_\tau$ .
$\mathcal{I}$	Set of all possible realisations of the vector of intakes $I$ .
$q$	A discrete probability distribution over the set of intakes $\mathcal{I}$ .
$i_\tau^{\max}$	The maximum value $I_\tau$ can take.
$p_\tau$	Success probability parameter of the binomial distribution of intake $I_\tau$ .
$\mathcal{U}$	Set of all probability distributions $q$ that are binomial.
$\mathcal{U}^p$	Set of vectors $p$ obtained from a distribution $q$ in $\mathcal{U}$ .
$\mathcal{U}_\alpha^p$	Set of $ps$ such that the probability under $p$ of total intake going above total spare capacity is no more than $\alpha$ .

Table 11: General model notation from Section 4.



## F.2 S&S Notation

Notation	Meaning
$y_\tau^{\max}$	Maximum number of jobs that can be pulled forward to $\tau$ under S&S algorithm.
$k$	Number of days after $\tau$ that we are currently considering pulling forward jobs to $\tau$ from.
$\mathcal{L}_P$	Partition of $\{1, \dots, L\}$ into pairs inside the Pull Half Forward Algorithm.
$r_\tau$	Number of baseline incomplete jobs on $\tau$ before accounting for intake.
$r_{(\tau_1, \tau_2)}$	Total number of baseline incomplete jobs on $\tau_1$ and $\tau_2$ before accounting for intake.
$\bar{r}_P$	Average number of baseline incomplete jobs per pair of days.
$\mathcal{P}_\tau^{\max}$	Set of probability vectors $p$ such that the probability $p_\tau$ for day $\tau$ is the highest it can be while $p$ is inside $\mathcal{U}^P$ .
$\mathcal{P}_\tau^{\text{ext}}$	Set of (extreme) probability vectors $p \in \mathcal{P}_\tau^{\max}$ such that the sum of $p$ is the largest.
$\mathcal{P}^{\text{ext}}$	Union of $\mathcal{P}_\tau^{\text{ext}}$ over $\tau = 1, \dots, L$ , set of probability vectors $p$ corresponding to the most extreme probability distributions $q$ .
$\mathcal{X}$	Set of algorithms $x$ used in generating a $y$ solution in the multi-period S&S algorithm.
$\mathcal{Y}$	Set of candidate $y$ solutions generated by S&S algorithm.
$C_p$	Expected rollover cost for the chosen $y$ given the distribution of intake is given by $p$ .
$p_y$	Probability vector $p$ that gives the highest value of $C_p$ for the solution $y$ .
$y^*$	$y$ solution in $\mathcal{Y}$ that gave the lowest maximum expected cost $C_{p_y}$ in S&S algorithm.
$x^*, p^*$	Algorithm and success probability that led to this minimum cost.

Table 12: Notation used in S&S algorithm (Section 4.3)

### F.3 CS/AO Notation

Notation	Meaning
$k$ (repeated dummy variable)	Index for the iteration of CS algorithm we are currently carrying out.
$k_{\max}$	Maximum number of iterations of CS algorithm allowed to run.
$\mathcal{U}^{p,k}$	Current subset of $\mathcal{U}^p$ being used at iteration $k$ of CS.
$y^k$	Pulling forward decision generated by solving outer problem at iteration $k$ of CS.
$p^k$	Probability vector generated by solving distribution separation problem at iteration $k$ of CS.
$\varepsilon$	Optimality tolerance of CS algorithm.
$t^k$	Objective value of problem obtained by solving outer problem at iteration $k$ of CS.
$\beta$	Minimum probability an intake must have of occurring in order to be used in the AO algorithm.
$\tilde{\mathcal{I}}$	Set of intakes with probability of occurring higher than $\beta$ .

Table 13: Notation used in CS/AO Algorithms (Section 4.4)

## F.4 Input Parameter and Results Notation

Notation	Meaning
$\mathcal{F}$	Set of pairs of days between which pulling forward is allowed.
$\mathcal{F}^+(c, D)$	Set of pairs of days between which pulling forward is feasible given $c$ and $D$ .
$n(i^{\max})$	Number of days with maximum intake higher than remaining capacity given $i^{\max}$ , $c$ , and $D$ .
$n_{\text{probs}}$	Number of values each probability in $p$ can take in our discretised ambiguity set.
$i^f$	Forecast of intake.
$p^f$	Probability vector resulting from $i^f$ assuming $i^f = i^{\max} p^f$ .
$\ \cdot\ _2$	$\ell_2$ norm.
$\gamma$	Maximum standardised distance from the forecast $p^f$ we allow a vector $p \in \mathcal{U}^p$ to be.

Table 14: Input parameter notation used in Section 5

Notation	Meaning
$f(y, p)$	Shorthand for expected rollover cost given pulling forward decision $y$ and distribution parameter $p$ .
$x$	An algorithm, namely in {S&S, CS, AO}.
$y^x, p^x$	$y, p$ solution obtained by algorithm $x$ .
$g_p(y^x, p^x)$	$p$ -gap of algorithm $x$ 's solution. The difference between the worst-case expected cost for $y^x$ and the expected cost obtained by the algorithm.
$z^*$	Overall optimal objective value.
$g_p(y^x)$	$y$ -gap. Difference between worst-case expected cost for $y^x$ over all distributions and the overall optimal objective value.

Table 15: Results analysis notation from Section 6