# The Core Conjecture of Hilton and Zhao II: a Proof

Yan Cao<sup>∗</sup> Guantao Chen† Guangming Jing‡ Songling Shan§

November 27, 2024

#### Abstract

A simple graph G with maximum degree  $\Delta$  is *overfull* if  $|E(G)| > \Delta |V(G)|/2$ . The core of G, denoted  $G_{\Delta}$ , is the subgraph of G induced by its vertices of degree  $\Delta$ . Clearly, the chromatic index of G equals  $\Delta + 1$  if G is overfull. Conversely, Hilton and Zhao in 1996 conjectured that if G is a simple connected graph with  $\Delta \geq 3$  and  $\Delta(G_{\Delta}) \leq 2$ , then  $\chi'(G) = \Delta + 1$  implies that G is overfull or  $G = P^*$ , where  $P^*$  is obtained from the Petersen graph by deleting a vertex. Cariolaro and Cariolaro settled the base case  $\Delta = 3$  in 2003, and Cranston and Rabern proved the next case  $\Delta = 4$  in 2019. In this paper, we give a proof of this conjecture for all  $\Delta \geq 4$ .

MSC (2010): Primary 05C15

Keywords: Overfull graph, Multifan, Kierstead path, Pseudo-multifan, Lollipop.

### 1 Introduction

Let G be a simple graph with maximum degree  $\Delta$ . The *core* of G, denoted  $G_{\Delta}$ , is the subgraph of G induced by its vertices of degree  $\Delta$ . If  $|E(G)| > \Delta \, |V(G)|/2$ , then G is overfull. Overfull graphs are class 2. The graph  $P^*$ , obtained from the Petersen graph by deleting one vertex, is also known to be class 2. Conversely, in 1996, Hilton and Zhao [\[5\]](#page-26-0) proposed the following conjecture.

<span id="page-0-0"></span>**Conjecture 1.1** (Core Conjecture). Let G be a simple connected graph with maximum degree  $\Delta \geq 3$  and  $\Delta(G_{\Delta}) \leq 2$ . Then G is class 2 implies that G is overfull or  $G = P^*$ .

<sup>∗</sup>Department of Mathematics, West Virginia University, Morgantown, WV 26506, USA. yacao@mail.wvu.edu.

<sup>†</sup>Department of Mathematics and Statistics, Georgia State University, Atlanta, GA 30302, USA. gchen@gsu.edu. Partially supported by NSF grant DMS-1855716.

<sup>‡</sup>Department of Mathematics, Augusta University, Augusta, GA 30912, USA. gjing@augusta.edu. Partially supported by NSF grant DMS-2001130.

<sup>§</sup>Department of Mathematics, Illinois State Univeristy, Normal, IL 61790, USA. sshan12@ilstu.edu. Partially supported by the NSF-AWM Mentoring Travel Grant 1642548 and by the New Faculty Initiative Grant of Illinois State University.

As a class 2 graph of maximum degree 2 is an odd cycle and odd cycles are overfull, if true, the Core Conjecture implies that for connected graphs G with  $\Delta(G_{\Delta}) \leq 2$ , determining whether G is class 2 can be done by checking whether  $|E(G)| > \Delta \lfloor |V(G)|/2 \rfloor$  if  $G \neq P^*$ . We call a connected class 2 graph G with  $\Delta(G_{\Delta}) \leq 2$  an HZ-graph. A first breakthrough of the Core Conjecture was achieved in 2003, when Cariolaro and Cariolaro [\[2\]](#page-26-1) settled the base case  $\Delta = 3$ . They proved that  $P^*$  is the only HZ-graph with maximum degree  $\Delta = 3$ , an alternative proof was given later by Král', Sereny, and Stiebitz (see  $[6, pp. 67–63]$ ). The next case  $\Delta = 4$  was recently solved by Cranston and Rabern [\[3\]](#page-26-3): they proved that the only HZ-graph with maximum degree  $\Delta = 4$  is the graph  $K_5$  with one edge removed. In this paper, we confirm the Core Conjecture for all HZ-graphs G with  $\Delta \geq 4$ . It worth mentioning that our proof implies a polynomial-time algorithm that, given  $G$  with maximum degree  $\Delta \geq 4$  and  $\Delta(G_{\Delta}) \leq 2$ , finds an optimal edge coloring of G.

<span id="page-1-0"></span>**Theorem 1.2.** Let G be a connected graph with maximum degree  $\Delta \geq 4$  and  $\Delta(G_{\Delta}) \leq 2$ . Then G is class 2 if and only if G is overfull.

Since every overfull graph is class 2, we will only prove the "only if" statement in Theorem [1.2.](#page-1-0) The remainder of the paper is organized as follows. In next section, we prove Theorem [1.2](#page-1-0) by applying Theorems [2.3](#page-2-0) to [2.5.](#page-2-1) In Section 3, we give necessary definitions and list results from [\[1\]](#page-26-4). Theorems [2.3](#page-2-0) to [2.5](#page-2-1) will be proved in Sections 4, 5, and 6, respectively.

# 2 Proof of Theorem [1.2](#page-1-0)

In this section, we prove Theorem [1.2](#page-1-0) by applying Theorems [2.3](#page-2-0) to [2.5.](#page-2-1) We start with some concepts. For two integers p and q, let  $[p, q] = \{i \in \mathbb{Z} : p \leq i \leq q\}$ . An edge k-coloring of G is a mapping  $\varphi$  from  $E(G)$  to [1, k], called *colors*, such that no two adjacent edges receive the same color. We denote by  $\mathcal{C}^k(G)$  the set of all edge k-colorings of G. The *chromatic* index  $\chi'(G)$  of G is the smallest k so that G has an edge k-coloring. The symbol  $\Delta$  is reserved for  $\Delta(G)$ , the maximum degree of G throughout this paper.

Let G be a graph,  $v \in V(G)$ , and  $i \geq 0$  be an integer. An *i*-vertex is a vertex of degree i in G, and an i-vertex from the neighborhood of v is called an i-neighbor of v. Define

$$
V_i = \{ w \in V(G) : d_G(w) = i \}, \qquad N_i(v) = N_G(v) \cap V_i, \quad \text{and} \quad N_i[v] = N_i(v) \cup \{ v \}.
$$

For  $X \subseteq V(G)$ , let  $N_G(X) = \bigcup_{x \in X} N_G(x)$  and  $N_i(X) = N_G(X) \cap V_i$ .

Let  $e \in E(G)$  and  $\varphi \in C^k(G-e)$  for some integer  $k \geq 0$ . The set of colors present at v is  $\varphi(v) = {\varphi(f) : f \text{ is incident to } v}$ , and the set of colors missing at v is  $\overline{\varphi}(v) = [1, k] \setminus \varphi(v)$ . If  $\overline{\varphi}(v) = {\alpha}$  is a singleton for some  $\alpha \in [1, k]$ , we also write  $\overline{\varphi}(v) = \alpha$ . For  $X \subseteq V(G)$ , let  $\overline{\varphi}(X) = \bigcup_{x \in X} \overline{\varphi}(x)$ . The set X is  $\varphi$ -elementary if  $\overline{\varphi}(x) \cap \overline{\varphi}(y) = \emptyset$  for any distinct  $x, y \in X$ .

An edge  $e \in E(G)$  is a critical edge of G if  $\chi'(G - e) < \chi'(G)$ , and G is edge  $\Delta$ -critical or simply  $\Delta$ -critical if G is connected,  $\chi'(G) = \Delta + 1$ , and every of its edge is critical. The following result by Hilton and Zhao in [\[4\]](#page-26-5) reveals certain properties of an HZ graph.

<span id="page-2-2"></span>**Lemma 2.1.** If G is an HZ-graph with maximum degree  $\Delta$ , then the following holds.

- (a) G is  $\Delta$ -critical and  $G_{\Delta}$  is 2-regular.
- (b)  $\delta(G) = \Delta 1$ , or  $\Delta = 2$  and G is an odd cycle.
- (c) Every vertex of G has at least two neighbors in  $G_{\Delta}$ .

Let  $\Delta \geq 4$  and let  $\mathcal{O}_{\Delta}$  be the set of all graphs obtained from two graphs  $H_1$  and  $H_2$ by adding all edges between  $V(H_1)$  and  $V(H_2)$ , where  $H_1$  is any 2-regular graph on  $n_1$ vertices,  $H_2$  is any  $(\Delta - 1 - n_1)$ -regular graph on  $(\Delta - 2)$  vertices, and  $n_1 \in [3, \Delta - 1]$  such that  $n_1 + (\Delta - 2)$  is odd. Stiebitz et al. showed that Conjecture [1.1](#page-0-0) is equivalent to the conjecture below.

<span id="page-2-5"></span>**Conjecture 2.2** ([\[6,](#page-26-2) Conjecture 4.10]). If G is an HZ-graph with maximum degree  $\Delta$ , then either  $G \in \mathcal{O}_\Delta$ , or  $\Delta = 2$  and G is an odd cycle, or  $\Delta = 3$  and  $G = P^*$ .

We will prove this equivalent form of the Core Conjecture for  $\Delta \geq 4$  by applying the following results.

<span id="page-2-3"></span><span id="page-2-0"></span>**Theorem 2.3.** If G is an HZ-graph with maximum degree  $\Delta \geq 4$ , then the following two statements hold.

- <span id="page-2-6"></span>(i) For any two adjacent vertices  $u, v \in V_{\Delta}$ ,  $N_{\Delta-1}(u) = N_{\Delta-1}(v)$ .
- (ii) For any  $r \in V_{\Delta}$ , there exist  $s \in N_{\Delta-1}(r)$  and  $\varphi \in C^{\Delta}(G rs)$  such that  $N_{\Delta-1}[r]$  is  $\varphi$ -elementary.

For an HZ-graph G with maximum degree  $\Delta \geq 4$ , each component of  $G_{\Delta}$  is a cycle by Lemma [2.1.](#page-2-2) So Theorem [2.3](#page-2-0) [\(i\)](#page-2-3) implies that  $N_{\Delta-1}(x) = N_{\Delta-1}(y)$  for any two vertices x, y from the same cycle of  $G_{\Delta}$ .

<span id="page-2-4"></span>**Theorem 2.4.** If G is an HZ-graph with maximum degree  $\Delta \geq 4$ , then for any two adjacent vertices  $x, y \in V_{\Delta-1}, N_{\Delta}(x) = N_{\Delta}(y)$ .

<span id="page-2-1"></span>**Theorem 2.5.** Let G be an HZ-graph with maximum degree  $\Delta \geq 7$  and  $u, r \in V_{\Delta}$ . If  $N_{\Delta-1}(u) \neq N_{\Delta-1}(r)$  and  $N_{\Delta-1}(u) \cap N_{\Delta-1}(r) \neq \emptyset$ , then  $|N_{\Delta-1}(u) \cap N_{\Delta-1}(r)| = \Delta - 3$ , i.e.  $|N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)| = |N_{\Delta-1}(r) \setminus N_{\Delta-1}(u)| = 1.$ 

<span id="page-2-7"></span>Corollary 2.6. If G is an HZ-graph with maximum degree  $\Delta \geq 7$  and there exist  $u, v \in V_{\Delta}$ such that  $N_{\Delta-1}(u) \neq N_{\Delta-1}(v)$ , then  $V_{\Delta-1}$  is an independent set in G.

**Proof.** Assume to the contrary that there exist  $x, y \in V_{\Delta-1}$  such that  $xy \in E(G)$ . By Lemma [2.1,](#page-2-2) there exists  $w \in N_{\Delta}(x)$ . By the assumption that there exist  $u, v \in V_{\Delta}$  such that  $N_{\Delta-1}(u) \neq N_{\Delta-1}(v)$ , there exists some  $w' \in V_{\Delta}$  such that  $N_{\Delta-1}(w) \neq N_{\Delta-1}(w')$ . We may further assume that the distance between  $w$  and  $w'$  in  $G$  is shortest among all pairs

of vertices  $w_1$  and  $w'_1$  such that  $w_1 \in N_\Delta(x)$  and  $N_{\Delta-1}(w_1) \neq N_{\Delta-1}(w'_1)$ . We claim that  $N_{\Delta-1}(w) \cap N_{\Delta-1}(w') \neq \emptyset$ . Let P be a shortest path connecting w and w' in G. By the choice of w and  $w'$ ,  $(V(P) \cap V_{\Delta}) \setminus \{w, w'\}$  contains no vertex  $w^*$  such that  $N_{\Delta-1}(w^*) = N_{\Delta-1}(w)$ . Consequently,  $V_{\Delta} \cap V(P) = \{w, w'\}$ . Since  $N_{\Delta-1}(w) \neq N_{\Delta-1}(w')$ , it follows that w and w' are not on the same cycle of  $G_{\Delta}$  and so  $ww' \notin E(G)$  by Theorem [2.3](#page-2-0) (i). Thus  $P - \{w, w'\}$ has at least one vertex. By Theorem [2.4,](#page-2-4) all vertices of  $P - \{w, w'\}$  have in G the same set of neighbors from  $V_{\Delta}$ . Thus, both w and w' are  $\Delta$ -neighbors of each vertex from  $P - \{w, w'\}$ and so  $N_{\Delta-1}(w) \cap N_{\Delta-1}(w') \neq \emptyset$ . By Theorem [2.5,](#page-2-1) we have  $|N_{\Delta-1}(w) \cap N_{\Delta-1}(w')| = \Delta - 3$ , which together with Theorem [2.4](#page-2-4) implies  $x, y \in N_{\Delta-1}(w) \cap N_{\Delta-1}(w')$ .

Let  $N_{\Delta-1}(w')\setminus N_{\Delta-1}(w) = \{z\}$ . We claim that  $N_{\Delta-1}(z) = \emptyset$ . For otherwise, let  $z' \in$  $N_{\Delta-1}(z)$ . Clearly  $z' \neq z$ . By Theorem [2.4,](#page-2-4)  $z' \in N_{\Delta-1}(w') \setminus N_{\Delta-1}(w)$ , giving a contradiction to  $N_{\Delta-1}(w')\setminus N_{\Delta-1}(w) = \{z\}.$  We then claim that  $N_{\Delta}(z) \subseteq N_{\Delta}(x)$ . For otherwise let  $w^* \in N_\Delta(z) \setminus N_\Delta(x)$ . As  $x \in N_{\Delta-1}(w')$  and  $x \notin N_{\Delta-1}(w^*)$ , it follows that  $w^* \neq w'$ . Since  $z \in N_{\Delta-1}(w^*) \cap N_{\Delta-1}(w')$ , it follows that  $|N_{\Delta-1}(w^*) \cap N_{\Delta-1}(w')| \geq \Delta - 3$  by Theorem [2.5](#page-2-1) (it can happen that  $N_{\Delta-1}(w^*) = N_{\Delta-1}(w')$ ). Thus  $N_{\Delta-1}(w^*) \cap N_{\Delta-1}(w')$  contains at least one of x and y as  $x, y \in N_{\Delta-1}(w')$ . As  $xy \in E(G)$ , we have  $x, y \in N_{\Delta-1}(w^*) \cap N_{\Delta-1}(w')$  by Theorem [2.4.](#page-2-4) This gives a contradiction to the choice of  $w^*$ . Therefore we have  $N_{\Delta-1}(z) = \emptyset$ and  $N_{\Delta}(z) \subseteq N_{\Delta}(x)$ . However,  $d_G(z) \leq |N_{\Delta}(x)| < |N_{\Delta}(x) \cup \{y\}| \leq d_G(x)$ , contradicting  $d_G(x) = d_G(z) = \Delta - 1$ . This completes the proof.  $\Box$ 

We now prove Conjecture [2.2](#page-2-5) for  $\Delta \geq 4$  as below.

#### **Theorem 2.7.** If G is an HZ-graph with maximum degree  $\Delta \geq 4$ , then  $G \in \mathcal{O}_{\Delta}$ .

*Proof.* Assume to the contrary that there exists an HZ-graph  $G$  with maximum degree  $\Delta \geq 4$  such that  $G \notin \mathcal{O}_{\Delta}$ . Let  $n = |V(G)|$ . First assume that  $N_{\Delta-1}(u) = N_{\Delta-1}(v)$  for every pair  $u, v \in V_{\Delta}$ . Then  $V_{\Delta}$ ,  $V_{\Delta-1}$  and the edges between them form a complete bipartite graph. Since  $G \notin \mathcal{O}_{\Delta}$ , it follows that n is even. Let  $r \in V_{\Delta}$ . The assumption above also implies that  $N_{\Delta-1}(r) = V_{\Delta-1}$ . By Theorem [2.3](#page-2-0) [\(ii\)](#page-2-6), there exist  $s \in N_{\Delta-1}(r) = V_{\Delta-1}$  and  $\varphi \in C^{\Delta}(G - rs)$  such that  $N_{\Delta-1}[r] = V_{\Delta-1} \cup \{r\}$  is  $\varphi$ -elementary, which thereby implies that  $V(G)$  is  $\varphi$ -elementary. Therefore, each color in  $\overline{\varphi}(N_{\Delta-1}[r])$  is missed at exactly one vertex in  $V(G)$ , showing that n is odd. This is a contradiction.

We now assume that there exist  $u, v \in V_{\Delta}$  such that  $N_{\Delta-1}(u) \neq N_{\Delta-1}(v)$ . We further assume that  $N_{\Delta-1}(u) \cap N_{\Delta-1}(v) \neq \emptyset$  (using the same argument to find u and v as for finding w and w' in the proof of Corollary [2.6\)](#page-2-7). By Theorem [2.3](#page-2-0) [\(i\)](#page-2-3), the cycle  $C_u$  containing u and the cycle  $C_v$  containing v from  $G_{\Delta}$  are distinct. Let  $w \in N_{\Delta-1}(u) \cap N_{\Delta-1}(v)$ . Then  $d_G(w) \geq |V(C_u)| + |V(C_v)| \geq 6$  by Theorem [2.3](#page-2-0) [\(i\)](#page-2-3). Thus  $\Delta = d_G(w) + 1 \geq 7$ . Applying Corollary [2.6,](#page-2-7) it follows that  $V_{\Delta-1}$  is an independent set of G.

Let  $A \subseteq V_{\Delta}$  be the set of all vertices a satisfying  $N_{\Delta-1}(a) = N_{\Delta-1}(u)$ , and let  $B \subseteq V_{\Delta}$ be the set of all vertices b satisfying  $N_{\Delta-1}(b) \neq N_{\Delta-1}(u)$  and  $N_{\Delta-1}(b) \cap N_{\Delta-1}(u) \neq \emptyset$ . Clearly  $u \in A$  and  $v \in B$ , so A and B are non-empty. Partition B into non-empty subsets  $B_1, B_2, \ldots, B_t$  such that for each  $i \in [1, t]$ , all vertices in  $B_i$  have the same neighborhood in  $V_{\Delta-1}$ . By Theorem [2.3](#page-2-0) [\(i\)](#page-2-3), each of  $A, B_1, B_2, \ldots, B_t$  induces a union of disjoint cycles in  $G_{\Delta}$ . So  $|A| \geq 3$  and  $|B_i| \geq 3$  for each  $i \in [1, t]$ .

Now we claim  $t \geq \Delta - 2$ . Assume otherwise  $t \leq \Delta - 3$ . Since for each  $i \in [1, t]$ ,  $|N_{\Delta - 1}(A)\rangle$  $N_{\Delta-1}(B_i)|=1$  by Theorem [2.5](#page-2-1) and  $|N_{\Delta-1}(A)|=\Delta-2$ , there exists  $z \in N_{\Delta-1}(A)$  such that  $z \notin N_{\Delta-1}(A) \setminus N_{\Delta-1}(B_i)$  for each  $i \in [1, t]$ , or equivalently,  $z \in N_{\Delta-1}(A) \cap (\bigcap_{i=1}^t N_{\Delta-1}(B_i)).$ Let  $z' \in N_{\Delta-1}(A) \setminus N_{\Delta-1}(B_1)$ . Then

$$
|A| + \sum_{1 \le i \le t} |B_i| = d_G(z) = d_G(z') \le |A| + \sum_{2 \le i \le t} |B_i|,
$$

achieving a contradiction. Hence  $t \geq \Delta - 2$ .

We now achieve a contradiction to the assumption  $\Delta \geq 7$  by counting the number of edges in G between  $N_{\Delta-1}(A)$  and  $A \cup B$ . Note that  $|N_{\Delta-1}(A)| = \Delta - 2$ . Since each vertex in B has exactly  $\Delta - 3$  neighbors in  $N_{\Delta - 1}(A)$  and  $|B_i| \geq 3$  for each  $i \in [1, t]$ , we have

$$
|E_G(A \cup B, N_{\Delta-1}(A))| = |A|(\Delta - 2) + |\cup_{i=1}^t B_i|(\Delta - 3) \ge 3(\Delta - 2) + 3t(\Delta - 3) \ge 3(\Delta - 2)^2.
$$

On the other hand, since  $N_{\Delta-1}(A)$  is an independent set and every vertex in it has degree  $\Delta - 1$  in G, we have

$$
|E_G(A \cap B, N_{\Delta - 1}(A))| = (\Delta - 1)(\Delta - 2).
$$

Since  $\Delta \geq 2$ , solving  $\Delta$  in  $(\Delta - 1)(\Delta - 2) \geq 3(\Delta - 2)^2$  gives  $\Delta \leq 2.5$ , achieving a desired  $\Box$ contradiction.

### 3 Definitions and previous results

In this section, we recall essential concepts from [\[1\]](#page-26-4) and list a number of results that we will use as lemmas in the proof of Theorems [2.3](#page-2-0) to [2.5.](#page-2-1)

Let G be a graph,  $e \in E(G)$ ,  $\varphi \in C^k(G - e)$  for some  $k \geq 0$ , and let  $\alpha, \beta \in [1, k]$ . Each component of  $G - e$  induced on edges colored by  $\alpha$  or  $\beta$  is either a path or an even cycle, which is called an  $(\alpha, \beta)$ -chain of  $G - e$  with respect to  $\varphi$ . Interchanging  $\alpha$  and  $\beta$  on an  $(\alpha, \beta)$ -chain C of G gives a new edge k-coloring, which is denoted by  $\varphi$ /C. This operation is called a Kempe change.

For  $x, y \in V(G)$ , if x and y are contained in the same  $(\alpha, \beta)$ -chain, we say x and y are  $(\alpha, \beta)$ -linked with respect to  $\varphi$ . Otherwise, they are  $(\alpha, \beta)$ -unlinked. If an  $(\alpha, \beta)$ -chain P is a path with one endvertex as x, we also denote it by  $P_x(\alpha, \beta, \varphi)$  and just write  $P_x(\alpha, \beta)$ if  $\varphi$  is understood. For a vertex u and an edge uv contained in  $P_x(\alpha, \beta, \varphi)$ , we write  $u \in P_x(\alpha, \beta, \varphi)$  and  $uv \in P_x(\alpha, \beta, \varphi)$ . If  $u, v \in P_x(\alpha, \beta, \varphi)$  such that u lies between x and v on P, then we say that  $P_x(\alpha, \beta, \varphi)$  meets u before v.

Let T be an alternating sequence of vertices and edges of G. We denote by  $V(T)$  the set of vertices contained in T, and by  $E(T)$  the set of edges contained in T. We simply write  $\overline{\varphi}(T)$  for  $\overline{\varphi}(V(T))$ . If  $V(T)$  is  $\varphi$ -elementary, then for a color  $\tau \in \overline{\varphi}(T)$ , we denote by  $\overline{\varphi}_T^{-1}$  $T^I(\tau)$ the unique vertex in  $V(T)$  at which  $\tau$  is missed. A coloring  $\varphi' \in C^k(G-e)$  is  $(T, \varphi)$ -stable if for every  $x \in V(T)$  and every  $f \in E(T)$ , it holds that  $\overline{\varphi}'(x) = \overline{\varphi}(x)$  and  $\varphi'(f) = \varphi(f)$ . Clearly,  $\varphi$  is  $(T, \varphi)$ -stable, and if  $\varphi_1 \in C^k(G-e)$  is  $(T, \varphi)$ -stable, and  $\varphi_2 \in C^k(G-e)$  is  $(T, \varphi_1)$ -stable, then  $\varphi_2$  is also  $(T, \varphi)$ -stable.

#### 3.1 Multifan

Let G be a graph,  $rs_1 \in E(G)$  and  $\varphi \in C^k(G - rs_1)$  for some  $k \geq 0$ . A multifan centered at r with respect to  $rs_1$  and  $\varphi$  is a sequence

$$
F_{\varphi}(r,s_1:s_p) := (r,rs_1,s_1,rs_2,s_2,\ldots,rs_p,s_p)
$$

with  $p \geq 1$  consisting of distinct vertices and edges such that for every edge  $rs_i$  with  $i \in [2, p]$ , there is a vertex  $s_j$  with  $j \in [1, i - 1]$  satisfying  $\varphi(rs_i) \in \overline{\varphi}(s_j)$ . The following result can be found in [\[6,](#page-26-2) Theorem 2.1].

<span id="page-5-2"></span><span id="page-5-0"></span>**Lemma 3.1.** Let G be a class 2 graph and  $F_{\varphi}(r, s_1 : s_p)$  be a multifan with respect to  $rs_1$ and  $\varphi \in C^{\Delta}(G - rs_1)$ . Then the following statements hold.

- <span id="page-5-1"></span>(a)  $V(F)$  is  $\varphi$ -elementary.
- (b) For any  $\alpha \in \overline{\varphi}(r)$  and any  $\beta \in \overline{\varphi}(s_i)$  with  $i \in [1, p]$ , r and  $s_i$  are  $(\alpha, \beta)$ -linked with respect to  $\varphi$ .

Let  $F_{\varphi}(r, s_1 : s_p)$  be a multifan. We call  $s_{\ell_1}, s_{\ell_2}, \ldots, s_{\ell_h}$ , a subsequence of  $s_2, \ldots, s_p$ , an  $\alpha$ -inducing sequence for some  $\alpha \in [1, k]$  with respect to  $\varphi$  and  $F$  if  $\varphi(rs_{\ell_1}) = \alpha \in \overline{\varphi}(s_1)$  and  $\varphi(rs_{\ell_i}) \in \overline{\varphi}(s_{\ell_{i-1}})$  for each  $i \in [2, h]$ . (By this definition,  $(r, rs_1, s_1, rs_{\ell_1}, s_{\ell_1}, \ldots, rs_{\ell_h}, s_{\ell_h})$ is also a multifan with respect to  $rs_1$  and  $\varphi$ .) A color in  $\overline{\varphi}(s_{\ell_i})$  for any  $i \in [1, h]$  is an  $\alpha$ -inducing color and is induced by  $\alpha$ . For  $\alpha_i \in \overline{\varphi}(s_{\ell_i})$  and  $\alpha_j \in \overline{\varphi}(s_{\ell_j})$  with  $i < j$  and  $i, j \in [1, h]$ , we write  $\alpha_i \prec \alpha_j$ . For convenience,  $\alpha$  itself is an  $\alpha$ -inducing color and is induced by  $\alpha$ , and  $\alpha \prec \beta$  for any  $\beta \in \overline{\varphi}(s_{\ell_i})$  and any  $i \in [1, h]$ . An  $\alpha$ -inducing color  $\beta$  is called a *last*  $\alpha$ *-inducing color* if there does not exist any  $\alpha$ -inducing color  $\delta$  such that  $\beta \prec \delta$ .

By Lemma [3.1](#page-5-0) (a), each color in  $\overline{\varphi}(F) \setminus \overline{\varphi}(r)$  is induced by a unique color in  $\overline{\varphi}(s_1)$ . Also if  $\alpha_1$  and  $\alpha_2$  are two distinct colors in  $\overline{\varphi}(s_1)$ , then an  $\alpha_1$ -inducing sequence is disjoint with an  $\alpha_2$ -inducing sequence. The following result is a consequence of Lemma [3.1](#page-5-0) (a).

<span id="page-5-3"></span>**Lemma 3.2** ([1, Lemma 3.2]). Let G be a class 2 graph and  $F_{\varphi}(r, s_1 : s_p)$  be a multifan with respect to rs<sub>1</sub> and  $\varphi \in C^{\Delta}(G - rs_1)$ . For any two colors  $\delta, \lambda$  with  $\delta \in \overline{\varphi}(s_i)$  and  $\lambda \in \overline{\varphi}(s_i)$ for some distinct  $i, j \in [1, p]$ , the following statements hold.

- <span id="page-6-1"></span><span id="page-6-0"></span>(a) If  $\delta$  and  $\lambda$  are induced by different colors from  $\overline{\varphi}(s_1)$ , then  $s_i$  and  $s_j$  are  $(\delta, \lambda)$ -linked with respect to  $\varphi$ .
- (b) If  $\delta$  and  $\lambda$  are induced by the same color from  $\overline{\varphi}(s_1)$  such that  $\delta \prec \lambda$  and  $s_i$  and  $s_j$  are  $(\delta, \lambda)$ -unlinked with respect to  $\varphi$ , then  $r \in P_{s_j}(\lambda, \delta, \varphi)$ .

By Lemma [2.1](#page-2-2) (a), every edge of an HZ graph is critical. For an HZ-graph  $G$  with maximum degree  $\Delta \geq 3$ , we let  $rs_1 \in E(G)$  with  $r \in V_{\Delta}$  and  $s_1 \in N_{\Delta-1}(r) := \{s_1, s_2, \ldots, s_{\Delta-2}\},\$ and  $\varphi \in \mathcal{C}^{\Delta}(G - rs_1)$ . Then we call  $(G, rs_1, \varphi)$  a *coloring-triple*. As  $\Delta$ -degree vertices in a multifan do not miss any color, for multifans in HZ-graphs, we add a further requirement in its definition as follows and we use this new definition in the remainder of this paper.

# Assumption. For multifans in an HZ-graph, all of its vertices except the center have degree  $\Delta - 1$ .

Let  $(G, rs_1, \varphi)$  be a coloring-triple and  $F := F_{\varphi}(r, s_1 : s_p)$  be a multifan. By its definition,  $|\overline{\varphi}(s_1)| = 2$ ,  $|\overline{\varphi}(s_i)| = 1$  for each  $i \in [2, p]$ , and so every color in  $\overline{\varphi}(F) \setminus \overline{\varphi}(r)$ is induced by one of the two colors in  $\overline{\varphi}(s_1)$ . We call F a typical multifan, denoted  $F_{\varphi}(r,s_1:s_{\alpha}:s_{\beta}):=(r,rs_1,s_1,rs_2,s_2,\ldots,rs_{\alpha},s_{\alpha},rs_{\alpha+1},s_{\alpha+1},\ldots,rs_{\beta},s_{\beta}),$  where  $\beta:=p$ , if  $\overline{\varphi}(r) = 1$  (recall we denote  $\overline{\varphi}(v)$  by a number if  $|\overline{\varphi}(v)| = 1$ ),  $\overline{\varphi}(s_1) = \{2, \Delta\}$ , and if  $|V(F)| \geq 3$ , then  $\varphi(rs_{\alpha+1}) = \Delta$  and  $\overline{\varphi}(s_{\alpha+1}) = \alpha + 2$  (if  $\beta > \alpha$ ), and for each  $i \in [2, \beta]$  with  $i \neq \alpha+1$ ,  $\varphi(rs_i) = i$  and  $\overline{\varphi}(s_i) = i+1$ . It is clear that  $s_2, \ldots, s_\alpha$  is the longest 2-inducing sequence and  $s_{\alpha+1}, \ldots, s_{\beta}$  is the longest  $\Delta$ -inducing sequence of  $F_{\varphi}(r, s_1 : s_{\alpha} : s_{\beta})$ . By relabelling vertices and colors if necessary, any multifan in an HZ-graph can be assumed to be a typical multifan, see Figure [1](#page-7-0) (a) for a depiction. If  $\alpha = \beta$ , then we write  $F_{\varphi}(r, s_1 : s_{\alpha})$ for  $F_{\varphi}(r, s_1 : s_{\alpha} : s_{\beta})$ , and call it a typical 2-inducing multifan.

#### 3.2 Kierstead path

Let G be a graph,  $e = v_0v_1 \in E(G)$ , and  $\varphi \in C^k(G-e)$  for some integer  $k \geq 0$ . A Kierstead path with respect to e and  $\varphi$  is a sequence  $K = (v_0, v_0v_1, v_1, v_1v_2, v_2, \ldots, v_{p-1}, v_{p-1}v_p, v_p)$ with  $p \geq 1$  consisting of distinct vertices and edges such that for every edge  $v_i v_{i+1}$  with  $i \in [1, p-1]$ , there exists  $j \in [0, i-1]$  satisfying  $\varphi(v_i v_{i+1}) \in \overline{\varphi}(v_j)$ .

A Kierstead path with at most 3 vertices is a multifan. We consider Kierstead paths with 4 vertices. Statement (a) below was proved in Theorem 3.3 from  $|6|$  and statement (b) is a consequence of  $(a)$ .

<span id="page-6-2"></span>**Lemma 3.3.** Let G be a class 2 graph,  $v_0v_1 \in E(G)$ , and  $\varphi \in C^{\Delta}(G - v_0v_1)$ . If  $K =$  $(v_0, v_0v_1, v_1, v_1v_2, v_2, v_2v_3, v_3)$  is a Kierstead path with respect to  $v_0v_1$  and  $\varphi$ , then the following statements hold.

(a) If  $\min\{d_G(v_1), d_G(v_2)\} < \Delta$ , then  $V(K)$  is  $\varphi$ -elementary.



<span id="page-7-0"></span>Figure 1: (a) A typical multifan  $F_{\varphi}(r, s_1 : s_{\alpha} : s_{\beta})$ , where  $\overline{\varphi}(r) = 1$  and  $\overline{\varphi}(s_1) = \{2, \Delta\};$  (b) A rotation centered at  $r$ , where a dashed line at a vertex indicates a color missing at the vertex; (c) A lollipop centered at r, where x can be the same as some  $s_{\ell}$  for  $\ell \in [\beta+1, \Delta-2]$ .

(b) For any two colors  $\alpha, \delta$  with  $\alpha \in \overline{\varphi}(v_0)$  and  $\delta \in \overline{\varphi}(v_3)$ , if  $\min\{d_G(v_1), d_G(v_2)\} < \Delta$  and  $\alpha \notin {\varphi}(v_1v_2), \varphi(v_2v_3)$ , then  $v_3$  and  $v_0$  are  $(\alpha, \delta)$ -linked with respect to  $\varphi$ .

#### 3.3 Pseudo-multifan

Let G be a graph,  $rs_1 \in E(G)$  and  $\varphi \in C^k(G-rs_1)$  for some  $k \geq 0$ . A multifan  $F_{\varphi}(r, s_1 : s_p)$ is maximum at r if  $|V(F)|$  is maximum among all multifans with respect to rs for all  $s \in$  $N_G(r)$  and all  $\varphi' \in C^k(G-rs)$ . A pseudo-multifan with respect to  $rs_1$  and  $\varphi$  is an alternating sequence  $S := S_{\varphi}(r, s_1 : s_t : s_p) := (r, rs_1, s_1, rs_2, s_2, \dots, rs_t, s_t, rs_{t+1}, s_{t+1}, \dots, s_{p-1}, rs_p, s_p)$ with  $t, p \geq 1$  of distinct vertices and edges satisfying the following conditions:

- (P1) the subsequence  $F := (r, rs_1, s_1, rs_2, s_2, \ldots, rs_t, s_t)$  is a maximum multifan at r.
- (P2)  $V(S)$  is  $\varphi'$ -elementary for every  $(F, \varphi)$ -stable  $\varphi' \in C^k(G rs_1)$ .

Every maximum multifan is a pseudo-multifan, and if  $S$  is a pseudo-multifan with respect to  $\varphi$  and a multifan F, then by the definition above, S is a pseudo-multifan under every  $(F, \varphi)$ -stable coloring  $\varphi'$ . We call a pseudo-multifan S typical (resp. typical 2-inducing) if the maximum multifan that is contained in  $S$  is typical (resp. typical 2-inducing).

Let  $(G, rs_1, \varphi)$  be a coloring-triple and  $i, j \in [2, \Delta - 2]$ . The shift from  $s_i$  to  $s_j$  is an operation that, for each  $\ell$  with  $\ell \in [i, j]$ , recolor  $rs_{\ell}$  by the color in  $\overline{\varphi}(s_{\ell})$ . We will apply a shift either on a sequence of vertices from a multifan or on a rotation.

<span id="page-8-0"></span>**Lemma 3.4.** Let  $(G, rs_1, \varphi)$  be a coloring-triple. Then for every typical pseudo-multifan  $S := S_{\varphi}(r, s_1 : s_t : s_p)$ , there exists a coloring  $\varphi' \in C^{\Delta}(G - rs_t)$  and a pseudo-multifan  $S^*$ centered at r with respect to rs<sub>t</sub> and  $\varphi'$  such that  $V(S^*) = V(S)$  and  $S^*$  is typical 2-inducing.

**Proof.** Let  $F = F_{\varphi}(r, s_1 : s_{\alpha} : s_{\beta})$  be the typical multifan contained in S, where  $s_{\beta} = s_t$ . If  $\beta = \alpha$ , then we are done. Thus we assume  $\beta \ge \alpha + 1 \ge 3$ . Let  $\varphi'$  be obtained from  $\varphi$  by uncoloring  $rs_\beta$ , shift from  $s_{\alpha+1}$  to  $s_{\beta-1}$  and coloring  $rs_1$  by  $\Delta$ . Now  $\overline{\varphi}'(s_{\beta}) = \{\beta, \beta + 1\}, F^* = (r, rs_{\beta}, s_{\beta}, rs_{\beta-1}, s_{\beta-1}, \ldots, rs_{\alpha+1}, s_{\alpha+1}, rs_1, s_1, \ldots, rs_{\alpha}, s_{\alpha})$  is a β-inducing multifan with respect to  $rs_\beta$  and  $\varphi'$ .

We next show that  $S^* = (F^*, rs_{t+1}, s_{t+1}, \ldots, rs_p, s_p)$  is a pseudo-multifan with respect to  $rs_{\beta}$  and  $\varphi'$ . Since  $|V(F^*)| = |V(F)|$ ,  $F^*$  is also a maximum multifan at r. Thus it suffices to show that for any  $(F^*, \varphi')$ -stable  $\varphi'' \in C^{\Delta}(G - rs_{\beta}), V(S^*)$  is  $\varphi''$ -elementary. Suppose to the contrary that there exists  $(F^*, \varphi')$ -stable  $\varphi'' \in C^{\Delta}(G - rs_{\beta})$  but  $V(S^*)$  is not  $\varphi''$ -elementary. As  $\varphi''$  is  $(F^*, \varphi')$ -stable, we can undo the operations we did before. Specifically, let  $\varphi'''$  be the coloring obtained from  $\varphi''$  by uncoloring  $rs_1$ , shift from  $s_{\alpha+1}$ to  $s_{\beta-1}$  and coloring  $rs_{\beta}$  by  $\beta$ . Then  $\varphi'''$  is  $(F, \varphi)$ -stable and  $\overline{\varphi}'''(S^*) = \overline{\varphi}''(S^*)$ . Thus,  $V(S^*)$  is not  $\varphi''$ -elementary implies that  $V(S^*)$  is not  $\varphi'''$ -elementary. Since  $V(S^*) = V(S)$ , this contradicts the assumption that  $V(S)$  is elementary under any  $(F, \varphi)$ -stable coloring. Therefore,  $S^*$  is a pseudo-multifan with respect to  $rs_\beta$  and  $\varphi'$ . By renaming colors and vertices, we can assume that  $F^*$  is typical 2-inducing and so  $S^*$  is typical 2-inducing.  $\Box$ 

Let  $(G, rs_1, \varphi)$  be a coloring-triple. A sequence of distinct vertices  $w_1, \ldots, w_t \in N_{\Delta-1}(r)$ form a rotation if  $\{w_1, \ldots, w_t\}$  is  $\varphi$ -elementary, and for each  $\ell$  with  $\ell \in [1, t]$ , it holds that  $\varphi(rw_\ell) = \overline{\varphi}(w_{\ell-1}),$  $\varphi(rw_\ell) = \overline{\varphi}(w_{\ell-1}),$  $\varphi(rw_\ell) = \overline{\varphi}(w_{\ell-1}),$  where  $w_0 := w_t$ . An example of a rotation is given in Figure 1 (b).

<span id="page-8-1"></span>**Lemma 3.5** ( [\[1,](#page-26-4) Theorem 2.5]). Let  $(G, rs_1, \varphi)$  be a coloring-triple,  $S := S_{\varphi}(r, s_1 : s_t : s_t)$  $s_{\Delta-2}$ ) be a pseudo-multifan with  $F := F_{\varphi}(r, s_1 : s_t)$  being the maximum multifan contained in it. Let  $j \in [t+1, \Delta-2]$  and  $\delta \in \overline{\varphi}(s_j)$ . Then

- <span id="page-8-3"></span>(a)  $\{s_{t+1}, \ldots, s_{\Delta-2}\}\$  can be partitioned into rotations with respect to  $\varphi$ .
- <span id="page-8-2"></span>(b)  $s_j$  and r are  $(1, \delta)$ -linked with respect to  $\varphi$ .
- (c) For every color  $\gamma \in \overline{\varphi}(F)$  with  $\gamma \neq 1$ , it holds that  $r \in P_y(\gamma, \delta) = P_{s_j}(\gamma, \delta)$ , where  $y=\overline{\varphi}_{F}^{-1}$  $\mathcal{F}_F^{-1}(\gamma)$ . Furthermore, for  $z \in N_G(r)$  such that  $\varphi(rz) = \gamma$ ,  $P_y(\gamma, \delta)$  meets z before r.
- <span id="page-8-4"></span>(d) For every  $\delta^* \in \overline{\varphi}(S) \setminus \overline{\varphi}(F)$  with  $\delta^* \neq \delta$ , it holds that  $P_y(\delta, \delta^*) = P_{s_j}(\delta, \delta^*)$ , where  $y = \overline{\varphi}_S^{-1}$  $\overline{S}^1(\delta^*)$ . Furthermore, either  $r \in P_{s_j}(\delta, \delta^*)$  or  $P_r(\delta, \delta^*)$  is an even cycle.

#### 3.4 Lollipop

If  $F = (a_1, \ldots, a_t)$  is a sequence, then for a new entry b,  $(F, b)$  denotes the sequence  $(a_1, \ldots, a_t, b)$ . Let  $(G, rs_1, \varphi)$  be a coloring-triple. A *lollipop* centered at r is a sequence  $L = (F, ru, u, ux, x)$  of distinct vertices and edges such that  $F = F_{\varphi}(r, s_1 : s_{\alpha} : s_{\beta})$  is a typical multifan,  $u \in N_\Delta(r)$  and  $x \in N_{\Delta-1}(u)$  $x \in N_{\Delta-1}(u)$  $x \in N_{\Delta-1}(u)$  with  $x \notin \{s_1, \ldots, s_\beta\}$  (see Figure 1 (c) for a depiction).

<span id="page-9-3"></span>**Lemma 3.6** ([\[1,](#page-26-4) Lemma 5.1]). Let  $(G, rs_1, \varphi)$  be a coloring-triple,  $F := F_{\varphi}(r, s_1 : s_{\alpha} : s_{\beta})$ be a typical multifan, and  $L := (F, ru, u, ux, x)$  be a lollipop centered at r such that  $\varphi(ru) =$  $\alpha + 1$  and  $\overline{\varphi}(x) = \alpha + 1$ . Then

<span id="page-9-4"></span>(a)  $\varphi(ux) \neq 1$  and  $ux \in P_r(1, \varphi(ux))$ .

If  $\varphi(ux) = \tau$  is a 2-inducing color with respect to  $\varphi$  and F, then the following holds.

- <span id="page-9-6"></span>(b) Let  $P_x(1,\tau)$  be the  $(1,\tau)$ -chain starting at x in  $G-rs_1-ux$ . Then  $P_x(1,\tau)$  ends at r.
- <span id="page-9-5"></span>(c) For any 2-inducing color  $\delta$  of F with  $\tau \prec \delta$ , we have  $r \in P_{s_1}(\delta, \Delta) = P_{s_{\delta-1}}(\delta, \Delta)$ .
- (d) For any  $\Delta$ -inducing color  $\delta$  of F, we have  $r \in P_{s_{\delta-1}}(\alpha+1,\delta) = P_{s_{\alpha}}(\alpha+1,\delta)$ , where  $s_{\Delta-1} = s_1$  if  $\delta = \Delta$ .
- (e) For any 2-inducing color  $\delta$  of F with  $\delta \prec \tau$ , we have  $r \in P_{s_\alpha}(\delta, \alpha+1) = P_{s_{\delta-1}}(\delta, \alpha+1)$ .

Let  $(G, rs_1, \varphi)$  be a coloring-triple. For a color  $\alpha \in [1, \Delta]$ , a sequence of  $Kempe (\alpha, *)$ *changes* is a sequence of Kempe changes that each involve the exchanging of the color  $\alpha$ with another color from  $[1, \Delta]$ .

<span id="page-9-0"></span>**Lemma 3.7** ([\[1,](#page-26-4) Lemma 5.2]). Let  $(G, rs_1, \varphi)$  be a coloring-triple,  $F := F_{\varphi}(r, s_1 : s_{\alpha} : s_{\beta})$ be a typical multifan, and  $L := (F, ru, u, ux, x)$  be a lollipop centered at r such that  $\varphi(ru) =$  $\alpha + 1$ . Then for  $w_1 \in \{s_{\beta+1}, \ldots, s_{\Delta-2}\}\$  with  $\varphi(rw_1) = \tau_1 \in [\beta + 2, \Delta - 1]$ , the following statements hold.

- <span id="page-9-1"></span>(1) If exists a vertex  $w \in V(G) \setminus (V(F) \cup \{w_1\})$  such that  $w \in P_r(1, \tau_1, \varphi')$  for every  $(F, \varphi)$ stable  $\varphi' \in C^{\Delta}(G - rs_1)$  with  $\varphi'(ru) = \alpha + 1$ , then there exists a sequence of distinct vertices  $w_1, \ldots, w_t \in \{s_{\beta+1}, \ldots, s_{\Delta-2}\}\$  satisfying the following conditions:
	- (a)  $\varphi(rw_{i+1}) = \overline{\varphi}(w_i) \in [\beta + 2, \Delta 1]$  for each  $i \in [1, t 1]$ ;
	- (b) r and w<sub>i</sub> are  $(1,\overline{\varphi}(w_i))$ -linked with respect to  $\varphi$  for each  $i \in [1,t]$ ;
	- $(c) \overline{\varphi}(w_t) = \tau_1.$
- <span id="page-9-2"></span>(2) If  $\overline{\varphi}(x) = \alpha + 1$  and there exists a vertex  $w \in V(G) \setminus (V(F) \cup \{w_1\})$  such that  $w \in P_r(1,\tau_1,\varphi')$  for every  $(L,\varphi)$ -stable  $\varphi' \in C^{\Delta}(G-rs_1)$  obtained from  $\varphi$  through

a sequence of Kempe  $(1,*)$ -changes not using r or x as endvertices, then there exists a sequence of distinct vertices  $w_1, \ldots, w_t \in \{s_{\beta+1}, \ldots, s_{\Delta-2}\}\$  satisfying the following conditions:

- (a)  $\varphi(rw_{i+1}) = \overline{\varphi}(rw_i) \in [\beta + 2, \Delta 1]$  for each  $i \in [1, t 1]$ ;
- (b) r and  $w_i$  are  $(1,\overline{\varphi}(w_i))$ -linked with respect to  $\varphi$  for each  $i \in [1,t-1]$ ;
- (c)  $\overline{\varphi}(w_t) = \tau_1$  or  $\overline{\varphi}(w_t) = \alpha + 1$ . If  $\overline{\varphi}(w_t) = \tau_1$ , then  $w_t$  and r are  $(1, \tau_1)$ -linked with respect to  $\varphi$ .

By the definition, the sequence  $w_1, \ldots, w_t$  in Lemma [3.7](#page-9-0) [\(1\)](#page-9-1) and in the case of Lemma 3.7 [\(2\)](#page-9-2) when  $\overline{\varphi}(w_t) = \tau_1$  form a rotation with the additional property that  $\overline{\varphi}(w_i) \in [\beta+2, \Delta-1]$  and r and  $w_i$  are  $(1,\overline{\varphi}(w_i))$ -linked for each  $i \in [1,t]$ . We call such a rotation a *stable rotation*. In the case of Lemma [3.7](#page-9-0) [\(2\)](#page-9-2) when  $\overline{\varphi}(w_t) = \alpha + 1$ , we call  $w_1, \ldots, w_t$  a near stable rotation. For  $u, v \in V(G)$ , we write  $u \sim v$  if u and v are adjacent in G, and write  $u \not\sim v$  otherwise.

<span id="page-10-0"></span>**Lemma 3.8** ([\[1,](#page-26-4) Corollary 2.7]). Let  $(G, rs_1, \varphi)$  be a coloring-triple,  $F := F_{\varphi}(r, s_1 : s_{\alpha})$ be a typical 2-inducing multifan, and  $L := (F, ru, u, ux, x)$  be a lollipop centered at r. If  $\varphi(ru) = \alpha + 1, \overline{\varphi}(x) = \alpha + 1, \text{ and } \varphi(ux) = \Delta, \text{ then } u \nsim s_1 \text{ and } u \nsim s_\alpha.$ 

<span id="page-10-1"></span>**Lemma 3.9** ( [\[1,](#page-26-4) Theorem 2.8]). Let  $(G, rs_1, \varphi)$  be a coloring-triple,  $F := F_{\varphi}(r, s_1 : s_{\alpha})$ be a typical 2-inducing multifan, and  $L := (F, ru, u, ux, x)$  be a lollipop centered at r. If  $\varphi(ru) = \alpha + 1$ ,  $\overline{\varphi}(x) = \alpha + 1$ , and  $\varphi(ux) = \mu \in \overline{\varphi}(F)$  is a 2-inducing color of F, then  $u \nsim s_{\mu-1}$  and  $u \nsim s_{\mu}$ .

Let G be a graph,  $rs_1 \in E(G)$  and  $\varphi \in C^{\Delta}(G - rs_1)$ . Let  $\alpha, \beta, \gamma, \tau \in [1, \Delta]$  and  $x, y \in V(G)$ . If P is an  $(\alpha, \beta)$ -chain containing both x and y such that P is a path, we denote by  $P_{[x,y]}(\alpha, \beta, \varphi)$  the subchain of P that has endvertices x and y.

Suppose  $|\overline{\varphi}(x) \cap {\alpha, \beta}| = 1$ . Then an  $(\alpha, \beta)$ -swap at x is just the Kempe change on  $P_x(\alpha, \beta, \varphi)$ . By convention, an  $(\alpha, \alpha)$ -swap at x does nothing at x. If also  $|\overline{\varphi}(y) \cap {\alpha, \beta}| = 1$ , then an  $(\alpha, \beta)$ -swap at both x and y is the Kempe change on  $P_x(\alpha, \beta, \varphi)$  if x and y are  $(\alpha, \beta)$ linked with respect to  $\varphi$ , and is obtained from  $\varphi$  by first doing an  $(\alpha, \beta)$ -swap at x and then doing an  $(\alpha, \beta)$ -swap at y if x and y are  $(\alpha, \beta)$ -unlinked with respect to  $\varphi$ . Suppose  $\beta_0 \in \overline{\varphi}(x)$ and  $\beta_1, \ldots, \beta_t \in \varphi(x)$  for colors  $\beta_0, \ldots, \beta_t \in [1, \Delta]$  for some integer  $t \geq 1$ . Then a

$$
(\beta_0, \beta_1) - (\beta_1, \beta_2) - \ldots - (\beta_{t-1}, \beta_t) - \text{swap}
$$

at x consists of t Kempe changes: let  $\varphi_0 = \varphi$ , then  $\varphi_i = \varphi_{i-1}/P_x(\beta_{i-1}, \beta_i, \varphi_{i-1})$  for each  $i \in [1, t]$ . Suppose the current color of an edge uv of G is  $\alpha$ , the notation  $uv : \alpha \to \beta$  means to recolor the edge uv using the color  $\beta$ .

We will use a matrix with two rows to denote a sequence of coloring operations taken based on  $\varphi$ . For example, the matrix below indicates three operations taken on the graph:

$$
\begin{bmatrix} P_{[a,b]}(\alpha,\beta,\varphi) & s_c : s_d & rs \\ \alpha/\beta & \text{shift} & \gamma \to \tau \end{bmatrix}.
$$

Step 1 Exchange  $\alpha$  and  $\beta$  on the  $(\alpha, \beta)$ -subchain  $P_{[a,b]}(\alpha, \beta, \varphi)$ .

Step 2 Based on the coloring obtained from Step 1, shift from  $s_c$  to  $s_d$  for vertices  $s_c, \ldots, s_d$ .

Step 3 Based on the coloring obtained from Step 2, do  $rs : \gamma \to \tau$ .

In the reminder, for simpler description, we may skip the phrase "with respect to  $\varphi$ " in related notation, which then needs to be understood with respect to the current edge coloring.

# 4 Proof of Theorem [2.3](#page-2-0)

We prove the following version of Theorem [2.3.](#page-2-0)

<span id="page-11-2"></span><span id="page-11-1"></span>**Theorem 4.1.** If G is an HZ-graph with maximum degree  $\Delta \geq 4$ , then for every vertex  $r \in V_{\Delta}$ , the following two statements hold.

- <span id="page-11-3"></span>(i) For every  $u \in N_{\Delta}(r)$ ,  $N_{\Delta-1}(r) = N_{\Delta-1}(u)$ .
- (ii) There exist  $s_1 \in N_{\Delta-1}(r)$  and  $\varphi \in C^{\Delta}(G rs_1)$  such that  $N_{\Delta-1}[r]$  is the vertex set of a typical 2-inducing pseudo-multifan with respect to rs<sub>1</sub> and  $\varphi$ . Consequently  $N_{\Delta-1}[r]$ is  $\varphi$ -elementary.

*Proof.* Let  $N_{\Delta-1}(r) = \{s_1, \ldots, s_{\Delta-2}\}\.$  We choose a vertex in  $N_{\Delta-1}(r)$ , say  $s_1$ , a coloring  $\varphi \in C^{\Delta}(G-r_{s_1})$  and a multifan F with respect to  $rs_1$  and  $\varphi$  such that F is maximum at r. That is,  $|V(F)|$  is maximum among all multifans with respect to  $rs_i$  for any  $i \in [1, \Delta-2]$  and any  $\varphi' \in C^{\Delta}(G - rs_i)$ . Assume that  $\overline{\varphi}(r) = 1$  and  $\overline{\varphi}(s_1) = \{2, \Delta\}$ , and  $F = F_{\varphi}(r, s_1 : s_p)$ is such a multifan. Furthermore, by relabeling vertices and colors, we assume that  $F$  is typical. As a maximum multifan at r is itself a pseudo-multifan, by Lemma [3.4,](#page-8-0) we assume that  $F_{\varphi}(r, s_1 : s_p) = F_{\varphi}(r, s_1 : s_\alpha)$  is a typical 2-inducing multifan, where  $\alpha = p$ .

Let  $u \in N_{\Delta}(r)$  and assume  $N_{\Delta-1}(r) \neq N_{\Delta-1}(u)$ . Roughly speaking, the main proof idea is the following. By assuming  $\varphi(ru) = \alpha + 1$  and  $\overline{\varphi}(x) = \alpha + 1$  for  $x \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$ , we will apply Lemmas [3.8](#page-10-0) and [3.9](#page-10-1) to show that u has at least two  $(\Delta - 1)$ -neighbors outside of  $N_{\Delta-1}(r)$ . By further applying Lemmas [3.8](#page-10-0) and [3.9,](#page-10-1) we can even find three  $(\Delta-1)$ -neighbors of u outside of  $N_{\Delta-1}(r)$ . A contradiction is then deduced at that point.

<span id="page-11-0"></span>**Claim 4.1.** We may assume that  $\varphi(ru) = \alpha + 1$ , which is the last 2-inducing color of  $F_{\varphi}(r,s_1:s_{\alpha}).$ 

*Proof of Claim [4.1.](#page-11-0)* Since  $F_{\varphi}(r, s_1 : s_{\alpha})$  is a maximum typical 2-inducing multifan,  $\varphi(ru) \in$  $\{\alpha+1,\Delta\}$ . Assume instead that  $\varphi(ru) = \Delta$ . If  $\alpha = 1$ , then we are done by exchanging the roles of 2 and  $\Delta$ . Thus we assume that  $\alpha \geq 2$ . Shift from  $s_2$  to  $s_{\alpha-1}$ , color  $rs_1$  by 2 and uncolor  $rs_\alpha$ . Then  $F^* = (r, rs_\alpha, s_\alpha, rs_{\alpha-1}, s_{\alpha-1}, \ldots, rs_1, s_1)$  is an  $\alpha$ -inducing multifan such

that  $\Delta$  is the last  $\alpha$ -inducing color. Now, relabeling colors and vertices in  $F^*$  by making  $F^*$  typical 2-inducing yields the desired assumption. □

<span id="page-12-0"></span>Claim 4.2. For any  $z \in N_{\Delta-1}(u) \setminus V(F)$  and any  $(F,\varphi)$ -stable  $\varphi' \in C^{\Delta}(G-rs_1)$ , if  $\varphi'(ru) = \alpha + 1$  and  $\overline{\varphi}'(z) = \alpha + 1$ , then  $\varphi'(uz) \in \overline{\varphi}'(F) \setminus \{1\}.$ 

*Proof of Claim [4.2.](#page-12-0)* Assume to the contrary that  $\varphi'(uz) \in \{1, \alpha + 2, ..., \Delta - 1\}$ . We first claim that  $\varphi'(uz) \neq 1$ . As otherwise,  $P_r(1, \alpha + 1, \varphi') = ruz$ , contradicting Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that r and  $s_{\alpha}$  are  $(1, \alpha + 1)$ -linked with respect to  $\varphi'$ . Let  $\varphi'(uz) = \tau \in [\alpha + 2, \Delta - 1]$ , and  $w_1 \in N_{\Delta-1}(r)$  such that  $\varphi'(rw_1) = \tau$ . By Lemma [3.6](#page-9-3) [\(a\)](#page-9-4),  $uz \in P_r(1, \tau, \varphi'')$  for every  $(L, \varphi')$ -stable  $\varphi'' \in C^{\Delta}(G - rs_1)$ , where  $L = (F, ru, u, uz, z)$  is a lollipop. Applying Lemma [3.7](#page-9-0) [\(2\)](#page-9-2) on L with u playing the role of  $w$ , we find a sequence of distinct vertices  $w_1, \ldots, w_t \in \{s_{\alpha+1}, \ldots, s_{\Delta-2}\}\$  that forms either a stable rotation or a near stable rotation.

Assume first that  $w_1, \ldots, w_t$  is a stable rotation, which in particular gives  $P_r(1, \tau, \varphi') =$  $P_{w_t}(1, \tau, \varphi')$ . By Lemma [3.6](#page-9-3) [\(a\)](#page-9-4),  $uz \in P_r(1, \tau, \varphi')$ . If  $P_r(1, \tau, \varphi')$  meets z before u, or equivalently,  $P_{wt}(1, \tau, \varphi')$  meets u before z, we do the following operations:

$$
\begin{bmatrix} P_{[r,z]}(1,\tau,\varphi') & ru & uz \\ 1/\tau & \alpha+1 \to \tau & \tau \to \alpha+1 \end{bmatrix}.
$$

Denote the new coloring by  $\varphi''$ . Now  $(r, rs_1, s_1, \ldots, s_\alpha)$  is a multifan, but  $\overline{\varphi}''(s_\alpha) = \overline{\varphi}''(r) =$  $\alpha + 1$ , giving a contradiction to Lemma [3.1](#page-5-0) [\(a\)](#page-5-2). Thus  $P_r(1, \tau, \varphi')$  meets u before z, or equivalently,  $P_{w_t}(1, \tau, \varphi')$  meets z before u. Shift from  $w_1$  to  $w_t$  to get  $\varphi''$ . Then  $P_r(1, \tau, \varphi'')$ meets z before u, giving back to the previous case as  $\varphi''$  is  $(F, \varphi')$ -stable.

Assume now that  $w_1, \ldots, w_t$  is a near stable rotation, i.e.,  $\overline{\varphi}'(w_t) = \alpha + 1$ . If  $z \neq w_t$ , then we shift from  $w_1$  to  $w_t$ , and do  $ru : \alpha + 1 \rightarrow \tau$ ,  $uz : \tau \rightarrow \alpha + 1$ . Denote the new coloring by  $\varphi''$ . As  $\varphi''$  is  $(F, \varphi')$ -stable and so is  $(F, \varphi)$ -stable, we see that  $F^* =$  $(F, rw_t, w_t, rw_{t-1}, w_{t-1}, \ldots, rw_1, w_1)$  is a multifan that contains more vertices than F does, showing a contradiction to the choice of F.

Thus we assume that  $z = w_t$ . Since  $\varphi'(rz) \neq \varphi'(uz) = \tau$ , we have  $t \geq 2$ . Note that  $uz \in P_r(1,\tau,\varphi') = P_w(1,\tau,\varphi')$  for some vertex  $w \in V(G) \setminus (V(F) \cup \{w_1,\ldots,w_t\})$ . If  $P_w(1, \tau, \varphi')$  meets u before z, we do the following operations:

$$
\begin{bmatrix} P_{[w,u]}(1,\tau,\varphi') & ru & uz \\ 1/\tau & \alpha+1 \to 1 & \tau \to \alpha+1 \end{bmatrix}.
$$

Denote the new coloring by  $\varphi''$ . Now  $(r, rs_1, s_1, \ldots, s_\alpha)$  is a multifan, but  $\overline{\varphi}''(s_\alpha) = \overline{\varphi}''(r) =$  $\alpha + 1$ , giving a contradiction to Lemma [3.1](#page-5-0) [\(a\)](#page-5-2). If  $P_w(1, \tau, \varphi')$  meets z before u, we do the following operations:

$$
\begin{bmatrix} P_{[w,z]}(1,\tau,\varphi') & w_1:w_t & rw_t = rz & ru & uz \\ 1/\tau & \text{shift} & \varphi'(rz) \to 1 & \alpha+1 \to \tau & \tau \to \alpha+1 \end{bmatrix}.
$$

Denote the new coloring by  $\varphi''$ . Now  $(r, rs_1, s_1, \ldots, s_\alpha)$  is a multifan, but  $\overline{\varphi}''(s_\alpha) = \overline{\varphi}''(r) =$  $\alpha + 1$ , giving a contradiction to Lemma [3.1](#page-5-0) [\(a\)](#page-5-2). □

By Claim [4.2,](#page-12-0)  $\tau \in \{2, \ldots, \alpha+1, \Delta\}$ . Applying Lemmas [3.8](#page-10-0) and [3.9,](#page-10-1) we have the following claim.

<span id="page-13-1"></span>Claim 4.3. Let  $z \in N_{\Delta-1}(u) \setminus V(F)$  and any  $(F, \varphi)$ -stable  $\varphi' \in C^{\Delta}(G - rs_1)$  such that  $\varphi'(ru) = \alpha + 1$  and  $\overline{\varphi}'(z) = \alpha + 1$ , and let  $\varphi'(uz) = \tau$ . Then  $\tau \in \overline{\varphi}'(F) \setminus \{1\}$ , and  $u \not\sim s_1, s_\alpha$ if  $\tau = \Delta$ ; and  $u \nsim s_{\tau-1}, s_{\tau}$  if  $\tau \in [2, \alpha + 1]$ .

<span id="page-13-0"></span>Claim 4.4. Suppose that  $N_{\Delta-1}(r) = N_{\Delta-1}(u)$  for every  $u \in N_{\Delta}(r)$ . Then for every  $(F, \varphi)$ stable coloring  $\varphi' \in C^{\Delta}(G - rs_1)$ ,  $N_{\Delta-1}[r]$  is  $\varphi'$ -elementary. In particular,  $N_{\Delta-1}[r]$  is the vertex set of a typical 2-inducing pseudo-multifan with respect to  $rs^*$  and  $\varphi^* \in C^{\Delta}(G - rs^*)$ for some  $s^* \in N_{\Delta-1}(r)$ .

Proof of Claim [4.4.](#page-13-0) Assume to the contrary that there exists an  $(F, \varphi)$ -stable coloring  $\varphi' \in$  $\mathcal{C}^{\Delta}(G - rs_1)$  such that  $N_{\Delta-1}[r]$  is not  $\varphi'$ -elementary. Since  $V(F)$  is  $\varphi'$ -elementary, there exists  $z \in N_{\Delta-1}[r] \setminus V(F)$  such that  $\overline{\varphi}'(z) \in \overline{\varphi}'(F)$  or there exists  $z^* \neq z$  with  $z^* \in$  $N_{\Delta-1}[r] \setminus V(F)$  such that  $\overline{\varphi}'(z) = \overline{\varphi}'(z^*)$ . Let  $\overline{\varphi}'(z) = \delta$ . If  $\delta \in \overline{\varphi}'(F)$ , then z and r are  $(1, \delta)$ -unlinked, so we do  $(\delta, 1) - (1, \alpha + 1)$ -swaps at z; if  $\overline{\varphi}'(z) = \overline{\varphi}'(z^*)$ , we may assume, without loss of generality, that z and r are  $(1, \delta)$ -unlinked, we again do  $(\delta, 1) - (1, \alpha + 1)$ swaps at z. In either case, we find an  $(F, \varphi')$ -stable coloring  $\varphi'' \in C^{\Delta}(G - rs_1)$  such that  $\overline{\varphi}''(z) = \alpha + 1$ . Since for any  $u \in N_{\Delta}(r)$ , it holds that  $N_{\Delta-1}(r) = N_{\Delta-1}(u)$ , we can choose  $u \in N_\Delta(r)$  such that  $\varphi''(ur) = \alpha+1$ , where  $\alpha+1$  is the last 2-inducing color of  $F_{\varphi''}(r, s_1 : s_\alpha)$ . Since  $N_{\Delta-1}(r) = N_{\Delta-1}(u)$ , we have  $uz \in E(G)$  and so  $L = (F_{\varphi''}(r, s_1 : s_\alpha), ru, u, uz, z)$ is a lollipop with respect to  $\varphi''$ . By Claim [4.3,](#page-13-1) u is not adjacent to at least one vertex in  $N_{\Delta-1}(r)$ , which in turn shows  $N_{\Delta-1}(r) \neq N_{\Delta-1}(u)$ , giving a contradiction.

Therefore, for every  $(F, \varphi)$ -stable coloring  $\varphi' \in C^{\Delta}(G - rs_1)$ , it holds that  $N_{\Delta-1}[r]$ is  $\varphi'$ -elementary. Consequently, there is a pseudo-multifan with vertex set  $N_{\Delta-1}[r]$ . By renaming colors and vertices from  $N_{\Delta-1}(r)$ , we can assume the pseudo-multifan with vertex set  $N_{\Delta-1}[r]$  is typical. By Lemma [3.4,](#page-8-0) we can further assume that the pseudo-multifan is typical 2-inducing.  $\Box$ 

By Claim [4.4,](#page-13-0) it suffices to only show Theorem [4.1](#page-11-1) [\(i\)](#page-11-2). Assume to the contrary that there exists  $u \in N_{\Delta}(r)$  such that  $N_{\Delta-1}(u) \setminus N_{\Delta-1}(r) \neq \emptyset$ .

<span id="page-13-2"></span>Claim 4.5. For every  $z \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$ , there is an  $(F, \varphi)$ -stable coloring  $\varphi' \in$  $\mathcal{C}^{\Delta}(G - rs_1)$  such that  $\varphi'(ru) = \alpha + 1$  and  $\overline{\varphi}'(z) = \alpha + 1$ .

*Proof of Claim [4.5.](#page-13-2)* By Claim [4.1,](#page-11-0) we assume  $\varphi(ru) = \alpha + 1$ . Let  $\overline{\varphi}(z) = \delta$ . If  $\delta = \alpha + 1$ , we simply let  $\varphi' = \varphi$ . So  $\delta \neq \alpha + 1$ . If  $\delta \in \overline{\varphi}(F)$ , we let  $\varphi'$  be obtained from  $\varphi$  by doing  $(\delta, 1) - (1, \alpha + 1)$ -swaps at z. This gives that  $\overline{\varphi}'(z) = \alpha + 1$ . By Lemma [3.1](#page-5-0) [\(b\)](#page-5-1),  $\varphi'$  is  $(F, \varphi)$ -stable and  $\varphi'(ru) = \varphi(ru) = \alpha + 1$ . Thus  $\varphi'$  is a desired coloring.

Assume now that  $\delta \in [\alpha+2,\Delta-1]$ . If there is an  $(F,\varphi)$ -stable  $\varphi'' \in C^{\Delta}(G-rs_1)$ with  $\varphi''(ru) = \alpha + 1$  such that  $z \notin P_r(1, \delta, \varphi'')$  (so z and r are  $(1, \delta)$ -unlinked), let  $\varphi'$  be obtained from  $\varphi''$  by doing  $(\delta, 1) - (1, \alpha + 1)$ -swaps at z. Since  $\varphi''(ru) = \alpha + 1$  and r and  $s_{\alpha}$  are  $(1, \alpha + 1)$ -linked with respect to  $\varphi''$  by Lemma [3.1](#page-5-0) [\(b\)](#page-5-1), it holds that  $\varphi'$  is  $(F, \varphi'')$ stable and so  $(F, \varphi)$ -stable with  $\varphi'(ru) = \varphi''(ru) = \alpha + 1$ . Thus,  $\varphi'$  is a desired coloring and we are done. Therefore every  $(F, \varphi)$ -stable  $\varphi'' \in C^{\Delta}(G - rs_1)$  with  $\varphi''(ru) = \alpha + 1$ satisfies  $z \in P_r(1, \delta, \varphi'')$ . Applying Lemma [3.7](#page-9-0) [\(1\)](#page-9-1) with z playing the role of w, there exists  $w_t \in N_{\Delta-1}(r) \setminus V(F)$  such that  $\overline{\varphi}(w_t) = \delta$  and  $w_t$  and  $r$  are  $(1, \delta)$ -linked with respect to  $\varphi$ . This is a contradiction by noting  $w_t \neq z$ , since  $\varphi$  is  $(F, \varphi)$ -stable but  $z \notin P_r(1, \delta, \varphi)$ .  $\Box$ 

<span id="page-14-0"></span>Claim 4.6.  $|N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)| \geq 2$ .

*Proof of Claim [4.6.](#page-14-0)* Let  $x \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$ . By Claim [4.5,](#page-13-2) we choose an  $(F, \varphi)$ -stable coloring from  $\mathcal{C}^{\Delta}(G - rs_1)$  and call it still  $\varphi$  such that  $\varphi(ru) = \alpha + 1$  and  $\overline{\varphi}(x) = \alpha + 1$ . By Claim [4.3,](#page-13-1)  $\varphi(ux) \in \{2, ..., \alpha + 1, \Delta\}$ . If  $|V(F)| \geq 3$ , then Claim [4.3](#page-13-1) gives that  $|N_{\Delta-1}(u)|$  $N_{\Delta-1}(r)| \geq 2$ . Thus we have  $V(F) = \{r, s_1\}$ . Consequently,  $\alpha + 1 = 2$ , and  $\varphi(ux) = \Delta$  by the fact that  $\varphi(ux) \in \{2, \Delta\}$ . We may assume further that  $N_{\Delta-1}(u) \setminus N_{\Delta-1}(r) = \{x\}.$ 

By Claim [4.3,](#page-13-1)  $u \nsim s_1$ . We consider two cases. Assume first that there exists an  $(F, \varphi)$ stable  $\varphi' \in C^{\Delta}(G-rs_1)$  such that  $N_{\Delta-1}[r]$  is not  $\varphi'$ -elementary. By exchanging the roles of 2 and  $\Delta$  if necessary, we may assume  $\varphi'(ru) = 2$ . Since  $V(F)$  is  $\varphi'$ -elementary, there exists  $z \in N_{\Delta-1}(r) \setminus V(F)$  such that  $\overline{\varphi}'(z) \in \overline{\varphi}'(F)$  or there exists  $z^* \neq z$  with  $z^* \in N_{\Delta-1}(r) \setminus V(F)$ such that  $\overline{\varphi}'(z) = \overline{\varphi}'(z^*)$ . Let  $\overline{\varphi}'(z) = \delta$ . If  $\delta \in \overline{\varphi}'(F)$ , then as r and z are  $(1, \delta)$ -unlinked, we do  $(\delta, 1) - (1, 2)$ -swaps at z; if  $\overline{\varphi}'(z) = \overline{\varphi}'(z^*)$ , we may assume, without loss of generality, that z and r are  $(1, \delta)$ -unlinked, we again do  $(\delta, 1) - (1, 2)$ -swaps at z. In either case, we find an  $(F, \varphi')$ -stable coloring  $\varphi'' \in C^{\Delta}(G - rs_1)$  with  $\varphi''(ru) = \varphi'(ru) = 2$  and  $\overline{\varphi}''(z) = 2$ . Note that  $z \in N_{\Delta-1}(u)$  since  $u \nsim s_1$ ,  $s_1 \neq z$ , and  $N_{\Delta-1}(u) \setminus N_{\Delta-1}(r) = \{x\}$ . By Claim [4.2,](#page-12-0)  $\varphi''(uz) \in \{2, \Delta\}$ , which implies  $\varphi''(uz) = \Delta$  by noting  $\varphi''(ru) = 2$ . Furthermore, we assume  $uz \in P_{s_1}(1, \Delta, \varphi'') = P_r(1, \Delta, \varphi'')$  (otherwise, after a  $(1, \Delta)$ -swap on the chain containing uz, we obtain a contradiction to Claim [4.2\)](#page-12-0). Since  $\varphi''(ru) = 2$  and  $\varphi''(uz) = \Delta$ ,  $\varphi''(ux) \neq 2, \Delta$ . Thus  $\varphi''(ux) \in \{1, 3, 4, \ldots, \Delta - 1\}$ , which implies  $\overline{\varphi}''(x) \neq 2$  by Claim [4.2.](#page-12-0) Let  $\overline{\varphi}''(x) = \tau$ and  $\varphi''(ux) = \lambda$ . Note that if  $\tau = \Delta$  then  $\lambda \neq 1$ , as  $uz \in P_{s_1}(1, \Delta, \varphi'') = P_r(1, \Delta, \varphi'')$ . Thus if  $\tau = \Delta$  or 1, we do  $(\tau, 1) - (1, 2)$ -swaps at x. As the color of ux is not  $\Delta$  after these swaps, we get a contradiction to Claim [4.2.](#page-12-0) Thus, we assume that  $\tau \in [3, \Delta - 1]$ , and that  $P_x(1, \tau, \varphi^{\prime\prime\prime}) = P_r(1, \tau, \varphi^{\prime\prime\prime})$  for any  $(L, \varphi^{\prime\prime})$ -stable coloring  $\varphi^{\prime\prime\prime}$ , where  $L = (F, ru, u, uz, z)$  is a lollipop. Let  $w_1 \in N_{\Delta-1}(r)$  such that  $\varphi''(rw_1) = \tau$ . Applying Lemma [3.7](#page-9-0) [\(2\)](#page-9-2) on L with x playing the role of w, we find a sequence of distinct vertices  $w_1, \ldots, w_t \in \{s_{\alpha+1}, \ldots, s_{\Delta-2}\}\$ that forms either a stable rotation or a near stable rotation. As x and r are  $(1, \tau)$ -linked, we conclude that  $w_1, \ldots, w_t$  form a near stable rotation and so  $\overline{\varphi}''(w_t) = 2$ . As  $\varphi''(uz) = \Delta$ ,  $\varphi''(ur) = 2$ , if  $w_t \neq z$ , then  $\varphi''(uw_t) \in \{1, 3, 4, \ldots, \Delta - 1\}$ . This gives a contradiction to Claim [4.2.](#page-12-0) Thus we assume that  $w_t = z$ . Notice that  $r \in P_{s_1}(2, \tau, \varphi'')$  by the maximality of

 $|V(F)|$ . Since  $r \in P_x(2, \tau, \varphi'')$  by Claim [4.2,](#page-12-0) we have  $r \in P_x(2, \tau, \varphi'') = P_{s_1}(2, \tau, \varphi'')$ . So  $w_t$ is  $(2, \tau)$ -unlinked with any of  $s_1, x$  and r with respect to  $\varphi''$ . We do a  $(2, \tau)$ -swap at  $w_t$  and then shift from  $w_1$  to  $w_t$ . This gives a coloring such that  $s_1$  and x are  $(2, \tau)$ -unlinked with respect to the coloring. Again, with respect to the current coloring, r and  $s_1$  are  $(2, \tau)$ linked by the maximality of  $|V(F)|$ . We do a  $(2, \tau)$ -swap at x to get a coloring  $\varphi'''$ . Note that  $\varphi'''(ru) = \varphi''(ru) = 2$ ,  $\varphi'''(ux) = \varphi''(ux) = \lambda$ ,  $\overline{\varphi}'''(x) = 2$ , and  $\varphi'''(uz) = \varphi''(uz) = \Delta$ . Therefore,  $\varphi'''(ux) = \lambda \in \{1, 3, 4, \dots, \Delta - 1\}$ , showing a contradiction to Claim [4.2.](#page-12-0)

Thus we assume that  $N_{\Delta-1}[r]$  is  $\varphi'$ -elementary for every  $(F, \varphi)$ -stable  $\varphi' \in C^{\Delta}(G-rs_1)$ . In particular,  $N_{\Delta-1}[r]$  is  $\varphi$ -elementary, and as  $|V(F)| = 2$  and F is maximum at r, we know that  $N_{\Delta-1}[r]$  is contained in a pseudo-multifan  $S = S_{\varphi}(r, s_1 : s_1 : s_{\Delta-2})$ . Let  $\delta \in \overline{\varphi}(S) \setminus \overline{\varphi}(F)$ . By Lemma [3.5](#page-8-1) [\(c\)](#page-8-2),  $\overline{\varphi}_S^{-1}(\delta)$  is  $(2,\delta)$ - and  $(\delta,\Delta)$ -linked with  $s_1$  and the corresponding chains contain the vertex r with respect to  $\varphi$ . Recall that  $\varphi(ru) = 2, \varphi(ux) =$  $\Delta$ , and  $\overline{\varphi}(x) = 2$ . Let  $\varphi'$  be obtained from  $\varphi$  by doing a  $(2, \delta) - (\delta, \Delta) - (\Delta, 1) - (1, 2)$ -swap at x. Since  $\varphi'$  is  $(F, \varphi)$ -stable,  $\varphi'(ru) = 2$ ,  $\varphi'(ux) = \delta$ , and  $\overline{\varphi}'(x) = 2$ , we get a contradiction to Claim [4.2.](#page-12-0)  $\Box$ 

<span id="page-15-0"></span>Claim 4.7. Let  $x, y \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$  be distinct, and  $\varphi' \in C^{\Delta}(G - rs_1)$  be any  $(F, \varphi)$ -stable coloring with  $\varphi'(ru) = \alpha + 1$ . Suppose  $\overline{\varphi}'(x) \in \overline{\varphi}'(F)$  and  $\overline{\varphi}'(x) \neq 1$ . Then  $\overline{\varphi}'(y) \notin \overline{\varphi}'(F)$ . Furthermore, y and r are  $(1, \overline{\varphi}'(y))$ -linked with respect to  $\varphi'$ .

*Proof of Claim [4.7.](#page-15-0)* The second part of the claim follows easily from the first part. Since otherwise, a  $(1, \overline{\varphi}'(y))$ -swap at y implies that 1 is missing at y, contradicting the first part.

Assume to the contrary that  $\overline{\varphi}'(x) \in \overline{\varphi}'(F)$  and  $\overline{\varphi}'(y) \in \overline{\varphi}'(F)$ . We claim that we may assume  $\overline{\varphi}'(x) = \overline{\varphi}'(y) = \alpha + 1$  or  $\overline{\varphi}'(x) = \alpha + 1$  and  $\overline{\varphi}'(y) = 1$ . By doing  $(\overline{\varphi}'(x), 1) - (1, \alpha + 1)$ swaps at x, we assume that  $\overline{\varphi}'(x) = \alpha + 1$ . Since  $1, \alpha + 1 \in \overline{\varphi}'(F)$ , we still have  $\overline{\varphi}'(y) \in \overline{\varphi}'(F)$ . If  $\overline{\varphi}'(y) = \alpha + 1$ , then we are done. Otherwise, doing a  $(1, \overline{\varphi}'(y))$ -swap at y gives a desired coloring. Let  $\varphi'(ux) = \tau$  and  $\varphi'(uy) = \lambda$ . We consider now two cases to finish the proof of Claim [4.7.](#page-15-0)

# <span id="page-15-1"></span>Case A:  $\overline{\varphi}'(x) = \overline{\varphi}'(y) = \alpha + 1$ .

By Claim [4.2,](#page-12-0)  $\tau, \lambda \in \overline{\varphi}'(F) \setminus \{1\}$ . Assume, without loss of generality, that  $\tau \neq \Delta$ . Then  $\tau \in \{2, ..., \alpha + 1\}$  is a 2-inducing color of F, since F is assumed to be typical 2-inducing. By Lemma [3.6](#page-9-3) [\(d\)](#page-9-5) that  $r \in P_{s_\alpha}(\alpha+1,\Delta) = P_{s_1}(\alpha+1,\Delta)$ , we know  $\lambda \neq \Delta$ . Thus  $\lambda \in \{2, ..., \alpha + 1\}$  is also a 2-inducing color. By symmetry between x and y, we assume  $\lambda \prec \tau$ . Shift from  $s_2$  to  $s_{\lambda-1}$ , uncolor  $rs_{\lambda}$ , then color  $rs_1$  by 2. Denote the resulting coloring by  $\varphi''$ . Now  $F^* = (r, rs_\lambda, s_\lambda, rs_{\lambda+1}, s_{\lambda+1}, \ldots, rs_\alpha, s_\alpha, rs_{\lambda-1}, s_{\lambda-1}, \ldots, rs_1, s_1)$  is a new multifan with respect to  $\varphi''$  that has the same vertex set as  $F_{\varphi'}(r, s_1 : s_\alpha)$ . In this new multifan  $F^*$ ,  $\lambda$  is itself a  $\lambda$ -inducing color,  $\tau$  is a  $(\lambda + 1)$ -inducing color, and  $\alpha + 1$  is the last  $(\lambda + 1)$ -inducing color. We can further assume that  $F^*$  is typical by relabeling colors and vertices. However,  $r \in P_y(\alpha + 1, \lambda, \varphi'')$ , shows a contradiction to Lemma [3.6](#page-9-3) [\(d\)](#page-9-5) that  $r \in P_{s_\alpha}(\alpha + 1, \lambda, \varphi'') = P_{s_\lambda}(\alpha + 1, \lambda, \varphi'').$ 

Case B:  $\overline{\varphi}'(x) = \alpha + 1$  and  $\overline{\varphi}'(y) = 1$ .

We assume that x and y are  $(1, \alpha + 1)$ -linked with respect to  $\varphi'$ . For otherwise, a  $(1, \alpha + 1)$ -swap at y reduces the problem to [Case A.](#page-15-1)

We show that  $\tau, \lambda \neq \Delta$ . If this is not the case, then by swapping colors along  $P_{[x,y]}(1, \alpha +$  $1, \varphi'$  and exchanging the roles of x and y if necessary, we assume that  $\tau \neq \Delta$  and  $\lambda = \Delta$ . Let  $\varphi''$  be obtained from  $\varphi'$  by a  $(1, \Delta)$ -swap at y. By Lemma [3.6](#page-9-3) [\(d\)](#page-9-5),  $r \in P_{s_1}(\alpha +$  $1, \Delta, \varphi'') = P_{s_\alpha}(\alpha + 1, \Delta, \varphi'')$ . Thus, we can do an  $(\alpha + 1, \Delta)$ -swap at y without affecting the coloring of  $F_{\varphi''}(r, s_1 : s_\alpha)$  and  $\varphi''(ru)$ . Thus, let  $\varphi^* = \varphi''/P_y(\alpha + 1, \Delta, \varphi'')$ . We see that  $P_r(1, \alpha + 1, \varphi^*) = ruy$ , showing a contradiction to Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that r and  $s_\alpha$  are  $(1, \alpha + 1)$ -linked with respect to  $\varphi^*$ .

Since  $\tau, \lambda \neq \Delta$ , both  $\tau$  and  $\lambda$  are 2-inducing colors of F by Claim [4.2.](#page-12-0) By swapping colors along  $P_{[x,y]}(1, \alpha+1, \varphi')$  and exchanging the roles of x and y if necessary, we assume  $\tau \prec \lambda$ . Note that  $r \in P_{s_1}(\lambda, \Delta, \varphi') = P_{s_{\lambda-1}}(\lambda, \Delta, \varphi')$  and  $r \in P_{s_1}(\alpha+1, \Delta, \varphi') = P_{s_{\alpha}}(\alpha+1, \Delta, \varphi')$ 1,  $\Delta, \varphi'$  by Lemma [3.6](#page-9-3) [\(c\)](#page-9-6) and [\(d\)](#page-9-5), respectively. Let  $\varphi''$  be obtained from  $\varphi'$  by doing a  $(1, \Delta) - (\Delta, \lambda) - (\lambda, 1) - (1, \Delta) - (\Delta, \alpha + 1)$ -swap at y. Note that  $\varphi''$  is  $(F, \varphi')$ -stable, and that  $P_r(1, \alpha + 1, \varphi'') = ruy$ , showing a contradiction to Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that r and  $s_\alpha$  are  $(1, \alpha + 1)$ -linked with respect to  $\varphi''$ . □

By Claim [4.5](#page-13-2) and Claim [4.6,](#page-14-0) we let  $x, y \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$  with  $x \neq y$ , and assume that  $\varphi(ru) = \alpha + 1$  and  $\overline{\varphi}(x) = \alpha + 1$ . By Claim [4.7,](#page-15-0) we also assume that  $\overline{\varphi}(y) =$  $\delta \in [\alpha + 2, \Delta - 1]$  and y and r are  $(1, \delta)$ -linked with respect to such a coloring  $\varphi$ . Let  $w_1 \in N_{\Delta-1}(r)$  such that  $\varphi(rw_1) = \delta$  and  $L = (F, ru, u, ux, x)$ . By Claim [4.7,](#page-15-0) for any Lstable  $\varphi' \in \mathcal{C}^{\Delta}(G-rs_1)$ , it holds that  $y \in P_r(1,\delta,\varphi')$ . Applying Lemma [3.7](#page-9-0) [\(2\)](#page-9-2) on L with y playing the role of w, we find a sequence of distinct vertices  $w_1, \ldots, w_t \in \{s_{\alpha+1}, \ldots, s_{\Delta-2}\}\$ that forms either a stable rotation or a near stable rotation. Since y and r are  $(1, \delta)$ -linked with respect to  $\varphi, w_1, \ldots, w_t$  is a near stable rotation, i.e.,  $\overline{\varphi}(w_t) = \alpha + 1$ .

<span id="page-16-0"></span>Claim 4.8.  $|N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)| \geq 3$ .

Proof of Claim [4.8.](#page-16-0) Let  $\varphi(ux) = \tau$  and  $\varphi(uy) = \lambda$ . Since  $\varphi(ru) = \alpha + 1$ , we have  $\alpha + 1 \notin$  $\{\tau, \lambda, \delta\}$ . By Claim [4.3,](#page-13-1)  $\tau \in \overline{\varphi}(F) \setminus \{1\}$ , and

$$
\begin{cases} s_1, s_\alpha \notin N_{\Delta - 1}(u) & \text{if } \tau = \Delta, \end{cases}
$$
 (1)

$$
(2) \quad s_{\tau-1}, s_{\tau} \notin N_{\Delta-1}(u) \quad \text{if } \tau \neq \Delta.
$$

<span id="page-16-2"></span><span id="page-16-1"></span>We then show that

$$
\begin{cases} s_1, s_\alpha \notin N_{\Delta - 1}(u) & \text{if } \lambda = \Delta, \end{cases}
$$
 (3)

$$
(4) \quad s_{\lambda-1}, s_{\lambda} \notin N_{\Delta-1}(u) \quad \text{if } \lambda \neq \Delta.
$$

To see this, let  $\varphi'$  be obtained from  $\varphi$  by first doing a  $(1, \alpha + 1)$ -swap at both x and w<sub>t</sub>, and then shift from  $w_1$  to  $w_t$ . Now,  $\overline{\varphi}'(r) = \delta$  and  $\varphi'(ux) = \varphi(ux) = \tau$ . Let  $\varphi'' =$  $\varphi'/P_y(\alpha+1,\delta,\varphi')$ . Note that  $\varphi''(ux) = \varphi'(ux) = \tau$  and  $\varphi''(uy) = \varphi'(uy) = \lambda$ . Applying Claim [4.2](#page-12-0) to the coloring  $\varphi''$ , we get  $\varphi''(uy) = \lambda \in \overline{\varphi}''(F) \setminus {\delta}.$  As  $\tau, \lambda, \delta, \alpha + 1 \in \overline{\varphi}''(F)$ and they are all distinct,  $|V(F)| \geq |\{\delta, \tau, \lambda, \alpha + 1\}| - 1 = 3$  $|V(F)| \geq |\{\delta, \tau, \lambda, \alpha + 1\}| - 1 = 3$ . Then (3) and ([4](#page-16-1)) follow from Claim [4.3.](#page-13-1) These two facts, together with ([1](#page-16-2)) and ([2](#page-16-2)), imply eithers<sub>1</sub>,  $s_{\alpha}, s_{\lambda-1}, s_{\lambda} \notin$  $N_{\Delta-1}(u)$ , ors<sub>1</sub>, s<sub>α</sub>, s<sub>τ</sub>-1, s<sub>τ</sub>  $\notin N_{\Delta-1}(u)$ . Note that s<sub>1</sub>  $\neq s_{\alpha}$  by  $|V(F)| \geq 3$ . We obtain  $|N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)| \geq 3$  from the above unless either  $\lambda = \alpha = 2$  or  $\tau = \alpha = 2$ .

Therefore we assume  $\alpha = 2$  and  $\{\lambda, \tau\} = \{2, \Delta\}$ . Furthermore, we may assume that  $|N_{\Delta-1}(u)\setminus N_{\Delta-1}(r)|=2$  $|N_{\Delta-1}(u)\setminus N_{\Delta-1}(r)|=2$  $|N_{\Delta-1}(u)\setminus N_{\Delta-1}(r)|=2$ , since (1) to ([4](#page-16-1)) imply that  $s_1, s_2 \notin N_{\Delta-1}(u)$ . Therefore  $N_{\Delta-1}(r)\setminus$  $\{s_1, s_2\} \subseteq N_{\Delta-1}(u)$ . In particular,  $w_t \in N_{\Delta-1}(u)$ . Since r and  $s_\alpha$  are  $(1, \alpha + 1)$ -linked with respect to  $\varphi$  and  $\overline{\varphi}(w_t) = \alpha + 1$ , it follows that  $\varphi(uw_t) \neq 1$ . This, together with the facts that  $\overline{\varphi}(F) = \{1, 2, 3, \Delta\}, \varphi(ru) = 3$ , and  $\{\lambda, \tau\} = \{2, \Delta\},$  implies that  $\varphi(uw_t) \in [4, \Delta - 1],$ showing a contradiction to Claim [4.2.](#page-12-0)  $\Box$ 

By Claim [4.8,](#page-16-0) let  $z \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$  with  $z \neq x, y$ . By Claim [4.7,](#page-15-0) we assume  $\overline{\varphi}(z) = \lambda \in [\alpha + 2, \Delta - 1],$  and z and r are  $(1, \lambda)$ -linked with respect to  $\varphi$ . Since  $\overline{\varphi}(y) = \delta$ , and also y and r are  $(1, \delta)$ -linked with respect to  $\varphi$ , we have  $\lambda \neq \delta$ .

Recall  $w_1, \ldots, w_t$  is a near stable rotation at r with  $\varphi(rw_1) = \delta =: \delta_1$ . Let  $\overline{\varphi}(w_i) = \delta_{i+1}$ for each  $i \in [1, t-1]$ . As z and r are  $(1, \lambda)$ -linked with respect to  $\varphi$  and  $w_i$  and r are  $(1, \delta_{i+1})$ -linked for each  $i \in [1, t-1], \lambda \neq \delta_i$  for each  $i \in [2, t]$ . Let  $\lambda_1 = \lambda$  and  $w_1^*$  be the neighbor of r such that  $\varphi(rw_1^*) = \lambda_1$ . For any  $(L, \varphi)$ -stable coloring  $\varphi' \in C^{\Delta}(G - rs_1)$ ,  $z \in P_r(1, \lambda, \varphi')$ . Applying Lemma [3.7](#page-9-0) [\(2\)](#page-9-2) on  $L = (F, ru, u, ux, x)$  and z, we find a sequence of distinct vertices  $w_1^*, \ldots, w_k^* \in \{s_{\alpha+1}, \ldots, s_{\Delta-2}\}\$  that forms either a stable rotation or a near stable rotation. If  $\overline{\varphi}(w_k^*) = \lambda_1$ , then since  $w_k^*$  and r are  $(1, \lambda_1)$ -linked, a  $(1, \lambda_1)$ -swap at z gives a contradiction to Claim [4.7.](#page-15-0) Thus  $\overline{\varphi}(w_k^*) = \alpha + 1$ . Let  $\overline{\varphi}(w_i^*) = \lambda_{i+1}$  for each  $i \in [1, k - 1].$ 

Recall that  $w_1^* \neq w_i$  for each  $i \in [1, t]$ . Furthermore, as  $w_i$  and r are  $(1, \delta_{i+1})$ -linked for each  $i \in [1, t-1]$  and  $w_j^*$  and  $r$  are  $(1, \lambda_{j+1})$ -linked for each  $j \in [1, k-1]$ ,  $w_1^* \neq w_i$  for each  $i \in [1, t]$  implies that  $\lambda_2 \notin \{\delta_1, \ldots, \delta_t\}$ . Consequently,  $w_2^* \neq w_i$  for each  $i \in [1, t]$ . Repeating the same process, we get  $w_j^* \neq w_i$  for each  $j \in [1, k]$  and each  $i \in [1, t]$ .

We claim that  $w_t$  and x are  $(1, \alpha + 1)$ -linked with respect to  $\varphi$ . For otherwise, first doing a  $(1, \alpha + 1)$ -swap at  $w_t$ , then shift from  $w_1$  to  $w_t$  gives a coloring  $\varphi'$  such that  $\varphi'(ru) = \varphi(ru) = \alpha + 1$ ,  $\overline{\varphi}'(y) = \overline{\varphi}'(r) = \delta_1$ , while  $\overline{\varphi}'(x) = \alpha + 1$ . Based on  $\varphi'$ , after doing a  $(1, \delta_1)$ -swap on all  $(1, \delta_1)$ -chains in  $G - rs_1$ , we obtain an  $(F, \varphi)$ -stable coloring  $\varphi''$ . However,  $\overline{\varphi}''(x) = \alpha + 1$  and  $\overline{\varphi}''(y) = 1$ , showing a contradiction to Claim [4.7.](#page-15-0) As  $w_t$  and x are  $(1, \alpha + 1)$ -linked, we do a sequence of Kempe changes around r from  $w_k^*$  to  $w_1^*$  as below: let  $\varphi_0 = \varphi$  and  $\lambda_{k+1} = \alpha + 1$ ,

$$
\varphi_j = \varphi_{j-1}/P_{w^*_{k-(j-1)}}(1, \lambda_{k+1-(j-1)}, \varphi_{j-1})
$$
 for each  $j \in [1, k]$ .

Note that

 $P_r(1, \lambda_{k-(j-1)}, \varphi_j) = rw_{k-(j-1)}^*$  for each  $j \in [1, k],$ 

and that  $\varphi_k$  is  $(F, \varphi)$ -stable,  $\varphi_k(ru) = \varphi(ru)$ ,  $\varphi_k(ux) = \varphi(ux)$ , and  $\overline{\varphi}_k(x) = \overline{\varphi}(x) = \alpha + 1$ ,

but z and r are  $(1, \lambda)$ -unlinked with respect to  $\varphi_k$ . Now doing a  $(1, \lambda)$ -swap at z gives a contradiction to Claim [4.7.](#page-15-0) This finishes the proof of Theorem [2.3.](#page-2-0) 口

# 5 Proof of Theorem [2.4](#page-2-4)

**Theorem 2.4.** If G is an HZ-graph with maximum degree  $\Delta \geq 4$ , then for any two adjacent vertices  $x, y \in V_{\Delta-1}$ ,  $N_{\Delta}(x) = N_{\Delta}(y)$ .

*Proof.* Assume to the contrary that  $N_{\Delta}(x) \neq N_{\Delta}(y)$ . Then there exists a vertex  $r \in$  $N_{\Delta}(x) \setminus N_{\Delta}(y)$ . Equivalently,  $x \in N_{\Delta-1}(r)$  and  $y \notin N_{\Delta-1}(r)$ . By Theorem [4.1](#page-11-1) [\(ii\)](#page-11-3), let  $s_1 \in N_{\Delta-1}(r)$  and  $\varphi \in C^{\Delta}(G - rs_1)$ , and  $F = F_{\varphi}(r, s_1 : s_{\alpha})$  be the typical 2-inducing multifan such that either  $V(F) = N_{\Delta-1}[r]$  or F is contained in a pseudo-multifan S with  $V(S) = N_{\Delta-1}[r]$ . Let  $N_{\Delta-1}(r) = \{s_1, \ldots, s_{\Delta-2}\}\.$  We consider two cases according to if  $x \in V(F)$  to finish the proof.

Assume first that  $x \notin V(F)$ . This implies that  $V(F) \neq N_{\Delta-1}[r]$ . Applying Theorem [4.1](#page-11-1) [\(ii\)](#page-11-3), it then follows that  $N_{\Delta-1}[r]$  is the vertex set of a typical 2-inducing pseudo-multifan. Let  $\overline{\varphi}(x) = \delta$  and  $\overline{\varphi}(y) = \lambda$ . Since  $V(S) = N_{\Delta-1}[r]$  is  $\varphi$ -elementary,  $\delta, \lambda \in \overline{\varphi}(S)$ . By Lemma [3.1](#page-5-0) [\(b\)](#page-8-3) or Lemma [3.5](#page-8-1) (b), we know that  $\overline{\varphi}_S^{-1}(\lambda)$  and r are  $(1, \lambda)$ -linked and x and r are  $(1, \delta)$ -linked. By doing a  $(\lambda, 1) - (1, \delta)$ -swap at y, we find  $(S, \varphi)$ -stable  $\varphi' \in C^{\Delta}(G - rs_1)$ such that  $\overline{\varphi}'(y) = \delta$ . Let  $\varphi'(xy) = \tau$ . Then  $P_x(\delta, \tau, \varphi') = xy$ , showing a contradiction to Lemma [3.5](#page-8-1) [\(c\)](#page-8-2) or [\(d\)](#page-8-4) depending on  $\tau \in \overline{\varphi}'(F)$  or  $\tau \in \overline{\varphi}'(S) \setminus \overline{\varphi}'(F)$ .

Assume then that  $x \in V(F)$ . We claim that we may assume  $x = s_1$ . Let  $x = s_i$  for some  $i \in [1, \alpha]$ , and  $\varphi'$  be obtained from  $\varphi$  by shift from  $s_2$  to  $s_{i-1}$ , uncoloring  $rs_i$ , and coloring  $rs_1$  by 2. The sequence  $F^* = (r, rs_i, s_i, rs_{i+1}, s_{i+1}, \ldots, rs_\alpha, s_\alpha, rs_{i-1}, s_{i-1}, \ldots, rs_1, s_1)$  is a multifan with respect to  $\varphi'$ . Since the shift and "changing" the uncolored edge operation like above is revertible, and  $V(S)$  and  $\overline{\varphi}(S)$  are kept unchanged under such an operation, we conclude that  $N_{\Delta-1}[r]$  is still the vertex set of a pseudo-multifan. By permuting the name of colors and the label of the vertices in  $N_{\Delta-1}(r)$ , we may assume that  $x = s_1$ . Still denote the current coloring by  $\varphi$ , the multifan by F, and the pseudo-multifan by S.

By doing a  $(1,\overline{\varphi}(y))$ -swap at y, we assume  $\overline{\varphi}(y) = 1$ . Let  $\varphi(s_1y) = \tau$ . By exchanging the roles of the color 2 and  $\Delta$  if necessary, we may assume that  $\varphi(s_1y)$  is either a 2-inducing color of F or is a color from  $\overline{\varphi}(S) \setminus \overline{\varphi}(F)$ . Let  $\varphi' = \varphi/P_y(1, \Delta, \varphi)$ . Now  $P_{s_1}(\tau, \Delta, \varphi') = s_1 y$ . This gives a contradiction to Lemma [3.2](#page-5-3) [\(b\)](#page-6-0) that  $s_1$  and  $\overline{\varphi}'^{-1}_F(\tau)$  are  $(\tau, \Delta)$ -linked if  $\tau$  is 2-inducing, and gives a contradiction to Lemma [3.5](#page-8-1) [\(c\)](#page-8-2) that  $s_1$  and  $\overline{\varphi}'^{-1}_S(\tau)$  are  $(\tau, \Delta)$ -linked if  $\tau \in \overline{\varphi}(S) \setminus \overline{\varphi}(F)$ . □

### 6 Proof of Theorem [2.5](#page-2-1)

**Theorem 2.5.** Let G be an HZ-graph with maximum degree  $\Delta \geq 7$  and  $u, r \in V_{\Delta}$ . If  $N_{\Delta-1}(u) \neq N_{\Delta-1}(r)$  and  $N_{\Delta-1}(u) \cap N_{\Delta-1}(r) \neq \emptyset$ , then  $|N_{\Delta-1}(u) \cap N_{\Delta-1}(r)| = \Delta - 3$ , i.e.

 $|N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)| = |N_{\Delta-1}(r) \setminus N_{\Delta-1}(u)| = 1.$ 

*Proof.* Assume to the contrary that there exist  $u, r \in N_\Delta$  such that  $1 \leq |N_{\Delta-1}(r)|$  $N_{\Delta-1}(u) \leq \Delta-4$ . By Theorem [4.1](#page-11-1) [\(ii\)](#page-11-3), there exist  $s_1 \in N_{\Delta-1}(r)$  and  $\varphi \in C^{\Delta}(G-rs_1)$  such that  $N_{\Delta-1}[r]$  is the vertex set of a typical 2-inducing pseudo-multifan. By this assumption of being typical, we have  $N_{\Delta-1}(r) = \{s_1, \ldots, s_{\Delta-2}\}, \overline{\varphi}(r) = 1$ , and  $\overline{\varphi}(s_1) = \{2, \Delta\}.$  Let  $x, y \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$  be two distinct vertices, and  $S := S_{\varphi}(r, s_1 : s_{\alpha} : s_{\Delta-2})$  be this pseudo-multifan with  $F_{\varphi}(r, s_1 : s_{\alpha})$  being the typical 2-inducing multifan contained in S. Since  $V(S) = N_{\Delta-1}[r]$  and  $V(S)$  is  $\varphi$ -elementary, it follows that  $\overline{\varphi}(S) = [1, \Delta]$ . We consider two cases.

### Case 1:  $V(S) \neq V(F)$ .

In this case, we will repeatedly apply Lemma [3.5](#page-8-1) [\(b\)](#page-8-3), [\(c\)](#page-8-2) or [\(d\)](#page-8-4). Assume first that for each  $i \in [1, \alpha], s_i \notin N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$ . Then by  $N_{\Delta-1}(r) \cap N_{\Delta-1}(u) \neq \emptyset$ , there exists  $w_1 \in \{s_{\alpha+1}, \ldots, s_{\Delta-2}\}\$  such that  $w_1 \in N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$ . Let  $\varphi(rw_1) = \delta_1$  and  $\overline{\varphi}(w_1) = \delta_2$ . Note that  $\delta_1, \delta_2 \in \overline{\varphi}(S) \setminus \overline{\varphi}(F)$ . We claim that we may assume  $\varphi(ux) = \delta_2$ . Otherwise, let  $\varphi(ux) = \delta^* \neq \delta_2$ . By Lemma [3.5](#page-8-1) [\(b\)](#page-8-3), [\(c\)](#page-8-2) or [\(d\)](#page-8-4) depending on what  $\overline{\varphi}(x)$  is, we can do a  $(\overline{\varphi}(x), \delta_2) - (\delta_2, \delta^*)$ -swap at x in getting an  $(S, \varphi)$ -stable coloring, still call it  $\varphi$  such that  $\varphi(ux) = \delta_2$ . Let  $\varphi(w_1u) = \tau$  and  $\varphi'$  be obtained from  $\varphi$  by doing a  $(\overline{\varphi}(x), \delta_1) - (\delta_1, \tau)$ -swap at x. By Lemma [3.5](#page-8-1) [\(b\)](#page-8-3), [\(c\)](#page-8-2) or [\(d\)](#page-8-4),  $\varphi'$  is  $(S, \varphi)$ -stable such that  $\varphi'(w_1u) = \varphi(w_1u) = \tau$ and  $\overline{\varphi}'(x) = \tau$ . However,  $P_{w_1}(\delta_2, \tau, \varphi') = w_1 u x = P_x(\delta_2, \tau, \varphi')$ , showing a contradiction to Lemma [3.5](#page-8-1) [\(b\)](#page-8-3), [\(c\)](#page-8-2) or [\(d\)](#page-8-4) (depending on if  $\tau = 1$ ,  $\tau \in \overline{\varphi}(F) \setminus \{1\}$  or  $\tau \in \overline{\varphi}(S) \setminus \overline{\varphi}(F)$ ) that  $w_1$  and  $\overline{\varphi}_S^{-1}(\tau)$  are  $(\delta_2, \tau)$ -linked with respect to  $\varphi'$ .

Assume now that there exists  $s_i \in N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$  for some  $i \in [1, \alpha]$ . By shift from  $s_2$  to  $s_{i-1}$ , uncoloring  $rs_i$ , and coloring  $rs_1$  by 2, we obtain a new multifan  $F^*$  =  $(r, rs_i, s_i, rs_{i+1}, s_{i+1}, \ldots, rs_{\alpha}, s_{\alpha}, rs_{i-1}, s_{i-1}, \ldots, rs_1, s_1)$ . By permuting the name of colors and the label of the vertices in  $N_{\Delta-1}(r)$  such that  $i+1$  is permuted to 2 and  $s_i$  is renamed as  $s_1$ , we assume that  $s_1 \in N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$  and  $F^*$  is a typical multifan.

Recall that  $x \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$ . Let  $\overline{\varphi}(x) = \lambda$ . By Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) or Lemma [3.5](#page-8-1) [\(b\)](#page-8-3), we know that  $\overline{\varphi}_S^{-1}(\lambda)$  and r are  $(1, \lambda)$ -linked. By doing a  $(1, \lambda)$ -swap at x if necessary, we assume  $\overline{\varphi}(x) = 1$ . By exchanging the roles of the colors 2 and  $\Delta$ , we assume that  $\varphi(s_1u)$  equals 1, or is a 2-inducing color of F, or is a color from  $\overline{\varphi}(S) \setminus \overline{\varphi}(F)$ . Note that by Lemma [3.5](#page-8-1) [\(c\)](#page-8-2), for a color  $\delta \in \overline{\varphi}(S) \setminus \overline{\varphi}(F)$ , and for any color  $\tau \in \overline{\varphi}(F)$ ,  $\overline{\varphi}_S^{-1}(\delta)$  and  $\overline{\varphi}_S^{-1}(\tau)$ are  $(\delta, \tau)$ -linked and  $r \in P_{\overline{\varphi}_S^{-1}(\delta)}(\delta, \tau, \varphi)$ .

Let  $\varphi(ux) = \tau$ . If  $\tau$  is a 2-inducing color of F or is from  $\overline{\varphi}(S) \setminus \overline{\varphi}(F)$ , we do  $(1, \Delta)$  –  $(\Delta, \tau)$  –  $(\tau, 1)$ -swaps at x. If  $\tau$  is a  $\Delta$ -inducing color of F, let  $\delta \in \overline{\varphi}(S) \setminus \overline{\varphi}(F)$ , we do  $(1,\delta) - (\delta,\tau) - (\tau,1) - (1,\Delta) - (\Delta,\delta) - (\delta,1)$ -swaps at x. In both cases, we let  $\varphi'$  be the resulting coloring. We have  $\varphi'(ux) = \Delta$  and  $\overline{\varphi}'(x) = 1$ . Since  $\varphi(s_1u) \neq \Delta, \tau$ , still  $\varphi'(s_1u)$ equals 1, or is a 2-inducing color of F, or is from  $\overline{\varphi}(S) \setminus \overline{\varphi}(F)$ .

Let  $\varphi'(s_1u) = \gamma$ . Since  $s_1$  and r are  $(1,\Delta)$ -linked with respect to  $\varphi', \gamma \neq 1$ . Thus,  $\gamma$  is a 2-inducing color of F, or is from  $\overline{\varphi}(S) \setminus \overline{\varphi}(F)$ . By Lemma [3.2](#page-5-3) [\(a\)](#page-6-1) or Lemma [3.5](#page-8-1) [\(c\)](#page-8-2),  $u \in P_x(1, \gamma, \varphi')$  (otherwise,  $s_1$  and x are  $(\gamma, \Delta)$ -linked after a  $(1, \gamma)$ -swap at x). Let  $\varphi'' = \varphi' / P_x(1, \gamma, \varphi')$ . Now  $\varphi''(s_1u) = 1$ ,  $\overline{\varphi}''(x) = \gamma$ , and  $K = (r, rs_1, s_1, s_1u, u, ux, x)$  is a Kierstead path with respect to  $rs_1$  and  $\varphi''$ . Let  $\delta \in \overline{\varphi}''(S) \setminus \overline{\varphi}''(F)$ . If  $\gamma \in \overline{\varphi}''(S) \setminus \overline{\varphi}''(F)$ , we do nothing. Otherwise, we do a  $(\gamma, \delta)$ -swap at x (by Lemma [3.5](#page-8-1) [\(c\)](#page-8-2), this swap does not end at any vertex of S). Denote by  $\varphi'''$  the resulting coloring. Since  $d_G(s_1) = \Delta - 1$ , in both cases, by Lemma [3.3](#page-6-2) (b), x and  $s_1$  are  $(2, \overline{\varphi}'''(x))$ -linked. Since  $\overline{\varphi}'''(x) \in \overline{\varphi}'''(S) \setminus \overline{\varphi}''(F)$ , we achieve a contradiction to Lemma [3.5](#page-8-1) [\(c\)](#page-8-2).

**Case 2:**  $V(S) = V(F)$ .

We claim that we may choose  $s_1$  such that  $s_1 \in N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$ . If  $s_1 \in N_{\Delta-1}(u) \cap$  $N_{\Delta-1}(r)$ , then we are done. Otherwise, let  $s_i \in N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$ . We shift from  $s_2$  to  $s_{i-1}$ , uncolor  $rs_i$  and color  $rs_1$  by 2. By relabeling colors and vertices, we may assume that  $s_1 \in N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$  and  $F^* = (r, rs_i, s_i, rs_{i+1}, s_{i+1}, \ldots, rs_\alpha, s_\alpha, rs_{i-1}, s_{i-1}, \ldots, rs_1, s_1)$ is a typical multifan. We let  $F_{\varphi}(r, s_1 : s_{\alpha} : s_{\Delta-2})$  be such a typical multifan.

For a coloring  $\psi \in C^{\Delta}(G - rs_1)$ , if  $\overline{\varphi}(s_1) = \overline{\psi}(s_1)$ ,  $\overline{\varphi}(F) = \overline{\psi}(F)$ , and some permutation of F is still a multifan with respect to rs<sub>1</sub> and  $\psi$ , we call  $\psi$  a near  $(F, \varphi)$ -stable coloring. As only colors in  $\overline{\varphi}(s_1)$  will be essential for the proof, we will not distinguish between  $\varphi$  and any near  $(F, \varphi)$ -stable coloring. As the vertex set of all the resulting multifans is always  $N_{\Delta-1}[r]$ , for a color  $\alpha \in [1, \Delta]$ , we use  $\overline{\psi}^{-1}(\alpha)$  to denote the vertex from  $N_{\Delta-1}[r]$  that misses  $\alpha$  with respect to  $\psi$ .

Let  $\psi \in C^{\Delta}(G - rs_1)$  be near  $(F, \varphi)$ -stable and  $F^*$  be the corresponding multifan. The following two facts will be used frequently in the proof without mentioning.

- Fact 1 For any  $i \in [2, \Delta]$ , since r and  $\overline{\psi}^{-1}(i)$  are  $(1, i)$ -linked by Lemma [3.1](#page-5-0) [\(b\)](#page-5-1), doing a  $(1, i)$ swap at vertices outside of  $V(F^*)$  gives an  $(F^*, \psi)$ -stable and so a near  $(F, \varphi)$ -stable coloring.
- Fact 2 For any 2-inducing color  $\tau$  and  $\Delta$ -inducing color  $\delta$  of  $F^*, \overline{\psi}^{-1}(\tau)$  and  $\overline{\psi}^{-1}(\delta)$  are  $(\tau, \delta)$ -linked by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Thus doing a  $(\tau, \delta)$ -swap at a vertex outside of  $V(F^*)$ or, when  $\tau \neq 2$  and  $\delta \neq \Delta$ , doing a  $(\tau, \delta)$ -swap at  $\overline{\psi}^{-1}(\tau)$  if  $r \notin P_{\overline{\psi}^{-1}(\tau)}(\tau, \delta, \psi)$  gives a near  $(F^*, \psi)$ -stable and so a near  $(F, \varphi)$ -stable coloring.

We denote by  $S(u; s_1, x, y)$  the star subgraph of G that is centered at u consisting of edges us<sub>1</sub>, ux, and uy. Recall that  $x, y \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$  are distinct vertices.

<span id="page-20-0"></span>Claim 6.1. We may assume that  $\overline{\varphi}(x) = 2$  and  $\overline{\varphi}(y) = \Delta$  or  $\overline{\varphi}(x) = \overline{\varphi}(y) = \Delta$ .

*Proof of Claim [6.1.](#page-20-0)* By doing  $(\overline{\varphi}(x), 1) - (1, 2)$ -swaps at x, we find  $(F, \varphi)$ -stable  $\varphi' \in C^{\Delta}(G$ rs<sub>1</sub>) such that  $\overline{\varphi}'(x) = 2$ . Now, let  $\overline{\varphi}'(y) = \lambda$ . If  $\lambda = 2$ , then doing  $(2, 1) - (1, \Delta)$ -swaps at both x and y, we find  $(F, \varphi')$ -stable  $\varphi'' \in C^{\Delta}(G - rs_1)$  such that  $\overline{\varphi}''(x) = \overline{\varphi}''(y) = \Delta$ . If  $\lambda \neq 2$ , by doing  $(\lambda, 1) - (1, \Delta)$ -swaps at y, we find  $(F, \varphi')$ -stable  $\varphi'' \in C^{\Delta}(G - rs_1)$  such that  $\overline{\varphi}''(x) = 2$  and  $\overline{\varphi}''(y) = \Delta$ . As  $\varphi''$  is  $(F, \varphi')$ -stable and  $\varphi'$  is  $(F, \varphi)$ -stable, it follows that  $\varphi''$  is  $(F, \varphi)$ -stable. So we can take  $\varphi''$  to be  $\varphi$ .  $\Box$ 

By Claim [6.1,](#page-20-0) we assume  $\overline{\varphi}(x) = 2$  and  $\overline{\varphi}(y) = \Delta$  or  $\overline{\varphi}(x) = \overline{\varphi}(y) = \Delta$  and so consider two cases below.

### <span id="page-21-2"></span>Subcase 2.1:  $\overline{\varphi}(x) = 2$  and  $\overline{\varphi}(y) = \Delta$ .

Note that by doing first a  $(2, 1)$ -swap at x, then a  $(1, \Delta)$ -swap at both x and y, and finally a (1, 2)-swap at y, we can always identify this current case with the case that  $\overline{\varphi}(x) = \Delta$ and  $\overline{\varphi}(y) = 2$ . Let  $\varphi(ux) = \tau$  and  $\varphi(uy) = \lambda$ . By exchanging the roles of the two colors 2 and  $\Delta$ , we consider two cases below: (A)  $\varphi(uy) = \lambda = 1$ ; and (B)  $\varphi(uy) = \lambda$  is 2-inducing. (When  $\varphi(uy)$  is  $\Delta$ -inducing, by assuming  $\overline{\varphi}(x) = \Delta$  and  $\overline{\varphi}(y) = 2$ , the argument will be symmetric to the argument for case (B) above.)

In both cases of (A) and (B), we do  $(\Delta, \lambda) - (\lambda, 1)$ -swaps at y and call the resulting coloring  $\varphi_1$  and the resulting multifan  $F_1$ . Note that  $\varphi_1$  is near  $(F, \varphi)$ -stable. Let  $\varphi_1(s_1u)$  = δ. The current coloring on  $S(u; s_1, x, y)$  is as shown in  $J_1$  of Figure [2.](#page-21-0)

<span id="page-21-1"></span>**Claim 6.2.** The color  $\varphi_1$  can be modified into a near  $(F_1, \varphi_1)$ -stable coloring such that the color on  $S(u; s_1, x, y)$  is as in  $J_2$  of Figure [2.](#page-21-0)



<span id="page-21-0"></span>Figure 2: Coloring of  $S(u; s_1, x, y)$ 

*Proof of Claim [6.2.](#page-21-1)* Since  $s_1$  and r are  $(1, \Delta)$ -linked by Lemma [3.1](#page-5-0) [\(b\)](#page-5-1), we know  $\varphi(s_1u)$  =  $\delta \neq 1$ . If  $u \in P_y(1, \delta, \varphi_1)$ , then a  $(1, \delta)$ -swap at y gives  $J_2$ . Thus, we assume  $u \notin P_y(1, \delta, \varphi_1)$ . This implies that  $\delta$  is  $\Delta$ -inducing with respect to  $F_1$  and  $\varphi_1$ . (Otherwise, after a  $(1, \delta)$ -swap at y,  $s_1$  and  $\overline{\varphi_1}^{-1}(\delta)$  are  $(\delta, \Delta)$ -unlinked, showing a contradiction to Lemma [3.2](#page-5-3) [\(a\)](#page-6-1).)

Let  $\varphi_2 = \varphi_1/P_y(1,\delta,\varphi_1)$  (see Figure [2\)](#page-21-0). We claim that  $\tau$  is 2-inducing with respect to  $F_1$  and  $\varphi_2$ . Otherwise  $\tau$  is 1 or is  $\Delta$ -inducing with respect to  $F_1$  and  $\varphi_2$ . We do  $(2, \tau) - (\tau, 1)$ -swaps at x and call the resulting coloring  $\varphi'_2$  and the resulting multifan  $F'_1$ . Again, as  $P_{s_1}(\delta, \Delta, \varphi_2') = s_1uy$ ,  $\delta$  is still a  $\Delta$ -inducing color of  $F'_1$  with respect to  $\varphi_2'$  by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Since  $\varphi'_2(ux) = 2$ , we must have  $u \in P_x(1, \delta, \varphi'_2)$ : otherwise, after a  $(1, \delta)$ swap at x,  $s_1$  and x are  $(2, \delta)$ -linked with respect to the current coloring, contradicting Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Now, let  $\varphi_2^*$  be obtained from  $\varphi_2'$  by doing a  $(1, \delta)$ -swap at both x and y. We get  $P_{s_1}(1, \Delta, \varphi_2^*) = s_1uy$ , showing a contradiction to Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that  $s_1$  and r are (1,  $\Delta)$ -linked with respect to  $\varphi_2^*.$ 

Thus  $\tau$  is 2-inducing with respect to  $F_1$  and  $\varphi_2$ . First, we let  $\varphi_3 = \varphi_2/P_x(1, 2, \varphi_2)$ . Note that  $u \notin P_x(1, \delta, \varphi_3)$  and  $u \notin P_y(1, \delta, \varphi_3)$ . Since otherwise, after a  $(1, \delta)$ -swap at both x and y, s<sub>1</sub> and y are  $(1, \Delta)$ -linked with respect to the current coloring, showing a contradiction to Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that  $s_1$  and r are  $(1, \Delta)$ -linked. Since  $\delta$  is  $\Delta$ -inducing and  $\tau$  is 2-inducing with respect to  $F_1$  and  $\varphi_3$ ,  $\overline{\varphi_3}^{-1}(\delta)$  and  $\overline{\varphi_3}^{-1}(\tau)$  are  $(\delta, \tau)$ -linked by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Then we let  $\varphi_4$  be obtained from  $\varphi_3$  by doing a  $(1, \delta)$ -swap at both x and y (see Figure [2\)](#page-21-0), and doing a  $(\tau, \delta)$ -swap at  $\overline{\varphi}_3^{-1}(\delta)$  (and so also at  $\overline{\varphi}_3^{-1}(\tau)$ ). Since  $\varphi_4$  is near  $(F_1, \varphi_3)$  stable, we let  $F_2$  be the resulting multifan. Note that  $\delta$  is a 2-inducing color and  $\tau$  is a  $\Delta$ -inducing color of  $F_2$  with respect to  $\varphi_4$ . As a consequence,  $u \in P_y(1, \delta, \varphi_4)$ . Since otherwise, after a  $(1, \delta)$ -swap at y, s<sub>1</sub> and y are  $(\delta, \Delta)$ -linked, contradicting Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). We then let  $\varphi_5$  be obtained from  $\varphi_4$  by doing a  $(1, \delta)$ -swap at both x, y (and so also u), and then a  $(1, 2)$ -swap at x. We obtain the desired coloring on  $S(u; s_1, x, y)$ .  $\Box$ 

By Claim [6.2,](#page-21-1) we let  $\varphi_2 \in C^{\Delta}(G - rs_1)$  be a near  $(F_1, \varphi_1)$ -stable coloring and  $F_2$  be a corresponding multifan such that under  $\varphi_2$  $\varphi_2$ , the color on  $S(u; s_1, x, y)$  is as in Figure 2  $J_2$ . Now  $K = (r, rs_1, s_1, s_1, u, uy, y)$  is a Kierstead path with respect to  $rs_1$  and  $\varphi_2$ . Since  $d_G(s_1) = \Delta - 1$ , by Lemma [3.3](#page-6-2) (b), y and  $s_1$  are  $(2, \delta)$ -linked. This implies that  $\delta$  must be a 2-inducing color of  $F_2$ , as otherwise,  $s_1$  and  $\overline{\varphi}_2^{-1}(\delta)$  should be  $(2,\delta)$ -linked. If  $\tau$  is  $\Delta$ -inducing of  $F_2$ , then as  $\overline{\varphi}_2^{-1}(\delta)$  and  $\overline{\varphi}_2^{-1}(\tau)$  are  $(\delta, \tau)$ -linked by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1), we do a  $(\delta, \tau)$ -swap at y. Again by Lemma [3.3](#page-6-2) (b), y and  $s_1$  are  $(2, \tau)$ -linked, showing a contradiction to Lemma [3.2](#page-5-3) [\(a\)](#page-6-1) that  $\overline{\varphi}_2^{-1}(\tau)$  and  $s_1$  are  $(2, \tau)$ -linked. Therefore,  $\tau$  is a 2-inducing color of F<sub>2</sub>. We first do  $(2, 1) - (1, \Delta)$  swaps at x, and let  $\varphi_3$  be the resulting coloring (see Figure [3\)](#page-23-0). At this step,  $\varphi_3(s_1u) = 1$ ,  $\varphi_3(uy) = \Delta$ ,  $\overline{\varphi}_3(y) = \delta$ , and y and  $s_1$  are  $(2,\delta)$ -linked with respect to  $\varphi_3$  by Lemma [3.3](#page-6-2) (b). Call this fact  $(*)$ .

Let  $\varphi_4 = \varphi_3/P_x(\tau, \Delta, \varphi_3)$  $\varphi_4 = \varphi_3/P_x(\tau, \Delta, \varphi_3)$  (see Figure 3) and  $F_3$  be the resulting multifan. Since  $\varphi_3^{-1}(\tau)$ appears before the edge with color  $\tau$  in  $F_2$ ,  $\tau$  is still a 2-inducing color of  $F_3$ . As  $s_1$  and r are  $(1, \Delta)$ -linked by Lemma [3.1](#page-5-0) [\(b\)](#page-5-1), we have  $u \in P_x(1, \tau, \varphi_4)$ . Let  $\varphi_5 = \varphi_4/P_x(1, \tau, \varphi_4)$ . The coloring of  $S(u; s_1, x, y)$  is now as in Figure [3](#page-23-0)  $J_3$ . Since  $2, \delta \notin \{1, \tau, \Delta\}$ , y and  $s_1$  are still  $(2, \delta)$ -linked with respect to  $\varphi_5$  by fact  $(*),$  which further implies that  $\delta$  is a 2-inducing color of  $F_3$  with respect to  $\varphi_5$ . Since  $\varphi_5$  is  $(F_3, \varphi_4)$ -stable,  $\tau$  is still a 2-inducing colors of  $F_3$  with respect to  $\varphi_5$ . We consider two cases to finish the remaining part of the proof.

#### <span id="page-22-0"></span>Subcase 2.1.1 :  $\tau \prec \delta$  in  $F_3$  with respect to  $\varphi_5$ .

Let  $s_i \in N_{\Delta-1}(r)$  such that  $\overline{\varphi}_5(s_i) = \delta$ . Since y and  $s_1$  are still  $(2,\delta)$ -linked with respect to  $\varphi_5$  and  $\delta$  is 2-inducing of  $F_3$ , by Lemma [3.2](#page-5-3) [\(b\)](#page-6-0),  $r \in P_{s_i}(2,\delta,\varphi_5)$ . We reach a contradiction through the following operations: (1) a  $(2, \delta)$ -swap at  $s_i$  (and so also at r); (2) a (1, 2)-swap at x and s<sub>i</sub>; and (3) shift from s<sub>j</sub> to s<sub>i</sub>, where we assume  $\varphi_5(rs_j) = \tau$  for some  $j \in [2, \Delta - 2]$  and  $s_j, s_{j+1}, \ldots, s_i$  is the 2-inducing sequence of  $F_3$  starting at  $s_j$  and ending at  $s_i$ . Denote the new coloring by  $\varphi_6$ . Now,  $\overline{\varphi}_6(r) = \tau$ ,  $\varphi_6(s_1u) = \tau$ ,  $\varphi_6(ux) = \Delta$ , and  $\overline{\varphi}_6(x) = 2$ , and  $K = (r, rs_1, s_1, s_1, u, u, u, x)$  is a Kierstead path with respect to  $rs_1$ and  $\varphi_6$ . Since  $d_G(s_1) = \Delta - 1$ , we get a contradiction to Lemma [3.3](#page-6-2) (a) that  $\{r, s_1, u, x\}$  is



<span id="page-23-0"></span>Figure 3: Coloring of  $S(u; s_1, x, y)$ 

 $\varphi_6$ -elementary.

Subcase 2.1.2:  $\delta \prec \tau$  in  $F_3$  with respect to  $\varphi_5$ .

We only show that by performing Kempe changes, we can find an  $(F_3, \varphi_5)$ -stable coloring such that the color on  $S(u; s_1, x, y)$  with respect to it is as given in Figure [3](#page-23-0)  $J_4$ . Then the proof follows the same ideas as in Case [Subcase 2.1.1](#page-22-0) by exchanging the roles of  $τ$  and  $δ$ . Based on the coloring in Figure [3](#page-23-0)  $J_3$ , do a  $(1, \delta)$ -swap at both x and y and denote the resulting coloring by  $\varphi_6$ .

<span id="page-23-1"></span>Claim 6.3.  $u \in P_u(1, \tau, \varphi_6)$ .

*Proof of Claim [6.3.](#page-23-1)* Assume to the contrary that  $u \notin P_y(1, \tau, \varphi_6)$ . Under this assumption, it must be the case that  $u \in P_r(1,\tau,\varphi_6)$  (otherwise, performing a  $(\delta,\Delta)$ -swap at x and a  $(1,\tau)$ swap at u shows that  $s_1$  and y are  $(1, \Delta)$ -linked, showing a contradiction to Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that  $s_1$  and r are  $(1, \Delta)$ -linked). Since  $u \in P_r(1, \tau, \varphi_6)$ , let  $\varphi_7$  be obtained by doing a  $(1, \tau)$ swap at y and  $(\delta, \Delta)$ -swap at x, and let  $F_3^*$  be the resulting multifan. Then  $P_{s_1}(\tau, \Delta, \varphi_7)$  =  $s_1uy$ , implying that  $\tau$  is a  $\Delta$ -inducing color of  $F_3^*$  by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Note that  $\delta$  is still a 2-inducing color of  $F_3^*$  as the only operation that changes the color sequence of  $F_3$  was the  $(\delta, \Delta)$ -swap we did to get  $\varphi_7$  from  $\varphi_6$ . Thus,  $\overline{\varphi}_7^{-1}(\delta)$  and  $\overline{\varphi}_7^{-1}(\tau)$  are  $(\delta, \tau)$ -linked by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Also, since  $\tau$  is  $\Delta$ -inducing and  $\delta$  is 2-inducing of  $F_3^*$ , we know  $u \notin P_y(\tau, \delta, \varphi_7)$ . Since otherwise, after a  $(\tau, \delta)$ -swap at y, s<sub>1</sub> and y are  $(\delta, \Delta)$ -linked, showing a contradiction to Lemma [3.2](#page-5-3) [\(a\)](#page-6-1).

Let  $\varphi_8 = \varphi_7/P_y(\tau, \delta, \varphi_7)$ . Now  $P_y(\delta, \Delta, \varphi_8) = yux$ . Note also that  $u \in P_{\overline{\varphi_8}^{-1}(\delta)}(\delta, \tau, \varphi_8) =$  $P_{\overline{\varphi}_8^{-1}(\tau)}(\delta, \tau, \varphi_8)$ . For otherwise, after a  $(\tau, \delta)$ -swap at  $u$ ,  $s_1$  and  $y$  are  $(\delta, \Delta)$ -linked, showing a contradiction to the fact that  $\delta$  is still a 2-inducing color of the resulting multifan. Note that  $P_x(\delta, \Delta, \varphi_8) = xuy$ . Let  $\varphi_9 = \varphi_8/P_x(\delta, \Delta, \varphi_8)$ . Now  $\overline{\varphi_9}^{-1}(\delta)$  and  $\overline{\varphi_9}^{-1}(\tau)$  are  $(\delta, \tau)$ -unlinked. However, since  $\varphi_9$  is  $(F_3^*, \varphi_8)$ -stable,  $\tau$  is still a  $\Delta$ -inducing color and  $\delta$  is 2-inducing of  $F_3^*$ with respect to  $\varphi_9$ , we get a contradiction to Lemma [3.2](#page-5-3) [\(a\)](#page-6-1).  $\Box$ 

Thus by Claim [6.3,](#page-23-1)  $u \in P_y(1, \tau, \varphi_6)$ . Do a  $(1, \tau)$ -swap at y (and u), and denote the resulting coloring by  $\varphi_7$ . Note that  $u \in P_x(1, \delta, \varphi_7)$  (as otherwise, after a  $(1, \delta)$ -swap at x, s<sub>1</sub> and x are  $(1, \Delta)$ -linked, showing a contradiction to Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that s<sub>1</sub> and r are  $(1, \Delta)$ -linked). Let  $\varphi_8 = \varphi_7/P_x(1, \delta, \varphi_7)$ . Now with respect to  $\varphi_8$ , we have the coloring in Figure [3](#page-23-0)  $J_4$ . By the definition,  $\varphi_8$  is  $(F_3, \varphi_5)$ -stable so we still have  $\delta \prec \tau$  in  $F_3$  with respect to  $\varphi_8$ . The remaining proof follows the same ideas as in Case [Subcase 2.1.1.](#page-22-0)

Subcase 2.2:  $\overline{\varphi}(x) = \Delta$  and  $\overline{\varphi}(y) = \Delta$ .

<span id="page-24-0"></span>Claim 6.4. We may assume that  $|N_{\Delta-1}(u) \cap N_{\Delta-1}(r)| = \Delta - 4$ .

*Proof of Claim [6.4.](#page-24-0)* We may assume that x and y are  $(1, \Delta)$ -linked. For otherwise, performing  $(\Delta, 1) - (1, 2)$ -swaps at x reduces the problem to [Subcase 2.1.](#page-21-2) Assume to the contrary that  $|N_{\Delta-1}(u) \cap N_{\Delta-1}(r)| \leq \Delta - 5$ . Then there exists  $z \in N_{\Delta-1}(u) \setminus N_{\Delta-1}(r)$  such that  $z \neq x, y$ . Let  $\overline{\varphi}(z) = \lambda$ . If  $\lambda = 2$ , by exchanging the roles of x and z, we reduce the problem to [Subcase 2.1.](#page-21-2) Thus,  $\lambda \neq 2$ . Doing  $(\lambda, 1) - (1, 2)$ -swaps at z and exchanging the roles of x and z reduces the problem to [Subcase 2.1.](#page-21-2) □

<span id="page-24-1"></span>**Claim 6.5.** We may assume that  $F(r, s_1 : s_\alpha : s_{\Delta-2})$  is a typical multifan with two sequences. That is, F contains both 2-inducing sequence and  $\Delta$ -inducing sequence.

*Proof of Claim [6.5.](#page-24-1)* Recall that  $F_{\varphi}(r, s_1 : s_{\alpha} : s_{\Delta-2})$  is a typical multifan. As  $\Delta \geq 7$ ,  $|N_{\Delta-1}(u) \cap N_{\Delta-1}(r)| = \Delta - 4 \geq 3$  by Claim [6.4.](#page-24-0) If F is a typical 2-inducing multifan, then let  $s_i \in N_{\Delta-1}(u) \cap N_{\Delta-1}(r)$  such that  $s_i \neq s_1$  and that  $\overline{\varphi}(s_i)$  is not the last 2inducing color of F. Then we shift from  $s_2$  to  $s_{i-1}$ , uncolor  $rs_i$ , and color  $rs_1$  by 2. Now  $F^* = (r, rs_i, s_i, rs_{i+1}, s_{i+1}, \ldots, rs_{\Delta-2}, s_{\Delta-2}, rs_{i-1}, s_{i-1}, \ldots, rs_1, s_1)$  is a multifan with two sequences. By permuting the name of colors and the label of vertices in  $\{s_1, \ldots, s_{\Delta-2}\}\,$ , we can assume that  $F = F^*$  is a typical multifan with two sequences.  $\Box$ 

Let  $\varphi(s_1u) = \delta, \varphi(ux) = \tau$ , and  $\varphi(uy) = \lambda$ . By exchanging the roles of the two colors 2 and  $\Delta$ , we have two possibilities for  $\varphi(uy)$ : (A)  $\varphi(uy) = \lambda = 1$ ; and (B)  $\varphi(uy) = \lambda$  is 2inducing. (When  $\varphi(uy)$  is  $\Delta$ -inducing, we will first assume that  $\overline{\varphi}(x) = 2$  and  $\overline{\varphi}(y) = 2$  (by performing  $(\Delta, 1) - (1, 2)$ -swaps at both x and y). Then all the argument will be symmetric to the argument for the case (B) above.) We now consider two cases to finish the proof.

#### Subcase 2.2.1:  $\lambda$  is not the last 2-inducing color of F.

We first perform  $(\Delta, \lambda) - (\lambda, 1)$ -swaps at both x and y. Denote by  $\varphi_1$  the resulting coloring and  $F_1$  the corresponding multifan. Since  $\lambda$  is not the last 2-inducing color of F,  $F_1$  still has two sequences with respect to  $\varphi_1$ . The current coloring of  $S(u; s_1, x, y)$  is given in Figure [4](#page-25-0)  $L_1$ . Since  $s_1$  and r are  $(1, \Delta)$ -linked by Lemma [3.1](#page-5-0) [\(b\)](#page-5-1),  $\delta \neq 1$ . We next show  $u \in P_y(1, \delta, \varphi_1)$  that will lead to the coloring in Figure [4](#page-25-0)  $L_2$  after a  $(1, \delta)$ -swap at both x and y.

<span id="page-24-2"></span>Claim 6.6.  $u \in P_y(1, \delta, \varphi_1)$ .

*Proof of Claim [6.6.](#page-24-2)* Assume to the contrary that  $u \notin P_u(1, \delta, \varphi_1)$ . This implies that  $\delta$  is a  $\Delta$ -inducing color of  $F_1$  (since after doing a  $(1, \delta)$ -swap at y, s<sub>1</sub> and y are  $(\delta, \Delta)$ -linked). If



<span id="page-25-0"></span>Figure 4: Coloring of  $S(u; s_1, x, y)$ 

 $\tau$  is a  $\Delta$ -inducing of  $F_1$ , then we let  $\varphi_2$  be obtained by performing  $(1,2) - (2,\tau) - (1,\tau)$ swaps at both  $x$  and  $y$  based on the coloring of  $L_1$  in Figure [4.](#page-25-0) Now, we must have that  $u \in P_x(1, \delta, \varphi_2)$  or  $u \in P_y(1, \delta, \varphi_2)$  since  $\delta$  is either 2-inducing or  $\Delta$ -inducing with respect to  $F_1$ . Let  $\varphi_3$  be obtained from  $\varphi_2$  by performing a  $(1, \delta)$ -swap at both x and y. Then both  $K_1 = (r, rs_1, s_1, s_u, u, ux, x)$  and  $K_2 = (r, rs_1, s_1, s_u, u, uy, y)$  are Kierstead paths with respect to rs<sub>1</sub> and  $\varphi_3$ . Since  $d_G(s_1) = \Delta - 1$ , applying Lemma [3.3](#page-6-2) (b), x and s<sub>1</sub> are  $(\delta, \Delta)$ linked and y and  $s_1$  are  $(2, \delta)$ -linked. However, by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1),  $s_1$  and  $\overline{\varphi}_3^{-1}(\delta)$  are either  $(\delta, 2)$  or  $(\delta, \Delta)$ -linked, showing a contradiction.

Thus we assume that  $\tau$  is a 2-inducing color of  $F_1$ . Based on the coloring of  $S(u; s_1, x, y)$ as given in Figure [4](#page-25-0)  $L_1$ , we perform  $(1, \tau) - (\tau, \delta)$ -swaps at both x and y and let  $\varphi_2$  be the resulting coloring. Note that either  $\varphi_2(s_1u) = \delta$  or  $\varphi_2(s_1u) = \tau$ . If  $\varphi_2(s_1u) = \delta$ , then after doing a  $(1, \delta)$ -swap at both x and y, s<sub>1</sub> and y are  $(1, \Delta)$ -linked, which gives a contradiction to Lemma [3.1](#page-5-0) [\(b\)](#page-5-1) that  $s_1$  and r are  $(1, \Delta)$ -linked. Thus  $\varphi_2(s_1u) = \tau$ . We first do a  $(1, \delta)$ swap at both x and y. Then since  $\tau$  is a 2-inducing color of  $F_1$ ,  $u \in P_y(1, \tau, \varphi_2)$  (since otherwise, after doing a  $(1, \tau)$ -swap at y, s<sub>1</sub> and y are  $(\tau, \Delta)$ -linked, showing a contradiction to Lemma [3.2](#page-5-3) [\(a\)](#page-6-1)). Thus we do a  $(1, \tau)$ -swap at both x and y and let  $\varphi_3$  be the new coloring. Note that  $\delta$  is still  $\Delta$ -inducing and  $\tau$  is 2-inducing with respect to  $F_1$  and  $\varphi_3$ . Thus  $\overline{\varphi_3}^{-1}(\delta)$ and  $\overline{\varphi}_3^{-1}(\tau)$  are  $(\delta, \tau)$ -linked by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Let  $\varphi_4$  be obtained from  $\varphi_3$  by doing a  $(\delta, \tau)$ -swap at y, and let  $F_1^*$  be the resulting multifan. Then  $K = (r, rs_1, s_1, s_1u, u, uy, y)$  is a Kierstead path with respect to rs<sub>1</sub> and  $\varphi_4$ . Since  $d_G(s_1) = \Delta - 1$ , applying Lemma [3.3](#page-6-2) (b), y and s<sub>1</sub> are  $(2, \delta)$ -linked. Since  $\delta$  is still  $\Delta$ -inducing and  $\tau$  is 2-inducing with respect to  $F_1^*$  and  $\varphi_4$ , we achieve a contradiction to the fact that  $s_1$  and  $\overline{\varphi_4}^{-1}(\delta)$  are  $(2,\delta)$ -linked by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1). Therefore it must be the case  $u \in P_y(1, \delta, \varphi_1)$ .  $\Box$ 

Since  $u \in P_u(1, \delta, \varphi_1)$ , we perform a  $(1, \delta)$ -swap at both x and y gives  $L_2$  in Figure [4.](#page-25-0) Call the resulting coloring  $\varphi_2$ . Now  $K = (r, rs_1, s_1, s_u, u, uy, y)$  is a Kierstead path with respect to rs<sub>1</sub> and  $\varphi_2$ . Since  $d_G(s_1) = \Delta - 1$ , by Lemma [3.3](#page-6-2) (b), y and s<sub>1</sub> are  $(2, \delta)$ -linked. It deduces that  $\delta$  must be a 2-inducing color of  $F_1$  with respect to  $\varphi_2$ . Recall that  $F_1$  still has two sequences with respect to  $\varphi_2$ . Let  $\gamma$  be a  $\Delta$ -inducing color of  $F_1$ . Since  $\overline{\varphi_2}^{-1}(\delta)$ 

and  $\overline{\varphi}_2^{-1}(\gamma)$  are  $(\delta, \gamma)$ -linked by Lemma [3.2](#page-5-3) [\(a\)](#page-6-1), we do a  $(\delta, \gamma)$ -swap at y to get  $\varphi_3$ . Still,  $\delta$ is a 2-inducing color and  $\gamma$  is a  $\Delta$ -inducing color of the resulting multifan. By Lemma [3.3](#page-6-2) (b),  $s_1$  and y are  $(2, \gamma)$ -linked, showing a contradiction to the fact that  $s_1$  and  $\overline{\varphi}_3^{-1}(\gamma)$  are  $(2, \gamma)$ -linked.

#### Subcase 2.2.2:  $\lambda$  is the last 2-inducing color of F.

If  $\tau$  is 2-inducing, then  $\tau \prec \lambda$ . This gives back to the previous case by exchanging the roles of  $\tau$  and  $\lambda$ . If  $\tau$  is  $\Delta$ -inducing and  $\tau$  is not the last  $\Delta$ -inducing color, then by doing  $(\Delta, 1)$ −(1, 2)-swaps at x and y, a similar proof follows as in the previous case by exchanging the roles of 2 and  $\Delta$ . Thus  $\tau$  is the last  $\Delta$ -inducing color of F.

Let  $C_u$  be the cycle in  $G_{\Delta}$  containing u. By Theorem [4.1](#page-11-1) [\(i\)](#page-11-2), for every vertex on  $C_u$ , its  $(\Delta - 1)$ -neighborhood is  $N_{\Delta - 1}(u)$ . As  $|V(C_u)| \geq 3$ , there exist  $u^*, u' \in V(C_u) \setminus \{u\}$ such that one of  $\varphi(u^*y)$  and  $\varphi(u'y)$  is neither  $\tau$  nor  $\lambda$ . Assume that  $\varphi(u^*y) \notin {\tau, \lambda}$ . Letting  $u^*$  play the role of  $u$ , we reduce the problem to the previous case, finishing proof of Theorem [2.5.](#page-2-1) □

# <span id="page-26-4"></span>References

- <span id="page-26-1"></span>[1] Y. Cao, G. Chen, G. Jing, and S. Shan. The Core Conjecture of Hilton and Zhao I: Pseudo-multifan and Lollipop. *[arXiv:2108.03549](http://arxiv.org/abs/2108.03549)*, 2021.
- <span id="page-26-3"></span>[2] D. Cariolaro and G. Cariolaro. Colouring the petals of a graph. Electron. J. Combin., 10:# R6, 2003.
- <span id="page-26-5"></span>[3] D. W. Cranston and L. Rabern. The Hilton–Zhao Conjecture is True for Graphs with Maximum Degree 4. SIAM J. Discrete Math., 33(3):1228–1241, 2019.
- <span id="page-26-0"></span>[4] A. J. W. Hilton and C. Zhao. The chromatic index of a graph whose core has maximum degree two. Discrete Math., 101(1-3): 135–147, 1992.
- <span id="page-26-2"></span>[5] A. J. W. Hilton and C. Zhao. On the edge-colouring of graphs whose core has maximum degree two. J. Combin. Math. Combin. Comput., 21:97–108, 1996.
- [6] M. Stiebitz, D. Scheide, B. Toft, and L. M. Favrholdt. Graph Edge Coloring: Vizing's Theorem and Goldberg's Conjecture. Wiley Series in Discrete Mathematics and Optimization. John Wiley & Sons, Inc., Hoboken, NJ, 2012.