

A Note on Squares in Binary Words

Tero Harju

Department of Mathematics and Statistics

University of Turku, Finland

harju@utu.fi

August 11, 2021

Abstract

We consider words over a binary alphabet. A word w is overlap-free if it does not have factors (blocks of consecutive letters) of the form $uvuvu$ for nonempty u . Let $M(w)$ denote the number of positions that are middle positions of squares in w . We show that for overlap-free binary words, $2M(w) \leq |w| + 3$, and that there are infinitely many overlap-free binary words for which $2M(w) = |w| + 3$.

Keywords. Overlap-free words, squares, centre of a square, binary words.

1 Introduction

Overlap-free words were first studied by Axel Thue in two papers [5, 6] in 1906 and 1912. In particular he proved that there are infinite overlap-free binary words.

In the following we consider binary words, and choose the alphabet $\Sigma = \{0, 1\}$ for these words. An *overlap* is a word of the form $uvuvu$ where u is nonempty. A word w is said to be *overlap-free* if it has no overlaps as factors, i.e., consecutive blocks of letters. Equally well, we can presume in the definition that x is a letter. The most well known overlap-free words are the factors of the infinite *Thue-Morse word* \mathbf{t} that is obtained by iterating the morphism $\mu: \Sigma^* \rightarrow \Sigma^*$ with

$$\mu(0) = 01 \text{ and } \mu(1) = 10$$

on the initial word 0. Therefore

$$\mathbf{t} = 011010011001011010010110 \dots$$

The morphism μ preserves overlap-freeness, i.e., if w is overlap-free, then so is $\mu(w)$; see e.g. Chapter 2 of Lothaire [3] or Berstel and Séébold [1].

Let $|w|$ denote the length of w . An integer p with $1 \leq p < |w|$ is a *position* in w . It denotes the place after the prefix of length p . A nonempty word u is a *square at position* $p = |x|$ in $w = xy$ if there are, possibly empty, words x' and y' such that $x = x'u$ and $y = uy'$. In this case we say that p is a *centre of a square*.

Let ε denote the empty word. If $w = xuy$ then u is a *factor* of w . It is a *prefix* if $x = \varepsilon$, and a *suffix* if $y = \varepsilon$. The word w is said to be *bordered* if there exists a nonempty word v , with $v \neq w$, that is both a prefix and a suffix of w . For a word $w = uv$, we denote by $u = wv^{-1}$ the prefix of w when the suffix v is deleted.

2 On the maximum number of centres

We consider the number of centres of squares in binary words. We follow Harju and Kärki [2], and define

$$M(w) = \#\{p \mid p \text{ is a centre of a square in } w\}.$$

In [2] the authors count ‘frames’, i.e., unbordered squares. The number of these equals $M(w)$ since if uu is a square at position p then for the border v of u of minimal length also vv is a square at p .

In the general binary case the paper [2, Theorem 4] gives the minimum number of centres of squares:

Theorem 1. *For binary words of length $n \geq 3$,*

$$\min_{|w|=n} M(w) = \left\lceil \frac{n}{2} \right\rceil - 2.$$

The problem for $\max_{|w|=n} M(w)$ is trivial. Indeed, the unary word 0^n has a square at every position, i.e., $M(0^n) = n - 1$. On the other hand, in the ternary case, i.e., words over a three letter alphabet, the minimum is zero since there are (infinite) square-free words in this case. Here we shall consider the maximum problem for the binary overlap-free words.

The next technical lemma will be used for short prefixes x in the proof of Theorem 5.

Lemma 2. *Let s be the longest common suffix of the binary words x and w . Assume that xw and w are overlap-free. Then $xw(ws^{-1})$ is overlap-free.*

Proof. Assume that there is a square $xuxu$ as a prefix of xw such that $|xuxu| \geq |w|$. Then u is a prefix of w , i.e., $w = uy = vt$ for y and t with $xu = yv$; see Fig. 1.

The word u is a suffix of $yv (= xu)$ and a prefix of $w = vt (= uy)$. The occurrences of u cannot overlap in the overlap-free w . Therefore either (1) $u = v$ or (2) $v = uzu$ for some z .

(1) Suppose first that $u = v$ and thus that $y = x$ and $t = x$. Then x is a suffix of the common suffix s and $xw(wx^{-1}) = xuxu$. Therefore $xuxu$ does not extend to an overlapping factor $xuxua$ of $xw(ws^{-1})$.

(2) Assume then that $v = uzu$ and hence that $y = zut$ and $w = uzut$. Then $xu = yv = zutuzu$, and finally, $xw = xuy = zutuzuzut$ which contains an overlap $uzuzu$ (since $u \neq \varepsilon$); a contradiction with the overlap-freeness of xw . \square




Figure 1: Overlapping factor: $xu = yv$.

In particular, if in the above $s = \varepsilon$ and xw and wv are overlap-free then also xwv is overlap-free.

The next lemma was proved in [2, Lemma 10]. We note that in the Thue-Morse word \mathbf{t} there are no consecutive centres of squares. This can be seen by looking at the short factors of \mathbf{t} . Indeed, the unbordered squares in \mathbf{t} are of the form uu for $u \in \{0, 1, 01, 10\}$; see Pansiot [4].

Lemma 3. *Let w be a factor of even length $n \geq 4$ of the Thue-Morse word starting at an even position. Then all squares are at even positions, and hence $M(w) = n/2 - 1$.*

We show in the next theorem that if an overlap-free word satisfies the maximality condition $2M(w) = |w| + 3$ then w has the form $w = auu$ (or symmetrically uua). In Theorem 5 we then construct overlap-free words for which $M(w)$ reaches the upper bound.

We call the words 001001 and 110110 *prefix special* and their reversed words 100100 and 011011 *suffix special*. These special words can occur only as a prefix (resp., suffix) of an overlap-free word since they cannot be extended to the left (resp., right) without violating overlap-freeness.

Theorem 4. *For overlap-free binary words w , we have $M(w) \leq \left\lceil \frac{|w|}{2} \right\rceil + 1$, i.e.,*

$$2M(w) \leq |w| + 3. \quad (1)$$

Proof. A word of length n has $n - 1$ positions of which $\lfloor n/2 \rfloor$ are odd positions and $\lfloor n/2 \rfloor - 1$ or $\lfloor n/2 \rfloor$ even positions depending if n is even or odd. For the bound (1) we look for the positions where the consecutive positions p and $p + 1$ are centres of squares.

Let $|w| = n$. If w has a special prefix 001001 or 110110 then the positions 3 and 4 are centres of squares. Similarly if w has a suffix 100100 or 011011 then the positions $n - 4$ and $n - 3$ are centres of squares.

Suppose then that $5 \leq p \leq n - 5$, and assume that there are minimal length, and thus unbordered, squares u and v at positions p and $p + 1$. By appealing to symmetry, we can assume that $|u| \geq |v|$. To avoid the overlap ccc (with $c \in \{0, 1\}$) that happens when $|u| = 1 = |v|$, necessarily $u = au'b$ for different letters a, b . Also, $|u| \geq 3$ since if $u = ab$ then $uu = abab$ yields that $v = ba$ and there is an overlap $ababa$ in w .

(A) Assume first that uu is not a prefix of w .

Claim 1. The square uu is a suffix of w .

Indeed, $\beta = auu$ is a factor of w since buu would be an overlapping factor. Hence ab is a prefix of u .

If $u = abb$ then $\beta = aabbabb$ ends in a special word. In this case there does not exist a square at $p + 1$. (There is one at $p - 1$.) Therefore $|u| \geq 4$.

In order for u to avoid ab as a border, bb must be a suffix of u . Hence $\beta = a(abxabb)(abxabb)$ for some x . It follows that β is a suffix of w since βa and βb have overlapping factors, $(abxabb)(abxabb)a$ and bbb , respectively. This proves Claim 1.

Claim 2. $w = \beta$.

We have $x \neq \varepsilon$ since if $x = \varepsilon$, then $\beta = aababbababb$ has an overlapping factor $babab$. Then x has a prefix a in order for u to avoid abb as a border. So far we have $\beta = a(abayabb)(abayabb)$. In particular, $vv = baba$ is the minimal square at $p + 1$. Now, either $y = \varepsilon$ or, in order to avoid the factor $babab$, ab is a prefix of y , i.e., $\beta = a(abaabzabb)(abaabzabb)$ for a word z . In both cases, the prefix is $aabaab$ that can occur only at the beginning of w . Hence $\beta = w$, and therefore

$$w = auu = a(abaabzabb)(abaabzabb).$$

Let $|u| = 2k$ or $|u| = 2k + 1$.

The positions 1 and 2 of u are not centres of squares in u , and hence, by itself, u can have at most $k - 1$ odd or even centres of squares. However, the position 3 of w is a centre of $aabaab$ and the position $p + 1$ of w is centre of vv . Together with the centres of uu and aa , we have $M(w) \leq 2k + 2$, and so $2M(w) \leq |w| + 3$ where $|w| = 4k + 1$.

(B) The case where uu is not a suffix of w is symmetric to the case (A). In this case, w has the form uuu .

(C) If $w = uu$ then w cannot have special words in the prefix or in the suffix. Hence $M(w) \leq |w|/2 + 1$, i.e., $2M(w) \leq |w| + 2$.

Finally, suppose that w does not have centres of squares p for $5 \leq p \leq |w| - 5$. Then the consecutive centres of squares are in the special words. E.g., by considering a possible factorisation $w = (00100)11x00(11011)$ with special words at the ends, we obtain $M(w) \leq (|w| - 10)/2 + 6 = |w|/2 + 1$, and so $2M(w) \leq |w| + 2$. \square

By the proof of Theorem 4, the bound $\left\lceil \frac{|w|}{2} \right\rceil + 1$ can be obtained only by overlap-free words of odd length.

Theorem 5. *There are infinitely many overlap-free binary words w such that*

$$M(w) = \left\lceil \frac{|w|}{2} \right\rceil + 1.$$

Proof. The the upper bound for $M(w)$ is given in Lemma 4. For equality we rely on the factors of the Thue-Morse word \mathbf{t} . Let α_n denote the factor of \mathbf{t} of length $3 \cdot 2^n$ that starts at position 5, e.g.,

$$\alpha_1 = 100110, \text{ and } \alpha_2 = 100110010110.$$

Then $|\mu(\alpha_n)| = 2|\alpha_n| = |\alpha_{n+1}|$, and $\mu(\alpha_n) = 1001 \cdot 0110 \cdots 0110 \cdot 1001$, where $\mu(\alpha_n)$ starts after $\mu(0110)$ at position 9 of \mathbf{t} . Hence $\alpha_{n+1} = 1001 \cdot 1001 \cdots 0110$ is a conjugate of $\mu(\alpha_n)$. In particular, as a factor of \mathbf{t} , $1001\mu(\alpha_n)$ is overlap-free for all n .

We show inductively that $\alpha_n\alpha_n$ is overlap-free for all n . To begin with $\alpha_1\alpha_1 = 100110100110$ is overlap-free. For $n \geq 1$, we have

$$\begin{aligned} \alpha_{n+1}\alpha_{n+1} &= 1001 \mu(\alpha_n)(1001)^{-1}(1001) \mu(\alpha_n)(1001)^{-1} \\ &= 1001 \mu(\alpha_n)\mu(\alpha_n)(1001)^{-1} \\ &= 1001 \mu(\alpha_n\alpha_n)(1001)^{-1}, \end{aligned}$$

where $1001 \mu(\alpha_n)$ is overlap-free and, by the induction hypothesis, so is $\mu(\alpha_n)\mu(\alpha_n)$ since μ preserves overlap-freeness. Lemma 2 applied to $1001 \mu(\alpha_n\alpha_n)$ gives that $1001 \mu(\alpha_n\alpha_n)(1001)^{-1}$ is overlap-free. Therefore $\alpha_{n+1}\alpha_{n+1}$ is overlap-free.

Denote by $\bar{\alpha}$ the prefix of α of length $|\alpha| - 1$, i.e., truncate the last letter of α . Let

$$w_n = 00\alpha_n\bar{\alpha}_n = 0(0\bar{\alpha}_n)(0\bar{\alpha}_n)$$

By Lemma 2, $0\alpha_n\bar{\alpha}_n$ is overlap-free. Also, $00\alpha_n\bar{\alpha}_n$ has a special prefix 001001 , and hence w_n is overlap-free.

By Lemma 3, $M(\alpha_n\alpha_n) = (2|\alpha_n|/2 - 1) + 1 = |\alpha_n| - 1$ where the $+1$ comes from the middle position. The centres of squares have even positions in

$\alpha_n \alpha_n$. Therefore also $M(\alpha_n \bar{\alpha}_n) = |\alpha_n| - 1$. The squares 001001 and 00 are at the consecutive positions 3 and 4 of w_n , and in the middle portion we have $(0\bar{\alpha}_n)(0\bar{\alpha}_n)$ next to the squares 00 and 1010 (of even positions). Since $|w_n| = 2|\alpha_n| + 1$, we have

$$M(w_n) = |\alpha_n| - 1 + 3 \text{ and so } 2M(w_n) = |w_n| + 3$$

as required. □

References

- [1] J. Berstel and P. Séébold. A characterization of overlap-free morphisms. *Discrete Appl. Math.*, 46(3):275–281, 1993.
- [2] T. Harju and T. Kärki. On the number of frames in binary words. *Theor. Comput. Sci.*, 412(39):5276–5284, 2011.
- [3] M. Lothaire. *Combinatorics on Words*, volume 17 of *Encyclopedia of Mathematics*. Addison-Wesley, Reading, Massachusetts, 1983.
- [4] J. J. Pansiot. The Morse sequence and iterated morphisms. *Inf. Process. Lett.*, 12:68–70, 1981.
- [5] A. Thue. Über unendliche Zeichenreihen. *Norske Vid. Selsk. Skr., I Mat.-nat. Kl. Christiania*, 7:1–22, 1906.
- [6] A. Thue. Über die gegenseitige Lage gleicher Teile gewisser Zeichenreihen. *Norske Vid. Selsk. Skr., I Mat.-nat. Kl. Christiania*, 1:1–67, 1912.