

# Hessian estimates for the conjugate heat equation coupled with the Ricci flow

Hong Huang

## Abstract

In this short note we obtain some local and global upper bounds for the Hessian of a positive solution to the conjugate heat equation coupled with the Ricci flow.

**Key words:** Hessian estimates; conjugate heat equation; Ricci flow

**AMS2020 Classification:** 53E20, 58J35

## 1 Introduction

In the literature there are extensive researches on derivative estimates for solutions to the heat equation or conjugate heat equation coupled with the Ricci flow; see for example [1], [3], [4], [5], [6], [7], [8], [9], [10], [11], [13], [14], [15], and [17].

In this short note we obtain some local and global upper bounds for the Hessian of a positive solution to the conjugate heat equation coupled with the Ricci flow. I was inspired by Han-Zhang [9]. To state our results we first introduce some notations. Fix  $T > 0$ . Let  $(M, g(t))$ ,  $t \in [0, T]$ , be a (not necessarily complete) solution to Hamilton's Ricci flow

$$\frac{\partial g(t)}{\partial t} = -2Ric(g(t))$$

on a manifold  $M$  (without boundary) of dimension  $n$ . For any points  $x, y \in M$ , let  $d(x, y, t)$  be the distance between  $x$  and  $y$  w.r.t.  $g(t)$ . For  $(x_0, t_0) \in M \times (0, T]$ ,  $r > 0$ , and  $T' > 0$ , let

$$Q_{r, T'}(x_0, t_0) := \{(x, t) \in M \times [0, T] \mid d(x, x_0, t) \leq r, t_0 - T' \leq t \leq t_0\}$$

be a parabolic cube.

Consider a smooth positive solution  $u$  to the conjugate heat equation  $(\frac{\partial}{\partial t} + \Delta_{g(t)} - R_{g(t)})u = 0$  coupled with the Ricci flow on  $M \times [0, T]$ , where  $R_{g(t)}$  is the scalar curvature of the metric  $g(t)$ . We have the following global upper bound for  $\text{Hess } u$ .

**Theorem 1.1.** *Let  $(M^n, g(t))$ ,  $t \in [0, T]$ , be a Ricci flow on a compact manifold,  $u$  be a positive solution to the conjugate heat equation coupled with the Ricci flow on  $M \times [0, T]$  with  $0 < u \leq A$ . Then we have*

$$\nabla^2 u \leq u \left( \frac{18}{T-t} + C \right) \left( 1 + \log \frac{A}{u} \right) g(t) \quad \text{on } M \times [0, T],$$

and, in particular,

$$\Delta u \leq u \left( \frac{18n}{T-t} + C \right) \left( 1 + \log \frac{A}{u} \right) \quad \text{on } M \times [0, T],$$

where  $C$  depends on  $n$ , the upper bounds of  $|Rm|$ ,  $|\nabla Ric|$ , and  $|\nabla^2 R|$  on  $M \times [0, T]$ .

We also get a local upper bound for  $\text{Hess } u$ .

**Theorem 1.2.** *Let  $(M^n, g(t))$ ,  $t \in [0, T]$ , be a (not necessarily complete) Ricci flow,  $u$  be a positive solution to the conjugate heat equation coupled with the Ricci flow on  $M \times [0, T]$  with  $0 < u \leq A$ . Let  $x_0 \in M$ . Assume that the parabolic cube  $Q_{4r, T}(x_0, T)$  is compact. Then we have*

$$\nabla^2 u \leq u \left( \frac{C_0}{T} + \frac{C_0}{r^2} + C_2 \right) \left( 1 + \log \frac{A}{u} \right)^2 g(t) \quad \text{on } Q_{r, \frac{T}{2}}(x_0, \frac{T}{2}),$$

where  $C_0$  is a universal constant, and  $C_2$  depends on  $n$ , the upper bounds of  $|Rm|$ ,  $|\nabla Ric|$ , and  $|\nabla^2 R|$  on  $Q_{4r, T}(x_0, T)$ .

In Section 2 we derive local gradient estimates for a positive solution to the conjugate heat equation coupled with the Ricci flow. In Section 3 we prove Theorems 1.1 and 1.2.

## 2 Gradient estimates

Let  $(M^n, g(t))$ ,  $t \in [0, T]$ , be a (not necessarily complete) Ricci flow, and  $u$  be a positive solution to the conjugate heat equation

$$\left( \frac{\partial}{\partial t} + \Delta_{g(t)} - R_{g(t)} \right) u = 0$$

coupled with the Ricci flow on  $M \times [0, T]$ . Sometimes we'll write  $\partial_t$  for  $\frac{\partial}{\partial t}$ , and  $u_t$  for  $\frac{\partial u}{\partial t}$ . Let  $f = \log u$ . Then

$$f_t = -\Delta f - |\nabla f|^2 + R.$$

Fix  $\alpha > 1$ . Let  $\tau = T - t$ , and

$$F = \tau \left( \frac{|\nabla u|^2}{u^2} + \alpha \frac{u_t}{u} - \alpha R \right) = \tau (|\nabla f|^2 + \alpha f_t - \alpha R).$$

Let  $x_0 \in M$ . Assume that the parabolic cube  $Q_{2r, T}(x_0, T)$  is compact, and  $-K_0 \leq Ric \leq K_0$ ,  $|\nabla R| \leq K_1$ , and  $\Delta R \leq K_2$  on  $Q_{2r, T}(x_0, T)$ .

**Lemma 2.1.** *With the above assumption, for any  $\varepsilon > 0$  we have*

$$\begin{aligned} (\Delta + \partial_t)F &\geq -2\langle \nabla f, \nabla F \rangle + \frac{2\tau}{n + \varepsilon}(f_t + |\nabla f|^2 - R)^2 - (|\nabla f|^2 + \alpha f_t - \alpha R) \\ &\quad - 2\tau[2 - \alpha|K_0|\nabla f|^2 - 2\tau(\alpha - 1)K_1|\nabla f| - \frac{n(n + \varepsilon)}{2\varepsilon}\alpha^2\tau K_0^2 - \alpha\tau K_2] \end{aligned}$$

on  $Q_{2r,T}(x_0, T)$ .

**Proof.** We have

$$\begin{aligned} \Delta|\nabla f|^2 &= 2|\nabla^2 f|^2 + 2\langle \nabla f, \nabla \Delta f \rangle + 2\text{Ric}(\nabla f, \nabla f), \\ \Delta f_t &= (\Delta f)_t - 2\langle \text{Ric}, \nabla^2 f \rangle, \\ \tau \Delta f &= \tau(\alpha - 1)(f_t - R) - F, \\ (|\nabla f|^2)_t &= 2\langle \nabla f, \nabla f_t \rangle + 2\text{Ric}(\nabla f, \nabla f), \end{aligned}$$

and

$$\begin{aligned} &\Delta F \\ &= \tau[2|\nabla^2 f|^2 + 2\langle \nabla f, \nabla \Delta f \rangle + 2\text{Ric}(\nabla f, \nabla f) + \alpha((\Delta f)_t - 2\langle \text{Ric}, \nabla^2 f \rangle) - \alpha\Delta R] \\ &= 2\tau|\nabla^2 f|^2 - 2\langle \nabla f, \nabla F \rangle + 2\tau(\alpha - 1)\langle \nabla f, \nabla f_t \rangle - 2\tau(\alpha - 1)\langle \nabla R, \nabla f \rangle \\ &\quad + 2\tau\text{Ric}(\nabla f, \nabla f) + \alpha\tau(-f_{tt} - (|\nabla f|^2)_t + R_t - 2\langle \text{Ric}, \nabla^2 f \rangle) - \alpha\tau\Delta R \\ &= 2\tau|\nabla^2 f|^2 - 2\langle \nabla f, \nabla F \rangle - F_t - \frac{F}{\tau} - 2\alpha\tau\langle \text{Ric}, \nabla^2 f \rangle + 2\tau(2 - \alpha)\text{Ric}(\nabla f, \nabla f) \\ &\quad - 2\tau(\alpha - 1)\langle \nabla R, \nabla f \rangle - \alpha\tau\Delta R. \end{aligned}$$

See p. 61 of [5].

Note that

$$|\langle \text{Ric}, \nabla^2 f \rangle| \leq |\text{Ric}||\nabla^2 f| \leq \frac{\alpha(n + \varepsilon)}{4\varepsilon}|\text{Ric}|^2 + \frac{\varepsilon}{\alpha(n + \varepsilon)}|\nabla^2 f|^2$$

for any  $\varepsilon > 0$ , and

$$|\nabla^2 f|^2 \geq \frac{1}{n}(\Delta f)^2 = \frac{1}{n}(f_t + |\nabla f|^2 - R)^2,$$

then the lemma follows.  $\square$

**Proposition 2.2.** *Let  $(M^n, g(t))$ ,  $t \in [0, T]$ , be a (not necessarily complete) Ricci flow and  $u$  be a positive solution to the conjugate heat equation coupled with the Ricci flow on  $M \times [0, T]$ . Let  $x_0 \in M$ . Suppose that  $Q_{2r,T}(x_0, T)$  is compact and  $-K_0 \leq \text{Ric} \leq K_0$ ,  $|\nabla R| \leq K_1$ ,  $\Delta R \leq K_2$  on  $Q_{2r,T}(x_0, T)$ . For any  $\alpha > 1$  and  $\varepsilon > 0$ , we have*

$$\frac{|\nabla u|^2}{u^2} + \alpha \frac{u_t}{u} - \alpha R \leq \frac{(n + \varepsilon)\alpha^2}{2(T - t)} + C(r^{-2} + r^{-1} + 1) \quad \text{on } Q_{r,T}(x_0, T) \setminus \{(x, T) \mid x \in M\},$$

where the constant  $C$  depends on  $n, \alpha, \varepsilon, K_0, K_1$  and  $K_2$ .

**Proof.** As in [12], choose a smooth cutoff function  $\psi : [0, \infty) \rightarrow [0, 1]$  with  $\psi = 1$  on the interval  $[0, 1]$ ,  $\psi = 0$  on  $[2, \infty)$ , and

$$\psi' \leq 0, \quad |\psi'|^2 \leq C_0\psi, \quad \psi'' \geq -C_0,$$

where  $C_0$  is a universal constant. Let  $\phi(x, t) = \psi(\frac{d(x, x_0, t)}{r})$ . Suppose that the maximum of the function  $\phi F$  is positive, otherwise the result follows trivially. Assume that  $\phi F$  achieves its positive maximum at the point  $(x_1, t_1)$ . Then  $(x_1, t_1) \in Q_{2r, T}(x_0, T)$  with  $t_1 \neq T$ . By Calabi's trick [2] we may assume that  $\phi F$  is smooth at  $(x_1, t_1)$ . Let  $\tau_1 = T - t_1$ . We compute at the point  $(x_1, t_1)$  using Lemma 2.1,

$$\begin{aligned} 0 &\geq (\Delta + \partial_t)(\phi F) \\ &\geq \tau_1 \phi \frac{2}{n + \varepsilon} (f_t + |\nabla f|^2 - R)^2 - CF \sqrt{\phi} r^{-1} |\nabla f| - \phi \frac{F}{\tau_1} \\ &\quad - C\tau_1 \phi |\nabla f|^2 - C\tau_1 \phi - CF(r^{-2} + r^{-1} + 1), \end{aligned}$$

where the constant  $C$  depends on  $n, \alpha, \varepsilon, K_0, K_1$  and  $K_2$ ; compare [5], [13], and [16]. Then we proceed as in [12] and [5].  $\square$

The following corollary is a slight improvement of Lemma 4.1 in [5] in the Ricci flow case.

**Corollary 2.3.** *Suppose that  $(M^n, g(t))$ ,  $t \in [0, T]$ , is a complete Ricci flow with  $-K_0 \leq Ric \leq K_0$ ,  $|\nabla R| \leq K_1$ ,  $\Delta R \leq K_2$  on  $M \times [0, T]$ , and that  $u$  is a positive solution to the conjugate heat equation coupled with the Ricci flow on  $M \times [0, T]$ . For any  $\alpha > 1$  and  $\varepsilon > 0$ , at  $(x, t) \in M \times [0, T)$  we have*

$$\frac{|\nabla u|^2}{u^2} + \alpha \frac{u_t}{u} - \alpha R \leq \frac{(n + \varepsilon)\alpha^2}{2(T - t)} + C,$$

where the constant  $C$  depends on  $n, \alpha, \varepsilon, K_0, K_1$  and  $K_2$ .

**Proposition 2.4.** *Let  $(M^n, g(t))$ ,  $t \in [0, T]$ , be a (not necessarily complete) Ricci flow and  $u$  be a positive solution to the conjugate heat equation coupled with the Ricci flow on  $M \times [0, T]$ . Let  $x_0 \in M$ . Suppose that  $Q_{2r, T}(x_0, T)$  is compact and  $-K_0 \leq Ric \leq K_0$ ,  $|\nabla R| \leq K_1$ ,  $\Delta R \leq K_2$  on  $Q_{2r, T}(x_0, T)$ . For any  $\alpha > 1$  and  $\varepsilon > 0$ , at  $(x, t) \in Q_{r, T}(x_0, T)$  with  $t \neq T$  we have*

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} + \alpha \frac{u_t}{u} - \alpha R &\leq \frac{n\alpha^2}{T - t} + \frac{C\alpha^2}{r^2} (r\sqrt{K_0} + \frac{\alpha^2}{\alpha - 1}) + C\alpha^2 K_0 \\ &\quad + \frac{n\alpha^2}{\alpha - 1} (|2 - \alpha|K_0 + \frac{\alpha - 1}{2}K_1) + n\alpha^2 K_0 + \alpha\sqrt{n(\alpha - 1)K_1} + \alpha\sqrt{n\alpha K_2}, \end{aligned}$$

where the constant  $C$  depends only on  $n$ .

**Proof.** In Lemma 2.1 we let  $\varepsilon = n$  and get

$$\begin{aligned} (\Delta + \partial_t)F &\geq -2\langle \nabla f, \nabla F \rangle + \frac{\tau}{n} (f_t + |\nabla f|^2 - R)^2 - (|\nabla f|^2 + \alpha f_t - \alpha R) \\ &\quad - \tau(2|2 - \alpha|K_0 + (\alpha - 1)K_1)|\nabla f|^2 - n\alpha^2\tau K_0^2 - (\alpha - 1)\tau K_1 - \alpha\tau K_2. \end{aligned}$$

Then we proceed as in [13] and [16].  $\square$

### 3 Hessian estimates

Let  $(M^n, g(t))$ ,  $t \in [0, T]$ , be a Ricci flow and  $u$  be a smooth positive solution to the conjugate heat equation coupled with the Ricci flow on  $M \times [0, T]$ .

**Lemma 3.1.** *Let  $u$  be a positive solution to the conjugate heat equation coupled with the Ricci flow with  $0 < u \leq A$ , and  $f = \log \frac{u}{A}$ . Let  $u_{ij}$  denote the Hessian of  $u$  in local coordinates, and  $v_{ij} := \frac{u_{ij}}{u(1-f)}$ . In local coordinates we have*

$$\begin{aligned} (\partial_t + \Delta - \frac{2f}{1-f} \nabla f \cdot \nabla) v_{ij} &= \frac{|\nabla f|^2 + Rf}{1-f} v_{ij} + \frac{1}{u(1-f)} [2R_{kijl} u_{kl} + R_{il} u_{jl} + R_{jl} u_{il} \\ &\quad + 2(\nabla_i R_{jl} + \nabla_j R_{il} - \nabla_l R_{ij}) \nabla_l u - u \nabla_i \nabla_j R - \nabla_i R \nabla_j u - \nabla_j R \nabla_i u - Ru_{ij}]. \end{aligned}$$

Here we adopt the usual convention on summations. For example,  $R_{kijl} u_{kl}$  means  $g^{ab} g^{pq} R_{aijp} u_{bq}$ . Note also that our convention on the curvature tensor  $R_{ijkl}$  is the same as that of R. Hamilton, but is different from that of Han-Zhang [9].

**Proof.** The proof is similar to that of Lemma 3.1 in [9].  $\square$

**Lemma 3.2.** *Let  $u$  be a positive solution to the conjugate heat equation coupled with the Ricci flow with  $0 < u \leq A$ , and  $f = \log \frac{u}{A}$ . For a function  $h$  we let  $h_i$  denote the 1-form  $dh$  in local coordinates. We also let  $w_{ij} := \frac{u_i u_j}{u^2(1-f)^2}$  denote the 2-tensor  $\frac{du \otimes du}{u^2(1-f)^2}$  in local coordinates. In local coordinates we have*

$$\begin{aligned} (\partial_t + \Delta - \frac{2f}{1-f} \nabla f \cdot \nabla) w_{ij} &= \frac{2(|\nabla f|^2 + Rf)}{1-f} w_{ij} + \frac{(Ru)_i u_j + u_i (Ru)_j}{u^2(1-f)^2} \\ &\quad + 2(v_{ki} + f w_{ki})(v_{kj} + f w_{kj}) + R_{ik} w_{kj} + R_{jk} w_{ki}, \end{aligned}$$

where  $v_{ij}$  is as in Lemma 3.1.

**Proof.** The proof is similar to that of Lemma 3.2 in [9].  $\square$

**Proof of Theorem 1.1.** Let  $V = (v_{ij})$ ,  $W = (w_{ij})$ ,  $w = \text{tr} W = g^{ij} w_{ij} = \frac{|\nabla f|^2}{(1-f)^2}$ , and

$$L = \partial_t + \Delta - \frac{2f}{1-f} \nabla f \cdot \nabla.$$

Then by Lemmas 3.1 and 3.2 we have

$$\begin{aligned} LV &= (1-f)wV + P, \\ LW &= 2(1-f)wW + 2(V + fW)^2 + Q, \end{aligned}$$

where  $P$  is a 2-tensor whose  $(i, j)$ -th component in local coordinates is given by

$$\begin{aligned} P_{ij} &= \frac{Rf}{1-f} v_{ij} + \frac{1}{u(1-f)} [2R_{kijl} u_{kl} + R_{il} u_{jl} + R_{jl} u_{il} \\ &\quad + 2(\nabla_i R_{jl} + \nabla_j R_{il} - \nabla_l R_{ij}) \nabla_l u - u \nabla_i \nabla_j R - \nabla_i R \nabla_j u - \nabla_j R \nabla_i u - Ru_{ij}], \end{aligned}$$

and  $Q$  is a 2-tensor whose  $(i, j)$ -th component in local coordinates is given by

$$Q_{ij} = \frac{2Rf}{1-f}w_{ij} + \frac{(Ru)_i u_j + u_i (Ru)_j}{u^2(1-f)^2} + R_{ik}w_{kj} + R_{jk}w_{ki}.$$

Now with the help of Theorem 10 in [7] and Corollary 2.3 here, we can proceed as in Han-Zhang [9].  $\square$

**Proof of Theorem 1.2.** With the help of Theorem 10 in [7], Proposition 2.2 here, and a space-time cutoff function, we can proceed as in Han-Zhang [9] with minor modifications.  $\square$

**Corollary 3.3.** *Let  $(M^n, g(t), t \in [0, T])$ , be a (not necessarily complete) Ricci flow,  $u$  be a positive solution to the conjugate heat equation coupled with the Ricci flow on  $M \times [0, T]$  with  $0 < u \leq A$ . Let  $x_0 \in M$ . Assume that the parabolic cube  $Q_{4r, T}(x_0, T)$  is compact. Then we have*

$$\nabla^2 u \leq u \left( \frac{C_0}{T-t} + \frac{C_0}{r^2} + C_2 \right) \left( 1 + \log \frac{A}{u} \right)^2 g(t) \quad \text{on } Q_{r, T}(x_0, T) \setminus \{(x, T) \mid x \in M\},$$

and, in particular,

$$\Delta u \leq u \left( \frac{C_0 n}{T-t} + \frac{C_0 n}{r^2} + C_2 \right) \left( 1 + \log \frac{A}{u} \right)^2 \quad \text{on } Q_{r, T}(x_0, T) \setminus \{(x, T) \mid x \in M\},$$

where  $C_0$  is a universal constant, and  $C_2$  depends on  $n$ , the upper bounds of  $|Rm|$ ,  $|\nabla Ric|$ , and  $|\nabla^2 R|$  on  $Q_{4r, T}(x_0, T)$ .

**Acknowledgements.** I would like to thank Professor Qi S. Zhang for helpful communications. I was partially supported by NSFC 12271040 and Beijing Natural Science Foundation (Z1900003).

## References

- [1] M. Bailesteanu, X. Cao, A. Pulemotov, Gradient estimates for the heat equation under the Ricci flow. *J. Funct. Anal.* 258 (2010), no. 10, 3517-3542.
- [2] E. Calabi, An extension of E. Hopf's maximum principle with an application to Riemannian geometry. *Duke Math. J.* 25 (1958), 45-56.
- [3] X. Cao, Differential Harnack estimates for backward heat equations with potentials under the Ricci flow. *J. Funct. Anal.* 255 (2008), no. 4, 1024-1038.
- [4] X. Cao, R. S. Hamilton, Differential Harnack estimates for time-dependent heat equations with potentials. *Geom. Funct. Anal.* 19 (2009), no. 4, 989-1000.

- [5] A. Chau, L.-F. Tam, C. Yu, Pseudolocality for the Ricci flow and applications. *Canad. J. Math.* 63 (2011), no. 1, 55-85.
- [6] B. Chow, S.-C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, D. Knopf, P. Lu, F. Luo, L. Ni, The Ricci flow: techniques and applications. Part II. Analytic aspects. *Mathematical Surveys and Monographs*, 144. American Mathematical Society, Providence, RI, 2008.
- [7] K. Ecker, D. Knopf, L. Ni, P. Topping, Local monotonicity and mean value formulas for evolving Riemannian manifolds. *J. Reine Angew. Math.* 616 (2008), 89-130.
- [8] R. S. Hamilton, The formation of singularities in the Ricci flow. *Surveys in differential geometry*, Vol. II, 7-136, Int. Press, Cambridge, MA, 1995.
- [9] Q. Han, Q. S. Zhang, An upper bound for Hessian matrices of positive solutions of heat equations. *J. Geom. Anal.* 26 (2016), no. 2, 715-749.
- [10] H. Huang, Local derivative estimates for the heat equation coupled to the Ricci flow, *Communications in Contemporary Mathematics*, Vol. 24 (2022), No. 06, 2150043.
- [11] S. Kuang, Qi S. Zhang, A gradient estimate for all positive solutions of the conjugate heat equation under Ricci flow. *J. Funct. Anal.* 255 (2008), no. 4, 1008-1023.
- [12] P. Li, S.-T. Yau, On the parabolic kernel of the Schrödinger operator. *Acta Math.* 156 (1986), no. 3-4, 153-201.
- [13] S. Liu, Gradient estimates for solutions of the heat equation under Ricci flow. *Pacific J. Math.* 243 (2009), no. 1, 165-180.
- [14] L. Ni, A note on Perelman's LYH type inequality, *Comm. Anal. Geom.* 14 (2006), 883-905.
- [15] G. Perelman, The entropy formula for the Ricci flow and its geometric applications, arXiv:math/0211159
- [16] J. Sun, Gradient estimates for positive solutions of the heat equation under geometric flow. *Pacific J. Math.* 253 (2011), no. 2, 489-510.
- [17] Q. S. Zhang, Some gradient estimates for the heat equation on domains and for an equation by Perelman. *Int. Math. Res. Not.* 2006, Art. ID 92314, 39 pp.

Laboratory of Mathematics and Complex Systems (Ministry of Education),  
 School of Mathematical Sciences, Beijing Normal University,  
 Beijing 100875, P.R. China  
 E-mail address: hhuang@bnu.edu.cn