

SHARP CRITICAL AND SUBCRITICAL TRACE TRUDINGER-MOSER AND ADAMS INEQUALITIES ON THE UPPER HALF SPACES

LU CHEN, GUOZHEN LU, QIAOHUA YANG, MAOCHUN ZHU

ABSTRACT. In this paper, we establish the sharp critical and subcritical trace Trudinger-Moser and Adams inequalities on the half spaces and prove the existence of their extremals through the method based on the Fourier rearrangement, harmonic extension and scaling invariance. These trace Trudinger-Moser and Adams inequalities can be considered as the borderline case of the Sobolev trace inequalities of first and higher orders. Furthermore, we show the existence of the least energy solutions for a class of bi-harmonic equations with nonlinear Neumann boundary condition associated with the trace Adams inequalities.

Keywords: Trace Trudinger-Moser inequality, Trace Adams inequality, Nonlinear Neumann boundary condition, Harmonic extension, Pohozaev identity, Ground state, Fourier rearrangement.

2010 MSC. 35J60, 35B33, 46E30.

1. INTRODUCTION

The main content of this paper is concerned with the problem of finding optimal trace Trudinger-Moser and Adams inequalities and existence of their extremals in unbounded domain. Sharp Trudinger-Moser inequalities and its high-order form (Adams inequalities) have been a matter of intensive research due to the importance of these inequalities in application to problems in analysis, PDEs, differential geometry, mathematical physics, etc.

The Trudinger inequality as the critical case of the Sobolev imbedding was obtained by Trudinger [53], Pohazaev [49] and Yudovic [57]. More precisely, Trudinger employed the power series expansion to prove that there exists $\alpha > 0$ such that

$$(1.1) \quad \sup_{\|\nabla u\|_n^n \leq 1, u \in W_0^{1,n}(\Omega)} \int_{\Omega} \exp(\alpha |u|^{\frac{n}{n-1}}) dx < \infty,$$

The first author was supported by NNSF (No.11901031); the second author was partly supported by a grant from the Simons Foundation; the third author was partially supported by NNSF (No.12071353); the fourth author was partially supported by NNSF (No. 12071185).

where $\Omega \subseteq \mathbb{R}^n$ is a smooth bounded domain and $W_0^{1,n}(\Omega)$ denotes the usual Sobolev space, i.e, the completion of $C_0^\infty(\Omega)$ with the norm

$$\|u\|_{W_0^{1,n}(\Omega)} = \left(\int_{\Omega} |\nabla u|^n dx \right)^{\frac{1}{n}}.$$

Subsequently, Moser [45] utilized the technique of the symmetry and rearrangement to give the sharp constants $\alpha_n = nw_{n-1}^{\frac{1}{n-1}}$ of the above inequality. Such an inequality in the sharp form of constant α_n is now called the *Trudinger-Moser* inequality. Since the Polyá-Szegö inequality on which the technique of the symmetry and rearrangement depends is not valid on the high-order Sobolev space, this adds much challenge to the research of high-order Trudinger-Moser inequalities, namely Adams type inequalities. Adams employed the method of representation formulas together the method of the rearrangement of convolutions by O'Neil [48] to establish the sharp high-order Trudinger-Moser inequality on any bounded domain. More precisely, he proved the following result.

Theorem A. ([4]) *Let Ω be an open and bounded set in \mathbb{R}^n . If m is a positive integer less than n , then there exists a constant $C_0 = C(n, m) > 0$ such that for any $u \in W_0^{m, \frac{n}{m}}(\Omega)$ and $\|\nabla^m u\|_{\frac{n}{m}} \leq 1$, then*

$$(1.2) \quad \frac{1}{|\Omega|} \int_{\Omega} \exp(\beta |u(x)|^{\frac{n}{n-m}}) dx \leq C_0,$$

for all $\beta \leq \beta(n, m)$, where

$$\beta(n, m) = \begin{cases} \frac{n}{\omega_{n-1}} \left[\frac{\pi^{\frac{n}{2}} 2^m \Gamma(\frac{m+1}{2})}{\Gamma(\frac{n-m+1}{2})} \right]^{\frac{n}{n-m}}, & \text{if } m \text{ is odd.} \\ \frac{n}{\omega_{n-1}} \left[\frac{\pi^{\frac{n}{2}} 2^m \Gamma(\frac{m}{2})}{\Gamma(\frac{n-m}{2})} \right]^{\frac{n}{n-m}}, & \text{if } m \text{ is even.} \end{cases}$$

Furthermore, the constant $\beta(n, m)$ is best possible in the sense that for any $\beta > \beta(n, m)$, the integral can be made as large as possible.

Later, Moser's results for the first order derivatives and Adams' result for the high order derivatives have been extended to the unbounded domains. The first order Trudinger-Moser inequality was proved in [8], [18] and [2] in the subcritical case, that is for any $\alpha < \alpha_n$, there exists a constant $C = C(\alpha, n)$ such that

$$(1.3) \quad \sup_{\|\nabla u\|_n^n \leq 1} \int_{\mathbb{R}^n} \Phi(\alpha |u|^{\frac{n}{n-1}}) dx \leq C \|u\|_n^n,$$

where $\Phi(t) = e^t - \sum_{j=0}^{n-2} t^j$.

Later, it is showed in [35, 51] that the exponent α_n becomes admissible if the Dirichlet norm $(\int_{\mathbb{R}^n} |\nabla u|^n dx)^{\frac{1}{n}}$ is replaced by Sobolev norm $\|u\|_{W^{1,n}(\mathbb{R}^n)} = (\int_{\mathbb{R}^n} (|u|^n + |\nabla u|^n) dx)^{\frac{1}{n}}$,

more precisely, they proved the following critical Trudinger-Moser inequality

$$(1.4) \quad \sup_{\|u\|_{W^{1,n}(\mathbb{R}^n)} \leq 1} \int_{\mathbb{R}^n} \Phi(\alpha_n |u|^{\frac{n}{n-1}}) dx \leq C_n < \infty.$$

All the proofs of both critical and subcritical Trudinger-Moser inequalities in the literature rely on rearrangement argument and the Polyá-Szegö inequality. The authors of [24], [25] used a symmetrization-free approach to give a simple proof for both critical and subcritical sharp Trudinger-Moser inequalities in $W^{1,n}(\mathbb{R}^n)$. It should be noted that this approach is surprisingly simple and can be easily applied to other settings where symmetrization argument does not work. Furthermore, they also develop this new approach to establish the global Trudinger-Moser inequalities on the entire Heisenberg group from the local one derived in [17] and fractional Adams inequalities in $W^{s,\frac{n}{s}}(\mathbb{R}^n)$ ($0 < s < n$) (see [24, 25, 27]). For more applications of the symmetrization-free approach, please also see [28, 31, 33, 54]. Moreover, the critical and subcritical Trudinger-Moser inequalities are equivalent as shown in [28].

As far as the existence of extremal functions of Trudinger-Moser inequality on bounded domains, the first breakthrough was due to Carleson and Chang [9] who established the existence on balls. (See also [20], [36] for smooth domains and the methods of using expansion formula of Dirichlet integrals [38], [40].)

Since the main purpose of this paper is to establish the trace Trudinger-Moser and Adams type inequalities on unbounded domains, we will only briefly review of the trace Trudinger-Moser and Adams inequalities on bounded domains. Let Ω be bounded domain in \mathbb{R}^n with smooth boundary and define the $W^{1,2}(\Omega)$ to be the Sobolev space, equipped with the norm

$$\|u\|_{W^{1,2}(\Omega)} = \left(\int_{\Omega} (|\nabla u|^2 + |u|^2) dx \right)^{\frac{1}{2}}.$$

As is well known, classical Sobolev trace embedding on bounded domain asserts that $W^{1,2}(\Omega) \subset L^q(\partial\Omega)$ for $1 \leq q \leq \frac{2(n-1)}{n-2}$ and $n > 3$. In the borderline case $n = 2$, the $W^{1,2}(\Omega) \subset L^q(\partial\Omega)$ for $1 \leq q < \infty$, but $W^{1,2}(\Omega) \not\subset L^\infty(\partial\Omega)$. To fill this gap, the authors of [3, 5, 16, 15, 34] studied the trace Trudinger-Moser inequalities and established the following

Theorem B. Let Ω be a sufficiently smooth bounded domain in \mathbb{R}^2 . For any $\alpha \leq \pi$, then there exists a positive constant C_n such that

$$(1.5) \quad \int_{\partial\Omega} \exp(\alpha |u|^2) d\sigma \leq C_n,$$

for any $u \in C^\infty(\Omega)$ with $\int_{\Omega} (|\nabla u|^2 + |u|^2) dx \leq 1$.

We should mention that the critical trace inequality of Trudinger-Moser type on balls in \mathbb{R}^2 for holomorphic functions with mean value zero was first established by Chang and Marshall [11]. Recently, the authors in [32] establish the Chang-Marshall type inequality to sufficiently smooth domains Ω in \mathbb{R}^n for all $n \geq 2$ for any Sobolev function u with mean

value $\int_{\Omega} u dx = 0$. We should also note that the author of [16] established Trudinger-Moser trace type inequalities on lower dimensional subsets of Ω with respect to measures supported on the subsets.

In a half space, a particular unbounded domain with the smooth boundary, sharp geometrical inequalities such as trace Sobolev inequalities on this domain have attracted much attention due to their importance in geometrical analysis and PDEs (see [1, 6, 19, 46, 55]). We recall some classical results that involve the sharp trace Sobolev inequalities on the half space. Sharp trace Sobolev inequalities for first order derivative of Escobar [19] assert that

Theorem C.([19]) Let $n \geq 3$. Then for any $U \in W^{1,2}(\overline{\mathbb{R}_+^n})$, there holds

$$(1.6) \quad 2 \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n-1}{2})} w_{n-1}^{\frac{1}{n-1}} \left(\int_{\partial \mathbb{R}_+^n} |U(x, 0)|^{\frac{2(n-1)}{n-2}} dx \right)^{\frac{n-2}{n-1}} \leq \int_{\mathbb{R}_+^n} |\nabla U(x, y)|^2 dx dy,$$

where $(x, y) \in \mathbb{R}^{n-1} \times \mathbb{R}^+$. Furthermore, the equality case is achieved by a harmonic extension of a function of the form $c(1 + |x - x_0|^2)^{-\frac{n-2}{2}}$, where c is a constant, $x \in \mathbb{R}^{n-1}$, x_0 is some fixed point in \mathbb{R}^{n-1} .

By conformal map, this inequality is in fact equivalent to the trace Sobolev inequalities on the ball (see Beckner [6]),

$$(1.7) \quad \frac{n-2}{2} w_{n-1}^{\frac{1}{n-1}} \left(\int_{\mathbb{S}^{n-1}} |f|^{\frac{2(n-1)}{n-2}} d\sigma \right)^{\frac{n-2}{n-1}} \leq \int_{\mathbb{B}^n} |\nabla v|^2 dx + \frac{n-2}{2} \int_{\mathbb{S}^{n-1}} |f|^2 d\sigma,$$

where v is a smooth extension of f to \mathbb{B}^n and $d\sigma$ is the surface measure on \mathbb{S}^{n-1} .

Motivated by the above works, we are concerned with the borderline case of trace Sobolev inequality, that is the sharp trace Trudinger-Moser inequalities in $W^{1,2}(\mathbb{R}_+^2)$. We first establish the subcritical and critical trace-type Trudinger-Moser inequalities on the half space. Our results read as

Theorem 1.1. *For $0 < \beta < \pi$, then there exists a positive constant C such that for all functions $U \in W^{1,2}(\overline{\mathbb{R}_+^2})$ with $\int_{\mathbb{R}_+^2} |\nabla U(x, y)|^2 dx dy \leq 1$, the following inequality holds*

$$(1.8) \quad \int_{\partial \mathbb{R}_+^2} (\exp(\beta |U(x, 0)|^2) - 1) dx \leq C \int_{\partial \mathbb{R}_+^2} |U(x, 0)|^2 dx.$$

Furthermore, the smallest C in (1.8) can be attained by some U , which is a harmonic extension of some even function $u \in W^{\frac{1}{2},2}(\mathbb{R})$. Moreover, the constant π is sharp in the sense that the inequality (1.8) fails if the constant β is replaced by any $\beta \geq \pi$.

Theorem 1.2. *There exists a positive constant C such that for all functions $U \in W^{1,2}(\overline{\mathbb{R}_+^2})$ with*

$$\int_{\mathbb{R}_+^2} |\nabla U(x, y)|^2 dx dy + \int_{\partial \mathbb{R}_+^2} |U(x, 0)|^2 dx \leq 1,$$

the following inequality holds

$$(1.9) \quad \int_{\partial\mathbb{R}_+^2} \exp(\pi|U(x, 0)|^2) dx \leq C.$$

Furthermore, the smallest C in (1.9) can be attained by some function U , which is a harmonic extension of some radial function $u \in W^{\frac{1}{2}, 2}(\mathbb{R})$. Moreover, the constant π is sharp in the sense that the inequality (1.9) fails if the constant π is replaced by any $\beta > \pi$.

Remark 1.3. The difference between Theorem 1.1 and Theorem 1.2 is that we assume only $\int_{\mathbb{R}_+^2} |\nabla U(x, y)|^2 dx dy \leq 1$ in Theorem 1.1 and assume

$$\int_{\mathbb{R}_+^2} |\nabla U(x, y)|^2 dx dy + \int_{\partial\mathbb{R}_+^2} |U(x, 0)|^2 dx \leq 1$$

in Theorem 1.2. This results in a **subcritical** trace Trudinger-Moser inequality in Theorem 1.1 and a **critical** trace Trudinger-Moser inequality in Theorem 1.2.

Recently, Ache and Chang [1] established the second order sharp trace Sobolev-inequalities (1.7) on the ball:

Theorem D. Let $u \in C^\infty(\mathbb{S}^{n-1})$ with $n > 4$. Suppose that v is a smooth extension of u to the unit ball \mathbb{B}^n which satisfies the Neumann boundary condition

$$\frac{\partial v}{\partial n}|_{\mathbb{S}^{n-1}} = -\frac{n-4}{2}u.$$

Then the following inequality holds

$$(1.10) \quad \begin{aligned} & 2 \frac{\Gamma(\frac{n+2}{2})}{\Gamma(\frac{n-4}{2})} w_{n-1}^{\frac{3}{n-1}} \left(\int_{\mathbb{S}^{n-1}} |u|^{\frac{2(n-1)}{n-4}} d\sigma \right)^{\frac{n-4}{n-1}} \\ & \leq \int_{\mathbb{B}^n} |\Delta v|^2 dx + 2 \int_{\mathbb{S}^{n-1}} |\tilde{\nabla} u|^2 d\sigma + \frac{n(n-4)}{2} \int_{\mathbb{S}^{n-1}} |u|^2 \sigma, \end{aligned}$$

where Δv is the Laplacian of v with respect to the Euclidean metric, and $\tilde{\nabla}$ is the gradient on the sphere.

We refer the reader to extensions to higher order derivatives trace Sobolev inequalities on half spaces in [56], [55] and [10].

By the Mobius transform, inequality (1.10) is equivalent to the following trace Sobolev inequality on the upper half space \mathbb{R}_+^n .

Theorem E ([46]) Let $n \geq 5$. Then for any $U \in W^{2,2}(\overline{\mathbb{R}_+^n})$ satisfying the Neumann boundary condition $\partial_y U(x, y)|_{y=0} = 0$, we have

$$(1.11) \quad 2 \frac{\Gamma(\frac{n+2}{2})}{\Gamma(\frac{n-4}{2})} w_n^{\frac{3}{n-1}} \left(\int_{\partial\mathbb{R}_+^n} |U(x, 0)|^{\frac{2(n-1)}{n-4}} dx \right)^{\frac{n-4}{n-1}} \leq \int_{\mathbb{R}_+^n} |\Delta U(x, y)|^2 dx dy.$$

Furthermore, the equality is achieved by a bi-harmonic extension of a function of the form $c(1 + |x - x_0|^2)^{-\frac{n-4}{2}}$, where c is a constant, $x \in \mathbb{R}^{n-1}$, x_0 is some fixed point in \mathbb{R}^{n-1} .

In this paper, we first establish the borderline case of the inequality (1.11) when $n = 4$. We employ the method based on the sharp Fourier rearrangement principle and bi-harmonic extension to establish the sharp trace Adams inequalities on the half space and the existence of their extremals. Both subcritical and critical trace Adams inequalities are proved for functions in $W^{2,2}(\overline{\mathbb{R}_+^4})$.

Our first result in the second order trace Adams inequality is the following subcritical inequality.

Theorem 1.4. *Let $0 < \beta < 12\pi^2$. Then there exists a positive constant C such that for all functions $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ satisfying the Neumann boundary condition $\partial_y U(x, y)|_{y=0} = 0$ with $\int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy \leq 1$, the following inequality holds:*

$$(1.12) \quad \int_{\partial\mathbb{R}_+^4} (\exp(\beta|U(x, 0)|^2) - 1) dx \leq C \int_{\partial\mathbb{R}_+^4} |U(x, 0)|^2 dx.$$

Furthermore, the smallest C in (1.12) can be attained by some function U , which is the bi-harmonic extension of some radial function $u \in W^{\frac{3}{2},2}(\mathbb{R}^3)$. Moreover, the constant $12\pi^2$ is sharp in the sense that the inequality fails if the constant β is replaced by any $\beta \geq 12\pi^2$.

Next, the following critical trace Adams inequality also holds.

Theorem 1.5. *There exists a positive constant C such that for all functions $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ satisfying the Neumann boundary condition $\partial_y U(x, y)|_{y=0} = 0$ with*

$$\int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy + \int_{\partial\mathbb{R}_+^4} |U(x, 0)|^2 dx \leq 1,$$

the following inequality holds

$$(1.13) \quad \int_{\partial\mathbb{R}_+^4} \exp(12\pi^2|U(x, 0)|^2) dx \leq C.$$

Moreover, the constant $12\pi^2$ is sharp in the sense that the inequality fails to hold uniformly for all $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ if the constant $12\pi^2$ is replaced by any $\beta > 12\pi^2$.

The second order Adams inequality with the exact growth was established in $W^{2,2}(\mathbb{R}^4)$ in [41] and then was extended to $W^{2,n}(\mathbb{R}^n)$ in [37] (see also the first order case in [22], and [42] and singular inequalities and under different norms with exact growth in [26] and [29]). A natural question is whether there exists trace Adams inequality with the exact growth in $W^{2,2}(\mathbb{R}_+^4)$. In this paper, we establish the following

Theorem 1.6. *There exists a positive constant C such that for all functions $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ satisfying the Neumann boundary condition $\partial_y U(x, y)|_{y=0} = 0$ with $\int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy = 1$, the following inequality holds*

$$(1.14) \quad \int_{\partial\mathbb{R}_+^4} \frac{(\exp(12\pi^2|U(x, 0)|^2) - 1)}{(1 + |U(x, 0)|)^2} dx \leq C \int_{\partial\mathbb{R}_+^4} |U(x, 0)|^2 dx.$$

Moreover, this inequality fails to hold uniformly for all $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ if the power 2 in the denominator of the left hand side is replaced by any $p < 2$.

Remark 1.7. The proof of Theorem 1.6 is based on bi-harmonic extension and fractional Adams inequalities with the exact growth in $W^{\frac{n}{2},2}(\mathbb{R}^n)$ for $n = 3$. Adams inequalities with the exact growth in $W^{\frac{n}{2},2}(\mathbb{R}^n)$ for even n has been established in [43] (see also [47]). However, the validity of this inequality for odd n still remain open. Using the fractional Hardy-Rellich inequalities established by Beckner in [7] together with the method of the reduction of orders (see also [47]), we present a simple proof for the fractional Trudinger-Moser-Adams inequality with the exact growth in $W^{\frac{n}{2},2}(\mathbb{R}^n)$ for all $n \geq 2$.

Theorem 1.8. There exists a positive constant C such that for all functions $u \in W^{\frac{n}{2},2}(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} |(-\Delta)^{\frac{n}{4}} u|^2 dx = 1$, the following inequality holds

$$(1.15) \quad \int_{\mathbb{R}^n} \frac{(\exp(\beta(n, \frac{n}{2})|u(x)|^2) - 1)}{(1 + |u(x)|)^2} dx \leq C \int_{\mathbb{R}^n} |u(x)|^2 dx.$$

Moreover, this inequality fails if the power 2 in the denominator is replaced by any $p < 2$.

Furthermore, we also establish the trace Adams inequalities on $W^{m,2}(\overline{\mathbb{R}_+^{2m}})$ ($m > 2$). The proof is based on the following technical lemma whose proof will be given in Section 5.

Lemma 1.9. Given a function $u \in W^{m-\frac{1}{2},2}(\mathbb{R}^{2m-1})$. If we assume that the function $U \in W^{m,2}(\overline{\mathbb{R}_+^{2m}})$ satisfying

$$(-\Delta)^m U = 0$$

on the upper half space \mathbb{R}_+^{2m} and the Neumann boundary conditions

$$(1.16) \quad \Delta^k U(x, y)|_{y=0} = \frac{\Gamma(m)\Gamma(m - \frac{1}{2} - k)}{\Gamma(m - 1 - k)\Gamma(m - \frac{1}{2})} \Delta_x^k u(x), \quad 0 \leq k \leq \left[\frac{m-1}{2} \right]$$

and

$$(1.17) \quad \partial_y \Delta^k U(x, y)|_{y=0} = 0, \quad 0 \leq k \leq \left[\frac{m-2}{2} \right],$$

then we have the following identity

$$\int_{\mathbb{R}_+^{2m}} |\nabla^m U(x, y)|^2 dx dy = \frac{\Gamma(m)\Gamma(\frac{1}{2})}{\Gamma(m - \frac{1}{2})} \int_{\mathbb{R}^{2m-1}} |(-\Delta)^{\frac{2m-1}{4}} u|^2 dx.$$

Once we have proved Lemma 1.9, we can establish the following both subcritical and critical trace Adams inequalities on $W^{m,2}(\overline{\mathbb{R}_+^{2m}})$ ($m > 2$) by a similar argument as for Theorems 1.4, 1.5 and 1.6 on $W^{2,2}(\overline{\mathbb{R}_+^4})$.

Theorem 1.10. Let $0 < \beta < \frac{\Gamma(m)\Gamma(\frac{1}{2})}{\Gamma(m - \frac{1}{2})} \beta(2m - 1, \frac{2m-1}{2}) = \widetilde{\beta}_m$. Then there exists a positive constant C such that for all functions $U \in W^{m,2}(\overline{\mathbb{R}_+^{2m}})$ satisfying the Neumann boundary

conditions (1.16) and (1.17) with $\int_{\mathbb{R}_+^{2m}} |\nabla^m U|^2 dx dy \leq 1$, the following inequality holds:

$$(1.18) \quad \int_{\partial\mathbb{R}_+^{2m}} (\exp(\beta|U(x,0)|^2) - 1) dx \leq C \int_{\partial\mathbb{R}_+^{2m}} |U(x,0)|^2 dx.$$

Moreover, the constant $\widetilde{\beta}_m$ is sharp in the sense that the inequality fails to hold uniformly if the constant β is replaced by any $\beta \geq \widetilde{\beta}_m$. Furthermore, equality in (1.12) holds if and only if U is a m -harmonic extension of some radial function u in $W^{\frac{2m-1}{2},2}(\mathbb{R}^{2m-1})$.

Theorem 1.11. *There exists a positive constant C such that for all functions $U \in W^{m,2}(\overline{\mathbb{R}_+^{2m}})$ satisfying the Neumann boundary conditions (1.16) and (1.17) with*

$$\int_{\mathbb{R}_+^{2m}} |\nabla^m U(x,y)|^2 dx dy + \int_{\partial\mathbb{R}_+^{2m}} |U(x,0)|^2 dx \leq 1,$$

the following inequality holds

$$(1.19) \quad \int_{\partial\mathbb{R}_+^{2m}} (\exp(\widetilde{\beta}_m|U(x,0)|^2) - 1) dx \leq C.$$

Moreover, the constant $\widetilde{\beta}_m$ is sharp in the sense that the inequality fails to hold uniformly if the constant $\widetilde{\beta}_m$ is replaced by any $\beta > \widetilde{\beta}_m$.

Theorem 1.12. *There exists a positive constant C such that for all functions $U \in W^{m,2}(\overline{\mathbb{R}_+^{2m}})$ satisfying the Neumann boundary condition (1.16) and (1.17) with*

$$\int_{\mathbb{R}_+^{2m}} |\nabla^m U(x,y)|^2 dx dy = 1,$$

the following inequality holds

$$(1.20) \quad \int_{\partial\mathbb{R}_+^{2m}} \frac{(\exp(\widetilde{\beta}_m|U(x,0)|^2) - 1)}{(1 + |U(x,0)|)^2} dx \leq C \int_{\partial\mathbb{R}_+^{2m}} |U(x,0)|^2 dx.$$

Moreover, this inequality fails if the power 2 in the denominator on the left hand side is replaced by any $p < 2$.

Finally, we are also concerned with the ground-state solutions to the equation associated with the trace Adams inequality in $W^{m,2}(\overline{\mathbb{R}_+^{2m}})$. For simplicity, we only consider the case $m = 2$. By the Euler-Lagrange multiplier theorem, extremals of the trace Adams supremum

$$\sup_{\int_{\mathbb{R}_+^4} |\Delta U(x,y)|^2 dx dy + \int_{\partial\mathbb{R}_+^4} |U(x,y)|^2 dx \leq 1} \int_{\partial\mathbb{R}_+^4} (\exp(12\pi^2 |U|^2) - 1) dx$$

must satisfy the following Euler-Lagrange equation with nonlinear Neumann boundary conditions: there exists some λ such that

$$(1.21) \quad \begin{cases} \Delta^2 U(x, y) = 0 \text{ for } (x, y) \in \mathbb{R}^3 \times \mathbb{R}^+, \\ (-\frac{\partial(\Delta U)}{\partial y} + U(x, y))|_{y=0} = \lambda U(x, y) \exp(12\pi^2 U^2(x, y))|_{y=0}, \\ \frac{\partial U}{\partial y}|_{y=0} = 0. \end{cases}$$

We are interested in the existence of ground-state solutions to equation (1.21) with some fixed λ . Define

$$E := \{U \in W^{2,2}(\overline{\mathbb{R}_+^4}) \cap L^2(\partial\mathbb{R}_+^4) : \partial_y U(x, y)|_{y=0} = 0\},$$

the corresponding functional of equation (1.21) is

$$(1.22) \quad I_\lambda(U) = \frac{1}{2} \int_{\mathbb{R}_+^4} |\Delta U|^2 dx dy + \frac{1}{2} \int_{\partial\mathbb{R}_+^4} |U|^2 dx - \frac{\lambda}{24\pi^2} \int_{\partial\mathbb{R}_+^4} (\exp(12\pi^2 |U|^2) - 1) dx.$$

It is easy to check that $I_\lambda \in C^1(E, \mathbb{R})$, and

$$(1.23) \quad \begin{aligned} I'_\lambda(U)V &= \int_{\mathbb{R}_+^4} \Delta U(x, y) \Delta V(x, y) dx dy + \int_{\partial\mathbb{R}_+^4} U(x, 0) V(x, 0) dx \\ &\quad - \lambda \int_{\partial\mathbb{R}_+^4} \exp(12\pi^2 |U(x, 0)|^2) U(x, 0) V(x, 0) dx, \quad U, V \in E. \end{aligned}$$

Define

$$M_\lambda = \inf\{I'_\lambda(U) = 0 \mid U \in E\}.$$

By using the sharp trace Adams inequalities (1.13) and variational method, we can prove the following result.

Theorem 1.13. *For any $0 < \lambda < 1$, there exists $V \in E$ such that $I_\lambda(V) = M_\lambda$, that is to say that the problem (1.21) has a ground state solution provided $0 < \lambda < 1$.*

This paper is organized as follows. In Section 2, we establish the sharp subcritical and critical trace Trudinger-Moser inequalities in \mathbb{R}^2 and existence of their extremal functions based on the harmonic extension. Section 3 is devoted to the proof of the sharp trace Adams inequalities and the existence of the extremals for subcritical trace Adams inequalities in \mathbb{R}^4 . In Section 4, we are concerned with the existence of a ground state solution to a class of problems with the nonlinear Neumann boundary condition on the half space by using the trace Adams inequality, sharp Fourier rearrangement principle and variational method. Section 5 is devoted to the proof of Lemma 1.9 which is the main technical part of establishing Theorems 1.10, 1.11 and 1.12 in the case $m > 2$ in \mathbb{R}^{2m} by adapting the same argument of proving Theorems 1.4 and 1.5 in the case $m = 2$. In Section 6, we give the proof of the fractional Adams inequalities with the exact growth in $W^{\frac{n}{2}, 2}(\mathbb{R}^n)$, which is needed in proving Theorem 1.6 and (1.20).

2. TRACE TRUDINGER-MOSER INEQUALITIES ON THE HALF SPACE \mathbb{R}_+^2

In this section, we consider the sharp subcritical and critical trace Trudinger-Moser inequalities and the existence of their extremal functions. The method is based on the harmonic extension, sharp subcritical fractional Adams inequalities in $W^{\frac{1}{2},2}(\mathbb{R})$ and existence of their extremals. First, we introduce some known results about the fractional Adams inequalities and the harmonic extension.

Lemma 2.1. ([21]) *For $0 < \beta < \pi$, then there exists a positive constant C such that for all functions $u \in W^{\frac{1}{2},2}(\mathbb{R})$ with $\|(-\Delta)^{\frac{1}{4}}u\|_2 = 1$, the following inequality holds.*

$$(2.1) \quad \int_{\mathbb{R}} (\exp(\beta|u|^2) - 1) dx \leq C \int_{\mathbb{R}} |u|^2 dx,$$

Moreover, the constant π is sharp in the sense that the inequality fails if the constant β is replaced by any $\beta \geq \pi$.

Lemma 2.2. *There exists extremals for the above subcritical fractional Adams inequality (2.1). Furthermore, all the extremals of inequality (2.1) must be radially symmetric with respect to some point $x_0 \in \mathbb{R}$.*

Remark 2.3. *The proof can be found in [13]. One can also use the technique of proof of Lemma 3.2 to prove Lemma 2.2, then we omit the details.*

Lemma 2.4. ([50],[52]) *Given a function $u \in W^{\frac{1}{2},2}(\mathbb{R})$. If we assume that function $U \in W^{2,2}(\overline{\mathbb{R}_+^2})$ satisfying*

$$-\Delta U(x, y) = 0$$

on the upper half space \mathbb{R}_+^2 and the boundary condition

$$U(x, y)|_{y=0} = u(x),$$

then we have the following identity

$$(2.2) \quad \int_{\mathbb{R}_+^2} |\nabla U(x, y)|^2 dx dy = \int_{\mathbb{R}} |(-\Delta)^{\frac{1}{4}}u|^2 dx.$$

Now, we are in the position to prove Theorem 1.1.

Proof of Theorem 1.1: For any $U \in W^{1,2}(\overline{\mathbb{R}_+^2})$, we define $V(x, y)$ as the harmonic extension of $U(x, 0)$, that is to say $V(x, y)$ satisfies that

$$-\Delta V(x, y) = 0$$

on the upper half space \mathbb{R}_+^2 and the boundary condition

$$V(x, y)|_{y=0} = U(x, y)|_{y=0}.$$

Through Green's representation formula, we can write

$$V(x, y) = \frac{1}{\pi} \int_{\mathbb{R}} \frac{y}{(|x - \tilde{x}| + y^2)} U(\tilde{x}, 0) d\tilde{x}.$$

According to the Dirichlet principle, we know that

$$(2.3) \quad \int_{\mathbb{R}_+^2} |\nabla V(x, y)|^2 dx dy \leq \int_{\mathbb{R}_+^2} |\nabla U(x, y)|^2 dx dy.$$

Combining (2.3) and (2.2) and Lemma 2.1, we obtain

$$(2.4) \quad \begin{aligned} \int_{\partial\mathbb{R}_+^2} (\exp(\beta|U(x, 0)|^2) - 1) dx &= \int_{\partial\mathbb{R}_+^2} (\exp(\beta|V(x, 0)|^2) - 1) dx \\ &\leq C \int_{\partial\mathbb{R}_+^2} |V(x, 0)|^2 dx = C \int_{\partial\mathbb{R}_+^2} |U(x, 0)|^2 dx. \end{aligned}$$

The sharpness of the above inequality can be deduced from the sharpness of the sharp subcritical inequalities (2.1). In fact, we can pick the test function $u_k \in W^{\frac{1}{2}, 2}(\mathbb{R})$ satisfying $\|(-\Delta)^{\frac{1}{4}} u_k\|_2 \leq 1$ such that

$$\lim_{k \rightarrow \infty} \frac{\int_{\mathbb{R}} (\exp(\pi|u_k|^2) - 1) dx}{\|u_k\|_2^2} = \infty.$$

Define $U_k(x, y)$ as the harmonic extension of u_k , through Lemma 2.4, we know that $\int_{\mathbb{R}_+^2} |\nabla U_k(x, y)|^2 dx dy = \int_{\mathbb{R}} |(-\Delta)^{\frac{1}{4}} u_k|^2 dx \leq 1$, then it follows that

$$\lim_{k \rightarrow \infty} \frac{\int_{\partial\mathbb{R}_+^2} (\exp(\pi|U_k|^2) - 1) dx}{\int_{\partial\mathbb{R}_+^2} |U_k(x, 0)|^2 dx} = \lim_{k \rightarrow \infty} \frac{\int_{\mathbb{R}} (\exp(\pi|u_k|^2) - 1) dx}{\|u_k\|_2^2} = \infty,$$

which implies the sharpness of the inequality (1.8).

Next, we show the existence of extremals for subcritical trace Trudinger-Moser inequalities. According to Lemma 2.2, we know for any $\beta < \pi$, there exists $u_0(x) \in W^{\frac{1}{2}, 2}(\mathbb{R})$ satisfying $\|(-\Delta)^{\frac{1}{4}} u_0\|_2^2 = 1$ such that

$$\frac{\int_{\mathbb{R}} [\exp(\beta|u_0|^2) - 1] dx}{\int_{\mathbb{R}} |u_0|^2 dx} = \sup_{u \in W^{\frac{1}{2}, 2}(\mathbb{R}), \|(-\Delta)^{\frac{1}{4}} u\|_2^2 = 1} \frac{\int_{\mathbb{R}} [\exp(\beta|u|^2) - 1] dx}{\int_{\mathbb{R}} |u|^2 dx}.$$

Define $U(x, y)$ as the harmonic extension of $u_0(x)$, through Lemma 2.4, we see that $U(x, y)$ is an extremal function for (1.8).

Finally, we prove that the extremals of the inequality (1.8) must be a harmonic extension of some radial function u in $W^{\frac{1}{2}, 2}(\mathbb{R})$. In fact, assume that $U(x, y)$ is the extremal function of the inequality (1.8), we define $W(x, y)$ as the harmonic extension of $U(x, 0)$ and can easily check that $W(x, y)$ is also the extremal function of the inequality with the

$$\int_{\mathbb{R}_+^2} |\nabla U|^2 dx dy = \int_{\mathbb{R}_+^2} |\nabla W|^2 dx dy = 1.$$

According to the Dirichlet principle, we know that $\int_{\mathbb{R}_+^2} |\nabla U|^2 dx dy = \int_{\mathbb{R}_+^2} |\nabla W|^2 dx dy$ if and only if $U = W$. Furthermore, it is also easy to check that $U(x, 0)$ is the extremal functions of the inequality (2.1). According to the Lemma 2.2, the $U(x, 0)$ must be radially

symmetric with respect to some point $x_0 \in \mathbb{R}$. Hence the extremals of the (1.8) must be a harmonic extension of some radial function.

Proof of Theorem 1.2: For any $U \in W^{1,2}(\overline{\mathbb{R}_+^2})$ satisfying

$$\int_{\mathbb{R}_+^2} |\nabla U(x, y)|^2 dx dy + \int_{\partial\mathbb{R}_+^2} |U(x, 0)|^2 dx \leq 1,$$

we define $V(x, y)$ as the harmonic extension of $U(x, 0)$. By the Dirichlet principle and Lemma 2.2, we derive that $\int_{\partial\mathbb{R}_+^2} (|(-\Delta)^{\frac{1}{4}} V(x, 0)|^2 + |V(x, 0)|^2) dx \leq 1$. With the help of the critical fractional Trudinger-Moser inequality in $W^{\frac{1}{2},2}(\mathbb{R})$ (see [23]), we conclude that

$$(2.5) \quad \int_{\partial\mathbb{R}_+^2} (\exp(\pi|U(x, 0)|^2) - 1) dx = \int_{\partial\mathbb{R}_+^2} (\exp(\pi|V(x, 0)|^2) - 1) dx \leq C,$$

which accomplishes the proof of the inequality (1.9). The existence of extremal of the inequality (1.9) relies on the existence of the critical fractional Trudinger-Moser inequality in $W^{\frac{1}{2},2}(\mathbb{R})$ which was recently established by Mancini and Martinazzi in [39]. The proof is similar to the proof of the existence of extremals of the subcritical trace Trudinger-Moser inequalities, we omit the details here.

3. TRACE ADAMS INEQUALITIES ON THE HALF SPACE \mathbb{R}_+^4

In this section, we are devoted to establishing the sharp trace Adams inequalities and the existence of their extremal functions. The method is based on the bi-harmonic, Fourier rearrangement of [30] and sharp fractional Adams inequalities. We note that the method of Fourier rearrangement to establish the existence of extremals to Adams inequalities have been used earlier in [12] and [14].

For this purpose, we need the following lemma.

Lemma 3.1. ([21]) *For $0 < \beta < 6\pi^2$, then there exists a positive constant C such that for all functions $u \in W^{\frac{3}{2},2}(\mathbb{R}^3)$ with $\|(-\Delta)^{\frac{3}{4}} u\|_2 = 1$, the following inequality holds.*

$$(3.1) \quad \int_{\mathbb{R}^3} (\exp(\beta|u|^2) - 1) dx \leq C \int_{\mathbb{R}^3} |u|^2 dx,$$

Moreover, the constant $6\pi^2$ is sharp in the sense that the inequality fails if the constant β is replaced by any $\beta \geq 6\pi^2$.

Next, we will prove the following

Lemma 3.2. *There exists an extremal for the above subcritical fractional Adams inequality (3.1). Furthermore, all the extremals of inequality (3.1) must be radially symmetric with respect to some point $x_0 \in \mathbb{R}^3$.*

Proof. Define the sharp constant μ_β by

$$\mu_\beta := \sup_{u \in W^{\frac{3}{2},2}(\mathbb{R}^3), \|(-\Delta)^{\frac{3}{4}} u\|_2 = 1} F_\beta(u),$$

where

$$F_\beta(u) := \frac{\int_{\mathbb{R}^3} (\exp(\beta|u|^2) - 1) dx}{\|u\|_2^2}.$$

We first employ the Fourier rearrangement tools to prove that there exists a radially maximizing sequence for μ_β . In fact, assume that (u_k) is a maximizing sequence for μ_β , that is

$$\|(-\Delta)^{\frac{3}{4}} u_k\|_2 = 1, \quad \lim_{k \rightarrow \infty} F_\beta(u_k) \rightarrow \mu_\beta.$$

Define u_k^\sharp by $u_k^\sharp = \mathcal{F}^{-1}\{(\mathcal{F}(u_k))^*\}$, where \mathcal{F} denotes the Fourier transform on \mathbb{R}^3 (with its inverse \mathcal{F}^{-1}) and f^* stands for the Schwarz symmetrization of f . Using the property of the Fourier rearrangement from [30], one can derive that

$$\|(-\Delta)^{\frac{3}{4}} u_k^\sharp\|_2 \leq \|(-\Delta)^{\frac{3}{4}} u_k\|_2, \quad \|u_k^\sharp\|_2 = \|u_k\|_2, \quad \|u_k^\sharp\|_q \geq \|u_k\|_q \quad (q > 2).$$

Hence, $\lim_{k \rightarrow \infty} F_\beta(u_k) \leq \lim_{k \rightarrow \infty} F_\beta(u_k^\sharp)$, which implies that (u_k^\sharp) is also the maximizing sequence for μ_β . Constructing a new function sequence (v_k) defined by $v_k(x) := u_k(\|u_k\|_2^{\frac{2}{3}} x)$ for $x \in \mathbb{R}^3$, one can easily verify that (v_k) is also a maximizing sequence for μ_β with $\|(-\Delta)^{\frac{3}{4}} v_k\|_2 = 1$ and $\|v_k\|_2 = 1$. Note (v_k) is bounded in $W^{\frac{3}{2},2}(\mathbb{R}^3)$, up to a sequence, we may assume that

$$v_k \rightharpoonup v \text{ in } W^{\frac{3}{2},2}(\mathbb{R}^3),$$

thus v satisfies that $\|v\|_2 \leq 1$ and $\|(-\Delta)^{\frac{3}{4}} v\|_2^2 \leq 1$. Since $W_r^{\frac{3}{2},2}(\mathbb{R}^3)$ (the collection of all radial functions in $W^{\frac{3}{2},2}(\mathbb{R}^3)$) can be compactly imbedded into $L^p(\mathbb{R}^3)$ for any $p > 2$, we can claim that

$$(3.2) \quad \lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} (\exp(\beta|v_k|^2) - 1 - \beta|v_k|^2) dx = \int_{\mathbb{R}^3} (\exp(\beta|v|^2) - 1 - \beta|v|^2) dx.$$

For simplicity, we define $\Psi(\tau) := \exp(\tau) - 1 - \tau$ and $\Phi(\tau) := \exp(\tau) - 1$ for $\tau > 0$, then we can rewrite (3.2) as

$$(3.3) \quad \lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} \Psi(\beta|u_k|^2) dx = \int_{\mathbb{R}^3} \Psi(\beta|u|^2) dx.$$

Hence, it follows from the mean value theorem and the convexity of the function Ψ that

$$(3.4) \quad \begin{aligned} & |\Psi(\beta|u_k|^2) - \Psi(\beta|u|^2)| \\ & \lesssim \Phi(\theta\beta|u_k|^2 + (1-\theta)\beta|u|^2)(|u| + |u_k|)|u_k - u| \\ & \lesssim (|u_k| + |u|)(\Phi(\beta|u_k|^2) + \Phi(\beta|u|^2))|u_k - u|, \end{aligned}$$

where $\theta \in [0, 1]$. This together with the fractional Adams inequality leads to

$$\begin{aligned}
(3.5) \quad & \left| \int_{\mathbb{R}^3} (\Psi(\beta|u_k|^2) - \Psi(\beta|u|^2)) dx \right| \\
& \lesssim \int_{\mathbb{R}^3} (|u_k| + |u|)(\Phi(\beta|u_k|^2) + \Phi(\beta|u|^2)) |u_k - u| dx \\
& \lesssim \| |u_k| + |u| \|_{L^a(\mathbb{R}^3)} \| \Phi(\beta|u_k|^2) + \Phi(\beta|u|^2) \|_{L^b(\mathbb{R}^n)} \| u_k - u \|_{L^c(\mathbb{R}^n)} \\
& \lesssim \| u_k - u \|_{L^c(\mathbb{R}^3)},
\end{aligned}$$

where the constants $b > 1$ sufficiently close to 1 and $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 1$. Note that $W^{\frac{3}{2},2}(\mathbb{R}^3)$ can be compactly imbedded into $L^r(\mathbb{R}^3)$ for any $r > 2$, we derive that

$$\lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} \Psi(\beta|u_k|^2) dx = \int_{\mathbb{R}^3} \Psi(\beta|u|^2) dx,$$

which accomplishes the proof of (3.2).

Then it follows from (3.2) that

$$\begin{aligned}
(3.6) \quad \mu_\beta &= F_\beta(v_k) + o(1) \\
&= \int_{\mathbb{R}^3} (\exp(\beta|v_k|^2) - 1) dx + o(1) \\
&= \beta + \int_{\mathbb{R}^3} (\exp(\beta|v_k|^2) - 1 - \beta|v_k|^2) dx + o(1) \\
&= \beta + \int_{\mathbb{R}^3} (\exp(\beta|v|^2) - 1 - \beta|v|^2) dx + o(1).
\end{aligned}$$

Next, we show $v \neq 0$. Indeed, one can pick u_0 in $W^{\frac{3}{2},2}(\mathbb{R}^3)$ satisfying $\|(-\Delta)^{\frac{3}{4}} u_0\|_2 = 1$ arbitrarily. Then, we have

$$\begin{aligned}
\mu_\beta \geq F_\beta(u_0) &= \frac{\int_{\mathbb{R}^3} (\exp(\beta|u_0|^2) - 1) dx}{\|u_0\|_2^2} \\
&= \frac{\sum_{j=1}^{\infty} \frac{\beta^j}{j!} \|u_0\|_{2j}^{2j}}{\|u_0\|_2^2} \\
&= \beta + \frac{\sum_{j=2}^{\infty} \frac{\beta^j}{j!} \|u_0\|_{2j}^{2j}}{\|u_0\|_2^2} > \beta.
\end{aligned}$$

Hence,

$$\begin{aligned}
\mu_\beta &\leq \beta + \frac{\int_{\mathbb{R}^3} (\exp(\beta|v|^2) - 1 - \beta|v|^2) dx}{\|v\|_2^2} \\
&= \frac{\int_{\mathbb{R}^3} (\exp(\beta|v|^2) - 1) dx}{\|v\|_2^2} = F_\beta(v).
\end{aligned}$$

Therefore, it remains to show $\|(-\Delta)^{\frac{3}{4}}v\|_2^2 = 1$. Recall that $\|(-\Delta)^{\frac{3}{4}}v\|_2^2 \leq 1$, it suffices to show that $\|(-\Delta)^{\frac{3}{4}}v\|_2^2 \geq 1$. Through the definition of μ_β , one can obtain that

$$\begin{aligned}
\mu_\beta &\geq F_\beta\left(\frac{v}{\|(-\Delta)^{\frac{3}{4}}v\|_2}\right) \\
&= \sum_{j=1}^{\infty} \frac{\beta^j \|v\|_{2j}^{2j}}{j! \|v\|_2^2} \|(-\Delta)^{\frac{3}{4}}v\|_2^{2-2j} \\
(3.7) \quad &\geq \beta + \frac{\beta^2 \|v\|_4^4}{2 \|v\|_2^2} \|(-\Delta)^{\frac{3}{4}}v\|_2^{-2} + \sum_{j=2}^{\infty} \frac{\beta^j \|v\|_{2j}^{2j}}{j! \|v\|_2^2} \\
&= F_\beta(v) + \frac{\beta^2 \|v\|_4^4}{2 \|v\|_2^2} (\|(-\Delta)^{\frac{3}{4}}v\|_2^{-2} - 1) \\
&\geq \mu_\beta + \frac{\beta^2 \|v\|_4^4}{2 \|v\|_2^2} (\|(-\Delta)^{\frac{3}{4}}v\|_2^{-2} - 1)
\end{aligned}$$

which implies that $\|(-\Delta)^{\frac{3}{4}}v\|_2^2 \geq 1$. Thus, v is a maximizer. Next, we prove that all the extremals of the inequality (3.1) must be radially symmetric with respect to some point $x_0 \in \mathbb{R}^3$. Assume that u is a maximizer for the inequality (3.1), we easily see that u^\sharp is also a maximizer for the inequality (3.1) with $\|(-\Delta)^{\frac{3}{4}}u^\sharp\|_2 = \|(-\Delta)^{\frac{3}{4}}u\|_2$ and $\|u^\sharp\|_{L^q} = \|u\|_{L^q}$ for even q . Using the property of the Fourier rearrangement from [30], we conclude that

$$u(x) = e^{i\alpha} u^\sharp(x - x_0) \quad \text{for any } x \in \mathbb{R}^3$$

with some constants $\alpha \in \mathbb{R}$ and $x_0 \in \mathbb{R}^3$. That is to say that u is radially symmetric and real valued up to translation and constant phase. Then we accomplish the proof of Lemma 3.2. \square

Lemma 3.3. ([46]) *Given a function $u \in W^{\frac{3}{2},2}(\mathbb{R}^3)$. If we assume that function $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ satisfying*

$$(-\Delta)^2 U(x, y) = 0$$

on the upper half space \mathbb{R}_+^4 and the boundary condition

$$U(x, 0) = u(x), \quad \partial_y U(x, y)|_{y=0} = 0,$$

then we have the following identity

$$(3.8) \quad \int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy = 2 \int_{\mathbb{R}^3} |(-\Delta)^{\frac{3}{4}}u|^2 dx.$$

Now, we are in the position to prove Theorem 1.4.

Proof of Theorem 1.4: For any $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ satisfying the Neumann boundary condition $\partial_y U(x, y)|_{y=0} = 0$, we define $V(x, y)$ as the bi-harmonic extension of $U(x, 0)$, that is to say $V(x, y) \in W^{2,2}(\overline{\mathbb{R}_+^4})$ satisfies that

$$(-\Delta)^2 V(x, y) = 0$$

on the upper half space \mathbb{R}_+^4 and the boundary condition

$$V(x, y)|_{y=0} = U(x, y)|_{y=0}, \quad \partial_y V(x, y)|_{y=0} = 0.$$

In fact, we can write

$$V(x, y) = c \int_{\partial\mathbb{R}_+^4} \frac{y^3}{(|x - \tilde{x}| + y^2)^3} U(\tilde{x}, 0) d\tilde{x}$$

by the Green formula. Furthermore, by the Dirichlet principle for bi-harmonic function, we derive that

$$(3.9) \quad \int_{\mathbb{R}_+^4} |\Delta V(x, y)|^2 dx dy \leq \int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy.$$

On the other hand, through Lemma 3.3, we get

$$(3.10) \quad \int_{\mathbb{R}_+^4} |\Delta V(x, y)|^2 dx dy = 2 \int_{\partial\mathbb{R}_+^4} |(-\Delta)^{\frac{3}{4}} V(x, 0)|^2 dx.$$

Combining (3.9) and (3.10) and Lemma 3.1, we conclude that

$$(3.11) \quad \begin{aligned} \int_{\partial\mathbb{R}_+^4} (\exp(\beta|U(x, 0)|^2) - 1) dx &= \int_{\partial\mathbb{R}_+^4} (\exp(\beta|V(x, 0)|^2) - 1) dx \\ &\leq C \int_{\partial\mathbb{R}_+^4} |V(x, 0)|^2 dx = C \int_{\partial\mathbb{R}_+^4} |U(x, 0)|^2 dx. \end{aligned}$$

The sharpness of the inequality (1.12) can be similarly deduced from the sharpness of the sharp subcritical inequalities (3.1) as done in the proof of sharpness of inequality (1.8).

Now, we prove the existence of extremals for the subcritical trace Adams inequalities. From Lemma 3.2, we see that there exists $u_0(x) \in W^{\frac{3}{2}, 2}(\mathbb{R}^3)$ such that

$$C = \sup_{u \in W^{\frac{3}{2}, 2}(\mathbb{R}^3), \|(-\Delta)^{\frac{3}{4}} u\|_2 = 1} \frac{\int_{\mathbb{R}^3} (\exp(\beta|u_0|^2) - 1) dx}{\int_{\mathbb{R}^3} |u_0|^2 dx}$$

for any $\beta < 6\pi^2$. Define $U(x, y)$ as the bi-harmonic extension of $u_0(x)$, it is easy to check that $U(x, y)$ is an extremal function for (1.12) through Lemma 3.3.

Finally, we show that the extremals of the inequality (1.12) must be a bi-harmonic extension of some radial function u in $W^{\frac{3}{2}, 2}(\mathbb{R}^3)$. In fact, assume that $U(x, y)$ is the extremal function of the inequality (1.12), It is easy to check that the bi-harmonic extension $V(x, y)$ is also the extremal function of the inequality with the

$$\int_{\mathbb{R}_+^4} |\Delta U|^2 dx dy = \int_{\mathbb{R}_+^4} |\Delta V|^2 dx dy = 1,$$

which implies that $\int_{\mathbb{R}_+^4} |\Delta U|^2 dx dy = \int_{\mathbb{R}_+^4} |\Delta V|^2 dx dy$ if and only if $U = V$ through Dirichlet principle. On the other hand, it is also easy to see that $U(x, 0)$ is the extremal functions of the inequality (3.1). Applying Lemma 3.2 again, we conclude that $U(x, 0)$ must be

radially symmetric with respect to some point $x_0 \in \mathbb{R}^3$. This proves that the extremals of inequality (1.12) must be a bi-harmonic extension of some radial function.

Proof of Theorem 1.5 and Theorem 1.6: For any $U \in W^{2,2}(\overline{\mathbb{R}_+^4})$ satisfying Neumann boundary condition $\partial_y U(x, y)|_{y=0} = 0$ with

$$\int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy + \int_{\partial\mathbb{R}_+^4} |U(x, 0)|^2 dx \leq 1,$$

we derive that

$$(3.12) \quad \int_{\mathbb{R}_+^4} |\Delta V(x, y)|^2 dx dy \leq \int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy,$$

where $V(x, y)$ is the bi-harmonic extension of $U(x, 0)$. This together with Lemma 3.3 and (3.12) yields that

$$\int_{\partial\mathbb{R}_+^4} (2|(-\Delta)^{\frac{3}{4}}V(x, 0)|^2 + |V(x, 0)|^2) dx \leq 1.$$

Combining the critical fractional Adams inequalities in $W^{\frac{3}{2},2}(\mathbb{R}^3)$, we derive that

$$(3.13) \quad \int_{\partial\mathbb{R}_+^4} (\exp(12\pi^2|U(x, 0)|^2) - 1) dx = \int_{\partial\mathbb{R}_+^4} (\exp(12\pi^2|V(x, 0)|^2) - 1) dx \leq C.$$

Similarly, one can deduce from the fractional Adams inequalities with the exact growth in $W^{\frac{2}{3}}(\mathbb{R}^3)$ (see Theorem 1.8) that

$$(3.14) \quad \int_{\partial\mathbb{R}_+^4} \frac{(\exp(12\pi^2|U(x, 0)|^2) - 1)}{1 + |U(x, 0)|^2} dx = \int_{\partial\mathbb{R}_+^4} (\exp(12\pi^2|V(x, 0)|^2) - 1) dx \leq C \int_{\partial\mathbb{R}_+^4} |U(x, 0)|^2 dx,$$

which gives the inequality (1.15).

4. EXISTENCE OF GROUND STATES TO THE BI-LAPLACIAN WITH NONLINEAR NEUMANN BOUNDARY CONDITIONS ON THE HALF SPACE \mathbb{R}_+^4

In this section, we consider the critical point of the functional $I_\lambda(U)$ on E (see (1.22)). For this purpose, we will first study the critical point of the following functional

$$J_\lambda(u) = \frac{1}{2} \left(2 \int_{\mathbb{R}^3} |(-\Delta)^{\frac{3}{4}}u|^2 dx + \int_{\mathbb{R}^3} |u|^2 dx \right) - \frac{\lambda}{24\pi^2} \int_{\mathbb{R}^3} (\exp(12\pi^2|u|^2) - 1) dx,$$

and we show that the functional $J_\lambda(u)$ has a least energy critical point provided $0 < \lambda < 1$, that is,

Proposition 4.1. $m_\lambda = \inf\{J_\lambda(u) : J'_\lambda(u) = 0, u \in W^{\frac{3}{2},2}(\mathbb{R}^3)\}$ can be achieved for any $0 < \lambda < 1$.

Proof of Proposition 4.1: Motivated by the Pohazoev identity for the functional J_λ , we introduce the functional

$$G_\lambda(u) = \|u\|_2^2 - \frac{\lambda}{12\pi^2} \int_{\mathbb{R}^3} (\exp(12\pi^2|u|^2) - 1) dx = (1 - \lambda)\|u\|_2^2 - \int_{\mathbb{R}^3} g_\lambda(u) dx,$$

where $g_\lambda(t)$ is defined as $g_\lambda(t) = \frac{\lambda}{12\pi^2} (\exp(12\pi^2|t|^2) - 1 - 12\pi^2 t^2)$ and the constrained minimization problem

$$(4.1) \quad \begin{aligned} A_\lambda &= \inf \left\{ \|(-\Delta)^{\frac{3}{4}} u\|_2^2 \mid u \in W^{\frac{3}{2},2}(\mathbb{R}^3), G_\lambda(u) = 0 \right\} \\ &= \inf \left\{ J_\lambda(u) \mid u \in W^{\frac{3}{2},2}(\mathbb{R}^3), G_\lambda(u) = 0 \right\} \leq m_\lambda. \end{aligned}$$

Set $M = \{u \in W^{\frac{3}{2},2}(\mathbb{R}^3), G_\lambda(u) = 0\}$, we point out that M is not empty. In fact, let $u_0 \in W^{\frac{3}{2},2}(\mathbb{R}^3)$ be compactly supported and define

$$h(s) := G_\lambda(su_0) = s^2(1 - \lambda)\|u_0\|_2^2 - \int_{\mathbb{R}^3} g_\lambda(su_0) dx, \forall s > 0.$$

From $\lim_{t \rightarrow 0} \frac{g_\lambda(t)}{t^2} = 0$ and $\lim_{t \rightarrow +\infty} \frac{g_\lambda(t)}{t^2} = \infty$, we have $h(s) > 0$ for $s > 0$ small enough and $h(s) < 0$ for $s > 0$ sufficiently large. Therefore, there exists $s_0 > 0$ satisfying $h(s_0) = 0$, which implies $s_0 u_0 \in M$.

Next, we show that A_λ can be attained by some function $u \in W^{\frac{3}{2},2}(\mathbb{R}^3) \setminus \{0\}$. For this, we need the following lemma:

Lemma 4.2. *There exists a radially minimizing sequence $\{u_k\}_k$ satisfying $\|u_k\|_2^2 = 1$ for A_λ .*

Proof. Assume that $\{u_k\}_k$ is a minimizing sequence for A_λ , that is $u_k \in M$ satisfying

$$\lim_{k \rightarrow \infty} \|(-\Delta)^{\frac{3}{4}} u_k\|_2^2 = A_\lambda.$$

Define u_k^\sharp by the Fourier rearrangement of u_k . Using the property of the Fourier rearrangement from [30], one can derive that

$$\|(-\Delta)^{\frac{3}{4}} u_k^\sharp\|_2 \leq \|(-\Delta)^{\frac{3}{4}} u_k\|_2, \quad \|u_k^\sharp\|_2^2 = \|u_k\|_2^2,$$

and

$$\int_{\mathbb{R}^3} (\exp(12\pi^2(u_k^\sharp)^2) - 1) dx \geq \int_{\mathbb{R}^3} (\exp(12\pi^2 u_k^2) - 1) dx.$$

Then it follows that

$$(1 - \lambda)\|u_k^\sharp\|_2^2 = (1 - \lambda)\|u_k\|_2^2 = \int_{\mathbb{R}^3} g_\lambda(u_k) dx \leq \int_{\mathbb{R}^3} g_\lambda(u_k^\sharp) dx.$$

Hence if we set

$$\gamma(t) = (1 - \lambda)\|tu_k^\sharp\|_2^2 - \int_{\mathbb{R}^3} g_\lambda(tu_k^\sharp) dx,$$

then we have $\gamma(1) \leq 0$ while $\gamma(t) > 0$ for $t > 0$ sufficiently small. Therefore, there exists $t_k \in (0, 1]$ such that $\gamma(t_k) = 0$, that is $t_k u_k^\# \in M$. We obtain

$$(4.2) \quad A_\lambda \leq I(t_k u_k^\#) = \|(-\Delta)^{\frac{3}{4}}(t_k u_k^\#)\|_2^2 \leq t_k^2 \|(-\Delta)^{\frac{3}{4}} u_k\|_2^2 \leq I(u_k) = A_\lambda + o(1).$$

This implies that $\{v_k\} = \{t_k u_k^\#\}_k$ is a radial minimizing sequence for A_λ . Let $\tilde{v}_k = u_k^\#(\|v_k\|_2^{\frac{1}{2}} x)$, direct computations yield that $\|\tilde{v}_k\|_2 = 1$, $\tilde{v}_k \in M$ and $\|(-\Delta)^{\frac{3}{4}} v_k\|_2 = \|(-\Delta)^{\frac{3}{4}} \tilde{v}_k\|_2$. \square

In order to study the attainability of A_λ , we also need the following compactness result.

Lemma 4.3 (Compactness). *Suppose that $g : \mathbb{R} \rightarrow [0, +\infty)$ is a continuous function and define*

$$G(u) = \int_{\mathbb{R}^3} g(u) dx.$$

Then for any $K > 0$, if

$$\lim_{t \rightarrow +\infty} \exp\left(-\frac{1}{K}|t|^2\right)g(t) < +\infty, \quad \lim_{t \rightarrow 0} |t|^{-2}g(t) = 0,$$

then for any bounded sequence $\{u_k\}_k \in W_{rad}^{\frac{3}{2},2}(\mathbb{R}^3)$ satisfying $\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^3} |(-\Delta)^{\frac{3}{4}} u_k|^2 dx < 6\pi^2 K$ and weakly converging to some u , we have that $G(u_k) \rightarrow G(u)$.

Proof. Note that $\{u_k\}_k$ is a radial sequence in $W^{\frac{3}{2},2}(\mathbb{R}^3)$, then by radial Lemma, there exists $\delta > 0$ such that

$$(4.3) \quad |u_k(r)| \lesssim \frac{1}{r^\delta}.$$

Hence $u_k(r) \rightarrow 0$ as $r \rightarrow \infty$ uniformly with respect to k . This together with the subcritical Adams inequality (3.1) yields that for any $\varepsilon > 0$, there exists $R > 0$ such that

$$(4.4) \quad \int_{\mathbb{R}^3 \setminus B_R} g(u_k) dx \lesssim \varepsilon \int_{\mathbb{R}^3} |u_k|^2 dx \lesssim \varepsilon, \quad \int_{\mathbb{R}^3 \setminus B_R} g(u) dx \lesssim \varepsilon.$$

On the other hand, since $\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^3} |(-\Delta)^{\frac{3}{4}} u_k|^2 dx < 6\pi^2 K$, then there exists some $\gamma > 0$ such that $\lim_{k \rightarrow +\infty} \int_{\mathbb{R}^3} |(-\Delta)^{\frac{3}{4}} u_k|^2 dx < \gamma < 6\pi^2 K$.

Through $\lim_{t \rightarrow +\infty} \exp\left(-\frac{1}{K}|t|^2\right)g(t) < +\infty$, we derive that for $\varepsilon > 0$, there exists L independent of k such that

$$\int_{|u_k| > L} g(u_k) dx \lesssim \varepsilon \int_{|u_k| > L} \exp\left(\frac{\gamma}{K} \left| \frac{u_k}{\|(-\Delta)^{\frac{3}{4}} u_k\|_2} \right|^2\right) dx$$

and

$$\int_{|u_k| > L} g(u) dx \lesssim \varepsilon \int_{|u_k| > L} \exp\left(\frac{\gamma}{K} \left| \frac{u}{\|(-\Delta)^{\frac{3}{4}} u\|_2} \right|^2\right) dx,$$

Applying the sharp subcritical Adams inequality (3.1), we derive that

$$(4.5) \quad \int_{|u_k|>L} g(u_k)dx \lesssim \varepsilon \int_{\mathbb{R}^3} |u_k|^2 dx \lesssim \varepsilon, \quad \int_{|u_k|>L} g(u)dx \lesssim \varepsilon.$$

Combining (4.4) and (4.5), we conclude that

$$(4.6) \quad \begin{aligned} \lim_{k \rightarrow \infty} |G(u_k) - G(u)| &\leq \int_{\mathbb{R}^3 \setminus B_R} (g(u_k) - g(u))dx + \int_{B_R} (g(u_k) - g(u))dx \\ &\lesssim \varepsilon + \lim_{k \rightarrow \infty} \left| \int_{B_R \cap \{|u_k|>L\}} g(u_k)dx - \int_{B_R \cap \{|u_k|>L\}} g(u)dx \right| \\ &\quad + \lim_{k \rightarrow \infty} \left| \int_{B_R \cap \{|u_k|\leq L\}} g(u_k)dx - \int_{B_R \cap \{|u_k|\leq L\}} g(u)dx \right| \\ &\lesssim \varepsilon + \lim_{k \rightarrow \infty} \left| \int_{B_R \cap \{|u_k|\leq L\}} g(u_k)dx - \int_{B_R \cap \{|u_k|\leq L\}} g(u)dx \right| \\ &\lesssim \varepsilon, \end{aligned}$$

which gives the proof by the Lebesgue dominated convergence theorem. \square

With this compactness result, we can prove the following

Lemma 4.4. *If $A_\lambda < \frac{1}{2}$, then A_λ can be attained and $A_\lambda = I_\lambda(u)$.*

Proof. Let $\{u_k\}_k$ is a radial minimizing sequence for A_λ , that is $u_k \in M$ satisfying

$$\lim_{k \rightarrow \infty} \|(-\Delta)^{\frac{3}{4}} u_k\|_2^2 = A_\lambda \quad \text{and} \quad \|u_k\|_2^2 = 1.$$

We also assume that $u_k \rightharpoonup u$ in $W^{\frac{3}{2},2}(\mathbb{R}^3)$. We first prove that $A_\lambda > 0$. By way of contradiction, we assume that $A_\lambda = 0$. That is $\lim_{k \rightarrow \infty} \|(-\Delta)^{\frac{3}{4}} u_k\|_2^2 = 0$, which implies that $u = 0$. Since

$$\lim_{t \rightarrow +\infty} \exp(-c|t|^2)g_\lambda(t) = 0 \quad \text{for any } c > 12\pi^2, \quad \lim_{t \rightarrow 0} |t|^{-2}g_\lambda(t) = 0.$$

It follows from the Lemma 4.3 that

$$\int_{\mathbb{R}^3} g_\lambda(u_k)dx = \int_{\mathbb{R}^3} g_\lambda(u)dx.$$

On the other hand, it follows from $u_k \in M$ and $\|u_k\|_2^2 = 1$ that

$$0 < (1 - \lambda) \leq \lim_{k \rightarrow \infty} (1 - \lambda)\|u_k\|_2^2 = \lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} g_\lambda(u_k)dx = \int_{\mathbb{R}^3} g_\lambda(u)dx,$$

which contradicts $u = 0$. This proves that $A_\lambda > 0$.

Now are in position to prove that if $A_\lambda < \frac{1}{2}$, then A_λ could be attained. Under the assumption of Lemma 3.2, we have $\lim_{k \rightarrow \infty} \|\Delta u_k\|_2^2 = A_\lambda < \frac{1}{2}$. Set $K = \frac{A_\lambda}{6\pi^2}$, obviously,

$$\lim_{t \rightarrow +\infty} \exp\left(-\frac{1}{K}|t|^2\right)g_\lambda(t) = 0, \quad \lim_{t \rightarrow 0} |t|^{-2}g_\lambda(t) = 0.$$

It follows from Lemma 4.3 that

$$\int_{\mathbb{R}^3} g_\lambda(u_k) dx = \int_{\mathbb{R}^3} g_\lambda(u) dx.$$

Hence

$$(1 - \lambda) = \lim_{k \rightarrow \infty} (1 - \lambda) \|u_k\|_2^2 = \lim_{k \rightarrow \infty} \int_{\mathbb{R}^3} g_\lambda(u_k) dx = \int_{\mathbb{R}^3} g_\lambda(u) dx$$

and

$$\|(-\Delta)^{\frac{3}{4}} u\|_2^2 \leq \lim_{k \rightarrow \infty} \|(-\Delta)^{\frac{3}{4}} u_k\|_2^2 = A_\lambda.$$

In order to show u is minimizer for A_λ , it suffices to show that $G_\lambda(u) = 0$. Since (4.7)

$$G_\lambda(u) = (1 - \lambda) \|u\|_2^2 - \int_{\mathbb{R}^3} g_\lambda(u) dx \leq \lim_{k \rightarrow \infty} (1 - \lambda) \|u_k\|_2^2 - \int_{\mathbb{R}^3} g_\lambda(u_k) dx = \lim_{k \rightarrow \infty} G_\lambda(u_k) = 0.$$

If we define

$$h(t) = G_\lambda(tu) = (1 - \lambda) \|tu\|_2^2 - \int_{\mathbb{R}^3} g(tu) dx,$$

then $h(1) \leq 0$ and from $\lim_{k \rightarrow \infty} g(t) = o(t^2)$, we deduce that $h(t) < 0$ for small $t > 0$. Consequently, there exists $s_0 \in (0, 1]$ such that $G_\lambda(s_0 u) = 0$. Then we have

$$A_\lambda \leq \|(-\Delta)^{\frac{3}{4}} s_0 u\|_2^2 = s_0^2 \|(-\Delta)^{\frac{3}{4}} u\|_2^2 \leq s_0^2 A_\lambda,$$

which proves that $s_0 = 1$ and $\|(-\Delta)^{\frac{3}{4}} u\|_2^2 = A_\lambda$. Then we accomplish the proof. \square

In order to obtain the attainability of A_λ , we introduce the fractional Adams ratio

$$C_{A,\lambda}^L = \sup \left\{ \frac{1}{\|u\|_2^2} \int_{\mathbb{R}^3} g_\lambda(u) dx \mid u \in W^{\frac{3}{2},2}(\mathbb{R}^3), \|(-\Delta)^{\frac{3}{4}} u\|_2^2 \leq L \right\}.$$

The fractional Adams threshold $R(g_\lambda)$ is defined as

$$R(g_\lambda) = \sup \{ L > 0 \mid C_{A,\lambda}^L < +\infty \}.$$

Obviously, according subcritical fractional Adams inequality (3.1), we know $R(g_\lambda) = \frac{1}{2}$ and $C_{A,\lambda}^* := C_{A,\lambda}^{R(g_\lambda)} = \infty$. By using this fact, we can claim that

$$(4.8) \quad A_\lambda < \frac{1}{2}.$$

Indeed, note that $C_{A,\lambda}^* = \infty$, hence there exists $u_0 \in W^{\frac{3}{2},2}(\mathbb{R}^3)$ such that

$$(1 - \lambda) < \frac{1}{\|u_0\|_2^2} \int_{\mathbb{R}^3} g_\lambda(u_0) dx, \quad \|(-\Delta)^{\frac{3}{4}} u_0\|_2^2 \leq \frac{1}{2}.$$

Hence $G_\lambda(u_0) = (1 - \lambda) \|u_0\|_2^2 - \int_{\mathbb{R}^3} g_\lambda(u_0) dx < 0$. Then there exists $s_0 \in (0, 1)$ such that $s_0 u_0 \in M$, which yields that

$$A_\lambda \leq \|(-\Delta)^{\frac{3}{4}} (s_0 u_0)\|_2^2 = s_0^2 \|(-\Delta)^{\frac{3}{4}} u_0\|_2^2 \leq \frac{1}{2} s_0^2 < \frac{1}{2}.$$

Then the claim is finished. Now, we come to

Completion of the proof of Proposition 4.1: Combining Lemmas 4.2, 4.4 and (4.8), we see that A_λ can be attained. Under a suitable change of scale, it is easy to see that m_λ can also be attained.

In view of Proposition 4.1, we can give the proof of Theorem 1.13.

Proof of Theorem 1.13: Let $v(x) \in W^{\frac{3}{2},2}(\mathbb{R}^3)$ be the minimum point of the functional J_λ on the submanifold $F := \{J'_\lambda(u) = 0 \mid u \in W^{\frac{3}{2},2}(\mathbb{R}^3)\}$ and define $V(x, y)$ as the biharmonic extension of $v(x)$, that is to say $V(x, y)$ satisfies that

$$(-\Delta)^2 V(x, y) = 0$$

on the upper half space \mathbb{R}_+^4 and the boundary condition.

$$V(x, 0) = v(x), \quad \partial_y V(x, y)|_{y=0} = 0.$$

We will show that $V(x, y)$ is the ground state of equation (1.21), that is to prove that $V(x, y)$ is the minimum point of the functional I_λ on the submanifold $\tilde{F} := \{I'_\lambda(U) = 0 \mid U \in E\}$.

Assume that $U(x, y)$ is any solution of equation (1.21), we only need to verify that $I_\lambda(V) \leq I_\lambda(U)$. Through equation (1.21), we easily see that $U(x, y)$ is the harmonic extension of $u(x) = U(x, 0)$ and $u(x)$ is the critical point of functional J_λ in $W^{\frac{3}{2},2}(\mathbb{R}^3)$. In view of Lemma 3.3, we derive that

$$(4.9) \quad \int_{\mathbb{R}_+^4} |\Delta V(x, y)|^2 dx dy = 2 \int_{\mathbb{R}^3} |(-\Delta)^{\frac{3}{4}} v(x)|^2 dx, \quad \int_{\mathbb{R}_+^4} |\Delta U(x, y)|^2 dx dy = 2 \int_{\mathbb{R}^3} |(-\Delta)^{\frac{3}{4}} u(x)|^2 dx.$$

Then it follows that $I_\lambda(V) = J_\lambda(v)$ and $I_\lambda(U) = J_\lambda(u)$. This together with v being the least energy critical point of the J_λ on F implies that

$$I_\lambda(V) = J_\lambda(v) \leq J_\lambda(u) = I_\lambda(U),$$

that is to say V is the ground-state solution for equation (1.21) with nonlinear Neumann boundary conditions on the half space.

5. PROOFS OF LEMMA 1.9 AND THEOREMS 1.10 AND 1.11

The main purpose of this section is to establish Lemma 1.9 which is the essential technical ingredient in proving Theorems 1.10 and 1.11 for $m > 2$ by adapting the same argument in establishing Theorems 1.4 and 1.5 in the case of $m = 2$.

We now give the proof of Lemma 1.9.

Proof. It is known that the solution $U(x, y)$ is unique. By using the the generalized Poisson kernel, it has been shown in [55] that the explicit formula of U is given by

$$U(x, y) = \pi^{-m+\frac{1}{2}} \frac{\Gamma(2m-1)}{\Gamma(m-\frac{1}{2})} \int_{\mathbb{R}^{2m-1}} \frac{y^{2m-1}}{(|x-\xi|^2+y^2)^{2m-1}} u(\xi) d\xi.$$

Furthermore, $U(x, y)$ satisfying the following Neumann boundary condition

$$(5.1) \quad \begin{aligned} \Delta^k U(x, y)|_{y=0} &= \frac{\Gamma(m)\Gamma(m - \frac{1}{2} - k)}{\Gamma(m - 1 - k)\Gamma(m - \frac{1}{2})} \Delta_x^k u, & 0 \leq k \leq \left\lfloor \frac{m-1}{2} \right\rfloor; \\ \partial_y \Delta^k U(x, y)|_{y=0} &= 0, & 0 \leq k \leq \left\lfloor \frac{m-2}{2} \right\rfloor; \\ \partial_y \Delta^{m-1} U(x, y)|_{y=0} &= (-1)^{m-1} \frac{\Gamma(m)\Gamma(\frac{1}{2})}{\Gamma(m - \frac{1}{2})} (-\Delta_x)^{m-\frac{1}{2}} u. \end{aligned}$$

By Green's formula and (5.1), we have

$$\begin{aligned} 0 &= \int_{\mathbb{R}_+^{2m}} U \Delta^m U dx dy \\ &= \int_{\mathbb{R}_+^{2m}} \Delta U \Delta^{m-1} U dx dy + \int_{\mathbb{R}^{2m-1}} U \partial_y \Delta^{m-1} U dx - \int_{\mathbb{R}^{2m-1}} \partial_y U \Delta^{m-1} U dx \\ &= \int_{\mathbb{R}_+^{2m}} \Delta U \Delta^{m-1} U dx dy + (-1)^m \frac{\Gamma(m)\Gamma(\frac{1}{2})}{\Gamma(m - \frac{1}{2})} \int_{\mathbb{R}^{2m-1}} u(x) (-\Delta_x)^{m-\frac{1}{2}} u(x) dx \end{aligned}$$

and

$$\int_{\mathbb{R}_+^{2m}} \Delta U \Delta^{m-1} U dx dy = (-1)^{m-2} \int_{\mathbb{R}_+^{2m}} |\nabla^m U|^2 dx dy.$$

Therefore,

$$\int_{\mathbb{R}_+^{2m}} |\nabla^m U|^2 dx dy = \frac{\Gamma(m)\Gamma(\frac{1}{2})}{\Gamma(m - \frac{1}{2})} \int_{\mathbb{R}^{2m-1}} u(x) (-\Delta_x)^{m-\frac{1}{2}} u(x) dx.$$

The desired result follows. \square

6. FRACTIONAL ADAMS INEQUALITIES WITH THE EXACT GROWTH IN $W^{\frac{n}{2},2}(\mathbb{R}^n)$

In this section, we shall prove the fractional Adams inequalities with the exact growth in $W^{\frac{n}{2},2}(\mathbb{R}^n)$ for all $n \geq 2$. For this purpose, we need the following lemmas.

Lemma 6.1. [7] *For any radial function $u \in W^{\frac{n}{2},2}(\mathbb{R}^n)$, one has*

$$\int_{\mathbb{R}^n} |(-\Delta)^{\frac{n}{4}} u|^2 dx \geq 2^{n-2} \Gamma^2\left(\frac{n}{2}\right) \int_{\mathbb{R}^n} \frac{|\nabla u|^2}{|x|^{n-2}} dx.$$

Lemma 6.2. *For any $u \in W^{\frac{n}{2},2}(\mathbb{R}^n)$, one has*

$$\int_{\mathbb{R}^{2m}} \frac{e^{\beta(n, \frac{n}{2})|u|^2} - 1}{1 + |u|^2} dx \leq 2 \left(e^{\beta(n, \frac{n}{2})} - 1 - 2\beta(n, \frac{n}{2}) \right) \|u\|_2^2 + 2 \int_{\mathbb{R}^{2m}} \frac{e^{\beta(n, \frac{n}{2})|u^\#|^2} - 1}{1 + |u^\#|^2} dx,$$

where $u^\# = \mathcal{F}^{-1}\{(\mathcal{F}u)^*\}$ denotes the Fourier rearrangement of u .

Proof. This lemma can be similarly proved as Lemma 2.1 in [47]. We omit the details. \square

With these two lemmas, we can give

Proof of Theorem 1.8. Assume $u \in W^{\frac{n}{2},2}(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} |(-\Delta)^{\frac{n}{4}} u(x)|^2 dx = 1$, then by the Fourier rearrangement inequality [30], there exists some radial function $u^\# \in W^{\frac{n}{2},2}(\mathbb{R}^n)$ such that

$$\int_{\mathbb{R}^n} |(-\Delta)^{\frac{n}{4}} u^\#|^2 dx \leq 1.$$

Combining this and Lemma 6.2, we can assume u is a radial function satisfying

$$\int_{\mathbb{R}^n} |(-\Delta)^{\frac{n}{4}} u|^2 dx = 1$$

and $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$.

In view of Lemma 6.1, we have

$$\begin{aligned} \int_{\mathbb{R}^n} |(-\Delta)^{\frac{n}{4}} u|^2 dx &\geq 2^{n-2} \Gamma^2\left(\frac{n}{2}\right) \int_{\mathbb{R}^n} \frac{|\nabla u|^2}{|x|^{n-2}} dx \\ &= 2^{n-2} \Gamma^2\left(\frac{n}{2}\right) \frac{2^{\frac{n+2}{2}} \pi^{\frac{n-1}{2}}}{(n-2)!!} \int_0^\infty |u'|^2 r dr \\ &= 2\pi \cdot \frac{2^{n-3} \Gamma^2\left(\frac{n}{2}\right) 2^{\frac{n+2}{2}} \pi^{\frac{n-1}{2}}}{\pi (n-2)!!} \int_0^\infty |u'|^2 r dr \\ &= 2\pi \left(A_n \int_0^\infty |u'|^2 r dr \right), \end{aligned}$$

where

$$A_n = \frac{2^{n-3} \Gamma^2\left(\frac{n}{2}\right) 2^{\frac{n+2}{2}} \pi^{\frac{n-1}{2}}}{\pi (n-2)!!}.$$

Denote $w(s) = \left(\frac{n}{2} A_n\right)^{1/2} u(s^{2/n})$, that is

$$u(r) = \left(\frac{n}{2} A_n\right)^{-1/2} w(r^{n/2}).$$

Then direct calculation gives

$$\begin{aligned} \int_0^\infty |w'(s)|^2 s ds &= \int_0^\infty \left| \left(\frac{n}{2} A_n\right)^{1/2} u'(s^{2/n}) \frac{2}{n} s^{\frac{2-n}{n}} \right|^2 s ds \\ &= A_n \int_0^\infty |u'(s^{2/n})|^2 s^{2/n} ds^{2/n} \\ &= A_n \int_0^\infty |u'(r)|^2 r dr. \end{aligned}$$

Therefore, it follows that

$$1 \geq 2\pi \left(A_n \int_0^\infty |u'|^2 r dr \right) = 2\pi \left(\int_0^\infty |w'(s)|^2 s ds \right)$$

and

$$\begin{aligned} \int_0^\infty |w(s)|^2 s ds &= \int_0^\infty \left| \left(\frac{n}{2} A_n \right)^{1/2} u(s^{2/n}) \right|^2 s ds \\ &= \frac{n A_n}{2} \int_0^\infty |u(r)|^2 r^{n/2} dr^{n/2} \\ &= \frac{n^2 A_n}{4} \int_0^\infty |u(r)|^2 r^{n-1} dr. \end{aligned}$$

Gathering the above estimate, we conclude that

$$\begin{aligned} \int_{\mathbb{R}^n} \frac{\exp(\beta(n, \frac{n}{2})u^2) - 1}{1 + |u|^2} dx &= \omega_{n-1} \int_0^\infty \frac{\exp\left(\beta(n, \frac{n}{2})\left(\left(\frac{n}{2}A_n\right)^{-1/2}w\left(r^{\frac{n}{2}}\right)\right)^2\right) - 1}{1 + \left|\left(\frac{n}{2}A_n\right)^{-1/2}w\left(r^{\frac{n}{2}}\right)\right|^2} r^{n-1} dr \\ &= \frac{\omega_{n-1}}{n} \int_0^\infty \frac{\exp\left(\frac{2\beta(n, \frac{n}{2})}{nA_n}w^2\left(r^{\frac{n}{2}}\right)\right) - 1}{1 + \left|\left(\frac{n}{2}A_n\right)^{-1/2}w\left(r^{\frac{n}{2}}\right)\right|^2} dr^n \\ &= \frac{\omega_{n-1}}{n} \int_0^\infty \frac{\exp(4\pi w^2(r^{\frac{n}{2}})) - 1}{1 + \left(\frac{n}{2}A_n\right)^{-1} |w\left(r^{\frac{n}{2}}\right)|^2} dr^n \\ &\leq A_n \omega_{n-1} \int_0^\infty \frac{\exp(4\pi w^2(s)) - 1}{1 + |w(s)|^2} s ds \\ &\leq c \int_{\mathbb{R}^2} w^2(|x|) dx \leq c \int_{\mathbb{R}^n} u^2 dx, \end{aligned}$$

which accomplishes the proof of (1.15).

Now, we prove the sharpness of $p \geq 2$. We adopt the test function sequence used in [21]:

$$\phi_{\varepsilon, r}(x) = \begin{cases} c_{n/2} |x|^{-n/2}, & \varepsilon r < |x| < r \\ 0, & \text{otherwise,} \end{cases}$$

where $c_{n/2} = \frac{1}{2^{n/2} \pi^{n/2}}$ and define the function on B_r by :

$$\widetilde{\phi}_{\varepsilon, r} = \phi_{\varepsilon, r}(x) - P_{n-1}^r \phi_\varepsilon,$$

where $P_{n-1}^r \phi_\varepsilon$ is the projection on the space of polynomials of degree up to $n-1$ on the ball B_r . Obviously, $\widetilde{\phi}_{\varepsilon, r}$ is orthogonal to every polynomial of order up to $n-1$ on the ball B_r and satisfies $|P_{n-1}^r \phi_\varepsilon| \leq C r^{-\frac{n}{2}}$.

By (100), (104) and (107) in [21], we have

$$\left\| \widetilde{\phi_{\varepsilon,r}} \right\|_2^2 \leq -\frac{\log(\varepsilon r)^n}{\beta(n, \frac{n}{2})} + b_r + C,$$

$$(6.1) \quad \left\| I_{n/2} \widetilde{\phi_{\varepsilon,r}} \right\|_2^2 \leq Cr^n$$

and

$$I_{n/2} \widetilde{\phi_{\varepsilon,r}}(x) \geq -\frac{\log(\varepsilon r)^n}{\beta(n, \frac{n}{2})} + b_r - C$$

on the ball $B_{\varepsilon r/2}$, where $b_r \leq C \log r$.

On the other hand, since

$$\begin{aligned} \left| \widetilde{\phi_{\varepsilon,r}} \right|^2 &= |\phi_{\varepsilon,r}(x) - P_{n-1}^r \phi_\varepsilon|^2 \\ &\geq C_1 |\phi_{\varepsilon,r}(x)|^2 - C_2 |P_{n-1}^r \phi_\varepsilon|^2 \\ &\geq C_1 |\phi_{\varepsilon,r}(x)|^2 - C_2 r^{-n/2}, \end{aligned}$$

and thus it follows that

$$(6.2) \quad \left\| \widetilde{\phi_{\varepsilon,r}} \right\|_2^2 \geq -C_1 \log(\varepsilon r)^n + C_2.$$

Set $\psi_{\varepsilon,r} = \frac{I_{n/2} \widetilde{\phi_{\varepsilon,r}}}{\left\| \widetilde{\phi_{\varepsilon,r}} \right\|_2}$, then on the ball $B_{\varepsilon r/2}$,

$$\begin{aligned} (\psi_{\varepsilon,r}(x))^2 &\geq \frac{\left(-\frac{\log(\varepsilon r)^n}{\beta(n, \frac{n}{2})} + b_r - C \right)^2}{-\frac{\log(\varepsilon r)^n}{\beta(n, \frac{n}{2})} + b_r + C} \\ &\geq -\frac{\log(\varepsilon r)^n}{\beta(n, \frac{n}{2})} + b_r \left(1 - \frac{C}{\log \frac{1}{\varepsilon^n}} \right) - C, \end{aligned}$$

which gives

$$\begin{aligned}
& \sup_{\|(-\Delta)^{n/4}u\|_2 \leq 1} \int_{\mathbb{R}^n} \frac{\exp\left(\beta\left(n, \frac{n}{2}\right)u^2\right) - 1}{(1+u)^p} dx \\
& \geq \int_{B_{r\varepsilon/2}} \frac{\exp\left(\beta\left(n, \frac{n}{2}\right)(\psi_{\varepsilon,r}(x))^2\right)}{(1+\psi_{\varepsilon,r}(x))^p} dx \\
& \geq \int_{B_{r\varepsilon/2}} \frac{\exp\left(\beta\left(n, \frac{n}{2}\right)\left(-\frac{\log(\varepsilon r)^n}{\beta\left(n, \frac{n}{2}\right)} + b_r\left(1 - \frac{C}{\log\frac{1}{\varepsilon^n}}\right) - C\right)\right)}{\left(-\frac{\log(\varepsilon r)^n}{\beta\left(n, \frac{n}{2}\right)} + b_r\left(1 - \frac{C}{\log\frac{1}{\varepsilon^n}}\right) - C\right)^{\frac{p}{2}}} dx \\
& \geq \frac{C}{\left(\log\frac{1}{(\varepsilon r)^n}\right)^{\frac{p}{2}}}.
\end{aligned}$$

On the other hand, from (6.1) and (6.2), we see that

$$\int_{\mathbb{R}^n} (\psi_{\varepsilon,r}(x))^2 dx = \frac{\|I_{n/2}\widetilde{\phi}_{\varepsilon,r}\|_2^2}{\|\widetilde{\phi}_{\varepsilon,r}\|_2^2} \geq \frac{Cr^n}{-\log(\varepsilon r)^n + C}.$$

Combining the above estimate, we conclude that if $p < 2$,

$$\begin{aligned}
\sup_{\|(-\Delta)^{n/4}u\|_2 \leq 1} \frac{1}{\|u\|_2^2} \int_{\mathbb{R}^n} \frac{\exp(\beta u^2) - 1}{(1+u)^p} dx & \geq \frac{1}{\|\psi_{\varepsilon,r}(x)\|_2^2} \int_{B_{r\varepsilon/2}} \frac{\exp(\beta(\psi_{\varepsilon,r}(x))^2)}{(1+\psi_{\varepsilon,r}(x))^p} dx \\
& \geq C \frac{\log\frac{1}{(\varepsilon r)^n}}{r^n \left(\log\frac{1}{(\varepsilon r)^n}\right)^{\frac{p}{2}}} \\
& = \frac{C}{r^n} \left(\log\frac{1}{(\varepsilon r)^n}\right)^{1-\frac{p}{2}} \rightarrow \infty
\end{aligned}$$

as $\varepsilon \rightarrow 0$, which accomplishes the proof of Theorem 1.8. □

REFERENCES

- [1] A. G. Ache, S.-Y. A. Chang, *Sobolev trace inequalities of order four*, Duke Math. J., **166** (2017), 2719-2748.
- [2] S. Adachi, K. Tanaka, *Trudinger type inequalities in \mathbb{R}^N and their best exponents*, Proc. Amer. Math. Soc., **128** (2000), 2051-2057.
- [3] D. Adams, *Traces of potentials*, Indiana Univ. Math. J., **22** (1973) 907-918.

- [4] D. Adams, *A sharp inequality of J. Moser for higher order derivatives*, Ann. of Math. **128** (1998), 383-398.
- [5] Adimurthi, S. L. Yadava, *Critical exponent problem in \mathbb{R}^2 with Neumann boundary condition*, Commun. Partial Differ. Equ., **15** (1990), 461-501.
- [6] W. Beckner, *Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality*, Ann. of Math., **138** (1993) 213-242.
- [7] W. Beckner, *Weighted inequalities and Stein-Weiss potentials*, Forum Math., **20** (2008), 587-606.
- [8] D. Cao, *Nontrivial solution of semilinear elliptic equation with critical exponent in \mathbb{R}^2* , Comm. Partial Differential Equations **17** (1992), 407-435.
- [9] L. Carleson and S. Y. A. Chang, *On the existence of an extremal function for an inequality of J. Moser*. Bull. Sci. Math. (2) 110 (1986), no. 2, 113-127.
- [10] J. Case, *Sharp weighted Sobolev trace inequalities and fractional powers of the Laplacian*. J. Funct. Anal. 279 (2020), no. 4, 108567, 33 pp.
- [11] S. Y. A. Chang and D. Marshall, *On a sharp inequality concerning the Dirichlet integral*. Amer. J. Math. 107 (1985), no. 5, 1015-1033.
- [12] L. Chen, G. Lu and C. Zhang, *Sharp weighted Trudinger-Moser-Adams inequalities on the whole space and the existence of their extremals*. Calc. Var. Partial Differential Equations 58 (2019), no. 4, Paper No. 132, 31 pp.
- [13] L. Chen, G. Lu and C. Zhang, *Maximizers for fractional Caffarelli-Kohn-Nirenberg and Trudinger-Moser inequalities on the fractional Sobolev spaces*, J. Geom. Anal. 31 (2021), no. 4, 3556-3582.
- [14] L. Chen, G. Lu and M. Zhu, *Existence and nonexistence of extremals for critical Adams inequalities in \mathbb{R}^4 and Trudinger-Moser inequalities in \mathbb{R}^2* . Adv. Math. 368 (2020), 107143, 61 pp.
- [15] P. Cherrier, *Problèmes de Neumann nonlinéaires sur les variétés riemanniennes*, C. R. Acad. Sci. ParisSr. I Math., **292**, 637-640 (in French).
- [16] A. Cianchi, *Moser-Trudinger trace inequalities*, Adv. Math., **217** (2008), 2005-2044.
- [17] W. S. Cohn, G. Lu, *Best constants for Moser-Trudinger inequalities on the Heisenberg group*, Indiana Univ. Math. J., **50** (2001) 1567-1591.
- [18] J. M. do Ó, *N -Laplacian equations in \mathbb{R}^N with critical growth*, Abstr. Appl. Anal., **2** (1997), 301-315.
- [19] J. Escobar, *Sharp constant in a Sobolev trace inequality*, Indiana Univ. Math. J., **37** (1988), 687-698.
- [20] M. Flucher, *Extremal functions for the Trudinger-Moser inequality in 2 dimensions*, Comment. Math. Helv., **67** (1992), 471-497.
- [21] L. Fontana, C. Morpurgo, *Sharp exponential integrability for critical Riesz potentials and fractional Laplacians on \mathbb{R}^n* , Nonlinear Anal. **167** (2018), 85-122.
- [22] S. Ibrahim, N. Masmoudi and K. Nakanishi, *Trudinger-Moser inequality on the whole plane with the exact growth condition*, J. Eur. Math. Soc., **17** (2015), 819-835.
- [23] S. Iula, A. Maalaoui and L. Martinazzi, *Critical points of a fractional Moser-Trudinger embedding in dimension 1*, Differ. Integr. Equ., **29** (2016), 455-492.
- [24] N. Lam, G. Lu, *Sharp Moser-Trudinger inequality on the Heisenberg group at the critical case and applications*, Adv. Math., **231** (2012), 3259-3287.
- [25] N. Lam, G. Lu, *A new approach to sharp Moser-Trudinger and Adams type inequalities: a rearrangement-free argument*, J. Differential Equations, **255** (2013), 298-325.
- [26] N. Lam, G. Lu, *Sharp singular Trudinger-Moser-Adams type inequalities with exact growth*. Geometric methods in PDE's, 43-80, Springer INdAM Ser., 13, Springer, Cham, 2015.
- [27] N. Lam, G. Lu and H. Tang, *Sharp subcritical Moser-Trudinger inequalities on Heisenberg groups and subelliptic PDEs*. Nonlinear Anal. 95 (2014), 77-92.
- [28] N. Lam, G. Lu and L. Zhang, *Equivalence of critical and subcritical sharp Trudinger-Moser-Adams inequalities*, Rev. Mat. Iberoam., **33** (2017), 1219-1246.
- [29] N. Lam, G. Lu and L. Zhang, *Sharp singular Trudinger-Moser inequalities under different norms*. Adv. Nonlinear Stud. 19 (2019), no. 2, 239-261.

- [30] E. Lenzmann, J. Sok, *A sharp rearrangement principle in Fourier space and symmetry results for PDEs with arbitrary order*, arXiv:1805.06294v1.
- [31] J. Li and G. Lu, *Critical and subcritical Trudinger-Moser inequalities on complete noncompact Riemannian manifolds*. Adv. Math. 389 (2021), Paper No. 107915.
- [32] J. Li and G. Lu, *The Chang-Marshall Trace Inequality for Sobolev functions in domains in higher dimensional space \mathbb{R}^n* , Preprint.
- [33] J. Li, G. Lu and M. Zhu, *Concentration-compactness principle for Trudinger-Moser inequalities on Heisenberg groups and existence of ground state solutions*, Calc. Var. Partial Differential Equations, **57** (2018), 57-84.
- [34] Y. X. Li, P. Liu, *A Moser-Trudinger inequality on the boundary of a compact Riemann surface*, Math.Z., **250**, (2005) 363-386.
- [35] Y. X. Li and B. Ruf, *A sharp Moser-Trudinger type inequality for unbounded domains in \mathbb{R}^n* , Indiana Univ. Math. J. **57** (2008), 451-480.
- [36] K. C. Lin, *Extremal functions for Moser's inequality*, Trans. Amer. Math. Soc., **348** (1996), 2663-2671.
- [37] G. Lu, H. Tang and M. Zhu, *Best constants for Adams' inequalities with the exact growth condition in R^n* , Adv. Nonlinear Stud., **15** (2015), 763-788.
- [38] A. Malchiodi, L. Martinazzi, *Critical points of the Moser-Trudinger functional on a disk*. (English summary) J. Eur. Math. Soc. (JEMS) 16 (2014), no. 5, 893-908.
- [39] G. Mancini, L. Martinazzi, *Extremals for fractional Moser-Trudinger inequalities in dimension 1 via harmonic extensions and commutator estimates*, Adv. Nonlinear Stud., **20** (2020), 599-632.
- [40] G. Mancini, L. Martinazzi, *The Moser-Trudinger inequality and its extremals on a disk via energy estimates*. Calc. Var. Partial Differential Equations 56 (2017), no. 4, Paper No. 94, 26 pp.
- [41] N. Masmoudi, F. Sani, *Adams' inequality with the exact growth condition in \mathbb{R}^4* , Comm. Pure Appl. Math., **67** (2014), 1307-1335.
- [42] N. Masmoudi, F. Sani, *Trudinger-Moser inequalities with the exact growth condition in RN and applications*, Comm. Partial Differential Equations, **40** (2015), 1408-1440.
- [43] N. Masmoudi, F. Sani, *Higher order Adams' inequality with the exact growth condition*, Commun. Contemp. Math., **20** (2018), Doi:10.1142/S0219199717500729.
- [44] V. Mazya, *Sobolev Spaces*, Springer Ser. Sov. Math., Springer-Verlag, Berlin, 1985, translated from Russian by T. O. Shaposhnikova.
- [45] J. Moser, *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. J., **20** (1970), 1077-1092.
- [46] Q. A. Ngô, V. H. Nguyen and Q. H. Pham, *Higher order Sobolev trace inequalities on balls revisited*, J. Funct. Anal., **278**, (2020), 52 pp.
- [47] V. H. Nguyen, *a simple proof of Adams type inequalities in \mathbb{R}^{2m}* , Discrete Contin. Dyn. Syst., **40**, (2020), 5755-5764.
- [48] R. O'Neil. *Convolution operators and $L(p, q)$ spaces*. Duke Math. J., 30:129-142, 1963.
- [49] S. I. Pohozaev, *On the Sobolev embedding theorem for $pl = n$* , in: Doklady Conference, Moscow Power Institut, Moscow (1965), 158-170.
- [50] W. Rudin, *Lectures on the edge-of-the-wedge theorem.*, Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 6. (1971), Published by the American Mathematical Society.
- [51] B. Ruf, *A sharp Moser-Trudinger type inequality for unbounded domains in \mathbb{R}^2* , J. Funct. Anal., **219** (2005), 340-367.
- [52] D. Siegel, E. O. Talvila, *Uniqueness for the n -dimensional half space Dirichlet problem*, Pacific J. Math., **175** (1996), no. 2, 571-587.
- [53] N. S. Trudinger, *On imbeddings into Orlicz spaces and some applications*, J. Math. Mech., **17** (1967), 473-483.

- [54] C. Zhang, L. Chen, *Concentration-compactness principle of singular Trudinger-Moser inequalities in \mathbb{R}^n and n -Laplace equations*, Adv. Nonlinear Stud., **18** (2018), 567-585.
- [55] Q. Yang, *Sharp Sobolev trace inequalities for higher order derivatives*, arXiv:1901.03945.
- [56] R. Yang, *On higher order extensions for the fractional Laplacian*, arXiv:1302.4413.
- [57] V. I. Yudovic, *Some estimates connected with integral operators and with solutions of elliptic equations*, Dokl. Akad. Nauk SSSR 138 (1961), 805-808.

SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081,
P. R. CHINA

Email address: chenlu5818804@163.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269, USA

Email address: guozhen.lu@uconn.edu

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN, 430072, P. R. CHINA

Email address: qhyang.math@gmail.com; qhyang.math@whu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, INSTITUTE OF APPLIED SYSTEM ANALYSIS, JIANGSU UNIVERSITY, ZHENJIANG, 212013, P. R. CHINA

Email address: zhumaochun2006@126.com