

# The $q$ -golden ratio, Catalan numbers, and an identity of Sauermann–Wigderson

Kevin Carde

2021-04-07

## Abstract

In this note, we present some basic properties of  $q$ -Fibonacci numbers and their relationship to the  $q$ -golden ratio and Catalan numbers. We then use this relationship to give a short proof of a combinatorial identity.

## 1 $q$ -Fibonacci numbers

**Definition 1.** *The  $q$ -Fibonacci numbers are defined recursively by  $F_0(q) = F_1(q) = 1$ ,  $F_n(q) = F_{n-1}(q) + qF_{n-2}(q)$ .*

Note that the original Fibonacci numbers are retrieved by setting  $q = 1$ . These  $q$ -Fibonacci numbers have a nice non-recursive form.

**Proposition 2.**

$$F_n(q) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} q^k.$$

*Proof.* We carry out a quick proof by induction. Clearly this holds for  $F_0(q) = F_1(q) = 1$ . Assuming the statement for  $n-1$  and  $n-2$ , we have

$$\begin{aligned} F_n(q) &= F_{n-1}(q) + qF_{n-2}(q) = \sum_k \left( \binom{n-1-k}{k} + \binom{n-2-(k-1)}{k-1} \right) q^k \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} q^k \end{aligned}$$

as desired. □

From this, by plugging in  $q = 1$ , we immediately have an identity relating the standard Fibonacci numbers to binomial coefficients:

$$F_n = \binom{n}{0} + \binom{n-1}{1} + \cdots + \binom{\lfloor \frac{n}{2} \rfloor}{\lfloor \frac{n}{2} \rfloor}.$$

## 2 The $q$ -golden ratio

It is well known that the the ratio of successive Fibonacci numbers approaches the golden ratio  $\varphi = \frac{1+\sqrt{5}}{2}$ . We prove the following  $q$ -analog:

**Theorem 3.** *For  $n \geq 0$ , we have as a power series around  $q = 0$*

$$\frac{F_{n+1}(q)}{F_n(q)} = \frac{1 + \sqrt{1+4q}}{2} + O(q^{n+1}).$$

*In particular, as  $n \rightarrow \infty$ , the ratio approaches  $\varphi(q) = \frac{1+\sqrt{1+4q}}{2}$ , which itself equals the golden ratio  $\varphi$  when  $q = 1$ .*

*Proof.* Before we begin, note that the theorem is equivalent (by taking reciprocals) to

$$\frac{F_n(q)}{F_{n+1}(q)} = -\frac{1 - \sqrt{1+4q}}{2q} + O(q^{n+1}).$$

The theorem itself will be a straightforward proof by induction on  $n$ . The base case of  $n = 0$  is immediate. We will now show that if the statement is true for  $n - 1$ , it is also true for  $n$ .

By the defining recurrence for the  $q$ -Fibonacci, we have

$$\frac{F_{n+1}(q)}{F_n(q)} = \frac{F_n(q) + qF_{n-1}(q)}{F_n(q)} = 1 + q\frac{F_{n-1}(q)}{F_n(q)}.$$

By the inductive hypothesis and the equivalent reciprocal formulation, this is equal to

$$1 + q\left(-\frac{1 - \sqrt{1+4q}}{2q} + O(q^n)\right) = \frac{1 + \sqrt{1+4q}}{2} + O(q^{n+1}),$$

as desired. □

The formula appearing in this theorem requires no guessing to discover. Ignoring the error term, the recurrence for the  $q$ -Fibonacci tells us that the  $q$ -golden ratio  $\varphi(q)$  should satisfy  $\varphi(q) = 1 + \frac{q}{\varphi(q)}$  or  $\varphi(q)^2 - \varphi(q) - q = 0$ , the  $q$ -analog of the standard quadratic satisfied by  $\varphi$ . By the quadratic formula and appropriate choice of signs, we immediately get  $\varphi(q) = \frac{1+\sqrt{1+4q}}{2}$ . This all parallels how one might discover and prove that  $F_{n+1}/F_n \rightarrow \varphi$  for the standard Fibonacci numbers.

## 3 Catalan numbers and the Sauermaann–Wigderson identity

In the proof, we took the reciprocal

$$\frac{F_n(q)}{F_{n+1}(q)} = -\frac{1 - \sqrt{1+4q}}{2q} + O(q^{n+1}).$$

Ignoring the error term, the right hand side is, upon replacing  $q$  by  $-q$ , the generating function for the Catalan numbers.

We therefore have the following:

**Corollary 4.** *The coefficients on  $1, q, q^2, \dots, q^n$  of  $F_n(q)/F_{n+1}(q)$  are, up to sign, the Catalan numbers:  $1, -1, 2, -5, 14, \dots, (-1)^n C_n$ .*

As a further corollary, we get a stronger version of an identity needed by Sauermann–Wigderson in [1]. (This strengthening was noted and proved differently in [2].)

**Corollary 5.** *Let  $m \leq n$ . Then*

$$\sum_{(m_1, \dots, m_t)} (-1)^t \binom{n-m_1}{m_1-1} \binom{n-m_2}{m_2} \cdots \binom{n-m_t}{m_t} = (-1)^m C_{m-1},$$

where the sum is over all tuples (of varying length) of positive integers summing to  $m$ .

*Proof.* Note that

$$\sum_k \binom{n-k}{k-1} q^k = \sum_k \binom{n-1-(k-1)}{k-1} q^k = q F_{n-1}(q).$$

The other binomial coefficients appearing in the original identity are of the form  $\binom{n-k}{k}$ , which are directly the coefficients in  $F_n(q) - 1$ . (The requirement of positive  $k$  means we need to subtract off the constant term.)

The desired quantity, therefore, is the  $q^m$  coefficient in

$$-q F_{n-1}(q) \left( 1 - (F_n(q) - 1) + (F_n(q) - 1)^2 - \dots \right) = -q \frac{F_{n-1}(q)}{F_n(q)}.$$

Because of the  $-q$  out front, we are seeking the negative of the  $q^{m-1}$  coefficient in  $\frac{F_{n-1}(q)}{F_n(q)}$ . By the  $q$ -golden ratio Catalan corollary, our final result is  $-(-1)^{m-1} C_{m-1} = (-1)^m C_{m-1}$ , as desired. □

## References

- [1] L. Sauermann and Y. Wigderson, *Polynomials that vanish to high order on most of the hypercube*, arXiv:2010.00077 [math.CO] (2020).
- [2] D. Zeilberger, *Two Quick Proofs of a Catalan Lemma Needed by Lisa Sauermann and Yuval Wigderson*, Personal Journal of Shalosh B. Ekhad and Doron Zeilberger (2020).