

A Combinatorial Interpretation for Sequence A345973 in OEIS

David Callan

August 10, 2021

Abstract

We give a combinatorial interpretation in terms of bicolored ordered trees for the sequence $(a_n)_{n \geq 1} = (1, 1, 1, 2, 3, 6, 10, 20, 36, 73, \dots)$, A345973 in OEIS, whose generating function satisfies the defining identity $\sum_{n \geq 1} a_n x^n = x + \frac{x^2}{\prod_{n \geq 1} (1 - a_n x^n)}$.

Our purpose is to give a combinatorial interpretation of the sequence $(a_n)_{n \geq 1} = (1, 1, 1, 2, 3, 6, 10, 20, 36, 73, \dots)$, A345973 in OEIS [1], with defining identity $\sum_{n \geq 1} a_n x^n = x + \frac{x^2}{\prod_{n \geq 1} (1 - a_n x^n)}$. Our counting sequence begins $(0, 1, 1, 2, \dots)$, differing only in the first term. All our trees have edges colored black or gray and size measured by number of black edges. Consider the set \mathcal{A} of such bicolored trees, unlabeled but rooted with ordered edges, that satisfy the following 4 conditions: For each nonleaf vertex v ,

1. v has at least two black child edges,
2. the gray child edges of v (if any) are right justified, that is, they lie to the right of all black child edges of v ,
3. the subtrees of v taken left to right have weakly decreasing sizes,

and for each leaf,

4. its parent edge (if it has one) is black.

The tree on the left in Figure 1 below is certainly not in \mathcal{A} because it violates each of the first three conditions with v taken as the root and it also violates the fourth condition due to the leaf terminating the gray edge, while the other two trees are in \mathcal{A} . Let \mathcal{A}_n denote the set of trees in \mathcal{A} of size n .




Figure 1

Clearly, \mathcal{A}_0 consists of the singleton tree (a root with no edges) and \mathcal{A}_1 is empty due to condition 1. So $|\mathcal{A}_0| = 1$ and $|\mathcal{A}_1| = 0$. Set $a_n = |\mathcal{A}_n|$ for $n \geq 2$ (leaving a_1 undefined for now).

Theorem. For $n \geq 2$, $a_n = A345973(n)$.

Proof. Suppose $T \in \mathcal{A}_n$ with $n \geq 2$. Say there are k black edges from the root. They contribute k to the size of T and $k \geq 2$ by condition 1. The subtrees of the root are all in \mathcal{A} since they inherit the 4 defining conditions. If there are no gray edges from the root, then either all subtrees of the root are singletons or the singleton subtrees (if any) are right justified by condition 3 and the non-singleton subtrees, taken left to right, have sizes (weakly decreasing by condition 3) $n_1 \geq n_2 \geq \dots \geq n_j$ summing to $n - k$, for some j with $1 \leq j \leq k$ (j depending on the number of singleton subtrees), and $n_j \geq 2$ by condition 1. If gray edges from the root are present, then the root has no leaf child by conditions 2, 3, and 4, the gray edges are right justified, and the list of sizes of all subtrees of the root forms a partition $n_1 \geq n_2 \geq \dots \geq n_j \geq 2$ of $n - k$ with $j > k$. Putting all this together, we obtain the recurrence for $n \geq 2$,

$$a_n = 1 + \sum_{k=2}^{n-2} \sum_{(n_1, n_2, \dots, n_j) \in \mathcal{P}_{n-k}^*} a_{n_1} a_{n_2} \dots a_{n_j}, \quad (1)$$

where \mathcal{P}_r^* is the set $\{(n_1, n_2, \dots, n_j) : j \geq 1, n_1 \geq n_2 \geq \dots \geq n_j \geq 2, n_1 + n_2 + \dots + n_j = r\}$ of partitions of r with all parts ≥ 2 . Replacing n by $n + 2$, (1) becomes

$$a_{n+2} = 1 + \sum_{k=2}^n \sum_{(n_1, n_2, \dots, n_j) \in \mathcal{P}_{n+2-k}^*} a_{n_1} a_{n_2} \dots a_{n_j}. \quad (2)$$

In (2), the summation index set $\cup_{k=2}^n \mathcal{P}_{n+2-k}^*$ consists of all partitions of integers between 2 and n inclusive with all parts ≥ 2 . By appending 1's to these partitions (if necessary) we get all partitions of n except the all-1s partition. Thus, by setting $a_1 = 1$, (2) becomes, adopting the frequency-of-parts notation for integer partitions,

$$a_{n+2} = \sum_{1^{i_1} 2^{i_2} \dots n^{i_n} \vdash n} a_1^{i_1} a_2^{i_2} \dots a_n^{i_n}, \quad (3)$$

valid for $n \geq 1$, where the all-1s partition in (3) produces the initial 1 on the right side of (2). This is precisely the recurrence obtained by equating coefficients of x^{n+2} in the defining generating function identity

$$\sum_{n \geq 1} a_n x^n = x + \frac{x^2}{\prod_{n \geq 1} (1 - a_n x^n)} \quad (4)$$

for [A345973](#) and (4) immediately implies $a_1 = 1$.

□

The trees in \mathcal{A}_n with no gray edges are the trees counted by [A346787](#) (offset because A346787 counts by number of vertices which of course differs by 1 from the number of edges).

References

- [1] The On-Line Encyclopedia of Integer Sequences, published electronically at <http://oeis.org>, 2021.

Department of Statistics, University of Wisconsin-Madison