On comparison of fractional Laplacians

Alexander I. Nazarov^{*}

Abstract. For s > -1, $s \notin \mathbb{N}_0$, we compare two natural types of fractional Laplacians $(-\Delta)^s$, namely, the restricted Dirichlet and the spectral Neumann ones. We show that for the quadratic form of their difference taken on the space $\widetilde{H}^s(\Omega)$ is positive or negative depending on whether the integer part of s is even or odd. For $s \in (0,1)$ and convex domains we prove also that the difference of these operators is positivity preserving on $\widetilde{H}^s(\Omega)$. This paper complements [10] and [11] where similar statements were proved for the spectral Dirichlet and the restricted Dirichlet fractional Laplacians.

1 Introduction

In recent decades a lot of efforts have been invested in studying nonlocal differential operators and nonlocal variational problems. Model operators here are various fractional Laplacian (FLs for the brevity) $(-\Delta)^s$, mainly for $s \in (0, 1)$.

Recall that the **spectral Dirichlet and Neumann** FLs are the *s*th powers of conventional Dirichlet and Neumann Laplacian in the sense of spectral theory. In a Lipschitz bounded domain Ω , they can be defined by corresponding quadratic forms

$$Q_s^{\mathrm{DSp}}[u] \equiv ((-\Delta_{\Omega})_{\mathrm{DSp}}^s u, u) := \sum_{j=1}^{\infty} \lambda_j^s |(u, \varphi_j)|^2;$$
$$Q_s^{\mathrm{NSp}}[u] \equiv ((-\Delta_{\Omega})_{\mathrm{NSp}}^s u, u) := \sum_{j=0}^{\infty} \mu_j^s |(u, \psi_j)|^2,$$

^{*}St.Petersburg Department of Steklov Institute, Fontanka, 27, St.Petersburg, 191023, Russia and St.Petersburg State University, Universitetskii pr. 28, St.Petersburg, 198504, Russia. E-mail: al.il.nazarov@gmail.com. Supported by RFBR grant 20-01-00630.

where λ_j , φ_j and μ_j , ψ_j are eigenvalues and (normalized) eigenfunctions of the Dirichlet and Neumann Laplacian in Ω , respectively. Notice that $\mu_0 = 0$ and $\psi_0 \equiv const$.

For $s \in (0, 1)$ the domains of these quadratic forms are the classical Sobolev–Slobodetskii spaces (see [15, Ch. 4] or [5])

$$\operatorname{Dom}(Q_s^{\mathrm{DSp}}) = \widetilde{H}^s(\Omega); \qquad \operatorname{Dom}(Q_s^{\mathrm{NSp}}) = H^s(\Omega) \tag{1}$$

(we recall that

$$\widetilde{H}^{s}(\Omega) = H^{s}(\Omega) \quad \text{if} \quad 0 < s < 1/2; \qquad \widetilde{H}^{s}(\Omega) \subsetneq H^{s}(\Omega) \quad \text{if} \quad s \ge 1/2,$$

see, e.g., [15, 4.3.2]).

The first equality in (1) is proved in [10, Lemma 1]; the proof of the second one is quite similar.

For s > 1 the domains of spectral quadratic forms are more complicated but the following relations are always true:

$$\widetilde{H}^{s}(\Omega) \subset \operatorname{Dom}(Q^{\operatorname{DSp}}_{s}); \qquad \widetilde{H}^{s}(\Omega) \subset \operatorname{Dom}(Q^{\operatorname{NSp}}_{s}).$$

On the other hand, the quadratic form of **restricted Dirichlet** FL is defined as follows:

$$Q_s^{\mathrm{DR}}[u] \equiv \left((-\Delta_{\Omega})_{\mathrm{DR}}^s u, u \right) := \int_{\mathbb{R}^n} |\xi|^{2s} |\mathcal{F}u(\xi)|^2 d\xi$$

where \mathcal{F} is the Fourier transform

$$\mathcal{F}u(\xi) = \frac{1}{(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^n} e^{-i\langle\xi,x\rangle} u(x) \, dx.$$

Corresponding domain is $\widetilde{H}^s(\Omega)$ for all s > 0.

For $s \in (0, 1)$ the following relation holds:

$$Q_s^{\mathrm{DR}}[u] = c_{n,s} \iint_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} \, dx \, dy,$$

where

$$c_{n,s} = 2^{2s-1} \pi^{-n/2} \frac{\Gamma(\frac{n+2s}{2})}{|\Gamma(-s)|}.$$

Remark 1 Notice that for $s \in (0, 1)$ the quadratic form of restricted Neumann (or regional) FL is

$$Q_s^{\mathrm{NR}}[u] \equiv \left((-\Delta_\Omega)_{\mathrm{NSp}}^s u, u \right) := c_{n,s} \iint_{\Omega \times \Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} \, dx \, dy.$$

For some other types of fractional Laplacians see, e.g., [12] and references therein.

The operators $(-\Delta_{\Omega})^s_{\text{DSp}}$ and $(-\Delta_{\Omega})^s_{\text{DR}}$ were compared in the sense of quadratic forms and in the pointwise sense in [10] $(s \in (0, 1))$ and [11] (for partial results see also [4], [6], [7], [13]).

Theorem 1 (Theorem 2 in [10] and Theorem 1 in [11]) Let s > -1and $s \notin \mathbb{N}_0$. Suppose that $u \in \tilde{H}^s(\Omega)$, $u \not\equiv 0$. Then the following relation holds:

$$\begin{split} &Q^{\mathrm{DSp}}_{s}[u] > Q^{\mathrm{DR}}_{s}[u], \quad \text{if} \quad 2k < s < 2k+1, \quad k \in \mathbb{N}_{0}; \\ &Q^{\mathrm{DSp}}_{s}[u] < Q^{\mathrm{DR}}_{s}[u], \quad \text{if} \quad 2k-1 < s < 2k, \quad k \in \mathbb{N}_{0}. \end{split}$$

Theorem 2 1. (Theorem 1 in [10]) Let $s \in (0, 1)$, and let $u \in \widetilde{H}^s(\Omega)$, $u \ge 0, u \not\equiv 0$. Then the following relation holds in the sense of distributions:

$$(-\Delta_{\Omega})^s_{\mathrm{DSp}}u > (-\Delta_{\Omega})^s_{\mathrm{DR}}u.$$

2. (Theorem 3 in [11]) Let $s \in (-1,0)$. Suppose that $u \in H^s(\Omega)$, $u \ge 0$ in the sense of distributions, $u \ne 0$. Then the following relation holds:

$$(-\Delta_{\Omega})^s_{\mathrm{DSp}}u < (-\Delta_{\Omega})^s_{\mathrm{DR}}u.$$

In this paper we prove similar results for the operators $(-\Delta_{\Omega})^s_{\text{DR}}$ and $(-\Delta_{\Omega})^s_{\text{NSp}}$. Since the domains of their quadratic forms are in general different, we consider them on the smaller domain $\widetilde{H}^s(\Omega)$.

Theorem 3 Let s > -1 and $s \notin \mathbb{N}_0$. Suppose that $u \in \widetilde{H}^s(\Omega)$, $u \not\equiv 0$. Then the following relation holds:

$$Q_s^{\text{DR}}[u] > Q_s^{\text{NSp}}[u], \quad \text{if} \quad s \in (2k, 2k+1), \quad k \in \mathbb{N}_0;$$
 (2)

$$Q_s^{\mathrm{DR}}[u] < Q_s^{\mathrm{NSp}}[u], \quad \text{if} \quad s \in (2k-1, 2k), \quad k \in \mathbb{N}_0.$$
(3)

¹For n = 1 and $s \le -\frac{1}{2}$ assume in addition that $(u, \mathbf{1}) = 0$.

²For s < 0 assume in addition that (u, 1) = 0.

Remark 2 Notice that a weaker inequality $Q_s^{\text{DSp}}[u] \ge Q_s^{\text{NSp}}[u]$ for $u \in \widetilde{H}^s(\Omega)$, $s \in (0, 1)$, is a particular case of the well-known Heinz inequality [8]. On the other hand, the inequality $Q_s^{\text{DR}}[u] \ge Q_s^{\text{NR}}[u]$ for $u \in \widetilde{H}^s(\Omega)$, $s \in (0, 1)$, is trivial.

Theorem 4 Suppose that Ω is convex. Let $s \in (0,1)$, and let $u \in \widetilde{H}^s(\Omega)$, $u \ge 0, u \ne 0$. Then the following relation holds in the sense of distributions:

$$(-\Delta_{\Omega})^{s}_{\mathrm{DR}}u > (-\Delta_{\Omega})^{s}_{\mathrm{NSp}}u \qquad in \quad \Omega.$$
(4)

The structure of our paper is as follows. In Section 2 we recall the basic facts on the generalized harmonic extensions related to fractional Laplacians of orders $\sigma \in (0,1)$ and $-\sigma \in (-1,0)$. Theorems 3 and 4 are proved in Section 3. Also we show that the assumption of convexity in Theorem 4 cannot be removed.

2 Fractional Laplacians as D-to-N and N-to-D operators

It is a common knowledge nowaday that some of FLs of order $\sigma \in (0, 1)$ are related to the so-called *harmonic extension in* $n + 2 - 2\sigma$ *dimensions* and to the generalized Dirichlet-to-Neumann map (notice that for $\sigma = \frac{1}{2}$ it was known long ago).

Let $u \in H^{\sigma}(\mathbb{R}^n)$ (in our consideration, we always assume that $u \in H^{\sigma}(\Omega)$ is extended by zero to \mathbb{R}^n). In the pioneering paper [2], it was shown that there exists a unique solution $w_{\sigma}^{\mathrm{DR}}(x, y)$ of the BVP in the half-space

$$-\operatorname{div}(y^{1-2\sigma}\nabla w) = 0$$
 in $\mathbb{R}^n \times \mathbb{R}_+;$ $w\big|_{y=0} = u,$

with finite energy (weighted Dirichlet integral)

$$\mathcal{E}_{\sigma}^{\mathrm{DR}}(w) = \int_{0}^{\infty} \int_{\mathbb{R}^n} y^{1-2\sigma} |\nabla w(x,y)|^2 \, dx dy,$$

and the relation

$$(-\Delta_{\Omega})^{\sigma}_{\mathrm{DR}}u(x) = -C_{\sigma} \cdot \lim_{y \to 0^+} y^{1-2\sigma} \partial_y w^{\mathrm{DR}}_{\sigma}(x,y)$$
(5)

holds in the sense of distributions and pointwise at every point of smoothness of u. Here the constant C_{σ} is given by

$$C_{\sigma} := \frac{4^{\sigma} \Gamma(1+\sigma)}{\Gamma(1-\sigma)}.$$

Moreover, the function $w_{\sigma}^{\mathrm{DR}}(x,y)$ minimizes $\mathcal{E}_{\sigma}^{\mathrm{DR}}$ over the set

$$\mathcal{W}^{\mathrm{DR}}_{\sigma}(u) = \Big\{ w(x,y) \, : \, \mathcal{E}^{\mathrm{DR}}_{\sigma}(w) < \infty \, , \ w \big|_{y=0} = u \Big\},$$

and the following equality holds:

$$Q_{\sigma}^{\mathrm{DR}}[u] = \frac{C_{\sigma}}{2\sigma} \cdot \mathcal{E}_{\sigma}^{\mathrm{DR}}(w_{\sigma}^{\mathrm{DR}}).$$
(6)

In [14] this approach was transferred to quite general situation. In particular, it was shown that for $u \in H^{\sigma}(\Omega)$ there is a unique solution $w_{\sigma}^{\text{NSp}}(x, y)$ of the BVP in the half-cylinder

 $-\mathrm{div}(y^{1-2\sigma}\nabla w) = 0 \quad \mathrm{in} \quad \Omega \times \mathbb{R}_+; \qquad w\big|_{y=0} = u, \qquad \partial_{\mathbf{n}} w\big|_{x \in \partial\Omega} = 0$

(here **n** is the unit vector of exterior normal to $\partial \Omega$) having finite energy

$$\mathcal{E}^{\mathrm{NSp}}_{\sigma}(w) = \int_{0}^{\infty} \int_{\Omega} y^{1-2\sigma} |\nabla w(x,y)|^2 \, dx dy,$$

and the relation

$$(-\Delta_{\Omega})^{\sigma}_{\mathrm{NSp}}u(x) = -C_{\sigma} \cdot \lim_{y \to 0^+} y^{1-2\sigma} \partial_y w^{\mathrm{NSp}}_{\sigma}(x,y).$$
(7)

holds in the sense of distributions on Ω and pointwise at every point of smoothness of u.

Moreover, the function $w_{\sigma}^{\text{NSp}}(x, y)$ minimizes $\mathcal{E}_{\sigma}^{\text{NSp}}$ over the set

$$\mathcal{W}^{\mathrm{NSp}}_{\sigma,\Omega}(u) = \Big\{ w(x,y) \, : \, \mathcal{E}^{\mathrm{NSp}}_{\sigma}(w) < \infty \, , \ w \big|_{y=0} = u \Big\},$$

and the following equality holds:

$$Q_{\sigma}^{\mathrm{NSp}}[u] = \frac{C_{\sigma}}{2\sigma} \cdot \mathcal{E}_{\sigma}^{\mathrm{NSp}}(w_{\sigma}^{\mathrm{NSp}}).$$
(8)

In a similar way, one can connect FLs of order $-\sigma \in (-1,0)$ with the generalized Neumann-to-Dirichlet map. It was done in [3] for the spectral Dirichlet FL and in [1] for the FL in \mathbb{R}^n (and therefore for the restricted Dirichlet FL). Variational characterization of these operators was given in [11]. We formulate this result for the operator³ $(-\Delta_{\Omega})_{\text{DR}}^{-\sigma}$.

Let $u \in \widetilde{H}^{-\sigma}(\Omega)$ (for n = 1 and $\sigma \geq \frac{1}{2}$ assume in addition that $(u, \mathbf{1}) = 0$). We consider the problem of minimizing the functional

$$\widetilde{\mathcal{E}}_{-\sigma}^{\mathrm{DR}}(w) = \mathcal{E}_{\sigma}^{\mathrm{DR}}(w) - 2\left(u, w\big|_{y=0}\right)$$

over the set $\mathcal{W}_{-\sigma}^{\mathrm{DR}}$, that is closure of smooth functions on $\mathbb{R}^n \times \overline{\mathbb{R}}_+$ with bounded support, with respect to $\mathcal{E}_{\sigma}^{\mathrm{DR}}(\cdot)$. We notice that by the result of [2] the duality $(u, w|_{y=0})$ is well defined.

If $n > 2\sigma$ (this is a restriction only for n = 1) then the minimizer is determined uniquely. Denote it by $w_{-\sigma}^{\text{DR}}(x, y)$. Then formulae (5) and (6) imply the relations

$$Q_{-\sigma}^{\mathrm{DR}}[u] = -\frac{2\sigma}{C_{\sigma}} \cdot \widetilde{\mathcal{E}}_{-\sigma}^{\mathrm{DR}}(w_{-\sigma}^{\mathrm{DR}}); \qquad (-\Delta_{\Omega})_{\mathrm{DR}}^{-\sigma}u(x) = \frac{2\sigma}{C_{\sigma}} w_{-\sigma}^{\mathrm{DR}}(x,0) \qquad (9)$$

(the second relation holds for a.a. $x \in \Omega$).

In case $n = 1 \leq 2\sigma$ the minimizer $w_{-\sigma}^{DR}(x, y)$ is defined up to an additive constant. However, by assumption $(u, \mathbf{1}) = 0$ the functional $\widetilde{\mathcal{E}}_{-\sigma}^{DR}(w_{-\sigma}^{DR})$ does not depend on the choice of the constant, and the first relation in (9) holds. The second equality in (9) also holds if we choose the constant such that $w_{-\sigma}^{DR}(x, 0) \to 0$ as $|x| \to \infty$.

Notice that the function $w_{-\sigma}^{\mathrm{DR}}$ solves the Neumann problem in the half-space

$$-\operatorname{div}(y^{1-2\sigma}\nabla w) = 0$$
 in $\mathbb{R}^n \times \mathbb{R}_+;$ $\lim_{y \to 0^+} y^{1-2\sigma} \partial_y w = -u$

(the boundary condition holds in the sense of distributions). So, we obtain the "dual" Caffarelli–Silvestre characterization of $(-\Delta_{\Omega})_{\rm DR}^{-\sigma}$ as the Neumannto-Dirichlet map.

Now we introduce the "dual" Stinga–Torrea characterization of $(-\Delta_{\Omega})_{NSp}^{-\sigma}$ in almost the same way as it was done in [11] for $(-\Delta_{\Omega})_{DSp}^{-\sigma}$. Namely, let

³We emphasize that $(-\Delta_{\Omega})_{\mathrm{DR}}^{-\sigma}$ is not inverse to $(-\Delta_{\Omega})_{\mathrm{DR}}^{\sigma}$.

 $u \in H^{-\sigma}(\Omega)$ and let $(u, \mathbf{1}) = 0$. Then the function $w_{-\sigma}^{NSp}(x, y)$ minimizing the functional

$$\widetilde{\mathcal{E}}_{-\sigma}^{NSp}(w) = \mathcal{E}_{\sigma}^{NSp}(w) - 2\left(u, w\Big|_{y=0}\right)$$

over the set

$$\mathcal{W}_{-\sigma,\Omega}^{\mathrm{NSp}}(u) = \Big\{ w(x,y) \, : \, \mathcal{E}_{\sigma}^{\mathrm{NSp}}(w) < \infty \Big\},$$

is defined up to an additive constant. By assumption $(u, \mathbf{1}) = 0$ the functional $\widetilde{\mathcal{E}}_{-\sigma}^{\mathrm{NSp}}(w_{-\sigma}^{\mathrm{NSp}})$ does not depend on the choice of the constant, and formulae (8) and (7) imply

$$Q_{-\sigma}^{NSp}[u] = -\frac{2\sigma}{C_{\sigma}} \cdot \widetilde{\mathcal{E}}_{-\sigma}^{NSp}(w_{-\sigma}^{NSp}); \qquad (-\Delta_{\Omega})_{NSp}^{-\sigma}u(x) = \frac{2\sigma}{C_{\sigma}}w_{-\sigma}^{NSp}(x,0) \quad (10)$$

(The second equality holds for a.a. $x \in \Omega$ if we choose the constant such that $w_{-\sigma}^{\text{NSp}}(x, y) \to 0$ as $y \to +\infty$).

Also the function $w_{-\sigma}^{\text{NSp}}$ solves the Neumann problem in the half-cylinder

$$-\mathrm{div}(y^{1-2\sigma}\nabla w) = 0 \quad \text{in} \quad \Omega \times \mathbb{R}_+; \quad \lim_{y \to 0^+} y^{1-2\sigma} \partial_y w = -u, \quad \partial_{\mathbf{n}} w \big|_{x \in \partial \Omega} = 0$$

(the boundary condition on the bottom holds in the sense of distributions).

3 Proof of main results

Proof of Theorem 3. We split the proof in three parts.

1. Let $s \in (0,1)$. For any $w \in \mathcal{W}_s^{\mathrm{DR}}(u)$ we have $w|_{\Omega \times \mathbb{R}_+} \in \mathcal{W}_{s,\Omega}^{\mathrm{NSp}}(u)$. Therefore, relations (6) and (8) provide

$$\begin{split} Q_s^{\mathrm{NSp}}[u] &= \frac{C_s}{2s} \cdot \inf_{w \in \mathcal{W}_{s,\Omega}^{\mathrm{NSp}}(u)} \mathcal{E}_s^{\mathrm{NSp}}(w) \leq \frac{C_s}{2s} \, \mathcal{E}_s^{\mathrm{NSp}}(w_s^{\mathrm{DR}}) \\ &\leq \frac{C_s}{2s} \, \mathcal{E}_s^{\mathrm{DR}}(w_s^{\mathrm{DR}}) = Q_s^{\mathrm{DR}}[u], \end{split}$$

and (2) follows with the large sign.

Finally, the equality in (2) implies $\nabla w_s^{\text{DR}} = 0$ on $(\mathbb{R}^n \setminus \Omega) \times \mathbb{R}_+$. Since any *x*-derivative of w_s^{DR} solves the same equation in the whole half-space $\mathbb{R}^n \times \mathbb{R}_+$, it should be zero everywhere that is impossible for $u \neq 0$.

2. Let $s \in (-1, 0)$. We define $\sigma = -s \in (0, 1)$ and construct the extension $w_{-\sigma}^{\text{DR}}$ as described in Section 2.

We again have $w_{-\sigma}^{\text{DR}}|_{\Omega \times \mathbb{R}_+} \in \mathcal{W}_{-\sigma,\Omega}^{\text{NSp}}(u)$. Therefore, relations (9) and (10) provide

$$\begin{split} -Q_{s}^{\mathrm{NSp}}[u] &= \frac{2\sigma}{C_{\sigma}} \cdot \inf_{w \in \mathcal{W}_{-\sigma,\Omega}^{\mathrm{NSp}}} \widetilde{\mathcal{E}}_{-\sigma}^{\mathrm{NSp}}(w) \leq \frac{2\sigma}{C_{\sigma}} \, \widetilde{\mathcal{E}}_{-\sigma}^{\mathrm{NSp}}(w_{-\sigma}^{\mathrm{DR}}) \\ &\leq \frac{2\sigma}{C_{\sigma}} \, \widetilde{\mathcal{E}}_{-\sigma}^{\mathrm{DR}}(w_{-\sigma}^{\mathrm{DR}}) = -Q_{s}^{\mathrm{DR}}[u], \end{split}$$

and (3) follows with the large sign. To complete the proof, we repeat the argument of the first part.

3. Now let $s > 1, s \notin \mathbb{N}$. We put $k = \lfloor \frac{s+1}{2} \rfloor$ and define for $u \in \widetilde{H}^s(\Omega)$

$$v = (-\Delta)^k u \in \widetilde{H}^{s-2k}(\Omega), \qquad s - 2k \in (-1,0) \cup (0,1).$$

Note that $v \neq 0$ if $u \neq 0$, and

$$(v, \mathbf{1}) = \mathcal{F}v(0) = |\xi|^{2k} \mathcal{F}u(\xi)|_{\xi=0} = 0.$$

Then we have

$$Q_s^{\text{DR}}[u] = Q_{s-2k}^{\text{DR}}[v], \qquad Q_s^{\text{NSp}}[u] = Q_{s-2k}^{\text{NSp}}[v],$$

and the conclusion follows from cases 1 and 2.

Proof of Theorem 4. We recall the representation formulae for w_s^{DR} and w_s^{NSp} , see [2] and [14], respectively:

$$w_s^{\mathrm{DR}}(x,y) = const \cdot \int_{\mathbb{R}^n} \frac{y^{2s} u(\xi) \, d\xi}{\left(|x-\xi|^2+y^2\right)^{\frac{n+2s}{2}}};$$
$$w_s^{\mathrm{NSp}}(x,y) = \sum_{j=0}^{\infty} (u,\psi_j)_{L_2(\Omega)} \cdot \mathcal{Q}_s(y\sqrt{\mu_j})\psi_j(x), \qquad \mathcal{Q}_s(\tau) = \frac{2^{1-s}\tau^s}{\Gamma(s)}\mathcal{K}_s(\tau),$$

where $\mathcal{K}_s(\tau)$ stands for the modified Bessel function of the second kind. First of all, these formulae imply for $u \ge 0$, $u \not\equiv 0$

$$\lim_{y \to +\infty} w_s^{\text{DR}}(x, y) = 0; \qquad \lim_{y \to +\infty} w_s^{\text{NSp}}(x, y) = (u, \psi_0)_{L_2(\Omega)} \cdot \psi_0(x) > 0;$$

the second relation follows from the asymptotic behavior (see, e.g., [14, (3.7)])

$$\mathcal{K}_s(\tau) \sim \Gamma(s) 2^{s-1} \tau^{-s}, \quad \text{as} \quad \tau \to 0;$$

 $\mathcal{K}_s(\tau) \sim \left(\frac{\pi}{2\tau}\right)^{\frac{1}{2}} e^{-\tau} \left(1 + O(\tau^{-1})\right) \quad \text{as} \quad \tau \to +\infty.$

Next, for $x \in \partial \Omega$ we derive by convexity of Ω

$$\partial_{\mathbf{n}} w_s^{\mathrm{DR}}(x,y) = const \cdot \int\limits_{\mathbb{R}^n} \frac{y^{2s} \langle (\xi-x), \mathbf{n} \rangle u(\xi) \, d\xi}{(|x-\xi|^2+y^2)^{\frac{n+2s+2}{2}}} < 0.$$

Thus, the difference $W(x, y) = w_s^{NSp}(x, y) - w_s^{DR}(x, y)$ has the following properties in the half-cylinder $\Omega \times \mathbb{R}_+$:

$$-\mathrm{div}(y^{1-2s}\nabla W) = 0; \qquad W\big|_{y=0} = 0; \qquad W\big|_{y=\infty} > 0; \qquad \partial_{\mathbf{n}}W\big|_{x\in\partial\Omega} > 0.$$

By the strong maximum principle, W > 0 in $\Omega \times \mathbb{R}_+$. Finally, we apply the boundary point principle (the Hopf–Oleinik lemma, see [9]) to the function $W(x, t^{\frac{1}{2s}})$ and obtain (cf. [10, Theorem 1])

$$\liminf_{y \to 0^+} y^{1-2\sigma} \partial_y W(x,y) = \liminf_{y \to 0^+} \frac{W(x,y)}{y^{2s}} = \liminf_{t \to 0^+} \frac{W(x,t^{\frac{1}{2s}})}{t} > 0, \quad x \in \Omega.$$

This completes the proof in view of (5) and (7).

Remark 3 For non-convex domains the relation (4) does not hold in general. We provide corresponding counterexample.

Put temporarily $\Omega = \Omega_1 \cup \Omega_2$ where $\Omega_1 \cap \Omega_2 = \emptyset$. If $u \ge 0$ is a smooth function supported in Ω_1 then easily $(-\Delta_{\Omega})^s_{NSp}u \equiv 0$ in Ω_2 . On the other hand, $w^{DR}_s(x,y) > 0$ for all $x \in \mathbb{R}^n$, y > 0, and the Hopf–Oleinik lemma gives $(-\Delta_{\Omega})^s_{DR}u < 0$ in Ω_2 .

Finally, if we join Ω_1 with Ω_2 by a small channel then the inequality $(-\Delta_{\Omega})^s_{\mathrm{DR}}u < (-\Delta_{\Omega})^s_{\mathrm{NSp}}u$ in Ω_2 holds by continuity.

References

 X. Cabré and Y. Sire, Nonlinear equations for fractional Laplacians. I: Regularity, maximum principles, and Hamiltonian estimates, AIHP – AN. 31 (2014), no. 1, 23–53.

- [2] L. Caffarelli and L. Silvestre, An extension problem related to the fractional Laplacian, Comm. PDEs. 32 (2007), no. 7-9, 1245–1260.
- [3] A. Capella, J. Dávila, L. Dupaigne and Y. Sire, Regularity of radial extremal solutions for some non-local semilinear equations, Comm. PDEs. 36 (2011), no. 8, 1353–1384.
- [4] Z.-Q. Chen and R. Song, Two-sided eigenvalue estimates for subordinate processes in domains, J. Funct. Anal. 226 (2005), 90–113.
- [5] E. Di Nezza, G. Palatucci and E. Valdinoci, *Hitchhiker's guide to the frac*tional Sobolev spaces, Bull. Sci. Math., **136** (2012), no. 5, 521–573.
- [6] M.M. Fall, Semilinear elliptic equations for the fractional Laplacian with Hardy potential, Nonlin. Analysis – TMA, **193** (2020), 111311, DOI 10.1016/j.na.2018.07.008. Arxiv preprint 1109.5530v4 (2012).
- [7] R. L. Frank and L. Geisinger, *Refined semiclassical asymptotics for frac*tional powers of the Laplace operator, J. reine und angew. Math. (Crelles Journal), **712** (2016), 1–37.
- [8] E. Heinz, Beiträge zur Störungstheorie der Spektralzerlegung, Math. Ann., 193 (1951), 415–438.
- [9] L. I. Kamynin and B. N. Himčenko, Theorems of Giraud type for second order equations with a weakly degenerate non-negative characteristic part, Sib. Math. J. 18 (1977), 76–91.
- [10] R. Musina and A.I. Nazarov, On fractional Laplacians, Comm. PDEs, 39 (2014), no. 9, 1780–1790.
- [11] R. Musina and A.I. Nazarov, On fractional Laplacians-2, Ann. Inst. Henri Poincaré – An. Nonlin., 33 (2016), no. 6, 1667–1673.
- [12] R. Musina and A.I. Nazarov, Strong maximum principles for fractional Laplacians, Proc. Roy. Soc. Edinburgh A. 149 (2019), no. 5, 1223–1240.
- [13] R. Servadei and E. Valdinoci, On the spectrum of two different fractional operators, Proc. Roy. Soc. Edinburgh A, 144 (2014), no. 4, 831–855.
- [14] P. R. Stinga and J. L. Torrea, Extension problem and Harnack's inequality for some fractional operators, Comm. PDEs. 35 (2010), no. 11, 2092–2122.
- [15] H. Triebel, Interpolation theory, function spaces, differential operators, Deutscher Verlag Wissensch., Berlin, 1978.