

# Some Results on $k$ -Critical $P_5$ -Free Graphs

Qingqiong Cai\*    Jan Goedgebeur †‡    Shenwei Huang§

August 12, 2021

## Abstract

A graph  $G$  is  $k$ -vertex-critical if  $G$  has chromatic number  $k$  but every proper induced subgraph of  $G$  has chromatic number less than  $k$ . The study of  $k$ -vertex-critical graphs for graph classes is an important topic in algorithmic graph theory because if the number of such graphs that are in a given hereditary graph class is finite, then there is a polynomial-time algorithm to decide if a graph in the class is  $(k - 1)$ -colorable.

In this paper, we prove that for every fixed integer  $k \geq 1$ , there are only finitely many  $k$ -vertex-critical  $(P_5, \text{gem})$ -free graphs and  $(P_5, \overline{P_3 + P_2})$ -free graphs. To prove the results we use a known structure theorem for  $(P_5, \text{gem})$ -free graphs combined with properties of  $k$ -vertex-critical graphs. Moreover, we characterize all  $k$ -vertex-critical  $(P_5, \text{gem})$ -free graphs and  $(P_5, \overline{P_3 + P_2})$ -free graphs for  $k \in \{4, 5\}$  using a computer generation algorithm.

**Keywords.** Graph coloring;  $k$ -critical graphs; forbidden induced subgraphs; computer generation algorithm.

## 1 Introduction

All graphs in this paper are finite and simple. We say that a graph  $G$  contains a graph  $H$  if  $H$  is isomorphic to an induced subgraph of  $G$ . A graph  $G$  is  $H$ -free if it does not contain  $H$ . For a family of graphs  $\mathcal{H}$ ,  $G$  is  $\mathcal{H}$ -free if  $G$  is  $H$ -free for every  $H \in \mathcal{H}$ . When  $\mathcal{H}$  consists of two graphs, we write  $(H_1, H_2)$ -free instead of  $\{H_1, H_2\}$ -free.

A  $q$ -coloring of a graph  $G$  is a function  $\phi : V(G) \rightarrow \{1, \dots, q\}$  such that  $\phi(u) \neq \phi(v)$  whenever  $u$  and  $v$  are adjacent in  $G$ . Equivalently, a  $q$ -coloring of  $G$  is a partition of  $V(G)$  into  $q$  independent sets. A graph is  $q$ -colorable if it admits a  $q$ -coloring. The chromatic number of a graph  $G$ , denoted by  $\chi(G)$ , is the minimum number  $q$  for which  $G$  is  $q$ -colorable. The clique number of  $G$ , denoted by  $\omega(G)$ , is the size of a largest clique in  $G$ .

A graph  $G$  is  $k$ -chromatic if  $\chi(G) = k$ . We say that  $G$  is  $k$ -critical if it is  $k$ -chromatic and  $\chi(G - e) < \chi(G)$  for any edge  $e \in E(G)$ . For instance,  $K_2$  is the only

\*College of Computer Science, Nankai University, Tianjin 300350, China. Supported by Natural Science Foundation of Tianjin (19JCQNJC14400).

†Department of Applied Mathematics, Computer Science and Statistics, Ghent University, 9000 Ghent, Belgium.

‡Department of Computer Science, KU Leuven Campus Kulak, 8500 Kortrijk, Belgium.

§The corresponding author. College of Computer Science, Nankai University, Tianjin 300350, China. Supported by the National Natural Science Foundation of China (11801284) and Natural Science Foundation of Tianjin (20JCYBJC01190).

2-critical graph and odd cycles are the only 3-critical graphs. A graph is *critical* if it is  $k$ -critical for some integer  $k \geq 1$ . Critical graphs were first defined and studied by Dirac [11, 12, 13] in the early 1950s, and then by Gallai and Ore [14, 15, 24] among many others, and more recently by Kostochka and Yancey [22].

A weaker notion of criticality is the so-called vertex-criticality. A graph  $G$  is  $k$ -vertex-critical if  $\chi(G) = k$  and  $\chi(G - v) < k$  for any  $v \in V(G)$ . For a set  $\mathcal{H}$  of graphs and a graph  $G$ , we say that  $G$  is  $k$ -vertex-critical  $\mathcal{H}$ -free if it is  $k$ -vertex-critical and  $\mathcal{H}$ -free. We are mainly interested in the following question.

**The meta question.** Given a set  $\mathcal{H}$  of graphs and an integer  $k \geq 1$ , are there only finitely many  $k$ -vertex-critical  $\mathcal{H}$ -free graphs?

This question is important in the study of algorithmic graph theory because of the following theorem, whose proof can be found in [4].

**Theorem 1 (Folklore).** *Given a set  $\mathcal{H}$  of graphs and an integer  $k \geq 1$ , if the set of all  $k$ -vertex-critical  $\mathcal{H}$ -free graphs is finite, then there is a polynomial-time algorithm to determine whether an  $\mathcal{H}$ -free graph is  $(k - 1)$ -colorable.*

As usual,  $P_t$  and  $C_s$  denote the path on  $t$  vertices and the cycle on  $s$  vertices, respectively. The complete graph on  $n$  vertices is denoted by  $K_n$ . For two graphs  $G$  and  $H$ , we use  $G + H$  to denote the *disjoint union* of  $G$  and  $H$ . For a positive integer  $r$ , we use  $rG$  to denote the disjoint union of  $r$  copies of  $G$ . The *complement* of  $G$  is denoted by  $\overline{G}$ . A *clique* (resp. *independent set*) in a graph is a set of pairwise adjacent (resp. nonadjacent) vertices. If a graph  $G$  can be partitioned into  $k$  independent sets  $S_1, \dots, S_k$  such that there is an edge between every vertex in  $S_i$  and every vertex in  $S_j$  for all  $1 \leq i < j \leq k$ ,  $G$  is called a *complete  $k$ -partite graph*; each  $S_i$  is called a *part* of  $G$ . If we do not specify the number of parts in  $G$ , we simply say that  $G$  is a *complete multipartite graph*. We denote by  $K_{n_1, \dots, n_k}$  the complete  $k$ -partite graph such that the  $i$ th part  $S_i$  has size  $n_i$ , for each  $1 \leq i \leq k$ .

In this paper, we continue to study  $k$ -vertex-critical graphs in the class of  $P_5$ -free graphs. Our research is mainly motivated by the following dichotomy result.

**Theorem 2 ([5]).** *Let  $H$  be a graph of order 4 and  $k \geq 5$  be a fixed integer. Then there are infinitely many  $k$ -vertex-critical  $(P_5, H)$ -free graphs if and only if  $H$  is  $2P_2$  or  $P_1 + K_3$ .*

This Theorem completely settles the finiteness question of  $k$ -vertex-critical  $(P_5, H)$ -free graphs for graphs of order 4. In [5], the authors also posed the natural question of which five-vertex graphs  $H$  lead to finitely many  $k$ -vertex-critical  $(P_5, H)$ -free graphs. It is known that there are exactly 13 5-vertex-critical  $(P_5, C_5)$ -free graphs [19], and that there are finitely many 5-vertex-critical  $(P_5, \text{banner})$ -free graphs [4, 20], and finitely many  $k$ -vertex-critical  $(P_5, \overline{P_5})$ -free graphs for every fixed  $k$  [10]. Hell and Huang proved that there are finitely many  $k$ -vertex-critical  $(P_6, C_4)$ -free graphs [18]. This was later generalized to  $(P_t, K_{r,s})$ -free graphs in the context of  $H$ -coloring [21]. This gives an affirmative answer for  $H = K_{2,3}$ . Apart from these, there seem to be very few results on the finiteness of  $k$ -vertex-critical graphs for  $k \geq 5$ . In this paper, we prove two new finiteness results beyond 5-vertex-criticality.

## 1.1 Our Contributions

We continue to study the subclasses of  $P_5$ -free graphs. In particular, we focus on  $(P_5, H)$ -free graphs when  $H$  has order 5. The *gem* graph is the graph consisting of an



**Figure 1:** The graphs gem (left) and  $\overline{P_3 + P_2}$  (right).

induced  $P_4$  with an additional vertex that is adjacent to every vertex on the  $P_4$ . In this paper, we prove that there are only finitely many  $k$ -vertex-critical  $(P_5, H)$ -free graphs for every fixed  $k \geq 1$  when  $H$  is gem or  $\overline{P_3 + P_2}$  (see Figure 1 for drawings of gem and  $\overline{P_3 + P_2}$ ). Moreover, we characterize all 4- and 5-vertex-critical  $(P_5, \text{gem})$ -free graphs and  $(P_5, \overline{P_3 + P_2})$ -free graphs by extending the computer generation algorithm from [17].

To prove the result on gem-free graphs, we used a known structure theorem for  $(P_5, \text{gem})$ -free graphs that is used to give an optimal  $\chi$ -binding function for the class combined with an inductive argument based on a careful analysis of the structure. For the  $\overline{P_3 + P_2}$ -free case, we performed a careful structural analysis combined with the repeated use of the pigeonhole principle based on the properties of  $k$ -vertex-critical graphs.

The remainder of the paper is organized as follows. We present some preliminaries in Section 2 and prove our new results in Section 3. Finally, we give some concluding remarks in Section 4.

## 2 Preliminaries

For general graph theory notation we follow [1]. Let  $G = (V, E)$  be a graph. If  $uv \in E$ , we say that  $u$  and  $v$  are *neighbors* or *adjacent*; otherwise  $u$  and  $v$  are *nonneighbors* or *nonadjacent*. The *neighborhood* of a vertex  $v$ , denoted by  $N_G(v)$ , is the set of neighbors of  $v$ . For a set  $X \subseteq V(G)$ , let  $N_G(X) = \bigcup_{v \in X} N_G(v) \setminus X$ . We shall omit the subscript whenever the context is clear. For  $X, Y \subseteq V$ , we say that  $X$  is *complete* (resp. *anticomplete*) to  $Y$  if every vertex in  $X$  is adjacent (resp. nonadjacent) to every vertex in  $Y$ . If  $X = \{x\}$ , we write “ $x$  is complete (resp. anticomplete) to  $Y$ ” instead of “ $\{x\}$  is complete (resp. anticomplete) to  $Y$ ”. If a vertex  $v$  is neither complete nor anticomplete to a set  $S$ , we say that  $v$  is *mixed* on  $S$ . We say that  $H$  is a *homogeneous set* if no vertex in  $V - H$  is mixed on  $H$ . More generally, we say that  $H$  is *homogeneous with respect to a subset*  $S \subseteq V$  if no vertex in  $S$  can be mixed on  $H$ . A vertex is *universal* in  $G$  if it is adjacent to all other vertices. Two nonadjacent vertices  $u$  and  $v$  are said to be *comparable* if  $N(v) \subseteq N(u)$  or  $N(u) \subseteq N(v)$ . A vertex subset  $K \subseteq V$  is a *clique cutset* if  $G - K$  has more components than  $G$  and  $K$  induces a clique. For  $S \subseteq V$ , the subgraph *induced* by  $S$ , is denoted by  $G[S]$ . A  $k$ -*hole* in a graph is an induced cycle  $H$  of length  $k \geq 4$ . If  $k$  is odd, we say that  $H$  is an *odd hole*. A  $k$ -*antihole* in  $G$  is a  $k$ -hole in  $\overline{G}$ . Odd antiholes are defined analogously.

We proceed with a few useful results that will be needed later. The first one is a folklore property of  $k$ -vertex-critical graphs.

**Lemma 1** (Folklore). *A  $k$ -vertex-critical graph cannot contain clique cutsets.*

Another folklore property of vertex-critical graphs is that such graphs contain no comparable vertices. A generalization of this property was presented in [5].

**Lemma 2** ([5]). *Let  $G$  be a  $k$ -vertex-critical graph. Then  $G$  has no two nonempty disjoint subsets  $X$  and  $Y$  of  $V(G)$  that satisfy all the following conditions.*

- $X$  and  $Y$  are anticomplete to each other.
- $\chi(G[X]) \leq \chi(G[Y])$ .
- $Y$  is complete to  $N(X)$ .

A graph  $G$  is *perfect* if  $\chi(H) = \omega(H)$  for each induced subgraph  $H$  of  $G$ . An *imperfect* graph is a graph that is not perfect. We conclude this section with the celebrated Strong Perfect Graph Theorem [9].

**Theorem 3** (Strong Perfect Graph Theorem [9]). *A graph is perfect if and only if it contains no odd holes or odd antiholes.*

### 3 New Results

#### 3.1 Gem-Free Graphs

**Theorem 4.** *For every fixed integer  $k \geq 1$ , there are finitely many  $k$ -vertex-critical  $(P_5, \text{gem})$ -free graphs.*

To prove Theorem 4, we need a structure theorem for the class of  $(P_5, \text{gem})$ -free graphs that was obtained in [8]. Let  $\mathcal{H}$  be the class of connected  $(P_5, \text{gem})$ -free graphs  $G$  such that  $V(G)$  can be partitioned into seven nonempty sets  $A_1, A_2, \dots, A_7$  such that:


- Each  $A_i$  induces a  $P_4$ -free graph.
- $A_1$  is complete to  $A_2 \cup A_5 \cup A_6$  and anticomplete to  $A_3 \cup A_4 \cup A_7$ .
- $A_3$  is complete to  $A_2 \cup A_4 \cup A_6$  and anticomplete to  $A_5 \cup A_7$ .
- $A_4$  is complete to  $A_5 \cup A_6$  and anticomplete to  $A_2 \cup A_7$ .
- $A_2$  is anticomplete to  $A_5 \cup A_6 \cup A_7$ .
- $A_5$  is anticomplete to  $A_6 \cup A_7$ .
- The vertex set of each component of  $G[A_7]$  is a homogeneous set of  $G$ .
- The adjacency between  $A_6$  and  $A_7$  is not specified but it is restricted by the fact that  $G$  is  $(P_5, \text{gem})$ -free.

See Figure 2 for an illustration.

For two graphs  $G$  and  $H$ , we say that  $G$  is an *substitution* of  $H$  if  $V(G)$  can be partitioned into  $|V(H)|$  nonempty subsets  $Q_v$ ,  $v \in V(H)$  so that  $Q_u$  and  $Q_v$  are complete if  $uv \in E(H)$  and anticomplete if  $uv \notin E(H)$  for every different vertices  $u, v \in V(H)$ . If each  $Q_v$  is  $P_4$ -free, then it is called a  *$P_4$ -free substitution*; if each  $Q_v$  is a clique, then it is called a *clique substitution*. To state the structural theorem, we also need 10 special graphs  $G_1, G_2, \dots, G_{10}$  each of which has order at most 9. We omit the drawing of these 10 graphs and refer the readers to [8]. Now we are ready to state the structure theorem obtained in [8].

**Lemma 3** ([8]). *Let  $G$  be a connected  $(P_5, \text{gem})$ -free graph that contains an induced  $C_5$ . Then either  $G \in \mathcal{H}$  or  $G$  is a  $P_4$ -free substitution of  $G_1, G_2, \dots, G_{10}$ .*

We also need a lemma from [6].



**Figure 2:** A diagram for graphs  $G \in \mathcal{H}$ . A thick line between two sets means that the two sets are complete. A normal line between two sets means that the edges are arbitrary (subject to the  $(P_5, \text{gem})$ -freeness). No line between two sets means that the two sets are anticomplete. Moreover, each  $A_i$  is  $P_4$ -free, and the vertex set of each component of  $G[A_7]$  is a homogeneous set of  $G$ .

**Lemma 4** ([6]). *Let  $G$  be a connected  $(P_5, \text{gem})$ -free graph and  $H$  a homogeneous set of  $G$  that is not a clique. Then there exists a connected induced subgraph  $G'$  of  $G$  with  $|G'| < |G|$  such that  $\omega(G') = \omega(G)$  and  $\chi(G') = \chi(G)$ .*

We are now ready to prove the main result in this subsection.

*Proof of Theorem 4.* Let  $G$  be a  $k$ -vertex-critical  $(P_5, \text{gem})$ -free graph. Then  $G$  is connected. We show that  $|G|$  is bounded by a function of  $k$ . If  $G$  contains a  $K_k$ , then  $G = K_k$ . So we may assume that  $G$  is  $K_k$ -free. Since  $G$  is not perfect, it follows from Theorem 3 that  $G$  contains an induced  $C_5$ . By Lemma 3,  $G \in \mathcal{H}$  or  $G$  is a  $P_4$ -expansion of  $G_1, G_2, \dots, G_{10}$ . If  $G$  is a  $P_4$ -free substitution of  $G_1, G_2, \dots, G_{10}$ , then it must be a clique expansion of  $G_1, G_2, \dots, G_{10}$  by Lemma 4. Since  $|G_i| \leq 9$  for  $1 \leq i \leq 10$  (see [8]), it follows that  $|G| \leq 9k$ .

Therefore, we assume that  $G \in \mathcal{H}$  in the following and can be partitioned into seven nonempty subsets  $A_1, A_2, \dots, A_7$  as described in the definition. Note that each  $A_i$  ( $1 \leq i \leq 5$ ) is a homogeneous set of  $G$ . Thus,  $A_i$  (for  $1 \leq i \leq 5$ ) is a clique by Lemma 4 and so  $|A_i| \leq k$ . So it remains to bound  $A_6$  and  $A_7$ . For the same reason, every component of  $G[A_7]$  is a clique and has size at most  $k$ .

**Claim 1.** *Let  $X \subseteq A_6$  be a subset such that  $X$  is homogeneous with respect to  $V(G) - A_7$ . Then  $|X| \leq 2^{2\omega^2(X)}$ .*

*Proof.* We prove this by induction on  $\omega(X)$ . Since  $A_3 \neq \emptyset$ , it contains a vertex, say  $t$ . The base case is  $\omega(X) = 1$ , i.e.,  $X$  is an independent set. If  $X$  contains two vertices  $x_1, x_2$ , then by Lemma 2 and our assumption there exist vertices  $y_1, y_2 \in A_7$  such that  $x_1y_1, x_2y_2 \in E$  but  $x_1y_2, x_2y_1 \notin E$ . Since every component of  $G[A_7]$  is a homogeneous set of  $G$ ,  $y_1y_2 \notin E$ . Then  $y_1, x_1, t, x_2, y_2$  induces a  $P_5$ . This proves that  $|X| \leq 1$ .

Suppose now that  $\omega := \omega(X) \geq 2$  and the claim is true for any subset  $Y$  such that  $Y$  is homogeneous with respect to  $V(G) - A_7$  with  $\omega(Y) < \omega$ . We first prove two useful facts about components of  $X$ .

All but at most one component of  $X$  are homogeneous sets of  $G$ . (1)

*Proof of (1).* Suppose not. Then there exist two components  $C$  and  $C'$  of  $X$  such that  $C$  and  $C'$  are not homogeneous sets of  $G$ . By our assumption on  $X$ , there are vertices  $v, v' \in A_7$  such that  $v$  is mixed on an edge  $ab \in C$  with  $va \notin E$  and  $vb \in E$ , and  $v'$  is mixed on an edge  $a'b' \in C'$  with  $v'a' \notin E$  and  $v'b' \in E$ .

If  $v, v'$  are in the same component of  $A_7$ , then  $vv' \in E$ . Since every component of  $A_7$  is a homogeneous set of  $G$ ,  $vb', v'b \in E$  and  $va', v'a \notin E$ . Then  $a, b, v, b', a'$  induces a  $P_5$ .

So  $v, v'$  lie in different components of  $A_7$ . If  $vb', v'b \notin E$ , then  $v, b, t, b', v'$  induces a  $P_5$ . So by symmetry we assume that  $vb' \in E$ . Then  $va' \in E$  since otherwise  $a, b, v, b', a'$  induces a  $P_5$ . Note that  $a, b, v, b', v'$  induces a  $P_5$  unless  $v'$  is adjacent to  $a$  or  $b$ . If  $v'b \in E$ , then  $v'a \in E$  since otherwise  $a, b, v', b', a'$  induces a  $P_5$ . This shows that  $v'a \in E$ . But now  $va'$  and  $v'a$  induce a  $2P_2$  and this together with  $t \in A_3$  gives an induced  $P_5$ . This completes the proof of (1).

There is at most one homogeneous component of  $X$  of size  $i$  for  $1 \leq i \leq \omega$ . (2)

*Proof of (2).* Suppose not. Then there are two homogeneous components  $C$  and  $C'$  of  $X$  of size  $i$  for some  $1 \leq i \leq k$ . By Lemma 4, we know that  $C$  and  $C'$  are two cliques of size  $i$ , and thus  $\chi(G[C]) = \chi(G[C']) = i$ . By Lemma 2, there exist  $a, a' \in A_7$  such that  $a \in N(C) \setminus N(C')$  and  $a' \in N(C') \setminus N(C)$ . Then  $a, c, t, c', a'$  induces a  $P_5$ , where  $c \in C, c' \in C'$ . This completes the proof of (2).

Let  $H$  be the possible non-homogeneous component of  $X$ . Then  $|X| \leq \omega^2 + |H|$ . We now bound  $H$ . Let  $K = \{v_1, v_2, \dots, v_s\}$  be a maximum clique of  $H$ . Clearly,  $s \leq \omega(X)$ . For every  $I \subseteq K$ , define

$$T_I = \{v \in V(H) \setminus V(K) : N_K(v) = I\}.$$

We show that  $T_I$  is homogeneous with respect to  $V(G) - A_7$ . First note that  $T_K = \emptyset$  by the choice of  $K$ . Suppose that  $T_\emptyset \neq \emptyset$ . Then by the connectivity of  $H$ , there is a vertex  $z \in T_\emptyset$  that has a neighbor  $y \in T_I$  for some  $I \neq \emptyset$ . Since  $y$  cannot be complete to  $K$ , there exist  $i, j$  such that  $y$  is adjacent to  $v_i$  but not to  $v_j$ . Then  $z, y, v_i, v_j$  induces a  $P_4$ , which contradicts the fact that  $A_6$  is  $P_4$ -free. So  $T_\emptyset = \emptyset$ .

Let  $I, J \subseteq K$  with  $I \neq J$ . If there exist  $i, j$  such that  $v_i \in I \setminus J$  and  $v_j \in J \setminus I$ , then  $T_I$  and  $T_J$  are complete: if  $t_i \in T_I$  and  $t_j \in T_J$  are not adjacent, then  $t_i, v_i, v_j, t_j$  induces a  $P_4$ . Now suppose that  $I \subseteq J$ . Since  $|I| < |J|$ , there exists an index  $j$  such that  $v_j \in J \setminus I$ . Since  $J \neq K$ , there is an index  $k$  such that  $v_k \notin J$ . Then if  $t_i \in T_I$  and  $t_j \in T_J$  are adjacent, then  $t_i, t_j, v_j, v_k$  induces a  $P_4$ . So  $T_I$  and  $T_J$  are anticomplete.

This proves that for each  $I \subseteq K$ ,  $T_I$  is homogeneous with respect to  $V(G) - A_7$ . Moreover,  $\omega(T_I) < \omega$ . By the inductive hypothesis, it follows that  $|T_I| \leq 2^{2(\omega-1)^2}$ . So

$$|H| = \sum_{I \subseteq K} |T_I| + |K| \leq 2^{2(\omega-1)^2} 2^\omega + \omega.$$

Thus,  $|X| \leq 2^{2(\omega-1)^2} 2^\omega + \omega + \omega^2 = 2^{2\omega^2 - 3\omega + 2} + (\omega + \omega^2)$ . Since  $\omega + \omega^2 \leq 2^{3\omega - 2}$ , it follows that  $|X| \leq 2^{2\omega^2}$ . This proves the claim.  $\square$

Applying Claim 1 to  $A_6$ , we have that  $|A_6| \leq 2^{2\omega^2(A_6)} \leq 2^{2k^2}$ . Finally, we bound  $A_7$ .

**Claim 2.**  $|A_7| \leq k2^{|A_6|}$ .

*Proof.* Recall that each component of  $A_7$  is a clique and so has size at most  $k$ . We now group the components by size and show that each group has at most  $2^{|A_6|}$  components. Suppose not. Then there are two components  $C$  and  $C'$  of  $A_7$  having the same size  $i$  with  $N(C) = N(C')$ , which contradicts Lemma 2. This proves the claim.  $\square$

Therefore,  $|G| = \sum_{i=1}^7 |A_i| \leq 5k + |A_6| + |A_7|$ , which is a function of  $k$ .  $\square$

### 3.2 $\overline{P_3 + P_2}$ -Free Graphs

**Theorem 5.** *For every fixed integer  $k \geq 1$ , there are finitely many  $k$ -vertex-critical  $(P_5, \overline{P_3 + P_2})$ -free graphs.*

*Proof.* Let  $G$  be a  $k$ -vertex-critical  $(P_5, \overline{P_3 + P_2})$ -free graph. We show that  $|G|$  is bounded by a function of  $k$ . If  $G$  contains a  $K_k$ , then  $G$  is isomorphic to  $K_k$  and thus  $|G| = k$ . So assume that  $G$  is  $K_k$ -free. Since  $G$  is imperfect,  $G$  contains an induced  $C_5$  by Theorem 3. Let  $Q = v_1, v_2, v_3, v_4, v_5$  be an induced  $C_5$ . For each  $1 \leq i \leq 5$ , we define

$$\begin{aligned} Z &= \{v \in V \setminus Q : N_Q(v) = \emptyset\}, \\ C_i &= \{v \in V \setminus Q : N_Q(v) = \{v_{i-1}, v_{i+1}\}\}, \\ Y_i &= \{v \in V \setminus Q : N_Q(v) = \{v_{i-2}, v_i, v_{i+2}\}\}, \\ T_i &= \{v \in V \setminus Q : N_Q(v) = \{v_{i-1}, v_i, v_{i+1}\}\}, \\ F_i &= \{v \in V \setminus Q : N_Q(v) = V(Q) \setminus \{v_i\}\}, \\ U &= \{v \in V \setminus Q : N_Q(v) = V(Q)\}. \end{aligned}$$

Let  $C = \cup_{1 \leq i \leq 5} C_i$ ,  $Y = \cup_{1 \leq i \leq 5} Y_i$ ,  $T = \cup_{1 \leq i \leq 5} T_i$  and  $F = \cup_{1 \leq i \leq 5} F_i$ . We choose  $Q$  so that  $|Y|$  is maximum.

**Claim 3.**  $V(G) = V(Q) \cup Z \cup C \cup Y \cup T \cup F \cup U$ .

*Proof.* Suppose that the claim is false. Then there is a vertex  $u$  such that  $\{v_i\} \subseteq N(u) \subseteq \{v_i, v_{i+1}\}$  for some  $1 \leq i \leq 5$ . Then  $u, v_i, v_{i+4}, v_{i+3}, v_{i+2}$  induces a  $P_5$ .  $\square$

We first bound  $U$ .

**Claim 4.** *For each  $1 \leq i \leq 5$ ,  $Y_i \cup U$  and  $F_i \cup U$  are cliques.*

*Proof.* Suppose that  $Y_i \cup U$  contains two nonadjacent vertices  $x$  and  $y$ , then  $\{x, y, v_{i-2}, v_i, v_{i+2}\}$  induces a  $\overline{P_3 + P_2}$ . The proof for  $F_i \cup U$  is completely analogous.  $\square$

By Claim 4, it follows that  $|U| < k$ .

**Claim 5.** *For each  $1 \leq i \leq 5$ ,  $C_i$ ,  $Y_i$  and  $F_i$  are independent sets.*

*Proof.* Note that for each vertex  $x \in C \cup Y \cup F$ , there is an index  $i$  such that  $x$  is not adjacent to  $v_i$  but adjacent to both  $v_{i+1}$  and  $v_{i-1}$ . If  $C_i$  contains two adjacent vertices  $x$  and  $y$ , then let  $j$  be such that  $x, y$  are not adjacent to  $v_j$  but adjacent to both  $v_{j-1}$  and  $v_{j+1}$ . Then  $\{x, y, v_{j-1}, v_j, v_{j+1}\}$  induces a  $\overline{P_3 + P_2}$ . The proof for  $Y_i$  and  $F_i$  is completely analogous.  $\square$

By Claim 4 and Claim 5,  $|Y_i| \leq 1$  and  $|F_i| \leq 1$ . So  $|Y| \leq 5$  and  $|F| \leq 5$ . Next we bound  $Z$ .

**Claim 6.**  $Z$  is anticomplete to  $C \cup T$ .

*Proof.* Let  $z \in Z$ . If  $z$  has a neighbor  $x \in C_i \cup T_i$ , then  $z, x, v_{i+1}, v_{i+2}, v_{i+3}$  induces a  $P_5$ .  $\square$

**Claim 7.** *Each vertex in  $Y \cup F$  is either complete or anticomplete to each component of  $Z$ .*

*Proof.* Let  $v \in Y \cup F$  be that  $v$  is adjacent to  $a$  but not adjacent to  $b$  with  $ab \in E(Z)$ . Note that there is an index  $i$  such that  $v, v_i, v_{i+1}$  is an induced  $P_3$ . Then  $b, a, v, v_i, v_{i+1}$  induces a  $P_5$ .  $\square$

**Claim 8.**  *$Z$  is an independent set.*

*Proof.* Suppose not. Let  $K$  be a component of  $Z$  containing an edge  $ab$ . By Claim 6,  $N(K) \subseteq Y \cup F \cup U$ . Since  $G$  has no clique cutset, it follows from Claim 7 and Claim 4 that there are two nonadjacent vertices  $x$  and  $y$  in  $Y \cup F$  that are complete to  $K$ . Let  $v_i \in V(Q)$  be a common neighbor of  $x$  and  $y$ . Then  $\{a, b, x, y, v_i\}$  induces a  $\overline{P_3 + P_2}$ .  $\square$

By Claim 6,  $N(z) \subseteq Y \cup F \cup U$  for each  $z \in Z$ . Note that  $|Y \cup F \cup U| \leq k + 10$  and so there are  $2^{10+k}$  possible neighborhoods for vertices in  $Z$ . By Lemma 2 and the pigeonhole principle,  $|Z| \leq 2^{10+k}$ .

Next we bound  $C$ .

**Claim 9.**  *$C_i$  and  $C_{i+1}$  are complete to each other.*

*Proof.* Let  $c_3 \in C_3$  and  $c_4 \in C_4$ . If  $c_3c_4 \notin E$ , then  $c_4, v_5, v_1, v_2, c_3$  induces a  $P_5$ .  $\square$

**Claim 10.**  *$C_i$  is complete to  $T_i$ .*

*Proof.* By symmetry, we prove for  $i = 1$ . If  $c_1 \in C_1$  and  $t_1 \in T_1$  are not adjacent, then  $\{c_1, t_1, v_5, v_1, v_2\}$  induces a  $\overline{P_3 + P_2}$ .  $\square$

**Claim 11.**  *$C_i$  is anticomplete to  $T_{i+1}$  and  $T_{i-1}$ .*

*Proof.* If  $c_1 \in C_1$  is adjacent to  $t_2 \in T_2$ , then  $\{v_5, c_1, v_1, v_2, t_2\}$  induces a  $\overline{P_3 + P_2}$ . The case for  $T_5$  is similar.  $\square$

**Claim 12.** *No vertex in  $T_{i+2}$  or  $T_{i-2}$  can be mixed on  $C_i$ .*

*Proof.* Suppose that  $t_3 \in T_3$  is mixed on  $C_1$ . Then there are vertices  $c_1, c'_1 \in C_1$  with  $t_3c_1 \in E$  and  $t_3c'_1 \notin E$ . Then  $c'_1, v_5, c_1, t_3, v_3$  induces a  $P_5$ . The case for  $T_4$  is similar.  $\square$

**Claim 13.** *For each  $1 \leq i \leq 5$ ,  $|C_i| \leq 2^{|Y \cup F \cup U|}$ .*

*Proof.* Suppose not. Then by the pigeonhole principle, there are two vertices  $c_1, c_2 \in C_1$  such that  $c_1$  and  $c_2$  have the same neighbors in  $Y \cup F \cup U$ . By Claim 6 and Claim 9-Claim 12, it follows that  $c_1$  and  $c_2$  have the same neighbors in  $Z \cup T \cup C_1 \cup C_2 \cup C_5$ . Since  $c_1$  and  $c_2$  are not comparable, there exist  $d_1 \in N(c_1) \setminus N(c_2)$  and  $d_2 \in N(c_2) \setminus N(c_1)$ . So  $d_1, d_2 \in C_3 \cup C_4$ . If  $d_1, d_2 \in C_3$ , then  $d_1, c_1, v_5, c_2, d_2$  induces a  $P_5$ . So we may assume that  $d_1 \in C_4$  and  $d_2 \in C_3$ .

Consider the induced five-cycle  $Q' = Q - \{v_1\} \cup \{c_1\}$ . Let  $Y'$  be the set of vertices in  $V(G) \setminus V(Q')$  whose neighborhood on  $Q'$  induces a  $P_2 + P_1$ . Note that  $d_1 \in Y' \setminus Y$ . By the choice of  $Q$ , there exists a vertex  $y \in Y \setminus Y'$ . Clearly,  $y \in Y_1 \cup Y_3 \cup Y_4$  and  $yc_1 \notin$

*E.* If  $y \in Y_1$ , then  $c_1, v_5, v_1, y, v_3$  induces a  $P_5$ . So  $y \in Y_3 \cup Y_4$ . By symmetry, we may assume that  $y \in Y_4$ . Note that  $yc_2 \notin E$  since otherwise  $\{y, c_2, v_1, v_2, v_5\}$  induces a  $\overline{P_3 + P_2}$ ,  $yd_1 \in E$  since otherwise  $y, v_1, v_5, d_1, v_3$  induces a  $P_5$ , and that  $yd_2 \notin E$  since otherwise  $\{y, d_1, d_2, v_2, v_3\}$  induces a  $\overline{P_3 + P_2}$ . But now  $c_2, d_2, d_1, y, v_1$  induces a  $P_5$ , a contradiction.  $\square$

By Claim 13,  $|C| \leq 5 \times 2^{k+10}$ . Finally, we bound  $T$ .

**Claim 14.** *For each  $1 \leq i \leq 5$ ,  $T_i$  is  $P_2 + P_1$ -free and thus induces a complete multipartite graph.*

*Proof.* If  $T_i$  contains three vertices  $x, y, z$  such that  $xy$  and  $z$  induce a disjoint union of an edge and a vertex, then  $\{x, y, z, v_{i-1}, v_{i+1}\}$  induces a  $\overline{P_3 + P_2}$ . It is well-known that a graph is a complete multipartite graph if and only if it is  $P_2 + P_1$ -free.  $\square$

**Claim 15.** *For each  $1 \leq i \leq 5$ ,  $T_i$  and  $T_{i+1}$  are complete.*

*Proof.* If  $t_1 \in T_1$  and  $t_2 \in T_2$  are not adjacent, then  $t_2, v_3, v_4, v_5, t_1$  induces a  $P_5$ .  $\square$

**Claim 16.**  $|T_i| \leq k2^{|Z \cup C \cup Y \cup F \cup U|}$ .

*Proof.* By Claim 14,  $T_1$  is a complete  $r$ -partite graph  $(X_1, \dots, X_r)$  for some  $r \leq k$  where  $X_i$  and  $X_j$  are complete for all  $1 \leq i < j \leq r$ . If  $|X_i| > 2^{|Z \cup C \cup Y \cup F \cup U|}$ , then by the pigeonhole principle there are two nonadjacent vertices  $x_1$  and  $x_2$  that have the same neighbors in  $Z \cup C \cup Y \cup F \cup U$ . Since  $x_1$  and  $x_2$  are not comparable, there are vertices  $y_1 \in N(x_1) \setminus N(x_2)$  and  $y_2 \in N(x_2) \setminus N(x_1)$ . By Claim 15,  $y_1, y_2 \in T_3 \cup T_4$ . Then  $v_4, y_1, x_1, v_1, x_2$  induces a  $P_5$ .  $\square$

By Claim 16,  $|T|$  is bounded by a function of  $k$  and so is  $|G|$ . This completes the proof.  $\square$

### 3.3 Complete Characterization for $k = 4$ and $k = 5$


We remark that the upper bound on the order of the  $k$ -vertex-critical graphs in Theorem 5 and Theorem 4 is far from being tight: the objective was to show the *existence* of such a bound and so we did not try to optimize the bound. This indicates that it may be possible to characterize all  $k$ -vertex-critical graphs for small values of  $k$ . In this subsection, we prove that this is indeed possible for  $k = 4$  and  $k = 5$ . In particular, we prove the following theorems.

**Theorem 6.** *There are exactly 3 4-vertex-critical  $(P_5, \text{gem})$ -free graphs and there are exactly 6 4-vertex-critical  $(P_5, \overline{P_3 + P_2})$ -free graphs.*

**Theorem 7.** *There are exactly 7 5-vertex-critical  $(P_5, \text{gem})$ -free graphs and there are exactly 20 5-vertex-critical  $(P_5, \overline{P_3 + P_2})$ -free graphs.*

*Proof of Theorem 6 and Theorem 7.* The proof uses the computer program for generating  $k$ -vertex-critical  $\mathcal{H}$ -free graphs developed in [17]. The program takes a positive integer  $k$  and a set  $\mathcal{H}$  of forbidden induced subgraphs as input and if it terminates, it outputs a complete list of  $k$ -vertex-critical  $\mathcal{H}$ -free graphs. The source code of this algorithm can be downloaded from [16]. We refer to [17] for more details about the generation algorithm. The correctness of this algorithm has extensively been tested in the literature, see [5, 7, 17] for example.

We extended the algorithm from [17] so it can also generate  $H$ -free graphs where  $H = \text{gem}$  or  $H = \overline{P_3 + P_2}$  and executed the algorithm in with the following inputs:



**Figure 3:** All 3 4-vertex-critical  $(P_5, \text{gem})$ -free graphs.

- $k = 4$  or  $k = 5$ ;
- $\mathcal{H} = \{P_5, \text{gem}\}$  or  $\mathcal{H} = \{P_5, \overline{P_3 + P_2}\}$ .

In each of these 4 cases the algorithm terminates within a few seconds and yields the graphs reported in Theorem 6, Theorem 7, and in Table 1.  $\square$

The counts of the vertex-critical graphs from Theorem 6 and Theorem 7 are listed in Table 1. The 3 4-vertex-critical  $(P_5, \text{gem})$ -free graphs from Theorem 6 are shown in Figure 3 and the 7 5-vertex-critical  $(P_5, \text{gem})$ -free graphs from Theorem 7 are shown in Figure 4.

The adjacency lists of all vertex-critical graphs from Theorem 6 and Theorem 7 can be found in the Appendix and these graphs also be inspected in the database of interesting graphs at the *House of Graphs* [2] by searching for the keywords “4-critical  $P_5$ Gem-free”, “5-critical  $P_5$ Gem-free”, “4-critical  $P_5$ co( $P_2+P_3$ )-free”, or “5-critical  $P_5$ co( $P_2+P_3$ )-free”.

Note that it follows from [3, 23] that there are exactly 12 4-vertex-critical  $P_5$ -free graphs. As an independent test for the correctness of our implementation, we took these 12 graphs and tested which of these graphs is also gem-free or  $\overline{P_3 + P_2}$ -free. This yielded exactly the same graphs as in Theorem 6. As another correctness test, we modified our program to generate all gem-free graphs and compared it with the known counts of gem-free graphs at the *On-Line Encyclopedia of Integer Sequences*<sup>1</sup>. Also here the results were in complete agreement.

Vertices	4	5	6	7	8	9	10	13	Total
4-vertex-critical $(P_5, \text{gem})$ -free graphs	1			1			1		3
5-vertex-critical $(P_5, \text{gem})$ -free graphs		1				3		3	7
4-vertex-critical $(P_5, \overline{P_3 + P_2})$ -free graphs	1		1	4					6
5-vertex-critical $(P_5, \overline{P_3 + P_2})$ -free graphs		1		1	4	14			20

**Table 1:** Counts of all  $k$ -vertex-critical  $(P_5, H)$ -free graphs for  $k \in \{4, 5\}$  and  $H \in \{\text{gem}, \overline{P_3 + P_2}\}$ .

<sup>1</sup><https://oeis.org/A079576>




Figure 4: All 7 5-vertex-critical  $(P_5, \text{gem})$ -free graphs.

## 4 Conclusion

In this paper we gave an affirmative answer to the problem posed in [5] for  $H \in \{\text{gem}, \overline{P_3 + P_2}\}$ . Moreover, we characterized all 4- and 5-vertex-critical  $(P_5, \text{gem})$ -free graphs and  $(P_5, \overline{P_3 + P_2})$ -free graphs by extending the computer generation algorithm from [17]. As future research it is natural to investigate the finiteness of  $k$ -vertex-critical graphs in the class of  $(P_5, H)$ -free graphs for other graphs  $H$  of order 5.

## References

- [1] J. A. Bondy and U. S. R. Murty. *Graph Theory*. Springer, 2008.
- [2] G. Brinkmann, K. Coolsaet, J. Goedgebeur, and H. Mélot. House of graphs: a database of interesting graphs. *Discrete Applied Mathematics*, 161:311–314, 2013. Available at <https://hog.grinvin.org/>.
- [3] D. Bruce, C. T. Hoàng, and J. Sawada. A certifying algorithm for 3-colorability of  $P_5$ -free graphs. In *Proceedings of the 20th International Symposium on Algorithms and Computation*, Lecture Notes in Computer Science 5878, pages 594–604, 2009.

- [4] Q. Cai, S. Huang, T. Li, and Y. Shi. Vertex-critical  $(P_5, \text{banner})$ -free graphs. In Yijia Chen, Xiaotie Deng, and Mei Lu, editors, *Frontiers in Algorithmics - 13th International Workshop, FAW 2019, Sanya, China, April 29–May 3, 2019, Proceedings*, volume 11458 of *Lecture Notes in Computer Science*, pages 111–120, 2019.
- [5] K. Cameron, J. Goedgebeur, S. Huang, and Y. Shi.  $k$ -critical graphs in  $P_5$ -free graphs. *Theoretical Computer Science*, 864:80–91, 2021.
- [6] K. Cameron, S. Huang, and O. Merkel. A bound for the chromatic number of  $(P_5, \text{gem})$ -free graphs. *Bulletin of the Australian Mathematical Society*, 100:182–188, 2019.
- [7] M. Chudnovsky, J. Goedgebeur, O. Schaudt, and M. Zhong. Obstructions for three-coloring graphs without induced paths on six vertices. *Journal of Combinatorial Theory, Series B*, 140:45–83, 2020.
- [8] M. Chudnovsky, T. Karthick, P. Maceli, and F. Maffray. Coloring graphs with no induced five-vertex path or gem. *Journal of Graph Theory*, 95:527–542, 2020.
- [9] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas. The strong perfect graph theorem. *Annals of Mathematics*, 164:51–229, 2006.
- [10] H. S. Dhaliwal, A. M. Hamel, C. T. Hoàng, F. Maffray, T. J. D. McConnell, and S. A. Panait. On color-critical  $(P_5, \text{co-}P_5)$ -free graphs. *Discrete Applied Mathematics*, 216:142–148, 2017.
- [11] G. A. Dirac. Note on the colouring of graphs. *Mathematische Zeitschrift*, 54:347–353, 1951.
- [12] G. A. Dirac. A property of 4-chromatic graphs and some remarks on critical graphs. *Journal of the London Mathematical Society*, 27:85–92, 1952.
- [13] G. A. Dirac. Some theorems on abstract graphs. *Proceedings of the London Mathematical Society*, 2:69–81, 1952.
- [14] T. Gallai. Kritische graphen I. *Publications of the Mathematical Institute of the Hungarian Academy of Sciences*, 8:165–92, 1963.
- [15] T. Gallai. Kritische graphen II. *Publications of the Mathematical Institute of the Hungarian Academy of Sciences*, 8:373–395, 1963.
- [16] J. Goedgebeur. Homepage of generator for  $k$ -critical  $\mathcal{H}$ -free graphs: <https://caagt.ugent.be/criticalpfree/>.
- [17] J. Goedgebeur and O. Schaudt. Exhaustive generation of  $k$ -critical  $\mathcal{H}$ -free graphs. *Journal of Graph Theory*, 87:188–207, 2018.
- [18] P. Hell and S. Huang. Complexity of coloring graphs without paths and cycles. *Discrete Applied Mathematics*, 216:211–232, 2017.
- [19] C. T. Hoàng, B. Moore, D. Recoskiez, J. Sawada, and M. Vatshelle. Constructions of  $k$ -critical  $P_5$ -free graphs. *Discrete Applied Mathematics*, 182:91–98, 2015.
- [20] S. Huang, T. Li, and Y. Shi. Critical  $(P_6, \text{banner})$ -free graphs. *Discrete Applied Mathematics*, 258:143–151, 2019.

- [21] M. Kamiński and A. Pstrucha. Certifying coloring algorithms for graphs without long induced paths. *Discrete Applied Mathematics*, 261:258–267, 2019.
- [22] A. V. Kostochka and M. Yancey. Ore’s conjecture on color-critical graphs is almost true. *Journal of Combinatorial Theory, Series B*, 109:73–101, 2014.
- [23] F. Maffray and G. Morel. On 3-colorable  $P_5$ -free graphs. *SIAM Journal on Discrete Mathematics*, 26:1682–1708, 2012.
- [24] O. Ore. *The Four Color Problem*. Academic Press, 1967.

## Appendix

Below we list the adjacency lists of all vertex-critical graphs from Theorem 6 and Theorem 7. These graphs can also be inspected in the database of interesting graphs at the *House of Graphs* [2] by searching for the keywords “4-critical P5Gem-free”, “5-critical P5Gem-free”, “4-critical P5co(P2+P3)-free”, and “5-critical P5co(P2+P3)-free”.

### Adjacency lists of 4-vertex-critical ( $P_5$ ,gem)-free graphs

1. {0: 1 2 3; 1: 0 2 3; 2: 0 1 3; 3: 0 1 2}
2. {0: 1 4 5; 1: 0 2 5 6; 2: 1 3 6; 3: 2 4 6; 4: 0 3 5; 5: 0 1 4; 6: 1 2 3}
3. {0: 1 4 5; 1: 0 2 5 6 7; 2: 1 3 7; 3: 2 4 6; 4: 0 3 5 6 7; 5: 0 1 4; 6: 1 3 4 8 9; 7: 1 2 4 8 9; 8: 6 7 9; 9: 6 7 8}

### Adjacency lists of 5-vertex-critical ( $P_5$ ,gem)-free graphs

1. {0: 1 2 3 4; 1: 0 2 3 4; 2: 0 1 3 4; 3: 0 1 2 4; 4: 0 1 2 3}
2. {0: 1 4 5 6; 1: 0 2 5 6 7 8; 2: 1 3 7 8; 3: 2 4 7 8; 4: 0 3 5 6; 5: 0 1 4 6; 6: 0 1 4 5; 7: 1 2 3 8; 8: 1 2 3 7}
3. {0: 1 4 5 6; 1: 0 2 5 6; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 7 8; 5: 0 1 4 6; 6: 0 1 2 5; 7: 2 3 4 8; 8: 2 3 4 7}
4. {0: 1 4 5 6 7; 1: 0 2 5 6 8; 2: 1 3 6 8; 3: 2 4 7 8; 4: 0 3 5 7; 5: 0 1 4 6 7; 6: 0 1 2 5 8; 7: 0 3 4 5; 8: 1 2 3 6}
5. {0: 1 4 5 6; 1: 0 2 5 6 7 8 12; 2: 1 3 7 12; 3: 2 4 6 7 8 12; 4: 0 3 5 8; 5: 0 1 4 6; 6: 0 1 3 5 9 10 11; 7: 1 2 3 12; 8: 1 3 4 9 10 11; 9: 6 8 10 11; 10: 6 8 9 11; 11: 6 8 9 10; 12: 1 2 3 7}
6. {0: 1 4 5 6 7 8; 1: 0 2 5 6; 2: 1 3 6 7 8 12; 3: 2 4 8 12; 4: 0 3 5 7 12; 5: 0 1 4 6 7 8; 6: 0 1 2 5; 7: 0 2 4 5 9 10 11; 8: 0 2 3 5 9 10 11 12; 9: 7 8 10 11; 10: 7 8 9 11; 11: 7 8 9 10; 12: 2 3 4 8}
7. {0: 1 4 5 6 7 8; 1: 0 2 5 6 12; 2: 1 3 6 7 8 12; 3: 2 4 8 12; 4: 0 3 5 7; 5: 0 1 4 6 7 8; 6: 0 1 2 5 12; 7: 0 2 4 5 9 10 11 12; 8: 0 2 3 5 9 10 11 12; 9: 7 8 10 11; 10: 7 8 9 11; 11: 7 8 9 10; 12: 1 2 3 6 7 8}

**Adjacency lists of 4-vertex-critical  $(P_5, \overline{P_3 + P_2})$ -free graphs**

1. {0: 1 2 3; 1: 0 2 3; 2: 0 1 3; 3: 0 1 2}
2. {0: 1 4 5; 1: 0 2 5; 2: 1 3 5; 3: 2 4 5; 4: 0 3 5; 5: 0 1 2 3 4}
3. {0: 1 4 5; 1: 0 2 5 6; 2: 1 3 6; 3: 2 4 6; 4: 0 3 5; 5: 0 1 4; 6: 1 2 3}
4. {0: 1 4 5 6; 1: 0 2 5 6; 2: 1 3 5; 3: 2 4 6; 4: 0 3 5; 5: 0 1 2 4; 6: 0 1 3}
5. {0: 1 4 5 6; 1: 0 2 5 6; 2: 1 3 5 6; 3: 2 4 6; 4: 0 3 5; 5: 0 1 2 4; 6: 0 1 2 3}
6. {0: 1 4 5 6; 1: 0 2 5 6; 2: 1 3 6; 3: 2 4 6; 4: 0 3 5; 5: 0 1 4; 6: 0 1 2 3}

**Adjacency lists of 5-vertex-critical  $(P_5, \overline{P_3 + P_2})$ -free graphs**

1. {0: 1 2 3 4; 1: 0 2 3 4; 2: 0 1 3 4; 3: 0 1 2 4; 4: 0 1 2 3}
2. {0: 1 4 5 6; 1: 0 2 5 6; 2: 1 3 5 6; 3: 2 4 5 6; 4: 0 3 5 6; 5: 0 1 2 3 4 6; 6: 0 1 2 3 4 5}
3. {0: 1 4 5 6 7; 1: 0 2 5 6 7; 2: 1 3 5 7; 3: 2 4 6 7; 4: 0 3 5 7; 5: 0 1 2 4 7; 6: 0 1 3 7; 7: 0 1 2 3 4 5 6}
4. {0: 1 4 5 6 7; 1: 0 2 5 6 7; 2: 1 3 6 7; 3: 2 4 6 7; 4: 0 3 5 6; 5: 0 1 4 6; 6: 0 1 2 3 4 5 7; 7: 0 1 2 3 6}
5. {0: 1 4 5 6 7; 1: 0 2 5 6 7; 2: 1 3 5 6 7; 3: 2 4 6 7; 4: 0 3 5 7; 5: 0 1 2 4 7; 6: 0 1 2 3 7; 7: 0 1 2 3 4 5 6}
6. {0: 1 4 5 6; 1: 0 2 5 6 7; 2: 1 3 6 7; 3: 2 4 6 7; 4: 0 3 5 6; 5: 0 1 4 6; 6: 0 1 2 3 4 5 7; 7: 1 2 3 6}
7. {0: 1 4 5 6; 1: 0 2 5 6 7 8; 2: 1 3 7 8; 3: 2 4 7 8; 4: 0 3 5 6; 5: 0 1 4 6; 6: 0 1 4 5; 7: 1 2 3 8; 8: 1 2 3 7}
8. {0: 1 4 5 6 7 8; 1: 0 2 5 6 7 8; 2: 1 3 6 7; 3: 2 4 7 8; 4: 0 3 5 6 8; 5: 0 1 4 6 8; 6: 0 1 2 4 5; 7: 0 1 2 3 8; 8: 0 1 3 4 5 7}
9. {0: 1 4 5 6 7 8; 1: 0 2 5 6 7; 2: 1 3 5 6 8; 3: 2 4 6 7 8; 4: 0 3 5 7 8; 5: 0 1 2 4 8; 6: 0 1 2 3 7 8; 7: 0 1 3 4 6; 8: 0 2 3 4 5 6}
10. {0: 1 4 5 6; 1: 0 2 5 6; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 7 8; 5: 0 1 4 6; 6: 0 1 2 5; 7: 2 3 4 8; 8: 2 3 4 7}
11. {0: 1 4 5 6 7; 1: 0 2 5 6 8; 2: 1 3 6 8; 3: 2 4 7 8; 4: 0 3 5 7; 5: 0 1 4 6 7; 6: 0 1 2 5 8; 7: 0 3 4 5; 8: 1 2 3 6}
12. {0: 1 4 5 6 7; 1: 0 2 5 6 7; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 7 8; 5: 0 1 4 6 8; 6: 0 1 2 5 8; 7: 0 1 2 3 4; 8: 2 3 4 5 6}
13. {0: 1 4 5 6 7; 1: 0 2 5 6 7; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 7 8; 5: 0 1 4 6; 6: 0 1 2 5; 7: 0 1 2 3 4 8; 8: 2 3 4 7}
14. {0: 1 4 5 6 7 8; 1: 0 2 5 6 7; 2: 1 3 6 7; 3: 2 4 7 8; 4: 0 3 5 7 8; 5: 0 1 4 6 8; 6: 0 1 2 5 8; 7: 0 1 2 3 4; 8: 0 3 4 5 6}

15. {0: 1 4 5 6 7 8; 1: 0 2 5 6 7; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 7 8; 5: 0 1 4 6; 6: 0 1 2 5 8; 7: 0 1 2 3 4 8; 8: 0 2 3 4 6 7}
16. {0: 1 4 5 6 7 8; 1: 0 2 5 6 7 8; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 7; 5: 0 1 4 6; 6: 0 1 2 5 8; 7: 0 1 2 3 4 8; 8: 0 1 2 3 6 7}
17. {0: 1 4 5 6 7; 1: 0 2 5 6 7 8; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 6; 5: 0 1 4 6; 6: 0 1 2 4 5; 7: 0 1 2 3 8; 8: 1 2 3 7}
18. {0: 1 4 5 6 7; 1: 0 2 5 6 7 8; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 6; 5: 0 1 4 6 8; 6: 0 1 2 4 5; 7: 0 1 2 3 8; 8: 1 2 3 5 7}
19. {0: 1 4 5 6 8; 1: 0 2 5 6 7 8; 2: 1 3 6 7; 3: 2 4 6 7; 4: 0 3 5 6 8; 5: 0 1 4 6 8; 6: 0 1 2 3 4 5; 7: 1 2 3 8; 8: 0 1 4 5 7}
20. {0: 1 4 5 6; 1: 0 2 5 6 7 8; 2: 1 3 6 7 8; 3: 2 4 7 8; 4: 0 3 5 6; 5: 0 1 4 6; 6: 0 1 2 4 5; 7: 1 2 3 8; 8: 1 2 3 7}