

# The sustainability probability for the critical Derrida–Retaux model

Xinxing Chen,<sup>1</sup> Yueyun Hu,<sup>2</sup> and Zhan Shi<sup>3</sup>

**Summary.** We are interested in the recursive model  $(Y_n, n \geq 0)$  studied by Collet, Eckmann, Glaser and Martin [9] and by Derrida and Retaux [12]. We prove that at criticality, the probability  $\mathbf{P}(Y_n > 0)$  behaves like  $n^{-2+o(1)}$  as  $n$  goes to infinity; this gives a weaker confirmation of predictions made in [9], [12] and [6]. Our method relies on studying the number of pivotal vertices and open paths, combined with a delicate coupling argument.

**Keywords.** Derrida–Retaux model, sustainability probability, pivotal vertex, open path.

**2010 Mathematics Subject Classification.** 60J80, 82B27.

## 1 Introduction

Let us start with a simple example.

**Example 1.1.** Let  $Y_0$  be a random variable with  $\mathbf{P}(Y_0 = 2) = \frac{1}{5} = 1 - \mathbf{P}(Y_0 = 0)$ . Consider the recurrence relation as follows: for all  $n \geq 0$ ,

$$Y_{n+1} = (Y_n + \hat{Y}_n - 1)^+,$$

---

<sup>1</sup>School of Mathematical Sciences, Shanghai Jiaotong University, 200240 Shanghai, China, [chenxinx@sjtu.edu.cn](mailto:chenxinx@sjtu.edu.cn) Partially supported by NSFC grants 11771286 and 11531001.

<sup>2</sup>LAGA, Université Sorbonne Paris Nord, 99 av. J-B Clément, F-93430 Villetaneuse, France, [yueyun@math.univ-paris13.fr](mailto:yueyun@math.univ-paris13.fr) Partially supported by ANR MALIN and ANR SWIWS.

<sup>3</sup>LPSM, Sorbonne Université Paris VI, 4 place Jussieu, F-75252 Paris Cedex 05, France, [zhan.shi@sorbonne-universite.fr](mailto:zhan.shi@sorbonne-universite.fr) Partially supported by ANR MALIN.

where  $\widehat{Y}_n$  is an independent copy of  $Y_n$ , and  $x^+ := \max\{x, 0\}$  for all  $x \in \mathbb{R}$ . Then, from the physics literature, one expects

$$(1.1) \quad \mathbf{P}(Y_n > 0) \sim \frac{4}{n^2},$$

$$(1.2) \quad \mathbf{E}(Y_n) \sim \frac{8}{n^2}, \quad n \rightarrow \infty.$$

[Notation:  $a_n \sim b_n$  means  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1$ .] See [9], [12] and [6] for explanations of (1.1), and [6] for explanations of (1.2).

However, from a mathematical point of view, (1.1) and (1.2) are mere conjectures. The only relevant rigorous result (see [6]) available is:

$$\limsup_{n \rightarrow \infty} n \mathbf{P}(Y_n > 0) < \infty.$$

The main result of our paper, stated in Theorem 1.2 below, will imply that

$$\mathbf{P}(Y_n > 0) = \frac{1}{n^{2+o(1)}}, \quad \mathbf{E}(Y_n) = \frac{1}{n^{2+o(1)}}, \quad n \rightarrow \infty.$$

We are going to see that this is valid for a large family of initial distributions of  $Y_0$ .  $\square$

Throughout the paper, we fix an integer  $m \geq 2$ , and let  $X_0 \geq 0$  be a random variable taking values in  $\mathbb{Z}_+ := \{0, 1, 2, \dots\}$  and satisfying  $\mathbf{E}(X_0 m^{X_0}) < \infty$ . We consider the following recurrence relation: for all  $n \geq 0$ ,

$$(1.3) \quad X_{n+1} = (X_n^{(1)} + \dots + X_n^{(m)} - 1)^+,$$

where  $X_n^{(i)}$ ,  $i \geq 1$ , are independent copies of  $X_n$ . The assumption  $\mathbf{E}(X_0 m^{X_0}) < \infty$  ensures that  $\mathbf{E}(X_n m^{X_n}) < \infty$  for all  $n \geq 0$ . The basic question is how the distribution of  $X_n$  behaves when  $n$  is sufficiently large.

The random recursive system  $(X_n, n \geq 0)$ , referred to as the Derrida–Retaux system or the Derrida–Retaux model, has been investigated by Derrida and Retaux [12] as a simplified hierarchical renormalization model in order to understand the depinning transition of a line in presence of strong disorder. Pinning models have an intensive literature both in mathematics and in theoretical physics; see for example [2], [3], [14], [15], [16], [17], and [11], [12], [23], [24]. We refer to [12] and [13] for more references and explanations of the recursion (1.3). The Derrida–Retaux system has also appeared in other contexts: in Collet et al. [9] as a spin glass

model, in Li and Rogers [22] as an iteration function of random variables, in Goldschmidt and Przykucki [18] and Curien and Hénard [10] as a parking scheme; it also belongs to one of the max-type recursion families in the seminal paper by Aldous and Bandyopadhyay [1].

The system is known for exhibiting a phase transition:<sup>4</sup>

**Theorem A. (Collet et al. [9])** *Assume  $\mathbf{E}(X_0 m^{X_0}) < \infty$ .*

(i) *If  $(m - 1)\mathbf{E}(X_0 m^{X_0}) > \mathbf{E}(m^{X_0})$ , then  $\lim_{n \rightarrow \infty} \frac{\mathbf{E}(X_n)}{m^n}$  exists and is positive.*

(ii) *If  $(m - 1)\mathbf{E}(X_0 m^{X_0}) \leq \mathbf{E}(m^{X_0})$ , then  $\sup_{n \geq 0} \mathbf{E}(X_n) \leq \frac{1}{m-1}$ .*

According to Theorem A, there is a dichotomy for the asymptotic behaviour of  $\mathbf{E}(X_n)$ : it either grows exponentially fast as  $n \rightarrow \infty$ , or is always bounded by the constant  $\frac{1}{m-1}$ . The system  $(X_n, n \geq 0)$  is said to be supercritical if  $(m - 1)\mathbf{E}(X_0 m^{X_0}) > \mathbf{E}(m^{X_0})$ , critical if  $(m - 1)\mathbf{E}(X_0 m^{X_0}) = \mathbf{E}(m^{X_0}) < \infty$ , and subcritical if  $(m - 1)\mathbf{E}(X_0 m^{X_0}) < \mathbf{E}(m^{X_0})$ .

The present paper focuses on the critical case:  $(m - 1)\mathbf{E}(X_0 m^{X_0}) = \mathbf{E}(m^{X_0}) < \infty$ . In order to insist on the criticality, we write  $(Y_n, n \geq 0)$  instead of  $(X_n, n \geq 0)$  for the critical Derrida–Retaux system. It is known (see [9], [6]) that

$$\mathbf{P}(Y_n \geq 1) \rightarrow 0, \quad n \rightarrow \infty.$$

We study the rate of decay of  $\mathbf{P}(Y_n \geq 1)$ .

**Theorem 1.2.** *If  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ , then for  $n \rightarrow \infty$ ,*

$$\mathbf{P}(Y_n \geq 1) = \frac{1}{n^{2+o(1)}}, \quad \mathbf{E}(Y_n) = \frac{1}{n^{2+o(1)}}.$$

More precisely, our Proposition 4.1 implies that if  $\mathbf{E}(Y_0^5 m^{Y_0}) < \infty$ , then  $\mathbf{E}(Y_n)$  is bounded from above by a constant multiple of  $\frac{(\log n)^2}{n^2}$  for all  $n \geq 2$ , whereas Proposition 5.1 yields that if  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ , then  $\mathbf{P}(Y_n \geq 1)$  is bounded from below by a constant multiple of  $\frac{1}{n^{2(\log n)^\nu}}$  for some  $\nu > 0$  and all  $n \geq 2$ .

The problem of evaluating  $\mathbf{P}(Y_n \geq 1)$  was discussed in [9], [12] and [6], and  $\mathbf{E}(Y_n)$  in [6]; it was expected, without precision on integrability condition on  $Y_0$ , that

$$(1.4) \quad \mathbf{P}(Y_n \geq 1) \sim \frac{c_*}{n^2}, \quad \mathbf{E}(Y_n) \sim \frac{c_{**}}{n^2}, \quad n \rightarrow \infty,$$

---

<sup>4</sup>It is crucial to assume that in the Derrida–Retaux model,  $X_0$  is integer-valued; it is an open problem to extend Theorem A without this assumption. See Derrida and Retaux [12] for more discussions.

with  $c_* := \frac{4}{(m-1)^2}$  and  $c_{**} := \frac{m}{m-1} c_*$ . In [4], it was realized that (1.4) should be valid only under the assumption  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ . When the latter condition fails,  $\mathbf{P}(Y_n \geq 1)$  and  $\mathbf{E}(Y_n)$  should behave differently. For example, for  $m = 2$ , if  $\mathbf{P}(Y_0 = k) \sim c_0 k^{-\alpha} 2^{-k}$ ,  $k \rightarrow \infty$ , for some  $\alpha \in (2, 4]$  (so  $\mathbf{E}(Y_0^3 m^{Y_0}) = \infty$ ) and  $c_0 \in (0, \infty)$ , then it is expected (see [13]) that

$$(1.5) \quad \mathbf{P}(Y_n \geq 1) \sim \frac{c(\alpha)}{n^2}, \quad n \rightarrow \infty$$

with  $c(\alpha) := \frac{\alpha(\alpha-2)}{2}$ . For discussions on other related questions, see [13].

Let us also mention that some continuous-time systems were studied in [12], [19] and [5], for which the analogue of the asymptotic equivalences of  $\mathbf{P}(Y_n \geq 1)$  was proved; see [12] and [19] for the analogue of (1.4), and [5] for the analogue of (1.5).

When a Derrida–Retaux system  $(X_n, n \geq 0)$  is subcritical, it is not hard to prove that  $\mathbf{P}(X_n \geq 1)$  and  $\mathbf{E}(X_n)$  decay exponentially (as  $n \rightarrow \infty$ ) if  $X_0$  satisfies an appropriate integrability condition.

The proof of Theorem 1.2 relies on the study of the important notions of *pivotal vertices* and *open paths* which we define in Section 2; these notions are closely related. The main advantage of studying the expected number of pivotal vertices is that it satisfies a nice identity. This is Theorem 2.2, valid for any Derrida–Retaux system (regardless of whether it is supercritical, critical or subcritical) and useful in the proof of both upper and lower bounds in Theorem 1.2. In the proof of the upper bound in Theorem 1.2, only a simple consequence (inequality (2.13)) of Theorem 2.2 is needed a couple of times. The proof of the lower bound relies on the full strength of Theorem 2.2 and goes actually beyond, in the sense that pivotal vertices are studied in a more complicated setting, and that the key step in the proof consists in establishing a weaker form of inequality (2.13) in the opposite direction.

The notion of open paths was already introduced in [4], and was known to be an important tool in the study of the Derrida–Retaux model. From methodological point of view, an interesting contribution of the paper is to introduce the notion of pivotal vertices, and to prove Theorem 2.2 on the expected number of pivotal vertices. Another notable contribution of the paper consists in studying the Derrida–Retaux system at intermediate generations. The idea of studying intermediate generations is quite natural in statistical physics; our in-depth descriptions (in Sections 4 and 5) of Derrida–Retaux systems at appropriate intermediate generations are among the key ingredients in the proof of Theorem 1.2. These descriptions require a careful analysis of open paths and pivotal vertices and of the moment

generating function.

The rest of the paper is as follows. In Section 2, we introduce pivotal vertices and open paths, and prove the aforementioned Theorem 2.2. Section 3 collects some preliminary estimates for the moment generating function of critical Derrida–Retaux systems; these estimates are either known, or are routinely proved by means of existing arguments. The upper and lower bounds in Theorem 1.2 are proved in Sections 4 and 5, respectively.

## 2 Pivotal vertices and open paths

Let  $(X_n, n \geq 0)$  be a Derrida–Retaux system defined via (1.3). There is a natural hierarchical representation of the system (see [8], [12], [20]).

Let us consider a family of random variables  $(X(v), v \in \mathbb{T})$ , indexed by a (reversed)  $m$ -ary tree  $\mathbb{T}$ , defined as follows. For any vertex  $v$  in the genealogical tree  $\mathbb{T}$ , let  $|v|$  denote the generation of  $v$  (so  $|v| = 0$  if the vertex  $v$  is in the initial generation). Let  $X(v)$ , for  $v \in \mathbb{T}$  with  $|v| = 0$ , be independent and identically distributed (i.i.d.) having the law of  $X_0$ . For any  $v \in \mathbb{T}$  with  $|v| \geq 1$ , let  $v^{(1)}, \dots, v^{(m)}$  denote the  $m$  parents of  $v$  in generation  $|v| - 1$ , and define

$$X(v) := (X(v^{(1)}) + \dots + X(v^{(m)}) - 1)^+.$$

By definition, for any  $n \geq 0$ , the random variables  $X(v)$ , for  $v \in \mathbb{T}$  with  $|v| = n$ , are i.i.d. having the law of  $X_n$ .

For  $n \geq 0$ , let  $\epsilon_n$  denote the first lexicographic vertex in the  $n$ -th generation of  $\mathbb{T}$ . Let  $\mathbb{T}_n$  denote the (reversed) subtree formed by all the ancestors (including  $\epsilon_n$  itself) of  $\epsilon_n$  in the first  $n$  generations. See Figure 1 for an example.

For  $v \in \mathbb{T}$  with  $|v| = 0$  and integer  $n \geq 0$ , we write  $v_n$  for the unique descendant of  $v$  in generation  $n$ , and call  $(v = v_0, v_1, v_2, \dots, v_n)$  the path in  $\mathbb{T}$  from  $v$  leading to  $v_n$ ,<sup>5</sup> for any integer  $n \geq 0$ , let  $\text{bro}(v_n)$  be the set of the “brothers” of  $v_n$ , i.e., the parents of  $v_{n+1}$  that are not  $v_n$ . Let

$$(2.1) \quad \xi_n(v) := \sum_{y \in \text{bro}(v_n)} X(y).$$

---

<sup>5</sup>Degenerate case: when  $|v| = 0$ , the path from  $v$  to  $v$  is reduced to the singleton  $v$ .




Figure 1: An example of  $\mathbb{T}_4$  with  $m = 2$ . Each vertex is represented by a square box; the number in the box represents the value of the Derrida–Retaux system at the vertex.

The path  $(v = v_0, v_1, v_2, \dots, v_n)$  is called *open* if

$$X(v) + \xi_0(v) + \xi_1(v) + \dots + \xi_i(v) \geq i + 1, \quad \forall 0 \leq i \leq n - 1.$$

Equivalently, the path is open if for  $u := v_i$  (for any  $1 \leq i \leq n$ ), we have  $X(u) = X(u^{(1)}) + \dots + X(u^{(m)}) - 1$  (with  $u^{(1)}, \dots, u^{(m)}$  denoting as before the parents of  $u$ ),<sup>6</sup> i.e.,  $X(u^{(1)}) + \dots + X(u^{(m)}) \geq 1$ .

For  $v \in \mathbb{T}$  and integer  $i \geq 0$ , let

$$N^{(i)}(v) := \text{number of open paths leading to } v,$$

$$(2.2) \quad \text{and starting from a vertex } u \text{ with } |u| = 0 \text{ such that } X(u) = i.$$

In particular,  $N^{(i)}(v) = \mathbf{1}_{\{X(v)=i\}}$  if  $|v| = 0$ .

For  $n \geq 0$ , we write

$$(2.3) \quad N_n^{(i)} := N^{(i)}(\epsilon_n), \quad N_n := \sum_{i=0}^{\infty} N_n^{(i)}, \quad X_n := X(\epsilon_n).$$

[The identification of  $X_n$  with  $X(\epsilon_n)$  should cause no confusion because only the individual law of each  $X_n$  matters.] See Figure 2 for an example.

**Definition 2.1.** Let  $v$  be a vertex with  $|v| = 0$ . For  $k, n \geq 0$  and  $\ell \geq 0$ , we say that  $v$  is a  $(k, \ell)_n$ -**pivotal vertex** if  $k + \xi_0(v) + \xi_1(v) + \dots + \xi_{n-1}(v) = n + \ell$  and

$$k + \xi_0(v) + \xi_1(v) + \dots + \xi_i(v) \geq i + 1, \quad \forall 0 \leq i \leq n - 1.$$

---

<sup>6</sup>Degenerate case: for  $|v| = 0$ , the path consisting of the singleton  $v$  is considered as open.



**Theorem 2.2.** For  $n \geq 0$  and  $\ell \geq 0$ ,

$$(2.9) \quad \sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) = \mu_n(\ell).$$

*Proof.* We prove (2.9) by induction on  $n$ . When  $n = 0$ , (2.9) holds since  $S_0^{(k,\ell)} = \mathbf{1}_{\{k=\ell\}}$ . Assume (2.9) holds for  $n$ . Let us prove it for  $n + 1$ . By the definition of pivotal vertices, for any  $k, \ell \geq 0$  and  $v \in \mathbb{T}_n$  with  $|v| = 0$ ,

$$(2.10) \quad \mathbf{1}_{\{v \in A_{n+1}^{(k,\ell)}\}} = \sum_{j=0}^{\ell+1} \mathbf{1}_{\{\xi_n(v)=j\}} \mathbf{1}_{\{v \in A_n^{(k,\ell+1-j)}\}}.$$

By symmetry,  $\mathbf{E}(S_{n+1}^{(k,\ell)}) = \sum_{v \in \mathbb{T}_{n+1}: |v|=0} \mathbf{E}(\mathbf{1}_{\{v \in A_{n+1}^{(k,\ell)}\}}) = m^{n+1} \mathbf{E}(\mathbf{1}_{\{\epsilon_0 \in A_{n+1}^{(k,\ell)}\}})$ . Putting these identities together, we get

$$\mathbf{E}(S_{n+1}^{(k,\ell)}) = m^{n+1} \sum_{j=0}^{\ell+1} \mathbf{E}(\mathbf{1}_{\{\xi_n(\epsilon_0)=j\}} \mathbf{1}_{\{\epsilon_0 \in A_n^{(k,\ell+1-j)}\}}).$$

Since  $\{\xi_n(\epsilon_0) = j\}$  and  $\{\epsilon_0 \in A_n^{(k,\ell+1-j)}\}$  are independent events, this leads to:

$$\begin{aligned} \mathbf{E}(S_{n+1}^{(k,\ell)}) &= m^{n+1} \sum_{j=0}^{\ell+1} \mathbf{P}(\xi_n(\epsilon_0) = j) \mathbf{E}(\mathbf{1}_{\{\epsilon_0 \in A_n^{(k,\ell+1-j)}\}}) \\ &= m \sum_{j=0}^{\ell+1} \mathbf{P}(\xi_n(\epsilon_0) = j) \mathbf{E}(S_n^{(k,\ell+1-j)}), \end{aligned}$$

which implies that<sup>7</sup>

$$\begin{aligned} \sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_{n+1}^{(k,\ell)}) &= m \sum_{k=0}^{\infty} \mu_0(k) \sum_{j=0}^{\ell+1} \mathbf{P}(\xi_n(\epsilon_0) = j) \mathbf{E}(S_n^{(k,\ell+1-j)}) \\ &= m \sum_{j=0}^{\ell+1} \mathbf{P}(\xi_n(\epsilon_0) = j) \sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell+1-j)}). \end{aligned}$$

---

<sup>7</sup>We can exchange the order of the sums because  $\sum_{k=0}^{\infty}$  is a finite sum:  $\mathbf{E}(S_{n+1}^{(k,\ell)}) = 0$  for  $k > (n+1) + \ell$ .



By the induction assumption, we get that  $\sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k, \ell+1-j)}) = \mu_n(\ell+1-j)$  (for  $j \leq \ell+1$ ), which is  $m^{-(\ell+1-j)} \mathbf{E}[(1 - (m-1)X_n)m^{X_n} \mathbf{1}_{\{X_n \leq \ell+1-j\}}]$ . So we have

$$\begin{aligned}
\sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_{n+1}^{(k, \ell)}) &= m^{-\ell} \sum_{j=0}^{\ell+1} m^j \mathbf{P}(\xi_n(\mathbf{e}_0) = j) \mathbf{E}[(1 - (m-1)X_n)m^{X_n} \mathbf{1}_{\{X_n \leq \ell+1-j\}}] \\
&= m^{-\ell} \mathbf{E}[(1 - (m-1)X_n)m^{X_n + \xi_n(\mathbf{e}_0)} \mathbf{1}_{\{X_n + \xi_n(\mathbf{e}_0) \leq \ell+1\}}] \\
(2.11) \quad &= m^{-\ell} \mathbf{E}[(1 - (m-1)X_n^{(1)})m^{X_n^{(1)} + \dots + X_n^{(m)}} \mathbf{1}_{\{X_n^{(1)} + \dots + X_n^{(m)} \leq \ell+1\}}],
\end{aligned}$$

the last equality being a consequence of the fact that the two-dimensional random vector  $(X_n, \xi_n(\mathbf{e}_0))$ , which is  $(X(\mathbf{e}_n), \xi_n(\mathbf{e}_0))$  by definition, has the same distribution as  $(X_n^{(1)}, X_n^{(2)} + \dots + X_n^{(m)})$ .

On the other hand, by symmetry,

$$\begin{aligned}
&\mathbf{E}[(1 - (m-1)X_{n+1})m^{X_{n+1}} \mathbf{1}_{\{X_{n+1} \leq \ell\}}] - \mathbf{P}(X_n^{(1)} + \dots + X_n^{(m)} = 0) \\
&= \sum_{u=1}^{\ell+1} \mathbf{E}[(1 - (m-1)(X_n^{(1)} + \dots + X_n^{(m)} - 1))m^{u-1} \mathbf{1}_{\{X_n^{(1)} + \dots + X_n^{(m)} = u\}}] \\
&= \sum_{u=1}^{\ell+1} \mathbf{E}[(1 - (m-1)(mX_n^{(1)} - 1))m^{u-1} \mathbf{1}_{\{X_n^{(1)} + \dots + X_n^{(m)} = u\}}] \\
&= \sum_{u=1}^{\ell+1} \mathbf{E}[(m - m(m-1)X_n^{(1)})m^{u-1} \mathbf{1}_{\{X_n^{(1)} + \dots + X_n^{(m)} = u\}}] \\
&= \sum_{u=1}^{\ell+1} \mathbf{E}[(1 - (m-1)X_n^{(1)})m^u \mathbf{1}_{\{X_n^{(1)} + \dots + X_n^{(m)} = u\}}].
\end{aligned}$$

Hence

$$\begin{aligned}
&\mathbf{E}[(1 - (m-1)X_{n+1})m^{X_{n+1}} \mathbf{1}_{\{X_{n+1} \leq \ell\}}] \\
&= \mathbf{E}[(1 - (m-1)X_n^{(1)})m^{X_n^{(1)} + \dots + X_n^{(m)}} \mathbf{1}_{\{X_n^{(1)} + \dots + X_n^{(m)} \leq \ell+1\}}].
\end{aligned}$$

Going back to (2.11), we get

$$\sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_{n+1}^{(k, \ell)}) = m^{-\ell} \mathbf{E}[(1 - (m-1)X_{n+1})m^{X_{n+1}} \mathbf{1}_{\{X_{n+1} \leq \ell\}}],$$

which is the desired inequality (2.9) for  $n+1$ . By induction, (2.9) holds for all  $n \geq 0$ . Theorem 2.2 is proved.  $\square$

**Remark 2.3.** When the Derrida–Retaux system is critical or subcritical, Theorem 2.2 has a simple consequence that is useful in the proof of the upper bound in Theorem 1.2. [Its most important application, however, will arrive in Lemma 5.2 as a technical ingredient in the proof of the lower bound in Theorem 1.2.] Assume  $(X_n, n \geq 0)$  is critical or subcritical:  $(m-1)\mathbf{E}(X_0 m^{X_0}) \leq \mathbf{E}(m^{X_0}) < \infty$ . This yields (see [9], [6]; a simple explanation is given at the beginning of Section 3) that  $(m-1)\mathbf{E}(X_n m^{X_n}) \leq \mathbf{E}(m^{X_n}) < \infty$  for all  $n \geq 0$ . In particular,

$$\mu_n(k) \geq m^{-k} \mathbf{E}([(m-1)X_n - 1]m^{X_n} \mathbf{1}_{\{X_n \geq k+1\}}) \geq 0,$$

for all  $k \geq 0$ . On the other hand, by definition,

$$\mu_n(k) \leq m^{-k} \mathbf{E}([1 - (m-1)X_n]m^{X_n} \mathbf{1}_{\{X_n=0\}}) = m^{-k} \mathbf{P}(X_n = 0) \leq m^{-k}.$$

So for critical or subcritical Derrida–Retaux systems, we have

$$(2.12) \quad 0 \leq \mu_n(k) \leq m^{-k}, \quad k \geq 0, n \geq 0.$$

Let us look at Theorem 2.2. By considering only the term  $k = 0$ , and since  $\mu_n(\cdot)$  is non-negative, the theorem implies that for integers  $n \geq 0$  and  $\ell \geq 0$ ,

$$\mu_0(0) \mathbf{E}(S_n^{(0,\ell)}) \leq \mu_n(\ell) \leq m^{-\ell},$$

the last inequality being from (2.12). Let  $N_n^{(0)}$  be as in (2.3). We have  $\mathbf{E}(N_n^{(0)} \mathbf{1}_{\{X_n=\ell\}}) = \mu_0(0) \mathbf{E}(S_n^{(0,\ell)})$  (see (2.8)). Hence

$$\mathbf{E}(N_n^{(0)} \mathbf{1}_{\{X_n=\ell\}}) \leq m^{-\ell}.$$

Consequently,

$$(2.13) \quad \mathbf{E}(N_n^{(0)} \mathbf{1}_{\{X_n \geq \ell\}}) \leq \frac{c_1}{m^\ell}, \quad n \geq 0, \ell \geq 0,$$

with  $c_1 := \frac{m}{m-1}$ . □

**Remark 2.4.** Let  $N_n$  denote, as in (2.3), the number of open paths from the initial generation to  $\mathbf{e}_n$ . When  $N_n \geq 1$ , we say that the system is sustainable until  $\mathbf{e}_n$ . We are going to see that in the study of the Derrida–Retaux system, it is important to understand the situation of sustainability (more important than to understand the situation  $X_n \geq 1$ ).

By definition,

$$\{X_n \geq 1\} \subset \{N_n \geq 1\} = \{X_{n-1}^{(1)} + \cdots + X_{n-1}^{(m)} \geq 1\},$$

where  $X_{n-1}^{(1)}, \dots, X_{n-1}^{(m)}$  are, as before, independent copies of  $X_{n-1}$ . Therefore,

$$\mathbf{P}(X_n \geq 1) \leq \mathbf{P}(N_n \geq 1) \leq m \mathbf{P}(X_{n-1} \geq 1).$$

In particular, for a critical system  $(Y_n, n \geq 0)$ , if  $\mathbf{E}(t^{Y_0}) < \infty$  (for some  $t > m$ ), then Theorem 1.2 says that the probability that the system is sustainable until  $\epsilon_n$  equals  $\frac{1}{n^{2+o(1)}}$  (for  $n \rightarrow \infty$ ).  $\square$

### 3 Preliminaries on the moment generating function

This section collects some estimates, in four lemmas below, for the moment generating function of Derrida–Retaux systems at criticality. All the results are either known, or are proved painlessly by means of existing methods, with the only exception of Lemma 3.2 (i) which is obtained as a consequence of (2.13).

Let  $(X_n, n \geq 0)$  be a Derrida–Retaux system defined via (1.3) (it will soon be assumed to be critical), satisfying  $\mathbf{E}(X_0 m^{X_0}) < \infty$ . Write

$$G_n(s) := \mathbf{E}(s^{X_n}),$$

for the moment generating function of  $X_n$ . The recursion in (1.3) means

$$(3.1) \quad G_{n+1}(s) = \frac{1}{s} G_n(s)^m + \left(1 - \frac{1}{s}\right) G_n(0)^m,$$

from which we get

$$G'_{n+1}(s) = \frac{m}{s} G'_n(s) G_n(s)^{m-1} - \frac{1}{s^2} G_n(s)^m + \frac{1}{s^2} G_n(0)^m.$$

Hence

$$(3.2) \quad s(s-1)G'_{n+1}(s) - G_{n+1}(s) = [m(s-1)G'_n(s) - G_n(s)]G_n(s)^{m-1}.$$

In particular,

$$m(m-1)G'_{n+1}(m) - G_{n+1}(m) = [m(m-1)G'_n(m) - G_n(m)]G_n(m)^{m-1}.$$

Iterating the identity leads to the following formula: for  $n \geq 1$ ,

$$m(m-1)G'_n(m) - G_n(m) = [m(m-1)G'_0(m) - G_0(m)] \prod_{j=0}^{n-1} G_j(m)^{m-1}.$$

In particular, it tells us that the sign of  $m(m-1)G'_n(m) - G_n(m)$  stays the same for all  $n \geq 0$ . For example, if  $(m-1)\mathbf{E}(X_0 m^{X_0}) \leq \mathbf{E}(m^{X_0})$ , then for all  $n \geq 0$ ,  $(m-1)\mathbf{E}(X_n m^{X_n}) \leq \mathbf{E}(m^{X_n})$ ; and since  $\mathbf{E}(X_n m^{X_n}) \geq \mathbf{E}(X_n) \mathbf{E}(m^{X_n})$  (by the FKG inequality), we get  $\mathbf{E}(X_n) \leq \frac{1}{m-1}$ ,  $\forall n \geq 0$ , which is the second part of Theorem A recalled in the introduction. Another consequence is that if  $(X_n, n \geq 0)$  is critical (resp. supercritical; subcritical), then so is  $(X_{n+k}, n \geq 0)$  for any integer  $k \geq 0$ .

We can extend (3.1) by bringing in also the number of open paths. For any vertex  $v \in \mathbb{T}$  with  $|v| \geq 1$ , let  $v^{(1)}, \dots, v^{(m)}$  denote as before its parents in generation  $|v| - 1$ . By definition of  $N^{(i)}(v)$  in (2.2), for any integer  $i \geq 0$ ,

$$\begin{aligned} X(v) &= [X(v^{(1)}) + \dots + X(v^{(m)}) - 1]^+, \\ N^{(i)}(v) &= [N^{(i)}(v^{(1)}) + \dots + N^{(i)}(v^{(m)})] \mathbf{1}_{\{X(v^{(1)}) + \dots + X(v^{(m)}) \geq 1\}}. \end{aligned}$$

With  $\Sigma_X = \Sigma_X(v) := X(v^{(1)}) + \dots + X(v^{(m)})$  and  $\Sigma_N = \Sigma_N(v, i) := N^{(i)}(v^{(1)}) + \dots + N^{(i)}(v^{(m)})$ , this can be rewritten as

$$(X(v), N^{(i)}(v)) = (\Sigma_X - 1, \Sigma_N) \mathbf{1}_{\{\Sigma_X \geq 1\}} + (0, 0) \mathbf{1}_{\{\Sigma_X = 0\}}.$$

Thus for  $s > 0$  and  $t \geq 0$  (with  $0^0 := 1$ ),

$$\begin{aligned} \mathbf{E}(s^{X(v)} t^{N^{(i)}(v)}) &= \mathbf{E}(s^{\Sigma_X - 1} t^{\Sigma_N} \mathbf{1}_{\{\Sigma_X \geq 1\}}) + \mathbf{P}(\Sigma_X = 0) \\ &= \mathbf{E}(s^{\Sigma_X - 1} t^{\Sigma_N}) - \frac{1}{s} \mathbf{E}(t^{\Sigma_N} \mathbf{1}_{\{\Sigma_X = 0\}}) + \mathbf{P}(\Sigma_X = 0). \end{aligned}$$

Define, for any integer  $i \geq 0$ ,

$$G_n(s, t) = G_n(s, t, i) := \mathbf{E}(s^{X_n} t^{N_n^{(i)}}), \quad s \geq 0, t \geq 0,$$

which is the joint moment generating function for the pair  $(X_n, N_n^{(i)})$  ( $N_n^{(i)}$  being from (2.3)). [In particular,  $G_n(s) = G_n(s, 1)$ .] Since  $(X(v^{(j)}), N^{(i)}(v^{(j)}))$ , for  $1 \leq j \leq m$ , are i.i.d. having the law of  $(X_{n-1}, N_{n-1}^{(i)})$ , we have

$$\begin{aligned} \mathbf{E}(s^{\Sigma_X - 1} t^{\Sigma_N}) &= \frac{1}{s} G_{n-1}(s, t)^m, \\ \mathbf{E}(t^{\Sigma_N} \mathbf{1}_{\{\Sigma_X = 0\}}) &= [\mathbf{E}(t^{N_{n-1}} \mathbf{1}_{\{X_{n-1} = 0\}})]^m = G_{n-1}(0, t)^m, \\ \mathbf{P}(\Sigma_X = 0) &= [\mathbf{P}(X_{n-1} = 0)]^m = G_{n-1}(0, 1)^m. \end{aligned}$$

Consequently, for  $n \geq 1$ ,  $s > 0$  and  $t \geq 0$ ,

$$(3.3) \quad G_n(s, t) = \frac{G_{n-1}(s, t)^m}{s} - \frac{G_{n-1}(0, t)^m}{s} + G_{n-1}(0, 1)^m.$$

This formula is useful in the proof of Lemma 3.2 below.

In the rest of this section, we deal with the Derrida–Retaux system at criticality, and denote it by  $(Y_n, n \geq 0)$ :  $(m-1)\mathbf{E}(Y_n m^{Y_n}) = \mathbf{E}(m^{Y_n})$  for all  $n \geq 0$ . We assume  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ , which implies  $\mathbf{E}(Y_n^3 m^{Y_n}) < \infty$  for all  $n \geq 0$ .

**Lemma 3.1.** *Assume  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ . Let  $G_n(s) := \mathbf{E}(s^{Y_n})$ . There exist positive constants  $c_2, c_3, c_4, c_5, c_6, c_7$  and  $c_8$  such that for  $n \geq 1$ ,*

- (i)  $\mathbf{E}(m^{Y_n}) \leq c_2$ ;
- (ii)  $\frac{1}{m-1} \leq \mathbf{E}(Y_n m^{Y_n}) \leq c_3$ ;
- (iii)  $\mathbf{E}(Y_n^2 m^{Y_n}) \leq c_4 n$ ;
- (iv)  $\frac{c_5}{n} \leq \mathbf{E}(m^{Y_n} \mathbf{1}_{\{Y_n \geq 1\}}) \leq \frac{c_6}{n}$ ;
- (v)  $c_7 n^2 \leq \prod_{i=0}^{n-1} G_i(m)^{m-1} \leq c_8 n^2$ .

Lemma 3.1 is known: (v) was proved in [9] (case  $m = 2$ ) and [6], whereas (i) follows from Theorem A (ii) (it is also a consequence of (v)); since the system is critical, we have  $\mathbf{E}(Y_n m^{Y_n}) = \frac{1}{m-1} \mathbf{E}(m^{Y_n})$ , so the second inequality in (ii) follows from (i); for (iii) and (iv), see [4] and [6], respectively.

The first four parts of Lemma 3.1 can be stated in a more compact way and may be extended as follows: Suppose  $\mathbf{E}(Y_0^j m^{Y_0}) < \infty$  for some integer  $j \geq 2$ . For any  $k \in \{0, 1, \dots, j\}$ , there exist some constants  $c_9 > 0$  and  $c_{10} > 0$  depending on  $k$ <sup>8</sup> such that for all integers  $n \geq 1$ ,

$$(3.4) \quad c_9 n^{k-1} \leq \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq 1\}}) \leq c_{10} n^{k-1}.$$

Plainly Lemma 3.1 corresponds to the case  $j = 3$ .

Let us prove (3.4); the proof of the forthcoming Lemmas 3.2 and 3.3 will be a refinement of the same argument. Only the case  $k \in \{2, 3, \dots, j\}$  needs to be proved because for  $k \in \{0, 1\}$ , both inequalities in (3.4) are already contained in Lemma 3.1. By (3.1),

$$sG_{n+1}(s) = G_n(s)^m + (s-1)G_n(0)^m.$$

---

<sup>8</sup>In Lemma 3.3 below, we are going to work under a stronger integrability assumption, and give dependence in  $k$  of the constants  $c_9$  and  $c_{10}$ .

On both sides, we differentiate  $k$  times with respect to  $s$  (since  $k \geq 2$ , the second term on the right-hand side, being affine in  $s$ , disappears after the second differentiation), and apply the general Leibniz rule to the first term on the right-hand side, to see that

$$(3.5) \quad sG_{n+1}^{(k)}(s) + kG_{n+1}^{(k-1)}(s) = \sum_{(k_1, \dots, k_m) \in \mathbb{Z}_+^m: k_1 + \dots + k_m = k} \frac{k!}{k_1! \cdots k_m!} \prod_{i=1}^m G_n^{(k_i)}(s).$$

We prove the upper bound in (3.4) (for all  $k \in \{2, 3, \dots, j\}$ ) by induction in  $k$ . Note that the case  $k = 2$  follows from Lemma 3.1 (iii), whereas the case  $k = 3$  is known from [6] (equation (19) and Proposition 1 there). Consider  $k \geq 4$  and suppose that the upper bound in (3.4) holds for all cases up to  $k - 1$ . Taking  $s = m$  in (3.5) we get that  $G_{n+1}^{(k)}(m) \leq G_n(m)^{m-1} G_n^{(k)}(m) + c_{11}n^{k-2}$  for some constant  $c_{11} > 0$  only depending on  $k$ . Iterating the inequality gives that for  $n \geq 2$ ,

$$G_n^{(k)}(m) \leq \left( G_1^{(k)}(m) + \sum_{\ell=1}^{n-1} \frac{c_{11} \ell^{k-2}}{\prod_{i=1}^{\ell} G_i(m)^{m-1}} \right) \prod_{i=1}^{n-1} G_i(m)^{m-1}.$$

Using Lemma 3.1 (v), this yields the upper bound in (3.4).

For the lower bound, we use the Cauchy–Schwarz inequality to see that for  $k \geq 2$ ,

$$(3.6) \quad \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq 1\}}) \geq \frac{[\mathbf{E}(Y_n^{k-1} m^{Y_n} \mathbf{1}_{\{Y_n \geq 1\}})]^2}{\mathbf{E}(Y_n^{k-2} m^{Y_n} \mathbf{1}_{\{Y_n \geq 1\}})}.$$

This, together with the upper bound in (3.4) which we have just proved, yield the lower bound by induction in  $k$ .

We now proceed to our second lemma, which focuses on the joint moment generating function of  $Y_n$  and  $N_n^{(0)}$ . The lemma will be useful in the proof of the upper bound in Theorem 1.2.

**Lemma 3.2.** *Assume  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ . Let  $N_n^{(0)}$  denote the number of open paths as in (2.3) for  $(Y_n)$ . There exist positive constants  $c_{12}, c_{13}, c_{14}, c_{15}, c_{16}$  and  $c_{17}$  such that for all  $n \geq 1$ ,*

- (i)  $\mathbf{E}(m^{Y_n} N_n^{(0)}) \leq c_{12}n$ ;
- (ii)  $c_{13}n^2 \leq \mathbf{E}(Y_n m^{Y_n} N_n^{(0)}) \leq c_{14}n^2$ ;
- (iii)  $\mathbf{E}(Y_n^2 m^{Y_n} N_n^{(0)}) \leq c_{15}n^3$ ;
- (iv)  $\mathbf{E}(m^{Y_n} (N_n^{(0)})^2) \leq c_{16}n^3$ ;
- (v)  $\mathbf{E}(Y_n m^{Y_n} (N_n^{(0)})^2) \leq c_{17}n^4$ .

*Proof.* (i) Let  $\mu_n(k) := m^{-k} \mathbf{E}([1 - (m-1)Y_n]m^{Y_n} \mathbf{1}_{\{X_n \leq k\}})$  as in (2.6). Since the system is critical, we have  $(m-1)\mathbf{E}(Y_n m^{Y_n}) = \mathbf{E}(m^{Y_n})$ , so  $\mu_n(k) = m^{-k} \mathbf{E}([(m-1)Y_n - 1]m^{Y_n} \mathbf{1}_{\{Y_n \geq k+1\}}]$ .

We have  $\mathbf{E}(m^{Y_n} N_n^{(0)}) = \sum_{k=0}^{\infty} m^k \mathbf{E}(N_n^{(0)} \mathbf{1}_{\{Y_n=k\}})$  by the Fubini–Tonelli theorem. Recall from (2.8) that  $\mathbf{E}(N_n^{(0)} \mathbf{1}_{\{Y_n=k\}}) = \mu_0(0) \mathbf{E}(S_n^{(0,k)})$ , which, by Theorem 2.2, is bounded by  $\mu_n(k)$ . Hence

$$\begin{aligned} \mathbf{E}(m^{Y_n} N_n^{(0)}) &\leq \sum_{k=0}^{\infty} \mathbf{E}([(m-1)Y_n - 1]m^{Y_n} \mathbf{1}_{\{Y_n \geq k+1\}}]) \\ &= \mathbf{E}([(m-1)Y_n - 1]m^{Y_n} \sum_{k=0}^{\infty} \mathbf{1}_{\{Y_n \geq k+1\}}]) \\ &= \mathbf{E}[Y_n((m-1)Y_n - 1)m^{Y_n} \mathbf{1}_{\{Y_n \geq 1\}}] \\ &\leq (m-1)\mathbf{E}(Y_n^2 m^{Y_n}). \end{aligned}$$

By Lemma 3.1 (iii), we have  $\mathbf{E}(Y_n^2 m^{Y_n}) \leq c_4 n$ . Thus  $\mathbf{E}(m^{Y_n} N_n^{(0)}) \leq (m-1)c_4 n$ .

(ii)–(v) The proofs of (ii), (iii), (iv) and (v) are similar, all originating from (3.3) (which we apply to the recursive system  $(Y_n)$  in lieu of  $(X_n)$ ). Let  $i \geq 0$  be an integer, and write  $G_n(s, t) := \mathbf{E}(s^{Y_n} t^{N_n^{(i)}})$ . Define

$$I_n(s) := \mathbf{E}(s^{Y_n} N_n^{(i)}) = \left. \frac{\partial G_n(s, t)}{\partial t} \right|_{t=1}.$$

It follows from (3.3) that

$$(3.7) \quad I_{n+1}(s) = \frac{m}{s} I_n(s) G_n(s)^{m-1} - \frac{m}{s} I_n(0) G_n(0)^{m-1},$$

where  $G_n(s) := \mathbf{E}(s^{Y_n}) = G_n(s, 1)$ . Multiplying by  $s$  on both sides of (3.7), and differentiating with respect to  $s$ , we get

$$I_{n+1}(s) + s I'_{n+1}(s) = m(m-1)G_n(s)^{m-2} G'_n(s) I_n(s) + mG_n(s)^{m-1} I'_n(s).$$

At criticality, we have  $m(m-1)G'_n(m) = G_n(m)$ , which yields, for  $n \geq 0$ ,

$$I_{n+1}(m) + m I'_{n+1}(m) = G_n(m)^{m-1} (I_n(m) + m I'_n(m)).$$

Iterating the identity, and noting that  $\mathbf{E}[m^{Y_n}(1 + Y_n)N_n^{(i)}] = I_n(m) + m I'_n(m)$ , we get, for

all  $i \geq 0$  and  $n \geq 1$ ,

$$\begin{aligned}
\mathbf{E}[m^{Y_n}(1 + Y_n)N_n^{(i)}] &= \mathbf{E}[m^{Y_0}(1 + Y_0)N_0^{(i)}] \prod_{j=0}^{n-1} G_j(m)^{m-1} \\
(3.8) \qquad \qquad \qquad &= (i + 1)m^i \mathbf{P}(Y_0 = i) \prod_{j=0}^{n-1} G_j(m)^{m-1}.
\end{aligned}$$

This formula is known (Proposition 5.1 in [4]). We take  $i = 0$  from now on.

Since  $c_7 n^2 \leq \prod_{j=0}^{n-1} G_j(m)^{m-1} \leq c_8 n^2$  (by Lemma 3.1 (v)), (ii) follows immediately.

To prove (iii), we note from (3.7) that

$$\frac{\partial^2 (sI_{n+1}(s))}{\partial s^2} = m \frac{\partial^2}{\partial s^2} [G_n(s)^{m-1} I_n(s)].$$

Taking  $s = m$ , we get

$$\begin{aligned}
2I'_{n+1}(m) + mI''_{n+1}(m) &= mG_n(m)^{m-1}I''_n(m) + 2m(m-1)G_n(m)^{m-2}G'_n(m)I'_n(m) \\
&\quad + m(m-1)G_n(m)^{m-2}G''_n(m)I_n(m) \\
&\quad + m(m-1)(m-2)G_n(m)^{m-3}G'_n(m)^2I_n(m).
\end{aligned}$$

On the left-hand side, the expression is at least  $mI''_{n+1}(m)$ , whereas on the right-hand side, for the third term, we use  $G_n(m)^{m-2} \leq G_n(m)^{m-1}$ . Since  $G_n(m) = m(m-1)G'_n(m)$  at criticality, this yields that

$$\begin{aligned}
I''_{n+1}(m) &\leq G_n(m)^{m-1}I''_n(m) \\
&\quad + G_n(m)^{m-1} \left( \frac{2}{m} I'_n(m) + (m-1)I_n(m)G''_n(m) + \frac{m-2}{m^2(m-1)} I_n(m) \right).
\end{aligned}$$

For the second expression on the right-hand side, we have, by Lemma 3.1 (i),  $G_n(m)^{m-1} \leq c_2^{m-1}$ , and by (ii) which has just been proved,  $I'_n(m) = \mathbf{E}(Y_n m^{Y_n-1} N_n^{(0)}) \leq \frac{c_{14}}{m} n^2$ , whereas by (i),  $I_n(m) = \mathbf{E}(m^{Y_n} N_n^{(0)}) \leq c_{12} n$ , and by Lemma 3.1 (iii),  $G''_n(m) \leq \mathbf{E}(Y_n^2 m^{Y_n}) \leq c_4 n$ . Consequently,

$$(3.9) \qquad \qquad \qquad I''_{n+1}(m) \leq G_n(m)^{m-1}I''_n(m) + c_{18} n^2,$$

where  $c_{18} := c_2^{m-1}(\frac{2}{m^2}c_{14} + (m-1)c_{12}c_4 + \frac{m-2}{m^2(m-1)}c_{12})$ . Iterating the inequality yields that for  $n \geq 1$  (with  $\prod_{\emptyset} := 1$ ),

$$I''_n(m) \leq I''_1(m) \prod_{k=1}^{n-1} G_k(m)^{m-1} + c_{18} \sum_{\ell=1}^{n-1} \ell^2 \prod_{k=\ell+1}^{n-1} G_k(m)^{m-1}.$$



By Lemma 3.1 (v),  $\prod_{k=\ell+1}^{n-1} G_k(m)^{m-1} \leq \frac{c_8 n^2}{c_7(\ell+1)^2}$ , this gives that for  $n \geq 1$ ,

$$I_n''(m) \leq I_1''(m)c_8 n^2 + \frac{c_{18}c_8}{c_7} \sum_{\ell=1}^{n-1} \ell^2 \frac{n^2}{(\ell+1)^2} \leq c_8 \left( \frac{c_{18}}{c_7} + I_1''(m) \right) n^3.$$

Together with (ii) and (i), this implies (iii).

To prove (iv), let us define

$$J_n(s) := \mathbf{E}(s^{Y_n} (N_n^{(0)})^2) = \frac{\partial^2}{\partial t^2} G_n(s, t) \Big|_{t=1} + I_n(s).$$

By (3.3),

$$\begin{aligned} J_{n+1}(m) &= G_n(m)^{m-1} J_n(m) + (m-1) G_n(m)^{m-2} I_n(m)^2 \\ &\quad - (m-1) G_n(0)^{m-2} I_n(0)^2 - G_n(0)^{m-1} J_n(0) \\ &\leq G_n(m)^{m-1} J_n(m) + (m-1) G_n(m)^{m-2} I_n(m)^2. \end{aligned}$$

For the last term on the right-hand side, we use  $G_n(m)^{m-2} \leq c_2^{m-2}$  (by Lemma 3.1 (i)), and  $I_n(m) = \mathbf{E}(m^{Y_n} N_n^{(0)}) \leq c_{12} n$  (by (i)). Therefore, with  $c_{19} := (m-1)c_2^{m-2}c_{12}^2$ ,

$$J_{n+1}(m) \leq G_n(m)^{m-1} J_n(m) + c_{19} n^2,$$

which is the analogue of (3.9) for  $J_n(m)$  in place of  $I_n''(m)$ . Since  $J_0(m) = \mathbf{E}(m^{Y_0} (N_0^{(0)})^2) = \mathbf{P}(Y_0 = 0)$ , this yields  $\mathbf{E}(m^{Y_n} (N_n^{(0)})^2) = J_n(m) \leq c_{16} n^3$  for some  $c_{16} > 0$  and all  $n \geq 1$ , exactly as (3.9) has implied (iii).

It remains to check (v), which is similar, with more computations. Define

$$K_n(s) := \mathbf{E}(Y_n s^{Y_n} (N_n^{(0)})^2) = s \frac{\partial}{\partial s} \frac{\partial^2}{\partial t^2} G_n(s, t) \Big|_{t=1} + s I_n'(s).$$

By (3.3) and the identity  $G_n'(m) = \frac{G_n(m)}{m(m-1)}$  at criticality, we get

$$\begin{aligned} K_{n+1}(m) &= G_n(m)^{m-1} K_n(m) - I_n(m)^2 G_n(m)^{m-2} + 2m(m-1) I_n(m) I_n'(m) G_n(m)^{m-2} \\ &\quad + J_n(0) G_n(0)^{m-1} + (m-1) I_n(0)^2 G_n(0)^{m-2}. \end{aligned}$$

On the right-hand side, we throw away the second term due to its negativity, use for the third term  $I_n(m) \leq c_{12} n$  (by (i)),  $I_n'(m) \leq \frac{c_{14}}{m} n^2$  (by (ii)) and  $G_n(m)^{m-2} \leq c_2^{m-2}$  (by Lemma 3.1 (i)), and for the last two terms,  $G_n(0) = \mathbf{P}(Y_n = 0) \leq 1$ ,  $J_n(0) \leq J_n(m) \leq c_{16} n^3$

(by (iv)), and  $I_n(0)^2 \leq I_n(m)^2 \leq c_{12}^2 n^2$  (by (i) again). This leads to, with  $c_{20} := 2(m-1)c_{12}c_{14}c_2^{m-2} + c_{16} + (m-1)c_{12}^2$ : for  $n \geq 1$ ,

$$K_{n+1}(m) \leq G_n(m)^{m-1}K_n(m) + c_{20} n^3,$$

which is once again the analogue of (3.9) for  $K_n(m)$  in place of  $I_n''(m)$ . This yields that  $\mathbf{E}(Y_n m^{Y_n} (N_n^{(0)})^2) = K_n(m) \leq c_{21} n^4$  for some  $c_{21} > 0$  and all  $n \geq 1$  exactly as (3.9) has implied (iii). The proof of Lemma 3.2 is complete.  $\square$

We have yet two more lemmas, Lemmas 3.3 and 3.4 to present in this section. Our interest is in Lemma 3.4, which (in the case  $k = 1$ ) will be used a couple of times in Section 5 in the proof of the lower bound in Theorem 1.2. The proof of Lemma 3.4 relies on Lemma 3.3.

**Lemma 3.3.** *Assume  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ . Let  $G_n(s) := \mathbf{E}(s^{Y_n})$  as before.*

(i) *There exists a constant  $c_{22} > 0$  such that for all integers  $n \geq 1$  and  $k \geq 1$ ,*

$$G_n^{(k)}(m) \leq k! n^{k-1} e^{c_{22}k}.$$

(ii) *There exist constants  $c_{23} \geq 1$  and  $c_{24} > 0$  such that for integers  $k \geq 1$  and  $n \geq e^{c_{23}k}$ ,*

$$G_n^{(k)}(m) \geq k! n^{k-1} e^{-c_{24}k}.$$

*Proof.* The inequality in (i) was proved in [7, Theorem 2]. Let us prove (ii). For fixed  $k$ , (ii) is simply the first inequality in (3.4), so we need to prove it only for  $k \geq 6$ .

We claim the existence of constants  $c_{23} \geq 1$  and  $c_{25} > 0$  such that for integers  $k \geq 2$  and  $n \geq \lceil e^{c_{23}k} \rceil$ ,

$$(3.10) \quad G_n^{(k)}(m) \geq k! n^{k-1} e^{-c_{25}(k-1)},$$

while keeping in mind that (3.10) is already proved for  $2 \leq k \leq 5$ . Our proof is based on an argument by induction in  $k$ .

Let  $k \geq 6$ . Assume (3.10) is valid for all derivatives of order  $\leq k-1$  at  $s = m$ . We need to prove it for  $G_n^{(k)}(m)$ . Let us recall the formula (3.5) at  $s = m$ :

$$mG_{n+1}^{(k)}(m) + kG_{n+1}^{(k-1)}(m) = \sum_{(k_1, \dots, k_m) \in \mathbb{Z}_+^m: k_1 + \dots + k_m = k} \frac{k!}{k_1! \dots k_m!} \prod_{i=1}^m G_n^{(k_i)}(m).$$

On the right-hand side, we only consider two special types of  $(k_1, \dots, k_m) \in \mathbb{Z}_+^m$  satisfying  $k_1 + \dots + k_m = k$ : (1) one of  $k_i$  is  $k$  (so all others vanish); (2) one of  $k_i$  is  $\ell$  (for some  $1 \leq \ell \leq \lfloor \frac{k}{2} \rfloor - 1$ ), another is  $k - \ell$  (and all others vanish). Hence, for  $k \geq 6$ ,

$$\begin{aligned} mG_{n+1}^{(k)}(m) + kG_{n+1}^{(k-1)}(m) &\geq mG_n^{(k)}(m)G_n(m)^{m-1} \\ &\quad + m(m-1) \sum_{\ell=1}^{\lfloor \frac{k}{2} \rfloor - 1} \frac{k!}{\ell!(k-\ell)!} G_n^{(\ell)}(m)G_n^{(k-\ell)}(m)G_n(m)^{m-2}. \end{aligned}$$

On the right-hand side, the term  $\ell = 1$  is special: we have  $G_n'(m) = \frac{G_n(m)}{m(m-1)}$  at criticality. This yields

$$\begin{aligned} mG_{n+1}^{(k)}(m) + kG_{n+1}^{(k-1)}(m) &\geq [mG_n^{(k)}(m) + kG_n^{(k-1)}(m)]G_n(m)^{m-1} \\ &\quad + m(m-1) \sum_{\ell=2}^{\lfloor \frac{k}{2} \rfloor - 1} \frac{k!}{\ell!(k-\ell)!} G_n^{(\ell)}(m)G_n^{(k-\ell)}(m)G_n(m)^{m-2}. \end{aligned}$$

Let  $2 \leq \ell \leq \lfloor \frac{k}{2} \rfloor - 1$ . By the induction assumption, if  $n \geq \lceil e^{c_{23}(k-2)} \rceil$ , then  $G_n^{(\ell)}(m)G_n^{(k-\ell)}(m) \geq \ell!(k-\ell)!n^{k-2}e^{-c_{25}(k-2)}$ . We also use the trivial inequalities  $m(m-1) \geq 1$  and  $G_n(m)^{m-2} \geq 1$ .

Consequently,

$$\begin{aligned} m(m-1) \sum_{\ell=2}^{\lfloor \frac{k}{2} \rfloor - 1} \frac{k!}{\ell!(k-\ell)!} G_n^{(\ell)}(m)G_n^{(k-\ell)}(m)G_n(m)^{m-2} &\geq \sum_{\ell=2}^{\lfloor \frac{k}{2} \rfloor - 1} k!n^{k-2}e^{-c_{25}(k-2)} \\ &= (\lfloor \frac{k}{2} \rfloor - 2)k!n^{k-2}e^{-c_{25}(k-2)}. \end{aligned}$$

Since  $k \geq 6$ , we have  $\lfloor \frac{k}{2} \rfloor - 2 \geq \frac{k}{7}$ . Hence, for  $k \geq 6$  and  $n \geq \lceil e^{c_{23}(k-2)} \rceil$ ,

$$mG_{n+1}^{(k)}(m) + kG_{n+1}^{(k-1)}(m) \geq [mG_n^{(k)}(m) + kG_n^{(k-1)}(m)]G_n(m)^{m-1} + \frac{k!k}{7}n^{k-2}e^{-c_{25}(k-2)}.$$

Iterating the inequality gives that, for  $n \geq n_0 + 1$  where  $n_0 = n_0(k) := \lceil e^{c_{23}(k-2)} \rceil$ ,

$$\begin{aligned} &mG_n^{(k)}(m) + kG_n^{(k-1)}(m) \\ &\geq \left( mG_{n_0}^{(k)}(m) + kG_{n_0}^{(k-1)}(m) + \sum_{i=n_0}^{n-1} \frac{k!k_i^{k-2}e^{-c_{25}(k-2)}}{\prod_{j=n_0}^i G_j(m)^{m-1}} \right) \prod_{j=n_0}^{n-1} G_j(m)^{m-1} \\ &\geq \sum_{i=n_0}^{n-1} \frac{k!k_i^{k-2}e^{-c_{25}(k-2)}}{\prod_{j=n_0}^i G_j(m)^{m-1}} \prod_{j=n_0}^{n-1} G_j(m)^{m-1}. \end{aligned}$$

By Lemma 3.1 (v),  $c_7 n^2 \leq \prod_{i=0}^{n-1} G_i(m)^{m-1} \leq c_8 n^2$ , so

$$mG_n^{(k)}(m) + kG_n^{(k-1)}(m) \geq \frac{k!k}{7} e^{-c_{25}(k-2)} \sum_{i=n_0}^{n-1} \frac{i^{k-2}}{(i+1)^2} \frac{c_7}{c_8} n^2.$$

If  $n \geq \lceil e^{c_{23}k} \rceil$ , then  $n_0 \leq \frac{n}{2}$  (recalling that  $c_{23} \geq 1$ ), so  $\sum_{i=n_0}^{n-1} \frac{i^{k-2}}{(i+1)^2} \geq \sum_{i=\lfloor n/2 \rfloor}^{n-1} \frac{i^{k-2}}{(i+1)^2} \geq c_{26} \frac{n^{k-3}}{k}$  for some constant  $c_{26} > 0$ . This yields, for  $n \geq \lceil e^{c_{23}k} \rceil$  (with  $c_{27} := \frac{c_7 c_{26}}{7c_8}$ )

$$(3.11) \quad mG_n^{(k)}(m) + kG_n^{(k-1)}(m) \geq c_{27} k! e^{-c_{25}(k-2)} n^{k-1}.$$

On the other hand, by (i),  $G_n^{(k-1)}(m) \leq e^{c_{22}(k-1)} (k-1)! n^{k-2}$  for some constant  $c_{22} > 0$ . Thus

$$kG_n^{(k-1)}(m) \leq k e^{c_{22}(k-1)} (k-1)! n^{k-2},$$

which is bounded by  $\frac{c_{27}}{2} k! e^{-c_{25}(k-2)} n^{k-1}$  if  $n \geq \lceil e^{c_{23}k} \rceil$  (with a large constant  $c_{23}$ ). Going back to (3.11), we obtain: for  $k \geq 6$  and  $n \geq \lceil e^{c_{23}k} \rceil$ ,

$$mG_n^{(k)}(m) \geq \frac{c_{27}}{2} k! e^{-c_{25}(k-2)} n^{k-1},$$

which is at least  $m k! e^{-c_{25}(k-1)} n^{k-1}$  if  $c_{25}$  is chosen to be sufficiently large.<sup>9</sup> As a consequence, (3.10) is valid for  $k \geq 6$ .

Lemma 3.3 is proved. □

**Lemma 3.4.** *Assume  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ . For any  $\lambda > 0$  and any integer  $k \geq 1$ ,*

$$(3.12) \quad \liminf_{n \rightarrow \infty} \frac{1}{n^{k-1}} \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) > 0.$$

*Proof.* We have, for  $k \geq 1$ ,

$$\mathbf{E}(Y_n^k m^{Y_n}) \leq \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) + (\lambda n)^{k-1} \mathbf{E}(Y_n m^{Y_n}).$$

By Lemma 3.1 (ii),  $\mathbf{E}(Y_n m^{Y_n}) \leq c_3$ . Thus

$$\mathbf{E}(Y_n^k m^{Y_n}) \leq \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) + c_3 (\lambda n)^{k-1}.$$

---

<sup>9</sup>The constants are chosen in this order: first  $c_{25}$  then  $c_{23}$ , while our presentation is in the opposite order.

On the other hand, by Lemma 3.3 (ii), for all sufficiently large integer  $k$  (say  $k \geq k_0$ ) and all integer  $n \geq e^{c_{23}k}$ ,

$$\mathbf{E}(Y_n^k m^{Y_n}) \geq 2c_3(\lambda n)^{k-1}.$$

Consequently, for  $k \geq k_0$  and  $n \geq e^{c_{23}k}$ ,

$$2c_3(\lambda n)^{k-1} \leq \mathbf{E}(Y_n^k m^{Y_n}) \leq \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) + c_3(\lambda n)^{k-1},$$

which implies that

$$(3.13) \quad \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) \geq c_3(\lambda n)^{k-1}, \quad k \geq k_0, n \geq e^{c_{23}k}.$$

In particular,

$$(3.14) \quad \liminf_{n \rightarrow \infty} \frac{1}{n^{k-1}} \mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) > 0,$$

which yields Lemma 3.4 when  $k \geq k_0$ .

To remove the restriction  $k \geq k_0$  for the validity of (3.14), we observe that by the Cauchy–Schwarz inequality,

$$\mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) \leq [\mathbf{E}(Y_n^{k+1} m^{Y_n})]^{1/2} [\mathbf{E}(Y_n^{k-1} m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}})]^{1/2}.$$

[A similar inequality was already used in (3.6).] Recall from (3.4) that  $\mathbf{E}(Y_n^{k+1} m^{Y_n}) \leq c_{10} n^k$ .

Hence, for  $k \geq 1$  and  $n \geq 1$ ,

$$(3.15) \quad \mathbf{E}(Y_n^{k-1} m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}}) \geq \frac{[\mathbf{E}(Y_n^k m^{Y_n} \mathbf{1}_{\{Y_n \geq \lambda n\}})]^2}{c_{10} n^k}.$$

Consequently, the inequality (3.14) applied to  $k = k_0$ , together with (3.15), yield that (3.14) is valid for  $k = k_0 - 1$ . Iterating the argument, we see that (3.14) remains valid for all  $k \geq 1$ .  $\square$

## 4 Proof of Theorem 1.2: upper bound

Let  $(Y_n, n \geq 0)$  be a Derrida–Retaux system satisfying  $\mathbf{E}(m^{Y_0}) = (m - 1)\mathbf{E}(Y_0 m^{Y_0}) < \infty$ ; so it is critical. The upper bound in Theorem 1.2 is a consequence of the following result in the particular case  $\ell = 0$ .

**Proposition 4.1.** *Assume  $\mathbf{E}(Y_0^5 m^{Y_0}) < \infty$ . There exists  $c_{28} > 0$  such that for integers  $n \geq 2$  and  $\ell \geq 0$ ,*

$$(4.1) \quad \mathbf{P}(Y_n \geq \ell + 1) \leq \frac{c_{28}}{m^\ell} \frac{(\log n)^2}{n^2}.$$

In the proof of Proposition 4.1, we are going to truncate the initial distribution of a system and compare it with the system itself. Let  $(X_n, n \geq 0)$  be a Derrida–Retaux system, with genealogical tree  $\mathbb{T}$ . Let  $B \subset \{0, 1, 2, \dots\}$ . Define

$$\widehat{X}_0 := X_0 \mathbf{1}_{\{X_0 \notin B\}}.$$

Let  $(\widehat{X}_n, n \geq 0)$  be a Derrida–Retaux system with initial distribution  $\widehat{X}_0$ , associated with genealogical tree  $\mathbb{T}$ , such that  $X(v) \geq \widehat{X}(v)$  for all  $v \in \mathbb{T}$ . In particular,  $\widehat{X}(v) = X(v) \mathbf{1}_{\{X(v) \notin B\}}$  if  $|v| = 0$ .

**Lemma 4.2.** *Fix  $n \geq 1$ . If there are no open paths in the system  $(X_k, k \geq 0)$  that lead to vertex  $\mathbf{e}_n$  and start from a vertex at the initial generation with a value lying in  $B$ , then  $X(\mathbf{e}_n) = \widehat{X}(\mathbf{e}_n)$ .*

*Proof of Lemma 4.2.* Consider the system  $(X_k, k \geq 0)$ . Let  $R_n$  denote the set of vertices (including  $\mathbf{e}_n$  itself) in all the open paths starting from the initial generation and leading to  $\mathbf{e}_n$ . Let  $R_{n,0}$  be the set of elements of  $R_n$  that are in the initial generation. [So  $N_n$ , defined in (2.3), is simply  $|R_{n,0}|$ , the cardinality of  $R_{n,0}$ .] Then the value of  $X(\mathbf{e}_n)$  does not depend on  $\{X(v), v \in \mathbb{T}_n \setminus R_{n,0} \text{ with } |v| = 0\}$ , and

$$(4.2) \quad X(\mathbf{e}_n) = \sum_{v \in R_{n,0}} X(v) - |R_n \setminus R_{n,0}|.$$

[If  $R_{n,0}$  is empty, then so is  $R_n$ , and (4.2) holds trivially.]

Assume that there are no open paths in the system  $(X_k, k \geq 0)$  that lead to vertex  $\mathbf{e}_n$  and start from a vertex at the initial generation with a value lying in  $B$ . Then for any  $v \in R_{n,0}$ , we have  $X(v) \notin B$ , so that  $\widehat{X}(v) = X(v)$ . This implies that  $\widehat{X}(v) \geq X(v)$  for  $v \in R_n$ . In particular,  $\widehat{X}(\mathbf{e}_n) \geq X(\mathbf{e}_n)$ .  $\square$

In order to prove Proposition 4.1, we work with a truncated version of the system  $(Y_n, n \geq 0)$ , the level of the truncation depending on an integer parameter  $M$  which is sufficiently

large (we write generically  $M \geq M_0$ ; the value of  $M_0$  may change from line to line). For integer  $M \geq M_0$ , define

$$(4.3) \quad \zeta(M) := -\log \mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 > M\}}) \in [1, \infty].$$

[Notation:  $\zeta(M) := \infty$  if  $\mathbf{P}(Y_0 > M) = 0$ . We have  $\lim_{M \rightarrow \infty} \zeta(M) = \infty$  under the assumption  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ .] Define

$$(4.4) \quad Z_0 = Z_0(M) := Y_0 \mathbf{1}_{\{Y_0 \leq M\zeta(M)\}}.$$

We consider a new Derrida–Retaux system  $(Z_n, n \geq 0)$  with initial distribution  $Z_0$ . Since  $Z_0 \leq Y_0$  and  $(Y_n, n \geq 0)$  is critical, the new system  $(Z_n, n \geq 0)$  is subcritical, or critical if  $Y_0 \leq M\zeta(M)$  a.s.

*Outline of the proof of Proposition 4.1.* The key ingredient in the proof is the following estimate (Lemma 4.6): if  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ , there exist constants  $c_{43} > 0$ ,  $c_{44} > 0$  and  $M_0 \geq 1$  such that for all integers  $n \geq M \geq M_0$ ,

$$(4.13) \quad \mathbf{P}(Z_n \geq \ell + 1) \leq c_{43} \left( \frac{1}{M^2} + \frac{1}{M} e^{-c_{44} \frac{n}{M}} \right) m^{-\ell}, \quad \ell \geq 0.$$

This will yield (4.1) by taking  $M$  to be (the integer part of) a suitable constant multiple of  $\frac{n}{\log n}$ , because we will see that the difference between  $\mathbf{P}(Y_n \geq \ell + 1)$  and  $\mathbf{P}(Z_n \geq \ell + 1)$  can be controlled under further integrability assumption  $\mathbf{E}(Y_0^5 m^{Y_0}) < \infty$ .

To prove (4.13), we define a deterministic set  $B_M$  of integers  $y$  of order of magnitude  $M$  such that conditionally on  $Z_M = y$ , “with reasonable probability”, there are at least (a constant multiple of)  $M^2$  open paths from the initial generation to generation  $M$ .

We distinguish two possible situations:

- Case 1: there are no open paths starting from generation  $M$  with a value lying in  $B_M$  leading to generation  $n + M$ ;
- Case 2: there are such open paths.

It is not hard to treat Case 2. The basic idea is that if there are such open paths, then by the construction of  $B_M$ , with “reasonable probability”, there are at least (a constant multiple of)  $M^2$  open paths leading to generation  $n + M$  and starting from the initial generation with value 0. We are going to see that thanks to (2.13), this leads to:

$$\mathbf{P}(Z_{n+M} \geq \ell, \text{ Case 2}) \leq \frac{c}{M^2} \frac{1}{m^\ell},$$

for some constant  $c > 0$ , all  $n \geq 0$  and  $\ell \geq 1$ .

To treat Case 1 which is more delicate, we define

$$W_0 = W_0(M) := Z_M \mathbf{1}_{\{Z_M \notin B_M\}},$$

and consider yet another Derrida–Retaux system  $(W_n, n \geq 0)$  with initial distribution  $W_0$ . The reason for which we are interested in this new system is that by Lemma 4.2, in Case 1, if  $Z_{n+M} = \ell$ , then  $W_n = \ell$ , so that for all  $n \geq 0$ ,

$$\mathbf{P}(Z_{n+M} = \ell, \text{ Case 1}) \leq \mathbf{P}(W_n = \ell) \leq \frac{\mathbf{E}(m^{W_n}) - 1}{m^\ell - 1}.$$

We need to control  $\mathbf{E}(m^{W_n})$ . This relies on subcriticality of the  $(W_i, i \geq 0)$ , and on a simple estimate (Lemma 4.5) on the moment generating function for all subcritical Derrida–Retaux systems satisfying a convenient condition for the initial distribution. We check by means of a technical lemma (Lemma 4.3) that  $W_{n_0}$  satisfies this condition, where  $n_0 := \lfloor c_{34}M \rfloor$  for some constant  $c_{34} > 0$  and all sufficiently large  $M$ .<sup>10</sup> This will yield that there exist constants  $c_{33} > 0$  and  $c_{35} \in (0, 1)$ , such that for all integer  $n \geq 0$ ,

$$\mathbf{E}(m^{W_{n+n_0}}) \leq 1 + \frac{c_{33}}{M} \left(1 - \frac{c_{35}}{M}\right)^n.$$

Consequently, for  $\ell \geq 1$  and  $n \geq 2n_0$ ,

$$\mathbf{P}(Z_{n+M} = \ell, \text{ Case 1}) \leq \frac{\mathbf{E}(m^{W_n}) - 1}{m^\ell - 1} \leq \frac{\frac{c_{33}}{M} (1 - \frac{c_{35}}{M})^{n-n_0}}{m^\ell - 1} \leq \frac{\frac{c_{33}}{M} (1 - \frac{c_{35}}{M})^{n/2}}{m^\ell - 1}.$$

Combining this with the bound for  $\mathbf{P}(Z_{n+M} \geq \ell, \text{ Case 2})$ , we see that for all  $n \geq 2n_0$  and  $\ell \geq 1$ ,

$$\mathbf{P}(Z_{n+M} \geq \ell) \leq \frac{c}{M^2} \frac{1}{m^\ell} + \sum_{j=\ell}^{\infty} \frac{\frac{c_{33}}{M} (1 - \frac{c_{35}}{M})^{n/2}}{m^j - 1}.$$

This will lead to (4.13). □

The rest of the section is devoted to the proof of Proposition 4.1. For the sake of clarity, it is divided into three steps and presented in distinct parts. The first step is about localization; it consists in understanding where the truncated Derrida–Retaux system  $(Z_n, n \geq 0)$  lies with good (weighted) probability conditioning on sustainability; this is then used to prove Lemma 4.3, the main result in the step. The second step gives (4.13) (which is Lemma 4.6), whereas the third step completes the proof of Proposition 4.1.

---

<sup>10</sup>It is not a surprise to consider the system at the intermediate generation  $n_0$ . A similar idea is going to be used in the proof of the lower bound in Section 5.



## 4.1 Step 1. Localization

The aim of Step 1 is to prove Lemma 4.3, which is the technical part of preparation to give an upper bound for  $\mathbf{P}(Z_n \geq \ell + 1)$  in Lemma 4.6. For  $n \geq 1$ , let  $N_n^{(0)}$  denote the number of open paths for the critical Derrida–Retaux system  $(Y_n)$  (replacing  $(X_n)$  by  $(Y_n)$  in (2.3)), and let  $L_n^{(0)}$  be the corresponding quantity for the truncated Derrida–Retaux system  $(Z_n)$ ; define the deterministic set  $B_n$  by

$$(4.5) \quad B_n := \{y \in [c_{29}n, c_{30}n] \cap \mathbb{Z} : \mathbf{P}(Z_n = y) > 0, \mathbf{P}(L_n^{(0)} \geq c_{31}n^2 \mid Z_n = y) \geq c_{32}\},$$

where the positive constants  $c_{29}$ ,  $c_{30}$ ,  $c_{31}$  and  $c_{32}$  are explicit but uninteresting:  $c_{29} := \min\{\frac{c_{13}}{5c_{12}}, \frac{5c_{15}}{2c_{13}}\}$ ,  $c_{30} := \frac{5c_{15}}{c_{13}}$ ,  $c_{31} := \min\{\frac{c_{13}}{5c_3}, \frac{5c_{17}}{2c_{13}}\}$ ,  $c_{32} := \frac{c_{13}^3}{250c_3c_{15}c_{17}} \min\{\frac{c_{13}}{5c_{12}}, \frac{5c_{15}}{2c_{13}}\}$  ( $c_3$  being from Lemma 3.1 (ii), and  $c_{12}$ ,  $c_{13}$ ,  $c_{15}$  and  $c_{17}$  from Lemma 3.2). Let

$$(4.6) \quad W_0 = W_0(M) := Z_M \mathbf{1}_{\{Z_M \notin B_M\}}.$$

We also consider a new Derrida–Retaux system  $(W_n, n \geq 0)$  with initial distribution  $W_0$ . Since  $W_0 \leq Z_M$ , and  $(Z_n)$  is subcritical or critical, it follows that  $(W_n, n \geq 0)$  is subcritical or critical. As such, we have three Derrida–Retaux systems: the critical system  $(Y_n, n \geq 0)$  which we are interested in, and the two associated systems  $(Z_n, n \geq 0)$  and  $(W_n, n \geq 0)$ . They are related in the following way: for all  $n \geq 0$ ,

$$W_n \leq Z_{n+M} \leq Y_{n+M}.$$

We mention that both systems  $(Z_n, n \geq 0)$  and  $(W_n, n \geq 0)$  depend on  $M$ .

The aim of the step is to prove the following estimate.

**Lemma 4.3.** *Assume  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ . There exist constants  $c_{33} \in (0, 1)$ ,  $c_{34} > 0$ ,  $c_{35} \in (0, 1)$ , such that for all sufficiently large integer  $M$ , with  $s = s(M) := m + \frac{c_{33}}{M}$  and  $n_0 = n_0(M) := \lfloor c_{34}M \rfloor$ ,*

$$(4.7) \quad \frac{m}{s} [\mathbf{E}(s^{W_{n_0}})]^{m-1} \leq 1 - \frac{c_{35}}{M}.$$

The proof of Lemma 4.3 relies on the following estimate.

**Lemma 4.4.** Assume  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ . There exists a constant  $c_{36} > 0$  such that for all sufficiently large integer  $M$ ,

$$\mathbf{E}(m^{Z_M} \mathbf{1}_{\{Z_M \in B_M\}}) \geq \frac{c_{36}}{M},$$

where  $B_M$  is the deterministic set of integers defined in (4.5).<sup>11</sup>

*Proof of Lemma 4.4.* Let  $b_1, b_2, b_3, b_4, b_5$  be positive constants whose values will be chosen later. Let us consider the quantity

$$q_n := \mathbf{E}(Z_n m^{Z_n} L_n^{(0)} \mathbf{1}_{\{L_n^{(0)} \in [b_1 n^2, b_2 n^2], Z_n \in [b_3 n, b_4 n]\}}).$$

Assume  $M \geq M_0$  ( $M_0$  being a sufficiently large integer such that  $\zeta(M) \geq 2$  for  $M \geq M_0$ ) and  $1 \leq n \leq M$ . We first recall that  $Z_0 = Y_0 \mathbf{1}_{\{Y_0 \leq M\zeta(M)\}}$ . Since  $Y_n \geq Y_0 - n \geq Y_0 - M$  (this is why we assume  $n \leq M$ ), we have, on the event  $\{Y_n \leq \frac{1}{2}M\zeta(M)\}$ ,  $Y_0 \leq Y_n + M \leq M\zeta(M)$ , so  $Z_n = Y_n$  and  $L_n^{(0)} = N_n^{(0)}$ . Consequently, if  $b_4 \leq \frac{\zeta(M)}{2}$ , then  $\{Y_n \in [b_3 n, b_4 n]\} \subset \{Z_n \in [b_3 n, b_4 n]\}$ , and

$$\begin{aligned} q_n &\geq \mathbf{E}(Y_n m^{Y_n} N_n^{(0)} \mathbf{1}_{\{N_n^{(0)} \in [b_1 n^2, b_2 n^2], Y_n \in [b_3 n, b_4 n]\}}) \\ &\geq \mathbf{E}(Y_n m^{Y_n} N_n^{(0)}) - \mathbf{E}(Y_n m^{Y_n} N_n^{(0)} \mathbf{1}_{\{N_n^{(0)} \notin [b_1 n^2, b_2 n^2]\}}) - \mathbf{E}(Y_n m^{Y_n} N_n^{(0)} \mathbf{1}_{\{Y_n \notin [b_3 n, b_4 n]\}}) \\ &=: q_{n,1} - q_{n,2} - q_{n,3}, \end{aligned}$$

with obvious notation. On the right-hand side, let us look at each of the three expressions  $q_{n,i}$ ,  $i \in \{1, 2, 3\}$ . The first term  $q_{n,1}$  is dealt with by Lemma 3.2 (ii), which says that

$$q_{n,1} \geq c_{13} n^2.$$

For the second term  $q_{n,2}$ , we note that on the event  $\{N_n^{(0)} \notin [b_1 n^2, b_2 n^2]\}$ , we have either  $N_n^{(0)} < b_1 n^2$ , in which case  $Y_n m^{Y_n} N_n^{(0)} \leq b_1 n^2 Y_n m^{Y_n}$ , or  $N_n^{(0)} > b_2 n^2$ , in which case  $Y_n m^{Y_n} N_n^{(0)} \leq \frac{1}{b_2 n^2} Y_n m^{Y_n} (N_n^{(0)})^2$ . Accordingly,

$$q_{n,2} \leq b_1 n^2 \mathbf{E}(Y_n m^{Y_n}) + \frac{1}{b_2 n^2} \mathbf{E}(Y_n m^{Y_n} (N_n^{(0)})^2) \leq c_3 b_1 n^2 + \frac{c_{17} n^4}{b_2 n^2},$$

---

<sup>11</sup>Since  $m^{Z_M} \mathbf{1}_{\{Z_M \in B_M\}} \leq m^{Z_M} \mathbf{1}_{\{Z_M \geq 1\}} \leq m^{Y_M} \mathbf{1}_{\{Y_M \geq 1\}}$ , it follows from Lemmas 4.4 and 3.1 (iv) that  $\mathbf{E}(m^{Z_M} \mathbf{1}_{\{Z_M \geq 1\}})$  is of order of magnitude  $\frac{1}{M}$ . So the meaning of Lemma 4.4 is as follows: if we work with the probability measure having a density proportional to  $m^{Z_M}$ , then conditionally on the event  $Z_M \geq 1$ , with probability greater than a certain constant (independent of  $M$ ),  $Z_M$  lies in  $B_M$ .

using Lemma 3.1 (ii) and Lemma 3.2 (v) for the last inequality. The third term  $q_{n,3}$  is handled in the same way:

$$q_{n,3} \leq b_3 n \mathbf{E}(m^{Y_n} N_n^{(0)}) + \frac{1}{b_4 n} \mathbf{E}(Y_n^2 m^{Y_n} N_n^{(0)}) \leq b_3 n c_{12} n + \frac{c_{15} n^3}{b_4 n},$$

the last inequality following from Lemma 3.2 (i) and (iii). Assembling these pieces yields that for  $1 \leq n \leq M$ ,

$$q_n \geq \left( c_{13} - c_3 b_1 - \frac{c_{17}}{b_2} - c_{12} b_3 - \frac{c_{15}}{b_4} \right) n^2 \geq \frac{c_{13}}{5} n^2,$$

if we choose  $b_2 := \frac{5c_{17}}{c_{13}}$ ,  $b_1 := \min\{\frac{c_{13}}{5c_3}, \frac{b_2}{2}\}$ ,  $b_4 := \frac{5c_{15}}{c_{13}}$  and  $b_3 := \min\{\frac{c_{13}}{5c_{12}}, \frac{b_4}{2}\}$  [ $M$  being large enough so that  $b_4 \leq \frac{\zeta(M)}{2}$ ].

Let us have another look at  $q_n := \mathbf{E}(Z_n m^{Z_n} L_n^{(0)} \mathbf{1}_{\{L_n^{(0)} \in [b_1 n^2, b_2 n^2], Z_n \in [b_3 n, b_4 n]\}})$ . On the event  $\{L_n^{(0)} \in [b_1 n^2, b_2 n^2], Z_n \in [b_3 n, b_4 n]\}$ , we have  $Z_n m^{Z_n} L_n^{(0)} \leq b_2 b_4 n^3 m^{Z_n}$ , so

$$q_n \leq b_2 b_4 n^3 \mathbf{E}(m^{Z_n} \mathbf{1}_{\{L_n^{(0)} \in [b_1 n^2, b_2 n^2], Z_n \in [b_3 n, b_4 n]\}}).$$

Since  $q_n \geq \frac{c_{13}}{5} n^2$ , this yields that for  $1 \leq n \leq M$ ,

$$\mathbf{E}(m^{Z_n} \mathbf{1}_{\{L_n^{(0)} \in [b_1 n^2, b_2 n^2], Z_n \in [b_3 n, b_4 n]\}}) \geq \frac{c_{37}}{n},$$

where  $c_{37} := \frac{c_{13}}{5b_2 b_4}$ . A fortiori,

$$\mathbf{E}(m^{Z_n} \mathbf{1}_{\{L_n^{(0)} \geq b_1 n^2, Z_n \in [b_3 n, b_4 n]\}}) \geq \frac{c_{37}}{n}.$$

On the other hand,

$$\begin{aligned} & \mathbf{E}(m^{Z_n} \mathbf{1}_{\{L_n^{(0)} \geq b_1 n^2, Z_n \in [b_3 n, b_4 n]\}}) \\ &= \mathbf{E}[m^{Z_n} \mathbf{1}_{\{Z_n \in [b_3 n, b_4 n]\}} \mathbf{P}(L_n^{(0)} \geq b_1 n^2 \mid Z_n)] \\ &\leq \mathbf{E}(m^{Z_n} \mathbf{1}_{\{Z_n \in [b_3 n, b_4 n]\}} \mathbf{1}_{\{\mathbf{P}(L_n^{(0)} \geq b_1 n^2 \mid Z_n) \geq b_5\}}) + b_5 \mathbf{E}(m^{Z_n} \mathbf{1}_{\{Z_n \in [b_3 n, b_4 n]\}}) \\ &\leq \mathbf{E}(m^{Z_n} \mathbf{1}_{\{Z_n \in [b_3 n, b_4 n]\}} \mathbf{1}_{\{\mathbf{P}(L_n^{(0)} \geq b_1 n^2 \mid Z_n) \geq b_5\}}) + \frac{b_5 c_3}{b_3 n}, \end{aligned}$$

the last inequality following from Lemma 3.1 (ii) (recalling that  $Z_0 \leq Y_0$ ). Consequently, for  $1 \leq n \leq M$ ,

$$\mathbf{E}(m^{Z_n} \mathbf{1}_{\{Z_n \in [b_3 n, b_4 n]\}} \mathbf{1}_{\{\mathbf{P}(L_n^{(0)} \geq b_1 n^2 \mid Z_n) \geq b_5\}}) \geq \frac{c_{37}}{n} - \frac{b_5 c_3}{b_3 n} = \frac{c_{37}}{2n},$$

if we choose  $b_5 := \frac{c_{37}b_3}{2c_3}$ . Let  $B_n := \{y \in [b_3n, b_4n] \cap \mathbb{Z} : \mathbf{P}(Z_n = y) > 0, \mathbf{P}(L_n^{(0)} \geq b_1n^2 \mid Z_n = y) \geq b_5\}$ . We have just proved that  $\mathbf{E}(m^{Z_n} \mathbf{1}_{\{Z_n \in B_n\}}) \geq \frac{c_{37}}{2n}$  for  $1 \leq n \leq M$ . This yields Lemma 4.4 by taking  $n = M$ .  $\square$

*Proof of Lemma 4.3.* (i) We first prove that there exist constants  $c_{33} \in (0, 1)$  and  $c_{38} \in (0, 1)$ , such that for any  $s \in [m, m + \frac{c_{33}}{M}]$ ,

$$(4.8) \quad \mathbf{E}(s^{W_0}) - (s-1)\mathbf{E}(W_0s^{W_0}) \geq c_{38}.$$

Let  $s \geq m$ . Since  $W_0 = Z_M \mathbf{1}_{\{Z_M \notin B_M\}}$  and  $y \geq c_{29}M$  for any  $y \in B_M$ ,

$$\begin{aligned} & [(s-1)\mathbf{E}(Z_Ms^{Z_M}) - \mathbf{E}(s^{Z_M})] - [(s-1)\mathbf{E}(W_0s^{W_0}) - \mathbf{E}(s^{W_0})] \\ &= \mathbf{E}\{[(s-1)Z_M - 1]s^{Z_M} + 1 \mid \mathbf{1}_{\{Z_M \in B_M\}}\} \\ &\geq ((m-1)c_{29}M - 1)\mathbf{E}(m^{Z_M} \mathbf{1}_{\{Z_M \in B_M\}}). \end{aligned}$$

We choose  $M$  to be larger than  $\frac{2}{(m-1)c_{29}}$ , which ensures that  $(m-1)c_{29}M - 1 \geq \frac{1}{2}(m-1)c_{29}M$ . Furthermore, we choose  $M$  to be sufficiently large so that we are entitled to apply Lemma 4.4 to have  $\mathbf{E}(m^{Z_M} \mathbf{1}_{\{Z_M \in B_M\}}) \geq \frac{c_{36}}{M}$ . Combining these inequalities, we get, for  $s \geq m$ ,

$$(4.9) \quad [(s-1)\mathbf{E}(Z_Ms^{Z_M}) - \mathbf{E}(s^{Z_M})] - [(s-1)\mathbf{E}(W_0s^{W_0}) - \mathbf{E}(s^{W_0})] \geq \frac{1}{2}(m-1)c_{29}c_{36}.$$

On the other hand, we claim the existence of constants  $c_{39} > 0$  and  $c_{40} \geq 1$  such that for all  $s \in [m, m + \frac{c_{39}}{M}]$ ,

$$(4.10) \quad (s-1)\mathbf{E}(Z_Ms^{Z_M}) - \mathbf{E}(s^{Z_M}) \leq c_{40}(s-m)M.$$

To prove our claim, let  $H_n(s) := \mathbf{E}(s^{Z_n})$  be the moment generating function of  $Z_n$ . We have, for  $s \geq m$ ,

$$\begin{aligned} (s-1)\mathbf{E}(Z_ns^{Z_n}) - \mathbf{E}(s^{Z_n}) &= s(s-1)H'_n(s) - H_n(s) \\ &\leq s(s-1)[H'_n(m) + (s-m)H''_n(s)] - H_n(m) \\ &\leq (s-m)(s+m-1)H'_n(m) + s(s-1)(s-m)H''_n(s), \end{aligned}$$

where, in the last inequality, we have used the fact that  $(m-1)\mathbf{E}(Z_nm^{Z_n}) - \mathbf{E}(m^{Z_n}) \leq 0$ . On the right-hand side, we have  $H'_n(m) \leq \frac{c_3}{m}$  (by Lemma 3.1 (ii) and the fact that  $Z_0 \leq Y_0$ ),

and use the existence of constants  $c_{39} > 0$  and  $c_{41} \geq 1$  (see Theorem 4 and Corollary 1 of [7]) such that  $\mathbf{E}(Z_n^2 s^{Z_n}) \leq c_{41} M$  for  $1 \leq n \leq M$  and  $s \in [m, m + \frac{c_{39}}{M}]$ ; this implies  $H_n''(s) \leq c_{41} M$ . Consequently, there exists a constant  $c_{40} \geq 1$  such that for all  $s \in [m, m + \frac{c_{39}}{M}]$  and  $1 \leq n \leq M$ ,

$$(4.11) \quad (s-1)\mathbf{E}(Z_n s^{Z_n}) - \mathbf{E}(s^{Z_n}) \leq c_{40}(s-m)M,$$

which obviously implies the claim in (4.10).

We choose  $c_{33} := \min\{\frac{1}{4c_{40}}(m-1)c_{29}c_{36}, c_{39}, \frac{1}{2}\}$ . If  $s \in [m, m + \frac{c_{33}}{M}]$ , then by (4.10) and (4.9),

$$(s-1)\mathbf{E}(W_0 s^{W_0}) - \mathbf{E}(s^{W_0}) \leq c_{40}(s-m)M - \frac{1}{2}(m-1)c_{29}c_{36} \leq -\frac{1}{4}(m-1)c_{29}c_{36}.$$

Setting  $c_{38} := \min\{\frac{1}{4}(m-1)c_{29}c_{36}, \frac{1}{2}\}$  yields (4.8).

(ii) We complete the proof of the lemma. Write  $F_n(t) = \mathbf{E}(t^{W_n})$  for  $t > 0$ . Let  $s := m + \frac{c_{33}}{M}$ . By (4.8),

$$F_0(s) - (s-1)sF_0'(s) \geq c_{38}.$$

By (3.2),  $F_{n+1}(s) - (s-1)sF_{n+1}'(s) = [F_n(s) - m(s-1)F_n'(s)]F_n(s)^{m-1} \geq F_n(s) - s(s-1)F_n'(s)$ , which implies that

$$F_n(s) - (s-1)sF_n'(s) \geq F_0(s) - (s-1)sF_0'(s) \geq c_{38}, \quad n \geq 0.$$

Using  $s(s-1) \geq m(m-1)$ , this yields:

$$F_n(s) \geq c_{38} + m(m-1)F_n'(s), \quad n \geq 0.$$

By convexity,  $F_n'(s) \geq \frac{F_n(s) - F_n(m)}{s-m} = \frac{M}{c_{33}}(F_n(s) - F_n(m))$ . Thus

$$F_n(s) \geq c_{38} + c_{42} M(F_n(s) - F_n(m)), \quad n \geq 0,$$

with  $c_{42} := \frac{m(m-1)}{c_{33}}$ . Since  $W_0 \leq Z_M \leq Y_M$ , we have  $F_n(m) \leq G_{n+M}(m)$ , and the latter is bounded, according to Lemma 3.1 (iv), by  $1 + \frac{c_6}{n+M}$ , a fortiori by  $1 + \frac{c_6}{n}$  (for  $n \geq 1$ ). Consequently,

$$F_n(s) \geq c_{38} + c_{42} M \left( F_n(s) - 1 - \frac{c_6}{n} \right), \quad n \geq 1;$$

in other words, we have

$$F_n(s) \leq \frac{c_{42}M(1 + \frac{c_6}{n}) - c_{38}}{c_{42}M - 1} = 1 + \frac{(1 - c_{38}) + c_{42}M\frac{c_6}{n}}{c_{42}M - 1}.$$

We now take  $n = n_0 := \lfloor c_{34}M \rfloor$ , with  $c_{34} := \max\{\frac{4c_6c_{42}}{c_{38}}, 2\}$ . This choice ensures that  $c_{42}M\frac{c_6}{n} \leq c_{42}M\frac{c_6}{\frac{1}{2}c_{34}M} \leq \frac{1}{2}c_{38}$ , so  $F_{n_0}(s) \leq 1 + \frac{1 - \frac{1}{2}c_{38}}{c_{42}M - 1}$ . This implies that

$$\frac{m}{s} F_{n_0}(s)^{m-1} = \frac{F_{n_0}(s)^{m-1}}{1 + \frac{c_{33}}{mM}} \leq \frac{1}{1 + \frac{c_{33}}{mM}} \left(1 + \frac{1 - \frac{1}{2}c_{38}}{c_{42}M - 1}\right)^{m-1}.$$

Recall that  $c_{42} = \frac{m(m-1)}{c_{33}}$ . We choose  $M_0$  so large that  $\frac{1}{1 + \frac{c_{33}}{mM}} \left(1 + \frac{1 - \frac{1}{2}c_{38}}{c_{42}M - 1}\right)^{m-1} \leq 1 - \frac{1}{3}c_{38}\frac{c_{33}}{mM}$  for all  $M \geq M_0$ . Consequently, for  $M \geq M_0$ ,

$$\frac{m}{s} F_{n_0}(s)^{m-1} \leq 1 - \frac{1}{3}c_{38}\frac{c_{33}}{mM},$$

proving Lemma 4.3. □

## 4.2 Step 2. Study of $\mathbf{P}(Z_n \geq \ell + 1)$

The goal in this step is to prove Lemma 4.6 below concerning  $\mathbf{P}(Z_n \geq \ell + 1)$ . We start with an upper bound for the moment generating function of an arbitrary Derrida–Retaux system. The bound is simple, but not sharp in general.

**Lemma 4.5.** *Let  $(X_n, n \geq 0)$  be a Derrida–Retaux system. Let  $s > m$  and  $\theta \in (0, 1)$ . If  $\frac{m}{s}[\mathbf{E}(s^{X_0})]^{m-1} \leq \theta$ , then for  $n \geq 0$ ,*

$$\mathbf{E}(s^{X_n}) \leq 1 + (s - m)\theta^n.$$

*Proof.* Let  $s > m$ . Consider the sequence  $(x_n, n \geq 0)$  defined by  $x_0 := \mathbf{E}(s^{X_0}) - 1$  and  $x_{i+1} := f(x_i)$  for  $i \geq 0$ , where  $f(x) := \frac{(1+x)^m}{s} - \frac{1}{s}$  for  $x \geq 0$ .

Since  $f'(x) = \frac{m}{s}(1+x)^{m-1}$ , we have, for all  $x \in [0, x_0]$ ,  $f'(x) \leq f'(x_0) \leq \theta$ . By convexity, for  $i \geq 0$ ,  $x_{i+1} = f(x_i) - f(0) \leq f'(x_i)x_i \leq \theta x_i$ . It follows that

$$(4.12) \quad x_n \leq \theta^n x_0, \quad n \geq 0.$$

On the other hand, by definition of the Derrida–Retaux system, for  $i \geq 0$ ,  $\mathbf{E}(s^{X_{i+1}}) = \frac{1}{s}[\mathbf{E}(s^{X_i})]^m + (1 - \frac{1}{s})[\mathbf{P}(X_i = 0)]^m \leq \frac{1}{s}[\mathbf{E}(s^{X_i})]^m + (1 - \frac{1}{s})$ , i.e.,  $\mathbf{E}(s^{X_{i+1}}) - 1 \leq f(\mathbf{E}(s^{X_i}) - 1)$ . Iterating the inequality gives that

$$\mathbf{E}(s^{X_n}) - 1 \leq x_n,$$

which, according to (4.12), is bounded by  $\theta^n x_0$ . By assumption,

$$\mathbf{E}(s^{X_0}) \leq \left(\frac{s\theta}{m}\right)^{1/(m-1)} \leq \left(\frac{s}{m}\right)^{1/(m-1)} \leq \frac{s}{m},$$

so  $x_0 = \mathbf{E}(s^{X_0}) - 1 \leq \frac{s}{m} - 1 \leq s - m$ . Lemma 4.5 follows immediately.  $\square$

We have now all the ingredients to prove the main result for the Derrida–Retaux system  $(Z_n, n \geq 0)$ , stated as follows.

**Lemma 4.6.** *Assume  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ . There exist constants  $c_{43} > 0$ ,  $c_{44} > 0$  and sufficiently large  $M_0$  such that for all integers  $n \geq M \geq M_0$ ,*

$$(4.13) \quad \mathbf{P}(Z_n \geq \ell + 1) \leq c_{43} \left( \frac{1}{M^2} + \frac{1}{M} e^{-c_{44} \frac{n}{M}} \right) m^{-\ell}, \quad \ell \geq 0.$$

*Proof.* Let  $M \geq M_0$ , where  $M_0$  is sufficiently large such that Lemma 4.3 applies to  $M \geq M_0$ . Let  $s = s(M) := m + \frac{c_{33}}{M}$  and  $n_0 = n_0(M) := \lfloor c_{34} M \rfloor$ . By Lemma 4.3,

$$\frac{m}{s} [\mathbf{E}(s^{W_{n_0}})]^{m-1} \leq 1 - \frac{c_{35}}{M}.$$

This entitles us to apply Lemma 4.5 to  $(W_{n+n_0}, n \geq 0)$ , to see that

$$\mathbf{E}(m^{W_{n+n_0}}) - 1 \leq \mathbf{E}(s^{W_{n+n_0}}) - 1 \leq \frac{c_{33}}{M} \left(1 - \frac{c_{35}}{M}\right)^n, \quad n \geq 0.$$

Consider the system  $(\tilde{Z}_n := Z_{n+M}, n \geq 0)$ . We define  $\tilde{L}_n^{(i)}$  the number of open paths at generation  $n$  of this system with initial value  $i$ , exactly as  $N_n^{(i)}$  for  $(X_n)$  in (2.3).<sup>12</sup> Let

$$\tilde{L}_n^{(B_M)} := \sum_{i \in B_M} \tilde{L}_n^{(i)}.$$

---

<sup>12</sup>Notation: The letter  $L$  is used for the number of open paths for  $(Z_n)$ , and  $\tilde{L}$  for  $(\tilde{Z}_n)$ .

Applying Lemma 4.2 to  $(\tilde{Z}_k, k \geq 0)$  and  $(W_k, k \geq 0)$ , we get, for any  $\ell \geq 1$ ,

$$\mathbf{P}(Z_{n+M} = \ell, \tilde{L}_n^{(B_M)} = 0) \leq \mathbf{P}(W_n = \ell) \leq \frac{\mathbf{E}(m^{W_n} - 1)}{m^\ell - 1}.$$

Hence, for  $n \geq 2n_0$ , we have, for  $\ell \geq 1$ ,

$$\begin{aligned} \mathbf{P}(Z_{n+M} = \ell, \tilde{L}_n^{(B_M)} = 0) &\leq \frac{1}{m^\ell - 1} \frac{c_{33}}{M} \left(1 - \frac{c_{35}}{M}\right)^{n/2} \\ &\leq \frac{1}{m^\ell - 1} \frac{c_{33}}{M} \exp\left(-\frac{c_{35}n}{2M}\right). \end{aligned}$$

We now look at the situation  $Z_{n+M} = \ell$  with  $\tilde{L}_n^{(B_M)} \geq 1$ . On the event  $\{\tilde{L}_n^{(B_M)} \geq 1\}$ , we choose a vertex  $v_M$  in generation  $M$  of the system  $(Z_n)$  such that  $Z(v_M) \in B_M$  and that there is an open path from  $v_M$  to generation  $n + M$ .<sup>13</sup> Call  $L^{(0)}(v_M)$  the number of open paths in the system  $(Z_n)$  from the initial generation to this particular vertex  $v_M$  at generation  $M$ , with initial value 0. Obviously,

$$\begin{aligned} L_{n+M}^{(0)} \mathbf{1}_{\{Z_{n+M}=\ell\}} &\geq L^{(0)}(v_M) \mathbf{1}_{\{\tilde{L}_n^{(B_M)} \geq 1, Z_{n+M}=\ell\}} \\ &\geq c_{31} M^2 \mathbf{1}_{\{\tilde{L}_n^{(B_M)} \geq 1, Z_{n+M}=\ell\}} \mathbf{1}_{\{L^{(0)}(v_M) \geq c_{31} M^2\}}. \end{aligned}$$

We take expectation on both sides. On the right-hand side, we first take conditional expectation with respect to the sigma-algebra  $\mathcal{F}_M^Z$  generated by the values of the system  $(Z_n)$  at generation  $M$ , and note that  $\tilde{L}_n^{(B_M)}$ ,  $Z_{n+M}$ , and  $v_M$  (on the event  $\{\tilde{L}_n^{(B_M)} \geq 1\}$ ) are all  $\mathcal{F}_M^Z$ -measurable. By definition of  $B_M$ ,  $\min_{y \in B_M} \mathbf{P}(L_M^{(0)} \geq c_{31} M^2 \mid Z_M = y) \geq c_{32}$ . This leads to, for all  $\ell \geq 1$ ,

$$\begin{aligned} \mathbf{E}(L_{n+M}^{(0)} \mathbf{1}_{\{Z_{n+M}=\ell\}}) &\geq c_{31} M^2 \mathbf{E}[\mathbf{1}_{\{\tilde{L}_n^{(B_M)} \geq 1, Z_{n+M}=\ell\}} \mathbf{P}(L^{(0)}(v_M) \geq c_{31} M^2 \mid \mathcal{F}_M^Z)] \\ &\geq c_{31} c_{32} M^2 \mathbf{P}(\tilde{L}_n^{(B_M)} \geq 1, Z_{n+M} = \ell), \end{aligned}$$

giving an upper bound for  $\mathbf{P}(\tilde{L}_n^{(B_M)} \geq 1, Z_{n+M} = \ell)$ . Consequently, for  $\ell \geq 1$ ,

$$\begin{aligned} \mathbf{P}(Z_{n+M} = \ell) &= \mathbf{P}(Z_{n+M} = \ell, \tilde{L}_n^{(B_M)} \geq 1) + \mathbf{P}(Z_{n+M} = \ell, \tilde{L}_n^{(B_M)} = 0) \\ &\leq \frac{\mathbf{E}(L_{n+M}^{(0)} \mathbf{1}_{\{Z_{n+M}=\ell\}})}{c_{31} c_{32} M^2} + \frac{1}{m^\ell - 1} \frac{c_{33}}{M} \exp\left(-\frac{c_{35}n}{2M}\right). \end{aligned}$$

---

<sup>13</sup>If there are several such vertices, we can take for example the leftmost one in the lexicographic order.



This yields that for  $\ell \geq 1$ ,

$$\mathbf{P}(Z_{n+M} \geq \ell) \leq \frac{\mathbf{E}(L_{n+M}^{(0)} \mathbf{1}_{\{Z_{n+M} \geq \ell\}})}{c_{31}c_{32}M^2} + \frac{c_{33}}{M} \exp\left(-\frac{c_{35}n}{2M}\right) \sum_{j=\ell}^{\infty} \frac{1}{m^j - 1}.$$

Since  $(Z_i, i \geq 0)$  is subcritical or critical, it follows from (2.13) that  $\mathbf{E}(L_{n+M}^{(0)} \mathbf{1}_{\{Z_{n+M} \geq \ell\}}) \leq \frac{c_1}{m^\ell}$ ; this implies the desired inequality (4.13) when  $n \geq (2c_{34} + 1)M$  (with  $M \geq M_0$ ).

The case  $M \leq n < (2c_{34} + 1)M$  (with  $M \geq M_0$ ) is easy: we have  $\mathbf{E}(m^{Z_n} \mathbf{1}_{\{Z_n \geq 1\}}) \leq \mathbf{E}(m^{Y_n} \mathbf{1}_{\{Y_n \geq 1\}}) \leq \frac{c_6}{n}$  (by Lemma 3.1 (iv)), thus for  $\ell \geq 0$ ,

$$\mathbf{P}(Z_n \geq \ell + 1) \leq \frac{\mathbf{E}(m^{Z_n} \mathbf{1}_{\{Z_n \geq 1\}})}{m^{\ell+1}} \leq \frac{c_6}{n} \frac{1}{m^{\ell+1}} \leq \frac{c_6}{M} \frac{1}{m^{\ell+1}},$$

which implies again the desired inequality (4.13) because  $n < (2c_{34} + 1)M$ . Lemma 4.6 is proved.  $\square$

### 4.3 Step 3. Proof of Proposition 4.1

Assume  $\mathbf{E}(Y_0^5 m^{Y_0}) < \infty$ . Let  $n$  and  $M := \lfloor c_{44} \frac{n}{\log n} \rfloor$  (where  $c_{44} > 0$  is the constant in Lemma 4.6) satisfy  $n \geq M \geq M_0$ , where  $M_0$  is such that Lemma 4.6 applies to  $n$  and  $M$  and that  $M \geq \frac{c_{44}}{2} \frac{n}{\log n}$ . By Lemma 4.6, for  $\ell \geq 0$ ,

$$\begin{aligned} \mathbf{P}(Z_n \geq \ell + 1) &\leq c_{43} \left( \frac{1}{M^2} + \frac{1}{M} e^{-c_{44} \frac{n}{M}} \right) m^{-\ell} \\ &\leq c_{43} \left( \frac{4(\log n)^2}{c_{44}^2 n^2} + \frac{2 \log n}{c_{44} n} e^{-\log n} \right) m^{-\ell} \\ (4.14) \quad &\leq \frac{c_{45}(\log n)^2}{n^2} \frac{1}{m^\ell}, \end{aligned}$$

with  $c_{45} := \frac{4c_{43}}{c_{44}^2} + \frac{2c_{43}}{c_{44}}$ .

Recall from (4.3) the definition:  $\zeta(M) = -\log \mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 > M\}})$ . By the Markov inequality,  $\mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 > M\}}) \leq \frac{c_{46}}{M^2}$ , where  $c_{46} := \mathbf{E}(Y_0^5 m^{Y_0}) < \infty$ . So  $\zeta(M) \geq \log\left(\frac{M^2}{c_{46}}\right)$  (and  $\zeta(M)$  can be possibly infinite). Let  $N_n^{(i)}$  be as before the number of open paths until generation  $n$  in the system  $(Y_n)$  with initial value  $i$  (replacing  $(X_n)$  by  $(Y_n)$  in (2.3)) and let

$$N_n^{(>M\zeta(M))} := \sum_{i > M\zeta(M)} N_n^{(i)}.$$

By (3.8) and the Fubini–Tonelli theorem,

$$\mathbf{E}((Y_n + 1)m^{Y_n} N_n^{(>M\zeta(M))}) = \mathbf{E}((Y_0 + 1)m^{Y_0} \mathbf{1}_{\{Y_0 > M\zeta(M)\}}) \prod_{i=0}^{n-1} G_i(m)^{m-1}.$$

By Lemma 3.1 (v),  $\prod_{i=0}^{n-1} G_i(m)^{m-1} \leq c_8 n^2$ , whereas

$$\mathbf{E}((Y_0 + 1)m^{Y_0} \mathbf{1}_{\{Y_0 > M\zeta(M)\}}) \leq 2 \mathbf{E}(Y_0 m^{Y_0} \mathbf{1}_{\{Y_0 > M\zeta(M)\}}) \leq \frac{2c_{46}}{(M\zeta(M))^4},$$

with  $c_{46} := \mathbf{E}(Y_0^5 m^{Y_0})$  as before. Hence

$$\mathbf{E}((Y_n + 1)m^{Y_n} N_n^{(>M\zeta(M))}) \leq \frac{2c_{46}}{(M\zeta(M))^4} c_8 n^2.$$

Therefore,

$$\begin{aligned} \mathbf{P}(Y_n \geq \ell + 1, N_n^{(>M\zeta(M))} \geq 1) &\leq m^{-(\ell+1)} \mathbf{E}((Y_n + 1)m^{Y_n} N_n^{(>M\zeta(M))}) \\ (4.15) \qquad \qquad \qquad &\leq m^{-(\ell+1)} \frac{2c_{46}}{(M\zeta(M))^4} c_8 n^2, \end{aligned}$$

Applying Lemma 4.2 to  $(Y_k, k \geq 0)$  and  $(Z_k, k \geq 0)$  yields that, for  $\ell \geq 0$ ,

$$\begin{aligned} \mathbf{P}(Y_n \geq \ell + 1) &\leq \mathbf{P}(Y_n \geq \ell + 1, N_n^{(>M\zeta(M))} \geq 1) + \mathbf{P}(Z_n \geq \ell + 1) \\ &\leq m^{-(\ell+1)} \frac{2c_{46}}{(M\zeta(M))^4} c_8 n^2 + \frac{c_{45}(\log n)^2}{n^2} \frac{1}{m^\ell}, \end{aligned}$$

by means of (4.15) and (4.14). Since  $\zeta(M) \geq \log(\frac{M^2}{c_{46}})$  and  $M := \lfloor \frac{c_{44}n}{\log n} \rfloor$ , this implies Proposition 4.1 for sufficiently large  $n$  (say  $n \geq n'_0$ ) and all  $\ell \geq 0$ . The case  $2 \leq n < n'_0$  follows simply from the Markov inequality and Lemma 3.1 (i).  $\square$

## 5 Proof of Theorem 1.2: lower bound

The lower bound in Theorem 1.2 is a simple consequence of the following proposition.

**Proposition 5.1.** *Assume  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ . Then for any  $\lambda > 0$ , there exists a constant  $\nu > 0$  such that for all sufficiently large integer  $n$ ,*

$$(5.1) \qquad \mathbf{P}(Y_n \geq \ell + 1) \geq \frac{1}{n^2(\log n)^\nu} \frac{1}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z}.$$

*Outline of the proof of Proposition 5.1.* The technical ingredient in the proof is stated as Lemma 5.2 below: there exist constants  $\theta > 0$  and  $c_{48} > 0$  such that for all sufficiently large integer  $n$ ,

$$(5.4) \quad \sum_{k=0}^{k_n(\theta)-1} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) \geq \frac{c_{48}}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z},$$

where

$$k_n(\theta) := \min \left\{ i \geq 0 : \mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 \geq i+1\}}) \leq \frac{\theta}{(\log n)^2} \right\},$$

and  $S_n^{(k,\ell)}$  denotes as before (see (2.7), with the system  $(X_n)$  replaced by  $(Y_n)$  there) the number of  $(k, \ell)_n$ -pivotal vertices in  $\mathbb{T}_n$ .

Let us first describe how (5.4) helps in the proof of Proposition 5.1. Using a simple relation between pivotal vertices and open paths, we deduce from (5.4) that for all sufficiently large  $n$ , there exist a constant  $\varrho > 0$  and a non-random integer  $i^* = i^*(n) \in [0, c_{55} \log \log n]$  such that

$$(5.11) \quad \mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell+1\}}) \geq \frac{1}{(\log n)^e} \frac{1}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z},$$

where, for any integer  $i \geq 0$ ,  $N_n^{(i)}$  denotes as before (see (2.3)) the number of open paths in  $\mathbb{T}_n$  to generation  $n$  with initial value  $i$ , whereas  $c_{55} > 0$  is a suitable constant. [This is stated as Proposition 5.4, which is a weaker form of inequality (2.13) in the opposite direction.]

Once (5.11) is established, we proceed by coupling  $(Y_i, i \geq 0)$  with some appropriate system  $(X_i, i \geq 0)$  (depending on  $n$ ) such that  $Y_i \leq X_i$  for all  $i \geq 0$ . A general inequality (Fact 5.6 below) borrowed from [4] for supercritical Derrida–Retaux systems says that

$$\mathbf{E}(X_i) \leq 3,$$

for all  $i$  smaller than a suitable constant multiple of  $n$ ; moreover, a useful coupling inequality (called the “bridge inequality”; see Fact 5.5 below), also borrowed from [4], says that

$$\mathbf{E}(N_n \mathbf{1}_{\{N_n \geq r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}) \leq \frac{2}{m^{k+\ell+1} \eta} \mathbf{E}(X_{n+k+\ell+1}),$$

for all  $n, k, \ell$  and  $r$  lying in suitable domains, where  $N_n$  denotes, as in (2.3), the number of open paths in  $\mathbb{T}_n$  to generation  $n$  for  $(Y_i)$ . Adjusting the parameters appropriately, this leads to the following estimate: for any  $a > 0$  and  $\lambda > 0$ , and all sufficiently large  $n$ ,

$$\mathbf{E}(N_n \mathbf{1}_{\{N_n \geq r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}) \leq \frac{1}{n^a} \frac{1}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z},$$

where  $r = r(n)$  is (approximately) a constant multiple of  $n^2 \log n$ . Together with (5.11), and using the trivial inequality  $N_n \geq N_n^{(i^*)}$ , this readily yields that

$$\mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{N_n^{(i^*)} < r\}} \mathbf{1}_{\{Y_n \geq \ell + 1\}}) \geq \frac{1}{(\log n)^e} \frac{1}{m^\ell} - \frac{1}{n^a} \frac{1}{m^\ell} \geq \frac{1}{2} \frac{1}{(\log n)^e} \frac{1}{m^\ell},$$

for all sufficiently large  $n$ , uniformly in  $\ell \in [0, \lambda n] \cap \mathbb{Z}$ . Consequently, for all sufficiently large  $n$  and uniformly in  $\ell \in [0, \lambda n] \cap \mathbb{Z}$ ,

$$\mathbf{P}(Y_n \geq \ell + 1) \geq \frac{1}{r} \mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{N_n^{(i^*)} < r\}} \mathbf{1}_{\{Y_n \geq \ell + 1\}}) \geq \frac{1}{2r} \frac{1}{(\log n)^e} \frac{1}{m^\ell},$$

from which Proposition 5.1 follows.

Let us now say some words about the proof of (5.4), which exploits the technique of pivotal vertices in Section 2 and extends some technique to a more general context. We write

$$\sum_{k=0}^{k_n(\theta)-1} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) = \sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) - \sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}).$$

By Theorem 2.2,  $\sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) = \mu_n(\ell)$ , which is greater than or equal to  $\frac{c_{47}}{m^\ell}$  for some constant  $c_{47} > 0$  (by Lemma 3.4; see (5.3) below). It remains to check that

$$\sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) \leq \frac{c_{47}}{2} \frac{1}{m^\ell}.$$

We make a decomposition of the system at an intermediate generation  $M = M(n) := \lfloor \frac{n}{1+c_{49}} \rfloor$ , where  $c_{49} > 0$  is a (large) constant.<sup>14</sup> Contributions to  $\mathbf{E}(S_n^{(k,\ell)})$  from the initial generation to generation  $M$  are controlled (via the forthcoming Lemma 5.3) using the number of open paths to generation  $M$ , whereas contributions from generation  $M$  to generation  $n$  are investigated by means of Proposition 4.1.

The rest of the section is devoted to the (rigorous) proof of Proposition 5.1, presented in three steps. The first step is the technical part of the section; it gives precision on the number of pivotal vertices as in (5.4) (Lemma 5.2). The second step gives a lower bound for  $\max_{i \in [0, c_{55} \log \log n]} \mathbf{E}(N_n^{(i)} \mathbf{1}_{\{Y_n \geq \ell + 1\}})$  as in (5.11) (Proposition 5.4). The third step completes the proof of Proposition 5.1.  $\square$

---

<sup>14</sup>It is natural to introduce such a decomposition at an intermediate generation  $M$ , because the system at generation  $M$  has “nicer” properties than at the initial generation, which allows to use some renormalization idea starting from generation  $M$ . A similar approach was already used in the proof of the upper bound (Proposition 4.1).

## 5.1 Step 1. Precision on the number of pivotal vertices

Let  $(Y_n, n \geq 0)$  be a critical Derrida–Retaux system:  $\mathbf{E}(m^{Y_0}) = (m-1)\mathbf{E}(Y_0 m^{Y_0}) < \infty$ . Recall from (2.6) that

$$(5.2) \quad \begin{aligned} \mu_n(k) &:= m^{-k} \mathbf{E}([1 - (m-1)Y_n]m^{Y_n} \mathbf{1}_{\{Y_n \leq k\}}) \\ &= m^{-k} \mathbf{E}([(m-1)Y_n - 1]m^{Y_n} \mathbf{1}_{\{Y_n \geq k+1\}}), \end{aligned}$$

the equality (5.2) being a consequence of the fact that the system is critical, as we have already observed before. We have also noted (see (2.12)) that

$$0 \leq \mu_n(k) \leq m^{-k}, \quad k \geq 0, n \geq 0.$$

On the other hand, by Lemma 3.4, if  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ , then for any  $\lambda > 0$ , there exists a constant  $c_{47} > 0$  such that for all sufficiently large integer  $n$  and all  $k \in [0, \lambda n] \cap \mathbb{Z}$ ,

$$(5.3) \quad \mu_n(k) \geq \frac{c_{47}}{m^k}.$$

Let  $S_n^{(k,\ell)}$  denote, as in (2.7), the number of  $(k, \ell)_n$ -pivotal vertices in  $\mathbb{T}_n$ . The aim of this step is to establish the following estimate.

**Lemma 5.2.** *Assume  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ . Let  $\lambda > 0$ . There exist constants  $\theta > 0$  and  $c_{48} > 0$  such that for all sufficiently large integer  $n$ ,*

$$(5.4) \quad \sum_{k=0}^{k_n(\theta)-1} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) \geq \frac{c_{48}}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z},$$

where

$$k_n(\theta) := \min \left\{ i \geq 0 : \mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 \geq i+1\}}) \leq \frac{\theta}{(\log n)^2} \right\}.$$

The proof of the lemma relies on a simple preliminary result, stated as follows.

**Lemma 5.3.** *For  $n \geq 0$  and  $i \geq k \geq 0$ ,*

$$\mathbf{E}[(Y_n + 1)m^{Y_n} N_n^{(i)}] \geq m^{i-k} \mathbf{P}(Y_0 = i) \sum_{\ell=0}^{\infty} (\ell + 1)m^\ell \mathbf{E}(S_n^{(k,\ell)}).$$

*Proof of Lemma 5.3.* Let  $n \geq 0$ ,  $i \geq k \geq 0$  and  $\ell \geq 0$  be integers. Consider  $\sum_{v \in A_n^{(k,\ell)}} \mathbf{1}_{\{Y(v)=i\}}$ . We have, by symmetry,  $\mathbf{E}(\sum_{v \in A_n^{(k,\ell)}} \mathbf{1}_{\{Y(v)=i\}}) = \mathbf{P}(Y_0 = i) \mathbf{E}(S_n^{(k,\ell)})$ . On the other hand,  $\sum_{v \in A_n^{(k,\ell)}} \mathbf{1}_{\{Y(v)=i\}} \leq N_n^{(i)} \mathbf{1}_{\{Y_n=i-k+\ell\}}$ . Hence

$$(5.5) \quad \mathbf{E}(N_n^{(i)} \mathbf{1}_{\{Y_n=i-k+\ell\}}) \geq \mathbf{P}(Y_0 = i) \mathbf{E}(S_n^{(k,\ell)}), \quad i \geq k \geq 0.$$

Writing

$$\begin{aligned} \mathbf{E}((Y_n + 1)m^{Y_n} N_n^{(i)}) &= \sum_{j=0}^{\infty} (j+1)m^j \mathbf{E}(N_n^{(i)} \mathbf{1}_{\{Y_n=j\}}) \\ &= \sum_{\ell=-(i-k)}^{\infty} (i-k+\ell+1)m^{i-k+\ell} \mathbf{E}(N_n^{(i)} \mathbf{1}_{\{Y_n=i-k+\ell\}}), \end{aligned}$$

this yields that

$$\begin{aligned} \mathbf{E}[(Y_n + 1)m^{Y_n} N_n^{(i)}] &\geq \sum_{\ell=-(i-k)}^{\infty} (i-k+\ell+1)m^{i-k+\ell} \mathbf{P}(Y_0 = i) \mathbf{E}(S_n^{(k,\ell)}) \\ &= m^{i-k} \mathbf{P}(Y_0 = i) \sum_{\ell=-(i-k)}^{\infty} (i-k+\ell+1)m^{\ell} \mathbf{E}(S_n^{(k,\ell)}). \end{aligned}$$

Since  $i - k \geq 0$ , this yields the desired inequality.  $\square$

We now have all the ingredients for the proof of Lemma 5.2.

*Proof of Lemma 5.2.* Fix  $\lambda > 0$ . We write

$$\sum_{k=0}^{k_n(\theta)-1} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) = \sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) - \sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}).$$

On the right-hand side,  $\sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)})$  equals  $\mu_n(\ell)$  (by Theorem 2.2), which is greater than or equal to  $\frac{c_{47}}{m^\ell}$  (by (5.3)) for all sufficiently large  $n$  and uniformly in  $\ell \in [0, \lambda n] \cap \mathbb{Z}$ . The proof of the lemma boils down to verifying that for some  $\theta > 0$  and all large  $n$ ,

$$(5.6) \quad \sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \mathbf{E}(S_n^{(k,\ell)}) \leq \frac{c_{47}}{2} \frac{1}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z}.$$

Let  $n > M \geq 0$  be integers. We extend the notion of pivotal vertices as follows: a vertex  $v$  with  $|v| = M$  is said to be  $(k, \ell)_{M,n}$ -pivotal if it is  $(k, \ell)_{n-M}$ -pivotal (in the sense of Definition 2.1) for the system  $(Y_{M+i}, i \geq 0)$ ; in other words,  $k + \xi_M(v) + \xi_{M+1}(v) + \cdots + \xi_{n-1}(v) = n - M + \ell$  and

$$k + \xi_M(v) + \xi_{M+1}(v) + \cdots + \xi_i(v) \geq i - M + 1, \quad \forall M \leq i \leq n - 1,$$

where, for  $i \geq M$ ,  $\xi_i(v) := \sum_{y \in \text{bro}(v_i)} Y(y)$ , with  $\text{bro}(v_i)$  denoting the set of the ‘‘brothers’’ of  $v_i$  (notation in agreement with (2.1) if  $M = 0$ ); here,  $v_i$  is the unique descendant of  $v$  at generation  $i$  (with  $v_i := v$  if  $i = M$ ). We write  $A_{M,n}^{(k,\ell)}$  for the set of all  $(k, \ell)_{M,n}$ -pivotal vertices in  $\mathbb{T}_n$  (so  $A_n^{(k,\ell)} = A_{0,n}^{(k,\ell)}$ ), and  $S_{M,n}^{(k,\ell)} := |A_{M,n}^{(k,\ell)}|$ . By definition,

$$\mathbf{1}_{\{\mathbf{e}_0 \in A_n^{(k,\ell)}\}} = \sum_{j=0}^{\infty} \mathbf{1}_{\{\mathbf{e}_0 \in A_M^{(k,j)}\}} \mathbf{1}_{\{\mathbf{e}_M \in A_{M,n}^{(j,\ell)}\}}.$$

A similar idea has already been used in (2.10). As in Section 2, this implies that

$$\mathbf{E}(S_n^{(k,\ell)}) = m^n \mathbf{E}(\mathbf{1}_{\{\mathbf{e}_0 \in A_n^{(k,\ell)}\}}) = m^n \sum_{j=0}^{\infty} \mathbf{P}(\mathbf{e}_0 \in A_M^{(k,j)}, \mathbf{e}_M \in A_{M,n}^{(j,\ell)});$$

by independence of  $\{\mathbf{e}_0 \in A_M^{(k,j)}\}$  and  $\{\mathbf{e}_M \in A_{M,n}^{(j,\ell)}\}$ , this leads to:

$$\mathbf{E}(S_n^{(k,\ell)}) = m^n \sum_{j=0}^{\infty} \mathbf{P}(\mathbf{e}_0 \in A_M^{(k,j)}) \mathbf{P}(\mathbf{e}_M \in A_{M,n}^{(j,\ell)}) = \sum_{j=0}^{\infty} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}).$$

Since  $S_{M,n}^{(j,\ell)} = 0$  for  $j > n - M + \ell$ , we obtain that

$$(5.7) \quad \mathbf{E}(S_n^{(k,\ell)}) = \sum_{j=0}^{n-M+\ell} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}).$$

As such, the proof of (5.6) will be complete if we are able to show that it is possible to choose an appropriate value of  $M = M(n)$  such that for some  $\theta > 0$ , all sufficiently large  $n$ , and all  $\ell \in [0, \lambda n] \cap \mathbb{Z}$ ,

$$(5.8) \quad \sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \sum_{j=0}^{n-M+\ell-1} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}) \leq \frac{c_{47}}{4m^\ell},$$

$$(5.9) \quad \sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \mathbf{E}(S_M^{(k,j=n-M+\ell)}) \mathbf{E}(S_{M,n}^{(j=n-M+\ell,\ell)}) \leq \frac{c_{47}}{4m^\ell}.$$

Let us first check (5.9). By the trivial inequalities  $\sum_{k=k_n(\theta)}^{\infty}(\cdots) \leq \sum_{k=0}^{\infty}(\cdots)$  and  $S_{M,n}^{(n-M+\ell,\ell)} \leq m^{n-M}$ , we get

$$\begin{aligned} \sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \mathbf{E}(S_M^{(k,n-M+\ell)}) \mathbf{E}(S_{M,n}^{(n-M+\ell,\ell)}) &\leq m^{n-M} \sum_{k=0}^{\infty} \mu_0(k) \mathbf{E}(S_M^{(k,n-M+\ell)}) \\ &= m^{n-M} \mu_M(n-M+\ell), \end{aligned}$$

the last equality being a consequence of Theorem 2.2. By Lemma 3.1 (iii), under the assumption  $\mathbf{E}(Y_0^3 m^{Y_0}) < \infty$ , we have  $\mathbf{E}(Y_j^2 m^{Y_j}) \leq c_4 j$  (for  $j \geq 1$ ), so by the Markov inequality, there exists a constant  $c_{49} > 0$  such that for  $j \geq 1$ ,

$$\max_{i \geq c_{49}j} m^i \mu_j(i) = m^{\lceil c_{49}j \rceil} \mu_j(\lceil c_{49}j \rceil) = \mathbf{E}[\left((m-1)Y_j - 1\right) m^{Y_j} \mathbf{1}_{\{Y_j \geq \lceil c_{49}j \rceil + 1\}}] \leq \frac{c_{47}}{4}.$$

We choose

$$(5.10) \quad M := \left\lfloor \frac{n}{1+c_{49}} \right\rfloor.$$

Then  $m^{n-M+\ell} \mu_M(n-M+\ell) \leq \frac{c_{47}}{4}$ , proving (5.9).

It remains to check (5.8). We bound  $\mathbf{E}(S_{M,n}^{(j,\ell)})$ . Let us write, for brevity,  $\xi_i := \xi_i(\mathbf{e}_0)$ ,  $i \geq 0$ . For  $0 \leq j < n-M+\ell$ ,

$$\begin{aligned} \mathbf{1}_{\{\mathbf{e}_M \in A_{M,n}^{(j,\ell)}\}} &= \sum_{i=0}^{j \wedge (n-M-1)} \sum_{r=1}^{n-M+\ell-j} \mathbf{1}_{\{\xi_M=0, \dots, \xi_{M+i-1}=0, \xi_{M+i}=r\}} \mathbf{1}_{\{\mathbf{e}_{M+i+1} \in A_{M+i+1,n}^{(j-i+r-1,\ell)}\}} \\ &\leq \sum_{i=0}^{j \wedge (n-M-1)} \sum_{r=1}^{n-M+\ell-j} \mathbf{1}_{\{\xi_{M+i}=r\}} \mathbf{1}_{\{\mathbf{e}_{M+i+1} \in A_{M+i+1,n}^{(j-i+r-1,\ell)}\}}. \end{aligned}$$

Hence

$$\begin{aligned} \mathbf{E}(S_{M,n}^{(j,\ell)}) &= m^{n-M} \mathbf{P}(\mathbf{e}_M \in A_{M,n}^{(j,\ell)}) \\ &\leq m^{n-M} \sum_{i=0}^{j \wedge (n-M-1)} \sum_{r=1}^{n-M+\ell-j} \mathbf{P}(\xi_{M+i} = r) \mathbf{P}(\mathbf{e}_{M+i+1} \in A_{M+i+1,n}^{(j-i+r-1,\ell)}) \\ &= \sum_{i=0}^{j \wedge (n-M-1)} \sum_{r=1}^{n-M+\ell-j} \mathbf{P}(\xi_{M+i} = r) m^{i+1} \mathbf{E}(S_{M+i+1,n}^{(j-i+r-1,\ell)}). \end{aligned}$$

Let us have a closer look at  $\mathbf{P}(\xi_{M+i} = r)$ , or rather, at  $\mathbf{P}(\xi_{M+i} \geq r)$ . By definition,  $\xi_{M+i}$  is the sum of  $(m-1)$  independent copies of  $Y_{M+i}$ , and  $Y_{M+i+1}$  is distributed as  $(Y_{M+i} +$



$\xi_{M+i} - 1)^+$ , with  $Y_{M+i}$  and  $\xi_{M+i}$  being independent. By Proposition 4.1,  $\mathbf{P}(Y_{M+i} \geq 1) \leq c_{28} \frac{(\log(M+i))^2}{(M+i)^2}$ , so

$$\mathbf{P}(\xi_{M+i} \geq 1) \leq (m-1) \mathbf{P}(Y_{M+i} \geq 1) \leq (m-1) c_{28} \frac{(\log(M+i))^2}{(M+i)^2}.$$

For  $r \geq 2$ , we use  $\mathbf{P}(\xi_{M+i} \geq r) \leq \frac{\mathbf{P}(Y_{M+i+1} \geq r-1)}{\mathbf{P}(Y_{M+i}=0)}$ ; since  $\mathbf{P}(Y_{M+i}=0) \geq \frac{1}{2}$  (for  $M \geq M_0$  with a sufficiently large  $M_0$ ), whereas  $\mathbf{P}(Y_{M+i+1} \geq r-1) \leq \frac{c_{28}}{m^{r-2}} \frac{(\log(M+i+1))^2}{(M+i+1)^2}$  (by Proposition 4.1 again), we get  $\mathbf{P}(\xi_{M+i} \geq r) \leq \frac{2c_{28}}{m^{r-2}} \frac{(\log(M+i+1))^2}{(M+i+1)^2}$ . Since  $M = M(n) := \lfloor \frac{n}{1+c_{49}} \rfloor$ , we obtain, for some constant  $c_{50} > 0$  and all integer  $r \geq 1$ ,

$$\mathbf{P}(\xi_{M+i} \geq r) \leq \frac{c_{50}}{m^r} \frac{(\log n)^2}{n^2}.$$

Hence, for  $0 \leq j \leq n - M + \ell - 1$ ,

$$\begin{aligned} \mathbf{E}(S_{M,n}^{(j,\ell)}) &\leq \sum_{i=0}^{j \wedge (n-M-1)} \sum_{r=1}^{n-M+\ell-j} \frac{c_{50}}{m^r} \frac{(\log n)^2}{n^2} m^{i+1} \mathbf{E}(S_{M+i+1,n}^{(j-i+r-1,\ell)}) \\ &= c_{50} \frac{(\log n)^2}{n^2} m^j \sum_{i=0}^{j \wedge (n-M-1)} \sum_{q=j-i}^{n-M+\ell-i-1} m^{-q} \mathbf{E}(S_{M+i+1,n}^{(q,\ell)}), \end{aligned}$$

with  $q := j - i + r - 1$ . Using  $\sum_{q=j-i}^{n-M+\ell-i-1} (\dots) \leq \sum_{q=0}^{n-M+\ell} (\dots)$ , we arrive at: for  $0 \leq j \leq n - M + \ell - 1$ ,

$$\mathbf{E}(S_{M,n}^{(j,\ell)}) \leq c_{50} \frac{(\log n)^2}{n^2} m^j \sum_{i=0}^{j \wedge (n-M-1)} \sum_{q=0}^{n-M+\ell} m^{-q} \mathbf{E}(S_{M+i+1,n}^{(q,\ell)}).$$

Recall from (5.3) that with our choice of  $M$  as in (5.10),

$$\mu_{M+i+1}(q) \geq c_{51} m^{-q},$$

for some constant  $c_{51} > 0$ , all sufficiently large  $n$ , all integers  $0 \leq i \leq n - M - 1$  and  $0 \leq q \leq (1 + \lambda)n$ ; in other words,  $m^{-q} \leq \frac{1}{c_{51}} \mu_{M+i+1}(q)$ . This implies that

$$\begin{aligned} \mathbf{E}(S_{M,n}^{(j,\ell)}) &\leq \frac{c_{50}}{c_{51}} \frac{(\log n)^2}{n^2} m^j \sum_{i=0}^{j \wedge (n-M-1)} \sum_{q=0}^{n-M+\ell} \mu_{M+i+1}(q) \mathbf{E}(S_{M+i+1,n}^{(q,\ell)}) \\ &\leq \frac{c_{50}}{c_{51}} \frac{(\log n)^2}{n^2} m^j \sum_{i=0}^j \sum_{q=0}^{\infty} \mu_{M+i+1}(q) \mathbf{E}(S_{M+i+1,n}^{(q,\ell)}). \end{aligned}$$

By Theorem 2.2 (applied to the system  $(Y_{k+M+i+1}, k \geq 0)$  which is critical), we have  $\sum_{q=0}^{\infty} \mu_{M+i+1}(q) \mathbf{E}(S_{M+i+1,n}^{(q,\ell)}) = \mu_n(\ell) \leq \frac{1}{m^\ell}$ . Hence, for all sufficiently large  $n$ , all  $0 \leq \ell \leq \lambda n$  and all  $0 \leq j \leq n - M + \ell - 1$ , with  $c_{52} := \frac{c_{50}}{c_{51}}$ ,

$$\mathbf{E}(S_{M,n}^{(j,\ell)}) \leq c_{52} \frac{(\log n)^2}{n^2} \frac{(j+1)m^j}{m^\ell}.$$

Since  $\mu_0(k) := m^{-k} \mathbf{E}([(m-1)Y_0 - 1]m^{Y_0} \mathbf{1}_{\{Y_0 \geq k+1\}}) \leq \frac{m-1}{m^k} \sum_{i=k+1}^{\infty} im^i \mathbf{P}(Y_0 = i)$ , we get, for all sufficiently large  $n$ , all  $k \geq 0$  and all  $0 \leq \ell \leq \lambda n$  (with  $c_{53} := (m-1)c_{52}$ )

$$\begin{aligned} \mu_0(k) & \sum_{j=0}^{n-M+\ell-1} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}) \\ & \leq \frac{c_{53}}{m^{k+\ell}} \frac{(\log n)^2}{n^2} \sum_{i=k+1}^{\infty} im^i \mathbf{P}(Y_0 = i) \sum_{j=0}^{n-M+\ell-1} (j+1)m^j \mathbf{E}(S_M^{(k,j)}) \\ & \leq \frac{c_{53}}{m^{k+\ell}} \frac{(\log n)^2}{n^2} \sum_{i=k+1}^{\infty} im^i \mathbf{P}(Y_0 = i) \sum_{j=0}^{\infty} (j+1)m^j \mathbf{E}(S_M^{(k,j)}). \end{aligned}$$

By Lemma 5.3,  $\mathbf{P}(Y_0 = i) \sum_{j=0}^{\infty} (j+1)m^j \mathbf{E}(S_M^{(k,j)}) \leq m^{k-i} \mathbf{E}[(Y_M + 1)m^{Y_M} N_M^{(i)}]$ . Thus

$$\mu_0(k) \sum_{j=0}^{n-M+\ell-1} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}) \leq \frac{c_{53}}{m^\ell} \frac{(\log n)^2}{n^2} \sum_{i=k+1}^{\infty} i \mathbf{E}[(Y_M + 1)m^{Y_M} N_M^{(i)}].$$

Recall from (3.8) that  $\mathbf{E}[(Y_M + 1)m^{Y_M} N_M^{(i)}] = (i+1)m^i \mathbf{P}(Y_0 = i) \prod_{j=0}^{M-1} G_j(m)^{m-1}$ , where  $G_n(s) := \mathbf{E}(s^{Y_n})$  as before. We have  $\prod_{j=0}^{M-1} G_j(m)^{m-1} \leq c_8 M^2 \leq c_8 n^2$  (see Lemma 3.1 (v)).

Therefore, with  $c_{54} := c_{53}c_8$ ,

$$\mu_0(k) \sum_{j=0}^{n-M+\ell-1} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}) \leq \frac{c_{54}}{m^\ell} (\log n)^2 \sum_{i=k+1}^{\infty} i(i+1)m^i \mathbf{P}(Y_0 = i).$$

Summing over all integers  $k \geq k_n(\theta)$  leads to:

$$\begin{aligned} & \sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \sum_{j=0}^{n-M+\ell-1} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}) \\ & \leq \frac{c_{54}}{m^\ell} (\log n)^2 \sum_{k=k_n(\theta)}^{\infty} \sum_{i=k+1}^{\infty} i(i+1)m^i \mathbf{P}(Y_0 = i) \\ & = \frac{c_{54}}{m^\ell} (\log n)^2 \sum_{i=k_n(\theta)+1}^{\infty} i(i+1)(i - k_n(\theta))m^i \mathbf{P}(Y_0 = i), \end{aligned}$$

the last equality being a consequence of the Fubini–Tonelli theorem. Since  $i(i+1)(i-k_n(\theta)) \leq i^2(i+1) \leq 2i^3$ , and  $\sum_{i=k_n(\theta)+1}^{\infty} 2i^3 m^i \mathbf{P}(Y_0 = i) = 2\mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 \geq k_n(\theta)+1\}})$ , which is bounded by  $\frac{2\theta}{(\log n)^2}$  by definition of  $k_n(\theta)$ , we see that, for all sufficiently large  $n$  and all  $0 \leq \ell \leq \lambda n$ ,

$$\sum_{k=k_n(\theta)}^{\infty} \mu_0(k) \sum_{j=0}^{n-M+\ell-1} \mathbf{E}(S_M^{(k,j)}) \mathbf{E}(S_{M,n}^{(j,\ell)}) \leq \frac{2c_{54} \theta}{m^\ell},$$

which is bounded by  $\frac{c_{47}}{4m^\ell}$  if  $\theta$  is chosen to satisfy  $\theta \leq \frac{c_{47}}{8c_{54}}$ . This yields (5.8), and completes the proof of Lemma 5.2.  $\square$

## 5.2 Step 2. A lower bound for $\max_i \mathbf{E}(N_n^{(i)} \mathbf{1}_{\{Y_n \geq \ell+1\}})$

Let  $(Y_n, n \geq 0)$  be a Derrida–Retaux system satisfying  $\mathbf{E}(m^{Y_0}) = (m-1)\mathbf{E}(Y_0 m^{Y_0}) < \infty$ . For any  $i \geq 0$ , let  $N_n^{(i)}$  denote as in (2.3) (associated with  $(Y_n)$ ) the number of open paths in  $\mathbb{T}_n$  to generation  $n$  with initial value  $i$ . The aim of this step is to show that Lemma 5.3 will give an interesting lower bound for  $\mathbf{E}(N_n^{(i)})$  for a certain  $i = i(n)$ .

**Proposition 5.4.** *Assume  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ . Let  $\lambda > 0$ . There exist constants  $\rho > 0$  and  $c_{55} > 0$  such that for all sufficiently large integer  $n$ , there exists a non-random integer  $i^* = i^*(n) \in [0, c_{55} \log \log n]$  satisfying*

$$(5.11) \quad \mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell+1\}}) \geq \frac{1}{(\log n)^\rho} \frac{1}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z}.$$

*Proof.* By assumption, there exists  $s_0 > m$  such that  $c_{56} := \mathbf{E}(Y_0^3 s_0^{Y_0}) < \infty$ .

Let  $\theta > 0$  be the constant in Lemma 5.2. For all integer  $k \geq 0$ ,

$$\mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 \geq k+1\}}) \leq c_{56} \left(\frac{m}{s_0}\right)^{k+1}.$$

We take  $k := \lfloor c_{55} \log \log n \rfloor =: a_n$  (with  $c_{55} := \frac{3}{\log(s_0/m)}$ ), so that  $\left(\frac{m}{s_0}\right)^{k+1} \leq \left(\frac{m}{s_0}\right)^{c_{55} \log \log n} = \frac{1}{(\log n)^3} \leq \frac{\theta}{2c_{56} (\log n)^2}$  (for sufficiently large  $n$ ). Hence for all sufficiently large  $n$ ,

$$\mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 \geq a_n+1\}}) \leq \frac{\theta}{2(\log n)^2}.$$

On the other hand,  $\mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{Y_0 \geq k_n(\theta)\}}) > \frac{\theta}{(\log n)^2}$  by definition. Thus for all sufficiently large  $n$ , we have  $k_n(\theta) \leq a_n$ , and

$$\mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{k_n(\theta) \leq Y_0 \leq a_n\}}) \geq \frac{\theta}{(\log n)^2} - \frac{\theta}{2(\log n)^2} = \frac{\theta}{2(\log n)^2}.$$

Since

$$\mathbf{E}(Y_0^3 m^{Y_0} \mathbf{1}_{\{k_n(\theta) \leq Y_0 \leq a_n\}}) = \sum_{i=k_n(\theta)}^{a_n} i^3 m^i \mathbf{P}(Y_0 = i) \leq a_n^4 m^{a_n} \max_{i \in [k_n(\theta), a_n] \cap \mathbb{Z}} \mathbf{P}(Y_0 = i),$$

this yields that for all sufficiently large  $n$ , there exists  $i^* = i^*(n) \in [k_n(\theta), a_n] \cap \mathbb{Z}$ , such that

$$\mathbf{P}(Y_0 = i^*) \geq \frac{\theta}{2(\log n)^2} \frac{1}{a_n^4 m^{a_n}} \geq \frac{1}{(\log n)^{c_{57}}},$$

with  $c_{57} := 3 + c_{55} \log m$ .

On the other hand, by Lemma 5.2, for all sufficiently large  $n$  and all  $0 \leq \ell \leq \lambda n$ , there exists  $k^* = k^*(n, \ell) \in [0, k_n(\theta) - 1] \cap \mathbb{Z}$ , such that  $\mu_0(k^*) \mathbf{E}(S_n^{(k^*, \ell)}) \geq \frac{c_{48}}{m^\ell k_n(\theta)}$ ; a fortiori  $\mathbf{E}(S_n^{(k^*, \ell)}) \geq \frac{c_{48}}{m^\ell k_n(\theta)}$ . Moreover, since  $i^* \geq k_n(\theta) > k^*$ , we are entitled to apply (5.5) to  $i = i^*$  and  $k = k^*$ , to see that  $\mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n = i^* - k^* + \ell\}}) \geq \mathbf{P}(Y_0 = i^*) \mathbf{E}(S_n^{(k^*, \ell)})$ . Consequently, for all sufficiently large  $n$ ,

$$\mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n = i^* - k^* + \ell\}}) \geq \mathbf{P}(Y_0 = i^*) \frac{c_{48}}{m^\ell k_n(\theta)} \geq \frac{1}{(\log n)^{c_{57}}} \frac{c_{48}}{m^\ell k_n(\theta)} \geq \frac{1}{(\log n)^{c_{57}}} \frac{c_{48}}{m^\ell a_n}.$$

Since  $i^* - k^* \geq 1$ , we have  $\mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n = i^* - k^* + \ell\}}) \leq \mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell + 1\}})$ ; hence for all sufficiently large  $n$ ,

$$\mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell + 1\}}) \geq \frac{1}{(\log n)^{c_{57}}} \frac{c_{48}}{m^\ell a_n} = \frac{1}{(\log n)^{c_{57}}} \frac{c_{48}}{m^\ell \lfloor c_{55} \log \log n \rfloor},$$

proving the proposition.  $\square$

### 5.3 Step 3. Proof of Proposition 5.1

The proof of Proposition 5.1 relies on two results borrowed from [4], stated below as Facts 5.5 and 5.6. Fact 5.5 is a bridge inequality connecting the number of open vertices in  $(Y_n, n \geq 0)$  and the expected value of an appropriate supercritical system  $(X_n, n \geq 0)$ . Fact 5.6 gives an upper bound for  $\mathbf{E}(X_n)$  when the system is “slightly supercritical”.

**Fact 5.5.** ([4, Theorem 4.2]; **The bridge inequality**)<sup>15</sup> *Let  $(X_n, n \geq 0)$  and  $(Y_n, n \geq 0)$  be Derrida–Retaux systems such that  $X_0 \geq Y_0$ . Let  $\eta > 0$ . If  $\mathbf{E}(X_0 - Y_0 | Y_0) \geq \eta$ , then for*

---

<sup>15</sup>The bridge inequality was stated in [4] for  $N_n^{(0)}$  instead of  $N_n$ , but the proof was valid for  $N_n$ . Furthermore,  $X_0$  and  $Y_0$  were assumed in [4] to satisfy  $\mathbb{P}(X_0 = k) \geq \mathbb{P}(Y_0 = k)$  for all integer  $k \geq 1$  in order to ensure the so-called “XY coupling”; the latter holds automatically in our setting, thanks to the additional assumption  $X_0 \geq Y_0$ .

all  $r \geq 0$  and all integers  $n \geq 0$ ,  $\ell \geq 0$  and  $k \in [0, \frac{rn}{2}]$ ,

$$\mathbf{E}(X_{n+k+\ell}) \geq \frac{m^{k+\ell} \eta}{2} \mathbf{E} \left[ N_n \mathbf{1}_{\{N_n \geq r\}} \mathbf{1}_{\{Y_n \geq \ell\}} \right].$$

**Fact 5.6.** ([4, Proposition 3.1 and Lemma 2.4]) *Assume  $\mathbf{E}(m^{X_0}) < (m-1)\mathbf{E}(X_0 m^{X_0}) < \infty$ . Let  $\delta_0 := (m-1)\mathbf{E}(X_0 m^{X_0}) - \mathbf{E}(m^{X_0}) > 0$ . There exist constants  $c_{58} \in (0, 1]$  and  $c_{59} > 0$ , depending only on  $m$ , such that if  $\delta_0 \in (0, c_{58}]$ , then*

$$\max_{0 \leq i \leq c_{59} \left( \frac{\mathbf{P}(X_0=0)}{\delta_0} \right)^{1/2}} \mathbf{E}(X_i) \leq 3.$$

*Proof of Proposition 5.1.* Let  $(Y_n, n \geq 0)$  be a critical system such that  $\mathbf{E}(t^{Y_0}) < \infty$  for some  $t > m$ . Let  $\lambda > 0$ .

Let  $n_2$  be such that Proposition 5.4 applies to all  $n \geq n_2$  (which is now taken for granted); in particular, Proposition 5.4 yields, for all  $n \geq n_2$ , the existence of  $i^* = i^*(n) \in [0, c_{55} \log \log n] \cap \mathbb{Z}$  such that

$$(5.12) \quad \mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell+1\}}) \geq \frac{1}{(\log n)^e} \frac{1}{m^\ell}, \quad \forall \ell \in [0, \lambda n] \cap \mathbb{Z}.$$

Let  $\eta = \eta(n) := \frac{c_{60}}{n^2}$ , where  $c_{60} := \frac{c_{59}^2 \mathbf{P}(Y_0=0)}{2(\lambda+2)^2 c_{61}}$ , with  $c_{61} := m(m-1)\mathbf{E}(m^{Y_0})$ . We can enlarge the value of  $n_2$ , if necessary, such that  $\eta < \min\{\frac{1}{2}, \frac{c_{58}}{c_{61}}\}$  for all  $n \geq n_2$ . Let  $U_0$  be a random variable independent of  $Y_0$  and such that  $\mathbf{P}(U_0 = 1) = \eta = 1 - \mathbf{P}(U_0 = 0)$ . Let  $X_0 := Y_0 + U_0$ . Then  $X_0 \geq Y_0$  and  $\mathbf{E}(X_0 - Y_0 | Y_0) = \eta$ . So we are entitled to apply Fact 5.5.

We also intend to apply Fact 5.6, so let us note that by definition,

$$\begin{aligned} \delta_0 &:= (m-1)\mathbf{E}(X_0 m^{X_0}) - \mathbf{E}(m^{X_0}) \\ &= (m-1)\mathbf{E}(Y_0 m^{Y_0}) \mathbf{E}(m^{U_0}) + (m-1)\mathbf{E}(m^{Y_0}) \mathbf{E}(U_0 m^{U_0}) - \mathbf{E}(m^{Y_0}) \mathbf{E}(m^{U_0}) \\ &= (m-1)\mathbf{E}(m^{Y_0}) \mathbf{E}(U_0 m^{U_0}) \\ &= c_{61} \eta, \end{aligned}$$

where  $c_{61} := m(m-1)\mathbf{E}(m^{Y_0})$  as before. Moreover,  $\mathbf{P}(X_0 = 0) = (1-\eta)\mathbf{P}(Y_0 = 0) \geq \frac{1}{2}\mathbf{P}(Y_0 = 0)$ . By Fact 5.6,

$$\max_{0 \leq i \leq (\lambda+2)n} \mathbf{E}(X_i) \leq 3.$$

Fix  $a > 0$ . Let  $r = r(n) := \frac{2n^2}{c_{60} \log m} \log\left(\frac{6n^{2+a}}{c_{60}}\right)$  and  $k = k(n) := \lfloor \frac{rn}{2} \rfloor$ . Let  $0 \leq \ell \leq \lambda n$ . Then  $n + k + \ell + 1 \leq (\lambda + 2)n$  (for  $n \geq n_2$ , with the value of  $n_2$  enlarged if need be), so  $\mathbf{E}(X_{n+k+\ell+1}) \leq 3$ . On the other hand, by Fact 5.5,

$$\mathbf{E}(X_{n+k+\ell+1}) \geq \frac{m^{k+\ell+1} \eta}{2} \mathbf{E}\left[N_n \mathbf{1}_{\{N_n \geq r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}\right],$$

which implies that

$$(5.13) \quad \mathbf{E}\left[N_n \mathbf{1}_{\{N_n \geq r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}\right] \leq \frac{2}{m^{k+\ell+1} \eta} \mathbf{E}(X_{n+k+\ell+1}) \leq \frac{6}{m^{k+\ell+1} \eta} \leq \frac{1}{n^a} \frac{1}{m^\ell},$$

recalling the definition of  $k$ , together with that of  $r$  and  $\eta$ . We write

$$\begin{aligned} \mathbf{E}\left[N_n^{(i^*)} \mathbf{1}_{\{N_n^{(i^*)} < r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}\right] &= \mathbf{E}[N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell+1\}}] - \mathbf{E}\left[N_n^{(i^*)} \mathbf{1}_{\{N_n^{(i^*)} \geq r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}\right] \\ &\geq \mathbf{E}[N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell+1\}}] - \mathbf{E}\left[N_n \mathbf{1}_{\{N_n \geq r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}\right]. \end{aligned}$$

By (5.12),  $\mathbf{E}(N_n^{(i^*)} \mathbf{1}_{\{Y_n \geq \ell+1\}}) \geq \frac{1}{(\log n)^e} \frac{1}{m^\ell}$ , whereas by (5.13),  $\mathbf{E}[N_n \mathbf{1}_{\{N_n \geq r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}] \leq \frac{1}{n^a} \frac{1}{m^\ell}$ . Therefore, for  $n \geq n_2$  and  $0 \leq \ell \leq \lambda n$ ,

$$\mathbf{E}\left[N_n^{(i^*)} \mathbf{1}_{\{N_n^{(i^*)} < r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}\right] \geq \frac{1}{(\log n)^e} \frac{1}{m^\ell} - \frac{1}{n^a} \frac{1}{m^\ell} \geq \frac{1}{2(\log n)^e} \frac{1}{m^\ell}.$$

[We need to enlarge the value of  $n_2$ , if necessary, to ensure that  $\frac{1}{n^a} \leq \frac{1}{2} \frac{1}{(\log n)^e}$  for all  $n \geq n_2$ .] On the other hand,  $\mathbf{E}[N_n^{(i^*)} \mathbf{1}_{\{N_n^{(i^*)} < r\}} \mathbf{1}_{\{Y_n \geq \ell+1\}}] \leq r \mathbf{P}(Y_n \geq \ell + 1)$ . Consequently, for  $n \geq n_2$ ,

$$\mathbf{P}(Y_n \geq \ell + 1) \geq \frac{1}{r} \frac{1}{2(\log n)^e} \frac{1}{m^\ell} = \frac{c_{60} \log m}{2n^2 \log\left(\frac{6n^{2+a}}{c_{60}}\right)} \frac{1}{2(\log n)^e} \frac{1}{m^\ell},$$

from which (5.1) follows. Proposition 5.1 is proved.  $\square$

## Acknowledgements

We are indebted to Victor Dagard, Bernard Derrida and Mikhail Lifshits for regular and stimulating discussions on the Derrida–Retaux model throughout the last few years. We also wish to thank two anonymous referees for their careful reading of the original manuscript and for their constructive comments.

## References

- [1] Aldous, D.J. and Bandyopadhyay, A. (2005). A survey of max-type recursive distributional equations. *Ann. Appl. Probab.* **15**, 1047–1110.
- [2] Alexander, K.S. (2008). The effect of disorder on polymer depinning transitions. *Commun. Math. Phys.*, **279**, 117–146.
- [3] Berger, Q., Giacomin, G. and Lacoïn, H. (2019). Disorder and critical phenomena: the  $\alpha = 0$  copolymer model. *Probab. Theory Related Fields* **174**, 787–819.
- [4] Chen, X., Dagard, V., Derrida, B., Hu, Y., Lifshits, M. and Shi, Z. (2021). The Derrida–Retaux conjecture on recursive models. *Ann. Probab.* **49**, 637–670.
- [5] Chen, X., Dagard, V., Derrida, B. and Shi, Z. (2020) The critical behaviors and the scaling functions of a coalescence equation. *J. Phys. A* **53**, 195202, 25 pp.
- [6] Chen, X., Derrida, B., Hu, Y., Lifshits, M. and Shi, Z. (2019). A max-type recursive model: some properties and open questions. In: *Sojourns in Probability Theory and Statistical Physics-III* (pp. 166–186). Springer, Singapore.
- [7] Chen, X. and Shi, Z. (2021). The stable Derrida–Retaux system at criticality. In: *In and out of equilibrium, 3, Celebrating Vladas Sidoravicius*, 239–264, *Progr. Probab.* **77**, Birkhäuser/Springer, Cham.
- [8] Collet, P., Eckmann, J.P., Glaser, V. and Martin, A. (1984). A spin glass with random couplings. *J. Statist. Phys.* **36**, 89–106.
- [9] Collet, P., Eckmann, J.P., Glaser, V. and Martin, A. (1984). Study of the iterations of a mapping associated to a spin-glass model. *Commun. Math. Phys.* **94**, 353–370.
- [10] Curien, N. and Hénard, O. (2019+). The phase transition for parking on Galton–Watson trees. arXiv: 1912.06012
- [11] Derrida, B., Hakim, V. and Vannimenus, J. (1992). Effect of disorder on two-dimensional wetting. *J. Statist. Phys.* **66**, 1189–1213.
- [12] Derrida, B. and Retaux, M. (2014). The depinning transition in presence of disorder: a toy model. *J. Statist. Phys.* **156**, 268–290.
- [13] Derrida, B. and Shi, Z. (2020). Results and conjectures on a toy model of depinning. *Moscow Math. J.* **20**, 695–709.
- [14] Giacomin, G. (2007). *Random Polymer Models*. Imperial College Press.
- [15] Giacomin, G. (2011). *Disorder and critical phenomena through basic probability models. École d’été Saint-Flour XL (2010)*, *Lecture Notes in Mathematics* **2025**, Springer, Heidelberg.
- [16] Giacomin, G. and Toninelli, F.L. (2006). Smoothing effect of quenched disorder on polymer depinning transitions. *Commun. Math. Phys.* **266**, 1–16.

- [17] Giacomin, G., Toninelli, F. and Lacoïn, H. (2010). Marginal relevance of disorder for pinning models. *Commun. Pure Appl. Math.*, **63**, 233–265.
- [18] Goldschmidt, C. and Przykucki, M. (2019). Parking on a random tree. *Combin. Probab. Comput.* **28**, 23–45.
- [19] Hu, Y., Mallein, B. and Pain, M. (2020). An exactly solvable continuous-time Derrida–Retaux model. *Commun. Math. Phys.* **375**, 605–651.
- [20] Hu, Y. and Shi, Z. (2018). The free energy in the Derrida–Retaux recursive model. *J. Statist. Phys.* **172**, 718–741.
- [21] Jordan, J. (2002). Almost sure convergence for iterated functions of independent random variables. *Ann. Appl. Probab.* **12**, 985–1000.
- [22] Li, D. and Rogers, T.D. (1999). Asymptotic behavior for iterated functions of random variables. *Ann. Appl. Probab.* **9**, 1175–1201.
- [23] Monthus, C. (2017). Strong disorder renewal approach to DNA denaturation and wetting: typical and large deviation properties of the free energy. *J. Statist. Mech. Theory Exper.* 2017(1), 013301.
- [24] Tang, L.H. and Chaté, H. (2001). Rare-event induced binding transition of heteropolymers. *Phys. Rev. Letters* **86**(5), 830.