

TOTALLY REAL BI-QUADRATIC FIELDS WITH LARGE PÓLYA GROUPS

JAITRA CHATTOPADHYAY AND ANUPAM SAIKIA

ABSTRACT. For an algebraic number field K with ring of integers \mathcal{O}_K , an important subgroup of the ideal class group Cl_K is the *Pólya group*, denoted by $Po(K)$, which measures the failure of the \mathcal{O}_K -module $\text{Int}(\mathcal{O}_K)$ of integer-valued polynomials on \mathcal{O}_K from admitting a regular basis. In this paper, we prove that for any integer $n \geq 2$, there are infinitely many totally real bi-quadratic fields K with $|Po(K)| = 2^n$. In fact, we explicitly construct such an infinite family of number fields. This extends an infinite family of bi-quadratic fields with Pólya group $\mathbb{Z}/2\mathbb{Z}$ given by the authors in [4]. This also provides an infinite family of bi-quadratic fields with class numbers divisible by 2^n .

1. INTRODUCTION

Let K be an algebraic number field with ring of integers \mathcal{O}_K and ideal class group Cl_K . Let

$$\text{Int}(\mathcal{O}_K) = \{f(X) \in K[X] : f(\alpha) \in \mathcal{O}_K \text{ for all } \alpha \in \mathcal{O}_K\}.$$

It is a well-known fact (cf. [1], [2]) that $\text{Int}(\mathcal{O}_K)$ is a free \mathcal{O}_K -module. An \mathcal{O}_K -basis $\{f_i\}_{i \geq 0}$ of $\text{Int}(\mathcal{O}_K)$ is said to be a regular basis if $\deg(f_i) = i$ for each $i \geq 0$. The existence of a regular basis leads to the definition of a Pólya field as follows.

Definition 1. (cf. [1], [16]) An algebraic number field K is said to be a Pólya field if the \mathcal{O}_K -module $\text{Int}(\mathcal{O}_K)$ has a regular basis.

It is quite interesting that the existence of a regular basis for K is closely governed by the structure of Cl_K . To describe the interplay between Cl_K and $\text{Int}(\mathcal{O}_K)$, we first introduce the following notation.

$$\Pi_q(K) = \begin{cases} \prod_{\substack{\mathfrak{p} \in \text{Spec}(\mathcal{O}_K) \\ N\mathfrak{p}=q}} \mathfrak{p} & ; \text{ if } q \text{ is a prime power} \\ \mathcal{O}_K & ; \text{ otherwise.} \end{cases}$$

Definition 2. (cf. [1], [12], [13] [16]) The Pólya group $Po(K)$ of K is defined to be the subgroup generated by the ideal classes $[\Pi_q(K)]$ in Cl_K .

2020 *Mathematics Subject Classification*: Primary 13F20; Secondary 11R29, 11R34.

Key words and phrases: Pólya field, Pólya group.

It is known (cf. [1]) that K is a Pólya field if and only if $Po(K)$ trivial. Thus the study of $\text{Int}(\mathcal{O}_K)$ is translated into the study of Cl_K . Moreover, when K/\mathbb{Q} is a finite Galois extension, it immediately follows from Definition 2 that $Po(K)$ is generated precisely by the ideal classes $[\Pi_q(K)]$, where q runs through the norms of ramified primes. Therefore, there is a close connection between the ramified primes in K and the Pólya group $Po(K)$. In particular, for quadratic fields, the following result is known due to the work of Hilbert.

Proposition 1. [7] *Let $K = \mathbb{Q}(\sqrt{d})$ be a quadratic field. Let ε be the fundamental unit of K and let r_K denote the number of ramified primes in K/\mathbb{Q} . Then*

$$|Po(K)| = \begin{cases} 2^{r_K-2} & ; \text{if } d > 0 \text{ and } N(\varepsilon) = 1 \\ 2^{r_K-1} & ; \text{otherwise.} \end{cases}$$

From Proposition 1, it is clear that the order of the Pólya group for a quadratic field is necessarily 1 or a power of 2. In [16], Zantema generalized this to finite Galois extensions K/\mathbb{Q} and recently, Maarefparvar and Rajaei extended the result for relative extensions of number fields in [11]. Leriche addressed the cubic field case in [8]. There has been a lot of studies on the Pólya group of bi-quadratic fields in recent times (cf. [4],[5],[6],[9],[10]). In [4], the authors determined the Pólya groups of bi-quadratic fields that are known to have an ‘‘Euclidean ideal class’’. The interested reader is encouraged to look into [3] and the references listed therein to know about Euclidean ideal class.

In this paper, using Zantema’s result, we explicitly construct an infinite family of totally real bi-quadratic fields with Pólya group having 2-rank n , for any integer $n \geq 1$. Our result extends the previously known family of bi-quadratic fields with Pólya group $\mathbb{Z}/2\mathbb{Z}$ (cf. Theorem 3 in [4]). More precisely, we prove the following theorem.

Theorem 1. *Let $n \geq 1$ be an integer. Then there exist infinitely many totally real bi-quadratic fields K with Pólya group $Po(K) \simeq (\mathbb{Z}/2\mathbb{Z})^n$.*

Since the Pólya group is a subgroup of the ideal class group, we immediately have the following corollary.

Corollary 1. *For an integer $n \geq 1$, there exist infinitely many totally real bi-quadratic fields K with 2-rank of Cl_K being n .*

2. PRELIMINARIES

In [16], Zantema made use of Galois cohomology to study $Po(K)$ and its relation with the ramified primes in K/\mathbb{Q} , when K is a finite Galois extension of \mathbb{Q} . Since the Galois group $G = \text{Gal}(K/\mathbb{Q})$ acts on the multiplicative group of units \mathcal{O}_K^* via the action $(\sigma, \alpha) \mapsto \sigma(\alpha)$, thus endowing \mathcal{O}_K^* with a G -module structure. Then Zantema’s result can be stated as follows.

Proposition 2. [16] *Let K be a finite Galois extension of \mathbb{Q} with Galois group $G = \text{Gal}(K/\mathbb{Q})$. Let p_1, \dots, p_m be the rational primes that ramify in K . Let the ramification index of p_i be e_i for*

each $i \in \{1, \dots, m\}$. Then there is an exact sequence

$$0 \rightarrow H^1(G, \mathcal{O}_K^*) \rightarrow \bigoplus_{i=1}^m \mathbb{Z}/e_i\mathbb{Z} \rightarrow \text{Po}(K) \rightarrow 0$$

of abelian groups.

The next two lemma help us understand the structure of the group $H^1(G, \mathcal{O}_K^*)$ via its subgroup of 2-torsion elements.

Lemma 1. [14] *Let K_1, K_2 and K_3 be the quadratic subfields of a bi-quadratic field K . Let $H[2]$ be the 2-torsion subgroup of $H^1(G, \mathcal{O}_K^*)$. Then the index of $H[2]$ in $H^1(G, \mathcal{O}_K^*)$ is ≤ 2 . The index is 2 if and only if the rational prime 2 is totally ramified in K/\mathbb{Q} and there exist $\alpha_i \in \mathcal{O}_{K_i}$ for each $i = 1, 2, 3$ with*

$$N(\alpha_1) = N(\alpha_2) = N(\alpha_3) = \pm 2.$$

Lemma 2. [16] *Let K_1, K_2 and K_3 be the quadratic subfields of a totally real bi-quadratic field K . For $i = 1, 2, 3$, let Δ_i be the square-free part of the discriminant of K_i . Let $u_i = z_i + t_i\sqrt{\Delta_i}$ be a fundamental unit of \mathcal{O}_{K_i} with $z_i > 0$. Let us define*

$$a_i = \begin{cases} N(u_i + 1) & ; \text{if } N(u_i) = 1, \\ 1 & ; \text{if } N(u_i) = -1. \end{cases}$$

Let $H[2]$ be the 2-torsion subgroup of $H^1(G, \mathcal{O}_K^)$. Then $H[2]$ is isomorphic to the subgroup of $\mathbb{Q}^*/(\mathbb{Q}^*)^2$ generated by the images of $\Delta_1, \Delta_2, \Delta_3, a_1, a_2$ and a_3 in $\mathbb{Q}^*/(\mathbb{Q}^*)^2$.*

The next lemma is very crucial for proving Theorem 1 and we furnish a detailed proof here.

Lemma 3. *Let $t \geq 2$ be an integer. For every unordered pair (i, j) with $1 \leq i \neq j \leq t$, let $\varepsilon_{ij} \in \{\pm 1\}$ be given. Then there exist infinitely many prime numbers p_1, \dots, p_t with $p_i \equiv 1 \pmod{8}$ for each i , and the Legendre symbol $\left(\frac{p_i}{p_j}\right) = \varepsilon_{ij}$ for all $i \neq j$.*

Proof. We prove by induction on t . We begin with $t = 2$. If $\varepsilon_{12} = 1$, we take $p_1 = 17$ and $p_2 \equiv 1 \pmod{8p_1}$. On the other hand, if $\varepsilon_{12} = -1$, we take $p_2 \equiv 3 \pmod{8p_1}$. By Dirichlet's theorem for primes in arithmetic progression, we obtain infinitely many choices for p_2 in both the cases. Therefore, the lemma holds for $t = 2$.

Now, suppose that we have prime numbers p_1, \dots, p_{t-1} such that $p_i \equiv 1 \pmod{8}$ for each $i \in \{1, \dots, t-1\}$ and $\left(\frac{p_i}{p_j}\right) = \varepsilon_{ij}$ for all $1 \leq i \neq j \leq t-1$. We wish to find a prime number p_t with $\left(\frac{p_t}{p_i}\right) = \varepsilon_{it}$ for all $1 \leq i \leq t-1$. For each integer $i \in \{1, \dots, t-1\}$, we choose integers n_i such that $1 \leq n_i \leq p_i - 1$ and $\left(\frac{n_i}{p_i}\right) = \varepsilon_{it}$. We consider the following system of congruences.

$$\begin{aligned}
X &\equiv 1 \pmod{8} \\
X &\equiv n_1 \pmod{p_1} \\
&\vdots \\
X &\equiv n_{t-1} \pmod{p_{t-1}}.
\end{aligned}$$

By the Chinese remainder theorem, there exists a unique integer, say x_0 , modulo $8p_1 \dots p_{t-1}$ satisfying the above system of congruences. Since $\gcd(x_0, 8p_1 \dots p_{t-1}) = 1$, by Dirichlet's theorem for primes in arithmetic progression, there exist infinitely many prime numbers ℓ such that $\ell \equiv x_0 \pmod{8p_1 \dots p_{t-1}}$. Then $\ell \equiv 1 \pmod{8}$ and $\left(\frac{\ell}{p_i}\right) = \left(\frac{x_0}{p_i}\right) = \left(\frac{n_i}{p_i}\right) = \varepsilon_{it}$ for each $i \in \{1, \dots, t-1\}$. Taking $p_t = \ell$, we conclude that there are infinitely many such prime numbers fulfilling the desired requirements. \square

The next lemma, due to Trotter [15], provides a useful criterion for a real quadratic field of a particular type to have a fundamental unit of negative norm.

Lemma 4. [15] *Let p_1, \dots, p_t be prime numbers with $p_i \equiv 1 \pmod{4}$ for each $i \in \{1, \dots, t\}$. Let $\varepsilon_{ij} = \left(\frac{p_i}{p_j}\right)$ for $1 \leq i \neq j \leq t$. Then each of the following two conditions is sufficient to ensure that the fundamental unit of the real quadratic field $\mathbb{Q}(\sqrt{p_1 \dots p_t})$ to have norm -1 .*

- (1) t is odd and $\varepsilon_{ij} = -1$ for all $1 \leq i \neq j \leq t$.
- (2) t is even, $\varepsilon_{12} = -1$, $\varepsilon_{1i} = 1$ for all $i > 2$ and $\varepsilon_{ij} = -1$ for all $2 \leq i \neq j \leq t$.

3. PROOF OF THEOREM 1

Let $t \geq 3$ be an odd integer and let $p_1 \dots p_t$ be prime numbers with $p_i \equiv 1 \pmod{8}$ and $\left(\frac{p_i}{p_j}\right) = -1$ for all $1 \leq i \neq j \leq t$. Lemma 3 allows us to choose such prime numbers. Let $K = \mathbb{Q}(\sqrt{2}, \sqrt{p_1 \dots p_t})$ with three quadratic subfields $K_1 = \mathbb{Q}(\sqrt{2})$, $K_2 = \mathbb{Q}(\sqrt{p_1 \dots p_t})$ and $K_3 = \mathbb{Q}(\sqrt{2p_1 \dots p_t})$. The only ramified primes in K/\mathbb{Q} are 2 and p_i 's. Moreover, since $K = K_1K_2$ and $K_1 \cap K_2 = \mathbb{Q}$, we conclude that the ramification indices of all these primes are 2.

Therefore, by Proposition 2, $H^1(G, \mathcal{O}_K^*)$ embeds into $\bigoplus_{i=1}^{t+1} \mathbb{Z}/2\mathbb{Z}$, where G is the Galois group $\text{Gal}(K/\mathbb{Q})$. Now, we prove that $|H^1(G, \mathcal{O}_K^*)| = 4$, which, by Proposition 2, implies that

$$|Po(K)| = \frac{\left| \bigoplus_{i=1}^{t+1} \mathbb{Z}/2\mathbb{Z} \right|}{|H^1(G, \mathcal{O}_K^*)|} = \frac{2^{t+1}}{4} = 2^{t-1}.$$

We note that the rational prime 2 is not totally ramified in K/\mathbb{Q} . Therefore, by Lemma 1, we get that $H[2] = H^1(G, \mathcal{O}_K^*)$. Following the notations of Lemma 2, we get that $\Delta_1 = 2$, $\Delta_2 = p_1 \dots p_t$ and $\Delta_3 = 2p_1 \dots p_t$. Using the notation $[x]$ to denote the image of a non-zero

element $x \in \mathbb{Q}$ in $\mathbb{Q}^*/(\mathbb{Q}^*)^2$, we see that $[\Delta_1], [\Delta_2], [\Delta_3] \in \langle [2], [p_1 \cdots p_t] \rangle$ in $\mathbb{Q}^*/(\mathbb{Q}^*)^2$. We note that $1 + \sqrt{2}$ is a fundamental unit of K_1 with negative norm. Also, by Lemma 4, the fundamental unit of K_2 has negative norm and consequently, $[a_1]$ and $[a_2]$ are trivial in $\mathbb{Q}^*/(\mathbb{Q}^*)^2$.

Now, let $u = z + w\sqrt{2p_1 \cdots p_t}$ be a fundamental unit of K_3 . If $N(u) = -1$, then $[a_3] \in \langle [2], [p_1 \cdots p_t] \rangle$ and hence $|H^1(G, \mathcal{O}_K^*)| = 4$. Otherwise, we assume that $N(u) = 1$. Then by Lemma 2, we have $a_3 = N(u + 1) = 2(z + 1)$. Since $z^2 - 1 = 2w^2p_1 \cdots p_t$ is even, we conclude that z is odd and therefore $\gcd(z - 1, z + 1) = 2$. Therefore, $w = 2w_1$ for some integer w_1 and hence

$$(1) \quad \frac{z-1}{2} \cdot \frac{z+1}{2} = 2w_1^2 p_1 \cdots p_t.$$

Since $\gcd\left(\frac{z-1}{2}, \frac{z+1}{2}\right) = 1$, from equation (1), we get that $\frac{z-1}{2} = a^2\alpha$ and $\frac{z+1}{2} = b^2\beta$, where a, b are relatively prime integers with $ab = k$ and α, β are relatively prime integers with $\alpha\beta = 2p_1 \cdots p_t$. Thus we have $[2(z+1)] = [4b^2\beta] = [\beta] \in \mathbb{Q}^*/(\mathbb{Q}^*)^2$. If $\beta = 1$ or 2 or $p_1 \cdots p_t$ or $2p_1 \cdots p_t$, then $[\beta] \in \langle [2], [p_1 \cdots p_t] \rangle$. We now prove that other choices cannot occur for β .

Case 1. $\beta = 2B$, where B is an integer dividing $p_1 \cdots p_t$ and consisting of an odd number of prime divisors. Then for a prime divisor ℓ of α , from the equation

$$(2) \quad 1 = \frac{z+1}{2} - \frac{z-1}{2} = b^2\beta - a^2\alpha,$$

we have

$$1 = \left(\frac{1}{\ell}\right) = \left(\frac{2b^2B - a^2\alpha}{\ell}\right) = \left(\frac{2}{\ell}\right) \left(\frac{B}{\ell}\right) = \left(\frac{B}{\ell}\right) = -1,$$

which is a contradiction.

Case 2. $\beta = 2B$, where B is an integer dividing $p_1 \cdots p_t$ and consisting of an even number of prime divisors. Then α consists of an odd number of prime divisors, since t is odd. Therefore, for an odd prime divisor ℓ of β , we obtain

$$1 = \left(\frac{1}{\ell}\right) = \left(\frac{2b^2B - a^2\alpha}{\ell}\right) = \left(\frac{-\alpha}{\ell}\right) = \left(\frac{\alpha}{\ell}\right) = -1,$$

which is a contradiction.

Case 3. $\alpha = 2A$, where A is an integer dividing $p_1 \cdots p_t$ and consisting of an even number of prime divisors. Then β consists of an odd number of prime factors. Therefore, for an odd prime divisor ℓ of α , we get

$$1 = \left(\frac{1}{\ell}\right) = \left(\frac{b^2\beta - 2a^2A}{\ell}\right) = \left(\frac{\beta}{\ell}\right) = -1,$$

which is a contradiction.

Case 4. $\alpha = 2A$, where A is an integer dividing $p_1 \cdots p_t$ and consisting of an odd number of prime divisors. Then for a prime divisor ℓ of β , we obtain

$$1 = \left(\frac{1}{\ell}\right) = \left(\frac{b^2\beta - 2a^2A}{\ell}\right) = \left(\frac{-1}{\ell}\right) \left(\frac{2}{\ell}\right) \left(\frac{A}{\ell}\right) = \left(\frac{A}{\ell}\right) = -1,$$

which is a contradiction.

Consequently, we have $[a_3] = [2(z+1)] = [\beta] \in \langle [2], [p_1 \cdots p_t] \rangle$ and hence by Lemma 1 and Lemma 2, we have $|H^1(G, \mathcal{O}_K^*)| = |H[2]| = |\langle [2], [p_1 \cdots p_t] \rangle| = 4$. Hence $|Po(K)| = 2^{t-1}$.

Now, we assume that t is even. By Lemma 3, we can choose prime numbers p_1, \dots, p_t such that $p_i \equiv 1 \pmod{8}$ for each i and they satisfy condition (2) of Lemma 4. Following the same notation as above, we see that $[\Delta_1], [\Delta_2], [\Delta_3], [a_1], [a_2] \in \langle [2], [p_1 \cdots p_t] \rangle$. If $N(u) = -1$, then we get that $|H^1(G, \mathcal{O}_K^*)| = |H[2]| = |\langle [2], [p_1 \cdots p_t] \rangle| = 4$. We now prove that for $N(u) = 1$ also, we have $[a_3] \in \langle [2], [p_1 \cdots p_t] \rangle$. For that, we observe that if $\beta = 1$ or 2 or $p_1 \cdots p_t$ or $2p_1 \cdots p_t$, then we are through. We prove that no other possibilities can occur.

Case 1. $p_1 \mid \beta$ and $p_2 \mid \alpha$. Then from equation (2), we get

$$1 = \left(\frac{1}{p_1}\right) = \left(\frac{-a^2\alpha}{p_1}\right) = \left(\frac{\alpha}{p_1}\right) = \left(\frac{p_2}{p_1}\right) = -1,$$

by the conditions on p_i 's. Therefore, this cannot occur.

Case 2. $p_1 \mid \alpha$ and $p_2 \mid \beta$. Then again from equation (2), we get

$$1 = \left(\frac{1}{p_1}\right) = \left(\frac{b^2\beta}{p_1}\right) = \left(\frac{\beta}{p_1}\right) = \left(\frac{p_2}{p_1}\right) = -1,$$

which is a contradiction

Case 3. $\beta = p_1 p_2 B$ or $2p_1 p_2 B$, where $B \mid p_1 \cdots p_t$ and consists of an even number of odd prime divisors. Then α also comprises with an even number of odd prime factors and therefore from equation (2), we obtain for any odd prime divisor ℓ of α that $1 = \left(\frac{1}{\ell}\right) = \left(\frac{2}{\ell}\right) \left(\frac{p_1}{\ell}\right) \left(\frac{p_2}{\ell}\right) \cdot 1 = -1$, which is a contradiction. Similarly, if B consists of an odd number of odd prime divisors, then so does α . Again, from equation (2), we get $1 = \left(\frac{1}{p_2}\right) = \left(\frac{-a^2\alpha}{p_2}\right) = \left(\frac{\alpha}{p_2}\right) = -1$, a contradiction.

Case 4. $p_1 p_2 \nmid \beta$. Then $p_1 p_2 \mid \alpha$ and considering the equation $-1 = a^2\alpha - b^2\beta$, we are essentially back to Case 3.

Hence we conclude that for t even as well, we have infinitely many bi-quadratic fields K with $|Po(K)| = 2^{t-1}$. Putting $n = t - 1$ completes the proof of theorem 1. \square

Acknowledgements. It is a pleasure for the first author to thank Indian Institute of Technology Guwahati for providing the financial support and necessary facilities to carry out this research work. The second author would like to thank MATRICS, SERB for their research grant.

REFERENCES

- [1] P.J. Cahen and J.L. Chabert, *Integer-valued polynomials. Mathematical Surveys and Monographs 48*, American Mathematical Society, (1997).
- [2] P. J. Cahen and J. L. Chabert, *What you should know about integer-valued polynomials*, Amer. Math. Monthly, **123** (2016), 311-337.

- [3] J. Chattopadhyay and S. Muthukrishnan, *Bi-quadratic fields having a non-principal Euclidean ideal class*, *J. Number Theory*, **204** (2019), 99-112.
- [4] J. Chattopadhyay and A. Saikia, *Non-Pólya bi-quadratic fields having an Euclidean ideal class*, *Communicated* (arXiv:2105.14436v1).
- [5] B. Heidaryan and A. Rajaei, *Biquadratic Pólya fields with only one quadratic Pólya subfield*, *J. Number Theory*, **143** (2014), 279-285.
- [6] B. Heidaryan and A. Rajaei, *Some non-Pólya biquadratic fields with low ramification*, *Rev. Mat. Iberoam.*, **33** (2017), 1037-1044.
- [7] D. Hilbert, *The Theory of Algebraic Number Fields (English summary) Translated from the German and with a preface by Iain T. Adamson. With an introduction by Franz Lemmermeyer and Norbert Schappacher*, Springer-Verlag, Berlin, (1998).
- [8] A. Leriche, *Cubic, quartic and sextic Pólya fields*, *J. Number Theory*, **133** (2013), 59-71.
- [9] A. Maarefparvar, *Existence of relative integral basis over quadratic fields and Pólya property*, *Acta Math. Hungar.*, **164** (2021), 593-598.
- [10] A. Maarefparvar, *Pólya group in some real biquadratic fields*, *J. Number Theory*, **228** (2021), 1-7.
- [11] A. Maarefparvar and A. Rajaei, *Relative Pólya group and Pólya dihedral extensions of \mathbb{Q}* , *J. Number Theory*, **207** (2020), 367-384.
- [12] A. Ostrowski, *Über ganzwertige Polynome in algebraischen Zahlkörpern*, *J. Reine Angew. Math.*, **149** (1919), 117-124.
- [13] G. Pólya, *Über ganzwertige Polynome in algebraischen Zahlkörpern*, *J. Reine Angew. Math.*, **149** (1919), 97-116.
- [14] C. B. Setzer, *Units over totally real $C_2 \times C_2$ fields*, *J. Number Theory*, **12** (1980), 160-175.
- [15] H. F. Trotter, *On the norms of units in quadratic fields*, *Proc. Amer. Math. Soc.*, **22** (1969), 198-201.
- [16] H. Zantema, *Integer valued polynomials over a number field*, *Manuscripta Math.*, **40** (1982) 155-203.

(Jaitra Chattopadhyay) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, GUWAHATI, GUWAHATI - 781039, ASSAM, INDIA

Email address, Jaitra Chattopadhyay: jaitra@iitg.ac.in, chat.jaitra@gmail.com

(Anupam Saikia) DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, GUWAHATI, GUWAHATI - 781039, ASSAM, INDIA

Email address, Anupam Saikia: a.saikia@iitg.ac.in