

# Comment on “An appropriate approach to pricing european-style options with the Adomian decomposition method”

Francisco M. Fernández\*  
 INIFTA, DQT, Sucursal 4, C. C. 16,  
 1900 La Plata, Argentina

## Abstract

We show that the Adomian decomposition method proposed by Ke et al [ANZIAM J. **59** (2018) 349] is just the Taylor series approach in disguise. The latter approach is simpler, more straightforward and yields a recurrence relation free from integrals.

In a recent paper published in this journal Ke et al [1] (KGZ from now on) applied the so called Adomian decomposition method (ADM) to the Black-Scholes model for the pricing of European options. In order to avoid the problem of differentiability of the solution to the partial differential equation at origin they moved the discontinuity at infinity by means of a suitable change of variables. The purpose of this Comment is the analysis of the implementation of the ADM proposed by KGZ.

Our starting point is the partial differential equation derived by KGZ after the change of variables:

$$\partial_z [zu(y, z)] = 2\partial_y^2 u(y, z) + y\partial_y u(y, z) + 2(k_1 - 1)z\partial_y u(y, z) - 2k_2 z^2 u(y, z), \quad (1)$$

---

\*fernande@quimica.unlp.edu.ar

where  $\partial_z$  and  $\partial_y$  denote partial derivatives with respect to  $z$  and  $y$ , respectively.

Here, we expand the solution  $u(y, z)$  in a Taylor series about  $z = 0$ :

$$u(y, z) = \sum_{j=0}^{\infty} f_j(y) z^j, \quad f_j(y) = \frac{1}{j!} \partial_z^j u(y, z) \Big|_{z=0}. \quad (2)$$

If we insert the expansion (2) into the differential equation (1) we obtain a recurrence relation for the functions  $f_j(y)$ :

$$\begin{aligned} f_n(y) &= \frac{1}{n+1} [2\partial_y^2 f_n(y) + y\partial_y f_n(y) + 2(k_1 - 1)\partial_y f_{n-1}(y) - 2k_2 f_{n-2}(y)], \\ n &= 0, 1, \dots, \quad f_{-1} = f_{-2} = 0. \end{aligned} \quad (3)$$

It is not difficult to verify that this recurrence relation enables one to obtain all the functions  $u_j(y, z) = f_j(y) z^j$  derived by KGZ by means of the ADM through a lengthier and more cumbersome procedure. We can show the connection between the Taylor and ADM solutions in an even clearer way.

If we multiply equation (3) by  $z^n$  and take into account that  $z^n/(n+1) = z^{-1} \int_0^z z^n dz$  we obtain

$$\begin{aligned} f_n(y) z^n &= \frac{1}{z} \int_0^z [2\partial_y^2 f_n(y) z^n + y\partial_y f_n(y) z^n + 2(k_1 - 1) z \partial_y f_{n-1}(y) z^{n-1} \\ &\quad - 2k_2 z^2 f_{n-2}(y) z^{n-2}] dz, \end{aligned} \quad (4)$$

that is exactly the equation derived by KGZ by means of the ADM if  $u_j(y, z) = f_j(y) z^j$ .

It is clear that the implementation of the ADM proposed by KGZ is simply the Taylor-series approach in disguise. The latter is by far simpler and more straightforward and yields the expression (3) that is free from the integral over  $z$ . It is worth adding that the solutions provided by this particular form of the ADM are valid for sufficiently small values of the expansion variable  $z$ . This is a well known limitation of any approach based on a power-series expansion.

## References

- [1] Z. Ke, J. Goard, and S-P. Zhu, "An appropriate approach to pricing european-style options with the Adomian decomposition method", ANZIAM

J. **59**, 349-369 (2018).