

ASYMPTOTIC MEAN-VALUE FORMULAS FOR SOLUTIONS OF GENERAL SECOND-ORDER ELLIPTIC EQUATIONS

PABLO BLANC, FERNANDO CHARRO, JUAN J. MANFREDI, AND JULIO D. ROSSI

ABSTRACT. We obtain asymptotic mean-value formulas for solutions of second-order elliptic equations. Our approach is very flexible and allows us to consider several families of operators obtained as an infimum, a supremum, or a combination of both infimum and supremum, of linear operators. The families of equations that we consider include well-known operators such as Pucci, Issacs, and k -Hessian operators.

1. INTRODUCTION

It is well-known that mean-value formulas characterize harmonic functions; in fact, a weaker statement, known as the asymptotic mean-value property, is enough to characterize harmonicity (see [7, 15, 23]). Furthermore, we have

$$\Delta u(x) = \lim_{\varepsilon \rightarrow 0} \frac{2(n+2)}{\varepsilon^2} \left(\int_{B_\varepsilon(x)} u(y) dy - u(x) \right),$$

from which we conclude that, when f is continuous, a function u satisfies the classical Poisson equation $\Delta u(x) = f(x)$ in Ω if and only if

$$u(x) = \int_{B_\varepsilon(x)} u(y) dy - \frac{\varepsilon^2}{2(n+2)} f(x) + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0 \quad (1.1)$$

for each $x \in \Omega$. We use the standard notation $o(\varepsilon^2)$ to denote a quantity such that $o(\varepsilon^2)/\varepsilon^2 \rightarrow 0$ as $\varepsilon \rightarrow 0$. Analogous results hold for sub- and supersolutions replacing equalities by appropriate inequalities.

The classic mean-value property can be seen as a nonlocal formulation of the local Laplace and Poisson equations. It has deep connections with other fundamental properties such as the maximum principle, which allows proving existence results by Perron's method or symmetry of solutions by the moving-planes method. In this sense, the mean-value property and the maximum principle are nothing more than a quantitative expression of the monotonicity inherent to the equation.

In our recent paper [4] we established asymptotic mean-value properties in the viscosity sense for solutions to the Monge-Ampère equation,

$$\det D^2 u(x) = f(x), \quad \text{in } \Omega, \quad (1.2)$$

which is elliptic only in the class of convex functions and, consequently, requires $f \geq 0$. Our results in [4] are based on the formula

$$n(\det D^2 u(x))^{1/n} = \inf_{\det A=1} \text{trace}(A^t D^2 u(x) A),$$

Key words and phrases. Mean-value formulas, viscosity solutions, k -Hessian equations, Issacs equations.

2020 *Mathematics Subject Classification:* 35J60, 35D40, 35B05.

P.B. partially supported by the Academy of Finland project no. 298641.

F.C. partially supported by grant MTM2017-84214-C2-1-P and PID2019-110712GB-I100 funded by MCIN/AEI/10.13039/501100011033 and by "ERDF A way of making Europe".

J.D.R. partially supported by CONICET grant PIP GI No 11220150100036CO (Argentina), PICT-2018-03183 (Argentina) and UBACyT grant 20020160100155BA (Argentina).

which holds whenever $D^2u(x) \geq 0$. Mean-value properties in the viscosity sense require more geometrical arguments adapted to the operator. In particular, our methods do not require explicit representation formulas, making them flexible enough to be applied to various nonlinear problems.

In this article, we establish asymptotic mean-value formulas for a wide array of fully nonlinear equations that includes degenerate operators such as the k -Hessians, [11], truncated Laplacians, [3], and prescribed eigenvalues of the Hessian, [5, 22]. The operators we discuss can be written as an infimum of linear operators with coefficients chosen from a given set, see below for examples. Of course, there are corresponding statements for equations involving a supremum or combinations of infimum and supremum.

Mean-value formulas hold under more lenient regularity conditions than the corresponding PDEs and can provide a simple and unified approach to nonlinear equations in non-Euclidean contexts such as Carnot groups. As a proof of concept, we prove mean-value formulas for Monge-Ampère operators in the Heisenberg group in Section 6 below. Other directions for applications of the asymptotic mean-value formulas below concern game-theoretic interpretations of the corresponding PDEs, and their numerical analysis. In this regard, there are convergent difference schemes for the normalized infinity Laplacian and p -Laplacian using mean-value formulas, see [21]. Hence, the mean-value formulas that we develop here could provide new numerical methods for the corresponding nonlinear equations.

Let us now describe our results. First, we consider differential operators of the form $F(D^2u)$, with $F : S^n(\mathbb{R}) \rightarrow \mathbb{R} \cup \{-\infty\}$ given by

$$F(M) = \inf_{A \in \mathcal{A}} \text{trace}(A^t M A). \quad (1.3)$$

Here, \mathcal{A} is a subset of $S_+^n(\mathbb{R})$, the set of symmetric positive semi-definite matrices. We observe in Lemma 4.1 below that the set \mathcal{A} is bounded if and only if the operator $F(M)$ is well-defined (finite) for all $M \in S^n(\mathbb{R})$. Therefore, \mathcal{A} determines the set of admissible solutions, functions for which $F(D^2u)$ is finite. We say that

$$u \in C^2(\Omega) \text{ is } \mathcal{A}\text{-admissible in } \Omega \text{ if } D^2u(x) \in \Gamma_{\mathcal{A}} \text{ for every } x \in \Omega, \quad (1.4)$$

where we define the cone

$$\Gamma_{\mathcal{A}} = \left\{ M \in S^n(\mathbb{R}) : F(M) > -\infty \right\}$$

(see [11] for a related notion of admissibility). This condition plays an analogous role to the convexity for the Monge-Ampère equation; in fact,

$$\inf_{A \in \mathcal{A}} \text{trace}(A^t D^2u(x) A) = \begin{cases} n (\det D^2u(x))^{1/n} & \text{if } D^2u(x) \geq 0 \\ -\infty & \text{otherwise,} \end{cases} \quad (1.5)$$

for the unbounded set $\mathcal{A} = \{A \in S_+^n(\mathbb{R}) : \det(A) = 1\}$, see [4].

Observe that F is an infimum of linear functions, which are continuous; therefore, it is upper semi-continuous. Even more, F is concave and continuous in $\Gamma_{\mathcal{A}}^\circ$. We assume that

$$F \text{ is lower semi-continuous in } \Gamma_{\mathcal{A}} \setminus \Gamma_{\mathcal{A}}^\circ. \quad (1.6)$$

so that we have that

$$F \text{ is continuous in } \Gamma_{\mathcal{A}}. \quad (1.7)$$

Condition (1.6) is not satisfied in Example 4.6 below and the mean-value formula fails.

To deal with operators of the form (1.3), where the class of matrices \mathcal{A} is unbounded, we consider matrices A such that $A \in \mathcal{A}$ and $A \leq \phi(\varepsilon)I$ with $\phi(\varepsilon)$ a positive function such that

$$\lim_{\varepsilon \rightarrow 0} \phi(\varepsilon) = +\infty \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \varepsilon \phi(\varepsilon) = 0 \quad (1.8)$$

(an example of such a function is $\phi(\varepsilon) = \varepsilon^{-\alpha}$ for $0 < \alpha < 1$). Notice that the condition $A \leq \phi(\varepsilon)I$ becomes less restrictive as $\varepsilon \rightarrow 0$ but is still enough to make the mean-value formula local, see Section 4. We obtain the following result.

Theorem 1.1. *Let $\phi(\varepsilon)$ be a positive function that satisfies (1.8), and $F : S^n(\mathbb{R}) \rightarrow \mathbb{R} \cup \{-\infty\}$ an operator of the form (1.3) that satisfies (1.6). Let the function $u \in C^2(\Omega)$ be \mathcal{A} -admissible. Then, for every $x \in \Omega$ we have*

$$F(D^2u(x)) = 2(n+2) \inf_{\substack{A \in \mathcal{A} \\ A \leq \phi(\varepsilon)I}} \int_{B_\varepsilon(0)} \frac{u(x+Ay) - u(x)}{\varepsilon^2} dy + o(1), \quad \text{as } \varepsilon \rightarrow 0. \quad (1.9)$$

As a consequence, we can characterize viscosity solutions to the equation $F(D^2u(x)) = f(x)$ in terms of an asymptotic mean-value formula in the viscosity sense. For the precise definition of the notion of viscosity solutions and viscosity mean-value formulas, see Section 2 below, [12], and [19]. Informally, an equation or a mean-value property hold in the viscosity sense when they hold with an appropriate inequality (instead of an equality) for smooth test functions that touch u from above or below at x .

The concept of a mean-value formula in the viscosity sense is weaker than a mean-value formula that holds pointwise. For the infinity Laplacian and the Monge-Ampère equation, there are instances of asymptotic mean-value formulas that hold in the viscosity sense but do not hold pointwise, see [19] and [4], respectively. An interesting question is, under what circumstances do the viscosity mean-value formulas outlined above hold pointwise. This question has an obvious affirmative answer for all equations for which viscosity solutions are known to be classical since, in that case, we can apply the pointwise formula for regular functions in (1.9). Nevertheless, there are examples of non-classical viscosity solutions for which mean-value formulas hold pointwise; see [2, 4, 18].

Theorem 1.2. *Let $\phi(\varepsilon)$ be a positive function that satisfies (1.8). Consider $f \in C(\Omega)$ and $F : S^n(\mathbb{R}) \rightarrow \mathbb{R} \cup \{-\infty\}$ defined as in (1.3) that satisfies (1.6). Then, a function $u \in C(\Omega)$ is a viscosity subsolution (respectively, supersolution) of*

$$F(D^2u(x)) = f(x) \quad \text{in } \Omega,$$

if and only if

$$u(x) \leq \inf_{\substack{A \in \mathcal{A} \\ A \leq \phi(\varepsilon)I}} \int_{B_\varepsilon(0)} u(x+Ay) dy - \frac{\varepsilon^2}{2(n+2)} f(x) + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0$$

(respectively, \geq) in the viscosity sense.

Remark 1.3. Analogous results hold for supremum operators

$$F(D^2u(x)) = \sup_{A \in \mathcal{A}} \text{trace}(A^t D^2u(x) A)$$

and operators with a combination of an infimum and a supremum.

A fundamental example of application of Theorems 1.1 and 1.2 are the k -Hessian operators, given by the elementary symmetric polynomials

$$\sigma_k(\lambda_1, \dots, \lambda_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_k}$$

evaluated in the eigenvalues of the Hessian, $\{\lambda_i(D^2u)\}_{1 \leq i \leq n}$. Here, $k = 1, 2, \dots, n$ and the cases $k = 1$ and $k = n$ correspond to the Laplacian and Monge-Ampère, respectively. For these operators to fit our framework we need to write them in the form

$$F_k(D^2u(x)) = k \sigma_k \left(\lambda_1(D^2u(x)), \dots, \lambda_n(D^2u(x)) \right)^{\frac{1}{k}},$$

according to the following characterization, which we prove in Section 5.

Lemma 1.4. *Let $k \geq 2$ and*

$$\Gamma_k = \left\{ \lambda \in \mathbb{R}^n : \sigma_j(\lambda) > 0 \text{ for all } j = 1, \dots, k \right\}.$$

Then, for every $M \in S^n(\mathbb{R})$ we have

$$\inf_{A \in \mathcal{A}_k} \text{trace}(A^t M A) = \begin{cases} k \sigma_k(\lambda(M))^{\frac{1}{k}} & \text{if } M \in \bar{\Gamma}_k, \\ -\infty & \text{otherwise,} \end{cases} \quad (1.10)$$

where

$$\mathcal{A}_k = \left\{ A \in S_+^n(\mathbb{R}) : \lambda_i^2(A) = \sigma_{k-1,i}(\gamma) \text{ with } \gamma = (\gamma_1, \dots, \gamma_n) \in \Gamma_k \text{ and } \sigma_k(\gamma) = 1 \right\},$$

and $\sigma_{k-1,i}(\gamma_1, \dots, \gamma_n) = \sigma_{k-1}(\gamma_1, \dots, \gamma_{i-1}, 0, \gamma_{i+1}, \dots, \gamma_n)$.

An important consequence of Lemma 1.4 is that for the k -Hessian operators, \mathcal{A} -admissibility is equivalent to the notion of k -convexity in [25, 26, 27], see Corollary 5.3. This means that

$$\Gamma_{\mathcal{A}_k} = \{M \in S^n(\mathbb{R}) : F(M) > -\infty\} = \{M \in S^n(\mathbb{R}) : F(M) \geq 0\} = \bar{\Gamma}_k$$

and

$$F \equiv 0 \quad \text{on } \partial\Gamma_{\mathcal{A}_k}. \quad (1.11)$$

A condition such as (1.11) could be used in place of (1.6) to prove the mean-value property. In fact, (1.11) may even appear more natural at first; for instance, it was used in [13] in relation to the existence and uniqueness of solutions to general fully nonlinear second-order PDEs. However, condition (1.11) implies (1.6), but not vice versa (see Remark 4.4 and Example 4.5 below), making condition (1.6) more general.

In this case Theorem 1.1 reads as follows, see Section 5 for details.

Theorem 1.5. *Let $\phi(\varepsilon)$ be a positive function that satisfies (1.8) and assume that $u \in C^2(\Omega)$ is k -convex, that is, $\sigma_j(\lambda(D^2u(x))) \geq 0$ for all $j = 1, \dots, k$, for every $x \in \Omega$. Then, for every $x \in \Omega$ we have*

$$u(x) = \inf_{\substack{A \in \mathcal{A}_k \\ A \leq \phi(\varepsilon)I}} \int_{B_\varepsilon(0)} u(x + Ay) dy - \frac{\varepsilon^2}{2(n+2)} k (\sigma_k(D^2u(x)))^{\frac{1}{k}} + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0, \quad (1.12)$$

with a corresponding viscosity characterization as in Theorem 1.2.

It is also possible to consider operators where the class of matrices \mathcal{A} depends on the point x . This happens naturally when we consider mean-value formulas for Monge-Ampère operators in the Heisenberg group in Section 6. In that case, the sets \mathcal{A} depend on the point x and are unbounded.

We also obtain a rich family of examples when considering operators for which $\mathcal{A}_x \subset S_+^n(\mathbb{R})$ is bounded for each $x \in \Omega$. In fact, we consider sup-inf operators where for every x , the supremum is taken over a subset \mathbb{A}_x of $\mathcal{P}(S_+^n(\mathbb{R}))$, the power set of $S_+^n(\mathbb{R})$. This is not done only for the sake of generality. It is motivated by examples that cannot be covered otherwise, such as prescribed eigenvalues of the Hessian and Isaacs operators. These operators can be degenerate elliptic since we do not impose any lower bounds on the eigenvalues of the matrices $A \in \mathcal{A}_x$. We provide a list of examples below which shows the flexibility of the approach.

Therefore, let us now consider differential operators of the form $F(x, D^2u(x))$ with $F : \Omega \times S^n(\mathbb{R}) \rightarrow \mathbb{R}$ given by

$$F(x, M) = \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t M A). \quad (1.13)$$

Here, $\mathbb{A}_x \subset \mathcal{P}(S_+^n(\mathbb{R}))$ (the power set of $S_+^n(\mathbb{R})$) is a non-empty subset for each $x \in \mathbb{R}^n$ and we assume that

$$\bigcup \mathbb{A}_x = \left\{ A \in S_+^n(\mathbb{R}) : A \in \mathcal{A} \text{ for some } \mathcal{A} \in \mathbb{A}_x \right\} \text{ is bounded.} \quad (1.14)$$

Observe that if \mathbb{A}_x contains only one element \mathcal{A}_x for each x , then (1.13) is equivalent to

$$F(x, M) = \inf_{A \in \mathcal{A}_x} \text{trace}(A^t M A), \quad (1.15)$$

with $\mathcal{A}_x \subset S_+^n(\mathbb{R})$ bounded for each $x \in \mathbb{R}^n$. On the other hand, if every \mathbb{A}_x is a set of singletons, (1.13) is equivalent to

$$F(x, M) = \sup_{A \in \mathcal{A}_x} \text{trace}(A^t M A), \quad (1.16)$$

where $\mathcal{A}_x = \{A \in S_+^n(\mathbb{R}) : A \in \mathcal{A} \text{ for some } \mathcal{A} \in \mathbb{A}_x\}$. One can also consider inf-sup operators of the form

$$F(x, M) = \inf_{\mathcal{A} \in \mathbb{A}_x} \sup_{A \in \mathcal{A}} \text{trace}(A^t M A) \quad (1.17)$$

with straightforward adaptations in the statements and proofs.

By Lemma 4.1, (1.14) is equivalent to the operator $M \mapsto F(x, M)$ being well-defined and finite for every $M \in S^n(\mathbb{R})$; in particular, every $u \in C^2(\Omega)$ is admissible. Moreover, we prove in Lemma 3.2 that $M \mapsto F(x, M)$ is Lipschitz continuous for every $x \in \Omega$.

We obtain the following counterparts of Theorems 1.1 and 1.2. Observe that when $\bigcup \mathbb{A}_x$ is bounded, the condition $A \leq \phi(\varepsilon)I$ in (1.9) becomes unnecessary because it is satisfied at every point for ε small enough.

Theorem 1.6. *Consider $u \in C^2(\Omega)$ and let $F : \Omega \times S^n(\mathbb{R}) \rightarrow \mathbb{R}$ be an operator of the form (1.13) that satisfies (1.14). Then, for every $x \in \Omega$ we have*

$$F(x, D^2 u(x)) = 2(n+2) \sup_{\mathcal{A} \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \int_{B_\varepsilon(0)} \frac{u(x+Ay) - u(x)}{\varepsilon^2} dy + o(1), \quad \text{as } \varepsilon \rightarrow 0. \quad (1.18)$$

We have a corresponding result in the line of Theorem 1.2, characterizing viscosity solutions to the equation $F(x, D^2 u(x)) = f(x)$ by an asymptotic mean-value formula in the viscosity sense.

Theorem 1.7. *Consider $f \in C(\Omega)$ and $F : S^n(\mathbb{R}) \rightarrow \mathbb{R}$ defined as in (1.13) that satisfies (1.14). Then, a function $u \in C(\Omega)$ is a viscosity subsolution (respectively, supersolution) of*

$$F(D^2 u(x)) = f(x) \quad \text{in } \Omega,$$

if and only if

$$u(x) \leq \sup_{\mathcal{A} \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \int_{B_\varepsilon(0)} u(x+Ay) dy - \frac{\varepsilon^2}{2(n+2)} f(x) + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0$$

(respectively, \geq) in the viscosity sense.

Let us now provide some examples of operators of the form (1.13), (1.15), and (1.16) and the corresponding sets \mathbb{A}_x and \mathcal{A}_x . In the sequel, we consider the eigenvalues of a matrix $M \in S^n(\mathbb{R})$ arranged in increasing order, that is, $\lambda_1(M) \leq \lambda_2(M) \leq \dots \leq \lambda_n(M)$.

(1) Operators of the form (1.13), (1.17) include the usual Isaacs operators (see [8]) defined as

$$F(x, M) = \sup_{\alpha \in \mathcal{A}} \inf_{\beta \in \mathcal{B}} \text{trace}(A_{\alpha\beta}^t M A_{\alpha\beta}) \quad \text{and} \quad F(x, M) = \inf_{\alpha \in \mathcal{A}} \sup_{\beta \in \mathcal{B}} \text{trace}(A_{\alpha\beta}^t M A_{\alpha\beta})$$

for a given family of matrices $\{A_{\alpha\beta}\}_{\alpha\in\mathcal{A},\beta\in\mathcal{B}}$ (and similarly in the inf-sup case). Here we take $\mathbb{A} = \{\{A_{\alpha\beta}\}_{\beta\in\mathcal{B}}\}_{\alpha\in\mathcal{A}}$ in (1.13), with \mathbb{A} independent of the point x . In fact, operators of the form (1.13) could be seen as Issacs operators, but the usual Isaacs condition

$$\inf_{\alpha\in\mathcal{A}} \sup_{\beta\in\mathcal{B}} \text{trace}(A_{\alpha\beta}^t M A_{\alpha\beta}) = \sup_{\beta\in\mathcal{B}} \inf_{\alpha\in\mathcal{A}} \text{trace}(A_{\alpha\beta}^t M A_{\alpha\beta})$$

does not seem to have a clear counterpart. A typical requirement in the literature is that the family $\{A_{\alpha\beta}\}_{\alpha\in\mathcal{A},\beta\in\mathcal{B}}$ is uniformly elliptic, i.e., it is uniformly bounded between two constants $0 < \theta < \Theta$. In this case, the resulting Issacs operator is uniformly elliptic. We want to emphasize that here we are only requiring $\cup\mathbb{A}_x$ to be bounded and, therefore, the operators (1.13) may be degenerate elliptic.

- (2) We consider the truncated Laplacians [3], defined as

$$\mathcal{P}_k^-(D^2u) = \sum_{i=1}^k \lambda_i(D^2u) \quad \text{and} \quad \mathcal{P}_k^+(D^2u) = \sum_{i=1}^k \lambda_{n+1-i}(D^2u),$$

for $k = 1, 2, \dots, n-1$ (for $k = n$ these operators coincide with the Laplacian). These degenerate operators appear naturally in geometric problems, when considering manifolds of partially positive curvature [24, 29], or mean curvature flow in arbitrary codimension [1], see also [13, 14, 10, 3] and the references therein. The operators \mathcal{P}_k^- and \mathcal{P}_k^+ are of the form (1.15) and (1.16), respectively, for the set

$$\mathcal{A} = \{A \in S_+^n(\mathbb{R}) : \lambda_1 = \dots = \lambda_{n-k} = 0 \text{ and } \lambda_{n-k+1} = \dots = \lambda_n = 1\}.$$

- (3) Operators of the form (1.13) allow us to consider degenerate operators such as the k -th smallest eigenvalue of the Hessian, given by the Courant–Fischer min-max principle

$$\lambda_k(D^2u(x)) = \max_V \left\{ \min_{v \in V, |v|=1} \langle D^2u(x)v, v \rangle : V \subset \mathbb{R}^n \text{ subspace, } \dim(V) = n - k + 1 \right\}. \quad (1.19)$$

We can write the operator $\lambda_k(D^2u)$ in the form (1.13) for the set

$$\mathbb{A} = \left\{ \{A \in S_+^n(\mathbb{R}) : \lambda_i(A) = 0 \text{ for } i \neq n, \lambda_n(A) = 1, \text{ and } v_n \in V\} : \right. \\ \left. V \subset \mathbb{R}^n \text{ is a subspace of dimension } n - k + 1 \right\},$$

where v_n is the eigenvector corresponding to $\lambda_n(A)$. The cases $k = 1$ and $k = n$ were studied in [5, 13, 22] in connection with the convex and concave envelope of a function; i.e., the unique viscosity solutions of $\lambda_1(D^2u(x)) = 0$ and $\lambda_n(D^2u(x)) = 0$ are, respectively, the convex and concave envelopes of $u|_{\partial\Omega}$ in Ω . These operators are of the form (1.15), (1.16) with the set of matrices

$$\mathcal{A} = \left\{ A \in S_+^n(\mathbb{R}) : \lambda_1(A) = \dots = \lambda_{n-1}(A) = 0 \text{ and } \lambda_n(A) = 1 \right\}.$$

- (4) When the mean-value formula involves averages over balls that are not centered at 0 but at $\varepsilon^2 v$ with $|v| = 1$, we obtain operators with first-order terms. For example, we have

$$\inf_{A \in \mathcal{A}_x} \int_{B_\varepsilon(\varepsilon^2 v)} u(x + Ay) dy - u(x) = \varepsilon^2 \inf_{A \in \mathcal{A}_x} \left\{ \frac{1}{2(n+2)} \text{trace}(A^t D^2u(x)A) + \langle Du(x), Av \rangle \right\} + o(\varepsilon^2),$$

as $\varepsilon \rightarrow 0$. We can also look for zero-order terms and consider mean-value properties like

$$\inf_{A \in \mathcal{A}_x} (1 - \alpha\varepsilon^2) \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x) = \varepsilon^2 \left\{ \frac{1}{2(n+2)} \inf_{A \in \mathcal{A}_x} \text{trace}(A^t D^2u(x)A) - \alpha u(x) \right\} + o(\varepsilon^2),$$

as $\varepsilon \rightarrow 0$.

(5) Finally, we consider extremal Pucci operators for given ellipticity constants $0 < \theta < \Theta$, defined as

$$\mathcal{M}_{\theta, \Theta}^-(D^2u) = \theta \sum_{\lambda_i(D^2u) > 0} \lambda_i(D^2u) + \Theta \sum_{\lambda_i(D^2u) < 0} \lambda_i(D^2u)$$

and

$$\mathcal{M}_{\theta, \Theta}^+(D^2u) = \Theta \sum_{\lambda_i(D^2u) > 0} \lambda_i(D^2u) + \theta \sum_{\lambda_i(D^2u) < 0} \lambda_i(D^2u).$$

These operators are of the form (1.15), (1.16) with

$$\mathcal{A}_{\theta\Theta} = \left\{ A \in S_+^n(\mathbb{R}) : \sqrt{\theta} \leq \lambda_i(A) \leq \sqrt{\Theta} \right\},$$

which is bounded uniformly in x . From the definition of $\mathcal{A}_{\theta\Theta}$, it is clear that $\mathcal{M}_{\theta, \Theta}^\pm$ are uniformly elliptic operators. We apply similar ideas to general fully nonlinear uniformly elliptic operators $F(x, D^2u(x))$ (with ellipticity constants $0 < \theta \leq \Theta$) by means of the characterization

$$F(x, M) = \sup_{N \in S^n(\mathbb{R})} \inf_{A \in \mathcal{A}_{\theta\Theta}} \left\{ \text{trace}(A^t M A) + F(x, N) - \text{trace}(A^t N A) \right\},$$

see Lemma 3.4.

The paper is organized as follows: In Section 2 we gather some definitions and preliminary results and in Section 3 we deal with concave and Isaacs operators, when the sets of coefficients are bounded. These include general uniformly elliptic operators and operators including lower-order terms. We consider bounded coefficients first for the sake of clarity, to establish the techniques and ideas before we dive into the unbounded case in Section 4. In Section 5 we study the k -Hessians. Finally, Section 6 is devoted to examples in the Heisenberg group, where the class \mathcal{A}_x is unbounded and naturally depends on x .

2. PRELIMINARIES

We begin by stating the definition of a viscosity solution to a fully nonlinear, second-order, elliptic PDE. We refer to [12] for general results on viscosity solutions. Given a continuous function

$$\mathcal{F} : \Omega \times \mathbb{R} \times \mathbb{R}^n \times S^n(\mathbb{R}) \rightarrow \mathbb{R},$$

we consider the PDE

$$\mathcal{F}(x, u(x), Du(x), D^2u(x)) = 0, \quad x \in \Omega. \quad (2.1)$$

Viscosity solutions use the monotonicity of \mathcal{F} in D^2u (ellipticity) in order to “pass derivatives to smooth test functions”.

Definition 2.1. *A lower semi-continuous function u is a viscosity supersolution of (2.1) if for every $\phi \in C^2$ such that ϕ touches u at $x \in \Omega$ strictly from below (that is, $u - \phi$ has a strict minimum at x with $u(x) = \phi(x)$), we have*

$$\mathcal{F}(x, \phi(x), D\phi(x), D^2\phi(x)) \leq 0.$$

An upper semi-continuous function u is a subsolution of (2.1) if for every $\phi \in C^2$ such that ϕ touches u at $x \in \Omega$ strictly from above (that is, $u - \phi$ has a strict maximum at x with $u(x) = \phi(x)$), we have

$$\mathcal{F}(x, \phi(x), D\phi(x), D^2\phi(x)) \geq 0.$$

Finally, u is a viscosity solution of (2.1) if it is both a super- and a subsolution.

We will also need the definition of an asymptotic mean-value formula in the viscosity sense. In the next definition, $M(u, \varepsilon)(x)$ stands for a mean-value operator (that depends on the parameter ε) applied to u at the point x . For example, we can take

$$M(u, \varepsilon)(x) = \inf_{A \in \mathbb{A}_x} \int_{B_\varepsilon(0)} u(x + Ay) dy - \frac{\varepsilon^2}{2(n+2)} f(x)$$

as in Theorem 1.2 and the next section.

Definition 2.2. *A continuous function u verifies*

$$u(x) \geq M(u, \varepsilon)(x) + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0,$$

in the viscosity sense if for every $\phi \in C^2$ such that $u - \phi$ has a strict minimum at the point $x \in \bar{\Omega}$ with $u(x) = \phi(x)$ (i.e., ϕ touches u at $x \in \Omega$ strictly from below), we have

$$\phi(x) \geq M(\phi, \varepsilon)(x) + o(\varepsilon^2).$$

Similarly, a continuous function u verifies

$$u(x) \leq M(u, \varepsilon)(x) + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0,$$

in the viscosity sense if for every $\phi \in C^2$ such that $u - \phi$ has a strict maximum at the point $x \in \bar{\Omega}$ with $u(x) = \phi(x)$ (i.e., ϕ touches u at $x \in \Omega$ strictly from above), we have

$$\phi(x) \leq M(\phi, \varepsilon)(x) + o(\varepsilon^2).$$

Next, we include a lemma that is related to the simplest mean-value property for the usual Laplacian. A proof can be found in [4].

Lemma 2.3. *Let M be a square matrix of dimension n . Then,*

$$\text{trace}(M) = \frac{n}{\varepsilon^2} \int_{\partial B_\varepsilon(0)} \langle My, y \rangle d\mathcal{H}^{n-1}(y) = \frac{n+2}{\varepsilon^2} \int_{B_\varepsilon(0)} \langle My, y \rangle dy.$$

Remark 2.4. We will use the solid mean

$$\text{trace}(M) = \frac{n+2}{\varepsilon^2} \int_{B_\varepsilon(0)} \langle My, y \rangle dy,$$

in our proofs of the mean-value formulas. However, if one uses the mean on spheres

$$\text{trace}(M) = \frac{n}{\varepsilon^2} \int_{\partial B_\varepsilon(0)} \langle My, y \rangle d\mathcal{H}^{n-1}(y)$$

one can obtain mean-value formulas of the type

$$\inf_{A \in \mathbb{A}_x} \int_{\partial B_\varepsilon(0)} u(x + Ay) dy - u(x) = \frac{\varepsilon^2}{2n} F(x, D^2u(x)) + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0, \quad (2.2)$$

for $u \in C^2(\Omega)$ and $F(x, D^2u)$ defined as in (1.15) (and accordingly for the rest of the cases).

3. BOUNDED OPERATORS

In this section we prove Theorems 1.6 and 1.7. We consider differential operators given by

$$F(x, D^2u(x)) = \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t D^2u(x) A), \quad (3.1)$$

where $\mathbb{A}_x \subset \mathcal{P}(S_+^n(\mathbb{R}))$ (the power set of $S_+^n(\mathbb{R})$) is a non-empty subset for each $x \in \mathbb{R}^n$ and

$$\bigcup \mathbb{A}_x = \left\{ A \in S_+^n(\mathbb{R}) : A \in \mathcal{A} \text{ for some } \mathcal{A} \in \mathbb{A}_x \right\} \quad \text{is bounded.} \quad (3.2)$$

Due to (3.2), the operator $F(x, M)$ is finite for every $M \in S^n(\mathbb{R})$.

Moreover, since the set of matrices $\cup \mathbb{A}_x$ is bounded, the mean-value formula (1.18) is local. In fact, for every $x \in \Omega$ and $y \in B_\varepsilon(0)$ there exists $C_x > 0$ such that $A \leq C_x I$ for every $A \in \cup \mathbb{A}_x$. We get

$$\text{dist}(x + Ay, x) = |Ay| \leq C_x \varepsilon \rightarrow 0$$

as $\varepsilon \rightarrow 0$. In particular, observe that for ε small enough $x + Ay \in \Omega$ for every $y \in B_\varepsilon(0)$ and hence the integrals

$$\int_{B_\varepsilon(0)} u(x + Ay) dy$$

are well-defined for integrable functions $u : \Omega \rightarrow \mathbb{R}$.

Remark 3.1. We can assume that the matrices $A \in \cup \mathbb{A}_x$ are symmetric and positive semi-definite without loss of generality. This is because, given $A \in M^{n \times n}(\mathbb{R})$, we can write its left polar decomposition $A = SQ$ with Q orthogonal and S positive semi-definite and symmetric. Then, for every $M \in M^{n \times n}(\mathbb{R})$, we have

$$\text{trace}(A^t M A) = \text{trace}(Q^t S^t M S Q) = \text{trace}(S^t M S Q Q^t) = \text{trace}(S^t M S)$$

and

$$\int_{B_\varepsilon(0)} u(x + Ay) dy = \int_{B_\varepsilon(0)} u(x + SQy) dy = \int_{B_\varepsilon(0)} u(x + Sz) dz.$$

In the next lemma we prove an explicit continuity estimate that we use in the proof of Theorem 1.6.

Lemma 3.2. *Consider the differential operator $F(x, D^2 u(x))$ given by (3.1). If $\cup \mathbb{A}_x$ is bounded, then the mapping $M \mapsto F(x, M)$ is Lipschitz continuous in M . In particular,*

$$\sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t (M \pm \eta I) A) \rightarrow \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t M A) \quad (3.3)$$

as $\eta \rightarrow 0$, for every $M \in S^n(\mathbb{R})$.

Proof. We fix $x \in \mathbb{R}^n$. For every $A \in \cup \mathbb{A}_x$, there exists $C_x > 0$ such that $A \leq C_x I$. Given $M, N \in S^n(\mathbb{R})$, we have

$$\begin{aligned} & \left| \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t M A) - \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t N A) \right| \\ & \leq \sup_{A \in \cup \mathbb{A}_x} |\text{trace}(A^t M A) - \text{trace}(A^t N A)| \leq \sup_{A \in \cup \mathbb{A}_x} |\text{trace}(A^t (M - N) A)| \leq n C_x^2 \|M - N\|, \end{aligned}$$

and the result follows. \square

We are now ready to prove Theorem 1.6.

Proof of Theorem 1.6. Given $x \in \Omega$, let us consider the paraboloid

$$P(z) = u(x) + \langle \nabla u(x), z - x \rangle + \frac{1}{2} \langle D^2 u(x)(z - x), (z - x) \rangle. \quad (3.4)$$

Since $u \in C^2(\Omega)$, we have

$$u(z) - P(z) = o(|z - x|^2) \quad \text{as } z \rightarrow x,$$

which means that for every $\eta > 0$, there exists $\delta > 0$ such that for every $z \in B_\delta(x)$,

$$P(z) - \frac{\eta}{2} |z - x|^2 \leq u(z) \leq P(z) + \frac{\eta}{2} |z - x|^2, \quad (3.5)$$

with equality only when $z = x$. For convenience, let us denote

$$P_\eta^\pm(z) = P(z) \pm \frac{\eta}{2} |z - x|^2.$$

Then,

$$\begin{aligned} \int_{B_\varepsilon(0)} (P_\eta^\pm(x + Ay) - P_\eta^\pm(x)) dy &= \frac{1}{2} \int_{B_\varepsilon(0)} (\langle A^t D^2 u(x) Ay, y \rangle \pm \eta |Ay|^2) dy \\ &= \frac{1}{2} \int_{B_\varepsilon(0)} \langle A^t (D^2 u(x) \pm \eta I) Ay, y \rangle dy = \frac{\varepsilon^2}{2(n+2)} \text{trace} (A^t (D^2 u(x) \pm \eta I) A), \end{aligned} \quad (3.6)$$

by Lemma 2.3.

On the other hand, since $\cup \mathbb{A}_x$ is bounded there exists $C_x > 0$ such that $A \leq C_x I$ for every $A \in \cup \mathbb{A}_x$. Then, $x + Ay \in B_\delta(x)$ for every $|y| \leq \varepsilon$ and $\varepsilon < \varepsilon_0$, where $\varepsilon_0 C_x \leq \delta$. Therefore, by (3.5), if $\varepsilon < \varepsilon_0$, then

$$P_\eta^-(x + Ay) \leq u(x + Ay) \leq P_\eta^+(x + Ay) \quad \text{for every } y \in B_\varepsilon. \quad (3.7)$$

Then,

$$\begin{aligned} \frac{\varepsilon^2}{2(n+2)} \text{trace} (A^t (D^2 u(x) - \eta I) A) \\ \leq \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x) \leq \frac{\varepsilon^2}{2(n+2)} \text{trace} (A^t (D^2 u(x) + \eta I) A) \end{aligned}$$

and the result follows by (3.3). \square

We now prove Theorem 1.7.

Proof of Theorem 1.7. First, assume that u is a viscosity solution and take a C^2 function ϕ that touches u at $x \in \Omega$ strictly from below. Then, as $u - \phi$ has a strict minimum at x with $u(x) = \phi(x)$, we have

$$F(x, D^2 \phi(x)) = \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t D^2 \phi(x) A) \leq f(x).$$

Now, since ϕ is C^2 , Theorem 1.6 gives

$$\sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \int_{B_\varepsilon(0)} \phi(x + Ay) dy - u(x) + o(\varepsilon^2) = \frac{\varepsilon^2}{2(n+2)} \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \text{trace}(A^t D^2 \phi(x) A) \leq \frac{\varepsilon^2}{2(n+2)} f(x),$$

proving that u satisfies

$$u(x) \geq \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \int_{B_\varepsilon(0)} u(x + Ay) dy - \frac{\varepsilon^2}{2(n+2)} f(x) + o(\varepsilon^2),$$

in the viscosity sense. An analogous computation reversing the inequalities shows that when a C^2 function ϕ touches u at $x \in \Omega$ strictly from above, u satisfies

$$u(x) \leq \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \int_{B_\varepsilon(0)} u(x + Ay) dy - \frac{\varepsilon^2}{2(n+2)} f(x) + o(\varepsilon^2)$$

in the viscosity sense.

Now, assume that the mean-value property holds and take a C^2 function ϕ that touches u at $x \in \Omega$ strictly from below. Then, as $u - \phi$ has a strict minimum at x with $u(x) = \phi(x)$, Theorem 1.6 yields

$$\frac{\varepsilon^2}{2(n+2)} F(x, D^2 \phi(x)) + o(\varepsilon^2) = \sup_{A \in \mathbb{A}_x} \inf_{A \in \mathcal{A}} \int_{B_\varepsilon(0)} \phi(x + Ay) dy - u(x) \leq \frac{\varepsilon^2}{2(n+2)} f(x) + o(\varepsilon^2).$$

Dividing by ε^2 and letting $\varepsilon \rightarrow 0$ we get that u is a viscosity supersolution to

$$F(x, D^2 u(x)) = f(x).$$

A similar argument reversing the inequalities shows that u is a viscosity subsolution. \square

We devote the rest of the section to discuss the examples mentioned in the introduction in more detail.

3.1. Isaacs Operators. As mentioned in the introduction, hypotheses (3.1) and (3.2) also cover degenerate operators such as the k -th smallest eigenvalue of the Hessian, given by the Courant–Fischer min-max principle

$$\lambda_k(D^2u(x)) = \max_V \left\{ \min_{v \in V, |v|=1} \langle D^2u(x)v, v \rangle : V \subset \mathbb{R}^n \text{ subspace of dimension } n - k + 1 \right\}. \quad (3.8)$$

To write this operator in the form (3.1), let $A_v = v \otimes v$ and note that $A_v^2 = A_v$ when $|v| = 1$. With this notation we have $\text{trace}(A_v^t M A_v) = \langle Mv, v \rangle$. Let G_{n-k+1} denote the set of all subspaces $V \subset \mathbb{R}^n$ of dimension $n - k + 1$. For each $V \in G_{n-k+1}$, we set $\mathcal{A}_V = \{v \otimes v : v \in V \text{ such that } |v| = 1\}$. Finally we define $\mathbb{A}_k = \{\mathcal{A}_V : V \in G_{n-k+1}\}$, or equivalently,

$$\mathbb{A}_k = \left\{ \{A \in S_+^n(\mathbb{R}) : \lambda_i(A) = 0 \text{ for } i \neq n, \lambda_n(A) = 1, \text{ and } v_n \in V\} : V \subset \mathbb{R}^n \text{ subspace of dim. } n - k + 1 \right\},$$

where v_n is the eigenvector corresponding to $\lambda_n(A)$. We can then write

$$\lambda_k(M) = \sup_V \left\{ \inf_{v \in V, |v|=1} \langle Mv, v \rangle \right\} = \sup_{\mathcal{A}_V \in \mathbb{A}_k} \inf_{A \in \mathcal{A}_V} \text{trace}(A^t M A).$$

This allows us to prove mean-value formulas for (3.8). In fact, Theorem 1.6 gives

$$\sup_{\mathcal{A}_V \in \mathbb{A}_k} \inf_{A \in \mathcal{A}_V} \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x) = \frac{\varepsilon^2}{2(n+2)} \lambda_k(D^2u(x)) + o(\varepsilon^2), \quad (3.9)$$

and the corresponding viscosity analogue following Theorem 1.7.

A variant of (3.9) is contained in [6],

$$\sup_{\dim(V)=n-k+1} \inf_{v \in V, |v|=1} \left\{ \frac{u(x + \varepsilon v) + u(x - \varepsilon v)}{2} \right\} - u(x) = \frac{\varepsilon^2}{2} \lambda_k(D^2u(x)) + o(\varepsilon^2). \quad (3.10)$$

In order to make the connection between both mean-value formulas, (3.9) and (3.10), let $A_v = v \otimes v$ with $|v| = 1$ as before and observe that

$$\begin{aligned} 2(n+2) \left(\int_{B_\varepsilon(0)} u(x + A_v y) dy - u(x) \right) &= \varepsilon^2 \langle D^2u(x)v, v \rangle + o(\varepsilon^2) \\ &= u(x + \varepsilon v) + u(x - \varepsilon v) - 2u(x) + o(\varepsilon^2), \end{aligned}$$

where the error estimate is uniform in v .

3.2. Uniformly elliptic fully nonlinear operators. Let us state the definition of a uniformly elliptic operator for completeness.

Definition 3.3. *An operator $F : \Omega \times S^n(\mathbb{R}) \rightarrow \mathbb{R}$ is uniformly elliptic with constants $0 < \theta \leq \Theta$ if and only if for every $M, N \in S^n(\mathbb{R})$ with $N \geq 0$,*

$$\theta \cdot \text{trace}(N) \leq F(x, M + N) - F(x, M) \leq \Theta \cdot \text{trace}(N) \quad (3.11)$$

for all $x \in \Omega$.

Here we modify the formulas from the previous section to obtain a mean-value formula for general uniformly elliptic operators. This is possible because every uniformly elliptic operator can be written as an Isaacs operator. We include a proof for the reader's convenience, see also [9].

Lemma 3.4. *Let*

$$\mathcal{A}_{\theta\Theta} = \left\{ A \in S^n(\mathbb{R}) : \sqrt{\theta}|\xi|^2 \leq \langle A\xi, \xi \rangle \leq \sqrt{\Theta}|\xi|^2 \quad \forall \xi \in \mathbb{R}^n \right\}.$$

An operator $F : \Omega \times S^n(\mathbb{R}) \rightarrow \mathbb{R}$ is uniformly elliptic with constants $0 < \theta \leq \Theta$ if and only if

$$\begin{aligned} F(x, M) &= \inf_{N \in S^n(\mathbb{R})} \sup_{A \in \mathcal{A}_{\theta\Theta}} \left\{ \text{trace}(A^t M A) + F(x, N) - \text{trace}(A^t N A) \right\}, \\ F(x, M) &= \sup_{N \in S^n(\mathbb{R})} \inf_{A \in \mathcal{A}_{\theta\Theta}} \left\{ \text{trace}(A^t M A) + F(x, N) - \text{trace}(A^t N A) \right\}. \end{aligned} \quad (3.12)$$

Proof. It is well known that an operator $F : \Omega \times S^n(\mathbb{R}) \rightarrow \mathbb{R}$ is uniformly elliptic if and only if,

$$\inf_{A \in \mathcal{A}_{\theta\Theta}} \text{trace}(A^t(M - N)A) \leq F(x, M) - F(x, N) \leq \sup_{A \in \mathcal{A}_{\theta\Theta}} \text{trace}(A^t(M - N)A)$$

for every $M, N \in S^n(\mathbb{R})$ and $x \in \mathbb{R}$. Since we have equalities when $M = N$, in particular (3.12) holds. \square

With this characterization, we can prove the following.

Theorem 3.5. *Consider $F : \Omega \times S^n(\mathbb{R}) \rightarrow \mathbb{R}$ uniformly elliptic and let $u \in C^2(\Omega)$. Then, for every $x \in \Omega$ we have*

$$\inf_{N \in S^n(\mathbb{R})} \sup_{A \in \mathcal{A}} \left(\int_{B_\varepsilon(0)} u(x + Ay) dy + F(x, N) - \text{trace}(A^t N A) \right) - u(x) = \frac{\varepsilon^2}{2(n+2)} F(x, D^2 u(x)) + o(\varepsilon^2),$$

as $\varepsilon \rightarrow 0$, where $\mathcal{A} = \left\{ A \in S_+^n(\mathbb{R}) : \sqrt{\theta}|\xi|^2 \leq \langle A\xi, \xi \rangle \leq \sqrt{\Theta}|\xi|^2, \quad \forall \xi \in \mathbb{R}^n \right\}$.

Proof of Theorem 3.5. Since the family of matrices $\mathcal{A}_{\theta\Theta}$ is uniformly elliptic, in particular it is bounded. Then, an analogous result to Lemma 3.2 can be obtained for the operators considered here and the proof follows in the same way as the one of Theorem 1.6. From the expression

$$F(x, D^2 u(x)) = \inf_{N \in S^n(\mathbb{R})} \sup_{A \in \mathcal{A}_{\theta\Theta}} \left\{ \text{trace}(A^t D^2 u(x) A) + F(x, N) - \text{trace}(A^t N A) \right\},$$

we obtain that for a smooth function u it holds that

$$\inf_{N \in S^n(\mathbb{R})} \sup_{A \in \mathcal{A}} \left(\int_{B_\varepsilon(0)} u(x + Ay) dy + F(x, N) - \text{trace}(A^t N A) \right) - u(x) = \frac{\varepsilon^2}{2(n+2)} F(x, D^2 u(x)) + o(\varepsilon^2),$$

as $\varepsilon \rightarrow 0$, where $\mathcal{A} = \left\{ A \in S_+^n(\mathbb{R}) : \sqrt{\theta}|\xi|^2 \leq \langle A\xi, \xi \rangle \leq \sqrt{\Theta}|\xi|^2, \quad \forall \xi \in \mathbb{R}^n \right\}$. \square

Remark 3.6. Observe that for some operators both Theorems 1.6 and 3.5 apply. Even then, the formulas that we get are different. For example, for Issacs operators of the form

$$F(x, D^2 u(x)) = \sup_{\alpha \in \mathcal{A}} \inf_{\beta \in \mathcal{B}} \text{trace}(A_{\alpha\beta}^t D^2 u(x) A_{\alpha\beta}),$$

we get two possible mean-value formulas, namely,

$$\inf_{\alpha \in \mathcal{A}} \sup_{\beta \in \mathcal{B}} \int_{B_\varepsilon(0)} u(x + A_{\alpha\beta} y) dy - u(x) = \frac{\varepsilon^2}{2(n+2)} F(x, D^2 u(x)) + o(\varepsilon^2),$$

and

$$\begin{aligned} \inf_{N \in S^n(\mathbb{R})} \sup_{A \in \mathcal{A}_{\theta\Theta}} \left(\int_{B_\varepsilon(0)} u(x + Ay) dy + F(x, N) - \text{trace}(A^t N A) \right) - u(x) \\ = \frac{\varepsilon^2}{2(n+2)} F(x, D^2 u(x)) + o(\varepsilon^2). \end{aligned}$$

3.3. Off-center means and equations involving lower-order terms. When the mean-value formula involves integrals in balls that are not centered at 0 but at $\varepsilon^2 v$ with $|v| = 1$, we obtain second-order operators with first-order terms,

$$\inf_{A \in \mathcal{A}_x} \int_{B_\varepsilon(\varepsilon^2 v)} \frac{u(x + Ay) - u(x)}{\varepsilon^2} dy = \inf_{A \in \mathcal{A}_x} \left\{ \frac{1}{2(n+2)} \text{trace}(A^t D^2 u(x) A) + \langle Du(x), Av \rangle \right\} + o(1), \quad (3.13)$$

as $\varepsilon \rightarrow 0$. Here we assume for simplicity that $\mathcal{A}_x \subset S_+^n(\mathbb{R})$ is bounded for each $x \in \Omega$; whenever \mathcal{A}_x is not bounded, we can restrict the argument to matrices that satisfy $A \leq \phi(\varepsilon)I$ for $\phi(\varepsilon)$ satisfying (1.8).

The mean-value formula (3.13) is a consequence of the fact that for a C^2 function we have

$$\begin{aligned} \int_{B_\varepsilon(\varepsilon^2 v)} u(x + Ay) dy - u(x) &= \int_{B_\varepsilon(0)} u(x + A(z + \varepsilon^2 v)) dz - u(x) \\ &= \int_{B_\varepsilon(0)} \left(\langle Du(x), A(z + \varepsilon^2 v) \rangle + \frac{1}{2} \langle D^2 u(x) A(z + \varepsilon^2 v), A(z + \varepsilon^2 v) \rangle \right) dz + o(\varepsilon^2) \\ &= \varepsilon^2 \langle Du(x), Av \rangle + \frac{1}{2} \int_{B_\varepsilon(0)} \langle D^2 u(x) Az, Az \rangle dz + o(\varepsilon^2) \\ &= \varepsilon^2 \left\{ \langle Du(x), Av \rangle + \frac{1}{2(n+2)} \text{trace}(A^t D^2 u(x) A) \right\} + o(\varepsilon^2). \end{aligned}$$

Notice that the remainder $o(\varepsilon^2)$ is independent of A because \mathcal{A}_x is a bounded set for each $x \in \Omega$.

In addition, we observe that when we center the average at $\varepsilon^\alpha v$ we obtain: for $\alpha > 2$ a pure second-order operator,

$$\inf_{A \in \mathcal{A}_x} \int_{B_\varepsilon(\varepsilon^\alpha v)} u(x + Ay) dy - u(x) = \frac{\varepsilon^2}{2(n+2)} \inf_{A \in \mathcal{A}_x} \text{trace}(A^t D^2 u(x) A) + o(\varepsilon^2),$$

as $\varepsilon \rightarrow 0$; and for $0 < \alpha < 2$ a pure first-order operator

$$\inf_{A \in \mathcal{A}_x} \int_{B_\varepsilon(\varepsilon^\alpha v)} u(x + Ay) dy - u(x) = \varepsilon^\alpha \inf_{A \in \mathcal{A}_x} \langle Du(x), Av \rangle + o(\varepsilon^\alpha),$$

as $\varepsilon \rightarrow 0$.

Also, we can look for zero-order terms and consider mean-value properties like

$$(1 - \alpha\varepsilon^2) \inf_{A \in \mathcal{A}_x} \int_{B_\varepsilon(0)} \frac{u(x + Ay) - u(x)}{\varepsilon^2} dy = \frac{1}{2(n+2)} \inf_{A \in \mathcal{A}_x} \text{trace}(A^t D^2 u(x) A) - \alpha u(x) + o(1),$$

as $\varepsilon \rightarrow 0$. Arguing as before, we have

$$\begin{aligned} (1 - \alpha\varepsilon^2) \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x) &= \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x) - \alpha\varepsilon^2 \int_{B_\varepsilon(0)} u(x + Ay) dy \\ &= \varepsilon^2 \left\{ \frac{1}{2(n+2)} \text{trace}(A^t D^2 u(x) A) - \alpha u(x) \right\} + o(\varepsilon^2). \end{aligned}$$

Similar mean-value formulas also hold for sup-inf operators with lower-order terms. We leave the details to the reader.

4. UNBOUNDED OPERATORS

In this section we deal with unbounded operators $F : S^n(\mathbb{R}) \rightarrow \mathbb{R} \cup \{-\infty\}$ given by

$$F(M) = \inf_{A \in \mathcal{A}} \text{trace}(A^t M A). \quad (4.1)$$

As mentioned in the introduction, we restrict the set of matrices where we compute the infimum in the mean-value property by considering matrices $A \in \mathcal{A}$ such that $A \leq \phi(\varepsilon)I$, with $\phi(\varepsilon)$ a positive function satisfying

$$\lim_{\varepsilon \rightarrow 0} \phi(\varepsilon) = +\infty \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \varepsilon \phi(\varepsilon) = 0. \quad (4.2)$$

The condition $A \leq \phi(\varepsilon)I$ becomes less restrictive as $\varepsilon \rightarrow 0$, but is still enough to make the mean-value formula (1.9) local. In fact, for every $x \in \Omega$ and $|y| \leq \varepsilon$,

$$\text{dist}(x + Ay, x) = |Ay| \leq \varepsilon \phi(\varepsilon) \leq \text{dist}(x, \partial\Omega)$$

for ε sufficiently small (since $\varepsilon \phi(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$).

Recall that we have assumed (1.6) and therefore

$$F \text{ is continuous in the cone } \Gamma_{\mathcal{A}} = \{M \in S^n(\mathbb{R}) : F(M) > -\infty\}.$$

Next, we prove that, as long as \mathcal{A} is unbounded, there are matrices $M \in S^n(\mathbb{R})$ for which $F(M) = -\infty$.

Lemma 4.1. *The operator*

$$F(x, M) = \inf_{A \in \mathcal{A}_x} \text{trace}(A^t M A)$$

is finite for every $M \in S^n(\mathbb{R})$ if and only if \mathcal{A}_x is bounded.

Proof. It is clear that if \mathcal{A}_x is bounded then $F(x, M)$ is finite for every $M \in S^n(\mathbb{R})$. To prove the converse, suppose that $F(x, M)$ is finite for every $M \in S^n(\mathbb{R})$ and \mathcal{A}_x is not bounded. Then there exists a sequence of matrices $A_k \in \mathcal{A}_x$ such that their largest eigenvalues $\lambda_n(A_k)$ diverge as $k \rightarrow \infty$. Write $A_k = Q_k^t D_k Q_k$, where Q_k is an orthogonal matrix and D_k is a diagonal matrix with diagonal entries $(D_k)_{jj} = \lambda_j(A_k)$. Let us assume that the eigenvalues of A_k satisfy $0 \leq \lambda_1(A_k) \leq \lambda_2(A_k) \leq \dots \leq \lambda_n(A_k)$. Extracting a subsequence, if needed, we may also assume that $Q_k \rightarrow Q_\infty$. Set $M = Q_\infty^t J Q_\infty$, where J is the $n \times n$ diagonal matrix with diagonal entries $\{0, 0, \dots, 0, -1\}$. We then have

$$\begin{aligned} \text{trace}(A_k^t M A_k) &= \text{trace}(J Q_\infty Q_k^t D_k^2 Q_k Q_\infty^t) = - (Q_\infty Q_k^t D_k^2 Q_k Q_\infty^t)_{nn} \\ &= - \sum_{j=1}^n (Q_\infty Q_k^t)_{nj} [\lambda_j(A_k)]^2 (Q_k Q_\infty^t)_{jn} = - \sum_{j=1}^n (Q_\infty Q_k^t)_{nj}^2 [\lambda_j(A_k)]^2 \\ &\leq - (Q_\infty Q_k^t)_{nn}^2 [\lambda_n(A_k)]^2, \end{aligned}$$

which tends to $-\infty$ since $(Q_\infty Q_k^t)_{nn} \rightarrow 1$ and $\lambda_n(A_k) \rightarrow \infty$ as $k \rightarrow \infty$ contradicting the fact that $F(x, M)$ is finite. \square

We can now proceed with the proof of Theorem 1.1. First, we prove a continuity lemma, analogous to Lemma 3.2. Here, however, we have the extra restriction $A \leq \phi(\varepsilon)I$ and the continuity of F from only one side. This is because the cone $\Gamma_{\mathcal{A}}$ is, in principle, neither open nor closed, and when $D^2u(x) \in \partial\Gamma_{\mathcal{A}}$ we can only use perturbations of the form $D^2u(x) + \eta I$.

Lemma 4.2. *For every $M \in \Gamma_{\mathcal{A}}$, we have*

$$\inf_{\substack{A \in \mathcal{A} \\ A \leq \phi(\varepsilon)I}} \text{trace}(A^t (M + \eta I) A) \rightarrow \inf_{A \in \mathcal{A}} \text{trace}(A^t M A) \quad \text{as } \varepsilon, \eta \searrow 0. \quad (4.3)$$

Proof. Since $\text{trace}(A^t (M + \eta I) A) \geq \text{trace}(A^t M A)$, we have

$$\inf_{\substack{A \in \mathcal{A} \\ A \leq \phi(\varepsilon)I}} \text{trace}(A^t (M + \eta I) A) \geq \inf_{A \in \mathcal{A}} \text{trace}(A^t M A).$$

Let us fix $M \in \Gamma_{\mathcal{A}}$ and $\delta > 0$. We consider $A_0 \in \mathcal{A}$ such that

$$\inf_{A \in \mathcal{A}} \text{trace}(A^t M A) + \delta/2 \geq \text{trace}(A_0^t M A_0).$$

Since $\phi(\varepsilon) \rightarrow \infty$ as $\varepsilon \rightarrow 0$ there exists ε_0 such that $A_0 \leq \phi(\varepsilon)I$ for every $\varepsilon < \varepsilon_0$. Let $\eta_0 > 0$ be such that $\text{trace}(A_0^t A_0)\eta_0 < \delta/2$, we get

$$\begin{aligned} \inf_{\substack{A \in \mathcal{A} \\ A \leq \phi(\varepsilon)I}} \text{trace}(A^t (M + \eta I) A) &\leq \text{trace}(A_0^t (M + \eta I) A_0) \\ &= \text{trace}(A_0^t M A_0) + \text{trace}(A_0^t A_0)\eta \leq \inf_{A \in \mathcal{A}} \text{trace}(A^t M A) + \delta/2 + \delta/2 \end{aligned}$$

for every $\varepsilon < \varepsilon_0$ and $\eta < \eta_0$. We have proved (4.3). \square

Lemma 4.2 allows us to obtain an upper bound for

$$\inf_{\substack{A \in \mathcal{A} \\ A \leq \phi(\varepsilon)I}} \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x). \quad (4.4)$$

To conclude the lower bound we need a different argument and it is at this point that we use that F is continuous in $\Gamma_{\mathcal{A}}$. The continuity of F is a necessary condition in order to have a mean-value property, see Example 4.6 below.

Proof of Theorem 1.1. From (4.2) it follows that $A \leq \phi(\varepsilon)I$ and $|y| \leq \varepsilon$ imply $x + Ay \in B_\delta(x)$ for $\varepsilon < \varepsilon_0$, where $\varepsilon \phi(\varepsilon) < \delta$ for every $\varepsilon < \varepsilon_0$ and δ is as in (3.5). Therefore, by (3.5), if $\varepsilon < \varepsilon_0$, then

$$u(x + Ay) \leq P_\eta^+(x + Ay) \quad \text{for every } y \in B_\varepsilon(0), \quad (4.5)$$

for

$$P_\eta^+(z) = P(z) + \frac{\eta}{2}|z - x|^2$$

with P given by (3.4). Then, following the proof of Theorem 1.6, we have

$$\int_{B_\varepsilon(0)} u(x + Ay) dy - u(x) \leq \frac{\varepsilon^2}{2(n+2)} \text{trace}(A^t (D^2 u(x) + \eta I) A). \quad (4.6)$$

By (4.3) this gives us an upper bound of (4.4).

To obtain a lower bound we use the continuity of F . Given x and $\eta > 0$ there exists $\delta > 0$ such that

$$F(D^2 u(z)) \geq F(D^2 u(x)) - \eta$$

for every $z \in B_\delta(x)$. Then, there exists $\varepsilon_0 > 0$ such that for every $\varepsilon < \varepsilon_0$, $y \in B_\varepsilon(0)$ and $A \in \mathcal{A}$ with $A \leq \phi(\varepsilon)I$ we have $x + Ay \in B_\delta(x)$. Let us fix $A \in \mathcal{A}$. We have

$$\text{trace}(A^t D^2 u(x + Ay) A) \geq F(D^2 u(x + Ay)) \geq F(D^2 u(x)) - \eta.$$

We consider $v(y) = u(x + Ay)$, and we observe that

$$\Delta v(y) = \text{trace}(A^t D^2 u(x + Ay) A) \geq F(D^2 u(x)) - \eta.$$

Therefore, from the mean-value formula for the Laplacian, we get

$$\frac{\varepsilon^2}{2(n+2)} (F(D^2 u(x)) - \eta) \leq \int_{B_\varepsilon(0)} v(y) dy - v(0) = \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x).$$

With this lower bound, taking infimums and the limit as $\eta \rightarrow 0$, we have completed the proof. \square

Remark 4.3. Our assumptions imply that $F(D^2u(x)) > -\infty$ (since u is assumed to be \mathcal{A} -admissible). Observe that in the case that $F(D^2u(x)) = -\infty$ by the upper bound in (4.6) we also get

$$\inf_{A \in \mathcal{A}} \int_{B_\varepsilon(0)} u(x + Ay) dy - u(x) = -\infty.$$

Observe that condition (1.7) is necessary. In Example 4.6 this assumption is not satisfied and the mean-value formula fails.

Remark 4.4. In some important examples we have

$$F(M) = 0 \quad \text{for every } M \in \partial\Gamma_{\mathcal{A}} \quad (4.7)$$

and

$$\Gamma_{\mathcal{A}} = \left\{ M \in S^n(\mathbb{R}) : \text{trace}(A^t M A) \geq 0 \text{ for all } A \in \mathcal{A} \right\}.$$

This holds for the Monge-Ampère equation, see (1.5), and more generally for the k -Hessians, see Lemma 5.1 and Remark 5.2 below. Condition (4.7) was used in [13] in relation to existence and uniqueness of solutions to general fully nonlinear second-order PDEs. We observe that condition (4.7) implies the continuity of F in $\Gamma_{\mathcal{A}}$ and therefore Theorem 1.1 can be proved assuming (4.7) instead of (1.7). In fact, if there exists $C \in \mathbb{R}$ such that the cone $\Gamma_{\mathcal{A}} = \{M : F(M) > -\infty\}$ can be written as

$$\Gamma_{\mathcal{A}} = \{M : F(M) \geq C\}$$

and $F \equiv C$ in $\partial\Gamma_{\mathcal{A}}$, then F is continuous; that is, (1.7) holds. This can be easily deduced from the fact that F is lower semi-continuous on $\partial\Gamma_{\mathcal{A}}$ because F attains its minimum at every point of the boundary.

In the following example we show that condition (4.7) is not necessary by exhibiting an operator that does not satisfy (4.7) for which the mean-value property holds. Here, F is not constant on the boundary of the associated cone; nevertheless Theorem 1.1 applies since F is continuous.

Example 4.5. If we consider

$$\mathcal{A} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & n \end{bmatrix} : n = 0, 1, 2, \dots \right\},$$

then we have

$$F(D^2u) = \inf_{A \in \mathcal{A}} \text{trace}(A^t D^2u A) = \begin{cases} u_{x_1 x_1} & \text{if } u_{x_2 x_2} \geq 0 \\ -\infty & \text{if } u_{x_2 x_2} < 0. \end{cases}$$

The equation $F(D^2u) = f$ is equivalent to have $u_{x_1 x_1} = f$ and $u_{x_2 x_2} \geq 0$. It is not a nice equation, in the sense that it is overdetermined. Observe, however, that F is continuous where it is finite and therefore our result applies to this case and we have a mean value formula for this operator.

Associated with this F , we have the closed cone


$$\Gamma = \left\{ M : M_{22} \geq 0 \right\} = \left\{ M : F(M) > -\infty \right\}.$$

It is interesting that in this case F is not constant in the boundary of Γ . Observe that both

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

belong to Γ . Even more, they belong to the boundary of Γ , in fact

$$\begin{bmatrix} 0 & 0 \\ 0 & -\delta \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 0 \\ 0 & -\delta \end{bmatrix}$$

FIGURE 1. The operator F in Example 4.6

are not in Γ for every $\delta > 0$. Hence, F is not constant on the boundary of the cone since we have

$$F\left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}\right) = 0 \quad \text{and} \quad F\left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}\right) = 1.$$

The second example shows that continuity of F is necessary for the validity of a mean-value formula.

Example 4.6. We provide an example showing that the continuity of F is a necessary condition for Theorem 1.1. We consider the set $\mathcal{A} = \{A_\delta\}_{\delta>0}$ where

$$A_\delta = \begin{bmatrix} \sqrt{2}/\sqrt{\delta} & 0 \\ 0 & 1/\delta \end{bmatrix}$$

and define

$$F(D^2u) = \inf_{A \in \mathcal{A}} \text{trace}(A^t D^2u A) = \inf_{\delta>0} \left(\frac{2}{\delta} u_{x_1 x_1} + \frac{1}{\delta^2} u_{x_2 x_2} \right).$$

We have

$$F(D^2u) = \begin{cases} -\infty & \text{if } u_{x_2 x_2} < 0 \\ -\infty & \text{if } u_{x_2 x_2} = 0 \text{ and } u_{x_1 x_1} < 0 \\ 0 & \text{if } u_{x_2 x_2} \geq 0 \text{ and } u_{x_1 x_1} \geq 0 \\ -\frac{u_{x_1 x_1}}{u_{x_2 x_2}} & \text{if } u_{x_2 x_2} > 0 \text{ and } u_{x_1 x_1} < 0, \end{cases}$$

see Figure 1. Observe that F is not continuous at the origin, even if we restrict the domain to the set where it is finite.

Consider

$$u(x_1, x_2) = -|x_1|^{5/2} + x_1^2 x_2^2 + x_2^{10}.$$

Observe that $u \in C^2$ and \mathcal{A} -admissible. In fact we have

$$u_{x_2 x_2}(x_1, x_2) = 2x_1^2 + 90x_2^8 \geq 0,$$

with equality only at the origin, with $u_{x_1 x_1}(0, 0) = 0$ (notice that $D^2u(0, 0) = 0$). Therefore, we have that $F(D^2u(x_1, x_2)) > -\infty$ for every $(x_1, x_2) \in \mathbb{R}^2$.

Let us show that the mean-value formula in Theorem 1.1 is not satisfied at $(0, 0)$, i.e.,

$$\inf_{\delta>0} \int_{B_\varepsilon(0)} u(A_\delta y) dy - u(0, 0) \neq \frac{\varepsilon^2}{2(n+2)} F(D^2u(0, 0)) + o(\varepsilon^2).$$

Since $D^2u(0,0) = 0$, we have $F(D^2u(0,0)) = 0$ and we only need to prove that

$$\inf_{\delta > 0} \int_{B_\varepsilon(0)} u(A_\delta y) dy \neq o(\varepsilon^2). \quad (4.8)$$

An explicit computation shows that

$$\int_{B_\varepsilon(0)} u(A_\delta y) dy = \int_{B_\varepsilon(0)} \left(\frac{-2^{5/4}|y_1|^{5/2}}{\delta^{5/4}} + \frac{2y_1^2 y_2^2}{\delta^3} + \frac{y_2^{10}}{\delta^{10}} \right) dy = -C_1 \frac{\varepsilon^{5/2}}{\delta^{5/4}} + C_2 \frac{\varepsilon^4}{\delta^3} + C_3 \frac{\varepsilon^{10}}{\delta^{10}}.$$

For $\delta = \varepsilon^{1/2}$ we have

$$\frac{\varepsilon^{5/2}}{\delta^{5/4}} = \varepsilon^{15/8}, \quad \frac{\varepsilon^4}{\delta^3} = \varepsilon^{5/2} \quad \text{and} \quad \frac{\varepsilon^{10}}{\delta^{10}} = \varepsilon^5$$

and we obtain

$$\inf_{\delta > 0} \int_{B_\varepsilon(0)} u(A_\delta y) dy \leq -C\varepsilon^{15/8},$$

which proves (4.8). Therefore the mean-value formula in Theorem 1.1 does not hold in this case.

5. k -HESSIAN OPERATORS

A relevant example where Theorem 1.1 applies are the k -Hessian operators. For $k = 2, \dots, n$ the k -Hessian operators are given by elementary symmetric polynomials in the eigenvalues of the Hessian, i.e.,

$$\sigma_k(\lambda_1, \dots, \lambda_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_k}.$$

We write the operators in the form

$$F_k(D^2u(x)) = k \sigma_k(\lambda(D^2u(x)))^{\frac{1}{k}}, \quad (5.1)$$

where $\lambda(D^2u(x)) = (\lambda_1(D^2u(x)), \dots, \lambda_n(D^2u(x)))$, see Lemma 1.4 below.

We recall some definitions and properties of elementary symmetric polynomials, see for example [28]. We define the cone

$$\Gamma_k = \left\{ \lambda \in \mathbb{R}^n : \sigma_j(\lambda) > 0 \text{ for all } j = 1, \dots, k \right\}.$$

With a slight abuse of notation we write $M \in \Gamma_k$ to denote that $\lambda(M) \in \Gamma_k$. We have

$$\bar{\Gamma}_k = \left\{ \lambda \in \mathbb{R}^n : \sigma_j(\lambda) \geq 0 \text{ for all } j = 1, \dots, k \right\}$$

and $\bar{\Gamma}_k^\circ = \Gamma_k$. Let us define

$$\sigma_{k-1,i}(\gamma_1, \dots, \gamma_n) = \sigma_{k-1}(\gamma_1, \dots, \gamma_{i-1}, 0, \gamma_{i+1}, \dots, \gamma_n)$$

and

$$\mathcal{A}_k = \left\{ A : \lambda_i^2(A) = \sigma_{k-1,i}(\gamma) \text{ with } \gamma = (\gamma_1, \dots, \gamma_n) \in \Gamma_k \text{ and } \sigma_k(\gamma) = 1 \right\}.$$

Then, we have $\mathcal{A}_k \subset S_+^n(\mathbb{R})$ since

$$\sigma_{k-1,i}(\gamma) > 0, \quad \forall \gamma \in \Gamma_k,$$

see [28]. Also observe that by the continuity of $\sigma_{k-1,i}(\gamma)$ we get $\sigma_{k-1,i}(\gamma) \geq 0$ for every $\gamma \in \bar{\Gamma}_k$.

Our goal is find the k -Hessian counterpart of formula (1.5), which holds for the Monge-Ampère operator ($k = n$). We do that in two steps, first show the result for numbers, then for matrices.

Lemma 5.1. For every $\mu = (\mu_1, \dots, \mu_n) \in \mathbb{R}^n$ we have

$$\inf_{\substack{\gamma \in \Gamma_k \\ \sigma_k(\gamma)=1}} \sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma) = \begin{cases} k \sigma_k(\mu)^{\frac{1}{k}} & \text{if } \mu \in \bar{\Gamma}_k, \\ -\infty & \text{otherwise.} \end{cases}$$

Remark 5.2. Note that

$$\inf_{\substack{\gamma \in \Gamma_k \\ \sigma_k(\gamma)=1}} \sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma) = 0 \quad \text{if } \mu \in \partial\Gamma_k,$$

since

$$\partial\Gamma_k = \left\{ M \in S^n(\mathbb{R}) : \sigma_k(\lambda(M)) = 0 \text{ and } \sigma_j(\lambda(M)) \geq 0 \text{ for all } j = 1, \dots, k-1 \right\}. \quad (5.2)$$

Proof of Lemma 5.1. From formula (1.3) in [11] (see also (xi) in [28]), we have that for all $\mu, \gamma \in \Gamma_k$,

$$k \sigma_k(\mu)^{\frac{1}{k}} \sigma_k(\gamma)^{\frac{k-1}{k}} \leq \sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma),$$

with an equality for $\gamma^* = \sigma_k(\mu)^{-\frac{1}{k}} \mu$, i.e.,

$$\sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma^*) = \sum_{i=1}^n \mu_i \frac{\sigma_{k-1,i}(\mu)}{\sigma_k(\mu)^{\frac{k-1}{k}}} = \frac{k \sigma_k(\mu)}{\sigma_k(\mu)^{\frac{k-1}{k}}} = k \sigma_k(\mu)^{\frac{1}{k}}.$$

For $\mu \in \partial\Gamma_k$, observe that $(\mu_1 + \varepsilon, \dots, \mu_n + \varepsilon) \in \Gamma_k$ for every $\varepsilon > 0$. We consider

$$\gamma_i^\varepsilon = (\mu_i + \varepsilon) \sigma_k(\mu_1 + \varepsilon, \dots, \mu_n + \varepsilon)^{-\frac{1}{k}}$$

and we have

$$k \sigma_k(\mu_1 + \varepsilon, \dots, \mu_n + \varepsilon)^{\frac{1}{k}} = \sum_{i=1}^n (\mu_i + \varepsilon) \sigma_{k-1,i}(\gamma^\varepsilon) \geq \sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma^\varepsilon),$$

since $\sigma_{k-1,i}(\gamma^\varepsilon) \geq 0$. To conclude observe that

$$k \sigma_k(\mu_1 + \varepsilon, \dots, \mu_n + \varepsilon)^{\frac{1}{k}} \rightarrow k \sigma_k(\mu)^{\frac{1}{k}}$$

as $\varepsilon \rightarrow 0$.

The only remaining case is when $\mu \notin \bar{\Gamma}_k$. Notice that if $a > 0$ is large enough such that $\mu_i + a > 0$ for every $i \in \{1, \dots, n\}$, we get $(\mu_1 + a, \dots, \mu_n + a) \in \Gamma_k$. Thus, there exists $a > 0$ such that $(\mu_1 + a, \dots, \mu_n + a) \in \partial\Gamma_k$. Observe that $\sigma_k(\mu_1 + a, \dots, \mu_n + a) = 0$ by (5.2). Let us fix one such value a and consider

$$\gamma_i^\varepsilon = (\mu_i + a + \varepsilon) \sigma_k(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon)^{-\frac{1}{k}}.$$

As before, we have

$$\sum_{i=1}^n (\mu_i + a + \varepsilon) \sigma_{k-1,i}(\gamma^\varepsilon) = k \sigma_k(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon)^{\frac{1}{k}}$$

and therefore,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma^\varepsilon) &= - \lim_{\varepsilon \rightarrow 0} (a + \varepsilon) \sum_{i=1}^n \sigma_{k-1,i}(\gamma^\varepsilon) + k \lim_{\varepsilon \rightarrow 0} \sigma_k(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon)^{\frac{1}{k}} \\ &= -a(n-k+1) \lim_{\varepsilon \rightarrow 0} \sigma_{k-1}(\gamma^\varepsilon) = -a(n-k+1) \lim_{\varepsilon \rightarrow 0} \left(\frac{\sigma_{k-1}(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon)}{\sigma_k(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon)^{\frac{k-1}{k}}} \right). \end{aligned}$$

Recall that

$$\lim_{\varepsilon \rightarrow 0} \sigma_k(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon) = \sigma_k(\mu_1 + a, \dots, \mu_n + a) = 0,$$

and if we have

$$\lim_{\varepsilon \rightarrow 0} \sigma_{k-1}(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon) = \sigma_{k-1}(\mu_1 + a, \dots, \mu_n + a) > 0$$

we can conclude that

$$\lim_{\varepsilon \rightarrow 0} \sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma^\varepsilon) = -\infty$$

as desired. If this is not the case we have to look at the rate of convergence in more detail.

We consider two cases. First we assume that $\mu_i + a \neq 0$ for some $i \in \{1, \dots, n\}$. In this case we have that $\sigma_j(\mu_1 + a, \dots, \mu_n + a) > 0$ for some $j \in \{1, \dots, k-1\}$ because otherwise

$$\sum_{i=1}^n (\mu_n + a)^2 = \sigma_1^2(\mu_1 + a, \dots, \mu_n + a) - 2\sigma_2(\mu_1 + a, \dots, \mu_n + a) = 0,$$

a contradiction. Let $l \in \{1, \dots, k-1\}$ be the largest integer such that $\sigma_l(\mu_1 + a, \dots, \mu_n + a) > 0$.

We have

$$\begin{aligned} \sigma_k(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon) &= \sum_{i=0}^k \binom{n-i}{n-k} \sigma_i(\mu_1 + a, \dots, \mu_n + a) \varepsilon^{k-i}, \\ &= \varepsilon^{k-l} \left[\binom{n}{n-k} \varepsilon^l + \binom{n-1}{n-k} \sigma_1(\mu_1 + a, \dots, \mu_n + a) \varepsilon^{l-1} + \dots \right. \\ &\quad \left. \dots + \binom{n-l}{n-k} \sigma_l(\mu_1 + a, \dots, \mu_n + a) \right] \\ &= \varepsilon^{k-l} \left[\binom{n-l}{n-k} \sigma_l(\mu_1 + a, \dots, \mu_n + a) + O(\varepsilon) \right] \end{aligned}$$

as $\varepsilon \rightarrow 0$ (in the first equality we have defined $\sigma_0(\mu_1 + a, \dots, \mu_n + a) = 1$). Similarly,

$$\sigma_{k-1}(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon) = \varepsilon^{k-1-l} \left[\binom{n-l}{n-k+1} \sigma_l(\mu_1 + a, \dots, \mu_n + a) + O(\varepsilon) \right]$$

as $\varepsilon \rightarrow 0$. Therefore,

$$\frac{\sigma_{k-1}(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon)}{\sigma_k(\mu_1 + a + \varepsilon, \dots, \mu_n + a + \varepsilon)^{\frac{k-1}{k}}} = \left(\frac{\binom{n-l}{n-k+1} \sigma_l(\mu_1 + a, \dots, \mu_n + a) + O(\varepsilon)}{\left[\binom{n-l}{n-k} \sigma_l(\mu_1 + a, \dots, \mu_n + a) + O(\varepsilon) \right]^{\frac{k-1}{k}}} \right) \varepsilon^{-\frac{l}{k}}$$

as $\varepsilon \rightarrow 0$ and we conclude that

$$\lim_{\varepsilon \rightarrow 0} \sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma^\varepsilon) = -\infty.$$

Finally, in the case that $\mu_i = -a$ for every $i \in \{1, \dots, n\}$ we consider $\gamma_b = (1, \dots, 1, b, 1/b, 0, \dots, 0)$ where $b > 0$ and $k-2$ coordinates are equal to 1. We have $\gamma_b \in \Gamma_k$, $\sigma_k(\gamma_b) = 1$ and

$$\sum_{i=1}^n \mu_i \sigma_{k-1,i}(\gamma_b) = -a \sum_{i=1}^n \sigma_{k-1,i}(\gamma_b) = -a(n-k+1) \sigma_{k-1}(\gamma_b) = -a(n-k+1) \left(k-2 + b + \frac{1}{b} \right)$$

which goes to $-\infty$ as $b \rightarrow \infty$. □

We are now ready to show Lemma 1.4, the matrix counterpart of Lemma 5.1.

Proof of Lemma 1.4. Given $M \in S^n(\mathbb{R})$ we consider $A \in \mathcal{A}_k$ such that both matrices are diagonal in the same basis. We have

$$\text{trace}(A^t M A) = \text{trace}(A A^t M) = \sum_{i=1}^n \lambda_i(A A^t) \lambda_i(M) = \sum_{i=1}^n \lambda_i^2(A) \lambda_i(M).$$

Then, by the definition of \mathcal{A}_k and Lemma 5.1 we get that

$$\inf_{A \in \mathcal{A}_k} \text{trace}(A^t M A) \leq \inf_{\substack{\gamma \in \Gamma_k \\ \sigma_k(\gamma)=1}} \sum_{i=1}^n \lambda_i(M) \sigma_{k-1,i}(\gamma) = \begin{cases} k \sigma_k(\lambda(M))^{\frac{1}{k}} & \text{if } M \in \bar{\Gamma}_k, \\ -\infty & \text{otherwise.} \end{cases}$$

Therefore, it only remains to prove that

$$\text{trace}(A^t M A) \geq k \sigma_k(\lambda(M))^{\frac{1}{k}} \quad (5.3)$$

for every $A \in \mathcal{A}_k$ and $M \in \bar{\Gamma}_k$. To that end we recall the following inequality by Marcus (see [20])

$$\text{trace}(X M) \geq \min_p \sum_{i=1}^n \lambda_i(M) \lambda_{p(i)}(X),$$

where p ranges over the permutations of the numbers $\{1, \dots, n\}$. Recalling that $A \in \mathcal{A}_k$ we obtain

$$\text{trace}(A^t M A) \geq \min_p \sum_{i=1}^n \lambda_i(M) \sigma_{k-1,p(i)}(\gamma)$$

for some $\gamma \in \Gamma_k$ such that $\sigma_k(\gamma) = 1$. Let \tilde{p} be the permutation where the minimum is attained and observe that

$$\sigma_{k-1,\tilde{p}(i)}(\gamma) = \sigma_{k-1,i}(\tilde{\gamma})$$

where $\tilde{\gamma}$ is such that $\tilde{\gamma}_i = \gamma_{\tilde{p}(i)}$. We have

$$\text{trace}(A^t M A) \geq \sum_{i=1}^n \lambda_i(M) \sigma_{k-1,i}(\tilde{\gamma}).$$

Observe that $\tilde{\gamma} \in \Gamma_k$ and $\sigma_k(\tilde{\gamma}) = 1$, hence, by Lemma 5.1 we have

$$\sum_{i=1}^n \lambda_i(M) \sigma_{k-1,i}(\tilde{\gamma}) \geq k \sigma_k(\lambda(M))^{\frac{1}{k}}$$

and (5.3) follows. \square

Recall that a function $u \in C^2(\Omega)$ is called k -convex whenever $D^2 u(x) \in \bar{\Gamma}_k$ for every $x \in \Omega$, see [25, 26, 27]. As a consequence of Lemma 1.4 we have the following result.

Corollary 5.3. *A function $u \in C^2(\Omega)$ is k -convex if and only if it is \mathcal{A}_k -admissible. In other words, we have*

$$\Gamma_{\mathcal{A}_k} = \bar{\Gamma}_k.$$

As a consequence of the Lemma 1.4 we can obtain Theorem 1.5 from Theorem 1.1.

Proof of Theorem 1.5. From Corollary 5.3, we get $\Gamma_{\mathcal{A}_k} = \bar{\Gamma}_k$, i.e., it is equivalent to be k -convex and \mathcal{A}_k -admissible. Also observe that F_k is continuous in \mathcal{A}_k . Therefore we are under the hypothesis of Theorem 1.1 and the result follows. \square

6. EXAMPLES IN THE HEISENBERG GROUP

Let $\mathbb{H} = (\mathbb{R}^3, *)$ be the Heisenberg group. We write a point $q \in \mathbb{H}$ as $q = (x, y, z)$. The point $\bar{q} = (x, y)$ is the horizontal projection of q . When convenient, we will also use the notation $q = (q_1, q_2, q_3) = (\bar{q}, q_3)$. The group operation is

$$q * q' = (x, y, z) * (x', y', z') = \left(x + x', y + y', z + z' + \frac{1}{2}(xy' - yx') \right).$$

The Korányi gauge is given by

$$|q|_K = ((x^2 + y^2)^2 + 16z^2)^{1/4}.$$

It induces a left-invariant metric $d(q, q') = |q^{-1} * q'|_K$ in \mathbb{H} . We also have a family of anisotropic dilations:

$$\rho_\lambda(x, y, z) = (\lambda x, \lambda y, \lambda^2 z), \quad \lambda > 0$$

that are group homomorphisms. The Korányi gauge and the Korányi metric are homogeneous with respect to the dilations

$$|\rho_\lambda(q)|_K = \lambda |q|_K, \quad d(\rho_\lambda(q), \rho_\lambda(q')) = \lambda d(q, q').$$

The open ball centered at q with radius $r > 0$ is a translation and dilation of the open ball centered at 0 of radius 1

$$B_r(q) = \{q' \in \mathbb{H}; d(q, q') < r\} = q * B_r(0) = q * \rho_r(B_1(0)).$$

Euclidean balls in 2 dimensions will be denoted by B_r^2 .

The vector fields

$$X = \partial_x - \frac{y}{2}\partial_z, \quad Y = \partial_y + \frac{x}{2}\partial_z, \quad Z = \partial_z$$

are left-invariant and form a basis for the Lie algebra of \mathbb{H} . The only non-trivial commuting relation is $Z = [X, Y]$. The horizontal tangent space at the point q is the plane generated by $X(q)$ and $Y(q)$

$$T_q = \text{span} \left\{ \left(1, 0, -\frac{y}{2} \right), \left(0, 1, \frac{x}{2} \right) \right\} = \left(\frac{y}{2}, -\frac{x}{2}, 1 \right)^\perp. \quad (6.1)$$

The horizontal gradient and the sub-Laplacian of a function $v : \mathbb{H} \rightarrow \mathbb{R}$ are respectively the vector field and function

$$\nabla_{\mathbb{H}} v = (Xv)X + (Yv)Y, \quad \Delta_{\mathbb{H}} v = (X^2 + Y^2)v.$$

The horizontal second derivatives are given by the non necessarily symmetric 2×2 matrix

$$\nabla_{\mathbb{H}}^2 v(q) = \begin{pmatrix} X^2 v(q) & XY v(q) \\ YX v(q) & Y^2 v(q) \end{pmatrix}.$$

The symmetrized horizontal second derivatives are $(\nabla_{\mathbb{H}}^2 v(q))^* = \frac{1}{2}(\nabla_{\mathbb{H}}^2 v(q) + (\nabla_{\mathbb{H}}^2 v(q))^t)$. Our starting point is the Taylor expansion for a function $v \in \mathcal{C}^2(\mathbb{H})$ at a point $q = (x, y, z) \in \mathbb{H}$ adapted to the Heisenberg group, which we take from Section 3 in [17],

$$v(p) = v(q) + \langle (\nabla_{\mathbb{H}}, Z)v(q), q^{-1} * p \rangle + \frac{1}{2} \langle (\nabla_{\mathbb{H}}^2 v(q))^* \overline{q^{-1} * p}, \overline{q^{-1} * p} \rangle + o(|q^{-1} * p|_K^2). \quad (6.2)$$

Next, we consider the horizontal average operator

$$\mathcal{A}_2(v, \varepsilon)(q) = \int_{B_{\varepsilon}^2(0)} v(q * (a, b, 0)) da db = \int_{B_1^2(0)} v\left(q + \varepsilon\left(a, b, \frac{1}{2}(xb - ya)\right)\right) da db.$$

From Proposition 2.3 in [17] we get

Lemma 6.1.

$$\mathcal{A}_2(v, \varepsilon)(q) - v(q) = \frac{\varepsilon^2}{8} \Delta_{\mathbb{H}} v(q) + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0.$$

By changing variables we obtain averages over general horizontal planes.

Lemma 6.2. *Let A be a 2×2 matrix and let $\varepsilon > 0$. We have*

$$\int_{B_1^2(0)} v(q * \varepsilon(A \cdot (a, b), 0)) da db - v(q) = \frac{\varepsilon^2}{8} \text{trace}(A^t (\nabla_{\mathbb{H}}^2 v(q))^* A) + o(\varepsilon^2 |A|^2)$$

as $\varepsilon \rightarrow 0$.

Proof. Set $p = q * \varepsilon(A \cdot (a, b), 0)$, where $(a, b) \in B_1^2(0)$. Observe that $q^{-1} * p = \varepsilon(A \cdot (a, b), 0)$ so that $q^{-1} * p = \varepsilon A \cdot (a, b)$. Writing the Taylor expansion (6.2) we get

$$\begin{aligned} v(p) &= v(q) + \langle (\nabla_{\mathbb{H}}, Z)v(q), (\varepsilon A \cdot (a, b), 0) \rangle \\ &\quad + \frac{\varepsilon^2}{2} \langle (\nabla_{\mathbb{H}}^2 v(q))^* \cdot A \cdot (a, b), A \cdot (a, b) \rangle + o(|\varepsilon A \cdot (a, b), 0|_K^2) \\ &= v(q) + \varepsilon \langle \nabla_{\mathbb{H}} v(q), A \cdot (a, b) \rangle + \frac{\varepsilon^2}{2} \langle (A^t \cdot \nabla_{\mathbb{H}}^2 v(q))^* \cdot A \cdot (a, b), (a, b) \rangle + o(|\varepsilon A \cdot (a, b)|^2). \quad \square \end{aligned}$$

We now state a version of Theorem 1.1 in the Heisenberg group for the expression

$$F(D^2 v(q)) = 2 (\det(\nabla_{\mathbb{H}}^2 v(q))^*)^{1/2}.$$

Recall that we have

$$\inf_{A \in \mathcal{A}} \text{trace}(A^t M A) = \begin{cases} 2 (\det M)^{1/2} & \text{if } M \geq 0 \\ -\infty & \text{otherwise,} \end{cases}$$

for the unbounded set $\mathcal{A} = \{A \in S_+^2(\mathbb{R}) : \det(A) = 1\}$, for example see [4]. We say that a function $v \in C^2$ is horizontally convex if $(\nabla_{\mathbb{H}}^2 v(q))^* \geq 0$.

Theorem 6.3. *Let $v \in C^2$ be horizontally convex and $\phi(\varepsilon)$ a positive function satisfying (1.8). Then, we have*

$$\inf_{\substack{\det(A)=1 \\ A \leq \phi(\varepsilon)I}} \int_{B_1^2(0)} v(q * (A(a, b), 0)) da db - v(q) = \frac{\varepsilon^2}{4} (\det(\nabla_{\mathbb{H}}^2 v(q))^*)^{1/2} + o(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0.$$

The proof of this theorem is a straightforward adaptation of the proof of Theorem 1.1 in [4]. Note that the two-dimensional averages are taken over ellipsoids in the horizontal tangent plane T_q defined in (6.1).

REFERENCES

- [1] L. Ambrosio, H. M. Soner, *Level set approach to mean curvature flow in arbitrary codimension*, J. Differential Geom. 43 (1996), 693–737.
- [2] A. R. Arroyo García; *Nonlinear Mean Value Properties related to the p -Laplacian*, Ph.D. thesis, Universitat Autònoma de Barcelona, (2017).
- [3] I. Birindelli, G. Galise, and H. Ishii; *Existence through convexity for the truncated Laplacians*, Math. Ann., 379, (2021), 909–950.
- [4] P. Blanc, F. Charro, J. D. Rossi, J. J. Manfredi; *A nonlinear Mean Value Property for the Monge-Ampère operator*, J. Convex Analysis, 28(2), (2021), 353–386.
- [5] P. Blanc and J. D. Rossi; *Games for eigenvalues of the Hessian and concave/convex envelopes*, Jour. Math. Pures Appl., 127, (2019), 192–215.
- [6] P. Blanc and J. D. Rossi; *An asymptotic mean-value formula for eigenvalues of the Hessian related to concave/convex envelopes*. Vietnam Jour. Math., 48(2), (2020), 335–344.
- [7] W. Blaschke; *Ein Mittelwertsatz und eine kennzeichnende Eigenschaft des logarithmischen Potentials*, Ber. Verh. Sächs. Akad. Wiss., Leipziger, 68, (1916), 3–7.
- [8] X. Cabré and L. Caffarelli; *Fully Nonlinear Elliptic Equations*, Amer. Math. Soc., Colloquium publications, Vol. 43 (1995).

- [9] X. Cabré and L. Caffarelli; *Interior $C^{2,\alpha}$ regularity theory for a class of nonconvex fully nonlinear elliptic equations*, Jour. Math. Pures Appl., 82(5), (2003), 573–612.
- [10] L. Caffarelli, Y. Y. Li, L. Nirenberg, *Some remarks on singular solutions of nonlinear elliptic equations. I*, J. Fixed Point Theory Appl., 5, (2009), 353–395.
- [11] L.A. Caffarelli, L. Nirenberg, and J. Spruck; *The Dirichlet problem for nonlinear second-order elliptic equations. III. Functions of the eigenvalues of the Hessian*, Acta Math., 155(3-4), (1985), 261–301.
- [12] M.G. Crandall, H. Ishii and P.L. Lions. *User’s guide to viscosity solutions of second order partial differential equations*. Bull. Amer. Math. Soc., 27, (1992), 1–67.
- [13] F. R. Harvey, H. B. Jr. Lawson, *Dirichlet duality and the nonlinear Dirichlet problem*, Comm. Pure Appl. Math., 62, (2009), 396–443.
- [14] F. R. Harvey, H. B. Jr. Lawson, *p -convexity, p -plurisubharmonicity and the Levi problem*, Indiana Univ. Math. J., 62, (2013), 149–169.
- [15] Ü. Kuran; *On the mean-value property of harmonic functions*, Bull. London Math. Soc., 4, (1972), 311–312.
- [16] M. Lewicka. *A Course on Tug-of-War Games with Random Noise*. Universitext. Springer, (2020).
- [17] M. Lewicka, J. J. Manfredi, and D. Ricciotti, *Random walks and random tug of war in the Heisenberg group*, Math. Annalen, 377(1-2), (2020), 797–846.
- [18] P. Lindqvist, and J. Manfredi; *On the mean-value property for the p -Laplace equation in the plane*, Proc. Amer. Math. Soc., 144(1), (2016), 143–149.
- [19] J. J. Manfredi, M. Parviainen, and J. D. Rossi, *An asymptotic mean-value characterization of p -harmonic functions*, Proc. Amer. Math. Soc., 138, (2010), 881–889.
- [20] M. Marcus, *An Eigenvalue for the Product of Normal Matrices*, The American Mathematical Monthly, 63(3), (1956), 173–174.
- [21] A. M. Oberman, *Finite difference methods for the Infinity Laplace and p -Laplace equations*, Journal of Computational and Applied Mathematics, Volume 254, (2013), pp. 65–80.
- [22] A. M. Oberman, and L. Silvestre, *The Dirichlet problem for the convex envelope*, Trans. Amer. Math. Soc., 363, (2011), 5871–5886.
- [23] I. Privaloff; *Sur les fonctions harmoniques*, Mat. Sb., 32(3), (1925), 464–471.
- [24] J. P. Sha, *p -convex Riemannian manifolds*, Invent. Math., 83(3), (1986), 437–447.
- [25] N.S. Trudinger, X.J. Wang; *Hessian measures I*, Topol. Meth. Nonlinear Anal., 10, (1997), 225–239.
- [26] N.S. Trudinger, X.J. Wang; *Hessian measures II*, Ann. Math., 150, (1999), 579–604.
- [27] N.S. Trudinger, X.J. Wang; *Hessian measures III*, J. Functional Analysis, 193, (2002), 1–23.
- [28] X. Wang, *The k -Hessian equation*, Geometric analysis and PDEs. Springer, Berlin, Heidelberg. (2009), 177–252.
- [29] H. Wu, *Manifolds of partially positive curvature*, Indiana Univ. Math. J., 36, (1987), 525–548.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ, PO BOX 35, FI-40014, JYVÄSKYLÄ, FINLAND

Email address: pblanc@dm.uba.ar

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, 656 W. KIRBY, DETROIT, MI 48202, USA

Email address: fcharro@wayne.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITTSBURGH, PITTSBURGH, PA 15260, USA.

Email address: manfredi@pitt.edu

DEPARTAMENTO DE MATEMÁTICA, FCEyN, UNIVERSIDAD DE BUENOS AIRES, PABELLÓN I, CIUDAD UNIVERSITARIA (1428), BUENOS AIRES, ARGENTINA.

Email address: jrossi@dm.uba.ar