

A NOTE ON SYZYGIES AND NORMAL GENERATION FOR TRIGONAL CURVES

MICHAEL HOFF

ABSTRACT. Let C be a trigonal curve of genus $g \geq 5$ and let T be the unique trigonal line bundle inducing a map $\pi : C \xrightarrow{3:1} \mathbb{P}^1$. This note provides a short and easy proof of the normal generation for the residual line bundle $K_C \otimes T^{-1}$ for curves of genus $g \geq 7$. Moreover, we compute the minimal free resolution of the embedded curve $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-n})^*)$ for the residual line bundle $K_C \otimes T^{-n}$ for $n \geq 1$ and $g \geq 3n + 4$.

1. INTRODUCTION

In this short note we show how to determine the minimal free resolution of a trigonal curve lying on a rational normal surface scroll following [Sch86]. This was studied in a general context (arithmetic Cohen-Macaulay and non-arithmetic Cohen-Macaulay divisors on rational normal scrolls) by [Nag99] or [Par14], but we could not find any application for trigonal curves¹. In particular, for a trigonal curve C of genus $g \geq 5$ with trigonal bundle T the shape of the minimal free resolution implies normal generation for line bundles of the form $K_C \otimes T^{-n}$ (by our knowledge, the latter seems to be unknown for $n = 1$ and small genus or for $n \geq 2$). We present a short and selfcontained proof (see also Remark 1.3).

We summarize known result about normal generation of line bundles on trigonal curves. Let C be a curve of genus g . A line bundle L on C is said to be normally generated if L is very ample and the embedded curve $C \subset \mathbb{P}(H^0(C, L)^*)$ is projectively normal (equivalently, the natural maps $S^m H^0(C, L) \rightarrow H^0(C, L^m)$ are surjective for all $m \geq 0$). In their seminal paper [GL86], Green and Lazarsfeld prove that a very ample line bundle L of degree $2g - k$ on C is normally generated as long as k is smaller than the Clifford index c of C ². For our purpose the following result is important.

Theorem. [GL86, Theorem 2] *There exists an explicit constant $N(c)$ such that if C is a curve of Clifford index c and of genus $g > N(c)$, then every very ample line bundle L with $h^1(C, L) \geq 2$ of degree $\deg(L) = 2g - 2 \cdot h^1(C, L) - c$ is normally generated.*

They furthermore showed the existence of non-special line bundles which are not normally generated, and classified not normally generated line bundles with $h^1(C, L) = 1$.

Date: August 16, 2021.

2010 Mathematics Subject Classification. 14H45, 14H51, 13D02 .

Key words and phrases. trigonal curve, normal generation, special divisors.

¹A trigonal curve is a non-hyperelliptic smooth curve C of genus g with a line bundle T inducing a $3 : 1$ morphism to \mathbb{P}^1 . For $g \geq 5$, the line bundle T is unique.

²For $g \geq 4$ the Clifford index is defined as $\min\{\deg L - 2 \cdot h^0(C, L) - 2 : L \in \text{Pic}(C), h^i(C, L) \geq 2, i = 0, 1\}$.

Example 1.1. For a trigonal curve C (\Rightarrow Clifford index 1) let T be the trigonal bundle. We set $L = K_C \otimes T^{-1}$ in the above theorem. Then $K_C \otimes T^{-1}$ is normally generated if $g > N(1) = 16$. We could not find any further result in the literature concerning the normal generation of the line bundle $K_C \otimes T^{-1}$ for trigonal curves of genus $16 \geq g \geq 7$ (the low genus cases are trivial by a Bézout argument). We close the remaining gap in the genus with our theorem and furthermore describe the whole minimal free resolution of $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-1})^*)$.

Given a projective variety $Y \subset \mathbb{P}^n$, we can resolve the structure sheaf of Y by free $\mathcal{O}_{\mathbb{P}^n}$ -modules, that is, a minimal free resolution

$$\mathcal{O}_Y \leftarrow F_0 \leftarrow F_1 \leftarrow \cdots \leftarrow F_m \leftarrow 0,$$

where $F_i = \bigoplus_j \mathcal{O}_{\mathbb{P}^n}(-j)^{\beta_{ij}}$. We call the numbers β_{ij} the *Betti numbers* of Y . The shape of the minimal free resolution will be encoded in the so-called *Betti table* $(\beta_{ij})_{ij}$ (see [Eis95] for an introduction).

Our main theorem is the following (see also Theorem 3.4 for its generalization).

Theorem 1.2. *Let C be a trigonal curve of genus $g \geq 7$ and let T be the unique trigonal bundle inducing the map $\pi : C \xrightarrow{3:1} \mathbb{P}^1$. The Betti table of a minimal free resolution of $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-1})^*) = \mathbb{P}^{g-3}$ has the following shape*

	0	1	...	$g-6$	$g-5$	$g-4$
0	1	0	...	0	0	0
1	0	*	...	*	*	0
2	0	*	...	*	0	0
3	0	0	...	0	*	*

where $*$ indicates nonzero entries. In particular, the embedded curve is projectively normal.

Remark 1.3. Given a trigonal curve C of genus g and a positive number n , we provide a sharp range for g and n where $K_C \otimes T^{-n}$ is very ample in Section 3.1. Hence the embedded curve $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-n})^*)$ is smooth. One can show that projective normality of $K_C \otimes T^{-n}$ follows already from the classical Castelnuovo lemma [Cas89]. We note that trigonal curves $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-n})^*)$ are extremal in the sense of Castelnuovo's bound and refer to [ACGH85, Chapter III, §2, pp 120] for a detailed analysis. Furthermore the statement of Theorem 1.2 also follows from [Nag99, Theorem 2.4 and Corollary 2.5] using the smoothness assumption. We recall that we present a new selfcontained proof where our method of the proof is exactly valid in the range for g and n where $K_C \otimes T^{-n}$ is very ample.

In [MS86], Martens and Schreyer classified non-special not normally generated line bundles on trigonal curves. They used the extrinsic geometry of the embedding $C \subset \mathbb{P}(H^0(C, L)^*)$ depending on the Maroni-invariant (see Section 2.1 for the definition).

Remark 1.4. In [LN13], Lange and Newstead determined all vector bundles on trigonal curves which compute all higher Clifford indices (see for the definition e.g., [LN10]). Their main theorem [LN13, Theorem 4.7] relies on the normal generation of $K_C \otimes T^{-1}$ and is thus true for all trigonal curves of genus $g \geq 5$ by our main theorem.

Remark 1.5. The nonzero Betti numbers in Theorem 1.2 are:

$$\begin{aligned} \beta_{j,j+1} &= j \cdot \binom{g-4}{j+1}, & \text{for } 1 \leq j \leq g-5 \text{ (first row)} \\ \beta_{j,j+2} &= (g-5-j) \cdot \binom{g-4}{j-1}, & \text{for } 1 \leq j \leq g-6 \text{ (second row)} \\ \beta_{j,j+3} &= (j-g+6) \cdot \binom{g-4}{j}, & \text{for } g-5 \leq j \leq g-4 \text{ (third row)} \end{aligned}$$

These are exactly the numbers as in [Nag99, Theorem 2.4]

In Section 2 we present the proof of Theorem 1.2 and introduce the necessary background. In Section 3 we give a bound for the very ampleness of $K_C \otimes T^{-n}$ in terms of n and the genus of the trigonal curve C . At the end we proof the generalization of Theorem 1.2.

2. PROOF OF THE MAIN THEOREM

Our proof follows the strategy of [Sch86]:

- $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-1})^*)$ lies on a rational normal surface X swept out by T ,
- resolve \mathcal{O}_C as an \mathcal{O}_X -module by direct sums of line bundles on X and take the minimal free resolution of each of these line bundles as $\mathcal{O}_{\mathbb{P}^{g-3}}$ -modules,
- a mapping cone construction induces a minimal free resolution of \mathcal{O}_C as an $\mathcal{O}_{\mathbb{P}^{g-3}}$ -module.

2.1. A rational normal surface associated to the trigonal bundle. Let C be a trigonal curve of genus $g \geq 5$. An important invariant associated to any trigonal curve - classically used to describe its Brill–Noether locus - is the so-called *Maroni-invariant*. It is defined as follows. Let $\pi : C \rightarrow \mathbb{P}^1$ be the map associated to the trigonal bundle T . The direct image $\pi_* K_C$ of the canonical bundle on C splits into a direct sum of line bundles

$$\pi_* K_C = \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(m) \oplus \mathcal{O}_{\mathbb{P}^1}(-2)$$

where m is the *Maroni-invariant*. It is known that

$$(2.1) \quad 0 < \frac{g-4}{3} \leq m \leq \frac{g-2}{2}.$$

and $a = g - 2 - m$. Furthermore, $m+2$ and $a+2$ describe the two jumps of the function $f : \mathbb{N} \rightarrow \{0, 1, 2\}$ with $f(n) := h^1(C, T^{n-1}) - h^1(C, T^n)$. Hence,

$$(2.2) \quad m+2 = \min\{n \in \mathbb{N} : h^0(C, T^n) > n+1\}.$$

Maroni [Mar46] introduced m as a geometrical invariant determining a smooth rational normal scroll which is swept out by the trigonal bundle in the canonical space $\mathbb{P}(H^0(C, K_C)^*)$.

We will explain this procedure for our purpose in the following setting. We assume that $g \geq 6$. Let $C \subset \mathbb{P}^{g-3} := \mathbb{P}(H^0(C, K_C \otimes T^{-1})^*)$ be embedded by the very ample linear series $|K_C \otimes T^{-1}|$ (see Section 3.1 and note that for $g = 5$, $K_C \otimes T^{-1}$ maps C to

a plane quintic with one node). We consider the scroll swept out by the unique pencil $|T|$, that is,

$$X = \bigcup_{D \in |T|} \overline{D} \subset \mathbb{P}^{g-3}$$

where \overline{D} is the linear span of the divisor D defined by the linear forms

$$H^0(C, K_C \otimes T^{-1}(-D)) \rightarrow H^0(C, K_C \otimes T^{-1}).$$

Since $H^0(C, K_C \otimes T^{-2})$ is $(g-4)$ -dimensional (by (2.1) and (2.2) for $g \geq 6$), the linear span \overline{D} is a line and X is a rational normal surface of degree $g-4$. In particular, X is the image of a projective bundle $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(e_1) \oplus \mathcal{O}_{\mathbb{P}^1}(e_2))$ where $e_1 = g-3-m$ and $e_2 = m-1 \geq 0$. We will explain this fact in the next section.

2.2. Resolution of the curve on the rational normal surface. Let C be a trigonal curve with Maroni-invariant m . Let X be the rational normal surface of degree $g-4$ as above and let $\pi : \mathcal{E} = \mathcal{O}_{\mathbb{P}^1}(e_1) \oplus \mathcal{O}_{\mathbb{P}^1}(e_2) \rightarrow \mathbb{P}^1$ be the rank two bundle on \mathbb{P}^1 with $e_1 = g-3-m$ and $e_2 = m-1$. By [Har81], if $e_1, e_2 > 0$, then

$$j : \mathbb{P}(\mathcal{E}) \rightarrow X \subset \mathbb{P}H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)) = \mathbb{P}^{g-3}$$

is an isomorphism. Otherwise it is a resolution of singularities. Since $R^i j_* \mathcal{O}_{\mathbb{P}(\mathcal{E})} = 0$, it is convenient to consider $\mathbb{P}(\mathcal{E})$ instead of X for cohomological considerations. Note that this may happen for $g = 6, 7$ and $m = 1$.

It is furthermore known that the Picard group $\text{Pic}(\mathbb{P}(\mathcal{E}))$ is generated by the ruling $R = [\pi^* \mathcal{O}_{\mathbb{P}^1}(1)]$ and the hyperplane class $H = [j^* \mathcal{O}_{\mathbb{P}^{g-3}}(1)]$ with intersection products

$$H^2 = e_1 + e_2 = g-4, \quad H \cdot R = 1, \quad R^2 = 0.$$

Since $C \subset X$ and T is base point free, we consider $C \subset \mathbb{P}(\mathcal{E})$ as a subvariety of the projectivised bundle. Note that

$$\mathcal{O}_{\mathbb{P}(\mathcal{E})}(H) \otimes \mathcal{O}_C = K_C \otimes T^{-1} \text{ and } \mathcal{O}_{\mathbb{P}(\mathcal{E})}(R) \otimes \mathcal{O}_C = T.$$

Since the curve C is a codimension one subvariety of $\mathbb{P}(\mathcal{E})$, the ideal sheaf $\mathcal{I}_{C/\mathbb{P}(\mathcal{E})}$ is generated by an element in $H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(aH - bR))$. Since the ruling on C has degree 3 ($\Rightarrow a = 3$) and the degree of $C \subset \mathbb{P}^{g-3}$ is $\deg(C) = 2g-5$ ($\Rightarrow b = g-7$), the resolution of \mathcal{O}_C as an $\mathcal{O}_{\mathbb{P}(\mathcal{E})}$ -module is

$$(2.3) \quad 0 \longleftarrow \mathcal{O}_C \longleftarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})} \longleftarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})}(-3H + (g-7)R) \longleftarrow 0$$

In the next step we will resolve the line bundles in the above resolution in terms of $\mathcal{O}_{\mathbb{P}^{g-3}}$ -modules. Therefore we recall the definition of *Eagon-Northcott* type resolutions whereby we restrict to our case.

We have a natural multiplication map

$$H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(R)) \otimes H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(H-R)) \longrightarrow H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(H)),$$

and the equations of the rational normal surface X are given by the 2×2 -minors of the matrix Φ . We define

$$F := H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(H-R)) \otimes \mathcal{O}_{\mathbb{P}^{g-3}} = \mathcal{O}_{\mathbb{P}^{g-3}}^{g-4}, \quad G := H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathbb{P}(\mathcal{E})}(R)) \otimes \mathcal{O}_{\mathbb{P}^{g-3}} = \mathcal{O}_{\mathbb{P}^{g-3}}^2$$

and regard Φ as a map $\Phi : F(-1) \rightarrow G$ (where we identify $G = G^*$).

For $b \geq -1$ we can resolve $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(aH+bR)$ as an $\mathcal{O}_{\mathbb{P}^{g-3}}$ -module by an Eagon-Northcott type complex $\mathcal{C}^b \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(a)$ where the j^{th} term of \mathcal{C}_j^b is defined as

$$\mathcal{C}_j^b = \begin{cases} \bigwedge^j F \otimes S_{b-j} G \otimes \mathcal{O}_{\mathbb{P}^r}(-j) & \text{for } 0 \leq j \leq b \\ \bigwedge^{j+1} F \otimes D_{j-b-1} G^* \otimes \mathcal{O}_{\mathbb{P}^r}(-j-1) & \text{for } j \geq b+1 \end{cases}$$

and whose differentials $\delta_j : \mathcal{C}_j^b \rightarrow \mathcal{C}_{j-1}^b$ are given by the multiplication by

$$\begin{aligned} \Phi &\in H^0(\mathbb{P}(\mathcal{E}), F^* \otimes G) && \text{for } j \neq b+1 \\ \bigwedge^2 \Phi &\in H^0(\mathbb{P}(\mathcal{E}), \bigwedge^2 F^* \otimes \bigwedge^2 G) && \text{for } j = b+1 \end{aligned}$$

(see [BE75]). The resolution in (2.3) induces a double complex of Eagon–Northcott type resolutions

$$(2.4) \quad \mathcal{C}^0 \leftarrow \mathcal{C}^{g-7} \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-3).$$

2.3. Mapping cone construction and examples. Since all maps in the double complex in (2.4) are minimal, the mapping cone $[\mathcal{C}^0 \leftarrow \mathcal{C}^{g-7} \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-3)]$ of this double complex induces a minimal free resolution of $C \subset \mathbb{P}^{g-3}$ (see [Eis95]).

Example 2.1. For a trigonal curve of genus $g = 7$ the double complex is

$$\begin{array}{ccccccc} 0 & \longleftarrow & \mathcal{O}_C & \longleftarrow & \mathcal{O}_{\mathbb{P}(\mathcal{E})} & \longleftarrow & \mathcal{O}_{\mathbb{P}(\mathcal{E})}(-3H) & \longleftarrow & 0 \\ & & & & \uparrow & & \uparrow & & \\ & & & & \mathcal{O}_{\mathbb{P}^4} & \longleftarrow & \mathcal{O}_{\mathbb{P}^4}(-3) & & \\ & & & & \uparrow & & \uparrow & & \\ & & & & \bigwedge^2 F \otimes \mathcal{O}_{\mathbb{P}^4}(-2) & \longleftarrow & \bigwedge^2 F \otimes \mathcal{O}_{\mathbb{P}^4}(-5) & & \\ & & & & \uparrow & & \uparrow & & \\ & & & & \bigwedge^3 F \otimes DG^* \otimes \mathcal{O}_{\mathbb{P}^4}(-3) & \longleftarrow & \bigwedge^3 F \otimes DG^* \otimes \mathcal{O}_{\mathbb{P}^4}(-6) & & \\ & & & & \uparrow & & \uparrow & & \\ & & & & 0 & & 0 & & \end{array}$$

In the mapping cone construction we sum up the terms on the diagonal in the double complex (with the right differentials). This yields

$$0 \leftarrow \mathcal{O}_C \leftarrow \mathcal{O}_{\mathbb{P}^4} \leftarrow \begin{array}{c} \bigwedge^2 F \otimes \mathcal{O}_{\mathbb{P}^4}(-2) \\ \oplus \\ \mathcal{O}_{\mathbb{P}^4}(-3) \end{array} \leftarrow \begin{array}{c} \bigwedge^3 F \otimes DG^* \otimes \mathcal{O}_{\mathbb{P}^4}(-3) \\ \oplus \\ \bigwedge^2 F \otimes \mathcal{O}_{\mathbb{P}^4}(-5) \end{array} \leftarrow \bigwedge^3 F \otimes DG^* \otimes \mathcal{O}_{\mathbb{P}^4}(-6) \leftarrow 0,$$

and the Betti table of the minimal free resolution of $C \subset \mathbb{P}^4$ is

	0	1	2	3
0	1	0	0	0
1	0	3	2	0
2	0	1	0	0
3	0	0	3	2

For $g \geq 8$ the construction of Section 2.2 yields the following double complex

$$\begin{array}{ccccccc}
0 & \longleftarrow & \mathcal{O}_C & \longleftarrow & \mathcal{O}_{\mathbb{P}(\mathcal{E})} & \longleftarrow & \mathcal{O}_{\mathbb{P}(\mathcal{E})}(-3H + (g-7)R) \longleftarrow 0 \\
& & & & \uparrow & & \uparrow \\
& & & & \mathcal{O}_{\mathbb{P}^{g-3}} & \longleftarrow & S_{g-7}G \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-3) \\
& & & & \uparrow & & \uparrow \\
& & & & \bigwedge^2 F \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-2) & \longleftarrow & F \otimes S_{g-8}G \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-4) \\
& & & & \uparrow & & \uparrow \\
& & & & \bigwedge^3 F \otimes DG^* \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-3) & & \vdots \\
& & & & \uparrow & & \\
& & & & \vdots & &
\end{array}$$

and the mapping cone induces the minimal free resolution

$$\begin{array}{ccccccc}
0 \leftarrow \mathcal{O}_C \leftarrow \mathcal{O}_{\mathbb{P}^{g-3}} \leftarrow & \bigwedge^2 F \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-2) & & \bigwedge^3 F \otimes DG^* \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-3) & & & \\
& \oplus & \longleftarrow & \oplus & \longleftarrow & \dots & \\
& S_{g-7}G \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-3) & & F \otimes S_{g-8}G \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-5) & & &
\end{array}$$

The shape of the minimal free resolution of C as an $\mathcal{O}_{\mathbb{P}^{g-3}}$ -module is given as stated in the main theorem. Indeed, we note that the second last map in $\mathcal{E}^{g-7} \otimes \mathcal{O}_{\mathbb{P}^{g-3}}(-3)$ is of degree 2 and occurs in the $(g-6)^{th}$ step in the mapping cone construction. The theorem follows.

Example 2.2. We end this section with a trigonal curve C of genus 6. The minimal free resolution of $C \subset \mathbb{P}^3 = \mathbb{P}(H^0(C, K_C \otimes T^{-1})^*)$ has the following shape

	0	1	2
0	1	0	0
1	0	1	0
2	0	0	0
3	0	2	0
4	0	0	1

We can still apply the above method to compute a minimal free resolution. Note that C has quartic generators induced by $\mathcal{E}^{-1} \otimes \mathcal{O}_{\mathbb{P}^3}(-3)$.

3. HIGHER RESIDUALS OF T WITH RESPECT TO THE CANONICAL BUNDLE

In this section we extend Theorem 1.2 to trigonal curves of genus g embedded by line bundles of the form $K_C \otimes T^{-n}$ for $n \geq 1$ and $g \geq 3n + 4$. We fix the notation of this section. For a trigonal curve C of genus $g \geq 5$ let T be the unique trigonal bundle and let m be the Maroni-invariant (and $a = g - 2 - m$).

3.1. Very ampleness of $K_C \otimes T^{-n}$. We have the following two lemmata.

Lemma 3.1. *For $n > m$ the line bundle $K_C \otimes T^{-n}$ does not separate points of the morphism $C \xrightarrow{3:1} \mathbb{P}^1$ induced by T . In particular, $K_C \otimes T^{-n}$ is not very ample.*

Proof. We may assume that $n = m + 1$ since $H^0(C, K_C \otimes T^{-n_1}) \subset H^0(C, K_C \otimes T^{-n_2})$ for $n_1 \geq n_2$. Let $D = p + q + r \in |T|$ be a divisor. There is a short exact sequence

$$0 \rightarrow T^{m+1} \rightarrow T^{m+1}(D) \rightarrow T^{m+1}(D)|_D \rightarrow 0.$$

The long exact sequence

$$0 \rightarrow H^0(C, T^{m+1}) \rightarrow H^0(C, T^{m+2}) \rightarrow \underbrace{\Gamma(T^{m+1}(D)|_D)}_{3\text{-dimensional}} \rightarrow H^1(C, T^{m+1}) \rightarrow H^1(C, T^{m+2}) \rightarrow 0$$

is induced by the global section functor. Since $m + 2 = \min\{n \in \mathbb{N} : h^0(C, T^n) > n + 1\}$, the difference is

$$h^0(C, T^{m+2}) - h^0(C, T^{m+1}) \geq 2.$$

Hence, by the long exact sequence $h^1(C, T^{m+1}) - h^1(C, T^{m+2}) \leq 1$ and for $p, q \in \text{Supp}(D)$

$$\begin{aligned} h^0(C, K_C \otimes T^{-m-1}) - h^0(C, K_C \otimes T^{-m-1}(-p-q)) &\leq h^0(C, K_C \otimes T^{-m-1}) - h^0(C, K_C \otimes T^{-m-2}) \\ &= h^1(C, T^{m+1}) - h^1(C, T^{m+2}) \\ &\leq 1 \end{aligned}$$

Thus, we can not find a section separating the point p and q . The lemma follows. \square

Lemma 3.2. [Har77, V, Theorem 2.17] or [MS86, Lemma 1, p. 176] *For $n < m$ the line bundle $K_C \otimes T^{-n}$ on C is very ample. The line bundle $K_C \otimes T^{-m}$ is generated by global sections and separates points of the morphism $C \xrightarrow{3:1} \mathbb{P}^1$ induced by T .*

Remark 3.3. For $g \geq 3m + 3$ the line bundle $K_C \otimes T^{-m}$ is very ample since $h^0(C, T^m(p+q)) = h^0(C, T^m) = m + 1$. Indeed, by [Mar46] or [MS86, Proposition 1] the Brill–Noether locus $W_{3m+2}^{m+1}(C)$ is empty for $g \geq 3m + 3$ and that implies $h^0(C, T^m(p+q)) = m + 1$. This can only happen in few cases since $g \leq 3m + 4$ always holds by the inequalities (2.1). Furthermore, the bound $g \geq 3m + 3$ is satisfied for our assumptions in the next section.

3.2. The minimal free resolution of $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-n})^*)$. We fix an integer $n \geq 1$. In order to ensure that the linear system $K_C \otimes T^{-n}$ on a curve C of genus g is very ample, we have to assume that n is less than or equal to the minimal possible Maroni-invariant of a curve of genus g . By (2.1), this yields the bound $n \leq \frac{g-4}{3}$, or equivalently

$$g \geq 3n + 4.$$

We use the same strategy as in the previous section to show that the curve C is projectively normal in $\mathbb{P}(H^0(C, K_C \otimes T^{-n})^*)$. We have the following generalisation of Theorem 1.2.

Theorem 3.4. *Let $n \geq 1$ be an integer. Let C be a trigonal curve of genus $g \geq 3n + 4$ and let T be the unique trigonal bundle. The Betti table of a minimal free resolution of $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-n})^*) = \mathbb{P}^{g-2n-1}$ has the following shape*

	0	1	...	$g - 3n - 3$	$g - 3n - 2$...	$g - 2n - 3$	$g - 2n - 2$
0	1	0	...	0	0	...	0	0
1	0	*	...	*	*	...	*	0
2	0	*	...	*	0	...	0	0
3	0	0	...	0	*	...	*	*

where $*$ indicates nonzero entries. In particular, the embedded curve is projectively normal.

The nonzero Betti numbers are given as in [Nag99, Theorem 2.4] for $c = g - 2n - 2$ and $p = g - 3n - 3$.

Proof. By Section 3.1, let $C \subset \mathbb{P}(H^0(C, K_C \otimes T^{-n})^*) =: \mathbb{P}^{g-2n-1}$ be a smooth embedded curve of genus g and of degree $\deg(C) = 2g - 3n - 2$. Note that $g - 2n - 1 \geq n + 3 \geq 4$. Let

$$X = \bigcup_{D \in |T|} \overline{D} \subset \mathbb{P}^{g-2n-1}$$

be the rational normal scroll swept out by the trigonal bundle. Since $n + 1 < m + 2$,

$$h^0(C, K_C \otimes T^{-n}) - h^0(C, K_C \otimes T^{-n-1}) = 2$$

by (2.2). Hence, X is a rational normal surface of degree $g - 2n - 2$. Let $\mathcal{E} = \mathcal{O}_{\mathbb{P}^1}(g - 2 - m - n) \oplus \mathcal{O}_{\mathbb{P}^1}(m - n)$ be a rank 2 bundle. Then the rational normal surface X is the image of $\mathbb{P}(\mathcal{E})$ as in Section 2.2. Since T is base point free, we consider $C \subset \mathbb{P}(\mathcal{E})$. We denote again H and R the generators of the Picard group of $\mathbb{P}(\mathcal{E})$ as in Section 2.2 such that $H^2 = g - 2n - 2$, $H.R = 1$ and $R^2 = 0$ as well as $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(H) \otimes \mathcal{O}_C = K_C \otimes T^{-n}$ and $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(R) \otimes \mathcal{O}_C = T$.

The resolution of \mathcal{O}_C as an $\mathcal{O}_{\mathbb{P}(\mathcal{E})}$ -module is

$$0 \longleftarrow \mathcal{O}_C \longleftarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})} \longleftarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})}(-aH + bR) \longleftarrow 0$$

for integers a, b since C is a divisor on $\mathbb{P}(\mathcal{E})$. We have

$$(aH - bR).R = 3 \text{ and } (aH - bR).H = \deg(C) = 2g - 3n - 2.$$

Hence, $a = 3$ and $b = g - 3n - 4 \geq 0$ and thus the $\mathcal{O}_{\mathbb{P}(\mathcal{E})}$ -modules in

$$0 \longleftarrow \mathcal{O}_C \longleftarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})} \longleftarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})}(-3H + (g - 3n - 4)R) \longleftarrow 0$$

can be resolved as $\mathcal{O}_{\mathbb{P}^{g-2n-1}}$ -modules by Eagon–Northcott type resolutions (see Section 2.2). As in Section 2.3, the mapping cone of the induced double complex

$$[\mathcal{C}^0 \leftarrow \mathcal{C}^{g-3n-4} \otimes \mathcal{O}_{\mathbb{P}^{g-2n-1}}(-3)]$$

is a minimal free resolution of $C \subset \mathbb{P}^{g-2n-1}$. The shape of its Betti table is given as in Theorem 3.4. Indeed, the length of \mathcal{C}^0 is $g-2n-3$ and the degree 2 syzygies in \mathcal{C}^{g-3n-4} are in the step $j = b + 1 = g - 3n - 3$. Since \mathcal{C}^{g-3n-4} is homologically shifted by 1 in the mapping cone construction, these syzygies appear in step $g - 3n - 2$ as stated in the theorem. \square

REFERENCES

- [ACGH85] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris. *Geometry of algebraic curves. Vol. I*, volume 267 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1985.
- [BE75] David A. Buchsbaum and David Eisenbud. Generic free resolutions and a family of generically perfect ideals. *Adv. Math.*, 18:245–301, 1975.
- [Cas89] G. Castelnuovo. Ricerche di geometria sulle curve algebriche. *Torino Atti*, 24:346–373, 1889.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [GL86] Mark Green and Robert Lazarsfeld. On the projective normality of complete linear series on an algebraic curve. *Invent. Math.*, 83:73–90, 1986.
- [Har77] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [Har81] J. Harris. A bound on the geometric genus of projective varieties. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 8(1):35–68, 1981.
- [LN10] Herbert Lange and Peter E. Newstead. Clifford indices for vector bundles on curves. In *Affine flag manifolds and principal bundles*, Trends Math., pages 165–202. Birkhäuser/Springer Basel AG, Basel, 2010.
- [LN13] Herbert Lange and P. E. Newstead. Bundles computing Clifford indices on trigonal curves. *Arch. Math.*, 101(1):21–31, 2013.
- [Mar46] Arturo Maroni. Le serie lineari speciali sulle curve trigonali. *Ann. Mat. Pura Appl. (4)*, 25:343–354, 1946.
- [MS86] G. Martens and F.-O. Schreyer. Line bundles and syzygies of trigonal curves. *Abh. Math. Semin. Univ. Hamb.*, 56:169–189, 1986.
- [Nag99] Uwe Nagel. Arithmetically Buchsbaum divisors on varieties of minimal degree. *Trans. Amer. Math. Soc.*, 351(11):4381–4409, 1999.
- [Par14] Euisung Park. On syzygies of divisors on rational normal scrolls. *Math. Nachr.*, 287(11-12):1383–1393, 2014.
- [Sch86] Frank-Olaf Schreyer. Syzygies of canonical curves and special linear series. *Math. Ann.*, 275:105–137, 1986.

UNIVERSITÄT DES SAARLANDES, CAMPUS E2 4, D-66123 SAARBRÜCKEN, GERMANY
 Email address: hahn@math.uni-sb.de