

Hom-Yang-Baxter equations and Hom-Yang-Baxter systems

Shengxiang Wang¹, Xiaohui Zhang², Shuangjian Guo^{3*}

1. School of Mathematics and Finance, Chuzhou University,
Chuzhou 239000, China
2. School of Mathematical Sciences, Qufu Normal University,
Qufu 273165, China
3. School of Mathematics and Statistics, Guizhou University of
Finance and Economics, Guiyang 550025, China

ABSTRACT

In this paper, we mainly present some new solutions of the Hom-Yang-Baxter equation from Hom-algebras, Hom-coalgebras and Hom-Lie algebras, respectively. Also, we prove that these solutions are all self-inverse and give some examples. Finally, we introduce the notion of Hom-Yang-Baxter systems and obtain two kinds of Hom-Yang-Baxter systems.

Key words: Hom-Yang-Baxter equation; Hom-Yang-Baxter system; Hom-algebra; Hom-coalgebra; Hom-Lie algebra.

2020 Mathematics Subject Classification: 16T25; 17A30; 17B38.

INTRODUCTION

The Yang-Baxter equation (YBE), which was first introduced by Yang, Baxter and McGuire [1, 28]. YBE has various forms in physics and plays an important role in many topics in mathematical physics, including quantum groups, quantum integrable systems, braided categories and invariants of knots and links.

Recently, a twisted Hom-type generalization of the YBE called Hom-Yang-Baxter equation (HYBE) was introduced in [29, 30, 31] by Yau. The HYBE states

$$(\alpha \otimes B) \circ (B \otimes \alpha) \circ (\alpha \otimes B) = (B \otimes \alpha) \circ (\alpha \otimes B) \circ (B \otimes \alpha),$$

*Correspondence: shuangjianguo@126.com

where α is an endomorphism of the vector space V , and $B : V \otimes V \rightarrow V \otimes V$ is a bilinear map that commutes with $\alpha \otimes \alpha$. In which Yau constructed several classes of solutions of the HYBE, generalizing the solutions of the YBE from Lie algebras and quasitriangular bialgebras. Later, Yau [32] extended the the classical Yang-Baxter equation to the classical Hom-Yang-Baxter equation (CHYBE) in a Hom-Lie algebra and studied the related algebraic structure.

The study of Hom-algebras can be traced back to Hartwig, Larsson and Silvestrov's work in [7], where the notion of Hom-Lie algebra in the context of q-deformation theory of Witt and Virasoro algebras [8] was introduced, which plays an important role in physics, mainly in conformal field theory. Hom-algebras and Hom-coalgebras were introduced by Makhlof and Silvestrov [15] as generalizations of ordinary algebras and coalgebras in the following sense: the associativity of the multiplication is replaced by the Hom-associativity and similar for Hom-coassociativity. They also defined the structures of Hom-bialgebras and Hom-Hopf algebras, and described some of their properties extending properties of ordinary bialgebras and Hopf algebras in [16, 17]. Many more properties and structures of Hom-Hopf algebras have been developed, see [4, 6, 12] and references cited therein.

In [29, 31] Yau proposed the definition of quasitriangular Hom-Hopf algebras and showed that each quasitriangular Hom-Hopf algebra yields a solution of the Hom-Yang-Baxter equation. Meanwhile, several classes of solutions of the Hom-Yang-Baxter equation were constructed from different respects, including those associated to Hom-Lie algebras [5, 25, 29, 30], Drinfeld (co)doubles [2, 34, 35] and Hom-Yetter-Drinfeld modules [3, 10, 13, 14, 18, 26, 33].

In [19, 20, 21], Nichita, Parashar and Popovici presented some solutions of the YBE from associative algebras, coassociative coalgebras and Lie algebras respectively. Depending on the above working, the motivation of constructing new solutions of HYBE is natural. The purpose of the present paper is to investigate how to construct solutions of HYBE from Hom-algebras, Hom-coalgebras and Hom-Lie algebras, and show two kinds of Hom-Yang-Baxter systems from Hom-algebras and Hom-coalgebras.

This paper is organized as follows. In Section 1, we recall some basic definitions about Hom-algebras, Hom-coalgebras and Hom-Lie algebras. In Section 2, we show two solutions of HYBE from Hom-algebras and prove that they are self-inverse (see Theorem 2.1 and Theorem 2.4). In Section 3, we present two solutions of HYBE from Hom-coalgebras and prove that they are self-inverse (see Theorem 3.1 and Theorem 3.4). In Section 4, we obtain a solution of HYBE from Hom-Lie algebras and prove that it is self-inverse (see Theorem 4.1). In Section 5, we introduce the notion of Hom-Yang-Baxter systems and present two kinds of Hom-Yang-Baxter systems (see Theorem 5.2 and Theorem 5.3).

1 PRELIMINARIES

Throughout this paper, k is a fixed field. Unless otherwise stated, all vector spaces, algebras, modules, maps and unadorned tensor products are over k . For a coalgebra C , the coproduct will be denoted by Δ . We adopt a Sweedler's notation $\Delta(c) = c_1 \otimes c_2$, for any $c \in C$, where the summation is understood. We refer to [22] for the Hopf algebra theory and terminology.

We now recall some useful definitions in [9, 15, 16, 17].

Definition 1.1. A Hom-algebra is a quadruple $(A, \mu, 1_A, \alpha)$ (abbr. (A, α)), where A is a k -linear space, $\mu : A \otimes A \rightarrow A$ is a k -linear map, $1_A \in A$ and α is an endomorphism of A , such that

$$\begin{aligned} (HA1) \quad & \alpha(aa') = \alpha(a)\alpha(a'); \quad \alpha(1_A) = 1_A, \\ (HA2) \quad & \alpha(a)(a'a'') = (aa')\alpha(a''); \quad a1_A = 1_Aa = \alpha(a) \end{aligned}$$

are satisfied for $a, a', a'' \in A$. Here we use the notation $\mu(a \otimes a') = aa'$.

Definition 1.2. A Hom-coalgebra is a quadruple $(C, \Delta, \varepsilon, \alpha)$ (abbr. (C, α)), where C is a k -linear space, $\Delta : C \rightarrow C \otimes C$, $\varepsilon : C \rightarrow k$ are k -linear maps, and α is an endomorphism of C , such that

$$\begin{aligned} (HC1) \quad & \alpha(c)_1 \otimes \alpha(c)_2 = \alpha(c_1) \otimes \alpha(c_2); \quad \varepsilon \circ \alpha = \varepsilon; \\ (HC2) \quad & \alpha(c_1) \otimes c_{21} \otimes c_{22} = c_{11} \otimes c_{12} \otimes \alpha(c_2); \quad \varepsilon(c_1)c_2 = c_1\varepsilon(c_2) = \alpha(c) \end{aligned}$$

are satisfied for $c \in C$.

Definition 1.3. A Hom-Lie algebra is a triple $(L, [\cdot, \cdot], \alpha)$ consisting of a linear space L , a bilinear map $[\cdot, \cdot] : L \otimes L \rightarrow L$ and an endomorphism $\alpha : L \rightarrow L$, such that

$$\begin{aligned} (HL1) \quad & [l, l'] = -[l', l], \\ (HL2) \quad & [\alpha(l), [l', l'']] + [\alpha(l'), [l'', l]] + [\alpha(l''), [l, l']] = 0 \end{aligned}$$

are satisfied for all $l, l', l'' \in L$.

2 Solutions of the HYBE from Hom-algebras

In this section, we will give two kinds of solutions of the HYBE from Hom-algebras and prove that these solutions are both self-inverse.

Theorem 2.1. Let $(A, \mu, 1_A, \alpha)$ be a Hom-algebra and $\lambda, \nu \in k$. Then

$$B : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto \lambda ab \otimes 1_A + \nu 1_A \otimes ab - \lambda \alpha(a) \otimes \alpha(b)$$

is a solution for HYBE.

Proof. We first show that B is compatible with the twist map α . For this, we take any $a, b \in A$ and calculate

$$\begin{aligned} (\alpha \otimes \alpha) \circ B(a \otimes b) &= (\alpha \otimes \alpha)(\lambda ab \otimes 1_A + \nu 1_A \otimes ab - \lambda \alpha(a) \otimes \alpha(b)) \\ &= \lambda \alpha(ab) \otimes 1_A + \nu 1_A \otimes \alpha(ab) - \lambda \alpha^2(a) \otimes \alpha^2(b), \\ B \circ (\alpha \otimes \alpha)(a \otimes b) &= \lambda \alpha(a) \alpha(b) \otimes 1_A + \nu 1_A \otimes \alpha(a) \alpha(b) - \lambda \alpha^2(a) \otimes \alpha^2(b). \end{aligned}$$

It follows that $(\alpha \otimes \alpha) \circ B = B \circ (\alpha \otimes \alpha)$, as desired.

Next we will verify that B satisfies the HYBE. In fact, for any $a, b, c \in A$, one may directly check that $B(a \otimes 1_A) = \nu 1_A \otimes \alpha(a)$. On the one hand, we have

$$\begin{aligned} &(\alpha \otimes B)(a \otimes b \otimes c) \\ &= \alpha(a) \otimes (\lambda bc \otimes 1_A + \nu 1_A \otimes bc - \lambda \alpha(b) \otimes \alpha(c)), \\ &(B \otimes \alpha) \circ (\alpha \otimes B)(a \otimes b \otimes c) \\ &= \lambda B(\alpha(a) \otimes bc) \otimes 1_A + \nu B(\alpha(a) \otimes 1_A) \otimes \alpha(bc) - \lambda B(\alpha(a) \otimes \alpha(b)) \otimes \alpha^2(c) \\ &= \lambda^2 \alpha(a)(bc) \otimes 1_A \otimes 1_A + \lambda \nu 1_A \otimes \alpha(a)(bc) \otimes 1_A - \lambda^2 \alpha^2(a) \otimes \alpha(bc) \otimes 1_A \\ &\quad + \nu^2 1_A \otimes \alpha^2(a) \otimes \alpha(bc) - \lambda^2 \alpha(ab) \otimes 1_A \otimes \alpha^2(c) - \lambda \nu 1_A \otimes \alpha(ab) \otimes \alpha^2(c) \\ &\quad + \lambda^2 \alpha^2(a) \otimes \alpha^2(b) \otimes \alpha^2(c), \\ &(\alpha \otimes B) \circ (B \otimes \alpha) \circ (\alpha \otimes B)(a \otimes b \otimes c) \\ &= \lambda^2 \alpha^2(a)(bc) \otimes B(1_A \otimes 1_A) + \lambda \nu 1_A \otimes B(\alpha(a)(bc) \otimes 1_A) - \lambda^2 \alpha^3(a) \otimes B(\alpha(bc) \otimes 1_A) \\ &\quad + \nu^2 1_A \otimes B(\alpha^2(a) \otimes \alpha(bc)) - \lambda^2 \alpha^2(ab) \otimes B(1_A \otimes \alpha^2(c)) - \lambda \nu 1_A \otimes B(\alpha(ab) \otimes \alpha^2(c)) \\ &\quad + \lambda^2 \alpha^3(a) \otimes B(\alpha^2(b) \otimes \alpha^2(c)) \\ &= \frac{\lambda^2 \nu \alpha^2(a) \alpha(bc) \otimes 1_A \otimes 1_A}{(1)} + \frac{\lambda \nu^2 1_A \otimes 1_A \otimes \alpha^2(a) \alpha(bc)}{(2)} - \frac{\lambda^2 \nu \alpha^3(a) \otimes 1_A \otimes \alpha^2(bc)}{(3)} \\ &\quad + \frac{\lambda \nu^2 1_A \otimes \alpha^2(a) \alpha(bc) \otimes 1_A}{(4)} + \frac{\nu^3 1_A \otimes 1_A \otimes \alpha^2(a) \alpha(bc)}{(5)} - \frac{\lambda \nu^2 1_A \otimes \alpha^3(a) \otimes \alpha^2(bc)}{(6)} \\ &\quad - \frac{\lambda^3 \alpha^2(ab) \otimes \alpha^3(c) \otimes 1_A}{(7)} - \frac{\lambda^2 \nu \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)}{(8)} + \frac{\lambda^3 \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)}{(9)} \\ &\quad - \frac{\lambda^2 \nu 1_A \otimes \alpha(ab) \alpha^{23}(c) \otimes 1_A}{(10)} - \frac{\lambda \nu^2 1_A \otimes 1_A \otimes \alpha^2(a) \alpha(bc)}{(11)} + \frac{\lambda^2 \nu 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)}{(12)} \\ &\quad + \frac{\lambda^3 \alpha^3(a) \otimes \alpha^2(bc) \otimes 1_A}{(13)} + \frac{\lambda^2 \nu \alpha^3(a) \otimes 1_A \otimes \alpha^2(bc)}{(14)} - \frac{\lambda^3 \alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)}{(15)} \\ &= \frac{\lambda^2 \nu \alpha^2(a) \alpha(bc) \otimes 1_A \otimes 1_A}{(1)} + \frac{\nu^3 1_A \otimes 1_A \otimes \alpha^2(a) \alpha(bc)}{(5)} - \frac{\lambda \nu^2 1_A \otimes \alpha^3(a) \otimes \alpha^2(bc)}{(6)} \\ &\quad - \frac{\lambda^3 \alpha^2(ab) \otimes \alpha^3(c) \otimes 1_A}{(7)} - \frac{\lambda^2 \nu \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)}{(8)} + \frac{\lambda^3 \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)}{(9)} \\ &\quad + \frac{\lambda^2 \nu 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)}{(12)} + \frac{\lambda^3 \alpha^3(a) \otimes \alpha^2(bc) \otimes 1_A}{(13)} - \frac{\lambda^3 \alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)}{(15)}. \end{aligned}$$

The last equality holds since (2)=(11), (3)=(14) and (4)=(10).

On the other hand, we have

$$\begin{aligned} &(B \otimes \alpha)(a \otimes b \otimes c) \\ &= \lambda ab \otimes 1_A \otimes \alpha(c) + \nu 1_A \otimes ab \otimes \alpha(c) - \lambda \alpha(a) \otimes \alpha(b) \otimes \alpha(c), \\ &(\alpha \otimes B) \circ (B \otimes \alpha)(a \otimes b \otimes c) \\ &= \lambda \alpha(ab) \otimes B(1_A \otimes \alpha(c)) + \nu 1_A \otimes B(ab \otimes \alpha(c)) - \lambda \alpha^2(a) \otimes B(\alpha(b) \otimes \alpha(c)) \\ &= \lambda^2 \alpha(ab) \otimes \alpha^2(c) \otimes 1_A + \lambda \nu \alpha(ab) \otimes 1_A \otimes \alpha^2(c) - \lambda^2 \alpha(ab) \otimes 1_A \otimes \alpha^2(c) \end{aligned}$$

$$\begin{aligned}
& + \lambda\nu 1_A \otimes (ab)\alpha(c) \otimes 1_A + \nu^2 1_A \otimes 1_A \otimes (ab)\alpha(c) - \lambda\nu 1_A \otimes \alpha(ab) \otimes \alpha^2(c) \\
& - \lambda^2 \alpha^2(a) \otimes \alpha(bc) \otimes 1_A - \lambda\nu \alpha^2(a) \otimes 1_A \otimes \alpha(bc) + \lambda^2 \alpha^2(a) \otimes \alpha^2(b) \otimes \alpha^2(c), \\
(B \otimes \alpha) \circ (\alpha \otimes B) \circ (B \otimes \alpha)(a \otimes b \otimes c) \\
& = \lambda^2 B(\alpha(ab) \otimes \alpha^2(c)) \otimes 1_A + \lambda\nu B(\alpha(ab) \otimes 1_A) \otimes \alpha^3(c) - \lambda^2 B(\alpha(ab) \otimes 1_A) \otimes \alpha^3(c) \\
& + \lambda\nu B(1_A \otimes (ab)\alpha(c)) \otimes 1_A + \nu^2 B(1_A \otimes 1_A) \otimes \alpha(ab)\alpha^2(c) - \lambda\nu B(1_A \otimes \alpha(ab)) \otimes \alpha^3(c) \\
& - \lambda^2 B(\alpha^2(a) \otimes \alpha(bc)) \otimes 1_A - \lambda\nu B(\alpha^2(a) \otimes 1_A) \otimes \alpha^2(bc) + \lambda^2 B(\alpha^2(a) \otimes \alpha^2(b)) \otimes \alpha^3(c) \\
& = \lambda^3 \alpha(ab)\alpha^2(c) \otimes 1_A \otimes 1_{A(16)} + \lambda^2 \nu 1_A \otimes \alpha(ab)\alpha^2(c) \otimes 1_{A(17)} - \lambda^3 \alpha^2(ab) \otimes \alpha^3(c) \otimes 1_{A(18)} \\
& - \lambda^2 \nu 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)_{(19)} + \lambda \nu^2 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)_{(20)} + \lambda^2 \nu \alpha(ab)\alpha^2(c) \otimes 1_A \otimes 1_{A(21)} \\
& + \lambda \nu^2 1_A \otimes \alpha(ab)\alpha^2(c) \otimes 1_{A(22)} - \lambda^2 \nu 1_A \otimes \alpha(ab)\alpha^2(c) \otimes 1_{A(23)} + \nu^3 1_A \otimes 1_A \otimes \alpha(ab)\alpha^2(c)_{(24)} \\
& - \lambda^2 \nu \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)_{(25)} - \lambda \nu^2 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)_{(26)} + \lambda^2 \nu 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)_{(27)} \\
& - \lambda^3 \alpha^2(a)\alpha(bc) \otimes 1_A \otimes 1_{A(28)} - \lambda \nu^2 1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_{A(29)} + \lambda^3 \alpha^3(a) \otimes \alpha^2(bc) \otimes 1_{A(30)} \\
& - \lambda \nu^2 1_A \otimes \alpha^3(a) \otimes \alpha^2(bc)_{(31)} + \lambda^3 \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)_{(32)} + \lambda^2 \nu 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)_{(33)} \\
& - \lambda^3 \alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)_{(34)} \\
& = -\lambda^3 \alpha^2(ab) \otimes \alpha^3(c) \otimes 1_{A(18)} + \lambda^2 \nu \alpha(ab)\alpha^2(c) \otimes 1_A \otimes 1_{A(21)} + \nu^3 1_A \otimes 1_A \otimes \alpha(ab)\alpha^2(c)_{(24)} \\
& - \lambda^2 \nu \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)_{(25)} + \lambda^2 \nu 1_A \otimes \alpha^2(ab) \otimes \alpha^3(c)_{(27)} + \lambda^3 \alpha^3(a) \otimes \alpha^2(bc) \otimes 1_{A(30)} \\
& - \lambda \nu^2 1_A \otimes \alpha^3(a) \otimes \alpha^2(bc)_{(31)} + \lambda^3 \alpha^2(ab) \otimes 1_A \otimes \alpha^3(c)_{(32)} - \lambda^3 \alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)_{(34)}.
\end{aligned}$$

The last equality holds since (16)=(28), (17)=(23), (19)=(33), (20)=(26) and (22)=(29).

Compare two expressions above, we have

$$\begin{aligned}
(1) &= (21), (5) = (24), (6) = (31), (7) = (18), \\
(8) &= (25), (9) = (32), (12) = (27), (13) = (30), (15) = (34).
\end{aligned}$$

It follows that

$$(\alpha \otimes B) \circ (B \otimes \alpha) \circ (\alpha \otimes B)(a \otimes b \otimes c) = (B \otimes \alpha) \circ (\alpha \otimes B) \circ (B \otimes \alpha)(a \otimes b \otimes c).$$

So B is a solution for HYBE. \square

Corollary 2.2. Let $(A, \mu, 1_A, \alpha)$ be a Hom-algebra and $\lambda, \nu \in k^*$. Assume that α is involutive, then the solution B in Theorem 2.1 is invertible, where the inverse is given by

$$B^{-1} : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto \frac{1}{\nu} ab \otimes 1_A + \frac{1}{\lambda} 1_A \otimes ab - \frac{1}{\lambda} \alpha(a) \otimes \alpha(b).$$

Proof. We first show that $B \circ B^{-1} = id_{A \otimes A}$. In fact, for any $a, b, c \in A$, we have

$$\begin{aligned}
B \circ B^{-1}(a \otimes b) &= \frac{1}{\nu} B(ab \otimes 1_A) + \frac{1}{\lambda} B(1_A \otimes ab) - \frac{1}{B} (\lambda \alpha(a) \otimes \alpha(b)) \\
&= 1_A \otimes \alpha(ab) + \frac{1}{\lambda} \{ \lambda \alpha(ab) \otimes 1_A + \nu 1_A \otimes \alpha(ab) - \lambda 1_A \otimes \alpha(ab) \} \\
&\quad - \frac{1}{\lambda} \{ \lambda \alpha(ab) \otimes 1_A + \nu 1_A \otimes \alpha(ab) - \lambda \alpha^2(a) \otimes \alpha^2(b) \}
\end{aligned}$$

$$= a \otimes b.$$

It follows that $B \circ B^{-1} = id_{A \otimes A}$. Similarly, one may check that $B^{-1} \circ B = id_{A \otimes A}$. So B^{-1} is the inverse of B . \square

Example 2.3. Let $\{x_1, x_2, x_3\}$ be a basis of a 3-dimensional linear space A . The following multiplication μ and the twist map α on A define a Hom-algebra:

$$\begin{aligned} \mu(x_1, x_1) &= x_1, \quad \mu(x_1, x_2) = x_2, \quad \mu(x_1, x_3) = lx_3, \\ \mu(x_2, x_1) &= x_2, \quad \mu(x_2, x_2) = x_2, \quad \mu(x_2, x_3) = lx_3, \\ \mu(x_3, x_1) &= lx_3, \quad \mu(x_3, x_2) = 0, \quad \mu(x_3, x_3) = 0, \\ \alpha(x_1) &= x_1, \quad \alpha(x_2) = x_2, \quad \alpha(x_3) = lx_3, \end{aligned}$$

where l is a parameter in k ([13]). It is easy to see that $1_A = x_1$. Therefore, by Theorem 2.1, the solution B of the HYBE for the Hom-algebra A satisfies

$$\begin{aligned} B(x_1 \otimes x_1) &= \nu x_1 \otimes x_1, \quad B(x_1 \otimes x_2) = \lambda x_2 \otimes x_1 + (\nu - \lambda)x_1 \otimes x_2, \\ B(x_1 \otimes x_3) &= \lambda lx_3 \otimes x_1 + l(\nu - \lambda)x_1 \otimes x_3, \quad B(x_2 \otimes x_1) = \nu x_1 \otimes x_2, \\ B(x_2 \otimes x_2) &= \lambda x_2 \otimes x_1 + \nu x_1 \otimes x_2 - \lambda x_2 \otimes x_2, \\ B(x_2 \otimes x_3) &= \lambda lx_3 \otimes x_1 + \nu lx_1 \otimes x_3 - \lambda lx_2 \otimes x_3, \\ B(x_3 \otimes x_1) &= \nu lx_1 \otimes x_3, \quad B(x_3 \otimes x_2) = -\lambda lx_3 \otimes x_2, \quad B(x_3 \otimes x_3) = -\lambda l^2 x_3 \otimes x_3, \end{aligned}$$

where λ, ν are two parameters in k .

Theorem 2.4. Let $(A, \mu, 1_A, \alpha)$ be a Hom-algebra and $\lambda, \nu \in k$. Then

$$B : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto \lambda ab \otimes 1_A + \nu 1_A \otimes ab - \nu \alpha(a) \otimes \alpha(b)$$

is also a solution for HYBE. If, in addition, $\lambda, \nu \in k^*$ and α is involutive, then B is invertible, where the inverse is given by

$$B^{-1} : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto \frac{1}{\nu} ab \otimes 1_A + \frac{1}{\lambda} 1_A \otimes ab - \frac{1}{\nu} \alpha(a) \otimes \alpha(b).$$

Proof. Similar to the proof of Theorem 2.1 and Corollary 2.2. \square

Example 2.5. Let $\{1, g, x, y\}$ be a basis of a 4-dimensional linear space H_4 . The following multiplication μ and the twist map α on H_4 define a Hom-algebra:

$$\begin{aligned} \mu(1, 1) &= 1, \quad \mu(1, g) = g, \quad \mu(1, x) = kx, \quad \mu(1, y) = ky, \\ \mu(g, 1) &= g, \quad \mu(g, g) = g, \quad \mu(g, x) = ky, \quad \mu(g, y) = kx, \\ \mu(x, 1) &= kx, \quad \mu(x, g) = ky, \quad \mu(x, x) = 0, \quad \mu(x, y) = 0, \\ \mu(y, 1) &= ky, \quad \mu(y, g) = -kx, \quad \mu(y, x) = 0, \quad \mu(y, y) = 0, \end{aligned}$$

$$\alpha(1) = 1, \alpha(g) = g, \alpha(x) = kx, \alpha(y) = ky,$$

where k is a parameter in k ([11]). By Theorem 2.4, the solution B of the HYBE for the Hom-algebra H_4 satisfies

$$\begin{aligned} B(1 \otimes 1) &= \lambda 1 \otimes 1, B(1 \otimes g) = \lambda g \otimes 1, B(1 \otimes x) = \lambda kx \otimes 1, B(1 \otimes y) = \lambda k1 \otimes y, \\ B(g \otimes 1) &= (\lambda - \nu)g \otimes 1 + \nu 1 \otimes g, B(g \otimes g) = (\lambda + \nu)1 \otimes 1 - \nu g \otimes g, \\ B(g \otimes x) &= \lambda ky \otimes 1 + \nu k1 \otimes y - \nu kg \otimes x, B(g \otimes y) = \lambda kx \otimes 1 + \nu k1 \otimes x - \nu kg \otimes y, \\ B(x \otimes 1) &= k(\lambda - \nu)x \otimes 1 + \nu k1 \otimes x, B(x \otimes g) = -\lambda ky \otimes 1 - \nu k1 \otimes y - \lambda kx \otimes g, \\ B(x \otimes x) &= -k^2x \otimes x, B(x \otimes y) = -k^2x \otimes y, B(y \otimes 1) = k(\lambda - \nu)y \otimes 1 + \nu k1 \otimes y, \\ B(y \otimes g) &= -\lambda kx \otimes 1 - \nu k1 \otimes x - \lambda ky \otimes g, B(y \otimes x) = -k^2y \otimes x, B(y \otimes y) = -k^2y \otimes y, \end{aligned}$$

where λ, ν are two parameters in k .

3 Solutions of the HYBE from Hom-coalgebras

In this section, we will show two kinds of solutions of the HYBE from Hom-coalgebras and prove that these solutions are both self-inverse.

Theorem 3.1. Let $(C, \Delta, \varepsilon, \alpha)$ be a Hom-coalgebra and $\lambda, \nu \in k$. Then

$$B : C \otimes C \rightarrow C \otimes C, a \otimes b \mapsto \lambda \varepsilon(a)b_1 \otimes b_2 + \nu \varepsilon(b)a_1 \otimes a_2 - \lambda \alpha(a) \otimes \alpha(b)$$

is a solution for HYBE.

Proof. We first show that B is compatible with the twist map α . For this, we take any $a, b \in C$ and calculate

$$\begin{aligned} (\alpha \otimes \alpha) \circ B(a \otimes b) &= (\alpha \otimes \alpha)(\lambda \varepsilon(a)b_1 \otimes b_2 + \nu \varepsilon(b)a_1 \otimes a_2 - \lambda \alpha(a) \otimes \alpha(b)) \\ &= \lambda \varepsilon(a)\alpha(b_1) \otimes \alpha(b_2) + \nu \varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) - \lambda \alpha^2(a) \otimes \alpha^2(b), \\ B \circ (\alpha \otimes \alpha)(a \otimes b) &= \lambda \varepsilon(\alpha(a))\alpha(b_1) \otimes \alpha(b_2) + \nu \varepsilon(\alpha(b))\alpha(a_1) \otimes \alpha(a_2) - \lambda \alpha^2(a) \otimes \alpha^2(b) \\ &= \lambda \varepsilon(a)\alpha(b_1) \otimes \alpha(b_2) + \nu \varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) - \lambda \alpha^2(a) \otimes \alpha^2(b). \end{aligned}$$

It follows that $(\alpha \otimes \alpha) \circ B = B \circ (\alpha \otimes \alpha)$, as desired.

Next we will verify that B satisfies the HYBE. For this, we take any $a, b, c \in C$ and calculate

$$\begin{aligned} (\alpha \otimes B)(a \otimes b \otimes c) &= \lambda \varepsilon(b)\alpha(a) \otimes c_1 \otimes c_2 + \nu \varepsilon(c)\alpha(a) \otimes b_1 \otimes b_2 - \lambda \alpha(a) \otimes \alpha(b) \otimes \alpha(c), \\ (B \otimes \alpha) \circ (\alpha \otimes B)(a \otimes b \otimes c) &= \lambda \varepsilon(b)B(\alpha(a) \otimes c_1) \otimes \alpha(c_2) + \nu \varepsilon(c)B(\alpha(a) \otimes b_1) \otimes \alpha(b_2) - \lambda B(\alpha(a) \otimes \alpha(b)) \otimes \alpha^2(c) \\ &= \lambda^2 \varepsilon(a)\varepsilon(b)c_{11} \otimes c_{12} \otimes \alpha(c_2) + \lambda \nu \varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) \otimes \alpha^2(c) \end{aligned}$$

$$\begin{aligned}
& -\lambda^2\varepsilon(b)\alpha^2(a) \otimes \alpha(c_1) \otimes \alpha(c_2) + \lambda\nu\varepsilon(a)\varepsilon(c)b_{11} \otimes b_{12} \otimes \alpha(b_2) \\
& + \nu^2\varepsilon(c)\alpha(a_1) \otimes \alpha(a_2) \otimes \alpha^2(b) - \lambda\nu\varepsilon(c)\alpha^2(a) \otimes \alpha(b_1) \otimes \alpha(b_2) \\
& - \lambda^2\varepsilon(a)\alpha(b_1) \otimes \alpha(b_2) \otimes \alpha^2(c) - \lambda\nu\varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) \otimes \alpha^2(c) \\
& + \lambda^2\alpha^2(a) \otimes \alpha^2(b) \otimes \alpha^2(c), \\
(\alpha \otimes B) \circ (B \otimes \alpha) \circ (\alpha \otimes B)(a \otimes b \otimes c) \\
& = \lambda^2\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes B(c_{12} \otimes \alpha(c_2)) + \lambda\nu\varepsilon(b)\alpha^2(a_1) \otimes B(\alpha(a_2) \otimes \alpha^2(c)) \\
& - \lambda^2\varepsilon(b)\alpha^3(a) \otimes B(\alpha(c_1) \otimes \alpha(c_2)) + \lambda\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes B(b_{12} \otimes \alpha(b_2)) \\
& + \nu^2\varepsilon(c)\alpha^2(a_1) \otimes B(\alpha(a_2) \otimes \alpha^2(b)) - \lambda\nu\varepsilon(c)\alpha^3(a) \otimes B(\alpha(b_1) \otimes \alpha(b_2)) \\
& - \lambda^2\varepsilon(a)\alpha^2(b_1) \otimes B(\alpha(b_2) \otimes \alpha^2(c)) - \lambda\nu\varepsilon(b)\alpha^2(a_1) \otimes B(\alpha(a_2) \otimes \alpha^2(c)) \\
& + \lambda^2\alpha^3(a) \otimes B(\alpha^2(b) \otimes \alpha^2(c)) \\
& = \frac{\lambda^3\varepsilon(a)\varepsilon(b)\alpha^2(c_1) \otimes \alpha(c_{21}) \otimes \alpha(c_{22})}{(1)} + \frac{\lambda^2\nu\varepsilon(a)\varepsilon(b)\alpha^2(c_1) \otimes \alpha(c_{21}) \otimes \alpha(c_{22})}{(2)} \\
& - \frac{\lambda^2\nu\varepsilon(a)\varepsilon(b)\alpha^2(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}{(3)} + \frac{\lambda^2\nu\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(4)} \\
& + \frac{\lambda\nu^2\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}{(5)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(6)} \\
& - \frac{\lambda^3\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(7)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(8)} \\
& + \frac{\lambda^3\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(9)} + \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}{(10)} \\
& + \frac{\lambda\nu^2\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}{(11)} - \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}{(12)} \\
& + \frac{\lambda\nu^2\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(13)} + \frac{\nu^3\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}{(14)} \\
& - \frac{\lambda\nu^2\varepsilon(c)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(b)}{(15)} - \frac{\lambda^2\nu\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(16)} \\
& - \frac{\lambda\nu^2\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(17)} + \frac{\lambda^2\nu\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(18)} \\
& - \frac{\lambda^3\varepsilon(a)\alpha^3(b) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(19)} - \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}{(20)} \\
& + \frac{\lambda^3\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(21)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(22)} \\
& - \frac{\lambda\nu^2\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}{(23)} + \frac{\lambda^2\nu\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(24)} \\
& + \frac{\lambda^3\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(25)} + \frac{\lambda^2\nu\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(26)} \\
& - \frac{\lambda^3\alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)}{(27)} \\
& = \frac{\lambda^3\varepsilon(a)\varepsilon(b)\alpha^2(c_1) \otimes \alpha(c_{21}) \otimes \alpha(c_{22})}{(1)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(8)} \\
& + \frac{\lambda^3\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(25)} + \frac{\nu^3\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}{(14)} \\
& + \frac{\lambda\nu^2\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}{(11)} - \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}{(20)} \\
& - \frac{\lambda\nu^2\varepsilon(c)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(b)}{(15)} + \frac{\lambda^2\nu\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(26)} \\
& - \frac{\lambda^3\varepsilon(a)\alpha^3(b) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(19)} + \frac{\lambda^3\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(21)} \\
& - \frac{\lambda^3\alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)}{(27)}.
\end{aligned}$$

The last equality holds since (2)=(3), (4)=(22), (5)=(23), (6)=(24), (7)=(9), (10)=(12), (13)=(17) and (16)=(18).

On the other hand, we have

$$(B \otimes \alpha)(a \otimes b \otimes c)$$

$$\begin{aligned}
&= \lambda\varepsilon(a)b_1 \otimes b_2 \otimes \alpha(c) + \nu\varepsilon(b)a_1 \otimes a_2 \otimes \alpha(c) - \lambda\alpha(a) \otimes \alpha(b) \otimes \alpha(c), \\
(\alpha \otimes B) \circ (B \otimes \alpha)(a \otimes b \otimes c) \\
&= \lambda\varepsilon(a)\alpha(b_1) \otimes B(b_2 \otimes \alpha(c)) + \nu\varepsilon(b)\alpha(a_1) \otimes B(a_2 \otimes \alpha(c)) - \lambda\alpha^2(a) \otimes B(\alpha(b) \otimes \alpha(c)) \\
&= \lambda^2\varepsilon(a)\alpha^2(b) \otimes \alpha(c_1) \otimes \alpha(c_2) + \lambda\nu\varepsilon(a)\varepsilon(c)\alpha(b_1) \otimes b_{21} \otimes b_{22} \\
&\quad - \lambda^2\varepsilon(a)\alpha(b_1) \otimes \alpha(b_2) \otimes \alpha^2(c) + \lambda\nu\varepsilon(b)\alpha^2(a) \otimes \alpha(c_1) \otimes \alpha(c_2) \\
&\quad + \nu^2\varepsilon(b)\varepsilon(c)\alpha(a_1) \otimes a_{21} \otimes a_{22} - \lambda\nu\varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) \otimes \alpha^2(c) \\
&\quad - \lambda^2\varepsilon(b)\alpha^2(a) \otimes \alpha(c_1) \otimes \alpha(c_2) - \lambda\nu\varepsilon(c)\alpha^2(a) \otimes \alpha(b_1) \otimes \alpha(b_2) \\
&\quad + \lambda^2\alpha^2(a) \otimes \alpha^2(b) \otimes \alpha^2(c), \\
(B \otimes \alpha) \circ (\alpha \otimes B) \circ (B \otimes \alpha)(a \otimes b \otimes c) \\
&= \lambda^2\varepsilon(a)B(\alpha^2(b) \otimes \alpha(c_1)) \otimes \alpha^2(c_2) + \lambda\nu\varepsilon(a)\varepsilon(c)B(\alpha(b_1) \otimes b_{21}) \otimes \alpha(b_{22}) \\
&\quad - \lambda^2\varepsilon(a)B(\alpha(b_1) \otimes \alpha(b_2)) \otimes \alpha^3(c) + \lambda\nu\varepsilon(b)B(\alpha^2(a) \otimes \alpha(c_1)) \otimes \alpha^2(c_2) \\
&\quad + \nu^2\varepsilon(b)\varepsilon(c)B(\alpha(a_1) \otimes a_{21}) \otimes \alpha(a_{22}) - \lambda\nu\varepsilon(b)B(\alpha(a_1) \otimes \alpha(a_2)) \otimes \alpha^3(c) \\
&\quad - \lambda^2\varepsilon(b)B(\alpha^2(a) \otimes \alpha(c_1)) \otimes \alpha^2(c_2) - \lambda\nu\varepsilon(c)B(\alpha^2(a) \otimes \alpha(b_1)) \otimes \alpha^2(b_2) \\
&\quad + \lambda^2B(\alpha^2(a) \otimes \alpha^2(b)) \otimes \alpha^3(c) \\
&= \frac{\lambda^3\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}{(28)} + \frac{\lambda^2\nu\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(29)} \\
&\quad - \frac{\lambda^3\varepsilon(a)\alpha^3(b) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(30)} + \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}{(31)} \\
&\quad + \frac{\lambda\nu^2\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}{(32)} - \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}{(33)} \\
&\quad - \frac{\lambda^3\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(34)} - \frac{\lambda^2\nu\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(35)} \\
&\quad + \frac{\lambda^3\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(36)} + \frac{\lambda^2\nu\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}{(37)} \\
&\quad + \frac{\lambda\nu^2\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(38)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(39)} \\
&\quad + \frac{\lambda\nu^2\varepsilon(b)\varepsilon(c)\alpha(a_{11}) \otimes \alpha(a_{12}) \otimes \alpha^2(a_2)}{(40)} + \frac{\nu^3\varepsilon(b)\varepsilon(c)\alpha(a_{11}) \otimes \alpha(a_{12}) \otimes \alpha^2(a_2)}{(41)} \\
&\quad - \frac{\lambda\nu^2\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}{(42)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(43)} \\
&\quad - \frac{\lambda\nu^2\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(44)} + \frac{\lambda^2\nu\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(45)} \\
&\quad - \frac{\lambda^3\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}{(46)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(47)} \\
&\quad + \frac{\lambda^3\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(48)} - \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}{(49)} \\
&\quad - \frac{\lambda\nu^2\varepsilon(c)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(b)}{(50)} + \frac{\lambda^2\nu\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(51)} \\
&\quad + \frac{\lambda^3\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(52)} + \frac{\lambda^2\nu\varepsilon(b)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(c)}{(53)} \\
&\quad - \frac{\lambda^3\alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)}{(54)} \\
&= \frac{\lambda^3\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}{(28)} - \frac{\lambda^3\varepsilon(a)\alpha^3(b) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(30)} \\
&\quad + \frac{\lambda\nu^2\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}{(32)} - \frac{\lambda^2\nu\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(39)} \\
&\quad + \frac{\nu^3\varepsilon(b)\varepsilon(c)\alpha(a_{11}) \otimes \alpha(a_{12}) \otimes \alpha^2(a_2)}{(41)} + \frac{\lambda^3\varepsilon(b)\alpha^3(a) \otimes \alpha^2(c_1) \otimes \alpha^2(c_2)}{(48)} \\
&\quad - \frac{\lambda^2\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}{(49)} - \frac{\lambda\nu^2\varepsilon(c)\alpha^2(a_1) \otimes \alpha^2(a_2) \otimes \alpha^3(b)}{(50)} \\
&\quad + \frac{\lambda^2\nu\varepsilon(c)\alpha^3(a) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(51)} + \frac{\lambda^3\varepsilon(a)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(c)}{(52)} \\
&\quad - \frac{\lambda^3\alpha^3(a) \otimes \alpha^3(b) \otimes \alpha^3(c)}{(54)}.
\end{aligned}$$

The last equality holds since (29)=(35), (31)=(33), (34)=(36), (37)=(46), (38)=(44), (40)=(42), (43)=(45) and (47)=(43). Compare the two expressions above, we have

$$(1) = (28), (8) = (39), (11) = (32), (14) = (41), (15) = (50), \\ (19) = (30), (20) = (49), (21) = (52), (25) = (48), (26) = (51), (27) = (54).$$

That is, we prove

$$(\alpha \otimes B) \circ (B \otimes \alpha) \circ (\alpha \otimes B)(a \otimes b \otimes c) = (B \otimes \alpha) \circ (\alpha \otimes B) \circ (B \otimes \alpha)(a \otimes b \otimes c),$$

as desired. So B is a solution for HYBE. \square

Corollary 3.2. Let $(C, \Delta, \varepsilon, \alpha)$ be a Hom-coalgebra and $\lambda, \nu \in k^*$. Assume that α is involutive, then the solution B in Theorem 3.1 is invertible, where the inverse is given by

$$B^{-1} : C \otimes C \rightarrow C \otimes C, \quad a \otimes b \mapsto \frac{1}{\nu}\varepsilon(a)b_1 \otimes b_2 + \frac{1}{\lambda}\varepsilon(b)a_1 \otimes a_2 - \frac{1}{\lambda}\alpha(a) \otimes \alpha(b)$$

Proof. We first show that $B \circ B^{-1} = id_{C \otimes C}$. In fact, for any $a, b, c \in C$, we have

$$\begin{aligned} B \circ B^{-1}(a \otimes b) &= \frac{1}{\nu}\varepsilon(a)B(b_1 \otimes b_2) + \frac{1}{\lambda}\varepsilon(b)B(a_1 \otimes a_2) - \frac{1}{\lambda}B(\alpha(a) \otimes \alpha(b)) \\ &= \frac{1}{\nu}\varepsilon(a)\{\lambda\varepsilon(b_1)b_{21} \otimes b_{22} + \nu\varepsilon(b_2)b_{11} \otimes b_{12} - \lambda\alpha(b_1) \otimes \alpha(b_2)\} \\ &\quad + \frac{1}{\lambda}\varepsilon(b)\{\lambda\varepsilon(a_1)a_{21} \otimes a_{22} + \nu\varepsilon(a_2)a_{11} \otimes a_{12} - \lambda\alpha(a_1) \otimes \alpha(a_2)\} \\ &\quad - \frac{1}{\lambda}\{\lambda\varepsilon(a)\alpha(b_1) \otimes \alpha(b_2) + \nu\varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) - \lambda\alpha^2(a) \otimes \alpha^2(b)\} \\ &= \varepsilon(a)\alpha(b_1) \otimes \alpha(b_2) + \frac{\nu}{\lambda}\varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) - \varepsilon(a)\alpha(b_1) \otimes \alpha(b_2) \\ &\quad - \frac{\nu}{\lambda}\varepsilon(b)\alpha(a_1) \otimes \alpha(a_2) + \alpha^2(a) \otimes \alpha^2(b) \\ &= a \otimes b. \end{aligned}$$

It follows that $B \circ B^{-1} = id_{C \otimes C}$. Similarly, one may check that $B^{-1} \circ B = id_{C \otimes C}$. So B^{-1} is the inverse of B . \square

Example 3.3. Let $\{1, a, a^2\}$ be a basis of a 3-dimensional linear space C . The following comultiplication Δ , the counit ε and the twist map α on C define a Hom-coalgebra ([27]):

$$\begin{aligned} \Delta(1) &= 1 \otimes 1, \quad \Delta(a) = a^2 \otimes a^2, \quad \Delta(a^2) = a \otimes a, \\ \varepsilon(1) &= 1, \quad \varepsilon(a) = 1, \quad \varepsilon(a^2) = 1, \\ \alpha(1) &= 1, \quad \alpha(a) = a^2, \quad \alpha(a^2) = a^2. \end{aligned}$$

Therefore, by Theorem 3.1, the solution B of the HYBE for the Hom-coalgebra C satisfies:

$$B(1 \otimes 1) = \nu 1 \otimes 1, \quad B(1 \otimes a) = \lambda a^2 \otimes a^2 + \nu 1 \otimes 1 - \lambda 1 \otimes a^2,$$

$$\begin{aligned}
B(1 \otimes a^2) &= \lambda a \otimes a + \nu 1 \otimes 1 - \lambda 1 \otimes a, & B(a \otimes 1) &= \lambda 1 \otimes 1 + \nu a^2 \otimes a^2 - \lambda a^2 \otimes 1, \\
B(a \otimes a) &= \nu a^2 \otimes a^2, & B(a \otimes a^2) &= \lambda a \otimes a + \nu a^2 \otimes a^2 - \lambda a^2 \otimes a, \\
B(a^2 \otimes 1) &= \lambda 1 \otimes 1 + \nu a \otimes a - \lambda a \otimes 1, & B(a^2 \otimes a) &= \nu a \otimes a, \\
B(a^2 \otimes a^2) &= \lambda a^2 \otimes a^2 + \nu a \otimes a - \lambda a \otimes a^2,
\end{aligned}$$

where λ, ν are two parameters in k .

Theorem 3.4. Let $(C, \Delta, \varepsilon, \alpha)$ be a Hom-coalgebra and $\lambda, \nu \in k$. Then

$$B : C \otimes C \rightarrow C \otimes C, \quad a \otimes b \mapsto \lambda \varepsilon(a) b_1 \otimes b_2 + \nu \varepsilon(b) a_1 \otimes a_2 - \nu \alpha(a) \otimes \alpha(b)$$

is also a solution for HYBE. If, in addition, $\lambda, \nu \in k^*$ and α is involutive, then B is invertible, where the inverse is given by

$$B^{-1} : C \otimes C \rightarrow C \otimes C, \quad a \otimes b \mapsto \frac{1}{\nu} \varepsilon(a) b_1 \otimes b_2 + \frac{1}{\lambda} \varepsilon(b) a_1 \otimes a_2 - \frac{1}{\nu} \alpha(a) \otimes \alpha(b).$$

Proof. Similar to the proof of Theorem 3.1 and Corollary 3.2. \square

Example 3.5. Let $\{1, g, x, y\}$ be a basis of a 4-dimensional linear space H_4 . The following comultiplication Δ , counit ε and the twist map α on H_4 define a Hom-coalgebra ([11]):

$$\begin{aligned}
\Delta(1) &= 1 \otimes 1, \Delta(g) = g \otimes g, \Delta(x) = kx \otimes 1 + g \otimes kx, \Delta(y) = ky \otimes g + 1 \otimes ky, \\
\varepsilon(1) &= 1, \varepsilon(g) = 1, \varepsilon(x) = 0, \varepsilon(y) = 0, \\
\alpha(1) &= 1, \alpha(g) = g, \alpha(x) = kx, \alpha(y) = ky,
\end{aligned}$$

where k is a parameter in k . By Theorem 3.4, the solution B of the HYBE for the Hom-algebra H_4 satisfies:

$$\begin{aligned}
B(1 \otimes 1) &= \lambda 1 \otimes 1, & B(1 \otimes g) &= \lambda g \otimes g + \nu 1 \otimes 1 - \nu 1 \otimes g, \\
B(1 \otimes x) &= \lambda kx \otimes 1 + \lambda kg \otimes x - \nu k1 \otimes x, & B(1 \otimes y) &= \lambda ky \otimes g + \lambda k1 \otimes y - \nu k1 \otimes y, \\
B(g \otimes 1) &= \lambda 1 \otimes 1 + \nu g \otimes g - \nu g \otimes 1, & B(g \otimes g) &= \lambda g \otimes g, \\
B(g \otimes x) &= \lambda kx \otimes 1 + \lambda kg \otimes x - \nu kg \otimes x, & B(g \otimes y) &= \lambda ky \otimes g + \nu k1 \otimes y - \nu kg \otimes y, \\
B(x \otimes 1) &= \nu kg \otimes x, & B(x \otimes g) &= \nu kx \otimes 1 + \nu kg \otimes x - \nu kx \otimes g, \\
B(x \otimes x) &= -k^2 x \otimes x, & B(x \otimes y) &= -k^2 x \otimes y, & B(y \otimes 1) &= \nu ky \otimes g + \nu k1 \otimes y - \nu ky \otimes 1, \\
B(y \otimes g) &= \nu ky \otimes g + \nu k1 \otimes y - \nu ky \otimes g, & B(y \otimes x) &= -k^2 y \otimes x, & B(y \otimes y) &= -k^2 y \otimes y,
\end{aligned}$$

where λ, ν are two parameters in k .

4 Solutions of the HYBE from Hom-Lie algebras

In this section, we obtain a new solution of the HYBE from Hom-Lie algebras and prove that this solution is self-inverse.

Theorem 4.1. Let $(L, [\cdot, \cdot], \alpha)$ be a Hom-Lie algebra, u is an α -invariant element in $Z(L)$ and $\lambda, \nu \in k$. Then

$$B : L \otimes L \rightarrow L \otimes L, \quad x \otimes y \mapsto \lambda[x, y] \otimes u - \nu\alpha(y) \otimes \alpha(x)$$

is a solution for HYBE, where $Z(L) = \{u \in L \mid [u, x] = 0, \forall x \in L\}$.

Proof. Obviously, B is compatible with the twist map α since z is α -invariant. Now we verify that B satisfy the HYBE. In fact, for any $x, y, z \in L$, one may directly check that $B(x \otimes u) = -\nu u \otimes \alpha(x)$, $B(u \otimes x) = -\nu\alpha(x) \otimes u$. On the one hand, we have

$$\begin{aligned} & (\alpha \otimes B)(x \otimes y \otimes z) \\ &= \lambda\alpha(x) \otimes [y, z] \otimes u - \nu\alpha(x) \otimes \alpha(z) \otimes \alpha(y), \\ & (B \otimes \alpha) \circ (\alpha \otimes B)(x \otimes y \otimes z) \\ &= \lambda B(\alpha(x) \otimes [y, z]) \otimes u - \nu B(\alpha(x) \otimes \alpha(z)) \otimes \alpha^2(y) \\ &= \lambda^2[\alpha(x), [y, z]] \otimes u \otimes u - \lambda\nu[\alpha(y), \alpha(z)] \otimes \alpha^2(x) \otimes u \\ &\quad - \lambda\nu[\alpha(x), \alpha(z)] \otimes u \otimes \alpha^2(y) + \nu^2\alpha^2(z) \otimes \alpha^2(x) \otimes \alpha^2(y), \\ & (\alpha \otimes B) \circ (B \otimes \alpha) \circ (\alpha \otimes B)(x \otimes y \otimes z) \\ &= \lambda^2[\alpha^2(x), [\alpha(y), \alpha(z)]] \otimes B(u \otimes u) - \lambda\nu[\alpha^2(y), \alpha^2(z)] \otimes B(\alpha^2(x) \otimes u) \\ &\quad - \lambda\nu[\alpha^2(x), \alpha^2(z)] \otimes B(u \otimes \alpha^2(y)) + \nu^2\alpha^3(z) \otimes B(\alpha^2(x) \otimes \alpha^2(y)) \\ &= \underline{-\lambda^2\nu[\alpha^2(x), [\alpha(y), \alpha(z)]] \otimes u \otimes u} + \lambda\nu^2[\alpha^2(y), \alpha^2(z)] \otimes u \otimes \alpha^3(x) \\ &\quad + \lambda\nu^2[\alpha^2(x), \alpha^2(z)] \otimes \alpha^3(y) \otimes u + \lambda\nu^2\alpha^3(z) \otimes [\alpha^2(x), \alpha^2(y)] \otimes u \\ &\quad - \nu^3\alpha^3(z) \otimes \alpha^3(y) \otimes \alpha^3(x). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} & (B \otimes \alpha)(x \otimes y \otimes z) \\ &= \lambda[x, y] \otimes u \otimes \alpha(z) - \nu\alpha(y) \otimes \alpha(x) \otimes \alpha(z), \\ & (\alpha \otimes B) \circ (B \otimes \alpha)(x \otimes y \otimes z) \\ &= \lambda[\alpha(x), \alpha(y)] \otimes B(u \otimes \alpha(z)) - \nu\alpha^2(y) \otimes B(\alpha(x) \otimes \alpha(z)) \\ &= -\lambda\nu[\alpha(x), \alpha(y)] \otimes \alpha^2(z) \otimes u - \lambda\nu\alpha^2(y) \otimes [\alpha(x), \alpha(z)] \otimes u \\ &\quad + \nu^2\alpha^2(y) \otimes \alpha^2(z) \otimes \alpha^2(x), \\ & (B \otimes \alpha) \circ (\alpha \otimes B) \circ (B \otimes \alpha)(x \otimes y \otimes z) \\ &= -\lambda\nu B([\alpha(x), \alpha(y)] \otimes \alpha^2(z)) \otimes u - \lambda\nu B(\alpha^2(y) \otimes [\alpha(x), \alpha(z)]) \otimes u \\ &\quad + \nu^2 B(\alpha^2(y) \otimes \alpha^2(z)) \otimes \alpha^3(x) \\ &= \underline{-\lambda^2\nu[[\alpha(x), \alpha(y)], \alpha^2(z)] \otimes u \otimes u} + \lambda\nu^2\alpha^3(z) \otimes [\alpha^2(x), \alpha^2(y)] \otimes u \\ &\quad - \underline{\lambda^2\nu[\alpha^2(y), [\alpha(x), \alpha(z)]] \otimes u \otimes u} + \lambda\nu^2[\alpha^2(x), \alpha^2(z)] \otimes \alpha^3(y) \otimes u \\ &\quad + \lambda\nu^2[\alpha^2(y), \alpha^2(z)] \otimes u \otimes \alpha^3(x) - \nu^3\alpha^3(z) \otimes \alpha^3(y) \otimes \alpha^3(x). \end{aligned}$$

According to the anti-symmetry and the Hom-Jacobi identity, we have

$$[\alpha^2(x), [\alpha(y), \alpha(z)]] = [[\alpha(x), \alpha(y)], \alpha^2(z)] + [\alpha^2(y), [\alpha(x), \alpha(z)]].$$

It follows that

$$(\alpha \otimes B) \circ (B \otimes \alpha) \circ (\alpha \otimes B)(x \otimes y \otimes z) = (B \otimes \alpha) \circ (\alpha \otimes B) \circ (B \otimes \alpha)(x \otimes y \otimes z).$$

That is, B is a solution for HYBE. \square

Corollary 4.2. Let $(L, [\cdot, \cdot], \alpha)$ be a Hom-Lie algebra, u is an α -invariant element in $Z(L)$ and $\lambda \in k$. Then

$$B : L \otimes L \rightarrow L \otimes L, \quad x \otimes y \mapsto \lambda[x, y] \otimes u - \alpha(y) \otimes \alpha(x)$$

is a solution for HYBE. If, in addition, α is involutive, then B is also invertible, where the inverse is given by

$$B^{-1} : L \otimes L \rightarrow L \otimes L, \quad x \otimes y \mapsto \lambda u \otimes [x, y] - \alpha(y) \otimes \alpha(x).$$

Furthermore, B^{-1} is also a solution for HYBE.

Proof. First, it is easy to see that B is a solution for HYBE by setting $\nu = 1$.

Next, show that $B \circ B^{-1} = id_{L \otimes L}$. In fact, for any $x, y, z \in L$, we have

$$\begin{aligned} B \circ B^{-1}(x \otimes y) &= \lambda B(u \otimes [x, y]) - B(\alpha(y) \otimes \alpha(x)) \\ &= -\lambda[\alpha(x), \alpha(y)] \otimes u - \lambda[\alpha(y), \alpha(x)] \otimes u + \alpha^2(y) \otimes \alpha^2(x) \\ &= x \otimes y. \end{aligned}$$

It follows that $B \circ B^{-1} = id_{L \otimes L}$. Similarly, one may check that $B^{-1} \circ B = id_{L \otimes L}$. So B^{-1} is the inverse of B . Further, similar to the proof of Theorem 4.1, one may calculate

$$\begin{aligned} &(\alpha \otimes B^{-1}) \circ (B^{-1} \otimes \alpha) \circ (\alpha \otimes B^{-1})(x \otimes y \otimes z) \\ &= -\lambda^2 u \otimes u \otimes [[\alpha(x), \alpha(y)], \alpha^2(z)] + \lambda \alpha^3(z) \otimes u \otimes [\alpha^2(x), \alpha^2(y)] \\ &\quad + \lambda u \otimes \alpha^3(y) \otimes [\alpha^2(x), \alpha^2(z)] + \lambda u \otimes [\alpha^2(y), \alpha^2(z)] \otimes \alpha^3(x) \\ &\quad - \alpha^3(z) \otimes \alpha^3(y) \otimes \alpha^3(x) \\ &= (B^{-1} \otimes \alpha) \circ (\alpha \otimes B^{-1}) \circ (B^{-1} \otimes \alpha)(x \otimes y \otimes z). \end{aligned}$$

So B^{-1} is a solution for HYBE.

Example 4.3. Let $(L, [\cdot, \cdot], \alpha)$ be a Hom-Lie algebra on 3-dimensional Euclidean E^3 with basis elements $\{e_1, e_2, e_3\}$, whose bracket $[\cdot, \cdot]$ is given by

$$[e_1, e_2] = e_1, \quad [e_1, e_3] = 0, \quad [e_2, e_3] = 0.$$

The twist map α is given by

$$\alpha(e_1) = e_1, \quad \alpha(e_2) = e_2, \quad \alpha(e_3) = -e_3.$$

Obviously, $e_3 \in Z(L)$. Therefore, by Theorem 4.1, the solution B of the HYBE for the Hom-Lie algebra L satisfies:

$$\begin{aligned} B(e_1 \otimes e_1) &= -\nu e_1 \otimes e_1, \quad B(e_1 \otimes e_2) = \lambda e_1 \otimes e_3 - \nu e_2 \otimes e_1, \quad B(e_1 \otimes e_3) = \nu e_3 \otimes e_1, \\ B(e_2 \otimes e_1) &= -\lambda e_1 \otimes e_3 - \nu e_1 \otimes e_2, \quad B(e_2 \otimes e_2) = -\nu e_2 \otimes e_2, \quad B(e_2 \otimes e_3) = \nu e_3 \otimes e_2, \\ B(e_3 \otimes e_1) &= \nu e_1 \otimes e_3, \quad B(e_3 \otimes e_2) = \nu e_2 \otimes e_3, \quad B(e_3 \otimes e_3) = -\nu e_3 \otimes e_3, \end{aligned}$$

where $\lambda, \nu \in k$.

In the final part of this section, we present a kind of solutions of CHYBE from Hom-Lie algebras. Recall from [32], Yau defined the CHYBE in a Hom-Lie algebra $(L, [\cdot, \cdot], \alpha)$ as

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{33}] = 0,$$

for $r \in L^{\otimes 2}$. Here the three brackets above are defined as

$$\begin{aligned} [r^{12}, r^{13}] &= [a_i, a_j] \otimes \alpha(b_i) \otimes \alpha(b_j), \\ [r^{12}, r^{23}] &= \alpha(a_i) \otimes [b_i, a_k] \otimes \alpha(a_k), \\ [r^{13}, r^{23}] &= \alpha(a_j) \otimes \alpha(a_k) \otimes [b_j, j_k], \end{aligned}$$

where $r^{12} = r \otimes 1 = a_i \otimes b_i \otimes 1$, $r^{13} = (\tau \otimes id)(1 \otimes r) = a_j \otimes 1 \otimes b_j$, $r^{23} = 1 \otimes r = 1 \otimes a_k \otimes b_k$.

Theorem 4.4. Let $(L, [\cdot, \cdot], \alpha)$ be a Hom-Lie algebra and u be an element in $Z(L)$. Then for any $x, y \in L$ and $m, n \in \mathbb{Z}$,

$$r = \alpha^m([x, y]) \otimes \alpha^n(u)$$

is a solution for CHYBE.

Proof. It is easy to see that $[r^{12}, r^{13}] = [r^{12}, r^{23}] = [r^{13}, r^{23}] = 0$ since u is an element in $Z(L)$.

5 Hom-Yang-Baxter systems

In this section, we extend the notion of Yang-Baxter systems to Hom-Yang-Baxter systems and present two kinds of Hom-Yang-Baxter systems.

Consider three vector spaces V, V', V'' , let $\alpha_V, \alpha_{V'}, \alpha_{V''}$ be three endomorphisms on V, V', V'' and $R : V \otimes V' \rightarrow V \otimes V', S : V \otimes V'' \rightarrow V \otimes V'', T : V' \otimes V'' \rightarrow V' \otimes V''$ be three linear maps. Then a Hom-Yang-Baxter commutator is a map $[R, S, T] : V \otimes V' \otimes V'' \rightarrow V \otimes V' \otimes V''$ defined by

$$[R, S, T] = R^{12} \circ S^{13} \circ T^{23} - T^{23} \circ S^{13} \circ R^{12},$$

where $R^{12} = R \otimes \alpha_V \nu$, $S^{13} = (\tau \otimes id) \circ (\alpha_{V'} \otimes T) \circ (\tau \otimes id)$, $T^{23} = \alpha_V \otimes T$.

Definition 5.1. Let V, V' be two vector spaces and $\alpha_V, \alpha_{V'}$ be two endomorphisms. A system of linear maps

$$W : V \otimes V \rightarrow V \otimes V, \quad Z : V' \otimes V' \rightarrow V' \otimes V', \quad X : V \otimes V' \rightarrow V \otimes V'$$

is called a Hom-Yang-Baxter system, if the following conditions are satisfied:

$$[W, W, W] = 0, \quad [Z, Z, Z] = 0, \quad [W, X, X] = 0, \quad [X, X, Z] = 0. \quad (5.1)$$

Theorem 5.2. Let $(A, \mu, 1_A, \alpha)$ be a Hom-algebra and $\lambda, \nu \in k$. Then the following is a Hom-Yang-Baxter system:

$$W : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto ab \otimes 1_A + \lambda 1_A \otimes ab - \alpha(b) \otimes \alpha(a),$$

$$Z : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto \nu ab \otimes 1_A + 1_A \otimes ab - \alpha(b) \otimes \alpha(a),$$

$$X : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto ab \otimes 1_A + 1_A \otimes ab - \alpha(b) \otimes \alpha(a).$$

Proof. It is sufficient to prove that the four equalities in Eq. (5.1) hold. Here we only verify the equality $[W, X, X] = 0$ and similar for other three equalities. In fact, for any $a, b, c \in A$, on the one side, we have

$$\begin{aligned} & W^{12} \circ X^{13} \circ X^{23}(a \otimes b \otimes c) \\ &= W^{12} \circ X^{13} \{ \alpha(a) \otimes bc \otimes 1_A + \alpha(a) \otimes 1_A \otimes bc - \alpha(a) \otimes \alpha(b) \otimes \alpha(c) \} \\ &= W^{12} \{ \alpha^2(a) \otimes \alpha(bc) \otimes 1_A + \alpha(a)(bc) \otimes 1_A \otimes 1_A + 1_A \otimes 1_A \otimes \alpha(a)(bc) \\ &\quad - \alpha(bc) \otimes 1_A \otimes \alpha^2(a) - \alpha(ab) \otimes \alpha^2(c) \otimes 1_A - 1_A \otimes \alpha^2(c) \otimes \alpha(ab) \\ &\quad + a^2(b) \otimes a^2(c) \otimes a^2(a) \} \\ &= \frac{\alpha^2(a)\alpha(bc) \otimes 1_A \otimes 1_A}{(1)} + \frac{\lambda 1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_A}{(2)} - \frac{\alpha^2(bc) \otimes \alpha^3(a) \otimes 1_A}{(3)} \\ &\quad + \frac{\alpha^2(a)\alpha(bc) \otimes 1_A \otimes 1_A}{(4)} + \frac{\lambda 1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_A}{(5)} - \frac{1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_A}{(6)} \\ &\quad + \frac{\lambda 1_A \otimes 1_A \otimes \alpha^2(a)\alpha(bc)}{(7)} - \frac{\alpha^2(bc) \otimes 1_A \otimes \alpha^3(a)}{(8)} - \frac{\lambda 1_A \otimes \alpha^2(bc) \otimes \alpha^3(a)}{(9)} \\ &\quad + \frac{1_A \otimes \alpha^2(bc) \otimes \alpha^3(a)}{(10)} - \frac{\alpha(ab)\alpha^2(c) \otimes 1_A \otimes 1_A}{(11)} - \frac{\lambda 1_A \otimes \alpha(ab)\alpha^2(c) \otimes 1_A}{(12)} \\ &\quad + \frac{\alpha^3(c) \otimes \alpha^2(ab) \otimes 1_A}{(13)} - \frac{\lambda 1_A \otimes \alpha^3(c) \otimes \alpha^2(ab)}{(14)} + \frac{\alpha^2(bc) \otimes 1_A \otimes \alpha^3(a)}{(15)} \\ &\quad + \frac{\lambda 1_A \otimes \alpha^2(bc) \otimes \alpha^3(a)}{(16)} - \frac{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}{(17)} \\ &= \frac{\alpha^2(a)\alpha(bc) \otimes 1_A \otimes 1_A}{(1)} + \frac{\lambda 1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_A}{(2)} - \frac{\alpha^2(bc) \otimes \alpha^3(a) \otimes 1_A}{(3)} \\ &\quad - \frac{1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_A}{(6)} + \frac{\lambda 1_A \otimes 1_A \otimes \alpha^2(a)\alpha(bc)}{(7)} + \frac{1_A \otimes \alpha^2(bc) \otimes \alpha^3(a)}{(10)} \\ &\quad + \frac{\alpha^3(c) \otimes \alpha^2(ab) \otimes 1_A}{(13)} - \frac{\lambda 1_A \otimes \alpha^3(c) \otimes \alpha^2(ab)}{(14)} - \frac{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}{(17)}. \end{aligned}$$

The last equality holds since (4)=(11), (5)=(12), (8)=(15) and (9)=(16).

On the other side, we have

$$X^{23} \circ X^{13} \circ W^{12}(a \otimes b \otimes c)$$

$$\begin{aligned}
&= X^{23} \circ X^{13} \{ ab \otimes 1_A \otimes \alpha(c) + \lambda 1_A \otimes ab \otimes \alpha(c) - \alpha(b) \otimes \alpha(a) \otimes \alpha(c) \} \\
&= X^{23} \{ (ab)\alpha(c) \otimes 1_A \otimes 1_A + 1_A \otimes 1_A \otimes (ab)\alpha(c) - \alpha^2(c) \otimes 1_A \otimes \alpha(ab) \\
&\quad + \lambda 1_A \otimes \alpha(ab) \otimes \alpha^2(c) - \alpha(bc) \otimes \alpha^2(a) \otimes 1_A - 1_{\otimes} \alpha^2(a) \otimes \alpha(bc) \\
&\quad + \alpha^2(c) \otimes \alpha^2(a) \otimes \alpha^2(b) \} \\
&= \frac{\alpha(ab)\alpha^2(c) \otimes 1_A \otimes 1_A}{(18)} + \frac{1_A \otimes 1_A \otimes \alpha(ab)\alpha^2(c)}{(19)} - \frac{\alpha^3(c) \otimes 1_A \otimes \alpha^2(ab)}{(20)} \\
&\quad + \frac{\lambda 1_A \otimes \alpha(ab) \otimes \alpha^2(c)}{(21)} + \frac{\lambda 1_A \otimes 1_A \otimes \alpha(ab)\alpha^2(c)}{(22)} - \frac{\lambda 1_A \otimes \alpha^3(c) \otimes \alpha^2(ab)}{(23)} \\
&\quad - \frac{a^2(bc) \otimes \alpha^3(a) \otimes 1_A}{(24)} - \frac{1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_A}{(25)} - \frac{1_A \otimes 1_A \otimes \alpha^2(a)\alpha(bc)}{(26)} \\
&\quad + \frac{1_A \otimes \alpha^2(bc) \otimes \alpha^3(a)}{(27)} + \frac{\alpha^3(c) \otimes \alpha^2(ab) \otimes 1_A}{(28)} + \frac{\alpha^3(c) \otimes 1_A \otimes \alpha^2(ab)}{(29)} \\
&\quad - \frac{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}{(30)} \\
&= \frac{\alpha(ab)\alpha^2(c) \otimes 1_A \otimes 1_A}{(18)} + \frac{\lambda 1_A \otimes \alpha(ab) \otimes \alpha^2(c)}{(21)} + \frac{\lambda 1_A \otimes 1_A \otimes \alpha(ab)\alpha^2(c)}{(22)} \\
&\quad - \frac{\lambda 1_A \otimes \alpha^3(c) \otimes \alpha^2(ab)}{(23)} - \frac{a^2(bc) \otimes \alpha^3(a) \otimes 1_A}{(24)} - \frac{1_A \otimes \alpha^2(a)\alpha(bc) \otimes 1_A}{(25)} \\
&\quad + \frac{1_A \otimes \alpha^2(bc) \otimes \alpha^3(a)}{(27)} + \frac{\alpha^3(c) \otimes \alpha^2(ab) \otimes 1_A}{(28)} - \frac{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}{(30)}.
\end{aligned}$$

The last equality holds since (19)=(26) and (20)=(29).

Compare the two expressions above, we have

$$\begin{aligned}
(1) &= (18), (2) = (21), (3) = (24), (6) = (25), \\
(7) &= (22), (10) = (27), (13) = (28), (14) = (23), (17) = (30).
\end{aligned}$$

It follows that $[W, X, X] = 0$, as desired. The proof is finished. \square

Theorem 5.3. Let $(C, \Delta, \varepsilon, \alpha)$ be a Hom-coalgebra and $\lambda, \nu \in k$. Then the following is a Hom-Yang-Baxter system:

$$\begin{aligned}
W : C \otimes C &\rightarrow C \otimes C, \quad a \otimes b \mapsto \lambda \varepsilon(a) b_1 \otimes b_2 + \varepsilon(b) a_1 \otimes a_2 - \alpha(b) \otimes \alpha(a), \\
Z : C \otimes C &\rightarrow C \otimes C, \quad a \otimes b \mapsto \varepsilon(a) b_1 \otimes b_2 + \nu \varepsilon(b) a_1 \otimes a_2 - \alpha(b) \otimes \alpha(a), \\
X : C \otimes C &\rightarrow C \otimes C, \quad a \otimes b \mapsto \varepsilon(a) b_1 \otimes b_2 + \varepsilon(b) a_1 \otimes a_2 - \alpha(b) \otimes \alpha(a).
\end{aligned}$$

Proof. We only prove $[X, X, Z] = 0$ and similar for other three equalities. In fact, for any $a, b, c \in C$, on the one side, we have

$$\begin{aligned}
&Z^{23} \circ X^{13} \circ X^{12}(a \otimes b \otimes c) \\
&= Z^{23} \circ X^{13} \{ \varepsilon(a) b_1 \otimes b_2 \otimes \alpha(c) + \nu \varepsilon(b) a_1 \otimes a_2 \otimes \alpha(c) - \alpha(b) \otimes \alpha(a) \otimes \alpha(c) \} \\
&= Z^{23} \{ \varepsilon(a) \alpha(c_1) \otimes \alpha^2(b) \otimes \alpha(c_2) + \varepsilon(a) \varepsilon(c) b_{11} \otimes \alpha(b_2) \otimes b_{12} \\
&\quad - \varepsilon(a) \alpha^2(c) \otimes \alpha(b_2) \otimes \alpha(b_1) + \frac{\varepsilon(b) \alpha(c_1) \otimes \alpha^2(a) \otimes \alpha(c_2)}{} \\
&\quad + \varepsilon(b) \varepsilon(c) a_{11} \otimes \alpha(a_2) \otimes a_{12} - \varepsilon(b) \alpha^2(c) \otimes \alpha(a_2) \otimes \alpha(a_1) \\
&\quad - \frac{\varepsilon(b) \alpha(c_1) \otimes \alpha^2(a) \otimes \alpha(c_2)}{} - \varepsilon(c) \alpha(b_1) \otimes \alpha^2(a) \otimes \alpha(b_2) \\
&\quad + \alpha^2(c) \otimes \alpha^2(a) \otimes \alpha^2(b) \} \\
&= \frac{\varepsilon(a) \varepsilon(b) \alpha^2(c_1) \otimes \alpha(c_{21}) \otimes \alpha(c_{22})}{(1)} + \frac{\nu \varepsilon(a) \alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}{(2)}
\end{aligned}$$

$$\begin{aligned}
& - \underline{\varepsilon(a)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(b)}_{(3)} + \underline{\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}_{(4)} \\
& + \underline{\nu\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}_{(5)} - \underline{\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}_{(6)} \\
& - \underline{\varepsilon(a)\alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}_{(7)} - \underline{\nu\varepsilon(a)\alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}_{(8)} \\
& + \underline{\varepsilon(a)\alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}_{(9)} + \underline{\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}_{(10)} \\
& + \underline{\nu\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}_{(11)} - \underline{\varepsilon(b)\varepsilon(c)\alpha(a_{11}) \otimes \alpha(a_{12}) \otimes \alpha^2(a_2)}_{(12)} \\
& - \underline{\varepsilon(b)\alpha^3(c) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(13)} - \underline{\nu\varepsilon(b)\alpha^3(c) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(14)} \\
& + \underline{\varepsilon(b)\alpha^3(c) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(15)} - \underline{\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}_{(16)} \\
& - \underline{\nu\varepsilon(c)\alpha^3(b) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(17)} + \underline{\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(18)} \\
& + \underline{\varepsilon(a)\alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}_{(19)} + \underline{\nu\varepsilon(b)\alpha^3(c) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(20)} \\
& - \underline{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}_{(21)} \\
= & \underline{\varepsilon(a)\varepsilon(b)\alpha^2(c_1) \otimes \alpha(c_{21}) \otimes \alpha(c_{22})}_{(1)} - \underline{\varepsilon(a)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(b)}_{(3)} \\
& + \underline{\nu\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}_{(5)} + \underline{\varepsilon(a)\alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}_{(9)} \\
& + \underline{\nu\varepsilon(b)\varepsilon(c)\alpha^2(a_1) \otimes \alpha(a_{21}) \otimes \alpha(a_{22})}_{(11)} - \underline{\varepsilon(a)\varepsilon(c)\alpha^2(b_1) \otimes \alpha(b_{21}) \otimes \alpha(b_{22})}_{(16)} \\
& - \underline{\nu\varepsilon(c)\alpha^3(b) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(17)} + \underline{\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(18)} \\
& - \underline{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}_{(21)}.
\end{aligned}$$

The last equality holds since (2)=(8), (4)=(6), (7)=(9),(10)=(12),(13)=(15) and (14)=(20).

On the other side, we have

$$\begin{aligned}
& X^{12} \circ X^{13} \circ Z^{23}(a \otimes b \otimes c) \\
= & X^{12} \circ X^{13} \{ \varepsilon(b)\alpha(a) \otimes c_1 \otimes c_2 + \nu\varepsilon(c)\alpha(a) \otimes b_1 \otimes b_2 - \alpha(a) \otimes \alpha(c) \otimes \alpha(b) \} \\
= & X^{12} \{ \varepsilon(a)\varepsilon(b)c_{21} \otimes \alpha(c_1) \otimes c_{22} + \underline{\varepsilon(b)\alpha(a_1) \otimes \alpha^2(c) \otimes \alpha(a_2)} \\
& - \varepsilon(b)\alpha(c_2) \otimes \alpha(c_1) \otimes \alpha^2(a) + \nu\varepsilon(a)\varepsilon(c)b_{21} \otimes \alpha(b_1) \otimes b_{22} \\
& + \nu\varepsilon(c)\alpha(a_1) \otimes \alpha^2(b) \otimes \alpha(a_2) - \nu\varepsilon(c)\alpha(b_2) \otimes \alpha(b_1) \otimes \alpha^2(a) \\
& - \varepsilon(a)\alpha(b_1) \otimes \alpha^2(c) \otimes \alpha(b_2) - \underline{\varepsilon(b)\alpha(a_1) \otimes \alpha^2(c) \otimes \alpha(a_2)} \\
& + \alpha^2(b) \otimes \alpha^2(c) \otimes \alpha^2(a) \} \\
= & \underline{\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}_{(22)} + \underline{\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}_{(23)} \\
& - \underline{\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}_{(24)} - \underline{\varepsilon(b)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(a)}_{(25)} \\
& - \underline{\varepsilon(b)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(a)}_{(26)} + \underline{\varepsilon(b)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(a)}_{(27)} \\
& + \underline{\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}_{(28)} + \underline{\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}_{(29)} \\
& - \underline{\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}_{(30)} + \underline{\nu\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(31)} \\
& + \underline{\nu\varepsilon(b)\varepsilon(c)\alpha(a_{11}) \otimes \alpha(a_{12}) \otimes \alpha^2(a_2)}_{(32)} - \underline{\nu\varepsilon(c)\alpha^3(b) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(33)} \\
& - \underline{\nu\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(34)} - \underline{\nu\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(35)} \\
& + \underline{\nu\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(36)} - \underline{\varepsilon(a)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(b)}_{(37)} \\
& - \underline{\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}_{(38)} + \underline{\varepsilon(a)\alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}_{(39)}
\end{aligned}$$

$$\begin{aligned}
& + \underline{\varepsilon(b)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(a)}_{(40)} + \underline{\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(41)} \\
& - \underline{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}_{(42)} \\
= & \underline{\varepsilon(a)\varepsilon(b)\alpha(c_{11}) \otimes \alpha(c_{12}) \otimes \alpha^2(c_2)}_{(22)} + \underline{\nu\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}_{(28)} \\
& + \underline{\nu\varepsilon(b)\varepsilon(c)\alpha(a_{11}) \otimes \alpha(a_{12}) \otimes \alpha^2(a_2)}_{(32)} - \underline{\nu\varepsilon(c)\alpha^3(b) \otimes \alpha^2(a_1) \otimes \alpha^2(a_2)}_{(33)} \\
& - \underline{\varepsilon(a)\alpha^2(c_1) \otimes \alpha^2(c_2) \otimes \alpha^3(b)}_{(37)} - \underline{\varepsilon(a)\varepsilon(c)\alpha(b_{11}) \otimes \alpha(b_{12}) \otimes \alpha^2(b_2)}_{(38)} \\
& + \underline{\varepsilon(a)\alpha^3(c) \otimes \alpha^2(b_1) \otimes \alpha^2(b_2)}_{(39)} + \underline{\varepsilon(c)\alpha^2(b_1) \otimes \alpha^2(b_2) \otimes \alpha^3(a)}_{(41)} \\
& - \underline{\alpha^3(c) \otimes \alpha^3(b) \otimes \alpha^3(a)}_{(42)}.
\end{aligned}$$

The last equality holds since (23)=(24), (25)=(40), (26)=(27),(29)=(30),(31)=(34) and (35)=(36). Compare the two expressions above, we have

$$\begin{aligned}
(1) & = (22), (3) = (37), (5) = (28), (9) = (39), \\
(11) & = (32), (17) = (33), (18) = (41), (16) = (38), (21) = (42).
\end{aligned}$$

It follows that $[X, X, Z] = 0$, as desired. The proof is finished. \square

ACKNOWLEDGEMENT

The paper is supported by the NSF of China (Nos. 11761017 and 11801304), Guizhou Provincial Science and Technology Foundation (No. [2020]1Y005), the Anhui Provincial Natural Science Foundation (No. 1908085MA03) and the Key University Science Research Project of Anhui Province (No. KJ2020A0711).

Data Availability Statement Our manuscripts does not include a data availability statement.

REFERENCES

- [1] R. J. Baxter. Partition function of the eight-vertex lattice model. *Ann. Phys.* 1972, 70(1):193-228.
- [2] Y. Chen, Z. Wang, L. Zhang. Quasitriangular Hom-Hopf algebras. *Colloq. Math.* 2014, 137(1):67-88.
- [3] Y. Chen, L. Zhang. The category of Yetter-Drinfel'd Hom-modules and the quantum Hom-Yang-Baxter equation. *J. Math. Phys.* 2014, 55(3):031702.
- [4] Y. Chen, X. Zhou. Separable and Frobenius monoidal Hom-algebras. *Colloq. Math.* 2014, 137(2):229-251.
- [5] X. Fang, W. Liu. Solutions of the BiHom-Yang-Baxter equation. *Sbornik: Mathematics* 2018, 209(6):901-918.

- [6] A. Gohr. On hom-algebras with surjective twisting. *J. Algebra* 2010, 324(7):1483-1491.
- [7] J. Hartwig, D. Larsson, S. Silvestrov. Deformations of Lie algebras using σ -derivations. *J. Algebra* 2006, 295(2):314-361.
- [8] N. Hu. q-Witt algebras, q-Lie algebras, q-holomorph structure and representations. *Algebr. Colloq* 1999, 6(1):51-70.
- [9] H. Li, T. Ma. A construction of the Hom-Yetter-Drinfeld category. *Colloq. Math.* 2014, 137(1):43-65.
- [10] L. Liu, B. Shen. Radford's biproducts and Yetter-Drinfeld modules for monoidal Hom-Hopf algebras. *J. Math. Phys.* 2014, 55(3):031701.
- [11] D. Lu, Y. Li, S. Guo. Crossed products of Hom-Hopf algebras. *Filomat* 2020, 34(4):1295-1313.
- [12] T. Ma, H. Li, T. Yang. Cobraided smash product Hom-Hopf algebras. *Colloq. Math.* 2014, 134(1): 75-92.
- [13] T. Ma, Y. Wang, L. Liu. Generalized Radford biproduct Hom-Hopf algebras and related braided tensor categories. *J. Math.* 2017, 37(6): 1161-1172.
- [14] T. Ma, H. Yang, L. Liu, Q. Chen. On unified Hom-Yetter-Drinfeld categories. *J. Geom. Phys.* 2019, 144(7): 81-107.
- [15] A. Makhlouf, S. Silvestrov. Hom-algebra structures. *J. Gen. Lie Theory Appl.* 2008, 2(2): 51-64.
- [16] A. Makhlouf, S. Silvestrov. Hom-Lie admissible Hom-coalgebras and Hom-Hopf algebras. *J. Gen. Lie Theory in Mathematics, Physics and beyond.* Springer-Verlag, Berlin, 2009, pp. 189-206.
- [17] A. Makhlouf, S. Silvestrov. Hom-algebras and Hom-coalgebras. *J. Algebra Appl.* 2010, 9(4): 553-589.
- [18] A. Makhlouf, F. Panaite. Yetter-Drinfeld modules for Hom-bialgebras. *J. Math. Phys.* 2014, 55(1): 013501.
- [19] F. F. Nichita. Self-inverse Yang-Baxter operators from (co)algebra structures. *J. Algebra* 1999, 218(2): 738-759.
- [20] F. F. Nichita, D. Parashar. Spectral-parameter dependent Yang-Baxter operators and Yang-Baxter systems from algebra structures. *Commun. Algebra* 2006, 34(6): 2713-2726.

- [21] F. F. Nichita, B. P. Popovici. Some results on the Yang-Baxter equations and applications. *Romanian Journal of Physics* 2010, 63(3): 641-650.
- [22] M. E. Sweedler. Hopf algebras. *Benjamin*, New York, 1969.
- [23] S. X. Wang, S. J. Guo. Symmetries and the u-condition in Hom-Yetter-Drinfeld categories. *J. Math. Phys.* 2014, 55(8): 081708.
- [24] S. X. Wang, S. J. Guo. Symmetric pairs and pseudosymmetries in Hom-Yetter-Drinfeld categories. *J. Algebra Appl.* 2017, 16(7): 1750125.
- [25] S. Wang, S. Guo. BiHom-Lie superalgebra structures and BiHom-Yang-Baxter equations. *Adv. Appl. Clifford Algebr.* 2020, 30(3), Art. 35, 18 pp.
- [26] Z. Wang, Y. Chen, L. Zhang. The antipode and Drinfel'd double of Hom-Hopf algebras. *Sci. Sin. Math.* 2012, 42(11): 1079-1093.
- [27] D. Yan, S. Wang, T. Ma. Four-angle Hopf modules for Hom-Hopf algebras. *arXiv:2006.15267v1*.
- [28] C. N. Yang. Some exact results for the many-body problem in one dimension with repulsive delta-function interaction. *Phys. Rev. Lett.* 1967, 19(23): 1312-1315.
- [29] D. Yau. The Hom-Yang-Baxter equation, Hom-Lie algebras, and quasi-triangular bialgebras. *J. Phys. A* 2009, 42(16): 165202.
- [30] D. Yau. The Hom-Yang-Baxter equation and Hom-Lie algebras. *J. Math. Phys.* 2011, 52(5): 053502.
- [31] D. Yau. Hom-quantum groups I: Quasi-triangular Hom-bialgebras. *J. Phys. A* 2012, 45(6):065203.
- [32] D. Yau. The classical Hom-Yang-Baxter equation and Hom-Lie bialgebras. *Int. Electron. J. Algebra* 2015, 17(17): 11-45.
- [33] M. You, S. Wang. Constructing new braided T -categories over monoidal Hom-Hopf algebras. *J. Math. Phys.* 2014, 55(11):111701.
- [34] X. Zhang, S. Guo, S. Wang. Drinfeld codoubles of Hom-Hopf algebras. *Adv. Appl. Clifford Algebr.* 2019, 29(2), Art. 36, 26 pp.
- [35] X. Zhang, W. Wang, X. Zhao. Smash coproducts of monoidal comonads and Hom-entwining structures. *Rocky Mountain J. Math.* 2019, 49(6): 2063-2105.