# Mean Test with Fewer Observation than Dimension and Ratio Unbiased Estimator for Correlation Matrix

Tiefeng Jiang

Ping Li

School of Statistics University of Minnesota 224 Church St SE Minneapolis, MN 55455 jiang040@umn.edu Cognitive Computing Lab Baidu Research 10900 NE 8th St. Bellevue, WA 98004 liping11@baidu.com

#### Abstract

Hotelling's T-squared test is a classical tool to test if the normal mean of a multivariate normal distribution is a specified one or the means of two multivariate normal means are equal. When the population dimension is higher than the sample size, the test is no longer applicable. Under this situation, in this paper we revisit the tests proposed by Srivastava and Du (2008), who revise the Hotelling's statistics by replacing Wishart matrices with their diagonal matrices. They show the revised statistics are asymptotically normal. We use the random matrix theory to examine their statistics again and find that their discovery is just part of the big picture. In fact, we prove that their statistics, decided by the Euclidean norm of the population correlation matrix, can go to normal, mixing chi-squared distributions and a convolution of both. Examples are provided to show the phase transition phenomenon between the normal and mixing chi-squared distributions. The second contribution of ours is a rigorous derivation of an asymptotic ratio-unbiased-estimator of the squared Euclidean norm of the correlation matrix.

## 1 Introduction

Among many statistical hypothesis testing problems, the Hotelling's  $T^2$  tests (Hotelling, 1931) are classic ones to study if the mean of a multivariate normal distribution is equal to the given one, or if the means of two normal distributions are equal. Let us quickly review the two problems. First, assume  $\{X_1, \dots, X_n\}$  is a random sample from a *p*-dimensional normal distribution  $N_p(\mu, \Sigma)$ , where  $\mu$  is the mean vector and  $\Sigma$  is the covariance matrix. Consider the test  $H_0: \mu = \mathbf{0}$  versus  $H_1: \mu \neq \mathbf{0}$ . The Hotelling  $T^2$  test statistic is defined by  $T^2 = (\bar{X} - \mu)^T \hat{\mathbf{S}}^{-1}(\bar{X} - \mu)$ , where  $\bar{X}$  and  $\hat{\mathbf{S}}$  are the sample mean and the sample covariance matrix. For the two population test, assume that  $\{X_{i1}, \dots, X_{in_i}\}$  for i = 1, 2 are two independent random samples from  $N_p(\mu_1, \Sigma)$  and  $N_p(\mu_2, \Sigma)$ , respectively. We aim to test  $H_0: \mu_1 = \mu_2$  versus  $H_1: \mu_1 \neq \mu_2$ . The corresponding Hotelling's  $T^2$  test statistic is given by  $\frac{n_1n_2}{n_1+n_2}(\bar{X}_1 - \bar{X}_2)^T \hat{\mathbf{S}}^{-1}(\bar{X}_1 - \bar{X}_2)$  where  $\bar{X}_i$  is the sample mean vector of the *i*-th sample and  $\hat{\mathbf{S}}$  is the pooled sample covariance matrix.

As a consequence of the likelihood ratio test, the above two Hotelling  $T^2$  tests have very nice properties. For example, for the one-sample test,  $T^2$  is invariant under linear transformations. Also,  $T^2$  is a uniformly most powerful test under the general linear group; see, for example, p. 211 from Muirhead (1982) or p. 190 from Anderson (2003).

Despite its important role in classical statistics, Hotelling's  $T^2$  have some limitations. First, in order to guarantee that the sample covariance matrix is invertible, the sample size have to be larger than the population dimension. These are not true for some modern data in which the population dimension is larger or even much larger than the sample size. For example, for DNA microarray data, thousands of gene expression levels are often measured on a small number of subjects. From http://genomics-pubs.princeton.edu/oncology/, which is a popular colon dataset, one can see p = 2000,  $n_1 = 22$  and  $n_2 = 40$ . Second, as Bai and Saranadasa (1996) have showed, the Hotelling two-sample test is inconsistent as the population dimension and the sample size are comparable.

To accommodate data with the feature of large p small n for the above one-sample and two-sample testing problems, Dempster (1958, 1960) study the so-called "non-exact" test. Bai and Saranadasa (1996) construct a new test statistic by removing the inverse matrix from the definition of  $T^2$  given earlier. Starting from this century, investigators begin to conceive new test statistics. Srivastava and Du (2008), Srivastava (2009), Chen and Qin (2010), and Dong et al. (2016) replace the inverse of the sample covariance matrix by mild quantities. Lopes et al. (2011) and Srivastava et al. (2016) use a sort of dimension reduction method to lower the population dimension and then use Hotelling's  $T^2$  tests. For data with a certain of sparse nature, Cai et al. (2014) establish tests by measuring the maximum componentwise mean difference of appropriately transformed observations. Other research related to sparsity can be seen from Zhong et al. (2013), Chen et al. (2014), Wang et al. (2015), Gregory et al. (2015) and Guo and Chen (2016). The researchers Biswas and Ghosh (2014), Chang et al. (2017), Chakraborty and Chaudhuri (2017), and Xue and Yao (2020) consider the above test under non-normal assumptions. Other contributions include Park and Ayyala (2013) and Wu et al. (2006) for the consideration of scale-invariant tests and Gretton et al. (2012) for a kernel-based discrepancy measure. Feng et al. (2016) work on a two-sample location problem via a multivariate-sign-based high-dimensional test. Xu et al. (2016) and Zhang et al. (2020a) construct test statistics for the two-sample test problem by  $l_q$ -norm. Recently, built upon Chen et al. (2011), the authors (Li et al., 2020) investigate the Hotelling's  $T^2$  by a ridge-regularized method. Zhang et al. (2020b) extend the tests by Wu et al. (2006) and Zhang et al. (2020a) through a modification of the test by Srivastava and Du (2008).

In this paper, we will revisit the procedure by Srivastava and Du (2008) for the onesample and two-sample problems of testing means. We will briefly review them next and state our motivation. Let  $\{X_1, \dots, X_n\}$  be a random sample from a *p*-dimensional normal distribution  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ with correlation matrix **R**. The sample mean and the sample covariance matrix defined by

$$\bar{\boldsymbol{X}} = \frac{1}{n} \sum_{i=1}^{n} \boldsymbol{X}_{i} \text{ and } \hat{\boldsymbol{S}} = \frac{1}{n} \sum_{i=1}^{n} (\boldsymbol{X}_{i} - \bar{\boldsymbol{X}}) (\boldsymbol{X}_{i} - \bar{\boldsymbol{X}})^{T}.$$
(1)

Let  $\hat{\mathbf{D}}$  be the diagonal matrix of  $\hat{\mathbf{S}}$  and  $\hat{\mathbf{R}}$  be the sample correlation matrix defined by  $\hat{\mathbf{R}} = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2}$ . For the testing problem  $H_0: \boldsymbol{\mu} = \mathbf{0}$  vs  $H_1: \boldsymbol{\mu} \neq \mathbf{0}$ , Srivastava and Du (2008) propose the following test statistic.

$$T_{SD} = \frac{n\bar{\mathbf{X}}^T\hat{\mathbf{D}}^{-1}\bar{\mathbf{X}} - p(n-1)(n-3)^{-1}}{\sqrt{2[\operatorname{tr}(\hat{\mathbf{R}}^2) - p^2(n-1)^{-1}]}}.$$
(2)

Under certain conditions, they show  $T_{SD}$  converges to a normal distribution.

For two population case, the test is  $H_0: \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2$  vs  $H_1: \boldsymbol{\mu}_1 \neq \boldsymbol{\mu}_2$ . Assume that  $\{\boldsymbol{X}_{i1}, \dots, \boldsymbol{X}_{in_i}\}$  for i = 1, 2 are two independent random samples from  $N_p(\boldsymbol{\mu}_1, \boldsymbol{\Sigma})$  and  $N_p(\boldsymbol{\mu}_2, \boldsymbol{\Sigma})$ , respectively. Let  $\bar{\boldsymbol{X}}_i$  be the sample mean for the *i*-th sample and  $\hat{\boldsymbol{S}}$  be the pooled sample covariance matrix defined by

$$\hat{\mathbf{S}} = \frac{1}{n_1 + n_2} \Big[ \sum_{j=1}^{n_1} (\mathbf{X}_{1j} - \bar{\mathbf{X}}_1) (\mathbf{X}_{1j} - \bar{\mathbf{X}}_1)^T + \sum_{j=1}^{n_2} (\mathbf{X}_{2j} - \bar{\mathbf{X}}_2) (\mathbf{X}_{2j} - \bar{\mathbf{X}}_2)^T \Big].$$
(3)

Assume  $\hat{\mathbf{D}}$  is the diagonal matrix of  $\hat{\mathbf{S}}$  and  $\hat{\mathbf{R}} = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2}$  is the pooled sample correlation matrix. Srivastava and Du (2008) consider the following statistic defined by

$$T'_{SD} = \frac{\frac{n_1 n_2}{n_1 + n_2} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2)^T \hat{\mathbf{D}}^{-1} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2) - \frac{(n_1 + n_2 - 2)p}{n_1 + n_2 - 4}}{\sqrt{2 \left[ \operatorname{tr}(\hat{\mathbf{R}}^2) - \frac{p^2}{n_1 + n_2 - 2} \right]}}.$$
(4)

Under certain conditions, they show  $T'_{SD}$  converges to a normal distribution. In fact, to improve the convergence speeds, they actually add a term  $c_{p,n} = 1 + \frac{\operatorname{tr}(\hat{\mathbf{R}}^2)}{p^{3/2}}$  under the squared roots in the denominators of  $T_{SD}$  and  $T'_{SD}$ , respectively. Based on their assumptions,  $c_{p,n}$ goes to one. However, we will study a more general case in which  $c_{p,n}$  may not go to one, and sometimes it even goes to infinity. This is the reason we dump the term  $c_{p,n}$  from both  $T_{SD}$  and  $T'_{SD}$ , respectively. Evidently,  $T_{SD}$  and  $T'_{SD}$  have a nice property of scale-invariance, i.e., they are not changed if data are multiplied by a constant. Also, they can be directly computed. Our motivation in this paper to reexamine the tests by Srivastava and Du (2008) has three folds. (a) The conditions to guarantee the central limit theorems of  $T_{SD}$  and  $T'_{SD}$  are somehow stringent; see (8). Also the sample size and population dimension have to satisfy that  $p = o(n^2)$ . To make the method more applicable, we hope to relax the condition imposed on  $\Sigma$  as well as that p on n. (b) We would like to understand an interesting observation by Zhang et al. (2020b) from their simulation: the distribution of  $T'_{SD}$  sometimes looks like a normal curve, and other times it looks like a chi-square curve. (c) We plan to give a rigorous proof of the major ingredient of this theory, that is, an asymptotic ratio-unbiased-estimator of  $tr(\mathbf{R}^2)$ , where **R** is the population correlation matrix of  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  aforementioned.

Now we state our findings. For (a), we have obtained the asymptotic distributions of  $T_{SD}$  and  $T'_{SD}$  for arbitrary  $\Sigma$  in Theorems 1 and 2, respectively. In particular, Theorem 1 holds for two extreme cases: the independent case with  $\mathbf{R} = \mathbf{I}_p$  and the most dependent case, i.e., all entries of the population vector are identical. Our theory says the restriction on pand n will be  $p = o(n^2)$  if the entries of the population vector are not far from independent. However, for dependent or very dependent case, our conclusion holds as long as p is not more than a polynomial order of n. For (b), we successfully understand the observation by Zhang et al. (2020b). In fact, there are indeed transition phenomena of the limiting distributions of  $T_{SD}$  and  $T'_{SD}$ . They are sometimes normal, mixing chi-squared or the sum of two independent random variables, one has normal distribution and the other has a mixing chi-squared distribution. This can be quickly seen from (11) and (17). As for (c), for arbitrary  $\Sigma$  we have proved rigorously that an asymptotic ratio-unbiased-estimator of  $\operatorname{tr}(\mathbf{R}^2)$  is  $\operatorname{tr}(\hat{\mathbf{R}}^2) - \frac{p(p-1)}{n-1}$  in Theorem 3. Especially the theorem is true for two extreme cases:  $\Sigma$  is diagonal or proportional to a matrix whose entries are all equal to 1. In addition, some of our partial calculations and heuristics indicate that the above phase transition from the Gaussian to mixing chi-squared distributions is possibly a universal phenomenon. We will present three testing procedures to justify this claim.

The above solution is conducted through the understanding of the sample correlation matrix  $\hat{\mathbf{R}}$ , a special random matrix, defined below (1). Unlike the Gaussian orhtogonal/unitary/symplectic ensemble or Wishart matrices,  $\hat{\mathbf{R}}$  lacks the orthogonal-invariant property. As a consequence, investigating the sample correlation matrix always cost extra energy than working on other popular matrices. We employ the machinery for  $\hat{\mathbf{R}}$  developed by Jiang (2004), Jiang (2009), Cai et al. (2013), Jiang and Yang (2013), and Fan and Jiang (2019). In particular, we extend the method conceived by Jiang (2004) and Fan and Jiang (2019) to prove a weak law of large numbers for tr( $\hat{\mathbf{R}}^2$ ) en route to obtain an asymptotic ratio-unbiased-estimator. More elaboration are provided in Section 4.

The remaining of the paper is organized as follows. The one-sample mean and twosample mean problems are studied in Sections 2 and 3, respectively. In Section 4, we give an asymptotic ratio-unbiased-estimator of  $tr(\mathbf{R}^2)$ , which is more applicable than the one by Srivastava and Du (2008) and Srivastava (2009) (their result lacks a rigorous proof although it has been used in literature). The concluding remarks and discussions are presented in Section 5, in which we particularly point out our findings on phase transitions between normal and mixing chi-squared distributions may also exist for some other testing procedures. Finally, the proofs are given in Section 6.

#### **2** One Sample Mean Test for Large p and Small n

Let  $\{X_1, \dots, X_n\}$  be a random sample from  $N_p(\mu, \Sigma)$ . Consider the test that

$$H_0: \boldsymbol{\mu} = \boldsymbol{0} \quad \text{versus} \quad H_1: \boldsymbol{\mu} \neq \boldsymbol{0}. \tag{5}$$

Let **D** be the diagonal matrix of  $\Sigma$ . Then the  $p \times p$  population correlation matrix is

$$\mathbf{R} = \mathbf{D}^{-1/2} \mathbf{\Sigma} \mathbf{D}^{-1/2}.$$
 (6)

Similarly, let  $\hat{\mathbf{D}}$  be the diagonal matrix of  $\hat{\mathbf{S}}$  from (1). Then the sample correlation matrix  $\hat{\mathbf{R}}$  is defined by

$$\hat{\mathbf{R}} = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2}.$$
(7)

Srivastava and Du (2008) and Srivastava (2009) obtain a result on  $T_{SD}$  from (2) as follows.

**Result 2.1** Assume  $n = O(p^{\zeta}), \frac{1}{2} < \zeta \leq 1$  and

$$0 < \lim_{p \to \infty} \frac{tr(\mathbf{R}^i)}{p} < \infty, \qquad i = 1, 2, 3, 4.$$
(8)

If  $\boldsymbol{\mu} = \mathbf{0}$  then  $T_{SD} \to N(0, 1)$  in distribution as  $p \to \infty$ .

A quick comment is that (8) holds automatically for i = 1 since all of the diagonal entries of **R** are equal to 1, and hence  $tr(\mathbf{R}) = p$ . In order to make approximation better, we now revise the statistic  $T_{SD}$  slightly. Set

$$T_{p,1} = \frac{n\bar{\boldsymbol{X}}^T\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} - pn(n-3)^{-1}}{\sqrt{2\left|\operatorname{tr}(\hat{\mathbf{R}}^2) - p(p-1)(n-1)^{-1}\right|}}.$$
(9)

We are doing so because, by Lemma 6.3, the major contribution of the mean of  $\operatorname{tr}(\hat{\mathbf{R}}^2)$  is  $\operatorname{tr}(\mathbf{R}^2) + p(p-1)(n-1)^{-1}$ . Also, by Theorem 3,  $\operatorname{tr}(\hat{\mathbf{R}}^2) - p(p-1)(n-1)^{-1} > 0$  as n and p are large enough. On the other hand, with probability one,  $\operatorname{tr}(\hat{\mathbf{R}}^2) - p(p-1)(n-1)^{-1} \neq 0$  because  $\operatorname{tr}(\hat{\mathbf{R}}^2)$  is a continuous function of Gaussian random variables. A discussion between  $T_{p,1}$  and  $T_{SD}$  will be elaborated shortly. Review the Frobenius norm, sometimes also called the Euclidean norm,  $\|\mathbf{A}\|_F := [\operatorname{tr}(\mathbf{A}^T \mathbf{A})]^{1/2}$  for any matrix  $\mathbf{A}$ . For mathematical rigor, we assume the sample sizes n depends on p. According to Hu and Bai (2016), the behavior of the statistic  $T_{SD}$  is not known if  $\Sigma$  has spikes. The following gives a complete characterization of the properties of  $T_{SD}$  and  $T_{p,1}$  in terms of the spikes of the correlation matrix  $\mathbf{R}$ .

**THEOREM 1** Suppose  $X_1, \dots, X_n$  is a random sample from  $N_p(\mu, \Sigma)$ . Let **R** be the correlation matrix as in (6) with eigenvalues  $\lambda_1 \geq \dots \geq \lambda_p \geq 0$ . Assume

(a)  $\lim_{p\to\infty} \frac{\lambda_i}{\|\mathbf{R}\|_F} = \rho_i \ge 0$  for all  $i \ge 1$ ;

(b)  $\lim_{p\to\infty} \frac{p}{n||\mathbf{R}||_F} = 0$  and  $\lim_{p\to\infty} \frac{p}{n^a} = 0$  for some constant a > 0. If  $\boldsymbol{\mu} = \mathbf{0}$ , then  $T_{SD} \to b\xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i (\xi_i^2 - 1)$  in distribution, where  $\xi_0, \xi_1, \xi_2, \cdots$  are i.i.d. N(0,1) and  $b = (1 - \sum_{i=1}^{\infty} \rho_i^2)^{1/2}$ . The same conclusion also holds for  $T_{p,1}$ . Condition (a) considers the possibility that **R** may have spikes. If  $\rho_1 = 0$ , then  $\rho_i = 0$ for every  $i \ge 1$  due to the monotonicity of  $\rho_i$ , and we say there are no spikes in **R**. In this situation we have  $T_{p,1} \to N(0,1)$  by Theorem 1. Using essentially the Fatou lemma, we have checked the given conditions actually imply  $\sum_{i=1}^{\infty} \rho_i^2 \le 1$ ; see the proof of Lemma 4.2. If  $\sum_{i=1}^{\infty} \rho_i^2 = 1$ , then  $T_{p,1}$  converges to a mixing chi-square distribution. If  $\sum_{i=1}^{\infty} \rho_i^2 \in (0,1)$ , then the asymptotic distribution is a blend of a Gaussian distribution and a mixing chisquared distribution. Condition (b) characterizes the restriction between the sample size nand population dimension p.

Observe that (8) with i = 3 implies that  $\lambda_1^3 \leq \operatorname{tr}(\mathbf{R}^3) = O(p)$ . Thus,  $\lambda_1 = O(p^{1/3})$ . Since all of the diagonal entries of  $\mathbf{R}$  are identical to 1 then  $\operatorname{tr}(\mathbf{R}^2) \geq p$ . This shows that  $\rho_1 = 0$  where  $\rho_1$  is from Theorem 1. Also, (8) with i = 2 implies that  $||\mathbf{R}||$  and  $\sqrt{p}$  have the same order, thus  $\frac{p}{n||\mathbf{R}||_F} \to 0$  as  $p \to \infty$ . This together with the condition " $n = O(p^{\zeta})$ ,  $\frac{1}{2} < \zeta \leq 1$ " implies (b) from Theorem 1. So our theorem is more general than Result 2.1 by Srivastava and Du (2008).

Let  $\mathbf{X}_1, \dots, \mathbf{X}_n$  be a random sample from an AR(1) model with  $\mathbf{R} = (\gamma^{|i-j|})$  and the absolute values of  $\gamma = \gamma_p$  staying away from 1. By using the Gersgorin disc theorem [see, e.g., p. 344 from Horn and Johnson (2012)], the largest eigenvalue or  $\mathbf{R}$  is of order O(1). Hence condition (a) of Theorem 1 holds with  $\rho_1 = 0$ . If condition (b) also holds, then both  $T_{SD}$  and  $T_{p,1}$  go to the standard normal distribution. The same conclusion is also valid for a banded correlation matrix  $\mathbf{R} = (r_{ij})$  with  $r_{ij} = 0$  for  $|j - i| \ge t$  where  $t = t_p = o(\sqrt{p})$ . In this case, the largest eigenvalue or  $\mathbf{R}$  is of order  $o(\sqrt{p})$ . Similar results can be obtained for other patterned matrices including Toeplitz matrices, Hankel matrices and symmetric circulant matrices; see, e.g., Brockwell et al. (2016).

Now let us look at some special features of Theorem 1. First, we do not need the population matrix  $\Sigma$  to be invertible as required in Hotelling's  $T^2$  test (Hotelling, 1931). The largest discrepancy between n and p from Srivastava and Du (2008) is that  $p = o(n^2)$ . Our range is that p can be at any polynomial order of n provided  $p = o(n ||\mathbf{R}||_F)$ . If the entries of the population vector are not very far from independence, in the sense that  $||\mathbf{R}||_F$ 

is in the order of  $\sqrt{p}$ , then the restriction  $p = o(n ||\mathbf{R}||_F)$  is reduced to  $p = o(n^2)$ . In the case that the entries of the population vector are very dependent such that  $||\mathbf{R}||_F$  is in the order of p, then p is allowed to take any polynomial order of n. To convince our readers for the dependent case and to make further discussions, we next study two extreme cases: independence and most dependence. The derivation of the results below does not use any techniques and steps from the proof of Theorem 1.

**PROPOSITION 1** Let  $X_1, \dots, X_n$  be a random sample from  $N_p(\mu, \Sigma)$  with  $\mu = 0$ . Assume  $n = n_p \to \infty$  as  $p \to \infty$ . The following hold.

(i) Let all of the  $p^2$  entries of  $\Sigma$  be identical. Then both  $T_{SD}$  and  $T_{p,1}$  converge to  $\frac{1}{\sqrt{2}}[\chi^2(1)-1]$  in distribution as  $p \to \infty$  regardless of the relative speeds of n and p.

(ii) Let  $\Sigma$  be a diagonal matrix whose diagonal entries are all positive, equivalently,  $\mathbf{R} = \mathbf{I}_p$ . Then

$$T_{SD} \rightarrow \begin{cases} \eta, & \text{if } p/n^2 \rightarrow 0; \\ \eta + \sqrt{h/2}, & \text{if } p/n^2 \rightarrow h; \\ \infty, & \text{if } p/n^2 \rightarrow \infty \end{cases}$$
(10)

in distribution, where  $\eta \sim N(0,1)$ . However,  $T_{p,1} \rightarrow N(0,1)$  as  $p \rightarrow \infty$  regardless of the speeds of n and p going to infinity.

In case (i) we see Theorem 1 holds without the assumption that  $p = o(n^a)$  for some constant a > 0. The conclusion for case (ii) says that the condition  $\lim_{p\to\infty} \frac{p}{n||\mathbf{R}||_F} = 0$ in Theorem 1 is sharp, and  $T_{p,1}$  is better than  $T_{SD}$ . In particular, we do not need the assumption  $\lim_{p\to\infty} \frac{p}{n||\mathbf{R}||_F} = 0$  to assure  $T_{p,1} \to N(0,1)$ . This is simply a coincidence because of the special structure of  $\Sigma$  or  $\mathbf{R}$ . The condition  $\lim_{p\to\infty} \frac{p}{n||\mathbf{R}||_F} = 0$  is essentially required at handling the denominator of  $T_{p,1}$  for arbitrary  $\mathbf{R}$ ; see (9) and Theorem 3. Figure 1 clearly shows the existence of shifts between the density curves of  $T_{SD}$  and N(0,1). However, with the ratio  $p/n^2$  becoming smaller, the shift diminishes gradually.



Figure 1: These pictures are based on Proposition 1, from  $10^5$  simulations. It shows the centers of  $T_{p,1}$  and N(0,1) are always close, and the existence of shifts between density curves of  $T_{SD}$  and N(0,1). The shifts are smaller as  $p/n^2$  become smaller.

Practically, given the population correlation  $\mathbf{R}$ , we need to determine if it has spikes and how many spikes. Interested readers are referred to Fan et al. (2020) and Morales-Jimenez et al. (2021) for procedures to do so. We next present some examples. Their verification are presented in Section 6.4.

**Example 2.1** Given  $r \in [0,1]$ , define  $\mathbf{A}_m := (1-r)\mathbf{I}_m + r\mathbf{J}_m$  for any  $m \ge 1$ , where  $\mathbf{I}_p$ and  $\mathbf{J}_p$  denote the  $p \times p$  identity matrix and the  $p \times p$  matrix of ones. Set  $\mathbf{R} = \mathbf{A}_p$ . Then  $\lambda_1 = 1 + (p-1)r$  and  $\lambda_2 = \cdots = \lambda_p = 1 - r$ . Assume  $r = r_p$  and  $\lim_{p\to\infty} \sqrt{p} \cdot r = c \ge 0$ . Then

$$T_p \to \begin{cases} \xi_0, & \text{if } c = 0; \\ \frac{1}{\sqrt{c^2 + 1}} \xi_0 + \frac{c}{\sqrt{2(c^2 + 1)}} (\xi_1^2 - 1), & \text{if } c \in (0, \infty); \\ \frac{1}{\sqrt{2}} \cdot (\xi_1^2 - 1), & \text{if } c = \infty \end{cases}$$
(11)

under condition  $p = o(n^2)$  for the case  $c \in [0, \infty)$  and under the conditions  $nr \to \infty$  and  $p = o(n^a)$  for some constant a > 0 for the case  $c = \infty$ . Here  $\xi_0$  and  $\xi_1$  are i.i.d. N(0, 1) and  $T_p$  stands for  $T_{SD}$  or  $T_{p,1}$ . The phase transition occurs between the normal and mixing chi-squared distributions as the entries of the correlation matrix change their values.

We make Figures 2 and 3 based on Example 2.1 to compare the performances of our  $T_{p,1}$ and  $T_{SD}$  by Srivastava and Du (2008). For different values of n, p, r, the statistics  $T_{p,1}$  and  $T_{SD}$  are simulated for 10<sup>5</sup> times. The elaboration is given below.

Figure 2 is designed for a small value of r with r = 0.1. Look at the first two pictures on the top row, that is, the ones with (n, p) = (10, 50) and (n, p) = (20, 100), respectively. Neither  $T_{p,1}$  nor  $T_{SD}$  perform well. But this is expected because the sample sizes are very small. However, by fixing the rate of p/n and let n and p grow gradually, the approximations become better and better. In particular our theoretical curves are always close to the empirical ones, whereas the normal ones stay farther or much farther from the empirical ones.

Figure 3 is designed for a big value of r with r = 0.5. Except the first two pictures on the top row, which correspond to (n, p) = (10, 50) and (n, p) = (20, 100), respectively, our theoretical curves match the empirical ones well. As the explanation aforementioned, that the two simulations do not behave well is understandable, simply because the sample sizes are very small. Our theoretical curves are constantly close to the empirical ones, the normal ones are nowhere close to the empirical ones.

In summary, the simulation indicates our approximation stated in Theorem 1 that  $(1 - \sum_{i=1}^{\infty} \rho_i^2)^{1/2} \xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i (\xi_i^2 - 1)$  to  $T_{SD}$  and  $T_{p,1}$ , respectively, outperforms the normal approximation from *Result 2.1*.



Figure 2: Example 2.1. For small r = 0.1, the empirical curves in blue are close to our theoretical curves in red. As sample size n becomes larger, our approximation becomes better; the normal approximation (black curve) is no longer valid.



Figure 3: Example 2.1. Compared to Figure 2 with r = 0.1, we take r = 0.5 here. The empirical curves in blue are close to our theoretical curves in red. The normal approximation (black curve) from *Result 2.1* is no longer valid.

**Example 2.2** For  $r \in (0,1)$ , set  $m = [p^r]$ . Recall  $\mathbf{A}_m$  from Example 2.1. Define  $\mathbf{R}$  by

$$\mathbf{R} = egin{pmatrix} \mathbf{A}_m & \mathbf{0} \ \mathbf{0} & \mathbf{I}_{p-m} \end{pmatrix}_{p imes p},$$

where the two "r" from " $m = [p^r]$ " and " $\mathbf{A}_m$ " are the same one. Assuming  $\boldsymbol{\mu} = \mathbf{0}$ , we have

$$T_p \to \begin{cases} \xi_0 & \text{if } 0 < r < \frac{1}{2}; \\ \frac{2}{\sqrt{5}}\xi_0 + \frac{1}{\sqrt{10}} \cdot (\xi_1^2 - 1), & \text{if } r = \frac{1}{2}; \\ \frac{1}{\sqrt{2}} \cdot (\xi_1^2 - 1), & \text{if } \frac{1}{2} < r < 1 \end{cases}$$
(12)

under condition  $p = o(n^2)$  for  $0 < r \le \frac{1}{2}$  and  $p = o(n^{1/(1-r)})$  for  $\frac{1}{2} < r < 1$ , where  $\xi_0$  and  $\xi_1$  are i.i.d. N(0,1) as  $r = \frac{1}{2}$ . Here  $T_p$  stands for  $T_{SD}$  or  $T_{p,1}$ . Obviously, there is a phase transition at  $r = \frac{1}{2}$  as r runs between 0 and 1. The phase transition appears at both relative sizes of n and p together with the change of the values of matrix entries.

**Example 2.3** Set  $m = [\log(p+2)]$ . Given  $\tau \ge 0$ , define integer  $p' = p - [\tau\sqrt{p}]$ ,  $\lambda_i = 1 + \tau 2^{-i} (1 - 2^{-m})^{-1} \sqrt{p}$  for  $1 \le i \le m$  and  $\lambda_i = 1$  for  $i = m + 1, \dots, p' - 1$ ,  $\lambda_{p'} = 1 + [\tau\sqrt{p}] - \tau\sqrt{p} \in [0,1]$ , and the rest of  $\lambda_i$  are identical to 0. By Lemma 4.3, there exists a correlation matrix **R** such that **R** has eigenvalues  $\lambda_i$ ,  $1 \le i \le p$ . Assuming  $\boldsymbol{\mu} = \mathbf{0}$ , we have

$$T_{p} \rightarrow \begin{cases} \xi_{0}, & \text{if } \tau = 0; \\ \sqrt{\frac{3}{\tau^{2}+3}}\xi_{0} + \sqrt{\frac{3\tau^{2}}{2(\tau^{2}+3)}} \sum_{i=1}^{\infty} \frac{1}{2^{i}}(\xi_{i}^{2}-1), & \text{if } \tau \in (0,\infty); \\ \sqrt{\frac{3}{2}} \sum_{i=1}^{\infty} \frac{1}{2^{i}}(\xi_{i}^{2}-1), & \text{if } \tau \to \infty \end{cases}$$
(13)

in distribution as  $p \to \infty$  under condition  $p = o(n^2)$ . Here  $\xi_0, \xi_1, \dots, \xi_d$  are i.i.d. N(0, 1). Here  $T_p$  stands for  $T_{SD}$  or  $T_{p,1}$ . The statistic  $T_p$  behaves like a rainbow, which has a Gaussian distribution at one end, and a mixing chi-squared distribution at the horizon, and a hybrid in between. An interesting remark is that, unlike the eigenvalues of a covariance matrix, the eigenvalues  $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_p \geq 0$  of the  $p \times p$  correlation matrix **R** cannot be arbitrary. By Lemma 4.3, they have to satisfy the so-called "majorization" property:  $\lambda_1 + \cdots + \lambda_k \geq k$  for every  $1 \leq k \leq p$  and  $\lambda_1 + \cdots + \lambda_p = p$ . Conversely, for any  $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_p \geq 0$  with the "majorization" property, there is always a correlation matrix with eigenvalues  $\lambda_i$ ,  $1 \leq i \leq p$ .

## **3** Two Sample Mean Test for Large p and Small n

We will study the two-sample mean testing problem in the high-dimensional setting in this section. As before, assume that  $\{X_{i1}, \dots, X_{in_i}\}$  for i = 1, 2 are two independent random samples with sizes  $n_1$  and  $n_2$ , and from *p*-variate normal distributions  $N_p(\mu_1, \Sigma)$  and  $N_p(\mu_2, \Sigma)$ , respectively. We wish to test

$$H_0: \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2 \quad \text{versus} \quad H_1: \boldsymbol{\mu}_1 \neq \boldsymbol{\mu}_2. \tag{14}$$

Let  $\bar{\mathbf{X}}_i$  be the sample mean of the *i*-th sample with i = 1, 2 and  $\hat{\mathbf{S}}$  be the pooled sample covariance matrix defined in (3). Set  $n = n_1 + n_2$ . Srivastava and Du (2008) consider the statistic  $T'_{SD}$  from (4). Recall  $\hat{\mathbf{D}}$  is the diagonal matrix of  $\hat{\mathbf{S}}$  in (3) and

$$\hat{\mathbf{R}} = \hat{\mathbf{R}}_p = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2} \tag{15}$$

is the pooled sample correlation matrix. Similar to the discussion before (9) on the modification of  $T_{SD}$ , we make a little change for  $T'_{SD}$  as follows. Define

$$T_{p,2} = \frac{\frac{n_1 n_2}{n_1 + n_2 - 1} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2)^T \hat{\mathbf{D}}^{-1} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2) - \frac{(n_1 + n_2 - 1)p}{n_1 + n_2 - 4}}{\sqrt{2 \left| \operatorname{tr}(\hat{\mathbf{R}}^2) - \frac{p(p-1)}{n_1 + n_2 - 2} \right|}}.$$
(16)

Srivastava and Du (2008) derive the limiting distribution of  $T'_{SD}$  under conditions similar to those from *Result 2.1* and an extra assumption " $\frac{n_1}{n_1+n_2} \rightarrow c \in (0,1)$ ". On the other hand, Hu and Bai (2016) ask for the properties of  $T'_{SD}$  as the variances of the entries of the population vector are quite different. We will provide a solution for  $T'_{SD}$  and  $T_{p,2}$  next. For mathematical rigor, we assume that both sample sizes  $n_1$  and  $n_2$  depend on p. **THEOREM 2** Let  $\{X_{i1}, \dots, X_{in_i}\}$  for i = 1, 2 be two independent random samples from  $N_p(\boldsymbol{\mu}_1, \boldsymbol{\Sigma})$  and  $N_p(\boldsymbol{\mu}_2, \boldsymbol{\Sigma})$ , respectively. Let **R** be as in (6) with eigenvalues  $\lambda_1 \geq \dots \geq \lambda_p \geq 0$ . Assume

(a)  $\lim_{p\to\infty} \frac{\lambda_i}{\|\mathbf{R}\|_F} = \rho_i \ge 0$  for all  $i \ge 1$ ;

(b)  $\lim_{p\to\infty} \frac{p}{(n_1+n_2)\|\mathbf{R}\|_F} = 0$  and  $\lim_{p\to\infty} \frac{p}{(n_1+n_2)^a} = 0$  for some constant a > 0.

If  $\boldsymbol{\mu}_1 = \boldsymbol{\mu}_2$ , then  $T_{p,2} \to b\xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i (\xi_i^2 - 1)$  in distribution, where  $\xi_0, \xi_1, \xi_2, \cdots$  are *i.i.d.* N(0,1) and  $b = (1 - \sum_{i=1}^{\infty} \rho_i^2)^{1/2}$ . The same conclusion also holds for " $T_{p,2}$ ".

The above result does not require condition "min $\{n_1, n_2\} \to \infty$ " or " $\frac{n_1}{n_1+n_2} \to \gamma \in (0, 1)$ ", which are commonly assumed in literature for the two-sample testing problem.

As discussed below Theorem 1, condition (8) implies that  $\rho_1 = 0$  and  $\frac{p}{n ||\mathbf{R}||_F} \to 0$  as  $p \to \infty$ . Condition " $n_1 + n_2 = O(p^{\zeta}), \frac{1}{2} < \zeta \leq 1$ " from (8) implies (b) from Theorem 2. So our theorem is more general than the conclusion for the two-sample test by Srivastava and Du (2008).

If **R** is AR(1) model, the banded model, the Toeplitz or Hankel matrices, please see the discussion below Theorem 1. They are also applied here.

Let  $\mathbf{X}_1, \dots, \mathbf{X}_n$  be a random sample from an AR(1) model with  $\mathbf{R} = (\gamma^{|i-j|})$  and the absolute values of  $\gamma = \gamma_p$  staying away from 1. By using the Gersgorin disc theorem [see, e.g., p. 344 from Horn and Johnson (2012)], the largest eigenvalue or  $\mathbf{R}$  is of order O(1). Hence condition (a) of Theorem 1 holds with  $\rho_1 = 0$ . If condition (b) also holds, then both  $T_{SD}$  and  $T_{p,1}$  go to the standard normal distribution. The same conclusion is also valid for a banded correlation matrix  $\mathbf{R} = (r_{ij})$  with  $r_{ij} = 0$  for  $|j - i| \ge t$  where  $t = t_p = o(\sqrt{p})$ . In this case, the largest eigenvalue or  $\mathbf{R}$  is of order  $o(\sqrt{p})$ . Similar results can be obtained for other patterned matrices including Toeplitz matrices, Hankel matrices and symmetric circulant matrices; see, e.g., Brockwell et al. (2016).

Recall the definition of  $T_{p,2}$  from (16). Let **D** be the diagonal matrix of  $\Sigma$ . Under some conditions including  $\|\mathbf{R}\|_F$  is of the order p, Zhang et al. (2020b) obtain the limiting distribution of a normalized  $(\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2)^T \mathbf{D}^{-1} (\bar{\mathbf{X}}_1 - \bar{\mathbf{X}}_2)$  as  $\frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i (\xi_i^2 - 1)$ . Although this quantity is not directly applicable because the unknown parameter matrix **D** is involved, it is indeed suggestive. Their result seems to be a special case of Theorem 2 with  $\sum_{i=1}^{\infty} \rho_i^2 = 1$ . In fact, under their assumptions, we have confirmed that  $\sum_{i=1}^{\infty} \rho_i^2 = 1$  in Lemma 6.37 from Section 6.4. Notice  $\sqrt{p} \leq ||\mathbf{R}||_F \leq p$  for any **R**. Srivastava and Du (2008) consider the extreme case that  $||\mathbf{R}||_F$  is of the scale of  $\sqrt{p}$ . This is essentially a weakly dependent situation with  $\rho_i = 0$  for each  $i \geq 1$ . According to Theorem 2, the limiting distribution is a normal. On the other hand, roughly speaking, Zhang et al. (2020b) intend to study another extreme case in which  $||\mathbf{R}||_F$  is of the scale p, and hence **R** is completely a spiked model. As explained above, the corresponding limiting distribution is a mixing chi-squared distribution. Our Theorem 2 handles the case for any **R** with  $||\mathbf{R}||_F$  running everywhere between  $\sqrt{p}$  and p, and the limiting distribution turns out to be, interestingly enough, a convolution of both.

Now we demonstrate our results by giving some examples.

**Example 3.1** Recall Example 2.1 and  $\mathbf{R} = \mathbf{A}_p$ . For this example, Zhang et al. (2020b) observe that "When r = 0.01, the histograms are quite symmetric and bell-shaped, indicating that a normal approximation as suggested by the theory of Srivastava and Du (2008) may be applied for approximating the null distribution of  $T_{SD}$ . However, when r = 0.5 and r = 0.9, the histograms are quite skewed, indicating that a normal approximation is no longer adequate." In fact, our Theorem 2 explains their inspection very accurately as follows. Assume  $r = r_p$  and  $\lim_{p\to\infty} \sqrt{p} \cdot r = c \ge 0$ . By changing "n" in Example 2.1 to " $n_1 + n_2$ ", we have

$$T_p \to \begin{cases} \xi_0, & \text{if } c = 0; \\ \frac{1}{\sqrt{c^2 + 1}} \xi_0 + \frac{c}{\sqrt{2(c^2 + 1)}} (\xi_1^2 - 1), & \text{if } c \in (0, \infty); \\ \frac{1}{\sqrt{2}} \cdot (\xi_1^2 - 1), & \text{if } c = \infty \end{cases}$$
(17)

under condition  $p = o((n_1 + n_2)^2)$  for the case  $c \in [0, \infty)$  and under the condition  $p = o((n_1 + n_2)^a)$  for some constant a > 0 for the case  $c = \infty$ , where  $\xi_0$  and  $\xi_1$  are i.i.d. N(0, 1). Here  $T_p$  stands for  $T'_{SD}$  or  $T_{p,2}$ . Obviously, there is a phase transition between c = 0 and  $c = \infty$ .



Figure 4: Example 3.1. Our theoretical curves always match the empirical ones well no matter the former is a normal-like or chi-square-like curve, even one of  $n_1$  and  $n_2$  being small. The normal curves by Srivastava and Du (2008) are farther from the empirical ones as correlation r increases.

Similar to Example 3.1, modifications of Examples 2.2 and 2.3 can also be constructed for the two-sample case. It is straightforward, so we omit the detail. Given a data set, the spikes of the population correlation  $\mathbf{R}$  have to be determined; see Fan et al. (2020) and Morales-Jimenez et al. (2021).

Based on Example 3.1, the statistic  $T_{p,2}$  from (16) is simulated for  $10^5$  times for each set of values of  $(n_1, n_2, p, r)$ . The value of r is chosen at 0, 1, 0, 5 and 0.9, designed for weakly dependence, dependence and strong dependence, respectively. Our theoretical curves are normal-like, mixing between N(0, 1) and chi-squared distribution and chi-squared-like curves. For weakly dependent case, by examining the four graphs in the first column, we see the normal approximation by Srivastava and Du (2008) becomes less and less accurate as all of the values of  $n_1, n_2, p$  increase. However, our theoretical curve always match the empirical ones well. For dependent and very dependent cases appeared in the second and third columns, respectively, the normal curves are nowhere close to the empirical ones. It is good to see our theoretical curves are always close to their empirical counterparts.

## 4 A Law of Large Numbers for Sample Correlation Matrices

Let  $X_1, \dots, X_n$  be a random sample from  $N_p(\mu, \Sigma)$  with correlation matrix  $\mathbf{R}$ . Let  $\hat{\mathbf{R}}$  be the sample correlation matrix defined in (7). One of the crucial steps in proving Theorems 1 and 2 is the use of an asymptotically ratio-unbiased-estimator of  $\operatorname{tr}(\mathbf{R}^2)$ . In their Lemma 3.2, Srivastava and Du (2008) state the following.

**Result 4.1** If  $n = O(p^{\zeta})$ ,  $0 < \zeta \leq 1$ , under the condition (8),  $(tr(\mathbf{R}^2) - p^2/n)/p$  converges to  $\lim_{p\to\infty} tr(\mathbf{R}^2)/p$  in probability as  $(n,p) \to \infty$  and thus can be considered as a consistent estimator of  $tr(\mathbf{R}^2)/p$  as  $(n,p) \to \infty$ .

The proof of the above result is give on pages 400-402 from Srivastava and Du (2008). It lacks a mathematical rigor. By using a method from the random matrix theory, we rigorously

obtain a more general result than the above in the following. Most efforts in the whole proofs of this paper are devoted to this result, which is rather involved. The reason is that we assume no independence at all among the entries of the population distribution, i.e.,  $\mathbf{R}$  is arbitrary.

**THEOREM 3** Let  $X_1, \dots, X_n$  be a random sample from  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  with correlation matrix  $\mathbf{R}$ . Let  $\hat{\mathbf{R}}$  be the sample correlation matrix defined in (7). If  $\lim_{p\to\infty} \frac{p}{n\|\mathbf{R}\|_F} = 0$  and  $\lim_{p\to\infty} \frac{p}{n^a} = 0$  for some constant a > 0, then

$$\frac{1}{tr(\mathbf{R}^2)} \left[ tr(\hat{\mathbf{R}}^2) - \frac{p(p-1)}{n-1} \right] \to 1$$

in probability as  $p \to \infty$ .

Unlike  $\operatorname{tr}(\mathbf{R}^2)$ ,  $\operatorname{tr}(\mathbf{R}^3)$  and  $\operatorname{tr}(\mathbf{R}^4)$  needed in *Result 4.1*, our conditions in the above theorem are imposed on  $\|\mathbf{R}\|_F = [\operatorname{tr}(\mathbf{R}^2)]^{1/2}$  only. Also, it is easy to see the difference between the assumptions on restrictions of n and p: *Result 4.1* asks  $n = O(p^{\zeta})$  for some  $0 < \zeta \leq 1$  and our Theorem 3 holds as long as n and p are in the order of a polynomial. We make some remarks on Theorem 3 next.

Naively, an obvious ratio-unbiased-estimator of  $\operatorname{tr}(\mathbf{R}^2)$  is its sample version  $\operatorname{tr}(\hat{\mathbf{R}}^2)$ . Theorem 3 indicates that it is not true for the high dimensional setting. Instead, the modified version " $\operatorname{tr}(\hat{\mathbf{R}}^2) - \frac{p(p-1)}{n-1}$ " is a correct one. By using the fact  $\operatorname{tr}(\hat{\mathbf{R}}^2) \ge p$ , we can see from Theorem 3 that  $\operatorname{tr}(\hat{\mathbf{R}}^2)$  is indeed a ratio-unbiased-estimator of  $\operatorname{tr}(\mathbf{R}^2)$  if p is as small as p = o(n), that is,  $\operatorname{tr}(\hat{\mathbf{R}}^2)/\operatorname{tr}(\mathbf{R}^2) \to 1$  in probability.

Sample correlation matrices are a special type of random matrices. The spectral distributions of eigenvalue values are studied in Jiang (2004) and Xiao and Zhou (2010). The central limit theorems for determinants under independent and correlated situations are obtained by Jiang and Yang (2013) and Jiang (2009), respectively. The Tracy-Widom law is derived by Bao et al. (2012). The central limit theorem for linear statistics of eigenvalues is understood by Gao et al. (2017).

An interesting fact is the derivation of Theorem 3. Through the whole proof, contrary to standard techniques to handle randomness, we do not use/assume any independence from

the population distribution  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ . In fact, in the most dependent case, that is, all of the p entries of  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  are identical, then  $\hat{\mathbf{R}} = \mathbf{R}$  and all of their  $p^2$  entries are equal to 1, so  $\operatorname{tr}(\hat{\mathbf{R}}^2) = \operatorname{tr}(\mathbf{R}^2) = p^2$ . One can see Theorem 3 trivially holds. On the other hand, for the most independent case, i.e., if  $\boldsymbol{\Sigma} = \mathbf{R} = \mathbf{I}_p$ , then  $\operatorname{tr}(\mathbf{R}^2) = p$ . Assuming  $p/n \to \rho \in (0, 1]$ , we have from Theorem 3 that  $\operatorname{tr}(\hat{\mathbf{R}}^2)/p \to 1 + \rho$ . This actually can be confirmed independently by a random matrix theory. In fact, let  $\mu_p$  be the empirical distribution of the eigenvalues of  $\hat{\mathbf{R}}$ . By Theorem 2 from Jiang (2004),  $\mu_p$  converges weakly to the so-called Marčhenko-Pastur law  $\mu$ . The second moment of  $\mu$  is equal to  $1 + \rho$  [Lemma 3.1 from Bai and Silverstein (2010)]. Consequently,  $\operatorname{tr}(\hat{\mathbf{R}}^2)/p = \int x^2 d\mu_p \to \int x^2 d\mu = 1 + \rho$ .

The proof of Theorem 3 is a bit technical and lengthy. We first give an accurate estimate of  $Etr(\hat{\mathbf{R}}^2)$  in Lemma 6.3. Then we only need to show that its variance go to zero. To do so, since  $\hat{\mathbf{R}}^2 = p + 2 \sum_{1 \le i < j \le p} \hat{r}_{ij}^2$ , we need to understand the covariance between two sample correlations, say,  $\hat{r}_{ij}^2$  and  $\hat{r}_{kl}^2$  for any  $1 \le i, j, k, l \le p$ . The precise result is given next.

**THEOREM 4** Assume  $m \geq 5$ . Let  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T \in \mathbb{R}^4; 1 \leq j \leq m\}$  be i.i.d. random vectors with distribution  $N_4(\mathbf{0}, \mathbf{R})$ , where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each i. Set

$$\hat{r}_{ij} = \frac{\sum_{k=1}^{m} X_{ik} X_{jk}}{(\sum_{k=1}^{m} X_{ik}^2)^{1/2} (\sum_{k=1}^{m} X_{jk}^2)^{1/2}}$$
(18)

for  $1 \leq i \leq j \leq 4$ . Then, for any  $N \geq 1$ ,  $Cov(\hat{r}_{12}^2, \hat{r}_{34}^2)$  is equal to

$$\begin{split} \varrho_{m,1} \cdot \sum_{1 \le i < j \le 4} r_{ij}^2 + \varrho_{m,2} \cdot r_{12}^2 r_{34}^2 &+ \varrho_{m,3} \cdot \left( r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} \right) \\ &+ \varrho_{m,4} \cdot \left( r_{13} r_{34} r_{41} + r_{23} r_{34} r_{42} \right) r_{12}^2 \\ &+ \varrho_{m,5} \cdot \left( r_{12} r_{23} r_{31} + r_{12} r_{24} r_{41} \right) r_{34}^2 \\ &+ \frac{\varrho_{m,6}}{m^{(N+1)/2}}, \end{split}$$

where  $\{\varrho_{m,i}; 3 \leq i \leq 5\}$  are quantities not depending on **R**,

$$|\varrho_{m,1}| \le Km^{-2}, \ |\varrho_{m,2}| \lor |\varrho_{m,3}| \lor |\varrho_{m,4}| \lor |\varrho_{m,5}| \le Km^{-1}, \ |\varrho_{m,6}| \le K$$

and K is a constant depending on N but not on m or  $\mathbf{R}$ .

The proof of Theorem 4 is rather involved. There are two reasons, one of them is that no assumption is imposed on  $\mathbf{R}$ , as a consequence, we are not able to use any techniques/theory related to independence. The second one is that sample correlation coefficients are more difficult to be handled than sample covariances. Look at (18) for  $\hat{r}_{12}$  by taking i = 1 and j = 2. From the law of large numbers,  $(1/m) \sum_{k=1}^m X_{1k}^2 \to 1$  and  $(1/m) \sum_{k=1}^m X_{2k}^2 \to 1$ . To understand  $\hat{r}_{12}$ , a naive idea is to replace the denominator in the expression of  $\hat{r}_{12}$  by m. Interestingly, this works for the derivation of the Marčhenko-Pastur law (Jiang, 2004) when  $\mathbf{R} = \mathbf{I}$ . Now we elaborate this more for a further discussion. By the Taylor expansion, write

$$\frac{1}{\sqrt{1+x}} = 1 - \frac{x}{2} + \frac{3}{8}x^2 + \dots + a_N x^N + o(x^{N+1})$$
(19)

as x is small. Jiang (2004) uses the above expansion with N = 0 and x taking  $(1/m) \sum_{k=1}^{m} (X_{1k}^2 - 1)$  and  $(1/m) \sum_{k=1}^{m} (X_{2k}^2 - 1)$ , respectively. If  $\mathbf{R} \neq \mathbf{I}$  but  $\mathbf{R} = \mathbf{A}_p$  as in Example 2.1, Fan and Jiang (2019) use the expansion (19) with N = 1. In the proof our Theorem 3, since no structure of  $\mathbf{R}$  is assumed, we have to use (19) for an arbitrary N. This is why the condition "lim  $\sup_{p\to\infty} \frac{p}{n^a} = 0$  for some constant a > 0" is imposed in Theorems 1 and 3. The new method here would be useful in the future for the study on sample correlation matrices. It is worthwhile to mention that handling sample correlation matrices is much more difficult than working on sample covariance matrices. The difference is obvious: the entries of a sample covariance matrix does not have the denominator as in the expression of  $\hat{r}_{ij}$  from (18). For instance, Bai and Saranadasa (1996) also investigate a ratio-unbiased-estimator similar to Theorem 3 but for covariance matrices, their argument is quick and short.

A byproduct of the proofs of Theorems 1 and 2 is the behavior of a quadratic form related to the diagonal matrix of **R**. It would be useful for other research from this point forward. Assume  $X_1, \dots, X_n$  is a random sample from  $N_p(\mu, \Sigma)$  with correlation matrix **R**. Review **D** is the diagonal matrix of  $\Sigma$  and  $\hat{\mathbf{D}}$  is the diagonal matrix of  $\hat{\mathbf{S}}$  in (3).

**LEMMA 4.1** Let  $\boldsymbol{\eta} \sim N_p(\mathbf{0}, \boldsymbol{\Sigma})$  and  $\boldsymbol{\eta}$  be independent of  $\hat{\mathbf{S}}$ . If  $\lim_{p \to \infty} \frac{p}{n \|\mathbf{R}\|_F} = 0$ , then

$$\frac{\boldsymbol{\eta}^T \hat{\mathbf{D}}^{-1} \boldsymbol{\eta}}{\sqrt{2 \operatorname{tr}(\mathbf{R}^2)}} = \frac{\boldsymbol{\eta}^T \mathbf{D}^{-1} \boldsymbol{\eta}}{\sqrt{2 \operatorname{tr}(\mathbf{R}^2)}} + o_p(1)$$

as  $p \to \infty$ .

Since  $X_1, \dots, X_n$  is a random sample from a multivariate normal distribution,  $\bar{X}$  is independent of  $\hat{S}$ , and hence  $\bar{X}$  is independent of  $\hat{D}$ . So the above conclusion holds if we take  $\eta = \sqrt{p}\bar{X}$ . This is actually the way we use this lemma in later proofs.

The limiting distributions in Theorems 1 and 2 essentially come from the following.

**LEMMA 4.2** For each  $p \ge 1$ , let  $a_{p,1} \ge \cdots \ge a_{p,p} \ge 0$  be constants with  $a_{p,1}^2 + \cdots + a_{p,p}^2 = 1$ . Suppose  $\lim_{p\to\infty} a_{p,i} = \rho_i \ge 0$  for each  $i \ge 1$ . Let  $\xi_1, \xi_2, \cdots$  be i.i.d. with distribution  $\chi^2(1) - 1$ . Then  $a_{p,1}\xi_1 + \cdots + a_{p,p}\xi_p$  converges to  $[2(1-\sum_{i=1}^{\infty}\rho_i^2)]^{1/2}\eta + \sum_{i=1}^{\infty}\rho_i\xi_i$  in distribution as  $p \to \infty$ , where  $\eta \sim N(0,1)$  and  $\eta$  is independent of  $\{\xi_i; i \ge 1\}$ .

The comment after Example 2.3 is based on the following fact. It is interesting useful in its own right.

**LEMMA 4.3** Let **M** be a  $p \times p$  correlation matrix, namely, **M** is non-negative definite and its diagonal entries are all equal to 1. Suppose **M** has eigenvalues  $\lambda_1 \geq \cdots \geq \lambda_p$ . Then  $\lambda_1 + \cdots + \lambda_k \geq k$  for  $k = 1, \cdots, p$  and  $\lambda_1 + \cdots + \lambda_p = p$ . Conversely, for any  $\tau_1 \geq \cdots \geq \tau_p \geq 0$ with  $\tau_1 + \cdots + \tau_k \geq k$  for each  $k = 1, \cdots, p$  and  $\tau_1 + \cdots + \tau_p = p$ , there always exists a correlation matrix with eigenvalues  $\tau_1, \cdots, \tau_p$ .

#### 5 Concluding Remarks and Discussion

1. In this paper we have studied one-sample and two-sample mean tests. For the multiple population case, it becomes the classical MANOVA problem. In the "large p, small n" situation, Srivastava and Fujikoshi (2006) consider the question by revising the classical F-test. They use functions of non-zero eigenvalues of pseudo-MANOVA random matrix as test statistics. Zhang et al. (2017) and Chen et al. (2019) study this case based on the  $L_2$ -type statistics. An extension of our work to the MANOVA case under "large p, small n" situation is meaningful. One direction is to modify the classical tests, such as the Anderson test, the Pillal test, the Roy test, the Wilks test and the Olson test, by a method similar to those

from Srivastava and Fujikoshi (2006). The classical statistics take the correlations among population coordinates into account. It has kind of self-normalization, which is favorable.

2. All results in this paper are based on random samples from a multivariate Gaussian distribution. Is it possible to generalize this to non-Gaussian cases? The corresponding results are highly demanded simply because many data are not sampled from a Gaussian distribution. Also, conclusions on non-Gaussian scenario may help us understand robustness of our tests.

3. We study the two-sample mean test by assuming the two populations have the same covariance matrices. If the two covariance matrices are not identical, this is the multivariate Behrens-Fisher problem. Various methods are proposed, for example, by Yao (1965), Johansen (1980), Nel and Van der Merwe (1986), Krishnamoorthy and Yu (2004), Chen and Qin (2010) and Chen et al. (2019). For one-dimensional case (p = 1), the method initiated by Welch (1947) is probably the most popular one. The author basically normalize the difference of sample means by its sample standard deviation. Srivastava et al. (2013) propose a statistic by replacing  $\hat{\mathbf{D}}$  from (16) with a hybrid of two samples. Similar to the study in this paper, its properties are needed to be understood, too.

4. As far as proofs go, we spend most of our energy proving Theorem 4. We do not use complicated technology. Instead, by we use brute force to compute mixed moments of multiple Gaussian random variables in combination with some random matrix theory. As a result, the argument is lengthy. It is possible to shorten the proof. To get the law of large numbers stated in Theorem 4, one may like to try Gaussian concentration inequalities; see, for example, Ledoux (2001) and Boucheron et al. (2013). An alternative way is to get the joint density function of  $\hat{r}_{12}$  and  $\hat{r}_{34}$  similar to that of the marginal density of  $\hat{r}_{12}$  treated at (26) through an hypergeometric function.

5. Theorems 1 and 2 present the null distributions for the one-sample and two-sample tests, respectively. We actually have tried to derive the non-null limiting distribution to explore power functions of the tests. The argument is also very involved. The current paper is already very lengthy, so we postpone and leave it as a future work.

#### Acknowledgment

The bulk of the work of Tiefeng Jiang was conducted during his visit at Baidu Research in 2019-2020. After completing an initial draft, we solicited comments from several colleagues and we appreciate their feedback.

### 6 Proofs

One of the main steps of proving Theorems 1 and 2 is a weak law of large numbers for  $\operatorname{tr}(\hat{\mathbf{R}}^2)$  (Theorem 3), where  $\hat{\mathbf{R}} = \hat{\mathbf{R}}_p$  is the sample correlation matrix defined in (7). To derive this, we need to study the mean and variance of  $\operatorname{tr}(\hat{\mathbf{R}}^2)$ . In Section 6.1, we will get an accurate estimate of  $E[\operatorname{tr}(\hat{\mathbf{R}}^2)]$  by an argument on hypergeometric functions. Then we will show  $\operatorname{Var}(\operatorname{tr}(\hat{\mathbf{R}}^2))$  go to zero. The proof of Theorem 3 will be completed once this step is established. This is a rather involved step and it will be understood step by step in Sections 6.2.1-6.2.6.

## 6.1 Evaluation of the Mean of the Frobenius Norm of a Sample Correlation matrix

In this section, we will work on the mean of  $tr(\hat{\mathbf{R}}^2)$ , where  $\hat{\mathbf{R}} = \hat{\mathbf{R}}_p$  is the sample correlation matrix defined in (7). The critical tool is the hypergeometric function. We first need a preliminary result as follows.

**LEMMA 6.1** There exists a constant C > 0 such that the following holds for all  $k \ge 2$ .

$$\sum_{n=1}^{\infty} \binom{n+k}{k}^{-1} \le Ck^{-1/4}.$$

**Proof of Lemma 6.1**. For given a > 0, set  $f_a(x) = (1 + ax^{-1})^{-x}$  for x > 0. We claim that  $f_a(x)$  is decreasing in  $x \in (0, \infty)$ . In fact, let  $g(x) = \log f_a(x)$ . Then

$$g'(x) = \left[ -x \log\left(1 + \frac{a}{x}\right) \right]' = -\log\left(1 + \frac{a}{x}\right) - x \frac{-\frac{a}{x^2}}{1 + \frac{a}{x}}.$$

Then

$$g'(x) = \log\left(1 - \frac{a}{x+a}\right) + \frac{a}{x+a} \le 0$$

for all x > 0 since  $\log(1 + y) \le y$  for all y > -1. Hence,  $f_a(x)$  is decreasing in  $x \in (0, \infty)$ for each a > 0. By the Stirling formula,  $m! = \sqrt{2\pi m} m^m e^{-m + \frac{\theta_m}{12m}}$  with  $\theta_m \in (0, 1)$  for each  $m \ge 1$ . Consequently,

$$\sqrt{2\pi m}m^m e^{-m} \le m! \le e \cdot \sqrt{2\pi m}m^m e^{-m}$$

for all  $m \ge 1$ . Therefore,

$$\binom{n+k}{k} = \frac{(n+k)!}{n!k!} \geq e^{-2} \cdot \frac{\sqrt{n+k}}{\sqrt{2\pi nk}} \cdot \frac{(n+k)^{n+k}e^{-(n+k)}}{k^k n^n e^{-(n+k)}} \\ \geq e^{-2} \frac{1}{\sqrt{2\pi k}} \cdot \frac{(n+k)^{n+k}}{k^k n^n}.$$

This implies that

$$\frac{1}{\binom{n+k}{k}} \le (2\pi e^2 k^{1/2}) \left(1 + \frac{k}{n}\right)^{-n} \left(1 + \frac{n}{k}\right)^{-k}.$$
(20)

For fixed k, since  $f_k(x)$  is decreasing in x, we have

$$\sum_{n=1}^{k} \left(1 + \frac{k}{n}\right)^{-n} \left(1 + \frac{n}{k}\right)^{-k} \leq \sum_{n=1}^{k} \left(1 + \frac{k}{n}\right)^{-n}$$
$$\leq \frac{1}{k+1} + \sum_{n=2}^{k} \int_{n-1}^{n} \left(1 + \frac{k}{x}\right)^{-x} dx$$
$$= \frac{1}{k+1} + \int_{1}^{k} \left(1 + \frac{k}{x}\right)^{-x} dx.$$
(21)

Again, use the fact that  $f_k(x)$  is decreasing to see

$$\int_{1}^{k} \left(1 + \frac{k}{x}\right)^{-x} dx = \int_{1}^{k^{1/4}} \left(1 + \frac{k}{x}\right)^{-x} dx + \int_{k^{1/4}}^{k} \left(1 + \frac{k}{x}\right)^{-x} dx$$
$$\leq \frac{k^{1/4}}{k+1} + k \cdot (1 + k^{3/4})^{-k^{1/4}}.$$

If  $k \ge 81$ , then  $k^{1/4} \ge 3$ . Consequently,  $k \cdot (1 + k^{3/4})^{-k^{1/4}} \le k \cdot (k^{3/4})^{-3} = k^{-5/4}$ . From (21) we see

$$\sum_{n=1}^{k} \left(1 + \frac{k}{n}\right)^{-n} \left(1 + \frac{n}{k}\right)^{-k} \le \frac{1}{k+1} + \frac{k^{1/4}}{k+1} + \frac{1}{k^{5/4}} \le \frac{3}{k^{3/4}}.$$
(22)

Now,  $(1+\frac{k}{n})^{-n} \leq 2^{-k}$  as  $n \geq k$  by the fact  $f_k(x)$  is decreasing. Also,  $(1+\frac{n}{k})^{-k} \leq (1+\frac{n}{k})^{-81}$  as  $k \geq 81$ . Hence

$$\sum_{n=k}^{\infty} \left(1 + \frac{k}{n}\right)^{-n} \left(1 + \frac{n}{k}\right)^{-k} \le 2^{-k} \sum_{n=k}^{\infty} \left(1 + \frac{n}{k}\right)^{-81} \le 2^{-k} k^{81} \sum_{n=1}^{\infty} \frac{1}{n^{81}}$$
(23)

as  $k \ge 81$ . This joined with (20), (22) and (23) yields that

$$\sum_{n=1}^{\infty} \frac{k^{1/4}}{\binom{n+k}{k}} \le k^{1/4} \cdot (2\pi e^2 k^{1/2}) \cdot \left(3k^{-3/4} + \zeta(81)2^{-k}k^{81}\right) \le C'$$

for all  $k \ge 81$ , where  $\zeta(81) = \sum_{n=1}^{\infty} \frac{1}{n^{81}} < \infty$  is the Riemann zeta function evaluated at 81 and C' is another numerical constant not depending on k. In summary,

$$\sup_{k \ge 81} \sum_{n=1}^{\infty} k^{1/4} \binom{n+k}{k}^{-1} < \infty.$$
(24)

Note that

$$\frac{\binom{n+i+1}{i+1}}{\binom{n+i}{i}} = \frac{n+i+1}{i+1} > 1$$

for any  $i \ge 1$  and  $n \ge 1$ . We know  $\binom{n+i}{i}$  is increasing in *i*, and hence

$$\sup_{2 \le k \le 80} \sum_{n=1}^{\infty} k^{1/4} \binom{n+k}{k}^{-1} \le 80^{1/4} \sum_{n=1}^{\infty} \binom{n+2}{2}^{-1} \le 2 \cdot 80^{1/4} \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty.$$

This and (24) imply the conclusion.

The following result quantifies the second moment of the sample correlation coefficient of a random sample of size m up to an error  $O(m^{-1/4})$ .

**LEMMA 6.2** Let  $\{(X_i, Y_i)^T; 1 \le i \le m\}$  be i.i.d. 2-dimensional normal random vectors with  $EX_1 = EY_1 = 0$ ,  $EX_1^2 = EY_1^2 = 1$  and  $Cov(X_1, Y_1) = r$ . Set

$$\hat{r}_m = \frac{\sum_{k=1}^m X_i Y_i}{(\sum_{k=1}^m X_i^2)^{1/2} (\sum_{k=1}^m Y_i^2)^{1/2}}.$$

Write

$$E(\hat{r}_m^2) = \frac{1}{m} + r^2 + b_m(r) \cdot r^2$$

for  $m \ge 4$ . Then  $\sup_{m \ge 4, |r| \le 1} |m^{1/4} b_m(r)| < \infty$ .

**Proof of Lemma 6.2.** If  $r = \pm 1$ , since  $EX_1 = EY_1 = 0$  and  $EX_1^2 = EY_1^2 = 1$ , we know  $Y_i = \pm X_i$  for each *i*. By the definition of  $\hat{r}_m$ , trivially,  $\hat{r}_m = \pm 1$ . This implies that

$$b_m(\pm 1) = -\frac{1}{m} \tag{25}$$

for each  $m \ge 1$ . In the following we always assume  $r^2 < 1$ . From Ghosh (1966) or p. 156 in Muirhead (1982),

$$E(\hat{r}_m^2) = 1 - \frac{m-1}{m} (1 - r^2) \cdot {}_2F_1\left(1, 1; \frac{1}{2}m + 1; r^2\right),$$
(26)

where  ${}_{2}F_{1}(a,b;c;z)$  is an hypergeometric function defined by

$$_{2}F_{1}(a,b;c;z) = \sum_{k=0}^{\infty} \frac{(a)_{k}(b)_{k}}{(c)_{k}k!} z^{k}, \quad |z| < 1,$$

and where  $(x)_0 = 1$  and  $(x)_k = x(x+1)\cdots(x+k-1)$  for  $k \ge 1$ ; see, for example, p. 20 from Muirhead (1982). Notice  $(1)_k = k!$  and

$$\left(\frac{1}{2}m+1\right)_k = \frac{m+2}{2} \cdot \frac{m+4}{2} \cdots \frac{m+2k}{2}.$$

Thus

$${}_{2}F_{1}\left(1,1;\frac{1}{2}m+1;r^{2}\right) = 1 + \sum_{k=1}^{\infty} \frac{2^{k}k!}{(m+2)(m+4)\cdots(m+2k)} r^{2k}$$
$$= 1 + r^{2} \sum_{k=1}^{\infty} \frac{2^{k}k!}{(m+2)(m+4)\cdots(m+2k)} r^{2k-2}.$$
 (27)

Evidently, the last sum is bounded by

$$\sum_{k=1}^{\infty} \frac{2^k k!}{(m+2)(m+4)\cdots(m+2k)}.$$

Set  $j = [\frac{m}{2}]$ , where [x] denotes the integer part of a real number  $x \ge 0$ . Then  $j \le \frac{m}{2}$ , or equivalently,  $m \ge 2j$ . It follows that

$$\frac{2^k k!}{(m+2)(m+4)\cdots(m+2k)} \le \frac{k!}{(j+1)(j+2)\cdots(j+k)} = \frac{j!k!}{(j+k)!}.$$

It follows that

$$\sum_{k=1}^{\infty} \frac{2^k k!}{(m+2)(m+4)\cdots(m+2k)} \le \sum_{k=1}^{\infty} \frac{1}{\binom{j+k}{j}} \le \frac{K_1}{j^{1/4}} \le \frac{K_2}{m^{1/4}}$$

as  $j \ge 2$ , or equivalently,  $m \ge 4$  by Lemma 6.1, where  $K_1$  and  $K_2$  are constants not depending on m. From this and (27), we are able to write

$$_{2}F_{1}\left(1,1;\frac{1}{2}m+1;r^{2}\right) = 1 + a_{m}r^{2},$$

where  $0 \le a_m \le K_2 m^{-1/4}$ . Substitute this back to (26), we get

$$E(\hat{r}_m^2) = 1 - \frac{m-1}{m}(1-r^2) \cdot (1+a_m r^2)$$
  
=  $1 - \frac{m-1}{m}(1-r^2) - \frac{m-1}{m}(1-r^2) \cdot a_m r^2$   
=  $\frac{1}{m} + r^2 + b_m(r)r^2$ ,

where

$$b_m(r) = -\frac{1}{m}r^2 - \frac{m-1}{m}(1-r^2) \cdot a_m r^2.$$

Obviously,  $\sup_{|r|\leq 1} |b_m(r)| \leq K_3 m^{-1/4}$  for every  $m \geq 4$ , where  $K_3$  is a constant not depending on m. This and (25) lead to the desired conclusion.

**LEMMA 6.3** Let  $X_1, \dots, X_n$  be a random sample from  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  with correlation matrix **R**. Let  $\hat{\mathbf{R}} = \hat{\mathbf{R}}_p$  be the sample correlation matrix defined in (7). If  $\lim_{p\to\infty} \frac{p}{n\|\mathbf{R}\|_F} = 0$  and  $\limsup_{p\to\infty} \frac{p}{n^a} = 0$  for some constant a > 0, then, as  $p \to \infty$ ,

$$E tr(\hat{\mathbf{R}}_{p}^{2}) = \frac{p(p-1)}{n-1} + tr(\mathbf{R}_{p}^{2}) \cdot \left[1 + O(m^{-1/4})\right].$$

**Proof of Lemma 6.3**. Set m = n-1. The notation  $W_p(m, \Sigma)$  represents the distribution of the Wishart matrix  $\mathbf{U}^T \mathbf{U}$ , where  $\mathbf{U}$  is an  $m \times p$  matrix whose rows are i.i.d. with distribution  $N_p(\mathbf{0}, \Sigma)$ . Then,  $n\hat{\mathbf{S}}$  has the Wishart distribution  $W_p(m, \Sigma)$ ; see, for example, Theorem 3.1.2 from Muirhead (1982). That is,  $n\hat{\mathbf{S}} \stackrel{d}{=} \mathbf{U}^T \mathbf{U}$ . We claim

$$\hat{\mathbf{R}}_{p} = (\hat{r}_{ij})_{p \times p} \stackrel{d}{=} \left( \frac{\mathbf{v}_{i}^{T} \mathbf{v}_{j}}{\|\mathbf{v}_{i}^{T}\| \cdot \|\mathbf{v}_{j}\|} \right)_{p \times p},\tag{28}$$

where the *m* rows of  $(\mathbf{v}_1, \dots, \mathbf{v}_p)_{m \times p}$  are i.i.d. with distribution  $N_p(\mathbf{0}, \mathbf{R})$ . In fact, write  $\mathbf{U} = (u_{ij}) = (\mathbf{u}_1, \dots, \mathbf{u}_p)$  where  $\mathbf{u}_i = (u_{1i}, \dots, u_{mi})^T \in \mathbb{R}^m$  for each *i*. Then  $\mathbf{u}_i^T \mathbf{u}_j$  is the

(i, j)-entry of  $\mathbf{U}^T \mathbf{U}$  and the diagonal entries are  $\|\mathbf{u}_i\|^2$  for  $1 \le i \le p$ . We know from (7) that  $\hat{\mathbf{R}}_p = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2}$ , where  $\hat{\mathbf{D}}$  is the diagonal matrix of  $\hat{\mathbf{S}}$ . Then

$$\hat{\mathbf{R}}_p = (\hat{r}_{ij})_{p \times p} \stackrel{d}{=} \left( \frac{\mathbf{u}_i^T \mathbf{u}_j}{\|\mathbf{u}_i^T\| \cdot \|\mathbf{u}_j\|} \right)_{p \times p}.$$

Notice  $(\mathbf{u}_i, \mathbf{u}_j) = (u_{ki}, u_{kj})_{1 \le k \le m}$  for any i < j and the *m* rows are i.i.d. bivariate normal random variables with mean vector **0** and covariance matrix

$$\begin{pmatrix} \sigma_{ii} & \sigma_{ij} \\ \sigma_{ij} & \sigma_{jj} \end{pmatrix}$$

where  $\Sigma := (\sigma_{ij})_{p \times p}$ . Write  $\mathbf{R}_p = (r_{ij})_{p \times p}$ . Then  $r_{ij} = \sigma_{ij} (\sigma_{ii} \sigma_{jj})^{-1/2}$  by definition. Observe

$$\frac{\mathbf{u}_{i}^{T}\mathbf{u}_{j}}{\|\mathbf{u}_{i}^{T}\|\cdot\|\mathbf{u}_{j}\|} = \frac{(\sigma_{ii}^{-1/2}\mathbf{u}_{i})^{T}(\sigma_{jj}^{-1/2}\mathbf{u}_{j})}{\|\sigma_{ii}^{-1/2}\mathbf{u}_{i}^{T}\|\cdot\|\sigma_{jj}^{-1/2}\mathbf{u}_{j}\|}.$$

Take  $\mathbf{v}_i = \sigma_{ii}^{-1/2} \mathbf{u}_i$  to obtain (28).

By Lemma 6.2,

$$E \operatorname{tr}(\hat{\mathbf{R}}_{p}^{2}) = p + 2 \sum_{1 \leq i < j \leq p} E(\hat{r}_{ij}^{2})$$
  
$$= p + 2 \sum_{1 \leq i < j \leq p} \left[\frac{1}{m} + r_{ij}^{2} + b_{m}(r_{ij})r_{ij}^{2}\right]$$
  
$$= \frac{p(p-1)}{m} + \operatorname{tr}(\mathbf{R}_{p}^{2}) + 2 \sum_{1 \leq i < j \leq p} b_{m}(r_{ij})r_{ij}^{2},$$

where  $\sup_{|r|\leq 1} |b_m(r)| \leq Km^{-1/4}$  and K is a constant not depending on m or r. Since

$$2\Big|\sum_{1 \le i < j \le p} b_m(r_{ij})r_{ij}^2\Big| \le (2K)m^{-1/4} \cdot \operatorname{tr}(\mathbf{R}_p^2),$$

the above two displays show that

$$E \operatorname{tr}(\hat{\mathbf{R}}_{p}^{2}) = \frac{p(p-1)}{n-1} + \operatorname{tr}(\mathbf{R}_{p}^{2}) \cdot \left[1 + O(m^{-1/4})\right]$$

as  $p \to \infty$ . The proof is completed.

#### 6.2 The Proofs of Theorems 3 and 4

#### 6.2.1 Mixing Moments of Gaussian Random Variables

The following lemma is a very useful tool to compute the mean of the product of Gaussian random variables.

**LEMMA 6.4** (Isserlis and Wick formula) Let  $n \ge 2$  be an even integer and  $(X_1, \dots, X_n)^T \in \mathbb{R}^n$  follows a normal distribution with mean vector **0**. Then

$$E(X_1X_2\cdots X_n) = \sum_p \prod_{(i,j)\in p} E(X_iX_j),$$

where the sum runs over every pairing p of  $\{1, 2, \dots, n\}$ , that is, all distinct ways of partitioning  $\{1, 2, \dots, n\}$  into pairs  $\{i, j\}$ , and the product is over the pairs contained in p. Sometimes we also use its equivalent form:

$$E(X_1X_2\cdots X_n) = \sum_{i=2}^n E(X_1X_i) \cdot E\left(\frac{X_2\cdots X_n}{X_i}\right).$$

A seemingly more general, but actually an equivalent formula of Lemma 6.4 is the following Lemma 6.5, which will be used only once in a comment after the proof of Lemma 6.7.

**LEMMA 6.5** (*Guiard*, 1986) Let  $X = (X_1, \dots, X_n)$  be n-dimensionally centralized normally distributed with  $Cov(X_i, X_j) = \sigma_{ij}$ . Let  $\{\alpha_i; 1 \leq i \leq n\}$  be positive integers. If  $\sum_{i=1}^n \alpha_i$  is odd, then  $\mathbb{E}(\prod_{i=1}^n X_i^{\alpha_i}) = 0$ . If  $\sum_{i=1}^n \alpha_i$  is even, then

$$\mathbb{E}\left(\prod_{i=1}^{n} X_{i}^{\alpha_{i}}\right) = \sum_{2\beta_{ii}+\sum_{j:j\neq i}\beta_{ij}=\alpha_{i},\forall i} \frac{\prod_{i=1}^{n} \alpha_{i}!}{\prod_{i=1}^{n} 2^{\beta_{ii}} \prod_{1\leq i\leq j\leq n} \beta_{ij}!} \prod_{1\leq i\leq j\leq n} \sigma_{ij}^{\beta_{ij}},$$

where  $\{\beta_{ij}, 1 \leq i \leq j \leq n\}$  are non-negative integers.

In Lemmas 6.6 and 6.7 below, two identities on mixing moments of Gaussian random variables will be presented.

**LEMMA 6.6** Let  $(X_1, X_2, X_3, X_4)^T \sim N_4(0, \mathbf{R})$ . Assume  $\mathbf{R} = (r_{ij})_{4 \times 4}$  with  $r_{ii} = 1$  for each *i*. Then

(i) 
$$E(X_1X_2X_3^2X_4^2) = r_{12} + 2r_{12}r_{34}^2 + 2r_{13}r_{23} + 2r_{14}r_{24} + 4r_{13}r_{24}r_{34} + 4r_{14}r_{23}r_{34};$$
  
(ii)  $E(X_1^2X_2^2X_3^2) = 1 + 2r_{12}^2 + 2r_{13}^2 + 2r_{23}^2 + 8r_{12}r_{23}r_{31}.$ 

**Proof of Lemma 6.6.** Let  $(\xi_1, \dots, \xi_6)^T$  be a multivariate normal with mean vector  $\mathbf{0}$ ,  $\operatorname{Cov}(\xi_i, \xi_j) = \sigma_{ij}$  and  $\operatorname{Var}(\xi_i) = 1$  for each *i*. Then, by Lemma 6.4,

$$E(\xi_{1}\xi_{2}\xi_{3}\xi_{4}\xi_{5}\xi_{6}) = \sigma_{12}\sigma_{34}\sigma_{56} + \sigma_{12}\sigma_{35}\sigma_{46} + \sigma_{12}\sigma_{36}\sigma_{45} + \sigma_{13}\sigma_{24}\sigma_{56} + \sigma_{13}\sigma_{25}\sigma_{46} + \sigma_{13}\sigma_{26}\sigma_{45} + \sigma_{14}\sigma_{23}\sigma_{56} + \sigma_{14}\sigma_{25}\sigma_{36} + \sigma_{14}\sigma_{26}\sigma_{35} + \sigma_{15}\sigma_{23}\sigma_{46} + \sigma_{15}\sigma_{24}\sigma_{36} + \sigma_{15}\sigma_{26}\sigma_{34} + \sigma_{16}\sigma_{23}\sigma_{45} + \sigma_{16}\sigma_{24}\sigma_{35} + \sigma_{16}\sigma_{25}\sigma_{34}.$$

Now take  $\xi_3 = \xi_4$  and  $\xi_5 = \xi_6$  to see

$$E(\xi_{1}\xi_{2}\xi_{3}^{2}\xi_{5}^{2}) = \sigma_{12} + \sigma_{12}\sigma_{35}^{2} + \sigma_{12}\sigma_{35}^{2} + \sigma_{13}\sigma_{23} + \sigma_{13}\sigma_{25}\sigma_{35} + \sigma_{13}\sigma_{25}\sigma_{35} + \sigma_{13}\sigma_{23} + \sigma_{13}\sigma_{25}\sigma_{35} + \sigma_{13}\sigma_{25}\sigma_{35} + \sigma_{15}\sigma_{23}\sigma_{35} + \sigma_{15}\sigma_{23}\sigma_{35} + \sigma_{15}\sigma_{25} + \sigma_{15}\sigma_{23}\sigma_{35} + \sigma_{15}\sigma_{23}\sigma_{35} + \sigma_{15}\sigma_{25} + \sigma_{12}\sigma_{23}\sigma_{35} + \sigma_{15}\sigma_{25} + \sigma_{12}\sigma_{25}\sigma_{35} + \sigma_{15}\sigma_{25} + \sigma_{12}\sigma_{25}\sigma_{35} + \sigma_{15}\sigma_{25}\sigma_{35} + \sigma$$

This says that

$$E(X_1X_2X_3^2X_4^2) = r_{12} + 2r_{12}r_{34}^2 + 2r_{13}r_{23} + 2r_{14}r_{24} + 4r_{13}r_{24}r_{34} + 4r_{14}r_{23}r_{34}.$$
 (29)

We obtain (i). Now, take  $X_2 = X_1$  in (29). By using the fact  $r_{12} = 1$  and by changing "2" to "1" to the indices of r we obtain

$$E(X_1^2 X_3^2 X_4^2) = 1 + 2r_{34}^2 + 2r_{13}^2 + 2r_{14}^2 + 4r_{13}r_{14}r_{34} + 4r_{14}r_{13}r_{34}$$
$$= 1 + 2r_{13}^2 + 2r_{14}^2 + 2r_{34}^2 + 8r_{13}r_{14}r_{34}.$$

In the above and change "3" to "2" and "4" to "3" to see

$$E(X_1^2 X_2^2 X_3^2) = 1 + 2r_{12}^2 + 2r_{13}^2 + 2r_{23}^2 + 8r_{12}r_{13}r_{23}.$$

The proof is completed.

**LEMMA 6.7** Suppose the 4-dimensional random vector  $(X_1, X_2, X_3, X_4)^T \sim N_4(0, \mathbf{R})$ . Assume  $\mathbf{R} = (r_{ij})_{4 \times 4}$  with  $r_{ii} = 1$  for each *i*. Then

$$E(X_1^2 X_2^2 X_3^2 X_4^2) = 1 + 2 \sum_{1 \le i < j \le 4} r_{ij}^2 + 4(r_{12}^2 r_{34}^2 + r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2) + 8(r_{12} r_{23} r_{31} + r_{12} r_{24} r_{41} + r_{23} r_{34} r_{42} + r_{13} r_{34} r_{41}) + 16(r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} + r_{13} r_{32} r_{24} r_{41}).$$

In particular, take  $X_3 = X_1$  and  $X_4 = X_2$  to see  $E(X_1^4 X_2^4) = 9 + 72r_{12}^2 + 24r_{12}^4$ .

Proof of Lemma 6.7. By Lemma 6.4,

$$E(X_1^2 X_2^2 X_3^2 X_4^2) = E(X_1^2) \cdot E(X_2^2 X_3^2 X_4^2) + 2E(X_1 X_2) \cdot E(X_1 X_2 X_3^2 X_4^2) + 2E(X_1 X_3) \cdot E(X_1 X_3 X_2^2 X_4^2) + 2E(X_1 X_4) \cdot E(X_1 X_4 X_2^2 X_3^2).$$

By Lemma 6.6(i),

$$E(X_1X_2X_3^2X_4^2) = r_{12} + 2r_{12}r_{34}^2 + 2r_{13}r_{23} + 2r_{14}r_{24} + 4r_{13}r_{24}r_{34} + 4r_{14}r_{23}r_{34}.$$
 (30)

Similarly, we obtain  $E(X_1X_3X_2^2X_4^2)$  and  $E(X_1X_4X_2^2X_3^2)$  by exchanging the roles of " $X_2$ " and " $X_3$ " and exchanging the roles of " $X_2$ " and " $X_4$ ", respectively, from (30). By Lemma 6.6(ii),

$$E(X_1^2 X_2^2 X_3^2) = 1 + 2r_{12}^2 + 2r_{13}^2 + 2r_{23}^2 + 8r_{12}r_{13}r_{23}.$$

Hence,

$$E(X_1^2 X_2^2 X_3^2 X_4^2) = 1 + 2r_{23}^2 + 2r_{24}^2 + 2r_{34}^2 + 8r_{23}r_{24}r_{34} + 2r_{12}(r_{12} + 2r_{12}r_{34}^2 + 2r_{13}r_{23} + 2r_{14}r_{24} + 4r_{13}r_{24}r_{34} + 4r_{14}r_{23}r_{34}) + 2r_{13}(r_{13} + 2r_{13}r_{24}^2 + 2r_{12}r_{23} + 2r_{14}r_{34} + 4r_{12}r_{34}r_{24} + 4r_{14}r_{23}r_{24}) + 2r_{14}(r_{14} + 2r_{14}r_{23}^2 + 2r_{13}r_{34} + 2r_{12}r_{24} + 4r_{13}r_{24}r_{23} + 4r_{12}r_{34}r_{23}).$$

Sorting them out, we have

$$\begin{split} E(X_1^2 X_2^2 X_3^2 X_4^2) &= 1 + 2 \sum_{1 \le i < j \le 4} r_{ij}^2 + 4(r_{12}^2 r_{34}^2 + r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2) + \\ & 8(r_{12} r_{23} r_{31} + r_{12} r_{24} r_{41} + r_{23} r_{34} r_{42} + r_{13} r_{34} r_{41}) + \\ & 16(r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} + r_{13} r_{32} r_{24} r_{41}). \end{split}$$

The proof is completed.

Lemma 6.7 studies  $E(X_1^2 X_2^2 \cdots X_n^2)$  for n = 4, which is sufficient for our purpose. It is interesting to see the formula for  $n \ge 5$  by pure curiosity. If we argue the same way as in the proof of Lemma 6.7 through Lemma 6.4, the sorting procedure would be messy. However, Lemma 6.5 provides a way to do so by figuring out the non-negative integer solutions of the system equations  $2\beta_{ii} + \sum_{j:j\neq i} \beta_{ij} = 2$  for  $1 \le i \le n$ .

The following is a key step to study the covariance between two squared sample covariances stated in Lemma 6.9.

**LEMMA 6.8** Let  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T \in \mathbb{R}^4; 1 \leq j \leq m\}$  be i.i.d. random vectors with distribution  $N_4(\mathbf{0}, \mathbf{R})$ , where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each *i*. Set

$$B_1 = \frac{1}{m} \sum_{j=1}^m X_{1j} X_{2j} \quad and \quad B_2 = \frac{1}{m} \sum_{j=1}^m X_{3j} X_{4j}.$$

Then,

**Comment**. We now conduct an independent check of the accuracy of Lemma 6.8(iv) for two special cases. In the above result, take  $X_{1j} = X_{2j} = X_{3j} = X_{4j}$ , then  $r_{ij} = 1$  for all i, j. Consequently, we get from Theorem 6.8(iv) that

$$E\left[\frac{1}{m}\sum_{j=1}^{m}(X_{1j}^2-1)\right]^4 = E\left[(B_1-r_{12})^2(B_2-r_{34})^2\right] = \frac{12}{m^2} + \frac{48}{m^3}.$$

Notice  $\{X_{1j}; 1 \leq j \leq m\}$  are i.i.d. N(0,1). Then  $\sum_{j=1}^{m} (X_{1j}^2 - 1)$  is a sum of i.i.d. random variables with  $E(X_{11}^2 - 1)^2 = 2$  and  $E(X_{11}^2 - 1)^4 = 60$ . By a classical formula (see, for example, p. 69 from Durrett (2019)),

$$E\left[\frac{1}{m}\sum_{j=1}^{m} (X_{1j}^2 - 1)\right]^4 = \frac{1}{m^4} \left\{ m \cdot E(X_{11}^2 - 1)^4 + 3m(m-1) \cdot \left[E(X_{11}^2 - 1)^2\right]^2 \right\}$$
$$= \frac{12}{m^2} + \frac{48}{m^3}.$$

So Theorem 6.8(iv) recovers the case for  $r_{ij} = 1$  for all i, j. On the other hand, assume the two 2-dimensional random vectors  $(X_{11} X_{21})^T$  and  $(X_{31} X_{41})^T$  are independent, that is,  $r_{13} = r_{14} = r_{23} = r_{24} = 0$ . By Theorem 6.8(iv),

$$E[(B_1 - r_{12})^2 (B_2 - r_{34})^2] = \frac{1}{m^2} (1 + r_{12}^2) (1 + r_{34}^2).$$

On the other hand, by independence and Theorem 6.8(i),

$$E[(B_1 - r_{12})^2 (B_2 - r_{34})^2] = E[(B_1 - r_{12})^2] \cdot E[(B_2 - r_{34})^2] = \frac{1}{m^2} (1 + r_{12}^2)(1 + r_{34}^2)$$

So this independent check indicates that Theorem 6.8(iv) holds for the case  $r_{13} = r_{14} = r_{23} = r_{24} = 0$ .

**Proof of Lemma 6.8**. First, by Lemma 6.4,

$$E(X_{11}^2 X_{21}^2) = 1 + 2r_{12}^2, \quad E(X_{11}^3 X_{21}) = 3r_{12},$$
(31)

$$E(X_{11}^2 X_{21} X_{31}) = r_{23} + 2r_{12}r_{13}, \quad E(X_{11} X_{21} X_{31} X_{41}) = r_{12}r_{34} + r_{13}r_{24} + r_{14}r_{23}.$$
(32)

In fact, the two middle identities from the four in the above can be deduced immediately from the last one by taking  $X_{11} = X_{21} = X_{31}$  and  $X_{11} = X_{21}$ , respectively.

(i) Notice  $mB_1$  and  $mB_2$  are sums of i.i.d. random variables with mean  $r_{12}$  and  $r_{34}$ , respectively. Also,  $X_{ij} \sim N(0, 1)$  for each i, j. Then  $E[(B_1 - r_{12})^2] = \frac{1}{m} \text{Var}(X_{11}X_{21})$ . Since  $\text{Cov}(X_{11}, X_{21}) = E(X_{11}X_{21}) = r_{12}$ , we have from (31) that  $\text{Var}(X_{11}X_{21}) = 1 + r_{12}^2$ . So (i) follows.

(ii) By independence, 
$$E[(B_1 - r_{12})(B_2 - r_{34})] = \frac{1}{m} \text{Cov}(X_{11}X_{21}, X_{31}X_{41})$$
. From (32),  
 $\text{Cov}(X_{11}X_{21}, X_{31}X_{41}) = r_{12}r_{34} + r_{13}r_{24} + r_{14}r_{23} - r_{12}r_{34} = r_{13}r_{24} + r_{14}r_{23}$ . (33)

(iii) Write

$$(B_2 - r_{34})^2 = \frac{1}{m^2} \sum_{j=1}^m (X_{3j} X_{4j} - r_{34})^2 + \frac{2}{m^2} \sum_{1 \le k < l \le m} (X_{3k} X_{4k} - r_{34}) (X_{3l} X_{4l} - r_{34}).$$
(34)

By independence, the covariance between  $B_1 - r_{12}$  and any term from the last sum is zero. This implies

$$E[(B_1 - r_{12})(B_2 - r_{34})^2] = \frac{1}{m^2} E[(X_{11}X_{21} - r_{12})(X_{31}X_{41} - r_{34})^2].$$
(35)

Use expansion  $(X_{31}X_{41} - r_{34})^2 = X_{31}^2 X_{41}^2 - 2r_{34}X_{31}X_{41} + r_{34}^2$  to see that the last expectation in (35) is equal to  $E[(X_{11}X_{21} - r_{12})(X_{31}^2X_{41}^2 - 2r_{34}X_{31}X_{41})]$ , which is again equal to

$$E(X_{11}X_{21}X_{31}^2X_{41}^2) - 2r_{34}E(X_{11}X_{21}X_{31}X_{41}) - r_{12}E(X_{31}^2X_{41}^2) + 2r_{12}r_{34}E(X_{31}X_{41})$$

$$= E(X_{11}X_{21}X_{31}^2X_{41}^2) - 2r_{34}(r_{12}r_{34} + r_{13}r_{24} + r_{14}r_{23}) - r_{12}(1 + 2r_{34}^2) + 2r_{12}r_{34}^2$$

$$= E(X_{11}X_{21}X_{31}^2X_{41}^2) - 2r_{34}(r_{12}r_{34} + r_{13}r_{24} + r_{14}r_{23}) - r_{12}$$
(36)

by (31) and (32). From Lemma 6.6(i), we see that

$$E(X_{11}X_{21}X_{31}^2X_{41}^2) = r_{12} + 2r_{12}r_{34}^2 + 2r_{13}r_{23} + 2r_{14}r_{24} + 4r_{13}r_{24}r_{34} + 4r_{14}r_{23}r_{34}.$$
 (37)

Plug this into the previous display, we arrive at

$$E[(B_1 - r_{12})(B_2 - r_{34})^2] = \frac{1}{m^2} (2r_{13}r_{23} + 2r_{14}r_{24} + 2r_{13}r_{24}r_{34} + 2r_{14}r_{23}r_{34})$$

(iv) By (i),

$$E[(B_1 - r_{12})^2 (B_2 - r_{34})^2]$$

$$= \operatorname{Cov}((B_1 - r_{12})^2, (B_2 - r_{34})^2) + E(B_1 - r_{12})^2 \cdot E(B_2 - r_{34})^2$$

$$= \operatorname{Cov}((B_1 - r_{12})^2, (B_2 - r_{34})^2) + \frac{1}{m^2}(1 + r_{12}^2)(1 + r_{34}^2).$$
(38)
Similar to (34), we have

$$(B_1 - r_{12})^2 = \frac{1}{m^2} \sum_{j=1}^m (X_{1j} X_{2j} - r_{12})^2 + \frac{2}{m^2} \sum_{1 \le k < l \le m} (X_{1k} X_{2k} - r_{12}) (X_{1l} X_{2l} - r_{12}).$$
(39)

Recall the *m* random variables  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T; 1 \leq j \leq m\}$  are i.i.d., thus each term from the double sums in (34) and (39) is a product of two independent random variables. This implies that  $E[(X_{3k}X_{4k} - r_{34})(X_{3l}X_{4l} - r_{34})] = 0$  for any k < l, and the term  $(X_{1j}X_{2j} - r_{12})^2$ is uncorrelated to  $(X_{3k}X_{4k} - r_{34})(X_{3l}X_{4l} - r_{34})$  for any j, k, l with k < l. By the same spirit, it is easy to check that  $(X_{1k}X_{2k} - r_{12})(X_{1l}X_{2l} - r_{12})$  and  $(X_{3k_1}X_{4k_1} - r_{34})(X_{3l_1}X_{4l_1} - r_{34})$ are uncorrelated for any k < l and  $k_1 < l_1$  as long as  $(k, l) \neq (k_1, l_1)$ . These yield

$$Cov((B_{1} - r_{12})^{2}, (B_{2} - r_{34})^{2})$$

$$= \frac{1}{m^{3}}Cov((X_{11}X_{21} - r_{12})^{2}, (X_{31}X_{41} - r_{34})^{2}) + \frac{4}{m^{4}}\sum_{1 \le k < l \le m} Cov((X_{11}X_{21} - r_{12})(X_{12}X_{22} - r_{12}), (X_{31}X_{41} - r_{34})(X_{32}X_{42} - r_{34}))$$

$$= \frac{1}{m^{3}}Cov((X_{11}X_{21} - r_{12})^{2}, (X_{31}X_{41} - r_{34})^{2}) + \frac{2(m-1)}{m^{3}}Cov((X_{11}X_{21} - r_{12})(X_{12}X_{22} - r_{12}), (X_{31}X_{41} - r_{34})(X_{32}X_{42} - r_{34})). (40)$$

For brevity of notation, let  $(X_1, X_2, X_3, X_4) \in \mathbb{R}^4$  and  $(Y_1, Y_2, Y_3, Y_4) \in \mathbb{R}^4$  be i.i.d. random vectors with distribution  $N_4(\mathbf{0}, \mathbf{R})$ . Then the last covariance in (40) is identical to

$$E[(X_1X_2 - r_{12})(Y_1Y_2 - r_{12})(X_3X_4 - r_{34})(Y_3Y_4 - r_{34})]$$
  
=  $\{E[(X_1X_2 - r_{12})(X_3X_4 - r_{34})]\}^2$   
=  $r_{13}^2r_{24}^2 + r_{14}^2r_{23}^2 + 2r_{13}r_{32}r_{24}r_{41}$  (41)

by independence and (33). Now we calculate the first covariance in (40). In fact,

$$\operatorname{Cov}\left((X_{11}X_{21} - r_{12})^{2}, (X_{31}X_{41} - r_{34})^{2}\right)$$

$$= \operatorname{Cov}\left((X_{1}X_{2} - r_{12})^{2}, (X_{3}X_{4} - r_{34})^{2}\right)$$

$$= \operatorname{Cov}\left(X_{1}^{2}X_{2}^{2}, X_{3}^{2}X_{4}^{2}\right) - 2r_{12}\operatorname{Cov}\left(X_{1}X_{2}, X_{3}^{2}X_{4}^{2}\right) - 2r_{34}\operatorname{Cov}\left(X_{1}^{2}X_{2}^{2}, X_{3}X_{4}\right) + 4r_{12}r_{34}\operatorname{Cov}\left(X_{1}X_{2}, X_{3}X_{4}\right)$$

$$(42)$$

since  $(X_1X_2 - r_{12})^2 = X_1^2X_2^2 - 2r_{12}X_1X_2 + r_{12}^2$  and  $(X_3X_4 - r_{34})^2 = X_3^2X_4^2 - 2r_{34}X_3X_4 + r_{34}^2$ . From (31) and Lemma 6.7,

$$Cov(X_1^2 X_2^2, X_3^2 X_4^2) = E(X_1^2 X_2^2 X_3^2 X_4^2) - E(X_1^2 X_2^2) \cdot E(X_3^2 X_4^2)$$
  
=  $2(r_{13}^2 + r_{14}^2 + r_{23}^2 + r_{24}^2) + 4(r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2)$   
+ $8(r_{12} r_{23} r_{31} + r_{12} r_{24} r_{41} + r_{23} r_{34} r_{42} + r_{13} r_{34} r_{41})$   
+ $16(r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} + r_{13} r_{32} r_{24} r_{41}).$ 

Also, by (37),

$$Cov (X_1 X_2, X_3^2 X_4^2) = E(X_1 X_2 X_3^3 X_4^2) - r_{12}(1 + 2r_{34}^2)$$
  
=  $2r_{13}r_{23} + 2r_{14}r_{24} + 4r_{13}r_{24}r_{34} + 4r_{14}r_{23}r_{34}.$ 

By exchanging "1" to "3" and exchanging "2" and "4" in the above, we obtain

$$\operatorname{Cov}\left(X_{1}^{2}X_{2}^{2}, X_{3}X_{4}\right) = 2r_{13}r_{14} + 2r_{23}r_{24} + 4r_{13}r_{24}r_{12} + 4r_{23}r_{14}r_{12}.$$

Combining the above computations with (33) and (42), we arrive at

$$Cov \left( (X_{11}X_{21} - r_{12})^2, (X_{31}X_{41} - r_{34})^2 \right)$$
  
=  $2(r_{13}^2 + r_{14}^2 + r_{23}^2 + r_{24}^2) + 4(r_{13}^2r_{24}^2 + r_{14}^2r_{23}^2)$   
+ $8(r_{12}r_{23}r_{31} + r_{12}r_{24}r_{41} + r_{23}r_{34}r_{42} + r_{13}r_{34}r_{41})$   
+ $16(r_{12}r_{23}r_{34}r_{41} + r_{12}r_{24}r_{43}r_{31} + r_{13}r_{32}r_{24}r_{41})$   
 $-2r_{12}(2r_{13}r_{23} + 2r_{14}r_{24} + 4r_{13}r_{24}r_{34} + 4r_{14}r_{23}r_{34}))$   
 $-2r_{34}(2r_{13}r_{14} + 2r_{23}r_{24} + 4r_{13}r_{24}r_{12} + 4r_{23}r_{14}r_{12}))$   
 $+4r_{12}r_{34}(r_{13}r_{24} + r_{14}r_{23}).$ 

A careful cancellation leads to

$$Cov((X_{11}X_{21} - r_{12})^2, (X_{31}X_{41} - r_{34})^2)$$
  
=  $2(r_{13}^2 + r_{14}^2 + r_{23}^2 + r_{24}^2) + 4(r_{13}^2r_{24}^2 + r_{14}^2r_{23}^2)$   
 $+4(r_{12}r_{23}r_{31} + r_{12}r_{24}r_{41} + r_{23}r_{34}r_{42} + r_{13}r_{34}r_{41})$   
 $+4r_{12}r_{23}r_{34}r_{41} + 4r_{12}r_{24}r_{43}r_{31} + 16r_{13}r_{32}r_{24}r_{41}.$ 

By rewriting  $\frac{2(m-1)}{m^3} = \frac{2}{m^2} - \frac{2}{m^3}$ , we see from the above, (40) and (41) that  $\begin{aligned} & \operatorname{Cov}((B_1 - r_{12})^2, (B_2 - r_{34})^2) \\ &= \frac{1}{m^3} \Big[ 2(r_{13}^2 + r_{14}^2 + r_{23}^2 + r_{24}^2) + 4(r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2) \\ &+ 4(r_{12} r_{23} r_{31} + r_{12} r_{24} r_{41} + r_{23} r_{34} r_{42} + r_{13} r_{34} r_{41}) \\ &+ 4r_{12} r_{23} r_{34} r_{41} + 4r_{12} r_{24} r_{43} r_{31} + 16r_{13} r_{32} r_{24} r_{41} \Big] \\ &+ \Big( \frac{2}{m^2} - \frac{2}{m^3} \Big) (r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2 + 2r_{13} r_{32} r_{24} r_{41}), \end{aligned}$ 

which is again equal to

$$\frac{2}{m^2} \left[ r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2 + 2r_{13} r_{32} r_{24} r_{41} \right] \\ + \frac{1}{m^3} \left[ 2(r_{13}^2 + r_{14}^2 + r_{23}^2 + r_{24}^2) + 2(r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2) \right] \\ + 4(r_{12} r_{23} r_{31} + r_{12} r_{24} r_{41} + r_{23} r_{34} r_{42} + r_{13} r_{34} r_{41}) \\ + 4r_{12} r_{23} r_{34} r_{41} + 4r_{12} r_{24} r_{43} r_{31} + 12r_{13} r_{32} r_{24} r_{41} \right].$$

This and (38) imply the desired conclusion.

Now we compute the covariance of two squared sample covariances.

**LEMMA 6.9** Let  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T \in \mathbb{R}^4; 1 \leq j \leq m\}$  be *i.i.d.* random vectors with distribution  $N_4(\mathbf{0}, \mathbf{R})$ , where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each *i*. Let  $B_1$  and  $B_2$  be defined as in Lemma 6.8. Then

$$Cov(B_1^2, B_2^2) = \frac{1}{m} (r_{12}r_{23}r_{34}r_{41} + r_{12}r_{24}r_{43}r_{31}) + \frac{\delta_m}{m^2} \sum_{1 \le i < j \le 4} r_{ij}^2,$$

where  $|\delta_m| \leq \kappa$  and  $\kappa$  is a numerical constant not depending on m or  $\mathbf{R}$ .

#### Proof of Lemma 6.9. Write

$$B_1^2 = (B_1 - r_{12})^2 + 2r_{12}(B_1 - r_{12}) + r_{12}^2$$
 and  $B_2^2 = (B_2 - r_{34})^2 + 2r_{34}(B_2 - r_{34}) + r_{34}^2$ 

It follows that

$$\operatorname{Cov}(B_1^2, B_2^2) = \operatorname{Cov}((B_1 - r_{12})^2, (B_2 - r_{34})^2) + 2r_{12} \cdot \operatorname{Cov}((B_1 - r_{12}), (B_2 - r_{34})^2) + 2r_{34} \cdot \operatorname{Cov}((B_1 - r_{12})^2, (B_2 - r_{34})) + r_{12}r_{34} \cdot \operatorname{Cov}(B_1, B_2).$$

By Lemma 6.8(i) & (iv),

$$Cov((B_1 - r_{12})^2, (B_2 - r_{34})^2)$$

$$= E[(B_1 - r_{12})^2(B_2 - r_{34})^2] - E[(B_1 - r_{12})^2] \cdot E[(B_2 - r_{34})^2]$$

$$= \frac{2}{m^2} [r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2 + 2r_{13} r_{32} r_{24} r_{41}] + \frac{2}{m^3} [(r_{13}^2 + r_{14}^2 + r_{23}^2 + r_{24}^2) + (r_{13}^2 r_{24}^2 + r_{14}^2 r_{23}^2) + 2(r_{12} r_{23} r_{31} + r_{12} r_{24} r_{41} + r_{23} r_{34} r_{42} + r_{13} r_{34} r_{41})$$

 $+2r_{12}r_{23}r_{34}r_{41}+2r_{12}r_{24}r_{43}r_{31}+6r_{13}r_{32}r_{24}r_{41}].$ 

Use  $x^2y^2 \leq x^2$  and  $|xyzu| \leq |xyz| \leq |xy| \leq x^2 + y^2$  for any  $x, y, z, u \in [-1, 1]$  to see that

$$\left|\operatorname{Cov}\left((B_1 - r_{12})^2, (B_2 - r_{34})^2\right)\right| \le \frac{K_1}{m^2} \sum_{1 \le i < j \le 4} r_{ij}^2$$

where  $K_1$  is a numerical constant not depending on m or  $\mathbf{R}$ . Same bounds hold with another numerical constant  $K_2$  for  $2r_{12} \cdot \text{Cov}((B_1 - r_{12}), (B_2 - r_{34})^2)$  and  $2r_{34} \cdot \text{Cov}((B_1 - r_{12})^2, (B_2 - r_{34}))$  by Lemma 6.8(iii). Finally, by Lemma 6.8(ii),

$$r_{12}r_{34} \cdot \operatorname{Cov}(B_1, B_2) = \frac{1}{m}(r_{12}r_{24}r_{43}r_{31} + r_{12}r_{23}r_{34}r_{41}).$$

The proof is finished by combining all of the estimates.

Review sample correlation coefficient  $\hat{r}_{ij}$  defined in Theorem 4, the denominator is not easy to be handled. By the heuristic of the law of large numbers and the Taylor expansion, we are able to express  $\hat{r}_{ij}^2$  as a polynomial of Gaussian random variables plus an error. The following result provides basic computations for this step. It will be used in the derivation of Lemma 6.12.

**LEMMA 6.10** Let  $\{(X_i, Y_i)^T; 1 \le i \le m\}$  be i.i.d. 2-dimensional normal random vectors with  $EX_1 = EY_1 = 0$ ,  $EX_1^2 = EY_1^2 = 1$  and  $Cov(X_1, Y_1) = r$ . Define

$$A = \frac{X_1^2 + \dots + X_m^2}{m}$$
 and  $B = \frac{Y_1^2 + \dots + Y_m^2}{m}$ 

The following statements hold for all  $m \ge 2$  with  $|\delta_m| < \kappa$  where  $\kappa$  is a constant not depending on m or r.

(i) 
$$E[(1-A)(1-B)] = \frac{2}{m}r^2$$
.  
(ii)  $E[(1-A)(1-B)^2] = -\frac{8}{m^2}r^2$ .  
(iii)  $E[(1-A)(1-B)^3] = \frac{12r^2}{m^2} + \frac{\delta_m}{m^3}$ .  
(iv)  $E[(1-A)^2(1-B)^2] = \frac{8r^4+4}{m^2} + \frac{\delta_m}{m^3}$ .  
(v)  $E[(1-A)^i(1-B)^j] = \frac{\delta_m}{m^3}$  for  $(i,j) = (0,5), (1,4), (2,3)$ .

Proof of Lemma 6.10. Write

$$1 - A = \frac{1}{m} \sum_{i=1}^{m} (1 - X_i^2) \quad \text{and} \quad 1 - B = \frac{1}{m} \sum_{i=1}^{m} (1 - Y_i^2).$$
(43)

(i) Easily,

$$\operatorname{Var}(X_1 Y_1) = E\left[ (X_1 Y_1 - r)^2 \right] = E\left( X_1^2 Y_1^2 \right) - r^2 = r^2 + 1;$$
(44)

$$E[(1 - X_1^2)(1 - Y_1^2)] = 1 - E(X_1^2) - E(Y_1^2) + E(X_1^2Y_1^2) = 2r^2$$
(45)

by using (31). Due to independence,

$$E[(1-A)(1-B)] = \frac{1}{m}E[(1-X_1^2)(1-Y_1^2)] = 2r^2.$$

(ii) Write

$$m^{2} \cdot (1-B)^{2} = \sum_{i=1}^{m} (1-Y_{i}^{2})^{2} + 2 \sum_{1 \le i < j \le m} (1-Y_{i}^{2})(1-Y_{j}^{2}).$$
(46)

Evidently,  $1 - X_k^2$  is uncorrelated to  $(1 - Y_i^2)(1 - Y_j^2)$  for any i, j, k with  $i \neq j$ . Consequently,

$$E[(1-A)(1-B)^2] = \frac{1}{m^2} E[(1-X_1^2)(1-Y_1^2)^2].$$
(47)

Use the fact  $(1 - Y_1^2)^2 = 1 - 2Y_1^2 + Y_1^4$  to see

$$E\left[(1 - X_1^2)(1 - Y_1^2)^2\right] = -2E\left[(1 - X_1^2)Y_1^2\right] + E\left[(1 - X_1^2)Y_1^4\right]$$
$$= 2E\left[(1 - X_1^2)(1 - Y_1^2)\right] + E\left(Y_1^4\right) - E\left(X_1^2Y_1^4\right).$$
(48)

Write  $X_1 = rY_1 + \sqrt{1 - r^2}Y_1'$ , where  $Y_1' \sim N(0, 1)$  and  $Y_1'$  is independent of  $X_1$ . Then

$$E(X_1^2Y_1^4) = r^2 E(Y_1^6) + (1 - r^2)E(Y_1'^2)E(Y_1^4) = 15r^2 + 3(1 - r^2) = 3 + 12r^2.$$

Notice  $E(Y_1^4) = 3$  and  $E(Y_1^6) = 15$ . By (45) and (48),

$$E\left[(1-X_1^2)(1-Y_1^2)^2\right] = 2 \cdot 2r^2 + 3 - (3+12r^2) = -8r^2.$$

Then (ii) follows from (47).

(iii) Write

$$m^{3}(1-B)^{3} = \sum_{i=1}^{m} (1-Y_{i}^{2})^{3} + 3 \sum_{1 \le i \ne j \le m} (1-Y_{i}^{2})^{2}(1-Y_{j}^{2}) + \sum_{1 \le i \ne j \ne k \le m} (1-Y_{i}^{2})(1-Y_{j}^{2})(1-Y_{k}^{2}) := U_{1} + U_{2} + U_{3}.$$
(49)

Notice  $1 - X_k^2$  is uncorrelated to  $U_3$  and  $1 - X_i^2$  is uncorrelated to  $(1 - Y_i^2)^2(1 - Y_j^2)$  if  $k = i \neq j$ . This gives

$$m^{4} \cdot E[(1-A)(1-B)^{3}]$$

$$= \sum_{i=1}^{m} E[(1-X_{i}^{2})(1-Y_{i}^{2})^{3}] + 3\sum_{1 \le i \ne j \le m} E[(1-X_{j}^{2})(1-Y_{i}^{2})^{2}(1-Y_{j}^{2})]$$

$$= mE[(1-X_{1}^{2})(1-Y_{1}^{2})^{3}] + 3m(m-1)E[(1-Y_{1}^{2})^{2}] \cdot E[(1-X_{2}^{2})(1-Y_{2}^{2})]$$

The product of the last two expectations is equal to  $2 \cdot (2r^2) = 4r^2$  by (45). This implies

$$E[(1-A)(1-B)^3] = \frac{12r^2}{m^2} + \frac{\delta_m}{m^3}$$

with  $|\delta_m| < \kappa$  for all  $m \ge 2$ , where  $\kappa$  here and later represents a constant not depending on m or r, and can be different from line to line.

(iv) Similar to (46),

$$m^{2} \cdot (1-A)^{2} = \sum_{i=1}^{m} (1-X_{i}^{2})^{2} + 2 \sum_{1 \le i < j \le m} (1-X_{i}^{2})(1-X_{j}^{2})$$
  
:=  $V_{1} + V_{2}.$  (50)

Trivially,  $(1 - X_i^2)^2$  is uncorrelated to  $(1 - Y_j^2)(1 - Y_k^2)$  for any i, j, k with  $j \neq k$ . The same is true if X and Y are switched. Thus

$$m^{4} \cdot \operatorname{Cov}\left((1-A)^{2}, (1-B)^{2}\right)$$
  
=  $m \operatorname{Cov}\left((1-X_{1}^{2})^{2}, (1-Y_{1}^{2})^{2}\right) + 2m(m-1)\operatorname{Cov}\left((1-X_{1}^{2})(1-X_{2}^{2}), (1-Y_{1}^{2})(1-Y_{2}^{2})\right).$ 

Easily, by independence and (45),

$$\operatorname{Cov}\left((1-X_1^2)(1-X_2^2),(1-Y_1^2)(1-Y_2^2)\right) = \left[E(1-X_1^2)(1-Y_1^2)\right]^2 = 4r^4$$

Evidently,  $E[(1-A)^2] = \frac{2}{m}$ . It follows that

$$E[(1-A)^{2}(1-B)^{2}] = \operatorname{Cov}((1-A)^{2}, (1-B)^{2}) + E[(1-A)^{2}] \cdot E[(1-B)^{2}]$$
  
$$= \frac{1}{m^{4}} [m \operatorname{Cov}((1-X_{1}^{2})^{2}, (1-Y_{1}^{2})^{2}) + 8m(m-1)r^{4}] + \frac{4}{m^{2}}$$
  
$$= \frac{8r^{4}+4}{m^{2}} + \frac{\delta_{m}}{m^{3}}.$$

(v) We will study the three cases one by one.

 $Case \ 1: \ (i,j) = (0,5). \ \text{Let} \ Z_1, \cdots, Z_m \text{ be i.i.d. random variables with mean zero. Write} \\ (Z_1 + \dots + Z_m)^5 = \sum_{i=1}^m Z_i^5 + c_1 \sum_{i \neq j} Z_i^4 Z_j + c_2 \sum_{i \neq j} Z_i^3 Z_j^2 + c_3 \sum_{i \neq j \neq k} Z_i^3 Z_j Z_k \\ + c_4 \sum_{i \neq j \neq k} Z_i^2 Z_j^2 Z_k + c_5 \sum_{i \neq j \neq k} Z_i^2 Z_j Z_k Z_l + c_6 \sum_{a \neq i \neq j \neq k \neq l} Z_a Z_i Z_j Z_k Z_l,$ 

where  $c_1, \dots, c_6$  are numerical coefficients not depending on m or r. By independence, only the first and third sums of the right hand side above are non-zero if the expectation is taken on both sides. The total number of the terms appeared in the two sums is  $m + 2 \cdot {m \choose 2} = m^2$ . Take  $Z_i = 1 - Y_i^2$  for each i to have

$$E[(1-B)^5] = \frac{\delta_m}{m^3}$$

for all  $m \geq 2$ .

Case 2: (i, j) = (1, 4). Similar to Case 1,

$$(Z_{1} + \dots + Z_{m})^{4} = \sum_{i=1}^{m} Z_{i}^{4} + c_{1} \sum_{i \neq j} Z_{i}^{3} Z_{j} + 6 \sum_{i < j} Z_{i}^{2} Z_{j}^{2} + c_{3} \sum_{i \neq j \neq k} Z_{i}^{2} Z_{j} Z_{k}$$
$$+ c_{4} \sum_{i \neq j \neq k \neq l} Z_{i} Z_{j} Z_{k} Z_{l}.$$
(51)

Take  $Z_i = 1 - Y_i^2$  for each *i*. Recall (43), each term  $1 - X_i^2$  is uncorrelated to any term in the last two sums. By the same argument as in *Case 1*, we know  $E[(1-A)(1-B)^4] = \frac{1}{m^3}\delta_m$  for all  $m \ge 2$ .

Case 3: (i, j) = (2, 3). Review (49) and (50). Easily,  $U_3$  is uncorrelated to both  $V_1$  and  $V_2$ ;  $U_1$  is uncorrelated to  $V_2$ . It follows that

$$m^5 \cdot \operatorname{Cov}((1-A)^2, (1-B)^3) = \operatorname{Cov}(U_1, V_1) + \operatorname{Cov}(U_2, V_1) + \operatorname{Cov}(U_2, V_2).$$

By independence and the same argument as before, it is easy to see that  $\operatorname{Cov}(U_1, V_1) = m\delta_m$ ,  $\operatorname{Cov}(U_2, V_1) = m^2\delta'_m$  and  $\operatorname{Cov}(U_2, V_2) = m^2\delta''_m$ , where  $|\delta_m| + |\delta'_m| + |\delta''_m| \leq \kappa$  and  $\kappa$  is a constant not depending on m or r. Consequently,

$$\operatorname{Cov}((1-A)^2, (1-B)^3) = \frac{\delta_m}{m^3}.$$
(52)

From (50) and (49), it is readily seen  $E[(1-A)^2] = \frac{2}{m}$  and  $E[(1-B)^3] = \frac{1}{m^2} \cdot E(1-Y_1^2)^3$ . The conclusion then follows from these facts, (52) and the formula

$$E[(1-A)^{2}(1-B)^{3}] = \operatorname{Cov}((1-A)^{2}, (1-B)^{3}) + E[(1-A)^{2}] \cdot E[(1-B)^{3}].$$

The proof is finished.

# 6.2.2 Evaluations of Covariances between Monomials of Gaussian Random Variables

Let  $(X_1, X_2, X_3, X_4)^T$  be a 4-dimensional random vector with distribution  $N_4(\mathbf{0}, \mathbf{R})$  where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each *i*. One of major tasks in this section is showing  $|\operatorname{Cov}(X_1^{\alpha}X_2^{\beta}, X_3^{\gamma}X_4^{\delta})| \leq C \sum_{1 \leq i < j \leq 4} r_{ij}^2$  for some constant *C* depending only on non-negative integers  $\alpha, \beta, \gamma, \delta$ . Review the *t*-norm  $||U||_t = [E(|U|^t)]^{1/t}$  for any random variable *U* and  $t \geq 1$ . In the following lemma, by convention we regard  $\prod_{i=1}^{0} a_i = 1$  for any  $a_i$ .

**LEMMA 6.11** Let  $\{X_1, \dots, X_k, Y_1, \dots, Y_l\}$  be random variables with  $k \ge 0, l \ge 0$  and  $k+l \ge 1$ . Let  $p_1, \dots, p_k, q_1, \dots, q_l$  be positive integers. Assume, for each i, (a)  $X_i = \frac{1}{\sqrt{p_i}} \sum_{j=1}^{p_i} \xi_{ij}$  where  $\{\xi_{i1}, \dots, \xi_{ip_i}\}$  are *i.i.d.* with mean 0, and (b)  $Y_i = \frac{1}{q_i} \chi^2(q_i)$ . Then

$$E\left[\prod_{i=1}^{k} |X_{i}|^{\alpha_{i}} \cdot \prod_{j=1}^{l} Y_{j}^{\beta_{j}}\right] < C \cdot \prod_{i=1}^{k} || |\xi_{i1}|^{\alpha_{i}} ||_{k+l}$$

for any  $\alpha_i \geq 2/(k+l)$ ,  $\beta_j > (2-q_j)/[2(k+l)]$ ,  $1 \leq i \leq k$  and  $1 \leq j \leq l$ , where C is a constant depending on k, l,  $\alpha_i$  and  $\beta_j$  but not depending on  $p_i$  or  $q_j$ .

**Proof of Lemma 6.11**. First we assume  $k \ge 1$  and  $l \ge 1$ . By applying the Hölder inequality to the product of k + l terms, we see

$$E\left[\prod_{i=1}^{k} |X_{i}|^{\alpha_{i}} \cdot \prod_{j=1}^{l} Y_{j}^{\beta_{j}}\right] \leq \prod_{i=1}^{k} \left(E|X_{i}|^{\alpha_{i}(k+l)}\right)^{1/(k+l)} \cdot \prod_{j=1}^{l} \left(EY_{j}^{\beta_{j}(k+l)}\right)^{1/(k+l)}.$$
(53)

By assumption,  $\alpha_i(k+l) \ge 2$  for each *i*. According to the Marcinkiewicz-Zygmund inequality [see, for example, the proof of Corollary 2 on p. 387 from Chow and Teicher (1997)],

$$E|X_i|^{\alpha_i(k+l)} \le C(\alpha_i, k, l) \cdot E\left(|\xi_{i1}|^{\alpha_i(k+l)}\right),\tag{54}$$

where  $C(\alpha_i, k, l)$  is a constant depending on  $\alpha_i, k, l$  only. As a result,

$$\prod_{i=1}^{k} \left( E|X_{i}|^{\alpha_{i}(k+l)} \right)^{1/(k+l)} \le C(\alpha_{1}, \cdots, \alpha_{k}, k, l) \cdot \prod_{i=1}^{k} \||\xi_{i1}|^{\alpha_{i}}\|_{k+l}.$$
(55)

Second, by recalling the density of  $\chi^2(m)$  is  $f(x) := \frac{1}{\Gamma(\frac{m}{2})2^{m/2}} x^{(m/2)-1} e^{-x/2}$  for x > 0, we have

$$E[\chi^{2}(m)^{\beta}] = \frac{1}{\Gamma(\frac{m}{2})2^{m/2}} \int_{0}^{\infty} x^{(m+2\beta)/2-1} e^{-x/2} dx$$
  
$$= \frac{\Gamma(\frac{m}{2}+\beta)}{\Gamma(\frac{m}{2})2^{-\beta}}$$
(56)

for any real number  $\beta > -m/2$ . From the fact that  $\lim_{x\to\infty} \frac{\Gamma(x+a)}{\Gamma(x)x^a} = 1$  for any number  $a \in \mathbb{R}$ , it is seen that  $E[\chi^2(m)/m]^\beta \to 1$  as  $m \to \infty$ . This implies that

$$\sup E\left[\frac{\chi^2(m)}{m}\right]^{\beta} = C_{\beta} < \infty, \tag{57}$$

where the supremum is taken over all m satisfying  $m \ge 1$  and  $m \ge 2(1 - \beta)$  and where  $C_{\beta}$  depends on  $\beta$  only. The reason we choose m such that  $\frac{m}{2} + \beta \ge 1$  is because we need  $\frac{m}{2} + \beta$  to stay away from the only singular point 0 of  $\Gamma(x)$  defined on  $[0, \infty)$ . It follows that

$$E(|Y_j|^{\beta_j(k+l)}) = E\left[\left(\frac{\chi^2(q_j)}{q_j}\right)^{\beta_j(k+l)}\right] \le C_{\beta_j(k+l)} < \infty$$

provided  $q_j \ge 2[1 - \beta_j(k+l)]$ , or equivalently,  $\beta_j \ge (2 - q_j)/[2(k+l)]$ . This asserts that

$$\prod_{j=1}^{l} \left( E Y_j^{\beta_j(k+l)} \right)^{1/(k+l)} = C(k, l, \beta_1, \cdots, \beta_l) < \infty$$
(58)

under the assumption  $\beta_j \ge (2 - q_j)/[2(k + l)]$  for each  $1 \le j \le l$ . This joined with (53) and (55) leads to the desired inequality.

If  $k \ge 1$  and l = 0, then (53)-(55) still hold. If k = 0 and  $l \ge 1$ , then (53) and (58) are also true. The desired statements are then derived.

Recall sample correlation coefficient  $\hat{r}_{ij}$  defined in Theorem 4 from Section 6.2, the next result provides an estimate for the dependency between two denominators.

**LEMMA 6.12** Let  $\{(X_i, Y_i)^T; 1 \le i \le m\}$  be i.i.d. 2-dimensional normal random vectors with  $EX_1 = EY_1 = 0$ ,  $EX_1^2 = EY_1^2 = 1$  and  $Cov(X_1, Y_1) = r$ . Write

$$E\left(\frac{m}{X_1^2 + \dots + X_m^2} \cdot \frac{m}{Y_1^2 + \dots + Y_m^2}\right) = 1 + \frac{4 + 2r^2}{m} + \frac{12 + 8r^2 + 8r^4}{m^2} + \frac{\delta_m}{m^3}.$$
 (59)

Then  $|\delta_m| \leq \kappa$  for all  $m \geq 11$ , where  $\kappa$  is a constant not depending on m or r.

Before we present the proof let us have a quick check for two cases with r = 0 and r = 1. Recall (56). We have  $E_{\chi^2(m)} = \frac{1}{m-2}$  and  $E_{\chi^2(m)^2} = \frac{1}{(m-2)(m-4)}$  for any  $m \ge 5$ . If r = 0, then  $\{X_i, Y_i; 1 \le i \le m\}$  are i.i.d. N(0, 1). Consequently the left hand side of (59) is identical to

$$m^{2} \left[ E \frac{1}{\chi^{2}(m)} \right]^{2} = \frac{m^{2}}{(m-2)^{2}} = 1 + \frac{4}{m} + \frac{12}{m^{2}} + \frac{32m - 48}{m^{2}(m-2)^{2}},$$

which corresponds to the right hand side of (59) with r = 0. If r = 1, then  $X_i = Y_i$  for each *i*. Thus the left hand side of (59) becomes

$$m^{2} \cdot E \frac{1}{\chi^{2}(m)^{2}} = \frac{m^{2}}{(m-2)(m-4)} = 1 + \frac{6}{m} + \frac{28}{m^{2}} + \frac{120m - 224}{m^{2}(m-2)(m-4)}$$

which is equal to the right hand side of (59) with r = 1.

### Proof of Lemma 6.12. Define

$$A = \frac{X_1^2 + \dots + X_m^2}{m}$$
 and  $B = \frac{Y_1^2 + \dots + Y_m^2}{m}$ 

Set  $A_1 = 1 - A$  and  $B_1 = 1 - B$ . By the formula  $x^{-1} = 1 + (1 - x) + (1 - x)^2 + (1 - x)^3 + (1 - x)^4 + (1 - x)^5 + x^{-1}(1 - x)^6$ , we write

$$\frac{1}{AB} = \left(\frac{A_1^6}{A} + \sum_{i=0}^5 A_1^i\right) \left(\frac{B_1^6}{B} + \sum_{i=0}^5 B_1^i\right).$$

We will expand  $(\sum_{i=0}^{5} A_{1}^{i})(\sum_{i=0}^{5} B_{1}^{i})$  and write it as the sum of  $A_{1}^{a}B_{1}^{b}$ , and furthermore break the sum into two sums, the first of which is for the terms with  $a + b \leq 5$  and the second of which is for those with  $a + b \geq 6$ . Therefore,

$$\frac{1}{AB} = 1 + (A_1 + B_1) + (A_1^2 + A_1B_1 + A_2^2) + (A_1^3 + A_1^2B_1 + A_1B_1^2 + B_1^3) + (A_1^4 + A_1^3B_1 + A_1^2B_1^2 + A_1B_1^3 + B_1^4) + (A_1^5 + A_1^4B_1 + A_1^3B_1^2 + A_1^2B_1^3 + A_1B_1^4 + B_1^5) + \epsilon_m,$$
(60)

where

$$\epsilon_m := \frac{A_1^6}{A} \sum_{i=0}^5 B_1^i + \frac{B_1^6}{B} \sum_{i=0}^5 A_1^i + \frac{A_1^6 B_1^6}{AB} + p_m$$

and where  $p_m = \sum A_1^a B_1^b$  with the sum running over all  $a \ge 1$  and  $b \ge 1$  satisfying  $6 \le a + b \le 10$ . Obviously, by counting the number of the terms on the right hand side of (60), we know the total number of the terms in the sum of  $p_m$  is 36 - (1 + 2 + 3 + 4 + 5 + 6) = 15. Notice

$$A \sim \frac{\chi^2(m)}{m}, \quad B \sim \frac{\chi^2(m)}{m}, \quad \sqrt{m}A_1 = \frac{1}{\sqrt{m}}\sum_{i=1}^m (1 - X_i^2), \quad \sqrt{m}B_1 = \frac{1}{\sqrt{m}}\sum_{i=1}^m (1 - Y_i^2).$$

Review the notation in Lemma 6.11. We first consider the term  $E(A_1^6B_1^iA^{-1})$ . If i = 0, take k = l = 1,  $\alpha_1 = 6$ ,  $\beta_1 = -1$  and  $q_1 = m$ . Then  $\alpha_1 \ge 2/(k+l) = 1$  and  $\beta_1 > (2-q_1)/[2(k+l)] = (2-m)/4$  as  $m \ge 7$ . Similarly, if  $i \ge 1$ , take k = 2, l = 1,  $\alpha_1 = 6$ ,  $\alpha_2 = i$ ,  $\beta_1 = -1$  and  $q_1 = m$ . It is always true that  $\alpha_1 \land \alpha_2 \ge 2/(k+l) = 2/3$ . Also,  $\beta_1 > (2-q_1)/[2(k+l)] = (2-m)/6$  if  $m \ge 9$ . A similar but easier check can be done for  $A_1^a B_1^b$  with  $a \ge 1$ ,  $b \ge 1$  and  $a + b \ge 6$ . It then follows from Lemma 6.11 that

$$E\left(\frac{A_1^6}{A}B_1^i\right) = \frac{\delta(m,i)}{m^3} \text{ and } E(A_1^aB_1^b) = \frac{\delta(m,a,b)'}{m^3}$$
 (61)

as  $m \ge 9$ , where  $|\delta(m, i)| + |\delta(m, a, b)'| \le \kappa$  for all  $1 \le i \le 5$  and integers  $a \ge 1$  and  $b \ge 1$ with  $a + b \ge 6$ . Here and later  $\kappa$  represents a constant not depending on m or r and can be different from line to line. Now we turn to look at the term  $A_1^6 B_1^6 A^{-1} B^{-1}$ . Take k = l = 2,  $\alpha_1 = \alpha_2 = 6$ ,  $\beta_1 = \beta_2 = -1$  and  $q_1 = q_2 = m$  to see that, for each i = 1, 2, we have  $\alpha_i \ge 2/(k+l) = 1/2$ and  $\beta_i > (2-q_i)/[2(k+l)] = (2-m)/8$  provided  $m \ge 11$ . From Lemma 6.11 again we obtain

$$E\frac{A_1^6 B_1^6}{AB} = \frac{\delta_m}{m^6}$$
(62)

with  $|\delta(m)| \leq \kappa$  as  $m \geq 11$ . Combining (61) and (62) and using the symmetry of A and B as well as that of  $A_i$  and  $B_i$ , we see that

$$E\epsilon_m = \frac{\delta_m}{m^3} \tag{63}$$

with  $|\delta_m| \leq \kappa$  for all  $m \geq 11$ .

Back to (60), let us examine the expectation of each monomial of  $A_1$  and  $B_1$  on the right hand side. By independence,

$$E(A_1^2) = \frac{1}{m^2} \sum_{i=1}^m E[(1 - X_i^2)^2] = \frac{2}{m}$$
(64)

since  $\operatorname{Var}(X_1^2) = 2$ . Recall

$$(t_1 + \dots + t_m)^3 = \sum_{i=1}^m t_i^3 + 3\sum_{1 \le i \ne j \le m} t_i^2 t_j + \sum_{1 \le i \ne j \ne k \le m} t_i t_j t_k$$

for any real numbers  $t_1, \dots, t_m$ . Take  $t_i = 1 - X_i^2$  and use independence to see

$$E(A_1^3) = \frac{1}{m^2} E[(1 - X_1^2)^3] = \frac{1}{m^2} [1 - 3E(X_1^2) + 3E(X_1^4) - E(X_1^6)] = -\frac{8}{m^2}$$

since  $E(X_1^2) = 1$ ,  $E(X_1^4) = 3$  and  $E(X_1^6) = 15$ . Review

$$(t_1 + \dots + t_m)^4 = \sum_{i=1}^m t_i^4 + c_1 \sum_{i \neq j} t_i^3 t_j + 6 \sum_{i < j} t_i^2 t_j^2 + c_3 \sum_{i \neq j \neq k} t_i^2 t_j t_k$$
$$+ c_4 \sum_{i \neq j \neq k \neq l} t_i t_j t_k t_l$$

for any  $t_1, \dots, t_m$ . By taking  $t_i = 1 - X_i^2$  and using independence, we get

$$E(A_1^4) = \frac{1}{m^3} \cdot E[(1 - X_1^2)^4] + \frac{6}{m^4} \cdot \binom{m}{2} \cdot E[(1 - X_1^2)^2(1 - X_2^2)^2]$$
  
=  $\frac{12}{m^2} + \frac{\delta_m}{m^3}$ 

with  $|\delta_m| \leq \kappa$  for all  $m \geq 1$  since  $E\left[(1-X_1^2)^2(1-X_2^2)^2\right] = [E(1-X_1^2)^2]^2 = 4$ . Now, by (60),

$$E\left(\frac{1}{AB} - \epsilon_m\right) = 1 + E\left(A_1^2 + A_1B_1 + B_1^2\right) + E\left(A_1^3 + A_1^2B_1 + A_1B_1^2 + B_1^3\right) + E\left(A_1^4 + A_1^3B_1 + A_1^2B_1^2 + A_1B_1^3 + B_1^4\right) + E\left(A_1^5 + A_1^4B_1 + A_1^3B_1^2 + A_1^2B_1^3 + A_1B_1^4 + B_1^5\right).$$

From Lemma 6.10, the expressions of  $EA_1^i$  for i = 2, 3, 4 above as well as the symmetry of A and B, we obtain

$$E\left(\frac{1}{AB} - \epsilon_m\right) = 1 + \left(\frac{4}{m} + \frac{2}{m}r^2\right) + \left(-\frac{16}{m^2} - \frac{16}{m^2}r^2\right) \\ + \left(\frac{24}{m^2} + \frac{24r^2}{m^2} + \frac{8r^4 + 4}{m^2}\right) + \frac{\delta_m}{m^3} \\ = 1 + \frac{4 + 2r^2}{m} + \frac{12 + 8r^2 + 8r^4}{m^2} + \frac{\delta_m}{m^3}$$

with  $|\delta_m| \leq \kappa$  for all  $m \geq 2$ . This and (63) conclude

$$E\frac{1}{AB} = 1 + \frac{4+2r^2}{m} + \frac{12+8r^2+8r^4}{m^2} + \frac{\delta_m}{m^3}$$

with  $|\delta_m| \leq \kappa$  for all  $m \geq 11$ .

Recall the notation  $(2i-1)!! = 1 \cdot 3 \cdots (2i-1)$  for any integer  $i \ge 1$ . We set (-1)!! = 1 by convention.

**LEMMA 6.13** Let X and Y be N(0,1). Assume they are jointly normal with covariance r. Then the following hold.

(i) For any integers  $i \ge 1$  and  $j \ge 1$ , there exists a polynomial f(x) depending on i and j such that f(0) = 1 and  $E(X^{2i-1}Y^{2j-1}) = (2i-1)!! \cdot (2j-1)!! \cdot rf(r^2)$ .

(ii) For any integers  $i \ge 0$  and  $j \ge 0$ , there exists a polynomial g(x) depending on i and j such that  $E(X^{2i}Y^{2j}) = (2i-1)!! \cdot (2j-1)!! + r^2g(r^2)$ .

**Proof of Lemma 6.13.** Let Z be a random variable with distribution N(0, 1) and Z be independent of X and Y. Then we may write Y = rX + r'Z, where  $r' = \sqrt{1 - r^2}$ .

(i) Easily,

$$Y^{2j-1} = \sum_{k=0}^{2j-1} {2j-1 \choose k} r^k r'^{2j-k-1} X^k Z^{2j-k-1}.$$

By independence and the fact  $E[N(0,1)^n] = 0$  for any odd number  $n \ge 1$ , we have

$$E(X^{2i-1}Y^{2j-1}) = \sum_{k=0}^{2j-1} {\binom{2j-1}{k}} r^k r'^{2j-k-1} E(X^{2i+k-1}Z^{2j-k-1})$$
  

$$= \sum_{l=1}^{j} {\binom{2j-1}{2l-1}} r^{2l-1} r'^{2(j-l)} E(X^{2(i+l-1)}) \cdot E(Z^{2(j-l)})$$
  

$$= r \sum_{l=1}^{j} {\binom{2j-1}{2l-1}} (r^2)^{l-1} (1-r^2)^{j-l} \cdot (2(i+l)-3)!! \cdot (2(j-l)-1)!!$$
  

$$:= rh(r^2),$$
(65)

where we set k = 2l - 1 in the second identity and the fact  $E[N(0, 1)^n] = (n - 1)!!$  for even number  $n \ge 1$  is used in the last display. Obviously the last term in (65) is obviously a polynomial. Then

$$h(0) = \sum_{l=1}^{j} \binom{2j-1}{2l-1} (r^2)^{l-1} (1-r^2)^{j-l} \cdot (2(i+l)-3)!! \cdot (2(j-l)-1)!! \Big|_{r=0}$$
  
=  $(2j-1) \cdot (2(i+1)-3)!! \cdot (2(j-1)-1)!!$   
=  $(2i-1)!! \cdot (2j-1)!!.$ 

Set  $f(x) = g(x)[(2i-1)!! \cdot (2j-1)!!]^{-1}$ . The desired conclusion follows.

(ii) Recall  $E(X^{2i}) = (2i-1)!!$  and  $E(Y^{2j}) = (2j-1)!!$ . The conclusion is obviously true if (i, j) = (0, 0), (i, j) = (0, 1) and (i, j) = (1, 0), in which cases g(x) = 0 for each  $x \in \mathbb{R}$ . We next assume  $i \ge 1$  and  $j \ge 1$ . By the same argument as in (i),

$$Y^{2j} = (rX + r'Z)^{2j} = \sum_{k=0}^{2j} {\binom{2j}{k}} r^k r'^{2j-k} X^k Z^{2j-k}.$$

Then  $E(X^{2i}Y^{2j})$  is equal to

$$\sum_{k=0}^{2j} {2j \choose k} r^k r'^{2j-k} E\left(X^{2i+k} Z^{2j-k}\right)$$

$$= \sum_{l=0}^{j} {2j \choose 2l} r^{2l} r'^{2(j-l)} E\left(X^{2(i+l)}\right) \cdot E\left(Z^{2(j-l)}\right)$$

$$= (1-r^2)^j E\left(X^{2i}\right) \cdot E\left(Z^{2j}\right) + r^2 \sum_{l=1}^{j} {2j \choose 2l} (r^2)^{l-1} (1-r^2)^{j-l} E\left(X^{2(i+l)}\right) \cdot E\left(Z^{2(j-l)}\right),$$

where we set k = 2l in the first identity and single out the term with l = 0 in the last step. Obviously the last sum is a polynomial of  $r^2$ , say,  $h_1(r^2)$ . Write  $(1 - r^2)^j = 1 + r^2 h_2(r^2)$  with the function  $h_2(x)$  being a polynomial. Consequently,

$$E(X^{2i}Y^{2j}) = [1 + r^2h_2(r^2)]E(X^{2i}) \cdot E(Z^{2j}) + r^2h_1(r^2)$$
$$= (2i - 1)!! \cdot (2j - 1)!! + r^2g(r^2),$$

where  $g(x) := (2i-1)!! \cdot (2j-1)!!h_2(x) + h_1(x)$  is a polynomial. The proof is completed.  $\Box$ 

With the help of Lemma 6.13, we will obtain Lemmas 6.14-6.16 in the following. They estimate the size of  $|Cov(X_1^{d_1}X_2^{d_2}, X_3^{d_3}X_4^{d_4})|$  for non-negative integers  $d_1, d_2, d_3, d_4$  with special requirements.

**LEMMA 6.14** Let  $(X_1, X_2, X_3, X_4)^T$  be a 4-dimensional random vector with distribution  $N_4(\mathbf{0}, \mathbf{R})$  where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each *i*. Then, there exists a constant C > 0 depending on *i*, *j*, *k*, *l* but not on  $\mathbf{R}$  such that  $|Cov(X_1^{2i}X_2^{2j}, X_3^{2k}X_4^{2l})| \leq C \sum_{1 \leq i < j \leq 4} r_{ij}^2$  for all non-negative integers *i*, *j*, *k* and *l*.

**Proof of Lemma 6.14**. If i + j = 0, then i = j = 0, and hence  $X_1^{2i}X_2^{2j} = 1$ . The conclusion obviously holds. The same is true if k + l = 0. So we assume  $i + j \ge 1$  and  $k + l \ge 1$  next. Write

$$\operatorname{Cov}(X_1^{2i}X_2^{2j}, X_3^{2k}X_4^{2l}) = E\left(X_1^{2i}X_2^{2j}X_3^{2k}X_4^{2l}\right) - E\left(X_1^{2i}X_2^{2j}\right) \cdot E\left(X_3^{2k}X_4^{2l}\right).$$
(66)

By Lemma 6.13(ii), there exists polynomials  $g_1(x)$  and  $g_2(x)$  for  $x \in \mathbb{R}$  such that

$$E(X_1^{2i}X_2^{2j}) = (2i-1)!! \cdot (2j-1)!! + r_{12}^2 \cdot g_1(r_{12}^2);$$
(67)

$$E(X_3^{2k}X_4^{2l}) = (2k-1)!! \cdot (2l-1)!! + r_{34}^2 \cdot g_2(r_{34}^2).$$
(68)

Set  $r = (r_{12}, r_{13}, r_{14}, r_{23}, r_{24}, r_{34})^T \in \mathbb{R}^6$ . Then, by Lemma 6.4,  $E(X_1^{2i}X_2^{2j}X_3^{2k}X_4^{2l})$  is a multivariate polynomial of  $r_{ij}, 1 \le i < j \le 4$ . So we are able to write

$$E\left(X_1^{2i}X_2^{2j}X_3^{2k}X_4^{2l}\right) = C_0 + \sum_{1 \le i < j \le 4} C_{ij}r_{ij} + c(r),$$
(69)

where  $C_0$  and  $C_{ij}$  are constants and c(r) is a linear combination of  $\prod_{1 \le i < j \le 4} r_{ij}^{\alpha_{ij}}$  with  $2 \le \sum_{1 \le i < j \le 4} \alpha_{ij} \le i + j + k + l$ . Next we will use the fact that (69) holds for every  $r_{ij} \in [-1, 1]$  and  $1 \le i < j \le 4$  to identify the values of  $C_0$  and every  $C_{ij}$ .

Set  $r_{ij} = 0$  for all  $i \neq j$ . Then  $X_1, X_2, X_3, X_4$  are i.i.d. N(0, 1). It follows from (69) that

$$C_0 = E(X_1^{2i}) \cdot E(X_2^{2j}) \cdot E(X_3^{2k}) \cdot E(X_4^{2l})$$
  
=  $(2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!!.$ 

Review (69). We claim that  $C_{ij} = 0$  for all  $1 \le i < j \le 4$ . Take  $r_{ij} = 0$  for all  $1 \le i < j \le 4$ except (i, j) = (1, 2). Then the three random variables  $(X_1, X_2)^T \in \mathbb{R}^2$ ,  $X_3$  and  $X_4$  are independent. By (69),

$$E(X_1^{2i}X_2^{2j}) \cdot E(X_3^{2k}) \cdot E(X_4^{2l})$$
  
=  $(2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!! + C_{12}r_{12} + c_1(r_{12}),$  (70)

where  $c_1(x) = c_2 x^2 + c_3 x^3 + \dots + c_p x^p$  with  $2 \le p \le i + j + k + l$ . Use the fact  $E(X_3^{2k}) = (2k-1)!!$ ,  $E(X_4^{2l}) = (2l-1)!!$  and (67) to see

$$E(X_1^{2i}X_2^{2j}) \cdot E(X_3^{2k}) \cdot E(X_4^{2l})$$
  
=  $[(2i-1)!! \cdot (2j-1)!! + r_{12}^2 \cdot g_1(r_{12}^2)] \cdot (2k-1)!! \cdot (2l-1)!!$ 

Observe there are no linear terms of  $r_{12}$  on the right hand side. Compare this with (70), we see  $C_{12} = 0$ . By symmetry,  $C_{ij} = 0$  for all  $1 \le i < j \le 4$ . Combining this, (66)-(69), we get

 $Cov(X_1^{2i}X_2^{2j}, X_3^{2k}X_4^{2l})$  is identical to

$$(2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!! + c(r)$$

$$- [(2i-1)!! \cdot (2j-1)!! + r_{12}^2 \cdot g_1(r_{12}^2)] \cdot [(2k-1)!! \cdot (2l-1)!! + r_{34}^2 \cdot g_2(r_{34}^2)]$$

$$:= c_2(r).$$
(71)

Observe that  $\operatorname{Cov}(X_1^{2i}X_2^{2j}, X_3^{2k}X_4^{2l}) = c_2(r)$  is a linear combination of  $\prod_{1 \leq i < j \leq 4} r_{ij}^{\alpha_{ij}}$  with  $2 \leq \sum_{1 \leq i < j \leq 4} \alpha_{ij} \leq i + j + k + l$ . Notice  $|r_{ij}| \leq 1$ , and hence  $|r_{ij}^{\alpha_{ij}}| \leq r_{ij}^2$  for all  $\alpha_{ij} \geq 2$  and  $\prod_{1 \leq i < j \leq 4} r_{ij}^{\alpha_{ij}} \leq |r_{kl}|^{\alpha_{kl}} |r_{uv}|^{\alpha_{uv}}$  for any  $1 \leq k < l \leq 4$ ,  $1 \leq u < v \leq 4$ ,  $(k, l) \neq (u, v)$  with  $\alpha_{kl} + \alpha_{uv} \geq 2$ . Use the formula  $|xy| \leq \frac{1}{2}(x^2 + y^2)$  to get

$$|r_{kl}|^{\alpha_{kl}} |r_{uv}|^{\alpha_{uv}} \le r_{kl}^2 + r_{uv}^2 \le \sum_{1 \le i < j \le 4} r_{ij}^2.$$

This says that there exists a constant C > 0 depending on i, j, k, l but not on **R** such that

$$|\operatorname{Cov}(X_1^{2i}X_2^{2j}, X_3^{2k}X_4^{2l})| = |c_2(r)| \le C \sum_{1 \le i < j \le 4} r_{ij}^2.$$
(72)

The proof is completed.

**LEMMA 6.15** Let  $(X_1, X_2, X_3, X_4)^T$  be the same as in Lemma 6.14. Then, there exists a constant C > 0 depending on i, j, k, l but not on  $\mathbf{R}$  such that  $|Cov(X_1^{2i-1}X_2^{2j-1}, X_3^{2k}X_4^{2l})| \leq C \sum_{1 \leq i < j \leq 4} r_{ij}^2$  for all non-negative integers i, j, k, l with  $i \geq 1$  and  $j \geq 1$ .

**Proof of Lemma 6.15**. If k+l=0, then k=l=0, and hence  $X_3^{2k}X_4^{2l}=1$ . The conclusion trivially holds. So we assume and  $k+l \ge 1$  next.

By Lemma 6.13(i), there exists a polynomial f(x) for  $x \in \mathbb{R}$  with f(0) = 1 and

$$E(X_1^{2i-1}X_2^{2j-1}) = (2i-1)!! \cdot (2j-1)!! \cdot r_{12}f(r_{12}^2)$$
(73)

for all integers  $i \ge 1$  and  $j \ge 1$ . Write

$$\operatorname{Cov}(X_1^{2i-1}X_2^{2j-1}, X_3^{2k}X_4^{2l}) = E(X_1^{2i-1}X_2^{2j-1}X_3^{2k}X_4^{2l}) - E(X_1^{2i-1}X_2^{2j-1}) \cdot E(X_3^{2k}X_4^{2l}).$$
(74)

_	

Then, by Lemma 6.4,

$$E\left(X_1^{2i-1}X_2^{2j-1}X_3^{2k}X_4^{2l}\right) = D_0 + \sum_{1 \le i < j \le 4} D_{ij}r_{ij} + d(r),$$
(75)

where  $D_0$  and  $D_{ij}$ 's are constants and d(r) is a linear combination of  $\prod_{1 \le i < j \le 4} r_{ij}^{\alpha_{ij}}$  with  $2 \le \sum_{1 \le i < j \le 4} \alpha_{ij} \le i + j + k + l - 1$ . We now evaluate the values of  $D_0$  and  $D_{ij}$ 's.

Take  $r_{ij} = 0$  for all  $i \neq j$ . Then  $X_1, X_2, X_3, X_4$  are i.i.d. with distribution N(0, 1). From (75), we see

$$D_0 = E(X_1^{2i-1}) \cdot E(X_2^{2j-1}) \cdot E(X_3^{2k}) \cdot E(X_4^{2l}) = 0$$

since  $E(X_1^{2i-1}) = 0$ . Then (75) becomes

$$E(X_1^{2i-1}X_2^{2j-1}X_3^{2k}X_4^{2l}) = \sum_{1 \le i < j \le 4} D_{ij}r_{ij} + d(r).$$
(76)

Take  $r_{ij} = 0$  for all  $1 \le i < j \le 4$  except (i, j) = (1, 2). Then the three random variables  $(X_1, X_2)^T \in \mathbb{R}^2$ ,  $X_3$  and  $X_4$  are independent. It follows from (73) and (76) that

$$(2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!! \cdot f(r_{12}^2)r_{12} = D_{12}r_{12} + d_1(r_{12})r_{12}$$

where  $d_1(x) = c_2 x^2 + c_3 x^3 + \dots + c_p x^p$  with  $2 \le p \le i + j + k + l - 1$ . Set  $d_2(x) = c_2 x + c_3 x^2 + \dots + c_p x^{p-1}$ . The above implies

$$(2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!! \cdot f(r_{12}^2) = D_{12} + d_2(r_{12}).$$

Take  $r_{12} = 0$ . Recall f(0) = 1. Then

$$D_{12} = (2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!!$$

Now we claim  $D_{ij} = 0$  for all  $(i, j) \neq (1, 2)$ . In fact, set  $r_{ij} = 0$  for all  $1 \leq i < j \leq 4$  except (i, j) = (1, 3). Then  $X_2$  is independent of  $(X_1, X_3, X_4)^T$ . Use the fact  $E(X_2^{2j-1}) = 0$  and independence to see from (76) that

$$0 = D_{13}r_{13} + d_3(r_{13}), (77)$$

where  $d_3(x) = a_2 x^2 + \cdots + a_p x^p$  with  $2 \le p \le i + j + k + l - 1$ . Divide both sides of (77) by  $r_{13}$  and then set  $r_{13} = 0$ , we get  $D_{13} = 0$ . Similarly,  $D_{ij} = 0$  for all  $(i, j) \ne (1, 2)$ . Therefore, by (76),

$$E\left(X_1^{2i-1}X_2^{2j-1}X_3^{2k}X_4^{2l}\right) = (2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!! \cdot r_{12} + d(r).$$
(78)

Since f(x) in (73) is a polynomial with f(0) = 1, we are able to write  $f(x) = 1 + xf_1(x)$  for each  $x \in \mathbb{R}$ , where  $f_1(x)$  is a polynomial. Hence,

$$E(X_1^{2i-1}X_2^{2j-1})$$
  
=  $(2i-1)!! \cdot (2j-1)!! \cdot r_{12} + (2i-1)!! \cdot (2j-1)!! \cdot r_{12}^3 f_1(r_{12}^2).$ 

Joining this by (68), (74) and (78), we have that  $Cov(X_1^{2i-1}X_2^{2j-1}, X_3^{2k}X_4^{2l})$  is equal to

$$\begin{aligned} &(2i-1)!! \cdot (2j-1)!! \cdot (2k-1)!! \cdot (2l-1)!! \cdot r_{12} + d(r) \\ &- \left[ (2i-1)!! \cdot (2j-1)!! \cdot r_{12} + (2i-1)!! \cdot (2j-1)!! \cdot r_{12}^3 f_1(r_{12}^2) \right] \\ &\cdot \left[ (2k-1)!! \cdot (2l-1)!! + r_{34}^2 \cdot g_2(r_{34}^2) \right] \\ &:= d_4(r), \end{aligned}$$

where  $g_2(x)$  is a polynomial. Observe there are no linear terms of  $r_{ij}$ ,  $1 \le i < j \le 4$  in the expression of  $d_4(r)$ . Consequently,  $d_4(r)$  is a linear combination of  $\prod_{1 \le i < j \le 4} r_{ij}^{\alpha_{ij}}$  with  $2 \le \sum_{1 \le i < j \le 4} \alpha_{ij} \le i + j + k + l - 1$ . The conclusion then follows from the same argument between (71) and (72).

**LEMMA 6.16** Let  $(X_1, X_2, X_3, X_4)^T$  be the same as in Lemma 6.14. Then, there exists a constant C > 0 depending on i, j, k, l but not on  $\mathbf{R}$  such that  $|Cov(X_1^{2i-1}X_2^{2j-1}, X_3^{2k-1}X_4^{2l-1})| \leq C \sum_{1 \leq i < j \leq 4} r_{ij}^2$  for all positive integers i, j, k, l.

Proof of Lemma 6.16. Write

$$\operatorname{Cov}(X_1^{2i-1}X_2^{2j-1}, X_3^{2k-1}X_4^{2l-1}) = E(X_1^{2i-1}X_2^{2j-1}X_3^{2k-1}X_4^{2l-1}) - E(X_1^{2i-1}X_2^{2j-1}) \cdot E(X_3^{2k-1}X_4^{2l-1}).$$
(79)

By Lemma 6.13(i),

$$E(X_3^{2k-1}X_4^{2l-1}) = (2k-1)!! \cdot (2l-1)!! \cdot r_{34}g(r_{34}^2)$$

for some polynomial g(x). This together with (73) implies that

$$\left| E\left(X_1^{2i-1}X_2^{2j-1}\right) \cdot E\left(X_3^{2k-1}X_4^{2l-1}\right) \right| \le C_1 |r_{12}r_{34}| \le C \sum_{1 \le i < j \le 4} r_{ij}^2, \tag{80}$$

where  $C_1$  is a constant depending on i, j, k, l but not on **R**. By Lemma 6.4,

$$E(X_1^{2i-1}X_2^{2j-1}X_3^{2k-1}X_4^{2l-1}) = B_0 + \sum_{1 \le i < j \le 4} B_{ij}r_{ij} + b(r),$$
(81)

where  $B_0$  and  $B_{ij}$ 's are constants and b(r) is a linear combination of  $\prod_{1 \le i < j \le 4} r_{ij}^{\alpha_{ij}}$  with  $2 \le \sum_{1 \le i < j \le 4} \alpha_{ij} \le i + j + k + l - 2$ . We claim that

$$B_0 = B_{ij} = 0 \tag{82}$$

for all  $1 \leq i < j \leq 4$ . In fact, take  $r_{ij} = 0$  for all  $1 \leq i < j \leq 4$ , then  $X_1, X_2, X_3, X_4$  are i.i.d. N(0, 1)-distributed random variables. The left hand side of (81) is zero because of independence. The right hand side of (81) is equal to  $B_0$ . This concludes  $B_0 = 0$ . Then (81) becomes

$$E\left(X_1^{2i-1}X_2^{2j-1}X_3^{2k-1}X_4^{2l-1}\right) = \sum_{1 \le i < j \le 4} B_{ij}r_{ij} + b(r).$$
(83)

For this identity we take  $r_{ij} = 0$  for all  $1 \le i < j \le 4$  except (i, j) = (1, 2). Then the three random variables  $(X_1, X_2)^T \in \mathbb{R}^2$ ,  $X_3$  and  $X_4$  are independent. By independence and the fact  $E(X_3^{2k-1}) = 0$ , the left hand side of (83) is zero. Hence, (83) is reduced to

$$0 = B_{12}r_{12} + b_1(r_{12}),$$

where  $b_1(x) = c_2 x^2 + c_3 x^3 + \dots + c_p x^p$  with  $2 \le p \le i + j + k + l - 2$ . Set  $b_2(x) = c_2 x + c_3 x^2 + \dots + c_p x^{p-1}$ . Then  $0 = B_{12} + b_2(r_{12})$ . Take  $r_{12} = 0$  to get  $B_{12} = 0$ . By symmetry, we know  $B_{ij} = 0$  for all  $1 \le i < j \le 4$ . So (82) has been verified. It follows that (81) is

reduced to  $E(X_1^{2i-1}X_2^{2j-1}X_3^{2k-1}X_4^{2l-1}) = b(r)$ . By employing the same argument between (71) and (72), we get

$$\left| E \left( X_1^{2i-1} X_2^{2j-1} X_3^{2k-1} X_4^{2l-1} \right) \right| \le C_2 \sum_{1 \le i < j \le 4} r_{ij}^2,$$

where  $C_2$  is a constant depending on i, j, k, l but not on **R**. This joined (79) and (80) yields the desired inequality.

The following fact supplies a convenient tool to handle the covariance between products of independent random variables. Recall  $\|\xi\|_q = (E|\xi|^q)^{1/q}$  for any random variable  $\xi$  and  $q \ge 1$ .

**LEMMA 6.17** For  $k \ge 2$ , let  $\{(U_i, V_i) \in \mathbb{R}^2; 1 \le i \le k\}$  be independent random vectors. Assume  $C := 3 \prod_{i=1}^k (1 + \|U_i\|_k)(1 + \|V_i\|_k) < \infty$  and  $K = C^k$ . Then

$$\left| Cov\left(\prod_{i=1}^{k} U_{i}, \prod_{i=1}^{k} V_{i}\right) \right| \leq K \cdot \sum_{i=1}^{k} |Cov(U_{i}, V_{i})|.$$

**Proof of Lemma 6.17**. The inequality holds obviously for k = 1. Assume now  $k \ge 2$ . By definition,

$$Cov(U_1U_2, V_1V_2) = E(U_1U_2V_1V_2) - E(U_1U_2) \cdot E(V_1V_2)$$
  
=  $E(U_1V_1) \cdot E(U_2V_2) - EU_1 \cdot EU_2 \cdot EV_1 \cdot EV_2.$  (84)

Write  $E(U_1V_1) = Cov(U_1, V_1) + EU_1 \cdot EV_1$  and  $E(U_2V_2) = Cov(U_2, V_2) + EU_2 \cdot EV_2$ . Plug the two identities into (84) to see

$$Cov(U_1U_2, V_1V_2) = Cov(U_1, V_1) \cdot Cov(U_2, V_2) + EU_2 \cdot EV_2 \cdot Cov(U_1, V_1) + EU_1 \cdot EV_1 \cdot Cov(U_2, V_2).$$
(85)

Since  $k \ge 2$ , we have from the Hölder inequality that  $E|\xi| \le ||\xi||_F \le ||\xi||_k$  for any random variable  $\xi$ . By the Cauchy-Schwartz inequality,  $E|U_1V_1| \le ||U_1||_2 \cdot ||V_1||_2 \le ||U_1||_k \cdot ||V_1||_k$ . As a result,  $|\text{Cov}(U_1, V_1)|$  is bounded by

$$E|U_1V_1| + E|U_1| \cdot E|V_1| \le 2||U_1||_k ||V_1||_k.$$
(86)

Set  $\tau_j = 3 \prod_{i=1}^{j} (1 + \|U_i\|_k) (1 + \|V_i\|_k)$  for  $j = 1, \cdots, k$ . We claim that  $\left| \operatorname{Cov} \left( \prod_{i=1}^{k} U_i, \prod_{i=1}^{k} V_i \right) \right| \le (\tau_1 \cdots \tau_k) \cdot \sum_{i=1}^{k} |\operatorname{Cov}(U_i, V_i)|.$ (87)

In fact, by applying (86) to (85), we see

$$|\operatorname{Cov}(U_{1}U_{2}, V_{1}V_{2})| \leq 2\|U_{1}\|_{k}\|V_{1}\|_{k} \cdot |\operatorname{Cov}(U_{2}, V_{2})| + \|U_{2}\|_{k}\|V_{2}\|_{k} \cdot |\operatorname{Cov}(U_{1}, V_{1})| + \|U_{1}\|_{k}\|V_{1}\|_{k} \cdot |\operatorname{Cov}(U_{2}, V_{2})| \leq \left[3\prod_{i=1}^{2}\left(1+\|U_{i}\|_{k}\right)\left(1+\|V_{i}\|_{k}\right)\right] \cdot \left[|\operatorname{Cov}(U_{1}, V_{1})| + |\operatorname{Cov}(U_{2}, V_{2})|\right].$$

$$(88)$$

So claim (87) holds for k = 2 due to the fact  $\tau_j \ge 1$  for each j. Now we assume  $k \ge 3$  and use induction to complete the proof. Assume

$$\left|\operatorname{Cov}\left(\prod_{i=1}^{j} U_{i}, \prod_{i=1}^{j} V_{i}\right)\right| \leq (\tau_{1} \cdots \tau_{j}) \cdot \sum_{i=1}^{j} |\operatorname{Cov}(U_{i}, V_{i})|$$

$$(89)$$

for some  $2 \leq j < k$ . By assumption,  $(U_{j+1}, V_{j+1})$  and  $(\prod_{i=1}^{j} U_i, \prod_{i=1}^{j} V_i)$  are independent. We obtain from (88) that

$$\left|\operatorname{Cov}\left(\prod_{i=1}^{j+1} U_{i}, \prod_{i=1}^{j+1} V_{i}\right)\right| \leq \left[3\left(1 + \|U_{j+1}\|_{k}\right)\left(1 + \|U_{1} \cdots U_{j}\|_{k}\right)\left(1 + \|V_{1} \cdots V_{j}\|_{k}\right)\right] \cdot \left[\left|\operatorname{Cov}(U_{j+1}, V_{j+1})\right| + \left|\operatorname{Cov}\left(\prod_{i=1}^{j} U_{i}, \prod_{i=1}^{j} V_{i}\right)\right|\right].$$
(90)

Since  $2 \le j < k$ , then we have from the Hölder inequality that

$$\|U_1\cdots U_j\|_k = \left[E\left(U_1\cdots U_j\cdot \underbrace{1\cdots 1}_{k-j}\right)\right]^{1/k} \le \prod_{i=1}^j \|U_i\|_k$$

The above also holds if the symbol "U" is replaced by "V". Thus,

$$3(1 + \|U_1 \cdots U_j\|_k)(1 + \|V_1 \cdots V_j\|_k)(1 + \|U_{j+1}\|_k)(1 + \|V_{j+1}\|_k) \le \tau_{j+1}.$$

By (89) and (90),

$$\begin{aligned} \left| \operatorname{Cov} \left( \prod_{i=1}^{j+1} U_i, \prod_{i=1}^{j+1} V_i \right) \right| &\leq \tau_{j+1} \cdot \left[ |\operatorname{Cov} (U_{j+1}, V_{j+1})| + (\tau_1 \cdots \tau_j) \cdot \sum_{i=1}^{j} |\operatorname{Cov} (U_i, V_i)| \right] \\ &\leq (\tau_1 \cdots \tau_{j+1}) \cdot \sum_{i=1}^{j+1} |\operatorname{Cov} (U_i, V_i)|. \end{aligned}$$

This confirms (87). The proof is completed by taking  $C = \tau_k$  and  $K = C^k$ .

### 6.2.3 Combinatorics

In this section we will work on some combinatorics problems. They will be used to evaluate covariances between squared sample correlations coefficients in Section 6.2.4. We always assume  $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_m, \gamma_1, \dots, \gamma_m, \delta_1, \dots, \delta_m$  are non-negative integers. Set  $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_m), \boldsymbol{\beta} = (\beta_1, \dots, \beta_m), \boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m)$  and  $\boldsymbol{\delta} = (\delta_1, \dots, \delta_m)$ .

**LEMMA 6.18** Let  $m \ge 2$ ,  $a \ge 0$  and  $b \ge 1$  be integers. Then the following hold with constant K depending on a and b but not m.

(i) Let  $N_1$  be the total number of non-negative integer solutions  $(x_1, \dots, x_m)$  of  $x_1 + \dots + x_m = a$ , then  $N_1 \leq Km^a$ .

(ii) Let  $N_2$  be the total number of non-negative integer solutions  $(x_1, \dots, x_m)$  of  $x_1 + \dots + x_m = a$  with  $x_1 \ge b$ . Then  $N_2 \le Km^{a-b}$ .

(iii) Given  $1 \leq n < m$  and  $c_1 \geq 1, \dots, c_n \geq 1$  with  $c_1 + \dots + c_n \leq a$ , let  $N_3$  the total number of non-negative integer solutions  $(x_1, \dots, x_m)$  of  $x_1 + \dots + x_m = a$  with  $x_i \geq c_i$  for  $1 \leq i \leq n$ . Then  $N_3 \leq Km^{a-c_1-\dots-c_n}$ .

A quick comment is that (ii) is a special case of (iii). We single it out because  $N_2$  has a much neater statement and it will be used very frequently.

**Proof of Lemma 6.18**. (i) If a = 0, the only non-negative integer solution of  $x_1 + \cdots + x_m = a$  is  $(0, \dots, 0)$ . Then  $N_1 = 1$  and the conclusion follows with any constant  $K \ge 1$ . We assume next that  $a \ge 1$ . It is well-known that

$$N_1 = \binom{m+a-1}{a} \le (m+a-1)^a \le (1+a)^a m^a.$$
(91)

(iii) Set  $y_i = x_i - c_i$  for  $i = 1, \dots, n$  and  $y_i = x_i$  for  $n+1 \le i \le m$ . Then  $N_3$  is equal to the total number of non-negative integer solutions  $(y_1, \dots, y_m)$  of  $y_1 + \dots + y_m = a - c_1 - \dots - c_n$ . From (i) we see  $N_3 \le Km^{a-c_1-\dots-c_n}$ . The proof is completed. The statement (ii) follows because it is a special case of (iii).  $\Box$ 

In the following when we say a non-negative integer solution  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  of a certain equation, we mean  $(\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_m, \gamma_1, \dots, \gamma_m, \delta_1, \dots, \delta_m)$  satisfies that equation with each of  $\{\alpha_i, \beta_i, \gamma_i, \delta_i; 1 \leq i \leq m\}$  being a non-negative integer.

**LEMMA 6.19** Let  $m \ge 4$  and  $\alpha, \beta, \gamma, \delta$  be non-negative integers. Let  $N_1$  be the total number of non-negative integer solutions of  $(\alpha, \beta, \gamma, \delta)$  satisfying

$$\sum_{i=1}^{m} \alpha_i = \alpha, \quad \sum_{i=1}^{m} \beta_i = \beta, \quad \sum_{i=1}^{m} \gamma_i = \gamma, \quad \sum_{i=1}^{m} \delta_i = \delta.$$
(92)

Set  $I_1 := \{1\} \cup \{2 \le i \le m; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies } (92) \text{ and } \alpha_i + \beta_i \ge 1\}$  and

$$I_2 := \{2,3\} \cup \{i \in \{1,4,5,\cdots,m\}; (\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta}) \text{ satisfies (92) and } \gamma_i + \delta_i \ge 1\}.$$

Let  $N_2$  be the total number of non-negative integer solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and  $I_1 \cap I_2 \neq \emptyset$ . Then, there exists a constant K depending on  $\alpha, \beta, \gamma, \delta$  but not m such that (i)  $N_1 \leq K \cdot m^{\alpha+\beta+\gamma+\delta}$ ; (ii)  $N_2 \leq K \cdot m^{\alpha+\beta+\gamma+\delta-1}$ .

**Proof of Lemma 6.19**. First, recall the fact that the total number of non-negative integer solutions of  $x_1 + \cdots + x_m = k$  for any non-negative integer k is  $\binom{m+k-1}{k}$ . Therefore, considering the four equations in (92) separately, the total numbers of non-negative integer solutions are

$$\binom{m+\alpha-1}{\alpha}, \ \binom{m+\beta-1}{\beta}, \ \binom{m+\gamma-1}{\gamma} \text{ and } \binom{m+\delta-1}{\delta},$$
 (93)

respectively.

(i) By (91) and (93),

$$N_{1} \leq \binom{m+\alpha-1}{\alpha} \binom{m+\beta-1}{\beta} \binom{m+\gamma-1}{\gamma} \binom{m+\delta-1}{\delta} \leq (1+\alpha)^{\alpha} (1+\beta)^{\beta} (1+\gamma)^{\gamma} (1+\delta)^{\delta} \cdot m^{\alpha+\beta+\gamma+\delta}.$$
(94)

The conclusion follows by taking  $K = (1 + \alpha)^{\alpha} (1 + \beta)^{\beta} (1 + \gamma)^{\gamma} (1 + \delta)^{\delta}$ .

(ii) If  $\alpha + \beta = 0$  and  $\gamma + \delta = 0$ , then  $I_1 = \{1\}$  and  $I_2 = \{2, 3\}$ , and hence  $I_1 \cap I_2 = \emptyset$ . Thus N = 0. The conclusion holds. So we assume next that either  $\alpha + \beta \ge 1$  or  $\gamma + \delta \ge 1$ . Notice  $I_1 \cap I_2 = A_1 \cup A_2 \cup A_3$ , where

$$A_{1} = \{i \in \{1\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \gamma_{i} + \delta_{i} \neq 0\};$$
  

$$A_{2} = \{\{i \in \{2, 3\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \alpha_{i} + \beta_{i} \neq 0\};$$
  

$$A_{3} = \{i \in \{4, 5, \cdots, m\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \alpha_{i} + \beta_{i} \neq 0 \text{ and } \gamma_{i} + \delta_{i} \neq 0\}.$$

Hence, if  $I_1 \cap I_2 \neq \emptyset$ , then either  $A_1 \neq \emptyset$ ,  $A_2 \neq \emptyset$  or  $A_3 \neq \emptyset$ . Let us consider the three scenarios one by one next.

Scenario 1:  $A_1 \neq \emptyset$ . In this situation,  $\gamma_1 + \delta_1 \geq 1$ . Consequently, either  $\gamma_1 \geq 1$ or  $\delta_1 \geq 1$ . Thus, taking b = 1 in Lemma 6.18(i) and (ii), we know the total number of non-negative integer solutions of  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and  $A_1 \neq \emptyset$  is bounded by

$$K_1 m^{\alpha} \cdot K_1 m^{\beta} \cdot K_1 m^{\gamma - 1} \cdot K_1 m^{\delta} + K_1 m^{\alpha} \cdot K_1 m^{\beta} \cdot K_1 m^{\gamma} \cdot K_1 m^{\delta - 1}$$
  
=  $2(K_1)^4 \cdot m^{\delta + \beta + \gamma + \delta - 1}$ 

where  $K_1$  here and below is a constant depending on  $\alpha, \beta, \gamma, \delta$  but not m.

Scenario 2:  $A_2 \neq \emptyset$ . In this situation, either  $\alpha_2 + \beta_2 \ge 1$  or  $\alpha_3 + \beta_3 \ge 1$ . Similar to the first case, the total number of non-negative integer solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and  $A_2 \neq \emptyset$  is bounded by  $2(K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-1} + 2(K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-1} = 4(K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-1}$ .

Scenario 3:  $A_3 \neq \emptyset$ . In this situation, there exists  $i \in \{4, 5, \dots, m\}$  such that  $\alpha_i + \beta_i \geq 1$ and  $\gamma_i + \delta_i \geq 1$ . For fixed i, if  $\alpha_i + \beta_i \geq 1$  and  $\gamma_i + \delta_i \geq 1$ , then one of the four cases must be true: (a)  $\alpha_i \geq 1$  and  $\gamma_i \geq 1$ ; (b)  $\alpha_i \geq 1$  and  $\delta_i \geq 1$ ; (c)  $\beta_i \geq 1$  and  $\gamma_i \geq 1$ ; (d)  $\beta_i \geq 1$  and  $\delta_i \geq 1$ . From Lemma 6.18(i) and (ii) again, we have that the total number of non-negative integer solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and (a) is dominated by  $K_1 m^{\alpha-1} \cdot K_1 m^{\beta} \cdot K_1 m^{\gamma-1} \cdot K_1 m^{\delta} = (K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-2}$ . By symmetry, the same inequality holds if "(a)" is replaced by (b), (c) and (d), respectively. In conclusion, for fixed i, the total number of non-negative integer solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and  $\alpha_i + \beta_i \geq 1$ and  $\gamma_i + \delta_i \geq 1$  is controlled by  $4(K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-2}$ . Now,  $i \in \{4, 5, \dots, m\}$  has at most m choices. Then the total number of non-negative integer solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and  $A_3 \neq \emptyset$  is bounded by  $m \cdot 4(K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-2} = 4(K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-1}$ . Finally, add the bounds up in the above three scenarios, we get  $N_2 \leq 10(K_1)^4 \cdot m^{\delta+\beta+\gamma+\delta-1}$ . The proof is completed by taking  $K = 10(K_1)^4$ .

**LEMMA 6.20** Assume  $m \geq 5$  and  $\alpha, \beta, \gamma, \delta$  are non-negative integers. Define

$$S = \{5 \le i \le m; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies } (92), \alpha_i + \beta_i \ge 1 \text{ and } \gamma_i + \delta_i \ge 1\}.$$

Then the following statements hold with a constant K depending on  $\alpha, \beta, \gamma, \delta$  but not m.

(i) The total number of solutions of  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and  $S \neq \emptyset$  is bounded by  $Km^{\alpha+\beta+\gamma+\delta-1}$ .

(ii) The total number of solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) with  $\gamma_1 + \delta_1 \geq 2$  is bounded by  $K \cdot m^{\alpha+\beta+\gamma+\delta-2}$ .

**Proof of Lemma 6.20.** (i) Since  $S \neq \emptyset$ , then there exists some  $5 \leq i \leq m$  such that  $\alpha_i + \beta_i \geq 1$  and  $\gamma_i + \delta_i \geq 1$ . According to Scenario 3 in the proof of Lemma 6.19, the total number of solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92),  $\alpha_i + \beta_i \geq 1$  and  $\gamma_i + \delta_i \geq 1$  is dominated by  $K \cdot m^{\delta+\beta+\gamma+\delta-2}$ , where K is a constant depending on  $\alpha, \beta, \gamma, \delta$  but not m. Noticing  $5 \leq i \leq m$ , then the desired number is bounded by  $(m-4) \cdot Km^{\delta+\beta+\gamma+\delta-2} \leq Km^{\delta+\beta+\gamma+\delta-1}$ .

(ii) Let K be a constant in Lemma 6.18(i) with  $a = \alpha$  or  $\beta$ . Also, the K satisfies Lemma 6.18(ii) with  $a \in \{\gamma, \delta\}$  and  $b \in \{1, 2\}$ . Since  $\gamma_1 + \delta_1 \geq 2$ , then one of the three cases must be true: (a)  $\gamma_1 \geq 2$ , (b)  $\delta_1 \geq 2$  or (c)  $\gamma_1 \geq 1$  and  $\delta_1 \geq 1$  simultaneously. By Lemma 6.18(ii), the total number of solutions  $(\boldsymbol{\gamma}, \boldsymbol{\delta})$  of the last two equations from (92) with  $\gamma_1 \geq 2$  is no more than  $Km^{\gamma+\delta-2}$ . The same holds if " $\gamma_1 \geq 2$ " is replaced by " $\delta_1 \geq 2$ ". Similarly, by Lemma 6.18(ii) again, the total number of solutions  $(\boldsymbol{\gamma}, \boldsymbol{\delta})$  of the last two equations from (92) satisfying  $\gamma_1 \geq 1$  and  $\delta_1 \geq 1$  is bounded by  $Km^{\gamma-1} \cdot Km^{\delta-1} = K^2m^{\gamma+\delta-2}$ . Consequently, the total number of solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) with  $\gamma_1 + \delta_1 \geq 2$ is bounded by

$$Km^{\alpha} \cdot Km^{\beta} \cdot (Km^{\gamma+\delta-2} + Km^{\gamma+\delta-2} + K^2m^{\gamma+\delta-2}) = (K^4 + 2K^2)m^{\alpha+\beta+\gamma+\delta-2}.$$

Therefore, the desired conclusion follows by regarding  $K^4 + 2K^2$  as new constant K.

**LEMMA 6.21** Assume  $m \geq 5$  and  $\alpha, \beta, \gamma, \delta$  are non-negative integers. Define

$$S = \{i \in \{3,4\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \alpha_i + \beta_i \ge 1\} \cup \{j \in \{1,2\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \gamma_j + \delta_j \ge 1\}.$$

Let  $T_{m,1}$  be the set of  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and |S| = 1. Let  $T_{m,2}$  be the set of  $(\alpha, \beta, \gamma, \delta)$ satisfying (92) and  $|S| \ge 2$ . Let  $T_{m,3}$  be the set of  $(\alpha, \beta, \gamma, \delta)$  satisfying (92), |S| = 1 and one of the following:

(1)  $\alpha_i + \beta_i \ge 1$  for some  $i \in \{1, 2\}$ ;

(2)  $\gamma_j + \delta_j \ge 1$  for some  $j \in \{3, 4\}$ ;

- (3)  $\alpha_k + \beta_k \ge 2$  for some  $5 \le k \le m$ ;
- (4)  $\gamma_l + \delta_l \ge 2$  for some  $5 \le l \le m$ ;

(5)  $\alpha_t + \beta_t = 1$  and  $\gamma_t + \delta_t = 1$  simultaneously for some  $5 \leq t \leq m$ . Then, there exists a constant K depending on  $\alpha, \beta, \gamma, \delta$  but not m such that  $|T_{m,1}| \leq Km^{\alpha+\beta+\gamma+\delta-1}$ ,  $|T_{m,2}| \leq Km^{\alpha+\beta+\gamma+\delta-2}$  and  $|T_{m,3}| \leq K \cdot m^{\alpha+\beta+\gamma+\delta-2}$ .

**Proof of Lemma 6.21.** If  $\alpha = \beta = \gamma = \delta = 0$ , then  $T_{m,1} = T_{m,2} = T_{m,3} = \emptyset$ , the conclusion obviously holds. So we assume next that at least one of the four numbers is positive. Note that the bounds in the conclusions are  $Km^{\alpha+\beta+\gamma+\delta-1}$  and  $Km^{\alpha+\beta+\gamma+\delta-2}$ . So, in case one of  $\{\alpha, \beta, \gamma, \delta\}$  is zero, say,  $\alpha = 0$ , any discussions below related to  $\alpha$  will disappear by convention. In the following we will prove the three conclusions one by one.

The bound for  $T_{m,1}$ . If |S| = 1, then one of the following four situations must occur: (a)  $\alpha_i \geq 1$  for some  $i \in \{3, 4\}$ ; (b)  $\beta_i \geq 1$  for some  $i \in \{3, 4\}$ ; (c)  $\gamma_j \geq 1$  for some  $j \in \{1, 2\}$ ; (d)  $\delta_j \geq 1$  for some  $j \in \{1, 2\}$ . If  $\alpha_i \geq 1$ , by Lemma 6.18(ii), the total number of non-negative integer solutions  $\boldsymbol{\alpha}$  of  $\alpha_1 + \cdots + \alpha_m = \alpha$  is no more than  $K_1 m^{\alpha-1}$ . Here and later  $K_1$ represents a constant depending on  $\alpha, \beta, \gamma, \delta$  but not m, and could be different from line to line. By Lemma 6.18(i), the total number of points  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and (a) is controlled by

$$K_1 m^{\alpha - 1} \cdot K_1 m^{\beta} \cdot K_1 m^{\gamma} \cdot K_1 m^{\delta} = (K_1)^4 \cdot m^{\alpha + \beta + \gamma + \delta - 1}.$$

Likewise the same bound holds if "(a)" is replaced by "(b)", "(c)" or "(d)". This implies  $|T_{m,1}|$  is dominated by the sum of the four bounds, that is,  $4(K_1)^4 \cdot m^{\alpha+\beta+\gamma+\delta-1}$ .

The bound for  $T_{m,2}$ . The assumption  $|S| \ge 2$  implies one of the following three statements must be true: (e)  $\alpha_3 + \beta_3 \ge 1$  and  $\alpha_4 + \beta_4 \ge 1$ ; (f)  $\gamma_1 + \delta_1 \ge 1$  and  $\gamma_2 + \delta_2 \ge 1$ ; (g)  $\alpha_i + \beta_i \ge 1$ for some  $i \in \{3, 4\}$  and  $\gamma_j + \delta_j$  for some  $j \in \{1, 2\}$ .

Under (e), one of the next four cases has to be true: (e1)  $\alpha_3 \ge 1$  and  $\alpha_4 \ge 1$ ; (e2)  $\alpha_3 \ge 1$ and  $\beta_4 \ge 1$ ; (e3)  $\alpha_4 \ge 1$  and  $\beta_3 \ge 1$ ; (e4)  $\beta_3 \ge 1$  and  $\beta_4 \ge 1$ . By Lemma 6.18(i) and (ii), the total number of points ( $\alpha, \beta, \gamma, \delta$ ) satisfying (92) and (e1) is no more than

$$K_1 m^{\alpha-2} \cdot K_1 m^{\beta} \cdot K_1 m^{\gamma} \cdot K_1 m^{\delta} = (K_1)^4 \cdot m^{\alpha+\beta+\gamma+\delta-2}.$$

By the same spirit, the total number of points  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and (e2) is controlled by

$$K_1 m^{\alpha-1} \cdot K_1 m^{\beta-1} \cdot K_1 m^{\gamma} \cdot K_1 m^{\delta} = (K_1)^4 \cdot m^{\alpha+\beta+\gamma+\delta-2}.$$

By similar discussions, the same conclusion above also holds if "(e2)" is replaced by "(e3)" and "(e4)", respectively, and " $(K_1)^4$  is replaced by another polynomial of  $K_1$ . In conclusion, by summing the four bounds corresponding to (e1) – (e4), we see that the total number of points ( $\alpha, \beta, \gamma, \delta$ ) satisfying (92) and (e) is no more than  $K_1 \cdot m^{\alpha+\beta+\gamma+\delta-2}$ . Similarly, the same conclusion holds if "(e)" is replaced by "(f)" and "(g)", respectively. The desired conclusion is then yielded by adding up the three bounds corresponding to (e), (f) and (g).

The bound for  $T_{m,3}$ . Let  $A_1$  be the set of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92), |S| = 1 and (1). Similarly, we define  $A_2, A_3, A_4, A_5$  with "(1)" is replaced by "(2)", "(3)", "(4)", "(5)", respectively. It suffices to show

$$|A_i| \le C_i \cdot m^{\alpha + \beta + \gamma + \delta - 2} \tag{95}$$

for i = 1, 2, 3, 4, 5, where  $C_i$  is a constant depending on  $\alpha, \beta, \gamma, \delta$  but not m.

We first look into  $A_1$  and  $A_2$ . Assuming |S| = 1 and (1), then there are two possibilities:  $\alpha_i + \beta_i \ge 1$  and  $\alpha_j + \beta_j \ge 1$  for a pair (i, j) with  $i, j \in \{1, 2, 3, 4\}$  and  $i \ne j$ ;  $\alpha_i + \beta_i \ge 1$  and  $\gamma_j + \delta_j \ge 1$  for a pair (i, j) with  $i, j \in \{1, 2\}$ . Review the analysis of case (e) and (g) above and the conclusion that the total number of points  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and (e) is no more than  $K_1 \cdot m^{\alpha+\beta+\gamma+\delta-2}$ . We know (95) is true for i = 1. By symmetry, (95) is also true for i = 2.

Now we work on  $A_3$  and  $A_4$ . For fixed  $k \in \{5, \dots, m\}$ , the assumption  $\alpha_k + \beta_k \ge 2$  hints that either  $\alpha_k \ge 2, \beta_k \ge 2$  or the third possibility that  $\alpha_k \ge 1$  and  $\beta_k \ge 1$ . On the other hand, the condition |S| = 1 implies that either  $\alpha_i + \beta_i \ge 1$  for some  $i \in \{3, 4\}$  or  $\gamma_j + \delta_j \ge 1$  for some  $j \in \{1, 2\}$ . In total we see  $3 \times 2 = 6$  scenarios. The only scenario we have not encountered so far comes from the combination  $\alpha_k \ge 2$  and  $\alpha_i + \beta_i \ge 1$  for some  $i \in \{3, 4\}$ . In this case, either  $\alpha_k \ge 2$  and  $\alpha_i \ge 1$  for some  $i \in \{3, 4\}$  or the second possibility  $\alpha_k \ge 2$  and  $\beta_i \ge 1$  for some  $i \in \{3, 4\}$ . By Lemma 6.18(ii) and (iii), the total number of points  $(\alpha, \beta)$  satisfying (92) and this combination is bounded by  $K_1(2m^{\alpha-3} \cdot m^\beta + m^{\alpha-2} \cdot (2m^{\beta-1}) = (4K_1)m^{\alpha+\beta-3}$ . By using this and earlier argument, we have the same bound for any of the six scenarios. Adding them up and noting k has m - 4 choices, we obtain (95) for i = 3. Similarly, (95) also holds for i = 4.

Finally we study  $A_5$ . Fix  $5 \le t \le m$ . Then the assumptions that  $\alpha_t + \beta_t = 1$  and  $\gamma_t + \delta_t = 1$  have four possibilities:  $\alpha_t = 1$  and  $\gamma_t = 1$ ;  $\alpha_t = 1$  and  $\delta_t = 1$ ;  $\beta_t = 1$  and  $\gamma_t = 1$ ;  $\beta_t = 1$  and  $\delta_t = 1$ . As aforementioned, the condition |S| = 1 implies that either  $\alpha_i + \beta_i \ge 1$  for some  $i \in \{3, 4\}$  or  $\gamma_j + \delta_j \ge 1$  for some  $j \in \{1, 2\}$ . So there are eight combinations with a common feature that the values of three different members of  $\{\alpha_i, \beta_i, \gamma_i, \delta_i; 1 \le i \le m\}$  are required to be at least 1. By Lemma 6.18 and the assumption that t has no more than m choices, we know (95) is true for i = 5.

After the verification of (95) for i = 1, 2, 3, 4, 5, we obtain the bound for  $T_{m,3}$ . Observe the three upper bounds for  $T_{m,1}$ ,  $T_{m,2}$  and  $T_{m,3}$  are involved with polynomials of  $K_1$ . We choose K to be the maximum of the three polynomials. The whole proof is completed.  $\Box$  **LEMMA 6.22** Assume  $m \ge 5$  and  $\alpha, \beta, \gamma, \delta$  are non-negative integers. Let S be the set of  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and one of the following holds:

(*i*)  $\alpha_i + \beta_i \ge 1$  for some  $i \in \{1, 2, 3\}$ ;

(*ii*)  $\gamma_i + \delta_i \ge 1$  for some  $i \in \{1, 2, 3\}$ ;

(iii)  $\alpha_i + \beta_i \geq 2 \text{ or } \gamma_i + \delta_i \geq 2 \text{ for some } 4 \leq i \leq m;$ 

(iv)  $\alpha_i + \beta_i \geq 1$  and  $\gamma_i + \delta_i \geq 1$  simultaneously for some  $1 \leq i \leq m$ .

Then  $|S| \leq Km^{\alpha+\beta+\gamma+\delta-1}$  for some constant K depending on  $\alpha, \beta, \gamma, \delta$  but not m.

**Proof of Lemma 6.22**. The proof is very similar to that of Lemma 6.21 and is even easier. We omit the details.  $\Box$ 

## 6.2.4 Evaluation of Covariances between Polynomials of Gaussian Random Variables

With the previous preparation, we are now ready to study covariances between polynomials of Gaussian random variables. The basic setting is that

Let 
$$m \ge 5$$
 and  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T \in \mathbb{R}^4; 1 \le j \le m\}$  be i.i.d. random vectors  
with distribution  $N_4(\mathbf{0}, \mathbf{R})$ , where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each  $i$ . (96)

In this section, K and  $K_1$  always represent constants depending on  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  but not m or  $\mathbf{R}$ , and can be different from line to line. Review the notation  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  before the statement of Lemma 6.18.

**LEMMA 6.23** Assume (96) holds. Let  $\{a, b, c, d, a_i, b_i, c_i, d_i; 1 \le i \le m\}$  be non-negative integers with  $a = \sum_{i=1}^{m} a_i$ ,  $b = \sum_{i=1}^{m} b_i$ ,  $c = \sum_{i=1}^{m} c_i$ ,  $d = \sum_{i=1}^{m} d_i$ . Define  $U_i = X_{i1}^{a_i} X_{i2}^{b_i}$  and  $V_i = X_{i3}^{c_i} X_{i4}^{d_i}$  for  $1 \le i \le m$ . If  $a_i + b_i$  and  $c_i + d_i$  are both even for each  $1 \le i \le m$ , then

$$\left| Cov\left(\prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i\right) \right| \le K \sum_{1 \le i < j \le 4} r_{ij}^2.$$

**Proof of Lemma 6.23.** If a+b=0, then  $U_i=1$  for each i, and the conclusion trivially holds. If c+d=0, then  $V_i=1$  for each i, and the conclusion is still valid. Now we assume that both  $a + b \ge 1$  and  $c + d \ge 1$ . For any random variable  $\xi$ , its  $L_s$ -norm  $\|\xi\|_s = (E|\xi|^s)^{1/s}$  is nondecreasing in  $s \ge 1$  by Hölder's inequality. Furthermore, by the same inequality, since  $X_{ij} \sim N(0,1)$  for each i, j, we have  $E(|X_{11}|^{2a_is}) \le E(|X_{11}|^{(2a+1)s})^{a_i/(2a+1)} \le 1 + E(|X_{11}|^{(2a+1)s})$ . A similar conclusion holds for  $E(|X_{11}|^{2b_is})$ . Consequently,

$$||U_i||_s^s = E(|X_{i1}|^{sa_i} \cdot |X_{i2}^{sb_i}|) \leq [E(|X_{11}|^{2a_is})]^{1/2} \cdot [E(|X_{11}|^{2b_is})]^{1/2}$$
  
$$\leq [1 + E(|X_{11}|^{(2a+1)s})] \cdot [1 + E(|X_{11}|^{(2b+1)s})]$$

Hence,

$$\max\{\|U_i\|_s, \|V_i\|_s; 1 \le i \le m\} \le K_1,$$
(97)

where  $K_1$  depends on a, b, c, d and  $s \ge 1$ . By definition,  $a = \sum_{i=1}^{m} a_i$ , hence  $|\{1 \le i \le m; a_i \ge 1\}| \le a$ . The same is also true for the analogue of b, c and d, respectively. Set  $\Psi = \{1 \le i \le m; a_i + b_i \ge 1 \text{ or } c_i + d_i \ge 1\}$ . For any  $i \in \Psi$ , either  $a_i \ge 1, b_i \ge 1, c_i \ge 1$  or  $d_i \ge 1$ , it follows that  $|\Psi| \le a + b + c + d$ . On the contrary, if  $i \notin \Psi$  then  $U_i = V_i = 1$ , therefore

$$\operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = \operatorname{Cov}\left(\prod_{i\in\Psi} U_{i}, \prod_{i\in\Psi} V_{i}\right).$$

Set  $k = |\Psi| \ge 1$ . Then  $C(k) := 3 \prod_{i=1}^{k} (1 + ||U_i||_k) (1 + ||V_i||_k) \le C(l)$  with l = a + b + c + dsince  $\|\cdot\|_s$  is non-decreasing in s. By Lemma 6.17, there exists a constant K > 0 depending on a, b, c, d but not m such that

$$\left|\operatorname{Cov}\left(\prod_{i\in\Psi}U_{i},\prod_{i\in\Psi}V_{i}\right)\right| \leq K\cdot\sum_{i\in\Psi}\left|\operatorname{Cov}(U_{i},V_{i})\right| \leq K\cdot|\Psi|\cdot\max_{1\leq i\leq m}|\operatorname{Cov}(U_{i},V_{i})|.$$

Use the fact  $|\Psi| \le a + b + c + d$  to see

$$\left|\operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right)\right| \leq (a+b+c+d)K \cdot \max_{1 \leq i \leq m} \left|\operatorname{Cov}(U_{i}, V_{i})\right|$$

For non-negative integers x and y with x + y being even, we know that x and y have to be both even or both odd. The conclusion then follows from Lemmas 6.14-6.16. **LEMMA 6.24** Assume the setting in (96). Define

$$A_i = \frac{1}{m} \sum_{j=1}^m X_{ij}^2, \ i = 1, 2, 3, 4.$$
(98)

For given non-negative integers  $\alpha, \beta, \gamma, \delta$  and  $q \in \{1, 2\}$ , we have

$$\left| Cov((X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{3q}X_{4q})^2 A_3^{\gamma} A_4^{\delta}) \right| \le K \sum_{1 \le i < j \le 4} r_{ij}^2.$$
<sup>(99)</sup>

Proof of Lemma 6.24. Write

$$(mA_1)^{\alpha} = \left(\sum_{j=1}^m X_{1j}^2\right)^{\alpha} = \sum \frac{\alpha!}{\alpha_1! \cdots \alpha_m!} X_{11}^{2\alpha_1} \cdots X_{1m}^{2\alpha_m};$$
(100)

$$(mA_2)^{\beta} = \left(\sum_{j=1}^m X_{2j}^2\right)^{\beta} = \sum \frac{\beta!}{\beta_1! \cdots \beta_m!} X_{21}^{2\beta_1} \cdots X_{2m}^{2\beta_m};$$
(101)

$$(mA_3)^{\gamma} = \left(\sum_{j=1}^m X_{3j}^2\right)^{\gamma} = \sum \frac{\gamma!}{\gamma_1! \cdots \gamma_m!} X_{31}^{2\gamma_1} \cdots X_{3m}^{2\gamma_m};$$
(102)

$$(mA_4)^{\delta} = \left(\sum_{j=1}^m X_{4j}^2\right)^{\delta} = \sum \frac{\delta!}{\delta_1! \cdots \delta_m!} X_{41}^{2\delta_1} \cdots X_{4m}^{2\delta_m},$$
(103)

where  $\alpha_i$ ,  $\beta_i$ ,  $\gamma_i$  and  $\delta_i$  are non-negative integers for each *i* satisfying

$$\sum_{i=1}^{m} \alpha_i = \alpha, \quad \sum_{i=1}^{m} \beta_i = \beta, \quad \sum_{i=1}^{m} \gamma_i = \gamma, \quad \sum_{i=1}^{m} \delta_i = \delta,$$

respectively. This restriction is exactly the same as (92). To avoid repetition in the future, once this restriction is used, we will always quote (92).

First, we consider the case q = 1. Notice

$$m^{\alpha+\beta+\gamma+\delta} \cdot \text{Cov}\left((X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{3q}X_{4q})^2 A_3^{\gamma} A_4^{\delta}\right)$$
(104)

is a linear combination of  $\mathcal{N}_1$  terms of the form

$$\operatorname{Cov}\left(\prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i\right)$$

with positive coefficients no more than  $\alpha!\beta!\gamma!\delta!$ , where

$$U_{1} = X_{11}^{2\alpha_{1}+2} X_{21}^{2\beta_{1}+2} \text{ and } U_{i} = X_{1i}^{2\alpha_{i}} X_{2i}^{2\beta_{i}};$$
  

$$V_{1} = X_{31}^{2\gamma_{1}+2} X_{41}^{2\delta_{1}+2} \text{ and } V_{i} = X_{3i}^{2\gamma_{i}} X_{4i}^{2\delta_{i}}$$
(105)

for  $2 \leq i \leq m$ . Here  $N_1$  is the total number of non-negative integer solutions of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$ satisfying the set of equations from (92). By Lemma 6.19(i),

$$N_1 \le K_1 \cdot m^{\alpha + \beta + \gamma + \delta}.$$
(106)

$$m^{\alpha+\beta+\gamma+\delta} \cdot \left| \operatorname{Cov}\left( (X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{3q}X_{4q})^2 A_3^{\gamma} A_4^{\delta} \right) \right|$$
  

$$\leq (K_1 \alpha! \beta! \gamma! \delta!) \cdot m^{\alpha+\beta+\gamma+\delta} \cdot \max \left| \operatorname{Cov}\left( \prod_{i=1}^m U_i, \prod_{i=1}^m V_i \right) \right|,$$
(107)

where the maximum is taken over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92). Note that

$$(2\alpha_1 + 2) + \sum_{i=2}^{m} 2\alpha_i = 2\alpha + 2; \quad (2\beta_1 + 2) + \sum_{i=2}^{m} 2\beta_i = 2\beta + 2;$$
  
$$(2\gamma_1 + 2) + \sum_{i=2}^{m} 2\gamma_i = 2\gamma + 2; \quad (2\delta_1 + 2) + \sum_{i=2}^{m} 2\delta_i = 2\delta + 2.$$
(108)

By Lemma 6.23,

$$\left|\operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right)\right| \leq K_{1} \sum_{1 \leq i < j \leq 4} r_{ij}^{2}.$$
(109)

Combining this with (107), we arrive at

$$m^{\alpha+\beta+\gamma+\delta} \cdot \left| \operatorname{Cov}\left( (X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{3q}X_{4q})^2 A_3^{\gamma} A_4^{\delta} \right) \right|$$
  
$$\leq K \cdot m^{\alpha+\beta+\gamma+\delta} \cdot \sum_{1 \leq i < j \leq 4} r_{ij}^2,$$
(110)

where  $K = K_1^2 \alpha! \beta! \gamma! \delta!$ . So (99) follows for the case q = 1.

For the case q = 2, we keep  $U_i$  in (105) unchanged but modify  $V_i$  such that  $V_2 = X_{32}^{2\gamma_2+2}X_{42}^{2\delta_2+2}$  and  $V_i = X_{3i}^{2\gamma_i}X_{4i}^{2\delta_i}$  for all  $i = 1, 3, \dots, m$ . By Lemma 6.23, (109) still holds. From (107) we then get (99) for the case q = 2. The proof is completed. **LEMMA 6.25** Assume the setting in (96). Let  $A_i$  be defined as in (98). Given non-negative integers  $\alpha, \beta, \gamma, \delta$ , set

$$I_m(a,b) = Cov((X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{3a}X_{4a})(X_{3b}X_{4b}) A_3^{\gamma} A_4^{\delta})$$

for integers  $a \ge 1$  and  $b \ge 1$ . Then

$$|I_m(a,b)| \le \begin{cases} K \sum_{1 \le i < j \le 4} r_{ij}^2 & \text{if } (a,b) = (1,2); \\ \frac{K}{m} \sum_{1 \le i < j \le 4} r_{ij}^2 & \text{if } (a,b) = (2,3). \end{cases}$$

**Proof of Lemma 6.25**. We will use the same notation as in the proof of Lemma 6.24. Review (92) and (100). We will consider the two cases for (a, b) separately, that is, (a, b) = (1, 2) or (a, b) = (2, 3).

Case 1: (a, b) = (1, 2). Set

$$U_1 = X_{11}^{2(\alpha_1+1)} X_{21}^{2(\beta_1+1)} \text{ and } U_i = X_{1i}^{2\alpha_i} X_{2i}^{2\beta_i} \text{ for } 2 \le i \le m;$$
(111)

$$V_1 = X_{31}^{2\gamma_1+1} X_{41}^{2\delta_1+1}, \quad V_2 = X_{32}^{2\gamma_2+1} X_{42}^{2\delta_2+1} \quad \text{and} \quad V_i = X_{3i}^{2\gamma_i} X_{4i}^{2\delta_i}$$
(112)

for  $3 \leq i \leq m$ . As before, let  $N_1$  be the total number of solutions  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  of (92) with a bound provided in (106). Then

$$m^{\alpha+\beta+\gamma+\delta} \cdot \operatorname{Cov}((X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{31}X_{41})(X_{32}X_{42}) A_3^{\gamma} A_4^{\delta})$$

is a linear combination of  $N_1$  terms of the form  $\operatorname{Cov}(\prod_{i=1}^m U_i, \prod_{i=1}^m V_i)$  with positive coefficients no more than  $\alpha!\beta!\gamma!\delta!$ . From the restriction in (92), we know  $\alpha_i \in \{0, \dots, \alpha\}$ ,  $\beta_i \in \{0, \dots, \beta\}, \gamma_i \in \{0, \dots, \gamma\}$  and  $\delta_i \in \{0, \dots, \delta\}$  for each *i*. By Lemma 6.23 and a discussion similar to (108), we obtain

$$\max \left| \operatorname{Cov}(\prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i) \right| \le K_1 \sum_{1 \le i < j \le 4} r_{ij}^2,$$

where the maximum is taken over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92). By (107), we obtain the bound for  $I_m(1, 2)$ .

*Case 2:* (a, b) = (2, 3). Again,

$$m^{\alpha+\beta+\gamma+\delta} \cdot \operatorname{Cov}\left((X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{32}X_{42})(X_{33}X_{43}) A_3^{\gamma} A_4^{\delta}\right)$$
(113)

is a linear combination of  $N_1$  terms of the form  $\text{Cov}(\prod_{i=1}^m U_i, \prod_{i=1}^m V_i)$  with positive coefficients no more than  $\alpha!\beta!\gamma!\delta!$ , where  $U_i$  is as in (111) and

$$V_1' = X_{31}^{2\gamma_1} X_{41}^{2\delta_1}, \quad V_2' = X_{32}^{2\gamma_2+1} X_{42}^{2\delta_2+1}, \quad V_3' = X_{33}^{2\gamma_3+1} X_{43}^{2\delta_3+1} \text{ and } \quad V_i' = X_{3i}^{2\gamma_i} X_{4i}^{2\delta_i}$$
(114)

for  $4 \leq i \leq m$ . Set

$$I_1 = \{1\} \cup \{2 \le i \le m; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \alpha_i + \beta_i \ge 1\};$$
  
$$I_2 = \{2, 3\} \cup \{i \in \{1, 4, 5, \cdots, m\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \gamma_i + \delta_i \ge 1\}.$$

Recalling (96),  $\{(U_i, V_i)^T; 1 \le i \le m\}$  are independent and  $m \ge 5$ . Reviewing the form of  $U_i$  from (111) and  $V'_i$  from (114), we see  $U_i = 1$  if  $\alpha_i + \beta_i = 0$  for  $2 \le i \le m$  and  $V'_i = 1$  if  $\gamma_i + \delta_i = 0$  for  $i \in \{1, 4, 5, \cdots, m\}$ . Consequently, if  $I_1 \cap I_2 = \emptyset$ , Then  $\alpha_2 = \beta_2 = \alpha_3 = \beta_3 = 0$  and  $\gamma_1 = \delta_1 = 0$ . This says that  $U_2 = U_3 = V'_1 = 1$ . Also, for each  $4 \le i \le m$ , the following have to be true:  $\alpha_i + \beta_i = 0$  if  $\gamma_i + \delta_i \ge 1$  and  $\gamma_i + \delta_i = 0$  if  $\alpha_i + \beta_i \ge 1$ . These imply  $\{U_i, V_i; 1 \le i \le m\}$  are independent, and hence  $\operatorname{Cov}(\prod_{i=1}^m U_i, \prod_{i=1}^m V_i') = 0$ . Thus, we only need to study the situation  $I_1 \cap I_2 \ne \emptyset$ . Let  $N_2$  be defined as in Lemma 6.19. Thus, the quantity from (113) is a linear combination of  $N_2$  terms of the form  $\operatorname{Cov}(\prod_{i=1}^m U_i, \prod_{i=1}^m V_i)$ . From Lemma 6.19,  $N_2 \le K_1 \cdot m^{\alpha+\beta+\gamma+\delta-1}$ . By Lemma 6.23 and applying the same argument of (108) to (114), we have

$$\max \left| \operatorname{Cov}(\prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i) \right| \le K_1 \sum_{1 \le i < j \le 4} r_{ij}^2$$

where the maximum is taken over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92). Combining all of these we get

$$m^{\alpha+\beta+\gamma+\delta} \cdot \left| \operatorname{Cov} \left( (X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{32}X_{42}) (X_{33}X_{43}) A_3^{\gamma} A_4^{\delta} \right) \right|$$
  
$$\leq K_1^2 (1+\alpha+\beta) \alpha! \beta! \gamma! \delta! \cdot m^{\alpha+\beta+\gamma+\delta-1} \cdot \sum_{1 \leq i < j \leq 4} r_{ij}^2.$$

This gives the bound for  $I_m(2,3)$ .

**LEMMA 6.26** Assume the setting in (96). Let  $A_i$  be defined as in (98). Given non-negative integers  $\alpha, \beta, \gamma, \delta$ , set

$$J_m(a,b) = Cov((X_{11}X_{21})(X_{12}X_{22})A_1^{\alpha}A_2^{\beta}, (X_{3a}X_{4a})(X_{3b}X_{4b})A_3^{\gamma}A_4^{\delta})$$
(115)

for integers  $a \ge 1$  and  $b \ge 1$ . Then

$$|J_m(1,2)| \le K \sum_{1 \le i < j \le 4} r_{ij}^2.$$

Proof of Lemma 6.26. Set

$$U_{1} = X_{11}^{2\alpha_{1}+1} X_{21}^{2\beta_{1}+1}, \quad U_{2} = X_{12}^{2\alpha_{2}+1} X_{22}^{2\beta_{2}+1} \text{ and } U_{i} = X_{1i}^{2\alpha_{i}} X_{2i}^{2\beta_{i}};$$
  

$$V_{1} = X_{31}^{2\gamma_{1}+1} X_{41}^{2\delta_{1}+1}, \quad V_{2} = X_{32}^{2\gamma_{2}+1} X_{42}^{2\delta_{2}+1} \text{ and } V_{i} = X_{3i}^{2\gamma_{i}} X_{4i}^{2\delta_{i}}$$
(116)

for  $3 \le i \le m$ . Let  $N_1$  be the total number of solutions  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92). From (106), we see  $N_1 \le K_1 \cdot m^{\alpha+\beta+\gamma+\delta}$ . Review the formulas between (100) and (103). We have

$$m^{\alpha+\beta+\gamma+\delta} \cdot \operatorname{Cov}\left((X_{11}X_{21})(X_{12}X_{22})A_1^{\alpha}A_2^{\beta}, (X_{31}X_{41})(X_{32}X_{42})A_3^{\gamma}A_4^{\delta}\right)$$
(117)

is a linear combination of  $N_1$  terms of the form  $\operatorname{Cov}(\prod_{i=1}^m U_i, \prod_{i=1}^m V_i)$  with positive coefficients no more than  $\alpha!\beta!\gamma!\delta!$ . Recall the notation  $J_m(a,b)$  and (117). We then have

$$m^{\alpha+\beta+\gamma+\delta} \cdot \left| J_m(1,2) \right| \le N_1 \cdot \left( \alpha!\beta!\gamma!\delta! \right) \cdot \max \left| \operatorname{Cov} \left( \prod_{i=1}^m U_i, \prod_{i=1}^m V_i \right) \right|,$$

where the maximum is taken over all  $\{\alpha_i, \beta_i, \gamma_i, \delta_i; 1 \le i \le m\}$  satisfying (92). By (106), we have

$$\left|J_m(1,2)\right| \le K_1 \cdot \max\left|\operatorname{Cov}\left(\prod_{i=1}^m U_i, \prod_{i=1}^m V_i\right)\right|,\tag{118}$$

where the maximum is taken over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92). By Lemma 6.23 and applying the same argument of (108) to (116), we have

$$\max |\text{Cov}(U_i, V_i)| \le K_1 \sum_{1 \le i < j \le 4} r_{ij}^2,$$
(119)
where the maximum is taken over all  $\{\alpha_i, \beta_i, \gamma_i, \delta_i; 1 \leq i \leq m\}$  satisfying (92). This and (118) conclude

$$|J_m(1,2)| \le K_1^2 \sum_{1 \le i < j \le 4} r_{ij}^2.$$

This proves the inequality for (a, b) = (1, 2).

**LEMMA 6.27** Assume the setting in (96). Give non-negative integers  $\alpha_i, \beta_i, \gamma_j, \delta_j$  for i = 1, 2 and j = 3, 4, set

$$U_{1} = X_{11}X_{21}, \quad U_{2} = X_{12}X_{22} \quad and \quad U_{i} = X_{1i}^{2\alpha_{i}}X_{2i}^{2\beta_{i}}, \quad i \in \{3, 4\};$$
  

$$V_{3} = X_{33}X_{43}, \quad V_{4} = X_{34}X_{44} \quad and \quad V_{i} = X_{3i}^{2\gamma_{i}}X_{4i}^{2\delta_{i}}, \quad i \in \{1, 2\}.$$
(120)

Define

$$S = \{i \in \{3,4\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \alpha_i + \beta_i \ge 1\} \cup \{i \in \{1,2\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) and } \gamma_i + \delta_i \ge 1\}.$$

Then the following hold.

(i) If  $S = \{1\}$  and  $\gamma_1 + \delta_1 = 1$ , then

$$Cov\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \begin{cases} 2(r_{12}r_{23}r_{31})r_{34}^{2}, & \text{if } \gamma_{1} = 1 \text{ and } \delta_{1} = 0; \\ 2(r_{12}r_{24}r_{41})r_{34}^{2}, & \text{if } \gamma_{1} = 0 \text{ and } \delta_{1} = 1. \end{cases}$$

(ii) If  $S = \{2\}$  and  $\gamma_2 + \delta_2 = 1$ , then

$$Cov\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \begin{cases} 2(r_{12}r_{23}r_{31})r_{34}^{2}, & \text{if } \gamma_{2} = 1 \text{ and } \delta_{2} = 0; \\ 2(r_{12}r_{24}r_{41})r_{34}^{2}, & \text{if } \gamma_{2} = 0 \text{ and } \delta_{2} = 1. \end{cases}$$

(iii) If  $S = \{3\}$  and  $\alpha_3 + \beta_3 = 1$ , then

$$Cov\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \begin{cases} 2(r_{13}r_{34}r_{41})r_{12}^{2}, & \text{if } \alpha_{3} = 1 \text{ and } \beta_{3} = 0; \\ 2(r_{23}r_{34}r_{42})r_{12}^{2}, & \text{if } \alpha_{3} = 0 \text{ and } \beta_{3} = 1. \end{cases}$$

(iv) If  $S = \{4\}$  and  $\alpha_4 + \beta_4 = 1$ , then

$$Cov\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \begin{cases} 2(r_{13}r_{34}r_{41})r_{12}^{2}, & \text{if } \alpha_{4} = 1 \text{ and } \beta_{4} = 0; \\ 2(r_{23}r_{34}r_{42})r_{12}^{2}, & \text{if } \alpha_{4} = 0 \text{ and } \beta_{4} = 1. \end{cases}$$

**Proof of Lemma 6.27**. By assumption,  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T \in \mathbb{R}^4; 1 \le j \le m\}$  are i.i.d. random vectors with distribution  $N_4(\mathbf{0}, \mathbf{R})$  where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each *i*. Thus,

$$EU_1 = EU_2 = r_{12}$$
 and  $EV_3 = EV_4 = r_{34}$ . (121)

(i) Under the case  $S = \{1\}$  and  $\gamma_1 + \delta_1 = 1$ , we know that  $\alpha_3 = \beta_3 = \alpha_4 = \beta_4 = \gamma_2 = \delta_2 = 0$  and that  $(\gamma_1, \delta_1)$  is equal to (1, 0) or (0, 1). Hence

$$U_{1} = X_{11}X_{21}, \ U_{2} = X_{12}X_{22}, \ U_{3} = 1, \ U_{4} = 1;$$
  
$$V_{1} = X_{31}^{2} \text{ or } X_{41}^{2}, \ V_{2} = 1, \ V_{3} = X_{33}X_{43}, \ V_{4} = X_{34}X_{44}.$$
 (122)

This implies that  $\{(U_1, V_1)^T, U_i, V_i; 2 \le i \le 4\}$  are independent. By (121) and by Lemma 6.4,

$$Cov(U_1, X_{31}^2) = E(X_{11}X_{21}X_{31}^2) - r_{12} = 2r_{13}r_{23};$$
$$Cov(U_1, X_{41}^2) = E(X_{11}X_{21}X_{41}^2) - r_{12} = 2r_{14}r_{24}.$$

Notice

$$\operatorname{Cov}(\xi_1\eta_1,\xi_2\eta_2) = E\xi_1 \cdot E\xi_2 \cdot \operatorname{Cov}(\eta_1,\eta_2)$$
(123)

if  $\xi_1$  and  $\xi_2$  are independent and  $\{\xi_1, \xi_2\}$  are independent of  $\{\eta_1, \eta_2\}$ . Note  $V_1 = X_{31}^2$  if  $(\gamma_1, \delta_1) = (1, 0)$  and  $V_1 = X_{41}^2$  if  $(\gamma_1, \delta_1) = (0, 1)$ . Then

$$\operatorname{Cov}\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \operatorname{Cov}(U_{1}, V_{1}) \cdot \prod_{i=2,3,4} (EU_{i} \cdot EV_{i})$$
$$= \begin{cases} 2(r_{12}r_{23}r_{31})r_{34}^{2}, & \text{if } \gamma_{1} = 1 \text{ and } \delta_{1} = 0; \\ 2(r_{12}r_{24}r_{41})r_{34}^{2}, & \text{if } \gamma_{1} = 0 \text{ and } \delta_{1} = 1. \end{cases}$$

(ii) Under the case  $S = \{2\}$  and  $\gamma_2 + \delta_2 = 1$ , we know that  $\alpha_3 = \beta_3 = \alpha_4 = \beta_4 = \gamma_1 = \delta_1 = 0$  and that  $(\gamma_2, \delta_2)$  is equal to (1, 0) or (0, 1). Hence

$$U_1 = X_{11}X_{21}, U_2 = X_{12}X_{22}, U_3 = 1, U_4 = 1;$$
  
 $V_1 = 1, V_2 = X_{32}^2 \text{ or } X_{42}^2, V_3 = X_{33}X_{43}, V_4 = X_{34}X_{44}$ 

Then,  $U_1$ ,  $(U_2, V_2)^T$ ,  $V_3$  and  $V_4$  are independent. By (121),

$$Cov(U_2, X_{32}^2) = E(X_{12}X_{22}X_{32}^2) - r_{12} = 2r_{13}r_{23};$$
  

$$Cov(U_2, X_{42}^2) = E(X_{12}X_{22}X_{42}^2) - r_{12} = 2r_{14}r_{24}.$$

By (123),

$$\operatorname{Cov}\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \operatorname{Cov}(U_{2}, V_{2}) \cdot \prod_{i=1,3,4}^{m} \left(EU_{i} \cdot EV_{i}\right)$$
$$= \begin{cases} 2(r_{12}r_{23}r_{31})r_{34}^{2}, & \text{if } \gamma_{2} = 1 \text{ and } \delta_{2} = 0; \\ 2(r_{12}r_{24}r_{41})r_{34}^{2}, & \text{if } \gamma_{2} = 0 \text{ and } \delta_{2} = 1. \end{cases}$$

(iii) Under the case  $S = \{3\}$  and  $\alpha_3 + \beta_3 = 1$ , we know that  $\alpha_4 = \beta_4 = \gamma_1 = \delta_1 = \gamma_2 = \delta_2 = 0$  and that  $(\alpha_3, \beta_3)$  is equal to (1, 0) or (0, 1). Hence

$$U_1 = X_{11}X_{21}, U_2 = X_{12}X_{22}, U_3 = X_{13}^2 \text{ or } X_{23}^2, U_4 = 1;$$
  
 $V_1 = 1, V_2 = 1, V_3 = X_{33}X_{43}, V_4 = X_{34}X_{44}.$ 

Then,  $U_1$ ,  $U_2$ ,  $(U_3, V_3)^T$  and  $V_4$  are independent. By (121),

$$Cov(X_{13}^2, V_3) = E(X_{33}X_{43}X_{13}^2) - r_{34} = 2r_{13}r_{14};$$
$$Cov(X_{23}^2, V_3) = E(X_{33}X_{43}X_{23}^2) - r_{34} = 2r_{23}r_{24}.$$

By (123),

$$\operatorname{Cov}\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \operatorname{Cov}(U_{3}, V_{3}) \cdot \prod_{i=1,2,4}^{m} \left(EU_{i} \cdot EV_{i}\right)$$
$$= \begin{cases} 2(r_{13}r_{34}r_{41})r_{12}^{2}, & \text{if } \alpha_{3} = 1 \text{ and } \beta_{3} = 0; \\ 2(r_{23}r_{34}r_{42})r_{12}^{2}, & \text{if } \alpha_{3} = 0 \text{ and } \beta_{3} = 1. \end{cases}$$

(iv) Under the case  $S = \{4\}$  and  $\alpha_4 + \beta_4 = 1$ , we know that  $\alpha_3 = \beta_3 = \gamma_1 = \delta_1 = \gamma_2 = \delta_2 = 0$  and that  $(\alpha_4, \beta_4)$  is equal to (1, 0) or (0, 1). Hence

$$U_1 = X_{11}X_{21}, U_2 = X_{12}X_{22}, U_3 = 1, U_4 = X_{14}^2 \text{ or } X_{24}^2;$$
  
 $V_1 = 1, V_2 = 1, V_3 = X_{33}X_{43}, V_4 = X_{34}X_{44}.$ 

Then,  $U_1$ ,  $U_2$ ,  $V_3$ ,  $(U_4, V_4)^T$  are independent. By (121),

$$Cov(X_{14}^2, V_4) = E(X_{34}X_{44}X_{14}^2) - r_{34} = 2r_{13}r_{14};$$
$$Cov(X_{24}^2, V_4) = E(X_{34}X_{44}X_{24}^2) - r_{34} = 2r_{23}r_{24}.$$

By (123),

$$\operatorname{Cov}\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right) = \operatorname{Cov}(U_{4}, V_{4}) \cdot \prod_{i=1,2,3}^{m} \left(EU_{i} \cdot EV_{i}\right)$$
$$= \begin{cases} 2(r_{13}r_{34}r_{41})r_{12}^{2}, & \text{if } \alpha_{4} = 1 \text{ and } \beta_{4} = 0; \\ 2(r_{23}r_{34}r_{42})r_{12}^{2}, & \text{if } \alpha_{4} = 0 \text{ and } \beta_{4} = 1. \end{cases}$$

The verification is finished.

Let  $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_m, \gamma_1, \dots, \gamma_m, \delta_1, \dots, \delta_m$  be non-negative integers, review the notation  $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_m), \boldsymbol{\beta} = (\beta_1, \dots, \beta_m), \boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m)$  and  $\boldsymbol{\delta} = (\delta_1, \dots, \delta_m)$ . For non-negative integers  $\alpha, \beta, \gamma, \delta$  and  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92), define

$$C(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta}) = \frac{\alpha!}{\alpha_1!\cdots\alpha_m!} \cdot \frac{\beta!}{\beta_1!\cdots\beta_m!} \cdot \frac{\gamma!}{\gamma_1!\cdots\gamma_m!} \cdot \frac{\delta!}{\delta_1!\cdots\delta_m!}.$$
 (124)

**LEMMA 6.28** Assume the setting in (96). Define  $\{U_i, V_i; 1 \le i \le m\}$  such that

$$U_{1} = X_{11}^{2\alpha_{1}+1}X_{21}^{2\beta_{1}+1}, \quad U_{2} = X_{12}^{2\alpha_{2}+1}X_{22}^{2\beta_{2}+1} \text{ and } U_{i} = X_{1i}^{2\alpha_{i}}X_{2i}^{2\beta_{i}};$$
  
$$V_{3} = X_{33}^{2\gamma_{3}+1}X_{43}^{2\delta_{3}+1}, \quad V_{4} = X_{34}^{2\gamma_{4}+1}X_{44}^{2\delta_{4}+1} \text{ and } V_{j} = X_{3j}^{2\gamma_{j}}X_{4j}^{2\delta_{j}};$$

for  $3 \leq i \leq m$  and  $j \in \{1, 2, \dots, m\} \setminus \{3, 4\}$ , where  $\alpha_i, \beta_i, \gamma_i, \delta_i$  satisfies (92). Define

$$S = \{i \in \{3, 4\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) with } \alpha_i + \beta_i \ge 1\} \cup \{j \in \{1, 2\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) with } \gamma_j + \delta_j \ge 1\}.$$

_		
г		1
		L

Obviously,  $S \subset \{1, 2, 3, 4\}$ . Then

$$\sum_{\substack{(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta}):|S|=1\\1\leq i< j\leq 4}} C(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta}) \cdot Cov\left(\prod_{i=1}^{m} U_{i},\prod_{i=1}^{m} V_{i}\right)$$
$$= \rho_{m,1} \sum_{1\leq i< j\leq 4} r_{ij}^{2} + \rho_{m,2} \cdot \left[(r_{12}r_{23}r_{31})r_{34}^{2} + (r_{12}r_{24}r_{41})r_{34}^{2}\right]$$
$$+ \rho_{m,3} \cdot \left[(r_{13}r_{34}r_{41})r_{12}^{2} + (r_{23}r_{34}r_{42})r_{12}^{2}\right]$$

where  $\max\{m^2|\rho_{m,1}|, m|\rho_{m,2}|, m|\rho_{m,3}|\} \leq K$  and  $\rho_{m,2}$  and  $\rho_{m,3}$  do not depend on **R**.

**Proof of Lemma 6.28**. First, |S| = 1 implies that  $S = \{1\}$ ,  $S = \{2\}$ ,  $S = \{3\}$  or  $S = \{4\}$ . We will first examine the case  $S = \{1\}$  next.

Assume now  $S = \{1\}$ . Then  $\gamma_1 + \delta_1 \ge 1$  and  $\alpha_3 = \beta_3 = \alpha_4 = \beta_4 = \gamma_2 = \delta_2 = 0$ . Hence

 $U_3 = U_4 = V_2 = 1$  and  $U_2, V_3, V_4$  are independent themselves and they are

also independent of  $\{(U_i, V_i)^T; i = 1, 5, 6, \cdots, m\}$  (125)

by the fact  $\{(U_i, V_i)^T; 1 \leq i \leq m\}$  are independent aforementioned. By Lemma 6.23 and applying the same argument of (108) to  $\{U_i, V_i, 1 \leq i \leq m\}$ , we obtain

$$\max |\operatorname{Cov}(U_i, V_i)| \le K_1 \sum_{1 \le i < j \le 4} r_{ij}^2,$$
(126)

where the maximum is taken over all  $\{\alpha_i, \beta_i, \gamma_i, \delta_i; 1 \leq i \leq m\}$  satisfying (92). Next we bound

$$\sum \left| \operatorname{Cov} \left( \prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i \right) \right|, \tag{127}$$

where the sum runs over  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfing (92) and  $S = \{1\}$ . When  $S = \{1\}$ , we know  $\gamma_1 + \delta_1 \geq 1$ . We will distinguish two cases:  $\gamma_1 + \delta_1 \geq 2$  and  $\gamma_1 + \delta_1 = 1$ . Recall the definition of  $T_{m,3}$  from Lemma 6.21. For the case  $\gamma_1 + \delta_1 = 1$ , we will divide it into another two case:  $T_{m,3}$  and  $T_{m,3}^c$ . The derivation of bounds for (127) under  $\gamma_1 + \delta_1 \geq 2$  and  $T_{m,3}^c$  are easier than that under  $T_{m,3}$ . We will take two steps next two handle the two cases.

Step 1. By Lemma 6.20(ii), the total number of solutions  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  of (92) with  $\gamma_1 + \delta_1 \geq 2$  is bounded by  $K_1 \cdot m^{\alpha+\beta+\gamma+\delta-2}$ . This joined with (126) implies that

$$\sum \left| \operatorname{Cov} \left( \prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i \right) \right| \le K_1 \cdot m^{\alpha + \beta + \gamma + \delta - 2} \sum_{1 \le i < j \le 4} r_{ij}^2, \tag{128}$$

where the sum runs over all  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92),  $S = \{1\}$  and  $\gamma_1 + \delta_1 \geq 2$ .

Review the definition of  $T_{m,3}$  in Lemma 6.21. We have  $|T_{m,3}| \leq K_1 \cdot m^{\alpha+\beta+\gamma+\delta-2}$ . This together with (126) yields

$$\sum \left| \operatorname{Cov} \left( \prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i \right) \right| \le K_1 \cdot m^{\alpha + \beta + \gamma + \delta - 2} \sum_{1 \le i < j \le 4} r_{ij}^2, \tag{129}$$

where the sum runs over all  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92),  $S = \{1\}$  and  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \in T_{m,3}$ .

Step 2. We now estimate (127) as the index  $(\alpha, \beta, \gamma, \delta)$  satisfies (92), the event  $S = \{1\}$  holds and  $(\alpha, \beta, \gamma, \delta) \notin T_{m,3}$ . Review the definition of  $T_{m,3}$  and expressions of  $U_i$  and  $V_i$ , under the new conditions,  $U_i$  and  $V_i$  take much simpler form:

$$U_{1} = X_{11}X_{21}, U_{2} = X_{12}X_{22}, U_{3} = 1, U_{4} = 1;$$
  

$$V_{1} = X_{31}^{2} \text{ or } X_{41}^{2}, V_{2} = 1, V_{3} = X_{33}X_{43}, V_{4} = X_{34}X_{44}.$$
(130)

Furthermore, if  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \notin T_{m,3}$ , then  $\alpha_k + \beta_k \leq 1$  for all  $5 \leq k \leq m$ ,  $\gamma_l + \delta_l \leq 1$  for all  $5 \leq l \leq m$  and the two identities  $\alpha_t + \beta_t = 1$  and  $\gamma_t + \delta_t = 1$  cannot occur at the same time for any  $5 \leq t \leq m$ . The key observation is that, if  $\alpha_t + \beta_t = 1$  then  $U_t \sim \chi^2(1)$  and  $V_t = 1$ . Similarly, if  $\gamma_t + \delta_t = 1$  then  $U_t = 1$  and  $V_t \sim \chi^2(1)$ . Therefore, the 2m - 8 random variables in  $\{U_i, V_i; 5 \leq i \leq m\}$  are independent random variables, each has mean 1. As used earlier,  $\{(U_i, V_i)^T; 1 \leq i \leq m\}$  are independent. This and the special structures in (130) imply that the 2m - 1 random quantities in  $\{(U_1, V_1)^T, U_i, V_i; 2 \leq i \leq m\}$  are independent. By Lemma 6.27(i) and (123),

$$\operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = \operatorname{Cov}\left(\prod_{i=1}^{4} U_{i}, \prod_{i=1}^{4} V_{i}\right)$$
$$= \begin{cases} 2(r_{12}r_{23}r_{31})r_{34}^{2}, & \text{if } \gamma_{1} = 1 \text{ and } \delta_{1} = 0; \\ 2(r_{12}r_{24}r_{41})r_{34}^{2}, & \text{if } \gamma_{1} = 0 \text{ and } \delta_{1} = 1. \end{cases}$$

This says that

$$\sum \operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = L_{m,1} \cdot \left[(r_{12}r_{23}r_{31})r_{34}^{2} + (r_{12}r_{24}r_{41})r_{34}^{2}\right],\tag{131}$$

where the sum runs over  $\Gamma$ , defined to be the set of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92), |S| = 1 with  $\gamma_1 + \delta_1 = 1$ , and  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \notin T_{m,3}$ ;  $L_{m,1} := 2|\Gamma|$ . Obviously,  $L_{m,1}$  does not depend on the matrix  $\mathbf{R} = (r_{ij})$ . By the bound on  $T_{m,1}$  in Lemma 6.21, we have  $L_{m,1} \leq K_1 m^{\alpha+\beta+\gamma+\delta-1}$ . Notice, if  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfies (92), the event  $S = \{1\}$  holds and  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \notin T_{m,3}$ , then any one from  $\{\alpha_i, \beta_i, \gamma_i, \delta_i; 1 \leq i \leq m\}$  is either 1 or 0. According to (124),  $C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) = \alpha!\beta!\gamma!\delta!$ . Then (131) becomes

$$\sum C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \operatorname{Cov} \Big( \prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i \Big) = \alpha! \beta! \gamma! \delta! \cdot L_{m,1} \cdot \big[ (r_{12}r_{23}r_{31})r_{34}^2 + (r_{12}r_{24}r_{41})r_{34}^2 \big].$$

Thus, combining this with (128) and (129) and using the trivial fact that  $C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \leq \alpha! \beta! \gamma! \delta!$ , we arrive at

$$\sum C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \cdot \operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = \tau_{m,1} \cdot \left[(r_{12}r_{23}r_{31})r_{34}^{2} + (r_{12}r_{24}r_{41})r_{34}^{2}\right] + \tau_{m,1}^{\prime}, (132)$$

where the sum runs over the set of  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and  $S = \{1\}$ , and where  $\tau_{m,1}$ does not depend on  $r_{ij}$  and

$$|\tau_{m,1}| \le K_1 m^{\alpha+\beta+\gamma+\delta-1} \text{ and } |\tau'_{m,1}| \le K_1 \cdot m^{\alpha+\beta+\gamma+\delta-2} \sum_{1\le i< j\le 4} r_{ij}^2.$$
 (133)

By applying the same argument as the derivation of (132) to  $S = \{2\}$  and using Lemma 6.27(ii), we get an analogue of (132) as the sum runs over the set of  $(\alpha, \beta, \gamma, \delta)$ satisfying (92) and  $S = \{2\}$ , where  $\tau_{m,1}$  and  $\tau'_{m1}$  will be replaced by two corresponding symbols but still satisfy (133).

By applying the same argument as the derivation of (132) to  $S = \{3\}$  and using Lemma 6.27(iii), we get

$$\sum \operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = \tilde{\tau}_{m,1} \cdot \left[(r_{13}r_{34}r_{41} + r_{23}r_{34}r_{42})r_{12}^{2}\right] + \tilde{\tau}_{m,1}^{\prime},$$
(134)

where the sum runs over the set of  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) and  $S = \{3\}$ , and the inequalities from (133) still hold as " $(\tau_{m,1}, \tau'_{m,1})$ " is replaced by " $(\tilde{\tau}_{m,1}, \tilde{\tau}'_{m,1})$ ".

By applying the same argument as the derivation of (132) to  $S = \{4\}$  and using Lemma 6.27(iv), we get an analogue of (134) as the sum runs over the set of  $(\alpha, \beta, \gamma, \delta)$ satisfying (92) and  $S = \{4\}$ , the quantities " $(\tilde{\tau}_{m,1}, \tilde{\tau}'_{m,1})$ " is replaced by " $(T_{m,1}, T''_{m1})$ ", and  $T_{m,1}$  does not depend on  $r_{ij}$  and

$$|T_{m,1}| \le K_1 m^{\alpha+\beta+\gamma+\delta-1}$$
 and  $|T'_{m,1}| \le K_1 \cdot m^{\alpha+\beta+\gamma+\delta-2} \sum_{1 \le i < j \le 4} r_{ij}^2$ .

The proof is completed by summing the above four upper bounds corresponding to  $S = \{1\}, \{2\}, \{3\}$  and  $\{4\}$ .

**LEMMA 6.29** Assume the setting in (96). Let  $J_m(a, b)$  be defined as in Lemma 6.26. Then

$$J_m(3,4) = \tau_{m,1}r_{12}^2r_{34}^2 + \tau_{m,2}\sum_{1 \le i < j \le 4} r_{ij}^2 + \tau_{m,3} \cdot \left[ (r_{12}r_{23}r_{31})r_{34}^2 + (r_{12}r_{24}r_{41})r_{34}^2 \right] \\ + \tau_{m,4} \cdot \left[ (r_{13}r_{34}r_{41})r_{12}^2 + (r_{23}r_{34}r_{42})r_{12}^2 \right],$$

where  $\max\{m|\tau_{m,1}|, m^2|\tau_{m,2}|, m|\tau_{m,3}|, m|\tau_{m,4}|\} \leq K$  and  $\tau_{m,3}$  and  $\tau_{m,4}$  do not depend on **R**.

Proof of Lemma 6.29. Set

$$U_{1} = X_{11}^{2\alpha_{1}+1} X_{21}^{2\beta_{1}+1}, \quad U_{2} = X_{12}^{2\alpha_{2}+1} X_{22}^{2\beta_{2}+1} \text{ and } U_{i} = X_{1i}^{2\alpha_{i}} X_{2i}^{2\beta_{i}};$$
  

$$V_{3} = X_{33}^{2\gamma_{3}+1} X_{43}^{2\delta_{3}+1}, \quad V_{4} = X_{34}^{2\gamma_{4}+1} X_{44}^{2\delta_{4}+1} \text{ and } V_{j} = X_{3j}^{2\gamma_{j}} X_{4j}^{2\delta_{j}}$$
(135)

for  $3 \leq i \leq m$  and  $j \in \{1, 2, \dots, m\} \setminus \{3, 4\}$ . Let  $N_1$  be the total number of solutions  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  for (92) with a bound provided in (106). Review the discussions between (100) and (105). We know that  $m^{\alpha+\beta+\gamma+\delta}J_m(3,4)$  is a linear combination of  $N_1$  terms of the form  $\operatorname{Cov}(\prod_{i=1}^m U_i, \prod_{i=1}^m V_i)$  with positive coefficients no more than  $\alpha!\beta!\gamma!\delta!$ . Define

$$S = \{i \in \{3, 4\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) with } \alpha_i + \beta_i \ge 1\} \cup \{j \in \{1, 2\}; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) with } \gamma_j + \delta_j \ge 1\};$$
  

$$S_1 = \{5 \le i \le m; (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \text{ satisfies (92) with either } \alpha_i + \beta_i \ge 1 \text{ or } \delta_i + \gamma_i \ge 1\}.$$

Similar to the proof of Lemma 6.23, we have

$$|S_1| \leq \alpha + \beta + \gamma + \delta. \tag{136}$$

We now estimate  $\operatorname{Cov}(\prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i)$  by differentiating three cases: |S| = 0, |S| = 1 and  $|S| \ge 2$ . Quickly, for the case |S| = 1, by reviewing  $C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  from (124), we have from Lemma 6.28 that

$$\sum_{\substack{(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta}):|S|=1\\1\leq i< j\leq 4}} C(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta}) \cdot \operatorname{Cov}\left(\prod_{i=1}^{m} U_{i},\prod_{i=1}^{m} V_{i}\right)$$

$$= \rho_{m,1} \sum_{1\leq i< j\leq 4} r_{ij}^{2} + \rho_{m,2} \cdot \left[(r_{12}r_{23}r_{31})r_{34}^{2} + (r_{12}r_{24}r_{41})r_{34}^{2}\right]$$

$$+ \rho_{m,3} \cdot \left[(r_{13}r_{34}r_{41})r_{12}^{2} + (r_{23}r_{34}r_{42})r_{12}^{2}\right], \qquad (137)$$

where  $\rho_{m,1} \leq Km^{-2}$  and  $\rho_{m,2} \vee \rho_{m,3} \leq m^{-1}K$ , and where  $\rho_{m,2}$  and  $\rho_{m,3}$  do not depend on **R**. To finish the proof, it remains to study the cases "|S| = 0" and " $|S| \geq 2$ ". This will be worked out in two steps.

**Step 1**: First, the condition |S| = 0 implies  $\alpha_3 = \alpha_4 = \beta_3 = \beta_4 = \gamma_1 = \gamma_2 = \delta_1 = \delta_2 = 0$ , and hence we have from (135) that

$$U_1 = X_{11}^{2\alpha_1+1} X_{21}^{2\beta_1+1}, \quad U_2 = X_{12}^{2\alpha_2+1} X_{22}^{2\beta_2+1}, \quad U_3 = 1, \\ U_4 = 1 \text{ and } U_i = X_{1i}^{2\alpha_i} X_{2i}^{2\beta_i};$$
  
$$V_1 = 1, \\ V_2 = 1, \\ V_3 = X_{33}^{2\gamma_3+1} X_{43}^{2\delta_3+1}, \quad V_4 = X_{34}^{2\gamma_4+1} X_{44}^{2\delta_4+1} \text{ and } V_i = X_{3i}^{2\gamma_i} X_{4i}^{2\delta_i}$$

for  $i = 5, \dots, m$ . By assumption (96),  $\{(X_{1i}, X_{2i}, X_{3i}, X_{4i})^T \in \mathbb{R}^4; 1 \leq i \leq m\}$  are i.i.d. random vectors with distribution  $N_4(\mathbf{0}, \mathbf{R})$ , where  $\mathbf{R} = (r_{ij})_{4\times 4}$  and  $r_{ii} = 1$  for each i. In particular,  $\{(U_i, V_i)^T; 1 \leq i \leq m\}$  are independent. As a consequence,  $U_1, U_2, V_3, V_4$  are themselves independent, and furthermore  $\{U_1, U_2, V_3, V_4\}$  are also independent of  $\{U_i, V_i; 5 \leq i \leq m\}$ . By (123),

$$\operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = EU_{1}EU_{2}EV_{3}EV_{4} \cdot \operatorname{Cov}\left(\prod_{i=5}^{m} U_{i}, \prod_{i=5}^{m} V_{i}\right)$$
$$= EU_{1}EU_{2}EV_{3}EV_{4} \cdot \operatorname{Cov}\left(\prod_{i\in S_{1}} U_{i}, \prod_{i\in S_{1}} V_{i}\right).$$
(138)

By definition of  $S_1$ , we see that  $\sum_{i \in S_1} \alpha_i = \alpha$ ,  $\sum_{i \in S_1} \beta_i = \beta$ ,  $\sum_{i \in S_1} \gamma_i = \gamma$  and  $\sum_{i \in S_1} \delta_i = \delta$ . By Lemma 6.23,

$$\left|\operatorname{Cov}\left(\prod_{i \in S_1} U_i, \prod_{i \in S_1} V_i\right)\right| \le K_1 \sum_{1 \le i < j \le 4} r_{ij}^2 \le 6K_1.$$

Bounds for  $EU_1, EU_2, EV_3, EV_4$  are given in Lemma 6.13. By the lemma, we see

$$\left| \operatorname{Cov} \left( \prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i} \right) \right| \leq K_{1} r_{12}^{2} r_{34}^{2} \cdot \left| \operatorname{Cov} \left( \prod_{i=5}^{m} U_{i}, \prod_{i=5}^{m} V_{i} \right) \right| \\ \leq K_{1} \cdot r_{12}^{2} r_{34}^{2}.$$
(139)

Let  $S_2$  be the set of solutions  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92) with  $\alpha_i + \beta_i \geq 1$  and  $\gamma_i + \delta_i \geq 1$ simultaneously for some  $5 \leq i \leq m$ . By Lemma 6.22, we have  $|S_2| \leq Km^{\alpha+\beta+\gamma+\delta-1}$ . Recall  $U_i = X_{1i}^{2\alpha_i}X_{2i}^{2\beta_i}$  and  $V_i = X_{3i}^{2\gamma_i}X_{4i}^{2\delta_i}$  for  $5 \leq i \leq m$ . If  $\alpha_i = \beta_i = 0$  then  $U_i = 1$ . Likewise,  $V_i = 1$  if  $\gamma_i = \delta_i = 0$ . This together with the fact  $\{(U_i, V_i); 1 \leq i \leq m\}$  are independent implies  $\prod_{i=5}^m U_i$  and  $\prod_{i=5}^m V_i$  are independent (hence their covariance is zero) if there is no  $i \in \{5, \dots, m\}$  such that  $\alpha_i + \beta_i \geq 1$  and  $\gamma_i + \delta_i \geq 1$  at the same time. Therefore, by (139),

$$\sum \left| \operatorname{Cov} \left( \prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i \right) \right| \le K_1 \cdot m^{\alpha + \beta + \gamma + \delta - 1} r_{12}^2 r_{34}^2 \tag{140}$$

where the sum runs over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and |S| = 0.

Step 2: Assume the index  $(\alpha, \beta, \gamma, \delta)$  satisfies that  $|S| \ge 2$ . Review the structures of  $U_i$  and  $V_i$  from (135), we have from Lemma 6.23 that

$$\max \left| \operatorname{Cov} \left( \prod_{i=1}^m U_i, \prod_{i=1}^m V_i \right) \right| \le K_1 \cdot \sum_{1 \le i < j \le 4} r_{12}^2,$$

where the sum runs over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92). Review the definition of  $T_{m,2}$  from Lemma (6.21), we have from the lemma that  $|T_{m,2}| \leq K_{13}m^{\alpha+\beta+\gamma+\delta-2}$ . It follows that

$$\sum \left| \operatorname{Cov} \left( \prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i \right) \right| \le K_1 m^{\alpha + \beta + \gamma + \delta - 2} \sum_{1 \le i < j \le 4} r_{12}^2, \tag{141}$$

where the sum runs over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and  $|S| \ge 2$ .

Finally, we add up the three bounds from (137), (140) and (141). By changing " $K_1$ " to " $\tau_{m,1}$ ", " $\rho_{m,1} + K_1$ " to " $\tau_{m,2}$ ", " $\rho_{m,2}$ " to " $\tau_{m,3}$ " and " $\rho_{m,3}$ " to " $\tau_{m,4}$ ", we complete the proof.

**LEMMA 6.30** Assume the setting in (96). Let all notation be the same as those in Lemma 6.26. Then

$$J_m(2,3) = \tau_{m,1} \sum_{1 \le i < j \le 4} r_{ij}^2 + \tau_{m,2} \cdot (r_{12}r_{23}r_{34}r_{41} + r_{12}r_{24}r_{43}r_{31}),$$

where  $\max\{m|\tau_{m,1}|, |\tau_{m,2}|\} \leq K$  and  $\tau_{m,2}$  does not depend on **R**.

## Proof of Lemma 6.30. Set

$$U_{1} = X_{11}^{2\alpha_{1}+1}X_{21}^{2\beta_{1}+1}, \quad U_{2} = X_{12}^{2\alpha_{2}+1}X_{22}^{2\beta_{2}+1} \text{ and } U_{i} = X_{1i}^{2\alpha_{i}}X_{2i}^{2\beta_{i}};$$
  

$$V_{2} = X_{32}^{2\gamma_{2}+1}X_{42}^{2\delta_{2}+1}, \quad V_{3} = X_{33}^{2\gamma_{3}+1}X_{43}^{2\delta_{3}+1} \text{ and } V_{j} = X_{3j}^{2\gamma_{j}}X_{4j}^{2\delta_{j}}$$
(142)

for  $3 \le i \le m$  and  $j \in \{1, 2, \dots, m\} \setminus \{2, 3\}$ . Review  $C(\alpha, \beta, \gamma, \delta)$  in (124). By (100)-(103) and (115),

$$J_{m}(2,3) = \operatorname{Cov}((X_{11}X_{21})(X_{12}X_{22})A_{1}^{\alpha}A_{2}^{\beta}, (X_{32}X_{42})(X_{33}X_{43})A_{3}^{\gamma}A_{4}^{\delta}) = \frac{1}{m^{\alpha+\beta+\gamma+\delta}}\sum C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \cdot \operatorname{Cov}(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}),$$
(143)

where the sum runs over all  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92). Let S be defined as in Lemma 6.22. By the lemma,  $|S| \leq K_1 m^{\alpha+\beta+\gamma+\delta-1}$ . By Lemma 6.23 and structures of  $U_i$  and  $V_i$  in (142),

$$\max \left| \operatorname{Cov} \left( \prod_{i=1}^{m} U_i, \prod_{i=1}^{m} V_i \right) \right| \le K_1 \cdot \sum_{1 \le i < j \le 4} r_{12}^2,$$

where the maximum is taken over all  $(\alpha, \beta, \gamma, \delta)$  satisfying (92). Combining the two facts, we obtain that

$$\sum_{(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta})\in S} \left| \operatorname{Cov}\left(\prod_{i=1}^{m} U_{i},\prod_{i=1}^{m} V_{i}\right) \right| \leq K_{1} m^{\alpha+\beta+\gamma+\delta-1} \sum_{1\leq i< j\leq 4} r_{12}^{2}.$$

As used before,  $1 \leq C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) \leq \alpha! \beta! \gamma!$  for any  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfies (92). We rewrite the above as

$$\frac{1}{m^{\alpha+\beta+\gamma+\delta}} \sum_{(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta})\in S} C(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta}) \cdot \operatorname{Cov}\left(\prod_{i=1}^{m} U_{i},\prod_{i=1}^{m} V_{i}\right) = \tau_{m,1} \sum_{1 \le i < j \le 4} r_{12}^{2}, \quad (144)$$

where  $|\tau_{m,1}| \le K_1 m^{-1}$ .

Now, if  $(\alpha, \beta, \gamma, \delta)$  satisfies (92) but not in S, then (i)  $\alpha_i + \beta_i = 0$  for each  $i \in \{1, 2, 3\}$ ; (ii)  $\gamma_i + \delta_i = 0$  for each  $i \in \{1, 2, 3\}$ ; (iii)  $\alpha_i + \beta_i \leq 1$  and  $\gamma_i + \delta_i \leq 1$  for each  $4 \leq i \leq m$ ; (iv) for each  $4 \leq i \leq m$ , if  $\alpha_i + \beta_i = 1$  then  $\gamma_i + \delta_i = 0$ , and if  $\gamma_i + \delta_i = 1$  then  $\alpha_i + \beta_i = 0$ . This implies that  $\{U_i, V_i; 4 \leq i \leq m\}$  are independent random variables and each of them is either 1 or  $\chi^2(1)$ . Keep in mind that  $\{(U_i, V_i); 1 \leq i \leq m\}$  are independent random variables and  $E(\chi^2(1)) = 1$ . Furthermore, it is readily seen from (142) that

$$U_1 = X_{11}X_{21}, U_2 = X_{12}X_{22} \text{ and } U_3 = 1;$$
  
 $V_1 = 1, V_2 = X_{32}X_{42}, V_3 = X_{33}X_{43}.$ 

Since  $\{(U_i, V_i)^T; 1 \le i \le m\}$  are independent by the assumption from (96), the three random quantities  $\{U_1, (U_2, V_2)^T, V_3\}$  are independent and they are independent of  $\{(U_i, V_i)^T; 4 \le i \le m\}$ . Thus, it follows from (123) that

$$\operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = E\left(\prod_{i=4}^{m} U_{i}\right) \cdot E\left(\prod_{i=4}^{m} V_{i}\right) \cdot \operatorname{Cov}\left(\prod_{i=1}^{3} U_{i}, \prod_{i=1}^{3} V_{i}\right)$$
$$= EU_{1} \cdot EV_{3} \cdot \operatorname{Cov}(U_{2}, V_{2}).$$

Recall  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T \in \mathbb{R}^4; 1 \leq j \leq m\}$  are i.i.d. random vectors with distribution  $N_4(\mathbf{0}, \mathbf{R})$ , where  $\mathbf{R} = (r_{ij})_{4 \times 4}$  and  $r_{ii} = 1$  for each *i*. Then  $EU_1 = r_{12}$ ,  $EV_3 = r_{34}$  and

$$\operatorname{Cov}(U_2, V_2) = E(X_{12}X_{22}X_{32}X_{42}) - r_{12}r_{34} = r_{13}r_{24} + r_{14}r_{23}$$

by Lemma 6.4. Thus,

$$\operatorname{Cov}\left(\prod_{i=1}^{m} U_{i}, \prod_{i=1}^{m} V_{i}\right) = r_{12}r_{24}r_{43}r_{31} + r_{12}r_{23}r_{34}r_{41}.$$

Recall (124),  $C(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta}) = \alpha! \beta! \gamma! \delta!$  in this case, that is,  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfies (92) but not in *S*. Let  $N_1$  be the total number of solutions  $(\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\delta})$  satisfying (92). From (106), we see  $N_1 \leq K_1 \cdot m^{\alpha+\beta+\gamma+\delta}$ . Therefore, there exists a constant  $\tau_{m,2}$  not depending  $r_{ij}$  and  $|\tau_{m,2}| \leq K_1$  such that

$$\frac{1}{m^{\alpha+\beta+\gamma+\delta}}\sum C(\boldsymbol{\alpha},\boldsymbol{\beta},\boldsymbol{\gamma},\boldsymbol{\delta})\cdot\operatorname{Cov}\Big(\prod_{i=1}^{m}U_{i},\prod_{i=1}^{m}V_{i}\Big)=\tau_{m,2}\big(r_{12}r_{23}r_{34}r_{41}+r_{12}r_{24}r_{43}r_{31}\big)$$

where the sum is taken over every  $(\alpha, \beta, \gamma, \delta)$  satisfying (92) and the restriction in  $S^c$ . By connecting this fact to (143) and (144), we get desired the conclusion.

## 6.2.5 A Study on Correlation Matrices

As needed in the proof of Lemma 6.34 later, we have to handle certain functions of the entries of sample correlation matrices. They are interesting on their own. Through the whole section, we assume  $\mathbf{R} = (r_{ij})_{p \times p}$  is a non-negative definite matrix with  $r_{ii} = 1$  for  $1 \leq i \leq p$ . Review the Frobenius norm  $\|\mathbf{R}\|_F = [\operatorname{tr}(\mathbf{R}^2)]^{1/2} = (\sum_{1 \leq i,j \leq p} r_{ij}^2)^{1/2}$ .

**LEMMA 6.31** Assume  $\{m_p; p \ge 1\}$  are positive constants with  $\lim_{p\to\infty} m_p = \infty$ . Define

$$W_1 = \sum_{1 \le i, j, k \le p} r_{ij} r_{jk} r_{ki} \quad and \quad W_2 = \sum_{1 \le i, j, k, l \le p} r_{ij} r_{jk} r_{kl} r_{li}.$$
  
Then  $\lim_{p \to \infty} \frac{W_i}{m \|\mathbf{R}\|_F^4} = 0$  for  $i = 1, 2$ .

**Proof of Lemma 6.31**. Let  $\lambda_1 \geq 0, \dots, \lambda_p \geq 0$  be the eigenvalues of **R**. Write  $\mathbf{R} = \mathbf{O}^T \operatorname{diag}(\lambda_1, \dots, \lambda_p) \mathbf{O}$ , where **O** is a  $p \times p$  orthogonal matrix. Recall the fact

$$\operatorname{tr}(\mathbf{R}^{s}) = \sum_{1 \le i_{1}, i_{2}, \cdots, i_{s} \le p} r_{i_{1}i_{2}} r_{i_{2}i_{3}} r_{i_{3}i_{4}} \cdots r_{i_{s}i_{1}}$$
(145)

for any integer  $s \ge 2$ . Easily,

$$W_1 = \operatorname{tr}(\mathbf{R}^3) = \lambda_1^3 + \dots + \lambda_p^3 \le \left(\lambda_1^2 + \dots + \lambda_p^2\right)^{3/2}$$

In addition,  $\|\mathbf{R}\|_F^2 \ge \sum_{i=1}^p r_{ii}^2 = p$ . Therefore,

$$\frac{|W_1|}{[\operatorname{tr}(\mathbf{R}^2)]^2} \le \frac{[\operatorname{tr}(\mathbf{R}^2)]^{3/2}}{[\operatorname{tr}(\mathbf{R}^2)]^2} = \frac{1}{[\operatorname{tr}(\mathbf{R}^2)]^{1/2}} \le \frac{1}{\sqrt{p}}$$

It follows that  $\lim_{p\to\infty} \frac{W_1}{m \|\mathbf{R}\|_F^4} = 0$ . Now prove the second conclusion. Note

$$W_2 = \operatorname{tr}(\mathbf{R}^4) = \lambda_1^4 + \dots + \lambda_p^4 \le (\lambda_1^2 + \dots + \lambda_p^2)^2.$$

Consequently,

$$\frac{|W_2|}{m[\operatorname{tr}(\mathbf{R}^2)]^2} \le \frac{[\operatorname{tr}(\mathbf{R}^2)]^2}{m[\operatorname{tr}(\mathbf{R}^2)]^2} = \frac{1}{m} \to 0$$

85

as  $p \to \infty$ . The proof is completed.

**LEMMA 6.32** Given integer  $\alpha \geq 0$ , define

$$S_m = \sum_{1 \le i,j,k,l \le p} r_{ij} r_{jk} r_{ki} r_{kl}^{\alpha}.$$

If  $\lim_{p\to\infty} \frac{p}{m\|\mathbf{R}\|_F} = 0$ , then  $\lim_{p\to\infty} \frac{S_m}{m\|\mathbf{R}\|_F^4} = 0$ .

**Proof of Lemma 6.32**. Set  $a_k = \sum_{l=1}^p r_{kl}^{\alpha}$  for  $k = 1, 2, \dots, p$ . Then  $|a_k| \leq p$  for each k. Define a  $p \times p$  matrix  $\mathbf{D} = \text{diag}(a_1, \dots, a_p)$ . Then the (k, i)-entry of  $\mathbf{RD}$  is  $r_{ki}a_i$ . It follows that

$$S_m = \sum_{1 \le i,j,k \le p} r_{ij} r_{jk} r_{ki} a_k = \sum_{1 \le i,j,k \le p} r_{ij} r_{jk} (\mathbf{RD})_{ki}$$

Therefore,  $S_m = \operatorname{tr}(\mathbf{RR}(\mathbf{RD})) = \operatorname{tr}(\mathbf{R}^3\mathbf{D})$ . Set  $(b_{ij})_{p \times p} = \mathbf{B} = \mathbf{R}^3$ . Then **B** is a non-negative definite matrix due to the fact that **R** is non-negative definite. Hence,  $b_{ii} \geq 0$  for each *i* and

$$|\operatorname{tr}(\mathbf{R}^{3}\mathbf{D})| = |\operatorname{tr}(\mathbf{B}\mathbf{D})| = \left|\sum_{i=1}^{p} b_{ii}a_{i}\right| \le p\sum_{i=1}^{p} b_{ii}.$$

This shows that  $|S_m| \leq p \cdot \operatorname{tr}(\mathbf{R}^3)$ . Let  $\lambda_1 \geq 0, \cdots, \lambda_p \geq 0$  be the eigenvalues of **R**. Easily,

$$\operatorname{tr}(\mathbf{R}^3) = \lambda_1^3 + \dots + \lambda_p^3 \le \left(\lambda_1^2 + \dots + \lambda_p^2\right)^{3/2}.$$

Therefore  $|S_m| \leq p \cdot [tr(\mathbf{R}^2)]^{3/2}$ . Consequently,

$$\frac{|S_m|}{m\,[\mathrm{tr}(\mathbf{R}^2)]^2} \le \frac{p}{m\,[\mathrm{tr}(\mathbf{R}^2)]^{1/2}} \to 0$$

by assumption. The proof is finished.

**LEMMA 6.33** Define  $\Lambda_p := \{(i, j, k, l); 1 \le i \ne j \le p \text{ and } 1 \le k \ne l \le p\},\$ 

$$V_{p,1} = \frac{1}{m \|\mathbf{R}\|_F^4} \cdot \sum_{(i,j,k,l) \in \Lambda_p} r_{ij} r_{jk} r_{kl} r_{li},$$
  
$$V_{p,2} = \frac{1}{m \|\mathbf{R}\|_F^4} \cdot \sum_{(i,j,k,l) \in \Lambda_p} (r_{ik} r_{kl} r_{li}) r_{ij}^2.$$

If  $\lim_{p\to\infty} \frac{p}{m\|\mathbf{R}\|_F} = 0$ , then  $\lim_{p\to\infty} V_{p,i} = 0$  for i = 1, 2.

**Proof of Lemma 6.33**. By assumption,  $r_{ii} = 1$  for any  $1 \le i \le p$ . Then

$$\sum_{1 \le i,j,k,l \le m} r_{ij} r_{jk} r_{kl} r_{li} = \sum_{1 \le i=j,k,l \le p} r_{ij} r_{jk} r_{kl} r_{li} + \sum_{1 \le i \ne j,k,l \le p} r_{ij} r_{jk} r_{kl} r_{li}$$
$$= \sum_{1 \le i,k,l \le p} r_{ik} r_{kl} r_{li} + \sum_{1 \le i \ne j,k=l \le p} r_{ij} r_{jk} r_{kl} r_{li} + \sum_{(i,j,k,l) \in \Lambda_p} r_{ij} r_{jk} r_{kl} r_{li}.$$

Now

$$\sum_{1 \le i \ne j, k=l \le m} r_{ij} r_{jk} r_{kl} r_{li} = \sum_{1 \le i \ne j, k \le m} r_{ij} r_{jk} r_{ki} = \sum_{1 \le i, j, k \le m} r_{ij} r_{jk} r_{ki} - \sum_{1 \le j, k \le m} r_{jk}^2.$$

Recall (145). The above two identities imply that

$$\sum_{(i,j,k,l)\in\Lambda_p} r_{ij}r_{jk}r_{kl}r_{li} = \operatorname{tr}(\mathbf{R}^4) - 2\operatorname{tr}(\mathbf{R}^3) + \operatorname{tr}(\mathbf{R}^2).$$

Then  $V_{p,1} \to 0$  by using the fact  $\|\mathbf{R}\|_F^2 \ge p$  and the conclusions for  $W_1$  and  $W_2$  in Lemma 6.31.

Now we prove  $V_{p,2} \to 0$ . Note that

$$\sum_{1 \le i,j,k,l \le p} (r_{ik}r_{kl}r_{li})r_{ij}^2 = \sum_{1 \le j,k,l \le p} r_{jk}r_{kl}r_{lj} + \sum_{1 \le i \ne j,k,l \le p} (r_{ik}r_{kl}r_{li})r_{ij}^2$$
$$= \operatorname{tr}(\mathbf{R}^3) + \sum_{1 \le i \ne j,k \le p} r_{ik}^2 r_{ij}^2 + \sum_{(i,j,k,l) \in \Lambda_p} (r_{ik}r_{kl}r_{li})r_{ij}^2.$$
(146)

Now

$$0 \le \sum_{1 \le i \ne j, k \le p} r_{ik}^2 r_{ij}^2 \le \sum_{i=1}^p \left(\sum_{j=1}^p r_{ij}^2\right)^2 \le \left(\sum_{i=1}^p \sum_{j=1}^p r_{ij}^2\right)^2 = \left[\operatorname{tr}(\mathbf{R}^2)\right]^2.$$

Hence,

$$\frac{1}{m \, [\operatorname{tr}(\mathbf{R}^2)]^2} \sum_{1 \le i \ne j, k \le p} r_{ik}^2 r_{ij}^2 \le \frac{1}{m} \to 0.$$

Also,  $\frac{1}{m[\operatorname{tr}(\mathbf{R}^2)]^2}\operatorname{tr}(\mathbf{R}^3) \to 0$  by using the conclusion for  $W_1$  in Lemma 6.31. Then the conclusion  $V_{p,2} \to 0$  follows from (146) and Lemma 6.32 with  $\alpha = 2$ .

**Proof of Lemma 4.3.** Write  $\mathbf{M} = (m_{ij})$ . Then  $m_{ii} = 1$  for each *i*. By Theorem 4.3.26 from Horn and Johnson (2012),  $\{\lambda_1, \dots, \lambda_p\}$  majorizes  $\{m_{11}, \dots, m_{pp}\}$ , the diagonal matrix of  $\mathbf{M}$ . By the definition of majorization,  $\lambda_1 + \dots + \lambda_k \ge m_{11} + \dots + m_{kk} = k$  for each  $1 \le k \le p$  and  $\lambda_1 + \dots + \lambda_p = m_{11} + \dots + m_{pp} = p$ .

On the other hand, by the definition of majorization, the p numbers  $\{1, \dots, 1\}$  majorizes  $\{\tau_1, \dots, \tau_p\}$ . By Theorem 4.3.32 from Horn and Johnson (2012), there exists a symmetric matrix  $\mathbf{B} = (b_{ij})_{p \times p}$  such that  $b_{ii} = 1$  for each i and that  $\mathbf{B}$  has non-negative eigenvalues  $\tau_1, \dots, \tau_p$ . By definition,  $\mathbf{B}$  is a correlation matrix.

## 6.2.6 The Proofs of Theorems 3 and 4

In this part, by using the preliminary results developed in Sections 6.2.1-6.2.5, we are now ready to prove the two main results Theorems 4 and 3 stated in Section 6.2.

**LEMMA 6.34** Assume the setting in (96) with  $\mathbf{R} = (r_{ij})_{4\times 4}$ . Recall  $A_i$  in (98). Define

$$B_1 = \frac{1}{m} \sum_{j=1}^m X_{1j} X_{2j}$$
 and  $B_2 = \frac{1}{m} \sum_{j=1}^m X_{3j} X_{4j}$ 

Given integer  $N \geq 1$ , the covariance between

$$\sum_{0 \le j,k \le N} (1 - A_1)^j (1 - A_2)^k B_1^2 \quad and \quad \sum_{0 \le j',k' \le N} (1 - A_3)^{j'} (1 - A_4)^{k'} B_2^2$$

is equal to

$$\begin{split} \varrho_{m,1} \cdot \sum_{1 \le i < j \le 4} r_{ij}^2 + \varrho_{m,2} \cdot r_{12}^2 r_{34}^2 &+ \varrho_{m,3} \cdot \left( r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} \right) \\ &+ \varrho_{m,4} \cdot \left[ (r_{13} r_{34} r_{41}) r_{12}^2 + (r_{23} r_{34} r_{42}) r_{12}^2 \right] \\ &+ \varrho_{m,5} \cdot \left[ (r_{12} r_{23} r_{31}) r_{34}^2 + (r_{12} r_{24} r_{41}) r_{34}^2 \right], \end{split}$$

where  $\{\varrho_{m,i}; 3 \leq i \leq 5\}$  do not depend on **R**,

$$|\varrho_{m,1}| \le Km^{-2}, \quad |\varrho_{m,2}| \lor |\varrho_{m,3}| \lor |\varrho_{m,4}| \lor |\varrho_{m,5}| \le Km^{-1}$$

and K is a constant depending on N but not on m or  $\mathbf{R}$ .

**Proof of Lemma 6.34**. For convenience, we use  $\Delta_m$  to denote the covariance between

$$\sum_{0 \le j,k \le N} (1 - A_1)^j (1 - A_2)^k B_1^2 \text{ and } \sum_{0 \le j',k' \le N} (1 - A_3)^{j'} (1 - A_4)^{k'} B_2^2.$$

Then

$$\Delta_m = \sum \operatorname{Cov} \left( (1 - A_1)^j (1 - A_2)^k B_1^2, (1 - A_3)^{j'} (1 - A_4)^{k'} B_2^2 \right), \tag{147}$$

where the sum runs over all non-negative integers j, j', k, k' such that  $0 \le j, k \le N$  and  $0 \le j', k' \le N$ . For each i = 1, 2, 3, 4, write

$$(1 - A_i)^l = 1 + \sum_{\alpha=1}^l (-1)^{\alpha} \binom{l}{\alpha} A_i^{\alpha}$$

for any  $l \ge 1$ . Trivially,  $\operatorname{Cov}(U_1 + h_1, U_2 + h_2) = \operatorname{Cov}(U_1, U_2)$  for any random variables  $U_1$ and  $U_2$  and constants  $h_1$  and  $h_2$ . Then the last covariance from (147) is

$$\operatorname{Cov}\left((1-A_1)^j(1-A_2)^k B_1^2, (1-A_3)^{j'}(1-A_4)^{k'} B_2^2\right)$$

= a finite linear combination of H terms of the form  $\operatorname{Cov}(B_1^2 A_1^{\alpha} A_2^{\beta}, B_2^2 A_3^{\gamma} A_4^{\delta})$ 

where the coefficients in the linear combination depend on  $\alpha, \beta, \gamma, \delta$  but not m or  $\mathbf{R}$ , and H := (j+1)(j'+1)(k+1)(k'+1). This and (147) imply that

 $\Delta_m = \text{a finite linear combination of } N' \text{ terms of the form } \operatorname{Cov}\left(B_1^2 A_1^{\alpha} A_2^{\beta}, B_2^2 A_3^{\gamma} A_4^{\delta}\right)$ (148)

where  $0 \leq \alpha + \beta \leq N$  and  $0 \leq \gamma + \delta \leq N$  and the coefficients in the linear combination depend on N but not on m or **R**, and N' is bounded by

$$\sum_{j,k \le N, j', k' \le N} (j+1)(j'+1)(k+1)(k'+1) \le (N+1)^4 \sum_{0 \le j, j', k, k' \le N} 1 \le (N+1)^8.$$

As  $\alpha + \beta = 0$  and  $\gamma + \delta = 0$ , the covariance becomes

$$\operatorname{Cov}(B_1^2, B_2^2) = \frac{1}{m} (r_{12}r_{23}r_{34}r_{41} + r_{12}r_{24}r_{43}r_{31}) + \frac{\delta_m}{m^2} \sum_{1 \le i < j \le 4} r_{ij}^2$$
(149)

by Lemma 6.9, where  $|\delta_m| \leq \kappa$  and  $\kappa$  is a numerical constant not depending on m, **R** or N. So we next only need to study  $\text{Cov}(B_1^2 A_1^{\alpha} A_2^{\beta}, B_2^2 A_3^{\gamma} A_4^{\delta})$  from (148) with an extra assumption that either  $\alpha + \beta \geq 1$  or  $\gamma + \delta \geq 1$ .

Write

$$m^{2} \cdot B_{1}^{2} = \sum_{j=1}^{m} (X_{1j} X_{2j})^{2} + 2 \sum_{1 \le k < l \le m} (X_{1k} X_{2k}) (X_{1l} X_{2l});$$
(150)

$$m^{2} \cdot B_{2}^{2} = \sum_{q=1}^{m} (X_{3q} X_{4q})^{2} + 2 \sum_{1 \le a < b \le m} (X_{3a} X_{4a}) (X_{3b} X_{4b}).$$
(151)

Then

$$m^{4} \cdot \operatorname{Cov}\left(B_{1}^{2}A_{1}^{\alpha}A_{2}^{\beta}, B_{2}^{2}A_{3}^{\gamma}A_{4}^{\delta}\right) = D_{1} + 2D_{2} + 2D_{3} + 4D_{4},$$
(152)

where

$$D_{1} = \sum_{1 \leq j,q \leq m} \operatorname{Cov} \left( (X_{1j}X_{2j})^{2} A_{1}^{\alpha} A_{2}^{\beta}, (X_{3q}X_{4q})^{2} A_{3}^{\gamma} A_{4}^{\delta} \right),$$
  

$$D_{2} = \sum_{1 \leq j \leq m, 1 \leq a < b \leq m} \operatorname{Cov} \left( (X_{1j}X_{2j})^{2} A_{1}^{\alpha} A_{2}^{\beta}, (X_{3a}X_{4a}) (X_{3b}X_{4b}) A_{3}^{\gamma} A_{4}^{\delta} \right),$$
  

$$D_{3} = \sum_{1 \leq q \leq m, 1 \leq k < l \leq m} \operatorname{Cov} \left( (X_{1k}X_{2k}) (X_{1l}X_{2l}) A_{1}^{\alpha} A_{2}^{\beta}, (X_{3q}X_{4q})^{2} A_{3}^{\gamma} A_{4}^{\delta} \right)$$

and

$$D_4 = \sum_{1 \le k < l \le m, 1 \le a < b \le m} \operatorname{Cov}((X_{1k}X_{2k})(X_{1l}X_{2l})A_1^{\alpha}A_2^{\beta}, (X_{3a}X_{4a})(X_{3b}X_{4b})A_3^{\gamma}A_4^{\delta}).$$

We next study the four terms in steps

Step 1: the estimate of  $D_1$ . Write

$$\sum_{\substack{1 \le j,q \le m \\ m}} \operatorname{Cov} \left( (X_{1j} X_{2j})^2 A_1^{\alpha} A_2^{\beta}, (X_{3q} X_{4q})^2 A_3^{\gamma} A_4^{\delta} \right)$$
  
= 
$$\sum_{j=1}^m \operatorname{Cov} \left( (X_{1j} X_{2j})^2 A_1^{\alpha} A_2^{\beta}, (X_{3j} X_{4j})^2 A_3^{\gamma} A_4^{\delta} \right)$$
  
+ 
$$\sum_{1 \le j \ne q \le m} \operatorname{Cov} \left( (X_{1j} X_{2j})^2 A_1^{\alpha} A_2^{\beta}, (X_{3q} X_{4q})^2 A_3^{\gamma} A_4^{\delta} \right).$$

Review  $\{(X_{1j}, X_{2j}, X_{3j}, X_{4j})^T \in \mathbb{R}^4; 1 \le j \le m\}$  are i.i.d. random vectors. It follows that

$$D_{1} = m \cdot \operatorname{Cov} \left( (X_{11}X_{21})^{2} A_{1}^{\alpha} A_{2}^{\beta}, (X_{31}X_{41})^{2} A_{3}^{\gamma} A_{4}^{\delta} \right) + m(m-1) \cdot \operatorname{Cov} \left( (X_{11}X_{21})^{2} A_{1}^{\alpha} A_{2}^{\beta}, (X_{32}X_{42})^{2} A_{3}^{\gamma} A_{4}^{\delta} \right).$$

By Lemma 6.24,

$$|D_1| \le K_1 m^2 \sum_{1 \le i < j \le 4} r_{ij}^2, \tag{153}$$

where  $K_1$  here and later denotes a constant depending on  $\alpha, \beta, \gamma, \delta$  but not m or  $\mathbf{R}$ , and can be different from line to line.

Step 2: the estimate of  $D_2$ . Write

$$D_{2} = \sum_{1 \le a < b \le m} \left( \sum_{j \in \{a,b\}} + \sum_{1 \le j \le m, j \notin \{a,b\}} \right) \operatorname{Cov} \left( (X_{1j} X_{2j})^{2} A_{1}^{\alpha} A_{2}^{\beta}, (X_{3a} X_{4a}) (X_{3b} X_{4b}) A_{3}^{\gamma} A_{4}^{\delta} \right)$$
$$= 2 \cdot \binom{m}{2} \cdot I_{m}(1,2) + (m-2) \binom{m}{2} \cdot I_{m}(2,3),$$

where

$$I_m(a,b) := \operatorname{Cov}((X_{11}X_{21})^2 A_1^{\alpha} A_2^{\beta}, (X_{3a}X_{4a})(X_{3b}X_{4b}) A_3^{\gamma} A_4^{\delta}).$$

By Lemma 6.25,

$$|D_2| \le K_1 m^2 \sum_{1 \le i < j \le 4} r_{ij}^2.$$
(154)

Step 3: the estimate of  $D_3$ . By switching the roles of " $(X_{1j}, X_{2j}, A_1, A_2, \alpha, \beta)$ " and " $(X_{3j}, X_{4j}, A_3, A_4, \gamma, \delta)$ " in Step 2, and using (154), we obtain

$$|D_3| \le K_1 m^2 \sum_{1 \le i < j \le 4} r_{ij}^2.$$
(155)

Step 4: the estimate of  $D_4$ . Rewrite

$$D_4 = \sum_{1 \le a < b \le m} \left( \sum_{\Gamma_1} + \sum_{\Gamma_2} + \sum_{\Gamma_3} \right) \operatorname{Cov} \left( (X_{1k} X_{2k}) (X_{1l} X_{2l}) A_1^{\alpha} A_2^{\beta}, (X_{3a} X_{4a}) (X_{3b} X_{4b}) A_3^{\gamma} A_4^{\delta} \right),$$

where

$$\Gamma_1 = \{(k,l) : (k,l) = (a,b)\}, \quad \Gamma_2 = \{(k,l) : 1 \le k < l \le m, |\{k,l\} \cap \{a,b\}| = 1\},$$
  
$$\Gamma_3 = \{(k,l) : 1 \le k < l \le m, \{k,l\} \cap \{a,b\} = \emptyset\}.$$

Given  $1 \le a < b \le m$ , it is easy to see  $|\Gamma_1| = 1$ ,  $|\Gamma_2| \le 2m$  and  $\Gamma_3 \le m^2$ . Set

$$J_m(a,b) = \operatorname{Cov}\left((X_{11}X_{21})(X_{12}X_{22})A_1^{\alpha}A_2^{\beta}, (X_{3a}X_{4a})(X_{3b}X_{4b})A_3^{\gamma}A_4^{\delta}\right)$$

for  $a \ge 1$  and  $b \ge 1$ . Then

$$D_4 = \frac{1}{2}m(m-1)\left[J_m(1,2) + |\Gamma_2| \cdot J_m(2,3) + |\Gamma_3| \cdot J_m(3,4)\right].$$

By Lemma 6.26,

$$|J_m(1,2)| \le K_1 \sum_{1 \le i < j \le 4} r_{ij}^2.$$

By Lemma 6.30,

$$J_m(2,3) = \tau_{m,1} \sum_{1 \le i < j \le 4} r_{ij}^2 + \tau_{m,2} \cdot \left( r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} \right)$$

where  $|\tau_{m,1}| \leq K_1 m^{-1}$ ,  $|\tau_{m,2}| \leq K_1$  and  $\tau_{m,2}$  does not depend on **R**. By Lemma 6.29,

$$J_m(3,4) = \tau'_{m,1}r_{12}^2r_{34}^2 + \tau'_{m,2}\sum_{1 \le i < j \le 4} r_{ij}^2 + \tau_{m,3} \cdot \left[ (r_{12}r_{23}r_{31})r_{34}^2 + (r_{12}r_{24}r_{41})r_{34}^2 \right] \\ + \tau_{m,4} \cdot \left[ (r_{13}r_{34}r_{41})r_{12}^2 + (r_{23}r_{34}r_{42})r_{12}^2 \right],$$

where  $|\tau'_{m,1}| \leq K_1 m^{-1}$ ,  $|\tau'_{m,2}| \leq K_1 m^{-2}$ ,  $|\tau_{m,3}| \vee |\tau_{m,4}| \leq K_1 m^{-1}$ , and  $\tau_{m,3}$  and  $\tau_{m,4}$  do not depend on **R**. Combining all of the above we get

$$\begin{split} D_4 &= \rho_{m,1} \cdot \sum_{1 \leq i < j \leq 4} r_{ij}^2 + \rho_{m,2} \cdot r_{12}^2 r_{34}^2 &+ \rho_{m,3} \cdot \left( r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} \right) \\ &+ \rho_{m,4} \cdot \left[ (r_{13} r_{34} r_{41}) r_{12}^2 + (r_{23} r_{34} r_{42}) r_{12}^2 \right] \\ &+ \rho_{m,5} \cdot \left[ (r_{12} r_{23} r_{31}) r_{34}^2 + (r_{12} r_{24} r_{41}) r_{34}^2 \right], \end{split}$$

where  $\{\rho_{m,i}, 1 \leq i \leq 5\}$  satisfy that

$$|\rho_{m,1}| \le K_1 m^2$$
,  $\max\{|\rho_{m,2}|, |\rho_{m,3}|, |\rho_{m,4}|, |\rho_{m,5}|\} \le K_1 m^3$ 

and  $\rho_{m,3}$ ,  $\rho_{m,4}$  and  $\rho_{m,5}$  do not depend on **R**. Through combining the estimates of  $D_1$ ,  $D_2$ ,  $D_3$ ,  $D_4$ and (152), we see  $\text{Cov}(B_1^2 A_1^{\alpha} A_2^{\beta}, B_2^2 A_3^{\gamma} A_4^{\delta})$  is equal to

$$\begin{split} \rho'_{m,1} \cdot \sum_{1 \le i < j \le 4} r_{ij}^2 + \rho'_{m,2} \cdot r_{12}^2 r_{34}^2 &+ \rho'_{m,3} \cdot \left( r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} \right) \\ &+ \rho'_{m,4} \cdot \left[ (r_{13} r_{34} r_{41}) r_{12}^2 + (r_{23} r_{34} r_{42}) r_{12}^2 \right] \\ &+ \rho'_{m,5} \cdot \left[ (r_{12} r_{23} r_{31}) r_{34}^2 + (r_{12} r_{24} r_{41}) r_{34}^2 \right], \end{split}$$

where  $\{\rho_{m,i}'; 1 \le i \le 5\}$  satisfy that

$$|\rho'_{m,1}| \le K_1 m^{-2}, \ |\rho'_{m,2}| \lor |\rho'_{m,3}| \lor |\rho'_{m,4}| \lor |\rho'_{m,5}| \le K_1 m^{-1},$$

and  $\rho'_{m,3}, \rho'_{m,4}$  and  $\rho'_{m,5}$  do not depend on **R**. Recalling (148), we arrive at that  $\Delta_m$  is equal to

$$\begin{split} \varrho_{m,1} \cdot \sum_{1 \le i < j \le 4} r_{ij}^2 + \varrho_{m,2} \cdot r_{12}^2 r_{34}^2 &+ \varrho_{m,3} \cdot \left( r_{12} r_{23} r_{34} r_{41} + r_{12} r_{24} r_{43} r_{31} \right) \\ &+ \varrho_{m,4} \cdot \left[ (r_{13} r_{34} r_{41}) r_{12}^2 + (r_{23} r_{34} r_{42}) r_{12}^2 \right] \\ &+ \varrho_{m,5} \cdot \left[ (r_{12} r_{23} r_{31}) r_{34}^2 + (r_{12} r_{24} r_{41}) r_{34}^2 \right], \end{split}$$

where

$$|\varrho_{m,1}| \le K_1 m^{-2}, \ |\varrho_{m,2}| \lor |\varrho_{m,3}| \lor |\varrho_{m,4}| \lor |\varrho_{m,5}| \le K_1 m^{-1},$$

and  $\varrho_{m,3}, \varrho_{m,4}$  and  $\varrho_{m,5}$  do not depend on **R**. The proof is completed.

We will first prove Theorem 4 and then prove 3.

**Proof of Theorem 4**. Recall the earlier notation that

$$A_{i} = \frac{1}{m} \sum_{j=1}^{m} X_{ij}^{2}, \qquad B_{1} = \frac{1}{m} \sum_{j=1}^{m} X_{1j} X_{2j}, \qquad B_{2} = \frac{1}{m} \sum_{j=1}^{m} X_{3j} X_{4j}$$

for i = 1, 2, 3, 4. Then

$$\hat{r}_{12} = \frac{B_1}{\sqrt{A_1 A_2}}$$
 and  $\hat{r}_{34} = \frac{B_2}{\sqrt{A_3 A_4}}$ . (156)

Given  $N \ge 1$ , write

$$\frac{1}{x} = 1 + (1 - x) + \dots + (1 - x)^N + \frac{1}{x}(1 - x)^{N+1}$$

for  $x \neq 0$ . Thus

$$\frac{1}{A_1A_2} = \left[\frac{(1-A_1)^{N+1}}{A_1} + \sum_{i=0}^N (1-A_1)^i\right] \cdot \left[\frac{(1-A_2)^{N+1}}{A_2} + \sum_{j=0}^N (1-A_2)^j\right]$$
$$= \epsilon_{m,1} + \sum_{0 \le i,j \le N} (1-A_1)^i (1-A_2)^j,$$

where

$$\epsilon_{m,1} = \frac{(1-A_1)^{N+1}(1-A_2)^{N+1}}{A_1A_2} + \sum_{j=0}^N \frac{(1-A_1)^{N+1}(1-A_2)^j}{A_1} + \sum_{i=0}^N \frac{(1-A_1)^i(1-A_2)^{N+1}}{A_2}.$$

Similarly,

$$\frac{1}{A_3 A_4} = \epsilon_{m,2} + \sum_{0 \le i,j \le N} (1 - A_3)^i (1 - A_4)^j$$

where

$$\epsilon_{m,2} = \frac{(1-A_3)^{N+1}(1-A_4)^{N+1}}{A_3A_4} + \sum_{j=0}^N \frac{(1-A_3)^{N+1}(1-A_4)^j}{A_3} + \sum_{i=0}^N \frac{(1-A_3)^i(1-A_4)^{N+1}}{A_4}.$$

By (156),

$$\operatorname{Cov}(\hat{r}_{12}^2, \hat{r}_{34}^2) = \operatorname{Cov}\left(\sum_{0 \le i, j \le N} (1 - A_1)^i (1 - A_2)^j B_1^2, \sum_{0 \le i, j \le N} (1 - A_3)^i (1 - A_4)^j B_2^2\right) + \operatorname{Cov}\left(\sum_{0 \le i, j \le N} (1 - A_1)^i (1 - A_2)^j B_1^2, \epsilon_{m,2} B_2^2\right) + \operatorname{Cov}\left(\epsilon_{m,1} B_1^2, \sum_{0 \le i, j \le N} (1 - A_3)^i (1 - A_4)^j B_2^2\right) + \operatorname{Cov}\left(\epsilon_{m,1} B_1^2, \epsilon_{m,2} B_2^2\right).$$
(157)

We claim that

the absolute value of each of the last three covariances in (157)  $\leq \frac{K_1}{m^{(N+1)/2}}$  (158)

where  $K_1$  is a constant depending on N but not m or **R**. In fact, by writing  $\bar{B}_1 = B_1 - r_{12}$  and  $\bar{B}_2 = B_2 - r_{34}$ , then  $B_1^2 = \bar{B}_1^2 + 2r_{12}\bar{B}_1 + r_{12}^2$  and  $B_2^2 = \bar{B}_2^2 + 2r_{34}\bar{B}_2 + r_{34}^2$ , and hence by linearity of the covariance, each of the last three covariances from (157) is a linear combination of N' terms of the form

$$r_{12}^{a}r_{34}^{b} \cdot E \frac{\bar{B}_{1}^{t_{1}}\bar{B}_{2}^{t_{2}}(1-A_{1})^{n_{1}}(1-A_{2})^{n_{2}}(1-A_{3})^{n_{3}}(1-A_{4})^{n_{4}}}{A_{1}^{s_{1}}A_{2}^{s_{2}}A_{3}^{s_{3}}A_{4}^{s_{4}}}$$
(159)

where N' depends only on N and all powers are non-negative integers with

$$a, b, t_i \in \{0, 1, 2\}$$
 and  $s_1 + s_2 + s_3 + s_4 \ge 1;$   
 $n_i \le N$  and  $n_1 + n_2 + n_3 + n_4 \ge N + 1$ 

for each possible *i*. The crucial observation is that  $n_1 + n_2 + n_3 + n_4 \ge N + 1$ . If some of  $\{t_1, t_2, n_1, n_2, n_3, n_4\}$  are zero, the corresponding terms simply disappear. Set

k = the count of positive values from  $\{t_1, t_2, n_1, n_2, n_3, n_4\};$ l = the count of positive values from  $\{s_1, s_2, s_3, s_4\}.$ 

Then  $k \ge 1$  and  $l \ge 1$ . Take

$$X_{i} = \begin{cases} \sqrt{m}\bar{B}_{i}, & i = 1, 2; \\ \sqrt{m}(1 - A_{i-2}), & i = 3, 4, 5, 6 \end{cases} \text{ and } \alpha_{i} = \begin{cases} t_{i}, & i = 1, 2; \\ n_{i-2}, & i = 3, 4, 5, 6 \end{cases}$$

and  $p_i = m$  for  $i = 1, \dots, 6$ . Furthermore, take  $Y_j = A_j, q_j = m$  and  $\beta_j = s_j$  for j = 1, 2, 3, 4. Easily,  $2/(k+l) \leq 1$  and  $(2-q_j)/[2(k+l)] < 0$  as  $m \geq 5$ . Observe

$$X_{1} = \frac{1}{\sqrt{m}} \sum_{j=1}^{m} (X_{1j} X_{2j} - r_{12}), \quad X_{2} = \frac{1}{\sqrt{m}} \sum_{j=1}^{m} (X_{3j} X_{4j} - r_{34})$$
$$X_{i+2} = \frac{1}{\sqrt{m}} \sum_{j=1}^{m} (1 - X_{ij}^{2}), \qquad Y_{j} \sim \frac{1}{m} \chi^{2}(m)$$

for i = 1, 2, 3, 4 and j = 1, 2, 3, 4. Set  $\xi_{1j} = X_{1j}X_{2j} - r_{12}, \xi_{2j} = X_{3j}X_{4j} - r_{34}$  and  $\xi_{i+2j} = 1 - X_{ij}^2$ for i = 1, 2, 3, 4. Notice  $X_{ij} \sim N(0, 1)$  and  $|r_{ij}| \leq 1$  for each i, j. By Lemma 6.11,

$$\left| r_{12}^{a} r_{34}^{b} \cdot E \frac{\bar{B}_{1}^{t_{1}} \bar{B}_{2}^{t_{2}} (1-A_{1})^{n_{1}} (1-A_{2})^{n_{2}} (1-A_{3})^{n_{3}} (1-A_{4})^{n_{4}}}{A_{1}^{s_{1}} A_{2}^{s_{2}} A_{3}^{s_{3}} A_{4}^{s_{4}}} \right| \leq \frac{K_{1}}{m^{(N+1)/2}},$$

where  $K_1$  is a constant depending on N but not on m or **R**. This confirms claim (158). The first covariance on the right hand side of (157) is studied in Lemma 6.34. Combining this lemma and (158), we finish the proof.

**LEMMA 6.35** Let  $X_1, \dots, X_n$  be a random sample from  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  with correlation matrix  $\mathbf{R}$ . Let  $\hat{\mathbf{R}}$  be defined in (7). Assume, for some a > 0,  $p \leq n^a$  for each  $p \geq 1$ . If  $\lim_{p\to\infty} \frac{p}{n\|\mathbf{R}\|_F} = 0$ , then  $Var(tr(\hat{\mathbf{R}}^2)) \cdot \|\mathbf{R}\|_F^{-2}$  goes to zero as  $p \to \infty$ .

**Proof of Lemma 6.35**. Set m = n - 1. By (28) from the proof of Lemma 6.3,

$$\hat{\mathbf{R}} = \hat{\mathbf{R}}_p = (\hat{r}_{ij})_{p \times p} \stackrel{d}{=} \left( \frac{\mathbf{v}_i^T \mathbf{v}_j}{\|\mathbf{v}_i^T\| \cdot \|\mathbf{v}_j\|} \right)_{p \times p},$$

where the *m* rows of  $(\mathbf{v}_1, \cdots, \mathbf{v}_p)_{m \times p}$  are i.i.d. with distribution  $N_p(\mathbf{0}, \mathbf{R})$ . Write  $\operatorname{tr}(\hat{\mathbf{R}}^2) = p + \sum_{1 \le i \ne j \le m} \hat{r}_{ij}^2$ . Then,

$$\operatorname{Var}\left(\operatorname{tr}(\hat{\mathbf{R}}^{2})\right) = \operatorname{Cov}\left(\sum_{1 \le i \ne j \le p} \hat{r}_{ij}^{2}, \sum_{1 \le k \ne l \le p} \hat{r}_{kl}^{2}\right) = \sum \operatorname{Cov}\left(\hat{r}_{ij}^{2}, \hat{r}_{kl}^{2}\right),$$
(160)

where the last sum runs over all  $(i, j, k, l) \in \Lambda_p$ , where

$$\Lambda_p := \left\{ (i, j, k, l); \ 1 \le i \ne j \le p \text{ and } 1 \le k \ne l \le p \right\}.$$

$$(161)$$

Review Theorem 4. We never impose any condition on the  $4 \times 4$  correlation matrix  $\mathbf{R}_{4\times 4}$ (not confuse the  $p \times p$  correlation matrix  $\mathbf{R}$  here) in the proposition. For example, if all of the entries of  $\mathbf{R}_{4\times 4}$  are equal to 1, then the four random variables are actually equal. Keeping this understanding in mind, by changing "(1, 2, 3, 4)" in Theorem 4 to "(i, j, k, l)" and taking N=7 in the proposition, we see  $\mathrm{Cov}(\hat{r}_{ij}^2,\hat{r}_{kl}^2)$  is equal to

$$\varrho_{p,1} \cdot \sum_{u,v \in \{i,j,k,l\}, u \neq v} r_{uv}^2 + \varrho_{p,2} \cdot r_{ij}^2 r_{kl}^2 + \varrho_{p,3} \cdot \left( r_{ij} r_{jk} r_{kl} r_{li} + r_{ij} r_{jl} r_{lk} r_{ki} \right) \\
+ \varrho_{p,4} \cdot \left[ (r_{ik} r_{kl} r_{li}) r_{ij}^2 + (r_{jk} r_{kl} r_{lj}) r_{ij}^2 \right] \\
+ \varrho_{p,5} \cdot \left[ (r_{ij} r_{jk} r_{ki}) r_{kl}^2 + (r_{ij} r_{jl} r_{li}) r_{kl}^2 \right] \\
+ \frac{\varrho_{p,6}}{m^4},$$
(162)

where the sum runs over the six pairs from  $\{i, j, k, l\}$ ,

$$|\varrho_{p,1}| \le Km^{-2}; \quad |\varrho_{p,2}| \lor |\varrho_{p,3}| \lor |\varrho_{p,4}| \lor |\varrho_{p,5}| \le Km^{-1}, \quad |\varrho_{p,6}| \le K,$$
(163)

 $\{\varrho_{p,i}; 3 \leq i \leq 5\}$  do not depend on **R**, and K is a constant not depending on m or **R**. Notice

$$\sum_{(i,j,k,l)\in\Lambda_p} \sum_{u,v\in\{i,j,k,l\}, u\neq v} r_{uv}^2 \le 6p^2 \sum_{1\le i,j\le p} r_{ij}^2 = 6p^2 \cdot \operatorname{tr}(\mathbf{R}^2).$$

Also,

$$\sum_{(i,j,k,l)\in\Lambda_p} r_{ij}^2 r_{kl}^2 \le \sum_{1\le i,j\le p} r_{ij}^2 \cdot \sum_{1\le k,l\le p} r_{kl}^2 = \left[ \operatorname{tr}(\mathbf{R}^2) \right]^2.$$

Recall  $\|\mathbf{R}\|_F^4 = [\operatorname{tr}(\mathbf{R}^2)]^2$ . The two facts together with (160) and (162) imply that

$$\frac{1}{\|\mathbf{R}\|_F^4} \cdot \operatorname{Var}\left(\operatorname{tr}(\hat{\mathbf{R}}^2)\right)$$

$$\leq (6K) \cdot \frac{p^2}{m^2 \cdot \operatorname{tr}(\mathbf{R}^2)} + \frac{K}{m} + \left(\sum_{i=3}^5 |m\varrho_{p,i}| \cdot |Q_{p,i}|\right) + \frac{Kp^4}{[\operatorname{tr}(\mathbf{R}^2)]^2 \cdot m^4}$$

where

$$Q_{p,3} = \frac{1}{m[\operatorname{tr}(\mathbf{R}^{2})]^{2}} \cdot \sum_{(i,j,k,l)\in\Lambda_{p}} \left( r_{ij}r_{jk}r_{kl}r_{li} + r_{ij}r_{jl}r_{lk}r_{ki} \right),$$
  

$$Q_{p,4} = \frac{1}{m[\operatorname{tr}(\mathbf{R}^{2})]^{2}} \cdot \sum_{(i,j,k,l)\in\Lambda_{p}} \left[ (r_{ik}r_{kl}r_{li})r_{ij}^{2} + (r_{jk}r_{kl}r_{lj})r_{ij}^{2} \right],$$
  

$$Q_{p,5} = \frac{1}{m[\operatorname{tr}(\mathbf{R}^{2})]^{2}} \cdot \sum_{(i,j,k,l)\in\Lambda_{p}} \left[ (r_{ij}r_{jk}r_{ki})r_{kl}^{2} + (r_{ij}r_{jl}r_{li})r_{kl}^{2} \right].$$

Now, by assumption,  $p = o(m \|\mathbf{R}\|_F)$ , hence

$$\frac{p^2}{m^2 \cdot \operatorname{tr}(\mathbf{R}^2)} \to 0 \text{ and } \frac{Kp^4}{[\operatorname{tr}(\mathbf{R}^2)]^2 \cdot m^4} \to 0.$$

Because of (163), to prove the conclusion, it suffices to show  $\lim_{p\to\infty} Q_{p,i} = 0$  for i = 3, 4, 5. Recall (161). By switching "k" and "l" in  $r_{ij}r_{jl}r_{lk}r_{ki}$ , switching "k" and "l" in  $(r_{jk}r_{kl}r_{lj})r_{ij}^2$ and switching "k" and "l" in  $(r_{ij}r_{jl}r_{li})r_{kl}^2$ , respectively, we obtain

$$Q_{p,3} = \frac{2}{m[\operatorname{tr}(\mathbf{R}^{2})]^{2}} \cdot \sum_{(i,j,k,l)\in\Lambda_{p}} r_{ij}r_{jk}r_{kl}r_{li},$$

$$Q_{p,4} = \frac{2}{m[\operatorname{tr}(\mathbf{R}^{2})]^{2}} \cdot \sum_{(i,j,k,l)\in\Lambda_{p}} (r_{ik}r_{kl}r_{li})r_{ij}^{2},$$

$$Q_{p,5} = \frac{2}{m[\operatorname{tr}(\mathbf{R}^{2})]^{2}} \cdot \sum_{(i,j,k,l)\in\Lambda_{p}} (r_{ij}r_{jk}r_{ki})r_{kl}^{2}.$$

By interchanging "(i, j)" with "(k, l)" in the last sum, we see  $Q_{p,4} = Q_{p,5}$ . Finally, we see from Lemma 6.33 that  $\lim_{p\to\infty} Q_{p,i} = 0$  for i = 3, 4, 5.

**Proof of Theorem 3**. Recall  $\hat{\mathbf{R}} = \hat{\mathbf{R}}_p$ . By Lemma 6.35,

$$\frac{\operatorname{tr}(\hat{\mathbf{R}}_p^2) - E \operatorname{tr}(\hat{\mathbf{R}}_p^2)}{\operatorname{tr}(\mathbf{R}_p^2)} \to 0$$
(164)

in probability as  $p \to \infty$ . By Lemma 6.3, under the assumption  $\limsup_{p\to\infty} \frac{p}{n^a} = 0$  for some constant a > 0, we have

$$E \operatorname{tr}(\hat{\mathbf{R}}_p^2) = \frac{p(p-1)}{n-1} + \operatorname{tr}(\mathbf{R}_p^2) \cdot \left[1 + O(m^{-1/4})\right].$$

This implies that

$$\frac{1}{\operatorname{tr}(\mathbf{R}_p^2)} \cdot \left[ E \operatorname{tr}(\hat{\mathbf{R}}_p^2) - \frac{p(p-1)}{n-1} - \operatorname{tr}(\mathbf{R}_p^2) \right] \to 0.$$

The proof is completed by adding this and that from (164).

## 6.3 The Proofs of Theorems 1 and 2 and Proposition 1

Let  $\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_n$  be i.i.d. *p*-dimensional random vectors with distribution  $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ . Let **D** be the diagonal matrix of  $\boldsymbol{\Sigma}$ . The  $p \times p$  population correlation matrix is defined by  $\mathbf{R} = \mathbf{D}^{-1/2} \boldsymbol{\Sigma} \mathbf{D}^{-1/2}$ . The sample mean is  $\bar{\boldsymbol{\xi}} = \frac{1}{n} (\boldsymbol{\xi}_1 + \dots + \boldsymbol{\xi}_n)$  and the sample covariance matrix is defined by

$$\hat{\mathbf{S}} = \frac{1}{n} \sum_{i=1}^{n} (\boldsymbol{\xi}_i - \bar{\boldsymbol{\xi}}) (\boldsymbol{\xi}_i - \bar{\boldsymbol{\xi}})^T.$$
(165)

Review  $W_p(m, \Sigma)$  stands for the distribution of the Wishart matrix  $\mathbf{U}^T \mathbf{U}$  for any  $m \geq 1$ , where  $\mathbf{U}$  is an  $m \times p$  matrix whose rows are i.i.d. with distribution  $N_p(\mathbf{0}, \Sigma)$ . Then,  $n\hat{\mathbf{S}}$ has the Wishart distribution  $W_p(n-1, \Sigma)$ ; see, for example, Theorem 3.1.2 from Muirhead (1982). Let  $\hat{\mathbf{D}}$  be the diagonal matrix of  $\hat{\mathbf{S}}$ . Then  $\hat{\mathbf{R}} := \hat{\mathbf{D}}^{-1/2}\hat{\mathbf{S}}\hat{\mathbf{D}}^{-1/2}$  is the sample correlation matrix generated by  $\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_n$ . Before proving Theorems 1 and 2, we first will reduce the test statistic appearing in Theorem 1 to a simple form. Recall we assume *n* depends on *p* and sometimes write  $n_p$  if there is any possible confusion. Also, the Frobenius norm  $\|\mathbf{R}\|_F = [\operatorname{tr}(\mathbf{R}^2)]^{1/2}$  and the notation  $o_p(1)$  representing a random variable converging to 0 in probability.

**Proof of Lemma 4.1**. We need to show

$$\frac{\boldsymbol{\eta}^T \hat{\mathbf{D}}^{-1} \boldsymbol{\eta} - \boldsymbol{\eta}^T \mathbf{D}^{-1} \boldsymbol{\eta}}{\sqrt{\operatorname{tr}(\mathbf{R}^2)}} \to 0$$
(166)

in probability as  $p \to \infty$ . To do so, it suffices to prove that both its mean and variance converging to 0.

Step 1: the mean of random variable from (166). Write  $\boldsymbol{\eta} = \boldsymbol{\Sigma}^{1/2} \boldsymbol{\theta}$  where  $\boldsymbol{\theta} \sim N_p(\mathbf{0}, \boldsymbol{I}_p)$ and  $\boldsymbol{\Sigma}^{1/2}$  is a non-negative definite matrix satisfying  $\boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{1/2} = \boldsymbol{\Sigma}$ . By assumption,  $\boldsymbol{\theta}$  is independent of  $\hat{\mathbf{S}}$ . In particular,  $\boldsymbol{\theta}$  is independent of  $\hat{\mathbf{D}}$ , the diagonal matrix of  $\hat{\mathbf{S}}$ . Notice

$$\boldsymbol{\eta}^{T} \hat{\mathbf{D}}^{-1} \boldsymbol{\eta} = \boldsymbol{\theta}^{T} \left( \boldsymbol{\Sigma}^{1/2} \hat{\mathbf{D}}^{-1} \boldsymbol{\Sigma}^{1/2} \right) \boldsymbol{\theta} \text{ and } \boldsymbol{\eta}^{T} \mathbf{D}^{-1} \boldsymbol{\eta} = \boldsymbol{\theta}^{T} \left( \boldsymbol{\Sigma}^{1/2} \mathbf{D}^{-1} \boldsymbol{\Sigma}^{1/2} \right) \boldsymbol{\theta}.$$
(167)

For any  $p \times p$  symmetric matrix **A** with eigenvalues  $\lambda_1, \dots, \lambda_p$ , by the orthogonal invariance of  $N_p(\mathbf{0}, \mathbf{I}_p)$ , we know  $\boldsymbol{\theta}^T \mathbf{A} \boldsymbol{\theta}$  and  $\lambda_1 \theta_1^2 + \dots + \lambda_p \theta_p^2$  have the same distribution, where  $\theta_1, \dots, \theta_p$  are i.i.d. N(0, 1)-distributed random variables. Consequently,

$$E(\boldsymbol{\theta}^T \mathbf{A} \boldsymbol{\theta}) = \operatorname{tr}(\mathbf{A}) \text{ and } \operatorname{Var}(\boldsymbol{\theta}^T \mathbf{A} \boldsymbol{\theta}) = 2 \operatorname{tr}(\mathbf{A}^2).$$
 (168)

It follows from independence and conditioning on  $\hat{\mathbf{D}}$  that

$$E(\boldsymbol{\eta}^T \hat{\mathbf{D}}^{-1} \boldsymbol{\eta}) = E \operatorname{tr} \left( \boldsymbol{\Sigma}^{1/2} \hat{\mathbf{D}}^{-1} \boldsymbol{\Sigma}^{1/2} \right) = \operatorname{tr} \left[ \boldsymbol{\Sigma}^{1/2} E(\hat{\mathbf{D}}^{-1}) \boldsymbol{\Sigma}^{1/2} \right]$$
(169)

by linearity of expectations and traces, where  $E(\hat{\mathbf{D}}^{-1})$  is the entry-wise expectation of the diagonal matrix  $\hat{\mathbf{D}}^{-1}$ . Set  $\boldsymbol{\Sigma} = (\sigma_{ij})_{p \times p}$ . Then  $\mathbf{D} = \text{diag}(\sigma_{11}, \cdots, \sigma_{pp})$ . Set m = n - 1. It is known

$$n\hat{\mathbf{S}} \stackrel{d}{=} \sum_{j=1}^{m} \hat{\boldsymbol{\xi}}_{j} \hat{\boldsymbol{\xi}}_{j}^{T}$$
(170)

and  $\hat{\mathbf{S}}$  is independent of  $\bar{\boldsymbol{\xi}} = \frac{1}{n} (\boldsymbol{\xi}_1 + \cdots + \boldsymbol{\xi}_n)$ , where  $\hat{\boldsymbol{\xi}}_1, \cdots, \hat{\boldsymbol{\xi}}_m$  are i.i.d.  $N_p(\mathbf{0}, \boldsymbol{\Sigma})$ -distributed random vectors; see, for example, Theorem 3.1.2 from Muirhead (1982). Write  $\hat{\boldsymbol{\xi}}_j = (\xi_{1j}, \cdots, \xi_{pj})^T$  for each j. Then the (i, i)-entry of  $\hat{\boldsymbol{\xi}}_j \hat{\boldsymbol{\xi}}_j^T$  is equal to  $\xi_{ij}^2$ . As a result,

the 
$$(i, i)$$
-entry of  $n\hat{\mathbf{S}}$  is  $\sum_{j=1}^{m} \xi_{ij}^2 \sim \sigma_{ii}^2 \cdot \chi^2(m)$  (171)

for each  $1 \leq i \leq p$ . Since  $\hat{\mathbf{D}} = \text{diag}(s_{11}, \cdots, s_{pp})$  is the diagonal matrix of  $\hat{\mathbf{S}} := (s_{ij})_{p \times p}$ , we know  $ns_{ii}/\sigma_{ii} \sim \chi^2(m)$  for each *i*. It is known that

$$E\frac{1}{\chi^2(k)} = \frac{1}{k-2}$$
 and  $\operatorname{Var}\left(\frac{1}{\chi^2(k)}\right) = \frac{2}{(k-2)^2(k-4)}$  (172)

for any integer  $k \geq 3$ . Therefore,

$$E\frac{1}{s_{ii}} = \frac{n}{(m-2)\sigma_{ii}} \quad \text{and} \quad \operatorname{Var}\left(\frac{1}{s_{ii}}\right) = \frac{2n^2}{(m-2)^2(m-4)\sigma_{ii}^2}.$$
 (173)

It follows that  $E(\hat{\mathbf{D}}^{-1}) = \frac{n}{m-2}\mathbf{D}^{-1}$ . Observe  $tr(\mathbf{\Sigma}^{1/2}\mathbf{D}^{-1}\mathbf{\Sigma}^{1/2}) = tr(\mathbf{D}^{-1/2}\mathbf{\Sigma}\mathbf{D}^{-1/2}) = tr(\mathbf{R}) = p$ . From (169), we have

$$E(\boldsymbol{\eta}^T \hat{\mathbf{D}}^{-1} \boldsymbol{\eta}) = \frac{n}{m-2} \cdot \operatorname{tr} \left( \boldsymbol{\Sigma}^{1/2} \mathbf{D}^{-1} \boldsymbol{\Sigma}^{1/2} \right) = \frac{np}{n-3}.$$
 (174)

Similarly, we have from (167) that

$$E(\boldsymbol{\eta}^T \mathbf{D}^{-1} \boldsymbol{\eta}) = E(\boldsymbol{\theta}^T \boldsymbol{\Sigma}^{1/2} \mathbf{D}^{-1} \boldsymbol{\Sigma}^{1/2} \boldsymbol{\theta}) = \operatorname{tr}(\boldsymbol{\Sigma}^{1/2} \mathbf{D}^{-1} \boldsymbol{\Sigma}^{1/2}) = p.$$

Therefore,

$$E(\boldsymbol{\eta}^T \hat{\mathbf{D}}^{-1} \boldsymbol{\eta}) - E(\boldsymbol{\eta}^T \mathbf{D}^{-1} \boldsymbol{\eta}) = \frac{np}{m-2} - p = \frac{3p}{m-2}$$

It follows that

$$\frac{1}{\sqrt{2\operatorname{tr}(\mathbf{R}^2)}} \left[ E(\boldsymbol{\eta}^T \hat{\mathbf{D}}^{-1} \boldsymbol{\eta}) - E(\boldsymbol{\eta}^T \mathbf{D}^{-1} \boldsymbol{\eta}) \right] = \frac{\sqrt{4.5} \, p}{(m-2)\sqrt{\operatorname{tr}(\mathbf{R}^2)}} \to 0 \tag{175}$$

by the assumption  $\lim_{p\to\infty} \frac{p}{m \|\mathbf{R}\|_F} = 0.$ 

Step 2: the variance of random variable from (166). Set  $\mathbf{B} = \Sigma^{1/2} (\hat{\mathbf{D}}^{-1} - \mathbf{D}^{-1}) \Sigma^{1/2}$ . It is seen from (167) that

$$\boldsymbol{\theta}^T \mathbf{B} \boldsymbol{\theta} = \boldsymbol{\eta}^T \hat{\mathbf{D}}^{-1} \boldsymbol{\eta} - \boldsymbol{\eta}^T \mathbf{D}^{-1} \boldsymbol{\eta}.$$
(176)

Recall the formula  $\operatorname{Var}(v) = E\operatorname{Var}(v|\mathbf{B}) + \operatorname{Var}(E(v|\mathbf{B}))$  for any random variable v. Then, by the independence between  $\boldsymbol{\theta}$  and  $\mathbf{B}$  as well as (168),

$$\operatorname{Var}(\boldsymbol{\theta}^{T} \mathbf{B} \boldsymbol{\theta}) = 2E \operatorname{tr}(\mathbf{B}^{2}) + \operatorname{Var}(\operatorname{tr}(\mathbf{B})).$$
(177)

Our focus next will be the evaluation of the two terms.

Step 3: the evaluation of  $Etr(\mathbf{B}^2)$  from (177). Let us consider the last two terms one by one. First,

$$\begin{aligned} \operatorname{tr}(\mathbf{B}^2) &= \operatorname{tr} \big( \mathbf{\hat{D}}^{-1} - \mathbf{D}^{-1} \big) \mathbf{\Sigma} (\hat{\mathbf{D}}^{-1} - \mathbf{D}^{-1}) \mathbf{\Sigma}^{1/2} \big) \\ &= \operatorname{tr} \big( (\hat{\mathbf{D}}^{-1} - \mathbf{D}^{-1}) \mathbf{\Sigma} (\hat{\mathbf{D}}^{-1} - \mathbf{D}^{-1}) \mathbf{\Sigma} \big). \end{aligned}$$

Let  $\mathbf{Q}(i, j)$  denote the (i, j)-entry of a matrix  $\mathbf{Q}$ . For any matrices  $\mathbf{Q}_1, \mathbf{Q}_2, \mathbf{Q}_3, \mathbf{Q}_4$ , we have  $\operatorname{tr}(\mathbf{Q}_1\mathbf{Q}_2\mathbf{Q}_3\mathbf{Q}_4) = \sum \mathbf{Q}_1(i, j)\mathbf{Q}_2(j, k)\mathbf{Q}_3(k, l)\mathbf{Q}_4(l, i)$ , where the sum runs over all possible indices i, j, k, l. It follows that

$$\operatorname{tr}(\mathbf{B}^{2}) = \sum_{1 \leq i,j \leq p} \sigma_{ij}^{2} \left(\frac{1}{s_{ii}} - \frac{1}{\sigma_{ii}}\right) \left(\frac{1}{s_{jj}} - \frac{1}{\sigma_{jj}}\right)$$
$$= \sum_{1 \leq i,j \leq p} r_{ij}^{2} \left(\frac{\sigma_{ii}}{s_{ii}} - 1\right) \left(\frac{\sigma_{jj}}{s_{jj}} - 1\right)$$
(178)

since  $r_{ij} = \sigma_{ij} (\sigma_{ii} \sigma_{jj})^{-1/2}$ . By (173),

$$\frac{\sigma_{ii}}{s_{ii}} - 1 = \frac{\sigma_{ii}}{s_{ii}} - E\frac{\sigma_{ii}}{s_{ii}} + \frac{3}{m-2}$$

Therefore,

$$E\left(\frac{\sigma_{ii}}{s_{ii}}-1\right)\left(\frac{\sigma_{jj}}{s_{jj}}-1\right) = \operatorname{Cov}\left(\frac{\sigma_{ii}}{s_{ii}},\frac{\sigma_{jj}}{s_{jj}}\right) + \frac{9}{(m-2)^2}.$$
(179)

The fact from (171) implies that  $ns_{ii} = X_1^2 + \cdots + X_m^2$  and  $ns_{jj} = Y_1^2 + \cdots + Y_m^2$ , where  $(X_1, Y_1)^T, \cdots, (X_m, Y_m)^T$  are i.i.d. 2-dimensional normal random vectors with  $EX_1 = EY_1 = 0$ ,  $EX_1^2 = \sigma_{ii}$  and  $EY_1^2 = \sigma_{jj}$  and  $Cov(X_1, Y_1) = \sigma_{ij}$ . Recall  $\mathbf{R} = (r_{ij})_{p \times p}$  with  $r_{ij} = \sigma_{ij}/\sqrt{\sigma_{ii}\sigma_{jj}}$ . Then  $Cov(X_1/\sqrt{\sigma_{ii}}, Y_1/\sqrt{\sigma_{jj}}) = r_{ij}$ . By Lemma 6.12, we have

$$\frac{m^2}{n^2} \cdot E\left(\frac{\sigma_{ii}}{s_{ii}} \cdot \frac{\sigma_{ii}}{s_{jj}}\right) = 1 + \frac{4 + 2r_{ij}^2}{m} + \frac{12 + 8r_{ij}^2 + 8r_{ij}^4}{m^2} + \frac{\delta_m(i,j)}{m^3},$$

where  $\max_{1 \le i,j \le p} |\delta_m(i,j)| \le C$  for all  $m \ge 11$ , where C is a constant not depending on m or  $\mathbf{R} = (r_{ij})$ . This and (173) conclude that  $\operatorname{Cov}(\frac{\sigma_{ii}}{s_{ii}}, \frac{\sigma_{jj}}{s_{jj}})$  is identical to

$$\left[ 1 + \frac{4 + 2r_{ij}^2}{m} + \frac{12 + 8r_{ij}^2 + 8r_{ij}^4}{m^2} + \frac{\delta_m(i,j)}{m^3} \right] \cdot \left(\frac{m+1}{m}\right)^2 - \frac{m+1}{m-2} \cdot \frac{m+1}{m-2}$$

$$= \left[ 1 + \frac{4 + 2r_{ij}^2}{m} + \frac{\delta_m(i,j)'}{m^2} \right] \cdot \left( 1 + \frac{2}{m} + \frac{1}{m^2} \right) - \left[ 1 + \frac{6}{m} + \frac{21m - 24}{m(m-2)^2} \right]$$

$$= \frac{2r_{ij}^2}{m} + \frac{\delta_m(i,j)''}{m^2},$$

$$(180)$$

where

$$\max_{1 \le i,j \le p} \left\{ |\delta_m(i,j)'|, |\delta_m(i,j)''| \right\} \le K_1$$
(181)

for all  $m \ge 11$  and  $K_1$  here and later represents a constant not depending on m or  $r_{ij}$ , and can be different from line to line. This, (178) and (179) conclude

$$E \operatorname{tr}(\mathbf{B}^{2}) = \frac{2}{m} \Big( \sum_{1 \le i, j \le p} r_{ij}^{4} \Big) + \sum_{1 \le i, j \le p} r_{ij}^{2} \Big[ \frac{\delta_{m}(i, j)'}{m^{2}} + \frac{9}{(m-2)^{2}} \Big]$$
  
$$\leq \frac{2}{m} \operatorname{tr}(\mathbf{R}^{2}) + \frac{K_{1}}{m^{2}} \operatorname{tr}(\mathbf{R}^{2})$$
  
$$\leq \frac{3}{m} \operatorname{tr}(\mathbf{R}^{2})$$
(182)

as *m* is sufficiently large, which is guaranteed as  $p \to \infty$  since  $\lim_{p\to\infty} n_p = \infty$  and  $m = n_p - 1$ . In the second step above we use the fact  $\sum_{1 \le i,j \le p} r_{ij}^4 \le \sum_{1 \le i,j \le p} r_{ij}^2 = \operatorname{tr}(\mathbf{R}^2)$ .

Step 4: the evaluation of  $Var(tr(\mathbf{B}))$  from (177). Note

$$\operatorname{tr}(\mathbf{B}) = \operatorname{tr}\left(\boldsymbol{\Sigma}^{1/2}(\hat{\mathbf{D}}^{-1} - \mathbf{D}^{-1})\boldsymbol{\Sigma}^{1/2}\right) = \operatorname{tr}\left((\hat{\mathbf{D}}^{-1} - \mathbf{D}^{-1})\boldsymbol{\Sigma}\right)$$
$$= \sum_{i=1}^{m} \left(\frac{1}{s_{ii}} - \frac{1}{\sigma_{ii}}\right)\sigma_{ii}.$$

Recall  $\frac{ns_{ii}}{\sigma_{ii}} \sim \chi^2(m)$  for each *i*. It then follows from (173) and (181) that

$$\operatorname{Var}(\operatorname{tr}(\mathbf{B})) = \operatorname{Var}\left(\sum_{i=1}^{m} \frac{\sigma_{ii}}{s_{ii}}\right)$$
$$= \sum_{i=1}^{m} \operatorname{Var}\left(\frac{\sigma_{ii}}{s_{ii}}\right) + 2\sum_{1 \le i < j \le p} \operatorname{Cov}\left(\frac{\sigma_{ii}}{s_{ii}}, \frac{\sigma_{jj}}{s_{jj}}\right)$$
$$\le m \cdot \frac{2n^2}{(m-2)^2(m-4)} + 2\sum_{1 \le i < j \le p} \left[\frac{2r_{ij}^2}{m} + \frac{\delta_m(i,j)''}{m^2}\right].$$

Thus,

$$\operatorname{Var}(\operatorname{tr}(\mathbf{B})) \le 3 + \frac{2}{m}\operatorname{tr}(\mathbf{R}^2) + \frac{K_1 p^2}{m^2}$$

as m is sufficiently large.

Finally, combining the analysis of the two terms from (177) in *Step 3* and *Step 4*, we eventually obtain

$$\operatorname{Var}(\boldsymbol{\theta}^{T} \mathbf{B} \boldsymbol{\theta}) \leq \frac{6}{m} \operatorname{tr}(\mathbf{R}^{2}) + 3 + \frac{2}{m} \operatorname{tr}(\mathbf{R}^{2}) + K_{1} \frac{p^{2}}{m^{2}}$$
$$= 3 + \frac{8}{m} \operatorname{tr}(\mathbf{R}^{2}) + K_{1} \frac{p^{2}}{m^{2}}$$

as p is sufficiently large. Easily,  $tr(\mathbf{R}^2) \ge p$ . It follows that

$$\operatorname{Var}\left(\frac{\boldsymbol{\theta}^{T}\mathbf{B}\boldsymbol{\theta}}{\sqrt{2\operatorname{tr}(\mathbf{R}^{2})}}\right) = \frac{\operatorname{Var}(\boldsymbol{\theta}^{T}\mathbf{B}\boldsymbol{\theta})}{2\operatorname{tr}(\mathbf{R}^{2})} \leq \frac{3}{p} + \frac{4}{m} + K_{1}\frac{p^{2}}{m^{2}\operatorname{tr}(\mathbf{R}^{2})} \to 0$$

since  $\lim_{p\to\infty} \frac{p}{m \|\mathbf{R}\|_F} = 0$  and  $\|\mathbf{R}\|_F^2 = \operatorname{tr}(\mathbf{R}^2)$ . This joined (175) concludes (166).

**Proof of Lemma 4.2.** First, by the monotone property of  $a_{p,i}$ , we obtain  $\rho_1 \ge \rho_2 \ge \cdots$ . Moreover,  $1 = a_{p,1}^2 + \cdots + a_{p,p}^2 \ge a_{p,1}^2 + \cdots + a_{p,i}^2 \ge ia_{p,i}^2$  for any  $1 \le i \le p$ . This implies that  $0 \le a_{p,i} \le i^{-1/2}$  for each  $1 \le i \le p$ . Take  $p \to \infty$  to obtain  $0 \le \rho_i \le i^{-1/2}$  for each  $i \ge 1$ . Also, by using the fact  $1 \ge a_{p,1}^2 + \cdots + a_{p,i}^2$  for any  $1 \le i \le p$ , and letting  $p \to \infty$  first and then  $i \to \infty$ , we get  $\sum_{i=1}^{\infty} \rho_i^2 \le 1$ .

We first handle a trivial case:  $\rho_1 = 0$ . By monotonicity,  $\rho_i = 0$  for each  $i \ge 1$ . Notice  $E\xi_1 = 0$  and  $\operatorname{Var}(\xi_1) = 2$ . Thus,  $s_n^2 := \operatorname{Var}(a_{p,1}\xi_1 + \cdots + a_{p,p}\xi_p) = 2(a_{p,1}^2 + \cdots + a_{p,p}^2) = 2$ . Easily,

$$\frac{1}{s_n^3} \sum_{i=1}^p E(|a_{p,i}\xi_i|^3) = \frac{E(|\xi_1|^3)}{2\sqrt{2}} \sum_{i=1}^p a_{p,i}^3 \le \frac{E(|\xi_1|^3)}{2\sqrt{2}} \cdot a_{p,1} \cdot \sum_{i=1}^p a_{p,i}^2,$$

which goes to zero by the assumption  $\lim_{p\to\infty} a_{p,1} = \rho_1 = 0$ . The desired result follows from the Lyapunov central limit theorem. From now on, we assume  $\rho_1 > 0$ .

In the following a useful fact will be derived first. For each  $p \ge 1$ , let  $b_{p,1} \ge b_{p,2} \ge \cdots \ge 0$ be constants satisfying  $\sum_{i=1}^{\infty} b_{p,i}^2 \le 1$ . We claim that

$$\prod_{i=m}^{\infty} \left[ e^{-tb_{p,i}} \left( 1 - 2tb_{p,i} \right)^{-1/2} \right] = e^{\gamma_{p,m}} \cdot \exp\left( t^2 \sum_{i=m}^{\infty} b_{p,i}^2 \right)$$
(183)

for all  $m \ge 16$  and |t| < 1, where  $\sup_{p\ge 1} |\gamma_{p,m}| \le \frac{8}{\sqrt{m}}$ . In fact, write  $\log(1-x) = -\sum_{i=1}^{\infty} \frac{1}{i} x^i := -x - \frac{1}{2} x^2 - B(x)$  for |x| < 1. Then

$$|B(x)| \le \sum_{i=3}^{\infty} \frac{1}{i} |x|^i \le \sum_{i=3}^{\infty} |x|^i \le \frac{|x|^3}{1-|x|} \le 2|x|^3$$
(184)

if  $|x| \leq \frac{1}{2}$ . By the same argument as that in the beginning, we know  $0 \leq b_{p,i} \leq \frac{1}{\sqrt{i}}$  for each  $i \geq 1$  and  $p \geq 1$ . Observe

$$\prod_{i=m}^{\infty} \left[ e^{-tb_{p,i}} \left( 1 - 2tb_{p,i} \right)^{-1/2} \right] = \prod_{i=m}^{\infty} \exp\left[ -tb_{p,i} - \frac{1}{2} \log\left( 1 - 2tb_{p,i} \right) \right]$$
$$= \prod_{i=m}^{\infty} \exp\left[ t^2 b_{p,i}^2 + \frac{1}{2} B(2tb_{p,i}) \right].$$

By the monotone property,  $\max_{i\geq m} |2tb_{p,i}| = 2|t|b_{p,m} \leq \frac{2|t|}{\sqrt{m}} \leq \frac{1}{2}|t|$  for  $m \geq 16$ . This and (184) say that

$$\sum_{i=m}^{\infty} \frac{1}{2} |B(2tb_{p,i})| \le 8|t|^3 \sum_{i=m}^{\infty} b_{p,i}^3 \le \frac{8|t|^3}{\sqrt{m}} \sum_{i=m}^{\infty} b_{p,i}^2 \le \frac{8}{\sqrt{m}}$$

for any t with |t| < 1 and  $p \ge 1$ . These lead to (183). In two steps next we will apply (183) to  $a_{p,1}\xi_1 + \cdots + a_{p,p}\xi_p$  and its limit stated in the lemma, respectively. The limit case goes first.

Step 1. Set  $b = [2(1 - \sum_{i=1}^{\infty} \rho_i^2)]^{1/2}$  and  $X = b\eta + \sum_{i=1}^{\infty} \rho_i \xi_i$ , where  $\eta \sim N(0, 1)$  and  $\eta$  is independent of  $\{\xi_i; i \ge 1\}$ . Then, by independence and the fact  $E \exp(t\chi^2(1)) = (1 - 2t)^{-1/2}$  for  $t < \frac{1}{2}$ , we see

$$Ee^{tX} = \left(\prod_{i=1}^{\infty} Ee^{t\rho_i\xi_1}\right) \cdot Ee^{tb\eta} = e^{b^2t^2/2} \cdot \prod_{i=1}^{\infty} \left[e^{-t\rho_i} \left(1 - 2t\rho_i\right)^{-1/2}\right]$$
(185)

for t with  $|t\rho_i| < \frac{1}{2}$  for each  $i \ge 1$ , which holds as  $|t| < \frac{1}{2\rho_1}$ . Take  $b_{p,i} = \rho_i$  for all  $i \ge 1$  and  $p \ge 1$  in (183) to see

$$\prod_{i=m}^{\infty} \left[ e^{-t\rho_i} \left( 1 - 2t\rho_i \right)^{-1/2} \right] = e^{\gamma_m} \cdot \exp\left( t^2 \sum_{i=m}^{\infty} \rho_i^2 \right)$$
(186)

for  $m \ge 16$  and |t| < 1, where  $|\gamma_m| \le \frac{8}{\sqrt{m}}$ . This and (185) especially indicate  $Ee^{tX} < \infty$  for every  $|t| < \frac{1}{2}$ . Recall  $\sum_{i=1}^{\infty} \rho_i^2 \le 1$ , by sending  $m \to \infty$  we see the left hand side of (186) goes to 1. Therefore,

$$\prod_{i=1}^{m-1} \left[ e^{-t\rho_i} (1 - 2t\rho_i)^{-1/2} \right] \to \prod_{i=1}^{\infty} \left[ e^{-t\rho_i} (1 - 2t\rho_i)^{-1/2} \right]$$
(187)

as  $m \to \infty$  for every  $|t| < \frac{1}{2}$ .

Step 2. Evidently,

$$E^{t(a_{p,1}\xi_{1}+\dots+a_{p,p}\xi_{p})} = \prod_{i=1}^{p} Ee^{ta_{p,i}\xi_{1}} = \prod_{i=1}^{p} \left[ e^{-ta_{p,i}} \left( 1 - 2ta_{p,i} \right)^{-1/2} \right]$$
(188)

provided  $|t| < \frac{1}{2a_{p,1}}$ . In particular, this holds if  $|t| < \frac{1}{2}$ . Now, by taking  $b_{p,i} = a_{p,i}$  for  $1 \le i \le p$  and  $b_{p,i} = 0$  for i > p from (183), we obtain

$$E^{t(a_{p,1}\xi_1 + \dots + a_{p,p}\xi_p)} = e^{\gamma_{p,m}} \cdot \exp\left(t^2 \sum_{i=m}^p a_{p,i}^2\right) \cdot \prod_{i=1}^{m-1} \left[e^{-ta_{p,i}} \left(1 - 2ta_{p,i}\right)^{-1/2}\right]$$

for any *m* with  $16 \leq m \leq p$  and  $|t| < \frac{1}{2}$ , where  $\sup_{p\geq 1} |\gamma_{p,m}| \leq \frac{8}{\sqrt{m}}$ . Consequently, if  $16 \leq m \leq p$  and  $|t| < \frac{1}{2}$  then

$$E^{t(a_{p,1}\xi_1 + \dots + a_{p,p}\xi_p)} \le e^{8/\sqrt{m}} \cdot \exp\left[t^2 \left(1 - \sum_{i=1}^{m-1} a_{p,i}^2\right)\right] \cdot \prod_{i=1}^{m-1} \left[e^{-ta_{p,i}} \left(1 - 2ta_{p,i}\right)^{-1/2}\right]$$
(189)

by the assumption  $a_{p,1}^2 + \cdots + a_{p,p}^2 = 1$ , and

$$E^{t(a_{p,1}\xi_1 + \dots + a_{p,p}\xi_p)} \ge e^{-8/\sqrt{m}} \cdot \exp\left[t^2 \left(1 - \sum_{i=1}^{m-1} a_{p,i}^2\right)\right] \cdot \prod_{i=1}^{m-1} \left[e^{-ta_{p,i}} \left(1 - 2ta_{p,i}\right)^{-1/2}\right].$$
 (190)

With the two steps established above, we are now ready to complete the proof. Recall the assumption  $\lim_{p\to\infty} a_{p,i} = \rho_i$  for each  $i \ge 1$ . For fixed  $m \ge 16$  we send  $p \to \infty$  and then send  $m \to \infty$  in (189) and (190), we have from (187) and then (185) that

$$E^{t(a_{p,1}\xi_{1}+\dots+a_{p,p}\xi_{p})} \to \exp\left[t^{2}\left(1-\sum_{i=1}^{\infty}\rho_{i}^{2}\right)\right] \cdot \prod_{i=1}^{\infty}\left[e^{-t\rho_{i}}\left(1-2t\rho_{i}\right)^{-1/2}\right] = Ee^{tX}$$

as  $p \to \infty$  for  $|t| < \frac{1}{2}$ . The desired conclusion then follows from the uniqueness of the moment generating function.

Recall F(1, m) stands for the F-distribution with degrees of freedoms 1 and m.

**LEMMA 6.36** Let  $m = m_p \to \infty$  as  $p \to \infty$ . For each  $p \ge 1$ , let  $X_{p,1}, \dots, X_{p,p}$  be i.i.d. with distribution F(1,m). Then  $(2p)^{-1/2}[X_{p,1} + \dots + X_{p,p} - mp(m-2)^{-1}] \to N(0,1)$  in distribution as  $p \to \infty$ .

**Proof of Lemma 6.36**. First, by the property of *F*-distribution,

$$EX_{p,1} = \frac{m}{m-2}$$
 and  $Var(X_{p,1}) = \frac{2m^2(m-1)}{(m-2)^2(m-4)}$ 

for  $m \geq 5$ . By definition, we write  $X_{p,1} = \frac{m\xi_0^2}{\xi_1^2 + \dots + \xi_m^2}$ , where  $\xi_0, \xi_1, \dots, \xi_m$  are i.i.d. N(0, 1). Then

$$E(X_{p,1} - EX_{p,1})^{4} = E\left[\left(\frac{m}{\xi_{1}^{2} + \dots + \xi_{m}^{2}} - 1\right)\xi_{0}^{2} + \xi_{0}^{2} - \frac{m}{m-2}\right]^{4}$$
  
$$\leq 3^{3}E\left[\left(\frac{m}{\xi_{1}^{2} + \dots + \xi_{m}^{2}} - 1\right)^{4}\xi_{0}^{8}\right] + 3^{3}E(\xi_{0}^{8}) + 3^{3}\left(\frac{m}{m-2}\right)^{4}.$$

By using the Cauchy-Schwartz inequality twice,

$$E\left[\left(\frac{m}{\xi_{1}^{2}+\dots+\xi_{m}^{2}}-1\right)^{4}\xi_{0}^{8}\right] \leq \left\{E\left[\left(\frac{\xi_{1}^{2}+\dots+\xi_{m}^{2}-m}{\xi_{1}^{2}+\dots+\xi_{m}^{2}}\right)^{8}\right]\right\}^{1/2} \cdot \left(E\xi_{0}^{16}\right)^{1/2} \\ \leq K \cdot \left[E\left(\xi_{1}^{2}+\dots+\xi_{m}^{2}-m\right)^{16}\right]^{1/4} \cdot \left[E\frac{1}{(\xi_{1}^{2}+\dots+\xi_{m}^{2})^{16}}\right]^{1/4},$$

where K here and later is a constant free of m and p, and can be different from line to line. By using the Marcinkiewicz-Zygmund inequality [see, for example, the proof of Corollary 2 on p. 387 from Chow and Teicher (1997)],  $E(\xi_1^2 + \cdots + \xi_m^2 - m)^{16} \leq Km^8$ . Furthermore, take  $\beta = -16$  in (57) to see  $E[(\xi_1^2 + \cdots + \xi_m^2)^{-16}] \leq Km^{-16}$  for all  $m \geq 34$ . Combining all of the above calculation, we see  $E(X_{p,1} - EX_{p,1})^4 \leq K$  as  $m \geq 34$ . Notice  $Var(X_{p,1}) \rightarrow 2$  as  $p \rightarrow \infty$ . Then

$$\frac{1}{(p \operatorname{Var}(X_{p,1}))^2} \sum_{i=1}^p E(X_{p,i} - EX_{p,i})^4 = O\left(\frac{1}{p}\right) \to 0$$

as  $p \to \infty$ . By the Lyapunov CLT, we obtain the desired result.

Proof of Theorem 1. First, by Theorem 3,

$$\frac{1}{\operatorname{tr}(\mathbf{R}^2)} \left[ \operatorname{tr}(\hat{\mathbf{R}}^2) - \frac{p(p-1)}{n-1} \right] \to 1$$
(191)

in probability as  $p \to \infty$ . In the following we will use this fact twice to show  $T_{SD}$  and  $T_{p,1}$  are equivalent. First, it follows from the assumption  $\lim_{p\to\infty} \frac{p}{n \|\mathbf{R}\|_F} = 0$  that

$$\frac{\operatorname{tr}(\hat{\mathbf{R}}^2) - p^2(n-1)^{-1}}{\operatorname{tr}(\mathbf{R}^2)} = \frac{\operatorname{tr}(\hat{\mathbf{R}}^2) - p(p-1)(n-1)^{-1}}{\operatorname{tr}(\mathbf{R}^2)} - \frac{p(n-1)^{-1}}{\operatorname{tr}(\mathbf{R}^2)} = 1 + o_p(1).$$

As a consequence,

$$H_p := \left[\frac{\operatorname{tr}(\hat{\mathbf{R}}^2) - p^2(n-1)^{-1}}{\operatorname{tr}(\mathbf{R}^2)}\right]^{-1/2} = 1 + o_p(1).$$

Review (2). We have

$$T_{SD} = \frac{[n\bar{\boldsymbol{X}}^T\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} - pn(n-3)^{-1}] + p(n-3)^{-1}}{\sqrt{2\mathrm{tr}(\mathbf{R}^2)}} \cdot H_p$$
  
=  $\frac{[n\bar{\boldsymbol{X}}^T\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} - pn(n-3)^{-1}]}{\sqrt{2\mathrm{tr}(\mathbf{R}^2)}} \cdot [1 + o_p(1)] + o_p(1)$  (192)

by the assumption  $\frac{p}{n \|\mathbf{R}\|_F} \to 0$  and the notation  $\|\mathbf{R}\|_F^2 = \operatorname{tr}(\mathbf{R}^2)$ . By (9),

$$T_{p,1} = \frac{n\bar{\mathbf{X}}^T\hat{\mathbf{D}}^{-1}\bar{\mathbf{X}} - pn(n-3)^{-1}}{\sqrt{2\left|\operatorname{tr}(\hat{\mathbf{R}}^2) - p(p-1)(n-1)^{-1}\right|}}.$$

It follows from (191) that

$$T_{p,1} = \frac{[n\bar{\boldsymbol{X}}^T\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} - pn(n-3)^{-1}]}{\sqrt{2\mathrm{tr}(\mathbf{R}^2)}} \cdot [1 + o_p(1)].$$

Comparing this with (192), we obtain

$$T_{SD} = T_{p,1} \cdot [1 + o_p(1)] + o_p(1).$$
(193)

So to finish the proof, by using the Slutsky lemma, it suffices to prove that  $T_{p,1} \to (1 - \sum_{i=1}^{\infty} \rho_i^2)^{1/2} \xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i(\xi_i^2 - 1)$  in distribution.

Set m = n - 1. Then

$$\sqrt{n}\bar{\boldsymbol{X}} \sim N_p(\boldsymbol{0}, \boldsymbol{\Sigma}), \ n\hat{\boldsymbol{S}} \sim W_p(m, \boldsymbol{\Sigma}), \ \text{and} \ \bar{\boldsymbol{X}} \ \text{and} \ \hat{\boldsymbol{S}} \ \text{are independent},$$
(194)

where  $W_p(m, \Sigma)$  is the Wishart distribution defined after (165); see, for example, Theorem 3.1.2 from Muirhead (1982). This implies

$$\hat{\mathbf{D}} \stackrel{d}{=}$$
 the diagonal matrix of  $\frac{1}{n} W_p(m, \boldsymbol{\Sigma})$ . (195)

In particular,  $\bar{\boldsymbol{X}}$  is independent of  $\hat{\mathbf{D}}$ . By Lemma 4.1 and assumption  $\lim_{p\to\infty} \frac{p}{m\|\mathbf{R}\|_F} = 0$ ,

$$\frac{n\bar{\boldsymbol{X}}^T\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}}}{\sqrt{2\operatorname{tr}(\mathbf{R}^2)}} = \frac{n\bar{\boldsymbol{X}}^T\mathbf{D}^{-1}\bar{\boldsymbol{X}}}{\sqrt{2\operatorname{tr}(\mathbf{R}^2)}} + o_p(1)$$

as  $p \to \infty$ , where **D** is the diagonal matrix of the population covariance matrix  $\Sigma$ . Now we will analyze the behavior of  $n\bar{X}^T \mathbf{D}^{-1}\bar{X}$  in two steps. Once they are established, the limiting distribution of  $T_{p,1}$  will be identified quickly.

Step 1. The exact distribution of  $n\bar{\boldsymbol{X}}^T \mathbf{D}^{-1}\bar{\boldsymbol{X}}$ . By assumption,  $\lambda_1, \dots, \lambda_p$  are the eigenvalues of  $\mathbf{R} = \mathbf{D}^{-1/2} \boldsymbol{\Sigma} \mathbf{D}^{-1/2}$ . We claim that

$$n\bar{\boldsymbol{X}}^{T}\mathbf{D}^{-1}\bar{\boldsymbol{X}} \stackrel{d}{=} \sum_{i=1}^{p} \lambda_{i}\xi_{i}^{2}, \qquad (196)$$

where  $\xi_1, \dots, \xi_p$  are i.i.d. N(0, 1) and where " $\stackrel{d}{=}$ " means both sides of "=" have the same distribution. We show (196) next.
Since  $\sqrt{n}\bar{\boldsymbol{X}} \sim N_p(\boldsymbol{0}, \boldsymbol{\Sigma})$ , we are able to write  $\sqrt{n}\bar{\boldsymbol{X}} = \boldsymbol{\Sigma}^{1/2}\boldsymbol{\xi}$ , where  $\boldsymbol{\Sigma}^{1/2}$  is a non-negative definite matrix satisfying  $\boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{1/2} = \boldsymbol{\Sigma}$  and  $\boldsymbol{\xi} \sim N_p(\boldsymbol{0}, \mathbf{I}_p)$ . It follows that

$$n\bar{\boldsymbol{X}}^{T}\boldsymbol{\mathrm{D}}^{-1}\bar{\boldsymbol{X}} = \boldsymbol{\xi}^{T} \big(\boldsymbol{\Sigma}^{1/2}\boldsymbol{\mathrm{D}}^{-1}\boldsymbol{\Sigma}^{1/2}\big)\boldsymbol{\xi}.$$
(197)

Since **AB** and **BA** have the same eigenvalues for any square matrix **A** and **B**, it is easy to see that  $\lambda_1, \dots, \lambda_p$  are also the eigenvalues of  $\Sigma^{1/2} \mathbf{D}^{-1} \Sigma^{1/2}$ . Write  $\Sigma^{1/2} \mathbf{D}^{-1} \Sigma^{1/2} = \mathbf{O}^T \operatorname{diag}(\lambda_1, \dots, \lambda_p) \mathbf{O}$  for some orthogonal matrix **O**. Then by the orthogonal invariance property, we know  $\mathbf{O}\boldsymbol{\xi} \stackrel{d}{=} \boldsymbol{\xi}$ . Consequently,

$$\boldsymbol{\xi}^{T} \left( \boldsymbol{\Sigma}^{1/2} \mathbf{D}^{-1} \boldsymbol{\Sigma}^{1/2} \right) \boldsymbol{\xi} = (\mathbf{O} \boldsymbol{\xi})^{T} \operatorname{diag}(\lambda_{1}, \cdots, \lambda_{p}) (\mathbf{O} \boldsymbol{\xi}) \stackrel{d}{=} \sum_{i=1}^{p} \lambda_{i} \xi_{i}^{2}$$
(198)

where  $\xi_1, \dots, \xi_p$  are i.i.d. N(0, 1). We then get (196).

Step 2: the limiting distribution of  $n\bar{\mathbf{X}}^T\mathbf{D}^{-1}\bar{\mathbf{X}}$ . By (196) and the fact  $\operatorname{tr}(\mathbf{R}) = \lambda_1 + \cdots + \lambda_p = p$ , we know

$$\frac{n\bar{\boldsymbol{X}}^T\mathbf{D}^{-1}\bar{\boldsymbol{X}}-p}{\sqrt{2\operatorname{tr}(\mathbf{R}^2)}} \stackrel{d}{=} \frac{1}{\sqrt{2}}\sum_{i=1}^p \frac{\lambda_i}{\|\mathbf{R}\|_F} (\xi_i^2-1).$$

Set  $a_{p,i} = \frac{\lambda_i}{\|\mathbf{R}\|_F}$  for  $1 \le i \le p$ . Then  $a_{p,1} \ge \cdots \ge a_{p,p} \ge 0$  and  $a_{p,1}^2 + \cdots + a_{p,p}^2 = 1$ . By assumption (a),  $\lim_{p\to\infty} a_{p,i} = \rho_i \ge 0$  for each  $i \ge 1$ . From Lemma 4.2,

$$\frac{1}{\sqrt{2}} \sum_{i=1}^{p} \frac{\lambda_i}{\|\mathbf{R}\|_F} (\xi_i^2 - 1) \to \left(1 - \sum_{i=1}^{\infty} \rho_i^2\right)^{1/2} \xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i (\xi_i^2 - 1)$$

in distribution as  $p \to \infty$ , where  $\xi_0 \sim N(0, 1)$  and  $\xi_0$  is independent of  $\{\xi_i; i \ge 1\}$ . Therefore,

$$\frac{n\bar{\boldsymbol{X}}^{T}\mathbf{D}^{-1}\bar{\boldsymbol{X}}-p}{\sqrt{2\operatorname{tr}(\mathbf{R}^{2})}} \to \left(1-\sum_{i=1}^{\infty}\rho_{i}^{2}\right)^{1/2}\xi_{0}+\frac{1}{\sqrt{2}}\sum_{i=1}^{\infty}\rho_{i}(\xi_{i}^{2}-1)$$
(199)

in distribution.

With *Step 1* and *Step 2* completed, let us now proceed to finish the proof. In fact, by assumption (b),

$$\frac{1}{\sqrt{2}\|\mathbf{R}\|_F} \left[ p - pn(n-3)^{-1} \right] = -\frac{3p}{\sqrt{2}(n-3)\|\mathbf{R}\|_F} \to 0.$$

Summing this and (199), we see from the Slutsky lemma that

$$\frac{n\bar{\boldsymbol{X}}^{T}\boldsymbol{\mathrm{D}}^{-1}\bar{\boldsymbol{X}} - pn(n-3)^{-1}}{\sqrt{2}\|\boldsymbol{\mathrm{R}}\|_{F}} \to \left(1 - \sum_{i=1}^{\infty}\rho_{i}^{2}\right)^{1/2}\xi_{0} + \frac{1}{\sqrt{2}}\sum_{i=1}^{\infty}\rho_{i}(\xi_{i}^{2} - 1)$$

in distribution. Combine this with (191) to see

$$T_{p,1}$$
,  $\rightarrow \left(1 - \sum_{i=1}^{\infty} \rho_i^2\right)^{1/2} \xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i(\xi_i^2 - 1)$ 

in distribution as  $p \to \infty$ . The proof is completed.

Now we begin to prove Proposition 1. As far as the proofs go, parts (i) and (ii) from the lemma have different natures, to make the presentation clear, we will handle the two parts separately.

**Proof of Proposition 1(i).** Assume  $\boldsymbol{\mu} = 0$ . Then  $\mathbf{X}_1, \dots, \mathbf{X}_n$  are i.i.d. with distribution  $N_p(\mathbf{0}, \boldsymbol{\Sigma})$ , where all of the  $p^2$  entries of  $\boldsymbol{\Sigma}$  are identical to  $\sigma^2$ . For this reason, we write  $\boldsymbol{X}_i = \xi_i (1, \dots, 1)^T \in \mathbb{R}^p$  for  $i = 1, \dots, n$ , where  $\xi_1, \dots, \xi_n$  are i.i.d.  $N(0, \sigma^2)$ . By (1) and (7),  $\bar{\boldsymbol{X}} = \frac{1}{n} \sum_{i=1}^n \boldsymbol{X}_i$ ,

$$\hat{\mathbf{S}} = \frac{1}{n} \sum_{i=1}^{n} (\mathbf{X}_i - \bar{\mathbf{X}}) (\mathbf{X}_i - \bar{\mathbf{X}})^T$$
 and  $\hat{\mathbf{R}} = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2}$ ,

where  $\hat{\mathbf{D}}$  is the diagonal matrix of  $\hat{\mathbf{S}}$ . Set  $\bar{\xi} = \frac{1}{n} \sum_{i=1}^{n} \xi_i$  and  $W = \frac{1}{n} \sum_{i=1}^{n} (\xi_i - \bar{\xi})^2$ . Then  $\sqrt{n}\bar{\xi}/\sigma \sim N(0,1)$ ,  $nW/\sigma^2 \sim \chi^2(n-1)$ , and  $\bar{\xi}$  and W are independent. Since  $\bar{\mathbf{X}} = \bar{\xi}(1,\cdots,1)^T \in \mathbb{R}^p$ , we see  $\mathbf{X}_i - \bar{\mathbf{X}} = (\xi_i - \bar{\xi})^T (1,\cdots,1)^T \in \mathbb{R}^p$ . Hence,  $\hat{\mathbf{S}} = W \cdot \mathbf{J}$ , where  $\mathbf{J}$  is a  $p \times p$  matrix whose entries are all equal to 1. It follows  $\hat{\mathbf{D}} = W \cdot \mathbf{I}_p$  and  $\hat{\mathbf{R}} = \mathbf{J}$ . In particular,  $\operatorname{tr}(\hat{\mathbf{R}}^2) = p^2$ . Consequently,

$$n\bar{\boldsymbol{X}}^{T}\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} = \frac{np}{n-1} \cdot \frac{(\sqrt{n}\bar{\xi}/\sigma)^{2}}{nW/[(n-1)\sigma^{2}]} \stackrel{d}{=} \frac{np}{n-1} \cdot F_{1,n-1}.$$
 (200)

By (2),

$$T_{SD} = \frac{n\bar{\boldsymbol{X}}^T\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} - p(n-1)(n-3)^{-1}}{\sqrt{2[\operatorname{tr}(\hat{\mathbf{R}}^2) - p^2(n-1)^{-1}]}} \stackrel{d}{=} \frac{\frac{n}{n-1}\cdot F_{1,n-1} - \frac{n-1}{n-3}}{\sqrt{2[1-(n-1)^{-1}]}}.$$
 (201)

Notice  $F_{1,n-1} \to \chi^2(1)$  as  $n \to \infty$ . We see from the Slutsky lemma that  $T_{SD} \to \frac{1}{\sqrt{2}} \cdot [\chi^2(1) - 1]$ as  $p \to \infty$ . Finally, by (9) and (200),

$$T_{p,1} = \frac{n\bar{\mathbf{X}}^T\hat{\mathbf{D}}^{-1}\bar{\mathbf{X}} - pn(n-3)^{-1}}{\sqrt{2\left|\operatorname{tr}(\hat{\mathbf{R}}^2) - p(p-1)(n-1)^{-1}\right|}} \stackrel{d}{=} \frac{\frac{np}{n-1}\cdot F_{1,n-1} - pn(n-3)^{-1}}{\sqrt{2\left|p^2 - p(p-1)(n-1)^{-1}\right|}}.$$

As a consequence,

$$T_{p,1} \stackrel{d}{=} \frac{\frac{n}{n-1} \cdot F_{1,n-1} - n(n-3)^{-1}}{\sqrt{2\{1 - (p-1)/[(n-1)p]\}}} \to \frac{1}{\sqrt{2}} \cdot \left[\chi^2(1) - 1\right].$$

The proof is completed.

## **Proof of Proposition 1(ii)**. Recall

$$T_{SD} = \frac{n\bar{\boldsymbol{X}}^T\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} - p(n-1)(n-3)^{-1}}{\sqrt{2[\operatorname{tr}(\hat{\mathbf{R}}^2) - p^2(n-1)^{-1}]}}$$

Since  $T_{SD}$  is scale-invariant, without loss of generality, we assume  $\Sigma = \mathbf{I}_p$ . By (194),

 $\sqrt{n}\bar{\boldsymbol{X}} \sim N_p(\boldsymbol{0}, \mathbf{I}_p), \ n\hat{\mathbf{S}} \sim W_p(m, \mathbf{I}_p), \ \text{and} \ \bar{\boldsymbol{X}} \ \text{and} \ \hat{\mathbf{S}} \ \text{are independent},$ 

where m := n - 1 and  $W_p(m, \Sigma)$  is the Wishart distribution defined after (165). Also, by (195),  $\hat{\mathbf{D}}$  and the diagonal matrix of  $\frac{1}{n}W_p(m, \mathbf{I}_p)$  have the same distribution. By definition of  $W_p(m, \mathbf{I}_p)$ , its p diagonal entries are i.i.d.  $\chi^2(m)$ . Therefore,  $n\bar{\mathbf{X}}^T\hat{\mathbf{D}}^{-1}\bar{\mathbf{X}}$  is a sum of pi.i.d. random variables with distribution  $n \cdot \frac{N(0,1)^2}{\chi^2(m)}$ , where the numerator and denominator are independent. Thus, we are able to write

$$n\bar{\boldsymbol{X}}^{T}\hat{\boldsymbol{D}}^{-1}\bar{\boldsymbol{X}} = \frac{n}{m}\sum_{i=1}^{p}X_{p,i},$$
(202)

where  $X_{p,1}, \dots, X_{p,p}$  are i.i.d. with distribution F(1,m). By Lemma 6.36,

$$U_p := \frac{1}{\sqrt{2p}} \left( \sum_{i=1}^p X_{p,i} - \frac{mp}{m-2} \right) \to N(0,1)$$
(203)

as  $p \to \infty$ , regardless of the speeds of n and p going to infinity. Solve  $\sum_{i=1}^{p} X_{p,i}$  in terms of  $U_p$  to see  $\sum_{i=1}^{p} X_{p,i} = \sqrt{2p}U_p + \frac{mp}{m-2}$ . By plugging this into (202), we obtain

$$n\bar{\boldsymbol{X}}^{T}\hat{\boldsymbol{D}}^{-1}\bar{\boldsymbol{X}} = \frac{n}{m}\sqrt{2p}U_{p} + \frac{np}{m-2}.$$
(204)

г		
н		L
ь.		

Consequently,

$$n\bar{\boldsymbol{X}}^{T}\hat{\mathbf{D}}^{-1}\bar{\boldsymbol{X}} - p(n-1)(n-3)^{-1} = \frac{n}{m}\sqrt{2p}U_{p} + \frac{np}{m-2} - \frac{mp}{m-2}$$
$$= \sqrt{2p}U_{p} + \frac{\sqrt{2p}}{m}U_{p} + \frac{p}{m-2}.$$

It follows that

$$\frac{1}{\sqrt{2p}} \left[ n\bar{\boldsymbol{X}}^T \hat{\mathbf{D}}^{-1} \bar{\boldsymbol{X}} - p(n-1)(n-3)^{-1} \right] = U_p + \frac{1}{m} U_p + \frac{\sqrt{p}}{\sqrt{2}(m-2)}.$$
(205)

On the other hand,  $\hat{\mathbf{R}} = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2}$  by (7). Write  $n\hat{\mathbf{S}} = (\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_p)^T (\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_p)$ , where  $\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_p$  are i.i.d.  $N_m(0, \mathbf{I}_m)$ . Set  $\mathbf{e}_i = \frac{\boldsymbol{\xi}_i}{\|\boldsymbol{\xi}_1\|}$  for  $i = 1, \dots, p$ . Then  $\{\mathbf{e}_1, \dots, \mathbf{e}_p\}$ are i.i.d. uniformly distributed on the *m*-dimensional sphere; see, for example, p. 38 from Muirhead (1982). Also,  $\hat{\mathbf{R}} = (\mathbf{e}_i^T \mathbf{e}_j)$ . Set  $V_p := \frac{1}{p} \sum_{1 \le i < j \le p} \left[ m(\mathbf{e}_i' \mathbf{e}_j)^2 - 1 \right]$ . We claim

$$\sup_{p \ge 2} \operatorname{Var}(V_p) \le \frac{3}{2}.$$
(206)

In fact, by Lemma 11 from Cai et al. (2013) or (22) in Lemma 4.1 from Cai and Jiang (2012),  $\{\mathbf{e}_i^T \mathbf{e}_j; 1 \le i < j \le p\}$  are pairwise i.i.d. Thus,  $\operatorname{Var}(V_p) = \frac{1}{p^2} \cdot \frac{1}{2}p(p-1) \cdot m^2 \cdot \operatorname{Var}((\mathbf{e}_1'\mathbf{e}_2)^2)$ . Since  $\mathbf{e}_1$  and  $\mathbf{e}_2$  are independent, by Theorems 1.5.6 and 1.5.7 from Muirhead (1982), we know  $\mathbf{e}_1^T \mathbf{e}_2 \stackrel{d}{=} \xi_1(\xi_1^2 + \dots + \xi_m^2)^{-1/2}$ , where  $\xi_1, \dots, \xi_m$  are i.i.d. N(0, 1). In particular, if n = 2then m = 1,  $\mathbf{e}_1^T \mathbf{e}_2$  is a symmetric Bernoulli random variable, and hence  $(\mathbf{e}_1^T \mathbf{e}_2)^2 = 1$  and  $\operatorname{Var}((\mathbf{e}_1^T \mathbf{e}_2)) = 0$ . Now, for  $n \ge 3$ ,

$$\operatorname{Var}((\mathbf{e}_{1}^{\prime}\mathbf{e}_{2})^{2}) = \operatorname{Var}\left(\frac{\xi_{1}^{2}}{\xi_{1}^{2} + \dots + \xi_{m}^{2}}\right) \leq E\frac{\xi_{1}^{4}}{(\xi_{1}^{2} + \dots + \xi_{m}^{2})^{2}} = \frac{3}{m(m+2)}$$

by taking  $a_1 = 2$  and other  $a_i = 0$  in Lemma 2.4 from Jiang (2012). The inequality is true for all  $m \ge 1$ . Combining this and the earlier expression of  $Var(V_p)$ , we obtain (206). In particular, by the Chebyshev inequality, (206) indicates

$$\frac{V_p}{m} \to 0 \tag{207}$$

as  $p \to \infty$  regardless of the speeds of n and p going to infinity. Write

$$\sum_{1 \le i < j \le p} (\mathbf{e}'_i \mathbf{e}_j)^2 = \frac{p}{m} V_p + \frac{p(p-1)}{2m}.$$

Since  $\hat{\mathbf{R}}$  is a symmetric matrix whose diagonal entries are all equal to 1. The above implies

$$tr(\hat{\mathbf{R}}^2) = p + \frac{2p}{m}V_p + \frac{p(p-1)}{m}.$$
(208)

Consequently, we have from (207) that

$$\frac{2}{p} \cdot \left[ \text{tr}(\hat{\mathbf{R}}^2) - \frac{p^2}{m} \right] = 2 + \frac{4}{m} V_p - \frac{2}{m} \to 2$$
(209)

in probability. It follows from this and (205) that

$$T_{SD} = \frac{\sqrt{2p}}{\sqrt{p(2+4m^{-1}V_p - 2m^{-1})}} \cdot \left[U_p + \frac{1}{m}U_p + \frac{\sqrt{p}}{\sqrt{2}(m-2)}\right].$$

By (207) and the Slutsky lemma,  $\frac{\sqrt{2p}}{\sqrt{p(2+4m^{-1}V_p-2m^{-1})}} \rightarrow 1$  in probability. Therefore, by (203),

$$U_p + \frac{1}{m}U_p + \frac{\sqrt{p}}{\sqrt{2}(m-2)} \rightarrow \begin{cases} \xi_0, & \text{if } p/n^2 \to 0; \\ \xi_0 + \sqrt{h/2}, & \text{if } p/n^2 \to h; \\ \infty, & \text{if } p/n^2 \to \infty \end{cases}$$

in distribution, where  $\xi_0 \sim N(0, 1)$ . So we get part (ii) of Proposition 1.

Finally, as for  $T_{p,1}$ , in lieu of (9),

$$T_{p,1} = \frac{n\bar{\mathbf{X}}^T\hat{\mathbf{D}}^{-1}\bar{\mathbf{X}} - pn(n-3)^{-1}}{\sqrt{2\left|\operatorname{tr}(\hat{\mathbf{R}}^2) - p(p-1)(n-1)^{-1}\right|}} = \frac{\frac{n}{m}\sqrt{2p}U_p}{\sqrt{2p + \frac{4p}{m}V_p}} = \frac{\frac{n}{m}U_p}{\sqrt{1 + \frac{2}{m}V_p}}$$

by (204) and (208). This implies  $T_{p,1} \to N(0,1)$  as  $p \to \infty$  by (203) and (207), regardless of the speeds of n and p going to infinity.

**Proof of Theorem 2**. Recall

$$\bar{\mathbf{X}}_{i} = \frac{1}{n_{i}} \sum_{j=1}^{n_{i}} \mathbf{X}_{ij} \text{ for } i = 1, 2;$$
$$\hat{\mathbf{S}} = \frac{1}{n_{1} + n_{2}} \Big[ \sum_{j=1}^{n_{1}} (\mathbf{X}_{1j} - \bar{\mathbf{X}}_{1}) (\mathbf{X}_{1j} - \bar{\mathbf{X}}_{1})^{T} + \sum_{j=1}^{n_{2}} (\mathbf{X}_{2j} - \bar{\mathbf{X}}_{2}) (\mathbf{X}_{2j} - \bar{\mathbf{X}}_{2})^{T} \Big].$$

The  $p \times p$  matrix  $\hat{\mathbf{D}}$  is the diagonal matrix of  $\hat{\mathbf{S}}$  and  $\hat{\mathbf{R}} = \hat{\mathbf{D}}^{-1/2} \hat{\mathbf{S}} \hat{\mathbf{D}}^{-1/2}$  is the  $p \times p$  pooled sample correlation matrix. In particular, all of the diagonal entries of  $\hat{\mathbf{R}}$  are 1. Thus,

 $\|\mathbf{R}\|_F \ge \sqrt{p}$ . Now,  $\bar{\mathbf{X}}_i$  is independent of  $\sum_{j=1}^{n_i} (\mathbf{X}_{ij} - \bar{\mathbf{X}}_i) (\mathbf{X}_{ij} - \bar{\mathbf{X}}_i)^T$  for each i = 1, 2; see, for example, Theorem 3.1.2 from Muirhead (1982). Second, by assumption, the two samples are independent. Therefore, the three random vectors  $\bar{\mathbf{X}}_1$ ,  $\bar{\mathbf{X}}_2$  and  $\hat{\mathbf{S}}$  are independent. In particular,  $\bar{\mathbf{X}}_1$  and  $\bar{\mathbf{X}}_2$  are independent of  $\hat{\mathbf{D}}$ . By assumption  $\boldsymbol{\mu}_1 = \boldsymbol{\mu}_2$ , we obtain

$$ar{oldsymbol{X}}_1 - ar{oldsymbol{X}}_2 \sim N_p \Big( oldsymbol{0}, \Big( rac{1}{n_1} + rac{1}{n_2} \Big) oldsymbol{\Sigma} \Big)$$

Hence,

$$\sqrt{\frac{n_1 n_2}{n_1 + n_2}} \left( \bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2 \right) \sim N_p(\boldsymbol{0}, \boldsymbol{\Sigma}).$$
(210)

Furthermore, for each i,  $\sum_{j=1}^{n_i} (\mathbf{X}_{ij} - \bar{\mathbf{X}}_i) (\mathbf{X}_{ij} - \bar{\mathbf{X}}_i)^T \sim W_p(n_i - 1, \boldsymbol{\Sigma})$ ; see, for example, Theorem 3.1.2 from Muirhead (1982). Therefore,  $(n_1 + n_2)\hat{\mathbf{S}} \sim W_p(n_1 + n_2 - 2, \boldsymbol{\Sigma})$  by independence. Consequently,

$$\hat{\mathbf{D}}_1 := \frac{n_1 + n_2}{n_1 + n_2 - 1} \hat{\mathbf{D}} \stackrel{d}{=} \text{the diagonal matrix of } \frac{1}{m} W_p(m - 1, \mathbf{\Sigma}), \tag{211}$$

where  $m := n_1 + n_2 - 1$ . As explained earlier, the left hand side of (210) is independent of  $\hat{\mathbf{D}}_1$ . In particular, by replacing " $\boldsymbol{\eta}$ " and "n" from (174) with " $\sqrt{\frac{n_1n_2}{n_1+n_2}} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2)$ " and "m", respectively, we obtain

$$E\left[\frac{n_{1}n_{2}}{n_{1}+n_{2}-1}\left(\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}\right)^{T}\hat{\boldsymbol{D}}^{-1}\left(\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}\right)\right]$$
  
=  $E\left[\frac{n_{1}n_{2}}{n_{1}+n_{2}}\left(\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}\right)^{T}\hat{\boldsymbol{D}}_{1}^{-1}\left(\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}\right)\right]$   
=  $\frac{(n_{1}+n_{2}-1)p}{n_{1}+n_{2}-4}.$  (212)

This is the reason how come up with the numerator of  $T_{p,2}$  defined in (16). The assumption  $\lim_{p\to\infty} \frac{p}{(n_1+n_2)^a} = 0$  implies  $n_1 + n_2 \to \infty$  as  $p \to \infty$ . Thus we know from assumption (b) that  $\lim_{p\to\infty} \frac{p}{m ||\mathbf{R}||_F} = 0$ . Replacing "n" with "m" in Lemma 4.1, we have from (210) and (211) that

$$\frac{\frac{n_{1}n_{2}}{n_{1}+n_{2}} (\bar{\boldsymbol{X}}_{1} - \bar{\boldsymbol{X}}_{2})^{T} \hat{\boldsymbol{D}}_{1}^{-1} (\bar{\boldsymbol{X}}_{1} - \bar{\boldsymbol{X}}_{2}) - p}{\sqrt{2 \operatorname{tr}(\mathbf{R}^{2})}} \\
= \frac{\frac{n_{1}n_{2}}{n_{1}+n_{2}} (\bar{\boldsymbol{X}}_{1} - \bar{\boldsymbol{X}}_{2})^{T} \mathbf{D}^{-1} (\bar{\boldsymbol{X}}_{1} - \bar{\boldsymbol{X}}_{2}) - p}{\sqrt{2 \operatorname{tr}(\mathbf{R}^{2})}} + o_{p}(1), \quad (213)$$

Ξ

where **D** is the diagonal matrix of  $\Sigma$ ,  $\mathbf{R} = \mathbf{D}^{-1/2} \Sigma \mathbf{D}^{-1/2}$  is the population correlation matrix. By (196) and (210),

$$\frac{n_1 n_2}{n_1 + n_2} \left( \bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2 \right)^T \mathbf{D}^{-1} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2) \stackrel{d}{=} \sum_{i=1}^p \lambda_i \xi_i^2,$$
(214)

where  $\lambda_1 \geq \cdots \geq \lambda_p \geq 0$  are the eigenvalues of **R** and  $\xi_1, \cdots, \xi_p$  are i.i.d. N(0, 1). Similar to the argument as in *Step 2* in the proof of Theorem 1, we see

$$\frac{1}{\|\mathbf{R}\|_F} \Big( -p + \sum_{i=1}^p \lambda_i \xi_i^2 \Big) \to \sqrt{2} \Big( 1 - \sum_{i=1}^\infty \rho_i^2 \Big)^{1/2} \xi_0 + \sum_{i=1}^\infty \rho_i (\xi_i^2 - 1)$$

in distribution, where  $\xi_0 \sim N(0, 1)$  and  $\xi_0$  is independent of  $\xi_1, \xi_2, \cdots$ . Combining this with (213) and (214), we have

$$\frac{\frac{n_{1}n_{2}}{n_{1}+n_{2}-1} \left(\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}\right)^{T} \hat{\mathbf{D}}^{-1} (\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}) - p}{\sqrt{2 \operatorname{tr}(\mathbf{R}^{2})}} \\
= \frac{\frac{n_{1}n_{2}}{n_{1}+n_{2}} \left(\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}\right)^{T} \hat{\mathbf{D}}_{1}^{-1} (\bar{\boldsymbol{X}}_{1}-\bar{\boldsymbol{X}}_{2}) - p}{\sqrt{2 \operatorname{tr}(\mathbf{R}^{2})}} \\
\rightarrow \left(1-\sum_{i=1}^{\infty} \rho_{i}^{2}\right)^{1/2} \xi_{0} + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_{i} (\xi_{i}^{2}-1) \tag{215}$$

in distribution. Evidently,

$$\frac{1}{\sqrt{2\mathrm{tr}(\mathbf{R}^2)}} \left[ p - \frac{(n_1 + n_2 - 1)p}{n_1 + n_2 - 4} \right] = \frac{1}{\sqrt{2}} \frac{-3p}{(n_1 + n_2 - 4)} \|\mathbf{R}\|_F \to 0$$

by the assumption  $\lim_{p\to\infty} \frac{p}{(n_1+n_2)\|\mathbf{R}\|_F} = 0$ . Add the left hand sides of the above two assertions to obtain

$$\frac{\frac{n_1 n_2}{n_1 + n_2 - 1} \left( \bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2 \right)^T \hat{\mathbf{D}}^{-1} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2) - \frac{(n_1 + n_2 - 1)p}{n_1 + n_2 - 4}}{\sqrt{2 \operatorname{tr}(\mathbf{R}^2)}}$$

$$\rightarrow \left( 1 - \sum_{i=1}^{\infty} \rho_i^2 \right)^{1/2} \xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i (\xi_i^2 - 1)$$
(216)

in distribution. Next we will replace  $tr(\mathbf{R}^2)$  by its ratio-unbiased-estimator.

Recall  $(n_1 + n_2)\hat{\mathbf{S}} \sim W_p(n_1 + n_2 - 2, \boldsymbol{\Sigma})$ , and  $\hat{\mathbf{D}}$  is the diagonal matrix of  $\hat{\mathbf{S}}$  and  $\hat{\mathbf{R}} = \hat{\mathbf{D}}^{-1/2}\hat{\mathbf{S}}\hat{\mathbf{D}}^{-1/2}$ . Set  $\kappa = \frac{n_1 + n_2}{n_1 + n_2 - 1}$ . Then  $\kappa \hat{\mathbf{S}} \sim \frac{1}{n_1 + n_2 - 1}W_p(n_1 + n_2 - 2, \boldsymbol{\Sigma});$  $\hat{\mathbf{R}} = (\kappa \hat{\mathbf{D}})^{-1/2}(\kappa \hat{\mathbf{S}})(\kappa \hat{\mathbf{D}})^{-1/2}$  and  $\kappa \hat{\mathbf{D}}$  is the diagonal matrix of  $\kappa \hat{\mathbf{S}}$ . The essential assumption from Theorem 3 is that, in its own notation, **R** is the sample correlation matrix obtained from  $\hat{\mathbf{S}} \sim \frac{1}{n} W_p(n-1, \boldsymbol{\Sigma})$ . Replace " $\hat{\mathbf{S}}$ " and "n" from Theorem 3 with " $\kappa \hat{\mathbf{S}}$ " and " $n_1 + n_2 - 1$ ", respectively. Assumption (b) indicates  $\lim_{p \to \infty} \frac{p}{(n_1 + n_2 - 1)^{\parallel} \mathbf{R} \parallel_F} = 0$  and  $\lim_{p \to \infty} \frac{p}{(n_1 + n_2 - 1)^a} = 0$ . Then by Theorem 3,

$$\frac{1}{\mathrm{tr}(\mathbf{R}^2)} \Big[ \mathrm{tr}(\hat{\mathbf{R}}^2) - \frac{p(p-1)}{n_1 + n_2 - 2} \Big] \to 1$$
(217)

in probability. This together with (216) and the Slutsky lemma yields

$$T_{p,2} = \frac{\frac{n_1 n_2}{n_1 + n_2 - 1} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2)^T \hat{\mathbf{D}}^{-1} (\bar{\boldsymbol{X}}_1 - \bar{\boldsymbol{X}}_2) - \frac{(n_1 + n_2 - 1)p}{n_1 + n_2 - 4}}{\sqrt{2 \left| \operatorname{tr}(\hat{\mathbf{R}}^2) - \frac{p(p-1)}{n_1 + n_2 - 2} \right|}} \\ \rightarrow \left( 1 - \sum_{i=1}^{\infty} \rho_i^2 \right)^{1/2} \xi_0 + \frac{1}{\sqrt{2}} \sum_{i=1}^{\infty} \rho_i (\xi_i^2 - 1)$$

in distribution as  $p \to \infty$ .

Finally, by the same argument as deriving (193), we see  $T'_{SD} = T_{p,2} \cdot [1 + o_p(1)] + o_p(1)$ . So the conclusion for  $T'_{SD}$  follows from the above and the Slutsky lemma. The proof is completed.

## 6.4 A Lemma and Verification of (11), (12) and (13)

The following lemma is used in the discussion after Theorem 2.

**LEMMA 6.37** Let  $\{\mathbf{R}_p; p \ge 1\}$  be non-negative definite matrices whose diagonal entries are all equal to 1. Let  $\lambda_{p,1} \ge \cdots \ge \lambda_{p,p} \ge 0$  be the eigenvalues of  $\mathbf{R}_p$ . Assume condition "C4" stated in Theorem 2 from Zhang et al. (2020b) hold, that is,  $\lim_{p\to\infty} \frac{\lambda_{p,i}}{\|\mathbf{R}_p\|_F} = \rho_i$  for all  $i \ge 1$  with  $\rho_1 > 0$  and  $\lim_{p\to\infty} \sum_{i=1}^p \frac{\lambda_{p,i}}{\|\mathbf{R}_p\|_F} = \sum_{i=1}^\infty \rho_i < \infty$ . Then  $\sum_{i=1}^\infty \rho_i^2 = 1$ .

**Proof of Lemma 6.37**. Notice  $\sum_{i=1}^{p} \lambda_{p,i} = \operatorname{tr}(\mathbf{R}_p) = p$  because all of the diagonal entries of  $\mathbf{R}_p$  are identical to 1. The assumption  $\lim_{p\to\infty} \sum_{i=1}^{p} \frac{\lambda_{p,i}}{\|\mathbf{R}_p\|_F} = \sum_{i=1}^{\infty} \rho_i < \infty$  implies that

$$\frac{p}{\|\mathbf{R}_p\|_F} \to \sum_{i=1}^{\infty} \rho_i \tag{218}$$

as  $p \to \infty$ . Set  $a_{p,i} = \frac{\lambda_{p,i}}{\|\mathbf{R}_p\|_F}$ . Then  $\lim_{p\to\infty} a_{p,i} = \rho_i$  and  $a_{p,1}^2 + \cdots + a_{p,p}^2 = 1$  by the definition of  $\|\mathbf{R}_p\|_F$ . For any  $K \ge 1$ , write

$$a_{p,1}^2 + \dots + a_{p,K}^2 = 1 - \sum_{i=K+1}^p a_{p,i}^2.$$
 (219)

We claim that  $\lim_{K\to\infty} \limsup_{p\to\infty} \sum_{i=K+1}^p a_{p,i}^2 = 0$ . If this is true, by letting  $p \to \infty$  first and then sending  $K \to \infty$ , then  $\sum_{i=1}^{\infty} \rho_i^2 = 1$ . We now prove the claim. In fact, write

$$\sum_{i=K+1}^{p} a_{p,i} = \frac{p}{\|\mathbf{R}_p\|_F} - \sum_{i=1}^{K} a_{p,i}.$$

For fixed  $K \ge 1$ , let  $p \to \infty$  and use (218) to have

$$\lim_{p \to \infty} \sum_{i=K+1}^{p} a_{p,i} = \left(\sum_{i=1}^{\infty} \rho_i\right) - \sum_{i=1}^{K} \rho_i = \sum_{i=K+1}^{\infty} \rho_i.$$

Use the assumption  $\sum_{i=1}^{\infty} \rho_i < \infty$  and let  $K \to \infty$  to see  $\lim_{K \to \infty} \lim_{p \to \infty} \sup_{p \to \infty} \sum_{i=K+1}^{p} a_{p,i} = 0$ . The claim is then verified since  $a_{p,i} \leq 1$  for all  $1 \leq i \leq p$ .

In the following the notation " $A_p \sim B_p$ " means that  $A_p/B_p \to 1$  as  $p \to \infty$ . The Verification of (11). Review Example 2.1. We have

$$\mathbf{R} = \begin{pmatrix} 1 & r & r & \cdots & r \\ r & 1 & r & \cdots & r \\ \vdots & & & & \\ r & r & r & \cdots & 1 \end{pmatrix}_{p \times p}$$

Easily,  $\lambda_1 = 1 + (p-1)r$  and  $\lambda_2 = \cdots = \lambda_p = 1 - r$ . To maintain **R** to be non-negative definite, all eigenvalues have to be non-negative, that is,  $-\frac{1}{p-1} \leq r \leq 1$ . Recall  $r = r_p$  satisfies that  $\lim_{p\to\infty} \sqrt{p} \cdot r = c$ . Now we consider three cases:  $c = 0, c \in (0, \infty)$  and  $c = \infty$ .

Case 1: c = 0. In this case  $\|\mathbf{R}\|_F^2 = [1 + (p-1)r]^2 + (p-1)(1-r)^2 \sim p$  as  $p \to \infty$ , thus  $\|\mathbf{R}\|_F \sim \sqrt{p}$ . Easily,  $\frac{p}{n\|\mathbf{R}\|_F} = O(\frac{\sqrt{p}}{n}) \to 0$  provided  $p = o(n^2)$ . Also,  $\rho_1 = \lim_{p \to \infty} \frac{\lambda_1}{\|\mathbf{R}\|_F} = 0$ , hence  $\lambda_i = 0$  for every  $i \ge 1$ . By Theorem 1, both  $T_{SD}$  and  $T_{p,1}$  converge to N(0, 1) in distribution.

Case 2:  $c \in (0, \infty)$ . In this case  $\|\mathbf{R}\|_F^2 = [1 + (p-1)r]^2 + (p-1)(1-r)^2 \sim (c^2+1)p$  as  $p \to \infty$ , hence  $\|\mathbf{R}\|_F \sim \sqrt{c^2+1} \cdot \sqrt{p}$ . Readily,

$$\frac{p}{n \|\mathbf{R}\|_F} = O\left(\frac{\sqrt{p}}{n}\right) \to 0 \quad \text{and} \quad \rho_1 = \lim_{p \to \infty} \frac{\lambda_1}{\|\mathbf{R}\|_F} \to \frac{c}{\sqrt{c^2 + 1}};$$
$$\rho_i = \lim_{p \to \infty} \frac{\lambda_i}{\|\mathbf{R}\|_F} = 0 \quad \text{for every} \quad i \ge 2$$

provided  $p = o(n^2)$ . By Theorem 1, under condition  $p = o(n^2)$ , we know both  $T_{SD}$  and  $T_{p,1}$  converge to

$$\frac{1}{\sqrt{c^2+1}}\xi_0 + \frac{c}{\sqrt{2(c^2+1)}}(\xi_1^2 - 1)$$

in distribution, where  $\xi_0$  and  $\xi_1$  are i.i.d. N(0, 1).

Case 3:  $c = \infty$ . In this case  $\|\mathbf{R}\|_F^2 = [1 + (p-1)r]^2 + (p-1)(1-r)^2 \sim (pr)^2$  as  $p \to \infty$ . Therefore,  $\|\mathbf{R}\|_F \sim pr$ . Notice  $\lim_{p\to\infty} p/(n\|\mathbf{R}\|_F) = \lim_{p\to\infty} 1/(nr_p) = 0$ ,  $\rho_1 = \lim_{p\to\infty} \lambda_1/\|\mathbf{R}\|_F = 1$  and  $\rho_i = 0$  for  $i \ge 2$  provided  $nr_p \to \infty$ . So, under conditions  $nr_p \to \infty$  and  $p = o(n^a)$  for some constant a > 0, we have  $T_{p,1} \to \frac{1}{\sqrt{2}}[\chi^2(1) - 1]$ .

The Verification of (12). Recall  $r \in (0,1)$ ,  $m = [p^r]$  and  $\mathbf{A}_m$  from Example 2.1. By definition,

$$\mathbf{R} = egin{pmatrix} \mathbf{A}_m & \mathbf{0} \ \mathbf{0} & \mathbf{I}_{p-m} \end{pmatrix}.$$

Then the largest eigenvalues of **R** is  $\lambda_1 = 1 + (m-1)r$  and the rest of them are  $\lambda_2 = \cdots = \lambda_m = 1 - r$  and  $\lambda_{m+1} = \cdots = \lambda_p = 1$ . Then

$$\|\mathbf{R}\|_F^2 = [1 + (m-1)r]^2 + (m-1)(1-r)^2 + (p-m) = m^2r^2 + p + O(m).$$

Thus,

$$\|\mathbf{R}\|_{F} = \begin{cases} \sqrt{p} \cdot [1 + o(1)], & \text{if } 0 < r < \frac{1}{2}; \\ \sqrt{5p/4} \cdot [1 + o(1)], & \text{if } r = \frac{1}{2}; \\ rp^{r} \cdot [1 + o(1)], & \text{if } \frac{1}{2} < r < 1. \end{cases}$$

Also,

$$\rho_1 = \lim_{p \to \infty} \frac{\lambda_1}{\|\mathbf{R}\|_F} = \begin{cases} 0, & \text{if } 0 < r < \frac{1}{2}; \\ 1/\sqrt{5}, & \text{if } r = \frac{1}{2}; \\ 1, & \text{if } \frac{1}{2} < r < 1. \end{cases}$$

Obviously,  $\rho_i = 0$  for  $i \ge 2$ . At last,

$$\frac{p}{n \|\mathbf{R}\|_F} = \begin{cases} O(\sqrt{p}/n), & \text{if } 0 < r \le \frac{1}{2}; \\ O(p^{1-r}/n), & \text{if } \frac{1}{2} < r < 1. \end{cases}$$

Assume  $\mu = 0$ . Then, we have from Theorem 1 that both  $T_{SD}$  and  $T_{p,1}$  go to

$$\begin{cases} N(0,1), & \text{if } 0 < r < \frac{1}{2}; \\ \frac{2}{\sqrt{5}}N(0,1) + \frac{1}{\sqrt{10}} \cdot [\chi^2(1) - 1], & \text{if } r = \frac{1}{2}; \\ \frac{1}{\sqrt{2}} \cdot [\chi^2(1) - 1], & \text{if } \frac{1}{2} < r < 1 \end{cases}$$

in distribution under condition  $p = o(n^2)$  for  $0 < r \le \frac{1}{2}$  and  $p = o(n^{1/(1-r)})$  for  $\frac{1}{2} < r < 1$ , where the random variables N(0, 1) and  $\chi^2(1)$  are independent as  $r = \frac{1}{2}$ .

The Verification of (13). Recall  $m = [\log(p+2)]$ . Given  $\tau \ge 0$ , we have  $p' = p - [\tau \sqrt{p}]$ ,  $\lambda_i = 1 + \tau 2^{-i} (1 - 2^{-m})^{-1} \sqrt{p}$  for  $1 \le i \le m$  and  $\lambda_i = 1$  for  $i = m + 1, \dots, p' - 1$ ,  $\lambda_{p'} = 1 + [\tau \sqrt{p}] - \tau \sqrt{p} \in [0, 1]$  and the rest of  $\lambda_i$  are zero. Obviously,  $\lambda_1 + \dots + \lambda_k \ge k$  for each  $i = 1, \dots, p' - 1$  and  $\lambda_1 + \dots + \lambda_{p'}$  is identical to

$$\sum_{i=1}^{m} \left[ 1 + \tau 2^{-i} (1 - 2^{-m})^{-1} \sqrt{p} \right] + (p' - m - 1) \cdot 1 + (1 + [\tau \sqrt{p}] - \tau \sqrt{p})$$
  
=  $(m + \tau \sqrt{p}) + (p - [\tau \sqrt{p}] - m - 1) + 1 + [\tau \sqrt{p}] - \tau \sqrt{p}$   
=  $p$ ,

where we use the fact  $\sum_{i=1}^{m} 2^{-i} (1 - 2^{-m})^{-1} = 1$ . Lemma 4.3 tells us that there exists a

correlation matrix **R** such that **R** has eigenvalues  $\lambda_i$ ,  $1 \leq i \leq p$ . Now,

$$\|\mathbf{R}\|_{F}^{2} = \sum_{i=1}^{m} \left[1 + \tau 2^{-i} \left(1 - 2^{-m}\right)^{-1} \sqrt{p}\right]^{2} + (p' - m - 1) \cdot 1^{2} + \left(1 + \left[\tau \sqrt{p}\right] - \tau \sqrt{p}\right)^{2}$$
$$= \tau^{2} p \left(1 - 2^{-m}\right)^{-2} \sum_{i=1}^{m} \frac{1}{4^{i}} + p + O\left(\sqrt{p}\right)$$
$$= \left(\frac{\tau^{2}}{3} + 1\right) p + o(p)$$

since  $\sum_{i=1}^{m} \frac{1}{4^i} \to \frac{1}{3}$  as  $p \to \infty$ . In particular,  $\lim_{p\to\infty} \frac{p}{n \|\mathbf{R}\|_F} = 0$  provided  $p = o(n^2)$ . Easily, for each  $i \ge 1$ ,

$$\rho_i := \lim_{p \to \infty} \frac{\lambda_i}{\|\mathbf{R}\|_F} = \lim_{p \to \infty} \frac{1 + \tau 2^{-i} (1 - 2^{-m})^{-1} \sqrt{p}}{\sqrt{(\frac{\tau^2}{3} + 1)p + o(p)}} \to \sqrt{\frac{3\tau^2}{\tau^2 + 3}} \cdot \frac{1}{2^i}$$

as  $p \to \infty$ . Now,

$$\sum_{i=1}^{\infty} \rho_i^2 = \frac{3\tau^2}{\tau^2 + 3} \sum_{i=1}^{\infty} \frac{1}{4^i} = \frac{\tau^2}{\tau^2 + 3}.$$

Assuming  $\mu = 0$ , by Theorem 1, we see that  $T_{p,1} \to \sqrt{\frac{3}{\tau^2+3}} \xi_0 + \sqrt{\frac{3\tau^2}{2(\tau^2+3)}} \sum_{i=1}^{\infty} \frac{1}{2^i} (\xi_i^2 - 1)$  in distribution, where  $\xi_0, \xi_1, \dots, \xi_d$  are i.i.d. N(0, 1).

## References

- Theodore W Anderson. An Introduction to Multivariate Statistical Analysis. John Wiley & Sons, 2003.
- Zhidong Bai and Hewa Saranadasa. Effect of high dimension: by an example of a two sample problem. *Statistica Sinica*, pages 311–329, 1996.
- Zhidong Bai and Jack W Silverstein. Spectral analysis of large dimensional random matrices, volume 20. Springer, 2010.
- Zhigang Bao, Guangming Pan, and Wang Zhou. Tracy-widom law for the extreme eigenvalues of sample correlation matrices. *Electronic Journal of Probability*, 17:1–32, 2012.

- Munmun Biswas and Anil K Ghosh. A nonparametric two-sample test applicable to high dimensional data. *Journal of Multivariate Analysis*, 123:160–171, 2014.
- Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. Concentration inequalities: A nonasymptotic theory of independence. Oxford university press, 2013.
- Peter J Brockwell, Peter J Brockwell, Richard A Davis, and Richard A Davis. *Introduction* to time series and forecasting. Springer, 2016.
- T Tony Cai and Tiefeng Jiang. Phase transition in limiting distributions of coherence of high-dimensional random matrices. *Journal of Multivariate Analysis*, 107:24–39, 2012.
- T Tony Cai, Jianqing Fan, and Tiefeng Jiang. Distributions of angles in random packing on spheres. *Journal of Machine Learning Research*, 14:1837, 2013.
- T Tony Cai, Weidong Liu, and Yin Xia. Two-sample test of high dimensional means under dependence. Journal of the Royal Statistical Society: Series B: Statistical Methodology, pages 349–372, 2014.
- Anirvan Chakraborty and Probal Chaudhuri. Tests for high-dimensional data based on means, spatial signs and spatial ranks. *The Annals of Statistics*, 45(2):771–799, 2017.
- Jinyuan Chang, Chao Zheng, Wen-Xin Zhou, and Wen Zhou. Simulation-based hypothesis testing of high dimensional means under covariance heterogeneity. *Biometrics*, 73(4): 1300–1310, 2017.
- Lin S Chen, Debashis Paul, Ross L Prentice, and Pei Wang. A regularized hotelling's t 2 test for pathway analysis in proteomic studies. *Journal of the American Statistical Association*, 106(496):1345–1360, 2011.
- Song Xi Chen and Ying-Li Qin. A two-sample test for high-dimensional data with applications to gene-set testing. *The Annals of Statistics*, 38(2):808–835, 2010.
- Song Xi Chen, Jun Li, and Ping-Shou Zhong. Two-sample tests for high dimensional means with thresholding and data transformation. *arXiv preprint arXiv:1410.2848*, 2014.

- Song Xi Chen, Jun Li, and Ping-Shou Zhong. Two-sample and anova tests for high dimensional means. *The Annals of Statistics*, 47(3):1443–1474, 2019.
- Yuan Shih Chow and Henry Teicher. Probability theory: Independence, interchangeability, martingales. 1997.
- Arthur P Dempster. A high dimensional two sample significance test. The Annals of Mathematical Statistics, pages 995–1010, 1958.
- Arthur P Dempster. A significance test for the separation of two highly multivariate small samples. *Biometrics*, 16(1):41–50, 1960.
- Kai Dong, Herbert Pang, Tiejun Tong, and Marc G Genton. Shrinkage-based diagonal hotelling's tests for high-dimensional small sample size data. *Journal of Multivariate Analysis*, 143:127–142, 2016.
- Rick Durrett. *Probability: theory and examples*, volume 49. Cambridge university press, 2019.
- Jianqing Fan and Tiefeng Jiang. Largest entries of sample correlation matrices from equicorrelated normal populations. *The Annals of Probability*, 47(5):3321–3374, 2019.
- Jianqing Fan, Jianhua Guo, and Shurong Zheng. Estimating number of factors by adjusted eigenvalues thresholding. Journal of the American Statistical Association, pages 1–10, 2020.
- Long Feng, Changliang Zou, and Zhaojun Wang. Multivariate-sign-based high-dimensional tests for the two-sample location problem. *Journal of the American Statistical Association*, 111(514):721–735, 2016.
- Jiti Gao, Xiao Han, Guangming Pan, and Yanrong Yang. High dimensional correlation matrices: The central limit theorem and its applications. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 79(3):677–693, 2017.

- B. K. Ghosh. Asymptotic expansions for the moments of the distribution of correlation coefficient. *Biometrika*, 53(1/2):258–262, 1966.
- Karl Bruce Gregory, Raymond J Carroll, Veerabhadran Baladandayuthapani, and Soumendra N Lahiri. A two-sample test for equality of means in high dimension. Journal of the American Statistical Association, 110(510):837–849, 2015.
- Arthur Gretton, Dino Sejdinovic, Heiko Strathmann, Sivaraman Balakrishnan, Massimiliano Pontil, Kenji Fukumizu, and Bharath K Sriperumbudur. Optimal kernel choice for largescale two-sample tests. In Advances in neural information processing systems, pages 1205– 1213. Citeseer, 2012.
- V Guiard. A general formula for the central mixed moments of the multivariate normal distribution. Statistics: A Journal of Theoretical and Applied Statistics, 17(2):279–289, 1986.
- Bin Guo and Song Xi Chen. Tests for high dimensional generalized linear models. Journal of the Royal Statistical Society. Series B (Statistical Methodology), pages 1079–1102, 2016.
- Roger A Horn and Charles R Johnson. *Matrix analysis*. Cambridge university press, 2012.
- Harold Hotelling. The generalization of student's ratio. The Annals of Mathematical Statistics, 2(3):360–378, 1931.
- Jiang Hu and Zhidong Bai. A review of 20 years of naive tests of significance for highdimensional mean vectors and covariance matrices. *Science China Mathematics*, 59(12): 2281–2300, 2016.
- Tiefeng Jiang. The limiting distributions of eigenvalues of sample correlation matrices. Sankhyā: The Indian Journal of Statistics, pages 35–48, 2004.
- Tiefeng Jiang. A variance formula related to a quantum conductance problem. *Physics* Letters A, 373(25):2117–2121, 2009.

- Tiefeng Jiang and Fan Yang. Central limit theorems for classical likelihood ratio tests for high-dimensional normal distributions. *The Annals of Statistics*, 41(4):2029–2074, 2013.
- Søren Johansen. The Welch-James approximation to the distribution of the residual sum of squares in a weighted linear regression. *Biometrika*, 67(1):85–92, 1980.
- K Krishnamoorthy and Jianqi Yu. Modified nel and van der merwe test for the multivariate behrens-fisher problem. *Statistics & probability letters*, 66(2):161–169, 2004.
- Michel Ledoux. *The concentration of measure phenomenon*. Number 89. American Mathematical Soc., 2001.
- Haoran Li, Alexander Aue, Debashis Paul, Jie Peng, and Pei Wang. An adaptable generalization of hotelling's  $t^2$  test in high dimension. *The Annals of Statistics*, 48(3):1815–1847, 2020.
- Miles Lopes, Laurent Jacob, and Martin J Wainwright. A more powerful two-sample test in high dimensions using random projection. *Advances in Neural Information Processing Systems*, 24:1206–1214, 2011.
- David Morales-Jimenez, Iain M Johnstone, Matthew R McKay, and Jeha Yang. Asymptotics of eigenstructure of sample correlation matrices for high-dimensional spiked models. *Statistica Sinica*, 31(2):571, 2021.
- Robb J Muirhead. Aspects of multivariate statistical theory. John Wiley & Sons, 1982.
- DG Nel and CA Van der Merwe. A solution to the multivariate behrens-fisher problem. Communications in Statistics-Theory and Methods, 15(12):3719–3735, 1986.
- Junyong Park and Deepak Nag Ayyala. A test for the mean vector in large dimension and small samples. *Journal of Statistical Planning and Inference*, 143(5):929–943, 2013.
- Muni S Srivastava. A test for the mean vector with fewer observations than the dimension under non-normality. *Journal of Multivariate Analysis*, 100(3):518–532, 2009.

- Muni S Srivastava and Meng Du. A test for the mean vector with fewer observations than the dimension. *Journal of Multivariate Analysis*, 99(3):386–402, 2008.
- Muni S Srivastava and Yasunori Fujikoshi. Multivariate analysis of variance with fewer observations than the dimension. *Journal of Multivariate Analysis*, 97(9):1927–1940, 2006.
- Muni S Srivastava, Shota Katayama, and Yutaka Kano. A two sample test in high dimensional data. *Journal of Multivariate Analysis*, 114:349–358, 2013.
- Radhendushka Srivastava, Ping Li, and David Ruppert. RAPTT: An exact two-sample test in high dimensions using random projections. *Journal of Computational and Graphical Statistics*, 25(3):954–970, 2016.
- Lan Wang, Bo Peng, and Runze Li. A high-dimensional nonparametric multivariate test for mean vector. Journal of the American Statistical Association, 110(512):1658–1669, 2015.
- Bernard L Welch. The generalization of 'student's' problem when several different population variances are involved. *Biometrika*, 34(1-2):28–35, 1947.
- Yujun Wu, Marc G Genton, and Leonard A Stefanski. A multivariate two-sample mean test for small sample size and missing data. *Biometrics*, 62(3):877–885, 2006.
- Han Xiao and Wang Zhou. Almost sure limit of the smallest eigenvalue of some sample correlation matrices. *Journal of Theoretical Probability*, 23(1):1–20, 2010.
- Gongjun Xu, Lifeng Lin, Peng Wei, and Wei Pan. An adaptive two-sample test for highdimensional means. *Biometrika*, 103(3):609–624, 2016.
- Kaijie Xue and Fang Yao. Distribution and correlation-free two-sample test of highdimensional means. *The Annals of Statistics*, 48(3):1304–1328, 2020.
- Ying Yao. An approximate degrees of freedom solution to the multivariate behrens fisher problem. *Biometrika*, 52(1/2):139–147, 1965.

- Jin-Ting Zhang, Jia Guo, and Bu Zhou. Linear hypothesis testing in high-dimensional oneway MANOVA. *Journal of Multivariate Analysis*, 155:200–216, 2017.
- Jin-Ting Zhang, Jia Guo, Bu Zhou, and Ming-Yen Cheng. A simple two-sample test in high dimensions based on l 2-norm. Journal of the American Statistical Association, 115(530): 1011–1027, 2020a.
- Liang Zhang, Tianming Zhu, and Jin-Ting Zhang. A simple scale-invariant two-sample test for high-dimensional data. *Econometrics and Statistics*, 14:131–144, 2020b.
- Ping-Shou Zhong, Song Xi Chen, and Minya Xu. Tests alternative to higher criticism for high-dimensional means under sparsity and column-wise dependence. *The Annals of Statistics*, 41(6):2820–2851, 2013.