STAR TRANSPOSITION GRAY CODES FOR MULTISET PERMUTATIONS

PETR GREGOR, ARTURO MERINO, AND TORSTEN MÜTZE

ABSTRACT. Given integers $k \geq 2$ and $a_1, \ldots, a_k \geq 1$, let $a := (a_1, \ldots, a_k)$ and $n := a_1 + \cdots + a_k$. An a-multiset permutation is a string of length n that contains exactly a_i symbols i for each $i=1,\ldots,k$. In this work we consider the problem of exhaustively generating all a-multiset permutations by star transpositions, i.e., in each step, the first entry of the string is transposed with any other entry distinct from the first one. This is a far-ranging generalization of several known results. For example, it is known that permutations $(a_1 = \cdots = a_k = 1)$ can be generated by star transpositions, while combinations (k=2) can be generated by these operations if and only if they are balanced $(a_1 = a_2)$, with the positive case following from the middle levels theorem. To understand the problem in general, we introduce a parameter $\Delta(a) :=$ $n-2\max\{a_1,\ldots,a_k\}$ that allows us to distinguish three different regimes for this problem. We show that if $\Delta(a) < 0$, then a star transposition Gray code for a-multiset permutations does not exist. We also construct such Gray codes for the case $\Delta(a) > 0$, assuming that they exist for the case $\Delta(a) = 0$. For the case $\Delta(a) = 0$ we present some partial positive results. Our proofs establish Hamilton-connectedness or Hamilton-laceability of the underlying flip graphs, and they answer several cases of a recent conjecture of Shen and Williams. In particular, we prove that the middle levels graph is Hamilton-laceable.

1. Introduction

Permutations and combinations are two of the most fundamental classes of combinatorial objects. Specifically, k-permutations are all linear orderings of $[k] := \{1, \ldots, k\}$, and their number is k!. Moreover, (α, β) -combinations are all β -element subsets of [n] where $n := \alpha + \beta$, and their number is $\binom{n}{\alpha} = \binom{n}{\beta}$. Permutations and combinations are generalized by so-called multiset permutations, and in this paper we consider the task of listing them such that any two consecutive objects in the list differ by particular transpositions, i.e., by swapping two elements. Such a listing of objects subject to a 'small change' operation is often referred to as $Gray\ code\ [Sav97,\ Rus16]$. One of the standard references for algorithms that efficiently generate various combinatorial objects, including permutations and combinations, is Knuth's book [Knu11] (see also [NW75]).

1.1. **Permutation generation.** There is a vast number of Gray codes for permutation generation, most prominently the Steinhaus-Johnson-Trotter algorithm [Tro62, Joh63], which generates

⁽Petr Gregor) Department of Theoretical Computer Science and Mathematical Logic, Charles University, Prague, Czech Republic

⁽Arturo Merino) DEPARTMENT OF MATHEMATICS, TU BERLIN, GERMANY

⁽Torsten Mütze) Department of Computer Science, University of Warwick, United Kingdom & Department of Theoretical Computer Science and Mathematical Logic, Charles University, Prague, Czech Republic

E-mail addresses: gregor@ktiml.mff.cuni.cz, merino@math.tu-berlin.de, torsten.mutze@warwick.ac.uk. Key words and phrases. Gray code, permutation, combination, transposition, Hamilton cycle.

An extended abstract of this paper was accepted for presentation at the 39th International Symposium on Theoretical Aspects of Computer Science (STACS 2022).

This work was supported by Czech Science Foundation grant GA 19-08554S, and by German Science Foundation grant 413902284.

all k-permutations by adjacent transpositions, i.e., swaps of two neighboring entries of the permutation; see Figure 1 (a). In this work, we focus on $star\ transpositions$, i.e., swaps of the first entry of the permutation with any later entry. An efficient algorithm for generating permutations by star transpositions was found by Ehrlich, and it is described as Algorithm E in Knuth's book [Knu11, Section 7.2.1.2]; see Figure 1 (b). For any permutation generation algorithm based on transpositions, we can define the $transposition\ graph$ as the graph with vertex set [k], and an edge between i and j if the algorithm uses transpositions between the ith and jth entry of the permutation. Clearly, the transposition graph for adjacent transpositions is a path, whereas the transposition graph for star transpositions is a star (hence the name 'star transposition'). In fact, Kompel'maher and Liskovec [KL75], and independently Slater [Sla78], showed that all k-permutations can be generated for any transposition tree on [k]. Transposition Gray codes for permutations with additional restrictions were studied by Compton and Williamson [CW93] and by Shen and Williams [SW13].

Several known algorithms for permutation generation use operations other than transpositions. Specifically, Zaks [Zak84] presented an algorithm for generating permutations by prefix reversals; see Figure 1 (c). Moreover, Corbett [Cor92] showed that all k-permutations can be generated by cyclic left shifts of any prefix of the permutation by one position; see Figure 1 (d). Another notable result is Sawada and Williams' recent solution [SW20] of the Sigma-Tau problem, proving that all k-permutations can be generated by cyclic left shifts of the entire permutation by one position or transpositions of the first two elements; see Figure 1 (e).

All of the aforementioned results can be seen as explicit constructions of Hamilton paths in the Cayley graph of the symmetric group, generated by different sets of generators (transpositions, reversals, or shifts). It




FIGURE 1. Gray codes for 4-permutations (SJT=Steinhaus-Johnson-Trotter).

is an open problem whether the Cayley graph of the symmetric group has a Hamilton path for any set of generators [RS93]. This is a special case of the well-known open problem whether any connected Cayley graph has a Hamilton path, or even more generally, whether this is the case for any vertex-transitive graph [Lov70].

1.2. Combination generation and the middle levels conjecture. In a computer, (α, β) -combinations can be conveniently represented by bitstrings of length $n := \alpha + \beta$, where the ith bit is 1 if the element i is in the set and 0 otherwise. For example, the (5,3)-combination $\{1,6,7\}$ is represented by the string 10000110.

In the 1980s, Buck and Wiedemann [BW84] conjectured that all (α, α) -combinations can be generated by star transpositions for every $\alpha \geq 1$, i.e., in every step we swap the first bit of the bitstring representation with a later bit. Figure 2 (a) shows such a star transposition Gray code for (4,4)-combinations. Buck and Wiedemann's conjecture was raised independently by Havel [Hav83], as a question about the existence of a Hamilton cycle through the middle two




FIGURE 2. Star transposition Gray codes for (a) (4,4)- and (b) (2,2,2)-multiset permutations. The strings are arranged in clockwise order, starting at 12 o'clock, with the first entry on the inner track, and the last entry on the outer track. As every star transposition changes the first entry, the color on the inner track changes in every step.

levels of the $(2\alpha-1)$ -dimensional hypercube. This conjecture became known as *middle levels* conjecture, and it attracted considerable attention in the literature and made its way into popular books [Win04, DG12], until it was answered affirmatively by Mütze [Müt16]; see also [GMN18]. Similarly to permutations, there are also many known methods for generating general (α, β) -combinations that use operations other than star transpositions, see [TL73, EM84, Cha89, JM95, RW09]. In particular, (α, β) -combinations can be generated by adjacent transpositions if and only if $\alpha = 1$, $\beta = 1$, or α and β are both odd [BW84, EHR84, Rus88].

1.3. **Multiset permutations.** Shen and Williams [SW21] proposed a far-ranging generalization of the middle levels conjecture that connects permutations and combinations. Their conjecture is about multiset permutations. For integers $k \geq 2$ and $a_1, \ldots, a_k \geq 1$, an (a_1, \ldots, a_k) -multiset permutation is a string over the alphabet $\{1, \ldots, k\}$ that contains exactly a_i occurrences of the symbol i. We refer to the sequence $\mathbf{a} := (a_1, \ldots, a_k)$ as the frequency vector, as it specifies the frequency of each symbol. The length of a multiset permutation is $n_{\mathbf{a}} := a_1 + \cdots + a_k$, and if the context is clear we omit the index and simply write $n = n_{\mathbf{a}}$. If all symbols appear equally often, i.e., $a_1 = \cdots = a_k = \alpha$, we use the abbreviation $\alpha^k := (a_1, \ldots, a_k)$. For example 123433153 is a (2, 1, 4, 1)-multiset permutation, and 331232142144 is a 3^4 -multiset permutation.

Clearly, multiset permutations are a generalization of permutations and combinations. Specifically, k-permutations are 1^k -multiset permutations, and (α, β) -combinations are (α, β) -multiset permutations (up to shifting the symbol names $1, 2 \mapsto 0, 1$). Stachowiak [Sta92] showed that




FIGURE 3. Star transposition graphs G(a) for several small multiset permutations a. Vertices are colored according to the first entry of the multiset permutations, and these color classes form independent sets. In G(2,1,1), an odd cycle is highlighted.

 (a_1, \ldots, a_k) -multiset permutations can be generated by adjacent transpositions if and only if at least two of the a_i are odd.

Shen and Williams [SW21] conjectured that all α^k -multiset permutations can be generated by star transpositions, for any $\alpha \geq 1$ and $k \geq 2$. We state their conjecture in terms of Hamilton cycles in a suitably defined graph, as follows. We write $\Pi(\mathbf{a}) = \Pi(a_1, \ldots, a_k)$ for the set of all (a_1, \ldots, a_k) -multiset permutations. Moreover, we let $G(\mathbf{a}) = G(a_1, \ldots, a_k)$ denote the graph on the vertex set $\Pi(\mathbf{a}) = \Pi(a_1, \ldots, a_k)$ with an edge between any two multiset permutations that differ in a star transposition, i.e., in swapping the first entry of the multiset permutation with any entry at positions $2, \ldots, n$ that is distinct from the first one. Figure 3 shows various examples of the graph $G(\mathbf{a})$. When denoting specific multiset permutations we sometimes omit commas and brackets for brevity, for example $1312214 \in \Pi(3, 2, 1, 1)$.

Conjecture 1 ([SW21]). For any $\alpha \geq 1$ and $k \geq 2$, the graph $G(\alpha^k)$ has a Hamilton cycle.

In this and the following statements, the single edge G(1,1) is also considered a cycle, as it gives a cyclic Gray code. Note that $G(a_1,\ldots,a_k)$ is vertex-transitive if and only if $a_1 = \cdots = a_k =: \alpha$. In this case, Conjecture 1 is an interesting instance of the aforementioned conjecture of Lovász [Lov70] on Hamilton paths in vertex-transitive graphs.

Evidence for Conjecture 1 comes from the results mentioned in Sections 1.1 and 1.2 on generating permutations by star transpositions and the solution of the middle levels conjecture, respectively, formulated in terms of the graph G(a) below. These known results settle the boundary cases $\alpha = 1$ and $k \geq 2$, and $\alpha \geq 1$ and k = 2, respectively, of Conjecture 1.

Theorem 2 (Ehrlich; [KL75]; [Sla78]). For any $k \geq 2$, the graph $G(1^k)$ has a Hamilton cycle.

Theorem 3 ([Müt16, GMN18]). For any $\alpha \geq 1$, the graph $G(\alpha, \alpha)$ has a Hamilton cycle.

In their paper, Shen and Williams also provided an ad-hoc solution for the first case of their conjecture that is not covered by Theorems 2 and 3, namely a Hamilton cycle in G(2,2,2), which is displayed in Figure 2 (b).

We approach Conjecture 1 by tackling the following even more general question: For which frequency vectors $\mathbf{a} = (a_1, \dots, a_k)$ does the graph $G(\mathbf{a})$ have a Hamilton cycle? By renaming symbols, we may assume w.l.o.g. that the entries of the vector \mathbf{a} are non-increasing, i.e.,

$$a_1 \ge a_2 \ge \dots \ge a_k. \tag{1}$$

We can thus think of the vector \boldsymbol{a} as an integer partition of n.

1.4. Our results. For any $i \in [n]$ and $c \in [k]$, we write $\Pi(\mathbf{a})^{i,c}$ for the set of all multiset permutations from $\Pi(\mathbf{a})$ whose ith symbol equals c. Note that every star transposition changes the first entry; see the inner track of each of the two wheels in Figure 2. As a consequence, $G(\mathbf{a})$ is a k-partite graph with partition classes $\Pi(\mathbf{a})^{1,1}, \ldots, \Pi(\mathbf{a})^{1,k}$; see Figure 3. Moreover, the partition class $\Pi(\mathbf{a})^{1,1}$ is a largest one because of (1). This k-partition of the graph $G(\mathbf{a})$ is a potential obstacle for the existence of Hamilton cycles and paths. Specifically, if one partition class is larger than all others combined, then there cannot be a Hamilton cycle, and if the size difference is more than 1, then there cannot be a Hamilton path.

We capture this by defining a parameter $\Delta(a)$ for any integer partition $a = (a_1, \dots, a_k)$ as

$$\Delta(\boldsymbol{a}) := n - 2a_1 = \left(\sum_{i=2}^k a_i\right) - a_1. \tag{2}$$

We will see that if $\Delta(\boldsymbol{a}) < 0$, then the partition class $\Pi(\boldsymbol{a})^{1,1}$ of the graph $G(\boldsymbol{a})$ is larger than all others combined, excluding the existence of Hamilton cycles. On the other hand, if $\Delta(\boldsymbol{a}) \geq 0$, then every partition class of the graph $G(\boldsymbol{a})$ is at most as large as all others combined (equality holds if $\Delta(\boldsymbol{a}) = 0$), which does not exclude the existence of a Hamilton cycle. The cases with $\Delta(\boldsymbol{a}) = 0$ lie on the boundary between the two regimes, and they are the hardest in terms of proving Hamiltonicity. These cases can be seen as generalizations of the middle levels conjecture, namely the case $\boldsymbol{a} = (\alpha, \alpha)$ captured by Theorem 3, which also satisfies $\Delta(\boldsymbol{a}) = 0$.

Theorem 4. For any integer partition $\mathbf{a} = (a_1, \dots, a_k)$ with $\Delta(\mathbf{a}) < 0$ the graph $G(\mathbf{a})$ does not have a Hamilton cycle, and it does not have a Hamilton path unless $\mathbf{a} = (2, 1)$.

For k = 2 symbols, the condition $\Delta(\mathbf{a}) < 0$ is equivalent to $a_1 > a_2$, i.e., there is no star transposition Gray code for 'unbalanced' combinations.

We now discuss the cases $\Delta(\boldsymbol{a}) \geq 0$. Our first main goal is to reduce all cases with $\Delta(\boldsymbol{a}) > 0$ to cases with $\Delta(\boldsymbol{a}) = 0$. For doing so, it is helpful to consider stronger notions of Hamiltonicity. Specifically, we consider Hamilton-connectedness and Hamilton-laceability, which have been heavily studied (see [Sim78, CQ81, HL87, HCH00, AQ01, Ara06]). A graph is called *Hamilton-connected* if there is a Hamilton path between any two distinct vertices. A bipartite graph is called *Hamilton-laceable* if there is a Hamilton path between any pair of vertices from the two partition classes. In general, the graphs $G(\boldsymbol{a})$ are not bipartite, so we say that $G(\boldsymbol{a})$ is 1-laceable if there is a Hamilton path between any vertex in $\Pi(\boldsymbol{a})^{1,1}$ and any vertex not in $\Pi(\boldsymbol{a})^{1,1}$, i.e., between any vertex with first symbol 1 and any vertex with first symbol distinct from 1.

This approach is inspired by the following result of Tchuente [Tch82], who strengthened Theorem 2 considerably.

Theorem 5 ([Tch82]). For any $k \geq 4$, the graph $G(1^k)$ is Hamilton-laceable.

The key insight is that proving a stronger property makes the proof easier and shorter, because the inductive statement is more powerful and flexible; see Section 1.5.2 below. Encouraged by this, we raise the following conjecture about graphs $G(\mathbf{a})$ with $\Delta(\mathbf{a}) = 0$. It is another natural and far-ranging generalization of the middle levels conjecture, which we support by extensive computer experiments and by proving some special cases.

Conjecture 6. For any integer partition $\mathbf{a} = (a_1, \dots, a_k)$ with $\Delta(\mathbf{a}) = 0$ the graph $G(\mathbf{a})$ is Hamilton-1-laceable, unless $\mathbf{a} = (2, 2)$.

The exceptional graph G(2,2) mentioned in this conjecture is a 6-cycle; see Figure 3. Assuming the validity of this conjecture, we settle all cases $G(\boldsymbol{a})$ with $\Delta(\boldsymbol{a}) > 0$ in the strongest possible sense. While being a conditional result, the main purpose of this theorem is to reduce all cases $\Delta(\boldsymbol{a}) \geq 0$ to the boundary cases $\Delta(\boldsymbol{a}) = 0$.

Theorem 7. Conditional on Conjecture 6, for any integer partition $\mathbf{a} = (a_1, \dots, a_k)$ with $\Delta(\mathbf{a}) > 0$ the graph $G(\mathbf{a})$ is Hamilton-connected, unless $\mathbf{a} = (1, 1, 1)$ or $\mathbf{a} = 1^k$ for $k \ge 4$, and possibly unless $\mathbf{a} = (\alpha, \alpha, 1)$ for $\alpha \ge 3$.

The dependence of Theorem 7 on Conjecture 6 can be captured more precisely. Specifically, $G(\mathbf{a})$ with $\Delta(\mathbf{a}) > 0$ is shown to be Hamilton-connected, assuming that $G(\mathbf{b})$ with $\Delta(\mathbf{b}) = 0$ is Hamilton-1-laceable for all integer partitions \mathbf{b} that are majorized componentwise by \mathbf{a} .

The three exceptions mentioned in Theorem 7 are well understood: Specifically, G(1,1,1) is a 6-cycle; see Figure 3. Furthermore, $G(1^k)$ for $k \geq 4$ is Hamilton-laceable by Theorem 5. Lastly, we will show that $G(\alpha, \alpha, 1)$ for $\alpha \geq 3$ satisfies a variant of Hamilton-laceability, which also guarantees a Hamilton cycle. In fact, we believe that $G(\alpha, \alpha, 1)$ is Hamilton-connected, but we cannot prove it.

We provide the following evidence for Conjecture 6. First of all, with computer help we verified that $G(\boldsymbol{a})$ is indeed Hamilton-1-laceable for all integer partitions $\boldsymbol{a} \neq (2,2)$ with $\Delta(\boldsymbol{a}) = 0$ that satisfy $n \leq 8$, i.e., for $\boldsymbol{a} \in \{(1,1),(2,1,1),(3,3),(3,2,1),(3,1,1,1),(4,4),(4,3,1),(4,2,2),(4,2,1,1),(4,1,1,1)\}$. Furthermore, we prove the case of k=2 symbols unconditionally. Note that for k=2, Hamilton-1-laceability is the same as Hamilton-laceability. Recall that $G(\alpha,\alpha)$ is isomorphic to the subgraph of the $(2\alpha-1)$ -dimensional hypercube induced by the middle two levels, so the following result is a considerable strengthening of Theorem 3, the middle levels theorem.

Theorem 8. For any $\alpha \geq 3$, the graph $G(\alpha, \alpha)$ is Hamilton-laceable.

We also have the following (unconditional) result for k=3 symbols.

Theorem 9. For any $\alpha \geq 2$, the graph $G(\alpha, \alpha - 1, 1)$ has a Hamilton cycle.

Lastly, we consider integer partitions $\mathbf{a} = (a_1, \dots, a_k), k \geq 3$, with $\Delta(\mathbf{a}) \geq 0$ and an upper bound on the part size, i.e., $a_1 \leq \alpha$ for some constant α . By the remarks after Theorem 7, the inductive proof of the theorem for such integer partitions only relies on Conjecture 6 being satisfied for integer partitions with the same upper bound on the part size. For any fixed bound α , there are only finitely many such partitions with $\Delta(\mathbf{a}) = 0$ that can be checked by computer. For example, for $\alpha = 4$ these are $\mathbf{a} \in \{(2,1,1),(3,2,1),(3,1,1,1),(4,3,1),(4,2,2),(4,2,1,1),(4,1,1,1,1)\}$. This yields the following (unconditional) result.

Theorem 10. For $\alpha \in \{2,3,4\}$ and any integer partition $\mathbf{a} = (a_1, \ldots, a_k)$ with $\Delta(\mathbf{a}) > 0$ and $a_1 = \alpha$, the graph $G(\mathbf{a})$ is Hamilton-connected.

In words, Theorem 10 settles all integer partitions a whose largest part a_1 is at most 4. In particular, this settles the cases $\alpha \in \{2, 3, 4\}$ and $k \geq 2$ of Shen and Williams' Conjecture 1 in a rather strong sense.

- 1.5. **Proof ideas.** In this section, we give a high-level overview of the main ideas and techniques used in our proofs.
- 1.5.1. The case $\Delta(\boldsymbol{a}) < 0$. The main idea for proving Theorem 4 is that if $\Delta(\boldsymbol{a}) < 0$, then the partition class $\Pi(\boldsymbol{a})^{1,1}$ of the graph $G(\boldsymbol{a})$ is larger than all others combined, which excludes the existence of a Hamilton cycle. To exclude the existence of a Hamilton path, we show that the size difference is strictly more than 1, unless $\boldsymbol{a} = (2,1)$. Note that the graph G(2,1) is the path on three vertices, so in this case the size difference is precisely 1. These arguments are based on straightforward algebraic manipulations involving multinomial coefficients.
- 1.5.2. The case $\Delta(\boldsymbol{a}) > 0$. To prove Theorem 7, it is convenient to think of an integer partition $\boldsymbol{a} = (a_1, \ldots, a_k)$ as an infinite non-increasing sequence (a_1, a_2, \ldots) , with only k nonzero entries at the beginning. Given two such integer partitions $\boldsymbol{a} = (a_1, a_2, \ldots)$ and $\boldsymbol{b} = (b_1, b_2, \ldots)$, we write $\boldsymbol{b} \prec \boldsymbol{a}$ if $b_i \leq a_i$ for all $i \geq 1$. Integer partitions with the partial order \prec form a lattice, which is the sublattice of the infinite lattice $\mathbb{N}^{\mathbb{N}}$ cut out by the hyperplanes defined by (1); see Figure 4. The cover relations in this lattice are given by decrementing any of the a_i for which $a_i > a_{i+1}$. We write $\boldsymbol{b} \prec \boldsymbol{a}$ for partitions $\boldsymbol{a} \neq \boldsymbol{b}$ if $\boldsymbol{b} \prec \boldsymbol{a}$ and there is no $\boldsymbol{c} \notin \{\boldsymbol{a}, \boldsymbol{b}\}$ with $\boldsymbol{b} \prec \boldsymbol{c} \prec \boldsymbol{a}$.

In this lattice of integer partitions, the hyperplane defined by $\Delta(\mathbf{a}) = 0$ separates the cases where Hamiltonicity is impossible, which lie on the side of the hyperplane where $\Delta(\mathbf{a}) < 0$ (Theorem 4), from the cases where Hamiltonicity can be established more easily, which lie on the side of the hyperplane where $\Delta(\mathbf{a}) > 0$ (Theorem 7). The cases $\Delta(\mathbf{a}) = 0$ on the hyperplane are the hardest ones (Conjecture 6).

Our proof of Theorem 7 proceeds by induction in this partition lattice and establishes the Hamiltonicity of G(a) by using the Hamiltonicity of G(b) for all integer partitions $b \prec a$, where Conjecture 6 serves as the base case of the induction. This is based on the observation that fixing one of the symbols at positions $2, \ldots, n$ in G(a) yields subgraphs that are isomorphic to G(b) for $b \prec a$.

Specifically, for any $\mathbf{a} = (a_1, \dots, a_k)$, $i = 2, \dots, n$, and $c \in [k]$, the subgraph of $G(\mathbf{a})$ induced by the vertex set $\Pi(\mathbf{a})^{i,c}$ is isomorphic to $G(\mathbf{b})$ where \mathbf{b} is the partition obtained from \mathbf{a} by decreasing a_i by 1 (and possibly sorting the resulting numbers non-increasingly); see Figure 5.

Moreover, for any $\mathbf{b} \prec \mathbf{a}$ we have $\Delta(\mathbf{b}) = \Delta(\mathbf{a}) - 1$ or $\Delta(\mathbf{b}) = \Delta(\mathbf{a}) + 1$. In particular, if $\Delta(\mathbf{a}) > 0$, then we have $\Delta(\mathbf{b}) \geq 0$. For example, the vertex set of $G(\mathbf{a})$ for $\mathbf{a} = (3, 2, 2)$ ($\Delta(\mathbf{a}) = 1$) can be partitioned into one copy of $G(\mathbf{b})$ for $\mathbf{b} = (2, 2, 2)$ (if the fixed symbol is c = 1; $\Delta(\mathbf{b}) = 2$) and two copies of $G(\mathbf{b}')$ for $\mathbf{b}' = (3, 2, 1)$ (if the fixed symbol is c = 2 or c = 3; $\Delta(\mathbf{b}') = 0$). Therefore, we may construct a Hamilton path in G(3, 2, 2) by gluing together paths in each of these three subgraphs which exist by induction.




FIGURE 5. Decomposing G(2,1,1) into three subgraphs by fixing the last symbol.

While conceptually simple, implementing this

idea incurs considerable technical obstacles, in particular for some of the graphs G(a) with $\Delta(a) =$




FIGURE 4. The lattice of integer partitions $\mathbf{a} = (a_1, \dots, a_k)$ with largest part $a_1 \leq 3$. The coordinates are projected into three dimensions depending on which value is increased.

1, i.e., instances that are very close to the hyperplane $\Delta(a) = 0$. The proof is split into several interdependent lemmas, and it is the technically most demanding part of our paper.

Theorem 10 follows immediately from the inductive proof of Theorem 7 and by settling finitely many cases with computer help.

To further illustrate the ideas outlined before, we close this section by reproducing Tchuente's proof of Theorem 5. To prove that $G(1^k)$ is Hamilton-laceable, we proceed by induction on k. The induction basis k=4 can be checked by straightforward case analysis. For the induction step, we assume that $G(1^{k-1})$, $k \geq 5$, is Hamilton-laceable, and we prove that $G(1^k)$ is also




FIGURE 6. Illustration of the proof of Theorem 5. The left hand side shows the general schematic partitioning of the graph $G(1^k)$ into blocks, each of which is a copy of $G(1^{k-1})$, by fixing symbols at position $\hat{\imath}$. The right hand side shows a concrete example for k=7.

Hamilton-laceable. Note that $1^{k-1} \prec 1^k$ and $\Delta(1^k) = k - 2 = \Delta(1^{k-1}) + 1$. The following arguments are illustrated in Figure 6. The partition classes of the graph $G(1^k)$ are given by the parity of the permutations, i.e., by the number of inversions. Therefore, we consider two distinct permutations x and y of [k] with opposite parity, and we need to show how to connect them by a Hamilton path in $G(1^k)$. As $x \neq y$, there is a position $\hat{i} > 1$ in which x and y differ, i.e., $x_i \neq y_i$, and this is the position that we will fix to different symbols. Specifically, we choose a permutation π of [k] such that $\pi_1 = x_i$ and $\pi_k = y_i$. The permutation π captures the order in which we will fix symbols at position \hat{i} . We then choose permutations u^j, v^j of $[k], j = 1, \dots, k$, satisfying $u^1 = x$, $v^k = y$, and such that u^j is obtained from v^{j-1} by a star transposition of the symbol π_i at position 1 with the symbol π_{i-1} at position $\hat{\imath}$, for all $j=2,\ldots,k$. Moreover, we choose u^j and v^j such that the parity of the number of inversions after removing the symbol π_i is opposite. Specifically, for u^j this parity is the same as for $u^1 = x$ if and only if π_i has the same parity as π_1 , and for v^j this parity is the same as for $v^k = y$ if and only if π_i has the same parity as π_k . For each $j=1,\ldots,k$, we now consider the permutations whose \hat{i} th entry equals π_i (formally, this is the set $\Pi(1^k)^{\hat{i},\pi_j}$). Clearly, the subgraph of $G(1^k)$ induced by these permutations is isomorphic to $G(1^{k-1})$. In other words, by removing the \hat{i} th entry and renaming entries to $1, \ldots, k-1$, we obtain permutations of [k-1]. Consequently, by induction there is a path in $G(1^k)$ that visits all permutations whose îth entry equals π_i and that connects u^j to v^j . The concatenation of those k paths obtained by induction is the desired Hamilton path in $G(1^k)$ from x to y, which completes the induction step.

Note that the constraints imposed on the permutations π and $u^j, v^j, j = 1, \ldots, k$, in this proof are very mild, and leave a lot of room for modifications to construct many different Hamilton paths, possibly so as to satisfy some additional conditions.

1.5.3. The case $\Delta(\mathbf{a}) = 0$. Our proofs of Theorems 8 and 9 build on ideas introduced in the papers [GMN18, MMM21].

Specifically, the first step in proving Theorem 8 is to build a cycle factor in the graph $G(\alpha, \alpha)$, i.e., a collection of disjoint cycles in the graph that together visit all vertices. We then choose vertices x and y from the two partition classes of the graph that we want to connect by a Hamilton path. In this we can take into account automorphisms of $G(\alpha, \alpha)$, i.e., for proving laceability only certain pairs of vertices x and y in the two partition classes have to be considered. In the next step, we join a small subset of cycles from the factor, including the ones containing x and y, to a short path between x and y. This is achieved by taking the symmetric difference of the edge set of the cycle factor with a carefully chosen path P from x to y that alternately uses edges on one of the cycles from the factor and edges that go between two such cycles; see Figure 7 (a)+(b). In the last step, we join the remaining cycles of the factor to the path between x and y, until we end with a Hamilton path from x to y. Each such joining is achieved by taking the symmetric difference of the cycle factor with a suitably chosen 6-cycle; see Figure 7 (b)+(c).




Figure 7. Strategy of the proof of Theorem 8

It was shown in [GMN18] that the cycles of the aforementioned cycle factor in $G(\alpha, \alpha)$ are bijectively equivalent to plane trees with α vertices, and the joining operations via 6-cycles can be interpreted combinatorially as local change operations between two such plane trees. To prove Theorem 9, we first generalize the construction of this cycle factor in the graph $G(\alpha, \alpha)$ to a cycle factor in any graph $G(\mathbf{a})$, $\mathbf{a} = (a_1, \dots, a_k)$, with $\Delta(\mathbf{a}) = 0$. It turns out that the cycles of this generalized factor can be interpreted combinatorially as vertex-labeled plane trees, where exactly a_i vertices have the label i for $i = 2, \dots, k$. If $\mathbf{a} = (\alpha, \alpha)$, then all vertex labels are the same, so we can consider the trees as unlabeled. Moreover, the joining 6-cycles in $G(\alpha, \alpha)$ generalize nicely to joining 12- or 6-cycles in $G(\mathbf{a})$ with $\Delta(\mathbf{a}) = 0$, and they correspond to local change operations involving sets T of labeled plane trees with $|T| \in \{2, 3, 5, 6\}$, depending on the location of vertex labels. For proving Theorem 9, we consider the special case $\mathbf{a} = (\alpha, \alpha - 1, 1)$, i.e., exactly one vertex in the plane trees is labeled differently from all other vertices, and we show that there is a choice of joining cycles so that the symmetric difference with the cycle factor yields a Hamilton cycle in the graph $G(\mathbf{a})$.

1.6. Outline of this paper. In Section 2 we collect definitions and observations that will be used throughout this paper. The proof of Theorem 4 is presented in Section 3. In Section 4, we prove Theorems 7 and 10. The proofs of Theorems 8 and 9 are presented in Section 5. This section can be read independently from the previous sections, but it relies on results from the previous papers [GMN18, MMM21]. We conclude in Section 6 with some open questions and possible directions for future work.

2. Preliminaries

The following definitions and lemmas will be used repeatedly in this paper.

2.1. Fixing symbols. For any sequence of non-negative integers $\mathbf{a} = (a_1, a_2, \dots, a_k)$, we let $\varphi(a)$ be the partition obtained by sorting the sequence \mathbf{a} non-increasingly. For instance, we have $\varphi((2,3,1,1,4)) = (4,3,2,1,1)$.

As mentioned before, $\Pi(\boldsymbol{a})^{i,c}$ is the set of all multiset permutations from $\Pi(\boldsymbol{a})$ whose ith symbol equals c. We write $G(\boldsymbol{a})^{i,c} := G(\boldsymbol{a})[\Pi(\boldsymbol{a})^{i,c}]$ for the subgraph of $G(\boldsymbol{a})$ induced by the vertex set $\Pi(\boldsymbol{a})^{i,c}$. We observed before that $\Pi(a)^{1,c}$ is an independent set for all $c \in [k]$, and that $G(\boldsymbol{a})^{i,c}$, $i=2,\ldots,n$, is isomorphic to $G(\boldsymbol{b})$ for $\boldsymbol{b}:=\varphi(\boldsymbol{a}') \prec \boldsymbol{a}$, $\boldsymbol{a}':=(a_1,\ldots,a_{c-1},a_c-1,a_{c+1},\ldots,a_k)$. We also allow repeating this operation and define $G(\boldsymbol{a})^{(i,j),(c,d)}:=(G(\boldsymbol{a})^{i,c})^{j,d}$ for $i,j\in[n]$ with $i\neq j$ and $c,d\in[k]$ with $c\neq d$ or $a_c=a_d\geq 2$.

Our first lemma follows directly from the definition (2).

Lemma 11. If $\mathbf{b} \prec \mathbf{a}$, then we have $\Delta(\mathbf{b}) = \Delta(\mathbf{a}) - 1$, except if \mathbf{a} and \mathbf{b} differ in the largest part, in which case $\Delta(\mathbf{b}) = \Delta(\mathbf{a}) + 1$.

2.2. Partition lemmas. The next lemma quantifies the size of the k partition classes of the graph G(a) discussed in Section 1.4.

Lemma 12. For any integer partition $\mathbf{a} = (a_1, \dots, a_k)$, the graph $G(a_1, \dots, a_k)$ is k-partite, with the partition classes $\Pi(\mathbf{a})^{1,1}, \dots, \Pi(\mathbf{a})^{1,k}$ given by fixing the first symbol, and the size of the ith partition class is

$$s_i := |\Pi(\boldsymbol{a})^{1,i}| = \binom{n-1}{a_1, \dots, a_{i-1}, a_i - 1, a_{i+1}, \dots, a_k} = \frac{(n-1)!}{a_1! \cdots a_{i-1}! (a_i - 1)! a_{i+1}! \cdots a_k!}.$$
 (3)

Moreover, we have $s_1 \ge \cdots \ge s_k$, in particular, the first partition class is a largest one. Lastly, if $\Delta(\mathbf{a}) < 0$, then the first partition class is larger than all others combined, if $\Delta(\mathbf{a}) = 0$ then the first partition class has the same size as all others combined, and if $\Delta(\mathbf{a}) > 0$ then every partition class is smaller than all others combined.

Proof. As every star transposition changes the first symbol, the sets $\Pi(\mathbf{a})^{1,1}, \ldots, \Pi(\mathbf{a})^{1,k}$ are independent sets in the graph $G(\mathbf{a})$. The multinomial coefficient in (3) describes the number of ways of arranging n-1 symbols at positions $2, \ldots, n$, with only a_i-1 occurrences of the symbol i remaining (one symbol i is fixed at the first position). This proves the first part of the lemma. The monotonicity $s_1 \geq \cdots \geq s_k$ follows immediately from (1) and (3). Using the definition (3), the equation $s_1 > \sum_{i=2}^k s_i$ can be rearranged to $2a_1 > n$, which by the definition (2) is equivalent to $\Delta(\mathbf{a}) < 0$. Reversing the direction of inequalities or replacing them by equality completes the proof of the second part of the lemma.

The following result complements Lemma 12 by showing that some of the graphs G(a) are not only k-partite, but also bipartite even for $k \geq 3$ distinct symbols. An *inversion* in a permutation is a pair of entries where the first entry is larger than the second.

Lemma 13. For any integer partition $\mathbf{a} = (a_1, \dots, a_k)$ with $k \geq 3$ parts, the graph $G(\mathbf{a})$ is not bipartite unless $a_1 = \dots = a_k = 1$. If $a_1 = \dots = a_k = 1$, then the two partition classes are given by the parity of the number of inversions of the permutations, and they have the same size.

Proof. Note that if $\mathbf{b} \prec \mathbf{a}$, then $G(\mathbf{b})$ is a subgraph of $G(\mathbf{a})$. Specifically, if $b_i < a_i$, then we may fix $a_i - b_i$ occurrences of i on one of the positions $2, \ldots, n_{\mathbf{a}}$ in the graph $G(\mathbf{a})$. Consequently,

unless $a_1 = \cdots = a_k = 1$, the graph G(2, 1, 1) is a subgraph of $G(\boldsymbol{a})$. As G(2, 1, 1) contains a 7-cycle (highlighted in Figure 3), a graph containing it as a subgraph is not bipartite.

We proceed to show that $G(1^k)$ is bipartite. This follows directly from the observation that every transposition of two entries of a permutation changes the parity of the number of inversions. Moreover, permuting the first two entries is a bijection between permutations with an even or odd number of inversions, so both partition classes have the same size.

The next lemma shows that Hamilton paths in $G(\mathbf{a})$ with $\Delta(\mathbf{a}) = 0$ have a very rigid structure.

Lemma 14. Let $\mathbf{a} = (a_1, \dots, a_k)$ be an integer partition with $\Delta(\mathbf{a}) = 0$. Then on every Hamilton path in $G(\mathbf{a})$ whose end vertices differ in the first entry, every second vertex of the path is from $\Pi(\mathbf{a})^{1,1}$, and every other vertex on the path is not from $\Pi(\mathbf{a})^{1,1}$.

In terms of star transpositions, this means that the symbol 1 is swapped in every step, either in or out of the first position.

Proof. By Lemma 12, the partition class $\Pi(\boldsymbol{a})^{1,1}$ of the graph $G(\boldsymbol{a})$ has the same size as all others combined. Consequently, one of the end vertices of any Hamilton path must be in this largest class. By the assumption that the other end vertex is not in $\Pi(\boldsymbol{a})^{1,1}$, the Hamilton path is forced to visit $\Pi(\boldsymbol{a})^{1,1}$ in every second step.

2.3. **Hamiltonicity notions.** In our proofs we will use Hamilton-connectedness, Hamilton-laceability, and Hamilton-1-laceability heavily, and also other strengthenings of the notion of containing a Hamilton cycle. It will be convenient to introduce shorthand notations for graphs satisfying these properties.

To this end, we let \mathcal{G} denote the family of all graphs $G(\mathbf{a})$ for any integer partition $\mathbf{a} = (a_1, \ldots, a_k)$, where $k \geq 2$ and $a_1 \geq \cdots \geq a_k \geq 1$. We also define the following subsets of \mathcal{G} :

- \mathcal{P} All graphs in \mathcal{G} that have a Hamilton path, i.e., a path that visits every vertex exactly once.
- \mathcal{C} All graphs in \mathcal{G} that have a Hamilton cycle, i.e., a cycle that visits every vertex exactly once.
- \mathcal{E} All graphs in \mathcal{G} such that for every edge of the graph, there is a Hamilton cycle containing this edge. ¹
- \mathcal{L} All graphs in \mathcal{G} that are Hamilton-laceable, i.e., bipartite graphs such that for any two vertices from the two partition classes, there is a Hamilton path between them.
- \mathcal{L}_1 All graphs in \mathcal{G} that are Hamilton-1-laceable, i.e., for which there is a Hamilton path between any vertex in $\Pi(\boldsymbol{a})^{1,1}$ and any vertex not in $\Pi(\boldsymbol{a})^{1,1}$. Recall that $\Pi(\boldsymbol{a})^{1,1}$ are the multiset permutations whose first symbol is 1.
- \mathcal{H} All graphs in \mathcal{G} that are Hamilton-connected, i.e., for which there is a Hamilton path between any two distinct vertices.

To handle the graphs $G(\boldsymbol{a})$ with $\boldsymbol{a}=(\alpha,\alpha,1)$ in our proofs, we need yet another Hamiltonicity notion. While we conjecture that these graphs $G(\boldsymbol{a})$ are Hamilton-connected, we are unable to prove this based on Conjecture 6. This is because all integer partitions \boldsymbol{b} with $\boldsymbol{b} \prec \boldsymbol{a}$, namely $\boldsymbol{b}=(\alpha,\alpha-1,1)$ and $\boldsymbol{b}=(\alpha,\alpha)$ satisfy $\Delta(\boldsymbol{a})=0$, which makes these graphs $G(\boldsymbol{a})$

position			$(\hat{\imath} > 1)$			
1	$\hat{\imath}$	1	$\hat{\imath}$	1	$\hat{\imath}$	$\ell \in \{3,4\}$
1	ℓ	1	ℓ	1	2	s p
1	1	ℓ	1	2	1	$m{t}$ q
2	ℓ	2	ℓ	2	1	
2	2	ℓ	2	1	2	
	-	-	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	•	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$

FIGURE 8. Definition of \mathcal{L}_{12} .

particularly difficult to deal with. We will content ourselves by showing that $G(\mathbf{a})$ satisfies the following variant of Hamilton-laceability, which we denote by \mathcal{L}_{12} .

¹This property is called 'positively Hamiltonian' in [HH72].

Roughly speaking, the set \mathcal{L}_{12} contains the graphs from \mathcal{G} that admit Hamilton paths between pairs of vertices x and y with particular combinations of first symbols s and t, subject to the constraint that another position $\hat{i} > 1$ contains a particular combination of values p and q. The required combinations of symbols are summarized in Figure 8, and they are formally defined as follows. For $\ell \in \{3,4\}$ and any $s \in \{1,2\}$ and $t \in \{1,2,\ell\}$ we define

$$p_{\ell}(s,t) := \begin{cases} \ell & \text{if } t \in \{s,\ell\}, \\ t & \text{otherwise,} \end{cases} \quad \text{and} \quad q_{\ell}(s,t) := s.$$
 (4)

The set \mathcal{L}_{12} contains all graphs in \mathcal{G} for which there is a Hamilton path between any vertex x with first symbol $s \in \{1, 2\}$ and any vertex y distinct from x with first symbol $t \in \{1, 2, \ell\}$, $\ell \in \{3, 4\}$, for which there is a position $\hat{i} > 1$ such that $(x_i, y_i) = (p_{\ell}(s, t), q_{\ell}(s, t))$.

Even though the vertices of $G(\alpha, \alpha, 1)$ have no symbol 4, we still allow $\ell = 4$ in this definition, as this graph may arise for example from $G(\alpha, \alpha, 1, 1)$ by fixing the symbol 3, yielding a graph with symbols $\{1, 2, 4\}$ that is isomorphic to $G(\alpha, \alpha, 1)$.

The obvious containment relations that follow from these definitions are shown in Figure 9. It is not clear whether $\mathcal{L}_1 \subseteq \mathcal{E}$, because for $k \geq 3$ a graph in \mathcal{L}_1 has edges between the partition classes where the first symbol is distinct from 1. It is also not clear whether $\mathcal{L} \subseteq \mathcal{L}_1$, because of the graphs $G(1^k)$, $k \geq 3$, which by Lemma 13 are bipartite but these partition classes are *not* given by fixing the first symbol. Recall that we consider a single edge, i.e., the graph G(1,1), as a cycle (otherwise $\mathcal{L} \subseteq \mathcal{E}$ and $\mathcal{L}_1 \subseteq \mathcal{C}$ would be violated).




FIGURE 9. Hamiltonicity notions ordered by inclusion.

3. Proof of Theorem 4

In this section we present the proof of Theorem 4, implementing the idea outlined in Section 1.5.1.

Proof of Theorem 4. If $\Delta(\mathbf{a}) < 0$, then by Lemma 12, the partition class $\Pi(\mathbf{a})^{1,1}$ of the graph $G(\mathbf{a})$ is larger than all others combined, so there cannot be a Hamilton cycle. This proves the first part of the theorem.

To prove the second part, we first show that if $\Delta(\boldsymbol{a}) < 0$, then the sizes of the partition classes defined in (3) satisfy $s_1 - \sum_{i=2}^k s_i > 1$, unless $\boldsymbol{a} = (2,1)$. Clearly, if one partition class of $G(\boldsymbol{a})$ is by strictly more than 1 larger than all others combined, then there cannot be a Hamilton path. Note also that the graph G(2,1) is the path on three vertices, which indeed has a Hamilton path. We have seen that $\Delta(\boldsymbol{a}) < 0$ implies $s_1 - \sum_{i=2}^k s_i > 0$, and we now argue that this difference equals 1 if and only if $\boldsymbol{a} = (2,1)$. We clearly have

$$\binom{n}{a_1, \dots, a_k} \ge \binom{n}{a_1}. \tag{5}$$

Moreover, from (3) we see that

$$s_1 = \frac{a_1}{n} \binom{n}{a_1, \dots, a_k}. \tag{6}$$

Combining these observations we get

$$1 = s_1 - \sum_{i=2}^k s_i = 2s_1 - \sum_{i=1}^k s_i \stackrel{\text{(3)}}{=} 2s_1 - \binom{n}{a_1, \dots, a_k} \stackrel{\text{(6)}}{=} \frac{2a_1 - n}{n} \binom{n}{a_1, \dots, a_k}$$

$$\stackrel{\text{(5)}}{\geq} \frac{2a_1 - n}{n} \binom{n}{a_1} = (2a_1 - n) \cdot \frac{n-1}{a_1} \cdot \frac{n-2}{a_1 - 1} \cdots \frac{n-a_1 + 1}{2}.$$

The resulting product of fractions is non-decreasing, i.e., we have $\frac{n-1}{a_1} \le \frac{n-2}{a_1-1} \le \cdots \le \frac{n-a_1+1}{2}$ because $n \ge a_1 + 1$. We thus obtain the inequality

$$1 \ge (2a_1 - n) \left(\frac{n-1}{a_1}\right)^{a_1 - 1} \tag{7}$$

As $2a_1 - n = -\Delta(\mathbf{a})$ and $\Delta(\mathbf{a}) < 0$ we have $2a_1 - n > 0$, and as this is an integer we have $2a_1 - n \ge 1$. As $n \ge a_1 + 1$ we also have $\frac{n-1}{a_1} \ge 1$. From (7) we conclude that $2a_1 - n = 1$ and $\frac{n-1}{a_1}=1$, which implies $a_1=2$ and n=3, i.e., $\boldsymbol{a}=(2,1)$. This completes the proof.

4. Proofs of Theorems 7 and 10

In this section we prove Theorems 7 and 10. The proofs are split into several auxiliary lemmas, which we present first. We then describe the computer experiments we did for settling the base cases of our induction proofs. We then present the proofs of Theorems 7 and 10, using the base cases and the lemmas. Finally, we present the proofs of all auxiliary lemmas.

4.1. Auxiliary lemmas. We will use the following auxiliary lemmas; see Figure 10. These lemmas establish the Hamiltonicity of G(a) for all integer partition a with $\Delta(a) \geq 1$, conditional on the Hamiltonicity of G(b) for various integer partitions $b \prec a$ that satisfy $\Delta(b) \geq 0$. The main technical achievement here is to partition all possible cases of frequency vectors a with $\Delta(a) \geq 1$ into disjoint cases, such that the induction proofs of Theorems 7 and 10 which apply these lemmas only rely on previously established cases.

Lemma 15 covers most graphs G(a) with $\Delta(a) \geq 2$, while three special cases of these graphs are covered by Lemmas 17, 18, and 22. Similarly, Lemma 16 covers most graphs G(a)with $\Delta(a) = 1$, while three special cases of these are covered by Lemmas 19, 20 and 21.

Lemma 15. Let $\mathbf{a} = (a_1, \dots, a_k)$ be an integer partition with $\Delta(\mathbf{a}) \geq 2$ with $\mathbf{a} \neq (1, 1, \dots, 1)$, $a \neq (2, 1, 1, ..., 1)$ and $a \notin \{(\alpha, \alpha, 2), (\alpha, \alpha, 1, 1) \mid \alpha \geq 3\}$. If for all $b \prec a$ we have $G(b) \in \mathcal{H}$, then we have $G(\mathbf{a}) \in \mathcal{H}$.

Lemma 16. Let $\mathbf{a} = (a_1, \dots, a_k)$ be an integer partition with $\Delta(\mathbf{a}) = 1$ with $\mathbf{a} \notin \{(1, 1, 1), (3, 1, 1, 1, 1)\}$ and $\mathbf{a} \notin \{(\alpha, \alpha, 1), (\alpha, \alpha - 1, 2), (\alpha, \alpha - 1, 1, 1) \mid \alpha \geq 2\}$. If for all $\mathbf{b} \prec \mathbf{a}$ with $\Delta(\mathbf{b}) = 0$ we have $G(b) \in \mathcal{L}_1$, and for the unique $b \prec a$ with $\Delta(b) = 2$ we have that all $c \prec b$ satisfy $G(c) \in \mathcal{H}$, then we have $G(\mathbf{a}) \in \mathcal{H}$.

Lemma 17. Consider the integer partition $\mathbf{a} = (\alpha, \alpha, 2)$ for $\alpha \geq 3$, which satisfies $\Delta(\mathbf{a}) = 2$. If $\mathbf{b} = (\alpha, \alpha, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}_{12}$ and $\mathbf{b} = (\alpha, \alpha - 1, 2) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{H}$, then we have $G(\boldsymbol{a}) \in \mathcal{H}$.

Lemma 18. Consider the integer partition $\mathbf{a} = (\alpha, \alpha, 1, 1)$ for $\alpha \geq 3$, which satisfies $\Delta(\mathbf{a}) = 2$. If $\mathbf{b} = (\alpha, \alpha, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}_{12}$ and $\mathbf{b} = (\alpha, \alpha - 1, 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{H}$, then we have $G(\mathbf{a}) \in \mathcal{H}$.




FIGURE 10. Relations between our results and auxiliary lemmas, and the different Hamiltonicity notions we establish for them. Arrows indicate dependencies in the proof of Theorem 7.

Lemma 19. Consider the integer partition $\mathbf{a} = (\alpha, \alpha - 1, 2)$ for $\alpha \geq 3$, which satisfies $\Delta(\mathbf{a}) = 1$. If $\mathbf{b} = \varphi((\alpha, \alpha - 2, 2)) \prec \mathbf{a}$ and $\mathbf{b} = (\alpha, \alpha - 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{L}_1$, and if $\mathbf{c} = (\alpha - 1, \alpha - 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{L}_{12}$ and $\mathbf{c} = \varphi((\alpha - 1, \alpha - 2, 2)) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{H}$, then we have $G(\mathbf{a}) \in \mathcal{H}$.

Lemma 20. Consider the integer partition $\mathbf{a} = (\alpha, \alpha - 1, 1, 1)$ for $\alpha \geq 3$, which satisfies $\Delta(\mathbf{a}) = 1$. If $\mathbf{b} = (\alpha, \alpha - 2, 1, 1) \prec \mathbf{a}$ and $\mathbf{b} = (\alpha, \alpha - 1, 1) \prec \mathbf{a}$ satisfy $G(\mathbf{b}) \in \mathcal{L}_1$, and if $\mathbf{c} = (\alpha - 1, \alpha - 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{L}_{12}$ and $\mathbf{c} = (\alpha - 1, \alpha - 2, 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{H}$, then we have $G(\mathbf{a}) \in \mathcal{H}$.

Lemma 21. Consider the integer partition $\mathbf{a} = (\alpha, \alpha, 1)$ for $\alpha \geq 3$, which satisfies $\Delta(\mathbf{a}) = 1$. If $\mathbf{b} = (\alpha, \alpha) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}$ and $\mathbf{b} = (\alpha, \alpha - 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}_1$, then we have $G(\mathbf{a}) \in \mathcal{L}_{12}$.

Lemma 22. Consider the integer partition $\mathbf{a} = (2, 1^{k-1})$ for $k \geq 5$, which satisfies $\Delta(\mathbf{a}) \geq 2$. Then we have $G(\mathbf{a}) \in \mathcal{H}$.

4.2. **Base cases.** We performed computer experiments to settle the base cases for our inductive proofs, and also for collecting evidence for Conjecture 6. The results of these experiments are summarized in Table 1. The second-to-last column contains the number of non-isomorphic pairs of vertices of the graph that need to be checked when testing for \mathcal{L}_1 or \mathcal{H} .

We restrict our attention to integer partitions \mathbf{a} with $\Delta(\mathbf{a}) = 0$ and $\Delta(\mathbf{a}) = 1$. The results in Table 1 confirm Conjecture 6 for all integer partitions $\mathbf{a} \neq (2,2)$ with $\Delta(\mathbf{a}) = 0$ that satisfy $n \leq 8$. Some of these results will be used in our proofs of Theorems 7–10, as indicated in the last column of the table. The last three instances in Table 1 were too large to test for Hamilton-connectedness, but they are Hamilton-connected by Theorem 10.

4.3. **Proof of Theorem 7.** With the help of Lemmas 15–22, proving Theorem 7 is relatively straightforward.

Proof of Theorem 7. This proof is illustrated in Figure 10. Conjecture 6 asserts that for all integer partitions $\mathbf{a} \neq (2,2)$ with $\Delta(\mathbf{a}) = 0$ we have $G(\mathbf{a}) \in \mathcal{L}_1$, and we now assume that this

n	Partition a	$\Delta(a)$	Hamiltonicity of $G(\boldsymbol{a})$	Vertices	Vertex pairs	Used in proof of
2	(1,1)	0	\mathcal{H}	2	1	
3	(1,1,1)	1	\mathcal{E} but not \mathcal{L}	6	2	
4	(2,2)	0	\mathcal{E} but not \mathcal{L}	6	2	
	(2,1,1)	0	\mathcal{L}_1 but not \mathcal{H}	12	6	Thm. 9+10
5	(2,2,1)	1	\mathcal{H}	30	23	Thm. 7+10
	(2,1,1,1)	1	$\mid \mathcal{H} \mid$	60	31	Thm. 7
6	(3,3)	0	$\mathcal{L}_1 = \mathcal{L}$ but not \mathcal{H}	20	3	Thm. 8+10
	(3,2,1)	0	\mathcal{L}_1 but not \mathcal{H}	60	26	Thm. 9+10
	(3,1,1,1)	0	\mathcal{L}_1 but not \mathcal{H}	120	16	Thm. 10
7	(3,3,1)	1	\mathcal{H}	140	38	Thm. 10
	(3,2,2)	1	$\mid \mathcal{H} \mid$	210	65	
	(3,2,1,1)	1	$\mid \mathcal{H} \mid$	420	209	
	(3,1,1,1,1)	1	$\mid \mathcal{H} \mid$	840	79	Thm. 7
8	(4,4)	0	$\mathcal{L}_1 = \mathcal{L}$ but not \mathcal{H}	70	4	Thm. 8+10
	(4,3,1)	0	\mathcal{L}_1 but not \mathcal{H}	280	40	Thm. 10
	(4,2,2)	0	\mathcal{L}_1 but not \mathcal{H}	420	32	Thm. 10
	(4,2,1,1)	0	\mathcal{L}_1 but not \mathcal{H}	840	100	Thm. 10
	(4,1,1,1,1)	0	\mathcal{L}_1 but not \mathcal{H}	1680	36	Thm. 10
9	(4,4,1)	1	\mathcal{H}	630	53	Thm. 10
	(4,3,2)	1	$\mid \mathcal{H} \mid$	1260	219	
	(4,3,1,1)	1	$\mid \mathcal{H} \mid$	2520	347	
	(4,2,2,1)	1	\mathcal{C} but nothing more tested	3780	565	
	(4,2,1,1,1)	1	\mathcal{C} but nothing more tested	7560	685	
	(4,1,1,1,1,1)	1	\mathcal{C} but nothing more tested	15120	173	

Table 1. Base cases settled by computer.

is indeed the case. We show that under this assumption the following eleven statements hold. Consider the following statements about integer partitions a with $\Delta(a) = 1$:

- (1a) For $\boldsymbol{a} = (1, 1, 1)$ we have $G(\boldsymbol{a}) \in \mathcal{E}$;
- (1b) For $\mathbf{a} \in \{(2,1,1,1), (2,2,1), (3,1,1,1,1)\}$ we have $G(\mathbf{a}) \in \mathcal{H}$;
- (1c) For any $\mathbf{a} \in \{(\alpha, \alpha, 1) \mid \alpha \geq 3\}$ we have $G(\mathbf{a}) \in \mathcal{L}_{12}$;
- (1d) For any $\boldsymbol{a} \in \{(\alpha, \alpha 1, 2) \mid \alpha \geq 3\}$ we have $G(\boldsymbol{a}) \in \mathcal{H}$;
- (1e) For any $\mathbf{a} \in \{(\alpha, \alpha 1, 1, 1) \mid \alpha \geq 3\}$ we have $G(\mathbf{a}) \in \mathcal{H}$;
- (1f) For any **a** that satisfies $\mathbf{a} \notin \{(1,1,1), (3,1,1,1,1)\}$ and $\mathbf{a} \notin \{(\alpha,\alpha,1), (\alpha,\alpha-1,2), (\alpha,\alpha-1,1,1) \mid \alpha \geq 2\}$ we have $G(\mathbf{a}) \in \mathcal{H}$.

Consider the following statements about integer partitions a with $\Delta(a) \geq 2$:

- (2a) For any $\mathbf{a} = 1^k$, $k \geq 4$, we have $G(\mathbf{a}) \in \mathcal{L}$;
- (2b) For any $a = (2, 1^{k-1}), k \ge 5$, we have $G(a) \in \mathcal{H}$;
- (2c) For any $\mathbf{a} \in \{(\alpha, \alpha, 2) \mid \alpha \geq 3\}$ we have $G(\mathbf{a}) \in \mathcal{H}$;
- (2d) For any $\mathbf{a} \in \{(\alpha, \alpha, 1, 1) \mid \alpha \geq 3\}$ we have $G(\mathbf{a}) \in \mathcal{H}$;
- (2e) For any \boldsymbol{a} that satisfies $\boldsymbol{a} \neq (1, 1, \dots, 1)$, $\boldsymbol{a} \neq (2, 1, 1, \dots, 1)$ and $\boldsymbol{a} \notin \{(\alpha, \alpha, 2), (\alpha, \alpha, 1, 1) \mid \alpha \geq 3\}$ we have $G(\boldsymbol{a}) \in \mathcal{H}$.

Note that the cases considered in these statements are mutually exclusive and cover all possibilities.

Recall that G(1,1,1) is a 6-cycle, so (1a) holds trivially and unconditionally. Also, (1b) holds unconditionally by Table 1. Moreover, (2a) and (2b) hold unconditionally by Theorem 5 and Lemma 22, respectively. Statement (1c) holds by Lemma 21, using that for $\mathbf{a} = (\alpha, \alpha, 1)$ with $\alpha \geq 3$, the partition $\mathbf{b} = (\alpha, \alpha) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}$ by Theorem 8, and the partition $\mathbf{b} = (\alpha, \alpha - 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}_1$ by Conjecture 6.

We prove the remaining statements by induction on $n = \sum_{i=1}^{k} a_i$.

Statement (1d) holds by Lemma 19, using that for $\mathbf{a} = (\alpha, \alpha - 1, 2)$ with $\alpha \geq 3$, the partitions $\mathbf{b} = \varphi((\alpha, \alpha - 2, 2)) \prec \mathbf{a}$ and $\mathbf{b} = (\alpha, \alpha - 1, 1) \prec \mathbf{a}$ satisfy $G(\mathbf{b}) \in \mathcal{L}_1$ by Conjecture 6, the partition $\mathbf{c} = (\alpha - 1, \alpha - 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{L}_{12}$ if $\alpha \geq 4$ by induction and (1c) and $G(\mathbf{c}) = G(2, 2, 1) \in \mathcal{H} \subseteq \mathcal{L}_{12}$ if $\alpha = 3$ by Table 1, and the partition $\mathbf{c} = \varphi((\alpha - 1, \alpha - 2, 2)) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{H}$ if $\alpha \geq 4$ by induction and (1d) and $G(\mathbf{c}) = G(2, 2, 1) \in \mathcal{H}$ if $\alpha = 3$.

Statement (1e) holds by Lemma 20, using that for $\mathbf{a} = (\alpha, \alpha - 1, 1, 1)$ with $\alpha \geq 3$, the partitions $\mathbf{b} = (\alpha, \alpha - 2, 1, 1) \prec \mathbf{a}$ and $\mathbf{b} = (\alpha, \alpha - 1, 1) \prec \mathbf{a}$ satisfy $G(\mathbf{b}) \in \mathcal{L}_1$ by Conjecture 6, the partition $\mathbf{c} = (\alpha - 1, \alpha - 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{L}_{12}$ if $\alpha \geq 4$ by induction and (1c) and $G(\mathbf{c}) = G(2, 2, 1) \in \mathcal{H}$ if $\alpha = 3$ by Table 1, and the partition $\mathbf{c} = (\alpha - 1, \alpha - 2, 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{c}) \in \mathcal{H}$ if $\alpha \geq 4$ by induction and (1e) and $G(\mathbf{c}) = G(2, 1, 1, 1) \in \mathcal{H}$ if $\alpha = 3$.

Statement (2c) holds by Lemma 17, using that for $\mathbf{a} = (\alpha, \alpha, 2)$ with $\alpha \geq 3$, the partition $\mathbf{b} = (\alpha, \alpha, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}_{12}$ by induction and (1c), and the partition $\mathbf{b} = (\alpha, \alpha - 1, 2) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{H}$ by induction and (1d).

Statement (2d) holds by Lemma 18, using that for $\mathbf{a} = (\alpha, \alpha, 1, 1)$ with $\alpha \geq 3$, the partition $\mathbf{b} = (\alpha, \alpha, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{L}_{12}$ by induction and (1c), and the partition $\mathbf{b} = (\alpha, \alpha - 1, 1, 1) \prec \mathbf{a}$ satisfies $G(\mathbf{b}) \in \mathcal{H}$ by induction and (1e).

To prove statement (1f), consider a partition $\mathbf{a}=(a_1,\ldots,a_k)$ satisfying the conditions of (1f). Note that $n=\sum_{i=1}^k a_i \geq 9$, as the only partitions with $n\in\{2,\ldots,8\}$ and $\Delta(\mathbf{a})=1$ are $\mathbf{a}\in\{(1,1,1),(2,2,1),(2,1,1,1),(3,3,1),(3,2,2),(3,2,1,1),(3,1,1,1,1)\}$, which are excluded by the conditions of (1f). From (2) and $\Delta(\mathbf{a})=1$ we obtain $a_1=(n-1)/2\geq 4$ and $a_1+1=\sum_{i=2}^k a_i$. The latter equation implies $a_2< a_1-1$, as otherwise $a_1+1\geq (a_1-1)+\sum_{i=3}^k a_i$, or equivalently, $\sum_{i=3}^k a_i \leq 2$, i.e., we would have $\mathbf{a}\in\{(\alpha,\alpha,1),(\alpha,\alpha-1,2),(\alpha,\alpha-1,1,1)\}$ for $\alpha:=a_1\geq 4$, which is impossible by the conditions of (1f). We thus obtain $a_2\leq a_1-2$ and $\sum_{i=3}^k a_i\geq 3$, in particular $k\geq 3$.

We now check that the preconditions of Lemma 16 are met. Consider any partition $\mathbf{b} \prec \mathbf{a}$ that has the same first entry as \mathbf{a} . By Lemma 11 we have $\Delta(\mathbf{b}) = 0$. From $n_{\mathbf{a}} \geq 9$ we obtain that $n_{\mathbf{b}} \geq 8$, implying that $\mathbf{b} \neq (2,2)$. Consequently, by Conjecture 6 we have $G(\mathbf{b}) \in \mathcal{L}_1$.

Now consider the partition $\mathbf{b} = (a_1 - 1, a_2, \dots, a_k)$, which satisfies $\Delta(\mathbf{b}) = \Delta(\mathbf{a}) + 1 = 2$ by Lemma 11. We show that all $\mathbf{c} \prec \mathbf{b}$ satisfy $G(\mathbf{c}) \in \mathcal{H}$. From $n_{\mathbf{a}} \geq 9$ we obtain that $n_{\mathbf{b}} \geq 8$ and $n_{\mathbf{c}} \geq 7$. Clearly, any $\mathbf{c} \prec \mathbf{b}$ satisfies $\Delta(\mathbf{c}) \in \{1,3\}$ by Lemma 11. Specifically, the partition $\mathbf{c} = (a_1 - 2, a_2, \dots, a_k)$ satisfies $\Delta(\mathbf{c}) = 3$, but we have $\mathbf{c} \neq 1^k$ for $k \geq 4$ because of $a_1 - 2 \geq 2$, so by induction and $(2\mathbf{b}) + (2\mathbf{e})$ we have $G(\mathbf{c}) \in \mathcal{H}$. Moreover, any other partition $\mathbf{c} \prec \mathbf{b}$ that has the same first entry as \mathbf{b} satisfies $\Delta(\mathbf{c}) = 1$, but we have $\mathbf{c} \neq (1, 1, 1)$ and $\mathbf{c} \neq (\alpha, \alpha, 1)$ for $\alpha \geq 3$ because of $\sum_{i=3}^k a_i \geq 3$, so by induction and $(1\mathbf{b}) + (1\mathbf{d}) + (1\mathbf{e}) + (1\mathbf{f})$ we have $G(\mathbf{c}) \in \mathcal{H}$.

We argued that the preconditions for applying Lemma 16 are met, and this lemma therefore yields $G(\mathbf{a}) \in \mathcal{H}$, proving (1f).

It remains to prove statement (2e). Consider a partition $\mathbf{a} = (a_1, \dots, a_k)$ satisfying the conditions of (2e). Note that $n = \sum_{i=1}^k a_i \ge 6$, as there are no partitions with $n \in \{2, \dots, 5\}$ and $\Delta(\mathbf{a}) = 2$. Note that $a_1 \ge a_2 \ge 2$ by the assumptions $\mathbf{a} \ne (1, 1, \dots, 1)$ and $\mathbf{a} \ne (2, 1, 1, \dots, 1)$ in (2e). Moreover, if $\mathbf{a} \notin \{(2, 2, 2), (2, 2, 1, 1)\}$ then we have $a_1 > a_2$ by the assumptions $\mathbf{a} \notin \{(\alpha, \alpha, 2), (\alpha, \alpha, 1, 1) \mid \alpha \ge 3\}$. From the equation $a_1 + 2 = \sum_{i=2}^k a_i$ and $a_1 \ge a_2$ we obtain

 $\sum_{i=3}^k a_i \geq 2$, in particular $k \geq 3$. Moreover, if $\mathbf{a} \notin \{(2,2,2),(2,2,1,1)\}$ then we can use the strict inequality $a_1 > a_2$ instead and obtain $\sum_{i=3}^k a_i \geq 3$.

We now verify that the preconditions of Lemma 15 are met, i.e., we consider all partitions $b \prec a$. From $n_a \geq 5$ we obtain that $n_b \geq 4$. If $a_1 > a_2$ then consider the partition $b = (a_1 - 1, a_2, \dots, a_k)$, which satisfies $\Delta(\mathbf{b}) = \Delta(\mathbf{a}) + 1 \geq 3$ by Lemma 11, but $\mathbf{b} \neq 1^k$ for $k \geq 4$ because of $a_2 \geq 1$ 2. Consequently, by induction and (2b)+(2e) we have $G(b) \in \mathcal{H}$. Now suppose that $a \notin$ $\{(2,2,2),(2,2,1,1)\}$ and consider any partition $b \prec a$ that has the same first entry as a, then by Lemma 11 we have $\Delta(\mathbf{b}) = \Delta(\mathbf{a}) - 1 \ge 1$, but we have $\mathbf{b} \ne (1, 1, 1)$ and $\mathbf{b} \ne (\alpha, \alpha, 1)$ for $\alpha \ge 3$ because of $\sum_{i=3}^k a_i - 1 \ge 2$, and $\mathbf{b} \ne 1^k$ for $k \ge 4$ and $\mathbf{b} \ne (2, 1^{k-1})$ for $k \ge 5$ because of $a_2 \ge 2$, so by induction and (1b)+(1d)+(1e)+(1f)+(2b)+(2c)+(2d)+(2e) we have $G(b) \in \mathcal{H}$. Lastly, if $a \in \{(2,2,2),(2,2,1,1)\}$ then there two partitions $b \prec a$, namely $b \in \{(2,2,1),(2,1,1,1)\}$, and both satisfy $G(\mathbf{b}) \in \mathcal{H}$ by (1b).

As the preconditions of Lemma 15 are satisfied, the lemma yields $G(a) \in \mathcal{H}$, proving (2e). This completes the proof of Theorem 7.

4.4. Proof of Theorem 10.

Proof of Theorem 10. From Table 1 we see that Conjecture 6 holds for all ten integer partitions $a \neq (2,2)$ with $\Delta(a) = 0$ whose first entry a_1 is at most 4. Specifically, these are the partitions $\boldsymbol{a} \in \{(1,1), (2,1,1), (3,3), (3,2,1), (3,1,1,1), (4,4), (4,3,1), (4,2,2), (4,2,1,1), (4,1,1,1,1)\}.$ For any integer partition a with $a_1 \leq 4$, we have that all integer partitions $b \prec a$ also satisfy $b_1 \leq 4$. Consequently, the statements (1a)-(1f) and (2a)-(2e) in the proof of Theorem 7 hold unconditionally for all such integer partitions. As $a_1 \in \{2,3,4\}$ we have $a \neq (1,1,1)$ and $a \neq 1^k$ for $k \geq 4$, and as $G(a) \in \mathcal{H}$ for $a \in \{(3,3,1),(4,4,1)\}$ by Table 1, we conclude that $G(a) \in \mathcal{H}$. \square

4.5. **Proofs of auxiliary lemmas.** To prove Lemmas 15–22, we follow the ideas outlined in Section 1.5.2, namely to partition the graph G(a) into subgraphs that are isomorphic to G(b)for integer partitions $b \prec a$, by fixing symbols at one or more of the positions $2, \ldots, n$. This allows us to construct a Hamilton path in the entire graph G(a) by gluing together the Hamilton paths obtained in the various subgraphs G(b) for $b \prec a$. The main difficulty is to ensure that the paths in the subgraphs G(b) under various constraints 'fit together', and that the constraints imposed on them at the gluing vertices are not too severe to rule out their existence. Four of these somewhat repetitive proofs are deferred to the appendix.

Proof of Lemma 15. As $\mathbf{a} \neq (1, 1, \dots, 1)$ and $\mathbf{a} \neq (2, 1, 1, \dots, 1)$, we know that $a_1 \geq a_2 \geq 2$. Moreover, from (2) we know that $a_1 + \Delta(\mathbf{a}) = \sum_{i=2}^k a_i$. Using $\Delta(\mathbf{a}) \geq 2$ and $a_1 \geq a_2$ in this equation yields $\sum_{i=3}^{k} a_i \geq 2$, in particular $k \geq 3$.

Let x, y be two distinct vertices in $G(\mathbf{a})$. As $x \neq y$ there is a position $\hat{\imath} > 1$ such that $x_{\hat{\imath}} \neq y_{\hat{\imath}}$. As $a_1 \geq a_2 \geq 2$ and $\sum_{i=3}^k a_i \geq 2$, there are two indices $i_1, i_2 \in [n] \setminus \{1, \hat{i}\}$ such that $x_{i_1} \neq x_{i_2}$. Similarly, there are two indices $i_3, i_4 \in [n] \setminus \{1, \hat{\imath}\}$ such that $y_{i_3} \neq y_{i_4}$.

We fix any permutation π on [k] such that $\pi_1 = x_i$ and $\pi_k = y_i$, and we choose a sequence of multiset permutations $u^j, v^j \in \Pi(\mathbf{a}), j = 1, \dots, k$, satisfying the following conditions; see

- (i) $u^1 = x$, $v^1_{i_1} = x_{i_2}$, and $v^k = y$, $u^k_{i_3} = y_{i_4}$; (ii) $u^j_1 = v^{j-1}_{\hat{\imath}} = \pi_{j-1}$, $u^j_{\hat{\imath}} = v^{j-1}_1 = \pi_j$, and $u^j_{\hat{\imath}} = v^{j-1}_{\hat{\imath}}$ for all $2 \le j \le k$ and $i \in [n] \setminus \{1, \hat{\imath}\}$.

As $u_1^j = \pi_{j-1}$ and $v_1^j = \pi_{j+1}$ by (ii), we have $u^j \neq v^j$ for all 1 < j < k. Moreover, by (i) and the choice of i_1, i_2 and i_3, i_4 we also have $u^1 \neq v^1$ and $u^k \neq v^k$, respectively.

For $j=1,\ldots,k$ we define $\boldsymbol{a}':=(a_1,\ldots,a_{\pi_j}-1,\ldots,a_k)$ and $\boldsymbol{b}:=\varphi(\boldsymbol{a}')\prec\boldsymbol{a}$, and we consider a Hamilton path P_j in the graph $G^{\hat{i},\pi_j}(\boldsymbol{a}) \simeq G(\boldsymbol{b})$ from u^j to v^j , which exists by the




FIGURE 11. Illustration of the proof of Lemma 15.

assumption $G(\mathbf{b}) \in \mathcal{H}$. By (ii), for all $2 \leq j \leq k$, the permutation u^j differs from v^{j-1} by a transposition of the entries at positions 1 and $\hat{\imath}$, implying that the concatenation $P_1P_2\cdots P_k$ is a Hamilton path in $G(\mathbf{a})$ from x to y.

The strategy for proving Lemma 17 and 18 is the same in the proof of Lemma 15, and these proofs can be found in the appendix. The crucial difference is that now one or two building blocks for partitions $\mathbf{b} \prec \mathbf{a}$ are only assumed to satisfy the weaker property $G(\mathbf{b}) \in \mathcal{L}_{12}$, and not $G(\mathbf{b}) \in \mathcal{H}$, so we have to work around this by imposing extra conditions on those building blocks.

The following lemma will be used in the proof of Lemma 16. It guarantees the existence of two multiset permutations π and ρ that are first and last vertices of Hamilton paths in subgraphs of G(a) obtained by fixing symbols, in such a way that π and ρ satisfy a number of extra conditions. In the proof of Lemma 16 we will encounter six different kinds of conditions, which are captured by the six cases in the lemma.

Lemma 23. For any odd $k \ge 3$ and $\ell := (k-1)/2$ and any $a, b, c \in [k]$ with $a \ne b$ and $c \ne 1$, there are permutations π, ρ on [k] satisfying any chosen of the following sets of conditions:

```
(oa) \pi_1 = 1, \rho_1 = a, \rho_k = b, \pi_{2j} \neq \rho_j and \pi_{2j+1} \neq \rho_{j+1} for all 1 \leq j \leq \ell;

(ob1) \pi_1 = 1, \pi_k = c, \rho_1 = 1, \pi_{2j} \neq \rho_j and \pi_{2j+1} \neq \rho_{j+1} for all 1 \leq j < \ell, \pi_{k-1} \neq \rho_k;

(ob2) \pi_1 = 1, \pi_2 = c, \rho_1 = 1, \pi_3 = \rho_k, \pi_{2j} \neq \rho_j and \pi_{2j+1} \neq \rho_{j+1} for all 1 \leq j \leq \ell.
```

For any even $k \geq 4$ and $\ell := k/2$ and any $a,b,c \in [k]$ with $a \neq b$ and $c \neq 1$, there are permutations π, ρ on [k] satisfying any chosen of the following sets of conditions:

(ea)
$$\pi_1 = 1$$
, $\rho_1 = a$, $\rho_k = b$, $\pi_3 = \rho_{k-1}$, $\pi_{2j} \neq \rho_j$ and $\pi_{2j+1} \neq \rho_{j+1}$ for all $1 \leq j < \ell$, $\pi_k \neq \rho_k$; (eb1) $\pi_1 = 1$, $\pi_k = c$, $\rho_1 = 1$, $\pi_3 = \rho_k$, $\pi_{2j} \neq \rho_j$ and $\pi_{2j+1} \neq \rho_{j+1}$ for all $1 \leq j < \ell$; (eb2) $\pi_1 = 1$, $\pi_k = c$, $\rho_1 = 1$, $\pi_{2j} \neq \rho_j$ and $\pi_{2j+1} \neq \rho_{j+1}$ for all $1 \leq j < \ell$, $\pi_k \neq \rho_k$.

Proof. For any of the given sets of conditions, we can define a graph H that captures the conditions on π and ρ as follows: We start with a complete bipartite graph with vertex partitions [k] and [k]. For any constraint of the form $\pi_i = \rho_j$ we remove both end vertices i and j from the graph. Similarly, for any pair of constraints of the form $\pi_i = x$ and $\rho_j = x$ we remove both end vertices i and j from the graph. Moreover, for any constraint of the form $\pi_i \neq \rho_j$ we remove the




FIGURE 12. Illustration of the graph H defined in the proof of Lemma 23. The pairs of vertices connected by thick solid lines are removed from the initial complete bipartite graph, and the thin edges are also removed (keeping the end vertices). From the two pairs of vertices connected by thick dashed lines, one pair of vertices is removed if a = 1 or b = 1, and both edges are removed if $a \neq 1$ and $b \neq 1$ (keeping the end vertices).

edge (i, j) from the graph (keeping the end vertices). Lastly, for any pair of constraints of the form $\pi_i = x$ and $\rho_j = y$ for $x \neq y$ we remove the edge (i, j) from the graph.

Note that there are permutations π and ρ satisfying one of the given sets of conditions, if and only if the corresponding graph H admits a perfect matching. Specifically, the matched pairs of vertices are the entries from π and ρ that will be assigned the same value.

The existence of a perfect matching in H defined before can be verified directly for $k \leq 5$; see Figure 12. For $k \geq 6$ we argue via Hall's theorem: We let s denote the sizes of the two partition classes of H. By the given equality constraints, at most two pairs of vertices are removed from

the initial complete bipartite graph, so we have $s \geq k-2 \geq 4$. Moreover, the given inequality constraints involve each π_i and ρ_j at most twice, so all vertices of H have degree at least s-2. Let A be a subset of vertices of one partition class, and let A' be the set of all of its neighbors in the other partition class of H. As all vertices of H have degree at least s-2, we know that $|A'| \geq s-2$ whenever |A| > 0, so it suffices to check Hall's condition for $|A| \geq s-1$. If |A'| < |A|, then there is a vertex on the same side of the partition as A' that is not connected to any vertex in A, i.e., this vertex has degree at most $s-|A| \leq s-(s-1)=1 < s-2$, a contradiction as $s \geq 4$. This proves that $|A'| \geq |A|$, so Hall's condition is satisfied and the graph H has a perfect matching. This completes the proof of the lemma.

We now outline our strategy for proving Lemma 16. In the preceding proof of Lemma 15, we split G(a) into subgraphs G(b) with $b \prec a$ by fixing one of the symbols. If we follow the same approach for an integer partition a as in Lemma 16 with $\Delta(a) = 1$, then by Lemma 11 all partitions $b \prec a$ satisfy $\Delta(b) = 0$, except the unique partition $b \prec a$ obtained from a by decreasing the first entry, which satisfies $\Delta(b) = 2$ (by the assumptions of Lemma 16, the first entry of a is the unique largest one). All subgraphs G(b) with $\Delta(b) = 0$ are not Hamiltonconnected, but only 1-laceable, so any Hamilton path in them starts with a vertex with first symbol 1 and ends with a vertex with first symbol distinct from 1. When constructing a Hamilton path in G(a), we process those subgraphs in pairs, so that the path through any pair starts and ends with a vertex with first symbol 1, and every such pair is surrounded by vertices with fixed symbol equal to 1. As a consequence, we need to further partition the unique subgraph G(b)with $\Delta(b) = 2$ by fixing yet another symbol, and split it into subgraphs G(c) with $c \prec b$, all of which satisfy $\Delta(c) \geq 1$ and $G(c) \in \mathcal{H}$ by the assumptions of the lemma, and those building blocks are inserted between the aforementioned pairs. The precise order in which the building blocks for a Hamilton path in G(a) are arranged depends on the parity of the number k of symbols, and on the start and end vertices of the desired Hamilton path, which results in several cases.

Proof of Lemma 16. As $\mathbf{a} \notin \{(1,1,1), (2,1,1,1), (2,2,1)\}$ we know that $a_1 \geq 3$, and combining this with $a_1 + \Delta(\mathbf{a}) = \sum_{i=2}^k a_i$, $\Delta(\mathbf{a}) = 1$ and $a_1 \geq a_2$ yields $\sum_{i=2}^k a_i \geq 4$ and $\sum_{i=3}^k a_i \geq 1$, in particular $k \geq 3$ (recall (2)). Moreover, we have $a_1 > a_2$ by the assumption that $\mathbf{a} \neq (\alpha, \alpha, 1)$ for $\alpha := (n-1)/2$.

Let x, y be two distinct vertices in G(a). If there is an index i > 1 with $x_i = 1$ and $y_i \neq 1$, then we let \check{i} be this index. Otherwise we fix an arbitrary index $\check{i} > 1$ with $x_{\check{i}} = y_{\check{i}} = 1$, which is possible as $a_1 \geq 3$.

We first consider the case that $k \geq 3$ is odd, and we introduce the abbreviation $\ell := (k-1)/2 \geq 1$. We now distinguish two cases, depending on the value of y_i .

Case (oa): $y_i = 1$. As $x \neq y$ there is a position $\hat{i} \in [n] \setminus \{1, \check{i}\}$ such that $x_i \neq y_i$, and by our choice of \check{i} , we can assume that $x_i, y_i > 1$.

As $k \geq 3$, $a_1 \geq 3$, and $\sum_{i=2}^k a_i \geq 4$, there are two indices $i_1, i_2 \in [n] \setminus \{1, \hat{i}, \check{i}\}$ such that $x_{i_1} \neq x_{i_2}$. Similarly, there are two indices $i_3, i_4 \in [n] \setminus \{1, \hat{i}, \check{i}\}$ such that $y_{i_3} \neq y_{i_4}$.

By Lemma 23 (oa) there are two permutations π, ρ on [k] satisfying the following conditions:

(1) $\pi_1 = x_i = 1$, $\rho_1 = x_i$, and $\rho_k = y_i$;

(2) $\pi_{2j} \neq \rho_j \text{ and } \pi_{2j+1} \neq \rho_{j+1} \text{ for all } 1 \leq j \leq \ell.$

We choose multiset permutations $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, ..., k$, and $\hat{u}^j, \hat{v}^j, \check{u}^j, \check{v}^j \in \Pi(\boldsymbol{a}), j = 1, ..., \ell$, satisfying the following conditions; see Figure 13 (oa):

(i)
$$u^1 = x$$
, $v_{i_1}^1 = x_{i_2}$, and $v^k = y$, $u_{i_3}^k = y_{i_4}$;

²The assumption $x_i, y_i > 1$ will not be used in this proof, but in the proof of Lemma 19.

- (ii) conditions (ϕ_i^1) , (ϕ_i^2) , and (ϕ_i^3) for all $1 \le j \le \ell$;
- (iii) condition (ϕ_j^4) for all $\ell < j < k$;
- (iv) condition $(\psi_{\ell+1})$,

where we use the following abbreviations:

```
(\phi_i^1): \ \hat{u}_1^j = v_i^j = 1, \ \hat{u}_i^j = v_1^j = \pi_{2j}, \ \hat{u}_i^j = v_i^j = \rho_j, \ \text{and} \ \hat{u}_i^j = v_i^j \ \text{for all} \ i \in [n] \setminus \{1, \check{\imath}, \hat{\imath}\};
```

 $(\phi_j^2): \ \check{u}_1^j = \hat{v}_i^j = \pi_{2j}, \ \check{u}_i^j = \hat{v}_1^j = \pi_{2j+1}, \text{ and } \check{u}_i^j = \hat{v}_i^j \text{ for all } i \in [n] \setminus \{1, \check{\imath}\};$ $(\phi_j^3): \ u_1^{j+1} = \check{v}_i^j = \pi_{2j+1}, \ u_i^{j+1} = \check{v}_1^j = 1, \ u_i^{j+1} = \check{v}_i^j = \rho_{j+1}, \text{ and } u_i^{j+1} = \check{v}_i^j \text{ for all } i \in [n]$

 $[n] \setminus \{1, \check{\imath}, \hat{\imath}\};$ $(\phi_j^4): u_1^{j+1} = v_{\hat{\imath}}^j = \rho_j, \ u_{\hat{\imath}}^{j+1} = v_1^j = \rho_{j+1}, \ u_{\check{\imath}}^{j+1} = v_{\check{\imath}}^j = 1, \ \text{and} \ u_i^{j+1} = v_i^j \ \text{for all} \ i \in [n] \setminus \{1, \check{\imath}, \hat{\imath}\};$

 $(\psi_i): u_{\tilde{i}}^j \neq v_{\tilde{i}}^j \text{ for some } \tilde{i} \in [n] \setminus \{1, \check{i}, \hat{i}\}.$

Observe that condition (ϕ_j^1) requires that both v^j and \hat{u}^j contain the symbols π_{2j} and ρ_j , and condition (ϕ_i^3) requires that both \check{v}^j and u^{j+1} contain the symbols π_{2j+1} and ρ_{j+1} , and this is possible for all $1 \leq j \leq \ell$ as π and ρ satisfy condition (2) from before, which ensures that these two symbols are distinct. Also note that $u^1 \neq v^1$ and $u^k \neq v^k$ by (i) and the choice of i_1, i_2 and i_3, i_4 , respectively, and $u_1^j \neq v_1^j$ for all 1 < j < k by (ii)+(iii)+(iv). Moreover, we have $\hat{u}_1^j \neq \hat{v}_1^j$ and $\check{u}_1^j \neq \check{v}_1^j$ for all $1 \leq j \leq \ell$ by (ii).

For j = 1, ..., k we define $b := (a_1 - 1, a_2, ..., a_k), c' := (a_1 - 1, ..., a_{\rho_i} - 1, ..., a_k),$ and $c := \varphi(c')$, which satisfy $c \prec b \prec a$ and $\Delta(b) = 2$ and $\Delta(c) = 1$, and we consider a Hamilton path P_i in $G^{(i,\hat{i}),(1,\rho_j)}(\boldsymbol{a}) \simeq G(\boldsymbol{c})$ from u^j to v^j , which exists by the assumption $G(\boldsymbol{c}) \in \mathcal{H}$.

For $j=1,\ldots,\ell$ we define $\boldsymbol{a}':=(a_1,\ldots,a_{\pi_{2j}}-1,\ldots,a_k)$ and $\boldsymbol{b}:=\varphi(\boldsymbol{a}')$, which satisfies $\boldsymbol{b}\prec\boldsymbol{a}$ and $\Delta(\boldsymbol{b}) = 0$, and we consider a Hamilton path \hat{P}_j in the graph $G^{i,\pi_{2j}}(\boldsymbol{a}) \simeq G(\boldsymbol{b})$ from \hat{u}^j to \hat{v}^j , which exists by the assumption $G(b) \in \mathcal{L}_1$, using that $\hat{u}^j \in \Pi(a)^{1,1}$ and $\hat{v}^j \notin \Pi(a)^{1,1}$ (recall that $\hat{u}_1^j = 1$ and $\hat{v}_1^j = \pi_{2j+1} \neq 1$ by (ii)).

Similarly, for $j=1,\ldots,\ell$ we define $\boldsymbol{a}':=(a_1,\ldots,a_{\pi_{2j+1}}-1,\ldots,a_k)$ and $\boldsymbol{b}:=\varphi(\boldsymbol{a}')$, and we consider a Hamilton path \check{P}_j in the graph $G^{\check{\imath},\pi_{2j+1}}(\boldsymbol{a}) \simeq G(\boldsymbol{b})$ from \check{u}^j to \check{v}^j , which exists by the assumption $G(\mathbf{b}) \in \mathcal{L}_1$, using that $\check{u}^j \notin \Pi(\mathbf{a})^{1,1}$ and $\check{v}^j \in \Pi(\mathbf{a})^{1,1}$ (recall that $\check{u}_1^j = \pi_{2j} \neq 1$ and $\check{v}_1^j = 1$ by (ii)).

The concatenation $(P_1\hat{P}_1\check{P}_1)(P_2\hat{P}_2\check{P}_2)\cdots(P_\ell\hat{P}_\ell\check{P}_\ell)P_{\ell+1}P_{\ell+2}\cdots P_k$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Case (ob): $y_i \neq 1$. We choose an index $\hat{i} \in [n] \setminus \{1, \check{i}\}$ for which $x_{\hat{i}} = 1$, and two indices $i_1, i_2 \in [n] \setminus \{1, \hat{i}, \check{i}\}$ such that $x_{i_1} \neq x_{i_2}$. We consider two subcases depending on the value of y_1 .

Case (ob1): $y_1 = 1$. By Lemma 23 (ob1) there are two permutations π, ρ on [k] satisfying the following conditions:

- (1) $\pi_1 = x_{\check{i}} = 1$, $\pi_k = y_{\check{i}}$, $\rho_1 = x_{\hat{i}} = 1$;
- (2) $\pi_{2j} \neq \rho_j$ and $\pi_{2j+1} \neq \rho_{j+1}$ for all $1 \leq j < \ell$, and $\pi_{k-1} \neq \rho_k$.

We choose multiset permutations $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, ..., k$, and $\hat{u}^j, \hat{v}^j, \check{u}^j, \check{v}^j \in \Pi(\boldsymbol{a})$, $j = 1, \dots, \ell$, satisfying the following conditions; see Figure 13 (ob1):

- (i) $u^1 = x$, $v_{i_1}^1 = x_{i_2}$, and $\check{v}^{\ell} = y$;
- (iia) conditions (ϕ_j^1) , (ϕ_j^2) , (ϕ_j^3) for all $1 \le j < \ell$, and condition (ϕ_ℓ^2) ; (iib) $\hat{u}_1^{\ell} = v_i^k = 1$, $\hat{u}_i^{\ell} = v_1^k = \pi_{k-1}$, $\hat{u}_i^{\ell} = v_i^k = \rho_k$, and $\hat{u}_i^{\ell} = v_i^k$ for all $i \in [n] \setminus \{1, \check{i}, \hat{i}\}$;
- (iii) condition (ϕ_i^4) for all $\ell \leq j < k$;
- (iv) condition (ψ_i) for $j \in \{\ell, k\}$.

The argument that all pairs (u^j, v^j) , $j = 1, \ldots, k$, and (\hat{u}^j, \hat{v}^j) and $(\check{u}^j, \check{v}^j)$, $j = 1, \ldots, \ell$, are distinct, is straightforward.




FIGURE 13. Illustration of the proof of Lemma 16 in the case when k is odd.

As before in case (oa), we consider Hamilton paths P_j from u^j to v^j for all $j=1,\ldots,k$, Hamilton paths \hat{P}_j and \check{P}_j from \hat{u}^j to \hat{v}^j , or from \check{u}^j to \check{v}^j , respectively, for all $j=1,\ldots,\ell$. The concatenation $(P_1\hat{P}_1\check{P}_1)(P_2\hat{P}_2\check{P}_2)\cdots(P_{\ell-1}\hat{P}_{\ell-1}\check{P}_{\ell-1})P_{\ell}P_{\ell+1}\cdots P_k(\hat{P}_{\ell}\check{P}_{\ell})$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Case (ob2): $y_1 \neq 1$. By Lemma 23 (ob2) there are two permutations π, ρ on [k] satisfying the following conditions:

- (1) $\pi_1 = x_i = 1$, $\pi_2 = y_i$, $\rho_1 = x_i = 1$, and $\pi_3 = \rho_k$;
- (2) $\pi_{2j} \neq \rho_j$ and $\pi_{2j+1} \neq \rho_{j+1}$ for all $1 \leq j \leq \ell$.

We choose two multiset permutations $\hat{u}^1, \hat{v}^1 \in \Pi(\boldsymbol{a})$ such that $\hat{u}^1_1 = 1, \ \hat{u}^1_{i_1} = x_{i_2}, \ \hat{u}^1_{\hat{i}} = 1,$ $\hat{u}_i^1 = \pi_2$, and $\hat{v}^1 = y$. We define $\mathbf{a}' := (a_1, \dots, a_{\pi_2} - 1, \dots, a_k)$ and $\mathbf{b} := \varphi(\mathbf{a}')$, which satisfies $\boldsymbol{b} \prec \boldsymbol{a}$ and $\Delta(\boldsymbol{b}) = 0$, and we consider a Hamilton path Q in the graph $G^{i,\pi_2}(\boldsymbol{a}) \simeq G(\boldsymbol{b})$ from \hat{u}^1 to \hat{v}^1 , which exists by the assumption $G(\boldsymbol{b}) \in \mathcal{L}_1$, using that $\hat{u}^1 \in \Pi(\boldsymbol{a})^{1,1}$ and $\hat{v}^1 \notin \Pi(\boldsymbol{a})^{1,1}$ (recall that $\hat{u}_1^1 = 1$ and $\hat{v}_1^1 = y_1 \neq 1$). Let z be the first vertex along this path from \hat{u}^1 to \hat{v}^1 for which $z_i = \pi_3$, and let z' be the predecessor of z on the path. By the definition of z we have $z_i' \neq \pi_3$, and consequently $z_1' = \pi_3$. By Lemma 14, we thus obtain $z_i' = z_1 = 1$. Let Q' be the subpath of Q from \hat{u}^1 to z', and let Q'' be the subpath of Q from z to \hat{v}^1 .

We choose multiset permutations $u^j, v^j \in \Pi(\mathbf{a}), j = 1, \ldots, k$, and $\hat{u}^j, \hat{v}^j \in \Pi(\mathbf{a}), j = 2, \ldots, \ell$, and $\check{u}^j, \check{v}^j \in \Pi(\boldsymbol{a}), j = 1, \dots, \ell$, satisfying the following conditions; see Figure 13 (ob2):

- (i) $u^1 = x$, and conditions (χ^1) and (χ^2_k) ;
- (ii) conditions (ϕ_j^1) and (ϕ_j^2) for all $1 < j \le \ell$, and condition (ϕ_j^3) for all $1 \le j \le \ell$;
- (iii) condition (ϕ_j^4) for all $\ell < j < k$;
- (iv) condition (ψ_i) for $j \in \{\ell + 1, k\}$,

where we introduce the abbreviations:

$$(\chi^1): v_1^1 = \hat{u}_i^1 = \pi_2, \ v_i^1 = \hat{u}_1^1 = 1, \ v_i^1 = \hat{u}_i^1, \ \text{and} \ \check{u}_1^1 = z_i' = \pi_2, \ \check{u}_i^1 = z_1' = \pi_3, \ \check{u}_i^1 = z_i' \ \text{for all} \ i \in [n] \setminus \{1, \check{i}\};$$

$$(\chi_i^2)$$
: $v_1^j = z_i = \pi_2$, $v_i^j = z_1 = 1$, $v_i^j = z_i$ for all $i \in [n] \setminus \{1, i\}$.

The argument that all pairs (u^j, v^j) , $j = 1, \ldots, k$, and (\hat{u}^j, \hat{v}^j) , $j = 2, \ldots, \ell$, and $(\check{u}^j, \check{v}^j)$, $j = 1, \dots, \ell$, are distinct, is straightforward.

As before in case (oa), we consider Hamilton paths P_j from u^j to v^j for all $j = 1, \ldots, k$, Hamilton paths \hat{P}_j from \hat{u}^j to \hat{v}^j for all $j=2,\ldots,\ell$, and Hamilton paths \dot{P}_j from \check{u}^j to \check{v}^j for all $j = 1, \ldots, \ell$.

The concatenation $(P_1Q'\check{P}_1)(P_2\hat{P}_2\check{P}_2)(P_3\hat{P}_3\check{P}_3)\cdots(P_\ell\hat{P}_\ell\check{P}_\ell)P_{\ell+1}\cdots P_kQ''$ is a Hamilton path in G(a) from x to y.

It remains to consider the case that $k \geq 4$ is even, and we introduce the abbreviation $\ell := k/2 \ge 2$. We now distinguish two cases, depending on the value of y_i .

Case (eb): $y_i \neq 1$. We choose an index $\hat{i} \in [n] \setminus \{1, \check{i}\}$ for which $x_{\hat{i}} = 1$, and two indices $i_1, i_2 \in [n] \setminus \{1, \hat{i}, \check{i}\}$ such that $x_{i_1} \neq x_{i_2}$. We consider two subcases depending on the value of y_1 .

Case (eb2): $y_1 \neq 1$. By Lemma 23 (eb2) there are two permutations π, ρ on [k] satisfying the following conditions:

- (1) $\pi_1 = x_{\check{i}} = 1$, $\pi_k = y_{\check{i}}$, and $\rho_1 = x_{\hat{i}} = 1$;
- (2) $\pi_{2j} \neq \rho_j$ and $\pi_{2j+1} \neq \rho_{j+1}$ for all $1 \leq j < \ell$, and $\pi_k \neq \rho_k$.

We choose multiset permutations $u^j, v^j \in \Pi(\mathbf{a}), j = 1, \ldots, k$, and $\hat{u}^j, \hat{v}^j \in \Pi(\mathbf{a}), j = 1, \ldots, \ell$, and $\check{u}^j, \check{v}^j \in \Pi(\boldsymbol{a}), j = 1, \dots, \ell - 1$, satisfying the following conditions; see Figure 14 (eb2):

- (i) $u^1 = x$, $v_{i_1}^1 = x_{i_2}$, and $\hat{v}^{\ell} = y$;
- (iia) conditions (ϕ_j^1) , (ϕ_j^2) , and (ϕ_j^3) for all $1 \le j < \ell$; (iib) $\hat{u}_1^{\ell} = v_i^k = 1$, $\hat{u}_i^{\ell} = v_1^k = \pi_k$, $\hat{u}_i^{\ell} = v_i^k = \rho_k$, and $\hat{u}_i^{\ell} = v_i^k$ for all $i \in [n] \setminus \{1, \check{i}, \hat{i}\}$;
- (iii) condition (ϕ_j^4) for all $\ell \leq j < k$;
- (iv) condition (ψ_j) for $j \in \{\ell, k\}$.

The argument that all pairs (u^j, v^j) , j = 1, ..., k, and (\hat{u}^j, \hat{v}^j) , $j = 1, ..., \ell$, and $(\check{u}^j, \check{v}^j)$, $j = 1, \dots, \ell - 1$, are distinct, is straightforward.

For j = 1, ..., k we define $b := (a_1 - 1, a_2, ..., a_k), c' := (a_1 - 1, ..., a_{\rho_i} - 1, ..., a_k),$ and $c := \varphi(c')$, which satisfy $c \prec b \prec a$ and $\Delta(b) = 2$ and $\Delta(c) = 1$, and we consider a Hamilton path P_j in $G^{(\check{\imath},\hat{\imath}),(1,\rho_j)}(\boldsymbol{a}) \simeq G(\boldsymbol{c})$ from u^j to v^j , which exists by the assumption $G(\boldsymbol{c}) \in \mathcal{H}$.




FIGURE 14. Illustration of the proof of Lemma 16 in the case when k is even.

For $j=1,\ldots,\ell$ we define $\mathbf{a}':=(a_1,\ldots,a_{\pi_{2j}}-1,\ldots,a_k)$ and $\mathbf{b}:=\varphi(\mathbf{a}')$, which satisfies $\mathbf{b}\prec\mathbf{a}$ and $\Delta(\mathbf{b})=0$, and we consider a Hamilton path \hat{P}_j in the graph $G^{\check{\imath},\pi_{2j}}(\mathbf{a})\simeq G(\mathbf{b})$ from \hat{u}^j to \hat{v}^j , which exists by the assumption $G(\mathbf{b})\in\mathcal{L}_1$, using that $\hat{u}^j\in\Pi(\mathbf{a})^{1,1}$ and $\hat{v}^j\notin\Pi(\mathbf{a})^{1,1}$ (recall that $\hat{u}^j_1=1$ and $\hat{v}^j_1=\pi_{2j+1}\neq 1$ for $j=1,\ldots,\ell-1$ by (ii) and $\hat{u}^\ell_1=1$ and $\hat{v}^\ell_1=y_1\neq 1$ by (iib) and (i), respectively).

Similarly, for $j = 1, ..., \ell - 1$ we define $\mathbf{a}' := (a_1, ..., a_{\pi_{2j+1}} - 1, ..., a_k)$ and $\mathbf{b} := \varphi(\mathbf{a}')$, and we consider a Hamilton path \check{P}_j in the graph $G^{\check{\imath},\pi_{2j+1}}(\mathbf{a}) \simeq G(\mathbf{b})$ from \check{u}^j to \check{v}^j , which exists by the assumption $G(\mathbf{b}) \in \mathcal{L}_1$, using that $\check{u}^j \notin \Pi(\mathbf{a})^{1,1}$ and $\check{v}^j \in \Pi(\mathbf{a})^{1,1}$ (recall that $\check{u}^j_1 = \pi_{2j} \neq 1$ and $\check{v}^j_1 = 1$ by (iia)).

The concatenation $(P_1\hat{P}_1\check{P}_1)(P_2\hat{P}_2\check{P}_2)\cdots(P_{\ell-1}\hat{P}_{\ell-1}\check{P}_{\ell-1})P_{\ell}P_{\ell+1}\cdots P_{k-1}P_k\hat{P}_{\ell}$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Case (eb1): $y_1 = 1$. By Lemma 23 (eb1) there are two permutations π, ρ on [k] satisfying the following conditions:

- (1) $\pi_1 = x_i = 1$, $\pi_k = y_i$, $\rho_1 = x_i = 1$, and $\pi_3 = \rho_k$;
- (2) $\pi_{2j} \neq \rho_j \text{ and } \pi_{2j+1} \neq \rho_{j+1} \text{ for all } 1 \leq j < \ell.$

We choose two multiset permutations $\hat{u}^1, \hat{v}^1 \in \Pi(\boldsymbol{a})$ such that $\hat{u}_1^1 = 1, \ \hat{u}_{i_1}^1 = x_{i_2}, \ \hat{u}_{\hat{i}}^1 = \rho_1 = 1, \ \hat{u}_{\hat{i}}^1 = \pi_2, \ \hat{v}_1^1 = \pi_k, \ \text{and} \ \hat{v}_{\hat{i}}^1 = \pi_2.$ We define $\boldsymbol{a}' := (a_1, \dots, a_{\pi_2} - 1, \dots, a_k)$ and $\boldsymbol{b} := \varphi(\boldsymbol{a}'), \ \text{which}$ satisfies $\boldsymbol{b} \prec \boldsymbol{a}$ and $\Delta(\boldsymbol{b}) = 0$, and we consider a Hamilton path Q in the graph $G^{\check{i},\pi_2}(\boldsymbol{a}) \simeq G(\boldsymbol{b})$ from \hat{u}^1 to \hat{v}^1 , which exists by the assumption $G(\boldsymbol{b}) \in \mathcal{L}_1$, using that $\hat{u}^1 \in \Pi(\boldsymbol{a})^{1,1}$ and $\hat{v}^1 \notin \Pi(\boldsymbol{a})^{1,1}$ (recall that $\hat{u}_1^1 = 1$ and $\hat{v}_1^1 = \pi_k \neq 1$). Let z be the first vertex along this path from \hat{u}^1 to \hat{v}^1 for which $z_{\hat{i}} = \pi_3$, and let z' be the predecessor of z on the path. By the definition of z we have $z'_{\hat{i}} \neq \pi_3$, and consequently $z'_1 = \pi_3$. By Lemma 14, we thus obtain $z'_{\hat{i}} = z_1 = 1$. Let Q' be the subpath of Q from \hat{u}^1 to z', and let Q'' be the subpath of Q from z to \hat{v}^1 .

We choose multiset permutations $u^j, v^j \in \Pi(\mathbf{a}), j = 1, ..., k$, and $\hat{u}^j, \hat{v}^j \in \Pi(\mathbf{a}), j = 2, ..., \ell-1$, and $\check{u}^j, \check{v}^j \in \Pi(\mathbf{a}), j = 1, ..., \ell$, satisfying the following conditions; see Figure 14 (eb1):

- (i) $u^1=x,\;\check{v}^\ell=y,$ and conditions $(\chi^1),\,(\chi^2_k),$ and $(\chi^3);$
- (ii) conditions (ϕ_j^1) and (ϕ_j^2) for all $1 < j < \ell$, and condition (ϕ_j^3) for all $1 \le j < \ell$;
- (iii) condition (ϕ_j^4) for all $\ell \leq j < k$;
- (iv) condition (ψ_j) for $j \in \{\ell, k\}$,

where we introduce the abbreviation:

$$(\chi^3): \ \check{u}_1^\ell = \hat{v}_{\check{i}}^1 = \pi_2, \ \check{u}_{\check{i}}^\ell = \hat{v}_1^1 = \pi_k, \ \check{u}_{\check{i}}^\ell = \hat{v}_i^1 \ \text{for all} \ i \in [n] \setminus \{1, \check{i}\}.$$

The argument that all pairs (u^j, v^j) , j = 1, ..., k, and (\hat{u}^j, \hat{v}^j) , $j = 2, ..., \ell - 1$, and $(\check{u}^j, \check{v}^j)$, $j = 1, ..., \ell$, are distinct, is straightforward.

As before in case (eb2), we consider Hamilton paths P_j from u^j to v^j for all $j=1,\ldots,k$, Hamilton paths \hat{P}_j from \hat{u}^j to \hat{v}^j for all $j=2,\ldots,\ell-1$, and Hamilton paths \check{P}_j from \check{u}^j to \check{v}^j for all $j=1,\ldots,\ell-1$. In addition, we consider a Hamilton path \check{P}_ℓ from \check{u}^ℓ to \check{v}^ℓ , which exists by the assumption $G(\boldsymbol{b}) \in \mathcal{L}_1$, using that $\check{u}^\ell \notin \Pi(\boldsymbol{a})^{1,1}$ and $\check{v}^\ell \in \Pi(\boldsymbol{a})^{1,1}$ (recall that $\check{u}^\ell_1 = \pi_2 \neq 1$ and $\check{v}^\ell_1 = y_1 = 1$ by (i)).

The concatenation $(P_1Q'\check{P}_1)(P_2\hat{P}_2\check{P}_2)(P_3\hat{P}_3\check{P}_3)\cdots(P_{\ell-1}\hat{P}_{\ell-1}\check{P}_{\ell-1})P_{\ell}P_{\ell+1}\cdots P_{k-1}P_kQ''\check{P}_{\ell}$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Case (ea): $y_i = 1$. As $x \neq y$ there is a position $\hat{i} \in [n] \setminus \{1, \check{i}\}$ such that $x_{\hat{i}} \neq y_{\hat{i}}$, and by our choice of \check{i} , we can assume that $x_{\hat{i}}, y_{\hat{i}} > 1$.

We fix two indices $i_1, i_2 \in [n] \setminus \{1, \hat{i}, \check{i}\}$ such that $x_{i_1} \neq x_{i_2}$, and two indices $i_3, i_4 \in [n] \setminus \{1, \hat{i}, \check{i}\}$ such that $y_{i_3} \neq y_{i_4}$.

By Lemma 23 (ea) there are two permutations π, ρ on [k] satisfying the following conditions:

- (1) $\pi_1 = x_{\tilde{i}} = 1$, $\rho_1 = x_{\hat{i}}$, $\rho_k = y_{\hat{i}}$, and $\pi_3 = \rho_{k-1}$;
- (2) $\pi_{2j} \neq \rho_j$ and $\pi_{2j+1} \neq \rho_{j+1}$ for all $1 \leq j < \ell$, and $\pi_k \neq \rho_k$.

We choose two multiset permutations $\hat{u}^1, \hat{v}^1 \in \Pi(\boldsymbol{a})$ as before in case (eb1), and we construct a Hamilton path Q from \hat{u}^1 to \hat{v}^1 and its subpaths Q' from \hat{u}^1 to z' and Q'' from z to \hat{v}^1 , as before.

³The assumption $x_{\hat{i}}, y_{\hat{i}} > 1$ will not be used in this proof, but in the proof of Lemma 20.

We choose multiset permutations $u^j, v^j \in \Pi(\mathbf{a}), j = 1, \dots, k$, and $\hat{u}^j, \hat{v}^j \in \Pi(\mathbf{a}), j = 2, \dots, \ell-1$, and $\check{u}^j, \check{v}^j \in \Pi(a), j = 1, \dots, \ell$, satisfying the following conditions; see Figure 14 (ea):

- (i) $u^1 = x$, $v^k = y$, $u^k_{i_3} = x_{i_4}$, and conditions (χ^1) , (χ^2_{k-1}) , and (χ^3) ; (iia) conditions (ϕ^1_j) and (ϕ^2_j) for all $1 < j < \ell$, and condition (ϕ^3_j) for all $1 \le j < \ell$;
- (iib) $u_1^k = \check{v}_{\check{i}}^\ell = \pi_k, \ u_{\check{i}}^k = \check{v}_1^\ell = 1, \ u_{\hat{i}}^k = \check{v}_{\hat{i}}^\ell = \rho_k, \ u_{\check{i}}^k = \check{v}_{\check{i}}^\ell \text{ for all } i \in [n] \setminus \{1, \check{i}, \hat{i}\};$
- (iii) condition (ϕ_i^4) for all $\ell \leq j < k-1$;
- (iv) condition (ψ_j) for $j \in \{\ell, k-1\}$.

The argument that all pairs (u^j, v^j) , $j = 1, \ldots, k$, and (\hat{u}^j, \hat{v}^j) , $j = 2, \ldots, \ell - 1$, and $(\check{u}^j, \check{v}^j)$, $j = 1, \dots, \ell$, are distinct, is straightforward.

As before in case (eb1), we consider Hamilton paths P_j from u^j to v^j for all $j = 1, \ldots, k$, Hamilton paths \hat{P}_j from \hat{u}^j to \hat{v}^j for all $j=2,\ldots,\ell-1$, and Hamilton paths \check{P}_j from \check{u}^j to \check{v}^j

The concatenation $(P_1Q'\check{P}_1)(P_2\hat{P}_2\check{P}_2)(P_3\hat{P}_3\check{P}_3)\cdots(P_{\ell-1}\hat{P}_{\ell-1}\check{P}_{\ell-1})P_{\ell}P_{\ell+1}\cdots P_{k-2}P_{k-1}Q''\check{P}_{\ell}P_k$ is a Hamilton path in G(a) from x to y.

This completes the proof of Lemma 16.

The strategy for proving Lemma 19 and 20 is the same as for the proof of Lemma 16, and these proofs can be found in the appendix. The crucial difference is that now one or two building blocks $c \prec b$ for the unique $b \prec a$ with $\Delta(b) = 2$ are only assumed to satisfy the weaker property $G(c) \in \mathcal{L}_{12}$, and not $G(c) \in \mathcal{H}$, so we have to work around this by imposing extra conditions on those building blocks.

Proof of Lemma 21. As the symbols 1 and 2 both appear with the same frequency α , it suffices to prove that G(a) has property \mathcal{L}_{12} for s=1. Let x,y be two distinct vertices in G(a) such that $x_1 = 1$ and $y_1 = t$ for which there is a position $\hat{i} > 1$ with $(x_{\hat{i}}, y_{\hat{i}}) = (p_3(1, t), q_3(1, t))$.

We first consider the case $t \in \{1,3\}$. In this case we have $(x_{\hat{i}},y_{\hat{i}})=(p_3(1,t),q_3(1,t))=(3,1)$ by (4). We choose multiset permutations $u^j, v^j \in \Pi(a), j = 1, 2, 3$, satisfying the following conditions; see Figure 15 (a):

- (i) $u^1 = x \text{ and } v^3 = y;$
- (ii) $u_1^2 = v_i^1 = 3$, $u_i^2 = v_1^1 = 2$, $u_1^3 = v_i^2 = 2$, $u_i^3 = v_1^2 = 1$, and $u_i^j = v_i^{j-1}$ for j = 2, 3 and all

Note that $u_1^1 \neq v_1^1$, $u_1^2 \neq v_1^2$, and $u_1^3 \neq v_1^3$ by (i)+(ii). Consequently, we have $u^j \neq v^j$ for j=1,2,3. We define $b := (\alpha, \alpha) \prec a$, and we consider a Hamilton path P_1 in the graph $G^{\hat{i},3}(a) \simeq G(b)$ from u^1 to v^1 , which exists by the assumption $G(b) \in \mathcal{L}$, using that u^1 and v^1 belong to the two distinct partition classes of the graph $G^{\hat{i},3}(\boldsymbol{a})$ by Lemma 12 (recall that $(u_1^1,v_1^1)=(1,2)$). We




FIGURE 15. Illustration of the proof of Lemma 21.

define $b := (\alpha, \alpha - 1, 1) \prec a$, and we consider a Hamilton path P_2 in the graph $G^{\hat{i},2}(a) \simeq G(b)$ from u^2 to v^2 , which exists by the assumption $G(b) \in \mathcal{L}_1$, using that $u^2 \notin \Pi(a)^{1,1}$ and $v^2 \in \Pi(a)^{1,1}$ (recall that $(u_1^2, v_1^2) = (3, 1)$). We define $\mathbf{a}' := (\alpha - 1, \alpha, 1)$ and $\mathbf{b} := \varphi(\mathbf{a}') = (\alpha, \alpha - 1, 1) \prec \mathbf{a}$, and we consider a Hamilton path P_3 in the graph $G^{\hat{i},1}(\boldsymbol{a}) \simeq G(\boldsymbol{b})$ from u^3 to v^3 , which exists by the assumption $G(\mathbf{b}) \in \mathcal{L}_1$, using that $u^3 \in \Pi(\mathbf{a})^{1,2}$ and $v^3 \notin \Pi(\mathbf{a})^{1,2}$ (recall that $(u_1^3, v_1^3) = (2, t)$ and $t \in \{1,3\}$). Note here that in $G^{\hat{i},1}(a)$, the symbol 2 is the most frequent one and takes the role of 1 in $G(\boldsymbol{b})$.

The concatenation $P_1P_2P_3$ is a Hamilton path in G(a) from x to y.

It remains to consider the case t=2. In this case we have $(x_{\hat{i}},y_{\hat{i}})=(p_3(1,t),q_3(1,t))=(2,1)$ by (4). We choose multiset permutations $u^j, v^j \in \Pi(a), j = 1, 2, 3$, satisfying the following conditions; see Figure 15 (b):

- (i) $u^1 = x \text{ and } v^3 = y$:
- (i) $u_1^2 = v_1^1 = 2$, $u_1^2 = v_1^1 = 3$, $u_1^3 = v_1^2 = 3$, $u_1^3 = v_1^2 = 1$, and $u_i^j = v_i^{j-1}$ for j = 2, 3 and all

Note that $u_1^1 \neq v_1^1$, $u_1^2 \neq v_1^2$, and $u_1^3 \neq v_1^3$ by (i)+(ii). Consequently, we have $u^j \neq v^j$ for j = 1, 2, 3. We consider a Hamilton path P_1 in $G^{\hat{\imath},2}(a)$ from u^1 to v^1 , which exists by the assumption $G(\alpha, \alpha - 1, 1) \in \mathcal{L}_1$, using that $u^1 \in \Pi(\boldsymbol{a})^{1,1}$ and $v^1 \notin \Pi(\boldsymbol{a})^{1,1}$ (recall that $(u_1^1, v_1^1) = (1, 3)$). We also consider a Hamilton path P_2 in $G^{\hat{i},3}(a)$ from u^2 to v^2 , which exists by the assumption $G(\alpha, \alpha) \in \mathcal{L}$, using that u^2 and v^2 belong to the two distinct partition classes of the graph $G^{\hat{i},3}(\boldsymbol{a})$ by Lemma 12 (recall that $(u_1^2, v_1^2) = (2, 1)$). Finally, we consider a Hamilton path P_3 in $G^{\hat{i},1}(\boldsymbol{a})$ from u^3 to v^3 , which exists by the assumption $G(\alpha, \alpha - 1, 1) \in \mathcal{L}_1$, using that $u^3 \notin \Pi(\boldsymbol{a})^{1,2}$ and $v^3 \in \Pi(a)^{1,2}$ (recall that $(u_1^3, v_1^3) = (3, 2)$).

The concatenation $P_1P_2P_3$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

The proof of Lemma 22 follows the same strategy as the proof of Theorem 5 presented in Section 1.5.2.

Proof of Lemma 22. We argue by induction on $k \geq 5$, using as a base case that $G(2,1,1,1) \in \mathcal{H}$ by Table 1. Let x, y be two distinct vertices in G(a). As $x \neq y$ there is a position $\hat{i} > 1$ such that $x_{\hat{i}} \neq y_{\hat{i}}$. As $k \geq 5$, there is an index $i_1 \in [n] \setminus \{1, \hat{i}\}$ such that $x_{i_1} \notin \{x_1, x_{\hat{i}}, y_{\hat{i}}\}$. Similarly, there is an index $i_k \in [n] \setminus \{1, \hat{\imath}\}$ such that $y_{i_k} \notin \{y_1, x_{\hat{\imath}}, y_{\hat{\imath}}, x_{i_1}\}$. We fix any permutation π on [k]such that $\pi_1 = x_i$, $\pi_2 = x_{i_1}$, $\pi_{k-1} = y_{i_k}$, and $\pi_k = y_i$. For convenience, we also define $\pi_0 := x_1$ and $\pi_{k+1} := y_1$. We let $\hat{j} \in [k]$ be such that $\pi_{\hat{j}} = 1$. Then we choose a sequence of multiset permutations $u^j, v^j \in \Pi(\mathbf{a}), j = 1, \dots, k$, satisfying the following conditions; see Figure 16:

- (i) $u^1 = x \text{ and } v^k = y$;
- (i) $u_1^{-} = x$ and $v_1^{-} = y$, (ii) $u_1^j = v_{\hat{i}}^{j-1} = \pi_{j-1}$, $u_{\hat{i}}^j = v_1^{j-1} = \pi_j$, and $u_i^j = v_i^{j-1}$ for all $2 \le j \le k$ and $i \in [n] \setminus \{1, \hat{i}\}$. (iii) $u_{\hat{i}}^j = v_{\hat{i}}^j$ for all $i \in [n] \setminus \{1, \hat{i}, i_{\hat{j}}\}$, where $i_{\hat{j}} \in [n] \setminus \{1, \hat{i}\}$ is chosen such that $u_{\hat{i}_{\hat{j}}}^j = \pi_{\hat{j}+1}$ and $v_{i_{\hat{j}}}^{\hat{j}} = \pi_{\hat{j}-1}$ (in particular, if $\hat{j} = 1$ then we have $i_{\hat{j}} = i_1$, and if $\hat{j} = k$ then we have $i_{\hat{j}} = i_k$).

Recall that $\pi_0 \neq \pi_2$ by the choice of i_1 , and $\pi_{k+1} \neq \pi_{k-1}$ by the choice of i_k . Consequently, using that $u_1^j = \pi_{j-1}$ and $v_1^j = \pi_{j+1}$ by (ii) we obtain that $u^j \neq v^j$ for all $1 \leq j \leq k$.

For j = 1, ..., k we define $\mathbf{a}' := (a_1, ..., a_{\pi_j} - 1, ..., a_k)$ and $\mathbf{b} := \varphi(\mathbf{a}')$. Note that $\mathbf{b} = (2, 1^{k-2})$ for all $j \in [k] \setminus \{\hat{j}\}$, whereas $b = 1^k$ for $j = \hat{j}$. In the first case, there is a Hamilton path P_i in the graph $G^{\hat{\imath},\pi_j}(\boldsymbol{a}) \simeq G(\boldsymbol{b}) = G((2,1^{k-2}))$ from u^j to v^j by induction. In the second case, there is a Hamilton path P_i in the graph $G^{i,1}(a) \simeq G(b) = G(1^k)$ from u^j to v^j by Theorem 5, using that u^{j} and v^{j} differ only in one transposition by (ii)+(iii), i.e., they belong to two distinct partition classes of the graph $G^{\hat{i},1}(\boldsymbol{a})$ by Lemma 13. The concatenation $P_1P_2\cdots P_k$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.




FIGURE 16. Illustration of the proof of Lemma 22.

5. Proof of Theorems 8 and 9

In this section we prove Theorems 8 and 9. For this, we focus entirely on integer partitions $\mathbf{a} = (a_1, \ldots, a_k)$ that satisfy $\Delta(\mathbf{a}) = 0$. By (2), this is equivalent to $a_1 = \sum_{i=2}^k a_i$, i.e., the first symbol appears equally often as all other symbols combined. Based on this observation, we will first reparametrize the problem for convenience, changing the meaning of the variables n, k, \mathbf{a} .

5.1. **Reparametrizing the problem.** The following notations are illustrated in Figure 17.

For any $n \geq 1$ and $k \geq 1$, we consider multiset permutations with n symbols 0, and with n non-zero symbols from the set $\{1,\ldots,k\}$. Specifically, the number of each non-zero symbol $i \in [k]$ is given by $a_i \geq 1$, where $n = a_1 + \cdots + a_k$ and $a_1 \geq a_2 \geq \ldots \geq a_k$. We write $\Pi_n(\mathbf{a})$, $\mathbf{a} := (a_1,\ldots,a_k)$, to denote those multiset permutations. For any $x \in \Pi_n(\mathbf{a})$, we write x^- for the string obtained from x by omitting the first entry, and we define $\Pi_n(\mathbf{a})^- := \{x^- \mid x \in \Pi_n(\mathbf{a})\}$. Given any $x \in \Pi_n(\mathbf{a})^-$, we can uniquely infer the omitted symbol, and we refer to it as the suppressed symbol, denoted $s(x) \in \{0,\ldots,k\}$.




FIGURE 17. The graph from Figure 5 in the reparametrized notation.

Clearly, $x, y \in \Pi_n(\mathbf{a})$ differ in a star transposition if and only if $x^-, y^- \in \Pi_n(\mathbf{a})^-$ differ in exactly one position. We also write $\Pi_n(\mathbf{a})^{-0}$ and $\Pi_n(\mathbf{a})^{-1}$ for all multiset permutations from $\Pi_n(\mathbf{a})^-$ with suppressed symbol 0 or suppressed symbol distinct from 0, respectively. We write $G_n(\mathbf{a})$ for the graph with vertex set $\Pi_n(\mathbf{a})^-$, with an edge between any two multiset permutations that

differ in exactly one position. By Lemma 14, any Hamilton cycle in $G_n(\boldsymbol{a})$ will alternately visit the vertex sets $\Pi_n(\boldsymbol{a})^{-0}$ and $\Pi_n(\boldsymbol{a})^{-1}$.

- 5.2. **Proof of Theorem 8.** We now consider the graph $G_n := G_n(n) \simeq G(n,n)$ for $n \geq 3$. The fact that it is Hamilton-laceable for n=3 and n=4 follows from Table 1. For the rest of this section, we will therefore assume that $n \geq 5$. We follow the proof strategy outlined in Section 1.5.3.
- 5.2.1. Translation into the hypercube. We introduce the abbreviations $B_n := \Pi_n(n)^{-1}$ and $B'_n := \Pi_n(n)^{-0}$ for the sets of bitstrings of length 2n-1 with exactly n-1 or n many 1s, respectively. Note that the graph G_n is the subgraph of the (2n-1)-dimensional hypercube induced by the 'middle levels' B_n and B'_n . In the following we will show that G_n , $n \ge 5$, is Hamilton-laceable.

We first reduce the number of different pairs of vertices $x \in B_n$, $y \in B'_n$ that we need to connect by a Hamilton path to only to n cases, one for each possible Hamming distance d := d(x, y), where $1 \le d \le 2n - 1$ is odd.

Lemma 24. For any two pairs of vertices $(x_1, y_1), (x_2, y_2) \in B_n \times B'_n$ with $d(x_1, y_1) = d(x_2, y_2)$ there is an automorphism φ of G_n such that $\varphi(x_1) = x_2$ and $\varphi(y_1) = y_2$.

Proof. Take an arbitrary permutation φ_1 of coordinates such that $\varphi_1(x_1) = x_2$ and define $z := \varphi_1(y_1)$. Since $d(x_2, y_2) = d(x_2, z) =: d$, both y_2 and z share with x_2 exactly (2n - 1 - d)/2 coordinates set to 1. Moreover, both y_2 and z have exactly (d+1)/2 coordinates set to 1 where x_2 has a 0. Clearly, there is a permutation φ_2 that maps these two sets of coordinates of z to the corresponding sets of y_2 and preserves the other coordinates. As $\varphi_2(z) = y_2$ and $\varphi_2(x_2) = x_2$, the desired automorphism is the composition of φ_1 and φ_2 .

5.2.2. Cycle factor construction. We now recall the construction of a cycle factor C_n in the graph G_n described in [MMM21]. For any bitstring x and any integer ℓ , we write $\sigma^{\ell}(x)$ for the bitstring obtained from x by cyclic left rotation by ℓ steps. If ℓ is negative, then this is a right rotation by $|\ell|$ steps. We write D_n for the set of all $Dyck\ words$, i.e., binary strings of length 2n-2 with n-1 many 0s and 1s, such that in every prefix, there are at least as many 1s as 0s. We also define $D:=\bigcup_{n\geq 1}D_n$. Any Dyck word x can be decomposed uniquely as $x=1\ u\ 0\ v$ for $u,v\in D$, and as $x=u\ 1\ v\ 0$ for $u,v\in D$. Moreover, any Dyck word $x=1\ u\ 0\ v\in D_n$ can be interpreted as an (ordered) rooted tree T with n vertices as follows; see Figure 18 (a): The leftmost child of the root of T leads to the subtree defined by u, and the remaining children of the root of T form the subtree defined by v. A tree rotation operation transforms the tree $x=1\ u\ 0\ v$ into $\rho(x):=u\ 1\ v\ 0$; see Figure 18 (a). For any rooted tree $x\in D$ we write $[x]:=\{\rho^i(x)\mid i\geq 0\}$ for the set of all trees obtained from x by 'forgetting' the root. We write T_n for the set of all plane trees with n vertices.

For any bitstring $x \in B_n$ there is a unique integer $\ell := \ell(x)$, $0 \le \ell \le 2n-2$, such that the first 2n-2 bits of $\sigma^{\ell}(x)$ are a Dyck word. Similarly, for any bitstring $y \in B'_n$ there is a unique integer $\ell := \ell(y)$, $0 \le \ell \le 2n-2$, such that the last 2n-2 bits of $\sigma^{\ell}(y)$ are a Dyck word. We refer to $\ell(x)$ as the *shift* of x or y, respectively. Moreover, we write $t(x) \in D_n$ for the first 2n-2 bits of $\sigma^{\ell}(x)$, and t(y) for the last 2n-2 bits of $\sigma^{\ell}(y)$, and we interpret them as a rooted trees with n vertices.

For any vertex $x \in B_n$ we define vertices $f(x), g(x) \in B'_n$ by considering the unique decomposition $\sigma^{\ell}(x) = t(x) \ 0 = 1 \ u \ 0 \ v \ 0$, where $0 \le \ell \le 2n - 2$ and $u, v \in D$, and by setting

$$f(x) := \sigma^{-\ell}(1 u 1 v 0) \in B'_n, \quad g(x) := \sigma^{-\ell}(1 u 0 v 1) \in B'_n.$$
 (8a)

Clearly, f(x) and g(x) are two distinct neighbors of x in the graph G_n . For any vertex $y \in B'_n$ we define vertices $f(y), g(y) \in B_n$ by considering the unique decomposition $\sigma^{\ell}(y) = 1 t(y) = 1 u 1 v 0$,




FIGURE 18. (a) Correspondence between cycles in C_n and tree rotation. The gray boxes highlight the Dyck substrings. Part (b) of the figure is explained in Section 5.3.1.

where $0 \le \ell \le 2n - 2$ and $u, v \in D$, and by setting

$$f(y) := \sigma^{-\ell}(0 u 1 v 0) \in B_n, \quad g(y) := \sigma^{-\ell}(1 u 0 v 0) \in B_n.$$
 (8b)

Clearly, f(y) and g(y) are two distinct neighbors of x in the graph G_n . The definition of these mappings is illustrated in Figures 18 (a) and 24 (a). For any $x \in B_n \cup B'_n$ we define $C(x) := (x, f(x), f^2(x), \ldots)$.

The following lemma was shown in [MMM21].

Lemma 25 ([MMM21]). Let $n \ge 1$.

- (i) For any $x \in B_n \cup B'_n$ we have g(f(x)) = x and f(g(x)) = x, i.e., f and g are inverse mappings. Consequently, $M := \{(x, f(x)) \mid x \in B_n\}$ and $N := \{(g(x), x) \mid x \in B_n\}$ are two disjoint perfect matchings in the graph G_n and $C_n := M \cup N = \{C(x) \mid x \in B_n \cup B'_n\}$ is a cycle factor of G_n .
- (ii) For any $x \in B_n$ we have $t(f(f(x))) = \rho(t(x))$ and $\ell(f(f(x))) = \ell(x) + 1 \mod (2n-1)$. In words, the next vertex from B_n that follows after x on the same cycle of C_n is obtained by rotating the tree t(x) and incrementing the shift.
- (iii) For any $x \in B_n$ we have $C(x) \cap B_n = \{\sigma^i(x'0) \mid 0 \le i \le 2n 2 \land x' \in [t(x)]\}$. Consequently, the cycles of C_n are in bijection to the set T_n of plane trees with n vertices.

Properties (ii) and (iii) are illustrated in Figures 18 (a) and 24 (a). One can check that the perfect matchings M and N defined in part (i) of the lemma are in fact the (n-2)-lexical and (n-1)-lexical matchings defined in [KT88] and used in [GMN18]. We will think of the cycles of \mathcal{C}_n as being oriented in the direction of applying f, oppositely to $g = f^{-1}$.

The following property of the cycle factor C_n was established in [GMN18].

Lemma 26 ([GMN18]). The mapping $h: B_n \cup B'_n \to B_n \cup B'_n$ defined by $(x_1, \ldots, x_{2n-1}) \mapsto (\overline{x_{2n-2}}, \overline{x_{2n-3}}, \ldots, \overline{x_1}, \overline{x_{2n-1}})$, where overline denotes complementation, is an automorphism of G_n that maps C_n onto itself.

5.2.3. Gluing cycles. We now describe how to join the cycles of the factor C_n to a single Hamilton cycle, by taking the symmetric difference with suitably chosen 6-cycles.




FIGURE 19. A gluing pair (x, y) with corresponding rooted trees.




FIGURE 20. The vertices x^i , y^i and the cycle C(x, y).

We consider two Dyck words $x, y \in D_n$ of the form

$$x = 110u0v, \quad y = 101u0v, \quad \text{with } u, v \in D.$$
 (9)

We refer to such a pair (x, y) as a *gluing pair*, and we write \mathcal{G}_n for the set of all gluing pairs. The rooted trees corresponding to x and y are shown in Figure 19. Note that the tree y is obtained from x by removing the leftmost edge that leads from the leftmost child of the root of x to a leaf, and reattaching this edge as the leftmost child of the root.

For any gluing pair $(x, y) \in \mathcal{G}_n$ we define $x^i := f^i(x 0)$ and $y^i := f^i(y 0)$ for $i \ge 0$. Using the definition (8), a straightforward calculation yields the vertices shown in Figure 20.

Note that

$$C(x,y) := (x^0, x^1, x^6, x^5, y^0, y^1)$$
(10)

is a 6-cycle in the graph G_n ; see Figure 20. This 6-cycle has the edges (x^0, x^1) and (x^6, x^5) in common with the cycle $C(x^0)$, and the edge (y^0, y^1) in common with the cycle $C(y^0)$. One can check that the symmetric difference of the edge sets $(C(x^0) \cup C(y^0)) \triangle C(x, y)$ is a single cycle on the vertex set of $C(x^0) \cup C(y^0)$. In other words, the cycle C(x, y) glues together two cycles from the cycle factor to a single cycle. For any set of gluing pairs $\mathcal{F} \subseteq \mathcal{G}_n$ we write $C(\mathcal{F}) := \{C(x, y) \mid (x, y) \in \mathcal{F}\}.$

The following results about the cycle factor C_n and the set of gluing pairs G_n were proved in [GMN18]; see this paper for illustrations.

Lemma 27 ([GMN18]). For any two gluing pairs $(x,y), (x',y') \in \mathcal{G}_n$, the 6-cycles C(x,y) and C(x',y') are edge-disjoint. For any two gluing pairs $(x,y), (x,y') \in \mathcal{G}_n$, the two pairs of edges that the two 6-cycles C(x,y) and C(x,y') have in common with the cycle $C(x^0)$ are not interleaved.

Lemma 28 ([GMN18]). For any $n \geq 4$, there is a set $\mathcal{T}_n \subseteq \mathcal{G}_n$ of gluing pairs of cardinality $|\mathcal{T}_n| = |\mathcal{C}_n| - 1 = |T_n| - 1$, such that $\{([x], [y]) \mid (x, y) \in \mathcal{T}_n\}$ is a spanning tree on the set of plane trees T_n , implying that the symmetric difference $\mathcal{C}_n \triangle C(\mathcal{T}_n)$ is a Hamilton cycle in G_n .

5.2.4. Alternating path. In the following, we describe how to join two vertices $x \in B_n$, $y \in B'_n$ with Hamming distance d(x,y) = d in the graph G_n via some cycles from C_n to a short path between x and y.

We say that a path P between two vertices x and y of G_n is C_n -alternating if it alternately uses edges and non-edges of C_n , starting and ending with an edge of C_n , and the symmetric difference $P \triangle C_n$ yields a path between x and y that contains all vertices of the intersected cycles; see Figure 21. Note that the resulting path will then also contain all vertices of P. In other words, the path P glues together all intersected cycles to a single path between x and y, while the non-intersected cycles are unchanged. Note that P is allowed to intersect a single cycle of C_n multiple times. The C_n -alternating paths we use in our proof are obtained as subpaths




FIGURE 21. Example of a C_n -alternating path between x and y.

of one fixed C_n -alternating path between two complementary vertices with maximum possible Hamming distance 2n-1.

Lemma 29. For $n \geq 5$, let P be the path defined in Figure 22 between the two vertices $x_1 \in B_n$ and $y_n \in B'_n$. For every i = 1, ..., n, the subpath of P between $x_1 \in B_n$ and $y_i \in B'_n$ is a C_n -alternating path and $d(x_1, y_i) = 2i - 1$.




FIGURE 22. Definition of the C_n -alternating path P in Lemma 29. The gray boxes highlight the Dyck substrings of the path vertices from B_n . The corresponding rooted trees are shown to the right, as well as the shifts and the Hamming distance from the first vertex x_1 . The subtrees s_i are stars with i vertices rooted at the center vertex.

Proof. By counting the positions in which x_i and y_i differ from x_1 , it is straightforward to check that $d(x_1, x_i) = 2i - 2$ and $d(x_1, y_i) = 2i - 1$ for all i = 1, ..., n. Note that the Hamming distance from x_1 increases monotonically along P, with the only exception being the step being from x'

to y', which satisfy $d(x_1, x') = 2n - 4$ and $d(x_1, y') = 2n - 5$. Next, we observe that every edge (x_i, y_i) for i = 1, ..., n and the edge (x', y') belong to a cycle from C_n . Specifically, by (8a) we have $y_i = f(x_i)$ for all i = 1, ..., n - 4 and $y_i = g(x_i) = f^{-1}(x_i)$ for i = n - 3, ..., n, and y' = f(x'). Furthermore, all other edges of P are not on a cycle from C_n , which can be verified by (8a).

We now determine which cycles of C_n the path P intersects. For this we consider the rooted trees $t(x_i)$ for i = 1, ..., n and t(x'); see Figure 22. For any $i \ge 1$ we define the rooted tree $s_i := (10)^{i-1} \in D_i$, which is the star with i vertices rooted at the center vertex. With this abbreviation we obtain

$$t(x_1) = s_{n-3} 110010,$$
 $t(x_{n-3}) = 110 s_{n-3} 010,$ $t(x_n) = 10110 s_{n-3} 0,$
 $t(x_i) = 1 s_i 0 s_{n-i-3} 110010$ for $i = 2, ..., n-4,$ $t(x_{n-1}) = 1100 s_{n-3} 10,$ (11)
 $t(x_{n-2}) = t(x') = s_n.$




FIGURE 23. The intersection of the path P and subpaths $Q \subseteq P$ with the cycles C_2 and C_3 .

Note that $[t(x_1)] = [t(x_{n-3})] = [t(x_n)]$ is the same underlying plane tree, so the vertices x_1 , x_{n-3} and x_n lie on the same cycle of \mathcal{C}_n by Lemma 25 (iii). Similarly, $t(x_{n-2}) = t(x') = s_n$, so the vertices x_{n-2} and x' lie on the same cycle of \mathcal{C}_n . From (11) we see that the tree $t(x_1)$ has diameter 3 and $t(x_{n-2})$ has diameter 2, so these two cycles are distinct. We claim that the remaining vertices x_i for $i = 2, \ldots, n-4$ and x_{n-1} all lie on their own cycle of \mathcal{C}_n that is also distinct from the previous two cycles. To see this, we use again Lemma 25 (iii), noting that $t(x_i)$ and $t(x_{n-1})$ have diameter 4, $[t(x_i)] \neq [t(x_j)]$ for any two distinct indices $i, j \in \{2, \ldots, n-4\}$, and moreover $t(x_{n-1})$ has two adjacent vertices of degree 2 unlike $t(x_i)$ for $i = 2, \ldots, n-4$.

Consequently, the entire path P intersects the cycle $C_3 := C(x_1) = C(x_{n-3}) = C(x_n)$ three times, the cycle $C_2 := C(x_{n-2}) = C(x')$ twice, and every other (intersected) cycle only once.

We now check that the symmetric differences $P \triangle C_2$, $P \triangle C_3$, and $P \triangle (C_2 \cup C_3)$ yield a single path containing all vertices of C_2 , C_3 , or $C_2 \cup C_3$, respectively. For $P \triangle C_2$ this can be seen easily, by noting that the first edge (x_{n-2}, y_{n-2}) of P intersecting C_2 is a backward edge of C_2 (i.e., $y_{n-2} = g(x_{n-2}) = f^{-1}(x_{n-2})$), whereas the second edge (x', y') of P intersecting C_2 is a forward edge of C_2 (i.e., y' = f(x')). The intersection pattern between P and P and P and P is shown in Figure 23 (a). This figure shows the cycle P and the corresponding shifts P and that lie on this cycle, the corresponding rooted trees P and the corresponding shifts P are given by Lemma 25 (ii). The shifts P and P and P and P and P are given by the definition of P, and result in the shown location of the three edges of P on P and P are given by the definition of P and result in the shown location of the three edges of P and P are given by the definition of P and result in the shown location of the three edges of P and P are given by the definition of P and result in the shown location of the three edges of P and P are given by the definition of P and result in the shown location of the three edges of P and P are given by the definition of P and result in the shown location of the three edges of P and P are given by the definition of P and P are given by the definition of P and P are given by the definition of P and P are given by the definition of P and P are given by the definition of P and P are given by P are given by P and P are given by P are given by P and P are given by P an

So far we have shown that the path P is C_n -alternating. However, the statement of the lemma is stronger, and asserts that every subpath $Q \subseteq P$ between x_1 and y_i for i = 1, ..., n is also C_n -alternating. The only non-trivial cases to consider are when Q intersects C_2 or C_3 twice, which happens precisely when Q ends at y_{n-1} , y_{n-2} or y_{n-3} , and those can be checked to work in Figure 23 (c).

This completes the proof of the lemma.

5.2.5. Proof of Theorem 8. We are now in position to present the proof of Theorem 8.

Proof of Theorem 8. We need to show that $G_n = G_n(n) \simeq G(n,n)$, $n \geq 3$, is Hamilton-laceable. The cases n=3 and n=4 are covered by Table 1, so we will assume that $n\geq 5$. Let d be an odd integer with $1\leq d\leq 2n-2$. Moreover, let \mathcal{C}_n be the cycle factor in G_n defined in Lemma 25 (i), and let Q be the path given by Lemma 29 between vertices $x\in B_n$ and $y\in B'_n$ with d(x,y)=d. Now consider the cyclically shifted path $Q':=\sigma^{-4}(Q)$, connecting $x':=\sigma^{-4}(x)$ and $y':=\sigma^{-4}(y)$, which also satisfy d(x',y')=d. By Lemma 24, the theorem is proved by exhibiting a Hamilton path from x' to y', which we will do in the following. All except possibly the last vertex of Q have a 0-bit at position 2n-5; see the highlighted column in Figure 22. It follows that all except possibly the last vertex of Q' have a 0-bit at their last position. As Q is \mathcal{C}_n -alternating by Lemma 29 and the definition of f in (8) is invariant under cyclic rotation of bitstrings, the path Q' is also \mathcal{C}_n -alternating. Let r denote the number of cycles of \mathcal{C}_n intersected by the path Q'. Then the symmetric difference $Q' \triangle \mathcal{C}_n$ is a path between x' and y' plus a set of $|\mathcal{C}_n| - r$ cycles, and together they visit all vertices of G_n .

Now consider a set $\mathcal{T}_n \subseteq \mathcal{G}_n$ of gluing pairs with the properties guaranteed by Lemma 28 and the corresponding 6-cycles $\mathcal{S} := C(\mathcal{T}_n) = \{C(x,y) \mid (x,y) \in \mathcal{T}_n\}$. We consider the image $\mathcal{S}' := h(\mathcal{S})$ of those 6-cycles under the automorphism h stated in Lemma 26. As this automorphism maps \mathcal{C}_n onto itself, we have that $\mathcal{C}_n \triangle \mathcal{S}'$ is a Hamilton cycle in G_n . From

Figure 20 we see that for every 6-cycle $C(x,y) \in \mathcal{S}$, the last bit of all vertices $x^0, \ldots, x^6, y^0, y^1$ is 0, and by the definition of h, the last bit of the corresponding vertices in $h(\mathcal{S}) = \mathcal{S}'$ is 1. Consequently, all edges between these vertices are disjoint from the path Q'. As r cycles of \mathcal{C}_n are already joined by Q', we can discard r-1 of the 6-cycles from the set \mathcal{S}' to obtain a set \mathcal{S}'' of 6-cycles of cardinality $|\mathcal{S}''| = |\mathcal{S}'_n| - (r-1) = |\mathcal{C}_n| - r$ such that $(Q' \triangle \mathcal{C}_n) \triangle \mathcal{S}''$ is a Hamilton path in G_n between x' and y'. This completes the proof of the theorem.

- 5.3. **Proof of Theorem 9.** The basic strategy for proving Theorem 9 is very similar to the proof of Theorem 8. We first construct a cycle factor in the graph, and we then join its cycles to a single Hamilton cycle. The construction of the cycle factor and of the gluing cycles is achieved by carefully generalizing the constructions described in Sections 5.2.2 and 5.2.3. These two steps work for any graph $G_n(\mathbf{a})$, and only the last step of constructing a Hamilton cycle is done specifically for the case $\mathbf{a} = (n-1,1)$ required for Theorem 9.
- 5.3.1. A cycle factor in $G_n(\mathbf{a})$. For any $n \geq 1$ and any integer partition \mathbf{a} of n, we now describe how to construct a cycle factor $C_n(\mathbf{a})$ in $G_n(\mathbf{a})$.

Given any string x over the alphabet $\{0, 1, \ldots, k\}$, we write \widehat{x} for the bitstring obtained from x by replacing all non-zero symbols by 1. The mappings ℓ , t and f defined in Section 5.2.2 on bitstrings can be generalized to operate on multiset permutations in the natural way. Specifically, for any $x \in \Pi_n(\mathbf{a})^-$, we define $\ell(x) := \ell(\widehat{x})$. Moreover, t(x) is the substring of x of length 2n-2 starting at position $\ell(x) + 1$ (modulo the length 2n-1). Similarly, $f(x) \in \Pi_n(\mathbf{a})^-$ is obtained by considering the position i in which \widehat{x} and $f(\widehat{x})$ differ, and by replacing the symbol at position i in x by the suppressed symbol s(x).

For example, for $x = 00230 \in \Pi_3(1,1,1)^-$ we have s(x) = 1, $\hat{x} = 00110$, $\ell(\hat{x}) = 2$ and $t(\hat{x}) = 1100$. Consequently, we obtain $\ell(x) = 2$ and t(x) = 2300. Moreover, $f(\hat{x}) = 10110$ differs in the first bit from \hat{x} , and therefore f(x) = s(x)0230 = 10230.

For any $x \in \Pi_n(\mathbf{a})^-$ we define the cycle $C(x) := (x, f(x), f^2(x), \ldots)$, and we define the cycle factor $C_n(\mathbf{a}) := \{C(x) \mid x \in \Pi_n(\mathbf{a})^-\}$.

We also define $D_n(a) := \{x \mid x \ 0 \in \Pi_n(a)^{-1} \land \widehat{x} \in D_n\}$. We can interpret any $x \in D_n(a)$ as a vertex-labeled rooted tree with n vertices, which has precisely a_i vertices labeled i for all $i = 1, \ldots, k$ (recall that $n = \sum_{i=1}^k a_i$); see Figures 18 and 24. For this recall the interpretation of the Dyck word $\widehat{x} \in D_n$ as a rooted tree with n vertices, which can be expressed iteratively as follows: We start by adding a root vertex. We then read \widehat{x} from left to right, and for every 1-bit encountered we add a new rightmost child below the current vertex and we move to this vertex, and for every 0-bit encountered we move from the current vertex to its parent, without adding any new vertices. Now the tree corresponding to x is obtained as follows: We start by adding a root vertex with label $s(x \ 0)$. We then read the string x from left to right, and for every non-zero symbol i > 0 encountered we add a new rightmost child with label i below the current vertex and we move to this vertex, and for every symbol 0 encountered we move from the current vertex to its parent, without adding any new vertices.

Tree rotations also generalize straightforwardly to the labeled setting. Specifically, for $x = b u 0 v \in D_n(\mathbf{a})$ with $\widehat{u}, \widehat{v} \in D$ and $a, b \in [k]$ such that a = s(x 0) we define $\rho(x) := u a v 0$, which corresponds to rotating the tree together with its vertex labels (note that $s(\rho(x) 0) = b$).

With these definitions the properties asserted in Lemma 25 about the factor C_n generalize straightforwardly to the labeled version $C_n(\mathbf{a})$. Specifically, if $x = \Pi_n(\mathbf{a})^{-1}$, then the tree t(f(f(x))) is obtained from t(x) by a labeled tree rotation ρ ; see Figure 18 (b) (and the shift is incremented, as before). Consequently, the cycles of $C_n(\mathbf{a})$ are in bijective correspondence with vertex-labeled plane trees with n vertices, exactly a_i of which have the label i for all $i = 1, \ldots, k$.




FIGURE 24. Construction of the cycle factor $C_n(a)$ in the graph $G_n(a)$ (right hand side) from the cycle factor C_n in the graph G_n (left hand side).

For example, the cycle factor $C_3(1,1,1)$ shown in Figure 24 has three cycles which correspond to all plane trees with three vertices with vertex labels 1,2,3, namely the path P_3 on three vertices with either 1,2, or 3 as the center vertex. The cycle factor $C_3(2,1)$ on the other hand, has only two cycles, corresponding to P_3 with the label 2 either at the center vertex or at a leaf. For comparison, the cycle factor $C_3(3)$ has only a single cycle corresponding to P_3 with all vertices labeled 1.

5.3.2. Labeled gluing cycles. We generalize the construction of gluing cycles described in Section 5.2.3 to the labeled case. Specifically, we consider six Dyck words $x_1, x_2, x_3, x_4, y_1, y_2 \in D_n(\mathbf{a})$ of the form

$$x_{1} = b c 0 u 0 v, y_{1} = b 0 c u 0 v,$$

$$x_{2} = c b 0 u 0 v, y_{2} = b 0 a u 0 v,$$

$$x_{3} = b a 0 u 0 v,$$

$$x_{4} = a b 0 u 0 v, \text{with } \hat{u}, \hat{v} \in D \text{ and } a, b, c \in [k].$$

$$(12)$$

Note that $s(x_1) = s(x_2) = s(y_1) = a$ and $s(x_3) = s(x_4) = s(y_2) = c$. We refer to such a 6-tuple $(x_1, x_2, x_3, x_4, y_1, y_2)$ as a gluing tuple, and we write $\mathcal{G}_n(a)$ for the set of all gluing tuples. Note

that by this definition we have $s(x_1) = s(x_2) = s(y_1) = a$ and $s(x_3) = s(x_4) = s(y_2) = c$. Also note that $\widehat{x_1} = \widehat{x_2} = \widehat{x_3} = \widehat{x_4} =: x$ and $\widehat{y_1} = \widehat{y_2} =: y$, and that (x, y) is a gluing pair as defined in (9), i.e., the rooted trees corresponding to $x_1, x_2, x_3, x_4, y_1, y_2$, which are shown in Figure 25, are vertex-labeled versions of the two trees shown in Figure 19. Observe also that some of the values a, b, c may coincide, which leads to different possible coincidences between the x_j or y_j ; see Figure 25.

For a gluing tuple $(x_1, x_2, x_3, x_4, y_1, y_2) \in \mathcal{G}_n(\boldsymbol{a})$ we define $x_j^i := f^i(x_j 0)$ for j = 1, 2, 3, 4 and $i \geq 0$, and $y_j^i := f^i(y_j 0)$ for j = 1, 2 and $i \geq 0$. It can be verified from Figure 25 that the sequence of vertices

$$C(x_1, x_2, x_3, x_4, y_1, y_2) := (x_1^0, x_1^1, x_2^6, x_2^5, y_1^0, y_1^1, x_3^0, x_3^1, x_4^6, x_4^5, y_2^0, y_2^1)$$

$$(13)$$




FIGURE 25. Definition of the cycle $C(x_1, x_2, y_1, x_3, x_4, y_2)$.

is cyclic, and any two consecutive vertices differ in one position. This sequence is obtained by applying the flip sequence of the 6-cycle in (10) two times. For any set of gluing tuples $\mathcal{F} \subseteq \mathcal{G}_n(\boldsymbol{a})$ we write $C(\mathcal{F}) := \{C(x_1, x_2, x_3, x_4, y_1, y_2) \mid (x_1, x_2, x_3, x_4, y_1, y_2) \in \mathcal{F}\}$. Note that $C(x_1, x_2, x_3, x_4, y_1, y_2)$ is a 12-cycle if $a \neq c$ and a 6-cycle if a = c. Moreover, this cycle can be used to join either 2, 3, 5 or 6 different cycles from $\mathcal{C}_n(\boldsymbol{a})$, depending on the coincidences between the values a, b, c, as shown in Figure 25.

By Lemma 27, any two distinct cycles from $C(\mathcal{G}_n(\boldsymbol{a}))$ are edge-disjoint, and if two of them intersect one cycle of $\mathcal{C}_n(\boldsymbol{a})$ twice each, then the two pairs of edges on this cycle are not interleaved. Note that the cycles $C(x_1, x_2, x_3, x_4, y_1, y_2)$ and $C(x_3, x_4, x_1, x_2, y_2, y_1)$ have the exact same set of vertices and edges, so they are the same cycle.

5.3.3. Proof of Theorem 9. The following lemma, proved in [MMM21], strengthens Lemma 28 from before. To state the lemma, we need a few more definitions. As before, we write $s_n := (10)^{n-1} \in D_n$ for the star with n vertices, rooted at the center vertex. Given a (rooted or plane) tree x, the potential of a vertex of x is the sum of distances from that vertex to all other vertices of x. Moreover, the potential of x, denoted by $\varphi(x)$, is the minimum potential of all vertices of x. For example, $\varphi(s_n) = n - 1$.

Lemma 30 ([MMM21]). For any $n \ge 4$, there is a set $\mathcal{T}_n \subseteq \mathcal{G}_n$ of gluing pairs of cardinality $|\mathcal{T}_n| = |\mathcal{C}_n| - 1 = |T_n| - 1$, such that $\{([x], [y]) \mid (x, y) \in \mathcal{T}_n\}$ is a spanning tree on the set of plane trees T_n , implying that the symmetric difference $\mathcal{C}_n \triangle C(\mathcal{T}_n)$ is a Hamilton cycle in G_n . Moreover, for every gluing pair $(x, y) \in \mathcal{T}_n$ we have $\varphi(y) = \varphi(x) - 1$, and for every plane tree $T \in T_n \setminus \{[s_n]\}$ there is exactly one gluing pair $(x, y) \in \mathcal{T}_n$ with T = [x] and $\varphi(y) = \varphi(x) - 1$.

The spanning tree described in this lemma is illustrated in Figure 26 for n = 7. We are now ready to present the proof of Theorem 9.

Proof of Theorem 9. The graphs G(2,1,1) and G(3,2,1) have a Hamilton cycle by Table 1. In the remainder of the proof we construct a Hamilton cycle in $G_n(n-1,1) \simeq G(n,n-1,1)$ for $n \geq 4$. We define $\mathbf{a} := (n-1,1)$.

We start with the cycle factor $C_n(a)$ constructed in Section 5.3.1. Its cycles are in bijective correspondence to plane trees that have a single vertex labeled 2, and all other vertices labeled 1. We say that this label-2 vertex is marked, and we refer to these trees as marked plane trees. We construct a set of gluing tuples $\mathcal{T}'_n \subseteq \mathcal{G}_n(a)$ such that $C_n(a) \triangle C(\mathcal{T}'_n)$ is a Hamilton cycle in $G_n(a)$.

We let $\mathcal{T}_n \subseteq \mathcal{G}_n$ be the set of gluing pairs given by Lemma 30. The set $\mathcal{T}'_n \subseteq \mathcal{G}_n(\boldsymbol{a})$ is constructed from \mathcal{T}_n by considering the potential of the trees of these gluing pairs. Note that the path on n vertices has the highest potential $\hat{p} := \lfloor n^2/4 \rfloor$, whereas the star $[s_n]$ has the lowest potential $\check{p} := n-1$. Roughly speaking, we consider all plane trees with decreasing potential $p = \hat{p}, \hat{p} - 1, \ldots, \check{p}$, and we add gluing tuples from $\mathcal{G}_n(\boldsymbol{a})$ derived from gluing pairs of \mathcal{T}_n step by step, such that all cycles of $\mathcal{C}_n(\boldsymbol{a})$ corresponding to marked plane trees with potential at least p are joined together.

Formally, for $p = \hat{p}, \hat{p} - 1, \ldots, \check{p}$, we inductively construct a set $\mathcal{T}'_{n,p} \subseteq \mathcal{G}_n(\boldsymbol{a})$ of gluing tuples such that $\mathcal{C}_n(\boldsymbol{a}) \triangle C(\mathcal{T}'_{n,p})$ is a cycle factor in $G_n(\boldsymbol{a})$ with the property that every vertex is contained in a cycle together with all vertices $\{\tilde{x}\ 0 \mid \tilde{x} \in [x]\}$ for some $x \in D_n(\boldsymbol{a})$ with $\varphi(x) \leq p$. Note that there are precisely two marked plane trees with minimal potential $p = \check{p} = n - 1$, namely the star $[s_n]$ with either the center vertex or a leaf marked, i.e., these are $[s'_n]$ and $[s''_n]$ where $s'_n := (10)^{n-1} \in D_n(\boldsymbol{a})$ and $s''_n := (20)(10)^{n-2} \in D_n(\boldsymbol{a})$. We will show that $s'_n \ 0, s''_n \ 0 \in \Pi_n(\boldsymbol{a})^{-1}$ lie on the same cycle, and these conditions imply that $\mathcal{C}_n(\boldsymbol{a}) \triangle C(\mathcal{T}'_{n,\check{p}})$ is a Hamilton cycle in $G_n(\boldsymbol{a})$.




FIGURE 26. Illustration of Lemma 30 and the proof of Theorem 9. The figure shows all plane trees with 7 vertices, arranged in levels according to their potential. The vertices with minimum potential are highlighted. The number next to each tree specifies the number of distinct plane trees obtained by marking a single vertex (in each orbit of symmetric vertices, one is drawn black and the others white). Each gluing pair $(x,y) \in \mathcal{T}_n \subseteq \mathcal{G}_n$ is visualized by indicating the placement of roots at the plane trees [x] and [y] by a small filled or non-filled arrow, respectively. The bubbles indicate gluing tuples from $\mathcal{T}'_n \subseteq \mathcal{G}_n(n-1,1)$ that join the 5 marked plane trees with the mark placed at one vertex within the bubbles in each tree. An example of two gluing tuples that are used for the four marked paths with potential 12 is depicted in the box.

The induction basis is $\mathcal{T}'_{n,\hat{p}} := \emptyset$. For the induction step $p \to p-1$, suppose that $\mathcal{T}'_{n,p}$, $\hat{p} \ge p > \check{p}$, with the properties as stated before is given. Then $\mathcal{T}'_{n,p-1}$ is obtained by adding gluing tuples from $\mathcal{G}_n(\boldsymbol{a})$ to the set $\mathcal{T}'_{n,p}$ as follows.

We consider each plane tree T with n vertices and potential p, and we consider the unique gluing pair $(x,y) \in \mathcal{T}_n$ with T = [x] and $\varphi(y) = \varphi(x) - 1$. We then consider all rooted trees obtained from x by marking one vertex, i.e., the set $X' := \{x' \in D_n(\mathbf{a}) \mid \hat{x'} = x\}$. Two marked rooted trees $x', x'' \in X'$ are equivalent, if x' 0 and x'' 0 lie on the same cycle in the factor $\mathcal{C}_n(\mathbf{a}) \triangle C(\mathcal{T}'_{n,p})$, and we partition the set X' into equivalence classes $X' = X'_1 \cup \cdots \cup X'_r$ accordingly. Note that x', x'' can be equivalent either because these trees have the same underlying marked plane tree, i.e., [x'] = [x''], or because the cycles C(x', 0) and C(x'', 0) have been joined previously.

By (9) we have x = 110u0v and y = 101u0v for some $u, v \in D$. We consider the three trees $x_3, x_2, x_1 \in D_n(\mathbf{a})$ obtained from x by marking the root of x, or the leftmost descendants in distance 1 or 2 from the root, respectively, and the two trees $y_2, y_1 \in D_n(\mathbf{a})$ obtained from y by marking the root of y, or its second child from left to right, respectively (these are the five trees marked by (*) in Figure 25). By these definitions we have

$$x_1 = 120u0v,$$
 $y_1 = 102u0v,$
 $x_2 = 210u0v,$ $y_2 = 101u0v,$
 $x_3 = 110u0v.$

We then add the gluing tuple $(x_1, x_2, x_3, x_3, y_1, y_2)$ to the set $\mathcal{T}'_{n,p-1}$. The corresponding 12-cycle $C(x_1, x_2, x_3, x_3, y_1, y_2)$ intersects five cycles from the factor $\mathcal{C}_n(\boldsymbol{a})$, corresponding to the marked plane trees $[x_1]$, $[x_2]$, $[x_3]$, $[y_1]$, and $[y_2]$; see Figure 26. There is one special case, namely if v = 10u, in which case $[x_2] = [x_3]$, and then the 12-cycle intersects only four cycles from the factor $\mathcal{C}_n(\boldsymbol{a})$, but one can check that the symmetric difference is still a single cycle; see Figure 27 (a).

In addition to the aforementioned gluing tuple, which is added in every case, we add further gluing tuples to the set $\mathcal{T}'_{n,p-1}$, depending on X'. Specifically, for every equivalence class X'_i , $i \in [r]$, that contains none of x_1, x_2, x_3 , we select one tree $x' \in X'_i$, and we let $y' \in D_n(\mathbf{a})$ be the string obtained from x' by swapping the second and third entry, i.e., x' = 110u0v, y' = 101u0v with $\widehat{u}, \widehat{v} \in D$ such that the concatenation uv contains a single occurrence of 2. We then add the




FIGURE 27. Verification of multiple intersections in the proof of Theorem 9.

gluing tuple (x', x', x', x', y', y') to the set $\mathcal{T}'_{n,p}$. The corresponding 6-cycle C(x', x', x', x', y', y') intersects two cycles from the factor $\mathcal{C}_n(\boldsymbol{a})$, corresponding to the marked plane trees [x'] and [y'].

By the construction of $\mathcal{T}'_{n,p-1}$, each equivalence class X'_i , $i \in [r]$, contains at most two of the x_i , and if this happens then $X'_i \cap \{x_1, x_2, x_3\} = \{x_2, x_3\}$, and $[x_2] = [\tilde{y}_1]$ and $[x_3] = [\tilde{y}_2]$ for some gluing pair $(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{x}_3, \tilde{y}_1, \tilde{y}_2) \in \mathcal{T}'_{n,p-1}$; see Figure 27 (b). One can check in the figure that the double intersection between the two gluing cycles and the two cycles from the factor still yields a single cycle in the symmetric difference.

By the above steps and the induction hypothesis for p, we have achieved that in $C_n(\mathbf{a}) \triangle C(T'_{n,p-1})$, every vertex is contained in a cycle together with all vertices $\{\tilde{x} \mid \tilde{x} \in [x]\}$ for some $x \in D_n(\mathbf{a})$ with $\varphi(x) \leq p-1$, i.e., the induction hypothesis holds for p-1. In the last step of the construction for $p = \check{p} + 1$ we added a gluing tuple $(x_1, x_2, x_3, x_3, y_1, y_2)$ where $y_1 = s'_n$, $y_2 = s'_n$, and thus we glued both cycles $[s''_n]$ and $[s'_n]$ into one.

This completes the proof of the theorem.

6. Open questions

We conclude this paper with the following open questions.

- We believe that the techniques used in Section 5.3 to prove that $G(\alpha, \alpha 1, 1)$ has a Hamilton cycle are in principle suitable to prove that G(a) has a Hamilton cycle for all a with $\Delta(a) = 0$, which would be an important first step towards a proof of Conjecture 6. In particular, the construction of the cycle factor $C_n(a)$ and gluing tuples $G_n(a)$ described in Sections 5.3.1 and 5.3.2, respectively, are fully general. The main difficulty in combining these ingredients lies in the fact that some gluing cycles join more than two cycles from the factor (namely 3, 5, or 6 cycles) to a single cycle, and in this case the resulting interactions between different gluing cycles seem to be hard to control (recall Figure 27).
- We conjecture that $G(\alpha, \alpha, 1)$ for $\alpha \geq 2$ is Hamilton-connected, just as all other graphs $G(\boldsymbol{a})$ with $\Delta(\boldsymbol{a}) > 0$ covered by Theorem 7. However, we are unable to prove this based on Conjecture 6; recall the discussion in Section 2.3. We confirmed this conjecture with computer help for $\alpha = 2, 3, 4$; see Table 1. Proving the conjecture would streamline our proof of Theorem 7 considerably, as it would make Lemmas 17–20 redundant, which build on top of Lemma 21 in the induction proof. To prove that $G(\alpha, \alpha, 1)$ is Hamilton-connected, it may help to establish a Hamiltonicity property for graphs $G(\boldsymbol{a})$ with $\Delta(\boldsymbol{a}) = 0$ and $k \geq 3$ that is stronger than \mathcal{L}_1 . Specifically, in addition to a Hamilton path between any vertex in $\Pi(\boldsymbol{a})^{1,1}$ and any vertex not in $\Pi(\boldsymbol{a})^{1,1}$, we may also ask for a Hamilton path between any two distinct vertices in $\Pi(\boldsymbol{a})^{1,1}$. We checked by computer whether $G(\boldsymbol{a})$ has this stronger property for $\boldsymbol{a} \in \{(2,1,1),(3,2,1),(3,1,1,1),(4,3,1),(4,2,2),(4,2,1,1),(4,1,1,1,1)\}$, and it was satisfied in all cases except for $\boldsymbol{a} = (2,1,1)$.
- While the proofs presented in this paper are constructive, they are far from yielding efficient algorithms for computing the corresponding Gray codes. Ideally, one would like algorithms whose running time is polynomial in n per generated multiset permutation of length n. Such algorithms are known for the Hamilton cycles mentioned in Theorems 2 and 3, see [Knu11, Section 7.2.1.2] and [MN20], respectively. An interesting direction could be to explore greedy algorithms for generating multiset permutations by star transpositions, which may yield much simpler constructions to start with, cf. [Wil13, SW16, CSW21].
- Knuth raised the question whether there are star transposition Gray codes for (α, α) combinations whose flip sequence can be partitioned into $2\alpha 1$ blocks, such that each
 block is obtained from the previous one by adding +1 modulo $2\alpha 1$. This problem is a
 strengthening of the middle levels conjecture, and it was answered affirmatively in [MMM21].




FIGURE 28. Star transposition Gray code for (2, 2, 2)-multiset permutations with 5-fold cyclic symmetry.

The Gray code for (4,4)-combinations shown in Figure 2 (a) has such a 7-fold cyclic symmetry. We can ask more generally: Are there star transposition Gray codes for multiset permutations whose flip sequence can be partitioned into n-1 blocks, such that each block is obtained from the previous one by adding +1 modulo n-1? Figure 28 shows an ad-hoc solution for (2,2,2)-multiset permutations with 5-fold cyclic symmetry.

• A more general version of the problem considered in this paper is the following: We consider an alphabet $\{1,\ldots,k\}$ of size $k\geq 2$, and frequencies $a_1,\ldots,a_k\geq 1$ that specify that symbol i appears exactly a_i times for all $i=1,\ldots,k$. Moreover, there is an additional integer parameter s with $1\leq s\leq k-1$ that has the following significance. The objects to be generated are all pairs (S,x), where S is a set or string of s distinct symbols, and s is a string of the remaining s0 symbols, where s1 is a set or string of s2. A star transposition swaps one symbol from s2 with one symbol from s3 that is currently not in s4, and the question is whether there is a star transposition Gray code for all those objects.

Note that multiset permutations considered in this paper are the special case when s=1. Andrea Sportiello suggested this problem with $a_1=\cdots=a_k=\alpha$ (uniform frequency) and S being a set as a generalization of the middle levels conjecture $(s=1, k=2, a_1=a_2=\alpha)$. Moreover, Ajit A. Diwan suggested this problem with $a_1=\cdots=a_k=\alpha$ (uniform frequency) and a set S of size s=k-1. Note that the uniform frequency case is particularly interesting, as the underlying flip graph for this problem is vertex-transitive if and only if $a_1=\cdots=a_k$ (recall Lovász' conjecture [Lov70]).

ACKNOWLEDGEMENTS

We thank Jiří Fink for inspiring brain-storming sessions in the early phases of this research project, and we thank Aaron Williams for interesting discussions about star transposition Gray codes.

References

- [AQ01] B. Alspach and Y. Qin. Hamilton-connected Cayley graphs on Hamiltonian groups. European J. Combin., 22(6):777-787, 2001.
- [Ara06] T. Araki. Hyper Hamiltonian laceability of Cayley graphs generated by transpositions. *Networks*, 48(3):121–124, 2006.
- [BW84] M. Buck and D. Wiedemann. Gray codes with restricted density. Discrete Math., 48(2-3):163-171, 1984.
- [Cha89] P. Chase. Combination generation and graylex ordering. Congr. Numer., 69:215–242, 1989. Eighteenth Manitoba Conference on Numerical Mathematics and Computing (Winnipeg, MB, 1988).
- [Cor92] P. F. Corbett. Rotator graphs: An efficient topology for point-to-point multiprocessor networks. IEEE Transactions on Parallel and Distributed Systems, 3:622–626, 1992.
- [CQ81] C. C. Chen and N. F. Quimpo. On strongly Hamiltonian abelian group graphs. In Combinatorial mathematics, VIII (Geelong, 1980), volume 884 of Lecture Notes in Math., pages 23–34. Springer, Berlin-New York, 1981.
- [CSW21] B. Cameron, J. Sawada, and A. Williams. A Hamilton cycle in the k-sided pancake network. arXiv:2103.09256, 2021.
- [CW93] R. C. Compton and S. G. Williamson. Doubly adjacent Gray codes for the symmetric group. *Linear and Multilinear Algebra*, 35(3-4):237–293, 1993.
- [DG12] P. Diaconis and R. Graham. *Magical mathematics*. Princeton University Press, Princeton, NJ, 2012. The mathematical ideas that animate great magic tricks, With a foreword by Martin Gardner.
- [EHR84] P. Eades, M. Hickey, and R. C. Read. Some Hamilton paths and a minimal change algorithm. *J. Assoc. Comput. Mach.*, 31(1):19–29, 1984.
- [EM84] P. Eades and B. McKay. An algorithm for generating subsets of fixed size with a strong minimal change property. *Inform. Process. Lett.*, 19(3):131–133, 1984.
- [GMN18] P. Gregor, T. Mütze, and J. Nummenpalo. A short proof of the middle levels theorem. *Discrete Analysis*, 2018:8:12 pp., 2018.
- [Hav83] I. Havel. Semipaths in directed cubes. In *Graphs and other combinatorial topics (Prague, 1982)*, volume 59 of *Teubner-Texte Math.*, pages 101–108. Teubner, Leipzig, 1983.
- [HCH00] S.-Y. Hsieh, G.-H. Chen, and C.-W. Ho. Hamiltonian-laceability of star graphs. *Networks*, 36(4):225–232, 2000.
- [HH72] C. A. Holzmann and F. Harary. On the tree graph of a matroid. SIAM J. Appl. Math., 22:187–193, 1972.
- [HL87] F. Harary and M. Lewinter. Hypercubes and other recursively defined Hamilton laceable graphs. volume 60, pages 81–84. 1987. Eighteenth Southeastern International Conference on Combinatorics, Graph Theory, and Computing (Boca Raton, Fla., 1987).
- [JM95] T. Jenkyns and D. McCarthy. Generating all k-subsets of $\{1 \cdots n\}$ with minimal changes. Ars Combin., 40:153-159, 1995.
- [Joh63] S. Johnson. Generation of permutations by adjacent transposition. Math. Comp., 17:282–285, 1963.
- [KL75] V. L. Kompel'maher and V. A. Liskovec. Sequential generation of permutations by means of a transposition basis. *Kibernetika (Kiev)*, (3):17–21, 1975.
- [Knu11] D. E. Knuth. The Art of Computer Programming. Vol. 4A. Combinatorial Algorithms. Part 1. Addison-Wesley, Upper Saddle River, NJ, 2011.
- [KT88] H. A. Kierstead and W. T. Trotter. Explicit matchings in the middle levels of the Boolean lattice. Order, 5(2):163–171, 1988.
- [Lov70] L. Lovász. Problem 11. In Combinatorial Structures and Their Applications (Proc. Calgary Internat. Conf., Calgary, Alberta, 1969). Gordon and Breach, New York, 1970.
- [MMM21] A. I. Merino, O. Mička, and T. Mütze. On a combinatorial generation problem of Knuth. In D. Marx, editor, Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021, pages 735-743. SIAM, 2021.
- [MN20] T. Mütze and J. Nummenpalo. A constant-time algorithm for middle levels Gray codes. *Algorithmica*, 82(5):1239–1258, 2020.
- [Müt16] T. Mütze. Proof of the middle levels conjecture. Proc. Lond. Math. Soc., 112(4):677–713, 2016.
- [NW75] A. Nijenhuis and H. Wilf. *Combinatorial algorithms*. Academic Press, New York-London, 1975. Computer Science and Applied Mathematics.
- [RS93] F. Ruskey and C. D. Savage. Hamilton cycles that extend transposition matchings in Cayley graphs of S_n . SIAM J. Discrete Math., 6(1):152-166, 1993.
- [Rus88] F. Ruskey. Adjacent interchange generation of combinations. J. Algorithms, 9(2):162–180, 1988.

- [Rus16] F. Ruskey. Combinatorial Gray code. In M.-Y. Kao, editor, Encyclopedia of Algorithms, pages 342–347. Springer, 2016.
- [RW09] F. Ruskey and A. Williams. The coolest way to generate combinations. Discrete Math., 309(17):5305-5320, 2009.
- C. D. Savage. A survey of combinatorial Gray codes. SIAM Rev., 39(4):605-629, 1997. [Sav97]
- [Sim78] G. J. Simmons. Almost all n-dimensional rectangular lattices are Hamilton-laceable. In Proceedings of the Ninth Southeastern Conference on Combinatorics, Graph Theory, and Computing (Florida Atlantic Univ., Boca Raton, Fla., 1978), Congress. Numer., XXI, pages 649-661. Utilitas Math., Winnipeg, Man., 1978.
- [Sla78] P. J. Slater. Generating all permutations by graphical transpositions. Ars Combin., 5:219–225, 1978.
- G. Stachowiak. Hamilton paths in graphs of linear extensions for unions of posets. SIAM J. Discrete [Sta92] Math., 5(2):199-206, 1992.
- [SW13] X. S. Shen and A. Williams. A 'hot potato' Gray code for permutations. Electronic Notes in Discrete Mathematics, 44:89-94, 2013.
- [SW16] J. Sawada and A. Williams. Greedy flipping of pancakes and burnt pancakes. Discrete Appl. Math., 210:61-74, 2016.
- [SW20] J. Sawada and A. Williams. Solving the sigma-tau problem. ACM Trans. Algorithms, 16(1):Art. 11, 17 pp., 2020.
- [SW21] X. S. Shen and A. Williams. A k-ary middle levels conjecture. In Proceedings of the 23rd Thailand-Japan Conference on Discrete and Computational Geometry, Graphs, and Games, 2021.
- [Tch82] M. Tchuente. Generation of permutations by graphical exchanges. Ars Combin., 14:115–122, 1982.
- [TL73] D. Tang and C. Liu. Distance-2 cyclic chaining of constant-weight codes. IEEE Trans. Computers, C-22:176–180, 1973.
- [Tro62] H. F. Trotter. Algorithm 115: Perm. Commun. ACM, 5(8):434-435, 1962.
- [Wil13] A. Williams. The greedy Gray code algorithm. In Algorithms and Data Structures - 13th International Symposium, WADS 2013, London, ON, Canada, August 12-14, 2013. Proceedings, pages 525-536, 2013.
- [Win04] P. Winkler. Mathematical puzzles: a connoisseur's collection. A K Peters, Ltd., Natick, MA, 2004.
- S. Zaks. A new algorithm for generation of permutations. BIT, 24(2):196–204, 1984. [Zak84]

Appendix A. Proofs of Lemmas 17–20

Proof of Lemma 17. Let x, y be two distinct vertices in G(a). As $x \neq y$ there is a position $\hat{i} > 1$ such that $x_i \neq y_i$. As the symbols 1 and 2 both appear with the same frequency α , we may assume w.l.o.g. that either $(x_{\hat{i}}, y_{\hat{i}}) = (1, 2)$ or $(x_{\hat{i}}, y_{\hat{i}}) = (1, 3)$.

We first consider the case $(x_{\hat{i}}, y_{\hat{i}}) = (1, 2)$. As $\alpha \geq 3$, there are two indices $i_1, i_2 \in [n] \setminus \{1, \hat{i}\}$ such that $x_{i_1} \neq 2$ and $x_{i_2} = 2$, in particular $x_{i_1} \neq x_{i_2}$. Similarly, there are two indices $i_3, i_4 \in [n] \setminus \{1, \hat{i}\}$ such that $y_{i_3} \neq 1$ and $y_{i_4} = 1$, in particular $y_{i_3} \neq y_{i_4}$. We choose multiset permutations $u^j, v^j \in \Pi(\boldsymbol{a}), j=1,2,3$, satisfying the following conditions; see Figure 29 (a):

- (i) $u^1 = x$, $v^1_{i_1} = x_{i_2} = 2$, and $v^3 = y$, $u^3_{i_3} = y_{i_4} = 1$; (ii) $u^2_1 = v^1_{\hat{\imath}} = 1$, $u^2_{\hat{\imath}} = v^1_1 = 3$, $u^3_1 = v^2_{\hat{\imath}} = 3$, $u^3_{\hat{\imath}} = v^2_1 = 2$, and $u^j_i = v^{j-1}_i$ for j = 2, 3 and all $i \in [n] \setminus \{1, \hat{\imath}\};$
- (iii) $(u_{i_1}^2, v_{i_1}^2) = (2, 1).$

Note that $u^1 \neq v^1$ and $u^3 \neq v^3$ holds by (i) and the choice of i_1, i_2 and i_3, i_4 , respectively, and $u_1^2 \neq v_1^2$ by (ii). Consequently, we have $u^j \neq v^j$ for j = 1, 2, 3.

We consider a Hamilton path P_1 in $G^{\hat{i},1}(\boldsymbol{a})$ from u^1 to v^1 , and a Hamilton path P_3 in $G^{\hat{i},2}(\boldsymbol{a})$ from u^3 to v^3 , which exist by the assumption $G(\alpha, \alpha - 1, 2) \in \mathcal{H}$. We also consider a Hamilton path P_2 in $G^{i,3}(\boldsymbol{a})$ from u^2 to v^2 , which exists by the assumption $G(\alpha,\alpha,1)\in\mathcal{L}_{12}$, using that $(u_1^2, v_1^2) = (1, 2)$ and $(u_{i_1}^2, v_{i_1}^2) = (2, 1) = (p_3(1, 2), q_3(1, 2))$ by (ii)+(iii) and (4). The concatenation $P_1P_2P_3$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

It remains to consider the case $(x_i, y_i) = (1, 3)$. We define $t := y_1 \in \{1, 2, 3\}$ and $p := p_3(2, t) \in$ $\{1,3\}$. As $\alpha \geq 3$, there are two indices $i_1, i_2 \in [n] \setminus \{1,\hat{i}\}$ such that $x_{i_1} \neq x_{i_2}$. Similarly, there




FIGURE 29. Illustration of the proof of Lemma 17.

is an index $i_3 \in [n] \setminus \{1, \hat{i}\}$ such that $y_{i_3} = 2$. We choose multiset permutations $u^j, v^j \in \Pi(\boldsymbol{a})$, j = 1, 2, 3, satisfying the following conditions; see Figure 29 (b):

- (i) $u^1=x, v^1_{i_1}=x_{i_2}$, and $v^3=y$; (ii) $u^2_1=v^1_{\hat{i}}=1, \ u^2_{\hat{i}}=v^1_1=2, \ u^3_1=v^2_{\hat{i}}=2, \ u^3_{\hat{i}}=v^2_1=3, \ \text{and} \ u^j_i=v^{j-1}_i \ \text{for} \ j=2,3 \ \text{and} \ \text{all}$
- (iii) $u_{i_3}^3 = p$.

Note that $u^1 \neq v^1$ by (i) and the choice of $i_1, i_2, u_1^2 \neq v_1^2$ by (ii), and $u^3 \neq v^3$ by (iii), the choice of i_3 and the fact that $p \neq 2$. Consequently, we have $u^j \neq v^j$ for j = 1, 2, 3.

We consider a Hamilton path P_1 in $G^{\hat{i},1}(a)$ from u^1 to v^1 , and a Hamilton path P_2 in $G^{\hat{i},2}(a)$ from u^2 to v^2 , which exist by the assumption $G(\alpha, \alpha - 1, 2) \in \mathcal{H}$. We also consider a Hamilton path P_3 in $G^{\hat{i},3}(\boldsymbol{a})$ from u^3 to v^3 , which exists by the assumption $G(\alpha,\alpha,1)\in\mathcal{L}_{12}$, using that $(u_1^3,v_1^3)=(2,t)$ and $(u_{i_3}^3,v_{i_3}^3)=(p,2)=(p_3(2,t),q_3(2,t))$ by (i)–(iii) and (4). The concatenation $P_1P_2P_3$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Proof of Lemma 18. Let x, y be two distinct vertices in G(a). As $x \neq y$ there is a position $\hat{i} > 1$ such that $x_i \neq y_i$. As the symbols 1 and 2 both appear with the same frequency α , and the symbols 3 and 4 both appear with the same frequency 1, we may assume w.l.o.g. that either $(x_{\hat{i}}, y_{\hat{i}}) = (1, 2), (x_{\hat{i}}, y_{\hat{i}}) = (1, 3), \text{ or } (x_{\hat{i}}, y_{\hat{i}}) = (3, 4).$

We first consider the case $(x_i, y_i) = (1, 2)$. Let i_2 be the unique index such that $x_{i_2} = 4$, and let i_4 be the unique index such that $y_{i_4} = 3$. As $\alpha \geq 3$ and hence $n \geq 2\alpha + 2 \geq 8$, there is an index $i_1 \in [n] \setminus \{1, \hat{i}, i_2\}$ and an index $i_3 \in [n] \setminus \{1, \hat{i}, i_4, i_1\}$, and they will satisfy $x_{i_1} \neq 4$ and $y_{i_3} \neq 3$. We choose multiset permutations $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, 2, 3, 4$, satisfying the following conditions; see Figure 30 (a):

- $\begin{array}{l} \text{(i)} \ \ u^1=x, \ v^1_{i_1}=x_{i_2}=4, \ \text{and} \ v^4=y, \ u^4_{i_3}=y_{i_4}=3; \\ \text{(ii)} \ \ u^2_1=v^1_{\hat{\imath}}=1, \ u^2_{\hat{\imath}}=v^1_1=3, \ u^3_1=v^2_{\hat{\imath}}=3, \ u^3_{\hat{\imath}}=v^2_1=4, \ u^4_1=v^3_{\hat{\imath}}=4, \ u^4_{\hat{\imath}}=v^3_1=2, \ \text{and} \\ \ \ u^j_i=v^{j-1}_{\hat{\imath}} \ \text{for} \ j=2,3,4 \ \text{and} \ \text{all} \ i\in[n]\setminus\{1,\hat{\imath}\}; \\ \text{(iii)} \ \ (u^2_{i_1},v^2_{i_1})=(4,1), \ (u^3_{i_3},v^3_{i_3})=(2,3). \end{array}$

Note that $u^1 \neq v^1$ and $u^4 \neq v^4$ by (i) and the choice of i_1, i_2 and i_3, i_4 , respectively, and $u_1^j \neq v_1^j$ for j=2,3 by (ii). Consequently, we have $u^j\neq v^j$ for j=1,2,3,4.

We consider a Hamilton path P_1 in $G^{\hat{i},1}(\boldsymbol{a})$ from u^1 to v^1 , and a Hamilton path P_4 in $G^{\hat{i},2}(\boldsymbol{a})$ from u^4 to v^4 , which exist by the assumption $G(\alpha, \alpha - 1, 1, 1) \in \mathcal{H}$. We also consider a Hamilton path P_2 in $G^{\hat{i},3}(\boldsymbol{a})$ from u^2 to v^2 , which exists by the assumption $G(\alpha,\alpha,1)\in\mathcal{L}_{12}$, using that $(u_1^2, v_1^2) = (1, 4)$ and $(u_{i_1}^2, v_{i_1}^2) = (4, 1) = (p_4(1, 4), q_4(1, 4))$ by (ii)+(iii) and (4). Note here that in $G^{\hat{i},3}(\boldsymbol{a})$, the symbol 4 takes the role of 3 in $G(\alpha,\alpha,1)$. Finally, we consider a




FIGURE 30. Illustration of the proof of Lemma 18.

Hamilton path P_3 in $G^{\hat{i},4}(\boldsymbol{a})$ from u^3 to v^3 , which exists by the assumption $G(\alpha,\alpha,1) \in \mathcal{L}_{12}$, using that $(u_1^3, v_1^3) = (3, 2)$ and $(u_{i_3}^3, v_{i_3}^3) = (2, 3) = (q_3(2, 3), p_3(2, 3))$ by (ii)+(iii) and (4). The concatenation $P_1P_2P_3P_4$ is a Hamilton path in G(a) from x to y.

We now consider the case $(x_{\hat{i}}, y_{\hat{i}}) = (1, 3)$. We define $t := y_1 \in \{1, 2, 4\}$ and $p := p_4(2, t) \in$ $\{1,4\}$. Let $i_1,i_2\in[n]\setminus\{1,\hat{\imath}\}$ be such that $x_{i_1}\neq 2$ and $x_{i_2}=2$. Similarly, let $i_3\in[n]\setminus\{1,\hat{\imath}\}$ be such that $y_{i_3}=2$. We choose multiset permutations $u^j,v^j\in\Pi(a),\ j=1,2,3,4$, satisfying the following conditions; see Figure 30 (b):

(i)
$$u^1=x, \ v^1_{i_1}=x_{i_2}=2, \ \text{and} \ v^4=y;$$

(ii) $u^2_1=v^1_{\hat{i}}=1, \ u^2_{\hat{i}}=v^1_1=4, \ u^3_1=v^2_{\hat{i}}=4, \ u^3_{\hat{i}}=v^2_1=2, \ u^4_1=v^3_{\hat{i}}=2, \ u^4_{\hat{i}}=v^3_1=3 \ \text{and} \ u^j_i=v^{j-1}_i \ \text{for} \ j=2,3,4 \ \text{and} \ \text{all} \ i\in[n]\setminus\{1,\hat{i}\};$
(iii) $(u^2_{i_1},v^2_{i_1})=(2,1) \ \text{and} \ u^4_{i_3}=p.$

Note that $u^1 \neq v^1$ by (i) and the choice of $i_1, i_2, u_1^j \neq v_1^j$ for j = 2, 3 by (ii), and $u^4 \neq v^4$ by (iii), the choice of i_3 and the fact that $p \neq 2$. Consequently, we have $u^j \neq v^j$ for j = 1, 2, 3, 4.

We consider a Hamilton path P_1 in $G^{\hat{\imath},1}(\boldsymbol{a})$ from u^1 to v^1 , and a Hamilton path P_3 in $G^{\hat{\imath},2}(\boldsymbol{a})$ from u^3 to v^3 , which exist by the assumption $G(\alpha, \alpha - 1, 1, 1) \in \mathcal{H}$. We also consider a Hamilton path P_2 in $G^{\hat{i},4}(\boldsymbol{a})$ from u^2 to v^2 , which exists by the assumption $G(\alpha,\alpha,1)\in\mathcal{L}_{12}$, using that $(u_1^2, v_1^2) = (1, 2)$ and $(u_{i_1}^2, v_{i_1}^2) = (2, 1) = (p_3(1, 2), q_3(1, 2))$ by (ii)+(iii) and (4). Finally, we consider a Hamilton path P_3 in $G^{\hat{\imath},3}(\boldsymbol{a})$ from u^4 to v^4 , which exists by the assumption $G(\alpha,\alpha,1)\in$ \mathcal{L}_{12} , using that $(u_1^4, v_1^4) = (2, t)$ and $(u_{i_3}^4, v_{i_3}^4) = (p, 2) = (p_4(2, t), q_4(2, t))$ by (i)–(iii) and (4). The concatenation $P_1P_2P_3P_4$ is a Hamilton path in G(a) from x to y.

It remains to consider the case $(x_i, y_i) = (3, 4)$. We define $t := y_1 \in \{1, 2, 3\}$ and $p := p_3(2, t) \in$ $\{1,3\}$. Moreover, we define $t':=x_1\in\{1,2,4\}$ and $p':=p_4(1,t')\in\{2,4\}$. Let $i_1\in[n]\setminus\{1,\hat{i}\}$ be such that $x_{i_1} = 1$, and let $i_3 \in [n] \setminus \{1, \hat{i}\}$ be such that $y_{i_3} = 2$. We choose multiset permutations $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, 2, 3, 4$, satisfying the following conditions; see Figure 30 (c):

(i)
$$u^1 = x \text{ and } v^4 = y;$$

(ii) $u_1^2 = v_i^1 = 3$, $u_i^2 = v_1^1 = 1$, $u_1^3 = v_i^2 = 1$, $u_i^3 = v_1^2 = 2$, $u_1^4 = v_i^3 = 2$, $u_i^4 = v_1^3 = 4$ and $u_i^j = v_i^{j-1}$ for j = 2, 3, 4 and all $i \in [n] \setminus \{1, \hat{\imath}\}$; (iii) $v_{i_1}^1 = p'$ and $u_{i_3}^4 = p$.

(iii)
$$v_{i_1}^1 = p'$$
 and $u_{i_3}^4 = p$.

Note that $u_1^j \neq v_1^j$ for j=2,3 by (ii), and $u^1 \neq v^1$ and $u^4 \neq v^4$ by (iii), the choice of i_1,i_3 and the fact that $p' \neq 1$ and $p \neq 2$, respectively. Consequently, we have $u^j \neq v^j$ for j = 1, 2, 3, 4.

We consider a Hamilton path P_2 in $G^{\hat{\imath},1}(\boldsymbol{a})$ from u^2 to v^2 , and a Hamilton path P_3 in $G^{\hat{\imath},2}(\boldsymbol{a})$ from u^3 to v^3 , which exist by the assumption $G(\alpha, \alpha - 1, 1, 1) \in \mathcal{H}$. We also consider a Hamilton path P_1 in $G^{\hat{\imath},3}(\boldsymbol{a})$ from u^1 to v^1 , which exists by the assumption $G(\alpha, \alpha, 1) \in \mathcal{L}_{12}$, using that $(u_1^1, v_1^1) = (t', 1)$ and $(u_{i_1}^1, v_{i_1}^1) = (1, p') = (q_4(1, t'), p_4(1, t'))$ by (i)–(iii) and (4). Finally, we consider a Hamilton path P_4 in $G^{\hat{\imath},4}(\boldsymbol{a})$ from u^4 to v^4 , which exists by the assumption $G(\alpha, \alpha, 1) \in \mathcal{L}_{12}$, using that $(u_1^4, v_1^4) = (2, t)$ and $(u_{i_3}^4, v_{i_3}^4) = (p, 2) = (p_3(2, t), q_3(2, t))$ by (i)–(iii) and (4). The concatenation $P_1P_2P_3P_4$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Proof of Lemma 19. We proceed exactly as in the proof of Lemma 16 for k = 3, choosing indices i, \hat{i} and distinguishing cases (oa), (ob1) and (ob2) depending on the values of y_i and y_1 , as before.

Case (oa): $y_i = 1$. By the assumption that $x_i, y_i > 1$ we can now assume w.l.o.g. that $x_i = 2$ and $y_i = 3$. We define the permutations $\pi := (1,2,3)$ and $\rho := (2,1,3)$. We set $t := y_1 \in \{1,2,3\}$ and $p := p_3(1,t) \in \{2,3\}$, and we choose $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1,2,3$, and $\hat{u}^1, \hat{v}^1, \check{u}^1, \check{v}^1 \in \Pi(\boldsymbol{a})$, as in case (oa) in the proof of Lemma 16 for k = 3, subject to the conditions (ii)–(vi) as before and the following condition (i') instead of (i); see Figure 31 (oa):

(i')
$$u^1=x,\,v^1_{i_1}=x_{i_2},$$
 and $v^3=y,\,u^3_{i_3}=p,$ where i_3 is chosen such that $y_{i_3}=1.$

For j=1,2,3 we consider a Hamilton path P_j in $G^{(\check{\imath},\hat{\imath}),(1,\rho_j)}(\boldsymbol{a})$ from u^j to v^j . For j=1,2 such a path exists by the assumption $G(\varphi(\alpha-1,\alpha-2,2))\in\mathcal{H}$. For j=3 such a path exists by the assumption $G(\alpha-1,\alpha-1,1)\in\mathcal{L}_{12}$, using that $(u_1^3,v_1^3)=(1,t)$ and $(u_{i_3}^3,v_{i_3}^3)=(p,1)=(p_3(1,t),q_3(1,t))$ by (i') and (4).

We also consider a Hamilton path \hat{P}_1 in $G^{\check{\imath},\pi_2}(\boldsymbol{a}) = G^{\check{\imath},2}(\boldsymbol{a})$ from \hat{u}^1 to \hat{v}^1 , which exists by the assumption $G(\varphi(\alpha,\alpha-2,2)) \in \mathcal{L}_1$, and a Hamilton path \check{P}_1 in $G^{\check{\imath},\pi_3}(\boldsymbol{a}) = G^{\check{\imath},3}(\boldsymbol{a})$ from \check{u}^1 to \check{v}^1 , which exists by the assumption $G(\alpha,\alpha-1,1) \in \mathcal{L}_1$.

The concatenation $P_1\hat{P}_1\check{P}_1P_2P_3$ is a Hamilton path in G(a) from x to y.

Case (ob1): $y_i \neq 1$ and $y_1 = 1$. We let $\pi = (\pi_1, \pi_2, \pi_3)$ be the permutation defined by $\pi_1 = x_i = 1$ and $\pi_3 = y_i$, and we define $\rho := (1, 2, 3)$.




Figure 31. Illustration of the proof of Lemma 19.

We choose $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, 2, 3$, and $\hat{u}^1, \hat{v}^1, \check{u}^1, \check{v}^1 \in \Pi(\boldsymbol{a})$, as in case (ob1) in the proof of Lemma 16 for k = 3, subject to the conditions (i)–(iii) as before and the following condition (iv') instead of (iv); see Figure 31 (ob1):

(iv') $u_{\tilde{i}}^3 = 3$ and $v_{\tilde{i}}^3 = 2$ for some $\tilde{i} \in [n] \setminus \{1, \check{i}, \hat{i}\}.$

We consider Hamilton paths P_1 , P_2 , P_3 and \hat{P}_1 , \check{P}_1 as before. The path P_3 from u^3 to v^3 exists by the assumption $G(\alpha - 1, \alpha - 1, 1) \in \mathcal{L}_{12}$, using that $(u_1^3, v_1^3) = (2, \pi_2)$ and $(u_i^3, v_i^3) = (3, 2) = (p_3(2, \pi_2), q_3(2, \pi_2))$ by (iv') and (4).

The concatenation $P_1P_2P_3\hat{P}_1\check{P}_1$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Case (ob2): $y_i \neq 1$ and $y_1 \neq 1$. We let $\pi = (\pi_1, \pi_2, \pi_3)$ be the permutation defined by $\pi_1 = x_i = 1$ and $\pi_2 = y_i$, and we define $\rho := \pi$. We choose $\hat{u}^1, \hat{v}^1 \in \Pi(\boldsymbol{a})$ as in case (ob2) in the proof of Lemma 16 for k = 3, and we construct the Hamilton path Q and its subpaths Q' and Q'' exactly as before. Also, we choose $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, 2, 3$, and $\check{u}^1, \check{v}^1 \in \Pi(\boldsymbol{a})$, as in the previous proof, subject to conditions (i)–(iii) as before and the following condition (iv') instead of (iv); see Figure 31 (ob2):

(iv') If $\pi_2 = 3$, we require that $u_{\tilde{i}}^2 = 3$ and $v_{\tilde{i}}^2 = 2$ for some $\tilde{i} \in [n] \setminus \{1, \check{i}, \hat{i}\}$ and condition (ψ_3) . If $\pi_3 = 3$, on the other hand, we require that $u_{\tilde{i}}^3 = 3$ and $v_{\tilde{i}}^3 = 2$ for some $\tilde{i} \in [n] \setminus \{1, \check{i}, \hat{i}\}$ with $z_{\tilde{i}} = 2$ and condition (ψ_2) .

We consider Hamilton paths P_1, P_2, P_3 and \check{P}_1 as before. If $j \in \{2, 3\}$ is such that $\pi_j = 3$, then the path P_j from u^j to v^j exists by the assumption $G(\alpha - 1, \alpha - 1, 1) \in \mathcal{L}_{12}$, using that $(u_1^j, v_1^j) = (2, 2)$ (clearly, $\pi_3 = 2$ if $\pi_2 = 3$ and $\pi_2 = 2$ if $\pi_3 = 3$) and $(u_{\tilde{i}}^j, v_{\tilde{i}}^j) = (3, 2) = (p_3(2, 2), q_3(2, 2))$ by (iv') and (4).

The concatenation $P_1Q'\check{P}_1P_2P_3Q''$ is a Hamilton path in G(a) from x to y.

Proof of Lemma 20. We proceed exactly as in the proof of Lemma 16 for k = 4, choosing indices \tilde{i}, \hat{i} and distinguishing cases (eb2), (eb1) and (ea) depending on the values of $y_{\tilde{i}}$ and y_1 , as before.

Case (eb2): $y_i \neq 1$ and $y_1 \neq 1$. If $y_i = 2$ we define the permutations $\pi := (1, 3, 4, y_i) = (1, 3, 4, 2)$ and $\rho := (1, 2, 3, 4)$, whereas if $y_i \in \{3, 4\}$ we define $\pi := (1, 7 - y_i, 2, y_i)$ and $\rho := (1, 3, 4, 2)$. We choose $u^j, v^j \in \Pi(\mathbf{a}), j = 1, 2, 3, 4$, and $\hat{u}^j, \hat{v}^j \in \Pi(\mathbf{a}), j = 1, 2$, and $\check{u}^1, \check{v}^1 \in \Pi(\mathbf{a})$, as in case (eb2) in the proof of Lemma 16 for k = 4, subject to the conditions (i)–(iii) as before and the following condition (iv') instead of (iv); see Figure 32 (eb2):

(iv') For $j \in \{2,3\}$ such that $\rho_j = 3$ we require that $u_{\tilde{i}}^j = 4$, $v_{\tilde{i}}^j = 2$, $u_{\tilde{i}}^{j+1} = 2$, and $v_{\tilde{i}}^{j+1} = 3$ for some $\tilde{i} \in [n] \setminus \{1, \tilde{i}, \hat{i}\}$, and condition (ψ_4) if j = 2.

For j=1,2,3,4 we consider a Hamilton path P_j in $G^{(\check{\imath},\hat{\imath}),(1,\rho_j)}(\boldsymbol{a})$ from u^j to v^j . For the two values of j with $\rho_j \in \{1,2\}$ such a path exists by the assumption $G(\alpha-1,\alpha-2,1,1) \in \mathcal{H}$. For the two values of j with $\rho_j \in \{3,4\}$ such a path exists by the assumption $G(\alpha-1,\alpha-1,1) \in \mathcal{L}_{12}$, using that $(u_1^j,v_1^j)=(2,4)$ and $(u_{\bar{\imath}}^j,v_{\bar{\imath}}^j)=(4,2)=(p_4(2,4),q_4(2,4))$ if $\rho_j=3$ and $(u_1^j,v_1^j)=(3,2)$ and $(u_{\bar{\imath}}^j,v_{\bar{\imath}}^j)=(2,3)=(q_3(2,3),p_3(2,3))$ if $\rho_j=4$, by (iv') and (4).

For j = 1, 2 we consider a Hamilton path \hat{P}_j in $G^{\check{\imath}, \pi_{2j}}(\boldsymbol{a})$ from \hat{u}^j to \hat{v}^j and a Hamilton path \check{P}_1 in $G^{\check{\imath}, \pi_3}(\boldsymbol{a})$ from \check{u}^1 to \check{v}^1 , which exist by the assumptions $G(\alpha, \alpha - 2, 1, 1) \in \mathcal{L}_1$ and $G(\alpha, \alpha - 1, 1) \in \mathcal{L}_1$.

The concatenation $P_1\hat{P}_1\check{P}_1P_2P_3P_4\hat{P}_2$ is a Hamilton path in G(a) from x to y.

Case (eb1): $y_i \neq 1$ and $y_1 = 1$. If $y_i = 2$ we define the permutations $\pi := (1, 3, 4, y_i) = (1, 3, 4, 2)$ and $\rho := (1, 3, 2, 4)$, whereas if $y_i \in \{3, 4\}$ we define $\pi := (1, 2, 7 - y_i, y_i)$ and $\rho := (1, y_i, 2, 7 - y_i)$. We choose $\hat{u}^1, \hat{v}^1 \in \Pi(\boldsymbol{a})$ as in case (eb1) in the proof of Lemma 16 for k = 4, and we construct the Hamilton path Q and its subpaths Q' and Q'' exactly as before. Also, we choose $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, 2, 3, 4$, and $\check{u}^j, \check{v}^j \in \Pi(\boldsymbol{a}), j = 1, 2$, as in the previous proof,




FIGURE 32. Illustration of the proof of Lemma 20.

subject to the conditions (i)–(iii) as before and the following condition (iv') instead of (iv); see Figure 32 (eb1):

(iv') We require that $u_{\tilde{i}}^2 = 2$, $v_{\tilde{i}}^2 = \pi_3$, $u_{\tilde{i}}^4 = \pi_2$, and $v_{\tilde{i}}^4 = \pi_4$ for some $\tilde{i} \in [n] \setminus \{1, \check{i}, \hat{i}\}$ with $z_{\tilde{i}} = \pi_4$. We consider Hamilton paths P_1, P_2, P_3, P_4 and \check{P}_1, \check{P}_2 as before. The paths $P_i, j \in \{2, 4\}$, from u^j to v^j exist by the assumption $G(\alpha - 1, \alpha - 1, 1) \in \mathcal{L}_{12}$, using that $(u_1^2, v_1^2) = (\pi_3, 2)$ and $(u_{\tilde{i}}^2, v_{\tilde{i}}^2) = (2, \pi_3) = (q_{\pi_3}(2, \pi_3), p_{\pi_3}(2, \pi_3)), \text{ and that } (u_1^4, v_1^4) = (2, \pi_2) \text{ and } (u_{\tilde{i}}^4, v_{\tilde{i}}^4) = (\pi_2, \pi_4),$ which equals $(p_3(2, \pi_2), q_3(2, \pi_2))$ if $\pi_2 = 3$ and $(q_{\pi_4}(\pi_2, 2), p_{\pi_4}(\pi_2, 2))$ if $\pi_2 = 2$ by (iv') and (4). The concatenation $P_1Q'\dot{P}_1P_2P_3P_4Q''\dot{P}_2$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.

Case (ea): $y_i = 1$. By the assumption that $x_i, y_i > 1$ and the fact that the symbols 3 and 4 both occur exactly once, we can now assume w.l.o.g. that $(x_{\hat{i}}, y_{\hat{i}}) \in \{(2,3), (3,4)\}$, and we treat these two cases separately.

Case (ea1): $(x_{\hat{i}}, y_{\hat{i}}) = (2, 3)$. We define the permutations $\pi := (1, 3, 4, 2)$ and $\rho := (2, 1, 4, 3)$, and we also define $t := y_1 \in \{1, 2, 4\}$ and $p := p_4(2, t) \in \{1, 4\}$. We fix indices $i_1, i_2, i_3 \in$ $[n] \setminus \{1, \check{i}, \hat{i}\}$ with $x_{i_1} \neq x_{i_2}$ and $y_{i_3} = 2$.

We choose $\hat{u}^1, \hat{v}^1 \in \Pi(a)$ as in case (ea) in the proof of Lemma 16 for k=4, and we construct the Hamilton path Q and its subpaths Q' and Q'' exactly as before. Also, we choose $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, 2, 3, 4, \text{ and } \check{u}^j, \check{v}^j \in \Pi(\boldsymbol{a}), j = 1, 2, \text{ as in the previous proof, subject to the}$ conditions (ii)-(iii) as before and the following condition (i') instead of (i) and (iv') instead of (iv); see Figure 32 (ea1):

- (i') $u^1=x, \ v^4=y, \ u_{i_3}^4=p,$ and conditions $(\chi^1), \ (\chi_3^2),$ and $(\chi^3);$ (iv') We require that $u_{\tilde{\imath}}^3=3$ and $v_{\tilde{\imath}}^3=1$ for some $\tilde{\imath}\in [n]\setminus\{1,\check{\imath},\hat{\imath}\}$ with $z_{\tilde{\imath}}=1.$

We consider Hamilton paths P_1, P_2, P_3, P_4 and P_1, P_2 as before. The paths $P_j, j \in \{3, 4\}$, from u^j to v^j exist by the assumption $G(\alpha - 1, \alpha - 1, 1) \in \mathcal{L}_{12}$, using that $(u_1^3, v_1^3) = (1, 3)$ and $(u_{\tilde{i}}^3, v_{\tilde{i}}^3) = (3, 1) = (p_3(1, 3), q_3(1, 3)) \text{ and } (u_1^4, v_1^4) = (2, t) \text{ and } (u_{i_3}^4, v_{i_3}^4) = (p, 2) = (p_4(2, t), q_4(2, t))$ by (i')+(iv') and (4).

Case (ea2): $(x_i, y_i) = (3, 4)$. This case is treated very differently than case (ea2) in the proof of Lemma 16. We define the permutations $\pi := (1,2,3,4)$ and $\rho := (3,1,2,4)$, and we also define $t := y_1 \in \{1, 2, 3\} \text{ and } p := p_3(2, t) \in \{1, 3\}, \text{ and } t' := x_1 \in \{1, 2, 4\} \text{ and } p' := p_4(2, t') \in \{1, 4\}.$ We fix indices $i_1, i_3 \in [n] \setminus \{1, \check{i}, \hat{i}\}$ with $x_{i_1} = 2$ and $y_{i_3} = 2$.

We choose two multiset permutations $\check{u}^1, \check{v}^1 \in \Pi(\boldsymbol{a})$ such that $\check{u}^1_1 = 1, \, \check{u}^1_{\hat{i}} = \rho_2 = 1, \, \check{u}^1_{\check{i}} = \pi_3 = 3,$ $\check{v}_1^1 = \pi_4 = 4$, and $\check{v}_{\check{i}}^1 = \pi_3 = 3$. We consider a Hamilton path Q in the graph $G^{\check{i},\pi_3}(a) = G^{\check{i},3}(a)$ from \check{u}^1 to \check{v}^1 , which exists by the assumption $G(\alpha, \alpha - 1, 1) \in \mathcal{L}_1$, using that $\check{u}^1 \in \Pi(\boldsymbol{a})^{1,1}$ and $\check{v}^1 \notin \Pi(a)^{1,1}$ (recall that $\check{u}_1^1 = 1$ and $\check{v}_1^1 = 4 \neq 1$). Let z be the first vertex along this path from \check{u}^1 to \check{v}^1 for which $z_{\hat{i}} = \pi_2 = 2$, and let z' be the predecessor of z on the path. By the definition of z we have $z_i' \neq \pi_2 = 2$, and consequently $z_1' = \pi_2 = 2$. By Lemma 14, we thus obtain $z'_{\hat{i}} = z_1 = 1$. Let Q' be the backward subpath of Q from z' to \check{u}^1 , and let Q'' be the backward subpath of Q from \check{v}^1 to z.

We choose multiset permutations $u^j, v^j \in \Pi(\boldsymbol{a}), j = 1, 2, 3, 4, \text{ and } \hat{u}^j, \hat{v}^j \in \Pi(\boldsymbol{a}), j = 1, 2,$ satisfying the following conditions; see Figure 32 (ea2):

- $\begin{array}{l} \text{(i)} \ \ u^1=x, \ v^1_{i_1}=p', \ \text{and} \ v^4=y, \ u^4_{i_3}=p; \\ \text{(ii)} \ \ \hat{v}^1_1=z'_{\check{i}}=\pi_3=3, \ \hat{v}^1_{\check{i}}=z'_1=\pi_2=2, \ \hat{v}^1_{\check{i}}=z'_i, \ u^2_1=\check{u}^1_{\check{i}}=\pi_3=3, \ u^2_{\check{i}}=\check{u}^1_1=1, \ u^2_{\check{i}}=\check{u}^1_{\check{i}}, \\ \hat{v}^2_1=\check{v}^1_{\check{i}}=\pi_3=3, \ \hat{v}^2_{\check{i}}=\check{v}^1_1=\pi_4=4, \ \hat{v}^2_{\check{i}}=\check{v}^1_{\check{i}}, \ \text{and} \ u^3_1=z_{\check{i}}=\pi_3=3, \ u^3_{\check{i}}=z_1=1, \ u^3_{\check{i}}=z_i \end{array}$ for all $i \in [n] \setminus \{1, \check{i}\};$
- (iii) conditions (ϕ_1^1) , (ϕ_2^1) and (ϕ_3^4) .

The argument that all pairs (u^j, v^j) , j = 1, 2, 3, 4, and (\hat{u}^j, \hat{v}^j) , j = 1, 2, are distinct, is straightforward.

For j = 1, 2, 3, 4 we define $\mathbf{b} := (a_1 - 1, a_2, a_3, a_4) = (\alpha - 1, \alpha - 1, 1, 1), \mathbf{c}' := (a_1 - 1, \dots, a_{\rho_j} - 1, \dots, a_4),$ and $\mathbf{c} := \varphi(\mathbf{c}') \in \{(\alpha - 1, \alpha - 1, 1) \ (\alpha - 1, \alpha - 2, 1, 1)\},$ which satisfy $\mathbf{c} \prec \mathbf{b} \prec \mathbf{a}$ and $\Delta(\mathbf{b}) = 2$ and $\Delta(\mathbf{c}) = 1$, and we consider a Hamilton path P_j in $G^{(\tilde{\imath}, \hat{\imath}), (1, \rho_j)}(\mathbf{a}) \simeq G(\mathbf{c})$ from u^j to v^j . For $j \in \{2, 3\}$ these paths exist by the assumption $G(\alpha - 1, \alpha - 2, 1, 1) \in \mathcal{H}$. For $j \in \{1, 4\}$ these paths exist by the assumption $G(\alpha - 1, \alpha - 1, 1) \in \mathcal{L}_{12}$, using that $(u_1^1, v_1^1) = (t', 2)$ and $(u_{i_1}^1, v_{i_1}^1) = (2, p') = (q_4(2, t'), p_4(2, t'))$ and $(u_1^4, v_1^4) = (2, t)$ and $(u_{i_3}^4, v_{i_3}^4) = (p, 2) = (p_3(2, t), q_3(2, t))$ by (i) and (4).

For j = 1, 2 we define $\mathbf{a}' := (a_1, \dots, a_{\pi_{2j}} - 1, \dots, a_4)$ and $\mathbf{b} := \varphi(\mathbf{a}') \in \{(\alpha, \alpha - 2, 1, 1), (\alpha, \alpha - 1, 1)\}$, which satisfies $\mathbf{b} \prec \mathbf{a}$ and $\Delta(\mathbf{b}) = 0$, and we consider a Hamilton path \hat{P}_j in the graph $G^{\check{\imath}, \pi_{2j}}(\mathbf{a}) \simeq G(\mathbf{b})$ from \hat{u}^j to \hat{v}^j , which exists by the assumption $G(\mathbf{b}) \in \mathcal{L}_1$, using that $\hat{u}^j \in \Pi(\mathbf{a})^{1,1}$ and $\hat{v}^j \notin \Pi(a)^{1,1}$ (recall that $\hat{u}^j_1 = 1$ and $\hat{v}^j_1 = 3 \neq 1$ by (ii)+(iii)).

The concatenation $P_1\hat{P}_1Q'P_2\hat{P}_2Q''P_3P_4$ is a Hamilton path in $G(\boldsymbol{a})$ from x to y.