Zeros of quasi-orthogonal q-Laguerre polynomials

Pinaki Prasad Kar^a, Priyabrat Gochhayat^b

Department of Mathematics, Sambalpur University, Sambalpur, 768019, Odisha, India ^aE-mail: pinakipk@gmail.com, ^bE-mail: pgochhayat@gmail.com

August 18, 2021

Abstract

We investigate the interlacing of zeros of polynomials of different degrees within the sequences of q-Laguerre polynomials $\left\{\tilde{L}_n^{(\delta)}(z;q)\right\}_{n=0}^{\infty}$ characterized by $\delta \in (-2,-1)$. The interlacing of zeros of quasi-orthogonal polynomials $\tilde{L}_n^{(\delta)}(z;q)$ with those of the orthogonal polynomials $\tilde{L}_m^{(\delta+t)}(z;q), m, n \in \mathbb{N}, t \in \{1,2\}$ is also considered. New bounds for the least zero of the (order 1) quasi-orthogonal q-Laguerre polynomials are derived.

MSC: 33D15; 33D45

Keywords: Zeros, Interlacing, Stieltjes interlacing, Quasi-orthogonal polynomials, *q*-Laguerre polynomials.

1 Introduction

Let $\{t_n(z)\}_{n=0}^{\infty}$, with $\deg(t_n) = n, n \in \mathbb{N}$, be a sequence of orthogonal polynomials and w(z) be the corresponding weight function on a support (c, d). Then a well known fact about the zeros of $t_n(z)$ is that, they are real, simple and lie in (c, d). Moreover, the zeros of $t_n(z)$ and $t_{n-1}(z)$ interlace, if there is exactly one zero of $t_{n-1}(z)$ in between any two consecutive zeros of $t_n(z)$. Similarly, the zeros of the two equal degree polynomials $t_n(z)$ and $u_n(z)$ are interlacing if there is exactly one zero of $u_n(z)$ (equivalently, $t_n(z)$) in between any two consecutive zeros of $t_n(z)$ (equivalently, $u_n(z)$). Another interlacing property given by Stieltjes (cf. [19, Theorem 3.3.3]), usually called Stieltjes interlacing, states that, for $m, n \in \mathbb{N}$ with m < n - 1, if there exists m open intervals with endpoints at the consecutive zeros of $t_n(z)$ then there is exactly one zero of $u_m(z)$ in each of the m open intervals. It follows that, if there is at least a zero of $u_m(z)$ lies outside the interval $(t_{1,n}, t_{n,n})$; where $t_{1,n}$ and $t_{n,n}$, respectively, are the smallest and the largest zero of $t_n(z)$, the Stieltjes interlacing will not retain between the zeros of $u_m(z)$ and $t_n(z)$ with n - 1 > m.

The departure of zeros of polynomials from the interval of orthogonality leads to the following definition of quasi-orthogonality:

Definition 1. Suppose $\{t_n(z)\}_{n=0}^{\infty}$ is a sequence of polynomials and $\deg(t_n) = n, n \in \mathbb{N}$, and w(z) is the corresponding weight function on a support [c, d]. Then $\{t_n(z)\}_{n=0}^{\infty}$ is called as quasi-orthogonal sequence of order r < n if

$$\int_{c}^{d} z^{i} t_{n}(z) w(z) dz \begin{cases} = 0; & i = 0, 1, \cdots, n - r - 1, \\ \neq 0; & i = n - r. \end{cases}$$
(1)

If $t_n(z)$ is a quasi-orthogonal polynomial of order r with respect to w(z) > 0 on (c, d), then there are minimum (n - r) real and distinct zeros of $t_n(z)$ in (c, d). In (1), letting r = 0 we find that $\{t_n(z)\}_{n=0}^{\infty}$ is an orthogonal polynomial sequence. Riesz [17] was the first mathematician who studied quasi-orthogonality of order 1 while order 2 was derived by Fejér [11]. Following which, Shohat [18], Chihara [3], Dickinson [4], Draux [5], etc. considered various general cases. For more information on quasi-orthogonality, we refer the works by Brezinski et al. [1, 2], Joulak [13] and the references therein. Recent contribution on the interlacing of zeros of different quasiorthogonal polynomials found in [6, 7, 8, 9, 10, 12]. More recently, in the context of q-calculus, the quasi-orthogonality of classical orthogonal polynomials on q-linear and q-quadratic lattices is discussed by Tcheutia et al. [20] while Kar et al. in [14] considered the polynomials that are not present in the q-Askey scheme of hypergeometric polynomials to study their quasi-orthogonality. Pertinently, very less known on quasi-orthogonality of q-polynomials.

In this work, we study the interlacing properties of zeros of *q*-Laguerre polynomials. Therefore, it is worthy to mention the orthogonality and quasi-orthogonality of these polynomials.

The (monic) q-Laguerre polynomials of degree n is defined as

$$\tilde{L}_{n}^{(\delta)}(z;q) = \frac{(-1)^{n}(q^{\delta+1};q)_{n}}{q^{n(\delta+n)}} \, _{1}\phi_{1} \begin{pmatrix} q^{-n} \\ q^{\delta+1};q,-q^{\delta+n+1}z \end{pmatrix}, \qquad \delta > -1$$

These are orthogonal with respect to the weight $\frac{z^{\delta}}{(-z;q)_{\infty}}$ on $(0,\infty)$. For any value of δ , the sequence of q-Laguerre polynomials is defined by the three term recurrence relation

$$\tilde{L}_{n}^{(\delta)}(z;q) = (z-a_{n})\tilde{L}_{n-1}^{(\delta)}(z;q) - \frac{(1-q^{n-1})(1-q^{\delta+n-1})}{q^{2\delta+4n-5}}\tilde{L}_{n-2}^{(\delta)}(z;q); \ n = 2, 3, \cdots,$$
(2)

where

$$a_n = \frac{1 - q^n + q(1 - q^{\delta + n - 1})}{q^{\delta + 2n - 1}},$$
(3)

and $\tilde{L}_{0}^{(\delta)}(z;q) \equiv 1$, $\tilde{L}_{1}^{(\delta)}(z;q) = z - \frac{1 - q^{\delta+1}}{q^{\delta+1}}$.

Tcheutia et al. in [20, Theorem 2.17], proved that for fixed $\delta \in \mathbb{R}$ with $\delta \in (-k-1, -k)$, $\left\{\tilde{L}_{n}^{(\delta)}(z;q)\right\}_{n=0}^{\infty}$ is quasi-orthogonal of order k with respect to $\frac{z^{\delta+k}}{(-z;q)_{\infty}}$. There are minimum n-k real and distinct zeros in $(0,\infty)$. Throughout this paper we denote $\{z_{i,n}\}_{i=1}^{n}, \{y_{i,n}\}_{i=1}^{n}$ and $\{x_{i,n}\}_{i=1}^{n}$, respectively, the zeros of $\tilde{L}_{n}^{(\delta)}(z;q), \tilde{L}_{n}^{(\delta+1)}(z;q)$ and $\tilde{L}_{n}^{(\delta+2)}(z;q)$ for fixed $\delta \in (-2, -1)$ and 0 < q < 1.

The following result obtained by substituting δ by $\delta + 1$ in [20, Theorem 2.18] is used in our work.

Lemma 1. The zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+1)}(z;q)$ are interlacing as

$$z_{1,n} < 0 < y_{1,n} < z_{2,n} < y_{2,n} < \dots < z_{n,n} < y_{n,n}.$$
(4)

Remark 1. Replacing n by n + 1 in [20, Theorem 2.18], we also obtain

$$z_{1,n+1} < 0 < y_{1,n} < z_{2,n+1} < y_{2,n} < \dots < z_{n,n+1} < y_{n,n} < z_{n+1,n+1}.$$
(5)

In the next section, we show that the zeros of the q-Laguerre polynomials of consecutive degree are not interlacing. The interlacing results between the zeros of the quasi-orthogonal and the orthogonal q-Laguerre polynomials of the same, consecutive and non-consecutive degree with different parameter values are also discussed in section 2. In section 3, we examine the common zeros of polynomials that are not co-prime. The last section is devoted to the derivation of the inner and outer bound for the only negative zero of the quasi-orthogonal q-Laguerre polynomial.

$\begin{array}{ll} \textbf{2} \quad \text{Zeros of } \tilde{L}_n^{(\delta)}(z;q) \text{ and } \tilde{L}_m^{(\delta+t)}(z;q) \text{ for } m \leq n+1,m,n \in \mathbb{N} \\ \text{ and } t \in \{0,1,2\} \end{array}$

Our first theorem proves that the interlacing does not take place between the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n+1}^{(\delta)}(z;q)$.

Theorem 1. The zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n+1}^{(\delta)}(z;q)$ are placed in the following order:

$$z_{1,n} < z_{1,n+1} < 0 < z_{2,n+1} < z_{2,n} < \dots < z_{n,n} < z_{n+1,n+1}.$$

Proof. Evaluating the mixed *q*-contiguous relation [16, (4.14)]

$$\tilde{L}_{n+1}^{(\delta)}(z;q) = -\frac{1-q^{\delta+n+1}}{q^{\delta+2n+1}}\tilde{L}_{n}^{(\delta)}(z;q) + z\tilde{L}_{n}^{(\delta+1)}(z;q)$$

at the consecutive zeros $z_{i,n}$ and $z_{i+1,n}$, $i \in \{1, \dots, n-1\}$, of $\tilde{L}_n^{(\delta)}(z;q)$, we obtain

$$\tilde{L}_{n+1}^{(\delta)}(z_{i,n};q)\tilde{L}_{n+1}^{(\delta)}(z_{i+1,n};q) = z_{i,n}z_{i+1,n}\tilde{L}_{n}^{(\delta+1)}(z_{i,n};q)\tilde{L}_{n}^{(\delta+1)}(z_{i+1,n};q).$$
(6)

Now from (4), $z_{1,n}z_{2,n} < 0$ and $z_{i,n}z_{i+1,n} > 0$ for $i \in \{2, \dots, n-1\}$ while for $i \in \{1, \dots, n-1\}$, $\tilde{L}_{n}^{(\delta+1)}(z_{i,n};q)\tilde{L}_{n}^{(\delta+1)}(z_{i+1,n};q) < 0$. We conclude from (6) that $\tilde{L}_{n+1}^{(\delta)}(z_{1,n};q)\tilde{L}_{n+1}^{(\delta)}(z_{2,n};q) > 0$ and $\tilde{L}_{n+1}^{(\delta)}(z_{i,n};q)\tilde{L}_{n+1}^{(\delta)}(z_{i+1,n};q) < 0$ for each $i \in \{2, \dots, n-1\}$. This implies that $\tilde{L}_{n+1}^{(\delta)}(z;q)$ has an even number of zeros in $(z_{1,n}, z_{2,n})$ and an odd number of zeros in $(z_{i,n}, z_{i+1,n})$ for each $i \in \{2, \dots, n-1\}$. Hence, $\tilde{L}_{n+1}^{(\delta)}(z;q)$ has minimum n-2 zeros in $(z_{2,n}, z_{n,n})$, plus its least zero $z_{1,n+1} < 0$ and from (4) and (5), its greatest zero $z_{n+1,n+1} > y_{n,n} > z_{n,n}$. Thus, we have located n zeros of $\tilde{L}_{n+1}^{(\delta)}(z;q)$ in $(z_{1,n}, z_{2,n})$ where $z_{1,n} < 0 < z_{2,n}$ for any $n \in \mathbb{N}$, we must have either no zero or two zeros in this single interval. Because we have precisely one negative zero of $\tilde{L}_{n+1}^{(\delta)}(z;q)$ for $n \in \mathbb{N}$, the only option is $z_{1,n} < z_{1,n+1} < 0 < z_{2,n+1} < z_{2,n}$. This proves the theorem.

It is clear from Theorem 1 that, the zeros of the quasi-orthogonal polynomials $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n+1}^{(\delta)}(z;q)$ are interlacing in the interval $(0,\infty)$. Furthermore, the zeros of $z\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n+1}^{(\delta)}(z;q)$ interlace.

The next result shows that, the Stieltjes interlacing fails to occur between the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$.

Theorem 2. For $n \ge 3, n \in \mathbb{N}$, if $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$ are co-prime, then the zeros of $z(z - a_n)\tilde{L}_{n-2}^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta)}(z;q)$ interlace, where a_n is given in (3). However, the Stieltjes interlacing will not take place between the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$.

Proof. Assume that $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$ are co-prime. Then from (2), $\tilde{L}_n^{(\delta)}(a_n;q) \neq 0$. Evaluate (2) at the consecutive zeros $z_{i,n}$ and $z_{i+1,n}$, $i \in \{2, \dots, n-1\}$, of $\tilde{L}_n^{(\delta)}(z;q)$ to get

$$\tilde{L}_{n-1}^{(\delta)}(z_{i,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{i+1,n};q) = \frac{(1-q^{n-1})^2(1-q^{\delta+n-1})^2}{q^{4\delta+8n-10}(z_{i,n}-a_n)(z_{i+1,n}-a_n)}.$$
(7)

For *n* substituted by n-1 in Theorem 1, we have the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-1}^{(\delta)}(z;q)$ interlace in $(0,\infty)$. So that for every $i \in \{2, \cdots, n-1\}$, the numerator of left-hand side of (7) is negative. Also the right-hand side of (7) is positive unless $a_n \in (z_{i,n}, z_{i+1,n})$ for each $i \in \{2, \cdots, n-1\}$. Hence we deduce from (7) that, for every $i \in \{2, \cdots, n-1\}$, $\tilde{L}_{n-2}^{(\delta)}(z_{i,n};q)$ and $\tilde{L}_{n-2}^{(\delta)}(z_{i+1,n};q)$ have different sign except may be for one pair $z_{j,n}, z_{j+1,n}$, with $z_{j,n} < a_n < z_{j+1,n}, 2 \le j \le n-1$. Since n-2 intervals are there with endpoints at the consecutive positive zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$ has exactly n-3 simple positive zeros, the zeros of $\tilde{L}_{n-2}^{(\delta)}(z;q)$, together with the point a_n , interlace with the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ in $(0,\infty)$. The given interlacing result holds true because from Theorem 1, we have $z_{1,n-2} < z_{1,n} < 0 < z_{2,n}$. Further, from Theorem 1, $z_{1,n-2} < z_{1,n-1} <$ $z_{1,n} < 0 < z_{2,n} < z_{2,n-1} < z_{2,n-2}$, the smallest zero $z_{1,n-2}$, of $\tilde{L}_{n-2}^{(\delta)}(z;q)$, lies outside $(z_{1,n}, z_{n,n})$, which shows that the Stieltjes interlacing does not hold between the zeros of the quasi-orthogonal q-Laguerre polynomials $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$.

In Lemma 1, we have for fixed $\delta \in (-2, -1)$, the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+1)}(z;q)$ are interlacing. Upon replacing n by n-1 in Remark 1, we see that the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-1}^{(\delta+1)}(z;q)$ interlace. The interlacing properties between the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta+1)}(z;q)$ are discussed in the next theorem. Note that, for fixed $\delta \in (-2, -1)$ and $k \ge 1$, the sequence $\left\{\tilde{L}_n^{(\delta+k)}(z;q)\right\}_{n=0}^{\infty}$ is orthogonal in $(0,\infty)$ because $\delta + k > -1$.

Theorem 3. For $n \ge 3, n \in \mathbb{N}$, the zeros of $\tilde{L}_{n-2}^{(\delta+1)}(z;q)$, together with the point $\frac{1-q^{\delta+n}}{q^{\delta+2n-1}} = b_n$ (say), interlace with those of $\tilde{L}_n^{(\delta)}(z;q)$ in $(0,\infty)$ provided these two polynomials do not have any common zeros.

Proof. Using [16, (4.14) and (4.16)] and [14, (10)], we get

$$\tilde{L}_{n}^{(\delta)}(z;q) = (z-b_{n})\,\tilde{L}_{n-1}^{(\delta)}(z;q) - \left(\frac{1-q^{n-1}}{q^{\delta+2n-2}}\right)z\tilde{L}_{n-2}^{(\delta+1)}(z;q).$$
(8)

From (8), $z_{i,n} \neq b_n$ for any $i \in \{1, \dots, n\}$, if not, this would contradict our supposition. Evaluating (8) at the consecutive positive zeros $z_{i,n}$ and $z_{i+1,n}$, $i \in \{2, \dots, n-1\}$, of $\tilde{L}_n^{(\delta)}(z;q)$ yields

$$\frac{\tilde{L}_{n-1}^{(\delta)}(z_{i,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{i+1,n};q)}{\tilde{L}_{n-2}^{(\delta+1)}(z_{i,n};q)\tilde{L}_{n-2}^{(\delta+1)}(z_{i+1,n};q)} = \frac{(1-q^{n-1})^2 z_{i,n} z_{i+1,n}}{q^{2\delta+4n-4}(z_{i,n}-b_n)(z_{i+1,n}-b_n)}.$$
(9)

Clearly, (4) shows that, both $z_{i,n}$ and $z_{i+1,n}$ are positive for each $i \in \{2, \dots, n-1\}$. Thus, the righthand side of (9) is positive if and only if for each $i \in \{2, \dots, n-1\}$, $b_n \notin (z_{i,n}, z_{i+1,n})$. Also from Theorem 1 we have for every $n \ge 2, n \in \mathbb{N}$ and $i \in \{2, \dots, n-1\}$, $\tilde{L}_{n-1}^{(\delta)}(z_{i,n}; q)\tilde{L}_{n-1}^{(\delta)}(z_{i+1,n}; q) < 0$. If $b_n \notin (z_{i,n}, z_{i+1,n})$, for every $i \in \{2, \dots, n-1\}$, then we deduce from (9) that the zeros of $\tilde{L}_{n-2}^{(\delta+1)}(z; q)$ and $\tilde{L}_n^{(\delta)}(z; q)$ interlace in $(0, \infty)$. Additionally, since the point b_n is not in $(z_{2,n}, z_{n,n})$, the zeros of $(z - b_n)\tilde{L}_{n-2}^{(\delta+1)}(z; q)$ and the positive zeros of $\tilde{L}_n^{(\delta)}(z; q)$ interlace. On the other hand, if $b_n \in (z_{i,n}, z_{i+1,n})$, then we have $b_n \in (z_{j,n}, z_{j+1,n})$ for one j with $2 \le j \le n - 1$. This implies, in $(z_{j,n}, z_{j+1,n}), \tilde{L}_{n-2}^{(\delta+1)}(z; q)$ has no sign change but it changes sign in each of the remaining n - 3intervals with endpoints at the consecutive positive zeros of $\tilde{L}_n^{(\delta)}(z; q)$. Finally, for i = 1, (9) becomes

$$\frac{\tilde{L}_{n-1}^{(\delta)}(z_{1,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{2,n};q)}{\tilde{L}_{n-2}^{(\delta+1)}(z_{1,n};q)\tilde{L}_{n-2}^{(\delta+1)}(z_{2,n};q)} = \frac{(1-q^{n-1})^2 z_{1,n} z_{2,n}}{q^{2\delta+4n-4}(z_{1,n}-b_n)(z_{2,n}-b_n)}.$$
(10)

Since $b_n \in (z_{j,n}, z_{j+1,n})$ for one $j, j \in \{2, \dots, n-1\}, b_n \notin (z_{1,n}, z_{2,n})$. We know from Theorem 1 that $z_{1,n-1} < z_{1,n} < 0 < z_{2,n} < z_{2,n-1}$. This implies, $\tilde{L}_{n-1}^{(\delta)}(z_{1,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{2,n};q) > 0$. Hence from (10), $\tilde{L}_{n-2}^{(\delta+1)}(z_{1,n};q)\tilde{L}_{n-2}^{(\delta+1)}(z_{2,n};q) < 0$. Consequently, $\tilde{L}_{n-2}^{(\delta+1)}(z;q)$ has one positive zero less than $z_{2,n}$. Thus, in each of the cases, we have the zeros of $(z - b_n)\tilde{L}_{n-2}^{(\delta+1)}(z;q)$ interlace with the positive zeros of $\tilde{L}_n^{(\delta)}(z;q)$, if these two polynomials do not have any common zeros.

For the rest of the results in this section we study the interlacing results for the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-k}^{(\delta+2)}(z;q)$, where $\delta \in (-2,-1)$ and $k \in \{0,1,2\}$. We now show a necessary and sufficient condition for the zeros of the same degree polynomials $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ to be interlacing.

Theorem 4. For $n \ge 2, n \in \mathbb{N}$, the zeros of the same degree q-Laguerre polynomials $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ interlace if and only if the smallest positive zero of $\tilde{L}_n^{(\delta)}(z;q)$, that is, $z_{2,n} > -\frac{1-q^{\delta+1}}{q^{\delta+n+1}} = c_n$ (say).

Proof. The sequence $\left\{\tilde{L}_n^{(\delta)}(z;q)\right\}_{n=0}^{\infty}$ satisfies the mixed q-contiguous relation (obtained using [16, (4.12) and (4.15)])

$$z\tilde{L}_{n}^{(\delta+2)}(z;q) = (z-c_{n})\,\tilde{L}_{n}^{(\delta+1)}(z;q) - \frac{1-q^{\delta+n+1}}{q^{\delta+2n+1}}\tilde{L}_{n}^{(\delta)}(z;q).$$
(11)

We have from (4) that $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+1)}(z;q)$ are co-prime for each fixed $\delta \in (-2,-1)$. Hence, it follows from (11) that $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ can have at most one common zero at $z = c_n$. If $c_n < z_{2,n}$, then since the zeros of $\tilde{L}_n^{(\delta+2)}(z;q)$ lie in $(0,\infty)$ and $z_{2,n} > 0$, the polynomials $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ are co-prime. Evaluating (11) at the consecutive zeros $z_{i,n}$ and $z_{i+1,n}, i \in \{1, \dots, n-1\}$, of $\tilde{L}_n^{(\delta)}(z;q)$, gives

$$\frac{\tilde{L}_{n}^{(\delta+1)}(z_{i,n};q)\tilde{L}_{n}^{(\delta+1)}(z_{i+1,n};q)}{\tilde{L}_{n}^{(\delta+2)}(z_{i,n};q)\tilde{L}_{n}^{(\delta+2)}(z_{i+1,n};q)} = \frac{z_{i,n}z_{i+1,n}}{(z_{i,n}-c_{n})(z_{i+1,n}-c_{n})}.$$
(12)

From (4), $z_{1,n}z_{2,n} < 0$ while for each $i \in \{2, \dots, n-1\}$, $z_{i,n}z_{i+1,n} > 0$. Further, for every $i \in \{1, \dots, n-1\}$, $\tilde{L}_n^{(\delta+1)}(z_{i,n};q)$ $\tilde{L}_n^{(\delta+1)}(z_{i+1,n};q) < 0$. Since $z_{1,n} < 0 < c_n < z_{2,n}$, we conclude from (12) that $\tilde{L}_n^{(\delta+2)}(z;q)$ changes sign in $(z_{i,n}, z_{i+1,n})$ for each $i \in \{1, \dots, n-1\}$. Consequently,

 $\tilde{L}_n^{(\delta+2)}(z;q)$ has an odd number of zeros in $(z_{i,n}, z_{i+1,n})$ for any $i \in \{1, \dots, n-1\}$. We note from [16, Theorem 3] that,

$$0 < y_{1,n} < x_{1,n} < y_{2,n} < x_{2,n} < \dots < y_{n,n} < x_{n,n}.$$
(13)

Now from (4) and (13), we have $z_{n,n} < x_{n,n}$, where $x_{n,n}$ is the largest zero of $\tilde{L}_n^{(\delta+2)}(z;q)$. Therefore, the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ are interlacing.

If $c_n > z_{2,n}$, then since we have $z_{1,n} < 0 < z_{2,n} < c_n$ so from (12), we deduce that $\tilde{L}_n^{(\delta+2)}(z_{1,n};q)$ and $\tilde{L}_n^{(\delta+2)}(z_{2,n};q)$ have same sign. Thus, $\tilde{L}_n^{(\delta+2)}(z;q)$ has an even number of zeros in $(z_{1,n}, z_{2,n})$ which confirms that the zeros of the polynomials $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ are not interlacing. This completes the proof of the theorem.

In the next theorem we derive an interlacing result between the zeros of the above considered polynomials involving the point c_n .

Theorem 5. For $n \ge 2, n \in \mathbb{N}$, let $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ be co-prime and $z_{2,n} < c_n$, where c_n is defined in Theorem 4. Then the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ interlace, if $c_n \notin (x_{i,n}, x_{i+1,n})$ for each $i \in \{1, \dots, n-1\}$. Otherwise, the zeros of $\tilde{L}_n^{(\delta)}(z;q)$, together with the point c_n , interlace with those of $z \tilde{L}_n^{(\delta+2)}(z;q)$.

Proof. Evaluating (11) at the consecutive zeros $x_{i,n}$ and $x_{i+1,n}$, $i \in \{1, \dots, n-1\}$, of $\tilde{L}_n^{(\delta+2)}(z;q)$, we obtain

$$(x_{i,n} - c_n)(x_{i+1,n} - c_n)\tilde{L}_n^{(\delta+1)}(x_{i,n};q)\tilde{L}_n^{(\delta+1)}(x_{i+1,n};q) = \left(\frac{1 - q^{\delta+n+1}}{q^{\delta+2n+1}}\right)^2 \tilde{L}_n^{(\delta)}(x_{i,n};q)\tilde{L}_n^{(\delta)}(x_{i+1,n};q).$$
(14)

It is clear from (13) that, $\tilde{L}_n^{(\delta+1)}(x_{i,n};q)\tilde{L}_n^{(\delta+1)}(x_{i+1,n};q) < 0$ for every $i \in \{1, \dots, n-1\}$. If for each $i \in \{1, \dots, n-1\}$, $c_n \notin (x_{i,n}, x_{i+1,n})$, then the left-hand side of (14) is negative, whence $\tilde{L}_n^{(\delta)}(x_{i,n};q)\tilde{L}_n^{(\delta)}(x_{i+1,n};q) < 0$. Since $\tilde{L}_n^{(\delta+2)}(z;q)$ has positive zeros and the least zero of $\tilde{L}_n^{(\delta)}(z;q)$ is negative, the only possible arrangement of zeros is

$$z_{1,n} < 0 < x_{1,n} < z_{2,n} < x_{2,n} < \dots < z_{n,n} < x_{n,n} < c_n,$$

which shows the first part of the theorem.

On the other hand, if $c_n \in (x_{j,n}, x_{j+1,n})$ for one $j, j \in \{1, \dots, n-1\}$, then we deduce from (14) that, $\tilde{L}_n^{(\delta)}(x_{j,n};q)\tilde{L}_n^{(\delta)}(x_{j+1,n};q) > 0$ for one $j, j \in \{1, \dots, n-1\}$. This shows that $\tilde{L}_n^{(\delta)}(z;q)$ has n-2 sign changes between the consecutive zeros of $\tilde{L}_n^{(\delta+2)}(z;q)$ in $(0,\infty)$ except for the one interval $(x_{j,n}, x_{j+1,n})$ containing the point c_n . Since $\tilde{L}_n^{(\delta)}(z;q)$ has n-1 positive zeros and n-2 sign changes in $(0,\infty)$, no zero of $\tilde{L}_n^{(\delta)}(z;q)$ lies in $(x_{j,n}, x_{j+1,n})$ that contains c_n and one zero of $\tilde{L}_n^{(\delta)}(z;q)$ is either $< x_{1,n}$ or $> x_{n,n}$. From (4) and (13), we have $z_{n,n} < y_{n,n} < x_{n,n}$. Hence, the only option is that the least positive zero $z_{2,n}$ of $\tilde{L}_n^{(\delta)}(z;q)$ is $< x_{1,n}$. Since exactly one zero of $\tilde{L}_n^{(\delta)}(z;q)$ is negative, therefore, the zeros of $(z-c_n)\tilde{L}_n^{(\delta)}(z;q)$ and $z\tilde{L}_n^{(\delta+2)}(z;q)$ interlace. This proves the theorem.

We analyze the interlacing properties between the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-1}^{(\delta+2)}(z;q)$ in the next result.

Theorem 6. For $n \geq 2, n \in \mathbb{N}$, the zeros of $z \tilde{L}_{n-1}^{(\delta+2)}(z;q)$ and $\tilde{L}_n^{(\delta)}(z;q)$ interlace.

Proof. Using [16, (4.12) and (4.14)], we obtain

$$\left(\frac{1-q^{\delta+n+1}}{1-q^n}\right)\tilde{L}_n^{(\delta)}(z;q) = \frac{q^n(1-q^{\delta+1})}{(1-q^n)}\tilde{L}_n^{(\delta+1)}(z;q) + z\tilde{L}_{n-1}^{(\delta+2)}(z;q).$$
(15)

Evaluating (15) at the consecutive zeros $z_{i,n}$ and $z_{i+1,n}$, $i \in \{1, \dots, n-1\}$, of $\tilde{L}_n^{(\delta)}(z;q)$ yields

$$\frac{q^{2n}(1-q^{\delta+1})^2}{(1-q^n)^2}\tilde{L}_n^{(\delta+1)}(z_{i,n};q)\tilde{L}_n^{(\delta+1)}(z_{i+1,n};q) = z_{i,n}z_{i+1,n}\tilde{L}_{n-1}^{(\delta+2)}(z_{i,n};q)\tilde{L}_{n-1}^{(\delta+2)}(z_{i+1,n};q).$$
 (16)

For any $i \in \{1, \dots, n-1\}$, (4) implies $\tilde{L}_n^{(\delta+1)}(z_{i,n};q)\tilde{L}_n^{(\delta+1)}(z_{i+1,n};q) < 0$. Also for i = 1, $z_{i,n}z_{i+1,n} < 0$ while $z_{i,n}z_{i+1,n} > 0$ for $i = 2, \dots, n-1$. Therefore, from (16), we deduce that $\tilde{L}_{n-1}^{(\delta+2)}(z;q)$ does not change sign in $(z_{1,n}, z_{2,n})$ and changes sign in $(z_{i,n}, z_{i+1,n})$ for every $i \in \{2, \dots, n-1\}$. Thus, we have minimum n-2 zeros of $\tilde{L}_{n-1}^{(\delta+2)}(z;q)$ in $(z_{2,n}, z_{n,n})$ and in $(z_{1,n}, z_{2,n})$, there is either no zero or two zeros of $\tilde{L}_{n-1}^{(\delta+2)}(z;q)$. Since $\tilde{L}_{n-1}^{(\delta+2)}(z;q)$ has exactly n-1 zeros, it has no zero in $(z_{1,n}, z_{2,n})$. Hence, the remaining one zero must be larger than $z_{n,n}$. Since $z_{1,n} < 0 < z_{2,n}$, the given interlacing property holds true. This proves the theorem.

Remark 2. From (4), $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+1)}(z;q)$ are co-prime. This implies, $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-1}^{(\delta+2)}(z;q)$ cannot be co-prime. Otherwise, any common zero of these polynomials is also a zero of $\tilde{L}_n^{(\delta+1)}(z;q)$.

Next theorem shows the interlacing between the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$.

Theorem 7. For $n \ge 3, n \in \mathbb{N}$, the zeros of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ interlace in $(0,\infty)$. Moreover, the zeros of $z(z + c_{n-1})\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ and $\tilde{L}_n^{(\delta)}(z;q)$ interlace, where c_n is given in Theorem 4.

Proof. Using (8) and [16, (4.14) and (4.15)], we get

$$\left(\frac{1-q^{n-1}}{1-q^{\delta+n}}\right)z^2\tilde{L}_{n-2}^{(\delta+2)}(z;q) = \frac{q^{\delta+2n}}{1-q^{\delta+n}}c_n\tilde{L}_n^{(\delta)}(z;q) + (z+c_{n-1})\tilde{L}_{n-1}^{(\delta)}(z;q).$$
(17)

Since $\tilde{L}_{n}^{(\delta)}(z;q)$ and $\tilde{L}_{n-1}^{(\delta)}(z;q)$ are co-prime in $(0,\infty)$, from (17), the only possible common zero of $\tilde{L}_{n}^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ in $(0,\infty)$ is $-c_{n-1}$. But for $\delta \in (-2,-1)$ and 0 < q < 1, $-c_{n-1} < 0$ and all the zeros of $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ are positive. This implies, the polynomials $\tilde{L}_{n}^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ are co-prime and hence, $z_{i,n} \neq -c_{n-1}$ for any $i \in \{1, \dots, n\}$. Evaluating (17) at the consecutive zeros $z_{i,n}$ and $z_{i+1,n}, i \in \{2, \dots, n-1\}$, of $\tilde{L}_{n}^{(\delta)}(z;q)$, we have

$$\frac{\tilde{L}_{n-2}^{(\delta+2)}(z_{i,n};q)\tilde{L}_{n-2}^{(\delta+2)}(z_{i+1,n};q)}{\tilde{L}_{n-1}^{(\delta)}(z_{i,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{i+1,n};q)} = \frac{(1-q^{\delta+n})^2(z_{i,n}+c_{n-1})(z_{i+1,n}+c_{n-1})}{(1-q^{n-1})^2 z_{i,n}^2 z_{i+1,n}^2}.$$
(18)

We know from Theorem 1 that, $\tilde{L}_{n-1}^{(\delta)}(z_{i,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{i+1,n};q) < 0$ for $i \in \{2, \dots, n-1\}$. For $\delta \in (-2, -1), -c_{n-1} < 0$, so $-c_{n-1} < z_{2,n}$ and hence for any $i \in \{2, \dots, n-1\}$, the right-hand side of (18) is positive. Therefore, we deduce from (18) that, for each $i \in \{2, \dots, n-1\}$, $\tilde{L}_{n-2}^{(\delta+2)}(z_{i,n};q)$ and $\tilde{L}_{n-2}^{(\delta+2)}(z_{i+1,n};q)$ have different sign. This means, the zeros of $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ and $\tilde{L}_{n}^{(\delta)}(z;q)$ interlace in $(0,\infty)$.

Now evaluating (17) at the zeros $z_{1,n}$ and $z_{2,n}$, of $\tilde{L}_n^{(\delta)}(z;q)$, we obtain

$$\frac{\tilde{L}_{n-2}^{(\delta+2)}(z_{1,n};q)\tilde{L}_{n-2}^{(\delta+2)}(z_{2,n};q)}{\tilde{L}_{n-1}^{(\delta)}(z_{1,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{2,n};q)} = \frac{(1-q^{\delta+n})^2(z_{1,n}+c_{n-1})(z_{2,n}+c_{n-1})}{(1-q^{n-1})^2 z_{1,n}^2 z_{2,n}^2}.$$
(19)

Since the zeros of $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ lie in $(z_{2,n}, z_{n,n})$, no zero of $\tilde{L}_{n-2}^{(\delta+2)}(z;q)$ are in $(z_{1,n}, z_{2,n})$ which means it does not change sign in $(z_{1,n}, z_{2,n})$. From Theorem 1, $\tilde{L}_{n-1}^{(\delta)}(z_{1,n};q)\tilde{L}_{n-1}^{(\delta)}(z_{2,n};q) > 0$. Therefore, from (19) we conclude that $-c_{n-1} \notin (z_{1,n}, z_{2,n})$ and since $-c_{n-1} < 0$, it follows that $-c_{n-1} < z_{1,n}$. Thus, the only possibility of configuration of zeros is as

$$-c_{n-1} < z_{1,n} < 0 < z_{2,n} < x_{1,n-2} < \dots < x_{n-2,n-2} < z_{n,n}$$

This follows the result.

3 Common zeros of $ilde{L}_n^{(\delta)}(z;q)$

In this section, we deal with the q-Laguerre polynomials that are not co-prime. In particular, we discuss the interlacing properties between the zeros of polynomials considered in Theorem 2 and Theorem 5.

Theorem 8. For $n \ge 3, n \in \mathbb{N}$, if $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$ are not co-prime, then a_n is a common zero of these polynomials and the zeros of $\tilde{L}_{n-2}^{(\delta)}(z;q)$ interlace with the non-common zeros of $\tilde{L}_n^{(\delta)}(z;q)$ in $(0,\infty)$, where a_n is given in (3).

Proof. If $\tilde{L}_{n}^{(\delta)}(z;q)$ and $\tilde{L}_{n-2}^{(\delta)}(z;q)$ have common zeros in $(0,\infty)$, then from (2), a_{n} is the only common zero of these polynomials since $\tilde{L}_{n}^{(\delta)}(z;q)$ and $\tilde{L}_{n-1}^{(\delta)}(z;q)$ are co-prime in $(0,\infty)$ by Theorem (1). Suppose $z'_{1,n-2} < z'_{2,n-2} < \cdots < z'_{n-2,n-2}$ are the n-2 non-common zeros of $\tilde{L}_{n}^{(\delta)}(z;q)$ in $(0,\infty)$. Then a_{n} belongs to one of the intervals $(z'_{i,n-2}, z'_{i+1,n-2})$, $i \in \{1, \cdots, n-3\}$. Evaluating (2) at the consecutive non-common zeros $z'_{i,n-2}$ and $z'_{i+1,n-2}$, $i \in \{1, \cdots, n-3\}$, of $\tilde{L}_{n}^{(\delta)}(z;q)$, such that $a_{n} \notin (z'_{i,n-2}, z'_{i+1,n-2})$, we obtain

$$\frac{(1-q^{n-1})^2 \left(1-q^{\delta+n-1}\right)^2}{q^{4\delta+8n-10}} \tilde{L}_{n-2}^{(\delta)}(z'_{i,n-2};q) \tilde{L}_{n-2}^{(\delta)}(z'_{i+1,n-2};q) = \left(z'_{i,n-2}-a_n\right) \left(z'_{i+1,n-2}-a_n\right) \tilde{L}_{n-1}^{(\delta)}(z'_{i,n-2};q) \tilde{L}_{n-1}^{(\delta)}(z'_{i+1,n-2};q).$$
(20)

Since the positive zeros of $\tilde{L}_{n}^{(\delta)}(z;q)$ and $\tilde{L}_{n-1}^{(\delta)}(z;q)$ are interlacing, we deduce from (20) that $\tilde{L}_{n-2}^{(\delta)}(z;q)$ has an odd number of zeros in each of the intervals $(z'_{i,n-2}, z'_{i+1,n-2})$, $i \in \{1, \dots, n-3\}$, that does not contain a_n . Since $\tilde{L}_{n-2}^{(\delta)}(z;q)$ has n-4 non-common zeros in $(0,\infty)$, there are at most n-4 such intervals. Therefore, $a_n = z_{j,n}$ where $j \in \{3, \dots, n-1\}$. Hence, the n-3 zeros of $\tilde{L}_{n-2}^{(\delta)}(z;q)$ (including a_n) interlace with the n-2 non-common zeros of $\tilde{L}_n^{(\delta)}(z;q)$ in $(0,\infty)$. This proves the theorem.

Theorem 9. Suppose $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ are not co-prime and the smallest positive zero of $\tilde{L}_n^{(\delta)}(z;q)$, that is, $z_{2,n} < c_n$ for $n \ge 2, n \in \mathbb{N}$, where c_n is given in Theorem 4. Then the above polynomials have a common zero at $z = c_n$ and the positive zeros of $\tilde{L}_n^{(\delta)}(z;q)$ interlace with the non-common zeros of $\tilde{L}_n^{(\delta+2)}(z;q)$.

Proof. We have by (4), $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+1)}(z;q)$ are co-prime. Therefore, from (11), c_n is the only common zero of $\tilde{L}_n^{(\delta)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ in $(0,\infty)$. If $x'_{1,n-1} < x'_{2,n-1} < \cdots < x'_{n-1,n-1}$ are the n-1 non-common zeros of $\tilde{L}_n^{(\delta+2)}(z;q)$, then c_n lies in at most one interval $(x'_{i,n-1}, x'_{i+1,n-1})$, $i \in \{1, \cdots, n-2\}$. Evaluating (11) at the consecutive non-common zeros $x'_{i,n-1}$ and $x'_{i+1,n-1}$, $i \in \{1, \cdots, n-2\}$, of $\tilde{L}_n^{(\delta+2)}(z;q)$ such that $c_n \notin (x'_{i,n-1}, x'_{i+1,n-1})$, yields

$$(x'_{i,n-1} - c_n) (x'_{i+1,n-1} - c_n) \tilde{L}_n^{(\delta+1)} (x'_{i,n-1}; q) \tilde{L}_n^{(\delta+1)} (x'_{i+1,n-1}; q) = \left(\frac{1 - q^{\delta+n+1}}{q^{\delta+2n+1}}\right)^2 \tilde{L}_n^{(\delta)} (x'_{i,n-1}; q) \tilde{L}_n^{(\delta)} (x'_{i+1,n-1}; q).$$
(21)

We have from (13) that, the zeros of $\tilde{L}_n^{(\delta+1)}(z;q)$ and $\tilde{L}_n^{(\delta+2)}(z;q)$ interlace. Thus, we deduce from (21) that $\tilde{L}_n^{(\delta)}(z;q)$ has an odd number of zeros in $(x'_{i,n-1}, x'_{i+1,n-1})$ for each $i \in \{1, \dots, n-2\}$, that does not contain the point c_n . From Theorem 5, we have $z_{2,n} < x_{1,n}$ when $c_n \in (x_{1,n}, x_{n,n})$. Since $\tilde{L}_n^{(\delta)}(z;q)$ has n-3 non-common zeros in $(x_{1,n}, x_{n,n})$, there are at most n-3 such intervals. Hence, $c_n = x_{j,n}$ where $j \in \{2, \dots, n-1\}$. Thus, the n-2 non-common zeros of $\tilde{L}_n^{(\delta+2)}(z;q)$. This completes the proof.

We illustrate Theorem 8 and Theorem 9 by the following numeric examples using *Mathematica* 12 :

Example 1. If we take q = 0.997, n = 26 and $\delta = -1.121695$, the polynomials $\tilde{L}_{24}^{(-1.121695)}(z; 0.997)$ and $\tilde{L}_{26}^{(-1.121695)}(z; 0.997)$ have a common zero in $(0, \infty)$, namely, $a_{26} = 0.167473$. Further, the zeros of $\tilde{L}_{24}^{(-1.121695)}(z; 0.997)$ interlace with the non-common zeros of $\tilde{L}_{26}^{(-1.121695)}(z; 0.997)$ in $(0, \infty)$.

Example 2. When q = 0.94, n = 26 and $\delta = -1.92598$, then $c_{26} = 0.278236$ is a common zero of $\tilde{L}_{26}^{(-1.92598)}(z; 0.94)$ and $\tilde{L}_{26}^{(-1.92598+2)}(z; 0.94)$. Moreover, the positive zeros of $\tilde{L}_{26}^{(-1.92598)}(z; 0.94)$ interlace with the non-common zeros of $\tilde{L}_{26}^{(-1.92598+2)}(z; 0.94)$.

4 Bounds for the smallest zero of $ilde{L}_n^{(\delta)}(z;q)$

We derive the inner and outer bounds for the negative zero of $\tilde{L}_n^{(\delta)}(z;q)$. To find the bounds, we make use of the following contiguous relations:

$$z^{3}\tilde{L}_{n-2}^{(\delta+3)}(z;q) = \frac{(1-q^{\delta+n+1})}{(1-q^{n-1})} b_{n}\left(z-A_{n}\right)\tilde{L}_{n-1}^{(\delta)}(z;q) + \left(z-A_{n-\delta-2}\right)\tilde{L}_{n}^{(\delta)}(z;q)$$
(22)

and

$$z^{4}\tilde{L}_{n-2}^{(\delta+4)}(z;q) = \frac{A_{n-\delta-2} b_{n}((1-q^{n})+q(1-q^{\delta+n+1}))}{q^{n}(1-q^{\delta+1})} (z-B_{n}) \tilde{L}_{n-1}^{(\delta)}(z;q) + \left(z^{2} + \frac{(1+q)(1-q^{\delta+2})}{q^{\delta+n+1}}z - \frac{A_{n-\delta-2}(1-q^{\delta+3})}{q^{\delta+n+1}}\right) \tilde{L}_{n}^{(\delta)}(z;q), \quad (23)$$

where $A_n = \frac{(1-q^{\delta+1})(1-q^{\delta+2})}{q^{\delta+1}(1-q^{\delta+n+1})}$, $B_n = \frac{(1-q^{\delta+1})(1-q^{\delta+3})}{q^{\delta+1}((1-q^n)+q(1-q^{\delta+n+1}))}$ and b_n is given in Theorem 3.

Clearly, (22) and (23) are the application of [16, (4.14) and (4.15)] in (17).

Theorem 10. For $n \geq 2, n \in \mathbb{N}$, the negative zero $z_{1,n}$ of $\tilde{L}_n^{(\delta)}(z;q)$ satisfies

$$-c_{n-1} < B_n < z_{1,n} < A_n < 0, (24)$$

where A_n, B_n and c_n are respectively given in (22), (23) and in Theorem 4.

Proof. At $z = z_{1,n}$, (23) gives

$$z_{1,n}^{4}\tilde{L}_{n-2}^{(\delta+4)}(z_{1,n};q) = \frac{A_{n-\delta-2} b_n((1-q^n) + q(1-q^{\delta+n+1}))}{q^n(1-q^{\delta+1})} (z_{1,n} - B_n) \tilde{L}_{n-1}^{(\delta)}(z_{1,n};q).$$
(25)

For all n and δ , we have

$$\lim_{z \to -\infty} \tilde{L}_n^{(\delta)}(z;q) = \infty.$$
(26)

This implies that $(z_{1,n} - B_n)$ must be positive since all other factors in (25) are positive, that means

$$B_n < z_{1,n}.\tag{27}$$

Similarly, we have from (22) at $z = z_{1,n}$ that

$$z_{1,n}^{3}\tilde{L}_{n-2}^{(\delta+3)}(z_{1,n};q) = \frac{(1-q^{\delta+n+1})}{(1-q^{n-1})} b_n \left(z_{1,n} - A_n\right) \tilde{L}_{n-1}^{(\delta)}(z_{1,n};q).$$
(28)

Using (4) and (26) we observe that, the factor $(z_{1,n} - A_n)$ must be negative since the left-hand side of (28) is negative while the first and final factor on the right-hand side are positive, so that

$$z_{1,n} < A_n. \tag{29}$$

Now for $n \ge 2$, we see that

$$(1 - q^n) + q(1 - q^{n-2}) > 0,$$

which implies

$$(1-q^n) + q(1-q^{\delta+n+1}) > q^{n-1}(1-q^{\delta+3}).$$

Suitable interpretation of the above inequality leads to

$$-c_{n-1} < B_n. \tag{30}$$

Therefore, from (27), (29) and (30), we have the inequality (24).

Remark 3. By using the fact as in [15, (14.21.18)], when we take limit as $q \rightarrow 1$ in (24), it recovers the bounds, obtained by Driver and Muldoon in [10], for the negative zero of the quasi-orthogonal Laguerre polynomials of order 1.

Using *Mathematica 12*, we examine the inner and outer bounds for the smallest zero of $\tilde{L}_n^{(\delta)}(z;q)$ for different values of the parameters q, n and δ such that $\delta \in (-2, -1)$ which are listed in Table 1.

q	n	δ	B_n	$z_{1,n}$	A_n
0.23	2	-1.1	-0.11032	-0.110294	-0.10681
		-1.81	-0.505545	-0.448837	-0.20526
	7	-1.1	-0.104315	-0.104286	-0.100269
		-1.81	-0.46738	-0.40912	-0.169571
	12	-1.1	-0.104311	-0.104283	-0.100265
		-1.81	-0.467357	-0.409097	-0.169552
0.89	2	-1.1	-0.00598206	-0.00597785	-0.00580805
		-1.81	-0.0360942	-0.0294349	-0.0152326
	7	-1.1	-0.00219276	-0.00219076	-0.0020879
		-1.81	-0.0114906	-0.00913766	-0.0038382
	12	-1.1	-0.0016198	-0.00161831	-0.00153785
		-1.81	-0.0083227	-0.00661085	-0.00270734

Table 1: Inner and outer bounds of $z_{1,n}$ for different values of q, n and δ .

Acknowledgments

Present investigation is supported under OURIIP, Govt. of Odisha, India, Sanction Number - 1040/69/OSHEC.

References

- [1] C. Brezinski, K. Driver, M. Redivo-Zaglia, Quasi-orthogonality with applications to some families of classical orthogonal polynomials, Appl. Numer. Math. 48 (2004) 157–168.
- [2] C. Brezinski, K. Driver, M. Redivo-Zaglia, Zeros of quadratic quasi-orthogonal order 2 polynomials, Appl. Numer. Math. 135 (2019) 143–145.
- [3] T. S. Chihara, On quasi-orthogonal polynomials, Proc. Amer. Math. Soc. 8 (1957) 765–767.
- [4] D. Dickinson, On quasi-orthogonal polynomials, Proc. Amer. Math. Soc. 12 (1961) 185– 194.
- [5] A. Draux, On quasi-orthogonal polynomials, J. Aprox. Theory 62(1) (1990) 1–14.
- [6] K. Driver, A. Jooste, Interlacing of zeros of quasi-orthogonal Meixner polynomials, Quaest. Math. 40 (2017) 477–490.
- [7] K. Driver, K. Jordaan, Zeros of Jacobi polynomials $P_n^{(\alpha,\beta)}$, $-2 < \alpha, \beta < -1$, Numer. Algor. 79(4) (2018) 1075–1085.
- [8] K. Driver, K. Jordaan, Zeros of quasi-orthogonal Jacobi polynomials, SIGMA, 12, 042, 13 pages (2016).

- [9] K. Driver, M. E. Muldoon, Zeros of quasi-orthogonal ultraspherical polynomials, 27 (2016) 930–944.
- [10] K. Driver, M. E. Muldoon, Interlacing properties and bounds for the quasi-orthogonal Laguerre polynomials, Comput. Methods Funct. Theorey 15 (2015) 645—654.
- [11] L. Fejér, Mechanische Quadraturen mit Positiven Cotesschen Zahlen, Math. Z. 37 (1933) 287–309.
- [12] A. Jooste, K. Jordaan, F. Toókos, On the zeros of Meixner polynomials, Numer. Math. 124 (2013) 57–71.
- [13] H. Joulak, A contribution to quasi-orthogonal polynomials and associated polynomials, Appl. Numer. Math. 54(1) (2005) 65–78.
- [14] P.P. Kar, K. Jordaan, P. Gochhayat, M.K. Nangho, Quasi-orthogonality and zeros of some $_2\phi_2$ and $_3\phi_2$ polynomials, J. Comput. Appl. Math. (2020) 112910, https://doi.org/10.1016/j.cam.2020.112910
- [15] R. Koekoek, P. A. Lesky, R. F. Swarttouw, Hypergeometric Orthogonal Polynomials and Their *q*-Analogues, Springer Monographs in Mathematics, Springer Verlag, Berlin (2010).
- [16] D.S. Moak, The q-analogue of the Laguerre polynomials, J. Math. Anal. 81 (1981) 20–47.
- [17] M. Riesz, Sur le probleme des moments, III, Ark. Mat. Astron. Fys. 17(16) (1923) 1–52.
- [18] J.A. Shohat, On mechanical quadratures, in particular, with positive coefficients, Trans. Amer. Math. Soc. 42(3) (1937) 461–496.
- [19] G. Szegö, Orthogonal Polynomials, American Mathematical Society Colloquium Publications, Vol. XXIII, (1939).
- [20] D.D. Tcheutia, A.S. Jooste, W. Koepf, Quasi-Orthogonality of Some Hypergeometric and *q*-Hypergeometric Polynomials, SIGMA, 14, 051, 26 pages (2018).