

ACTADS

SOPHIE KRIZ

ABSTRACT. In this paper, I introduce a new generalization of the concept of an operad, further generalizing the concept of an *opetope* introduced by Baez and Dolan, who used this for the definition of their version of non-strict n -categories. Opetopes arise from iterating a certain construction on operads called the $+$ -construction, starting with monoids. The first step gives rise to plain operads, i.e. operads without symmetries. The permutation axiom in a symmetric operad, however, is an additional structure resulting from permutation of variables, independent of the structure of a monoid. Even though we can apply the $+$ -construction to symmetric operads, there is the possibility of introducing a completely different kind of permutations on the higher levels by again permuting variables without regard to the structure on the previous levels. Defining and investigating these structures is the main purpose of this paper. The structures obtained in this way is what I call n -actads. In n -actads with $n > 1$, the permutations on the different levels give rise to a certain special kind of n -fold category. I also explore the concept of iterated algebras over an n -actad (generalizing an algebra and module over an operad), and various types of iterated units. I give some examples of algebras over 2-actads, and show how they can be used to construct certain new interesting homotopy types of operads. I also discuss a connection between actads and ordinal notation.

CONTENTS

1. Introduction	3
2. Plain Bases and Actads	10
3. Actads and Bases	19
4. Algebras over an n -Actad	26
5. Modules of Algebras over n -Actads	29
6. Iterated Algebras	31
7. R -Unital Actads and Bases	37
8. Monads	42
9. Case study of a 2-actad algebra	46
10. Examples of New Operads	54
11. Ordinals	60
References	63

1. INTRODUCTION

Operads are a central concept of modern algebraic topology. The term was coined by J. P. May [27]. Operads are used as an approach to infinite loop space theory. They are a more streamlined alternative to the PROPs of Adams and MacLane, Lawvere theories, and the concepts of Boardman and Vogt. An operad \mathcal{C} consists of sets (or spaces) $\mathcal{C}(n)$ where $\mathcal{C}(n)$ is the set of all operations in n variables in a type of algebraic structure, which it describes. The basic operations of an operad correspond to substituting the results of operations (in different variables) to the input variables of another operation. One also typically includes a unit (identity) operation and commutativity in the broader sense, which means a symmetric group action by permuting variables. All this structure is included in May's original definition of an *operad* ([27], Definition 1.1, which also included the assumption that $\mathcal{C}(0) = *$). The concept without commutativity was defined by May ([27], Definition 3.12) as *non- Σ operads*. Some authors also speak of *plain operads*. In contrast, operads where permuting variables is allowed are sometimes called *symmetric operads*. Another variant is an *S-sorted operad* (or *multicategory*), which is defined in the same way as an operad, but with an "object set" S .

In the original context of infinite loop space theory, May [27] used the *little n -cube operad* which naturally acts on an n -fold loop space. By passing to the limit, an infinite loop space is shown to be an E_∞ -space, which means an algebra over an operad \mathcal{C} where each $\mathcal{C}(n)$ is a contractible space with free Σ_n -action. This is one way of capturing the notion of a commutative monoid "up to all possible homotopies."

In a completely different context, operads also later appeared in algebra, where instead of operations on a set or a space, we axiomatize multilinear operations on a vector space. In this fashion, for example, commutative and associative algebras, as well as Lie and Poisson algebras can be axiomatized. Ginzburg and Kapranov ([17]) discovered a striking phenomenon of *Koszul duality* of operads and their algebras, analogous to previously known concept of Koszul duality of algebras and modules [32].

Passing from vector spaces to their chain complexes, we can also talk about chain-level E_∞ -operads and their algebras. Hinich and Schechtman [21] noticed that the cochain complex of a topological space has the structure of an E_∞ -algebra (which they called a *May algebra*). Mandell [24] proved that for well-behaved spaces (for example simply connected with finitely generated homotopy groups), their p -completed homotopy

type can be recovered from their E_∞ cochain complex with coefficients in $\overline{\mathbb{F}}_p$.

In some sense, the ideas of the topological and algebraic contexts are combined in *multiplicative infinite loop space theory* (cf. [29, 13]), which generalizes certain structures of multilinear algebra to stable homotopy theory, i.e. to the context of spectra (see [2], Part III).

Baez and Dolan (see [3]) introduced the *+construction* (unrelated to Quillen's *+construction*), which is a procedure used to pass from monoids to plain operads. Essentially, for an S -sorted operad \mathcal{O} , there is an operad \mathcal{O}^+ sorted over the set of operations of \mathcal{O} , whose algebras are operads over \mathcal{O} . Applying this procedure to monoids defines plain operads, and iterating defines *opetopes*, which Baez and Dolan [3] used to axiomatize non-strict n -categories. Variants and generalizations of these concepts were introduced by C. Hermida, M. Makkai, and J. Power [18, 19, 20], E. Cheng [8, 9, 10], and T. Palm [30, 31]. Additional references include M. Zawadowski [38, 39, 40] and M. Fiore [15].

The *+construction* was further generalized to Cartesian monads by Leinster [23], and interpreted combinatorially by Kock, Joyal, Batanin, and Mascari [22], using the calculus of polynomial functors of Gambino and Hyland [16]. An even more conceptual interpretation in this direction was given by Szawiel and Zawadowski [34, 35, 36]. For another approach using syntactic methods, see Curien, Thanh, and Mimram [12]. Computer implementations of polytopes are given in [22, 14].

However, opetopes do not explain how symmetric operads arise from monoids. While the *+construction* can be applied to symmetric operads, there is an additional possibility of introducing permutations on each higher level by permuting variables without regard to the structures on the previous levels. The interplay of these permutations is the concept I explore in this paper. I call the resulting structures *actads*. Note that the possibility of permuting variables randomly does not fit within the picture of any reasonable concept of a higher category: it is a truly new element. Perhaps for that reason, actads have so far defied the conceptual approaches described above, and the only rigorous definition I was able to make starts from scratch.

As it turns out, the higher permutations must be handled quite carefully. We must specify a delicate order of variables on the previous levels which must be preserved for the higher permutations to work. Permutations on the different levels cannot just be mixed into an ordinary category, but must form a special kind of n -fold category. Because

of this, a particularly canonical model of the structure without permutations (which I call *plain actads*) must be introduced. Plain n -actads form a category equivalent to a version of n -opetopes ([3]), a combinatorial model of which was given in [22]. However, we need an ordered model which will be introduced from scratch.

In this paper, I index the actads so that monoids are 0-actads and operads are 1-actads. The operations of an actad are indexed by an n -base. The 0-base is just 1 point, the 1-base is the set of natural numbers. Elements of the 2-base are, roughly speaking, based trees. Drawing, in the place of each node of a tree, a triangle whose vertex is the node, and whose base contains the successor nodes, one can visualize a 3-base as a tree drawn in this fashion, where each triangle can be, recursively (but only finitely many times), subdivided into a tree drawn in a similar fashion. Remembering all these subdivisions is a key part of the structure of an element of the 3-base. For $n > 3$, n -bases are more difficult to visualize.

To explain how the induction procedure works in more detail, I find it easier to first only discuss associativity. I call the concept only encoding associativity a *plain actad*. Plain actads are not categories, but merely sets. I define a plain n -actad using an indexing set B_n called the *plain n -base*. For example, the elements of B_2 are *planar trees*, which means that the successors to each node are linearly ordered. In addition to the B_n 's, there are maps F_n , which map B_n into the set of non-empty finite sequences in B_{n-1} , and G_n , which simply maps B_n into B_{n-1} . Now, the F_n 's can be thought of as assigning to an element of B_n the “set of its entries.” For example, F_2 can be visualized as giving the degrees of internal nodes of a planar tree. In addition, G_n can be thought of as assigning to an element of B_n its “total arity”, which is an element of B_{n-1} . Finally, we can define a composition operation by taking an element $x \in B_n$ and replacing one of its “entries” (i.e. one of the terms z of $F_n(x)$) with some $y \in B_n$ such that $G_n(y) = z$. For example, we begin with $B_0 = \{*\}$ and $B_1 = \mathbb{N}$ (where we identify a natural number n with a planar tree with 1 triangle and n prongs). Composition of $n, m \in \mathbb{N} = B_1$ is then defined to always give $n + m - 1$ and can be visualized in the diagram (4), below. Note that we always have $F_1(n) = \underbrace{(*, \dots, *)}_n$ and $G_1(n) = *$ for every n , so all $n, m \in B_1$ are composable in all ways.

Now, a plain n -actad is a system $\mathcal{C}(z)$ indexed by $z \in B_n$. The structure of a plain n -actad is defined by composition operations in \mathcal{C} “fibered over B_n ”, i.e. where the indices are elements involved in

a composition in B_n . Finally, we can derive the associativity axioms from associativity properties of the composition in B_n .

To advance the induction, I define B_{n+1} as the free plain n -actad on B_n . In other words, we no longer “execute” the composition operations on B_n , but simply keep track of which ones were made, subject to the associativity axiom. In this fashion, we can inductively define the B_n ’s, and thus we can define plain n -actads. I treat plain n -actads below in Section 2.

Just as in the case of operads, capturing a commutativity axiom complicates the structure substantially. In the case of an operad \mathcal{C} , a commutativity property is expressed by introducing a Σ_n -action on $\mathcal{C}(n)$. Then this turns B_1 into a groupoid \mathcal{B}_1 , making Σ_n the automorphism group of $n \in \mathbb{N}$. From a philosophical point of view, n really stands for the word

$$(1) \quad a^n = a \cdots a$$

in the free monoid on one element a , and the permutations interchange (independently of the monoid structure) the n copies of a . In this paper, I develop the corresponding additional structure \mathcal{B}_2 , and eventually \mathcal{B}_n . Recall that the elements of B_2 are planar trees. \mathcal{B}_2 should incorporate the “1-permutations” induced from \mathcal{B}_1 by (1), which are isomorphisms of trees. Notice that a 1-permutation can change the tree, so it may no longer be an automorphism, although it is always an isomorphism (so 1-permutations form a groupoid). For example, see the picture of a 1-permutation in \mathcal{B}_2 in (24). But now it seems that any natural generalization should introduce permutations of nodes of the tree of the same arity (call them 2-permutations), regardless of the structure of the tree (just as in (1), the permutations are completely independent to the product operation of the free monoid). For example, see (25).

We need to answer the question of how the 1-permutations and 2-permutations interact. Putting them both together into a single category seems to fail. In other words, a random permutation of nodes of the same arity in a tree does not seem to induce, in any meaningful way, a permutation on leaves. Nor does there really seem to be a natural structure of a 2-category: Both 1-permutations and 2-permutations act on objects. Thus, we are led to a bicategory structure (see [5]), where the $\{1, 2\}$ -morphisms are permutations of nodes of equal arity, on which we have prescribed the same permutation of successors. This, fortunately, is a rather special type of bicategory. In addition to the $\{1\}$ -morphisms and $\{2\}$ -morphisms (we may in the future omit the braces to simplify notation) forming groupoids, every 1-morphism

and 2-morphism with the same source complete to a unique $\{1, 2\}$ -morphism. In this paper, I call this type of bicategory *cube-like*, and generalize it to all n . Then, \mathcal{B}_n is, in particular, a cube-like n -fold category. Along with the appropriate axioms expressing compatibility with composition, this is my implementation of the commutativity axiom. Cube-like n -fold categories fibered over \mathcal{B}_n with composition satisfying the properties of \mathcal{B}_n 's composition are what I call *n-actads*. I treat this in Section 3.

The question of units is another issue. There is, of course, a natural concept of unit that can be considered a part of associativity, which generates two unit axioms. However, there is more to the story. Even in operads (which, recall, are 1-actads), we encounter another concept, namely that of a *based operad*, which also contains a manifestation of the unit of a monoid (which is a 0-actad). More precisely, a based operad $\mathcal{C} = (\mathcal{C}(n))$ is indexed by $n \in \mathbb{N}_0$, and we are given a base point $*$ $\in \mathcal{C}(0)$. Based operads play a crucial role in infinite loop space theory (for example, in the Approximation Theorem [27]).

Similarly, for n -actads, beside the “ordinary” unit notion there is a notion which includes carry-overs from k -actad units for $k < n$. I call this “recursive” unit axiom *R-unitality*. It is interesting to note that in the case of n -actads, the commutativity axiom becomes substantially richer in the *R-unital* case. This is because for $n \geq 3$, in the non-unital case, an element of an n -actad can be only non-trivially composed with elements of lower arity, with which it cannot be permuted by an $\{n\}$ -morphism. In the *R-unital* context, this is not necessarily the case. I treat the *R-unital* case in Section 7 below.

The original application of operads comes from their algebras. In addition, the notion of algebras prompted the notion of a module over an operad and its algebras, which was invented by May, and is treated, for example, in [4]. We would like to generalize these notions to n -actads. Indeed, there are directly analogous notions for n -actads, and I treat these concepts in Section 4. An important example is the *commutative* (in the narrower sense) $(n+1)$ -actad \mathcal{B}_{n+1} whose algebras can be thought of as strictly commutative n -actads. By “freeing up” the $(n+1)$ -morphisms, we are able to obtain the *associative* $(n+1)$ -actad \mathcal{A}_{n+1} , whose algebras are precisely n -actads.

For $n \geq 2$, the immediate thought is that this idea should be “iterated.” However, implementing that idea is not obvious. Before we can iterate, we must for example answer the question of what happens to n in the iterated algebras. The answer to these questions are given in Section 6, and are summarized as follows:

For an n -actad \mathcal{C} , it turns out that there exists a notion of an *iterated \mathcal{C} - I -algebra* for every subset $I \subseteq \{0, \dots, n\}$, which consists of $|I|$ cube-like $(n-1)$ -fold categories. An axiomatization of these notions is made using multisorted algebras (a definition of which can be found in [6]). The concept essentially allows one to label the entries of a tree with the elements of I (for example, see (36)). In the case of $I = \{0\}$ and $I = \{0, n\}$, the concept gives the notions of algebras and modules, respectively. I treat this in Section 6.

What are some interesting examples and applications of n -actads? So far, we have been working in the ground category of sets when discussing actads and their algebras. One can of course also work with spaces or simplicial sets without any difficulties (as is done with operads in [27]). This is where one may expect the most interesting topological examples to reside. At present, finding such examples is restricted by the fact that, for $n \geq 2$, the structures are unfamiliar, but we do present some examples of the new structure here. One interesting example is the E_∞ - n -actad $E\mathcal{A}_n$, which is the Čech resolution of the associative n -actad \mathcal{A}_n . Note that since we have a canonical map of n -actads

$$(2) \quad \mathcal{A}_n \rightarrow E\mathcal{A}_n,$$

$E\mathcal{A}_n$ -algebras are, in particular, $(n-1)$ -actads.

For $n = 2$, I give some concrete examples related to this case. I describe the free $E\mathcal{A}_2$ -algebra $A_{E\mathcal{A}_2}(X_2)$ on the groupoid X_2 fibered over \mathcal{B}_1 where $X_2(2)$ is the groupoid given by Σ_2 acting on itself freely and $X_2(n)$ is empty over $n \neq 2$. I also give a sufficient condition for the classifying space of a groupoid to be algebras over the operad $A_{E\mathcal{A}_2}(X_2)$. In some sense, this is an analogue of the result of May [28] that the classifying space of a permutative category is an algebra over the Barratt-Eccles E_∞ -operad, which is $E\mathcal{A}_1$.

In Section 9, I actually construct $E\mathcal{A}_2$ -algebras which are new operads (in the sense that they, as far as I know, are not equivalent to operads which have been previously studied). For a spectrum E (see [2] for an introduction the concept of spectra in algebraic topology) with a map $E \rightarrow H\mathbb{Z}$ (the Eilenberg-Mac Lane spectrum), we can obtain a map of E_∞ -spaces $E_0 \rightarrow \mathbb{Z}$. I show that this structure (also when E_0 is replaced with any E_∞ -space), in particular, leads to a structure of an $E\mathcal{A}_2$ -algebra, and show that in certain cases, the resulting example is non-trivial even as an operad, (via (2)), in the sense that an E_∞ -operad does not map into it.

In fact, algebras over a based version $\bar{\Xi}_{E_0}$ of this operad can be characterized as follows: For a map of spectra $E \rightarrow F \rightarrow H\mathbb{Z}$, let $\psi : F_0 \rightarrow \mathbb{Z}$ be the corresponding map of E_∞ -spaces. Then $\bar{\Xi}_{E_0}$ -algebras

are canonically equivalent to spaces of the form $\psi^{-1}(-1)$. Note that $\psi^{-1}(-1)$ is homotopy equivalent to $\psi^{-1}(0)$, which is an infinite loop space, but the homotopy equivalence is non-canonical when the map of spectra $E \rightarrow H\mathbb{Z}$ does not split.

While this is only one example of a homotopical structure obtained from the 2-morphisms of actads, its non-triviality can be thought of as higher-categorical in nature (since it comes from a lack of canonicity). Thus, our example may provide a clue to the nature of the kinds of applications the new structure of an n -actad may have.

There is yet an entirely different aspect of the story of plain actads. Kock, Joyal, Batanin, Mascari ([22]) discuss a machine interpretation and also a “fixed point” of the $+$ -construction. It is possible to pursue this further. Our particular model of plain actads suggests a close connection with the Veblen hierarchy of ordinal notation. From this point of view, it is possible to iterate the opetope construction transfinitely and we show that the level of complexity of the first fixed point of that construction is the Feferman-Schütte ordinal Γ . More specifically, recalling the fact that the elements of B_2 can be identified with planar trees, one is reminded of the Veblen hierarchy of ordinals, where, letting φ_β denote the Veblen functions (see [37]), $\varphi_1(\alpha) = \omega^\alpha$ and $\varphi_\beta(\alpha)$ is the α 'th fixed point of $\varphi_{\beta'}$ for $\beta' < \beta$. (This is shifted by 1 from the usual convention, which is more natural for my point of view. The shift disappears for $\beta \geq \omega$.) The first fixed point of this hierarchy (the first ordinal with $\varphi_\Gamma(0) = \Gamma$) is the Feferman-Schütte ordinal Γ [37]. It is particularly well known that trees can be used for a notation of ordinals below $\varepsilon = \varphi_2(0)$ which suggests a connection with plain actads. In Section 11, I indeed construct an onto function

$$(3) \quad B_n \rightarrow \varphi_n(0).$$

This prompts the idea of generalizing B_n to the case when n is an ordinal, which is carried out readily. Thus, we can also use B_α , for $\alpha < \Gamma$, as notation for countable ordinals below Γ . The map I constructed is not a bijection (for example even for trees, we have $1 + \omega = \omega$). This could be remedied by taking a proper subset of B_n , but we do not undertake this task in the present paper. Since actads are algebraic structures, even the map (3) adds to visualization of ordinals between ε and Γ .

Acknowledgment: I am grateful to the referees for many helpful suggestions for improving the presentation of this paper.

2. PLAIN BASES AND ACTADS

In this section, we will introduce plain actads. By the results of [22], (following the concept of Gambino and Hyland [16]), the category of plain n -actads is equivalent to n -opetopes without units, (a unital version of this statement also holds, see Section 7). However, plain actads give a canonical ordered model of the category which we will need when introducing actads. As we described in the introduction, in the process of defining plain n -actads, we must also define the plain n -base and the maps F_n 's and the G_n 's. We will define a set B_n , also called the *plain n -base* and plain n -actads inductively. Plain n -actads will be universal algebras sorted on B_n .

In general, (see [6]), for a set S , an S -sorted universal algebra is defined to be the collection of the data of a set X with a map $X \rightarrow S$ together with a set O_n for each $n \geq 0$ of n -ary operations (specific to the type of algebra we are considering) together with a map $O_n \rightarrow S^{n+1}$ (specifying which operations are allowed with given sources and target), that is required to satisfy certain prescribed identities, (also specific to the kind of algebra considered) where in compositions, the output of an operation is plugged into an input over the same $s \in S$ (i.e. compositions are

$$O_n \times_{S^n} (O_{k_1} \times \cdots \times O_{k_n}) \rightarrow O_{k_1 + \cdots + k_n}.$$

Homomorphisms of a given kind of S -sorted algebras are maps over S (i.e. commuting with the given maps to S) which are compatible with the operations. For example, a group acting on a set forms a $\{0, 1\}$ -sorted universal algebra $\varphi : X \rightarrow \{0, 1\}$ where the group is $\varphi^{-1}(0)$ and the set is $\varphi^{-1}(1)$. S -sorted algebras are equivalent to finitary monads in the category (Set/S) of *Sets* over S . (The proof follows from the methods of Chapter 3 of [1].) This means a monad M such that for any $X \rightarrow S$, $M(X \rightarrow S)$ is the colimit in Set/S of $M(K \rightarrow S)$ with $K \subseteq X$ finite.

For a map $f : S \rightarrow T$, a T -sorted algebra specifies an S -sorted algebra by allowing those operations and identifies which are allowed after composing with f . For a finitary monad M in Set/T , we can describe the monad f^*M in Set/S as RML where $L : Set/S \rightarrow Set/T$ is the forgetful functor and $R : Set/T \rightarrow Set/S$ is its right adjoint, given by $R(X) = X \times_T S$. In particular, an unsorted universal algebra can be made S -sorted by pulling back along the map $S \rightarrow *$. Thus S -sorted operads (or multicategories) are defined. Of course, in multicategories, we often consider multifunctors, which are more general

morphisms over a map of sets $S_1 \rightarrow S_2$ preserving the operations in the obvious sense. In general, for a S -sorted universal algebra pulled back via a map $f : S \rightarrow T$, we have a notion of a morphism over any map $S_1 \rightarrow S_2$ over T .

Plain n -actads are universal algebras sorted in B_n , this means that a plain n -actad is a set X over B_n (i.e. a map $X \rightarrow B_n$) with certain operations (which, as it will turn out, are binary) applicable to elements that map to certain elements of B_n , specific to the operation, and certain identities among iterated operations. For $n = 0$, the plain 0-base is just a point,

$$B_0 = \{*\}.$$

To describe the operations in more detail, we must also say more about the maps. For $n \geq 1$, additional structure on B_n is needed, which we will define inductively. Write

$$T_{B_{n-1}} = \{(a_1, \dots, a_k) \mid a_i \in B_{n-1}, k \geq 1\}.$$

We have maps

$$F_n : B_n \rightarrow T_{B_{n-1}},$$

(let m_x be the length of the sequence $F_n(x)$ for $x \in B_n$, and write

$$F_n(x) = (f_1(x), \dots, f_{m_x}(x)),$$

and

$$G_n : B_n \rightarrow B_{n-1}.$$

Suppose B_{n-1} and $(n-1)$ -actads have already been defined. We then have the following structure: We have an operation

$$x \circ_i^n y,$$

for $x, y \in B_n$, $1 \leq i \leq m_x$, and $G_n(y) = f_i(x)$, such that

$$m_{x \circ_i^n y} = m_x + m_y - 1.$$

To illustrate what we are after, before making exact definitions, let us consider what these structures look like for $n = 0, 1, 2$. For $n = 0$, only consists of a single B_0 point, so one can only compose an element with itself leading to picture that is not very interesting:

•


However, for $n = 1$ and $n = 2$, the pictures are much more revealing. I will use the following pictures to express 1- and 2-compositions. For $n = 1$, B_1 is the set of all natural numbers, and we see the standard


composition picture for planar trees (identifying a natural number n with the planar tree with n prongs). Here, we are replacing one of the “prongs” of a planar tree at the end of one of the trees with the “stem” of a different tree. The “prongs” are visualized as ordered “left to right.”

(4) 


It is important to note that, for operads, any two trees can be composed. This is because for every tree x , we have $G_1(x) = *$ and $F_1(x) = (*, \dots, *)$. So, for every $1 \leq i \leq m_x$, $f_i(x) = G_1(x) = *$. So every two trees, every possible f_i will agree with G_1 .

For $n = 2$, though, the composition is quite different. In the picture, I express 2-composition by putting a tree with the same arity inside one of the “triangles.” Letting

$x =$ 

$y =$ 

(where the numbers denote the ordering of the entries of x and y), we can compose the two trees via $x \circ_3^2 y$ (see the below picture)



The induction data of B_n also includes increasing functions

$$\varphi_n^{(x,i,y)} : \{1, \dots, m_x\} \setminus \{i\} \rightarrow \{1, \dots, m_x + m_y - 1\}$$

$$\psi_n^{(x,i,y)} : \{1, \dots, m_y\} \rightarrow \{1, \dots, m_x + m_y - 1\}$$

expressing how the elements of the two objects of B_n involved in the composition are shuffled in the resulting object. The purpose of these functions is to formulate an analogue of the associativity axiom. For example, we see that for $n = 1$, where the elements are prongs, the prongs of y “stay together” in the shuffle even though the indexing is shifted. For $n = 2$, instead of prongs, the elements are triangles. They are also ordered “left to right” (and in case of a tie, “top to bottom”), but we see that in this ordering, both shuffles φ and ψ are non-trivial, i.e. the “triangles” of y do not “stay together” after the composition.

The associativity property for composition we require states that for $1 \leq i < j \leq m_x$, $G_n(z) = f_j(x)$,

$$(5) \quad (x \circ_i^n y) \circ_{\varphi_n^{(x,i,y)}(j)}^n z = (x \circ_j^n z) \circ_i^n y.$$

For $n = 1$, we have

$$\varphi_1^{(x,i,y)}(j) = j + m_y - 1 \text{ for } i < j \leq m_x$$

$$\psi_1^{(x,i,y)}(j) = j + i - 1 \text{ for } 1 \leq j \leq m_y.$$

To make a rigorous definition of plain n -actads, we need to formulate a few more properties of composition and these functions. For $j < i$, we must have

$$\varphi_n^{(x,i,y)}(j) = j.$$

(We shall sometimes omit the superscript n when it is clear). We will also have, for $i < j < k$ and any n , the additional axioms

$$(6) \quad \varphi_n^{(x \circ_j^n z, i, y)}(\varphi_n^{(x, j, z)}(k)) = \varphi_n^{(x \circ_i^n y, \varphi_n^{(x, i, y)}(j), z)}(\varphi_n^{(x, i, y)}(k)),$$

$$(7) \quad \varphi_n^{(x, i, y)}(j) = \varphi_n^{(x \circ_k^n t, i, y)}(j).$$

Suppose we have defined B_n (and all associated structure).

Definition 1. A plain n -actad has the data of a set \mathcal{C} with a map

$$\mathcal{C} \rightarrow B_n,$$

and, writing $\mathcal{C}(x)$ for the inverse image of and element $x \in B_n$, composition operators

$$\gamma_{n,i} : \mathcal{C}(x) \times \mathcal{C}(y) \rightarrow \mathcal{C}(x \circ_i^n y)$$

for $x, y \in B_n$, with $1 \leq i \leq m_x$, $G_n(y) = f_i(y)$. We require that these composition operators satisfy the condition that, for $1 \leq i < j \leq m_x$, $z \in B_n$, $G_n(z) = f_j(x)$, the following diagram commutes:

$$(8) \quad \begin{array}{ccc} \mathcal{C}(x) \times \mathcal{C}(y) \times \mathcal{C}(z) & \xrightarrow{\gamma_{n,i} \times Id} & \mathcal{C}(x \circ_i^n y) \times \mathcal{C}(z) \\ \downarrow T & & \downarrow \gamma_{n, \varphi_n^{(x, i, y)}(j)} \\ \mathcal{C}(x) \times \mathcal{C}(z) \times \mathcal{C}(y) & & \mathcal{C}((x \circ_i^n y) \circ_{\varphi_n^{(x, i, y)}(j)}^n z) \\ \downarrow \gamma_{n,j} \times Id & & \downarrow = \\ \mathcal{C}(x \circ_j^n z) \times \mathcal{C}(y) & \xrightarrow{\gamma_{n,i}} & \mathcal{C}((x \circ_j^n z) \circ_i^n y), \end{array}$$

where $T : \mathcal{C}(x) \times \mathcal{C}(y) \times \mathcal{C}(z) \rightarrow \mathcal{C}(x) \times \mathcal{C}(z) \times \mathcal{C}(y)$ is the permutation switching the coordinates of $\mathcal{C}(z)$ and $\mathcal{C}(y)$. (We shall also sometimes write γ_i instead of $\gamma_{n,i}$ when the value of n is clear.)

We shall now inductively define B_n . For $n = 1$, let $B_1 = \mathbb{N}$. In addition, define

$$F_1 : B_1 \rightarrow \{(a_1, \dots, a_k) \mid a_i \in B_0 = \{*\}\},$$

$$G_1 : B_1 \rightarrow B_0$$

by $F_1(n) = \underbrace{(*, *, \dots, *)}_{n \text{ times}}$, $G_1(n) = \{*\}$. Then, for $x, y \in B_1$, let

$$x \circ_i^1 y = x + y - 1.$$

Note again that this works for every i because the coordinates of $F_1(x)$ are all $*$, and $G_1(y) = *$ for every y .

Suppose we have B_n and all the maps associated with it. By Definition 1 above, we have the notion of plain n -actads. Given this, define

$$\begin{aligned} B_{n+1} &= \{((x_1, x_2, \dots, x_k), (i_1, i_2, \dots, i_{k-1})) \mid k \geq 1, x_\ell \in B_n, \\ &G_n(x_j) = f_{i_{j-1}}((\dots (x_1 \circ_{i_1}^n x_2) \circ_{i_2}^n x_3 \dots) \circ_{i_{j-2}}^n x_{j-1}), \\ &1 \leq i_j \leq m_{(\dots (x_1 \circ_{i_1}^n x_2) \circ_{i_2}^n x_3 \dots) \circ_{i_{j-1}}^n x_j} = m_{x_1} + \dots + m_{x_j} - (j - 1), \\ &i_1 \leq i_2 \leq \dots \leq i_{k-1}\}. \end{aligned}$$

In addition, define

$$\begin{aligned} G_{n+1}((x_1, x_2, \dots, x_k), (i_1, i_2, \dots, i_{k-1})) &= \\ &= (\dots ((x_1 \circ_{i_1}^n x_2) \circ_{i_2}^n x_3) \dots) \circ_{i_{k-1}}^n x_k, \\ F_{n+1}((x_1, x_2, \dots, x_k), (i_1, i_2, \dots, i_{k-1})) &= \\ &= (x_1, x_2, \dots, x_k), \end{aligned}$$

and

$$m_{((x_1, x_2, \dots, x_k), (i_1, i_2, \dots, i_{k-1}))} = k.$$

Then we have the following

Lemma 2. B_{n+1} is the free plain n -actad on $Id : B_n \rightarrow B_n$, where

$$\begin{aligned} &((x_1, x_2, \dots, x_k), (i_1, i_2, \dots, i_{k-1})) = \\ &\gamma_{n, i_{k-1}}(\gamma_{n, i_{k-2}}(\dots \gamma_{n, i_2}(\gamma_{n, i_1}(x_1, x_2), x_3), x_4) \dots), x_k). \end{aligned}$$

Proof. The free plain n -actad on B_n is

$$\begin{aligned} T_n &= \{\gamma_{n, i_1}(\gamma_{n, i_2}(\dots \gamma_{n, i_k}(x, y_k), \dots, y_2), y_1) \mid i_k, \dots, i_1 \in \mathbb{N}, \\ &x, y_1, \dots, y_k \in B_n \text{ composable}\} / \sim \end{aligned}$$

where \sim is the smallest equivalence relation containing compositions of the γ 's differing only in one pair of consecutive γ 's, replaced by another according to Diagram (8). We have, by definition,

$$\begin{aligned} B_{n+1} &= \{\gamma_{n, j_1}(\gamma_{n, j_2}(\dots \gamma_{n, j_k}(x, x_k), \dots, x_2), x_1) \mid j_k \leq \dots \leq j_1, \\ &x, x_1, \dots, x_k \in B_n \text{ composable}\}. \end{aligned}$$

We want

$$B_{n+1} = T_n.$$

We have a canonical map

$$(9) \quad B_{n+1} \rightarrow T_n$$

Suppose $i < j$. Then we have, by Diagram (8),

$$\gamma_i(\gamma_j(x, z), y) = \gamma_{\varphi_n^{(x, i, y)}(j)}(\gamma_i(x, y), z).$$

and $\varphi_n^{(x, i, y)}(j) \geq j > i$. So, for every

$$(10) \quad \gamma_{n, i_1}(\gamma_{n, i_2}(\cdots \gamma_{n, i_k}(x, y_k), \dots, y_2), y_1) \in T_n,$$

we can use $\varphi_n^{(x, i_\ell, y)}(i_{\ell+1})$ to ‘switch’ the order of i_ℓ and $i_{\ell'}$ whenever $i_\ell < i_{\ell+1}$. So, there exist $j_1 \geq \cdots \geq j_k$ such that

$$\begin{aligned} & \gamma_{n, i_1}(\gamma_{n, i_2}(\cdots \gamma_{n, i_k}(x, y_k), \dots, y_2), y_1) = \\ & = \gamma_{n, j_1}(\gamma_{n, j_2}(\cdots \gamma_{n, j_k}(x, y_{\sigma(k)}), \dots, y_{\sigma(2)}), y_{\sigma(1)}) \end{aligned}$$

where σ is a suitable permutation. (Though we note that it is not necessarily true that $\{i_1, \dots, i_k\} = \{j_1, \dots, j_k\}$. This gives a left inverse of (9), so (9) is onto. However, then we need to know that we get the same answer regardless of the order of ‘switches’. In proving this, the axioms (6), (7) will come into play.

Suppose $i < j < k$. Performing switches in two different ways, we get

$$\begin{aligned} & \gamma_i(\gamma_j(\gamma_k(x, t), z), y) = \gamma_{\varphi_n^{(\gamma_k(x, t), i, y)}(j)}(\gamma_i(\gamma_k(x, t))) = \\ & = \gamma_{\varphi_n^{(\gamma_k(x, t), i, y)}(j)}(\gamma_{\varphi_n^{(x, i, y)}(k)}(\gamma_i(x, y), t), z) = \\ & \gamma_{\varphi_n^{(\gamma_i(x, y), \varphi_n^{(\gamma_k(x, t), i, y)}(j), z)}(\varphi_n^{(x, i, y)}(k))}(\gamma_{\varphi_n^{(\gamma_k(x, t), i, y)}(j)}(\gamma_i(x, y), z), t) \end{aligned}$$

and

$$\begin{aligned} & \gamma_i(\gamma_j(\gamma_k(x, t), z), y) = \gamma_i(\gamma_{\varphi_n^{(x, j, z)}(k)}(\gamma_j(x, z), t), y) = \\ & = \gamma_{\varphi_n^{(\gamma_j(x, z), i, y)}(\varphi_n^{(x, j, z)}(k))}(\gamma_i(\gamma_j(x, z), y), t) = \\ & = \gamma_{\varphi_n^{(\gamma_j(x, z), i, y)}(\varphi_n^{(x, j, z)}(k))}(\gamma_{\varphi_n^{(x, i, y)}(j)}(\gamma_i(x, y), z), t). \end{aligned}$$

Thus, we must have

$$(11) \quad \varphi_n^{(\gamma_j(x, z), i, y)}(\varphi_n^{(x, j, z)}(k)) = \varphi_n^{(\gamma_i(x, y), \varphi_n^{(\gamma_k(x, t), i, y)}(j), z)}(\varphi_n^{(x, i, y)}(k)),$$

$$(12) \quad \varphi_n^{(x, i, y)}(j) = \varphi_n^{(\gamma_k(x, t), i, y)}(j),$$

which are (6) and (7). Conversely, (6) and (7) guarantee that two switches have the same result, i.e.

$$(13) \quad \varphi_n^{(\gamma_j(x, z), i, y)}(\varphi_n^{(x, j, y)}(k)) = \varphi_n^{(\gamma_i(x, y), \varphi_n^{(x, i, y)}(j), z)}(\varphi_n^{(x, i, y)}(k)).$$

Now, we will use induction to prove that we always get the same result regardless of the order we perform switches of consecutive pairs $\gamma_i \gamma_j$, for $i < j$. Let

$$i = \min\{i_1, \dots, i_k\},$$

where i_1, \dots, i_k are as in (10). If i occurs only once in the sequence (i_1, \dots, i_k) , then let n_i be the distance from i to the end of the sequence. If there exist $\ell_1 < \dots < \ell_m \in \{1, \dots, k\}$ such that

$$i = i_{\ell_1}, \dots, i_{\ell_m},$$

then let n_i be the distance from i_{ℓ_m} to the end of the sequence.

If i is at the end of the sequence (i.e. $i = i_k$ and $n_i = 0$), then repeat this process for (i_1, \dots, i_{k-1}) . If the smallest i in every sequence is at the end, (i_1, \dots, i_k) is already in order. So, then we are done.

Suppose we know that the result is independent of the order we take the switches if $n_i = p$. Then suppose $n_i = p + 1$.

The strategy is to show that without loss of generality, all the possible switches from $j < k$ to $j > k$ to the right of i can be done first before i is moved to the right. Then the order of the switches to the right of i does not matter by the induction hypothesis.

Consider the position when the last switch to the right of i before i is moved to the right has been performed. Assume first that the situation immediately to the right of i is

$$(14) \quad i < j > k.$$

Thus, the next position will be

$$(15) \quad j' > i < k.$$

Now if any more switches to the right of i happen before i is moved again, then equivalently, those switches could have been made first, and then the move from (14) to (15).

Thus, either all the possible switches to the right of i have been performed before (14) was reached (which is what we were trying to assume), or the position right before i was moved to the right was

$$(16) \quad i < j < k.$$

Now i is moved to the right producing

$$\bar{j} > i < k.$$

By the induction hypothesis, the order of the remaining moves does not matter, so we may as well continue to

$$\bar{j} < \bar{k} > i,$$

$$\bar{k} > \bar{j} > i.$$

But by axioms (6) and (7), those are equivalent to

$$i < k' > j$$

$$(17) \quad k'' > i < j$$

$$k'' > j'' > i$$

In the sequence (17), one more move was executed to the right of i before i was moved.

□

Note that this proof is analogous to the proof of the presentation of Σ_n by the ‘‘Yang-Baxter relations’’ $a_i a_{i+1} a_i = a_{i+1} a_i a_{i+1}$ (along with $a_i a_j = a_j a_i$ for $j > i + 1$) where $a_i = (i, i + 1)$ in the cycle notation.

Then, the functions $\varphi_{n+1}^{(x,j,y)}$, $\psi_{n+1}^{(x,j,y)}$ for

$$\begin{aligned} x &= ((x_1, x_2, \dots, x_k), (i_1, i_2, \dots, i_{k-1})) \\ y &= ((y_1, y_2, \dots, y_\ell), (\iota_1, \iota_2, \dots, \iota_{\ell-1})). \end{aligned}$$

are determined as follows:

For a plain n -actad \mathcal{C} , and for $\zeta_i \in \mathcal{C}(x_i)$, $\zeta_\iota \in \mathcal{C}(y_\iota)$, we have

$$\begin{aligned} &\gamma_{n,i_{k-1}}(\dots \gamma_{n,i_j}(\gamma_{n,i_{j-1}}(\dots \gamma_{n,i_2}(\gamma_{n,i_1}(\zeta_1, \zeta_2), \zeta_3) \dots, \zeta_{j-1}), \\ &\quad \gamma_{n,\iota_{\ell-1}}(\dots \gamma_{n,\iota_2}(\gamma_{n,\iota_1}(\zeta_1, \zeta_2), \zeta_3), \dots, \zeta_\ell), \dots x_k) = \\ &\quad \gamma_{n,\lambda_{k+\ell-1}}(\dots \gamma_{n,\lambda_2}(\gamma_{n,\lambda_1}(\mu_1, \mu_2), \mu_3) \dots, \mu_{\lambda_{k+\ell-1}}), \end{aligned}$$

with

$$\begin{aligned} \lambda_1 &\leq \dots \leq \lambda_{k+\ell-1} \\ \zeta_s &= \mu_{\varphi_{n+1}^{(x,j,y)}(s)} \\ \zeta_t &= \mu_{\psi_{n+1}^{(x,j,y)}(t)}, \end{aligned}$$

for $s \in \{1, \dots, k\}$, $t \in \{1, \dots, \ell\}$. Finally, also define

$$\begin{aligned} &((x_1, \dots, x_k), (i_1, \dots, i_{k-1})) \circ_j^{n+1} ((y_1, \dots, y_\ell), (\iota_1, \dots, \iota_{\ell-1})) = \\ (18) \quad &= \gamma_{n,i_{k-1}}(\dots \gamma_{n,i_j}(\gamma_{n,i_{j-1}}(((x_1, \dots, x_k), (i_1, \dots, i_{k-1})), \\ &\quad ((y_1, \dots, y_\ell), (\iota_1, \dots, \iota_{\ell-1}))), x_{j+1}), \dots, x_k). \end{aligned}$$

Axioms (5) and (6) then follow from the fact that one can similarly define, directly analogously, a composition of $((x_1, \dots, x_k), (i_1, \dots, i_{k-1}))$ with two appropriate elements

$$((y_1, \dots, y_{\ell_1}), (\iota_1, \dots, \iota_{\ell_1-1}))$$

and

$$((z_1, \dots, z_{\ell_2}), (\kappa_1, \dots, \kappa_{\ell_2-1}))$$

at $1 \leq j_1 < j_2 \leq m_x$, and observe that it is equal, by definition, to the compositions in the two different orders, and similarly for the shuffles.

Axiom (7) follows from the fact that, by Lemma 2, composing at a j , as in (18) does not affect the shuffles of compositions at $j' < j$.

3. ACTADS AND BASES

Using this, we will define the n -base \mathcal{B}_n , and then n -actads. Unlike the B_n 's, the \mathcal{B}_n 's are not sets. \mathcal{B}_n and n -actads are n -fold categories. (Recall, these are more general than n -categories. For example, $n = 2$ gives the notion of bi-categories, see [5], which strictly contains the notion of 2-categories.) One can define an n -fold category inductively as a category internal in $(n-1)$ -fold categories (i.e. where both objects and morphisms are $(n-1)$ -fold categories, and the maps involved in the axioms of the category are morphisms of $(n-1)$ -fold categories, i.e. $(n-1)$ -fold functors). We start the induction with 1-fold categories. Note that n -fold categories still have pullbacks.

This means that it involves sets of “ S -morphisms” for S any subset of $\{1, \dots, n\}$. In particular, “ \emptyset -morphisms” are actually the objects, which for \mathcal{B}_n are the plain n -bases, and for n -actads, they are the plain n -actads. To discuss this further, let us introduce some notation:

Suppose \mathcal{C} is an n -fold category and $S \subseteq \{1, \dots, n\}$. For

$$(19) \quad \mathbb{I} : S \rightarrow \{0, 1\},$$

we have the source-target map

$$(20) \quad \mathcal{S}_{\mathbb{I}} : S - \text{Mor}(\mathcal{C}) \rightarrow (\emptyset - \text{Mor}(\mathcal{C})) =: \text{Obj}(\mathcal{C}).$$

(Then 0 stands for source and 1 stands for target.) In addition, let $\underline{0} : S \rightarrow \{0, 1\}$ be the constant 0 function.

More generally, for $T \subseteq S$, $\mathbb{I} : S \setminus T \rightarrow \{0, 1\}$, we have a source-target map

$$\mathcal{S}_{\mathbb{I}}^T : S - \text{Mor}(\mathcal{C}) \rightarrow T - \text{Mor}(\mathcal{C}).$$

Call an n -fold category \mathcal{C} *cube-like* if for all $S \subseteq \{1, \dots, n\}$, for all $x \in \text{Obj}(\mathcal{C})$, and

$$\phi_i \in \{i\} - \text{Mor}(\mathcal{C})$$

(for any $i \in S$), such that $\mathcal{S}_{\underline{0}}(\phi_i) = x$, there exists a unique $\phi \in S - \text{Mor}(\mathcal{C})$ such that

$$\mathcal{S}_{\underline{0}}^{\{i\}}(\phi) = \phi_i.$$

Being cube-like is a very substantial restriction on n -fold categories, essentially still a concept of (1-fold) categories, since the condition only involves morphisms between objects. In this paper, all n -fold categories we will deal with will be cube-like, unless we specify otherwise. Also, in all of our examples of cube-like n -fold categories, the 1-subcategories

of $\{i\}$ -morphisms will be groupoids. We shall call such cube-like categories *invertible cube-like*. In particular, \mathcal{B}_n and the n -actads will be invertible cube-like n -fold categories. An n -fold functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is called *fibred* if for every $x \in \text{Obj}(\mathcal{C})$, for every $\phi \in S - \text{Mor}(\mathcal{D})$ with $\mathcal{S}_{\underline{0}}(\phi) = F(x)$, there exists a unique $\psi \in S - \text{Mor}(\mathcal{C})$ such that $\mathcal{S}_{\underline{0}}(\psi) = x$, $F(\psi) = \phi$, and all S -morphisms of \mathcal{C} are given this way. (Note that other variants of this concept exist).

Also, define

Definition 3. Define the category I by

$$\text{Obj}(I) = \{0, 1\}$$

$$\text{Mor}(I) = \{(0 \rightarrow 0), (0 \rightarrow 1), (1 \rightarrow 1)\}.$$

Then let $\text{Set}_{(n)}$ be the n -fold category of sets, defined as follows: For a subset $S \subseteq \{1, \dots, n\}$, let the set of S -morphisms is

$$S - \text{Mor}(\text{Set}_{(n)}) = \{F \mid F \text{ is a functor } I^S \rightarrow \text{Set}\}.$$

Now, we have a Lemma:

Lemma 4. (1) Let \mathcal{D} be an n -fold category. A fibred n -fold functor $F : \mathcal{C} \rightarrow \mathcal{D}$ produces (in a bijective fashion) an n -fold functor $\Phi : \mathcal{D} \rightarrow \text{Set}_{(n)}$: for some $S \subseteq \{1, \dots, n\}$, $\psi \in S - \text{Mor}(\mathcal{D})$,

$$(21) \quad \Phi(\psi)(\mathbb{I}) = \mathcal{S}_{\mathbb{I}}(F^{-1}(\psi)),$$

(recall (19) for the definition of \mathbb{I}) with

$$\Phi(\psi) : \text{Mor}(I^{|S|}) \rightarrow \text{Mor}(\text{Set})$$

defined by the fibred property of F .

(2) If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a fibred n -fold functor and \mathcal{D} is cube-like, then \mathcal{C} is cube-like.

Proof. First we will discuss (1). Starting with a fibred n -fold functor $F : \mathcal{C} \rightarrow \mathcal{D}$, define Φ by (21). Then the fact that Φ is an n -fold functor follows directly from the properties of composition in an n -fold category.

Given an n -fold functor $\Phi : \mathcal{D} \rightarrow \text{Set}_{(n)}$, define, for $S \subseteq \{1, \dots, n\}$,

$$(22) \quad S - \text{Mor}(\mathcal{C}) = \coprod_{\psi \in S - \text{Mor}(\mathcal{D})} \Phi(\psi)$$

with $F : \mathcal{C} \rightarrow \mathcal{D}$ given by projection. Then F is a fibred n -fold functor by the functoriality properties of F . Additionally, the constructions of Φ from F and F from Φ are obviously inverse to each other.

(2) is immediate from (22), since in a fibered n -fold functor, for $S \subseteq \{1, \dots, n\}$, we can lift the $\{i\}$ -morphisms with $i \in S$ with a common source uniquely to $\{i\}$ -morphisms with a given common source, which then complete uniquely to an S -morphism both downstairs and upstairs. \square

We will also need a way to construct an n -fold category from sequences of morphisms of an $(n-1)$ -fold category for the induction step from \mathcal{B}_n to \mathcal{B}_{n+1} . (This is a generalization of $T_{\mathcal{B}_{n-1}}$ from the case of plain n -actads and n -bases.)

Definition 5. *Suppose C is an $(n-1)$ -fold category. Define the n -fold category T_C in the following way: Suppose $S \subseteq \{1, \dots, n-1\}$. Then define the S -morphisms by*

$$\begin{aligned}
 S - Mor(T_C) &= \{(\alpha_1, \dots, \alpha_k) \mid k \geq 1, \alpha_i \in S - Mor(C)\} \\
 (S \cup \{n\}) - Mor(T_C) &= \\
 \{f = ((\alpha_1, \dots, \alpha_k), \sigma, (\beta_1, \dots, \beta_k)) \\
 (23) \quad &: (\alpha_1, \dots, \alpha_k) \rightarrow (\beta_1, \dots, \beta_k) \mid k \geq 1, \alpha_i \in S - Mor(C) \\
 &\quad \sigma : \{1, \dots, k\} \rightarrow \{1, \dots, k\} \text{ permutation, } \forall i \\
 &\quad \beta_i = \alpha_{\sigma(i)}\}.
 \end{aligned}$$

For $f \in S \cup \{n\} - Mor(T_C)$ in (23), we will sometimes denote the permutation σ by σ_f . It suffices to have \mathcal{B}_n as an $(n-1)$ -fold category, with an $(n-1)$ -fold functor

$$\mathcal{F}_n : \mathcal{B}_n \rightarrow T_{\mathcal{B}_{n-1}}.$$

Then the n -fold category structure of \mathcal{B}_n is induced by \mathcal{F}_n . Explicitly, for $S \subseteq \{1, \dots, n-1\}$, $x, y \in S - Mor(\mathcal{B}_n)$,

$$(S \cup \{n\}) - Mor(\mathcal{B}_n)(x, y) = (S \cup \{n\}) - Mor(T_{\mathcal{B}_{n-1}})(\mathcal{F}(x), \mathcal{F}(y)).$$

Then for $S \subseteq \{1, \dots, n\}$, $x \in S - Mor(\mathcal{B}_n)$, denote again the length of $\mathcal{F}_n(x)$ by m_x and


$$\mathcal{F}_n(x) =: (f_1(x), \dots, f_{m_x}(x)).$$

Thereby, \mathcal{F}_n becomes an n -fold functor. We will also have an $(n-1)$ -fold functor

$$\mathcal{G}_n : \mathcal{B}_n \rightarrow \mathcal{B}_{n-1}$$


given by applying the \mathcal{B}_{n-1} -composition, similarly as in our definition of G_n .

An example of a $\{1\}$ -morphism in \mathcal{B}_2 is the map that takes the tree on the left to the tree on the right by




(24)

Call it ϑ . An example of a $\{2\}$ -morphism in \mathcal{B}_2 is the map that takes the tree on the left to the tree on the right by



(25)

Call it ϱ . Then we have a $\{1, 2\}$ -morphism in \mathcal{B}_2 given by the following diagram:



where the maps on the left are ϱ , and the maps on the bottom are ϑ .

To capture the properties of the composition, define cube-like n -fold categories $Comp_n$, $Comp_n^2$ as follows: For $S \subseteq \{1, \dots, n-1\}$, define

$$(26) \quad \begin{aligned} & S - Mor(Comp_n) = \\ & \{(x, i, y) \mid x, y \in S - Mor(\mathcal{B}_n), 1 \leq i \leq m_x, \mathcal{G}_n(y) = f_i(x)\}, \end{aligned}$$

$$\begin{aligned} (S \cup \{n\}) - Mor(Comp_n) &= \{(x, i, y) \rightarrow (x, i', y), f, g \mid 1 \leq i \leq m_x, \\ & x, y \in S - Mor(\mathcal{B}_n), \mathcal{G}_n(y) = f_i(x), \\ & f : x \rightarrow x, g : y \rightarrow y, f, g \in (S \cup \{n\}) - Mor(\mathcal{B}_n), i' = \sigma_f(i)\}, \end{aligned}$$

$$\begin{aligned} S - Mor(Comp_n^2) &= \{(x, i, j, y, z) \mid x, y, z \in S - Mor(\mathcal{B}_n), \\ & 1 \leq i \neq j \leq m_x, \mathcal{G}_n(y) = f_i(x), \mathcal{G}_n(z) = f_j(x)\}, \end{aligned}$$

$$\begin{aligned} (S \cup \{n\}) - Mor(Comp_n^2) &= \{(x, i, j, y, z) \rightarrow (x, i', j', y, z), f, g, h \mid \\ & 1 \leq i \neq j \leq m_x, \\ & x, y, z \in S - Mor(\mathcal{B}_n), \mathcal{G}_n(y) = f_i(x), \mathcal{G}_n(z) = f_j(x), \\ & f : x \rightarrow x, g : y \rightarrow y, h : z \rightarrow z, f, g, h \in S \cup \{n\} - Mor(\mathcal{B}_n), \\ & i' = \sigma_f(i), j' = \sigma_f(j)\}. \end{aligned}$$

Note that we then have, by definition, projection n -fold functors

$$\begin{aligned} Comp_n &\rightarrow \mathcal{B}_n \times \mathcal{B}_n, \\ Comp_n^2 &\rightarrow \mathcal{B}_n \times \mathcal{B}_n \times \mathcal{B}_n. \end{aligned}$$

We are also given (as a part of the induction data) increasing functions

$$\begin{aligned} \varphi_n^{(x,i,y)} &: \{1, \dots, m_x\} \rightarrow \{1, \dots, m_x + m_y - 1\}, \\ \psi_n^{(x,i,y)} &: \{1, \dots, m_x\} \rightarrow \{1, \dots, m_x + m_y - 1\} \end{aligned}$$

for $(x, i, y) \in S - Mor(Comp_n)$ for some $S \subseteq \{1, \dots, n-1\}$, and a composition n -fold functor

$$(27) \quad \Phi_n : Comp_n \rightarrow \mathcal{B}_n.$$

Then define

$$\Phi_n^1, \Phi_n^2 : Comp_n^2 \rightarrow Comp_n,$$

with

$$\begin{aligned} \Phi_n^1(x, i, j, y, z) &= (\Phi_n(x, i, y), \varphi_n^{(x,i,y)}(j), z) \\ \Phi_n^2(x, i, j, y, z) &= (\Phi_n(x, j, y), \varphi_n^{(x,i,y)}(i), z). \end{aligned}$$

On $(S \cup \{n\}) - \text{Mor}(Comp_n^2)$, Φ_n , Φ_n^1 , and Φ_n^2 are defined by the wreath products of the permutations such that

$$(28) \quad \begin{array}{ccc} Comp_n^2 & \xrightarrow{\Phi_n^1} & Comp_n \\ \Phi_n^2 \downarrow & & \downarrow \Phi_n \\ Comp_n & \xrightarrow{\Phi_n} & \mathcal{B}_n \end{array}$$

commutes strictly. (Note that then the $\{n\}$ -morphisms of \mathcal{B}_n are permutations.)

We will construct the \mathcal{B}_n 's inductively. Suppose we are given \mathcal{B}_n and all of the associated maps. Then take a cube-like n -fold category \mathcal{C} with a fibered n -fold functor

$$\mathcal{C} \rightarrow \mathcal{B}_n.$$

For $S \subseteq \{1, \dots, n\}$, denote the preimage of an S -morphism x by $\mathcal{C}(x)$. Then let

$$\begin{aligned} Comp_{\mathcal{C}} &= (\mathcal{C} \times \mathcal{C}) \times_{(\mathcal{B}_n \times \mathcal{B}_n)} Comp_n \\ Comp_{\mathcal{C}}^2 &= (\mathcal{C} \times \mathcal{C} \times \mathcal{C}) \times_{(\mathcal{B}_n \times \mathcal{B}_n \times \mathcal{B}_n)} Comp_n^2. \end{aligned}$$

Definition 6. A cube-like n -fold category \mathcal{C} with a fibered n -fold functor $\mathcal{C} \rightarrow \mathcal{B}_n$ is called an n -actad provided that there exist n -fold functors

$$\begin{aligned} \Gamma &: Comp_{\mathcal{C}} \rightarrow \mathcal{C}, \\ \Gamma^1, \Gamma^2 &: Comp_{\mathcal{C}}^2 \rightarrow Comp_{\mathcal{C}}, \end{aligned}$$

such that the following axioms hold:

(1)

$$\Gamma^1 = (\Gamma \times Id_{\mathcal{C}}) \times_{Comp_n \times \mathcal{B}_n} \Phi_n^1$$

(2)

$$\Gamma^2 = (Id_{\mathcal{C}} \times \Gamma) \times_{\mathcal{B}_n \times Comp_n} \Phi_n^2$$

(3)

$$\begin{array}{ccc} Comp_{\mathcal{C}}^2 & \xrightarrow{\Gamma^1} & Comp_{\mathcal{C}} \\ \Gamma^2 \downarrow & & \downarrow \Gamma \\ Comp_{\mathcal{C}} & \xrightarrow{\Gamma} & \mathcal{C} \end{array}$$

strictly commutes.

Let \mathcal{C}_n be the category of n -actads, and let \mathcal{D}_n be the category of n -fold fibered categories over \mathcal{B}_n . Then we have a forgetful fibered functor

$$\mathcal{C}_n \rightarrow \mathcal{D}_n.$$

Then the left adjoint to this functor is called a free n -actad.

Suppose that we are given \mathcal{B}_n , the n -actads and all of the associated maps.

Suppose \mathcal{C} is an n -fold category and $S \subseteq \{1, \dots, n\}$. Recall that for $\mathbb{I} : S \rightarrow \{0, 1\}$, we have the source-target map (20).

We then define the $(n+1)$ -base \mathcal{B}_{n+1} as a cube-like n -fold category by letting, for $S \subseteq \{1, \dots, n\}$, $S\text{-Mor}(\mathcal{B}_{n+1})$ be the free plain n -actad on

$$\mathcal{S}_{\underline{0}} : S\text{-Mor}(\mathcal{B}_n) \rightarrow B_n,$$

($B_n = \text{Obj}(\mathcal{B}_n)$). One checks readily that this defines an invertible cube-like n -fold category. (In fact, because of the invertibility, $\underline{0}$ could be equivalently replaced by any $\mathbb{I} : S \rightarrow \{0, 1\}$.)

\mathcal{B}_{n+1} then becomes an invertible cube-like $(n+1)$ -fold category by the device described under Definition 5, and satisfies (28) with n replaced by $n+1$. Then Φ_{n+1} (see (27)) on S -morphisms with $S \subseteq \{1, \dots, n\}$ is defined by plain n -actad composition of S -morphisms in \mathcal{B}_n . Again, Φ_{n+1} on $S \cup \{n+1\}$ are defined by wreath products of permutations (see the comments under (27)).

It is tempting to conclude that \mathcal{B}_{n+1} is the free n -actad on \mathcal{B}_n , but that is false because for $S \subseteq \{1, \dots, n\}$, not all S -morphisms of \mathcal{B}_n can be lifted to \mathcal{B}_{n+1} via \mathcal{G}_{n+1} . This is due to the fact that additional structure is being introduced, where not all isomorphisms downstairs will preserve it: for example, for $n=2$, not every 1-permutation of the prongs of a planar tree comes from an actual isomorphism of trees. Thus, \mathcal{B}_{n+1} is, in fact, not a fibered n -fold category over \mathcal{B}_n (via \mathcal{G}_n), and consequently not an n -actad. On the other hand, $\mathcal{G}_{n+1} : \mathcal{B}_{n+1} \rightarrow \mathcal{B}_n$ satisfies the *uniqueness* axiom of a fibered n -fold category, and in fact, the following is true (see also Section 8 below):

Lemma 7. *Let \mathcal{U} be the forgetful functor from the category of fibered n -fold categories over \mathcal{B}_n to the category of n -fold categories over \mathcal{B}_n (and n -fold functors). Let \mathcal{V} be the left adjoint to \mathcal{U} (“left Kan extension”). Then $\mathcal{B}_{n+1}^- = \mathcal{V}\mathcal{B}_{n+1}$ is the free n -actad on \mathcal{B}_n on $\text{Id} : \mathcal{B}_n \rightarrow \mathcal{B}_n$.*

□

(Note that in the case of $n = 1$, 1-actads are simply operads.)

4. ALGEBRAS OVER AN n -ACTAD

The next thing we will discuss is algebras over an n -actad \mathcal{C} . For operads, which are 1-actads, an algebra over an operad \mathcal{C} is an X with

$$\mathcal{C}(n) \times X^n \rightarrow X$$

satisfying the appropriate axioms (associativity, and, if we choose, commutativity in the broader sense and unitality), and a module over an operad \mathcal{C} and an algebra X is a Y with

$$\mathcal{C}(n) \times X^{n-1} \times Y \rightarrow Y$$

satisfying similar axioms.

Now, take an $(n-1)$ -fold category X . Suppose we also have a fibered $(n-1)$ -fold functor

$$\Xi : X \rightarrow \mathcal{B}_{n-1}$$

(then, in particular, X is cube-like). For $S \subseteq \{1, \dots, n-1\}$, $a \in S - \text{Mor}(\mathcal{B}_{n-1})$, let the inverse image of Ξ of a again be called $X(a)$.

Suppose we have $S \subseteq \{1, \dots, n-1\}$. Then define the extra structure for X needed to be n -fold category, by insisting that the map

$$(29) \quad (S \cup \{n\} - \text{Mor}(X)) \xleftarrow{\text{Id}} (S - \text{Mor}(X))$$

be a bijection. With this extra structure, X is an n -fold category.

Let \mathcal{C} be an n -actad. Then, by definition, we have an n -fold functor

$$\mathcal{C} \longrightarrow \mathcal{B}_n \xrightarrow{\mathcal{F}_n} T_{\mathcal{B}_{n-1}}.$$

Also, we have

$$T_{\Xi} : T_X \rightarrow T_{\mathcal{B}_{n-1}}.$$

So we can form the pullback $\mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X$.

Definition 8. X is called an algebra over \mathcal{C} provided that the following holds: There exists an $(n-1)$ -fold functor

$$(30) \quad \Theta : \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X \rightarrow X,$$

such that the diagram

$$\begin{array}{ccc}
 \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X & \xrightarrow{\Theta} & X \\
 \pi \downarrow & & \downarrow \Xi \\
 \mathcal{C} & & \mathcal{B}_{n-1} \\
 \downarrow & & \downarrow \\
 \mathcal{B}_n & \xrightarrow{\mathcal{G}_n} & \mathcal{B}_{n-1}
 \end{array}$$

commutes where π is the projection, and the map $\mathcal{C} \rightarrow \mathcal{B}_n$ is the fibered n -functor given by definition. Note that though some of the functors in this diagram are n -fold functors, this diagram is of $(n-1)$ -fold functors. We require the following conditions to hold:

- (1) *Equivariance:*

By the above comment on the n -fold categorical structure of X , We have that $\mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X$ is already an n -fold category. Then we require (30) to be an n -fold functor.

- (2) *Associativity:*

First, fix $x, y \in S\text{-Mor}(\mathcal{B}_n)$, $S \subseteq \{1, \dots, n-1\}$, $i \in \{1, \dots, m_x\}$, with $f_i(x) = \mathcal{G}_n(y)$. Or, in other words, $(x, i, y) \in \text{Comp}_n$. Then write

$$\begin{aligned}
 x_j &= f_j(x), \\
 y_k &= f_k(y).
 \end{aligned}$$

Suppose we have

$$\xi \in \mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\}.$$

Recall that elements of $\mathcal{C}(x)$ are S -morphisms of \mathcal{B}_n over x . Then define

$$\Omega(\xi) = (z_1, \dots, z_{m_x+m_y-1})$$

by

$$\begin{aligned}
 z_{\phi_n^{(x,i,y)}(j)} &= x_j, \\
 z_{\psi_n^{(x,i,y)}(k)} &= y_k.
 \end{aligned}$$

Then

$$\Omega : \mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\} \rightarrow T_{\mathcal{B}_{n-1}}.$$

We also have

$$T_{\Xi} : T_X \rightarrow T_{\mathcal{B}_{n-1}}.$$

So, we can write

$$(\mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\}) \times_{T_{\mathcal{B}_{n-1}}} T_X.$$

Then define

$$\begin{aligned} \tilde{\Theta}_{(x,i,y)} : (\mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\}) \times_{T_{\mathcal{B}_{n-1}}} T_X &\rightarrow \\ &\rightarrow \mathcal{C}(x) \times_{T_{\mathcal{B}_{n-1}}} \prod_{1 \leq j \leq m_x} X(x_j) \end{aligned}$$

by

$$(31) \quad \begin{aligned} \tilde{\Theta}_{(x,i,y)} &= (\varpi, Id_{\mathcal{C}(x) \times \prod_{1 \leq j \leq m_x, j \neq i} X(x_j)} \times \\ &\Theta |_{\mathcal{C}(y) \times_{T_{\mathcal{B}_{n-1}}} \prod_{i \leq k \leq m_y} X(y_k)}). \end{aligned}$$

where, again,

$$\varpi : (\mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\}) \times_{T_{\mathcal{B}_{n-1}}} T_X \rightarrow \mathcal{C}(x)$$

is the projection to the first coordinate.

Now, let

$$\tilde{\Theta} = \prod_{(x,i,y) \in \text{Comp}_n} \tilde{\Theta}_{(x,i,y)}.$$

Then, since

$$\text{Comp}_{\mathcal{C}} = (\mathcal{C} \times \mathcal{C}) \times_{(\mathcal{B}_n \times \mathcal{B}_n)} \text{Comp}_n,$$

we have

$$\tilde{\Theta} : \text{Comp}_{\mathcal{C}} \times_{T_{\mathcal{B}_{n-1}}} T_X \rightarrow \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X.$$

Then

$$(32) \quad \begin{array}{ccc} \text{Comp}_{\mathcal{C}} \times_{T_{\mathcal{B}_{n-1}}} T_X & \xrightarrow{\Gamma \times Id} & \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X \\ \tilde{\Theta} \downarrow & & \downarrow \Theta \\ \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X & \xrightarrow{\Theta} & X \end{array}$$

must strictly commute. Actually, $\tilde{\Theta}$ can be extended uniquely and naturally to $S \cup \{n\}$ -morphisms, and (32) is automatically a diagram of n -fold functors.

Again, at $n = 1$, this simply corresponds to the concept of algebras over an operad.

Examples: 1: Let \mathcal{A}_n be the n -actad given by taking, for $S \subseteq \{1, \dots, n-1\}$,

$$S - \text{Mor}(\mathcal{A}_n) = (S \cup \{n\}) - \text{Mor}(\mathcal{B}_n),$$

and $(S \cup \{n\}) - Mor(\mathcal{A}_n)$ is the n -composable pairs of $(S \cup \{n\}) - Mor(\mathcal{B}_n)$, i.e. the pairs

$$(\sigma, \tau) \in ((S \cup \{n\}) - Mor(\mathcal{B}_n))^2$$

such that $S_n \tau = T_n \sigma \in S - Mor(\mathcal{B}_n)$, where S_n, T_n denote the n -source and n -target of an $S \cup \{n\}$ -morphism. Then there is an equivalence of categories between \mathcal{A}_n -algebras and $(n - 1)$ -actads. More specific examples of \mathcal{A}_n -algebras will be given in later sections.

2: Note that, of course, \mathcal{B}_n is an n -actad via $Id : \mathcal{B}_n \rightarrow \mathcal{B}_n$. We call \mathcal{B}_n -algebras n -commutative $(n - 1)$ -actads, (in the narrower sense).

5. MODULES OF ALGEBRAS OVER n -ACTADS

Suppose X, M are $(n - 1)$ -fold categories and we have fibered $(n - 1)$ -fold functors

$$\Xi_X : X \rightarrow \mathcal{B}_{n-1}$$

$$\Xi_M : M \rightarrow \mathcal{B}_{n-1}.$$

Then let, again, $X(x)$ and $M(x)$ denote the inverse images under Ξ_X and Ξ_M of $x \in \mathcal{B}_{n-1}$ respectively. For $S \subseteq \{1, \dots, n - 1\}$, define $T_{X,M}$ by

$$(33) \quad \begin{aligned} S - Mor(T_{X,M}) = \{ & ((\iota, k), (x_1, \dots, x_{\iota-1}, m, x_{\iota+1}, \dots, x_k)) \mid \\ & 1 \leq \iota \leq k \in \mathbb{N}, \\ & x_j \in S - Mor(X), m \in S - Mor(M) \}, \end{aligned}$$

$$\begin{aligned} (S \cup \{n\}) - Mor(T_{X,M}) = \{ & ((\sigma, \iota, k), (x_1, \dots, x_{\iota-1}, m, x_{\iota+1}, \dots, x_k)) \mid \\ & 1 \leq \iota \leq k \in \mathbb{N} \\ & x_j \in S - Mor(X), m \in S - Mor(M), \sigma \in \Sigma_k \}. \end{aligned}$$

For an $\eta = ((\iota, k), (x_1, \dots, x_{\iota-1}, m, x_{\iota+1}, \dots, x_k)) \in S - Mor(T_{X,M})$, write $\iota_\eta = \iota$.

Definition 9. *Suppose \mathcal{C} is an n -actad and X is a \mathcal{C} -algebra. Then M is a (\mathcal{C}, X) -module provided that the following holds: There exists an $(n - 1)$ -fold functor*

$$(34) \quad \Theta : \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_{X,M} \rightarrow M$$

such that

$$\begin{array}{ccc}
 \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_{X,M} & \xrightarrow{\Theta} & M \\
 \pi \downarrow & & \downarrow \Xi_M \\
 \mathcal{C} & & \\
 \downarrow & & \\
 \mathcal{B}_n & \xrightarrow{g} & \mathcal{B}_{n-1},
 \end{array}$$

where π is the projection, and the map $\mathcal{C} \rightarrow \mathcal{B}_n$ is the n -functor given by the definition. Note that though some of the functors in this diagram are n -fold functors, this diagram is of $(n-1)$ -fold functors. We also require the following conditions to hold:

(1) *Equivariance:*

Suppose we have $S \subseteq \{1, \dots, n-1\}$. Then define the extra structure for M needed to be n -fold category, by insisting that the map

$$(S \cup \{n\} - \text{Mor}(M)) \xleftarrow{\text{Id}} (S - \text{Mor}(M))$$

be an isomorphism. With this extra structure, M is an n -fold category. We have that $\mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_{X,M}$ is already an n -fold category. Then (34) must be an n -fold functor.

(2) *Associativity:*

We will use the symbols $x, y, S, i, x_j, y_k, \Omega$ in the same way as in Axiom (2) in Section 4. We have

$$T_{\Xi_X} : T_X \rightarrow T_{\mathcal{B}_{n-1}},$$

We define

$$\Upsilon : T_{X,M} \rightarrow T_{\mathcal{B}_{n-1}}$$

by

$$\begin{aligned}
 & \Upsilon(x_1, \dots, x_{i-1}, m, x_{i+1}, \dots, x_k) = \\
 & = (\Xi_X(x_1), \dots, \Xi_X(x_{i-1}), \Xi_M(m), \Xi_X(x_{i+1}), \dots, \Xi_X(x_k)).
 \end{aligned}$$

So, we can write

$$(\mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\}) \times_{T_{\mathcal{B}_{n-1}}} T_{X,M}.$$

Then define

$$\begin{aligned}
 \tilde{\Theta}_{(x,i,y)} : (\mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\}) \times_{T_{\mathcal{B}_{n-1}}} T_{X,M} & \rightarrow \\
 & \rightarrow \mathcal{C}(x) \times_{T_{\mathcal{B}_{n-1}}} \prod_{1 \leq j \leq m_x} M(x_j)
 \end{aligned}$$

by

$$\begin{aligned} \tilde{\Theta}_{(x,i,y)} &= (\varpi, Id_{\mathcal{C}(x) \times \prod_{1 \leq j \leq m_x, j \neq i} XM_j(x_j)} \times \\ &\Theta |_{\mathcal{C}(y) \times_{T_{\mathcal{B}_{n-1}}} \prod_{i \leq k \leq m_y} XM_k(y_k)}). \end{aligned}$$

where, for $\eta \in T_{X,M}$ on $\mathcal{C}(x) \times \mathcal{C}(y) \times \{(x, i, y)\} \times_{T_{\mathcal{B}_{n-1}}} \{\eta\}$,

$$\begin{aligned} XM_j &= X \text{ for } j \neq \iota \\ &= M \text{ for } j = \iota. \end{aligned}$$

Then

$$(35) \quad \begin{array}{ccc} \text{Comp}_{\mathcal{C}} \times_{T_{\mathcal{B}_{n-1}}} T_{X,M} & \xrightarrow{\Gamma \times Id} & \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_{X,M} \\ \tilde{\Theta} \downarrow & & \downarrow \Theta \\ \mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_{X,M} & \xrightarrow{\Theta} & M \end{array}$$


must strictly commute.

We can also define a concept of a (\mathcal{C}, X) -algebra M analogously by modifying the definition so as to allow multiple m_j 's in (33).

6. ITERATED ALGEBRAS

There is a concept of an *I-iterated algebra* over an n -actad for any subset $I \in \{0, \dots, n\}$, of which the concepts of algebra and algebra with a module introduced in the last two sections are special cases for $I = \{n\}$ and $I = \{0, n\}$, respectively. We will use these constructions later to define unital actads and R -units. An I -iterated algebra consists of $|I|$ cube-like $(n-1)$ -fold categories, with rules on where they can be “plugged in into an element of a given n -actad.” The ways of plugging in the α 'th model, for an $\alpha \in I$, follow, roughly, the pattern of an α -actad algebra. However, the precise axiomatization is delicate, in particular, on morphisms. The only axiomatization I could work out uses multisorted algebras [6]. One must also design an appropriate multisorted version of the n -base, to keep track of the structure describing how the different models are being plugged in.

In the below picture, I illustrate how the 0-, 1-, and 2-models are plugged in in the composition for the 0-, 1-, and 2-models of a $\{0, 1, 2\}$ -iterated algebra over a 2-actad.



(the solid nodes mark the places where trees that will be labeled a number ≥ 0 are composed).

We shall start with the concept of the I -pointed n -base \mathcal{B}_n^I , a cube-like n -fold category, which will be defined inductively.

We start the induction with $\mathcal{B}_0^{\{0\}} = \mathcal{B}_0^0 = \mathcal{B}_0$. For $k \geq 0$, $0 \leq \alpha_1 < \dots < \alpha_k \leq n$, $I = \{\alpha_1, \dots, \alpha_k\}$, $\mathcal{B}_n^I = \mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$ is an n -fold category with a n -fold functor

$$\mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \rightarrow \mathcal{B}_n$$

and with some additional maps. First, $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$ comes with additional data of n -fold functor

$$\wp_n : \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \rightarrow \{1, \dots, k\}$$

as part of the induction data, where on the right hand side, all morphisms are identities. We shall also write

$$\mathcal{B}_n^I(i) = \mathcal{B}_n^{\alpha_1, \dots, \alpha_k}(i) = \wp_n^{-1}(i).$$

We will construct $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$ (and the associated structure) inductively on n . The induction step from $(n-1)$ to n depends in a crucial way on whether $\alpha_k < n$ or $\alpha_k = n$. Essentially, if $\alpha_k < n$, the induction step from $\mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_k}$ is similar as for \mathcal{B}_n . If $\alpha_k = n$, the induction step to $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$ will be from $\mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_{k-1}}$ to $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$. The rules for plugging in the new model are essentially to plug into one of the places

of the previous one, but the $(S \cup \{n\})$ -morphisms must remember their positions.

In more detail, if $\alpha_k < n$, then we have an $(n - 1)$ -fold functor over $\{1, \dots, k\}$

$$\begin{array}{ccc} \mathcal{F}_n^{\alpha_1, \dots, \alpha_k} : \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} & \longrightarrow & T_{\mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_k}} \\ & \searrow & \downarrow \\ & & \{1, \dots, k\}, \end{array}$$

where in the vertical arrow, an S -morphism (x_1, \dots, x_m) with $S \subseteq \{1, \dots, n - 1\}$ is mapped to

$$(37) \quad \max_{i \in \{1, \dots, m\}} \wp_{n-1}(x_i).$$

Then, like in the case of n -bases, $\mathcal{F}_n^{\alpha_1, \dots, \alpha_k}$ induces an n -fold category structure on $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$, and thus becomes an n -fold functor. Again, we also will have an $(n - 1)$ -fold functor

$$\mathcal{G}_n : \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \rightarrow \mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_k}$$

as part of the induction data.

Definitions of $Comp_n^{\alpha_1, \dots, \alpha_k}$ and $Comp_n^{2; \alpha_1, \dots, \alpha_k}$ then proceed precisely analogously to the definitions of $Comp_n$ and $Comp_n^2$ in Section 3, except for the fact that we are working over $\{1, \dots, k\}$. The analogue of Diagram (28) becomes a diagram of the form

$$(38) \quad \begin{array}{ccc} Comp_n^{2; \alpha_1, \dots, \alpha_k} & \xrightarrow{\Phi_n^{1; \alpha_1, \dots, \alpha_k}} & Comp_n^{\alpha_1, \dots, \alpha_k} \\ \Phi_n^{2; \alpha_1, \dots, \alpha_k} \downarrow & & \downarrow \\ Comp_n^{\alpha_1, \dots, \alpha_k} & \longrightarrow & \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \end{array}$$

over $\{1, \dots, k\}$.

Now, suppose $\alpha_k = n$. This case must be treated in more detail. Define a map

$$\varpi : \{1, \dots, k\} \rightarrow \{1, \dots, k - 1\}$$

by

$$\varpi(i) = i, \text{ for } i < k$$

$$\varpi(k) = k - 1.$$

Definition 10. Suppose $\mathcal{C} \rightarrow \{1, \dots, k-1\}$ is a $(n-1)$ -fold category over $\{1, \dots, k-1\}$. Then define an n -fold category $T_{\mathcal{C}}^{\bullet} \rightarrow \{1, \dots, k\}$ over $\{1, \dots, k\}$ as follows: For $i \in \{1, \dots, k-1\}$,

$$T_{\mathcal{C}}^{\bullet}(i) = T_{\mathcal{C}(i)}.$$

For $S \subseteq \{1, \dots, n-1\}$,

$$\begin{aligned} S - \text{Mor}(T_{\mathcal{C}}^{\bullet}(k)) = \\ \{((x_1, \dots, x_{\ell}), b) \mid \ell \geq 1, b \in \{1, \dots, \ell\}, \forall j \exists i_j \in \{1, \dots, k-1\} \\ \text{such that } x_j \in S - \text{Mor}(\mathcal{C}(i_j)), i_b = k-1, \} \end{aligned}$$

$$\begin{aligned} (S \cup \{n\}) - \text{Mor}(T_{\mathcal{C}}^{\bullet}(k)) = \\ \{f = (((x_1, \dots, x_{\ell}), b), \sigma, (y_1, \dots, y_{\ell})) \\ : ((x_1, \dots, x_{\ell}), b) \rightarrow ((y_1, \dots, y_{\ell}), \sigma(b)) \mid \ell \geq 1, \\ b \in \{1, \dots, \ell\}, \sigma : \{1, \dots, \ell\} \rightarrow \{1, \dots, \ell\} \text{ a permutation,} \\ \sigma(b) = b, \forall i y_i = x_{\sigma(i)}\}. \end{aligned}$$

We have an obvious forgetful n -fold functor

$$\begin{array}{ccc} T_{\mathcal{C}}^{\bullet} & \longrightarrow & T_{\mathcal{C}} \\ \downarrow & & \downarrow \\ \{1, \dots, k\} & \xrightarrow{\varpi} & \{1, \dots, k-1\} \end{array}$$

forgetting b .

Then we are also given a $(n-1)$ -fold functor

$$(39) \quad \mathcal{F}_n^I : \mathcal{B}_n^I = \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \rightarrow T_{\mathcal{B}_{n-1}}^{\bullet, \alpha_1, \dots, \alpha_{k-1}}$$

as part of the induction data. Like in the previous cases, \mathcal{F}_n^I becomes an n -fold functor. In addition, like in the previous cases, we have a $(n-1)$ -fold functor

$$\mathcal{G}_n^I : \mathcal{B}_n^I \rightarrow \mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_{k-1}},$$

with the following diagram

$$\begin{array}{ccc} \mathcal{B}_n^I & \xrightarrow{\mathcal{G}_n} & \mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_{k-1}} \\ \downarrow & & \downarrow \\ \{1, \dots, k\} & \xrightarrow{\varpi} & \{1, \dots, k-1\}. \end{array}$$

Then for $S \subseteq \{1, \dots, n-1\}$, $x \in S - \text{Mor}(\mathcal{B}_n^I)$, denote

$$\mathcal{F}_n^I(x) =: (f_1(x), \dots, f_{m_x}(x)).$$

Now, we will use these functors to define the n -fold categories $\text{Comp}_n^{\alpha_1, \dots, \alpha_k}$, $\text{Comp}_n^{2, \alpha_1, \dots, \alpha_k}$ similarly to the case of n -bases: For $S \subseteq \{1, \dots, n-1\}$, define

$$\begin{aligned} S - \text{Mor}(\text{Comp}_n^{\alpha_1, \dots, \alpha_k}) = \\ \{(x, i, y) | x, y \in S - \text{Mor}(\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}), 1 \leq i \leq m_x, \mathcal{G}_n(y) = f_i(x)\} \end{aligned}$$

$$\begin{aligned} (S \cup \{n\}) - \text{Mor}(\text{Comp}_n^{\alpha_1, \dots, \alpha_k}) = \{((x, i, y) \rightarrow (x, i', y), f, g) | 1 \leq i \leq m_x, \\ x, y \in S - \text{Mor}(\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}), \mathcal{G}_n(y) = f_i(x), \\ f : x \rightarrow x, g : y \rightarrow y, f, g \in (S \cup \{n\}) - \text{Mor}(\mathcal{B}_n), i' = \sigma_f(i)\}. \end{aligned}$$

$$\begin{aligned} S - \text{Mor}(\text{Comp}_n^{2, \alpha_1, \dots, \alpha_k}) = \{(x, i, j, y, z) | x, y, z \in S - \text{Mor}(\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}), \\ 1 \leq i \neq j \leq m_x, \mathcal{G}_n(y) = f_i(x), \mathcal{G}_n(z) = f_j(x)\}. \end{aligned}$$

$$\begin{aligned} (S \cup \{n\}) - \text{Mor}(\text{Comp}_n^{2, \alpha_1, \dots, \alpha_k}) = \{((x, i, j, y, z) \rightarrow (x, i', j', y, z), f, g, h) | \\ 1 \leq i \neq j \leq m_x, \\ x, y, z \in S - \text{Mor}(\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}), \mathcal{G}_n(y) = f_i(x), \mathcal{G}_n(z) = f_j(x), \\ f : x \rightarrow x, g : y \rightarrow y, h : z \rightarrow z, f, g, h \in S \cup \{n\} - \text{Mor}(\mathcal{B}_n), \\ i' = \sigma_f(i), j' = \sigma_f(j)\}. \end{aligned}$$

Then define maps

$$\Pi_1 : \text{Comp}_n^{\alpha_1, \dots, \alpha_k} \rightarrow \{1, \dots, k\}$$

$$\Pi_2 : \text{Comp}_n^{2, \alpha_1, \dots, \alpha_k} \rightarrow \{1, \dots, k\}$$

by

$$\Pi_1(x, i, y) = \varpi(x)$$

$$\Pi_2(x, i, j, y, z) = \varpi(x).$$

Also, like before, we have

$$\begin{aligned} \text{Comp}_n^{\alpha_1, \dots, \alpha_k} &\rightarrow \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \times \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \\ \text{Comp}_n^{2, \alpha_1, \dots, \alpha_k} &\rightarrow \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \times \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \times \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \\ \varphi_n^{(x, i, y)} : \{1, \dots, m_x\} &\rightarrow \{1, \dots, m_x + m_y - 1\} \\ \Phi_n : \text{Comp}_n^{\alpha_1, \dots, \alpha_k} &\rightarrow \mathcal{B}_n^{\alpha_1, \dots, \alpha_k}, \end{aligned}$$

and

$$\Phi_n^1, \Phi_n^2 : \text{Comp}_n^{2, \alpha_1, \dots, \alpha_k} \rightarrow \text{Comp}_n^{\alpha_1, \dots, \alpha_k},$$

where the diagram analogous to (38) commutes.

Then suppose we are given the $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$ and all of their associated structure described above.

To construct, for $\{\alpha_1, \dots, \alpha_k\} \subseteq \{0, \dots, n\}$, $\mathcal{B}_{n+1}^{\alpha_1, \dots, \alpha_k}$ from $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$, we can proceed precisely analogously to the end of Section 3: For $S \subseteq \{1, \dots, n\}$, $S - \text{Mor}(\mathcal{B}_{n+1}^{\alpha_1, \dots, \alpha_k})$ is the $(n+1)$ -dimensional free plain actad sorted over $\{1, \dots, k\}$ (i.e. plain actad in the category of sets over $\{1, \dots, k\}$) on

$$\mathcal{S}_{\underline{0}} : S - \text{Mor}(\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}) \rightarrow B_n \times \{1, \dots, k\}.$$

Again, this is an invertible cube-like category and $\underline{0}$ could be equivalently replaced by any $\mathbb{I} : S \rightarrow \{0, 1\}$. We let

$$\mathcal{B}_{n+1}^{\alpha_1, \dots, \alpha_k, n+1} = \mathcal{B}_{n+1}^{\alpha_1, \dots, \alpha_k} \times_{T_{\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}}} T_{\mathcal{B}_n^{\alpha_1, \dots, \alpha_k}}^\bullet.$$

(In particular, by definition, $\mathcal{B}_n^0 = \mathcal{B}_n$ for all n).

Now, we will define iterated algebras. Let \mathcal{C} be an n -actad. Let $0 \leq \alpha_1 < \dots < \alpha_k \leq n$. Let

$$\mathcal{C}^{\alpha_1, \dots, \alpha_k} = \mathcal{C} \times_{\mathcal{B}_n} \mathcal{B}_n^{\alpha_1, \dots, \alpha_k}.$$

$$\text{Comp}_{\mathcal{C}}^{\alpha_1, \dots, \alpha_k} = (\mathcal{C} \times \mathcal{C}) \times_{(\mathcal{B}_n \times \mathcal{B}_n)} \text{Comp}_n^{\alpha_1, \dots, \alpha_k}$$

$$\text{Comp}_{\mathcal{C}}^{2; \alpha_1, \dots, \alpha_k} = (\mathcal{C} \times \mathcal{C} \times \mathcal{C}) \times_{(\mathcal{B}_n \times \mathcal{B}_n \times \mathcal{B}_n)} \text{Comp}_n^{2; \alpha_1, \dots, \alpha_k}.$$

(Note that we have a forgetful n -fold functor $\mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \rightarrow \mathcal{B}_n$.)

Definition 11. *Suppose \mathcal{D} is an $(n-1)$ -fold category over $\mathcal{B}_{n-1} \times \{1, \dots, k\}$. Then, for $i = 1, \dots, k$, let $\mathcal{D}(i)$ be the sub- $(n-1)$ -fold categories that are fibered over $\mathcal{B}_{n-1} \times \{i\}$. Then let $T_{\mathcal{D}}^k$ be the n -fold category with*

$$\begin{aligned} S - \text{Mor}(T_{\mathcal{D}}^k) &= \{(x_1, \dots, x_k) \mid \forall j \exists i_j \in \{1, \dots, k-1\} \\ &\text{such that } x_j \in S - \text{Mor}(\mathcal{D}(i_j))\} \end{aligned}$$

$$\begin{aligned} S \cup \{n\} - \text{Mor}(T_{\mathcal{D}}^k) &= \{f = ((x_1, \dots, x_k), \sigma, (y_1, \dots, y_k)) \\ &: (x_1, \dots, x_k) \rightarrow (y_1, \dots, y_k) \mid \end{aligned}$$

$$\sigma : \{1, \dots, k\} \rightarrow \{1, \dots, k\} \text{ a permutation,}$$

$\forall j \ x_j \in S - \text{Mor}(\mathcal{D}(i_j)), y_j \in S - \text{Mor}(\mathcal{D}(\sigma(i_j))), \text{ and } y_j = x_{\sigma(i)}$
for $S \subseteq \{1, \dots, k-1\}$.

An iterated $(\alpha_1, \dots, \alpha_k)$ - \mathcal{C} algebra is a k -tuple of fibered $(n-1)$ -fold categories $X = (X_1, \dots, X_k)$ over \mathcal{B}_{n-1} together with an $(n-1)$ -fold functor

$$\Theta : \mathcal{C}^{\alpha_1, \dots, \alpha_k} \times_{T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k} T_X^k \rightarrow X$$

with

$$\begin{array}{ccc} \mathcal{C}^{\alpha_1, \dots, \alpha_k} \times_{T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k} T_X^k & \xrightarrow{\Theta} & X \\ & \searrow & \downarrow \\ & & \{1, \dots, k\}. \end{array}$$

Again, there are equivariance and associativity axioms. Equivariance, again, says that Θ is an n -fold functor if we let (29) be a bijection. Associativity is expressed by the commutativity of a diagram of n -fold:

$$\begin{array}{ccc} \text{Comp}_{\mathcal{C}}^{\alpha_1, \dots, \alpha_k} \times_{T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k} T_X^k & \xrightarrow{\Gamma \times Id} & \mathcal{C}^{\alpha_1, \dots, \alpha_k} \times_{T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k} T_X^k \\ \tilde{\Theta} \downarrow & & \downarrow \Theta \\ \mathcal{C}^{\alpha_1, \dots, \alpha_k} \times_{T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k} T_X^k & \xrightarrow{\Theta} & X \end{array}$$

where $\tilde{\Theta}$ is defined analogously to (31).

As an example, note that the cases of $\mathcal{B}_n^{\{n\}}$ and $\mathcal{B}_n^{\{0, n\}}$ give the notions of algebras and modules (respectively) over n -actads.

7. R -UNITAL ACTADS AND BASES

A unit axiom can be added to the concept of an n -actad (or a plain n -actad) without difficulty. Perhaps the most surprising aspect of units is how diverse they become for $n > 1$: In the plain case, the unit is expressed by the natural inclusion

$$(40) \quad B_{n-1} \subseteq B_n, \quad n \geq 1.$$

(given by the fact that B_n is the free plain $(n-1)$ -actad on $Id : B_{n-1} \rightarrow B_{n-1}$.)

We introduce the notation $1_y^n \in B_n$ for the image of every $y \in B_{n-1}$. Then $\forall x = ((x_1, \dots, x_\ell), (i_1, \dots, i_{\ell-1}))$

- (1) $x \circ_k^n 1_{x_k}^n = x$
- (2) $\forall y \in B_n$ such that $G_n(y) = x$,

$$1_x^n \circ_1^n y = y.$$

Then, we can define a unital plain actad as follows:

Definition 12. *Suppose we are given a plain n -actad \mathcal{C} . Then \mathcal{C} is a unital plain actad provided that for $x \in B_{n-1}$, $y \in B_n$, $z \in \mathcal{C}(y)$, we are given an element $1_{n,x} \in \mathcal{C}(1_x^n)$ such that*

(1)

$$\gamma_{n,j}(z, 1_{n,x}) = z$$

(2)

$$\gamma_{n,j}(1_{n,x}, z) = z,$$

if $G_n(y) = x$.

For $n = 0$, there is also a unit $1_{0,*}$, even though there is no -1 base.

A unital 0-actad is a monoid (instead of just a semigroup).

For actads, the story is analogous. Here, only the case of $n > 0$ is new. By the definition of \mathcal{B}_{n+1} from \mathcal{B}_n at the end of Section 3, we also obtain a canonical inclusion $(n-1)$ -fold functor of $(n-1)$ -fold categories

$$(41) \quad \mathcal{I}_n : \mathcal{B}_{n-1} \rightarrow \mathcal{B}_n.$$

In fact, it becomes an n -fold functor if we make all the $\{n\}$ -morphisms identities in the source. In fact, (41) is a section of the $(n-1)$ -fold functor \mathcal{G}_n . (In particular, in the n -actad case, note that units have $\{i\}$ -automorphisms for $i < n$.)

The left unit property for \mathcal{B}_n then can be expressed by introducing a left unit functor

$$\mathcal{H}_n : \mathcal{B}_n \rightarrow \text{Comp}_n$$

by

$$x \mapsto (\mathcal{I}_n \mathcal{G}_n x, 1, x)$$

(see (26)). Then the diagram

$$(42) \quad \begin{array}{ccc} \mathcal{B}_n & \xrightarrow{\mathcal{H}_n} & \text{Comp}_n \\ & \searrow \text{Id} & \downarrow \Phi_n \\ & & \mathcal{B}_n \end{array}$$

commutes, and we call it the left unit property. For the right unit axiom, we must remember where the unit is being inserted. Recalling the notation of the last section, we may define a right unit n -fold functor

$$\mathcal{K}_n : \mathcal{B}_n^0 \rightarrow \text{Comp}_n$$

by

$$\mathcal{K}_n(x, b) = (x, b, \mathcal{I}_n f_b(x)).$$

Then the right unit axiom is the commutativity of the diagram

$$(43) \quad \begin{array}{ccc} \mathcal{B}_n^0 & \xrightarrow{\mathcal{K}_n} & \text{Comp}_n \\ & \searrow & \downarrow \Phi_n \\ & & \mathcal{B}_n \end{array}$$

where the diagonal n -fold functor is the projection.

Definition 13. For an n -actad \mathcal{C} , let

$$\mathcal{C}^0 = \mathcal{C} \times_{\mathcal{B}_n} \mathcal{B}_n^0.$$

Then \mathcal{C} is called a unital n -actad if there exists an n -fold functor lift

$$\begin{array}{ccc} & & \mathcal{C} \\ & \nearrow \mathcal{I}_n^{\mathcal{C}} & \downarrow \\ \mathcal{B}_{n-1} & \xrightarrow{\mathcal{I}_n} & \mathcal{B}_n \end{array}$$

such that, letting

$$\mathcal{H}_n^{\mathcal{C}} : \mathcal{C} \rightarrow \text{Comp}_n^{\mathcal{C}}$$

be defined by

$$x \mapsto (\mathcal{I}_n^{\mathcal{C}} \mathcal{G}_n |x|, 1, x),$$

where $|x|$ denotes the projection of x to \mathcal{B}_n , and

$$\mathcal{K}_n^{\mathcal{C}} : \mathcal{C}^0 \rightarrow \text{Comp}_n^{\mathcal{C}}$$

be defined by

$$\mathcal{K}_n^{\mathcal{C}} = (x, b, \mathcal{I}_n^{\mathcal{C}} |f_b(x)|),$$

the following diagrams above (42) and (43) commute:

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\mathcal{H}_n^{\mathcal{C}}} & \text{Comp}_n \\ & \searrow \text{Id} & \downarrow \Gamma \\ & & \mathcal{C} \end{array}$$

$$\begin{array}{ccc} \mathcal{C}^0 & \xrightarrow{\mathcal{K}_n^{\mathcal{C}}} & \text{Comp}_n \\ & \searrow \pi & \downarrow \Gamma \\ & & \mathcal{C} \end{array}$$

where π is the projection.

But now there is also a version of unitality of n -actads (or plain n -actad) which includes units coming from i -actads (or plain i -actads) for all $i < n$. I call this concept *R-unitality* where R stands for “recursive.”

To give an example, a based operad (which, recall, is over \mathbb{N}_0 instead of \mathbb{N}) is a unital operad \mathcal{C} with a base point $* \in \mathcal{C}(0)$. In a way, this comes from the monoid unit, since the free monoid on one generator contains $1 = a^0$, corresponding to the 0 index. It is interesting to note that the base point of an operad is not subject to any axioms. Yet, it is an important feature, adding a rich structure of “degeneracies” given by plugging in the base point. This structure is crucial in the Approximation Theorem in infinite loop space theory [27].

To define R -unital plain actads, we will define, simultaneously, the *R-unital plain n -base* $B_n^R \supset B_n$ with maps

$$\begin{aligned} F_n^R : B_n^R &\rightarrow T_{B_{n-1}^R}^R \\ G_n^R : B_n^R &\rightarrow B_{n-1}^R \end{aligned}$$

and composition \circ^n . Here T_X^R is defined the same way as T_X (recall Definition 5), except that an empty sequence is allowed. In fact, there will be a distinguished subset

$$B_n^{R,0} \subseteq B_n^R$$

such that

$$x \in B_n^{R,0} \Rightarrow m_x = 0.$$

Now an *R-unital plain actad* \mathcal{C} is defined the same as a unital plain n -actad with B_n replaced by B_n^R , together with a section

$$B_n^{R,0} \rightarrow \mathcal{C}$$

over B_{n-1}^R (using G_n^R).

Then B_{n+1}^R is defined as the free plain R -unital n -actad over $Id : B_n^R \rightarrow B_n^R$, modulo the relations

$$(44) \quad \gamma_{n,i}(x, y) = x \circ_i^n y, \text{ for } x \in B_n^R, y \in B_n^{R,0}.$$

One defines, inductively,

$$(45) \quad B_{n+1}^{R,0} = B_n^{R,0} \coprod \{1_x^n \mid x \in B_{n-1}^R\}$$

(we want to begin the induction with $B_0^R = \{*, *\}$, $B_1^R = \mathbb{N}_0$). Note that m_x in B_{n+1} does not necessarily equal m_x in B_n . We emphasize that in the second term on the right hand side of (45) we mean the R -unital n -actad units, *not* the units $1_x^n \in B_n^R$.

The reason to keep these units separate is to be able to plug into 1_x^n , in \circ^{n+1} : we have $1_x^n \in B_{n+1}^R$ via the inclusion $B_n^R \subset B_{n+1}^R$ with $m_{1_x^n} = 1$. To explain this, in our definition of trees in B_2^R , there is no

reason to prohibit plugging in a tree with 1 prong into the triangle with one prong (“lozenge”) 1_1^1 via \circ^2 . However, plugging in the R-unit 1_1^1 via \circ^2 eliminates the lozenge from the tree, which is a new operation.

The structure of B_n^R is not as complicated as it may seem. In fact, one readily proves by induction

Lemma 14. *The canonical inclusion $B_n \subseteq B_n^R \setminus B_n^{R,0}$ is a bijection.*

□

Now \mathcal{B}_n^R , *R-unital n-actads*, (and all associated functors) can be defined completely analogously to, \mathcal{B}_n , *n-actads*, via an inductive definition of \mathcal{B}_n^R (and the associated functors) and then replacing \mathcal{B}_n^R while extending the formula (37) to *S*-morphisms, with $S \subseteq \{1, \dots, n\}$. This is then a generalization of (unital) *n-actads*.

In this approach, we keep permutations as the $\{n\}$ -morphisms; it is possible to create a larger category generated by these morphisms and *n*-compositions with the various *i*-units, $i < n$, but then we no longer have an invertible cube-like *n*-fold category.

Note that an R-unital 1-actad is the same thing as a based operad. 1-morphisms in \mathcal{B}_2 together with $? \circ_i^2 1_1^2$, $? \circ_i^2 1_0^1$ (with the obvious compatibility relations, noting that $B_2^{R,0} = \{1_1^2, 1_0^1\}$), generate a larger category of trees, whose opposite is equivalent to a tree analogue of a category of finite sets and injections.

Now we can define higher analogues of a based algebra over a based operad: An *R-unital algebra* over an R-unital *n-actad* \mathcal{C} is a fibered $(n - 1)$ -fold category

$$X \rightarrow \mathcal{B}_{n-1}^R$$

together with a section $(n - 1)$ -fold functor

$$\begin{array}{ccc} & & X \\ & \nearrow & \downarrow \\ \mathcal{B}_{n-1}^{R,0} & \xrightarrow{\subset} & \mathcal{B}_{n-1}^R \end{array}$$

which satisfies equivariance and associativity (and unit) properties analogous to those described in Section 4, in addition to the R-unitality axiom, which states that if

$$(y, (x_1, \dots, x_m)) \in S - Mor(\mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X)$$

with $x_i = \iota \in S - Mor(\mathcal{B}_{n-1}^{R,0})$, $S \subseteq \{1, \dots, n - 1\}$, then

$$\Theta(y, (x_1, \dots, x_m)) = \Theta(\gamma_{n,i}(y, \iota), (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_m)).$$

(recall that Θ denotes the functor mapping $\mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X \rightarrow T_X$ in (30)).

8. MONADS

A key point of the infinite loop space theory of May [27] is the close connection between operads and monads. A monad (for an introduction see [7], Vol. 2, Ch.4) in a category A is a functor $C : A \rightarrow A$ which satisfies monoid-like properties with respect to composition. Its *algebra* is an object X together with a natural transformation $\Theta : CX \rightarrow X$ which satisfies left module-like axioms with respect to composition.

For a universal algebra in the category of sets, its associated monad C assigns to a set X the free algebra CX of the given type on X . For an operad \mathcal{C} (in the unbased context), the associated monad C has a particularly simple form [27]:

$$CX = \coprod_{n \geq 1} \mathcal{C}(n) \times_{\Sigma_n} X^n$$

(on the right hand side, the subscript denotes taking orbits).

In this section, we will describe the monads associated to n -actads and plain n -actads. Let us discuss the plain case first. Let \mathcal{C} be a plain n -actad. In this case, we define a monad C in the category of sets over B_{n-1} . (In particular, we are dealing with multisorted algebra in the sense of [6].) Consider a set $\Xi : X \rightarrow B_{n-1}$ over B_{n-1} . Again, we write, for $y \in B_{n-1}$, $X(y)$ for $\Xi^{-1}(y)$. Then, for $y \in B_{n-1}$,

$$CX(y) := \coprod_{z \in B_n, G_n(z)=y} \prod_{i=1}^{m_z} X(f_i(z)).$$

In the case of an n -actad \mathcal{C} , we will construct the monad C describing \mathcal{C} -algebras in the category of $(n-1)$ -fold categories fibered over \mathcal{B}_{n-1} . We could, alternatively, describe a monad in sets over B_{n-1} , considering the $\{i\}$ -morphisms as unary operations. While this point of view also seems interesting, we do not follow it in the present paper.

Suppose \mathcal{C} is an n -actad. Let X be an $(n-1)$ -fold category with a fibered $(n-1)$ -fold functor

$$\Xi : X \rightarrow \mathcal{B}_{n-1}.$$

(Thus, X is an invertible cube-like $(n-1)$ -fold category.) Now we will construct the free \mathcal{C} -algebra CX on X . There are n -fold fibered functors

$$\begin{aligned} T_\Xi : T_X &\rightarrow T_{\mathcal{B}_{n-1}} \\ \Lambda : \mathcal{C} &\rightarrow T_{\mathcal{B}_{n-1}}. \end{aligned}$$

We can get a Λ since there is an n -fold fibered functor

$$\mathcal{C} \rightarrow \mathcal{B}_n,$$

and we can compose with

$$\mathcal{F}_n : \mathcal{B}_n \rightarrow T_{\mathcal{B}_{n-1}}.$$

Suppose $S \subseteq \{1, \dots, n-1\}$. Then first define an $(n-1)$ -fold category C_0X with an $(n-1)$ -fold functor $\Phi : C_0X \rightarrow \mathcal{B}_{n-1}$ by

$$S - \text{Mor}(C_0X) = S - \text{Mor}(\mathcal{C} \times_{T_{\mathcal{B}_{n-1}}} T_X)/$$

$$/(x, y) \sim (\tilde{f}(x), \hat{f}(y)), \text{ where } x \in S - \text{Mor}(\mathcal{C}), y \in S - \text{Mor}(T_X)$$

$$T_{\Xi}(y) = \Lambda(x) = \alpha \in T_{\mathcal{B}_{n-1}}, (f : \alpha \rightarrow \beta) \in S - \text{Mor}(T_{\mathcal{B}_{n-1}}),$$

for some β , and \tilde{f} is the lift of f using Λ , and

\hat{f} is the lift of f using T_{Ξ} .

The trouble is that the $(n-1)$ -fold functor Φ satisfies the uniqueness, but not the existence properties of a fibered $(n-1)$ -fold functor, due to the same property of the $(n-1)$ -fold functor

$$\mathcal{G}_n : \mathcal{B}_n \rightarrow \mathcal{B}_{n-1}.$$

Let \mathcal{A} be a k -fold category. The forgetful functor

$\mathcal{U}_{\mathcal{A}} : \text{fibered } k\text{-fold categories over } \mathcal{A} \rightarrow k\text{-fold categories over } \mathcal{A}$

(on both sides, k -fold functors are morphisms) has a left adjoint $K_{\mathcal{A}}$.

Definition 15. For $k = n-1$, $\mathcal{A} = \mathcal{B}_{n-1}$, the monad C in $(n-1)$ -fold categories fibered over \mathcal{B}_{n-1} associated with an n -actad \mathcal{C} is given by

$$CX = K_{\mathcal{B}_{n-1}}C_0X.$$

Then CX as defined is the free \mathcal{C} -algebra on X .

Now, one can also make a generalization of this construction to iterated algebras. Suppose \mathcal{C} is an n -actad. Then we have that $\mathcal{C}^{\alpha_1, \dots, \alpha_k}$ is a multisorted n -actad over $\{1, \dots, k\}$ (by which we mean a n -actad in the category of sets over $\{1, \dots, k\}$). From this point of view, we can just repeat the above construction over $\{1, \dots, k\}$.

More explicitly, let $X = (X_1, \dots, X_k)$ be a k -tuple of $(n-1)$ -fold categories with a fibered $(n-1)$ -fold functor

$$\Xi_j : X_j \rightarrow \mathcal{B}_{n-1}.$$

Now let X also be an $(n-1)$ -fold category, where the objects are k -tuples of objects of X_j 's, in the same order, and the S -morphisms are

k -tuples of S -morphisms of the X_j 's also in the same order. So, we also have an $(n-1)$ -fold fibered functor

$$\Xi = (\Xi_1, \dots, \Xi_k) : X \rightarrow \mathcal{B}_{n-1} \times \{1, \dots, k\}.$$

(Thus, X is an invertible cube-like $(n-1)$ -fold category.) Now, we will define the free $\mathcal{C}^{\alpha_1, \dots, \alpha_k}$ category $C^{\alpha_1, \dots, \alpha_k} X$ on X .

Then, there are n -fold, fibered functors

$$\begin{aligned} T_\Xi &: T_X^k \rightarrow T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k \\ \Lambda &: \mathcal{C}^{\alpha_1, \dots, \alpha_k} \rightarrow T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k. \end{aligned}$$

The construction Λ is as follows. First, by definition, we have that

$$\mathcal{C}^{\alpha_1, \dots, \alpha_k} = \mathcal{C} \times_{\mathcal{B}_n} \mathcal{B}_n^{\alpha_1, \dots, \alpha_k}.$$

Since we have that \mathcal{C} is an n -actad in the normal sense, we have a fibered n -fold functor

$$\mathcal{C} \rightarrow \mathcal{B}_n.$$

Then, we would need a fibered m -fold functor for every m , for every $k \leq m$ with some $1 \leq \alpha_1 < \dots < \alpha_k \leq m$,

$$h_m : \mathcal{B}_m^{\alpha_1, \dots, \alpha_k} \rightarrow \mathcal{B}_m \times \{1, \dots, k\} :$$

For $m = 1$, we have that k must equal 1. In addition, we also must have $\alpha_1 = 1$. However, for $k = 1$, we have that $\mathcal{B}_1^1 = \mathcal{B}_1$. Then we have the identity, which is a fibered functor,

$$\mathcal{B}_m^{\alpha_1, \dots, \alpha_k} \rightarrow \mathcal{B}_m = \mathcal{B}_m \times \{1, \dots, k\}.$$

Suppose are given this data for $m-1$. Fix some $1 \leq \alpha_1 < \dots < \alpha_k \leq m$. If $\alpha_k < m-1$, then we have

$$(46) \quad \mathcal{B}_m^{\alpha_1, \dots, \alpha_k} \xrightarrow{\mathcal{F}_m^{\alpha_1, \dots, \alpha_k}} T_{\mathcal{B}_{m-1}^{\alpha_1, \dots, \alpha_k}} \longrightarrow T_{\mathcal{B}_{m-1}} \times \{1, \dots, k\}$$

(where the second functor of (46) is given by applying h_{m-1} to the entries of $T_{\mathcal{B}_{m-1}^{\alpha_1, \dots, \alpha_k}}$ and taking the maximal element of $\{1, \dots, k\}$ for the second coordinate, as in (37)). If $\alpha_k = m$, then we have

$$\mathcal{B}_M^{\alpha_1, \dots, \alpha_{k-1}, m} \xrightarrow{\mathcal{F}_M^{\alpha_1, \dots, \alpha_{k-1}, n}} T_{\mathcal{B}_{m-1}}^{\bullet, \alpha_1, \dots, \alpha_{k-1}}.$$

Now, by Definition (10), we have

$$T_{\mathcal{B}_{m-1}}^{\alpha_1, \dots, \alpha_{k-1}} \rightarrow \{1, \dots, k\}$$

and we can construct

$$T_{\mathcal{B}_{m-1}}^{\alpha_1, \dots, \alpha_{k-1}} \rightarrow T_{\mathcal{B}_{m-1}}^{\alpha_1, \dots, \alpha_{k-1}} \rightarrow T_{\mathcal{B}_{n-1}}$$

where the first map is the forgetful functor and the second map is induced by the natural functor

$$\mathcal{B}_{m-1}^{\alpha_1, \dots, \alpha_{k-1}} \rightarrow \mathcal{B}_{m-1}.$$

This, we can construct a fibered functor

$$(47) \quad \mathcal{B}_m^{\alpha_1, \dots, \alpha_k} \rightarrow T_{\mathcal{B}_{m-1}} \times \{1, \dots, k\}$$

Then we can use the fact that

$$\mathcal{F}_m : \mathcal{B}_m \rightarrow T_{\mathcal{B}_{m-1}}.$$

is a fibered functor, to get a lift of every S -morphism of $T_{\mathcal{B}_{m-1}}$ to \mathcal{B}_m . Using this lifting by composing it with the fibered functor of (46) or (47), we get a fibered functor

$$h_m : \mathcal{B}_m^{\alpha_1, \dots, \alpha_k} \rightarrow \mathcal{B}_m \times \{1, \dots, k\}.$$

(This concludes the inductive definition of h_m .)

At the moment, we use this for $m = n - 1$. We have

$$h_{n-1} : \mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_k} \rightarrow \mathcal{B}_{n-1} \times \{1, \dots, k\}.$$

Then we have

$$\mathcal{C}^{\alpha_1, \dots, \alpha_k} \longrightarrow \mathcal{B}_n \times_{\mathcal{B}_n} \mathcal{B}_n^{\alpha_1, \dots, \alpha_k} = \mathcal{B}_n^{\alpha_1, \dots, \alpha_k}$$

Then, we will compose with $\mathcal{F}_n^{\alpha_1, \dots, \alpha_k}$ to get a map

$$\mathcal{B}_n^{\alpha_1, \dots, \alpha_k} \rightarrow T_{\mathcal{B}_{n-1}^{\alpha_1, \dots, \alpha_k}} \rightarrow T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k.$$

Using this, we get our Λ .

We still want the free iterated \mathcal{C} -algebra on X which we denote by $C^{\alpha_1, \dots, \alpha_k} X$ to be an $(n - 1)$ -fold category. So, again, suppose $S \subseteq \{1, \dots, n - 1\}$. Then define first $C_0^{\alpha_1, \dots, \alpha_k} X$ by

$$S - \text{Mor}(C_0^{\alpha_1, \dots, \alpha_k} X) = S - \text{Mor}(\mathcal{C}^{\alpha_1, \dots, \alpha_k} \times_{T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k} T_X^k) /$$

$/(x, y) \sim (\tilde{f}(x), \hat{f}(y))$, where $x \in S - \text{Mor}(\mathcal{C}^{\alpha_1, \dots, \alpha_k})$, $y \in S - \text{Mor}(T_X^k)$

$$T_{\Xi}(y) = \Lambda(x) = \gamma \in T_{\mathcal{B}_{n-1}}, (f : \gamma \rightarrow \delta) \in S - \text{Mor}(T_{\mathcal{B}_{n-1} \times \{1, \dots, k\}}^k),$$

for some δ , and \tilde{f} is the lift of f using Λ , and

$$\hat{f} \text{ is the lift of } f \text{ using } T_{\Xi}^k.$$

Then we have

$$C^{\alpha_1, \dots, \alpha_k} X = K_{\mathcal{B}_{n-1} \times \{1, \dots, k-1\}} C_0^{\alpha_1, \dots, \alpha_k} X.$$

Now these are analogous unital and R-unital versions of these concepts imposing additional identifications for the units, analogous to the concept of based algebras over an operad [27]. We omit the details.

9. CASE STUDY OF A 2-ACTAD ALGEBRA

When applying operads to topology, we do not, of course, work in the base category of sets, but rather in topological spaces or simplicial sets. All our definitions in this setting work essentially without change. In particular, geometric realizations of fibered n -fold functors are still fibered. In the case of spaces, we require all structure maps to be continuous. In the case of simplicial sets, we can think of a simplicial object in any of the structures we considered so far, which is considered as a category where morphisms are homomorphisms (i.e. maps preserving all structure). This makes sense, since all the structures we considered can be axiomatized as (multisorted) universal algebras. In particular, functors of singular sets and geometric realizations linking the structures in the topological and simplicial context exist and behave in the usual way. One source of examples is the Čech resolution. For a space X , EX denotes the simplicial space

$$EX : \Delta^{Op} \rightarrow Spaces,$$

with $EX_n = X^{n+1}$ (faces are given by taking an $(n+1)$ -tuple of elements of X to an n -tuple that is the original $(n+1)$ -tuple missing a coordinate, and degeneracies are given by taking an n -tuple to an $(n+1)$ -tuple which is the same except with one of the coordinates copied twice). We often identify EX with its geometric realization $|EX|$.

The Čech resolution tends to preserve algebraic structures, since the geometric realization of simplicial spaces preserves finite products (at least when we are working in the compactly generated category [33]). For example, considering the associative n -actad we introduced in Section 4, we obtain the n -actad $E\mathcal{A}_n$, which we call the E_∞ - n -actad. For example, $E\mathcal{A}_1$ is the Barratt-Eccles operad [26]. As shown in [26, 27], the classifying space BA of a permutative category A is an $E\mathcal{A}_1$ -algebra, hence an E_∞ -space. (Recall that the classifying space (or nerve) of a category is the simplicial space where BA_n consists of composable n -tuples of morphisms in A , meaning objects for $n=0$, faces are given by compositions or forgetting the first or last morphism,

and degeneracies are insertions of identities [27]). For a group G considered as a category with one object, we have an identification

$$BG \cong EG/G$$

where G acts diagonally on the right hand side. The bijection is given by homogenization of coordinates

$$(g_1, \dots, g_n) \mapsto (1, g_1, g_1g_2, \dots, g_1 \cdots g_n).$$

A permutative category is a symmetric monoidal category that is strictly associative where the commutativity isomorphism is a strict involution.

Recall the n -actad \mathcal{A}_n described in Example 1 at the end of Section 4. One may therefore ask what interesting $E\mathcal{A}_n$ -algebras there are, what they mean in homotopy theory, and how they arise. In this paper, I give some examples for $n = 2$. Note, however, that for any n , inclusion of the simplicial 0-stage gives a morphism of n -actads

$$(48) \quad \mathcal{A}_n \rightarrow E\mathcal{A}_n.$$

Therefore, $E\mathcal{A}_n$ -algebras are, in particular, $(n - 1)$ -actads, and we can study their algebras. For example, $E\mathcal{A}_2$ -algebras are, in fact, operads.

In this section, we will study the free $E\mathcal{A}_2$ -algebra on a given groupoid $X \rightarrow \mathcal{B}_1$, and in the simplest non-trivial case of X (the free “binary” case) will give a condition on a category which guarantees that its classifying space is an algebra over the free $E\mathcal{A}_2$ -algebra $A_{E\mathcal{A}_2}X$ on X . In some sense, this can be thought of as an $E\mathcal{A}_2$ analogue of the results of [26] on the structure of the Barratt-Eccles operad.

In a way, we can think of the operad $A_{E\mathcal{A}_2}X$, for a groupoid X , as describing algebras with operations from X which have commutativity properties given by X . However, these operations do not have any associativity properties. Arranging these operations in a planar tree (as in the reverse Polish notation on some old calculators), the $E\mathcal{A}_2$ -algebra property says that the order of the operations “does not matter in the E_∞ -sense.” We see from this example that, even for $n = 2$, which is the next stage beyond the well-studied case of $n = 1$, the world of n -actads becomes very rich.

To say things more precisely, suppose \mathcal{A} is a 2-actad. Then suppose X is a groupoid with a fibered functor

$$X \rightarrow \mathcal{B}_1.$$

Then, as noted in Section 8 above, the free \mathcal{A} -algebra on X is the operad

$$A_{\mathcal{A}}X(n) = \Sigma_n \times_{1-Mor(\mathcal{B}_2)} \coprod_{T \in B_2, m_{G_2}(T)=n} (\mathcal{A}(T) \times_{Aut_2(T)} \prod_{i=1}^{m_T} X(f_i(T))).$$

For $T \in B_2 = Obj(\mathcal{B}_2)$, we have that $E\mathcal{A}_2(T) = \{T\} \times Aut_2(T)$ (where $Aut_2(T) = \{2\} - Mor(T, T)$). So, for every X ,

$$\begin{aligned} A_{E\mathcal{A}_2}X(n) &= \\ \Sigma_n \times_{1-Mor(\mathcal{B}_2)} \coprod_{T \in B_2, m_{G_2}(T)=n} \{T\} \times Aut_2(T) \times_{Aut_2(T)} \prod_{i=1}^{m_T} X(f_i(T)) &= \\ \Sigma_n \times_{1-Mor(\mathcal{B}_2)} \coprod_{T \in B_2, m_{G_2}(T)=n} \{T\} \times \prod_{i=1}^{m_T} X(f_i(T)) &= \\ \Sigma_n \times_{1-Mor(\mathcal{B}_2)} \coprod_{T \in B_2, m_{G_2}(T)=n} \prod_{i=1}^{m_T} X(f_i(T)). \end{aligned}$$

As an example of this, let, for a set S , X_2^S be the groupoid fibered over \mathcal{B}_1

$$X_2^S(n) = \emptyset, \text{ if } n \neq 2,$$

and let

$$X_2^S(2) = S \times \Sigma_2$$

be the free Σ_2 -set on S . Then let \mathcal{C} be the underlying operad of the free $E\mathcal{A}_2$ -algebra on X_2^S . Then

$$\mathcal{C}(n) \cong B_2(n-1)_{\text{binary}} \times \Sigma_n \times B(\Sigma_{n-1}^S),$$

where Σ_{n-1}^S is the groupoid given by Σ_{n-1} acting on S^{n-1} , and $B_2(n)_{\text{binary}}$ are the elements of $B_2(n)$ which are of the form

$$((2, 2, \dots, 2), (i_1, i_2, \dots, i_{n-1}))$$

for some n . (Note that $B_2(n)_{\text{binary}}$ can also be interpreted as the set of binary trees with $n+1$ leaves.)

Let Y be a groupoid with a map

$$\Xi : S \rightarrow \text{Funct}(Y \times Y, Y).$$

Let $proj_i^2 : S \times S \rightarrow S$ be the coordinate projections for $i \in \{1, 2\}$, $proj_i^3 : S \times S \times S \rightarrow S$ be the coordinate projections for $i \in \{1, 2, 3\}$. When the source is obvious, we will simply write $proj_i$. We also have maps

$$\chi_\ell : S \times S \rightarrow \text{Funct}(Y \times Y \times Y, Y)$$

$$\chi_r : S \times S \rightarrow \text{Func}(Y \times Y \times Y, Y)$$

given by $\chi_\ell = (\Xi \circ \text{proj}_2) \circ ((\Xi \circ \text{proj}_1) \times \text{Id})$ and $\chi_r = (\Xi \circ \text{proj}_1) \circ (\text{Id} \times (\Xi \circ \text{proj}_2))$. Note that we apply the argument only to proj_2 and proj_1 , e.g.

$$\chi_\ell((s, t)) = (\Xi(t)) \circ (\Xi(s) \times \text{Id})$$

(we will continue to use this convention later). Then let $T : Y \times Y \rightarrow Y \times Y$ be the functor that switches coordinates. We also have maps induced by permutations of operations

$$\chi_{\ell,\ell}, \chi_{r,\ell}, \chi_{\ell,r}, \chi_{r,r}, \chi_u : S \times S \times S \rightarrow \text{Func}(Y \times Y \times Y \times Y, Y)$$

which are given by

$$\chi_{\ell,\ell} = (\Xi \circ \text{proj}_3) \circ (\Xi \circ \text{proj}_2 \times \text{Id}) \circ (\Xi \circ \text{proj}_1 \times \text{Id} \times \text{Id})$$

$$\chi_{r,\ell} = (\Xi \circ \text{proj}_3) \circ (\Xi \circ \text{proj}_2 \times \text{Id}) \circ (\text{Id} \times \Xi \circ \text{proj}_1 \times \text{Id})$$

$$\chi_{\ell,r} = (\Xi \circ \text{proj}_3) \circ (\text{Id} \times \Xi \circ \text{proj}_2) \circ (\text{Id} \times \Xi \circ \text{proj}_1 \times \text{Id})$$

$$\chi_{r,r} = (\Xi \circ \text{proj}_3) \circ (\text{Id} \times \Xi \circ \text{proj}_2) \circ (\text{Id} \times \text{Id} \times \Xi \circ \text{proj}_1)$$

$$\chi_u = (\Xi \circ \text{proj}_2) \circ ((\Xi \circ \text{proj}_1) \times (\Xi \circ \text{proj}_3)).$$

Consider the following additional structure for a groupoid Y :

(1) We have isomorphisms

$$(49) \quad \alpha_\ell : \chi_\ell \rightarrow \chi_\ell \circ T,$$

$$(50) \quad \alpha_r : \chi_r \rightarrow \chi_r \circ T$$

with the following commuting diagrams:

$$\begin{array}{ccc} \chi_\ell & \xrightarrow{\alpha_\ell} & \chi_\ell \circ T \\ & \searrow & \downarrow \alpha_\ell \circ T \\ & & \text{Id}_{\chi_\ell \circ T \circ T} \\ & & \downarrow = \\ & & \chi_\ell \end{array} \quad \begin{array}{ccc} \chi_r & \xrightarrow{\alpha_r} & \chi_r \circ T \\ & \searrow & \downarrow \alpha_r \circ T \\ & & \text{Id}_{\chi_r \circ T \circ T} \\ & & \downarrow = \\ & & \chi_r \end{array}$$

- (2) The following diagram (and the ones similar to it, with a different $\chi_{?,?}$'s) commute:

$$\begin{array}{ccc}
\chi_{\ell,\ell} & \xrightarrow{Id} & \chi_{\ell,\ell} \\
(\Xi \circ \text{proj}_3)(\alpha_\ell \times Id) \downarrow & & \downarrow \alpha_\ell \circ (\Xi \circ \text{proj}_1) \times Id \times Id \\
\chi_{\ell,\ell} \circ (T \times Id) & & \chi_{\ell,\ell} \circ (Id \times T) \\
\alpha_\ell \circ (\Xi \circ \text{proj}_1) \times Id \times Id \downarrow & & \downarrow (\Xi \circ \text{proj}_3)(\alpha_\ell \times Id) \\
\chi_{\ell,\ell} \circ (Id \times T) \circ (T \times Id) & & \chi_{\ell,\ell} \circ (T \times Id) \circ (Id \times T) \\
(\Xi \circ \text{proj}_3)(\alpha_\ell \times Id) \downarrow & & \downarrow \alpha_\ell \circ (\Xi \circ \text{proj}_1) \times Id \times Id \\
\chi_{\ell,\ell} \circ (T \times Id) \circ (Id \times T) \circ (T \times Id) & \xrightarrow{=} & \chi_{\ell,\ell} \circ (T \times Id) \circ (Id \times T) \circ (T \times Id)
\end{array}$$

- (3) The following diagram commutes:

$$\begin{array}{ccc}
\chi_u & \xrightarrow{Id} & \chi_u \\
\alpha_\ell \circ (Id \times Id \times (\Xi \circ \text{proj}_3)) \downarrow & & \downarrow \alpha_r \circ ((\Xi \circ \text{proj}_1) \times Id \times Id) \\
\chi_u \circ (T \times Id) & & \chi_u \circ (Id \times T) \\
\alpha_r \circ ((\Xi \circ \text{proj}_1) \times Id \times Id) \downarrow & & \downarrow \alpha_\ell \circ (Id \times Id \times (\Xi \circ \text{proj}_3)) \\
\chi_u \circ (Id \times T) \circ (T \times Id) & & \chi_u \circ (T \times Id) \circ (Id \times T) \\
\alpha_\ell \circ (Id \times Id \times (\Xi \circ \text{proj}_3)) \downarrow & & \downarrow \alpha_r \circ ((\Xi \circ \text{proj}_1) \times Id \times Id) \\
\chi_u \circ (T \times Id) \circ (Id \times T) \circ (T \times Id) & \xrightarrow{=} & \chi_u \circ (Id \times T) \circ (T \times Id) \circ (Id \times T).
\end{array}$$

Then we claim the following

Theorem 16. *Given the structure described the above axioms (1)-(3), the classifying space on morphisms BY is an algebra over the operad $A_{E\mathcal{A}_2}(X_2^S)$.*

The Theorem will be proved at the end of this Section. To prove the Theorem, we first will prove some results about $B_2(n)_{\text{binary}}$.

As a warm-up case, consider the usual presentation of the symmetric group:

$$(51) \quad \Sigma_n \cong \langle a_1, \dots, a_{n-1} | a_i^2, a_i a_{i+1} a_i a_{i+1} a_i a_{i+1}, a_i a_j a_i a_j \ j > i + 1 \rangle.$$

Here a_i are understood as switches of consecutive terms in an n -element sequence. This can be proved for 4 directly from using the Cayley

graph. Then, one uses induction and the isotropy groups of elements of $\{1, \dots, n\}$ in the standard action of Σ_n to prove it for every n .

In the case of an element of $B_2(n)_{\text{binary}}$, the terms are the triangles (we shall say “nodes”) of a binary tree. Thus, they do not naturally form a sequence but could be thought of as forming themselves a tree where edges are between triangles which are attached. For a given binary tree, we will present the group on its nodes in terms of generators which correspond to such edges (of which there are $n - 1$).

Concretely, we claim

Theorem 17. *We have*

$$(52) \quad \begin{aligned} 2 - \text{Mor}(\mathcal{B}_2(n)_{\text{binary}}) &\cong \langle a_1, \dots, a_{n-1} | a_i^2, \\ &a_i a_j a_i a_j a_i a_j \text{ when } i \neq j \text{ and } a_i \text{ and } a_j \text{ share a node,} \\ &(a_i a_j a_k a_j)^2 \text{ when } i < j < k \text{ and } a_i, a_j, a_k \text{ share a node,} \\ &a_i a_j a_i a_j \text{ when } a_i \text{ and } a_j \text{ do not share a node} \rangle. \end{aligned}$$

(Here we denote by $2 - \text{Mor}(\mathcal{B}_2(n)_{\text{binary}})$ the full subcategory of $2 - \text{Mor}(\mathcal{B}_2)$ on binary trees.)

Comment: When a_i, a_j, a_k share a node with i, j, k different, the relation $(a_i a_j a_k a_j)^2$ follows from the relations (52) regardless of the order of i, j, k . This is by the fact that the relations (52) involving a_i, a_j, a_k generate a copy of Σ_4 , which follows by examining the Cayley graph.

Proof. If $n = 4$, this can be proved drawing the Cayley graph. For $n \geq 5$, suppose we have a binary tree

$$((x_1, \dots, x_k), (i_1, \dots, i_{k-1})) = ((2, \dots, 2), (i_1, \dots, i_{k-1})).$$

For lower n , assume the claim as an induction hypothesis. Then suppose we can fix $x_{i-1}, x_i, x_{i+1}, x_{i+2}, x_{i+3}$ with x_i connected to both x_{i+1} and x_{i+2} . Then let a, b, c, d be the 2-morphisms that switch x_{i-1} and x_i , x_i and x_{i+2} , x_{i+2} and x_{i+3} and x_i and x_{i+1} respectively. In a picture, this looks like

$$(53) \quad \begin{array}{ccccccc} & & x_{i-1} & \xrightarrow{a} & x_i & \xrightarrow{b} & x_{i+2} & \xrightarrow{c} & x_{i+3} \\ & & \bullet & & \bullet & & \bullet & & \bullet \\ & & & & \downarrow d & & \nearrow \text{bdb} & & \\ & & & & \bullet & & & & \\ & & & & x_{i+1} & & & & \end{array}$$

where the dots represent the x_j 's. Let

$$(54) \quad G_i = \langle a, b, c, d | a^2, b^2, c^2, d^2, (ab)^3, (bc)^3, (ad)^3, (db)^3, (daba)^2, (ac)^2, (cd)^2 \rangle$$

(in our notation here, we rename the x_i 's after each switch, making the above compositions possible). Then let $G_{i,1}$ be the subgroup of G_i that is generated by a, b, d . Let $G_{i,2}$ be the subgroup of G_i that is generated by bdb, b, c .

$$\kappa_i : G_{i,1} \rightarrow G_{i,2}$$

with

$$(55) \quad \begin{aligned} \kappa_i(a) &= b \\ \kappa_i(b) &= c \\ \kappa_i(d) &= bdb. \end{aligned}$$

So, in other words, κ_i shifts x_i to x_{i+2} and takes the permutations around x_i to those around x_{i+2} .

First, note that by the case $n = 4$, $G_{i,1}$ is the symmetric group, and also can be presented only in the relations (54) including a, b, d . We want κ_i to be an isomorphism, so we have to check the relations of G_i for a, b , and d for $\kappa_i(a), \kappa_i(b)$, and $\kappa_i(d)$. It is obvious that $(\kappa_i(a))^2 = b^2$, $(\kappa_i(b))^2 = c^2$ are 1. We have

$$(\kappa_i(d))^2 = (bdb)^2 = bddb = bb = 1.$$

Next, we have

$$(\kappa_i(a)\kappa_i(d))^3 = (bbdb)^3 = (db)^3 = 1.$$

Next, we can commute d and c since they permute separate pairs of x_j 's.

$$\begin{aligned} (\kappa_i(d)\kappa_i(a)\kappa_i(b)\kappa_i(a))^2 &= \\ bdbbcbdbbcb &= bdcacb = bddccb = bb = 1. \end{aligned}$$

Now we need to check the Yang-Baxter relations. It is obvious that

$$\kappa_i(a)\kappa_i(b)\kappa_i(a)\kappa_i(b)\kappa_i(a)\kappa_i(b) = bcbcbc = 1.$$

We have

$$\begin{aligned} (\kappa_i(d)\kappa_i(b))^3 &= (bdb)^3 = \\ = bd(bcbd)^2bc &= bd(cbcd)^2bc = bd(cbdc)^2bc = \\ bdc(bd)^2cbc &= bc(d(bd)^2)c = (bc)^3 = 1 \end{aligned}$$

So, we have every relation. Thus, κ_i is a homomorphism. On the other hand, all of the relations (54) are valid in the symmetric group, and thus the image of κ_i must be a symmetric group and κ_i must be an isomorphism.

Now suppose we have a binary tree in $B_2(n)_{\text{binary}}$. Represent its triangles as nodes x_i , and edges a_i between them where the triangles are attached. Consider a degree three node which is farthest from the root. Using the move (53), we can switch one edge in a way that this node moves farther from the root, and the relations for the new tree are equivalent to the relations of the old tree. This is because all the relations which change occur in presentations of Σ_4 subgroups, which stay isomorphic, as we just proved. (In the end, we also need a variant of the move (53) where the edge c is absent, but that case is clear since we already know the group is Σ_4).

By repeating these moves, the tree can be turned into a sequence, in which case we are reduced to (51). □

Proof of Theorem 16. Analogous to the argument of [26] that for a permutative category A , BA is an $E\mathcal{A}_1$ -algebra. Consider n composable p -tuples of Y -morphisms

$$\begin{array}{ccccccc} X_{1,0} & \xrightarrow{f_{1,1}} & X_{1,1} & \xrightarrow{f_{1,2}} & \dots & \xrightarrow{f_{1,p}} & X_{1,p} \\ & & \vdots & & \vdots & & \\ X_{n,0} & \xrightarrow{f_{n,1}} & X_{n,1} & \xrightarrow{f_{n,2}} & \dots & \xrightarrow{f_{n,p}} & X_{n,p}. \end{array}$$

Choose a tree $T \in B_2(n)_{\text{binary}}$ with $p + 1$ decorations T_0, \dots, T_p of its nodes by elements of S . Apply the operations corresponding to T_i to the objects $X_{1,i}, \dots, X_{n,i}$, and apply the same operations, composed with the appropriate composition of the coherences (49),(50), to the morphisms $f_{1,i}, \dots, f_{n,i}$. The coherence diagrams (1)-(3) (given in this section, directly before the statement of Theorem 16) guarantee consistency of these compositions, by Theorem 17, thereby describing an $A_{E\mathcal{A}_2}(X_2^S)$ -action on BA . We construct maps

$$A_{E\mathcal{A}_2}X_2^S(n) \times (BY)^n \rightarrow BY.$$

□

10. EXAMPLES OF NEW OPERADS

In this section, I will describe a method for using $E\mathcal{A}_2$ algebras to construct new examples of operads.

First, suppose X is an operad. Then we have a functor

$$X \rightarrow \mathbb{N},$$

(where the morphisms in \mathbb{N} are identities) since we have a projection functor $\mathcal{B}_1 \rightarrow \mathbb{N}$.

Then define, for $m, n \in \mathbb{N}, k \in \mathbb{N}_0$,

$$S_k(n, m) = \{(T_0, \dots, T_k) \mid T_j \in B_2, m = m_{T_j}, G(T_j) = n, \\ \text{there exists permutations } \sigma_j : \{1, \dots, m\} \rightarrow \{1, \dots, m\} \\ \text{such that, for } j = 1, \dots, k, f_i(T_0) = f_{\sigma_j(i)}(T_j)\}.$$

This means that if $(T_0, \dots, T_k) \in S_k(n, m)$, then the $f_i(T_j)$'s are the same as the $f_i(T_0)$'s, up to permutation. Then we can define monads M_k in the category of spaces over \mathbb{N} of the form

$$M_k X(n) = \coprod_m \coprod_{(T_0, \dots, T_k) \in S_k(n, m)} \underbrace{\Sigma_n \times \dots \times \Sigma_n}_{k+1} \times \prod_{i=1}^{m_{T_0}} X(f_i(T_0)),$$

for $k \in \mathbb{N}_0$. The Σ_n factors correspond to permutations of “prongs” of the trees T_0, \dots, T_k (as would arise in the free operad on \mathcal{B}_1 ; see Sections 4, 8). Monad composition is defined by 2-composition of trees following the order of the natural numbers $1, \dots, m_{T_0}$ in the 0'th coordinate, and the order specified by the permutation on $S_k(n, m)$ in the other coordinates. This 2-composition is “twisted” by the 1-permutation of prongs in the Σ_n factors (meaning that the successors are switched according to the permutation of the prongs of the inserted tree). The permutations in the resulting $S_k(n, m)$, as well as in the Σ_n factors, are then determined as appropriate versions of the wreath product (to match entries which were originally matched). This is obviously associative. We observe that for $k = 0$, algebras over this monad are just operads in the category of spaces, i.e. the monad takes $\coprod X(n) \times \Sigma_n$, and then applies the monad defining operads in monoids fibered over \mathcal{B}_1 , as considered in Section 4, 8.

Now let

$$(56) \quad \mathcal{M}_k(\ell_1, \dots, \ell_m; n) = \Sigma_m \times \coprod_{(T_0, \dots, T_k) \in S_k(n, m), \ell_i = m_{f_i(T_0)}} \underbrace{\Sigma_n \times \dots \times \Sigma_n}_{k+1}.$$

Obviously, by counting prongs, this is only non-empty when

$$(57) \quad n = \ell_1 + \dots + \ell_m - m + 1.$$

Again, the Σ_n 's come from the permutation parts of the T_i 's. As above, we have $(k + 1)$ copies of Σ_n because we have $(k + 1)$ T_j 's encoding permutations of "prongs." The extra Σ_m is so we have n (and not $n - 1$) 2-permutations. We have, for every k ,

$$\mathcal{M}_k = \underbrace{\mathcal{M}_0 \times \dots \times \mathcal{M}_0}_{k+1},$$

since the k extra Σ_m 's on the right hand side corresponds to the $S_k(n, m)$ (instead of $S_0(n, m)$) on the bottom of the coproduct in (56). Let

$$\mathcal{M} := \mathcal{M}_0.$$

Now, \mathcal{M} is an \mathbb{N} -sorted operad (or multi-category, see [13]) and, therefore, so is \mathcal{M}_n . In addition, we see that M_k is actually the monad associated with \mathcal{M}_k (Note, again, that the extra Σ_m in (56) prevents extra identifications). \mathcal{M}_0 -algebra are exactly operads, which are \mathcal{A}_2 -algebras.

In fact, every \mathcal{M} -algebra is the sequence of spaces (i.e., space over \mathbb{N}) of an \mathcal{A}_2 -algebra (i.e. operads). Thus, every $E\mathcal{M}$ -algebra is the underlying sequence of spaces of an $E\mathcal{A}_2$ -algebra. We should explain that this last statement is not 'if and only if.' The reason is that the 2-permutations in \mathcal{M}_k (or M_k) can occur between the "triangles" of different trees, T_i, T_j . While $m_{T_i} = m_{T_j} = m$, we have no functorial correspondence between the "prongs" of the trees T_i, T_j . Therefore, if we allowed these more general 2-permutations in 2-actads, there would be no consistent way of defining \circ^2 .

Now suppose we have a spectrum E (for example in the sense of May, e.g., see [26, 27, 28, 29], i.e. a sequence of spaces $E_n, n \in \mathbb{N}_0$, with homeomorphisms

$$E_n \cong \Omega E_{n+1}$$

where $\Omega E_{n+1} = \text{Map}_{\text{based}}(S^1, E_{n+1})$ denotes the based loop space). Suppose we have a morphism of spectra

$$E \rightarrow H\mathbb{Z}.$$

Then we have a map

$$(58) \quad E_0 \xrightarrow{\varphi} \mathbb{Z},$$

with $E_0(n) = \varphi^{-1}(n - 1)$. Now E_0 is an algebra over \mathcal{C}_∞ , the ∞ -little cubes operad. Now, let \mathcal{C} be an \mathbb{N} -sorted operad with

$$\mathcal{C}(\ell_1, \dots, \ell_m, n) = \mathcal{C}_\infty(m)$$

when (57) holds, and $\mathcal{C}(\ell_1, \dots, \ell_m, n) = \emptyset$ else. Then $(E_0(n))_{n \in \mathbb{N}}$ is a \mathcal{C} -algebra.

So, we have projections

$$\begin{array}{ccc} \mathcal{C} \times E\mathcal{M} & \xrightarrow{p_2} & E\mathcal{M} \\ p_1 \downarrow & & \\ \mathcal{C} & & \end{array}$$

Then let N be the monad associated with $\mathcal{C} \times E\mathcal{M}$. By (56), the symmetric group action on \mathcal{M} is free, so $E\mathcal{M}$ is an E_∞ -operad since the Čech resolution of a non-empty space is always (simplicially) contractible. Then the 2-sided bar construction of monads

$$B(EM, N, (E_0(n))_{n \in \mathbb{N}}) \simeq (E_0(n))_{n \in \mathbb{N}}$$

(see [27]) is an $E\mathcal{M}$ -algebra. So it is an $E\mathcal{A}_2$ -algebra. The same construction also works for a map of E_∞ -spaces $X \rightarrow \mathbb{Z}$ in place of E_0 . (Recall that infinite loop space of spectra are group-like E_∞ -spaces.) Thus, we have

Proposition 18. *Suppose*

$$X \rightarrow \mathbb{Z}$$

is a map of E_∞ -spaces. Then there exists an $E\mathcal{A}_2$ -algebra Ξ_X equivalent to $(X(n))_{n \in \mathbb{N}}$.

□

We will now prove that in some cases, these $E\mathcal{A}_2$ -algebras are non-trivial as operads (see (48)) in the sense that an E_∞ -operad does not map into them.

Let X be the free $E\mathcal{A}_1$ -algebra on $*$, with

$$X(n) = B\Sigma_{n-1}$$

(which comes with a natural Σ_n -action). Now, we will write

$$X_n = X(n+1) = B\Sigma_n = E\Sigma_n \times_{\Sigma_n} *$$

so that, in particular, $X_0 = X(1)$.

Theorem 19. *There does not exist an E_∞ -operad \mathcal{C} together with an operad morphism*

$$(59) \quad \mathcal{C} \rightarrow \Xi_X.$$

Proof. Suppose such a \mathcal{C} does exist. Since Ξ_X is an operad in spaces, we have a $\mathbb{Z}/2$ -equivariant map

$$(60) \quad \varsigma : \Xi_X(2) \times (\Xi_X(2) \times \Xi_X(2)) \rightarrow \Xi_X(4),$$

where on the right hand side, $\mathbb{Z}/2 \rightarrow \Sigma_4$ acts by sending the generator of $\mathbb{Z}/2$ to the permutation (13)(24). For $\mathbb{Z}/2$ -spaces Z and T , where Z has a trivial $\mathbb{Z}/2$ -action, we have an equivalence

$$Z \times_{\mathbb{Z}/2} (T \times EZ/2) \simeq Z \times (EZ/2 \times_{\mathbb{Z}/2} T),$$

or equivalently

$$(61) \quad (Z \times EZ/2) \times_{\mathbb{Z}/2} T \simeq Z \times (EZ/2 \times_{\mathbb{Z}/2} T)$$

By (61), we have

$$\varrho : (EZ/2 \times X_1) \times_{\mathbb{Z}/2} (X_1 \times X_1) \xrightarrow{\cong} X_1 \times (EZ/2 \times_{\mathbb{Z}/2} (X_1 \times X_1)).$$

We also have a map

$$\nu : EZ/2 \times_{\mathbb{Z}/2} X_3 \rightarrow EZ/2 \times_{EZ/2} X_3,$$

since, by construction, the Σ_n -action on $X(n)$ extends, for every n , to an $E\Sigma_n$ -action. Let

$$\mu = \nu \circ (EZ/2 \times \varsigma) \circ \varrho^{-1}.$$

So, we have the following diagram:

$$\begin{array}{ccc} (EZ/2 \times X_1) \times_{\mathbb{Z}/2} (X_1 \times X_1) & \xrightarrow{EZ/2 \times \varsigma} & EZ/2 \times_{\mathbb{Z}/2} X_3 \\ \downarrow \varrho & & \downarrow \nu \\ X_1 \times (EZ/2 \times_{\mathbb{Z}/2} (X_1 \times X_1)) & & EZ/2 \times_{EZ/2} X_3 \\ & \searrow \mu & \downarrow = \\ & & X_3 \end{array}$$

Additionally, by construction, the map μ comes from the E_∞ -algebra (concretely, \mathcal{C}_∞ -algebra) structure of X . Now, take homology of μ with coefficients in $\mathbb{Z}/2$, i.e.

$$H_k(\mu) : H_k(X_1 \times (EZ/2 \times_{\mathbb{Z}/2} (X_1 \times X_1))) \rightarrow H_k(X_3).$$

Now, let α be the generator of $H_0(X_1) = \mathbb{Z}/2$ and let e_k be the generator of $H_k(B\mathbb{Z}/2) \cong \mathbb{Z}/2$. Then we have

$$(62) \quad \alpha \otimes (e_k \otimes \alpha \otimes \alpha) \mapsto \alpha \cdot Q^k \alpha \neq 0.$$

where Q^k denotes the Dyer-Lashof operation (see [25, 11]).

Now, suppose we have a map of operads (59). Then we have the commutative diagram

$$\begin{array}{ccc} \mathcal{C}(4) & \longrightarrow & X(4) \\ \downarrow & & \downarrow \\ E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathcal{C}(4) & \longrightarrow & E\mathbb{Z}/2 \times_{\mathbb{Z}/2} X(4) \end{array}$$

commutes. Then the diagram

$$\begin{array}{ccc} \mathcal{C}(4) & \longrightarrow & X(4) \\ \downarrow & & \downarrow \\ E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathcal{C}(4) & \longrightarrow & E\mathbb{Z}/2 \times_{\mathbb{Z}/2} X(4) \\ \downarrow & \searrow & \downarrow \\ E\Sigma_4 \times_{\Sigma_4} \mathcal{C}(4) & & E\mathbb{Z}/2 \times_{E\mathbb{Z}/2} X(4) \\ \downarrow & & \downarrow \\ E\Sigma_4 \times_{\Sigma_4} X(4) & \longrightarrow & X(4) \\ \downarrow & & \downarrow \\ E\Sigma_4 \times_{\Sigma_4} X(4) & \longrightarrow & E\Sigma_4 \times_{E\Sigma_4} X(4) \end{array}$$

commutes. In particular, we have a diagram of the form

$$\begin{array}{ccc} E\mathbb{Z}/2 \times_{\mathbb{Z}/2} \mathcal{C}(4) & \longrightarrow & X(4) \\ \downarrow & \nearrow & \\ E\Sigma_4 \times_{\Sigma_4} \mathcal{C}(4), & & \end{array}$$

in other words,

$$\begin{array}{ccc} B\mathbb{Z}/2 & \longrightarrow & B\Sigma_3 \\ \downarrow & \nearrow & \\ B\Sigma_4 & & \end{array}$$

commutes. So, by taking π_1 , we get a commutative diagram of groups

$$\begin{array}{ccc}
\mathbb{Z}/2 & \xrightarrow{f} & \Sigma_3 \\
\downarrow g & \nearrow h & \\
\Sigma_4 & &
\end{array}$$

By definition, g takes the generator of $\mathbb{Z}/2$ to $\sigma = (13)(24)$, f is an inclusion (by (62)), and h is a group homomorphism. Then $f = h \circ g$. We have that $\sigma = \sigma_1 \circ \sigma_2$, where $\sigma_1 = (13)$ and $\sigma_2 = (24)$. Since h is a group homomorphism, it takes σ to $h(\sigma_1) \circ h(\sigma_2)$, and $h(\sigma_1)$, $h(\sigma_2)$ are conjugate in Σ_3 , since σ_1 , σ_2 are conjugate in Σ_4 . Thus, $h(\sigma_1)$, $h(\sigma_2)$ are both non-trivial. However, a non-trivial involution in Σ_3 is not a composition of two non-trivial involutions. Contradiction. \square

Let me add a few words on the topological interpretation of Ξ_X -algebras where Ξ_X is the operad associated with an E_∞ -space $X = (X_n)_{n \geq 0}$. First, let us discuss the case when $X = E_0$ for a spectrum E with a morphism of (connective) spectra

$$(63) \quad E \rightarrow H\mathbb{Z}.$$

This gives a morphism of E_∞ -spaces $\varphi : E_0 \rightarrow \mathbb{Z}$, and as above, we adopt the convention

$$X(n) = X_{n-1} = \varphi^{-1}(n-1),$$

for $n \in \mathbb{Z}$. In particular, X_0 is an E_∞ -space (over \mathbb{Z}) which is the 0-space of the fiber \tilde{E} of the morphism of spectra (63). While we set up our formalism in such a way that 0 is not included in the indexing of an operad, completely analogously, one can also construct a based operad $\bar{\Xi}_X$ indexed by $n \in \mathbb{N}_0$, with

$$\bar{\Xi}_X(0) \sim X_{-1}.$$

For Ξ_X -algebra Y , $Y \amalg X_{-1}$ is canonically an $\bar{\Xi}_X$ -algebra. Now we already mentioned the fact that when (63) splits, then $\bar{\Xi}_X$ -algebras are equivalent to E_∞ -spaces Y together with E_∞ -maps

$$\tilde{E}_0 \rightarrow Y.$$

For a general morphism (63), a splitting can be achieved by replacing E with the homotopy pushout \check{E} of the diagram

$$\begin{array}{ccc}
\tilde{E} & \longrightarrow & E \\
\downarrow & & \\
E & &
\end{array}$$

Then, X gets replaced with \check{X} where for $n \in \mathbb{Z}$, $\check{X}(n) = X = E_0$. Then, a $\bar{\Xi}_{\check{X}}$ -algebra Y is equivalent to a morphism of E_∞ -algebra $X \rightarrow Y$. However, if we denote the monads associated with the operads we considered by replacing the symbol Ξ with Θ , then for a $\bar{\Xi}_X$ -algebra Y ,

$$(64) \quad \check{Y} = B(\bar{\Theta}_{\check{X}}, \bar{\Theta}_X, Y).$$

gives a diagram of E_∞ -spaces

$$(65) \quad \begin{array}{ccc} X & \longrightarrow & \check{Y} \\ & \searrow & \downarrow \psi \\ & & \mathbb{Z}. \end{array}$$

Moreover, we get a canonical equivalence

$$\psi^{-1}(-1) \xrightarrow{\sim} Y.$$

Thus, we see that $\bar{\Xi}_X$ -algebras Y , where $\pi_0 Y$ is a group, are canonically identified, up to equivalence, with fibers of -1 in diagrams of E_∞ -spaces of the form (65). Such spaces are, of course, homotopically equivalent to the fibers $Y' = \psi^{-1}(0)$, which are equivalent to E_∞ -morphisms

$$X_0 \rightarrow Y'.$$

However, when (63) does not split, the homotopy equivalence $Y \simeq Y'$ is not canonical.

Finally, let us comment what happens when the E_∞ -space X is not the 0-space of a spectrum (i.e. is not group complete). Then $\bar{\Xi}_X$ does not make sense as above, and as the Ξ_X -algebra Y is necessarily unbased. However, if X^g is the infinite loop space associated with X (i.e. its group completion) then applying $B(\Theta_{X^g}, \Theta_X, ?)$ in the above sense gets us into the above situation. In particular, the π_0 of $Y_+ = Y \amalg \{*\}$ is still canonically a monoid, and its group completion is canonically equivalent to $\psi^{-1}(-1)$ in the diagram (65), where X is replaced with X^g .

11. ORDINALS

In this section, we will consider the plain non-unital n -base B_n only. Let Ord denote the set of ordinals. It is well known that ordinal numbers $0 < \alpha < \varepsilon$ can be written uniquely as a sum

$$\alpha = \omega^{\alpha_1} + \cdots + \omega^{\alpha_k},$$

for some $\alpha > \alpha_1 \geq \dots \geq \alpha_k$. Now, by definition, every tree in B_2 can be written as

$$(66) \quad z = \gamma_{1,i_\ell}(\dots \gamma_{1,i_2}(\gamma_{1,i_1}(z_0, z_1), z_2) \dots, z_\ell)$$

with $i_1 > \dots > i_\ell$, $z_0 \in B_1 = \mathbb{N}$, $z_1, \dots, z_\ell \in B_2$ (so all the z_1, \dots, z_ℓ are plugged directly into z_0). So if we define inductively a function $\Phi_2 : B_2 \rightarrow Ord$ by $\Phi_2(n) = n$ for $n \in \mathbb{N}$,

$$\begin{aligned} \Phi_2(z) = & (i_1 - 1) + \omega^{\Phi_2(z_1)} + (i_2 - i_1 - 1) + \omega^{\Phi_2(z_2)} + \dots \\ & + (i_\ell - i_{\ell-1} - 1) + \omega^{\Phi_2(z_\ell)} + (z_0 - i_\ell), \end{aligned}$$

then Φ_2 is onto ε . We could identify a subset on B_2 on which Φ_2 is bijective, but it will not be bijective on all of B_2 because of order of summation, and also non-cancellation, for example $1 + \omega = \omega$.

It is therefore natural to investigate how this fact generalizes to $n > 2$. We will focus on the surjectivity part here. For $n > 2$, we will introduce more arguments to the function

$$\Phi_n : B_n \rightarrow Ord,$$

for the purpose of induction.

To be more precise, let $\varphi_\alpha : Ord \rightarrow Ord$ be the functions with

$$\varphi_1(\beta) = \omega^{1+\beta},$$

and $\varphi_\alpha(\beta) =$ the β 'th fixed point of all $\varphi_{\alpha'}$'s where $\alpha' < \alpha + 1$. The φ_α 's are the Veblen functions with the subscripts shifted by 1, for $\alpha \geq 2$. For every $\alpha \in Ord$, φ_α is continuous and strictly increasing.

We will now define functions

$$\Phi_n : B_n \rightarrow Ord$$

inductively.

For $x \in B_1$, $\alpha_1, \dots, \alpha_{m_x} \in Ord$, define

$$\Phi_1(x; \alpha_1, \dots, \alpha_{m_x}) = \alpha_1 + \dots + \alpha_{m_x}.$$

Now, suppose we have defined $\Phi_{n-1}(z; \alpha_1, \dots, \alpha_{m_z})$, for every $z \in B_{n-1}$, $\alpha_1, \dots, \alpha_{m_z} \in Ord$. Fix $z \in B_n$, $\alpha_1, \dots, \alpha_{m_z} \in Ord$. We can always write

$$\begin{aligned} z = & ((z_0, z_1, \dots, z_\ell), (i_1, \dots, i_\ell)) = \\ = & \gamma_{n-1,i_\ell}(\gamma_{n-1,i_{\ell-1}}(\dots \gamma_{n-1,i_2}(\gamma_{n-1,i_1}(z_0, z_1), z_2), \dots), z_\ell) \end{aligned}$$

with $i_1 > \dots > i_\ell$, $z_0 \in B_{n-1}$, and $z_1, \dots, z_\ell \in B_n$. Let

$$\beta_{i,j} = \alpha_{1+m_{z_1}+\dots+m_{z_{i-1}+j}}.$$

Define inductively

$$\begin{aligned} \Phi_n(z; \alpha_1, \dots, \alpha_{m_z}) := & \\ \Phi_{n-1}(z_0; \underbrace{1, \dots, 1}_{m_{z_1}-1}, \varphi_{n-1}(\Phi_n(z_1; \beta_{1,1}, \dots, \beta_{1,m_{z_1}})), \underbrace{1, \dots, 1}_{m_{z_2}-1}, & \\ \varphi_{n-1}(\Phi_n(z_2; \beta_{2,1}, \dots, \beta_{2,m_{z_2}})), \underbrace{1, \dots, 1}_{m_{z_3}-1}, \dots, & \\ \underbrace{1, \dots, 1}_{m_{z_\ell}-1}, \varphi_{n-1}(\Phi_n(z_\ell; \beta_{\ell,1}, \dots, \beta_{\ell,m_{z_\ell}}))). & \end{aligned}$$

Note that not all of the arguments get used in the definition. However, I do not think that eliminating this “wastefulness” would extend the range of ordinals expressible by B_n .

Theorem 20. *Fix an n . Then for every $\beta \in \text{Ord}$, there exists some $z \in B_n$ and some $\alpha_i \in \{1\} \cup \text{Im}(\varphi_n)$ such that*

$$\beta = \Phi_n(z; \alpha_1, \dots, \alpha_{m_z}).$$

Proof. Induction. Suppose $n=1$. Then

$$\Phi_1(n; \alpha_1, \dots, \alpha_n) = \alpha_1 + \dots + \alpha_n.$$

Then for every $\beta \in \text{Ord}$, there exists an n and $\alpha_1, \dots, \alpha_n \in \{1\} \cup \{\omega^\alpha \mid \alpha \in \text{Ord}\} = \{1\} \cup \text{Im}(\varphi_1)$ such that $\beta = \alpha_1 + \dots + \alpha_n$.

Now, suppose for every $\beta \in \text{Ord}$ there is a $z \in B_{n-1}$ and $\alpha_1, \dots, \alpha_{m_z} \in \{1\} \cup \text{Im}(\varphi_{n-1})$ with

$$\beta = \Phi_{n-1}(z; \alpha_1, \dots, \alpha_{m_z}).$$

If $\alpha_1, \dots, \alpha_{m_z} = 1$, we are done. Suppose $\alpha_i = \varphi_{n-1}(\alpha'_i)$. If $\alpha_i = \alpha'_i$, then α_i is a fixed point of φ_{n-1} . So there exists an α''_i with

$$\alpha_i = \varphi_n(\alpha''_i).$$

So we are done. Suppose $\alpha'_i < \alpha_i$. Then we use the induction hypothesis on α'_i . \square

By Theorem 20, the image of the function

$$\Phi_n : B_n \rightarrow \text{Ord}$$

given by $\Phi_n(x) := \Phi_n(x; 1, \dots, 1)$ is $\varphi_n(0)$, for $n = 1, 2, \dots$

Comment 21. *It is readily possible to extend the description of B_n (and, in addition, \mathcal{B}_n) to B_α , (and \mathcal{B}_α) for $\alpha \in \text{Ord}$, by taking unions for limit ordinals α . Then also our definition of Φ_n extends to Φ_α , $\alpha \in \text{Ord}$. Therefore, the first ordinal γ for which*

$$\text{Im}(\Phi_\gamma) = \gamma$$

is the Feferman-Schütte ordinal $\gamma = \Gamma_0$. This can be compared with [22], where the authors consider the similar concept of opetopes and possible machine implementations.

REFERENCES

- [1] J. Adamek, J. Rosicky: *Locally presentable and accessible categories*, LMS Lecture Notes 189, Cambridge U. Press, 1994
- [2] J. F. Adams: *Stable Homotopy and Generalized Cohomology*, Univ. of Chicago Press, Chicago, Il. 1974
- [3] J. C. Baez, J. Dolan: Higher-dimensional algebra III. n -categories and the algebra of opetopes, *Adv. Math.* 135 (1998), no. 2, 145-206
- [4] C. Bagen, I. Moerdijk: On the derived category of an algebra over an operad, arXiv:0801.2664
- [5] J. Bénabou: *Introduction to Bicategories*, Lecture Notes in Mathematics 47, Springer (1967), pp. 1-77
- [6] G. Birkhoff, J. Lipson: Heterogeneous algebras, *J. Comb. Theory* 8 (1970) 115-133
- [7] F. Borceux: *Handbook of Categorical Algebra*, Cambridge University Press, 1994
- [8] E. Cheng: The category of opetopes and the category of opetopic sets *Theory and Appl. of Categories*, 11 (2003) 353-374
- [9] E. Cheng: Weak n -categories: opetopic and multitopic foundations, *J. Pure Appl. Algebra* 186 (2004), no. 2, 109-137
- [10] E. Cheng: Weak n -categories: comparing opetopic foundations, *J. Pure Appl. Algebra* 186 (2004), no. 3, 219-231
- [11] F. R. Cohen, T. J. Lada, J.P.May: *The homology of iterated loop spaces*, Lecture Notes in Mathematics, Vol. 533. Springer-Verlag, Berlin-New York, (1976), vii+490 pp.
- [12] P.-L. Curien, C. H. Thanh, S. Mimram: Syntactic approaches to opetopes, arXiv:1903.05848
- [13] A. D. Elmendorf, M. A. Mandell: Rings, modules, and algebras in infinite loop space theory, *Adv. Math.* 205 (2006), no. 1, 163-228

- [14] E. Finster: Implementing the Opetopes: Programming with Higher Dimensional Trees, preprint, 2018
- [15] M. Fiore, P. Saville: List objects with algebraic structure. 2nd International Conference on Formal Structures for Computation and Deduction, Art. No. 16, 18 pp., LIPIcs. Leibniz Int. Proc. Inform., 84, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2017.
- [16] N. Gambino, M. Hyland: Wellfounded trees and dependent polynomial functors, in: S. Berardi, M. Coppo, F. Damiani (Eds.), TYPES 2003, in: *Lecture Notes in Comput. Sci.*, 3085, Springer-Verlag, Heidelberg, (2004), 210-225
- [17] V. Ginzburg, M. Kapranov: Koszul duality for operads, *Duke Math. J.* 76 (1994), no. 1, 203-272
- [18] C. Hermida, M. Makkai, J. Power: On weak higher dimensional categories. I. 1. Category theory and its applications (Montreal, QC, 1997). *J. Pure Appl. Algebra* 154 (2000), no. 1-3, 221-246.
- [19] C. Hermida, M. Makkai, J. Power: On weak higher-dimensional categories. I. 2. *J. Pure Appl. Algebra* 157 (2001), no. 2-3, 247-277.
- [20] C. Hermida, M. Makkai, J. Power: On weak higher-dimensional categories. I. 3. *J. Pure Appl. Algebra* 166 (2002), no. 1-2, 83-104.
- [21] V. A. Hinich, V. V. Schechtman: On homotopy limit of homotopy algebras, *K-theory, arithmetic and geometry*, 240-264, Lecture Notes on Math. 1289, Springer, Berlin, 1987
- [22] J. Kock, A. Joyal, M. Batanin, J.-F. Mascari: Polynomial functors and opetopes, *Adv. Math.* 224 (2010), no. 6, 2690-2737
- [23] T. Leinster: *Higher operads, higher categories*, London Mathematical Society Lecture Note Series, Vol. 298. Cambridge University Press, Cambridge, (2004), xiv+433 pp.
- [24] M. A. Mandell: E_∞ algebras and p-adic homotopy theory, *Topology* 40 (2001), no. 1, 43-94
- [25] J. P. May: A general algebraic approach to Steenrod operations, *The Steenrod Algebra and its Applications* (Proc. Conf. to Celebrate N. E. Steenrod's Sixtieth Birthday, Battelle Memorial Inst., Columbus, Ohio, 1970) (F. P. Peterson, ed.), Lecture Notes in Mathematics, Vol. 168, Springer, Berlin, (1970), pp. 153-231
- [26] J. P. May: E_∞ -spaces, group completions and permutative categories, *London Math. Soc. Lecture Notes* 11 (1974) 61-94

- [27] J. P. May: *The geometry of iterated loop spaces*, Lectures Notes in Mathematics, Vol. 271. Springer-Verlag, Berlin-New York, (1972), viii+175 pp.
- [28] J. P. May: The spectra associated to permutative categories, *Topology* 17 (1978) 225-228
- [29] J. P. May: Multiplicative infinite loop space theory, *J. Pure Appl. Algebra* 26 (1982), no. 1, 1-69
- [30] T. Palm: Dendrotopic sets. Galois theory, Hopf algebras, and semiabelian categories, 411-461, Fields Inst. Commun., 43, Amer. Math. Soc., Providence, RI, 2004.
- [31] T. Palm: Dendrotopic sets for weak infinity-categories. Thesis (Ph.D.)–York University (Canada). 2003. 65 pp.
- [32] S. B. Priddy: Koszul resolutions, *Trans. A.M.S.* 152 (1970), 39-60
- [33] N. E. Steenrod: A convenient category of topological spaces, *Mich. Math. J.* 14 (1967) 133-152.
- [34] S. Szawiel, M. Zawadowski: Theories of analytic monads. *Math. Structures Comput. Sci.* 24 (2014), no. 6, e240604, 33 pp.
- [35] S. Szawiel, M. Zawadowski: The web monoid and opetopic sets, *J. Pure Appl. Algebra* 217 (2013), no. 6, 1105-1140
- [36] S. Szawiel, M. Zawadowski: Polynomial and Analytic Functors and Monads, revisited, arXiv:1506.04317
- [37] O. Veblen: Continuous Increasing Functions of Finite and Transfinite Ordinals, *Transactions of the American Mathematical Society*, 9 (1908): 280-292
- [38] M. Zawadowski: On ordered face structures and many-to-one computads, arXiv:0708.2659 [math.CT]
- [39] M. Zawadowski: On positive face structures and positive-to-one computads, arXiv:0708.2658 [math.GT]
- [40] M. Zawadowski: Lax monoidal fibrations. Models, logics, and higher-dimensional categories, 341-426, CRM Proc. Lecture Notes, 53, Amer. Math. Soc., Providence, RI, 2011.