

Improved regularity for the parabolic normalized p -Laplace equation

Pêdra D. S. Andrade, Makson S. Santos

August 20, 2021

Abstract

We derive regularity estimates for viscosity solutions to the parabolic normalized p -Laplace. By using approximation methods and scaling arguments for the normalized p -parabolic operator, we show that the gradient of bounded viscosity solutions is locally asymptotically Lipschitz continuous when p is sufficiently close to 2. In addition, we establish regularity estimates in Sobolev spaces.

Keywords: Parabolic normalized p -Laplace equation; Regularity theory; Estimates in Sobolev and Hölder spaces; Approximation methods.

MSC(2020): 35B65; 35K55; 35Q91.

1 Introduction

We study the regularity of viscosity solutions to the normalized p -Laplace equation

$$u_t(x, t) - \Delta_p^N u(x, t) = f(x, t) \quad \text{in } Q_1, \quad (1)$$

where $p \in (1, \infty)$, $f \in \mathcal{C}(Q_1) \cap L^\infty(Q_1)$ and $Q_1 := B_1 \times (-1, 0]$. We prove new and sharp regularity results for the viscosity solutions to (1), when the exponent p is *close* to 2. In particular, we obtain gains of regularity in Sobolev and Hölder spaces. We exploit the properties of the regularized equation

$$(u_\varepsilon)_t - \Delta u_\varepsilon - (p-2) \frac{D^2 u_\varepsilon D u_\varepsilon \cdot D u_\varepsilon}{|D u_\varepsilon|^2 + \varepsilon^2}, \quad (2)$$

to establish the integral estimates for the solutions of the homogeneous p -Laplace equation. For the regularity in Hölder spaces, we argue by approximation methods importing regularity from the *homogeneous heat equation* to equation (1).

The normalized p -Laplace operator is defined as

$$\begin{aligned}\Delta_p^N u &:= |Du|^{2-p} \Delta_p u = \Delta u + (p-2) \Delta_\infty^N u \\ &= \Delta u + (p-2) \left\langle D^2 u \frac{Du}{|Du|}, \frac{Du}{|Du|} \right\rangle,\end{aligned}$$

where $\Delta_p u := |Du|^{2-p} \operatorname{div}(|Du|^{p-2} Du)$ is the p -Laplace operator. Hence, the equation (1) can be seen as a uniformly parabolic model in nondivergence form, with constants $\min\{p-1, 1\}$ and $\max\{p-1, 1\}$. We also notice that this model has a singularity on the set $\{Du = 0\}$, which implies that we can not use directly the classic $C^{1,\alpha}$ regularity theory of viscosity solutions, as in [24, 25, 35, 19].

The normalized p -Laplace equation appears in many branches of mathematics, ranging from differential geometry to stochastic process. At the limit $p = 1$, the equation (1) is related to the level set formulation of the mean curvature flow in the work of L. C. Evans and J. Spruck [14]. For $1 < p < \infty$, Y. Peres and S. Sheffield in [28] introduced a stochastic approach for the normalized p -Laplace equation using the theory of tug-of-war games with noise. For the parabolic framework we refer to the work of J. J. Manfredi, M. Parviainen and J. D. Rossi [26]. As p approaches infinity, the equation (1) finds an application in image processing; see [12, 21]. It is worth noticing that [22, 13, 27] address the game theoretical characterization for the borderlines cases $p = 1$ and $p = \infty$.

The normalized p -Laplace equation has been extensively studied by many authors. The existence and uniqueness was established in [6, 10]. In particular, A. Banerjee and N. Garofalo in [6] and K. Does in [10] proved Lipschitz regularity with respect to the space variable. Since the model (1) is uniformly parabolic, Hölder regularity of the solutions follows from the Krylov-Safonov theory see, e.g. [23, 25].

In [20], T. Jin and L. Silvestre showed that solutions to the homogeneous case are of class $\mathcal{C}^{1+\alpha, \frac{1+\alpha}{2}}$. In that paper, the authors follow the strategy adopted in [18], for a class of fully nonlinear degenerate equations. For the equation with a bounded right-hand side, A. Attouchi and M. Parviainen in [3] also established that solutions are of class $\mathcal{C}^{1+\alpha, \frac{1+\alpha}{2}}$. More recently, the variable exponent case

was studied in [15], where Y. Fang and C. Zhang obtained Hölder regularity for the gradient of the solutions.

Besides the regularity in Hölder spaces, many authors have studied regularity estimates in Sobolev spaces. In [16], F. Høeg and P. Lindqvist proved $W_{loc}^{2,1;2}$ -regularity for the viscosity solutions, in the homogeneous setting, when $p \in (\frac{6}{5}, \frac{14}{5})$. This range for p comes from their own method, and it is not clear if it can be extended to $p \in (1, \infty)$ for $d \geq 3$. This result were improved by H. Dong, F. Peng, Y. Zhang and Y. Zhou in [11] where they proved that solutions are of class $W^{2,1;q}$ for $q < 2 + \delta_{n,p}$ with $\delta_{n,p} \in (0, 1)$, when $p \in (1, 2) \cup (2, 3 + \frac{2}{d-2})$.

The purpose of this paper is to study gains of the regularity of solutions to (1) when p is *close* to 2. In this scenario, our model can be seen as a perturbation of the heat equation, as in [30], and then we can implement the so-called *approximation methods*. The approximation methods were introduced by L. Caffarelli in the seminal paper [7] and has been applied in a more general settings in the works of E. Teixeira, J.M. Urbano and their collaborators. We refer to the reader the papers [1, 9, 33, 31, 32], just to cite a few. See also the survey [29].

We combine the approximation methods and intrinsic scaling of the p -parabolic operator to obtain the desired estimate in Hölder spaces. This is the content of our first result:

Theorem 1.1. *Let $u \in \mathcal{C}(Q_1)$ be a viscosity solution to (1), with $f \in \mathcal{C}(Q_1) \cap L^\infty(Q_1)$. Given $\alpha \in (0, 1)$, there exists $\varepsilon > 0$, to be determined later, such that if*

$$|p - 2| \leq \varepsilon,$$

then $u \in \mathcal{C}^{1+\alpha, \frac{1+\alpha}{2}}(Q_{1/2})$ and there exists a universal constant $C > 0$ such that

$$\|u\|_{\mathcal{C}^{1+\alpha, \frac{1+\alpha}{2}}(Q_{1/2})} \leq C (\|u\|_{L^\infty(Q_1)} + \|f\|_{L^\infty(Q_1)}).$$

Theorem 1.1 ensures that although solutions to (1) are of class $\mathcal{C}^{1+\beta, \frac{1+\beta}{2}}$ (β can be very small), the gradient of the solutions become almost Lipschitz continuous when p is sufficiently close to 2.

Remark 1.1. *The Theorem 1.1 can be extended to a more general class of equations, namely*

$$u_t(x, t) - |Du|^\gamma \Delta_p^N u(x, t) = f(x, t) \quad \text{in } Q_1, \quad (3)$$

provided

$$|\gamma| + |p - 2| \leq \varepsilon.$$

We refer to the reader to [17, 4, 2, 5] for more details.

Our next result concerns regularity in Sobolev spaces in the homogeneous setting

$$u_t(x, t) - \Delta_p^N u(x, t) = 0 \quad \text{in} \quad Q_1. \quad (4)$$

Here, we make use of the regularized equation (2) and the $W^{2,1;q}$ -regularity theory developed by Wang in [34] to prove our estimates. More precisely, we prove the following:

Theorem 1.2. *Let $u \in \mathcal{C}(Q_1)$ be a viscosity solution to (4). There exists $\varepsilon_1 > 0$, to be determined later, such that if*

$$|p - 2| \leq \varepsilon_1,$$

then $u \in W^{2,1;q}(Q_{1/2})$ for every $1 < q < \infty$. In addition, there exists a universal constant $C > 0$ such that

$$\|u\|_{W^{2,1;q}(Q_{1/2})} \leq C \|u\|_{L^\infty(Q_1)}.$$

The Theorem 1.2 provides higher integrability for the viscosity solutions of (4), with the trade off of losing the precise range on the values of p for which the estimate holds true.

Remark 1.2. *By Sobolev embeddings we observe that Theorem 1.2 is a stronger version of Theorem 1.1 for the case $f \equiv 0$, since ε_1 would not depend on the Hölder exponent, see Corollary 4.1.*

The remainder of this paper is organized as follows: In Section 2 we fix some notations and gather a few facts used throughout the paper. In Section 3, we establish an approximation that connects our model with the heat equation. The regularity in space is subject of Section 4. We conclude the paper with the time regularity in Section 5.

2 Notations and preliminary results

This section puts forward elementary notations and gathers a few results used in the paper.

2.1 Elementary notation

Initially, we introduce some standard notations which will be used throughout the paper. In what follows $B_r \subset \mathbb{R}^d$ denotes the *open ball* of radius r and centered at the origin. The *parabolic domain* is given by

$$Q_r := \{(x, t) \in \mathbb{R}^{d+1} : x \in B_r, t \in (-r^2, 0]\} \subset \mathbb{R}^{d+1}.$$

We define the *parabolic distance* between the points (x_1, t_1) and (x_2, t_2) by

$$d((x_1, t_1), (x_2, t_2)) := \sqrt{|x_1 - x_2|^2 + |t_1 - t_2|}.$$

Similarly, the distance between the sets U and V stands for

$$\text{dist}(U, V) := \inf\{d((x_1, t_1), (x_2, t_2)) : (x_1, t_1) \in U, (x_2, t_2) \in V\},$$

where U, V are subsets in \mathbb{R}^{n+1} .

Fix $1 \leq q \leq \infty$, the *parabolic Sobolev space* is defined as follows

$$W^{2,1;q}(Q_r) := \{u \in L^q(Q_r) : u_t, Du, D^2u \in L^q(Q_r)\}.$$

If $u \in W^{2,1;q}(Q_r)$, we define its norm to be

$$\|u\|_{W^{2,1;q}(Q_r)} = \left[\|u\|_{L^q(Q_r)}^q + \|u_t\|_{L^q(Q_r)}^q + \|Du\|_{L^q(Q_r)}^q + \|D^2u\|_{L^q(Q_r)}^q \right]^{\frac{1}{q}}.$$

We say that u belongs to $W_{loc}^{2,1;q}(Q_r)$, if $u \in W^{2,1;q}(Q')$ for every $Q' \Subset Q_r$, where $Q' \Subset Q_r$ means $\text{dist}(Q', \partial_p Q_r) > 0$, where ∂_p denotes the parabolic boundary of Q_r .

For $0 < \alpha < 1$, the *parabolic Hölder space* stands for $C^{\alpha, \frac{\alpha}{2}}(Q_r)$. We define its norm to be

$$\|u\|_{C^{\alpha, \frac{\alpha}{2}}(Q_r)} := \|u\|_{L^\infty(Q_r)} + [u]_{C^{\alpha, \frac{\alpha}{2}}(Q_r)},$$

where $[u]_{C^{\alpha, \frac{\alpha}{2}}(Q_r)}$ is the semi-norm denoted by

$$[u]_{C^{\alpha, \frac{\alpha}{2}}(Q_r)} := \sup_{\substack{(x_1, t_1), (x_2, t_2) \in Q_r \\ (x_1, t_1) \neq (x_2, t_2)}} \frac{|u(x_1, t_1) - u(x_2, t_2)|}{d((x_1, t_1), (x_2, t_2))^\alpha}.$$

We say that u is a α -Hölder continuous with respect to the spatial variable and $\frac{\alpha}{2}$ -Hölder continuous with respect to the temporal variable if its norm is finite.

Similarly, we say that $u \in \mathcal{C}^{1+\alpha, \frac{1+\alpha}{2}}(Q_r)$ if there exists the spatial gradient $Du(x, t)$ for every (x, t) in Q_r in the classical sense and its norm

$$\begin{aligned} \|u\|_{\mathcal{C}^{1+\alpha, \frac{1+\alpha}{2}}(Q_r)} &:= \|u\|_{L^\infty(Q_r)} + \|Du\|_{L^\infty(Q_r)} \\ &+ \sup_{\substack{(x_1, t_1), (x_2, t_2) \in Q_r \\ (x_1, t_1) \neq (x_2, t_2)}} \frac{|u(x_1, t_1) - u(x_2, t_2) - Du(x_1, t_1) \cdot (x_1 - x_2)|}{d((x_1, t_1), (x_2, t_2))^{1+\alpha}}. \end{aligned}$$

is finite. It means that Du is α -Hölder continuous and u is $\frac{1+\alpha}{2}$ -Hölder continuous with respect to the temporal variable. For more details see [8, 9] and the references therein.

Finally, we define the *oscillation* of the function u as

$$\text{osc}_{Q_1} u = \sup_{Q_1} u - \inf_{Q_1} u.$$

2.2 Preliminary notions

We start with the definition of viscosity solutions to (1).

Definition 2.1. *Let $1 < p < \infty$ and f is a continuous function in Q_1 . We say that $u \in \mathcal{C}(Q_1)$ is a viscosity subsolution to (1). If, for every $(x_0, t_0) \in Q_1$ and $\varphi \in \mathcal{C}^2(Q_1)$ such that $u - \varphi$ has a local maximum at (x_0, t_0) , we have*

$$\left\{ \begin{array}{ll} \varphi_t(x_0, t_0) - \Delta_p^N \varphi(x_0, t_0) \leq f(x_0, t_0), & \text{if } D\varphi(x_0, t_0) \neq 0, \\ \varphi_t(x_0, t_0) - \Delta\varphi(x_0, t_0) - (p-2)\lambda_{\max}(D^2\varphi(x_0, t_0)) \\ \leq f(x_0, t_0), & \text{if } D\varphi(x_0, t_0) = 0 \text{ and } p \geq 2, \\ \varphi_t(x_0, t_0) - \Delta\varphi(x_0, t_0) - (p-2)\lambda_{\min}(D^2\varphi(x_0, t_0)) \\ \leq f(x_0, t_0), & \text{if } D\varphi(x_0, t_0) = 0, 1 < p < 2. \end{array} \right.$$

Conversely, we say that $u \in \mathcal{C}(Q_1)$ is a viscosity supersolution to (1). If, $x_0 \in Q_1$ and $\varphi \in \mathcal{C}^2(Q_1)$ such that $u - \varphi$ has a local minimum at (x_0, t_0) , we have

$$\left\{ \begin{array}{ll} \varphi_t(x_0, t_0) - \Delta_p^N \varphi(x_0, t_0) \geq f(x_0, t_0), & \text{if } D\varphi(x_0, t_0) \neq 0, \\ \varphi_t(x_0, t_0) - \Delta\varphi(x_0, t_0) - (p-2)\lambda_{\min}(D^2\varphi(x_0, t_0)) \\ \geq f(x_0, t_0), & \text{if } D\varphi(x_0, t_0) = 0 \text{ and } p \geq 2, \\ \varphi_t(x_0, t_0) - \Delta\varphi(x_0, t_0) - (p-2)\lambda_{\max}(D^2\varphi(x_0, t_0)) \\ \geq f(x_0, t_0), & \text{if } D\varphi(x_0, t_0) = 0, 1 < p < 2. \end{array} \right.$$

Whenever u is a viscosity subsolution and supersolution to (1) in Q_1 , we say that u is a viscosity solution to the equation.

For a general definition of this notion, we refer to the reader [3]. In our arguments the scaled functions satisfy a variant of (1), namely:

$$u_t - \Delta u - (p-2) \left\langle D^2 u \frac{Du + \xi}{|Du + \xi|}, \frac{Du + \xi}{|Du + \xi|} \right\rangle = f \text{ in } Q_1, \quad (5)$$

where $\xi \in \mathbb{R}^d$ is arbitrary. On account of completeness, we proceed by stating a local compactness result used in this paper that ensures the convergence property for the sequences.

Lemma 2.1 (Compactness). *Let $u \in \mathcal{C}(Q_1)$ be a viscosity solution to (5). Then for all $r \in (0, 1)$, there exist constants $\beta \in (0, 1)$ and $C > 0$ such that if $\|f\|_{L^\infty(Q_1)} \leq 1$ and $\text{osc}_{Q_1} u \leq 1$ then we have*

$$\|u\|_{\mathcal{C}^{\beta, \beta/2}(Q_r)} \leq C.$$

For a proof of this result, we refer the reader to [3, Lemma 3.1]. We close this section with the scaling properties of our model. Throughout the paper, we require

$$\|u\|_{L^\infty(Q_1)} \leq 1 \quad \text{and} \quad \|f\|_{L^\infty(Q_1)} \leq \varepsilon, \quad (6)$$

for some ε to be determined. The conditions in (6) are not restrictive. In fact, consider the function

$$v(x, t) = \frac{u(\rho x, \rho^2 t)}{K},$$

for all $0 < \rho \ll 1$ and $K > 0$. Notice that v is also a viscosity solution to (1) in Q_1 , with the right-hand side

$$\tilde{f}(x, t) = \frac{\rho^2}{K} f(\rho x, \rho^2 t).$$

Hence, by choosing

$$K = \|u\|_{L^\infty(Q_1)} + \varepsilon^{-1} \|f\|_{L^\infty(Q_1)},$$

we can assume (6) without loss of generality.

3 Hölder estimates

This section is devoted to the proof of Theorem 1.1. As usual, constants stand for C may change from line to line, and depend only on the appropriate quantities.

3.1 Geometric Tangential Path

First, we provide an approximation lemma relating our model with the heat equation. This lemma plays a pivotal role in the paper.

Lemma 3.1 (Approximation Lemma). *Let $u \in \mathcal{C}(Q_1)$ be a normalized viscosity solution to (5) with $f \in L^\infty(Q_1) \cap \mathcal{C}(Q_1)$. Given $\delta > 0$ there exists $\varepsilon > 0$ such that, if*

$$\|f\|_{L^\infty(Q_1)} + |p - 2| < \varepsilon,$$

then we can find $h \in \mathcal{C}^{2,1}(Q_{7/9})$ such that

$$\sup_{Q_{7/9}} |u(x, t) - h(x, t)| < \delta.$$

Proof. We argue by contradiction. Suppose that the statement does not hold, then there are $\delta_0 > 0$ and sequences $(u_j)_{j \in \mathbb{N}}$, $(\xi_j)_{j \in \mathbb{N}}$, $(p_j)_{j \in \mathbb{N}}$ and $(f_j)_{j \in \mathbb{N}}$ satisfying

$$\|f_j\|_{L^\infty(Q_1)} + |p_j - 2| < \frac{1}{j},$$

$$(u_j)_t - \Delta u_j - (p_j - 2) \left\langle D^2 u_j \frac{Du_j + \xi_j}{|Du_j + \xi_j|}, \frac{Du_j + \xi_j}{|Du_j + \xi_j|} \right\rangle = f_j \text{ in } Q_1, \quad (7)$$

and for every $h \in \mathcal{C}^{2,1}(Q_{7/9})$ and for all $j \in \mathbb{N}$,

$$\sup_{(x,t) \in Q_{7/9}} |u_j(x, t) - h(x, t)| > \delta_0. \quad (8)$$

From Lemma 2.1, we have

$$u_j \in \mathcal{C}^{\beta, \frac{\beta}{2}}(Q_{8/9}) \quad \text{and} \quad \|u_j\|_{\mathcal{C}^{\beta, \frac{\beta}{2}}(Q_{8/9})} \leq C,$$

where $0 < \beta < 1$ and $C > 0$ are constants that do not depend on $j \in \mathbb{N}$. Applying the Arzelá-Ascoli Theorem, there exist a subsequence $(u_j)_{j \in \mathbb{N}}$ and a continuous function u_∞ so that $(u_j)_{j \in \mathbb{N}}$ converges uniformly to u_∞ in $Q_{8/9}$.

We now examine two different cases. We start by considering the case in which the sequence $(\xi_j)_{j \in \mathbb{N}}$ is bounded. Using local compactness of \mathbb{R}^d , up to a subsequence ξ_j converges to ξ_∞ . Evaluating the limit in (7) as j approaches infinity, we obtain that u_∞ satisfies

$$(u_\infty)_t - \Delta u_\infty = 0 \text{ in } Q_{8/9}. \quad (9)$$

Since u_∞ is a viscosity solution of (9), it follows that u_∞ is of class $\mathcal{C}^{2,1}$.

On the other hand, if the sequence $(\xi_j)_{j \in \mathbb{N}}$ is unbounded, we choose a subsequence ξ_j such that $|\xi_j|$ goes to infinity for every j . By taking $e_j := \frac{\xi_j}{|\xi_j|}$ we have $e_j \rightarrow e_\infty$, for some $e_\infty \in \mathbb{R}^d$. Hence, we may rewrite the equation (7) as follows:

$$(u_j)_t - \Delta u_j - (p_j - 2) \left\langle D^2 u_j \frac{Du_j |\xi_j|^{-1} + e_j}{|Du_j |\xi_j|^{-1} + e_j|}, \frac{Du_j |\xi_j|^{-1} + e_j}{|Du_j |\xi_j|^{-1} + e_j|} \right\rangle = f_j \text{ in } Q_1.$$

Applying the limit in the equation above, we get that u_∞ solves (9). Once again, we have that u_∞ belongs to $C^{2,1}(Q_{7/9})$. Finally, by taking $h = u_\infty$, we reach a contradiction. \square

3.2 Regularity in space

We proceed with the regularity in the spatial variable.

Proposition 3.1. *Let $u \in C(Q_1)$ be a normalized viscosity solution of*

$$u_t - \Delta u - (p - 2) \left\langle D^2 u \frac{Du + \xi}{|Du + \xi|}, \frac{Du + \xi}{|Du + \xi|} \right\rangle = f \text{ in } Q_1,$$

for any arbitrary vector $\xi \in \mathbb{R}^d$ and $f \in L^\infty(Q_1) \cap C(Q_1)$. Then, given $\alpha \in (0, 1)$, there exists $\varepsilon > 0$ such that

$$\|f\|_{L^\infty(Q_1)} + |p - 2| \leq \varepsilon,$$

we can find a constant $0 < \rho < 1$ and a sequence of affine functions $(\ell_n)_{n \in \mathbb{N}}$ of the form $\ell_n(x, t) := a_n + b_n \cdot x$ satisfying

$$\sup_{Q_{\rho^n}} |u(x, t) - \ell_n(x, t)| \leq \rho^{n(1+\alpha)},$$

$$|a_{n+1} - a_n| \leq C\rho^{n(1+\alpha)}$$

and

$$|b_{n+1} - b_n| \leq C\rho^{n\alpha}$$

for a constant $C > 0$ and for every $n \in \mathbb{N}$.

Proof. The result follows by induction argument. For simplicity, we split the proof into two steps.

Step 1. Take $\delta > 0$ to be determined later. Lemma 3.1 implies that there exists a function $h \in \mathcal{C}^{2,1}(Q_{8/9})$ satisfying

$$\sup_{Q_{7/9}} |u(x, t) - h(x, t)| \leq \delta.$$

Set

$$\ell(x, t) := h(0, 0) + Dh(0, 0) \cdot x.$$

Recall that $h(0, 0)$ and $Dh(0, 0)$ are uniformly bounded by a constant. Since $h \in \mathcal{C}^{2,1}(Q_{8/9})$, we have that there exists a constant $C > 0$ such that

$$\sup_{Q_\rho} |h(x, t) - \ell(x, t)| \leq C\rho^2,$$

where ρ is a parabolic distance between the points (x, t) and $(0, 0)$. As consequence from the triangular inequality, we obtain

$$\begin{aligned} \sup_{Q_\rho} |u(x, t) - \ell(x, t)| &\leq \sup_{Q_\rho} |u(x, t) - h(x, t)| + \sup_{Q_\rho} |h(x, t) - \ell(x, t)| \\ &\leq \delta + C\rho^2. \end{aligned}$$

Making universal choices, we define

$$\rho := \left(\frac{1}{2C}\right)^{\frac{1}{1-\alpha}} \quad \text{and} \quad \delta := \frac{\rho^{1+\alpha}}{2}.$$

Notice that the universal choice of δ determines the value of $\varepsilon > 0$ through the Lemma 3.1. Therefore

$$\sup_{Q_\rho} |u(x, t) - \ell(x, t)| \leq \rho^{1+\alpha}.$$

This completes the case $n = 1$.

Step 2. Assume that the case $n = k$ has been verified. We shall prove the case $n = k + 1$. First, let us introduce an auxiliary function $v_k : Q_1 \rightarrow \mathbb{R}^d$ defined by

$$v_k(x, t) := \frac{u(\rho^k x, \rho^{2k} t) - \ell_k(\rho^k x, \rho^{2k} t)}{\rho^{k(1+\alpha)}}.$$

Notice that v_k solves the following equation:

$$(v_k)_t - \Delta v_k - (p-2) \left\langle D^2 v_k \frac{Dv_k + \rho^{-k\alpha} b_k}{|Dv_k + \rho^{-k\alpha} b_k|}, \frac{Dv_k + \rho^{-k\alpha} b_k}{|Dv_k + \rho^{-k\alpha} b_k|} \right\rangle = f_k \text{ in } Q_1,$$

where $f_k := \frac{1}{\rho^{k(\alpha-1)}} f$. That means $f_k \in L^\infty(Q_1)$, if and only if, $\alpha < 1$. By induction hypothesis, we have that v_k is normalized function in $Q_{8/9}$. Hence

v_k satisfies the assumptions of Lemma 3.1, which ensures the existence of $\tilde{h} \in \mathcal{C}^{2,1}(Q_{7/9})$, so that

$$\sup_{Q_{7/9}} |u(x, t) - \tilde{h}(x, t)| < \delta.$$

As a consequence of Step 1, there exists an affine function $\tilde{\ell}$ such that

$$\sup_{Q_\rho} |v_k(x, t) - \tilde{\ell}(x, t)| \leq \rho^{1+\alpha}.$$

Defining $\ell_{k+1}(x, t) := \ell_k(x, t) + \rho^{k(1+\alpha)}\tilde{\ell}(\rho^{-k}x, \rho^{-2k}t)$ yields

$$\sup_{Q_{\rho^{k+1}}} |u(x, t) - \ell_{k+1}(x, t)| \leq \rho^{(k+1)(1+\alpha)}.$$

Also, the coefficients satisfy

$$|a_{k+1} - a_k| \leq C\rho^{k(1+\alpha)} \tag{10}$$

and

$$|b_{k+1} - b_k| \leq C\rho^{k\alpha} \tag{11}$$

for every $k \in \mathbb{N}$, and the Proposition is concluded. \square

Proof of Theorem 1.1. From (10) and (11), we conclude that the sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ are Cauchy sequences, consequently there are constants a_∞ and b_∞ such that

$$\lim_{n \rightarrow \infty} a_n = a_\infty \quad \text{and} \quad \lim_{n \rightarrow \infty} b_n = b_\infty.$$

Moreover, we have the estimates

$$|a_k - a_\infty| \leq C\rho^{k(1+\alpha)} \quad \text{and} \quad |b_k - b_\infty| \leq C\rho^{k\alpha}.$$

Given any $0 < \rho \ll 1$, let k be a natural number such that $\rho^{k+1} < r < \rho^k$.

Thus, we estimate from the previous computations

$$\begin{aligned} \sup_{Q_r} |u(x, t) - \ell_\infty(x, t)| &\leq \sup_{Q_{\rho^k}} |u(x, t) - \ell_k(x, t)| + \sup_{Q_{\rho^k}} |\ell_k(x, t) - \ell_\infty(x, t)| \\ &\leq \rho^{k(1+\alpha)} + |a_k - a_\infty| + \rho^k |b_k - b_\infty| \\ &\leq \rho^{k(1+\alpha)} + C\rho^{k(1+\alpha)} + C\rho^k \cdot \rho^{k\alpha} \\ &\leq C \left(\frac{1}{\rho}\right)^{(1+\alpha)} \rho^{(k+1)(1+\alpha)} \\ &\leq C_T^{(1+\alpha)}. \end{aligned}$$

To conclude the proof, we characterize the coefficients a_∞ and b_∞ . In fact, evaluating the limit in the following inequality

$$\sup_{Q_{\rho^n}} |u(x, t) - \ell_n(x, t)| \leq C\rho^{n(1+\alpha)}, \quad (12)$$

as n approaches infinity on $(x, t) = (0, 0)$, we obtain that $a_\infty = u(0, 0)$. From [3, Lemma A.1], we can conclude that $b_\infty = Du(0, 0)$. This finishes the proof. \square

3.3 Regularity in time

For account of completeness we present the proof of the regularity in the time variable, see also [3, Lemma 3.5].

Lemma 3.2 (Regularity in time). *Let $u \in \mathcal{C}(Q_1)$ be a normalized viscosity solution to (1). Assume that ρ is as in Proposition 3.1. Given $\alpha \in (0, 1)$, there exists ε such that if*

$$\|f\|_{L^\infty(Q_1)} + |p - 2| < \varepsilon,$$

then for every $t \in (-r^2, 0)$

$$|u(0, t) - u(0, 0)| \leq C|t|^{\frac{1+\alpha}{2}},$$

where $C > 0$ is a universal constant.

Proof. For $(x, t) \in Q_r$, we set

$$v(x, t) := u(x, t) - u(0, 0) - Du(0, 0) \cdot x.$$

Proposition 3.1 implies that

$$|v(x_1, t) - v(x_2, t)| \leq Cr^{1+\alpha},$$

for $x_1, x_2 \in B_r$, $t \in [-r^2, 0]$, and consequently we have

$$\text{osc}_{B_r} v(\cdot, t) \leq Cr^{1+\alpha} =: A.$$

We claim that $\text{osc}_{Q_r} v \leq CA + 4r^2\|f\|_{L^\infty(Q_1)}$. In fact, notice that v solves

$$\partial_t v - \Delta v - (p-2) \left\langle D^2 v \frac{Dv+b}{|Dv+b|}, \frac{Dv+b}{|Dv+b|} \right\rangle = f \quad \text{in } Q_r,$$

where $b := Du(0, 0)$. Employing the Lemma 2.2 in [3], we obtain

$$\text{osc}_{Q_r} v \leq Cr^{1+\alpha} + 4r^2\|f\|_{L^\infty(Q_1)}.$$

Therefore,

$$|u(0, t) - u(0, 0)| = |v(0, t)| \leq C|t|^{\frac{1+\alpha}{2}}.$$

□

4 Regularity in Sobolev spaces

In this section, we present the proof of Theorem 1.2. Here, we examine the homogeneous problem

$$u_t(x, t) - \Delta_p^N u(x, t) = 0 \quad \text{in } Q_1. \quad (13)$$

Let $u \in \mathcal{C}(Q_1)$ be a viscosity solution to (13). We consider the following regularized Dirichlet problem:

$$\begin{cases} (v_\varepsilon)_t - \Delta v_\varepsilon - (p-2) \frac{D^2 v_\varepsilon Dv_\varepsilon \cdot Dv_\varepsilon}{|Dv_\varepsilon|^2 + \varepsilon^2} = 0 & \text{in } Q_{3/4} \\ v_\varepsilon = u & \text{on } \partial Q_{3/4}. \end{cases} \quad (14)$$

It is well known that v_ε is a classical solution (in the interior), and the gradient of v_ε is uniformly bounded with respect to ε , see for instance [10].

Proof of Theorem 1.2. Consider the operator

$$F(D^2 u, x, t) := -\Delta u - (p-2) \frac{D^2 u Dv_\varepsilon \cdot Dv_\varepsilon}{|Dv_\varepsilon|^2 + \varepsilon^2}.$$

First, notice that F is a uniformly elliptic operator with constants $\lambda = \min(1, p-1)$ and $\Lambda = \max(1, p-1)$. In addition, the oscillation of F given by

$$\theta_F(x, t) := \sup_M \frac{|F(M, x, t) - F(M, 0, 0)|}{|M| + 1},$$

satisfies

$$\theta_F(x, t) \leq 2|p-2|.$$

It follows from (14) that v_ε solves

$$(v_\varepsilon)_t + F(D^2 v_\varepsilon, x, t) = 0.$$

Hence, by applying [35, Theorem 5.7], we assure that there exists a positive constant ε_1 , such that if

$$|p-2| \leq \varepsilon_1,$$

we have that $v_\varepsilon \in W_{loc}^{2,1;q}(Q_1)$ for all $q \in [1, \infty)$ and there is a constant $C > 0$ independent of ε_1 such that

$$\|v_\varepsilon\|_{W^{2,1;q}} \leq C. \quad (15)$$

Since $v_\varepsilon \rightarrow u$ uniformly in compact sets, see [16, Lemma 3.1], we obtain that u also belongs to $W_{loc}^{2,1;q}(Q_1)$ and satisfies the estimate (15). This finishes the proof. \square

The following result is a direct consequence of the Theorem 1.2.

Corollary 4.1. *Let $u \in \mathcal{C}(Q_1)$ be a normalized viscosity solution of (13). Then $u \in \mathcal{C}_{loc}^{1+\alpha, \frac{1+\alpha}{2}}(Q_1)$, with the estimate*

$$\|u\|_{\mathcal{C}^{1+\alpha, \frac{1+\alpha}{2}}(Q_1)} \leq C \|u\|_{L^\infty(Q_1)}.$$

where C is a positive constant and for every $\alpha \in (0, 1)$.

Proof. The corollary follows immediately from Theorem 1.2 with general Sobolev inequalities in Sobolev spaces. \square

Acknowledgement: PA was supported by CAPES - Brazil. MS was partially supported by CONACyT-MEXICO Grant A1-S-48577. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brazil (CAPES) - Finance Code 001.

References

- [1] Damião. J. Araújo, Eduardo. V. Teixeira, and José. M. Urbano. A proof of the $C^{p'}$ -regularity conjecture in the plane. *Adv. Math.*, 316:541–553, 2017.
- [2] Amal Attouchi. Local regularity for quasi-linear parabolic equations in non-divergence form. *Nonlinear Anal.*, 199:112051, 28, 2020.
- [3] Amal Attouchi and Mikko Parviainen. Hölder regularity for the gradient of the inhomogeneous parabolic normalized p -Laplacian. *Commun. Contemp. Math.*, 20(4):1750035, 27, 2018.

- [4] Amal Attouchi and Eero Ruosteenoja. Remarks on regularity for p -Laplacian type equations in non-divergence form. *J. Differential Equations*, 265(5):1922–1961, 2018.
- [5] Amal Attouchi and Eero Ruosteenoja. Gradient regularity for a singular parabolic equation in non-divergence form. *Discrete Contin. Dyn. Syst.*, 40(10):5955–5972, 2020.
- [6] Agnid Banerjee and Nicola Garofalo. Gradient bounds and monotonicity of the energy for some nonlinear singular diffusion equations. *Indiana Univ. Math. J.*, 62(2):699–736, 2013.
- [7] Luis A. Caffarelli. Interior a priori estimates for solutions of fully nonlinear equations. *Ann. of Math. (2)*, 130(1):189–213, 1989.
- [8] Michael G. Crandall, Maciej Kocan, and Andrzej Świąch. L^p -theory for fully nonlinear uniformly parabolic equations. *Comm. Partial Differential Equations*, 25(11-12):1997–2053, 2000.
- [9] João Vitor da Silva and Eduardo V. Teixeira. Sharp regularity estimates for second order fully nonlinear parabolic equations. *Math. Ann.*, 369(3-4):1623–1648, 2017.
- [10] Kerstin Does. An evolution equation involving the normalized p -Laplacian. *Commun. Pure Appl. Anal.*, 10(1):361–396, 2011.
- [11] Hongjie Dong, Fa Peng, Yi Ru-Ya Zhang, and Yuan Zhou. Hessian estimates for equations involving p -Laplacian via a fundamental inequality. *Adv. Math.*, 370:107212, 40, 2020.
- [12] Abderrahim Elmoataz, Matthieu Toutain, and Daniel Tenbrinck. On the p -Laplacian and ∞ -Laplacian on graphs with applications in image and data processing. *SIAM J. Imaging Sci.*, 8(4):2412–2451, 2015.
- [13] Lawrence C. Evans. The 1-Laplacian, the ∞ -Laplacian and differential games. In *Perspectives in nonlinear partial differential equations*, volume 446 of *Contemp. Math.*, pages 245–254. Amer. Math. Soc., Providence, RI, 2007.

- [14] Lawrence C. Evans and Joel Spruck. Motion of level sets by mean curvature. I [MR1100206 (92h:35097)]. In *Fundamental contributions to the continuum theory of evolving phase interfaces in solids*, pages 328–374. Springer, Berlin, 1999.
- [15] Yuzhou Fang and Chao Zhang. Gradient Hölder regularity for parabolic normalized $p(x, t)$ -Laplace equation. *J. Differential Equations*, 295:211–232, 2021.
- [16] Fredrik Arbo Høeg and Peter Lindqvist. Regularity of solutions of the parabolic normalized p -Laplace equation. *Adv. Nonlinear Anal.*, 9(1):7–15, 2020.
- [17] Cyril Imbert, Tianling Jin, and Luis Silvestre. Hölder gradient estimates for a class of singular of degenerate parabolic equations. *Adv. Nonlinear Anal.*, 8(1):845–867, 2019.
- [18] Cyril Imbert and Luis Silvestre. $C^{1,\alpha}$ regularity of solutions of some degenerate fully non-linear elliptic equations. *Adv. Math.*, 233:196–206, 2013.
- [19] Cyril Imbert and Luis Silvestre. An introduction to fully nonlinear parabolic equations. In *An introduction to the Kähler-Ricci flow*, volume 2086 of *Lecture Notes in Math.*, pages 7–88. Springer, Cham, 2013.
- [20] Tianling Jin and Luis Silvestre. Hölder gradient estimates for parabolic homogeneous p -Laplacian equations. *J. Math. Pures Appl. (9)*, 108(1):63–87, 2017.
- [21] Petri Juutinen and Bernd Kawohl. On the evolution governed by the infinity Laplacian. *Math. Ann.*, 335(4):819–851, 2006.
- [22] Robert V. Kohn and Sylvia Serfaty. A deterministic-control-based approach to motion by curvature. *Comm. Pure Appl. Math.*, 59(3):344–407, 2006.
- [23] N. V. Krylov. *Nonlinear elliptic and parabolic equations of the second order*, volume 7 of *Mathematics and its Applications (Soviet Series)*. D. Reidel Publishing Co., Dordrecht, 1987. Translated from the Russian by P. L. Buzytsky [P. L. Buzytskiĭ].

- [24] Nicolai V. Krylov and Mikhail V. Safonov. An estimate for the probability of a diffusion process hitting a set of positive measure. *Dokl. Akad. Nauk SSSR*, 245(1):18–20, 1979.
- [25] Nicolai V. Krylov and Mikhail V. Safonov. A property of the solutions of parabolic equations with measurable coefficients. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(1):161–175, 239, 1980.
- [26] Juan J. Manfredi, Mikko Parviainen, and Julio D. Rossi. An asymptotic mean value characterization for a class of nonlinear parabolic equations related to tug-of-war games. *SIAM J. Math. Anal.*, 42(5):2058–2081, 2010.
- [27] Yuval Peres, Oded Schramm, Scott Sheffield, and David B. Wilson. Tug-of-war and the infinity Laplacian. *J. Amer. Math. Soc.*, 22(1):167–210, 2009.
- [28] Yuval Peres and Scott Sheffield. Tug-of-war with noise: a game-theoretic view of the p -Laplacian. *Duke Math. J.*, 145(1):91–120, 2008.
- [29] Edgard A. Pimentel and Makson S. Santos. Asymptotic methods in regularity theory for nonlinear elliptic equations: a survey. In *PDE models for multi-agent phenomena*, volume 28 of *Springer INdAM Ser.*, pages 167–194. Springer, Cham, 2018.
- [30] Edgard A. Pimentel and Makson S. Santos. Improved regularity for the porous medium equation along zero level-sets. *To appear in Israel Journal of Mathematics*, 2020.
- [31] E. Teixeira. Universal moduli of continuity for solutions to fully nonlinear elliptic equations. *Arch. Ration. Mech. Anal.*, 211(3):911–927, 2014.
- [32] E. Teixeira and J.-M. Urbano. A geometric tangential approach to sharp regularity for degenerate evolution equations. *Anal. PDE*, 7(3):733–744, 2014.
- [33] Eduardo V. Teixeira. Regularity for quasilinear equations on degenerate singular sets. *Math. Ann.*, 358(1-2):241–256, 2014.
- [34] Lihe Wang. On the regularity theory of fully nonlinear parabolic equations. I. *Comm. Pure Appl. Math.*, 45(1):27–76, 1992.

- [35] Lihe Wang. On the regularity theory of fully nonlinear parabolic equations.
II. *Comm. Pure Appl. Math.*, 45(2):141–178, 1992.

PÊDRA D. S. ANDRADE

Department of Mathematics

Pontifical Catholic University of Rio de Janeiro – PUC-Rio

22451-900, Gávea, Rio de Janeiro-RJ, Brazil

`pedra.andrade@mat.puc-rio.br`

MAKSON S. SANTOS

Center of Investigations in Mathematics (CIMAT)

36000 Guanajuato Gto - MEXICO

`makson.santos@cimat.mx`