COUNTING 3-DIMENSIONAL ALGEBRAIC TORI OVER \mathbb{Q}

JUNGIN LEE

Abstract

In this paper we count the number $N_3^{\text{tor}}(X)$ of 3-dimensional algebraic tori over \mathbb{Q} whose Artin conductor is bounded above by X. We prove that $N_3^{\text{tor}}(X) \ll_{\varepsilon} X^{1+\frac{\log 2+\varepsilon}{\log \log X}}$, and this upper bound can be improved to $N_3^{\text{tor}}(X) \ll X(\log X)^4 \log \log X$ under the Cohen-Lenstra heuristics for p = 3. We also prove that for 67 out of 72 conjugacy classes of finite nontrivial subgroups of $\operatorname{GL}_3(\mathbb{Z})$, Malle's conjecture for tori over \mathbb{Q} holds up to a bounded power of $\log X$ under the Cohen-Lenstra heuristics for p = 3 and Malle's conjecture for quartic A_4 -fields.

1 Introduction

Throughout the paper, we assume that every number field is contained in a fixed algebraic closure $\overline{\mathbb{Q}}$ of \mathbb{Q} . Let K be a number field and D_K be the absolute value of the discriminant of K. By Hermite-Minkowski theorem, there are only finitely many number fields K such that D_K is bounded above by a given number. For an integer $n \geq 2$, let $N_n(X)$ be the number of degree n number fields K such that $D_K \leq X$. For $n \geq 2$ and a transitive subgroup $G \leq S_n$, let $N_n(X; G)$ be the number of degree n number fields K such that $D_K \leq X$ and the Galois group $\operatorname{Gal}(K^c/\mathbb{Q})$ is permutation-isomorphic to G in S_n . Here K^c denotes the Galois closure of K over \mathbb{Q} .

Counting number fields by discriminant (i.e. the asymptotics of the numbers $N_n(X)$ and $N_n(X;G)$) is one of the central problems in arithmetic statistics. Linnik's conjecture states that for each $n \ge 2$, there exists a constant $c_n > 0$ such that $N_n(X) \sim c_n X$ as $X \to \infty$. This conjecture has been proved only for $n \le 5$. The case n = 2 is easy, the case n = 3 was proved by Davenport-Heilbronn [16] and the cases n = 4, 5 were proved by Bhargava [6,7]. For general n, upper bound of $N_n(X)$ has been studied by Schmidt [30], Ellenberg-Venkatesh [19], Couveignes [15] and Lemke Oliver-Thorne [23].

Malle's conjecture [27] states that

$$N_n(X;G) \sim c_G X^{\frac{1}{a(G)}} (\log X)^{b(G)-1}$$
(1)

for some positive integers a(G), b(G) and a constant $c_G > 0$. The index of $g \in G \leq S_n$ is define by

 $\operatorname{ind}(g) := n - \operatorname{the number of orbits of } g \text{ on } \{1, 2, \cdots, n\}$

and let $a(G) := \min_{g \in G \setminus \{1\}} \operatorname{ind}(g)$. The number b(G) is defined to be the number of orbits \mathcal{C} of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -action on the conjugacy classes of G (via the cyclotomic character) such that the index of some (equivalently, all) $g \in \mathcal{C}$ is a(G). We refer [22, Section 1.1] for a summary on the known results on Malle's conjecture.

As a natural generalization of counting number fields by discriminant, the author [22] studied counting (algebraic) tori over \mathbb{Q} by Artin conductor. Let T be an n-dimensional tori over \mathbb{Q} with a splitting field L and $X^*(T) := \operatorname{Hom}_{\overline{\mathbb{Q}}}(T_{\overline{\mathbb{Q}}}, \mathbb{G}_{m,\overline{\mathbb{Q}}})$ be its character group. Then the Galois group $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ acts on $X^*(T)$ by conjugation, and this induces the representation

$$\rho_T : \operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to \operatorname{Aut}(X^*(T)) \cong \operatorname{GL}_n(\mathbb{Z}).$$

Its image $G_T := \operatorname{im}(\rho_T)$ is a finite subgroup of $\operatorname{GL}_n(\mathbb{Z})$ isomorphic to $\operatorname{Gal}(L/\mathbb{Q})$. Since the isomorphism $\operatorname{Aut}(X^*(T)) \cong \operatorname{GL}_n(\mathbb{Z})$ depends on the choice of the \mathbb{Z} -basis of $X^*(T)$, ρ_T and G_T are well-defined only up to conjugation. Let C(T) be the Artin conductor of the representation

$$\rho : \operatorname{Gal}(L/\mathbb{Q}) \to \operatorname{Aut}(X^*(T)_{\mathbb{Q}}) \cong \operatorname{GL}_n(\mathbb{Q})$$

induced by ρ_T . For a degree *n* number field $K, T = \mathbb{R}_{K/\mathbb{Q}} \mathbb{G}_m$ (Weil restriction of \mathbb{G}_m) is an *n*-dimensional torus over \mathbb{Q} whose splitting field is K^c and the Artin conductor is $C(\mathbb{R}_{K/\mathbb{Q}} \mathbb{G}_m) = D_K$ (see [22, Section 1.2]). This shows that counting tori over \mathbb{Q} of given dimension by Artin conductor is a generalization of counting number fields of given degree by discriminant.

Let $N_n^{\text{tor}}(X)$ be the number of the isomorphism classes of tori over \mathbb{Q} of dimension n such that $C(T) \leq X$. For a finite subgroup $H \neq 1$ of $\operatorname{GL}_n(\mathbb{Z})$, $N_n^{\text{tor}}(X; H)$ denotes the number of such tori T over \mathbb{Q} such that G_T is conjugate to H in $\operatorname{GL}_n(\mathbb{Z})$. The following two conjectures from [22] are analogues of Linnik's and Malle's conjectures for tori over \mathbb{Q} . Note that the second conjecture follows from a more general conjecture of Ellenberg and Venkatesh [18, Question 4.3], so it is not new. It is also remarkable that the second conjecture implies the first conjecture [22, Corollary 3.6].

Conjecture 1.1. ([22, Conjecture 3.1]) For every $n \ge 1$, there exists a constant $c_n > 0$ satisfying

$$N_n^{\text{tor}}(X) \sim c_n X (\log X)^{n-1}.$$
(2)

Conjecture 1.2. ([22, Conjecture 3.2]) For every $n \ge 1$ and a finite subgroup $1 \ne H \le \operatorname{GL}_n(\mathbb{Z})$,

$$N_n^{\text{tor}}(X; H) \sim c_H X^{\frac{1}{a(H)}} (\log X)^{b(H)-1}$$
 (3)

where the positive integers a(H), b(H) and a constant $c_H > 0$ depend only on H. For an $n \times n$ identity matrix I_n , the number a(H) is given by

$$a(H) := \min_{h \in H \setminus \{I_n\}} \operatorname{rank}(h - I_n)$$
(4)

and the number b(H) is given by the number of the orbits C of the action of $\operatorname{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on the conjugacy classes of H via the cyclotomic character such that $\operatorname{rank}(h - I_n) = a(H)$ for some (equivalently, all) $h \in C$.

The above conjectures are trivial for the case n = 1. For the 2-dimensional case, the following results are known. Here $H_{12,A}$ is a finite subgroup of $\operatorname{GL}_2(\mathbb{Z})$ which is isomorphic to the dihedral group D_6 of order 12. Such group $H_{12,A}$ is unique up to conjugation and satisfies $a(H_{12,A}) = 1$ and $b(H_{12,A}) = 2$. We also use the asymptotic notation $f(X) \ll_{\varepsilon} g(X, \varepsilon)$, which means that $f(X) \ll g(X, \varepsilon)$ for every $\varepsilon > 0$.

Proposition 1.3. (1) ([22, Proposition 4.1]) Conjecture 1.2 holds for every finite nontrivial subgroup of $\operatorname{GL}_2(\mathbb{Z})$ which is not conjugate to $H_{12,A}$.

(2) ([22, Theorem 4.9]) We have the followings:

$$X \ll N_2^{\text{tor}}(X; H_{12,A}) \ll_{\varepsilon} X^{1 + \frac{\log 2 + \varepsilon}{\log \log X}}$$
(5a)

$$X \log X \ll N_2^{\text{tor}}(X) \ll_{\varepsilon} X^{1 + \frac{\log 2 + \varepsilon}{\log \log X}}.$$
(5b)

(3) ([22, Theorem 4.10]) Under the assumption of Conjecture 1.4, we have

$$N_2^{\text{tor}}(X; H_{12,A}) \le N_2^{\text{tor}}(X) \ll_{\varepsilon} X(\log X)^{1+\varepsilon}.$$
(6)

In the above proposition, the following version of the Cohen-Lenstra heuristics was used. Denote by NF_2^+ (resp. NF_2^-) the set of all real (resp. imaginary) quadratic fields whose elements are ordered by the absolute values of their discriminants. For a number field K and a prime p, denote the size of the p-torsion subgroup of the class group of K by $h_p(K)$. For an odd prime p and a positive integer α , consider the following version of the Cohen-Lenstra heuristics.

- ([11, (C10)]) Conj⁺(p, α) : The average of $\prod_{0 \le i < \alpha} (h_p(K) p^i)$ for $K \in NF_2^+$ is $p^{-\alpha}$.
- ([11, (C6)]) Conj⁻(p, α) : The average of $\prod_{0 \le i < \alpha} (h_p(K) p^i)$ for $K \in NF_2^-$ is 1.

The above conjectures are true for p = 3 and $\alpha = 1$ by [16, Theorem 3], but they are still open for the other cases. In many cases the assumption of the conjectures for p = 3 and every $\alpha > 0$ improves the upper bounds for counting tori over \mathbb{Q} .

Conjecture 1.4. $\operatorname{Conj}^+(3, \alpha)$ and $\operatorname{Conj}^-(3, \alpha)$ are true for every positive integer α .

The purpose of this paper is to count the number of 3-dimensional tori over \mathbb{Q} by Artin conductor. First we classify the 3-dimensional tori over \mathbb{Q} and compute their Artin conductors in Section 3. The classification is much more complicated compared to the 2-dimensional case. After that, we estimate the magnitude of $N_3^{\text{tor}}(X; H)$ for each finite nontrivial subgroup H of $\text{GL}_3(\mathbb{Z})$ in Section 4. We do this for abelian H in Section 4.1 and non-abelian H in Section 4.2 and 4.3. The results can be summarized as follow.

Theorem 1.5. Let *H* be a finite nontrivial subgroup of $GL_3(\mathbb{Z})$.

- (1) (Proposition 4.1, 4.3) Conjecture 1.2 is true for every abelian H.
- (2) (Theorem 4.8) $X^{\frac{1}{a(H)}} \ll N_3^{\text{tor}}(X; H)$ for every H.
- (3) (Proposition 4.1, 4.3, 4.4, 4.9, 4.10) Under the assumption of Conjecture 1.4 and Malle's conjecture for quartic A_4 -fields, we have

$$N_3^{\text{tor}}(X; H) \ll_{\varepsilon} X^{\frac{1}{a(H)}} (\log X)^{6+\varepsilon}$$

for 67 out of 72 conjugacy classes of finite nontrivial subgroups of $GL_3(\mathbb{Z})$.

It is notable that the ratios of the upper and lower bounds of $N_3^{\text{tor}}(X; H)$ are $(\log X)^{O(1)}$ for most of the finite subgroups H of $\text{GL}_3(\mathbb{Z})$. Since we have

$$N_3^{\rm tor}(X) = \sum_H N_3^{\rm tor}(X;H)$$

where H runs through the conjugacy classes of finite nontrivial subgroups of $GL_3(\mathbb{Z})$, we can bound the size of $N_3^{tor}(X)$. The following theorem on the upper bound of $N_3^{tor}(X)$ is the main result of the paper.

Theorem 1.6. (Theorem 4.15)

(1) We have

 $N_3^{\text{tor}}(X) \ll_{\varepsilon} X^{1 + \frac{\log 2 + \varepsilon}{\log \log X}}.$ (7)

(2) Under the assumption of Conjecture 1.4, we have

$$N_3^{\text{tor}}(X) \ll X (\log X)^4 \log \log X.$$
(8)

2 Preliminaries

2.1 Known results on counting number fields

In this section, C denotes a positive constant which may change from line to line. First we give a list of known cases of Malle's conjecture which will be used in the sequel.

Proposition 2.1. The equation (1) holds for the following cases:

- (1) ([25]) $G \leq S_{|G|}, G$ abelian
- (2) ([16,9,6]) $S_3 \leq S_3, D_4 \leq S_4, S_4 \leq S_4$
- (3) ([28, Theorem 1.1]) $D_6 \cong S_3 \times C_2 \leq S_6$ (i.e. $N_6(X; D_6) \sim CX^{\frac{1}{2}}$)
- (4) ([31, Theorem 1]) $D_4 \leq S_8$ (i.e. $N_8(X; D_4) \sim CX^{\frac{1}{4}} (\log X)^2$).

There also has been some progress on counting number fields by an invariant other than the discriminant. For a positive integer $n \ge 2$ and a transitive subgroup $G \le S_n$, let $\operatorname{NF}_n(G)$ be the set of degree n number fields such that the Galois group $\operatorname{Gal}(K^c/\mathbb{Q})$ is permutation-isomorphic to G. Let I be an invariant of number fields such that for every X > 0, there are finitely many number fields K such that $I(K) \le X$. Denote by $N_n(X;G;I)$ the number of $K \in \operatorname{NF}_n(G)$ such that $I(K) \le X$. Sometimes we write $N_n(X;G;I)$ by $N_n^K(X;G;I)$ to clarify that I is an invariant of K. It is clear from the definition that $N_n(X;G;I) = N_n(X;G)$ if $I(K) = D_K$. Denote $\operatorname{NF}_2(C_2)$ (resp. $N_2(X;C_2;I)$) by NF_2 (resp. $N_2(X;I)$) for simplicity.

The next proposition is a collection of results on the asymptotics of $N_n(X;G;I)$ where I is an invariant other than the discriminant. The first two results are direct consequences of the work of Mäki [26] on the asymptotics of the number of abelian number fields with bounded conductor (cf. [22, Section 2.3]).

Proposition 2.2. (1) For $L_4 \in NF_4(C_4)$, denote its unique quadratic subfield by L_2 . Then

$$N_4(X; C_4; \frac{D_{L_4}}{D_{L_2}}) \sim CX^{\frac{1}{2}} \log X.$$
 (9)

(2) For $L_6 \in NF_6(C_6)$, denote its unique cubic (resp. quadratic) subfield by L_3 (resp. L_2). Then

$$N_6(X; C_6; \frac{D_{L_6}}{D_{L_2} D_{L_3}}) \sim C X^{\frac{1}{2}} (\log X)^2.$$
 (10)

(3) ([1, Theorem 1]) For $L_4 \in NF_4(D_4)$, denote its unique quadratic subfield by L_2 . Then

$$N_4(X; D_4; \frac{D_{L_4}}{D_{L_2}}) = CX \log X + O(X \log \log X).$$
(11)

For $L_6 \in NF_6(D_6)$, denote its unique cubic (resp. quadratic) subfield by L_3 (resp. L_2). Then

$$N_2^{\text{tor}}(X; H_{12,A}) = N_6(X; D_6; \frac{D_{L_6}}{D_{L_3}D_{L_2}})$$

(cf. [22, Section 4.1]) so Proposition 1.3 implies that

$$N_6(X; D_6; \frac{D_{L_6}}{D_{L_3} D_{L_2}}) \ll_{\varepsilon} X^{1 + \frac{\log 2 + \varepsilon}{\log \log X}} \ll_{\varepsilon} X^{1 + \varepsilon}$$
(12)

and this can be improved to

$$N_6(X; D_6; \frac{D_{L_6}}{D_{L_3} D_{L_2}}) \ll_{\varepsilon} X(\log X)^{1+\varepsilon}$$
(13)

under the assumption of Conjecture 1.4.

We also introduce a proposition which concerns the product distribution appears in counting number fields. It is useful when we consider the compositum of two linearly disjoint number fields. **Proposition 2.3.** Let $F_i(X) = \# \{s \in S_i : s \leq X\}$ (i = 1, 2) be the asymptotic distribution of some multi-set S_i consists of a sequence of elements of $\mathbb{R}_{\geq 1}$. Suppose that $F_i(X) \sim A_i X^{n_i} (\log X)^{r_i}$ for $A_i > 0$, $n_i > 0$ and $r_i \in \mathbb{Z}_{\geq 0}$. Consider the product distribution

$$P(X) := \# \{ (s_1, s_2) \in S_1 \times S_2 : s_1 s_2 \le X \}.$$

(1) ([33, Lemma 3.1]) If $n_1 = n_2 = n$, then

$$P(X) \sim A_1 A_2 \frac{r_1! r_2!}{(r_1 + r_2 + 1)!} n X^n (\log X)^{r_1 + r_2 + 1}.$$


(2) ([33, Lemma 3.2]) If $n_1 > n_2$, then there exists a constant C > 0 such that

$$P(X) \sim C X^{n_1} (\log X)^{r_1}.$$

2.2 Discriminants of number fields

For some Galois extensions of number fields, there are algebraic relations between the discriminants of their subextensions. We provide such relations for Galois extensions whose Galois group is isomorphic to one of the groups C_2^2 , S_3 , A_4 and S_4 . The formulas in the following proposition will be used frequently in Section 3.

Proposition 2.4. Let L/K be a Galois extension of number fields.



(1) Assume that $\operatorname{Gal}(L/K) \cong C_2^2$ and let K_1 , K_2 and K_3 be the quadratic subextensions of L/K. Then we have

$$D_L D_K^2 = D_{K_1} D_{K_2} D_{K_3}.$$
 (14)

(2) Assume that $\operatorname{Gal}(L/K) \cong S_3$, let L_3 be one of the cubic subextensions of L/K and L_2 be the unique quadratic subextension of L/K. Then we have

$$D_L D_K^2 = D_{L_3}^2 D_{L_2}. (15)$$

(3) Assume that Gal(L/K) ≅ A₄, let L₆ be one of the sextic subextensions of L/K, L₄ be one of the quartic subextensions of L/K and L₃ be the unique cubic subextension of L/K. (In this case L₃ is a subfield of L₆.) Then we have the following formulas:

$$D_L D_K^3 = D_{L_4}^3 D_{L_3} \tag{16a}$$

$$D_{L_6}D_K = D_{L_4}D_{L_3} (16b)$$

$$D_L D_K^2 = D_{L_6} D_{L_4}^2. (16c)$$

(4) Assume that $\operatorname{Gal}(L/K) \cong S_4$, let L_i (i = 3, 4) be one of the degree *i* subextensions of L/K and L_6 be the unique subextension of L/K such that $L_3 \subset L_6$, $\operatorname{Gal}(L/L_6) \cong C_2^2$ and the Galois closure of L_6/K is *L*. Then we have

$$D_{L_6} D_K = D_{L_4} D_{L_3}.$$
 (17)

- Proof. (1) By the conductor-discriminant formula [29, VII.11.9], we have $D_{L/K} = D_{K_1/K} D_{K_2/K} D_{K_3/K}$ where $D_{L/K} := N_{K/\mathbb{Q}}(\mathcal{D}_{L/K})$ is the absolute norm of the relative discriminant $\mathcal{D}_{L/K}$.
 - (2) $S_3 = C_3 \rtimes C_2$ is a Frobenius group so it is a consequence of [20, Theorem 4].
 - (3) $A_4 = C_2^2 \rtimes C_3$ is a Frobenius group so the first formula is a consequence of [20, Theorem 4]. For a number field M, denote its Dedekind zeta function by $\zeta_M(s)$ and denote the number of real and complex embeddings of M by r_M and s_M , respectively. Then the character theory of A_4 implies the relation

$$\zeta_{L_6}(s)\zeta_K(s) = \zeta_{L_4}(s)\zeta_{L_3}(s)$$
(18)

(cf. [14, Theorem 5.1]) and the functional equation of the Dedekind zeta function [29, Corollary VII.5.10] implies that

$$D_M^{s-\frac{1}{2}} \cdot \frac{\zeta_M(s)}{\zeta_M(1-s)} \cdot \left(\frac{\Gamma(\frac{s}{2})}{\Gamma(\frac{1-s}{2})}\right)^{r_M} \cdot \left(\frac{2^{1-2s}\Gamma(s)}{\Gamma(1-s)}\right)^{s_M} \cdot \pi^{(\frac{1-2s}{2}[M:\mathbb{Q}])} = 1$$
(19)

for any number field M.

Now denote $c := s_K + s_{L_6} - s_{L_3} - s_{L_4} \in \mathbb{Z}$. The equations (18) and (19) imply that

$$\left(\frac{D_K D_{L_6}}{D_{L_4} D_{L_3}}\right)^{s-\frac{1}{2}} = \left(\frac{2^{1-2s}\Gamma(s)\Gamma(\frac{1-s}{2})^2}{\Gamma(1-s)\Gamma(\frac{s}{2})^2}\right)^c.$$

Substituting $s = \frac{3}{2}$ into the above equation, we obtain the equation

$$\frac{D_K D_{L_6}}{D_{L_4} D_{L_3}} = \left(\frac{\frac{1}{4}\Gamma(\frac{3}{2})\Gamma(-\frac{1}{4})^2}{\Gamma(-\frac{1}{2})\Gamma(\frac{3}{4})^2}\right)^c = (-1)^c$$

and its left-hand side is positive. This proves the second formula, and the last formula comes from the first and second formulas.

(4) It can be proved as in the proof of the formula (16b), except that the character theory of A_4 is replaced by the character theory of S_4 (cf. [14, Theorem 5.1]).

For a prime p and a positive integer m, denote the exponent of p in m by $v_p(m)$. If M is a number field which is tamely ramified at p, the inertia group $I_{M,p}$ is cyclic so we can choose its generator $g_{M,p}$. The following proposition gives a description of the discriminant of compositum of two number fields. See [33, Section 2] for details.

Proposition 2.5. ([33, Theorem 2.2 and 2.3]) Let K_1 and K_2 be number fields such that $K_1^c \cap K_2^c = \mathbb{Q}$ and p be a prime such that both of K_1 and K_2 are tamely ramified at p. Suppose that $g_{K_1,p} = \prod_k c_k$ (product of disjoint cycles) and $g_{K_2,p} = \prod_l d_l$. Then

$$v_p(D_{K_1K_2}) = m_1m_2 - \sum_{k,l} \gcd(|c_k|, |d_l|),$$

where m_i is the degree of K_i and |c| denotes the length of the cycle c. If the least common multiple of $|c_k|$ and the least common multiple of $|d_l|$ are coprime, then we have

$$v_p(D_{K_1K_2}) = v_p(D_{K_1}) \cdot m_2 + v_p(D_{K_2}) \cdot m_1 - v_p(D_{K_1})v_p(D_{K_2}).$$

2.3 Analytic preliminaries

The following version of the Tauberian theorem is very useful for counting number fields. For instance, Wright [35] proved Malle's conjecture for abelian extensions by studying their associated Dirichlet series and applying the Tauberian theorem.

Proposition 2.6. ([17, Theorem III]) Let $f(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ $(a_n \ge 0)$ be a Dirichlet series which converges for $\operatorname{Re}(s) > a > 0$. Assume that

$$f(s) = \frac{g(s)}{(s-a)^w} + h(s)$$

in the domain of convergence, where g(s) and h(s) are holomorphic in $\operatorname{Re}(s) \ge a$, $g(a) \ne 0$ and w is a positive integer. Then we have

$$\sum_{n \le X} a_n \sim \frac{g(a)}{a\Gamma(w)} X^a (\log X)^{w-1},$$

where $\Gamma(w)$ denotes the Gamma function.

For $\alpha > \beta > 0$, denote $f \in \mathcal{M}(\alpha, \beta)$ if f(s) converges for $\operatorname{Re}(s) > \alpha$ and has a meromorphic continuation to $\operatorname{Re}(s) > \beta$ which is holomorphic except for a simple pole at $s = \alpha$. The next lemma will be used in the proof of Proposition 4.6.

Lemma 2.7. We have

$$g(s) := \prod_{p \equiv \pm 1 \pmod{7}} \left(1 + \frac{3}{p^s}\right) \in \mathcal{M}(1, \frac{1}{2}).$$

Proof. For a positive integer q > 1 and an integer a prime to q, define the Dirichlet series

$$B_{q,a}(s) := \prod_{p \equiv a \pmod{q}} (1 - \frac{1}{p^s})^{-1}$$

Let χ be a Dirichlet character modulo q and χ_0 be the principal character modulo q. By the answer of Johan Andersson to the MathOverflow question 28000 [2], we have

$$B_{q,a}(s) = L(s,\chi_0)^{\frac{1}{\varphi(q)}} \prod_{\chi \neq \chi_0} L(s,\chi)^{\frac{\chi(a)}{\varphi(q)}} A_{q,a}(s)$$

where $A_{q,a}(s)$ is holomorphic and non-vanishing for $\operatorname{Re}(s) > \frac{1}{2}$.

A direct computation shows that

$$g(s) = B_{7,1}(s)^3 B_{7,-1}(s)^3 h(s)$$

for

$$h(s) := \prod_{p \equiv \pm 1 \pmod{7}} \left(1 - \frac{6}{p^{2s}} + \frac{8}{p^{3s}} - \frac{3}{p^{4s}} \right),$$

which is holomorphic and non-vanishing in $\operatorname{Re}(s) > \frac{1}{2}$. Therefore it is enough to show that

$$B_{7,1}(s)^3 B_{7,-1}(s)^3 = L(s,\chi_0) \prod_{\chi \neq \chi_0} L(s,\chi)^{\frac{\chi(1) + \chi(-1)}{2}} A_{7,1}(s)^3 A_{7,-1}(s)^3 \in \mathcal{M}(1,\frac{1}{2}).$$

Since $A_{7,1}(s)^3 A_{7,-1}(s)^3$ is holomorphic and non-vanishing for $\operatorname{Re}(s) > \frac{1}{2}$ and $\overline{\chi(1) + \chi(-1)} \in \{0,1\}$, the following well-known properties of Dirichlet *L*-functions finish the proof.

- If $\chi \neq \chi_0$, then $L(s,\chi)$ extends to a holomorphic function on $\operatorname{Re}(s) > 0$ and $L(1,\chi) \neq 0$.
- If $\chi = \chi_0$, then $L(s, \chi_0)$ extends to a meromorphic function on $\operatorname{Re}(s) > 0$ which is holomorphic except for a simple pole at s = 1.

The next lemma follows directly from Wirsing's theorem [34, Satz 1]. For a positive integer n, denote by $\tau(n)$ the number of positive divisors of n.

Lemma 2.8. ([5, Lemma 2.1]) For any positive real number t, we have

$$\sum_{n \le X} \tau(n)^t \ll_t X (\log X)^{2^t - 1}.$$

Even though the lemma is given in [5] only for the case where t is an integer, its proof works for every positive real number t.

3 Classification of 3-dimensional tori over \mathbb{Q}

In this section, we provide a classification of the isomorphism classes of 3-dimensional tori over \mathbb{Q} , together with their Artin conductors. There are 73 conjugacy classes of finite subgroups of $\operatorname{GL}_3(\mathbb{Z})$, which are computed by Tahara [32] (and corrected by Ascher and Grimmer [3]). For each finite subgroup $H \neq 1$ of $\operatorname{GL}_3(\mathbb{Z})$, we compute tori T over \mathbb{Q} such that G_T is conjugate to H in $\operatorname{GL}_3(\mathbb{Z})$. The computations of T can be done as in [22, Example 2.1] so we omit them here. The next proposition, which summarizes [22, Section 2.2], will be frequently used for the computation of the Artin conductor C(T).

Proposition 3.1. (1) For a number field K, $C(\mathbb{R}_{K/\mathbb{Q}} \mathbb{G}_m) = D_K$.

- (2) If $1 \to T_1 \to T_2 \to T_3 \to 1$ is an exact sequence of tori over \mathbb{Q} , then $C(T_2) = C(T_1)C(T_3)$.
- (3) For an extension L/K of number fields,

$$T_{L/K} := \ker(\mathbf{R}_{L/\mathbb{Q}} \mathbb{G}_m \xrightarrow{\mathbf{N}_{L/K}} \mathbf{R}_{K/\mathbb{Q}} \mathbb{G}_m)$$

 $(N_{L/K} \text{ denotes the norm map})$ is a torus over \mathbb{Q} of dimension $[L:\mathbb{Q}] - [K:\mathbb{Q}]$ and

$$C(T_{L/K}) = \frac{D_L}{D_K}.$$
(20)

(4) Suppose that K_1 and K_2 are linearly disjoint number fields and let $L = K_1 K_2$. Then $T_{L/K_1} \cap T_{L/K_2}$ is a torus over \mathbb{Q} of dimension $([K_1 : \mathbb{Q}] - 1)([K_2 : \mathbb{Q}] - 1)$ and

$$C(T_{L/K_1} \cap T_{L/K_2}) = \frac{D_L}{D_{K_1} D_{K_2}}.$$
(21)

In each case, denote the splitting field of a torus T by L and identify $\operatorname{Gal}(L/\mathbb{Q})$ with G_T . Let C_m be the cyclic group of order m, D_m be the dihedral group of order 2m, S_m be the symmetric group of degree m and A_m be the alternating group of degree m. For simplicity, denote $D_i := D_{L_i}, D'_i := D_{L'_i}$, and so on. We warn the readers not to confuse the discriminant D_i and the dihedral group D_m . The following list gives the classification of 3-dimensional tori over \mathbb{Q} (except for the trivial one \mathbb{G}^3_m), together with their Artin conductors.

(i) $G_T \cong C_2$: T is one of the following types.

(a)
$$G_T = H_{2,a} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times T^2_{L/\mathbb{Q}} \text{ and } C(T) = D^2_L.$$

(b) $G_T = H_{2,b} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \right\rangle : T = \mathbb{G}_m^2 \times T_{L/\mathbb{Q}} \text{ and } C(T) = D_L.$ (c) $G_T = H_{2,c} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{R}_{L/\mathbb{Q}} \mathbb{G}_m \times T_{L/\mathbb{Q}} \text{ and } C(T) = D_L^2.$ (d) $G_T = H_{2,d} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times \mathbb{R}_{L/\mathbb{Q}} \mathbb{G}_m \text{ and } C(T) = D_L.$ (e) $G_T = H_{2,e} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L/\mathbb{Q}}^3 \text{ and } C(T) = D_L^3.$

(ii) $G_T \cong C_3$: T is one of the following types.

(a)
$$G_T = H_{3,a} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times T_{L/\mathbb{Q}} \text{ and } C(T) = D_L$$

(b) $G_T = H_{3,b} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \right\rangle : T = \mathbb{R}_{L/\mathbb{Q}} \mathbb{G}_m \text{ and } C(T) = D_L.$

(iii) $G_T = \langle g \rangle \cong C_4$: $L_4 = L$ has the unique quadratic subfield $L_2 = L^{g^2}$. T is one of the following types.

(a)
$$G_T = H_{4,a} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times T_{L_4/L_2} \text{ and } C(T) = \frac{D_4}{D_2}.$$

(b) $G_T = H_{4,b} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L_2/\mathbb{Q}} \times T_{L_4/L_2} \text{ and } C(T) = D_4.$
(c) $G_T = H_{4,c} := \left\langle \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = \left\{ v \in \mathbb{R}_{L_4/\mathbb{Q}} \mathbb{G}_m : v \cdot g^2 v = gv \cdot g^3 v \right\}.$ Since

$$1 \to T \to \mathcal{R}_{L_4/\mathbb{Q}} \mathbb{G}_m \xrightarrow{v \mapsto \frac{v \cdot g^2 \cdot v}{g v \cdot g^3 v}} T_{L_2/\mathbb{Q}} \to 1$$

is exact, we have $C(T) = \frac{D_4}{D_2}$. (d) $G_T = H_{4,d} := \left\langle \begin{pmatrix} -1 & 0 & -1 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle$: $T = T_{L_4/\mathbb{Q}}$ and $C(T) = D_4$.

(iv) $G_T = \langle g, h \rangle \cong C_2 \times C_2$: L has 3 quadratic subfields $L_1 = L^g$, $L_2 = L^h$ and $L_3 = L^{gh}$. Then $D_L = D_1 D_2 D_3$ by the equation (14). T is one of the following types.

(e)
$$G_T = H_{4,e} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_1/\mathbb{Q}} \times T^2_{L_3/\mathbb{Q}} \text{ and } C(T) = D_1 D_3^2.$$

(f) $C_T = H_{4,e} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{4,e} \times T_{4,e} \times T_{4,e} \text{ and } C(T) = D_1 D_3^2.$

(f)
$$G_T = H_{4,f} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\rangle : T = T_{L_1/\mathbb{Q}} \times T_{L_2/\mathbb{Q}} \times T_{L_3/\mathbb{Q}} \text{ and } C(T) = D_L$$

(g)
$$G_T = H_{4,g} := \left\langle \begin{pmatrix} 1 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times T_{L_2/\mathbb{Q}} \times T_{L_3/\mathbb{Q}} \text{ and } C(T) = D_2 D_3.$$

(h)
$$G_T = H_{4,h} := \left\langle \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L_1/\mathbb{Q}} \times T_{L/L_1} \text{ and } C(T) = D_L$$

(i)
$$G_T = H_{4,i} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times T_{L/L_1} \text{ and } C(T) = \frac{D_L}{D_1} = D_2 D_3.$$

(j)
$$G_T = H_{4,j} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{R}_{L_1/\mathbb{Q}} \mathbb{G}_m \times T_{L_2/\mathbb{Q}} \text{ and } C(T) = D_1 D_2.$$

(k)
$$G_T = H_{4,k} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_3/\mathbb{Q}} \times T_{L/L_2} \text{ and } C(T) = D_1 D_3^2.$$

(l)
$$G_T = H_{4,l} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 1 & 0 & -1 \\ -1 & -1 & 0 \end{pmatrix} \right\rangle$$
:

$$T = \left\{ (v_1, v_2) \in T_{L/L_1} \times T_{L/L_3} : v_1 \cdot hv_1 = v_2 \cdot hv_2 \right\}$$

Since

$$1 \to T \to T_{L/L_1} \times T_{L/L_3} \xrightarrow{(v_1, v_2) \mapsto \frac{v_1 \cdot nv_1}{v_2 \cdot hv_2}} T_{L_2/\mathbb{Q}} \to \mathbb{I}$$
 is exact, we have $C(T) = \left(\frac{D_L}{D_1} \cdot \frac{D_L}{D_3}\right) \cdot D_2^{-1} = D_L.$

(m)
$$G_T = H_{4,m} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ -1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \right\rangle :$$

$$T = \left\{ (v_1, v_2) \in T_{L/L_1} \times \mathcal{R}_{L_3/\mathbb{Q}} \mathbb{G}_m : v_1 \cdot hv_2 = v_2 \cdot hv_1 \right\}.$$

Since

$$1 \to T \to T_{L/L_1} \times \mathcal{R}_{L_3/\mathbb{Q}} \mathbb{G}_m \xrightarrow{(v_1, v_2) \mapsto \frac{v_1 \cdot nv_2}{v_2 \cdot hv_1}} T_{L_3/\mathbb{Q}} \to 1$$

is exact, we have $C(T) = \left(\frac{D_L}{D_1} \cdot D_3\right) \cdot D_3^{-1} = D_2 D_3.$

- (n) $G_T = H_{4,n} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 1 & -1 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L/\mathbb{Q}} \text{ so } C(T) = D_L.$
- (o) $G_T = H_{4,o} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & -1 & 1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \left\{ v \in \mathbf{R}_{L/\mathbb{Q}} \, \mathbb{G}_m : v \cdot gv = hv \cdot ghv \right\}.$ Since

$$1 \to T \to \mathcal{R}_{L/\mathbb{Q}} \mathbb{G}_m \xrightarrow{v \mapsto \frac{v \cdot gv}{hv \cdot ghv}} T_{L_1/\mathbb{Q}} \to 1$$

is exact, we have $C(T) = D_2 D_3$.

- (v) $G_T = \langle g \rangle \cong C_6$: $L_6 = L$ has the unique cubic subfield $L_3 = L^{g^3}$ and the unique quadratic subfield $L_2 = L^{g^2}$. T is one of the following types.
 - (a) $G_T = H_{6,a} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 1 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times (T_{L_6/L_2} \cap T_{L_6/L_3}) \text{ and } C(T) = \frac{D_6}{D_2 D_3}.$
 - (b) $G_T = H_{6,b} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & -1 \end{pmatrix} \right\rangle : T = T_{L_2/\mathbb{Q}} \times T_{L_3/\mathbb{Q}} \text{ and } C(T) = D_2 D_3.$

(c)
$$G_T = H_{6,c} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 1 \end{pmatrix} \right\rangle : T = T_{L_2/\mathbb{Q}} \times (T_{L_6/L_2} \cap T_{L_6/L_3}) \text{ and } C(T) = \frac{D_6}{D_3}.$$

(d)
$$G_T = H_{6,d} := \left\langle \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ -1 & 0 & 0 \end{pmatrix} \right\rangle : T = T_{L_6/L_3} \text{ and } C(T) = \frac{D_6}{D_3}.$$

(vi) $G_T = \langle g, h : g^3 = h^2 = (gh)^2 = 1 \rangle \cong S_3$: $L_6 = L$ has 3 isomorphic cubic subfields L^h, L^{gh} and L^{g^2h} (denoted by L_3) and the unique quadratic subfield $L_2 = L^g$. Then $D_6 = D_3^2 D_2$ by the equation (15). T is one of the following types.

(e)
$$G_T = H_{6,e} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L_2/\mathbb{Q}} \times (T_{L_6/L_2} \cap T_{L_6/L_3}) \text{ and } C(T) = D_2 D_3.$$

- (f) $G_T = H_{6,f} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times T_{L_3/\mathbb{Q}} \text{ and } C(T) = D_3.$
- (g) $G_T = H_{6,g} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\rangle : T = T_{L_2/\mathbb{Q}} \times T_{L_3/\mathbb{Q}} \text{ and } C(T) = D_2 D_3.$
- (h) $G_T = H_{6,h} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times (T_{L_6/L_2} \cap T_{L_6/L_3}) \text{ and } C(T) = D_3.$
- (i) $G_T = H_{6,i} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 \\ -1 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \right\rangle : T = T_{L_6/L_3} \text{ and } C(T) = D_2 D_3.$
- (j) $G_T = H_{6,j} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \right\rangle : T = \mathcal{R}_{L_3/\mathbb{Q}} \mathbb{G}_m \text{ and } C(T) = D_3.$

(vii) $G_T = \langle g, h : g^4 = h^2 = 1, gh = hg \rangle \cong C_4 \times C_2 :$

$$L_4 = L^h, L'_4 = L^{g^2h}, L''_4 = L^{g^2}$$

are quartic subfields of L and

$$L_2 = L^g, L'_2 = L^{\langle g^2, h \rangle}, L''_2 = L^{gh}$$

are quadratic subfields of L. Since $\operatorname{Gal}(L/L'_2) \cong C_2^2$ and $\operatorname{Gal}(L''_4/\mathbb{Q}) \cong C_2^2$, we have the following formulas by the equation (14):

$$D_L D_2^{\prime 2} = D_4 D_4^{\prime} D_4^{\prime \prime} \tag{22a}$$

$$D_4'' = D_2 D_2' D_2''. (22b)$$

T is one of the following types.

(a)
$$G_T = H_{8,a} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_2/\mathbb{Q}} \times T_{L_4'/L_2'} \text{ and } C(T) = D_2 \frac{D_4'}{D_2'}.$$

(b) $G_T = H_{8,b} := \left\langle \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle$: $T = T_{L/L_4} \cap T_{L/L_2''}$ and $C(T) = \frac{D_L}{D_4 D_2''} = D_2 \frac{D_4'}{D_2'}$ by the equations (22a) and (22b).

(viii) $G_T = \langle g_1, g_2, g_3 \rangle \cong C_2^3$: For $\{i, j, k\} = \{1, 2, 3\},\$

$$M_i = L^{g_i}, \ M_{ij} = L^{g_i g_j}, \ M = L^{g_1 g_2 g_3}$$

are quartic subfields of L and

$$K_i = L^{\langle g_i, g_j g_k \rangle}, \ K_{ij} = L^{\langle g_i, g_j \rangle}, \ K = L^{\langle g_1 g_2, g_1 g_3 \rangle}$$

are quadratic subfields of L. Since M_{ij}/\mathbb{Q} , M/\mathbb{Q} and L/K_3 are Galois extensions with Galois groups C_2^2 , we have the following formulas by the equation (14):

$$D_{M_{ij}} = D_{K_k} D_{K_{ij}} D_K \tag{23a}$$

$$D_M = D_{K_1} D_{K_2} D_{K_3} \tag{23b}$$

$$D_L D_{K_3}^2 = D_{M_3} D_{M_{12}} D_M. (23c)$$

 ${\cal T}$ is one of the following types.

- (c) $G_T = H_{8,c} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{K_1/\mathbb{Q}} \times T_{K/\mathbb{Q}} \times T_{K_2/\mathbb{Q}} \text{ and } C(T) = D_K D_{K_1} D_{K_2}.$
- (d) $G_T = H_{8,d} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{K_1/\mathbb{Q}} \times T_{M_{13}/K_{13}} \text{ and } C(T) = D_{K_1} \frac{D_{M_{13}}}{D_{K_{13}}} = D_K D_{K_1} D_{K_2}.$

(e)
$$G_T = H_{8,e} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 1 & 0 & -1 \\ -1 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle :$$

$$T = \left\{ (v_1, v_2) \in T_{M_{13}/K_{13}} \times T_{M/K_3} : v_1 \cdot g_2 v_1 = v_2 \cdot g_2 v_2 \right\}.$$

Since

 $1 \to T \to T_{M_{13}/K_{13}} \times T_{M/K_3} \xrightarrow{(v_1, v_2) \mapsto \frac{v_1 \cdot g_2 v_1}{v_2 \cdot g_2 v_2}} T_{K_2/\mathbb{Q}} \to 1$

is exact, we have $C(T) = \frac{D_{M_{13}}}{D_{K_{13}}} \cdot \frac{D_M}{D_{K_3}} \cdot D_{K_2}^{-1} = D_K D_{K_1} D_{K_2}$ by the equations (23a) and (23b).

(f) $G_T = H_{8,f} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 1 & -1 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L/K_{12}} \cap T_{L/M_3} \text{ and } C(T) = \frac{D_L}{D_{K_{12}} D_{M_3}} = D_K D_{K_1} D_{K_2}$ by the equations (23a), (23b) and (23c).


(ix) $G_T = \langle g, h : g^4 = h^2 = (gh)^2 = 1 \rangle \cong D_4$:

$$M = L^{g^2}, M_1 = L^h, M'_1 = L^{g^2h}, M_2 = L^{gh}, M'_2 = L^{g^3h}$$

are quartic subfields of $L(M'_i)$ are Galois conjugates of M_i so $D_{M_i} = D_{M'_i}$ and

$$K = L^g, K_1 = L^{\langle g^2, h \rangle}, K_2 = L^{\langle g^2, gh \rangle}$$

are quadratic subfields of L. Then we have the following lattice of subfields of L.



Since L/K_1 , L/K_2 and M/\mathbb{Q} are Galois extensions with Galois groups C_2^2 , we have the following formulas by the equation (14):

$$D_L D_{K_1}^2 = D_{M_1}^2 D_M \tag{24a}$$

$$D_L D_{K_2}^2 = D_{M_2}^2 D_M \tag{24b}$$

$$D_M = D_K D_{K_1} D_{K_2} \tag{24c}$$

$$\frac{D_{M_1}}{D_{K_1}} = \frac{D_{M_2}}{D_{K_2}}.$$
 (24d)

The last equation is a consequence of the equations (24a) and (24b) (cf. [1, Proposition 2.4].) T is one of the following types.

(g)
$$G_T = H_{8,g} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = T_{K/\mathbb{Q}} \times T_{M_2'/K_2} \text{ and } C(T) = \frac{D_K D_{M_2}}{D_{K_2}}$$

(h)
$$G_T = H_{8,h} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{G}_m \times T_{M_2/K_2} \text{ and } C(T) = \frac{D_{M_2}}{D_{K_2}}$$

(i)
$$G_T = H_{8,i} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = T_{K_2/\mathbb{Q}} \times T_{M_2/K_2} \text{ and } C(T) = D_{M_2}.$$

(j)
$$G_T = H_{8,j} := \left\langle \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{K_1/\mathbb{Q}} \times T_{M_2'/K_2} \text{ and } C(T) = \frac{D_{K_1}D_{M_2}}{D_{K_2}} = D_{M_1}.$$

- (k) $G_T = H_{8,k} := \left\langle \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle$: $T = T_{L/K_1} \cap T_{L/M'_2}$ and $C(T) = \frac{D_L}{D_{K_1} D_{M_2}} = \frac{D_K D_{M_2}}{D_{K_2}}$ by the equations (24b) and (24c).
- (l) $G_T = H_{8,l} := \left\langle \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \left\{ v \in \mathbf{R}_{M'_2/\mathbb{Q}} \, \mathbb{G}_m : v \cdot g^2 v = gv \cdot g^3 v \right\}.$ Since

$$1 \to T \to \mathbf{R}_{M_2'/\mathbb{Q}} \,\mathbb{G}_m \xrightarrow{v \mapsto \frac{v \cdot g^2 \cdot v}{g v \cdot g^3 v}} T_{K_2/\mathbb{Q}} \to 1$$


is exact, we have $C(T) = \frac{D_{M_2}}{D_{K_2}}$.

- (m) $G_T = H_{8,m} := \left\langle \begin{pmatrix} -1 & 0 & -1 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{M'_2/\mathbb{Q}} \text{ and } C(T) = D_{M_2}.$
- (n) $G_T = H_{8,n} := \left\langle \begin{pmatrix} -1 & 0 & -1 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle$: $T = T_{L/M'_2} \cap T_{L/K}$ and $C(T) = \frac{D_L}{D_{M_2}D_K} = D_{M_1}$ by the equations (24b), (24c) and (24d).

(x) $G_T = \langle g, h : g^6 = h^2 = 1, gh = hg \rangle \cong C_6 \times C_2 : L_6 = L^{g^3h}$ has the unique cubic subfield $L_3 = L^{\langle g^3, h \rangle}$ and the unique quadratic subfield $L_2 = L^{\langle g^2, gh \rangle}$. Also $L'_2 = L^g$ is a quadratic subfield of L such that $L = L_6 L'_2$. T is the following type.

(a)
$$G_T = H_{12,a} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_2'/\mathbb{Q}} \times (T_{L_6/L_3} \cap T_{L_6/L_2}) \text{ and } C(T) = D_2' \frac{D_6}{D_3 D_2}.$$

(xi) $G_T = \langle g, h : g^6 = h^2 = (gh)^2 = 1 \rangle \cong D_6$: Consider the following lattice of subfields of L.



Since L_2/\mathbb{Q} , L/L_6 and L/L_8 are Galois extensions with Galois groups S_3 , we have the following formulas by the equation (15):

$$D_2 = D_4^2 D_7 (25a)$$

$$\frac{D_1^2}{D_6^2} = \frac{D_L}{D_5} = \frac{D_3^2}{D_8^2}.$$
(25b)

T is one of the following types.

$$\begin{array}{l} \text{(b)} \ G_{T} = H_{12,b} := \left\langle \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 \end{pmatrix} \right\rangle : T = T_{L_{7}/\mathbb{Q}} \times (T_{L_{1}/L_{4}} \cap T_{L_{1}/L_{6}}) \text{ and } C(T) = \frac{D_{1}D_{7}}{D_{4}D_{6}}. \\ \text{(c)} \ G_{T} = H_{12,c} := \left\langle \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & -1 & -1 & 0 \end{pmatrix} \right\rangle : T = \mathbb{G}_{m} \times (T_{L_{3}/L_{4}} \cap T_{L_{3}/L_{8}}) \text{ and } C(T) = \frac{D_{3}}{D_{4}D_{8}} = \frac{D_{1}}{D_{4}D_{6}}. \\ \text{(d)} \ G_{T} = H_{12,d} := \left\langle \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\rangle : T = T_{L_{6}/\mathbb{Q}} \times (T_{L_{2}/L_{4}} \cap T_{L_{2}/L_{7}}) \text{ and } C(T) = \frac{D_{2}D_{6}}{D_{4}D_{7}} = D_{4}D_{6}. \\ \text{(e)} \ G_{T} = H_{12,e} := \left\langle \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L_{6}/\mathbb{Q}} \times (T_{L_{1}/L_{4}} \cap T_{L_{1}/L_{6}}) \text{ and } C(T) = \frac{D_{1}}{D_{4}}. \\ \text{(f)} \ G_{T} = H_{12,f} := \left\langle \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \right\rangle : T = T_{L_{6}/\mathbb{Q}} \times (T_{L_{1}/L_{4}} \cap T_{L_{1}/L_{6}}) \text{ and } C(T) = \frac{D_{1}}{D_{4}}. \\ \text{(g)} \ G_{T} = H_{12,g} := \left\langle \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L_{6}/\mathbb{Q}} \times (T_{L_{3}/L_{4}} \cap T_{L_{3}/L_{8}}) \text{ and } C(T) = \frac{D_{3}}{D_{4}D_{8}} = \frac{D_{1}}{D_{4}}. \\ \text{(h)} \ G_{T} = H_{12,h} := \left\langle \begin{pmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right\rangle : T = T_{L_{6}/\mathbb{Q}} \times (T_{L_{3}/L_{4}} \cap T_{L_{3}/L_{8}}) \text{ and } C(T) = \frac{D_{3}}{D_{4}D_{8}} = \frac{D_{1}}{D_{4}}. \\ \end{array}$$

(xii) $G_T = \left\langle g, h : g^3 = h^2 = (gh)^3 = 1 \right\rangle \cong A_4$: Let

$$L_6 = L^h, L'_6 = L^{g^{-1}hg}, L_4 = L^{ghg}, L'_4 = L^{gh} \text{ and } L_3 = L^{\langle h, g^{-1}hg \rangle}$$

We have $D_6 = D'_6$, $D_4 = D'_4$ and the following formulas by the equations (16b) and (16c):

$$D_6 = D_3 D_4 \tag{26a}$$

$$D_L = D_4^2 D_6.$$
 (26b)

 ${\cal T}$ is one of the following types.

- (i) $G_T = H_{12,i} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_6/L_3} \text{ and } C(T) = \frac{D_6}{D_3} = D_4.$
- (j) $G_T = H_{12,j} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 1 \\ 0 & -1 & 0 \\ 1 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L_4/\mathbb{Q}} \text{ and } C(T) = D_4.$

(k) $G_T = H_{12,k} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & -1 & -1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = \left\{ v \in T_{L/L'_6} : hv \cdot gv = g^2v \right\}.$ Since

$$1 \to T \to T_{L/L'_6} \xrightarrow{v \mapsto \frac{hv \cdot gv}{g^2 v}} T_{L'_4/\mathbb{Q}} \to 1$$

is exact, we have $C(T) = \frac{D_L}{D_6 D_4} = D_4.$

(xiii) $G_T = \langle g, h : g^4 = h^2 = (gh)^2 = 1 \rangle \times \langle i \rangle \cong D_4 \times C_2 : L_8 = L^{hgi}$ is an octic subfield of L,

$$L_4 = L^{\langle hg,i \rangle}, L'_4 = L^{\langle hg,g^2i \rangle}, L''_4 = L^{\langle gh,g^2i \rangle}, L''_4 = L^{\langle hgi,g^2i \rangle}$$

are quartic subfields of ${\cal L}$ and

$$L_2 = L^{\langle h, gi \rangle}, L'_2 = L^{\langle hg, g^2, i \rangle}, L''_2 = L^{\langle g, hi \rangle}$$

are quadratic subfields of L. L_4'' is a Galois conjugate of L_4' so $D_4'' = D_4'$. Since L_8/L_2' and L_4'''/\mathbb{Q} are Galois extensions with Galois groups C_2^2 , we have the following formulas by the equation (14):

$$D_8 D_2^{\prime 2} = D_4 D_4^{\prime} D_4^{\prime \prime \prime} \tag{27a}$$

$$D_4''' = D_2 D_2' D_2''. (27b)$$

T is one of the following types.

(a)
$$G_T = H_{16,a} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = T_{L_2''/\mathbb{Q}} \times T_{L_4'/L_2'} \text{ and } C(T) = D_2'' \frac{D_4'}{D_2'}.$$

(b) $G_T = H_{16,b} := \left\langle \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_8/L_4} \cap T_{L_8/L_2} \text{ and } C(T) = \frac{D_8}{D_4 D_2} = D_2'' \frac{D_4'}{D_2'}$ by the equations (27a) and (27b).

(xiv) $G_T = \langle g, h : g^3 = h^2 = (gh)^3 = 1 \rangle \times \langle i \rangle \cong A_4 \times C_2$: Let $L_{12} = L^{hi}$, $L_8 = L^{hg}$, $L_6 = L^{\langle h, ghg^{-1} \rangle}$, $L_6 = L^{\langle h, i \rangle}$, $L_4 = L^{\langle hg, i \rangle}$, $L_3 = L^{\langle h, ghg^{-1}, i \rangle}$ and $L_2 = L^{\langle g, h \rangle}$. T is one of the following types.

(a)
$$G_T = H_{24,a} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_6/L_3} \text{ and } C(T) = \frac{D_6}{D_3}.$$

(b)
$$G_T = H_{24,b} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 1 \\ 0 & -1 & 0 \\ 1 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_8/L_4} \cap T_{L_8/L_2} \text{ and } C(T) = \frac{D_8}{D_4 D_2}.$$

(c)
$$G_T = H_{24,c} := \left\langle \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & -1 & -1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = \left\{ v \in T_{L_{12}/L_6'} : v \cdot hgv = g^2v \right\}.$$
 Since

$$1 \to T \to T_{L_{12}/L_{6}'} \xrightarrow{v \mapsto \frac{v \cdot hgv}{g^{2}v}} T_{L_{8}/L_{4}} \cap T_{L_{8}/L_{2}} \to 1$$

is exact, we have $C(T) = \frac{D_{12}D_4D_2}{D_8D'_6}$.


One can prove that the number C(T) for each torus of (a), (b) and (c) are equal.

Lemma 3.2.
$$\frac{D_6}{D_3} = \frac{D_8}{D_4 D_2} = \frac{D_{12} D_4 D_2}{D_8 D_6'}.$$

Proof. We need to consider some more subfields of L for the proof. Let

$$L'_{12} = L^h, L''_{12} = L^i, L''_6 = L^{\langle h, ghg^{-1} \rangle} \text{ and } L'''_6 = L^{\langle hi, g^{-1}hg \rangle}.$$

Since $\langle \tilde{h}i, g^{-1}\tilde{h}g \rangle = \langle hi, ghg^{-1} \rangle$ for $\tilde{h} := g^{-1}hg$, L_6''' is a Galois conjugate of L_6 so $D_6''' = D_6$. Consider the following lattices of subfields.



By the above diagrams, we have the following formulas by the equations (14) and (16c):

$$D_L D_2^2 = D_{12}' D_8^2 \tag{28a}$$

$$D_{12}'' = D_6' D_4^2 \tag{28b}$$

$$D_L D_6^{\prime 2} = D_{12} D_{12}^{\prime} D_{12}^{\prime \prime} \tag{28c}$$

$$D_{12}D_3^2 = D_6^2 D_6'. (28d)$$

Let
$$c_1 = \frac{D_6}{D_3}$$
, $c_2 = \frac{D_8}{D_4 D_2}$ and $c_3 = \frac{D_{12} D_4 D_2}{D_8 D'_6}$. Then we have
 $c_2^2 = \frac{D_8^2}{D_4^2 D_2^2} \stackrel{(28a)}{=} \frac{D_L}{D'_{12} D_4^2} \stackrel{(28c)}{=} \frac{D_{12} D''_{12}}{D'_6^2 D_4^2} \stackrel{(28b)}{=} \frac{D_{12}}{D'_6} (= c_2 c_3) \stackrel{(28d)}{=} \frac{D_6^2}{D_3^2} (= c_1^2),$

which implies that $c_1 = c_2 = c_3$.

(xv) $G_T = \langle g, h : g^6 = h^2 = (gh)^2 = 1 \rangle \times \langle i \rangle \cong D_6 \times C_2$: Let $L_6 = L^{\langle hg, g^3 i \rangle}$, $L_3 = L^{\langle g^3, hg, i \rangle} \subset L_6$, $L_2 = L^{\langle hg, gi \rangle} \subset L_6$ and $L'_2 = L^{\langle g, hi \rangle}$. T is the following type.

(d)
$$G_T = H_{24,d} := \left\langle \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L'_2/\mathbb{Q}} \times (T_{L_6/L_3} \cap T_{L_6/L_2}) \text{ and } C(T) = D'_2 \frac{D_6}{D_3 D_2}.$$

(xvi) $G_T = \langle g, h : g^4 = h^2 = (gh)^3 = 1 \rangle \cong S_4$: Let

$$L_{12} = L^{ghg^{-1}}, L'_{12} = L^{g^{-1}hg}, L_8 = L^{hg}, L'_8 = L^{gh}, L_6 = L^{\langle h, g^2hg^2 \rangle}, L'_6 = L^{\langle ghg^{-1}, g^{-1}hg \rangle}, L''_6 = L^{ghg^{-1}}$$

and

$$L_4 = L^{\langle hg, ghg^{-1} \rangle}, L'_4 = L^{\langle gh, g^{-1}hg \rangle}, L_3 = L^{\langle h, g^2 \rangle}, L'_3 = L^{\langle g, hg^2h \rangle}, L_2 = L^{\langle g^2, gh \rangle}.$$

Since L'_i is a Galois conjugate of L_i for $i \in \{3, 4, 6, 8, 12\}$, we have $D'_i = D_i$ for these *i*. *T* is one of the following types.

(e)
$$G_T = H_{24,e} := \left\langle \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : T = T_{L_6'/L_3'} \text{ and } C(T) = \frac{D_6'}{D_3}.$$


(f) $G_T = H_{24,f} := \left\langle \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle : T = T_{L_6/L_3} \text{ and } C(T) = \frac{D_6}{D_3}.$

$$\begin{array}{l} \text{(g)} \ \ G_T = H_{24,g} := \left\langle \begin{pmatrix} 0 & -1 & 0 \\ 1 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : \ T = T_{L_8/L_4} \cap T_{L_8/L_2} \ \text{and} \ C(T) = \frac{D_8}{D_4 D_2}. \\ \text{(h)} \ \ G_T = H_{24,h} := \left\langle \begin{pmatrix} 0 & -1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \right\rangle : \ T = T_{L_4/\mathbb{Q}} \ \text{and} \ C(T) = D_4. \\ \text{(i)} \ \ G_T = H_{24,i} := \left\langle \begin{pmatrix} 1 & 2 & 1 & 0 \\ -2 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : \ T = T_{L_{12}/L_6} \cap T_{L_{12}/L_4} \ \text{and} \ C(T) = \frac{D_{12}}{D_6 D_4} \\ \text{(j)} \ \ G_T = H_{24,j} := \left\langle \begin{pmatrix} -1 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 \end{pmatrix} \right\rangle : \ T = \left\{ v \in T_{L_{12}'/L_6} : hv \cdot gv \cdot g^2v = 1 \right\}. \ \text{Since} \\ 1 \to T \to T_{L_{12}'/L_6} \xrightarrow{v \to hv \cdot gv \cdot g^2v} T_{L_8'/L_4'} \cap T_{L_8'/L_2} \to 1 \end{array}$$

is exact, we have
$$C(T) = \frac{D_{12}D_4D_2}{D_8D_6}$$
.

Lemma 3.3. $\frac{D_6}{D_3} = D_4 = \frac{D_{12}D_4D_2}{D_8D_6}.$

Proof. Let $L_{12}'' = L^{g^2}$, $L_{12}''' = L^{g^2hg^2h}$ and $L_6''' = L^{\langle g^2, hg^2h \rangle}$. Consider the following lattices of subfields.



By the above diagrams, we have the following formulas. Note that $L_6^c = L$ and $\operatorname{Gal}(L/L_6) \cong C_2^2$ so we can apply Proposition 2.4(4).

$$D_6 = D_3 D_4 \tag{29a}$$


$$D_{12}''D_2 = D_8 D_6''' \tag{29b}$$

$$\frac{D_{12}}{D_6} = \frac{D_{12}''}{D_6'''}.$$
(29c)

The first two formulas come from the equations (17) and (16b), and the last formula can be obtained exactly same as the equation (24d). Now the equation (29a) implies that $\frac{D_6}{D_3} = D_4$ and the equations (29b) and (29c) imply that $D_4 = \frac{D_{12}D_4D_2}{D_8D_6}$.

Lemma 3.4.
$$\frac{D_6''}{D_3} = \frac{D_8}{D_4 D_2} = \frac{D_{12}}{D_6 D_4}.$$

Proof. Let $L_6^{\prime\prime\prime\prime} = L^{\langle g^2, ghg^2h \rangle}$ and consider the following lattices of subfields. Since $(L_6^{\prime\prime\prime\prime\prime})^c = L$ and $\operatorname{Gal}(L/L_6^{\prime\prime\prime\prime}) \cong C_2^2$, we can apply Proposition 2.4(4).



By the above diagrams, we have the following formulas by the equations (14) and (17):

$$D_6^{\prime\prime\prime\prime} = D_3 D_4 \tag{30a}$$

$$D_{12}''D_3^2 = D_6''D_6'''D_6''''.$$
(30b)

Now we have

$$\frac{D_6''}{D_3} \stackrel{(30b)}{=} \frac{D_{12}'D_3}{D_6'''D_6''''} \stackrel{(30a)}{=} \frac{D_{12}''}{D_6'''D_4} \stackrel{(29c)}{=} \frac{D_{12}}{D_6D_4} \text{ and } \frac{D_{12}''}{D_6'''D_4} \stackrel{(29b)}{=} \frac{D_8}{D_4D_2}.$$

(xvii) $G_T = \langle g, h : g^4 = h^2 = (gh)^3 = 1 \rangle \times \langle i \rangle \cong S_4 \times C_2$: Let

$$L_{12} = L^{\langle ghg^{-1}, g^{-1}hgi \rangle}, L_8 = L^{\langle hg, ghg^{-1}i \rangle}, L_8' = L^{\langle gh, g^{-1}hg \rangle}, L_6 = L^{\langle g, hg^2hi \rangle}, L_6' = L^{\langle ghg^{-1}, g^{-1}hg, i \rangle}$$

and

$$L_{4} = L^{\langle hg, ghg^{-1}, i \rangle}, \ L'_{4} = L^{\langle gh, g^{-1}hg, i \rangle}, \ L_{3} = L^{\langle g, hg^{2}h, i \rangle}, \ L_{2} = L^{\langle gi, hi \rangle}, \ L'_{2} = L^{\langle g, h \rangle}.$$

 L'_4 is a Galois conjugate of L_4 so $D'_4 = D_4$. T is one of the following types.

(a)
$$G_T = H_{48,a} := \left\langle \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_6/L_3} \text{ and } C(T) = \frac{D_6}{D_3}.$$

(b) $G_T = H_{48,b} := \left\langle \begin{pmatrix} 0 & -1 & 0 \\ -1 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} -1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = T_{L_8/L_4} \cap T_{L_8/L_2} \text{ and } C(T) = \frac{D_8}{D_4 D_2}.$
(c) $G_T = H_{48,c} := \left\langle \begin{pmatrix} 1 & 1 & 0 \\ -1 & -1 & -1 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & -1 & -1 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & -1 \end{pmatrix} \right\rangle : T = \left\{ v \in T_{L_{12}/L_6'} : hgv \cdot g^2v = hv \right\}.$ Since

$$1 \to T \to T_{L_{12}/L_6'} \xrightarrow{v \mapsto \frac{hgv \cdot g^2 v}{hv}} T_{L_8'/L_4'} \cap T_{L_8'/L_2'} \to 1$$


is exact, we have $C(T) = \frac{D_{12}D_4D'_2}{D'_8D'_6}$.

Lemma 3.5. $\frac{D_6}{D_3} = \frac{D_8}{D_4 D_2} = \frac{D_{12} D_4 D_2'}{D_8' D_6'}.$

Proof. Let

$$L_{24} = L^i, L'_{12} = L^{\langle g^2, ghg^2 hi \rangle}, L''_6 = L^{\langle hg^2 h, gi \rangle} \text{ and } L'''_6 = L^{\langle g^2, ghg^2 h, i \rangle}.$$

Consider the following 3 lattices of subfields.



By the above diagrams, we have the following formulas by the equations (14) and (17):

$$D_{12}'D_2 = D_8 D_6'' \tag{31a}$$

 $D_6''' = D_4 D_3 \tag{31b}$

$$D_{12}'D_3^2 = D_6 D_6'' D_6'''. (31c)$$

These equations imply that

$$\frac{D_6}{D_3} \stackrel{(31c)}{=} \frac{D_{12}' D_3}{D_6'' D_6'''} \stackrel{(31a)}{=} \frac{D_8 D_3}{D_6'' D_2} \stackrel{(31b)}{=} \frac{D_8}{D_4 D_2}$$


Now we prove the second equality. Let

$$L'_{24} = L^{hg^2hi}, \ L''_{24} = L^{hg^2h}, \ L_{16} = L^{hg}, \ L''_{12} = L^{\langle hg^2h,i\rangle}, \ L'''_{12} = L^{\langle g^2,hg^2h\rangle}, \ L'''_{12} = L^{\langle g^{-1}hg,ghg^{-1}i\rangle}$$

and

$$L_8'' = L^{\left< hg, ghg^{-1} \right>}, \ L_8''' = L^{\left< hg, i \right>}, \ L_6'''' = L^{\left< g^2, hg^2h, i \right>}, \ L_4'' = L^{\left< gh, hg \right>}, \ L_2'' = L^{\left< gh, hg, i \right>}.$$

Since $L_{12}^{\prime\prime\prime\prime}$ is a Galois conjugate of L_{12} and $L_8^{\prime\prime}$ is a Galois conjugate of L_8^{\prime} , we have $D_{12}^{\prime\prime\prime\prime} = D_{12}$ and $D_8^{\prime\prime} = D_8^{\prime}$. Consider the following 6 lattices of subfields.



By the above diagrams, we have the following formulas by the equations (14) and (16c):

$$D_L D_4^{\prime\prime 2} = D_{24}^{\prime\prime} D_{16}^2 \tag{32a}$$

$$D_{24}D_2''^2 = D_{12}''D_8'''^2 \tag{32b}$$

$$D_L D_{12}^{\prime\prime 2} = D_{24} D_{24}^{\prime} D_{24}^{\prime\prime} \tag{32c}$$

$$D_{24}'D_6'^2 = D_{12}^2 D_{12}'' \tag{32d}$$

$$D_{16}D_4^2 = D_8 D_8' D_8''' \tag{32e}$$

$$D_4'' = D_2 D_2' D_2''. ag{32f}$$

These equations imply that

$$\begin{split} \left(\frac{D_8}{D_4 D_2}\right)^2 \stackrel{(32e)}{=} \frac{D_{16}^2 D_4^2}{D_8'^2 D_8''^2 D_2^2} \\ \stackrel{(32a)}{=} \frac{D_L D_4^2 D_4''^2}{D_{24}'' D_8''^2 D_8'''^2 D_2^2} \\ \stackrel{(32c)}{=} \frac{D_2 4 D_{24}' D_4'^2 D_8'''^2 D_2^2}{D_{12}'' D_8'' D_2'''^2 D_2^2} \\ \stackrel{(32b)}{=} \frac{D_{24}' D_4' D_4' D_4''^2}{D_{12}' D_8'^2 D_2' D_2'''^2} \\ \stackrel{(32d)}{=} \frac{D_{12}^2 D_4^2 D_4''^2}{D_8'' D_6'' D_2'' D_2'''^2} \\ \stackrel{(32f)}{=} \left(\frac{D_{12} D_4 D_2'}{D_8' D_6'}\right)^2, \end{split}$$

which proves the second equality.

4 Counting algebraic tori over \mathbb{Q} of dimension 3

In this section, we count the number of the isomorphism classes of 3-dimensional tori over \mathbb{Q} . We provide asymptotic upper and lower bounds of $N_3^{\text{tor}}(X; H)$ for each finite subgroup $H \neq 1$ of $\text{GL}_3(\mathbb{Z})$. See Table 1 for the computation of the numbers a(H) and b(H) for each H. This gives an asymptotic upper bound of $N_3^{\text{tor}}(X)$ which is close to the asymptotics predicted by Conjecture 1.1.

H	H	a(H)	b(H)	H	H	a(H)	b(H)	H	H	a(H)	b(H)
2	$H_{2,a}$	2	1	6	$H_{6,c}$	2	1	12	$H_{12,c}$	1	2
	$H_{2,b}$	1	1		$H_{6,d}$	2	1		$H_{12,d}$	1	2
	$H_{2,c}$	2	1		$H_{6,e}$	2	2		$H_{12,e}$	1	2
	$H_{2,d}$	1	1		$H_{6,f}$	1	1		$H_{12,f}$	1	1
	$H_{2,e}$	3	1		$H_{6,g}$	2	2		$H_{12,g}$	1	1
	$H_{3,a}$	1	1		$H_{6,h}$	1	1		$H_{12,h}$	1	1
5	$H_{3,b}$	1	1		$H_{6,i}$	2	2		$H_{12,i}$	2	2
	$H_{4,a}$	2	2		$H_{6,j}$	1	1		$H_{12,j}$	2	2
	$H_{4,b}$	2	1	8	$H_{8,a}$	1	1		$H_{12,k}$	2	2
	$H_{4,c}$	2	2		$H_{8,b}$	1	1	16	$H_{16,a}$	1	3
	$H_{4,d}$	2	1		$H_{8,c}$	1	3		$H_{16,b}$	1	3
	$H_{4,e}$	1	1		$H_{8,d}$	1	3	24	$H_{24,a}$	1	1
	$H_{4,f}$	2	3		$H_{8,e}$	1	3		$H_{24,b}$	1	1
	$H_{4,g}$	1	2		$H_{8,f}$	1	3		$H_{24,c}$	1	1
4	$H_{4,h}$	2	3		$H_{8,g}$	2	4		$H_{24,d}$	1	3
	$H_{4,i}$	1	2		$H_{8,h}$	1	2		$H_{24,e}$	2	4
	$H_{4,j}$	1	2		$H_{8,i}$	1	1		$H_{24,f}$	1	1
	$H_{4,k}$	1	1		$H_{8,j}$	1	1		$H_{24,g}$	2	4
	$H_{4,l}$	2	3		$H_{8,k}$	2	4		$H_{24,h}$	1	1
	$H_{4,m}$	1	2		$H_{8,l}$	1	2		$H_{24,i}$	2	4
	$H_{4,n}$	2	3		$H_{8,m}$	1	1		$H_{24,j}$	1	1
	$H_{4,o}$	1	2		$H_{8,n}$	1	1	48	$H_{48,a}$	1	2
6	$H_{6,a}$	2	3	12	$H_{12,a}$	1	1		$H_{48,b}$	1	2
	$H_{6,b}$	1	1		$H_{12,b}$	2	5		$H_{48,c}$	1	2

Table 1: Computations of a(H) and b(H)

Throughout this section, C denotes a positive constant which may change from line to line. Denote the set of positive squarefree integers by Sqf. For a prime p, denote by Sqf_p the set of positive integers which are squarefree outside p. We also introduce a notation which generalizes $N_n(X;G;I)$ defined in Section 2.1. Let J be an invariant of pairs of number fields such that for every X > 0, there are finitely many pairs of number fields (K_1, K_2) such that $J(K_1, K_2) \leq X$. Denote by $N_{n_1, n_2}^{K_1, K_2}(X;G_1, G_2;J) = N_{n_1, n_2}(X;G_1, G_2;J)$ the number of $(K_1, K_2) \in NF_{n_1}(G_1) \times NF_{n_2}(G_2)$ such that $J(K_1, K_2) \leq X$. When $n_i = 2$, we may omit the group $G_i = C_2$ as before.

4.1 Abelian case

In this section, we prove Conjecture 1.2 for every finite abelian subgroup $1 \neq H \leq \operatorname{GL}_3(\mathbb{Z})$. First we prove this except for the cases $H = H_{6,c}$ and $H = H_{6,d}$.

Proposition 4.1. Conjecture 1.2 holds for every finite abelian subgroup $1 \neq H \leq \operatorname{GL}_3(\mathbb{Z})$ which is not conjugate to $H_{6,c}$ or $H_{6,d}$.

Proof. By Proposition 2.1(1), the conjecture is true if $|H| \in \{2, 3\}$ or

$$H = H_{4,x} \ (x \in \{b, d, f, h, l, n\}).$$

By Propositions 2.1(1) and 2.3, the conjecture is true if H is one of

$$H_{4,x}$$
 ($x \in \{e, g, i, j, k, m, o\}$), $H_{6,b}$ and $H_{8,x}$ ($x \in \{c, d, e, f\}$).

For example, we have

$$N_{3}^{\text{tor}}(X; H_{4,e}) = \# \left\{ (L_{1}, L_{3}) \in \text{NF}_{2} \times \text{NF}_{2} : L_{1} \neq L_{3} \text{ and } D_{1}D_{3}^{2} \leq X \right\}$$
$$= N_{2,2}^{L_{1},L_{3}}(X; D_{1}D_{3}^{2}) - N_{2}^{L_{1}}(X; D_{1}^{3})$$
$$\sim CX - O(X^{\frac{1}{3}}) (\because \text{Proposition } 2.3)$$
$$\sim CX.$$

By Propositions 2.2(1)(2) and 2.3, the conjecture is true if H is one of

 $H_{4,a}, H_{4,c}, H_{6,a}, H_{8,a}, H_{8,b}$ and $H_{12,a}$.

For example, we have

$$N_3^{\text{tor}}(X; H_{8,a}) = \# \left\{ (L_2, L'_4) \in \operatorname{NF}_2 \times \operatorname{NF}_4(C_4) : L_2 \neq L'_2 \text{ and } D_2 \frac{D'_4}{D'_2} \leq X \right\}$$
$$= N_{2,4}^{L_2, L'_4}(X; C_4; D_2 \frac{D'_4}{D'_2}) - N_4(X; C_4)$$
$$=: A_1(X) - A_2(X)$$

where L'_2 is the unique quadratic subfield of L'_4 . Since $N_2(X) \sim CX$ and

$$N_4^{L'_4}(X; C_4; \frac{D'_4}{D'_2}) \sim CX^{\frac{1}{2}} \log X$$

by Proposition 2.2(1), we have $A_1(X) \sim CX$ by Proposition 2.3. Proposition 2.1(1) implies that $A_2(X) \sim CX^{\frac{1}{2}}$ so we have $N_3^{\text{tor}}(X; H_{8,a}) \sim CX$.

Now let L_6 be a cyclic sextic field with the cubic subfield L_3 and the quadratic subfield L_2 . Conjecture 1.2 for the remaining cases ($H_{6,c}$ and $H_{6,d}$) are equivalent to the asymptotic formula

$$N_6(X; C_6, \frac{D_6}{D_3}) \sim CX^{\frac{1}{2}}.$$
 (33)

In order to prove this, we need to use analytic methods. We give an explicit formula for the Dirichlet series of $\frac{D_6}{D_3}$ and apply Delange's Tauberian theorem (Proposition 2.6) to this.

Lemma 4.2.

$$\sum_{L_6 \in \text{NF}_6(C_6)} \left(\frac{D_6}{D_3}\right)^{-s} = \frac{h(s)g_1(s) - g_2(3s) - g_3(s) + 1}{2}$$
(34)

where

$$\begin{split} h(s) &:= \left(1 + \frac{1}{2^{6s}} + \frac{2}{2^{9s}}\right) \left(1 + \frac{1}{3^{3s}} + \frac{2}{3^{4s}} + \frac{2}{3^{5s}}\right),\\ g_1(s) &:= \prod_{p \equiv 1 \pmod{6}} \left(1 + \frac{2}{p^{2s}} + \frac{3}{p^{3s}}\right) \prod_{p \equiv -1 \pmod{6}} \left(1 + \frac{1}{p^{3s}}\right),\\ g_2(s) &:= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}}\right) \prod_{p \text{ odd}} \left(1 + \frac{1}{p^s}\right),\\ g_3(s) &:= \left(1 + \frac{2}{3^{4s}}\right) \prod_{p \equiv 1 \pmod{6}} \left(1 + \frac{2}{p^{2s}}\right). \end{split}$$

Proof. Denote the conductor of an abelian number field M by Cond(M). By the conductor-discriminant formula, we have $\text{Cond}(L_6) = \left(\frac{D_6}{D_2 D_3}\right)^{\frac{1}{2}}$. We also have

$$\operatorname{Cond}(L_6) = \operatorname{lcm}(\operatorname{Cond}(L_2), \operatorname{Cond}(L_3)) = \operatorname{lcm}(D_2, D_3^{\frac{1}{2}})$$

([26, (4)]) so

$$\frac{D_6}{D_3} = \operatorname{lcm}(D_2^3, D_2 D_3)$$

The Dirichlet series of quadratic fields and cyclic cubic fields are given by

$$\sum_{L_2 \in \mathrm{NF}_2} \frac{1}{D_2^s} = g_2(s) - 1$$

([10, Section 2]) and

$$\sum_{a_3 \in \mathrm{NF}_3(C_3)} \frac{1}{D_3^s} = \frac{g_3(s) - 1}{2}$$

([10, Section 3]). Let S_1 and S_2 be multisets of positive integers defined by

Ι

$$S_1 := \{D_2 : L_2 \in NF_2\} \cup \{1\}$$

$$S_2 := \{D_3 : L_3 \in NF_3(C_3)\} \cup \{D_3 : L_3 \in NF_3(C_3)\} \cup \{1\}$$

(union as multisets). Then we have $\sum_{n_1 \in S_1} \frac{1}{n_1^s} = g_2(s)$ and $\sum_{n_2 \in S_2} \frac{1}{n_2^s} = g_3(s)$ so

$$\sum_{L_6 \in \mathrm{NF}_6(C_6)} \left(\frac{D_6}{D_3}\right)^{-s} = \frac{1}{2} \sum_{\substack{(n_1, n_2) \in S_1 \times S_2 \\ n_1 \neq 1, n_2 \neq 1}} \frac{1}{\mathrm{lcm}(n_1^3, n_1 n_2)^s}$$
$$= \frac{1}{2} \sum_{\substack{(n_1, n_2) \in S_1 \times S_2 \\ (n_1, n_2) \in S_1 \times S_2}} \frac{1}{\mathrm{lcm}(n_1^3, n_1 n_2)^s} - \frac{1}{2} \sum_{n_1 \in S_1} \frac{1}{n_1^{3s}} - \frac{1}{2} \sum_{n_2 \in S_2} \frac{1}{n_2^s} + \frac{1}{2}$$
$$= \frac{h(s)g_1(s)}{2} - \frac{g_2(3s)}{2} - \frac{g_3(s)}{2} + \frac{1}{2}.$$

Proposition 4.3. Conjecture 1.2 holds for $H = H_{6,c}$ and $H = H_{6,d}$.

Proof. By [35, Theorem 1.1], $g_2(s) \in \mathcal{M}(1, \frac{1}{2})$ and $g_3(s) \in \mathcal{M}(\frac{1}{2}, \frac{1}{3})$. (See Section 2.3 for the definition of $\mathcal{M}(\alpha, \beta)$ for $\alpha > \beta > 0$.) Since

$$\frac{h(s)g_1(s) - g_3(s)}{g_3(s)} = \left(1 + \frac{1}{2^{6s}} + \frac{2}{2^{9s}}\right) \left(1 + \frac{3^{2s} + 2}{3^{5s} + 2 \cdot 3^s}\right) \prod_{p \equiv 1 \pmod{6}} \left(1 + \frac{3}{p^{3s} + 2p^s}\right) \prod_{p \equiv -1 \pmod{6}} \left(1 + \frac{1}{p^{3s}}\right) - 1$$

converges absolutely for $\operatorname{Re}(s) > \frac{1}{3}$ and

$$\lim_{s \to \frac{1}{2}} \frac{h(s)g_1(s) - g_3(s)}{g_3(s)} > 0,$$

we have $h(s)g_1(s) - g_3(s) \in \mathcal{M}(\frac{1}{2}, \frac{1}{3})$. Now Lemma 4.2 implies that

$$\sum_{L_6 \in \mathrm{NF}_6(C_6)} \left(\frac{D_6}{D_3}\right)^{-s} \in \mathcal{M}(\frac{1}{2}, \frac{1}{3})$$

so Proposition 2.6 finishes the proof.

4.2 Lower bound for non-abelian case

There are 39 conjugacy classes of finite non-abelian subgroups of $\operatorname{GL}_3(\mathbb{Z})$. By the computation of the numbers C(T) in Section 3 and Table 1, it is enough to consider the asymptotics of $N_3^{\operatorname{tor}}(X; H)$ when H is one of the following 16 subgroups of $\operatorname{GL}_3(\mathbb{Z})$:

• $H_{6,e}, H_{6,f}(S_3)$

•
$$H_{8,x}$$
 $(x \in \{g, h, i\})$ $(D_4), H_{16,a}$ $(D_4 \times C_2)$

- $H_{12,i}$ $(A_4), H_{24,b}$ $(A_4 \times C_2)$
- $H_{12,x}$ $(x \in \{b, c, d, f\})$ $(D_6), H_{24,d}$ $(D_6 \times C_2)$
- $H_{24,g}, H_{24,h}$ (S₄), $H_{48,b}$ (S₄ × C₂)

Proposition 4.4. Conjecture 1.2 holds for $H = H_{6,f}$, $H_{8,h}$, $H_{8,i}$, $H_{16,a}$, $H_{12,d}$ and $H_{24,h}$. Under the assumption of Malle's conjecture for quartic A_4 -fields, it also holds for $H = H_{12,i}$.

Proof. The cases $H_{6,f}$, $H_{8,i}$ and $H_{24,h}$ follow from Proposition 2.1(2), the case $H_{8,h}$ follows from Proposition 2.2(3) and the case $H_{12,d}$ follows from Propositions 2.1(2) and 2.3. When $G_T = H_{16,a}$, we have $C(T) = D_2'' \frac{D_4'}{D_2'}$ for $L_2'' \in NF_2$ and $L_4' \in NF_4(D_4)$ (with the unique quadratic subfield L_2') such that $L_2'' \cap L_4'' = \mathbb{Q}$. Therefore

$$\begin{split} N_3^{\text{tor}}(X; H_{16,a}) &= \# \left\{ (L_2'', L_4') \in \mathrm{NF}_2 \times \mathrm{NF}_4(D_4) : L_2'' \cap L_4'^c = \mathbb{Q} \text{ and } D_2'' \frac{D_4'}{D_2'} \leq X \right\} \\ &= N_{2,4}^{L_2'', L_4'}(X; D_4; D_2'' \frac{D_4'}{D_2'}) - \# \left\{ (L_2'', L_4') \in \mathrm{NF}_2 \times \mathrm{NF}_4(D_4) : L_2'' \subset L_4'^c \text{ and } D_2'' \frac{D_4'}{D_2'} \leq X \right\} \\ &\sim CX(\log X)^2 - O(X \log X) \\ &\sim CX(\log X)^2 \end{split}$$

by Propositions 2.2(3) and 2.3. The second statement is trivial because $N_3^{\text{tor}}(X; H_{12,i}) = N_4(X; A_4)$.

Remark 4.5. Unconditionally, we have

$$X^{\frac{1}{2}} \ll N_4(X; A_4) \ll_{\varepsilon} X^{\frac{1}{2} + \gamma + \varepsilon}$$

for $\gamma = 0.2784...$ by [4, Theorem 3] and [8, Theorem 1.4].

Let \mathcal{W} be the set of the 9 remaining cases, i.e.

 $\mathcal{W} := \{H_{6,e}, H_{8,g}, H_{12,b}, H_{12,c}, H_{12,f}, H_{24,b}, H_{24,d}, H_{24,g}, H_{48,b}\}.$

We provide upper and lower bounds of $N_3^{\text{tor}}(X; H)$ for each H in \mathcal{W} , which are summarized in Table 2 below. The upper bounds for $H = H_{12,b}$, $H_{12,c}$, $H_{24,d}$ are under the assumption of Conjecture 1.4 and the upper bound for $H_{24,g}$ is under the assumption of Malle's conjecture for the Galois group 12T8. The other upper and lower bounds are unconditional.

Н	Malle	Lower	Upper		
$H_{6,e}, H_{6,g}, H_{6,i}$	$X^{\frac{1}{2}}\log X$	$X^{\frac{1}{2}}$	$X^{\frac{1}{2}}(\log X)^2$		
$H_{8,g}, H_{8,k}$	$X^{\frac{1}{2}}(\log X)^3$	$X^{\frac{1}{2}}(\log X)^2$	$X^{\frac{3}{4}-\delta} \ (\delta > 0)$		
$H_{12,b}$	$X^{\frac{1}{2}}(\log X)^4$	$X^{\frac{1}{2}}\log X$	$X^{\frac{1}{2}}(\log X)^{6+\varepsilon}$		
$H_{12,c}$	$X \log X$	X	$X(\log X)^{1+\varepsilon}$		
$H_{12,f}, H_{12,g}, H_{12,h}$	X	X	X		
$H_{24,a}, H_{24,b}, H_{24,c}$	X	X	$X(\log X)^3 \log \log X$		
$H_{24,d}$	$X(\log X)^2$	$X \log X$	$X(\log X)^{2+\varepsilon}$		
$H_{24,e}, H_{24,g}, H_{24,i}$	$X^{\frac{1}{2}}(\log X)^3$	$X^{\frac{1}{2}}$	$X^{\frac{9}{10}}$		
$H_{48,a}, H_{48,b}, H_{48,c}$	$X \log X$	X	$X(\log X)^4 \log \log X$		

Table 2: Upper and lower bounds of $N_3^{\rm tor}(X;H)$

We concentrate on the lower bounds in this section. The result is satisfactory, in the sense that the asymptotic inequality $X^{\frac{1}{\alpha(H)}} \ll N_3^{\text{tor}}(X; H)$ holds for every finite subgroup $H \neq 1$ of $\text{GL}_3(\mathbb{Z})$. By Propositions 4.1, 4.3, 4.4 and Remark 4.5, it is enough to prove this for $H \in \mathcal{W}$. The case $H = H_{24,b}$ is considered separately because its proof relies on the work of Cohen and Thorne [14].

For a cyclic cubic field k, denote by $\mathcal{F}(k)$ the set of quartic A_4 -fields whose cubic resolvent is k. For every $K \in \mathcal{F}(k)$, there is $f(K) \in \text{Sqf}_2$ which satisfies $D_K = D_k f(K)^2$. For every $f \in \text{Sqf}_2$, denote

$$\mathcal{F}(k, f^2) := \left\{ K \in \mathcal{F}(k) : D_K = D_k f^2 \right\}.$$

Proposition 4.6. $N_3^{\text{tor}}(X; H_{24,b}) \gg X$.

Proof. Let *E* be the maximal real subfield of $\mathbb{Q}(\zeta_7)$. Then $E \in NF_3(C_3)$, $D_E = 49$ and $h_E = 1$. A prime $p \neq 7$ splits completely in *E* if and only if *p* is a cubic residue modulo 7, i.e. $p \equiv \pm 1 \pmod{7}$. Therefore

$$\frac{1}{3} + \sum_{K \in \mathcal{F}(E)} \frac{1}{f(K)^s} = \frac{1}{3} \left(1 + \frac{3}{2^{3s}} \right) \prod_{p \equiv \pm 1 \pmod{7}} \left(1 + \frac{3}{p^s} \right)$$

by [14, Theorem 1.4]. Ignoring the even parts, we obtain

$$\sum_{\gcd(f,14)=1} \frac{|\mathcal{F}(E,f^2)|}{f^s} = \sum_{\substack{K \in \mathcal{F}(E) \\ \gcd(f(K),14)=1}} \frac{1}{f(K)^s} \\ = \frac{1}{3} \prod_{p \equiv \pm 1 \pmod{7}} \left(1 + \frac{3}{p^s}\right) - \frac{1}{3} \\ \in \mathcal{M}(1,\frac{1}{2})$$
(35)

by Lemma 2.7.

Let f be a positive squarefree integer which is coprime to 14 and define $L_{2,f} := \mathbb{Q}(\sqrt{f})$ (so $D_{2,f} \in \{f, 4f\}$). Let $L_{4,f}$ be an element of $\mathcal{F}(E, f^2)$ and $L_{8,f} := L_{4,f}L_{2,f} \in \mathrm{NF}_8(A_4 \times C_2)$. For any prime p dividing f, the splitting type of p in $L_{4,f}$ is (1^21^2) or (2^2) by [14, Theorem 5.1] so $v_p(D_{8,f}) = 4$ by Proposition 2.5. Also $D_{8,f} \mid D_{4,f}^2 D_{2,f}^4$ so $v_7\left(\frac{D_{8,f}}{D_{4,f}D_{2,f}}\right) \leq 2$ and $v_2\left(\frac{D_{8,f}}{D_{4,f}D_{2,f}}\right) \leq 6$. These imply that $\frac{D_{8,f}}{D_{4,f}D_{2,f}} \leq cf$ (36)

for $c = 56^2$. Now we have

$$N_3^{\text{tor}}(X; H_{24,b}) = N_{4,2}^{L_4, L_2}(X; A_4; \frac{D_8}{D_4 D_2}) \ge \sum_{\substack{f \le \frac{X}{c} \\ \gcd(f, 14) = 1}} \left| \mathcal{F}(E, f^2) \right| \sim CX$$

by the relation (35), the inequality (36) and Proposition 2.6.

Lemma 4.7. Let L_2 , L_4 and L_8 be number fields as in (xvi) of Section 3. Then

$$v_p(D_8) \le v_p(D_4^3 D_2)$$

for every odd prime p. Here $v_p(m)$ denotes the exponent of p in m.

Proof. Assume that $v_p(D_8) > v_p(D_4^3D_2)$ for an odd prime p.

• $L_8 = L_4 L_2$ implies that $v_p(D_8) \le 2v_p(D_4) + 4v_p(D_2)$ so $v_p(D_2) = 1$. We also have

$$3v_p(D_4) + 1 < v_p(D_8) \le 2v_p(D_4) + 4$$

so $v_p(D_4) < 3$.

- The cubic resolvent of $L_4 \in NF_4(S_4)$ is $L_3 \in NF_3(S_3)$ and the quadratic resolvent of L_3 is L_2 . Therefore $v_p(D_4)$, $v_p(D_3)$ and $v_p(D_2)$ have the same parity so $v_p(D_3) = v_p(D_4) = 1$ and $v_p(D_6) = v_p(D_3D_4) = 2$.
- Let $c := v_p(D_8)$. Then $v_p(D_6'') = c 1$ and $v_p(D_{12}) = c + 1$ by Lemma 3.4. Since L/L_4 is a Galois extension with a Galois group S_3 , we have $D_L D_4^2 = D_{12}^2 D_8$ by Proposition 2.4(2) so $v_p(D_L) = 3c$.
- The relation $v_p(D_L) \ge 4v_p(D_6'')$ implies that $c \le 4$, which contradicts the assumption. This finishes the proof.

Theorem 4.8. For every $H \in \mathcal{W}$, the lower bound of $N_3^{\text{tor}}(X; H)$ is given as in Table 2. In particular we have

$$X^{\frac{1}{a(H)}} \ll N_3^{\text{tor}}(X;H)$$

for every finite nontrivial subgroup H of $GL_3(\mathbb{Z})$.

Proof. For each $H \in \mathcal{W}$, we use the notation as in Section 3.

• $H = H_{6,e}$: Since L_2 is the quadratic resolvent of $L_3 \in NF_3(S_3)$, we have $D_2 \leq D_3$ so

$$N_3^{\text{tor}}(X; H_{6,e}) = N_3^{L_3}(X; S_3; D_2 D_3) \ge N_3^{L_3}(X; S_3; D_3^2) \sim C X^{\frac{1}{2}}.$$

•
$$H = H_{8,g}$$
: Since $\left(\frac{D_K D_{M_2}}{D_{K_2}}\right)^2 \le D_M \left(\frac{D_{M_2}}{D_{K_2}}\right)^2 = D_L$ by the equation (24b), we have
 $N_3^{\text{tor}}(X; H_{8,g}) = N_8^L(X; D_4; \frac{D_K D_{M_2}}{D_{K_2}}) \ge N_8^L(X^2; D_4; D_L) \sim CX^{\frac{1}{2}} (\log X)^2$

by Proposition 2.1(4).

• $H = H_{12,b}$: Since L_7 is the quadratic resolvent of $L_4 \in NF_3(S_3)$, we have $D_7 \leq D_4$ so

$$N_{3}^{\text{tor}}(X; H_{12,b}) = N_{6}^{L_{1}}(X; D_{6}; \frac{D_{1}D_{7}}{D_{4}D_{6}})$$

$$\geq N_{6}^{L_{1}}(X; D_{6}; \frac{D_{1}}{D_{6}})$$

$$\geq \# \left\{ (L_{4}, L_{6}) \in \text{NF}_{3}(S_{3}) \times \text{NF}_{2} : D_{4}^{2}D_{6}^{2} \leq X \text{ and } L_{6} \not\subset L_{4}^{c} \right\}$$

$$\sim CX^{\frac{1}{2}} \log X - O(X^{\frac{1}{2}})$$

$$\sim CX^{\frac{1}{2}} \log X.$$

- $H \in \{H_{12,c}, H_{12,f}, H_{48,b}\}$: They can be proved as in the proof of [22, Theorem 4.9].
- $H = H_{24,b}$: See Proposition 4.6.
- $H = H_{24,d}$: Since D_6 has 3 subgroups of order 6, L_6^c has 3 quadratic subfields. For a constant M > 0 which satisfies $\# \{L'_2 \in NF_2 : D'_2 \le M\} = N_2(M) \ge 6$, we have

$$\begin{split} N_{3}^{\text{tor}}(X; H_{24,d}) &= \# \left\{ \left(L_{6}, L_{2}^{\prime} \right) \in \text{NF}_{6}(D_{6}) \times \text{NF}_{2} : D_{2}^{\prime} \frac{D_{6}}{D_{3} D_{2}} \leq X \text{ and } L_{2}^{\prime} \not\subset L_{6}^{c} \right\} \\ &\geq \sum_{\substack{L_{6} \in \text{NF}_{6}(D_{6}) \\ \frac{D_{6}}{D_{3} D_{2}} \leq \frac{X}{M}}} \left(N_{2} \left(X \frac{D_{3} D_{2}}{D_{6}} \right) - 3 \right) \\ &\geq \frac{1}{2} \sum_{\substack{L_{6} \in \text{NF}_{6}(D_{6}) \\ \frac{D_{6}}{D_{3} D_{2}} \leq \frac{X}{M}}} N_{2} \left(X \frac{D_{3} D_{2}}{D_{6}} \right) \\ &\gg \frac{X}{M} \log \frac{X}{M} \end{split}$$

by Propositions 1.3(2) and 2.3.

• $H = H_{24,g}$: By Lemma 4.7,

$$N_3^{\text{tor}}(X; H_{24,g}) = N_4^{L_4}(X; S_4; \frac{D_8}{D_4 D_2}) \ge N_4^{L_4}(CX; S_4; D_4^2) \sim CX^{\frac{1}{2}}.$$

4.3 Upper bound for non-abelian case

Now we move to the upper bounds. If H is one of $H_{12,b}$, $H_{12,c}$ and $H_{24,d}$, the Cohen-Lenstra heuristics for p = 3 improves the upper bound of $N_3^{\text{tor}}(X; H)$.

Proposition 4.9. (1) $N_3^{\text{tor}}(X; H) \ll_{\varepsilon} X^{a(H)\left(1 + \frac{\log 2 + \varepsilon}{\log \log X}\right)}$ for $H \in \{H_{12,b}, H_{12,c}, H_{24,d}\}$.

(2) Under the assumption of Conjecture 1.4, the upper bounds of $N_3^{\text{tor}}(X; H)$ for $H \in \{H_{12,b}, H_{12,c}, H_{24,d}\}$ are given as in Table 2.

Proof. The case $H = H_{12,c}$ follows from the equality $N_3^{\text{tor}}(X; H_{12,c}) = N_2^{\text{tor}}(X; H_{12,A})$ and Proposition 1.3. The function $f(X) = \frac{\log X}{\log \log X}$ is increasing on $X > e^e = 15.15...$ so

$$N_2^{\mathrm{tor}}(Y; H_{12,A}) \ll_{\varepsilon} Y X^{\frac{\log 2+\varepsilon}{\log \log X}}$$

for $X \ge Y \ge 16$. Since $N_3^{\text{tor}}(X; H_{24,d})$ is bounded above by the product distribution of $N_2^{\text{tor}}(X; H_{12,A})$ and $N_2(X)$, we have

$$N_3^{\text{tor}}(X; H_{24,d}) \le 2 \sum_{n \le X} N_2^{\text{tor}}(\frac{X}{n}; H_{12,A})$$
$$\ll_{\varepsilon} \sum_{n \le \frac{X}{16}} \frac{X}{n} X^{\frac{\log 2+\varepsilon}{\log \log X}} + \sum_{\frac{X}{16} < n \le X} N_2^{\text{tor}}(16; H_{12,A})$$
$$\ll_{\varepsilon} X^{1+\frac{\log 2+\varepsilon}{\log \log X}}.$$

If we assume Conjecture 1.4, then we have

$$N_3^{\text{tor}}(X; H_{24,d}) \le 2 \sum_{n \le X} N_2^{\text{tor}}(\frac{X}{n}; H_{12,A})$$
$$\ll_{\varepsilon} \sum_{n \le X} \frac{X}{n} (\log X)^{1+\varepsilon}$$
$$\ll_{\varepsilon} X (\log X)^{2+\varepsilon}.$$

Now consider the case $H = H_{12,b}$. Let $F, K, L, E, f, m_1, m_2, w(f)$ and g(X) be defined as in [22, Section 4.2]. Denote

$$A_1(X) := N_{3,2}^{F,K}(X; S_3; D_E \frac{D_F D_K^2}{m^2})$$

and

$$A_{2}(X;B) := \sum_{\substack{m_{1},m_{2} \in \text{Sqf} \\ (m_{1}m_{2},6)=1 \\ (m_{1},m_{2})=1}} \sum_{\substack{E \in \text{NF}_{2} \\ B \leq D_{E}f^{2} \leq X \\ B \leq D_{E}f < 2B}} \frac{X^{\frac{1}{2}}}{D_{E}f} h_{3}(E) \cdot 2^{w(f)}$$

for $B \leq X^{\frac{1}{2}}$. Following the arguments of [22, Section 4], one can show that

$$N_3^{\text{tor}}(X; H_{12,b}) \le A_1(\beta X)$$
 (37)

for $\beta := 2^9 3^3$,

$$A_1(X) \ll \sum_{i=0}^{\lfloor \log_2 X^{\frac{1}{2}} \rfloor} A_2(X; 2^i)$$
 (38)

and

$$A_{2}(X;B) \ll \frac{X^{\frac{1}{2}}}{B} \sum_{\substack{f \in \operatorname{Sqf}_{3} \\ f < 2B}} 2^{w(f)} \tau(f) g(\frac{2B}{f})$$
$$\ll_{\varepsilon} \frac{X^{\frac{1}{2}}}{B} \sum_{\substack{f \in \operatorname{Sqf}_{3} \\ f < 2B}} \tau(f)^{2} \frac{2B}{f} X^{\frac{1}{2}\left(\frac{\log 2+\varepsilon}{\log \log X}\right)} \quad ([22, \operatorname{Corollary} 4.4])$$
$$\ll_{\varepsilon} X^{\frac{1}{2}\left(1 + \frac{\log 2+\varepsilon}{\log \log X}\right)} \sum_{f < 2B} \frac{\tau(f)^{2}}{f}$$
(39)

for $B \leq X^{\frac{1}{2}}$. By summing over the intervals $f \in [2^j, 2^{j+1})$ for $0 \leq j \leq \log_2(2B)$, we have

$$\sum_{f<2B} \frac{\tau(f)^2}{f} \le \sum_{j=0}^{\lfloor \log_2(2B) \rfloor} \frac{1}{2^j} \sum_{f<2^{j+1}} \tau(f)^2 \\ \ll \sum_{j=0}^{\lfloor \log_2(2B) \rfloor} \frac{1}{2^j} 2^{j+1} (\log 2^{j+1})^3 \text{ (Lemma 2.8)} \\ \ll (\log B)^4.$$
(40)

The inequalities (37), (38), (39) and (40) imply that $N_3^{\text{tor}}(X; H_{12,b}) \ll_{\varepsilon} X^{\frac{1}{2} \left(1 + \frac{\log 2 + \varepsilon}{\log \log X}\right)}$.

If we assume Conjecture 1.4, then

$$A_{2}(X;B) \ll_{\varepsilon} \frac{X^{\frac{1}{2}}}{B} \sum_{\substack{f \in \operatorname{Sqf}_{3} \\ f < 2B}} 2^{w(f)} \tau(f) \cdot \frac{2B}{f} (\log B)^{1+\varepsilon} \quad ([22, \operatorname{Corollary} 4.8])$$

$$\ll_{\varepsilon} X^{\frac{1}{2}} (\log B)^{1+\varepsilon} \sum_{f < 2B} \frac{\tau(f)^{2}}{f}.$$
(41)

The inequalities (37), (38), (40) and (41) imply that $N_3^{\text{tor}}(X; H_{12,b}) \ll_{\varepsilon} X^{\frac{1}{2}} (\log X)^{6+\varepsilon}$.

We proceed by case-by-case analysis.

Proposition 4.10. The following estimates hold:

- (1) $N_3^{\text{tor}}(X; H_{6,e}) \ll X^{\frac{1}{2}} (\log X)^2.$
- (2) $N_3^{\text{tor}}(X; H_{12,f}) \ll X.$
- (3) $N_3^{\text{tor}}(X; H_{48,b}) \ll X(\log X)^4 \log \log X.$
- (4) $N_3^{\text{tor}}(X; H_{24,b}) \ll X(\log X)^3 \log \log X.$
- *Proof.* (1) By [13, Theorem 2.5], the number of $F \in NF_3(S_3)$ such that $D_F = D_E f^2$ for given $E \in NF_2$ and $f \in Sqf_3$ is bounded by $O(h_3(E) \cdot 2^{w(f)})$. This implies that

$$\begin{split} N_{3}^{\text{tor}}(X; H_{6,e}) &= N_{3}^{F}(X; S_{3}; D_{F} D_{E}) \\ &\ll \sum_{\substack{f \in \text{Sqf}_{3} \\ f < X^{\frac{1}{2}} \\ D_{E} \leq X^{\frac{1}{2}}}} h_{3}(E) 2^{w(f)} \\ &\ll \sum_{\substack{f < X^{\frac{1}{2}} \\ f \neq X^{\frac{1}{2}}}} \frac{X^{\frac{1}{2}}}{f} \tau(f) \ ([16, \text{Theorem 3}]) \\ &\ll X^{\frac{1}{2}} (\log X)^{2}. \end{split}$$

The last inequality can be deduced as in the inequality (40) above.

(2) By Proposition 2.1(3), we have

$$N_3^{\text{tor}}(X; H_{12,f}) = N_6^{L_1}(X; D_6; \frac{D_1}{D_4}) \le N_6^{L_1}(X^2; D_6; D_1) \sim CX$$

(3) Let $F \in NF_4(S_4)$, $K \in NF_2$ such that $K \cap F^c = \mathbb{Q}$ and $L = FK \in NF_8(S_4 \times C_2)$. Also let

$$S_{1} := \left\{ p > 3 : p \mid \gcd(D_{F}, D_{K}) \text{ and } p \text{ has splitting type } (1^{2}11), (1^{2}2) \text{ or } (1^{3}1) \text{ in } F \right\}$$

$$S_{2} := \left\{ p > 3 : p \mid \gcd(D_{F}, D_{K}) \text{ and } p \text{ has splitting type } (1^{2}1^{2}), (2^{2}) \text{ or } (1^{4}) \text{ in } F \right\}$$

$$m_{i} := \prod_{p \in S_{i}} p \quad (i = 1, 2).$$

Then Proposition 2.5 implies that

$$C(L) := \frac{D_L}{D_F D_K} = \frac{D_F D_K^3}{c m_1^2 m_2^4}$$

for a positive integer $c \leq \beta := 2^{12} 3^4$ (cf. [28, Table 3]) so

$$N_{3}^{\text{tor}}(X; H_{48,b}) = \# \{ (F, K) \in \text{NF}_{4}(S_{4}) \times \text{NF}_{2} : K \cap F^{c} = \mathbb{Q} \text{ and } C(L) \leq X \}$$

$$\leq N_{4,2}^{F,K}(\beta X; S_{4}; \frac{D_{F}D_{K}^{3}}{m_{1}^{2}m_{2}^{4}})$$

$$=: A_{1}(\beta X).$$
(42)

The number $A_1(X)$ can be bounded as in [22, Section 4.2].

• For given F, m_1 and m_2 , the number of $K \in NF_2$ which satisfies $m_1m_2 \mid D_K$ and $\frac{D_K}{m_1m_2} \leq \left(\frac{m_2X}{m_1m_2}\right)^{\frac{1}{3}}$ is at most $2\left(\frac{m_2X}{m_1m_2}\right)^{\frac{1}{3}}$

$$\left(\frac{m_2 X}{m_1 D_F}\right)^3$$
 is at most $2\left(\frac{m_2 X}{m_1 D_F}\right)^3$.

- For every $F \in NF_4(S_4)$ and its cubic resolvent $E \in NF_3(S_3)$, there is $f \in Sqf_2$ such that $D_F = D_E f^2$. By [14, Theorem 5.1], m_1 divides D_E and m_2 divides f.
- By [14, Theorem 1.4 and Proposition 6.4], the number of $F \in NF_4(S_4)$ such that $D_F = D_E f^2$ for given $E \in NF_3(S_3)$ and $f \in Sqf_2$ is bounded by $O(h_2(E) \cdot 3^{w(f)})$.

• If
$$\frac{m_2 X}{m_1 D_F} = \frac{m_2 X}{m_1 D_E f^2} \ge 1$$
, then $D_E f^2 \le \frac{m_2 X}{m_1} \le f X$ so $D_E f^2 \le X^2$.

Combining all of these, we obtain

$$A_1(X) \ll \sum_{i=0}^{\lfloor 2\log_2 X \rfloor} A_2(X; 2^i)$$
(43)

 for

$$A_{2}(X;B) := \sum_{\substack{m_{1},m_{2} \in \text{Sqf} \\ (m_{1}m_{2},6)=1 \\ (m_{1},m_{2})=1}} \sum_{\substack{E \in \text{NF}_{3}(S_{3}) \\ m_{1}|D_{E} \\ D_{E}f^{2} \leq \frac{m_{2}X}{m_{1}} \\ B \leq D_{E}f^{2} \leq 2B}} \left(\frac{m_{2}}{m_{1}} \cdot \frac{X}{B}\right)^{\frac{1}{3}} h_{2}(E) \cdot 3^{w(f)}.$$
(44)

Now we bound the right-hand side of the inequality (44).

• The inequality $B \leq D_E f^2 \leq \frac{m_2 X}{m_1}$ implies that $m_1 \leq \frac{m_2 X}{B}$. Therefore the sum $\sum_{m_1,m_2} \left(\frac{m_2}{m_1}\right)^{\frac{1}{3}}$ for given E and f is bounded by

$$\sum_{m_2|f} \sum_{m_1 \le \frac{m_2 X}{B}} \left(\frac{m_2}{m_1}\right)^{\frac{1}{3}} \ll \sum_{m_2|f} m_2^{\frac{1}{3}} \left(\frac{m_2 X}{B}\right)^{\frac{2}{3}}$$
$$= \left(\frac{X}{B}\right)^{\frac{2}{3}} \sum_{m_2|f} m_2$$
$$\ll \left(\frac{X}{B}\right)^{\frac{2}{3}} f \log \log X.$$
(45)

The last inequality is due to the classical upper bound $\sigma(n) \ll n \log \log n$ of Gronwall [21].

• The inequality (45) implies that

$$A_{2}(X;B) \ll \frac{X \log \log X}{B} \sum_{f < (2B)^{\frac{1}{2}}} 3^{w(f)} f \sum_{\substack{E \in \mathrm{NF}_{3}(S_{3}) \\ D_{E} < \frac{2B}{f^{2}}}} h_{2}(E)$$

$$\ll \frac{X \log \log X}{B} \sum_{f < (2B)^{\frac{1}{2}}} 3^{w(f)} f \cdot \frac{2B}{f^{2}} \ ([6, \text{ Theorem 5}])$$

$$\ll X \log \log X \sum_{f < (2B)^{\frac{1}{2}}} \frac{\tau(f)^{c}}{f} \ (c := \frac{\log 3}{\log 2})$$

$$\ll X \log \log X (\log B)^{3}.$$
(46)

The last inequality can be deduced using Lemma 2.8 as in the inequality (40).

By the inequalities (42), (43) and (46), we have

$$N_3^{\text{tor}}(X; H_{48,b}) \ll \sum_{i=0}^{\lfloor 2 \log_2(\beta X) \rfloor} X \log \log X (\log 2^i)^3 \ll X (\log X)^4 \log \log X.$$

(4) Its proof is similar to the proof of (3). Following the arguments above, we have

$$N_3^{\text{tor}}(X; H_{24,b}) \ll \sum_{i=0}^{\lfloor 2 \log_2(\beta X) \rfloor} A_3(\beta X; 2^i)$$

for

$$A_3(X;B) := \frac{X \log \log X}{B} \sum_{f < (2B)^{\frac{1}{2}}} 3^{w(f)} f \sum_{\substack{E \in \mathrm{NF}_3(C_3) \\ D_E < \frac{2B}{f^2}}} h_2(E)$$

By [8, Theorem 1.1], $h_2(E) \ll D_E^{\frac{3}{10}}$ so

$$A_{3}(X;B) \ll \frac{X \log \log X}{B} \sum_{f < (2B)^{\frac{1}{2}}} 3^{w(f)} f\left(\frac{2B}{f^{2}}\right)^{\frac{2}{5}} \\ \ll \frac{X \log \log X}{B^{\frac{1}{5}}} \sum_{f < (2B)^{\frac{1}{2}}} \frac{\tau(f)^{c}}{f^{\frac{3}{5}}} \ (c = \frac{\log 3}{\log 2}) \\ \ll X \log \log X (\log B)^{2}$$

$$(47)$$

(following the inequality (40)) and

$$N_3^{\text{tor}}(X; H_{24,b}) \ll \sum_{i=0}^{\lfloor 2 \log_2(\beta X) \rfloor} X \log \log X (\log 2^i)^2 \ll X (\log X)^3 \log \log X.$$

4

Remark 4.11. Cohen-Martinet heuristics for cyclic cubic fields and p = 2 [12, p. 128] implies that

$$\sum_{\substack{E \in \mathrm{NF}_3(C_3) \\ D_E < \frac{2B}{f^2}}} h_2(E) \ll \frac{B^{\frac{1}{2}}}{f}.$$

However this does not improve the upper bound of the inequality (47).

So far, we have proved that

$$N_3^{\text{tor}}(X; H) \ll X^{\frac{1}{a(H)}} (\log X)^{O(1)}$$

(under the assumption of Conjecture 1.4 and Malle's conjecture for quartic A_4 -fields) if H is not conjugate to $H_{8,g}$ ($H_{8,k}$) or $H_{24,g}$ ($H_{24,e}$, $H_{24,i}$). Unfortunately, our upper bounds for the cases $H = H_{8,g}$ and $H = H_{24,g}$ are much weaker.

Proposition 4.12. $N_3^{\text{tor}}(X; H_{8,g}) \ll X^{\frac{3}{4}-\delta}$ for some $\delta > 0$.

Proof. We follow the notation in (ix) of Section 3. Let m_L be the product of odd primes whose splitting type in L is (1^41^4) or (2^4) . By [1, Table 1], $v_p(\frac{D_M}{D_W^2}) = v_p(m_L^2)$ for every odd prime p so

$$\left(\frac{D_K D_{M_2}}{D_{K_2}}\right)^2 \stackrel{(24b)}{=} D_L \frac{D_K^2}{D_M} = \frac{D_L}{2^{\alpha} m_L^2} \tag{48}$$

for $\alpha = v_2(\frac{D_M}{D_K^2}) \le 6.$

For an odd squarefree integer q > 0, denote by n(X;q) the number of octic D_4 -fields L such that $D_L \leq X$ and $m_L = q$. By [31, Corollary 3], we have

$$n(X;q) \ll_{\varepsilon} \frac{X^{\frac{1}{4}} (\log X)^2}{q^{\frac{3}{2}-\varepsilon}} + \frac{X^{\frac{1}{4}-\delta}}{q^{1+\delta}}$$
(49)

for some constant $\delta > 0$. Now the equation (48) and the inequality (49) imply that

$$\begin{split} N_{3}^{\text{tor}}(X; H_{8,g}) &= N_{8}^{L}(X; D_{4}; \frac{D_{K}D_{M_{2}}}{D_{K_{2}}}) \\ &\leq N_{8}^{L}(2^{6}X^{2}; D_{4}; \frac{D_{L}}{m_{L}^{2}}) \\ &\leq \sum_{q \leq X^{\frac{1}{2}}} n(2^{6}X^{2}q^{2}; q) \\ &\ll_{\varepsilon} \sum_{q \leq X^{\frac{1}{2}}} \left(\frac{X^{\frac{1}{2}}(\log X)^{2}}{q^{1-\varepsilon}} + \frac{X^{\frac{1}{2}-2\delta}}{q^{\frac{1}{2}+3\delta}} \right) \\ &\ll_{\varepsilon} X^{\frac{1}{2}+\varepsilon} + X^{\frac{3}{4}-\frac{7}{2}\delta}. \end{split}$$

Lemma 4.13. Let D_i (i = 2, 3, 4, 6, 8, 12) be as in (xvi) of Section 3. Then for every prime $p \ge 5$,

(1) $v_p(\frac{D_8}{D_4D_2}) \ge v_p(D_3f)$ for $f \in \text{Sqf}_2$ such that $D_4 = D_3f^2$. (2) $v_p(\frac{D_{12}}{D_6D_4}) \ge \frac{2}{9}v_p(D_{12})$.

Proof. (1) Assume that $v_p(\frac{D_8}{D_4D_2}) < v_p(D_3f)$ for a prime $p \ge 5$.

- Since $v_p(\frac{D_8}{D_4D_2}) \ge v_p(\frac{D_3f^2}{D_2})$, we have $v_p(D_2) = 1$ and $v_p(f) = 0$.
- L_2 is the quadratic resolvent of L_3 so $v_p(D_3) \le 2$ is odd. Therefore $v_p(D_3) = v_p(D_4) = 1$.
- Now $v_p(D_8) < v_p(D_4D_2D_3f) = 4 = v_p(D_2^4)$, which is impossible since $L_2 \subset L_8$.
- (2) Since $v_p(D_{12}) \ge \max(2v_p(D_6), 3v_p(D_4))$, it is enough to show that at least one of $v_p(D_6) \ge \frac{9}{5}v_p(D_4)$ or $v_p(D_6) \le \frac{4}{3}v_p(D_4)$ holds. If $v_p(f) = 0$, then $v_p(D_6) = v_p(D_4D_3) = 2v_p(D_4)$. If $v_p(f) = 1$, then $v_p(D_6) = 2v_p(D_4) - 2$. Since L_4 is tamely ramified at p, $v_p(D_4) \le 3$ so $v_p(D_6) \le \frac{4}{3}v_p(D_4)$.

Up to conjugation, there are two transitive subgroups of S_{12} which are isomorphic to S_4 : 12T8 and 12T9. (These groups can be found online at LMFDB [24].) For a number field K of degree 12 which satisfies $\operatorname{Gal}(K^c/\mathbb{Q}) \cong S_4$, the Galois group $\operatorname{Gal}(K^c/\mathbb{Q})$ is 12T8 in S_{12} if and only if K has a quartic subfield. Malle's conjecture predicts that

$$\begin{cases} N_{12}(X; 12T8) \sim CX^{\frac{1}{5}} \\ N_{12}(X; 12T9) \sim CX^{\frac{1}{4}}. \end{cases}$$

Proposition 4.14. (1) $N_3^{\text{tor}}(X; H_{24,g}) \ll X(\log X)^3$.

(2) Under the assumption of Malle's conjecture for 12T8, we have $N_3^{\text{tor}}(X; H_{24,g}) \ll X^{\frac{9}{10}}$.

Proof. (1) By Lemma 4.13,

$$N_3^{\text{tor}}(X; H_{24,g}) \le N_4^{L_4}(\beta X; S_4; D_3 f)$$

for some constant $\beta > 0$. As in the proof of Proposition 4.10, we have

$$N_3^{\text{tor}}(X; H_{24,g}) \ll \sum_{f \le \beta X} 3^{w(f)} \sum_{\substack{E \in \text{NF}_3(S_3) \\ D_E \le \frac{\beta X}{f}}} h_2(E)$$
$$\ll \sum_{f \le X} \frac{3^{w(f)} \beta X}{f}$$
$$\ll X(\log X)^3 \text{ (Lemma 2.8).}$$

(2) By Lemma 4.13,

$$N_3^{\text{tor}}(X; H_{24,g}) \le \# \left\{ L_{12} : D_{12} \le \beta X^{\frac{9}{2}} \right\}$$

for some constant $\beta > 0$. Since L_{12} has a quartic subfield L_4 , the Galois group $\operatorname{Gal}(L_{12}^c/\mathbb{Q})$ is 12T8 so

$$N_3^{\text{tor}}(X; H_{24,g}) \le N_{12}(\beta X^{\frac{9}{2}}; 12T8) \ll X^{\frac{9}{10}}.$$

Summing up the results of this section, we obtain the following theorem. Since there is an asymptotic lower bound $X(\log X)^2 \ll N_3^{\text{tor}}(X)$ ([22, Section 3]), the ratio of the upper and lower bounds of $N_3^{\text{tor}}(X)$ is $O((\log X)^2 \log \log X)$ under the Cohen-Lenstra heuristics for p = 3.

Theorem 4.15. (1) We have

$$N_3^{\text{tor}}(X) \ll_{\varepsilon} X^{1 + \frac{\log 2 + \varepsilon}{\log \log X}}.$$
(50)

(2) Under the assumption of Conjecture 1.4, we have

$$N_3^{\text{tor}}(X) \ll X (\log X)^4 \log \log X.$$
(51)

Acknowledgments

The author is supported by a KIAS Individual Grant (MG079601) at Korea Institute for Advanced Study. We thank Ila Varma for sharing the preprint [31]. We also thank Frank Thorne and Chia-Fu Yu for their helpful comments.

References

- S. A. Altuğ, A. Shankar, I. Varma and K. H. Wilson, The number of D₄-fields ordered by conductor, J. Eur. Math. Soc. 23 (2021), no. 8, 2733–2785.
- [2] J. Andersson, Answer to MathOverflow question, https://mathoverflow.net/a/55739, 2011.
- [3] E. Ascher and H. Grimmer, Comment on a paper by Tahara on the finite subgroups of GL(3, Z), Nagoya Math. J. 48 (1972), 203.
- [4] A. M. Baily, On the density of discriminants of quartic fields, J. Reine Angew. Math. 315 (1980), 190–210.
- [5] W. D. Banks and F. Pappalardi, Values of the Euler function free of kth powers, J. Number Theory 120 (2006), 326–348.
- [6] M. Bhargava, The density of discriminants of quartic rings and fields, Ann. of Math. (2) 162 (2005), no. 2, 1031–1063.
- [7] M. Bhargava, The density of discriminants of quintic rings and fields, Ann. of Math. (2) 172 (2010), no. 3, 1559–1591.
- [8] M. Bhargava, A. Shankar, T. Taniguchi, F. Thorne, J. Tsimerman and Y. Zhao, Bounds on 2-torsion in class groups of number fields and integral points on elliptic curves, J. Amer. Math. Soc. 33 (2020), no. 4, 1087–1099.
- H. Cohen, F. Diaz y Diaz and M. Olivier, Enumerating quartic dihedral extensions of Q, Compos. Math. 133 (2002), no. 1, 65–93.
- [10] H. Cohen, F. Diaz y Diaz and M. Olivier, Counting discriminants of number fields, J. Théor. Nombres Bordeaux 18 (2006), no. 3, 573–593.
- [11] H. Cohen and H. W. Lenstra Jr., Heuristics on class groups of number fields, Number Theory, Noordwijkerhout 1983, Lecture Notes in Math. 1068, 33-62, Springer, Berlin, 1984.
- [12] H. Cohen and J. Martinet, Class groups of number fields: numerical heuristics, Math. Comp. 48 (1987), no. 177, 123–137.
- [13] H. Cohen and F. Thorne, Dirichlet series associated to cubic fields with given quadratic resolvent, Michigan Math. J. 63 (2014), no. 2, 253–273.
- [14] H. Cohen and F. Thorne, Dirichlet series associated to quartic fields with given cubic resolvent, Res. Number Theory 2 (2016), Art. 29, 40.
- [15] J.-M. Couveignes, Enumerating number fields, Ann. of Math. (2) 192 (2020), no. 2, 487–497.
- [16] H. Davenport and H. Heilbronn, On the density of discriminants of cubic fields. II, Proc. Roy. Soc. London Ser. A 322 (1971), no. 1551, 405–420.
- [17] H. Delange, Généralisation du théorème de Ikehara, Ann. Sci. Éc. Norm. Supér. (3) 71 (1954), 213–242.
- [18] J. S. Ellenberg and A. Venkatesh, Counting extensions of function fields with bounded discriminant and specified Galois group, in Geometric Methods in Algebra and Number Theory, Progr. Math. 235, 151-168, Birkhäuser Boston, Boston, MA, 2005.
- [19] J. S. Ellenberg and A. Venkatesh, The number of extensions of a number field with fixed degree and bounded discriminant, Ann. of Math. (2) 163 (2006), no. 2, 723–741.
- [20] C. Fieker and J. Klüners, Minimal discriminants for fields with small Frobenius groups as Galois groups, J. Number Theory 99 (2003), no. 2, 318–337.
- [21] T. H. Gronwall, Some asymptotic expressions in the theory of numbers, Trans. Amer. Math. Soc. 14 (1913), no. 1, 113–122.
- [22] J. Lee, Counting algebraic tori over Q by Artin conductor, arXiv:2104.02855.
- [23] R. J. Lemke Oliver and F. Thorne, Upper bounds on number fields of given degree and bounded discriminant, Duke Math. J. 171 (2022), no. 15, 3077–3087.
- [24] The LMFDB Collaboration, The L-functions and modular forms database, https://www.lmfdb.org, 2022.

- [25] S. Mäki, On the density of abelian number fields, Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes 54 (1985).
- [26] S. Mäki, The conductor density of abelian number fields, J. Lond. Math. Soc. (2) 47 (1993), no. 1, 18–30.
- [27] G. Malle, On the distribution of Galois groups II, Exp. Math. 13 (2004), 129–135.
- [28] R. Masri, F. Thorne, W.-L. Tsai and J. Wang, Malle's conjecture for $G \times A$ with $G = S_3, S_4, S_5$, arXiv:2004.04651.
- [29] J. Neukirch, Algebraic number theory, Grundlehren der Mathematischen Wissenschaften 322, Springer, Berlin, 1999.
- [30] W. M. Schmidt, Number fields of given degree and bounded discriminant, Astérisque 228 (1995), 189–195.
- [31] A. Shankar and I. Varma, Malle's Conjecture for Galois octic fields over \mathbb{Q} , in preparation.
- [32] K. Tahara, On the finite subgroups of GL(3, Z), Nagoya Math. J. 41 (1971), 169–209.
- [33] J. Wang, Malle's conjecture for $S_n \times A$ for n = 3, 4, 5, Compos. Math. 157 (2021), no. 1, 83–121.
- [34] E. Wirsing, Das asymptotische Verhalten von Summenuber multiplikative Funktionen, Math. Ann. 143 (1961), 75–102.
- [35] D. J. Wright, Distribution of discriminants of abelian extensions, Proc. Lond. Math. Soc. (3) 58 (1989), no. 1, 17–50.

JUNGIN LEE, CENTER FOR MATHEMATICAL CHALLENGES, KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL 02455, KOREA *E-mail address:* jilee.math@gmail.com