EXISTENTIALLY CLOSED W*-PROBABILITY SPACES

ISAAC GOLDBRING AND CYRIL HOUDAYER

Abstract. We study several model-theoretic aspects of W*-probability spaces, that is, σ -finite von Neumann algebras equipped with a faithful normal state. We first study the existentially closed W*-spaces and prove several structural results about such spaces, including that they are type III₁ factors that tensorially absorb the Araki-Woods factor R_{∞} . We also study the existentially closed objects in the restricted class of W*-probability spaces with Kirchberg's QWEP property, proving that R_{∞} itself is such an existentially closed space in this class. Our results about existentially closed probability spaces imply that the class of type III₁ factors forms a \forall_2 -axiomatizable class. We show that for $\lambda \in (0,1)$, the class of III $_{\lambda}$ factors is not \forall_2 -axiomatizable but is \forall_3 -axiomatizable; this latter result uses a version of Keisler's Sandwich theorem adapted to continuous logic. Finally, we discuss some results around elementary equivalence of III_{λ} factors. Using a result of Boutonnet, Chifan, and Ioana, we show that, for any $\lambda \in (0,1)$, there is a family of pairwise non-elementarily equivalent III_{λ} factors of size continuum. While we cannot prove the same result for III₁ factors, we show that there are at least three pairwise non-elementarily equivalent III₁ factors by showing that the class of full factors is preserved under elementary equivalence.

CONTENTS

1.	Introduction				
2.	Prel	٤			
	2.1.	Basic facts about W*-probability spaces	Ę		
	2.2.	Ocneanu ultraproducts of W*-probability spaces	8		
	2.3.	Automorphism group, fullness and w-spectral gap	Ç		
	2.4.	W*-probability spaces as metric structures	10		
3.	Exis	11			
	3.1 Relative existential closedness		11		

Goldbring was partially supported by NSF grant DMS-2054477. Houdayer is supported by Institut Universitaire de France.

	3.2.	3.2. Global existential closedness				
	3.3.	The case of QWEP factors				
	3.4.	Building W*-probability spaces by games				
	3.5.	Open questions				
4.	Other model-theoretic results					
	4.1.	Axio	matizability results	2 3		
	4.2.	First-	order theories of W*-probability spaces	26		
Αţ	pend	ix A.	Embeddings of AFD W*-probability spaces into			
11			ultraproducts	29		
Appendix B.			Keisler's Sandwich Theorem	33		
Re	References					

1. Introduction

The model-theoretic study of von Neumann algebras began in earnest with the series of papers [16], [17], and [18] by Farah, Hart and Sherman. There, a theory in a (continuous) first-order language was described for which there was an equivalence of categories between the models of the theory and the category of tracial von Neumann algebras for which the model-theoretic ultraproduct construction corresponded to the tracial ultraproduct construction. In the time since these papers appeared, there has been a very interesting interplay between model-theoretic and operator-algebraic techniques; recent examples of applications of model-theoretic techniques to problems about tracial von Neumann algebras can be found in the papers [4], [20], and [22].

That a model-theoretic study of a wider class of von Neumann algebras (that is, beyond the finite realm) should be possible is hinted at by the existence of the Ocneanu ultraproduct construction, which allows one to take the ultraproduct of a family of W^* -probability spaces, that is, σ -finite von Neumann algebras equipped with a faithful, normal state. (The relevant facts about W^* -probability spaces needed in this paper are summarized in Section 2.) Motivated by the Ocneanu ultraproduct, Farah and Hart, in an unpublished work, observed that the category of σ -finite von Neumann algebras forms a so-called *compact abstract theory* (or CAT), which is a logical framework predating the current incarnation of continuous first-order logic. The first person to axiomatize (in the sense of the previous paragraph) W^* -probability spaces in an appropriate continuous first-order language was Dabrowski [13]; in particular, the model-theoretic

ultraproduct construction for this class corresponds to the Ocneanu ultraproduct construction. Dabrowski's axiomatization is quite technical and uses a fair amount of modular theory. A simpler (but less descriptive) axiomatization was given by Hart, Sinclair, and the first author in [24].

Now that the class of W*-probability spaces has been established as an axiomatizable class in an appropriate language, it is only natural to begin a thorough model-theoretic study of this class. In this paper, we initiate this endeavor. Our main focus will be on studying the class of existentially closed W*-probability spaces. The notion of an existentially closed structure is the model-theoretic generalization of the notion of an algebraically closed field. Roughly speaking, given structures M and N (in some language) for which M is a substructure of \mathbb{N} , we say that \mathbb{M} is an existentially closed substructure of \mathbb{N} (or that \mathbb{M} is existentially closed in \mathbb{N}) if any existential fact about some elements of \mathbb{M} which is true in \mathbb{N} is also true in \mathbb{M} . Considering that we are in the setting of continuous logic, truth in M is really approximate truth. Thankfully, this syntactic definition of existentially closed substructure can be given a semantic reformulation that aligns much more with the operator-algebraic perspective: M is an existentially closed substructure of $\mathbb N$ if and only if there is an embedding of $\mathbb N$ into an ultrapower $\mathfrak{M}^{\mathfrak{U}}$ of \mathfrak{M} for which the restriction of the embedding to \mathfrak{M} is the usual diagonal embedding of M into its ultrapower. If M belongs to some class $\mathfrak C$ of structures, we say that the structure $\mathfrak M$ is existentially closed for $\mathfrak C$ if it is existentially closed in all extensions belonging to \mathfrak{C} .

The Robinsonian school of model theory encourages one to understand the class of models of a theory by understanding its class of existentially closed models, that is, the models of the theory which are existentially closed for the class of models of the theory. The study of existentially closed tracial von Neumann algebras was carried out in the papers [23], [15], [19], and [21]. In Section 3 of this paper, we carry out a systematic study of the class of W*-probability spaces. In Subsection 3.1, we describe some properties of a W*-probability space that are inherited by an existentially closed substructure. In particular, we show that if the W*-probability space (M, φ) is existentially closed in (N, ψ) and N is a type III_1 factor, then so is M. This aids us in our study of the class of existentially closed W*-probability spaces in Subsection 3.2, where we show that any such W*-probability space is necessarily a type III₁ factor, which generalizes the result that an existentially closed tracial von Neumann algebra is necessarily a type II₁ factor. Other facts about existentially closed tracial von Neumann algebras are generalized to this setting, such as they tensorially absorb the Araki-Woods factor R_{∞} (generalizing the fact that any existentially closed II_1 factor tensorially absorbs the hyperfinite II₁ factor R) and that every automorphism of an existentially closed W*-probability space is approximately inner.

If one restricts one's attention to the class of tracial von Neumann algebras that satisfy the conclusion of the Connes Embedding Problem, that is, that admit a trace-preserving embedding into the tracial ultrapower of the hyperfinite II₁ factor R, then one obtains the fact that R itself is an existentially closed element of this class. Since a tracial von Neumann algebra embeds into the tracial ultrapower of R precisely when it has Kirchberg's QWEP property [34], it is natural to restrict attention to the class of QWEP W*-probability spaces. In fact, a result of Ando, Haagerup, and Winslow [3] shows that this class of W*-probability spaces can be characterized by admitting an embedding (with expectation) into the Ocneanu ultrapower of the Araki-Woods factor R_{∞} , or, as we show below, in model-theoretic terms, is a model of the universal theory of R_{∞} . We show that R_{∞} is an existentially closed QWEP W*-probability space and prove a few further results about this class of structures. Subsection 3.4 is concerned with the technique of building W*-probability spaces by games, which is a technique (first introduced in the continuous setting in [21]) that is very useful when trying to build e.c. objects with extra properties. The section concludes with Subsection 3.5, which contains some open questions about existentially closed W*-probability spaces.

Section 4 contains two further collections of results about the model theory of W*-probability spaces. The first collection of results concerns the axiomatizability of various classes of type III factors. It is shown in [3] that, given any $\lambda \in (0,1]$, the Ocneanu ultraproduct of a family of type III_{\lambda} factors is again a type III_{λ} factor and a factor is of type III_{λ} if its Ocneanu ultrapower is as well. Model-theoretically, this implies that the class of W*-probability spaces whose underlying von Neumann algebra is a type III_{λ} factor forms an axiomatizable class. In Section 4.1, we show that the results of our analysis of existentially closed W*-probability spaces implies that the class of type III₁ W*-probability spaces necessarily has a \forall_2 -axiomatization, that is, has a set of axioms of the form $\sup_{x} \inf_{y \in \mathcal{Y}} \theta(x, y)$, where x and y are finite tuples of variables and θ is a quantifier-free formula. We show that for a fixed $\lambda \in (0,1)$, the class of III_{λ} factors cannot be axiomatized using two quantifiers but can be axiomatized using three quantifiers. In none of these cases do we provide explicit axiomatizations but instead use a "soft" criterion for establishing the existence of such axiomatizations given by the Keisler Sandwich theorem.

In Subsection 4.2, we study the notion of elementary equivalence of W^* -probability spaces. Two W^* -probability spaces are elementarily equivalent if they cannot be distinguished using a first-order sentence. Using the Keisler-Shelah theorem, this can be given a semantic reformulation, namely that they have isomorphic ultrapowers. We first show how the result of Boutonnet, Chifan, and Ioana [6] stating that McDuff's family [36] of pairwise non-isomorphic separable II_1 factors are in fact pairwise non-elementarily equivalent can be used to show

that there exist continuum many non-elementarily equivalent separable type III_{λ} W*-probability spaces for any $\lambda \in (0,1)$. We are currently unable to extend this result to include $\lambda = 1$ but are able to identify at least three non-elementarily equivalent separable type III_1 W*-probability spaces. In order to accomplish this, we show that the class of *non-full* III_{λ} factors (for fixed $\lambda \in (0,1]$) is an axiomatizable class, generalizing the theorem of Farah, Hart, and Sherman [18] that the class of type II_1 factors with property Gamma is axiomatizable. This subsection includes a number of interesting open questions about the study of elementary equivalence of W*-probability spaces.

There are two appendices in this paper. The first appendix contains results about embedding AFD W*-probability spaces into ultraproducts that are needed in various portions of the paper; most of the results in this appendix are unpublished results of Ando and the second author. The second appendix concerns Keisler's Sandwich Theorem, which is the main model-theoretic tool needed in our axiomatization results appearing in Subsection 4.1. Since the continuous logic version of this result has never appeared, we include a complete proof of the result here. Moreover, we present the result using ultrapowers rather than arbitrary elementary extensions in the interest of the operator-algebraic audience.

We have made every attempt to keep the model-theoretic prerequisites for this paper to a minimum and try to use "semantic" definitions and proofs whenever possible. That being said, on a few occasions, we need to refer to basic model-theoretic terminology, such as elementary equivalence, elementary embedding, or first-order formula. A short introduction aimed towards operator algebraists (albeit in the language of tracial von Neumann algebras) can be found in Subsections 2.1 and 2.3 of [4].

2. Preliminaries

2.1. **Basic facts about W*-probability spaces.** For every von Neumann algebra M, we denote by $\|\cdot\|_{\infty}$ its uniform norm, by M_* its predual, by $\mathfrak{Z}(M)$ its center, by $\mathfrak{S}_{\mathrm{fn}}(M)$ the set of faithful normal states on M, by $\mathfrak{U}(M)$ its unitary group, by $\mathrm{Ball}(M)$ its unit ball with respect to the uniform norm, by $\mathrm{Aut}(M)$ its automorphism group and by $(M, L^2(M), J, L^2(M)^+)$ its standard form (see [25]). Under the identification $M = (M_*)^*$, the ultraweak topology on M coincides with the weak-* topology on $(M_*)^*$. A linear map $\Phi: M \to N$ is said to be *normal* if it is ultraweakly continuous.

A W^* -probability space is a pair (M, φ) that consists of a σ -finite von Neumann algebra M endowed with a faithful normal state $\varphi \in \mathfrak{S}_{fn}(M)$. For every $x \in M$, write $\|x\|_{\varphi} = \varphi(x^*x)^{1/2}$ (resp. $\|x\|_{\varphi}^{\sharp} = \varphi(x^*x)^{1/2} + \varphi(xx^*)^{1/2}$). On uniformly

bounded sets, the topology induced by the norm $\|\cdot\|_{\varphi}$ (resp. $\|\cdot\|_{\varphi}^{\sharp}$) coincides the strong (resp. *-strong) operator topology. We denote by σ^{ϕ} the modular automorphism group associated with the state φ . By definition, the *centralizer* M_{φ} of the state φ is the fixed point algebra of (M, σ^{φ}) . The *continuous core* of M with respect to φ is the crossed product von Neumann algebra $c_{\varphi}(M) =$ $M \rtimes_{\sigma^{\varphi}} \mathbb{R}$. The natural inclusion $\pi_{\varphi} : M \to c_{\varphi}(M)$ and the strongly continuous unitary representation $\lambda_{\varphi}: \mathbb{R} \to c_{\varphi}(M)$ satisfy the covariance relation

$$\lambda_\phi(t)\pi_\phi(x)\lambda_\phi(t)^*=\pi_\phi(\sigma_t^\phi(x))\quad \text{ for all } x\in M \text{ and all } t\in \mathbb{R}.$$

Set $L_{\varphi}(\mathbb{R}) = \lambda_{\varphi}(\mathbb{R})'' \subset c_{\varphi}(M)$. There is a unique faithful normal conditional expectation $E_{L_{\varphi}(\mathbb{R})}: c_{\varphi}(M) \to L_{\varphi}(\mathbb{R})$ satisfying $E_{L_{\varphi}(\mathbb{R})}(\pi_{\varphi}(x)\lambda_{\varphi}(t)) = \varphi(x)\lambda_{\varphi}(t)$. The faithful normal semifinite weight defined by $f \mapsto \int_{\mathbb{R}} \exp(-s)f(s) ds$ on $L^{\infty}(\mathbb{R})$ gives rise to a faithful normal semifinite weight Tr_{φ} on $L_{\varphi}(\mathbb{R})$ via the Fourier transform. The formula $\operatorname{Tr}_{\varphi}=\operatorname{Tr}_{\varphi}\circ \mathsf{E}_{\mathsf{L}_{\varphi}(\mathbb{R})}$ extends it to a faithful normal semifinite trace on $c_{\varphi}(M)$. Define the *dual action* $\theta^{\varphi} : \mathbb{R} \curvearrowright c_{\varphi}(M)$ by the formula

$$\theta_s^{\phi}(\pi_{\phi}(x)\lambda_{\phi}(t)) = \exp(-ist)\,\pi_{\phi}(x)\lambda_{\phi}(t) \quad \text{ for all } x \in M \text{ and all } s,t \in \mathbb{R}.$$

Then $\theta^{\phi}_{\sigma}: \mathbb{R} \curvearrowright c_{\phi}(\mathbb{R})$ is a trace-scaling action in the sense that $Tr_{\phi} \circ \theta^{\phi}_{s} =$ $\exp(-s)\operatorname{Tr}_{\varphi}$ for every $s \in \mathbb{R}$.

Let $\psi \in \mathfrak{S}_{fn}(M)$ be any other faithful normal state. By Connes' Radon–Nikodym cocycle theorem [7, Théorème 1.2.1] (see also [42, Theorem VIII.3.3]), there is a *-strongly continuous map $u : \mathbb{R} \to \mathcal{U}(M) : t \mapsto u_t$ such that

- $\begin{array}{l} (1) \ u_{s+t} = u_s \sigma_s^\phi(u_t) \ \text{for all } s,t \in \mathbb{R}, \\ (2) \ \sigma_t^\psi(x) = u_t \sigma_t^\phi(x) u_t^* \ \text{for all } t \in \mathbb{R} \ \text{and all } x \in M. \end{array}$

Item (1) says that $u : \mathbb{R} \to \mathcal{U}(M)$ is a 1-cocycle for σ^{φ} while Item (2) says that σ^{φ} and σ^{ψ} are cohomologous. Then the *-isomorphism $\Pi_{\phi,\psi}:c_{\phi}(M)\to c_{\psi}(M)$: $\pi_{\varphi}(x)u_{t}\lambda_{\varphi}(t) \mapsto \pi_{\psi}(x)\lambda_{\psi}(t)$ satisfies $\Pi_{\varphi,\psi} \circ \pi_{\varphi} = \pi_{\psi}$, $\Pi_{\varphi,\psi} \circ \theta^{\varphi} = \theta^{\psi} \circ \Pi_{\varphi,\psi}$ and $\operatorname{Tr}_{\psi} \circ \Pi_{\varphi,\psi} = \operatorname{Tr}_{\varphi}$. Note however that $\Pi_{\varphi,\psi}$ does not map the subalgebra $L_{\omega}(\mathbb{R}) \subset c_{\omega}(M)$ onto the subalgebra $L_{\psi}(\mathbb{R}) \subset c_{\psi}(M)$. It follows that the triple $(c_{\varphi}(M), \theta^{\varphi}, Tr_{\varphi})$ does not depend on the choice of the faithful normal state $\varphi \in$ $\mathfrak{S}_{\mathrm{fn}}(M)$ and we simply denote it by $(c(M), \theta, \mathrm{Tr})$.

Assume now that M is a factor. The restriction of θ to the center $\mathcal{Z}(c(M))$ is called the *flow of weights*. By factoriality of M, the flow of weights $\theta : \mathbb{R} \curvearrowright$ $\mathcal{Z}(c(M))$ is ergodic.

- If $\theta : \mathbb{R} \curvearrowright \mathcal{Z}(c(M))$ corresponds to the translation action $\mathbb{R} \curvearrowright \mathbb{R}$, then M is semifinite, that is, M is of type I or II.
- If $\theta : \mathbb{R} \curvearrowright \mathfrak{I}(c(M))$ is periodic with period T > 0, then letting $\lambda =$ $\exp(-T)$, we say that M is of type III_{λ} .
- If $\theta : \mathbb{R} \curvearrowright \mathcal{Z}(c(M))$ has no period, then we say that M is of type III₀.

• If θ : $\mathbb{R} \curvearrowright \mathcal{Z}(c(M))$ is trivial, that is, $\mathcal{Z}(c(M)) = \mathbb{C}1$, then we say that M is of type III_1 .

Next, we define Connes' S-invariant S(M) as the intersection

$$S(M) = \bigcap_{\phi \in \mathfrak{S}_{fn}(M)} \sigma(\Delta_\phi)$$

where $\sigma(\Delta_{\phi})$ is the spectrum of the modular operator Δ_{ϕ} associated with the faithful normal state $\phi \in \mathfrak{S}_{fn}(M)$. Then $S(M) \setminus \{0\}$ is a closed multiplicative subgroup of \mathbb{R}_+^* that completely determines the type of M. When M is a type III factor, we have that:

- M is of type III_0 if and only if $S(M) = \{0, 1\}$;
- M is of type III $_{\lambda}$ if and only if $S(M) = \{0\} \cup \lambda^{\mathbb{Z}}$, for $\lambda \in (0, 1)$;
- M is of type III₁ if and only if $S(M) = [0, +\infty)$.

We also define Connes' T-invariant T(M) as the set of all $t \in \mathbb{R}$ for which σ_t^ϕ is an inner automorphism. By Connes' Radon–Nikodym cocycle theorem, the above definition does not depend on the choice of the faithful normal state $\phi \in \mathfrak{S}_{fn}(M)$. Note that T(M) is a subgroup of \mathbb{R} . In case M is not of type III₀, then T(M) is a closed subgroup of \mathbb{R} that completely determines the type of M. Indeed, we have that:

- M is semifinite if and only if $T(M) = \mathbb{R}$;
- M is of type III $_{\lambda}$ if and only if $T(M) = \log(\lambda)\mathbb{Z}$, for $\lambda \in (0, 1)$;
- M is of type III₁ if and only if $T(M) = \{0\}$.

We refer to [7, 42] for further details regarding the structure of type III factors.

Throughout this paper, for $\lambda \in (0,1)$, (R_λ,ϕ_λ) denotes the Powers factor of type III_λ endowed with its (unique) $\frac{2\pi}{|\log(\lambda)|}$ -periodic faithful normal state. By definition, we have

$$(\mathsf{R}_{\lambda}, \varphi_{\lambda}) \cong (\mathsf{M}_{2}(\mathbb{C}), \omega_{\lambda})^{\otimes \mathbb{N}}$$

where $\omega_{\lambda}:M_2(\mathbb{C})\to\mathbb{C}$ is defined by

$$\omega_{\lambda}\left(\begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix}\right) = \frac{\lambda}{1+\lambda}x_{11} + \frac{1}{1+\lambda}x_{22}.$$

By Connes' result [11] (see also [43, Theorem XVIII.1.1]), R_{λ_i} is the unique AFD factor of type III_{λ_i} . We also let R_{∞} denote the Araki-Woods factor. Combining Connes' result [12] and Haagerup's result [26] (see also [43, Theorem XVIII.4.16]), R_{∞} is the unique AFD factor of type III_1 and moreover we have $R_{\infty} \cong R_{\lambda_1} \otimes R_{\lambda_2}$ whenever $\log(\lambda_1)/\log(\lambda_2)$ is irrational.

We next clarify what we mean by an inclusion of W*-probability spaces.

Definition 2.1. For W*-probability spaces (M, φ) and (N, ψ) , we say that (M, φ) *embeds into* (N, ψ) , denoted $(M, \varphi) \hookrightarrow (N, \psi)$, if there exist a unital normal *-embedding $\iota : M \to N$ such that $\psi \circ \iota = \varphi$ and a faithful normal conditional expectation $E : N \to \iota(M)$ such that $\varphi \circ \iota^{-1} \circ E = \psi$.

In what follows, we identify M with $\iota(M)$, regard $M \subseteq N$ as a von Neumann subalgebra, and assume that $\iota: M \to N$ is simply given by $\iota(x) := x$. In that case, we say that $(M, \phi) \subseteq (N, \psi)$ is an *inclusion* of W*-probability spaces. By [42, Theorem IX.4.2], the following assertions are equivalent:

- (1) $(M, \varphi) \subseteq (N, \psi)$.
- (2) The modular automorphism group σ^{ψ} leaves the subalgebra $M \subseteq N$ globally invariant, $\psi|_{M} = \varphi$, and $\sigma^{\varphi} = \sigma^{\psi}|_{M}$.

In that case, $E:N\to M$ is the unique faithful normal conditional expectation such that $\phi\circ E=\psi.$ Moreover, we have

$$\begin{split} M_{\phi} &= \{x \in M \mid \forall t \in \mathbb{R}, \sigma_t^{\phi}(x) = x\} \\ &= \{x \in M \mid \forall t \in \mathbb{R}, \sigma_t^{\psi}(x) = x\} \quad (\text{since } \sigma^{\phi} = \sigma^{\psi}|_{M}) \\ &\subseteq N_{\psi}. \end{split}$$

2.2. Ocneanu ultraproducts of W*-probability spaces. Let I be any nonempty directed set and $\mathcal U$ any nonprincipal ultrafilter on I. Let $(M_i,\phi_i)_{i\in I}$ be any family of W*-probability spaces. Following [1], define

$$\begin{split} \ell^{\infty}(I,M_i) &= \left\{ (x_i)_i \in \prod_{i \in I} M_i \mid \sup_{i \in I} \|x_i\|_{\infty} < +\infty \right\} \\ \mathfrak{I}_{\mathcal{U}} &= \left\{ (x_i)_i \in \ell^{\infty}(I,M_i) \mid \lim_{i \to \mathcal{U}} \|x_i\|_{\phi_i}^{\sharp} = 0 \right\} \\ \mathfrak{M}^{\mathcal{U}} &= \left\{ (x_i)_i \in \ell^{\infty}(I,M_i) \mid (x_i)_i \, \mathfrak{I}_{\mathcal{U}} \subset \mathfrak{I}_{\mathcal{U}} \quad \text{and} \quad \mathfrak{I}_{\mathcal{U}} (x_i)_i \subset \mathfrak{I}_{\mathcal{U}} \right\}. \end{split}$$

Observe that $\mathfrak{I}_{\mathfrak{U}}\subseteq\mathfrak{M}^{\mathfrak{U}}$. The *multiplier algebra* $\mathfrak{M}^{\mathfrak{U}}$ is a C*-algebra and $\mathfrak{I}_{\mathfrak{U}}\subset\mathfrak{M}^{\mathfrak{U}}$ is a norm closed two-sided ideal. Following [37, §5.1], we define the *ultraproduct* von Neumann algebra by $\prod_{\mathfrak{U}}(M_i,\phi_i):=(M_i,\phi_i)^{\mathfrak{U}}:=\mathfrak{M}^{\mathfrak{U}}/\mathfrak{I}_{\mathfrak{U}}$. We note that the proof given in [37, 5.1] for the case when $I=\mathbb{N}$ and $\mathfrak{U}\in\mathfrak{g}(\mathbb{N})\setminus\mathbb{N}$ applies *mutatis* mutandis. We denote the image of $(x_i)_i\in\mathfrak{M}^{\mathfrak{U}}$ in $(M_i,\phi_i)^{\mathfrak{U}}$ by $(x_i)^{\mathfrak{U}}$.

We now focus on the particular case when $(M_i, \phi_i) = (M, \phi)$ for some fixed W^* -probability space (M, ϕ) . In that case, we write $(M, \phi)^{\mathfrak{U}} = (M^{\mathfrak{U}}, \phi^{\mathfrak{U}})$ for the ultraprower of (M, ϕ) . For every $x \in M$, the constant sequence $(x)_i$ lies in the multiplier algebra $\mathfrak{M}^{\mathfrak{U}}$. We then identify M with $(M + \mathfrak{I}_{\mathfrak{U}})/\mathfrak{I}_{\mathfrak{U}}$ and regard $M \subset M^{\mathfrak{U}}$ as a von Neumann subalgebra. The map $E_{\mathfrak{U}}: M^{\mathfrak{U}} \to M$ given by $E_{\mathfrak{U}}((x_i)^{\mathfrak{U}}) = \sigma$ -weak $\lim_{i \to \mathfrak{U}} x_i$ is a faithful normal conditional expectation.

Moreover, we have $\varphi \circ E_{\mathcal{U}} = \varphi^{\mathcal{U}}$. Thus, $(M, \varphi) \subseteq (M, \varphi)^{\mathcal{U}}$ is an inclusion of W*-probability spaces. Following [8, §2], set

$$\mathfrak{M}_{\mathfrak{U}} = \left\{ (x_i)_i \in \ell^{\infty}(I,M) \mid \lim_{i \to \mathfrak{U}} \|x_i \zeta - \zeta x_i\| = 0, \forall \zeta \in L^2(M) \right\}.$$

Define the *asymptotic centralizer* von Neumann algebra by $M_{\mathfrak{U}}=\mathfrak{M}_{\mathfrak{U}}/\mathfrak{I}_{\mathfrak{U}}$, which is a von Neumann subalgebra of $M^{\mathfrak{U}}$. By [8, Proposition 2.8] (see also [1, Proposition 4.35]), we have $M_{\mathfrak{U}}=(M'\cap M^{\mathfrak{U}})_{\psi^{\mathfrak{U}}}$ for every faithful normal state $\psi\in\mathfrak{S}_{fn}(M)$.

Now let $(Q,\psi)\subseteq (M,\phi)$ be any inclusion of W^* -probability spaces and denote by $E:M\to Q$ the unique faithful normal conditional expectation such that $\psi\circ E=\phi$. We have $\ell^\infty(I,Q)\subset \ell^\infty(I,M),\ \mathfrak{I}_\mathfrak{U}(Q)\subseteq \mathfrak{I}_\mathfrak{U}(M)$ and $\mathfrak{M}^\mathfrak{U}(Q)\subseteq \mathfrak{M}^\mathfrak{U}(M)$. We then identify $Q^\mathfrak{U}=\mathfrak{M}^\mathfrak{U}(Q)/\mathfrak{I}_\mathfrak{U}(Q)$ with $(\mathfrak{M}^\mathfrak{U}(Q)+\mathfrak{I}_\mathfrak{U}(M))/\mathfrak{I}_\mathfrak{U}(M)$ and regard $Q^\mathfrak{U}\subset M^\mathfrak{U}$ as a von Neumann subalgebra. Observe that the norm $\|\cdot\|_{\psi^\mathfrak{U}}$ on $Q^\mathfrak{U}$ is the restriction of the norm $\|\cdot\|_{\phi^\mathfrak{U}}$ to $Q^\mathfrak{U}$. Observe moreover that $(E(x_i))_i\in\mathfrak{I}_\mathfrak{U}(Q)$ for all $(x_i)_i\in\mathfrak{I}_\mathfrak{U}(M)$ and $(E(x_i))_i\in\mathfrak{M}^\mathfrak{U}(Q)$ for all $(x_i)_i\in\mathfrak{M}^\mathfrak{U}(M)$. Therefore, the mapping $E^\mathfrak{U}:M^\mathfrak{U}\to Q^\mathfrak{U}$ given by $E^\mathfrak{U}((x_i)^\mathfrak{U})=(E(x_i))^\mathfrak{U}$ is a well-defined conditional expectation satisfying $\psi^\mathfrak{U}\circ E^\mathfrak{U}=\phi^\mathfrak{U}$. Thus, $(Q,\psi)^\mathfrak{U}\subseteq (M,\phi)^\mathfrak{U}$ is an inclusion of W^* -probability spaces.

2.3. **Automorphism group, fullness and w-spectral gap.** Let M be any σ -finite von Neumann algebra. Recall that for every $\varphi \in (M_*)^+$, there is a unique vector $\xi_{\varphi} \in L^2(M)^+$ such that $\varphi(x) = \langle x \xi_{\varphi}, \xi_{\varphi} \rangle$ for all $x \in M$. The group Aut(M) of all automorphisms of M acts on M_* by $\theta(\varphi) = \varphi \circ \theta^{-1}$ for all $\theta \in Aut(M)$ and $\varphi \in M_*$. Following [8, 25], the u-topology on Aut(M) is the topology of pointwise norm convergence on M_* , meaning that a net $(\theta_i)_{i \in I}$ in Aut(M) converges to the identity id_M in the u-topology if and only if for all $\varphi \in M_*$ we have $\|\theta_i(\varphi) - \varphi\| \to 0$ as $i \to \infty$. This turns Aut(M) into a complete topological group. When M_* is separable, Aut(M) is Polish. Since the standard form of M is unique, the group Aut(M) also acts naturally on $L^2(M)$ and we have $\theta(\xi_{\varphi}) = \xi_{\theta(\varphi)}$ for every $\varphi \in M_*^+$. It follows that the u-topology coincides with the topology of pointwise norm convergence on $L^2(M)$.

We introduce the following terminology:

Definition 2.2. We say that $\theta \in Aut(M)$ is

- approximately inner if there exists a net $(u_i)_{i \in I}$ in $\mathcal{U}(M)$ such that $Ad(u_i) \to \theta$ in the \mathfrak{u} -topology.
- weakly inner in the sense of [35] if the automorphism $\theta \odot$ id of $M \odot M^{op}$ extends to an automorphism of the C*-algebra $C^*_{\lambda \cdot \rho}(M)$ generated by the standard representation $\lambda \cdot \rho : M \odot M^{op} \to \mathbf{B}(L^2(M))$.

Any approximately inner automorphism is weakly inner (see [2, Theorem 4.1]) but the converse is not true (see [35]).

We say that a σ -finite factor M is full if whenever $(u_i)_{i \in I}$ is a net in $\mathcal{U}(Q)$ such that $Ad(u_i) \to id_M$ in the u-topology, there exists a bounded net $(\lambda_i)_{i \in I}$ in \mathbb{C} such that $u_i - \lambda_i 1 \to 0$ *-strongly. When M is tracial, M is full if and only if M does not have property Gamma of Murray and von Neumann. It is known that a full factor is never of type III_0 (see [8, Theorem 2.12]). By [31, Corollary 3.7], if M is a full factor, then for any nonprincipal ultrafilter $\mathcal U$ on any directed set I, we have $M' \cap M^{\mathcal U} = \mathbb C1$. (The converse is also true and follows readily from the definitions.) It follows from the classification of amenable factors [11, 12, 26] that any factor that is amenable and full is necessarily of type I.

We say that an inclusion of von Neumann algebras $Q \subset M$ is with expectation if there exists a faithful normal conditional expectation $E: M \to Q$. Moreover, we say that Q has w-spectral gap in M if, for any nonprincipal ultrafilter $\mathcal U$ on any set I, we have $Q' \cap M^{\mathcal U} = (Q' \cap M)^{\mathcal U}$. By [31, Theorem 4.4], for any σ -finite full factor M and any σ -finite von Neumann algebra N, M has w-spectral gap in $M \otimes N$.

2.4. **W*-probability spaces as metric structures.** As mentioned in the introduction, Dabrowski [13] introduced a first-order language for axiomatizing W*-probability spaces. In this language, the sorts are given by operator norm balls centered at the origin of various natural number radii. The metric on each ball is given by the norm $\|\cdot\|_{\phi}^*$, a relative of the norm $\|\cdot\|_{\phi}^*$ used above, which has the advantage that the state is Lipschitz continuous on each sort. While one has the natural symbols for scalar multiplication, addition, and adjoint, multiplication is not uniformly continuous on each sort and thus Dabrowski uses "smeared" multiplication maps defined using modular theory. Finally, he includes function symbols for the modular automorphism group (for rational times to keep the language countable) as well as some auxiliary symbols needed to make the axiomatization work. In this language, it is possible to axiomatize a class of structures which, as a category, is equivalent to the category of W*-probability spaces with inclusions as defined above and for which the model-theoretic ultraproduct corresponds to the Ocneanu ultraproduct.

While quite explicit, Dabrowski's language is very technical and cumbersome. An alternate axiomatization is given by Hart, Sinclair, and the first author in [24]. There, the sorts are given by vectors of operator norm at most N that are K-bounded (in a sense akin to that used in bimodule theory). On these sorts, the metric is induced by the norm $\|\cdot\|_{\phi}^{\sharp}$ and then all symbols (including multiplication) are naturally uniformly continuous. A much simpler axiomatization in this language can be given which once again yields an equivalence of categories

capturing the Ocneanu ultraproduct except that an inclusion of models of this theory only corresponds to a normal, state-preserving embedding of von Neumann algebras. In order to recover the correspondence with the above notion of inclusion of W*-probability spaces, one must add the modular automorphism group to the language, which is a harmless move as the modular automorphism group is definable in this language, as shown in [24]. However, the definability of the automorphism group is given by an abstract argument using the Beth Definability Theorem and thus this axiomatization lacks the concrete flavor of Dabrowski's axiomatization.

In this paper, the specific first-order framework for studying W*-probability spaces is not important and the reader can feel free to keep either of these two approaches in mind.

3. Existentially closed W*-probability spaces

3.1. **Relative existential closedness.** We begin this subsection by officially defining what it means for a W*-probability space to be existentially closed in another:

Definition 3.1. If $(M, \phi) \subseteq (N, \psi)$, we say that (M, ϕ) is *existentially closed* (e.c.) in (N, ψ) if and only if

$$(M, \phi) \subseteq (N, \psi) \subseteq (M, \phi)^{\mathfrak{U}}$$

so that $(M, \phi) \subseteq (M, \phi)^{\mathcal{U}}$ is the diagonal inclusion.

Remarks 3.2.

- (1) As discussed in the introduction, the previous definition is not the usual "syntactic" definition of existential closedness. Stated in syntactic terms, (M,ϕ) is e.c. in (N,ψ) if and only if: for every existential formula θ with parameters from M, we have $\theta^{(M,\phi)}=\theta^{(N,\psi)}$. The above definition is convenient for operator algebraists who do not wish to understand the precise definition of existential sentence.
- The previous definition is rather vague as to the nature of the ultrafilter \mathcal{U} . If N in the previous definition is a separably acting von Neumann algebra, then \mathcal{U} can be taken to be any nonprincipal ultrafilter on \mathbb{N} . For general N, one needs to take a particular kind of ultrafilter (known as a *good ultrafilter*) on some potentially large index set (depending on the density character of the metric associated to $\|\cdot\|_{\Psi}$).

The following flexibility result allows us to change states when dealing with relatively existentially closed W*-probability spaces.

Proposition 3.3. Suppose that (M, φ) is e.c. in (N, ψ) . Then for any faithful normal state ρ on M, we have that (M, ρ) is e.c. in $(N, \rho \circ E)$, where $E : N \to M$ is the unique faithful normal conditional expectation such that $\varphi \circ E = \psi$.

Proof. By assumption, we have $(M,\phi)\subseteq (N,\psi)\subseteq (M,\phi)^{\mathfrak{U}}$ so that $(M,\phi)\subseteq (M,\phi)^{\mathfrak{U}}$ is the diagonal inclusion. Denote by $F:M^{\mathfrak{U}}\to N$ the unique conditional expectation such that $\psi\circ F=\phi^{\mathfrak{U}}$. Recall from above that $E_{\mathfrak{U}}:M^{\mathfrak{U}}\to M$ given by $E_{\mathfrak{U}}((x_i)^{\mathfrak{U}})=w\text{-lim}_{\mathfrak{U}}\,x_i$ is the canonical faithful normal conditional expectation. By definition, we have $\phi\circ E_{\mathfrak{U}}=\phi^{\mathfrak{U}}$. Since ϕ is faithful and since

$$\varphi \circ (\mathsf{E} \circ \mathsf{F}) = \psi \circ \mathsf{F} = \varphi^{\mathsf{U}} = \varphi \circ \mathsf{E}_{\mathsf{U}},$$

by uniqueness of the conditional expectation, we have $E \circ F = E_{\mathcal{U}}$.

Let now ρ be any faithful normal state on M. We then have $(\rho \circ E) \circ F = \rho \circ E_{\mathcal{U}} = \rho^{\mathcal{U}}$. With respect to the same inclusions and the same conditional expectations E and E, we have $(M,\rho) \subseteq (N,\rho \circ E)$ and $(N,\rho \circ E) \subseteq (M^{\mathcal{U}},(\rho \circ E) \circ F) = (M,\rho)^{\mathcal{U}}$ and so $(M,\rho) \subseteq (N,\rho \circ E) \subseteq (M,\rho)^{\mathcal{U}}$. This shows that (M,ρ) is e.c. in $(N,\rho \circ E)$. \square

Lemma 3.4. *Suppose that* (M, φ) *is e.c. in* (N, ψ) *. Then:*

- (1) *If* N *is a factor, then* M *is a factor.*
- (2) $T(M) \subseteq T(N)$.

Proof. By assumption, we have $(M,\phi)\subseteq (N,\psi)\subseteq (M,\phi)^{\mathfrak{U}}$ where $(M,\phi)\subseteq (M,\phi)^{\mathfrak{U}}$ is the diagonal inclusion.

- (1) Assume that N is factor. Since $\mathcal{Z}(M)\subseteq\mathcal{Z}(N)$, it follows that $\mathcal{Z}(M)=\mathbb{C}1$ and so M is a factor.
- (2) Let $t \in T(M)$. Then $\sigma_t^{\phi} \in Inn(M)$ and there exists $\mathfrak{u} \in \mathfrak{U}(M)$ such that $\sigma_t^{\phi} = Ad(\mathfrak{u})$. Since $\sigma_t^{\psi} = \sigma_t^{\phi^{\mathfrak{U}}}|_N$ and since $\sigma_t^{\phi^{\mathfrak{U}}} = Ad(\mathfrak{u})$, it follows that $\sigma_t^{\psi} = Ad(\mathfrak{u}) \in Inn(N)$. Therefore, $t \in T(N)$.

The next key result gives a necessary condition for an e.c. inclusion of W^* -probability spaces to be of type III_0 . Unlike other results in this paper, the next theorem is a purely type III von Neumann algebraic statement.

Theorem 3.5. Suppose that (M, ϕ) is e.c. in (N, ψ) . If M is a type III_0 factor and N is a factor, then N is a type III_0 factor.

Proof. By assumption, we have $(M,\phi)\subseteq (N,\psi)\subseteq (M,\phi)^{\mathfrak{U}}$ where $(M,\phi)\subseteq (M,\phi)^{\mathfrak{U}}$ is the diagonal inclusion. Fix faithful normal conditional expectations $E:N\to M$ and $F:M^{\mathfrak{U}}\to N$ so that $E\circ F:M^{\mathfrak{U}}\to M$ is the canonical faithful normal conditional expectation $E_{\mathfrak{U}}:M^{\mathfrak{U}}\to M$.

Since M is a type III $_0$ factor, by [7, Lemme 5.3.2], there exists a lacunary faithful normal strictly semifinite weight Φ with infinite multiplicity on M. Then the centralizer M_{Φ} is a type II $_{\infty}$ von Neumann algebra and there exists a unique faithful normal Φ -preserving conditional expectation $E_{\Phi}: M \to M_{\Phi}$. By [7, Théorème 5.3.1], there exist $0 < \lambda_0 < 1$ and $u \in M(\sigma^{\Phi}, (-\infty, \log(\lambda_0)])$ such that $uM_{\Phi}u^* = M_{\Phi}$ and M is generated by M_{Φ} and u. For the definition of the spectral subspace $M(\sigma^{\Phi}, (-\infty, \log(\lambda_0)])$, we refer to [1, Subsection 2.2]. Moreover, we canonically have $M = M_{\Phi} \rtimes_{\theta} \mathbb{Z}$ where $\theta \in Aut(M_{\Phi})$ is given by $\theta = Ad(u)|_{M_{\Phi}}$. Also, M_{Φ} has a diffuse center and $\theta|_{\mathbb{Z}(M_{\Phi})} \in Aut(\mathbb{Z}(M_{\Phi}))$ is ergodic. Letting $\tau = \Phi|_{M_{\Phi}}$, we have $\tau(\theta(x)) \leq \lambda_0 \tau(x)$ for every $x \in (M_{\Phi})_+$.

Set $\Psi = \Phi \circ E$. Following [1, Definition 4.25], we have $\Psi \circ F = \Phi \circ E \circ F = \Phi \circ E_{\mathfrak{U}} = \Phi^{\mathfrak{U}}$. Since Φ is lacunary, by [1, Proposition 4.27], the equality $(M^{\mathfrak{U}})_{\Phi^{\mathfrak{U}}} = (M_{\Phi})^{\mathfrak{U}}$ holds. Then [1, Proposition 6.23] shows that $M^{\mathfrak{U}}$ is generated by $(M_{\Phi})^{\mathfrak{U}}$ and \mathfrak{U} . Moreover, we canonically have $M^{\mathfrak{U}} = (M_{\Phi})^{\mathfrak{U}} \rtimes_{\theta^{\mathfrak{U}}} \mathbb{Z}$, where $\theta^{\mathfrak{U}} \in \text{Aut}((M_{\Phi})^{\mathfrak{U}})$ is given by $\theta^{\mathfrak{U}} = \text{Ad}(\mathfrak{U})|_{(M_{\Phi})^{\mathfrak{U}}}$. Letting $\tau^{\mathfrak{U}} = \Phi^{\mathfrak{U}}|_{(M_{\Phi})^{\mathfrak{U}}}$, we have $\tau^{\mathfrak{U}}(\theta^{\mathfrak{U}}(x)) \leq \lambda_0 \tau^{\mathfrak{U}}(x)$ for every $x \in ((M_{\Phi})^{\mathfrak{U}})_+$.

Note that $N\subseteq M^{\mathfrak{U}}$ is globally invariant under $\sigma^{\Phi^{\mathfrak{U}}}$ and $\sigma^{\Psi}=\sigma^{\Phi^{\mathfrak{U}}}|_{N}$. This implies that $M_{\Phi}\subseteq N_{\Psi}\subseteq (M_{\Phi})^{\mathfrak{U}}$ which further implies that the centralizer N_{Ψ} is a type II_{∞} von Neumann algebra with diffuse center. Observe that $N_{\Psi}=N\cap (M_{\Phi})^{\mathfrak{U}}$ and that $\theta^{\mathfrak{U}}(N_{\Psi})=\mathfrak{U}N_{\Psi}\mathfrak{U}^{*}=N_{\Psi}$. Then $N_{\Psi}\subseteq (M_{\Phi})^{\mathfrak{U}}$ is a \mathbb{Z} -globally invariant von Neumann subalgebra and we have $N_{\Psi}\rtimes_{\theta^{\mathfrak{U}}}\mathbb{Z}\subseteq N$.

Claim 3.6. The equality $N = N_{\Psi} \rtimes_{\theta^{\mathcal{U}}} \mathbb{Z}$ holds.

Recall that $M^{\mathfrak{U}}=(M_{\Phi})^{\mathfrak{U}}\rtimes_{\theta^{\mathfrak{U}}}\mathbb{Z}$ and denote by $E_{\Phi^{\mathfrak{U}}}:M^{\mathfrak{U}}\to (M_{\Phi})^{\mathfrak{U}}$ the canonical faithful normal $\Phi^{\mathfrak{U}}$ -preserving conditional expectation. We will show below that $E_{\Phi^{\mathfrak{U}}}(N)=N_{\Psi}$. Once this is proven, [41, Corollary 3.4] implies that

$$N\subseteq \left\{x\in (M_\Phi)^\mathfrak{U}\rtimes_{\theta^\mathfrak{U}}\mathbb{Z}\ |\ \forall j\in\mathbb{Z}, \mathsf{E}_{\Phi^\mathfrak{U}}(x\mathfrak{u}^{-j})\in \mathsf{N}_\Psi\right\}=\mathsf{N}_\Psi\rtimes_{\theta^\mathfrak{U}}\mathbb{Z}$$

and thus $N = N_{\Psi} \rtimes_{\theta^{\mathfrak{U}}} \mathbb{Z}$.

First observe that $N_{\Psi} = E_{\Phi^{\mathfrak{U}}}(N_{\Psi}) \subset E_{\Phi^{\mathfrak{U}}}(N)$. Next, let $x \in (M_{\Phi})^{\mathfrak{U}} \rtimes_{alg} \mathbb{Z}$ be any element in the algebraic crossed product. Denote by $\mathcal{F} \subset \mathbb{Z}$ the finite support of x. Since $\mathcal{Z}(M_{\Phi}) = L^{\infty}(X, \nu)$ is diffuse and since the action $\mathbb{Z} \curvearrowright \mathcal{Z}(M_{\Phi})$ is ergodic, it follows that the action $\mathbb{Z} \curvearrowright (X, \nu)$ is essentially free. Then [41, Lemma 3.1] (whose proof works for arbitrary diffuse probability spaces) implies that there exists a finite partition of unity $\sum_i p_i = 1$ with projections $p_i \in \mathcal{Z}(M_{\Phi})$ such that $p_i\theta^j(p_i) = 0$ for all i and all $j \in \mathcal{F} \setminus \{0\}$. Then we have

$$\sum_i p_i x p_i = \sum_i \sum_{j \in \mathcal{F}} p_i \, E_{\Phi^{\mathcal{U}}}(x u^{-j}) u^j \, p_i$$

$$\begin{split} &= \sum_i \sum_{j \in \mathcal{F}} p_i E_{\Phi^{\mathcal{U}}}(x u^{-j}) \theta^j(p_i) u^j \\ &= \sum_i \sum_{j \in \mathcal{F}} p_i \theta^j(p_i) E_{\Phi^{\mathcal{U}}}(x u^{-j}) u^j \quad (\text{since } \theta^j(p_i) \in \mathcal{Z}(M_\Phi) \subseteq \mathcal{Z}((M_\Phi)^\mathcal{U})) \\ &= \sum_i p_i \, E_{\Phi^{\mathcal{U}}}(x) \\ &= E_{\Phi^{\mathcal{U}}}(x). \end{split}$$

Thus, we have $E_{\Phi^{\mathfrak{U}}}(x) = \sum_{i} p_{i} x p_{i}$.

We use an idea in [41, Lemma 3.2]. For this, we choose a faithful normal state ρ on $M^{\mathfrak{U}}$ such that $\rho \circ \mathsf{E}_{\Phi^{\mathfrak{U}}} = \rho$. Then $(M_{\Phi})^{\mathfrak{U}} \subseteq M^{\mathfrak{U}}$ is globally invariant under the modular automorphism group σ^{ρ} . Since $\mathcal{Z}(M_{\Phi}) \subseteq \mathcal{Z}((M_{\Phi})^{\mathfrak{U}})$, it follows that $\mathcal{Z}(M_{\Phi})$ is contained in the centralizer $(M^{\mathfrak{U}})_{\rho}$. Let now $y \in N$ be any element. For every $n \geq 1$, we may choose $x_n \in (M_{\Phi})^{\mathfrak{U}} \rtimes_{\text{alg}} \mathbb{Z}$ so that $\|y - x_n\|_{\rho} \leq \frac{1}{2n}$. We then have $\|\mathsf{E}_{\Phi^{\mathfrak{U}}}(y - x_n)\|_{\rho} \leq \|y - x_n\|_{\rho} \leq \frac{1}{2n}$. The above reasoning shows that there exists a finite partition of unity $\sum_i p_i^n = 1$ with projections $p_i^n \in \mathcal{Z}(M_{\Phi})$ such that $\mathsf{E}_{\Phi^{\mathfrak{U}}}(x_n) = \sum_i p_i^n x p_i^n$. Since for every i, we have $p_i^n \in \mathcal{Z}(M_{\Phi}) \subset (M^{\mathfrak{U}})_{\rho}$ and $0 \leq p_i^n \leq 1$, we obtain

$$\begin{split} \| \sum_{i} p_{i}^{n}(x_{n} - y) p_{i}^{n} \|_{\rho}^{2} &= \rho(\sum_{i} p_{i}^{n}(x_{n} - y)^{*} p_{i}^{n}(x_{n} - y) p_{i}^{n}) \\ &\leq \rho(\sum_{i} p_{i}^{n}(x_{n} - y)^{*}(x_{n} - y) p_{i}^{n}) \\ &= \rho(\sum_{i} p_{i}^{n}(x_{n} - y)^{*}(x_{n} - y)) \\ &= \rho((x_{n} - y)^{*}(x_{n} - y)) \\ &= \|x_{n} - y\|_{\rho}^{2}. \end{split}$$

This implies that

$$\begin{split} \|E_{\Phi^{\mathfrak{U}}}(y) - \sum_{i} p_{i}^{n} y p_{i}^{n}\|_{\rho} &\leq \|E_{\Phi^{\mathfrak{U}}}(y - x_{n})\|_{\rho} + \|E_{\Phi^{\mathfrak{U}}}(x_{n}) - \sum_{i} p_{i}^{n} x_{n} p_{i}^{n}\|_{\rho} \\ &+ \|\sum_{i} p_{i}^{n} (x_{n} - y) p_{i}^{n}\|_{\rho} \\ &\leq \|x_{n} - y\|_{\rho} + 0 + \|x_{n} - y\|_{\rho} \\ &\leq \frac{1}{2n} + \frac{1}{2n} = \frac{1}{n}. \end{split}$$

For every $n \ge 1$, set $y_n := \sum_i p_i^n y p_i^n \in N$ (recall that $p_i^n \in \mathfrak{Z}(M_\Phi) \subseteq N$). Since $\lim_n \|E_{\Phi^{\mathfrak{U}}}(y) - y_n\|_{\rho} = 0$, the uniformly bounded sequence $(y_n)_{n \ge 1}$ converges strongly to $E_{\Phi^{\mathfrak{U}}}(y)$. Since $y_n \in N$ for every $n \ge 1$ and since N is strongly closed,

this implies that $E_{\Phi^{\mathfrak{U}}}(y) \in N$. This further implies that $E_{\Phi^{\mathfrak{U}}}(N) \subset N \cap (M_{\Phi})^{\mathfrak{U}} = N_{\Psi}$ and so $E_{\Phi^{\mathfrak{U}}}(N) = N_{\Psi}$. This finishes the proof of the claim.

By Claim 3.6, we have $N = N_{\Psi} \rtimes_{\theta^{\mathfrak{U}}} \mathbb{Z}$ where N_{Ψ} is a type II_{∞} von Neumann algebra with diffuse center. Since N is a factor, we have that $\theta^{\mathfrak{U}}|_{\mathfrak{Z}(N_{\Psi})} \in Aut(\mathfrak{Z}(N_{\Psi}))$ is necessarily ergodic. Moreover, we have $\tau^{\mathfrak{U}}(\theta^{\mathfrak{U}}(x)) \leq \lambda_0 \tau^{\mathfrak{U}}(x)$ for every $x \in (N_{\Psi})_+$. Then [7, Proposition 5.1.1] implies that N is a type III₀ factor.

Corollary 3.7. *Suppose that* (M, ϕ) *is e.c. in* (N, ψ) *. If* N *is a type* III_1 *factor, then so is* M.

Proof. Lemma 3.4(1) implies that M is a factor. Theorem 3.5 implies that M is not of type III_0 . Lemma 3.4(2) implies that $T(M) \subseteq T(N) = \{0\}$. Since M is not of type III_0 , this further implies that M has type III_1 .

3.2. **Global existential closedness.** As stated in the introduction, we say that (M, φ) is *existentially closed* (e.c.) if whenever $(M, \varphi) \subseteq (N, \psi)$, then (M, φ) is e.c. in (N, ψ) . An immediate consequence of Proposition 3.3 is the following:

Proposition 3.8. If M is a σ -finite von Neumann algebra, then for any two faithful, normal states ϕ_1 and ϕ_2 on M, we have that (M, ϕ_1) is e.c. if and only if (M, ϕ_2) is e.c.

By the previous proposition, it is sensible to call a σ -finite von Neumann algebra M existentially closed (e.c.) if (M, φ) is an e.c. W*-probability space for some (equivalently any) $\varphi \in \mathfrak{S}_{fn}(M)$.

The next result enumerates many important facts about e.c. W*-probability spaces.

Theorem 3.9. Suppose that M is an e.c. W*-probability space. Then the following assertions hold:

- (1) M is a type III_1 factor.
- (2) $M \otimes R_{\infty} \cong M$.
- (3) For any full factor Q, either Q is of type I or $M \ncong Q \otimes R_{\infty}$.
- (4) Any automorphism of M is approximately inner.
- (5) For every subfactor $N \subset M$ with expectation and with w-spectral gap, we have $(N' \cap M)' \cap M = N$.

Proof. (1) Choose any faithful normal state ρ on R_{∞} and denote by $(N, \psi) := (M, \phi) * (R_{\infty}, \rho)$ the corresponding free product von Neumann algebra. Since N is a type III_1 factor (see [44, Theorem 4.1]), Corollary 3.7 implies that M is a type III_1 factor.

(2) Fix $\lambda \in (0,1)$. Then $(M,\phi) \subseteq (M,\phi) \otimes (R_{\lambda},\phi_{\lambda}) \hookrightarrow (M,\phi)^{\mathcal{U}}$ with the composition being the diagonal embedding. In particular, $(R_{\lambda},\phi_{\lambda})$ embeds into

- $(M'\cap M^{\mathfrak{U}},\dot{\phi}^{\mathfrak{U}})$ where $\dot{\phi}^{\mathfrak{U}}=\phi^{\mathfrak{U}}|_{M'\cap M^{\mathfrak{U}}}$. This implies that $\lambda\in\sigma_{\mathfrak{p}}(\Delta_{\dot{\phi}^{\mathfrak{U}}})$. Then [1, Theorem 4.32] shows that $M\otimes R_{\lambda}\cong M$. Since this is true for every $\lambda\in(0,1)$, it follows that $M\otimes R_{\infty}\cong M$.
- (3) Assume that there exists a full factor Q such that $M = Q \otimes R_{\infty}$. Denote by $\alpha \in \operatorname{Aut}(M \otimes M)$ the flip automorphism defined by $\alpha(x \otimes y) = y \otimes x$. Regard $M \subseteq M \otimes M : x \mapsto x \otimes 1$ and set $N := (M \otimes M) \times_{\alpha} \mathbb{Z}/2\mathbb{Z}$. Then $M \subseteq N \hookrightarrow M^{\mathfrak{U}}$ with the composition being the diagonal embedding. Denote by $u = (u_n)^{\mathcal{U}} \in \mathcal{U}(N) \subset$ $\mathcal{U}(M^{\mathcal{U}})$ the canonical unitary implementing the action $\mathbb{Z}/2\mathbb{Z} \curvearrowright^{\alpha} M$. Then for every $x \in M$, we have $1 \otimes x = \alpha(x \otimes 1) = u(x \otimes 1)u^*$. This implies that $uMu^* \subseteq u$ $M' \cap M^{\mathfrak{U}}$. Since Q is a full factor, Q has w-spectral gap in $M = Q \otimes R_{\infty}$ and so $Q' \cap M^{\mathfrak{U}} = (Q' \cap M)^{\mathfrak{U}} = R^{\mathfrak{U}}_{\infty}$ (see [31, Corollary 3.7]). We obtain $\mathfrak{u}M\mathfrak{u}^* \subseteq R^{\mathfrak{U}}_{\infty}$ and so $M \subseteq \mathfrak{u}^*R^{\mathfrak{U}}_{\infty}\mathfrak{u} \subseteq M^{\mathfrak{U}}$. Fix a faithful normal state ψ on Q and consider the faithful normal conditional expectation $E: M \to R_{\infty}$ defined by $E = \psi \otimes id_{R_{\infty}}$. For every n, set $R_n=u_n^*R_\infty u_n\subseteq M$ and define the faithful normal conditional expectation $E_n: M \to R_n$ by the formula $E_n = Ad(u_n^*) \circ E \circ Ad(u_n)$. Up to changing the net, we may assume that for every $x \in M$, we have $E_n(x) - x \to 0$ *strongly as $n \to \infty$. For every n, we may choose a faithful normal state ϕ_n on R_n so that (R_n, φ_n) is an AFD W*-probability space (see Definition A.3). Therefore, we may find nets of normal ucp maps $S_i: M \to M_{n_i}(\mathbb{C})$ and $T_i: M_{n_i}(\mathbb{C}) \to M$ such that for every $x \in M$, we have $(T_i \circ S_i)(x) - x \to 0$ *-strongly as $i \to \infty$. Then [43, Theorem XV.3.1] implies that $M = Q \otimes R_{\infty}$ is amenable. Then Q is amenable and full and so Q is a type I factor.
- (4) Fix $\theta \in \text{Aut}(M)$ and denote by $N := M \rtimes_{\theta} \mathbb{Z}$ the corresponding crossed product von Neumann algebra. Then $M \subseteq N \hookrightarrow M^{\mathfrak{U}}$ with the composition being the diagonal embedding. Denote by $\mathfrak{u} = (\mathfrak{u}_n)^{\mathfrak{U}} \in \mathcal{U}(N) \subset \mathcal{U}(M^{\mathfrak{U}})$ the canonical unitary implementing the action $\mathbb{Z} \curvearrowright^{\theta} M$. Then for every $x \in M$, we have $\theta(x) = uxu^*$ and [2, Theorem 4.1 (iv) \Rightarrow (v) [2] shows that θ is weakly inner. Since $M \cong M \otimes R_{\infty}$, [35, Theorem [35] implies that [35] is approximately inner.
- (5) It suffices to show that $(N'\cap M)'\cap M\subseteq N$. Define the amalgamated free product von Neumann algebra $Q:=M*_N(L(\mathbb{Z})\otimes N)$ with respect to the natural faithful normal conditional expectations $E:M\to N$ and $\tau_\mathbb{Z}\otimes \mathrm{id}_N:L(\mathbb{Z})\otimes N\to N$. Then $M\subseteq Q\hookrightarrow M^\mathfrak{U}$ with the composition being the diagonal embedding. Denote by $\mathfrak{u}\in \mathfrak{U}(L(\mathbb{Z}))\subset \mathfrak{U}(M^\mathfrak{U})$ the canonical Haar unitary. Then we have $\mathfrak{u}\in N'\cap Q\subseteq N'\cap M^\mathfrak{U}=(N'\cap M)^\mathfrak{U}$ and so we may write $\mathfrak{u}=(\mathfrak{u}_n)^\mathfrak{U}$ where $\mathfrak{u}_n\in \mathfrak{U}(N'\cap M)$ for every \mathfrak{n} . Now take now $\mathfrak{b}\in (N'\cap M)'\cap M$ and note that $\mathfrak{bu}_n=\mathfrak{u}_n\mathfrak{b}$ for every \mathfrak{n} and so $\mathfrak{bu}=\mathfrak{ub}$. Since $L(\mathbb{Z})$ is diffuse, we have $L(\mathbb{Z})\not\preceq_{L(\mathbb{Z})\otimes N}N$ in the sense of Popa's intertwining theory (see [40,30]), whence [46, Proposition 3.3] implies that $L(\mathbb{Z})'\cap Q=L(\mathbb{Z})\otimes N$. Thus, we obtain $\mathfrak{b}\in M\cap (L(\mathbb{Z})\otimes N)=N$. \square
- **Remark 3.10.** All of the items in the previous theorem are appropriate generalizations of the corresponding facts about e.c. tracial von Neumann algebras.

Indeed, suppose that M is an e.c. tracial von Neumann algebra. Then the finite analog of (1) states that M is a II_1 factor, which was proven in [23], while the finite analog of (2) states that M is McDuff, that is, tensorially absorbs the hyperfinite II_1 factor R, which was also proven in [23]. The finite analog of (3) is that M is not a strongly McDuff factor, where a strongly McDuff factor is one that is isomorphic to a factor of the form $Q \otimes R$, where Q is a II_1 factor without property Gamma; this fact was proven in [4, Proposition 6.2.11]. The finite analogs of (4) and (5) have identical statements and were proven in [15] and [19, Proposition 5.16] respectively.

Proposition A.8 in the appendix states that the ultrapower of a type III_1 factor is always a prime factor, that is, cannot be written as the tensor product of diffuse factors. Combined with Theorem 3.9(2), we immediately obtain:

Corollary 3.11. *The class of e.c.* W^* *-probability spaces is not closed under ultrapowers.*

In particular, the class of e.c. W*-probability spaces is not axiomatizable. In model-theoretic language, this means:

Corollary 3.12. *The theory of* W^* *-probability spaces does not have a model companion.*

The analogous fact for tracial von Neumann algebras also holds and was the main result of [23].

3.3. The case of QWEP factors. In [15], the authors consider the e.c. elements of the class of tracial von Neumann algebras that admit a trace-preserving embedding into the tracial ultrapower $R^{\mathbb{U}}$ of the hyperfinite II_1 factor R. That this is a model-theoretically sensible thing to consider is substantiated by the basic fact that a tracial von Neumann algebra embeds into $R^{\mathbb{U}}$ if and only if it is a model of the *universal theory* of R, denoted $Th_{\forall}(R)$, consisting of all conditions of the form $\sup_x \phi(x) = 0$ with $\phi(x)$ a quantifier-free formula. As in the unrestricted case, any e.c. model of $Th_{\forall}(R)$ is a McDuff II_1 factor with only approximately inner automorphisms. In this case, however, one can name a concrete e.c. object, namely R itself. In fact, a positive solution to the Connes Embedding Problem is equivalent to the statement that R is an e.c. tracial von Neumann algebra.

In this subsection, we consider the analogous situation for W^* -probability spaces. To motivate the move that is to follow, we recall that a tracial von Neumann algebra embeds into $R^{\mathfrak{U}}$ if and only if it has Kirchberg's QWEP property [34]. Thus, it appears that the natural course of action to take in our current context is to consider restricting to the class of QWEP W^* -probability spaces. To see that, once again, this is a natural move from the model-theoretic perspective, we recall the main result of [3], which states that a von Neumann algebra has QWEP if and only if it embeds into $R^{\mathfrak{U}}_{\infty}$. (This is technically proven in the case that the von

Neumann algebra under consideration is separably acting and the ultrafilter is a nonprincipal ultrafilter on \mathbb{N} ; this result naturally extends to all QWEP von Neumann algebras by writing them as an increasing union of separably acting QWEP subalgebras and using an ultrafilter on a larger index set.) To see that this latter condition has model-theoretic meaning, we make the following observation. Recall from the introduction that two W*-probability spaces (M, φ) and (N, ψ) are elementarily equivalent if and only if there are ultrafilters $\mathcal U$ and $\mathcal V$ such that $(M, \varphi)^{\mathcal U} \cong (N, \psi)^{\mathcal V}$.

Proposition 3.13. For any type III_1 factor M and faithful normal states ϕ and ψ on M, we have that (M, ϕ) and (M, ψ) are elementarily equivalent.

Proof. The state space of $M^{\mathfrak{U}}$ is *strictly homogeneous* by [1, Theorem 4.20]. (We note that this result indeed holds for ultrapowers with respect to arbitrary countably incomplete ultrafilters.) Consequently, there is $\mathfrak{u} \in \mathcal{U}(M^{\mathfrak{U}})$ such that $\mathfrak{u}\phi^{\mathfrak{U}}\mathfrak{u}^* = \psi^{\mathfrak{U}}$; the inner automorphism $Ad(\mathfrak{u})$ thus yields that $(M^{\mathfrak{U}}, \phi^{\mathfrak{U}}) \cong (M^{\mathfrak{U}}, \psi^{\mathfrak{U}})$. \square

Remark 3.14. The previous proposition is false for type III_{λ} factors, $\lambda \in (0,1)$, even when only considering the universal theory. Indeed, if M is a type III_{λ} factor, $\lambda \in (0,1)$, and $\phi, \psi \in \mathfrak{S}_{fn}(M)$ are such that ϕ is periodic and ψ is not, then (M,ψ) cannot embed into the ultrapower of (M,ϕ) .

A particular consequence of Proposition 3.13 is that we may unambiguously speak of the universal theory $\operatorname{Th}_{\forall}(M)$ of any type III_1 factor M, by which we mean the unique common universal theory of (M,ϕ) for any $\phi \in \mathfrak{S}_{fn}(M)$. From this point of view, the main result of [3] can be reworded by saying that a W^* -probability space (M,ϕ) is QWEP if and only if it is a model of $\operatorname{Th}_{\forall}(R_{\infty})$.

Remark 3.15. Most of the results of the previous subsection continue to hold when restricted to the elementary class of QWEP W*-probability spaces. More specifically, the first four items of Theorem 3.9 as well as Corollary 3.12 hold when restricted to the class of QWEP W*-probability spaces. We do not know if item (5) of Theorem 3.9 holds in this restricted case as it is unknown if the amalgamated free product of QWEP von Neumann algebras remains QWEP. In that respect, it follows from [29, Corollary B] that the free product of QWEP von Neumann algebras remains QWEP. For other permanence properties, we refer to [38, Proposition 4.1].

As mentioned above, R is an e.c. member of the class of QWEP II_1 factors. We now prove the analogous fact in the setting of W*-probability spaces:

Theorem 3.16. The Araki–Woods factor R_{∞} is an e.c. QWEP W*-probability space.

Proof. Let N be an e.c. QWEP W*-probability space such that $R_{\infty} \subseteq N$. Since N is QWEP, there is an embedding $N \hookrightarrow R_{\infty}^{\mathfrak{U}}$. By Corollary A.7, up to conjugating

by a unitary, we may suppose that the composite embedding is the diagonal embedding. This shows that R_{∞} is e.c. in N and hence is an e.c. QWEP W*-probability space. (Note that we have used the fact that being e.c. does not depend on the choice of state when we conjugated by a unitary.)

3.4. **Building W*-probability spaces by games.** We now introduce a method for building W*-probability spaces first introduced in [21] (based on the discrete case presented in [27]). This method goes under many names, such as *Henkin constructions, model-theoretic forcing,* or *building models by games*.

We fix a countably infinite set C of distinct symbols that are to represent generators of a separable W*-probability space that two players (traditionally named \forall and \exists) are going to build together (albeit adversarially). The two players take turns playing finite sets of expressions of the form $|\theta(c) - r| < \epsilon$, where c is a tuple of variables from C, $\theta(c)$ is some atomic formula, and each player's move is required to extend (that is, contain) the previous player's move. The exact form an atomic formula depends on which language we are considering for describing W*-probability spaces, but in either case, they roughly correspond to expressions of the form $\varphi(p(c))$, where p is some expression involving the *algebra operations as well as modular automorphisms and φ is a generic symbol for the state. (In the case of Dabrowski's language, one is only allowed to use "smeared" multiplication in such expressions.) These plays of the game are called (open) *conditions*. The game begins with \forall 's move. Moreover, these conditions are required to be satisfiable, meaning that there should be some W*probability space (M, φ) and some tuple a from M such that $|\theta(a) - r| < \epsilon$ for each such expression in the condition. We play this game for countably many rounds. At the end of this game, we have enumerated some countable, satisfiable set of expressions. Provided that the players address a certain "dense" set of conditions infinitely often, they can ensure that the play is *definitive*, meaning that the final set of expressions yields complete information about all atomic formulae in the variables C (that is, for each atomic formula $\theta(c)$, there should be a unique r such that the play of the game implies that $\theta(c) = r$) and that this data describes a countable, dense *-subalgebra of a unique W*-probability space, which is called the *compiled structure*.

Definition 3.17. Given a property P of W*-probability spaces, we say that P is an *enforceable* property is there a strategy for \exists so that, regardless of player \forall 's moves, if \exists follows the strategy, then the compiled structure will have property P.

Fact 3.18.

(1) (Conjunction lemma [27, Lemma 2.4]) If P_n is an enforceable property for each $n \in \mathbb{N}$, then so is the conjunction $\bigwedge_n P_n$.

(2) ([27, Proposition 2.10] *Being e.c. is enforceable.*

Item (2) in the previous fact indicates the significance of this technique of building W^* -probability spaces in connection with the study of e.c. W^* -probability spaces.

Definition 3.19. A W*-probability space (M, φ) is said to be **enforceable** if the property of being isomorphic to (M, φ) is an enforceable property.

Clearly, if an enforceable W*-probability space exists, then it is unique.

One can relativize the above context by considering only QWEP W*-probability spaces. One can then speak of enforceable properties of QWEP W*-probability spaces and ask about the existence of the enforceable W*-probability space.

In the analogous game for II_1 factors, it was shown in [21, Theorem 5.2] that a positive solution to the Connes Embedding Problem is equivalent to the statement that R is the enforceable tracial von Neumann algebra and that, when restricted to the context of QWEP tracial von Neumann algebras, R is indeed the enforceable object. It is worth asking if the same is true in the case of QWEP W*-probability algebras. The answer is actually negative and rests on the following:

Proposition 3.20. There is no faithful normal state φ on R_∞ such that (R_∞, φ) embeds into all e.c. QWEP W*-probability spaces.

Proof. By contradiction, assume that there exists a faithful normal state ϕ on R_{∞} such that for any other faithful normal state ψ on R_{∞} we have $(R_{\infty}, \phi) \hookrightarrow (R_{\infty}, \psi)$.

It is a standard fact that there exists a faithful normal state ψ on R_{∞} for which $(R_{\infty})_{\psi}=\mathbb{C}$. Since $(R_{\infty},\phi)\hookrightarrow (R_{\infty},\psi)$, it follows that $(R_{\infty})_{\phi}=\mathbb{C}1$. Next, identify $(R_{\infty},\psi)=(R_{\lambda_1}\otimes R_{\lambda_2},\phi_{\lambda_1}\otimes \phi_{\lambda_2})$ for appropriately chosen λ_1 and λ_2 . Then ψ is an almost periodic state on R_{∞} in the sense that the corresponding modular operator Δ_{ψ} is diagonalizable on $L^2(R_{\infty})$. Since $(R_{\infty},\phi)\hookrightarrow (R_{\infty},\psi)$, it follows that ϕ is an almost periodic state and so $(R_{\infty})_{\phi}\neq\mathbb{C}1$ by [45, Lemma 2.1]. This is a contradiction.

Corollary 3.21. *There is no enforceable QWEP W*-probability space.*

Proof. Suppose, towards a contradiction, that (M, φ) is the enforceable QWEP W*-probability space. Then (M, φ) is e.c. and embeds into all e.c. QWEP W*-probability spaces (see [21, Section 6]). In particular (M, φ) embeds into (R_∞, ψ) for all faithful normal states ψ on R_∞ . This implies that $M \cong R_\infty$, and we obtain a contradiction with the previous proposition.

On the other hand, we do have the following proposition:

Proposition 3.22. The property of being approximately finite dimensional is an enforceable property of QWEP W*-probability spaces.

Proof. By the Conjunction Lemma, it suffices to show that, given any open condition Σ , any finite number c_1,\ldots,c_n of constants, and any $\varepsilon>0$, there are matrix units (e_{ij}) for some matrix algebra and some complex coefficients α^k_{ij} for $k=1,\ldots,n$ such that $\Sigma\cup\{\|c_k-\sum_{ij}\alpha^k_{ij}e_{ij}\|^\#_\phi<\varepsilon\}$ is itself a condition. However, this follows from the fact that Σ , being satisfiable in some QWEP W*-probability space, must also be satisfiable in R_∞ .

It is quite interesting that player \exists can always enforce the underlying von Neumann algebra of the compiled W*-probability space to be R_{∞} although there is no single state on R_{∞} that can be enforced.

An important class of e.c. structures is the class of finitely generic structures. We end this section by briefly discussing this class. First, since the class of W*-probability spaces has the joint embedding property (meaning that any two W*-probability spaces can be jointly embedded into a third), it follows from [21, Corollary 2.16] that, for each sentence σ in the language of W*-probability spaces, there is a unique real number r such that the property $\sigma = r$ is an enforceable property; in this case, we set σ^f to denote this unique r.

Definition 3.23. A W*-probability space (M, φ) is called *finitely generic* if it satisfies the following two properties:

- (1) For every sentence σ in the language of W*-probability spaces, we have that $\sigma^{(M,\phi)} = \sigma^f$.
- (2) For any W*-probability space (N, ψ) elementarily equivalent to (M, ϕ) for which $(M, \phi) \subseteq (N, \psi)$, we have that this inclusion is an elementary embedding.

The following was shown in [21, Section 3], generalizing the corresponding classical results:

Fact 3.24.

- (1) The property of being finitely generic is enforceable. In particular, finitely generic structures exist.
- (2) Finitely generic structures are e.c.

In [15, Corollary 6.4], it was shown that \Re is a finitely generic QWEP tracial von Neumann algebra. We prove an analogous result here:

Theorem 3.25. There is a faithful normal state φ on R_{∞} so that (R_{∞}, φ) is a finitely generic QWEP von Neumann algebra.

Proof. Simply apply the Conjunction Lemma to the fact that both being approximately finite dimensional is enforceable (Proposition 3.22) and being finitely generic is enforceable. \Box

3.5. **Open questions.** We end this section with some open questions:

As shown in the last subsection, the enforceable QWEP factor does not exist. As mentioned above, a positive solution to the Connes Embedding Problem is equivalent to the statement that R is the enforceable tracial von Neumann algebra. Due to the recent negative solution of the Connes Embedding Problem [32], we conclude that R is not the enforceable tracial von Neumann algebra. However, whether or not the enforceable tracial von Neumann algebra exists is an interesting open question. The analogous question for W*-probability spaces is also open:

Question 3.26. Does the enforceable QWEP W*-probability space exist?

In connection with Theorem 3.25, we ask if the analog of Proposition 3.8 holds for finitely generic structures:

Question 3.27. If (M, φ) is finitely generic for some $\varphi \in \mathfrak{S}_{fn}(M)$, do we have that (M, ψ) is finitely generic for *every* $\psi \in \mathfrak{S}_{fn}(M)$?

For the next question, recall that the class of W*-probability spaces whose underlying von Neumann algebra is a III_{λ} factor (for some fixed $\lambda \in (0,1)$) and whose state is a periodic state forms an inductive class. This result is clear from the fact that this class is closed under ultraproducts and ultraroots (see [1, Theorem 6.11]) and inductive limits. Explicit axioms for this class were given by Dabrowski in [13]. Let T_{λ} denote some collection of axioms for this class. Since T_{λ} is an inductive theory, it has e.c. models.

Question 3.28. What can we say about the class of e.c. models of T_{λ} ? If we restrict to QWEP such objects, is R_{λ} , with its unique periodic state, an e.c. model? Does being e.c. depend on the state?

We should note that the argument appearing in the proof of Theorem 3.9(2) above shows that e.c. models of T_{λ} tensorially absorb R_{λ} (equipped with its unique periodic state) and the same is true for the e.c. elements of the class of QWEP models of T_{λ} .

We also note that by the main result of [3], given any $\lambda \in (0, 1)$, a von Neumann algebra is QWEP if and only if it embeds into $R^{\mathfrak{U}}_{\lambda}$ with expectation. If one could

improve this result so that any QWEP model (M, φ) of T_{λ} embeds into $(R_{\lambda}, \varphi_{\lambda})^{\mathfrak{U}}$, then one could mimic the proof of 3.22 to show that being hyperfinite is an enforceable property with respect to the class of QWEP models of T_{λ} . Since being e.c. is also an enforceable property, one could conclude that $(R_{\lambda}, \varphi_{\lambda})$ is an e.c. element of the class of QWEP models of T_{λ} . In fact, we have the following:

Proposition 3.29. *The following statements are equivalent:*

- (1) Every QWEP model of T_{λ} embeds into $(R_{\lambda}, \varphi_{\lambda})^{\mathfrak{U}}$.
- (2) Hyperfiniteness is an enforceable property of QWEP models of T_{λ} .
- (3) $(R_{\lambda}, \varphi_{\lambda})$ is the enforceable QWEP model of T_{λ} .
- (4) $(R_{\lambda}, \varphi_{\lambda})$ is an e.c. element of the class of QWEP models of T_{λ} .

Whether or not the above statements indeed hold, by Corollary A.6, $(R_{\lambda}, \phi_{\lambda})$ is an e.c. model of $Th_{\forall}(R_{\lambda}, \phi_{\lambda})$.

In connection with the previous question, one might also ask the following:

Question 3.30. Suppose that (M, φ) is a type III_{λ} factor equipped with a periodic state. Is (M, φ) an e.c. model of T_{λ} if and only if M_{φ} is an e.c. II_{1} factor?

An even more basic question about models of T_{λ} arises:

Question 3.31. If M is a III_{λ} factor and ϕ and ψ are both periodic states on M, must (M, ϕ) and (M, ψ) have the same universal theory?

Our final question returns to the general study of e.c. W*-probability spaces:

Question 3.32. Are any two e.c. (QWEP) W^* -probability spaces elementarily equivalent?

The analogous question for II_1 factors remains unsettled but is presumed to have a negative answer.

4. Other model-theoretic results

4.1. **Axiomatizability results.** In [1], it was shown that, for any $\lambda \in (0, 1]$, the Ocneanu ultraproduct of a family of type III_{λ} factors is once again a type III_{λ} factor and a σ -finite von Neumann algebra is a type III_{λ} factor if the same is true of an Ocneanu ultrapower. By the "soft" test for axiomatizability (see [5, Proposition 5.14]), this says that the class of III_{λ} factors (again, for fixed $\lambda \in (0,1]$) is an axiomatizable class. However, this test does not give us explicit axioms for these axiomatizable classes. Nevertheless, it is possible to at least describe the quantifier complexity of such axiomatizations.

In order to state such quantifier complexity results and the test we use for obtaining them, we define a \forall_n -sentence to be one of the form

$$\sup_{x_1}\inf_{x_2}\cdots Q_{x_n}\theta(x_1,\ldots,x_n),$$

where each x_i is a finite block of variables, θ is a quantifier-free formula, and $Q = \sup$ if n is odd and $Q = \inf$ if n is even. The notion of a \exists_n -sentence is defined analogously, beginning with a block of inf quantifiers instead of sup quantifiers. By using dummy variables, we note that any \forall_n - or \exists_n -sentence is automatically both \forall_{n+1} and \exists_{n+1} . Finally, we say that an axiomatizable class is \forall_n -axiomatizable (resp. \exists_n -axiomatizable) if there is a set T of axioms for the class consisting solely of conditions of the form $\sigma = 0$ with σ a nonnegative \forall_n -sentence (resp. a \exists_n -sentence).

In classical logic, given an axioimatizable class, the *Keisler Sandwich Theorem* [33] can be used to prove the existence of an axiomatization of a particular kind of quantifier complexity. In our results below, we use (a modified version of) the Keisler Sandwich Theorem, adapted to the setting of continuous logic. Since neither the continuous version of the Keisler Sandwich Theorem, nor the variant presented here, have appeared in the literature before, we include a proof in Appendix B.

Definition 4.1. Given structures M and N (in the same language) and $n \ge 1$, an (M, N, n)-ultrapower sandwich is a chain of embeddings of the form

$$M \hookrightarrow N \hookrightarrow M^{\mathcal{U}_1} \hookrightarrow N^{\mathcal{U}_2} \hookrightarrow \cdots \hookrightarrow Q^{\mathcal{U}_{n-1}}$$

where:

- $\mathcal{U}_1, \ldots, \mathcal{U}_{n-1}$ are ultrafilters,
- Q = M if n is even and Q = N if n is odd, and
- all compositions $M^{\mathcal{U}_{k-1}} \hookrightarrow M^{\mathcal{U}_{k+1}}$ and $N^{\mathcal{U}_{k-1}} \hookrightarrow N^{\mathcal{U}_{k+1}}$ are the diagonal embeddings (which in particular implies that $\mathcal{U}_{k+1} \cong \mathcal{U}_{k-1} \otimes \mathcal{V}$ for some ultrafilter \mathcal{V}).

Theorem 4.2 (Keisler Sandwich Theorem (Ultrapower version)). *Suppose that* T *is a theory. Then:*

- (1) T is \forall_n -axiomatizable if and only if: whenever there is an (M, N, n)-ultrapower sandwich and $N \models T$, then $M \models T$.
- (2) T is \exists_n -axiomatizable if and only if: whenever there is an (M, N, n)-ultrapower sandwich and $M \models T$, then $N \models T$.

Note that an (M, N, 2)-ultrapower sandwich is simply a chain of embeddings $M \hookrightarrow N \hookrightarrow M^{\mathfrak{U}}$ such that the embedding $M \hookrightarrow M^{\mathfrak{U}}$ is the diagonal embedding.

Thus, an (M, N, 2)-ultrapower sandwich exists precisely when M is e.c. in N. By Theorem 4.2 and Corollary 3.7, we immediately have:

Theorem 4.3. *The class of type* III_1 *factors is* \forall_2 -axiomatizable.

Alternate proof of Theorem 4.3. There is an alternative to showing that a theory T is \forall_2 -axiomatizable and that is to show that the collection of models of the theory is closed under unions of chains. This can be established quite easily for the case of III₁ factors as follows. Let $(M_i, \phi_i)_{i \in I}$ be any increasing chain of type III₁ W*-probability spaces. Denote by (M, ϕ) the inductive limit of $(M_i, \phi_i)_{i \in I}$ in the language of W*-probability spaces. Then for every $i \in I$, we have $(M_i, \phi_i) \subseteq (M, \phi)$ and we denote by $E_i : M \to M_i$ the unique faithful normal conditional expectation such that $\phi_i \circ E_i = \phi$. Also, we have $\bigvee_{i \in I} M_i = M$.

For every $i \in I$, we may consider the trace-preserving inclusion of continuous cores $c_{\phi_i}(M_i) \subseteq c_{\phi}(M)$ with trace-preserving faithful normal conditional expectation $F_i : c_{\phi}(M) \to c_{\phi_i}(M_i)$. Since M_i is a type III_1 factor, $c_{\phi_i}(M_i)$ is a type II_{∞} factor. In order to show that M is a type III_1 factor, it suffices to show that $c_{\phi}(M)$ is a factor. Let $x \in \mathcal{Z}(c_{\phi}(M))$ be any central element. Then for every $i \in I$, we have $F_i(x) \in \mathcal{Z}(c_{\phi_i}(M_i))$ and so $F_i(x) \in \mathbb{C}1$. Since $\bigvee_{i \in I} c_{\phi_i}(M_i) = c_{\phi}(M)$, it follows that $F_i(x) \to x$ *-strongly as $i \to \infty$, which implies that $x \in \mathbb{C}1$.

The case of type III_{λ} factors for $\lambda \in (0, 1)$ is inherently more complicated:

Proposition 4.4. For any $\lambda \in (0,1)$, the axiomatizable class of type III_{λ} factors is not \forall_2 -axiomatizable.

Proof. Fix $\lambda \in (0,1)$. We construct an increasing sequence of W*-probability space type III_{λ} factors (M_n,ϕ_n) so that, setting $(M,\phi) := \bigvee_{n \in \mathbb{N}} (M_n,\phi_n)$, we have that M is a type III_1 factor.

Take $\lambda_1,\lambda_2\in(0,1)$ with $\log(\lambda_1)/\log(\lambda_2)$ irrational, whence $R_{\lambda_1}\otimes R_{\lambda_2}\cong R_{\infty}$. For every $n\in\mathbb{N}$, set $(M_n,\phi_n)=(R_{\lambda_1},\phi_{\lambda_1})\otimes(M_2(\mathbb{C}),\omega_{\lambda_2})^{\otimes\{0,\dots,n\}}$. Then $(M_n,\phi_n)_{n\in\mathbb{N}}$ is an increasing sequence of W*-probability spaces type III_{λ_1} factors. Indeed, observe that for every $n\in\mathbb{N}$, we have $M_n\cong R_{\lambda_1}$. Moreover, for every $n\in\mathbb{N}$, we have $\phi_{n+1}|_{M_n}=\phi_n$ and the linear mapping $E_n:M_{n+1}\to M_n$ defined by $E_n=\mathrm{id}_{M_n}\otimes\omega_{\lambda_2}$ is a faithful normal conditional expectation such that $\phi_n\circ E_n=\phi_{n+1}$.

However, we have $\bigvee_{n\in\mathbb{N}}(M_n,\phi_n)\cong(R_{\lambda_1}\otimes R_{\lambda_2},\phi_{\lambda_1}\otimes\phi_{\lambda_2})=:(M,\phi)\cong(R_\infty,\phi).$ Indeed, for every $n\in\mathbb{N}$, we have $\phi|_{M_n}=\phi_n$ and the linear mapping $F_n:M\to M_n$ defined by $F_n=\mathrm{id}_{M_n}\otimes\omega_{\lambda_2}^{\otimes\mathbb{N}\setminus\{0,\dots,n\}}$ is a faithful normal conditional expectation such that $\phi_n\circ F_n=\phi$. Thus, the inductive limit of W*-probability spaces $(M,\phi)=\bigvee_{n\in\mathbb{N}}(M_n,\phi_n)$ is a type III₁ factor.

We also note:

Proposition 4.5. *For any* $\lambda \in (0, 1]$ *, the class of type* III_{λ} *factors is not* \exists_2 -axiomatizable.

Proof. Fix $\lambda \in (0, 1]$ and suppose that M is a type III_{λ} factor that is not full (e.g. $M = R_{\lambda}$). Then there is a σ-finite von Neumann algebra N such that N is not a factor and yet $M \subseteq N \subseteq M^{\mathfrak{U}}$. By Theorem 4.2, the result follows.

While the class of type III_{λ} factors, $\lambda \in (0,1)$, cannot be axiomatized using two quantifiers, we now show that it can be axiomatized using three quantifiers. The key to proving this is the following:

Proposition 4.6. Suppose that we have an (M, N, 3)-sandwich with M a type III_{μ} factor and N a type III_{λ} factor, $\lambda, \mu \in (0, 1)$. Then $\lambda = \mu$.

Proof. Set $T_{\lambda} = \frac{2\pi}{|\log(\lambda)|}$ and $T_{\mu} = \frac{2\pi}{|\log(\mu)|}$. Choose a T_{λ} -periodic faithful normal state ψ on N and apply Proposition 3.3 to the inclusion $N \subseteq M^{\mathcal{U}_1} \subseteq N^{\mathcal{U}_2}$. Since $\psi^{\mathcal{U}_2}|_{M^{\mathcal{U}_1}}$ is T_{λ} -periodic, we have $\{0\} \cup \mu^{\mathbb{Z}} = S(M^{\mathcal{U}_1}) \subseteq \sigma(\Delta_{\psi^{\mathcal{U}_2}|_{M^{\mathcal{U}_1}}}) \subseteq \{0\} \cup \lambda^{\mathbb{Z}}$. Next, choose a T_{μ} -periodic faithful normal state φ on M and apply Proposition 3.3 to the inclusion $M \subseteq N \subseteq M^{\mathcal{U}_1}$. Since $\phi^{\mathcal{U}_1}|_N$ is T_{μ} -periodic, we have $\{0\} \cup \lambda^{\mathbb{Z}} = S(N) \subseteq \sigma(\Delta_{\omega^{\mathcal{U}_1}|_N}) \subseteq \{0\} \cup \mu^{\mathbb{Z}}$. This shows that $\mu = \lambda$. □

Corollary 4.7. *The class of type* III_{λ} *factors is both* \forall_3 *- and* \exists_3 *-axiomatizable.*

Proof. Consider a (M, N, 3)-sandwich. First suppose that N is a III $_{\lambda}$ factor. By Lemma 3.4, M is a factor. Since $M^{\mathfrak{U}_1}$ contains N with expectation, $M^{\mathfrak{U}_1}$ is type III, whence so is M. By Theorem 3.5, M is not of type III $_0$. If M is type III $_1$, then so is $M^{\mathfrak{U}_1}$; since N is e.c. in $M^{\mathfrak{U}_1}$, Corollary 3.7 implies that N is type III $_1$, a contradiction. Thus, M is type III $_{\mu}$ for some $\mu \in (0,1)$, whence $\lambda = \mu$ by Proposition 4.6. This shows that the class of III $_{\lambda}$ factors is \forall_3 -axiomatizable.

Now suppose that M is a III_{λ} factor. Since N is e.c. in the III_{λ} factor $M^{\mathfrak{U}_1}$, we see that N is a factor. Since N contains the type III factor M with expectation, we see that N has type III. By Corollary 3.7, N does not have type III_1 . Since N is e.c. in the III_{λ} factor $M^{\mathfrak{U}_1}$, N does not have type III_0 by Theorem 3.5. Thus, N has type III_{μ} for some $\mu \in (0,1)$, and thus $\lambda = \mu$ by Proposition 4.6. This shows that the class of III_{λ} factors is \exists_3 -axiomatizable.

- 4.2. **First-order theories of W*-probability spaces.** In this final subsection, we consider the task of counting the number of first-order theories of III_{λ} W*-probability spaces for $\lambda \in (0,1]$.
- In [6], Boutonnet, Chifan, and Ioana showed that there exist continuum many pairwise non-elementarily equivalent separable II₁ factors. More precisely, they showed that the family $(M_{\alpha})_{\alpha \in 2^{\mathbb{N}}}$ of separable II₁ factors constructed by McDuff in [36] provides such a continuum. We observe that their result can be easily

applied to construct such a continuum in the realm of type III_{λ} factors for $\lambda \in (0,1)$.

Theorem 4.8. Fix $\lambda \in (0, 1)$. Then there exist continuum many non-pairwise elementary equivalent separable type III_{λ} factors.

Proof. Consider the family $(M_{\alpha})_{\alpha\in 2^{\mathbb{N}}}$ of separable II₁ factors constructed by McDuff in [36]. For every $\alpha\in 2^{\mathbb{N}}$, M_{α} is a McDuff factor, that is, $M_{\alpha}\otimes R\cong M_{\alpha}$. This implies that $\lambda\in \mathcal{F}(M_{\alpha})$. Set $M_{\alpha}^{\infty}=M_{\alpha}\otimes B(\ell^2)$ and choose an automorphism $\theta_{\alpha}^{\lambda}\in Aut(M_{\alpha}^{\infty})$ such that $\tau\circ\theta_{\alpha}^{\lambda}=\lambda\tau$, where τ is any faithful normal semifinite trace on M_{α}^{∞} . Then $N_{\alpha}=M_{\alpha}^{\infty}\rtimes_{\theta_{\alpha}^{\lambda}}\mathbb{Z}$ is a type III_{λ} factor whose discrete core is isomorphic to M_{α}^{∞} (see [7, Théorème 4.4.1]).

Now suppose that $\alpha, \beta \in 2^{\mathbb{N}}$ are such that N_{α} is elementarily equivalent to N_{β} . By the Keisler-Shelah Theorem, there are ultrafilters \mathcal{U} and \mathcal{V} such that $(N_{\alpha})^{\mathcal{U}} \cong (N_{\beta})^{\mathcal{V}}$. By [1, Proposition 4.7], the discrete core of $(N_{\alpha})^{\mathcal{U}}$ (resp. $(N_{\beta})^{\mathcal{V}}$) is $(M_{\alpha}^{\infty})^{\mathcal{U}}$ (resp. $(M_{\beta}^{\infty})^{\mathcal{V}}$). Then [7, Th'eor'eme 4.4.1] implies that $M_{\alpha}^{\mathcal{U}} \otimes B(\ell^2) = (M_{\alpha}^{\infty})^{\mathcal{U}} \cong (M_{\beta}^{\infty})^{\mathcal{V}} = M_{\beta}^{\mathcal{V}} \otimes B(\ell^2)$. Since $M_{\alpha}^{\mathcal{U}}$ and $M_{\beta}^{\mathcal{V}}$ have full fundamental group, we have $M_{\alpha}^{\mathcal{U}} \cong M_{\beta}^{\mathcal{V}}$ and [6] further implies that $\alpha = \beta$.

The preceding question naturally raises the following:

Question 4.9. Do there exist continuum many non-pairwise elementary equivalent separable type III_1 factors?

While we cannot yet answer the above question, we can at least find three such factors. First, due to the recent negative solution of the Connes Embedding Problem [32], there must exist a non-QWEP type III_1 factor M. Consequently, M is not a model of $Th_\forall(R_\infty)$, and thus is not elementarily equivalent to R_∞ . To find a third theory of type III_1 factors, we recall that in [18, 3.2.2], property Gamma was shown to be an axiomatizable property and thus could be used to distinguish theories of II_1 factors. The correct generalization of property Gamma to our context is that of being non-full. Here, we show that the non-full type III_λ factors form an axiomatizable class and, as a consequence, that the non-full factors form a *local class*, that is, closed under ultrapowers and ultraroots (which is still sufficient for differentiating between theories):

Theorem 4.10. For $\lambda \in (0, 1]$, the class of non-full type III_{λ} factors is axiomatizable.

Proof. We use the aforementioned "soft" test for being axiomatizable, that is, we show that the class of non-full type III_{λ} factors is closed under ultraproducts and ultraroots. We first show the latter. Let \mathcal{U} be any nonprincipal ultrafilter and let M be any full type III_{λ} factor; we show that $M^{\mathcal{U}}$ is also full. To see this, let \mathcal{V} be any nonprincipal ultrafilter. Then we have $(M^{\mathcal{U}})^{\mathcal{V}} = M^{\mathcal{V} \otimes \mathcal{U}}$ by [2, Proposition

2.4]. Since M is a full factor, we have $M' \cap M^{v \otimes u} = \mathbb{C}1$. Since $M \subset M^u$, we have $(M^u)' \cap (M^u)^v \subseteq M' \cap (M^u)^v = M' \cap M^{v \otimes u} = \mathbb{C}1$. Since this holds true for any nonprincipal ultrafilter V, this further implies that M^u is a full factor.

Conversely, let $(M_i, \varphi_i)_{i \in I}$ be any family of non-full type III_λ factor W^* -probability spaces and $\mathcal U$ a nonprincipal ultrafilter on I. Then $(M^{\mathcal U}, \varphi^{\mathcal U}) = (M_i, \varphi_i)^{\mathcal U}$ is a type III_λ factor. We show that $M^{\mathcal U}$ is also non-full. It is well-known that we may assume that $\mathcal U$ is countably incomplete, meaning that there is some countable collection of sets from $\mathcal U$ that has empty intersection. As a result, this allows us to define a sequence $(\varepsilon_i)_{i \in I}$ of positive real numbers such that $\lim_{\mathcal U} \varepsilon_i = 0$. For every $i \in I$, there is an index set J_i for which there exists a sequence $(u_j^i)_{j \in J_i}$ in $\mathcal U(M_i)$ such that $\lim_i ||u_i^i \varphi_i - \varphi_i u_i^i|| = 0$, $\lim_i ||yu_i^i - u_i^i y||_{\varphi_i}^\sharp = 0$ for every $y \in M_i$ and $\lim_i \varphi_i(u_i^i) = 0$. Let $\mathcal F = \{X_1, \dots, X_m\} \subset M^{\mathcal U}$ be any finite subset. Write $X_k = (x_{k,i})^{\mathcal U}$ for every $1 \le k \le m$. For every $i \in I$, there exists $v_i = u_{j_i} \in \mathcal U(M_i)$ such that $||v_i \varphi_i - \varphi_i v_i|| \le \varepsilon_i$, $||v_i x_{k,i} - x_{k,i} v_i||_{\varphi_i}^\sharp \le \varepsilon_i$ and $||\varphi_i (v_i)|| \le \varepsilon_i$ for every $1 \le k \le m$. Then we have $(v_i)_{i \in I} \in \mathfrak M^{\mathcal U}(M_i, \varphi_i)$, $v_{\mathcal F} = (v_i)^{\mathcal U} \in \mathcal U(M^{\mathcal U})_{\varphi^{\mathcal U}}$ and $\varphi^{\mathcal U}(v_{\mathcal F}) = 0$ (see [28, Proposition 2.4] and [1, Lemma 4.36]). By construction, we have $Xv_{\mathcal F} = v_{\mathcal F} X$ for every $X \in \mathcal F$ and $\varphi^{\mathcal U}(v_{\mathcal F}) = 0$. Since this holds true for any finite subset $\mathcal F \subset M^{\mathcal U}$, this further implies that $M^{\mathcal U}$ is not full.

Corollary 4.11. The following three III_1 factors are pairwise non-elementarily equivalent:

- (1) R_{∞} .
- (2) Any non-QWEP III₁ factor.
- (3) Any full QWEP III₁ factor.

Remark 4.12. In [18], the authors provide explicit axioms for having property Gamma. It would be interesting to find explicit axioms for the class of non-full III_{λ} W*-probability spaces.

In [18, Theorem 4.3], it is shown that, for any separable II₁ factor M, there are continuum many pairwise nonisomorphic separable II₁ factors elementarily equivalent to M. We ask if the analogous result holds true for W*-probability spaces:

Question 4.13. For any separable W*-probability space (M, φ) , are there continuum many nonisomorphic separable (N, ψ) elementarily equivalent to (M, φ) ?

By Proposition 3.13, if M is any type III_1 factor, then (M,ϕ) is elementarily equivalent to (M,ψ) for all $\phi,\psi\in\mathfrak{S}_{fn}(M)$. In this sense, the previous question has a somewhat trivial positive solution for type III_1 factors. It would be more interesting to find continuum many nonisomorphic separable models of any given theory of W*-probability spaces whose underlying von Neumann algebras themselves are pairwise nonisomorphic.

APPENDIX A. EMBEDDINGS OF AFD W*-PROBABILITY SPACES INTO ULTRAPRODUCTS

Most of the results presented in this appendix are due to Ando and Houdayer (unpublished work).

For the ease of exposition, all ultrafilters in this appendix are assumed to be nonprincipal ultrafilters on \mathbb{N} . However, all of the results hold verbatim for countably incomplete ultrafilters on arbitrary index sets with only routine modifications of the proofs needed. These more general versions of the results are what are used throughout the main part of the paper.

Let (M, φ) be any type III_{λ} factor, where $\lambda \in (0, 1]$, endowed with a faithful normal state such the centralizer $(M^{\mathfrak{U}})_{\varphi^{\mathfrak{U}}}$ of the ultraproduct state $\varphi^{\mathfrak{U}}$ is a type II_1 factor.

- If M is of type III_{λ} with $\lambda \in (0, 1)$, we may take any $\frac{2\pi}{|\log(\lambda)|}$ -periodic faithful normal state φ on M.
- If M is of type III₁, we may take any faithful normal state φ on M (see [1, Proposition 4.24]).

First, we observe that any atomic (discrete) W*-probability space has a unique embedding into $(M, \phi)^{\mathfrak{U}}$ up to unitary conjugacy.

Lemma A.1. Let (P, ψ) be any atomic W^* -probability space. For every $i \in \{1, 2\}$, let $\pi_i : (P, \psi) \hookrightarrow (M, \phi)^{\mathfrak{U}}$ be any embedding. Then there exists $\mathfrak{u} \in \mathfrak{U}((M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}})$ such that $\pi_2 = Ad(\mathfrak{u}) \circ \pi_1$.

Proof. Write $P=\bigoplus_p P_p$ where each P_p is a type I factor. For every p, denote by $(e^p_{kl})_{k,l}$ a system of matrix units for P_p and denote by $(\lambda^p_k)_k$ positive reals such that $\sum_k \lambda^p_k = 1$ and $\frac{\psi(\cdot 1_{P_p})}{\psi(1_{P_p})} = \tau_{P_p}(diag(\lambda^p_k) \cdot)$. For all k,l,p and all $t \in \mathbb{R}$, we have $\sigma^\psi_t(e^p_{kl}) = (\lambda^p_k/\lambda^p_l)^{it}e^p_{kl}$.

Observe for every $i \in \{1,2\}$, since $\phi^{\mathfrak{U}} \circ \pi_i = \psi$ and since $\pi_i(P) \subset M^{\mathfrak{U}}$ is globally invariant under $\sigma^{\phi^{\mathfrak{U}}}$, we have $\sigma^{\phi^{\mathfrak{U}}}_t \circ \pi_i = \pi_i \circ \sigma^{\psi}_t$ for all $t \in \mathbb{R}$ by [7, Lemme 1.2.10]. Since $(M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}}$ is a type II_1 factor and since $\pi_1(e^p_{11}), \pi_2(e^p_{11}) \in (M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}}$ and $\phi^{\mathfrak{U}}(\pi_1(e^p_{11})) = \psi(e^p_{11}) = \phi^{\mathfrak{U}}(\pi_2(e^p_{11}))$, there exists a partial isometry $\nu_p \in (M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}}$ such that $\nu_p^* \nu_p = \pi_1(e^p_{11})$ and $\nu_p \nu_p^* = \pi_2(e^p_{11})$. If we let $\mathfrak{U} = \sum_p \sum_k \pi_2(e^p_{k1}) \nu_p \pi_1(e^p_{1k})$, we have $\mathfrak{U} \in \mathfrak{U}(M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}}$ and $\pi_2(e^p_{k1}) = \mathfrak{U}(e^p_{k1})\mathfrak{U}^*$ for all k, l, p. Therefore, we have $\pi_2 = \mathrm{Ad}(\mathfrak{U}) \circ \pi_1$.

Next, we prove that the unique embedding property into $(M, \varphi)^{\mathcal{U}}$ up to unitary conjugacy, is stable under taking increasing unions.

Lemma A.2. Let (P, ψ) be any separable W^* -probability space and $(P_n, \psi_n) \subseteq (P, \psi)$ any increasing sequence of W^* -probability subspaces such that $\bigvee_{n \in \mathbb{N}} (P_n, \psi_n) = (P, \psi)$. For every $i \in \{1, 2\}$, let $\pi_i : (P, \psi) \hookrightarrow (M, \phi)^{\mathfrak{U}}$ be any embedding. Assume that for every $n \in \mathbb{N}$, there exists $u_n \in \mathfrak{U}((M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}})$ such that $\pi_2(x) = u_n \pi_1(x) u_n^*$ for every $x \in P_n$.

Then there exists $u \in \mathcal{U}((M^{\mathcal{U}})_{\omega^{\mathcal{U}}})$ such that $\pi_2 = Ad(u) \circ \pi_1$.

Proof. For every $n \in \mathbb{N}$, write $u_n := (u_m^n)^{\mathfrak{U}}$ where $(u_m^n)_m \in \mathfrak{M}^{\mathfrak{U}}(M)$ and $u_m^n \in \mathfrak{U}(M)$ (see e.g. [30, Lemma 2.1]). For every $n \in \mathbb{N}$, denote by $X_n = \{y_k^n \mid k \in \mathbb{N}\} \subset P_n$ a $\|\cdot\|_{\psi_n}^\sharp$ -dense countable subset and set $X_n^{\leq n} := \{y_k^n \mid 0 \leq k \leq n\}$. For every $i \in \{1,2\}$ and every $b \in X_n$, write $\pi_i(b) = (b_m^i)^{\mathfrak{U}}$ where $(b_m^i)_m \in \mathfrak{M}^{\mathfrak{U}}(M)$.

For every $n \in \mathbb{N}$, define

$$\begin{split} F_n := \bigcap_{0 \leq k \leq n, b \in X_k^{\leq k}} \left\{ m \in \mathbb{N} \mid \|u_m^n \, b_m^1 \, (u_m^n)^* - b_m^2 \|_\phi^\sharp < \frac{1}{n+1} \right\} \\ G_n := \left\{ m \in \mathbb{N} \mid m \geq n \right\} \cap \left\{ m \in \mathbb{N} \mid \|u_m^n \phi - \phi u_m^n\| < \frac{1}{n+1} \right\} \cap \bigcap_{i=1}^n F_j. \end{split}$$

By construction and since $\mathcal U$ is a nonprincipal ultrafilter on $\mathbb N$, $(G_n)_{n\in\mathbb N}$ is a decreasing sequence of subsets of $\mathcal U$ such that $G_0=\mathbb N$ and $\bigcap_{n\in\mathbb N}G_n=\emptyset$. For every $m\in\mathbb N$, set $\nu_m=\mathfrak u_m^n\in\mathcal U(M)$ where $n\in\mathbb N$ is the unique integer such that $m\in G_n\setminus G_{n+1}$.

Let $n \in \mathbb{N}$. If $m \in G_n = \bigcup_{j \geq n} G_j \setminus G_{j+1}$, denote by $p \geq n$ the unique integer such that $m \in G_p \setminus G_{p+1}$. Since $\nu_m = \mathfrak{u}_m^p$, we have

- $\bullet \ \|\nu_m\phi-\phi\nu_m\|=\|u_m^p\phi-\phi u_m^p\|<\tfrac{1}{p+1}\le \tfrac{1}{n+1} \text{ and }$
- $\|v_m b_m^1 v_m^* b_m^2\|_{\phi}^{\sharp} = \|u_m^p b_m^1 (u_m^p)^* b_m^2\|_{\phi}^{\sharp} < \frac{1}{p+1} \le \frac{1}{n+1} \text{ for all } 0 \le k \le n$ and all $b \in X_{\nu}^{\le k}$.

This implies that

$$\begin{split} G_n &\subset \left\{ m \in \mathbb{N} \mid \|\nu_m \phi - \phi \nu_m\| < \frac{1}{n+1} \right\} \\ G_n &\subset \bigcap_{0 \leq k \leq n, b \in X_k^{\leq k}} \left\{ m \in \mathbb{N} \mid \|\nu_m \, b_m^1 \, \nu_m^* - b_m^2\|_\phi^\sharp < \frac{1}{n+1} \right\}. \end{split}$$

Since $G_n \in \mathcal{U}$, it follows that

$$\left\{ m \in \mathbb{N} \mid \|\nu_m \phi - \phi \nu_m\| < \frac{1}{n+1} \right\} \in \mathfrak{U}$$

$$\forall 0 \leq k \leq n, \forall b \in X_k^{\leq k}, \quad \left\{ m \in \mathbb{N} \mid \|\nu_m \, b_m^1 \, \nu_m^* - b_m^2\|_\phi^\sharp < \frac{1}{n+1} \right\} \in \mathfrak{U}.$$

Since this holds for every $n \in \mathbb{N}$, we obtain $\lim_{m \to \mathcal{U}} \| \nu_m \phi - \phi \nu_m \| = 0$. This implies that $(\nu_m)_m \in \mathfrak{M}^{\mathcal{U}}(M)$ and $u = (\nu_m)^{\mathcal{U}} \in \mathcal{U}((M^{\mathcal{U}})_{\phi^{\mathcal{U}}})$. This further implies that for every $k \in \mathbb{N}$ and every $b \in X_k^{\leq k}$, we have $\|u\pi_1(b)u^* - \pi_2(b)\|_{\phi^{\mathcal{U}}}^{\sharp} = \lim_{m \to \mathcal{U}} \|\nu_m b_m^1 \nu_m^* - b_m^2\|_{\phi}^{\sharp} = 0$. Since $\bigcup_{k \in \mathbb{N}} X_k^{\leq k} = \bigcup_{n \in \mathbb{N}} X_n$, since for every $n \in \mathbb{N}$, the set X_n is $\|\cdot\|_{\psi_n}^{\sharp}$ -dense in P_n , since $\bigvee_{n \in \mathbb{N}} P_n = P$ and since for every $i \in \{1, 2\}$, $\pi_i : P \hookrightarrow M^{\mathcal{U}}$ is a normal embedding, this finally implies that $\pi_2(x) = u\pi_1(x)u^*$ for every $x \in P$.

Definition A.3. We say that a W*-probability space (P, ψ) is approximately finite dimensional (AFD) if there exists an increasing sequence of finite dimensional W*-probability subspaces $(P_n, \psi_n) \subseteq (P, \psi)$ such that $\bigvee_{n \in \mathbb{N}} (P_n, \psi_n) = (P, \psi)$.

If (P, ψ) is an AFD W*-probability space, then ψ is necessarily an almost periodic state on P.

Examples A.4. Here are the main examples of AFD W*-probability spaces:

- (1) Every AFD tracial von Neumann algebra (M, τ) endowed with a tracial faithful normal state is an AFD W*-probability space.
- (2) For every $\lambda \in (0,1)$, endow the type III_{λ} Powers factor R_{λ} with its canonical $\frac{2\pi}{|\log(\lambda)|}$ -periodic faithful normal state ϕ_{λ} . Then $(R_{\lambda},\phi_{\lambda})$ is an AFD W*-probability space.
- (3) Endow the type III_1 Araki-Woods factor $R_{\infty}=R_{\lambda_1}\otimes R_{\lambda_2}$, where $\frac{\log(\lambda_1)}{\log(\lambda_2)}\notin\mathbb{Q}$, with the faithful normal state $\phi=\phi_{\lambda_1}\otimes\phi_{\lambda_2}$. Then (R_{∞},ϕ) is an AFD W*-probability space.
- (4) For every AFD type III_0 factor P, there exists a faithful normal state φ on M such that (M, φ) is an AFD W*-probability space (see [9, Theorem 1]).

We deduce that any AFD W*-probability space has a unique embedding into $(M, \varphi)^{\mathfrak{U}}$ up to unitary conjugacy.

Theorem A.5. Let (P, ψ) be any AFD W*-probability space. For every $i \in \{1, 2\}$, let $\pi_i : (P, \psi) \hookrightarrow (M, \phi)^{\mathfrak{U}}$ be any embedding. Then there exists $\mathfrak{u} \in \mathfrak{U}((M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}})$ such that $\pi_2 = Ad(\mathfrak{u}) \circ \pi_1$.

Proof. Let $(P_n, \psi_n) \subseteq (P, \psi)$ be an increasing sequence of finite dimensional W*-probability subspaces such that $\bigvee_{n \in \mathbb{N}} (P_n, \psi_n) = (P, \psi)$. For every $n \in \mathbb{N}$, Lemma A.1 implies that there exists $u_n \in \mathcal{U}((M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}})$ such that $\pi_2(x) = u_n \pi_1(x) u_n^*$ for every $x \in P_n$. Then Lemma A.2 implies that there exists $u \in \mathcal{U}((M^{\mathfrak{U}})_{\phi^{\mathfrak{U}}})$ such that $\pi_2(x) = u \pi_1(x) u^*$ for every $x \in P$.

As a straightforward consequence of Theorem A.5, we obtain the following unique embedding property for the Powers factors R_{λ} where $\lambda \in (0, 1)$.

Corollary A.6. Let $\lambda \in (0,1)$ and $\pi : (R_{\lambda}, \phi_{\lambda}) \hookrightarrow (R_{\lambda}, \phi_{\lambda})^{\mathfrak{U}}$ be any embedding. Then there exists $\mathfrak{u} \in \mathfrak{U}((R_{\lambda}^{\mathfrak{U}})_{\phi_{\lambda}^{\mathfrak{U}}})$ such that $Ad(\mathfrak{u}) \circ \pi : (R_{\lambda}, \phi_{\lambda}) \hookrightarrow (R_{\lambda}, \phi_{\lambda})^{\mathfrak{U}}$ is the diagonal embedding.

Combining Theorem A.5 with Connes-Størmer transitivity theorem, we obtain the following unique embedding property for the Araki-Woods factor R_{∞} .

Corollary A.7. Let ψ be any faithful normal state on R_{∞} and $\pi: (R_{\infty}, \psi) \hookrightarrow (R_{\infty}, \psi)^{\mathfrak{U}}$ any embedding. Then there exists $\mathfrak{u} \in \mathfrak{U}((R_{\infty}^{\mathfrak{U}})_{\psi^{\mathfrak{U}}})$ such that $Ad(\mathfrak{u}) \circ \pi: (R_{\infty}, \psi) \hookrightarrow (R_{\infty}, \psi)^{\mathfrak{U}}$ is the diagonal embedding.

Proof. Denote by $E: R_\infty^{\mathfrak{U}} \to \pi(R_\infty)$ the unique faithful normal conditional expectation such that $\psi \circ \pi^{-1} \circ E = \psi^{\mathfrak{U}}$. Choose a faithful normal state φ on R_∞ such that (R_∞, φ) is an AFD W*-probability space (see Example A.4(3)). Set $\varphi = \varphi \circ \pi^{-1} \circ E \in (R_\infty^{\mathfrak{U}})_*$. By [1, Theorem 4.20], there exists $w \in \mathfrak{U}(R_\infty^{\mathfrak{U}})$ such that $\varphi = \varphi^{\mathfrak{U}} \circ \mathrm{Ad}(w)$. (This result indeed holds for countably incomplete ultrafilters on arbitrary index sets.) Then $\pi_w = \mathrm{Ad}(w) \circ \pi : R_\infty \hookrightarrow R_\infty^{\mathfrak{U}}$ is an embedding such that $\varphi^{\mathfrak{U}} \circ \pi_w = \varphi^{\mathfrak{U}} \circ \mathrm{Ad}(w) \circ \pi = \varphi \circ \pi = \varphi$. Moreover, the faithful normal conditional expectation $E_w = \mathrm{Ad}(w) \circ E \circ \mathrm{Ad}(w^*) : R_\infty^{\mathfrak{U}} \to w\pi(R_\infty)w^*$ satisfies $\varphi \circ \pi_w^{-1} \circ E_w = \varphi \circ \pi^{-1} \circ \mathrm{Ad}(w^*) \circ \mathrm{Ad}(w) \circ E \circ \mathrm{Ad}(w^*) = \varphi^{\mathfrak{U}}$. Thus, $\pi_w : (R_\infty, \varphi) \hookrightarrow (R_\infty, \varphi)^{\mathfrak{U}}$ is an embedding of W*-probability spaces.

By Theorem A.5, there exists $v \in \mathcal{U}((R_\infty^{\mathcal{U}})_{\phi^\mathcal{U}})$ such that $Ad(v) \circ \pi_w : (R_\infty, \phi) \hookrightarrow (R_\infty, \phi)^\mathcal{U}$ is the diagonal embedding $\iota : (R_\infty, \phi) \hookrightarrow (R_\infty, \phi)^\mathcal{U}$. Set $u = vw \in \mathcal{U}(R_\infty^\mathcal{U})$. Then we have $Ad(u) \circ \pi = \iota$ and $\phi^\mathcal{U} \circ Ad(u) \circ \pi = \phi = \phi^\mathcal{U} \circ \iota$. Denote by $E_\mathcal{U} : R_\infty^\mathcal{U} \to \iota(R_\infty)$ the canonical faithful normal expectation. Note that $E_u = Ad(u) \circ E \circ Ad(u^*) : R_\infty^\mathcal{U} \to u\pi(R_\infty)u^*$ is another faithful normal conditional expectation onto $\iota(R_\infty) = u\pi(R_\infty)u^*$. Since

$$\begin{split} \phi \circ \pi^{-1} \circ Ad(u^*) \circ E_u &= \phi \circ \pi^{-1} \circ Ad(u^*) \circ Ad(u) \circ E \circ Ad(u^*) \\ &= \varphi \circ Ad(w^*) \circ Ad(v^*) \\ &= \phi^{\mathcal{U}} \\ &= \phi \circ \iota^{-1} \circ E_{\mathcal{U}}, \end{split}$$

we have $E_{\mathfrak{U}}=E_{\mathfrak{u}}=Ad(\mathfrak{u})\circ E\circ Ad(\mathfrak{u}^*).$ This further implies that

$$\begin{split} \psi^{\mathcal{U}} &= \psi \circ \iota^{-1} \circ E_{\mathcal{U}} \\ &= \psi \circ \pi^{-1} \circ Ad(u^*) \circ Ad(u) \circ E \circ Ad(u^*) \\ &= \psi \circ \pi^{-1} \circ E \circ Ad(u^*) \end{split}$$

$$= \psi^{\mathcal{U}} \circ Ad(\mathfrak{u}^*)$$

and so $u \in \mathcal{U}((R^{\mathcal{U}}_{\infty})_{\psi^{\mathcal{U}}})$, finishing the proof.

We may also apply Theorem A.5 to the structure of ultraproduct type III_1 factors. The next result extends [14, Theorem 4.5].

Proposition A.8. Let M be any σ -finite type III_1 factor. Then $M^{\mathfrak{U}}$ is a prime factor, that is, $M^{\mathfrak{U}} \ncong M_1 \otimes M_2$ for any diffuse factors M_1, M_2 .

Proof. By contradiction, assume that $M^{\mathfrak{U}}$ is not prime and write $M^{\mathfrak{U}}=M_1\otimes M_2$. For every $\mathfrak{i}=1,2$, choose a separable diffuse abelian von Neumann subalgebra $A_{\mathfrak{i}}\subset M_{\mathfrak{i}}$ with expectation. By Theorem A.5, there exists a unitary $w\in \mathcal{U}(M^{\mathfrak{U}})$ such that $wA_1w^*=A_2$.

For every i=1,2, choose a faithful normal state ϕ_i on M_i such that $A_i \subset (M_i)_{\phi_i}$ and define the faithful normal conditional expectation $E_2: M_1 \otimes M_2 \to M_2$ by $E_2 = \phi_1 \otimes id_{M_2}$. Choose a sequence of unitaries (\mathfrak{u}_n) in $\mathcal{U}(A_1)$ such that $\mathfrak{u}_n \to 0$ weakly. For all $i \in \{1,2\}$ and all $x_i, y_i \in M_i$, we have

$$\lim_n E_{A_2}((x_1\otimes x_2)(u_n\otimes 1)(y_1\otimes y_2))=\lim_n \phi_1(x_1u_ny_1)x_2y_2=0.$$

By strong density of linear combinations of elementary tensors in $M_1 \otimes M_2$ and since $u_n \in A_1 \subset (M_1)_{\phi_1}$, it follows that for all $x, y \in M_1 \otimes M_2 = M^{\mathfrak{U}}$, we have

$$\lim_n E_{A_2}(x(u_n \otimes 1)y) = 0.$$

Applying the above result to x = w and $y = w^*$ and since $wA_1w^* = A_2$, we obtain

$$1 = \lim_{n} \|w(u_n \otimes 1)w^*\|_{\phi_2} = \lim_{n} \|\mathsf{E}_{\mathsf{A}_2}(w(u_n \otimes 1)w^*)\|_{\phi_2} = 0.$$

This is a contradiction.

APPENDIX B. KEISLER'S SANDWICH THEOREM

In this section we prove Keisler's Sandwich Theorem (Theorem 4.2). To simplify the matter, we work in the traditional [0, 1]-valued version of continuous logic presented in [5]. We freely use the notation and terminology established in [5].

Fix a language L. Let L_{\exists} be the language obtained by adding a predicate P_{ϕ} for every existential L-formula ϕ . It is clear that an embedding between L_{\exists} -structures is an existential embedding of their L-reducts and, conversely, any existential embedding between L-structures is an embedding of their canonical expansions to L_{\exists} -structures.

Lemma B.1. Any restricted quantifier-free L_{\exists} -formula is equivalent to both an \forall_2 L-formula and a \exists_2 L-formula.

Proof. We prove the lemma by induction on the complexity of formulae. The main case is the connective $\dot{-}$. This follows from the following calculations:

•
$$(\sup_x \inf_y \varphi) \div (\inf_z \sup_w \psi) \equiv \sup_x \sup_z \inf_y \inf_w (\varphi \div \psi)$$
 and
• $(\inf_x \sup_y \varphi) \div (\sup_z \inf_w \psi) \equiv \inf_x \inf_z \sup_y \sup_w (\varphi \div \psi)$.

In what follows, given an L-structure N, $Th_{\forall_n}(N)$ denotes the closed conditions of the form $\sigma=0$, where σ is a \forall_n -sentence for which $\sigma^N=0$. Similarly, if T is a theory, we let T_{\forall_n} denote the collection of closed conditions $\sigma=0$, where σ is a \forall_n -sentence for which $T\models\sigma=0$. The corresponding notions with \forall_n replaced by \exists_n are defined analogously. It is routine to verify that $M\models Th_{\forall_n}(N)$ if and only if $N\models Th_{\exists_n}(M)$.

Theorem B.2. For L-structures M and N, the following are equivalent:

- (1) $M \models Th_{\forall_n}(N)$;.
- (2) There is an elementary extension N' of N for which there is an (M, N', n)-ultrapower sandwich.

Proof. We prove this by induction on \mathfrak{n} . The case $\mathfrak{n}=1$ is well-known.

Now suppose that n > 1. If S is an n-ultrapower sandwich, let S' denote the sandwich with the last element removed.

First suppose that there is an (M, N', n)-ultrapower sandwich S with N' an elementary extension of N. Then S is an (M, N', n-1)-ultrapower sandwich with respect to L_{\exists} . By induction, we have that M is a model of $Th_{\forall_{n-1}}(N)$ with respect to the language L_{\exists} . It is clear that $Th_{\forall_n}(N)$ with respect to L is contained in $Th_{\forall_{n-1}}(N)$ with respect to L_{\exists} , so $M \models Th_{\forall_n}(N)$, as desired.

Now suppose that $M \models Th_{\forall_n}(N)$. By the previous lemma, we have that $M \models Th_{\forall_{n-1}}(N)$ with respect to L_{\exists} . By induction, there is an elementary extension N' of N for which there is an (M, N', n-1)-ultrapower sandwich with respect to L_{\exists} , that is, the embeddings are all existential embeddings. In particular, the chain can be extended by one more element if the embeddings are not required to be existential. Thus, there is an (M, N', n)-ultrapower sandwich, as desired. \square

We can now prove the above promised sandwich theorems:

Corollary B.3. *For any* L*-theory* T*, the following are equivalent:*

(1) T is \forall_n -axiomatizable.

(2) Whenever there is an (M, N, n)-ultrapower sandwich with $N \models T$, we also have that $M \models T$.

Proof. First suppose that T is \forall_n -axiomatizable and consider an (M, N, n)-ultrapower sandwich with $N \models T$. Then by Theorem B.2, we have that $M \models Th_{\forall_n}(N)$. Since $N \models T$, we have that $T_{\forall_n} \subseteq Th_{\forall_n}(N)$. It follows that $M \models T_{\forall_n}$. Since T is \forall_n -axiomatizable, we have that $M \models T$, as desired.

Conversely, suppose that (2) holds and let $M \models T_{\forall_n}$. We wish to show that $M \models T$. Consider the set

$$\Sigma:=T\cup\{\sigma\geq\frac{\varepsilon}{2}\ :\ \sigma^M\geq\varepsilon,\ \sigma\text{ is a }\forall_n\text{-sentence}\}.$$

If Σ were unsatisfiable, then there would be σ_1,\ldots,σ_m and ε such that $\sigma_i^M \geq \varepsilon$ for all i and yet $T \models \max_{1 \leq i \leq m} (\sigma_i \ \dot{-} \frac{\varepsilon}{2}) = 0$. Since this latter sentence is still \forall_n , we have that it belongs to T_{\forall_n} , contradicting the fact that $M \models T_{\forall_n}$.

Let $N \models \Sigma$. Note then that $N \models T$. Moreover, $M \models Th_{\forall_n}(N)$. Indeed, if σ is a \forall_n sentence such that $\sigma^N = 0$, then $\sigma^M = 0$, else there is $\varepsilon > 0$ such that $\sigma^M \ge \varepsilon$, whence $\sigma^N \ge \frac{\varepsilon}{2}$, a contradiction.

By Theorem B.2, there is an elementary extension N' of N for which there is an (M, N', n)-ultrapower sandwich. By (2) and the fact that $N' \models T$, we have that $M \models T$, as desired.

Corollary B.4. *For any* L*-theory* T*, the following are equivalent:*

- (1) T is \exists_n -axiomatizable.
- (2) Whenever there is an (M, N, n)-ultrapower sandwich with $M \models T$, we also have that $N \models T$.

Proof. First suppose that T is \exists_n -axiomatizable and consider an (M, N, n)-ultrapower sandwich with $M \models T$. By Theorem B.2, we have that $M \models Th_{\forall_n}(N)$. Note then that $N \models Th_{\exists_n}(M)$. Indeed, if σ is an \exists_n -sentence and $\sigma^M = 0$, then $\sigma^N = 0$, else $\sigma^N \geq \varepsilon$. Since $\varepsilon \doteq \sigma$ is equivalent to a \forall_n -sentence, we get that $\varepsilon \doteq \sigma \in Th_{\forall_n}(N)$, whence $(\varepsilon \doteq \sigma)^M = 0$, contradicting that $\sigma^M = 0$. Since T is \exists_n -axiomatizable, we have that $N \models T$, as desired.

Now suppose that (2) holds and $M \models T_{\exists_n}$. We wish to show that $M \models T$. The exact same argument as in the previous theorem shows that there is $N \models T$ such that $\sigma^N \ge \varepsilon$ whenever σ is a \exists_n -sentence with $\sigma^M \ge \varepsilon$. It follows that $M \models Th_{\exists_n}(N)$. Arguing as above, we have that $N \models Th_{\forall_n}(M)$, so there is an (N, M', n)-ultrapower sandwich with M' an elementary extension of M. By (2) and the fact that $N \models T$, we have that $M' \models T$, whence $M \models T$, as desired. \square

REFERENCES

- [1] H. Ando, U. Haagerup, *Ultraproducts of von Neumann algebras*. J. Funct. Anal. **266** (2014), 6842–6913.
- [2] H. Ando, U. Haagerup, C. Houdayer, and A. Marrakchi, Structure of bicentralizer algebras and inclusions of type III factors. Math. Ann. **376** (2020), no. 3-4, 1145–1194.
- [3] H. Ando, U. Haagerup, C. Winsløow, *Ultraproducts, QWEP von Neumann algebras, and the Effros-Maréchal topology*. J. Reine Angew. Math. **715** (2016), 231–250.
- [4] S. Atkinson, I. Goldbring, S. Kunnawalkam Elayavalli, Factorial relative commutants and the generalized Jung property for II₁ factors. arXiv:2004.02293
- [5] I. Ben Yaacov, A. Berenstein, C. W. Henson, and A. Usvyatsov, *Model theory for metric structures*, Model theory with applications to algebra and analysis. Vol. 2, London Math. Soc. Lecture Note Ser., vol. 350, Cambridge Univ. Press, Cambridge, 2008, 315-427.
- [6] R. Boutonnet, I. Chifan, A. Ioana, II₁ factors with nonisomorphic ultrapowers. Duke Math. J. **166** (2017), 2023–2051.
- [7] A. Connes, *Une classification des facteurs de type* III. Ann. Sci. École Norm. Sup. 6 (1973), 133–252.
- [8] A. Connes, Almost periodic states and factors of type III₁. J. Funct. Anal. **16** (1974), 415–445.
- [9] A. Connes, On hyperfinite factors of type III_0 and Krieger's factors. J. Funct. Anal. 18 (1975), 318–327.
- [10] A. Connes, *Outer conjugacy classes of automorphisms of factors*. Ann. Sci. École Norm. Sup. **8** (1975), 383–419.
- [11] A. Connes, Classification of injective factors. Cases II_1 , II_{∞} , III_{λ} , $\lambda \neq 1$. Ann. of Math. 74 (1976), 73–115.
- [12] A. Connes, Factors of type III₁, property L'_{λ} , and closure of inner automorphisms. J. Operator Theory **14** (1985), 189–211.
- [13] Y. Dabrowski, Continuous model theories for von Neumann algebras, Journal of Functional Analysis 277 (2019), 108308.
- [14] J. Fang, L. Ge, W. Li, Central sequence algebras of von Neumann algebras. Taiwanese J. Math. **10** (2006), 187–200.
- [15] I. FARAH, I. GOLDBRING, B. HART, AND D. SHERMAN, Existentially closed II₁ factors, Fund. Math.233 (2016), 173–196.
- [16] I. Farah, B. Hart, Bradd, D. Sherman, Model theory of operator algebras I: stability. Bull. Lond. Math. Soc. 45 (2013), 825–838.
- [17] I. Farah, B. Hart, Bradd, D. Sherman, Model theory of operator algebras II: model theory. Israel Journal of Mathematics **201** (2014), 477–505.
- [18] I. Farah, B. Hart, Bradd, D. Sherman, Model theory of operator algebras III: elementary equivalence and II₁ factors. Bull. Lond. Math. Soc. **46** (2014), 609–628
- [19] I. Goldbring, *Spectral gap and definability*. To appear in the book Beyond First Order Model Theory Volume 2.
- [20] I. Goldbring, *On Popa's factorial commutant embedding problem*, Proc. of the American Math Soc. **148** (2020), 5007–5012.
- [21] I. Goldbring, *Enforceable operator algebras*, Journal of the Institute of Mathematics of Jussieu **20** (2021), 31–63.
- [22] I. Goldbring and B. Hart, The universal theory of the hyperfinite II_1 factor is not computable, arXiv 2006.05629.

- [23] I. Goldbring, B. Hart, and T. Sinclair, *The theory of tracial von Neumann algebras does not have a model companion*, Journal of Symbolic Logic **78** (2013), 1000–1004.
- [24] I. Goldbring, B. Hart, and T. Sinclair, *Correspondences, ultraproducts, and model theory*, arXiv 1809.0049.
- [25] U. Haagerup, The standard form of von Neumann algebras. Math. Scand. 37 (1975), 271–283.
- [26] U. Haagerup, Connes' bicentralizer problem and uniqueness of the injective factor of type III₁. Acta. Math. **69** (1986), 95–148.
- [27] W. Hodges, Building models by games, London Math Soc. Student Texts 2 (1985), Cambridge University Press.
- [28] C. Houdayer, *Gamma stability in free product von Neumann algebras*. Commun. Math. Phys. **336** (2015), 831–851.
- [29] C. Houdayer, Y. Isono, Free independence in ultraproduct von Neumann algebras and applications. J. London Math. Soc. **92** (2015), 163–177.
- [30] C. Houdayer, Y. Isono, Bi-exact groups, strongly ergodic actions and group measure space type III factors with no central sequence. Comm. Math. Phys. **348** (2016), 991–1015.
- [31] C. Houdayer, A. Marrakchi, P. Verraedt, Fullness and Connes' τ invariant of type III tensor product factors. J. Math. Pures Appl. **121** (2019), 113–134.
- [32] Z. Ji, A. Natarajan, T. Vidick, J. Wright and H. Yuen, $MIP^* = RE$, arxiv 2001.04383
- [33] H. J. Keisler, *Theory of models with generalized atomic formulas*, Journal of Symbolic Logic **25** (1960), 1–26.
- [34] E. Kirchberg, On non-semisplit extensions, tensor products and exactness of group C*-algebras, Inventiones Mathematicae **112** (1993), 449–489.
- [35] A. Marrakchi, Full factors, bicentralizer flow and approximately inner automorphisms. Invent. Math. **222** (2020), 375–398.
- [36] D. McDuff, *Uncountably many* II₁ *factors*. Ann. of Math. **90** (1969), 372–377.
- [37] A. Ocneanu, *Actions of discrete amenable groups on von Neumann algebras*. Lecture Notes in Mathematics, **1138**. Springer-Verlag, Berlin, 1985. iv+115 pp.
- [38] N. Ozawa, About the QWEP conjecture. Internat. J. Math. 15 (2004), 501–530.
- [39] S. Popa, Orthogonal pairs of *-subagebras in finite von Neumann algebras. J. Operator Theory **9** (1983), 253–268.
- [40] S. Popa, On a class of type II_1 factors with Betti numbers invariants. Ann. of Math. **163** (2006), 809–899.
- [41] Y. Suzuki, Complete descriptions of intermediate operator algebras by intermediate extensions of dynamical systems. Comm. Math. Phys. **375** (2020), 1273–1297.
- [42] M. Takesaki, *Theory of operator algebras*. II. Encyclopaedia of Mathematical Sciences, **125**. Operator Algebras and Non-commutative Geometry, 6. Springer-Verlag, Berlin, 2003. xxii+518 pp.
- [43] M. Takesaki, *Theory of operator algebras*. III. Encyclopaedia of Mathematical Sciences, **127**. Operator Algebras and Non-commutative Geometry, 8. Springer-Verlag, Berlin, 2003. xxii+548 pp.
- [44] Y. Ueda, Factoriality, type classification and fullness for free product von Neumann algebras. Adv. Math. **228** (2011), 2647–2671.
- [45] Y. Ueda, On type III_1 factors arising as free products. Math. Res. Lett. **18** (2011), 909–920.
- [46] Y. Ueda, Some analysis on amalgamated free products of von Neumann algebras in non-tracial setting. J. London Math. Soc. 88 (2013), 25–48.

Department of Mathematics, University of California, Irvine, 340 Rowland Hall (Bldg.# 400), Irvine, CA 92697-3875

Email address: isaac@math.uci.edu

 URL : http://www.math.uci.edu/~isaac

Université Paris-Saclay, Institut Universitaire de France, CNRS, Laboratoire de mathématiques d'Orsay, 91405 Orsay, FRANCE

 ${\it Email address}: \verb|cyril.houdayer@universite-paris-saclay.fr|\\$

 URL : https://cyrilhoudayer.com