

EQUILIBRIUM ON TOEPLITZ EXTENSIONS OF HIGHER DIMENSIONAL NONCOMMUTATIVE TORI

ZAHRA AFSAR, MARCELO LACA, JACQUI RAMAGGE, AND CAMILA F. SEHNEM

ABSTRACT. The C^* -algebra generated by the left-regular representation of \mathbb{N}^n twisted by a 2-cocycle is a Toeplitz extension of an n -dimensional noncommutative torus, on which each vector $r \in [0, \infty)^n$ determines a one-parameter subgroup of the gauge action. We show that the equilibrium states of the resulting C^* -dynamical system are parametrised by tracial states of the noncommutative torus corresponding to the restriction of the cocycle to the vanishing coordinates of r . These in turn correspond to probability measures on a classical torus whose dimension depends on a certain degeneracy index of the restricted cocycle. Our results generalise the phase transition on the Toeplitz noncommutative tori used as building blocks in recent work of Brownlowe, Hawkins and Sims, and of Afsar, an Huef, Raeburn and Sims.

1. INTRODUCTION

Suppose $\Theta = (\theta_{i,j})$ is an $n \times n$ antisymmetric matrix with real coefficients. The n -dimensional noncommutative torus \mathcal{A}_Θ is the universal C^* -algebra generated by unitaries U_1, \dots, U_n satisfying

$$U_j U_k = e^{-2\pi i \theta_{j,k}} U_k U_j \quad j, k = 1, 2, \dots, n.$$

The matrix Θ determines a circle-valued 2-cocycle $\sigma_\Theta: \mathbb{Z}^n \times \mathbb{Z}^n \rightarrow \mathbb{T}$ on \mathbb{Z}^n given by

$$\sigma_\Theta(x, y) := e^{-\pi i \langle x | \Theta y \rangle}, \quad (1.1)$$

and the noncommutative torus \mathcal{A}_Θ is canonically isomorphic to the twisted group C^* -algebra $C^*(\mathbb{Z}^n, \sigma_\Theta)$, see [28, 30]. That is, \mathcal{A}_Θ is also the universal C^* -algebra for unitary σ_Θ -representations of \mathbb{Z}^n . In particular, a concrete realisation of \mathcal{A}_Θ as a C^* -algebra of operators on a Hilbert space is provided by the usual left regular unitary σ_Θ -representation λ^{σ_Θ} of \mathbb{Z}^n on $\ell^2(\mathbb{Z}^n)$. This is defined on a canonical orthonormal basis vector by $\lambda_x^{\sigma_\Theta} \delta_y = e^{-\pi i \langle x | \Theta y \rangle} \delta_{x+y}$ for $x, y \in \mathbb{Z}^n$. The universal property defining \mathcal{A}_Θ holds for λ^{σ_Θ} because \mathbb{Z}^n is abelian, hence amenable. See, for example, [6] for the theory of twisted group algebras.

In a recent paper, Latrèmollière and Packer [23] considered noncommutative solenoids, which can be described as inductive limits of 2-dimensional noncommutative tori, in analogy with their commutative counterparts. Subsequently, Brownlowe, Hawkins and Sims [5] defined Toeplitz extensions of the noncommutative solenoids of [23] and studied their phase transitions under natural dynamics. The solenoidal extensions of [5] are constructed as direct limits of building blocks, each of which is a Toeplitz-type extension of a noncommutative 2-torus. These Toeplitz-type extensions are obtained by replacing one of the two canonical unitary generators in the presentation of a noncommutative 2-torus by an isometry.

Prompted by the constructions given in [5, 23], Afsar, an Huef, Raeburn and Sims [2] defined higher-rank versions of Toeplitz noncommutative solenoids, using Toeplitz-type extensions of

Date: 24 July 2021.

Key words and phrases. KMS state; Toeplitz noncommutative torus; twisted semigroup C^* -algebra.

M. Laca was supported by the Natural Sciences and Engineering Research Council of Canada, Discovery Grant RGPIN-2017-04052.

C. F. Sehnem was supported by the Marsden Fund of the Royal Society of New Zealand, grant No.18-VUW-056. This research was supported by an Australian Research Council Discovery Grant.

higher-dimensional noncommutative tori as building blocks. As in [5], each of these building blocks is obtained by replacing one of the two generating unitaries of a noncommutative 2-torus by a Nica-covariant isometric representation V of \mathbb{N}^k , and the other by a unitary representation U of \mathbb{Z}^d ; the commutation relations satisfied by U and V are then encoded in a $k \times d$ real matrix θ , see [2, Equation (2.1)]. The phase transition computed in [2] is parametrised by measures on the classical torus \mathbb{T}^d , which is the spectrum of the subalgebra $C^*(U_x \mid x \in \mathbb{Z}^d)$.

The present investigation was sparked by a question raised by Ian Putnam about the possibility of replacing the classical torus $C^*(U_x \mid x \in \mathbb{Z}^d)$ by a noncommutative one. Instead of viewing a higher-rank noncommutative torus as a C^* -algebra generated by two unitary representations, one of \mathbb{Z}^k and one of \mathbb{Z}^d , satisfying a commutation relation, we consider all the $n = k + d$ generators on the same footing. Thus, we regard an n -dimensional noncommutative torus as the twisted group algebra $C^*(\mathbb{Z}^n, \sigma)$ in which the twist is given by a circle-valued 2-cocycle σ . Since the circle-valued second cohomology of \mathbb{N}^n is the same as that of \mathbb{Z}^n by [19, Corollary 2.3], it seems natural to take the *reduced twisted semigroup C^* -algebra* $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ as the appropriate Toeplitz extension of the noncommutative torus associated to an extension of σ to \mathbb{Z}^n . The Toeplitz-type extensions of noncommutative tori from [2] can then be obtained as quotients of special cases of ours.

Our purpose here is two-fold. Firstly we aim to describe various realisations of our Toeplitz noncommutative tori, using product systems, semigroup crossed products, and generators and relations. Secondly, we aim to compute their phase transitions of KMS_β states with respect to one-parameter subgroups of the canonical dual action of \mathbb{T}^n . Along the way we also provide an explicit description of the center and the space of traces of higher-rank noncommutative tori. We do not address here the question of how to use our Toeplitz noncommutative tori as building blocks for generalisations of the higher-rank noncommutative solenoids defined in [2], leaving the issue for further work.

We begin in Section 2 with a brief discussion of 2-cocycles on \mathbb{N}^n and we introduce their twisted semigroup algebras using as a model the left regular representation twisted by a cocycle. Then we show that these twisted semigroup algebras have a universal property with respect to twisted isometric representations that satisfy certain $*$ -commuting relations, which we show to be equivalent to Nica-covariance. They can also be characterised as Nica–Toeplitz C^* -algebras of product systems over \mathbb{N}^n and as twisted semigroup crossed products. We then use the cocycle extension results of [19] to show that these twisted semigroup algebras are indeed extensions of rank- n noncommutative tori.

As with every discrete abelian group, the circle-valued second cohomology of \mathbb{Z}^n can be described in terms of symplectic bicharacters, which in this case are parametrised by antisymmetric matrices over \mathbb{R} ; with our conventions these matrices are determined up to even integers. So we assume that $\sigma = \sigma_\Theta$ comes from an antisymmetric matrix Θ as in (1.1), which allows us to give a presentation of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ in terms of generators and relations that are entirely analogous to those in the original presentation of rotation algebras and, more generally, of higher-rank noncommutative tori. Using this presentation we show in Proposition 2.15 that the building blocks from [2] are quotients of particular cases of our Toeplitz extensions of noncommutative tori, corresponding to antisymmetric matrices that have two diagonal blocks of zeros and such that the generating isometries corresponding to the second block are unitaries.

In Section 3 we consider the dynamics induced on $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ by a vector $r \in \mathbb{R}^n$ with k positive coordinates and $d = n - k$ vanishing coordinates. In order to compute the KMS_β states, we follow the strategy developed in [21] for the Bost–Connes system, showing that KMS_β states are parametrised by tracial states of a corner. The projection Q that defines the corner is the common orthogonal complement of the range projections of the generating isometries that are not fixed by the dynamics, that is, those corresponding to the nonvanishing coordinates of the vector r .

In Section 4 we show that the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ is actually isomorphic to the Toeplitz noncommutative torus associated to the $d \times d$ submatrix Θ_d corresponding to the vanishing coordinates of the vector r . One of our main technical results is Proposition 4.2, where we give a formula for KMS_β states in terms of tracial states of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ and a finite Euler product that depends on the cocycle. The tracial states of this rank- d Toeplitz noncommutative torus factor through the associated noncommutative torus. This led us to compute the tracial states of the higher-rank noncommutative torus \mathcal{A}_D associated to a $d \times d$ antisymmetric real matrix D . We show that these tracial states come from states of the center $Z(\mathcal{A}_D)$, which is a classical torus of dimension equal to a certain degeneracy index of D .

Our main results are in Section 5, starting with Theorem 5.1, where we give a parametrisation of the simplex of KMS_β states in terms of probability measures on a classical torus. As a corollary, for the special case of antisymmetric matrix with zero diagonal blocks, we recover the phase transition for the building blocks of [2].

In Section 6 we compute the equilibrium states at $\beta = \infty$ and $\beta = 0$. In addition to the usual notions of ground states and invariant traces, we also study the limiting cases, namely $\beta \rightarrow \infty$, which gives the KMS_∞ states, and the limits of KMS_β states as $\beta \rightarrow 0^+$, a class that we call KMS_{0^+} states. We show that in general not every invariant trace is such a KMS_{0^+} state.

Finally, in Section 7 we realise the Toeplitz noncommutative tori $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ as the Nica-Toeplitz algebra of a product system over \mathbb{N}^k in which the coefficient algebra is the Toeplitz noncommutative torus associated to the restriction of σ_Θ to \mathbb{N}^d . This allows us to compare our results to those obtained from the characterisation of KMS_β states given in [3].

Acknowledgments: This research was started during a visit of Z. Afsar, J. Ramagge and C. F. Sehnem to the University of Victoria and continued through visits of C. F. Sehnem and M. Laca to the University of Sydney. We would like to acknowledge the support of these institutions and to thank them for the hospitality.

2. TOEPLITZ EXTENSIONS OF NONCOMMUTATIVE TORI

We wish to consider the natural Toeplitz-type extensions of noncommutative tori. Since there are some nuances associated with passing from unitaries to isometries, it is more straightforward to introduce these extensions spatially, via concrete projective representations of \mathbb{N}^n . We begin with a very brief introduction to projective isometric representations of semigroups and twisted semigroup C^* -algebras.

2.1. Projective isometric representations.

Definition 2.1. Suppose P is a subsemigroup of a group G . A circle-valued semigroup 2-cocycle is a function $\sigma: P \times P \rightarrow \mathbb{T}$ such that

$$\sigma(p, q)\sigma(pq, r) = \sigma(p, qr)\sigma(q, r), \quad p, q, r \in P.$$

The set of all \mathbb{T} -valued 2-cocycles on P is a group under pointwise multiplication, which we denote by $Z^2(P, \mathbb{T})$.

Definition 2.2. Let $\sigma \in Z^2(P, \mathbb{T})$ be a 2-cocycle on P . An *isometric σ -representation* of P is a map $p \mapsto V_p$ into the semigroup of isometries on a Hilbert space \mathcal{H} such that

$$V_p V_q = \sigma(p, q) V_{p+q}, \quad p, q \in P.$$

The isometric σ -representations are multiplicative only up to scalars. If P is unital, then V_e is a multiple of the identity on \mathcal{H} , where e denotes the unit element of P . When σ is normalised, that is, $\sigma(e, e) = 1$, then $V_e = 1_{\mathcal{H}}$. In this case $\sigma(p, e) = \sigma(e, p) = 1$ for all $p \in P$.

One can generate an example of a 2-cocycle by starting with an arbitrary function $\lambda: P \rightarrow \mathbb{T}$ and defining $\sigma_\lambda(p, q) := \lambda(pq)\lambda(p)^{-1}\lambda(q)^{-1}$, for all $p, q \in P$. This is called a *coboundary*, and

is trivial in the sense that any σ_λ -representation can be transformed into a true isometric representation through multiplication by the scalar-valued function λ . We denote by $B^2(P, \mathbb{T})$ the subgroup of all coboundaries. A standard argument shows that the representations associated to two 2-cocycles that differ by a coboundary are equivalent, in the sense that one is a multiple of the other by a circle-valued function. So what is interesting is the group

$$H^2(P, \mathbb{T}) := Z^2(P, \mathbb{T})/B^2(P, \mathbb{T})$$

of cocycles modulo coboundaries, which is called the *second cohomology group* of P .

Next we verify that for each σ in $Z^2(P, \mathbb{T})$, there exists an isometric σ -representation of P .

Proposition 2.3. *Suppose σ is a circle-valued 2-cocycle on P and let $\{\delta_q \mid q \in P\}$ be the canonical orthonormal basis of $\ell^2(P)$. Then for each $p \in P$, the map L_p^σ defined by*

$$L_p^\sigma \delta_q := \sigma(p, q) \delta_{pq}, \quad (q \in P)$$

extends uniquely to an isometry on $\ell^2(P)$ such that

$$(L_p^\sigma)^* \delta_q = \begin{cases} \overline{\sigma(p, q)} \delta_{p^{-1}q} & \text{if } p \leq q, \\ 0 & \text{otherwise.} \end{cases} \quad (2.1)$$

Moreover, we have for all $p, q \in P$

$$L_p^\sigma L_q^\sigma = \sigma(p, q) L_{pq}^\sigma, \quad \text{and} \quad L_p^\sigma (L_p^\sigma)^* = 1_{pP},$$

where 1_{pP} is the multiplication operator by the characteristic function of the set $pP = \{pq \mid q \in P\}$.

Proof. Since P is a cancellative semigroup, L_p^σ maps the elements of the standard orthonormal basis one to one and onto those of $\{\delta_{pq} \mid q \in P\}$, multiplied by complex numbers of modulus 1. So L_p^σ extends uniquely by linearity and continuity to an isometric operator (also denoted L_p^σ) on $\ell^2(P)$ with range $\ell^2(pP)$. The second assertion follows because the formula given in the right-hand side of (2.1) defines a left inverse for L_p^σ on orthonormal basis vectors since $\sigma(p, q) \overline{\sigma(p, q)} = 1$.

To prove the last claim, we compute first

$$L_p^\sigma L_q^\sigma \delta_r = L_p^\sigma \sigma(q, r) \delta_{qr} = \sigma(p, qr) \sigma(q, r) \delta_{pqr},$$

and then

$$L_{pq}^\sigma \delta_r = \sigma(pq, r) \delta_{pqr}.$$

So $\sigma(p, q) L_{pq}^\sigma \delta_r = \sigma(p, q) \sigma(pq, r) \delta_{pqr} = L_p^\sigma L_q^\sigma \delta_r$ by the cocycle identity. \square

Definition 2.4. Let σ be a circle-valued 2-cocycle on P . The *left regular σ -representation* of P is the map $p \mapsto L_p^\sigma \in \ell^2(P)$. The *reduced semigroup C^* -algebra twisted by σ* is the C^* -subalgebra of $\mathcal{B}(\ell^2(P))$ generated by the operators L_p^σ with $p \in P$; it will be denoted by $\mathcal{T}_r(P, \sigma)$.

When the cocycle is understood and there is no risk of confusion, we simply refer to $\mathcal{T}_r(P, \sigma)$ as a (reduced) twisted semigroup C^* -algebra.

2.2. Toeplitz extensions of higher dimensional noncommutative tori. By analogy with the realisation of an n -dimensional noncommutative torus as a twisted group algebra of \mathbb{Z}^n , we define the Toeplitz noncommutative torus as a twisted semigroup C^* -algebra associated to a 2-cocycle on the semigroup \mathbb{N}^n . We will see below in Proposition 2.12 that this is indeed a natural extension of the corresponding noncommutative torus.

Definition 2.5. Let σ be a circle-valued 2-cocycle on \mathbb{N}^n . The associated n -dimensional *Toeplitz noncommutative torus* is the twisted semigroup C^* -algebra $\mathcal{T}_r(\mathbb{N}^n, \sigma)$.

The Toeplitz noncommutative torus $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ also has a universal property with respect to a certain class of twisted isometric representations. We define this next by adding extra relations modelled on the left regular σ -representation.

Definition 2.6. A covariant isometric σ -representation of \mathbb{N}^n on a Hilbert space \mathcal{H} is a map $w: \mathbb{N}^n \rightarrow \mathcal{B}(\mathcal{H})$ such that

- (1) $w_p^* w_p = 1$;
- (2) $w_p w_q = \sigma(p, q) w_{p+q}$;
- (3) $w_p^* w_q = \overline{\sigma(p, -p + (p \vee q))} \sigma(q, -q + (p \vee q)) w_{-p+(p \vee q)} w_{-q+(p \vee q)}^*$.

The third condition above may seem a bit unusual at first but it will be convenient in some computations. We show next that it is equivalent to the better-known Nica-covariance condition from [24, Section 3.3].

Lemma 2.7. An isometric σ -representation w of \mathbb{N}^n satisfies condition (3) of Definition 2.6 if and only if it is Nica-covariant, that is,

$$w_p w_p^* w_q w_q^* = w_{p \vee q} w_{p \vee q}^* \quad p, q \in \mathbb{N}^n.$$

In particular, the left regular σ -representation of \mathbb{N}^n is covariant in the sense of Definition 2.6.

Proof. Assume first (3) holds. Then

$$w_p w_p^* w_q w_q^* = \overline{\sigma(p, -p + (p \vee q))} \sigma(q, -q + (p \vee q)) w_p w_{-p+(p \vee q)} (w_{-q+(p \vee q)})^* w_q^* = w_{p \vee q} w_{p \vee q}^*,$$

where we used the cocycle to combine factors in the last equality. Conversely, assume that w is Nica covariant. Then

$$\begin{aligned} w_p^* w_q &= w_p^* w_p w_p^* w_q w_q^* w_q = w_p^* (w_{p \vee q} w_{p \vee q}^*) w_q \\ &= \overline{\sigma(p, -p + (p \vee q))} \sigma(q, -q + (p \vee q)) w_p^* (w_p w_{-p+(p \vee q)} w_{-q+(p \vee q)}^* w_q^*) w_q \\ &= \overline{\sigma(p, -p + (p \vee q))} \sigma(q, -q + (p \vee q)) w_{-p+(p \vee q)} w_{-q+(p \vee q)}^* \end{aligned}$$

Since $L_p^\sigma (L_p^\sigma)^*$ is the operator of multiplication by the characteristic function of $p + \mathbb{N}^n$, the left regular σ -representation L^σ satisfies Nica-covariance, hence also condition (3). \square

The following immediate consequence of Definition 2.6 will be needed in the following sections.

Lemma 2.8. Suppose $w: \mathbb{N}^n \rightarrow \mathcal{B}(\mathcal{H})$ is a covariant isometric σ -representation of \mathbb{N}^n , and suppose $x, y \in \mathbb{N}^n$ satisfy $x \vee y = x + y$. Then

$$w_x w_x^* w_y = w_y w_x w_x^*.$$

Proof. Using first property (3) and then twice property (2) of Definition 2.6, we obtain

$$\begin{aligned} w_x w_x^* w_y &= \overline{\sigma(x, y)} \sigma(y, x) w_x w_y w_x^* = \overline{\sigma(x, y)} \sigma(y, x) \sigma(x, y) w_{x+y} w_x^* \\ &= \overline{\sigma(x, y)} \sigma(y, x) \sigma(x, y) \overline{\sigma(y, x)} w_y w_x w_x^*. \end{aligned}$$

This completes the proof of the lemma because $\overline{\sigma(x, y)} \sigma(y, x) \sigma(x, y) \overline{\sigma(y, x)} = 1$. \square

It will be useful to have a description of $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ as the Nica-Toeplitz algebra of a compactly aligned product system over \mathbb{N}^n (see [14, Theorem 6.3]). This will allow us to apply results of [13] and to relate our notion of a Toeplitz noncommutative torus to the one considered by Afsar, an Huef, Raeburn and Sims [2]. In order to simplify the notation, we assume from now on that σ is normalized, that is, $\sigma(0, 0) = 1$. For each $p \in \mathbb{N}^n$, let $\mathcal{E}_p := \mathbb{C}$ as a complex vector space and we write $\lambda \delta_p$ for an element $\lambda \in \mathcal{E}_p$. We equip \mathcal{E}_p with the canonical structure of a correspondence over \mathbb{C} , that is, the left and right actions are given by multiplication in \mathbb{C} and the right inner product is simply $\langle \lambda \delta_p | \alpha \delta_p \rangle = \bar{\lambda} \alpha$ for all $\lambda, \alpha \in \mathcal{E}_p$. We will use the cocycle σ to define the multiplication maps of the product system $\mathcal{E}^\sigma = (\mathcal{E}_p)_{p \in \mathbb{N}^n}$. For $p, q \in \mathbb{N}^n$, we set

$$\begin{aligned} \mu_{p,q} &: \mathcal{E}_p \times \mathcal{E}_q \rightarrow \mathcal{E}_{p+q} \\ (\lambda \delta_p, \alpha \delta_q) &\mapsto \sigma(p, q) \lambda \alpha \delta_{p+q}. \end{aligned}$$

These maps are associative because σ is a cocycle. So \mathcal{E}^σ is a product system over \mathbb{N}^n , which is compactly aligned because $\mathbb{K}(\mathcal{E}_p) = \mathbb{C}$ for all $p \in \mathbb{N}^n$.

Proposition 2.9. *Let $\mathcal{NT}_{\mathcal{E}^\sigma}$ be the Nica–Toeplitz algebra of the product system $\mathcal{E}^\sigma = (\mathcal{E}_p)_{p \in \mathbb{N}^n}$ as above. The map which sends $\delta_p \in \mathcal{E}_p$ to the isometry L_p^σ in the left regular σ -representation induces an isomorphism $\mathcal{NT}_{\mathcal{E}^\sigma} \cong \mathcal{T}_r(\mathbb{N}^n, \sigma)$.*

Proof. Let ϕ_p denote the map $\lambda\delta_p \mapsto \lambda L_p^\sigma$. For $p, q \in \mathbb{N}^n$, we have

$$\phi_p(\delta_p)\phi_q(\delta_q) = L_p^\sigma L_q^\sigma = \sigma(p, q)L_{p+q}^\sigma = \phi_{p+q}(\delta_p \cdot \delta_q).$$

Hence $\phi = \{\phi_p\}_{p \in P}$ preserves the multiplication on \mathcal{E}^σ . It is also compatible with inner products because each L_p^σ is an isometry. Notice next that the homomorphism $\phi^{(p)}: \mathbb{K}(\mathcal{E}_p) \rightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma)$ induced by ϕ is given by $\lambda 1_{\mathcal{E}_p} \in \mathbb{K}(\mathcal{E}_p) \mapsto \lambda L_p^\sigma (L_p^\sigma)^*$. Hence Lemma 2.7 gives

$$\phi^{(p)}(1_{\mathcal{E}_p})\phi^{(q)}(1_{\mathcal{E}_q}) = L_p^\sigma (L_p^\sigma)^* L_q^\sigma (L_q^\sigma)^* = L_{p \vee q}^\sigma (L_{p \vee q}^\sigma)^* = \phi^{p \vee q}(1_{\mathcal{E}_{p \vee q}}).$$

This proves that ϕ is Nica covariant and therefore it induces a surjective homomorphism $\mathcal{NT}_{\mathcal{E}^\sigma} \rightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma)$, which we still denote by ϕ . To see that ϕ is an isomorphism, notice that

$$\prod_{i=1}^k (1_{\ell^2(\mathbb{N}^n)} - L_{p_i}^\sigma (L_{p_i}^\sigma)^*) \delta_0 = \delta_0$$

whenever $p_1, p_2, \dots, p_k \in \mathbb{N}^n \setminus \{0\}$. Because \mathbb{N}^n is abelian, we can apply [13, Theorem 5.1] (see also [13, Example 5.6](1)) to deduce that ϕ is injective. \square

Corollary 2.10. *Let σ be a 2-cocycle on \mathbb{N}^n and let $w: \mathbb{N}^n \rightarrow B$ be a covariant isometric σ -representation in a C^* -algebra B . Then there is a homomorphism $\rho: \mathcal{T}_r(\mathbb{N}^n, \sigma) \rightarrow B$ that sends the isometry L_p^σ to w_p for all $p \in \mathbb{N}^n$. Moreover, ρ is faithful if and only if*

$$\prod_{i=1}^n (1_{\mathcal{H}} - w_{e_i} w_{e_i}^*) \neq 0. \quad (2.2)$$

As a consequence, relations (1)–(3) from Definition 2.6 form a presentation of $\mathcal{T}_r(\mathbb{N}^n, \sigma)$.

Proof. By Lemma 2.7, an isometric σ -representation of \mathbb{N}^n satisfies relation (3) of Definition 2.6 if and only if it is Nica covariant. Hence the first assertion in the statement follows from an application of the universal property of $\mathcal{NT}_{\mathcal{E}^\sigma}$ combined with the isomorphism $\mathcal{T}_r(\mathbb{N}^n, \sigma) \cong \mathcal{NT}_{\mathcal{E}^\sigma}$ from Proposition 2.9.

Since \mathbb{N}^n is abelian, it follows from [13, Theorem 5.1] that a representation of $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ is faithful if and only if

$$\prod_{i=1}^k (1 - w_{p_i} w_{p_i}^*) \neq 0 \quad \text{whenever } p_1, p_2, \dots, p_k \in \mathbb{N}^n \setminus \{0\}. \quad (2.3)$$

Now every projection of the form $1 - w_p w_p^*$ with $p \neq 0$ dominates a projection $1 - w_{e_i} w_{e_i}^*$ for some $i \in \{1, \dots, n\}$, and so (2.2) and (2.3) are equivalent. This establishes the equivalence in the second statement. The last assertion then follows because the left regular σ -representation of \mathbb{N}^n in $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ satisfies relations (1)–(3) of Definition 2.6. \square

The C^* -algebra $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ can also be realised as a twisted semigroup crossed product. Following [20] we consider the characteristic function $\mathbb{1}_{p+\mathbb{N}^n}$ of the cone with vertex $p \in \mathbb{N}^n$. We define $B_{\mathbb{N}^n}$ to be the C^* -subalgebra of $\ell^\infty(\mathbb{N}^n)$ generated by all the projections $\mathbb{1}_{p+\mathbb{N}^n}$ for $p \in \mathbb{N}^n$. There is a canonical action $\beta = \{\beta_p \mid p \in \mathbb{N}^n\}$ of \mathbb{N}^n on $B_{\mathbb{N}^n}$ by injective endomorphisms with hereditary range. Precisely, the endomorphism β_p is defined on a generator by

$$\beta_p(\mathbb{1}_{q+\mathbb{N}^n}) := \mathbb{1}_{p+q+\mathbb{N}^n}.$$

Proposition 2.11. *Let $(B_{\mathbb{N}^n}, \mathbb{N}^n, \beta)$ be as above and let σ be a 2-cocycle on \mathbb{N}^n . Then $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ is isomorphic to the twisted crossed product $B_{\mathbb{N}^n} \rtimes_{\beta, \sigma} \mathbb{N}^n$ via an isomorphism that identifies the canonical generating isometries.*

Proof. The result follows from [13, Theorem 4.3] since the product system \mathcal{E}^σ has one-dimensional fibres. \square

In analogy to what happens with the semigroup C^* -algebra $\mathcal{T}_\lambda(\mathbb{N}^n)$ (see, for example, [24]), there is a canonical *gauge action* γ of \mathbb{T}^n on $\mathcal{T}_r(\mathbb{N}^n, \sigma)$. It is implemented spatially by the unitaries $\{U_z : z \in \mathbb{T}^n\}$ defined by $U_z \delta_p = z^p \delta_p$ for $p \in \mathbb{N}^n$, where $z^p = \prod_{i=1}^n z_i^{p_i}$. The gauge action $z \mapsto \gamma_z := \text{Ad}_{U_z}$ can also be obtained through the relations (1)–(3) by stating that γ_z is the automorphism that sends an isometry L_p^σ to $z^p L_p^\sigma$. This action gives rise to a faithful conditional expectation

$$E: \mathcal{T}_r(\mathbb{N}^n, \sigma) \rightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma)^\gamma = \overline{\text{span}}\{L_p^\sigma (L_p^\sigma)^* \mid p \in \mathbb{N}^n\}$$

given by $E(b) = \int_{\mathbb{T}^n} \gamma_\lambda(b) d\lambda$ and $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ is the closed linear span of the set $\{L_p^\sigma (L_q^\sigma)^* : p, q \in \mathbb{N}^n\}$.

The Toeplitz noncommutative torus $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ has a canonical n -dimensional noncommutative torus as a quotient. We show this in the next result, where we also identify the canonical generating set for the kernel of the quotient map.

Proposition 2.12. *Every 2-cocycle $\sigma \in Z^2(\mathbb{N}^n, \mathbb{T})$ has an extension to a 2-cocycle $\tilde{\sigma}$ on \mathbb{Z}^n , and this extension is unique up to coboundaries. Moreover, the map that sends an isometry $L_p^\sigma \in \mathcal{T}_r(\mathbb{N}^n, \sigma)$ to the unitary $\lambda_p^{\tilde{\sigma}} \in C_r^*(\mathbb{Z}^n, \tilde{\sigma})$ determines an exact sequence*

$$0 \longrightarrow \mathcal{I} \longrightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma) \longrightarrow C_r^*(\mathbb{Z}^n, \tilde{\sigma}) \longrightarrow 0,$$

where \mathcal{I} is the ideal of $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ generated by the projections

$$\{1 - L_{e_j}^\sigma (L_{e_j}^\sigma)^* \mid j = 1, \dots, n\}.$$

Proof. Since \mathbb{N}^n is abelian and $\mathbb{Z}^n = \mathbb{N}^n - \mathbb{N}^n$ is the group of right (and left) quotients of \mathbb{N}^n , 2-cocycles on \mathbb{N}^n extend to 2-cocycles on \mathbb{Z}^n , uniquely up to coboundaries, by [19, Corollary 2.3]. More precisely, the restriction of 2-cocycles from \mathbb{Z}^n to \mathbb{N}^n induces an isomorphism of the second cohomology group $H^2(\mathbb{Z}^n, \mathbb{T})$ onto $H^2(\mathbb{N}^n, \mathbb{T})$.

Let σ be a 2-cocycle on \mathbb{N}^n and let $\tilde{\sigma}$ be a 2-cocycle on \mathbb{Z}^n extending σ . It follows that the map that sends $p \in \mathbb{N}^n$ to the unitary $\lambda_p^{\tilde{\sigma}} \in C_r^*(\mathbb{Z}^n, \tilde{\sigma})$ is a covariant isometric σ -representation of \mathbb{N}^n . This gives a homomorphism $\phi: \mathcal{T}_r(\mathbb{N}^n, \sigma) \rightarrow C_r^*(\mathbb{Z}^n, \tilde{\sigma})$ mapping L_p^σ to $\lambda_p^{\tilde{\sigma}}$, which is surjective because the unitaries $\lambda_p^{\tilde{\sigma}}$ for $p \in \mathbb{N}^n$ generate $C_r^*(\mathbb{Z}^n, \tilde{\sigma})$ as a C^* -algebra.

Let \mathcal{I} be the ideal of $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ generated by the projections $1 - L_{e_j}^\sigma (L_{e_j}^\sigma)^*$ for $j = 1, \dots, n$. Then ϕ vanishes on \mathcal{I} since it vanishes on the projection $1 - L_{e_j}^\sigma (L_{e_j}^\sigma)^*$ for each $j \in \{1, \dots, n\}$. Hence it factors through a homomorphism

$$\dot{\phi}: \mathcal{T}_r(\mathbb{N}^n, \sigma)/\mathcal{I} \rightarrow C_r^*(\mathbb{Z}^n, \tilde{\sigma}).$$

So it remains to show that $\dot{\phi}$ is injective. To do so, we invoke the description of $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ as the Nica–Toeplitz algebra of the product system \mathcal{E}^σ arising from Proposition 2.9. The quotient map $\dot{\phi}$ gives a representation of \mathcal{E}^σ in $C_r^*(\mathbb{Z}^n, \tilde{\sigma})$. In this case \mathcal{E}^σ is a product system of Hilbert bimodules since $\mathbb{K}(\mathcal{E}_p) = \mathbb{C}$ for all $p \in \mathbb{N}^n$. We claim that the canonical representation of \mathcal{E}^σ in the quotient $\mathcal{T}_r(\mathbb{N}^n, \sigma)/\mathcal{I}$ is Cuntz–Pimsner covariant on $\langle\langle \mathcal{E}_p \mid \mathcal{E}_p \rangle\rangle = \mathbb{C}$ for all $p \in \mathbb{N}^n$, where $\langle\langle \cdot \mid \cdot \rangle\rangle$ stands for the left inner product of a Hilbert bimodule. Indeed, it suffices to show that for all $p \in P$ the image of the projection L_p^σ under the quotient map is a unitary. We can write $p = \sum_{j=1}^n p_j e_j$, so that an application of the Nica covariance relation in $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ yields

$$L_p^\sigma (L_p^\sigma)^* = \prod_{j=1}^n L_{p_j e_j}^\sigma (L_{p_j e_j}^\sigma)^*. \quad (2.4)$$

Now each isometry $L_{p_j e_j}^\sigma$ has the form $a_j(L_{e_j}^\sigma)^{p_j}$ for some scalar $a_j \in \mathbb{C}$ of modulus 1. So the image of $L_{p_j e_j}^\sigma$ in $\mathcal{T}_r(\mathbb{N}^n, \sigma)/\mathcal{I}$ is a unitary for all $j = 1, 2, \dots, n$. Hence so is the image of L_p^σ by (2.4). This proves the claim.

Next we observe that \mathcal{E}^σ is simplifiable in the sense of [31, Definition 3.7] since for all $p, q \in \mathbb{N}^n$ one has

$$\langle\langle \mathcal{E}_p | \mathcal{E}_p \rangle\rangle \langle\langle \mathcal{E}_q | \mathcal{E}_q \rangle\rangle = \mathbb{C} = \langle\langle \mathcal{E}_{p \vee q} | \mathcal{E}_{p \vee q} \rangle\rangle.$$

Since the canonical representation of \mathcal{E}^σ in the quotient $\mathcal{T}_r(\mathbb{N}^n, \sigma)/\mathcal{I}$ is Cuntz–Pimsner covariant on $\langle\langle \mathcal{E}_p | \mathcal{E}_p \rangle\rangle$ for all $p \in \mathbb{N}^n$, it follows from [31, Lemma 3.11] that the fixed-point algebra of the canonical gauge action of \mathbb{T}^n on $\mathcal{T}_r(\mathbb{N}^n, \sigma)/\mathcal{I}$ is a copy of \mathbb{C} . The induced conditional expectation onto this copy is faithful because \mathbb{Z}^n is amenable. Now the homomorphism $\dot{\phi}: \mathcal{T}_r(\mathbb{N}^n, \sigma)/\mathcal{I} \rightarrow C_r^*(\mathbb{Z}^n, \tilde{\sigma})$ is gauge-compatible and clearly injective onto the fixed-point algebra \mathbb{C} , being a nonzero representation. Hence $\dot{\phi}$ is an isomorphism by [12, Proposition 19.8], completing the proof of the proposition. \square

Remark 2.13. It may be helpful to compare the conclusion of Proposition 2.12 with the analogous result in the theory of (untwisted) semigroup C^* -algebras. When σ is the trivial cocycle, that is, $\sigma(p, q) = 1$ for all $p, q \in \mathbb{N}^n$, $\mathcal{T}_r(\mathbb{N}^n, \sigma)$ is simply the Toeplitz C^* -algebra of \mathbb{N}^n . Proposition 2.12 gives a presentation for the (reduced) group C^* -algebra $C_r^*(\mathbb{Z}^n)$. This presentation may be obtained as an application of [8, Theorem 6.7] given that $C_r^*(\mathbb{Z}^n)$ is the boundary quotient $\partial\mathcal{T}_\lambda(\mathbb{N}^n)$ of $\mathcal{T}_\lambda(\mathbb{N}^n)$.

2.3. Symplectic bicharacters and antisymmetric matrices. The second cohomology group of a discrete abelian group G has a convenient parametrisation in terms of a specific subgroup of cocycles. Let $\sigma^*(x, y) := \overline{\sigma(y, x)}$ for $x, y \in G$ and recall that a 2-cocycle σ on G is a *symplectic bicharacter* if it is a bicharacter as a function $\sigma: G \times G \rightarrow \mathbb{T}$ and satisfies $\sigma^* = \sigma$. Denote by $X^2(G, \mathbb{T})$ the group of symplectic bicharacters on G . By [27, Proposition 3.2], the homomorphism $\sigma \mapsto \sigma\sigma^*$ of $Z^2(G, \mathbb{T})$ to $X^2(G, \mathbb{T})$ is surjective and has kernel precisely $B^2(G, \mathbb{T})$, so it induces an isomorphism of $H^2(G, \mathbb{T})$ onto $X^2(G, \mathbb{T})$.

The symplectic bicharacters on \mathbb{Z}^n are precisely the 2-cocycles associated to $n \times n$ antisymmetric matrices over \mathbb{R} . For each $n \times n$ antisymmetric matrix $\Theta = (\theta_{i,j})$ over \mathbb{R} , there is an associated symplectic bicharacter σ_Θ on \mathbb{Z}^n given by

$$\sigma_\Theta(x, y) = e^{-\pi i \langle x | \Theta y \rangle} \quad (x, y \in \mathbb{Z}^n),$$

and the map $\Theta \mapsto \sigma_\Theta$ is a group homomorphism. To retrieve the matrix from a given symplectic bicharacter σ , simply let $e^{\pi i \theta_{j,k}} := \sigma(e_k, e_j)$ for $1 \leq j \leq k \leq n$, which determines $\theta_{j,k}$ up to an additive even integer.

Recall that, at the level of twisted group algebras, $C^*(\mathbb{Z}^n, \sigma_\Theta)$ is canonically isomorphic to the noncommutative torus \mathcal{A}_Θ . Indeed, if $x \mapsto v_x$ is the canonical unitary σ_Θ -representation of \mathbb{Z}^n in $C^*(\mathbb{Z}^n, \sigma_\Theta)$, then the generators v_{e_j} for $j = 1, 2, \dots, n$ satisfy the defining commutation relations of \mathcal{A}_Θ . Conversely, if U_1, U_2, \dots, U_n are the canonical unitary generators of \mathcal{A}_Θ , and for each $x = (x_1, x_2, \dots, x_n) \in \mathbb{Z}^n$ we define

$$\bar{v}_x := e^{-\pi i \langle x - x_1 e_1 | \Theta x_1 e_1 \rangle} e^{-\pi i \langle x - x_1 e_1 - x_2 e_2 | \Theta x_2 e_2 \rangle} \dots e^{-\pi i \langle x_n e_n | \Theta x_{n-1} e_{n-1} \rangle} U_1^{x_1} U_2^{x_2} \dots U_n^{x_n}, \quad (2.5)$$

then $x \mapsto \bar{v}_x$ is a unitary σ_Θ -representation of \mathbb{Z}^n in \mathcal{A}_Θ .

Next we show that, in analogy to what happens for \mathcal{A}_Θ , the Toeplitz noncommutative torus $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ can be characterised as the universal C^* -algebra generated by n isometries subject to certain commutation relations.

Proposition 2.14. *Let Θ be an $n \times n$ antisymmetric real matrix and let $\sigma_\Theta(p, q) = e^{-\pi i \langle p | \Theta q \rangle}$ for $p, q \in \mathbb{N}^n$ be the associated symplectic bicharacter. Let $\{w_j \mid j \in \{1, \dots, n\}\}$ be isometries in*

a C^* -algebra B satisfying the relations

$$\begin{cases} w_j w_k = e^{-2\pi i \theta_{j,k}} w_k w_j & j, k = 1, 2, \dots, n; \\ w_j^* w_k = e^{2\pi i \theta_{j,k}} w_k w_j^* & j \neq k. \end{cases} \quad (\mathcal{R}_\Theta)$$

Then the map that sends $L_{e_j}^{\sigma_\Theta}$ to w_j induces a homomorphism from $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ to B that maps an isometry $L_p^{\sigma_\Theta}$ to the product

$$\sigma_\Theta(p - p_1 e_1, e_1)^{p_1} \sigma_\Theta(p - p_2 e_2 - p_1 e_1, e_2)^{p_2} \cdots \sigma_\Theta\left(p - \sum_{i=1}^{n-1} p_i e_i, e_{n-1}\right)^{p_{n-1}} w_1^{p_1} w_2^{p_2} \cdots w_n^{p_n}. \quad (2.6)$$

This homomorphism is an isomorphism if and only if $\prod_{j=1}^n (1 - w_j w_j^*) \neq 0$.

Proof. The first relation in (\mathcal{R}_Θ) implies that the map from \mathbb{N}^n to B given by (2.6) is an isometric σ_Θ -representation of \mathbb{N}^n . The second relation in (\mathcal{R}_Θ) implies that such a representation satisfies condition (3) of Definition 2.6, and hence is indeed a covariant σ_Θ -representation. The homomorphism from $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ to B and the condition for it to be an isomorphism now follow from Corollary 2.10. \square

In [2], Afsar, an Huef, Raeburn, and Sims consider certain Toeplitz extensions of noncommutative tori and use them as building blocks to construct their Toeplitz noncommutative solenoids. We will see below that these building blocks are quotients of special cases of our Toeplitz noncommutative tori.

Let k and d be nonnegative integers with $n = k + d$ and let $\Lambda \in M_{k,d}([0, \infty))$ be a $k \times d$ matrix. The C^* -algebra B_Λ considered in [2] is the universal C^* -algebra generated by a Nica covariant representation $v: \mathbb{N}^k \rightarrow B_\Lambda$ of \mathbb{N}^k and a unitary representation $u: \mathbb{Z}^d \rightarrow B_\Lambda$ of \mathbb{Z}^d , satisfying the commuting relations

$$u_x v_p = e^{2\pi i \langle p | \Lambda x \rangle} v_p u_x, \quad x \in \mathbb{Z}^d, p \in \mathbb{N}^k.$$

Proposition 2.15. *Suppose k and d are nonnegative integers with $n = k + d$. For each rectangular $k \times d$ matrix $\Lambda \in M_{k,d}([0, \infty))$ define $\Theta \in M_{k+d}(\mathbb{R})$ by consider the matrix*

Then the C^ -algebra B_Λ associated to Λ in [2] is canonically isomorphic to the quotient of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ by the ideal generated by the projections $1 - L_{e_j}^{\sigma_\Theta} (L_{e_j}^{\sigma_\Theta})^*$ for $j = k + 1, k + 2, \dots, k + d$.*

Proof. Let \mathcal{I}_d be the ideal of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ generated by the projections

$$\{1 - L_{e_j}^{\sigma_\Theta} (L_{e_j}^{\sigma_\Theta})^* \mid j = k + 1, k + 2, \dots, k + d\}.$$

For $p \in \mathbb{N}^n \setminus 0_k \times \mathbb{N}^d$, we write \bar{v}_p for the image of the isometry $L_p^{\sigma_\Theta}$ in the quotient $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)/\mathcal{I}_d$. Then the isometries $\{\bar{v}_p \mid p \in \mathbb{N}^k \times 0_d\}$ commute among themselves because the upper diagonal block of Θ consists of zeros. Also, the representation $p \in \mathbb{N}^k \cong \mathbb{N}^k \times 0_d \mapsto \bar{v}_p$ is Nica covariant by Proposition 2.9. If $j \in \{k + 1, \dots, k + d\}$, let \bar{u}_j denote the image of $L_{e_j}^{\sigma_\Theta}$ under the quotient map. Thus $\{\bar{u}_j \mid j = k + 1, \dots, k + d\}$ is a set of commuting unitaries. Hence there is a unique unitary representation $\bar{u}: \mathbb{Z}^d \cong 0_k \times \mathbb{Z}^d \rightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)/\mathcal{I}_d$ mapping a canonical generator $e_j \in 0_k \times \mathbb{Z}^d$ to \bar{u}_j , $j = k + 1, k + 2, \dots, k + d$. In addition, we have for $p \in \mathbb{N}^k \times 0_d$ and $j = k + 1, k + 2, \dots, k + d$,

$$\bar{u}_j \bar{v}_p = e^{-\pi i \langle e_j | \Theta p \rangle} \bar{v}_{e_j + p} = e^{-2\pi i \langle e_j | \Theta p \rangle} \bar{v}_p \bar{u}_j = e^{2\pi i \langle p | \Lambda e_j \rangle} \bar{v}_p \bar{u}_j.$$

This guarantees that the isometries $\{\bar{v}_p \mid p \in \mathbb{N}^k \times 0_d\}$ and the unitaries $\{\bar{u}_x \mid x \in 0_k \times \mathbb{Z}^d\}$ satisfy the commuting relations required in [2, Section 2]. We then get a homomorphism from B_Λ onto $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)/\mathcal{I}_d$ induced by the maps $\mathbb{N}^k \times 0_d \ni p \mapsto \bar{v}_p$, $0_k \times \mathbb{Z}^d \ni x \mapsto \bar{u}_x$. For the inverse map, we use the description of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ as a universal C^* -algebra generated by n isometries satisfying the relations (\mathcal{R}_Θ) given in Proposition 2.14. Notice that the second relation in (\mathcal{R}_Θ) is satisfied because $u_{e_j} v_{e_i} = e^{-2\pi i \langle e_j | \Theta e_i \rangle} v_{e_i} u_{e_j}$ in B_Λ implies

$$u_{e_j}^* v_{e_i} = e^{2\pi i \langle e_j | \Theta e_i \rangle} v_{e_i} u_{e_j}^*.$$

Thus the map that sends $L_{e_i}^{\sigma_\Theta}$ to v_{e_i} for $i = 1, \dots, k$ and $L_{e_j}^{\sigma_\Theta}$ to $u_{e_{j-k}}$ for $j = k+1, \dots, k+d$ gives a homomorphism from $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ onto B_Λ . The resulting homomorphism $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta) \rightarrow B_\Lambda$ factors through the quotient $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)/\mathcal{I}_d$, giving an inverse map for the canonical map $B_\Lambda \rightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)/\mathcal{I}_d$ built above. This finishes the proof. \square

In contrast to [2], our approach to Toeplitz noncommutative tori does not require either of the subsets $\{L_p^{\sigma_\Theta} \mid p \in \mathbb{N}^k \times 0_d\}$ and $\{L_x^{\sigma_\Theta} \mid x \in 0_k \times \mathbb{N}^d\}$ to consist of mutually commuting isometries. So our antisymmetric matrix Θ can have nontrivial diagonal blocks.

We also do not require any of the isometries to be unitaries, as is the case for the generators u_{e_1}, \dots, u_{e_d} considered in [2]. Nevertheless, for the dynamics considered in the following sections, KMS states necessarily factor through the quotient of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ in which the generating isometries that are fixed by the dynamics become unitaries, see Lemma 3.1 below. This confirms and explains the insight behind the choice made in [2, Section 2] where a subset of generators are chosen to be unitaries from the onset, and the dynamics acts nontrivially only on the others.

3. DYNAMICS AND KMS STATES

We fix an $n \times n$ antisymmetric matrix $\Theta = (\theta_{ij})$ throughout this section and we let $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ be the associated n -dimensional Toeplitz noncommutative torus. When the cocycle in question is understood and there is no chance of confusion, we will omit it from the notation of the left regular σ_Θ -representation of \mathbb{N}^n , simply writing L_p instead of $L_p^{\sigma_\Theta}$ for a canonical generating isometry. So $\{L_p \mid p \in \mathbb{N}^n\}$ is the canonical set of isometries on $\ell^2(\mathbb{N}^n)$ satisfying

$$L_p L_q = \sigma_\Theta(p, q) L_{p+q} = e^{-\pi i \langle p | \Theta q \rangle} L_{p+q}, \quad \text{and} \quad L_p L_p^* L_q L_q^* = L_{p \vee q} L_{p \vee q}^*,$$

and thus also condition (3) of Definition 2.6.

Let $r \in \mathbb{R}^n$ and let $\langle p | r \rangle = \sum_{j=1}^n p_j r_j$ be the usual scalar product of p and r . For each $t \in \mathbb{R}$ the map

$$u_t^r: \delta_q \mapsto e^{i \langle q | r \rangle t} \delta_q \quad (q \in \mathbb{N}^n)$$

extends to a (diagonal) unitary operator u_t^r on $\ell^2(\mathbb{N}^n)$ and $\{u_t^r \mid t \in \mathbb{R}\}$ is a strongly continuous one-parameter unitary group. The automorphism group of $\mathcal{B}(\ell^2(\mathbb{N}^n))$ obtained by conjugation with $\{u_t^r \mid t \in \mathbb{R}\}$ satisfies

$$u_t^r L_p (u_t^r)^* = e^{i \langle p | r \rangle t} L_p \quad (p \in \mathbb{N}^n),$$

and hence restricts to a strongly continuous one-parameter automorphism group $\{\alpha_t^r \mid t \in \mathbb{R}\}$ of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. This automorphism group is characterized by its action on generators:

$$\alpha_t^r(L_p) = e^{i \langle p | r \rangle t} L_p \quad (p \in \mathbb{N}^n).$$

An alternative way to obtain the dynamics is to observe that the vector r determines a group homomorphism $t \mapsto e^{i(r)t} = (e^{ir_1 t}, e^{ir_2 t}, \dots, e^{ir_n t})$ from \mathbb{R} to \mathbb{T}^n , and hence $\alpha_t^r = \gamma_{e^{i(r)t}}$ is simply the composition of this homomorphism with the canonical gauge action γ of \mathbb{T}^n on $\mathcal{T}_r(\mathbb{N}^n, \sigma)$.

We wish to study the KMS states for the dynamics associated to the vector r . Let $\beta \in \mathbb{R}$ and recall that a state φ of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ is a KMS_β state for the automorphism group $\{\alpha_t^r \mid t \in \mathbb{R}\}$ of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ if it satisfies the KMS_β condition

$$\varphi(AB) = \varphi(B\alpha_{i\beta}^r(A)) \tag{3.1}$$

for A α^r -analytic and $B \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. It is well known that it suffices to verify this for A and B in an α^r -invariant set of analytic elements with dense linear span, such as the generating elements $L_p L_q^*$ with $p, q \in \mathbb{N}^n$, see [26, Proposition 8.12.3].

It is immediate that $e^{-\langle p | r \rangle \beta} = \varphi(L_p L_p^*) \leq 1$ for every KMS_β state φ and each $p \in \mathbb{N}^n$. Hence, if $r_j < 0$ for some $j \in \{1, 2, \dots, n\}$, then there is no KMS_β state for $\beta > 0$, and similarly, if $r_j > 0$ for some $j \in \{1, 2, \dots, n\}$, then there is no KMS_β state for $\beta < 0$. It follows that for a KMS_β state to exist at nonzero β , each r_j must have the same sign as β . Here we choose to work

with nonnegative inverse temperatures. From now on we fix a vector $r \in [0, \infty)^n$, which we often omit from the notation, writing α instead of α^r when there is no risk of confusion.

We begin by verifying that the generating isometries corresponding to the vanishing coordinates of r become unitary operators in the GNS representation of any KMS_β state.

Lemma 3.1. *Let φ_β be a KMS_β state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Let $(\mathcal{H}_{\varphi_\beta}, \pi_{\varphi_\beta})$ denote the associated GNS representation. If $p \in \mathbb{N}^n$ and $\langle p | r \rangle = 0$, then $\pi_{\varphi_\beta}(L_p)$ is unitary. Moreover, φ_β factors through the quotient of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ modulo the ideal generated by the projections $\{1 - L_{e_j} L_{e_j}^* \mid r_j = 0\}$.*

Proof. Set $Q_p := 1 - L_p L_p^*$. Then Q_p is analytic, and by the KMS condition,

$$\varphi_\beta(Q_p) = 1 - \varphi_\beta(L_p L_p^*) = 1 - e^{-\langle p | r \rangle \beta} = 1 - 1 = 0.$$

We know that the elements of the form $L_p L_q^*$ have dense linear span in $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ and satisfy $\alpha_z(L_p L_q^*) = e^{i\langle p-q | r \rangle z} L_p L_q^*$. Hence the hypotheses of [17, Lemma 2.2] are satisfied, and so we conclude that φ factors through the quotient of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ by the ideal generated by the projections $\{Q_p \mid \langle p | r \rangle = 0\}$. This implies that $\varphi_\beta(B^* Q_p B) = 0$ for every $B \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$, which implies that $\pi_{\varphi_\beta}(L_p)$ is unitary as wished. In order to complete the proof of the lemma, notice that the ideal generated by the projections $\{Q_p \mid \langle p | r \rangle = 0\}$ coincides with the one generated by $\{1 - L_{e_j} L_{e_j}^* \mid r_j = 0\}$. \square

Given $d \in \mathbb{N}$, we denote by 0_d the zero element of \mathbb{N}^d , so that

$$0_d = \underbrace{(0, \dots, 0)}_{d \text{ times}}.$$

We shall assume without loss of generality that all the nonzero coordinates of r appear at the beginning. Thus we fix the notation $r = (r_1, r_2, \dots, r_k, 0_d)$ to indicate a vector in $[0, \infty)^n$ with strictly positive first k coordinates and zeros for the remaining d coordinates. This includes the two extreme cases of $r = 0$, in which case we have $k = 0$, and of strictly positive r , in which case $k = n$.

Proposition 3.2. *Let $n = k + d$ with $k, d \in \mathbb{N}$ and let $E^{(k)} := E^{\mathbb{T}^k \times \{1_d\}}$ denote the conditional expectation associated to the restriction of the gauge action of \mathbb{T}^n to the closed subgroup $\mathbb{T}^k \times \{1_d\}$. Then*

- (1) $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta) = \overline{\text{span}}\{L_p L_x L_y^* L_q^* \mid p, q \in \mathbb{N}^k \times 0_d, x, y \in 0_k \times \mathbb{N}^d\}$;
- (2) $E^{(k)}(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)) = \overline{\text{span}}\{L_p L_x L_y^* L_p^* \mid p \in \mathbb{N}^k \times 0_d, x, y \in 0_k \times \mathbb{N}^d\}$.

Proof. That the elements of the form $L_p L_x L_y^* L_q^*$ with $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$ span a dense $*$ -subalgebra of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ is a consequence of relations (1)–(3) of Definition 2.6. The second assertion follows from part (1) because $E^{(k)}$ is a contraction that satisfies $E^{(k)}(L_p L_x L_y^* L_q^*) = \delta_{p,q} L_p L_x L_y^* L_p^*$, where $\delta_{p,q}$ is the Kronecker delta. \square

Proposition 3.3. *Let $n = k + d$ with $k, d \in \mathbb{N}$ and let α be the dynamics determined by $r = (r_1, \dots, r_k, 0_d)$. Let $0 < \beta < \infty$ and suppose that φ is a KMS_β state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Then φ restricts to a trace on the C^* -subalgebra $C^*(L_x : x \in 0_k \times \mathbb{N}^d)$ and satisfies*

$$\varphi(L_p L_x L_y^* L_q^*) = \delta_{p,q} e^{-\beta \langle p | r \rangle} \varphi(L_x L_y^*) \quad (3.2)$$

for all $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$, where $\delta_{p,q}$ is the Kronecker delta.

Proof. Suppose that φ is a KMS_β state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Since the elements in $C^*(L_x : x \in 0_k \times \mathbb{N}^d)$ are fixed points of α , the KMS_β condition implies that the restriction of φ to $C^*(L_x : x \in 0_k \times \mathbb{N}^d)$ is a trace.

We show next that φ satisfies (3.2). Let $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$. In case $p = q$, the KMS_β condition with $A = L_p$ and $B = L_x L_y^* L_q^*$ gives (3.2) because $L_p^* L_q = 1$. So suppose $p \neq q$. We aim to prove that

$$\varphi(L_p L_x L_y^* L_q^*) = 0.$$

For this, we write $p' := (p \vee q) - p$ and $q' := (p \vee q) - q$. Notice that at least one of the numbers p', q' is nonzero. Because φ is a state and so preserves adjoints, we may assume without loss of generality that $q' \neq 0$. Applying the KMS condition (3.1) with $A = L_p L_x L_y^*$ and $B = L_q^*$ and using that σ_Θ is a circle-valued function we get

$$|\varphi(L_p L_x L_y^* L_q^*)| = e^{-\beta\langle p|r\rangle} |\varphi(L_q^* L_p L_x L_y^*)| = e^{-\beta\langle p|r\rangle} |\varphi(L_{q'} L_{p'}^* L_x L_y^*)|.$$

Since $p' \vee x = x + p'$, we have $L_{p'}^* L_x = \sigma_\Theta(x, p')^2 L_x L_{p'}^*$. Thus

$$|\varphi(L_p L_x L_y^* L_q^*)| = e^{-\beta\langle p|r\rangle} |\varphi(L_{q'} L_x L_{p'}^* L_y^*)| = e^{-\beta\langle p|r\rangle} |\varphi(L_{q'} L_x L_y^* L_{p'}^*)|. \quad (3.3)$$

Observe that $p' \vee q' = p' + q'$ and $p' \vee y = p' + y$. Hence the same argument as above gives

$$|\varphi(L_{q'} L_x L_y^* L_{p'}^*)| = e^{-\beta\langle q'|r\rangle} |\varphi(L_{q'} L_x L_y^* L_{p'}^*)|.$$

Continuing this process, we see that for each $l \in \mathbb{N}$

$$|\varphi(L_{q'} L_x L_y^* L_{p'}^*)| = e^{-l\beta\langle q'|r\rangle} |\varphi(L_{q'} L_x L_y^* L_{p'}^*)|. \quad (3.4)$$

Substituting (3.4) into (3.3) we arrive at

$$|\varphi(L_p L_x L_y^* L_q^*)| = e^{-\beta\langle p|r\rangle} e^{-l\beta\langle q'|r\rangle} |\varphi(L_{q'} L_x L_y^* L_{p'}^*)|.$$

Since $q' \neq 0$, it follows that $\lim_{l \rightarrow \infty} e^{-l\beta\langle q'|r\rangle} = 0$. Therefore $\varphi(L_p L_x L_y^* L_q^*) = 0$ as wished. \square

Lemma 3.4. *Let $n = k + d$ with $k, d \in \mathbb{N}$. The product*

$$Q := \prod_{j=1}^k (1 - L_{e_j} L_{e_j}^*) \quad (3.5)$$

is a projection in $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ satisfying

- (1) $QL_p = 0 = L_p^* Q$ for every $p \in \mathbb{N}^k \times 0_d \setminus \{0\}$;
- (2) $QL_p^* L_x L_y^* L_p = L_p^* L_x L_y^* L_p Q = QL_p^* L_x L_y^* L_p Q$ for every $x, y \in 0_k \times \mathbb{N}^d$ and $p \in \mathbb{N}^k \times 0_d$;
- (3) $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q = \overline{\text{span}}\{QL_x L_y^* Q \mid x, y \in 0_k \times \mathbb{N}^d\}$.

Proof. By definition, Q is the product of a finite collection of commuting projections, thus is a projection as well. In order to prove (1), let $0 \neq p \in \mathbb{N}^k \times 0_d$ and let j be the first nonzero coordinate of p . Then the product QL_p contains a factor of the form $(1 - L_{e_j} L_{e_j}^*) L_{e_j}$ and hence vanishes. Taking adjoints shows that $L_p^* Q$ vanishes too.

In order to prove (2), notice first that $e_i \vee x = e_i + x$ for every $x \in 0_k \times \mathbb{N}^d$ and every $i = 1, 2, \dots, k$. Thus by Lemma 2.8 the projection $L_{e_i} L_{e_i}^*$ commutes with L_x , with every L_y^* , and with every $L_x L_y^*$ for $y \in 0_k \times \mathbb{N}^d$. Thus Q commutes with $L_x L_y^*$ for all $x, y \in 0_k \times \mathbb{N}^d$. Now take $p \in \mathbb{N}^k \times 0_d$ and notice that Definition 2.6(3) applied twice gives

$$L_p^* L_x L_y^* L_p = \sigma_\Theta(x, p)^2 \sigma_\Theta(p, y)^2 L_x L_y^*.$$

So Q commutes with $L_p^* L_x L_y^* L_p$ because it commutes with $L_x L_y^*$. This completes the proof of part (2).

For part (3), we use that

$$\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta) = \overline{\text{span}}\{L_p L_x L_y^* L_q^* \mid p, q \in \mathbb{N}^k \times 0_d, x, y \in 0_k \times \mathbb{N}^d\}.$$

So the products $QL_pL_xL_y^*L_q^*Q$ span a dense subset of the corner. If $p, q \in \mathbb{N}^k \times 0_d$ are not both zero, and if $x, y \in 0_k \times \mathbb{N}^d$, then $QL_pL_xL_y^*L_q^*Q = 0$ because of part (1). Hence the corner is the closed linear span of the products $QL_xL_y^*Q$ with $x, y \in 0_k \times \mathbb{N}^d$. \square

Lemma 3.5. *Let $0 < \beta < \infty$ and suppose that φ is a KMS_β state for $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Let Q be the projection from (3.5). Then $\varphi(Q) > 0$ and the sum $Z(\beta) := \sum_{p \in \mathbb{N}^k \times 0_d} e^{-\beta\langle p|r \rangle}$ satisfies $Z(\beta) = \varphi(Q)^{-1}$. Moreover, the map $\omega_\varphi: X \mapsto Z(\beta)\varphi(X)$ for $X \in Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ is a tracial state of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ and φ can be reconstructed from ω_φ by*

$$\varphi(X) = \frac{1}{Z(\beta)} \sum_{p \in \mathbb{N}^k \times 0_d} e^{-\beta\langle p|r \rangle} \omega_\varphi(QL_p^*XL_pQ) \quad \text{for all } X \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta). \quad (3.6)$$

Proof. Expanding the product defining Q , we have

$$Q = \sum_{J \subseteq \{1, \dots, k\}} (-1)^{|J|} \prod_{j \in J} L_{e_j} L_{e_j}^* \quad (3.7)$$

since $\{L_p L_p^* \mid p \in \mathbb{N}^n\}$ are commuting projections. Let $i, j \in \{1, \dots, k\}$ with $i \neq j$, so that $L_{e_i} L_{e_i}^*$ commutes with L_{e_j} by Lemma 2.8. Then

$$\varphi(L_{e_i} L_{e_i}^* L_{e_j} L_{e_j}^*) = \varphi(L_{e_j} L_{e_j}^* L_{e_i} L_{e_i}^*) = e^{-\beta(r_j + r_i)}$$

by the KMS condition. An analogous computation shows that for each $J \subset \{1, \dots, k\}$, we have

$$\varphi\left(\prod_{j \in J} L_{e_j} L_{e_j}^*\right) = e^{-\beta \sum_{j \in J} r_j}.$$

It follows that

$$\varphi(Q) = \sum_{J \subseteq \{1, \dots, k\}} (-1)^{|J|} \varphi\left(\prod_{j \in J} L_{e_j} L_{e_j}^*\right) = \sum_{J \subseteq \{1, \dots, k\}} (-1)^{|J|} e^{-\beta \sum_{j \in J} r_j} = \prod_{j=1}^k (1 - e^{-\beta r_j}), \quad (3.8)$$

which is obviously positive and equal to $Z(\beta)^{-1}$ by an Euler product like expansion.

If $p \neq l$, then

$$L_p Q L_p^* L_l Q L_l^* = \overline{\sigma_\Theta(p, p \vee l - p)} \sigma_\Theta(l, p \vee l - l) L_p Q L_{p \vee l - p} L_{p \vee l - l}^* Q L_l^* = 0$$

by Lemma 3.4(1) because $p \vee l - p$ and $p \vee l - l$ cannot both be zero. Hence the sum of mutually orthogonal projections

$$P_q := \sum_{p \leq q} L_p Q L_p^*$$

is a projection itself and satisfies

$$\varphi(P_q) = \sum_{p \leq q} \varphi(L_p Q L_p^*) = \sum_{p \leq q} e^{-\beta\langle p|r \rangle} \varphi(L_p^* L_p Q) = \sum_{p \leq q} e^{-\beta\langle p|r \rangle} \varphi(Q).$$

It follows that $\varphi(P_q) \rightarrow 1$ as $q \rightarrow \infty$ in the directed set $\mathbb{N}^k \times 0_d$ since $\sum_{p \leq q} e^{-\beta\langle p|r \rangle} \rightarrow Z(\beta) = \varphi(Q)^{-1}$ by the first part of the proof. Hence an application of the Cauchy–Schwarz inequality implies that $\varphi(P_q X P_q) \rightarrow \varphi(X)$ as $q \rightarrow \infty$ for all $X \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. Thus by the KMS condition we have

$$\begin{aligned} \varphi(X) &= \sum_{p, l \in \mathbb{N}^k \times 0_d} \varphi(L_p Q L_p^* X L_l Q L_l^*) \\ &= \sum_{p \in \mathbb{N}^k \times 0_d} \varphi(L_p Q L_p^* X L_p Q L_p^*) \\ &= \sum_{p \in \mathbb{N}^k \times 0_d} e^{-\beta\langle p|r \rangle} \varphi(Q L_p^* X L_p Q) \end{aligned}$$

We then get the reconstruction formula (3.6) by replacing the summand $\varphi(QL_p^*XL_pQ)$ by $\frac{1}{Z(\beta)}\omega_\varphi(QL_p^*XL_pQ)$. \square

Proposition 3.6. *Let $0 < \beta < \infty$ and let Q be the projection from (3.5). For each tracial state ω of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$, define*

$$T_\beta(\omega)(X) := \frac{1}{Z(\beta)} \sum_{l \in \mathbb{N}^k \times 0_d} e^{-\beta\langle l|r \rangle} \omega(QL_l^*XL_lQ), \quad X \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta). \quad (3.9)$$

Then T_β is an affine weak* homeomorphism of the tracial state space of the corner onto the KMS $_\beta$ state space of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$.

Proof. Let ω be a tracial state on the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$. Clearly $T_\beta(\omega)$ is a state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. First we verify that $T_\beta(\omega)$ satisfies (3.2). Let $X = L_pL_xL_y^*L_q^*$, where $p, q \in \mathbb{N}^k \times 0_d$, $x, y \in 0_k \times \mathbb{N}^d$. Given $l \in \mathbb{N}^k \times 0_d$, Lemma 3.4(1) implies that if the elements $l \vee p - l$ and $l \vee q - l$ are not both zero, then $QL_l^*XL_lQ = 0$. Since $l \vee p - l = 0 = l \vee q - l$ is equivalent to $l \geq p \vee q$, we may restrict the sum to $l \geq p \vee q$ and simplify $L_l^*L_p = \sigma_\Theta(p, l-p)L_{l-p}^*$ using Definition 2.6 to get

$$T_\beta(\omega)(X) = \frac{1}{Z(\beta)} \sum_{\substack{l \in \mathbb{N}^k \times 0_d \\ l \geq p \vee q}} \frac{\sigma(p, l-p)}{\sigma(q, l-q)} e^{-\beta\langle l|r \rangle} \omega(QL_{l-p}^*L_xL_y^*L_{l-q}Q). \quad (3.10)$$

When $p = q$, a change of the index of summation to $m = l - p$ gives

$$\begin{aligned} T_\beta(\omega)(X) &= \frac{1}{Z(\beta)} \sum_{\substack{l \in \mathbb{N}^k \times 0_d \\ l \geq p}} e^{-\beta\langle l|r \rangle} \omega(QL_{l-p}^*L_xL_y^*L_{l-p}Q) \\ &= \frac{1}{Z(\beta)} \sum_{\substack{m \in \mathbb{N}^k \times 0_d \\ m \geq 0}} e^{-\beta\langle p|r \rangle} e^{-\beta\langle m|r \rangle} \omega(QL_m^*L_xL_y^*L_mQ) \\ &= e^{-\beta\langle p|r \rangle} T_\beta(\omega)(L_xL_y^*), \end{aligned} \quad (3.11)$$

To see that $T_\beta(\omega)(X) = 0$ if $p \neq q$, it suffices to show $QL_{l-p}^*L_xL_y^*L_{l-q}Q$ vanishes for all $l \geq p \vee q$, $l \in \mathbb{N}^k \times 0_d$. By Lemma 2.8 the projection $L_{l-p}L_{l-p}^*$ commutes $L_sL_t^*$ with $s, t \in 0_k \times \mathbb{N}^d$. In addition, it follows from Lemma 3.4(2) that $L_{l-p}^*L_xL_y^*L_{l-p}$ commutes with Q . So

$$QL_{l-p}^*L_xL_y^*L_{l-q}Q = QL_{l-p}^*L_xL_y^*L_{l-p}L_{l-p}^*L_{l-q}Q = QL_{l-p}^*L_xL_y^*L_{l-p}QL_{l-p}^*L_{l-q}Q.$$

Since $p \neq q$ implies that at least one of the elements $((l-p) \vee (l-q)) - (l-p)$ and $((l-p) \vee (l-q)) - (l-q)$ is nonzero, Lemma 3.4(1) yields $QL_{l-p}^*L_{l-q}Q = 0$ and hence $QL_{l-p}^*L_xL_y^*L_{l-q}Q = 0$. Therefore $T_\beta(\omega)(X) = 0$ when $p \neq q$, as asserted.

We aim to prove next that $T_\beta(\omega)$ satisfies the KMS $_\beta$ condition for the dynamics α . Let $p, q, a, b \in \mathbb{N}^k \times 0_d$ and $x, y, s, t \in 0_k \times \mathbb{N}^d$ and consider the elements $X = L_pL_xL_y^*L_q^*$ and $Y = L_aL_sL_t^*L_b^*$, so that $\alpha_{i\beta}(X) = e^{-\beta\langle p-q|r \rangle}X$. In order to conclude that $T_\beta(\omega)$ is a KMS $_\beta$ state it suffices to show that $T_\beta(\omega)(XY)e^{\beta\langle p|r \rangle} = e^{\beta\langle q|r \rangle}T_\beta(\omega)(YX)$. We may assume that $p - q + a - b = 0$, for otherwise both sides vanish because we have shown that $T_\beta(\omega)$ satisfies (3.2). By definition

$$T_\beta(XY) = \sum_{l \in \mathbb{N}^k \times 0_d} \frac{e^{-\beta\langle l|r \rangle}}{Z(\beta)} \omega(QL_l^*L_pL_xL_y^*L_q^*L_aL_sL_t^*L_b^*L_lQ).$$

The argument leading to (3.10) shows that we may restrict the summation to $l \geq p \vee b$, in which case $L_l^* L_p = \sigma_\Theta(p, l-p) L_{l-p}^*$ and similarly $L_b^* L_l = \sigma_\Theta(b, l-b) L_{l-b}$ by Definition 2.6(3). Thus

$$\begin{aligned}
T_\beta(XY) &= \sum_{\substack{l \in \mathbb{N}^k \times 0_d \\ l \geq p \vee b}} \frac{e^{-\beta \langle l | r \rangle}}{Z(\beta)} \frac{\sigma(p, l-p)}{\sigma(b, l-b)} \omega(Q L_{l-p}^* L_x L_y^* L_q^* L_a L_s L_t^* L_{l-b} Q) \\
&= \sum_{\substack{l \in \mathbb{N}^k \times 0_d \\ l \geq p \vee b}} \frac{e^{-\beta \langle l | r \rangle}}{Z(\beta)} \frac{\sigma_\Theta(p, l-p)}{\sigma_\Theta(b, l-b)} \omega(Q L_{l-p}^* L_x L_y^* (L_{l-p} L_{l-p}^*) L_q^* L_a (L_{l-b} L_{l-b}^*) L_s L_t^* L_{l-b} Q) \\
&= \sum_{\substack{l \in \mathbb{N}^k \times 0_d \\ l \geq p \vee b}} \frac{e^{-\beta \langle l | r \rangle}}{Z(\beta)} \frac{\sigma_\Theta(p, l-p)}{\sigma_\Theta(b, l-b)} \omega((Q L_{l-p}^* L_x L_y^* L_{l-p} Q) L_{l-p}^* L_q^* L_a L_{l-b} (Q L_{l-b}^* L_s L_t^* L_{l-b} Q)),
\end{aligned} \tag{3.12}$$

where we have used that $L_{l-p} L_{l-p}^*$ commutes with $L_x L_y^*$, that $L_{l-b} L_{l-b}^*$ commutes with $L_s L_t^*$, and that Q commutes with $L_{l-p}^* L_x L_y^* L_{l-p}$ and with $L_{l-b}^* L_s L_t^* L_{l-b}$. Notice also that the product in the middle simplifies to a scalar, namely

$$L_{l-p}^* L_q^* L_a L_{l-b} = \frac{\sigma_\Theta(a, l-b)}{\sigma_\Theta(q, l-p)} L_{q+l-p}^* L_{a+l-b} = \frac{\sigma_\Theta(a, l-b)}{\sigma_\Theta(q, l-p)}$$

because $q-p = a-b$. When we substitute this in the formula and change the index of summation to $m = l-p$ we get

$$\begin{aligned}
e^{\beta \langle p | r \rangle} T_\beta(XY) &= \\
&= \sum_{\substack{m \in \mathbb{N}^k \times 0_d \\ m+p \geq b}} \frac{e^{-\beta \langle m | r \rangle}}{Z(\beta)} \frac{\sigma_\Theta(p, m) \sigma_\Theta(a, p+m-b)}{\sigma_\Theta(b, m+p-b) \sigma_\Theta(q, m)} \omega((Q L_m^* L_x L_y^* L_m Q) (Q L_{m+p-b}^* L_s L_t^* L_{m+p-b} Q)).
\end{aligned} \tag{3.13}$$

Exchanging now the roles of X and Y and carrying out a computation like (3.12) gives

$$\begin{aligned}
T_\beta(YX) &= \\
&= \sum_{\substack{l \in \mathbb{N}^k \times 0_d \\ l \geq a \vee q}} \frac{e^{-\beta \langle l | r \rangle}}{Z(\beta)} \frac{\sigma_\Theta(a, l-a)}{\sigma_\Theta(q, l-q)} \omega((Q L_{l-a}^* L_s L_t^* L_{l-a} Q) L_{l-a}^* L_b^* L_p L_{l-q} (Q L_{l-q}^* L_x L_y^* L_{l-q} Q))
\end{aligned}$$

Substituting the scalar $L_{l-a}^* L_b^* L_p L_{l-q} = \overline{\sigma_\Theta(b, l-a)} \sigma_\Theta(p, l-q)$ in the middle and changing the index of summation to $m = l-q$, we get

$$\begin{aligned}
e^{\beta \langle q | r \rangle} T_\beta(YX) &= \\
&= \sum_{\substack{m \in \mathbb{N}^k \times 0_d \\ m+q \geq a}} \frac{e^{-\beta \langle m | r \rangle}}{Z(\beta)} \frac{\sigma_\Theta(a, m+q-a) \sigma_\Theta(p, m)}{\sigma_\Theta(q, m) \sigma_\Theta(b, q+m-a)} \omega((Q L_{m+q-a}^* L_s L_t^* L_{m+q-a} Q) (Q L_m^* L_x L_y^* L_m Q)).
\end{aligned} \tag{3.14}$$

Since $q-a = p-b$, so that $m+q \geq a$ iff $m+p \geq b$, and since ω is a trace on the corner, the two series in (3.13) and (3.14) are the same, term by term, which shows $e^{\beta \langle p | r \rangle} T_\beta(\omega)(XY) = e^{\beta \langle q | r \rangle} T_\beta(\omega)(YX)$. This completes the proof that T_β maps tracial states of $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ to KMS $_\beta$ states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$.

Notice that T_β is surjective by Lemma 3.5 and injective because the restriction of $T_\beta(\omega)$ to $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ is equal to $Z(\beta)\omega$. Clearly it is also an affine map; its inverse is given by the map $\varphi \mapsto \omega_\varphi$ of Lemma 3.5, which is obviously weak* continuous. We then conclude that T_β is a weak* homeomorphism, as it is a bijection with continuous inverse between compact Hausdorff spaces. This completes the proof of the proposition. \square

4. KMS $_{\beta}$ STATES AND TRACES ON NONCOMMUTATIVE TORI

Our first goal in this section is to show that the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})Q$ is isomorphic to the C^* -subalgebra of $\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})$ generated by $\{L_x \mid x \in 0_k \times \mathbb{N}^d\}$, which is itself isomorphic to the Toeplitz noncommutative torus $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ associated to the restriction Θ_d of Θ to the last d coordinates. We then show that the traces of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ factorize through its canonical quotient \mathcal{A}_{Θ_d} , and in fact come from states of its center, which is a classical torus of dimension equal to the degeneracy index of Θ_d .

Lemma 4.1. *Let Q be the projection from (3.5) and denote by Θ_d the lower right $d \times d$ corner of Θ . Then $C^*(L_x : x \in 0_k \times \mathbb{N}^d)$ is canonically isomorphic to the Toeplitz noncommutative torus $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$, and the map $\rho_Q : C^*(L_x : x \in 0_k \times \mathbb{N}^d) \rightarrow Q\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})Q$ given by the compression $X \mapsto QXQ$ is an isomorphism.*

Proof. The set of isometries $\{L_x \mid x \in 0_k \times \mathbb{N}^d\}$ satisfies relations (1)–(3) from Definition 2.6. Using the obvious identification $\mathbb{N}^d \cong 0_k \times \mathbb{N}^d$, we obtain a covariant isometric σ_{Θ_d} -representation of \mathbb{N}^d in $C^*(L_x : x \in 0_k \times \mathbb{N}^d)$. By Corollary 2.10, this gives a homomorphism from $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ onto $C^*(L_x : x \in 0_k \times \mathbb{N}^d)$ mapping $L_x^{\sigma_{\Theta_d}}$ to L_x . It is faithful because $\prod_{j=k+1}^n (1 - L_{e_j} L_{e_j}^*) \neq 0$. Notice that Q is the projection of $\ell^2(\mathbb{N}^n)$ onto the subspace $\ell^2(0_k \times \mathbb{N}^d)$, which is invariant for $C^*(L_x : x \in 0_k \times \mathbb{N}^d)$ and hence $\rho_Q : C^*(L_x : x \in 0_k \times \mathbb{N}^d) \rightarrow Q\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})Q$ is an isomorphism. \square

The isomorphisms from the lemma above allow us to express our characterisation of KMS $_{\beta}$ states in terms of traces on $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$.

Proposition 4.2. *Let ρ_Q be the isomorphism from Lemma 4.1 and identify $C^*(L_x \mid x \in 0_k \times \mathbb{N}^d)$ with $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ canonically. For each tracial state τ of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ there is a KMS $_{\beta}$ state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta}), \alpha)$ determined by*

$$T_{\beta}(\tau \circ \rho_Q^{-1})(L_p L_x L_y^* L_q^*) = \delta_{p,q} \tau(L_x L_y^*) \prod_{j=1}^k \frac{e^{-\beta r_j p_j} (1 - e^{-\beta r_j})}{1 - e^{-\beta r_j + 2\pi i \langle \Theta(x-y) | e_j \rangle}}, \quad (4.1)$$

where $x, y \in 0_k \times \mathbb{N}^d \cong \mathbb{N}^d$. The map $\tau \mapsto T_{\beta}(\tau \circ \rho_Q^{-1})$ is an affine weak* homeomorphism of the tracial state space of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ onto the simplex of KMS $_{\beta}$ states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta}), \alpha)$.

Proof. The (tracial) states of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})Q$ come from the (tracial) states of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ via the map $\tau \mapsto \tau \circ \rho_Q^{-1}$. Combining this with the map T_{β} from Proposition 3.6 we see that $\tau \mapsto T_{\beta}(\tau \circ \rho_Q^{-1})$ is an affine weak* homeomorphism of the tracial state space of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ onto the simplex of KMS $_{\beta}$ states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta}), \alpha)$.

In order to write $T_{\beta}(\tau \circ \rho_Q^{-1})$ in (3.9) in terms of the tracial state τ of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$, we first use Definition 2.6(3) and the fact that σ_{Θ} is a symplectic bicharacter to write

$$QL_m^* L_x L_y^* L_m Q = \sigma_{\Theta}(x, m)^2 \sigma_{\Theta}(m, y)^2 QL_x L_y^* Q = \sigma_{\Theta}(x - y, m)^2 QL_x L_y^* Q$$

for all $m \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$. Next, we use (3.11) with $\tau \circ \rho_Q^{-1}$ playing the role of ω , observing that $\rho_Q^{-1}(QL_x L_y^* Q) = L_x L_y^*$, and we conclude that

$$T_{\beta}(\tau \circ \rho_Q^{-1})(L_p L_x L_y^* L_q^*) = \tau(L_x L_y^*) \frac{\delta_{p,q} e^{-\beta \langle p | r \rangle}}{Z(\beta)} \sum_{m \in \mathbb{N}^k \times 0_d} e^{-\beta \langle m | r \rangle} \sigma_{\Theta}(x - y, m)^2. \quad (4.2)$$

The rest of the proof is a computation to obtain an Euler product formula for the series involving the cocycle. Observe that $\langle m | r \rangle = \langle r | m \rangle \geq 0$ and that $\sigma_{\Theta}(x - y, m)^2 = e^{-2\pi i \langle x-y | \Theta m \rangle} =$

$e^{2\pi i \langle \Theta(x-y) | m \rangle}$. The series from (4.2) then becomes

$$\begin{aligned} \sum_{m \in \mathbb{N}^k \times 0_d} e^{-\beta \langle m | r \rangle} \sigma_{\Theta}(x-y, m)^2 &= \sum_{m \in \mathbb{N}^k \times 0_d} e^{\langle -\beta r + 2\pi i \Theta(x-y) | m \rangle} \\ &= \sum_{m \in \mathbb{N}^k \times 0_d} e^{\langle -\beta r + 2\pi i \Theta(x-y) | m \rangle} \\ &= \sum_{m \in \mathbb{N}^k \times 0_d} e^{\sum_{j=1}^k \langle -\beta r + 2\pi i \Theta(x-y) | e_j \rangle \langle e_j | m \rangle} \\ &= \sum_{m \in \mathbb{N}^k \times 0_d} \prod_{j=1}^k \left(e^{\langle -\beta r + 2\pi i \Theta(x-y) | e_j \rangle} \right)^{m_j}. \end{aligned}$$

To simplify the notation, we define $A_j := e^{\langle -\beta r + 2\pi i \Theta(x-y) | e_j \rangle}$ and notice that $|A_j| < 1$ because $\langle \beta r | e_j \rangle > 0$ for $1 \leq j \leq k$. Then

$$\sum_{m \in \mathbb{N}^k \times 0_d} e^{-\beta \langle m | r \rangle} \sigma_{\Theta}(x-y, m)^2 = \sum_{m \in \mathbb{N}^k \times 0_d} \prod_{j=1}^k A_j^{m_j} = \prod_{j=1}^k \sum_{n=0}^{\infty} A_j^n = \prod_{j=1}^k \frac{1}{1 - A_j}.$$

Thus we obtain (4.1) by substituting into (4.2) the above expression and the usual Euler product expansion for $Z(\beta)^{-1}$ from (3.8), and replacing $e^{-\beta \langle p | r \rangle}$ by the product $\prod_{j=1}^k e^{-\beta r_j p_j}$. \square

The remainder of this section is dedicated to provide a concrete description of the space of tracial states of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$. Since our considerations are general, we momentarily adjust the notation and consider a generic $d \times d$ matrix D . We begin by showing that the traces of $\mathcal{T}_r(\mathbb{N}^d, \sigma_D)$ factor through the noncommutative torus \mathcal{A}_D .

Lemma 4.3. *Let $D \in M_d(\mathbb{R})$ be an antisymmetric matrix and let \mathcal{A}_D be the associated noncommutative d -torus, with generating unitaries U_j , for $j = 1, \dots, d$. The map $L_{e_j} \mapsto U_j$ extends to a surjective homomorphism $\pi: \mathcal{T}_r(\mathbb{N}^d, \sigma_D) \rightarrow \mathcal{A}_D$ which, in turn, induces an affine homeomorphism $\tau \mapsto \tau \circ \pi$ from the space of tracial states of \mathcal{A}_D onto the space of tracial states of $\mathcal{T}_r(\mathbb{N}^d, \sigma_D)$.*

Proof. By Proposition 2.14, the map that sends an isometry $L_{e_j} \in \mathcal{T}_r(\mathbb{N}^d, \sigma_D)$ to the unitary $U_j \in \mathcal{A}_D$ for each $1 \leq j \leq d$ induces a surjective homomorphism $\pi: \mathcal{T}_r(\mathbb{N}^d, \sigma_D) \rightarrow \mathcal{A}_D$. Hence the map that sends a tracial state τ of \mathcal{A}_D to the composite $\tau \circ \pi$ is an injective affine weak* continuous map from the tracial state space of \mathcal{A}_D into the tracial state space of $\mathcal{T}_r(\mathbb{N}^d, \sigma_D)$. To see that this map is a homeomorphism, it suffices to show that it is also surjective as the underlying spaces are compact and Hausdorff.

Suppose ψ is a tracial state on $\mathcal{T}_r(\mathbb{N}^d, \sigma_D)$ and let (H_{ψ}, π_{ψ}) be its GNS representation. By Lemma 3.1 each $\pi_{\psi}(L_x)$ is a unitary operator and by Proposition 2.14 the collection of unitaries $\{\pi_{\psi}(L_{e_j}) \mid j = 1, \dots, d\}$ satisfies the defining relations of \mathcal{A}_D . So by the universal property of \mathcal{A}_D , there is a homomorphism $\rho: \mathcal{A}_D \rightarrow \pi_{\psi}(\mathcal{T}_r(\mathbb{N}^d, \sigma_D))$ such that $\rho(U_j) = \pi_{\psi}(L_{e_j})$ for each $j = 1, \dots, d$. Thus

$$\psi(L_{e_j}) = \langle \pi_{\psi}(L_{e_j}) \xi_{\psi}, \xi_{\psi} \rangle = \langle \rho(U_j) \xi_{\psi}, \xi_{\psi} \rangle = \langle (\rho \circ \pi)(L_{e_j}) \xi_{\psi}, \xi_{\psi} \rangle,$$

showing that ψ factors through the homomorphism $\pi: \mathcal{T}_r(\mathbb{N}^d, \sigma_D) \rightarrow \mathcal{A}_D$ and finishing the proof of the lemma. \square

Following [32, Section 1], see also [28], we say that an antisymmetric, real, $d \times d$ matrix D is *nondegenerate* if whenever $x \in \mathbb{Z}^d$ and $\langle x | Dy \rangle \in \mathbb{Z}$ for all $y \in \mathbb{Z}^d$, then $x = 0$. The noncommutative torus \mathcal{A}_D is simple if and only if D is nondegenerate. See [32, Theorem 3.7] and also [28, Theorem 1.9].

There is always a canonical tracial state on \mathcal{A}_D , which is given by the conditional expectation $E^{\gamma}: \mathcal{A}_D \rightarrow \mathcal{A}_D^{\gamma} = \mathbb{C}$, where γ is the canonical gauge action of \mathbb{T}^d on \mathcal{A}_D . This is the unique

tracial state of \mathcal{A}_D when D is nondegenerate (see, for example, [28, Theorem 1.9]). In general, we consider the subgroup

$$H := \{x \in \mathbb{Z}^d \mid \langle x \mid Dy \rangle \in \mathbb{Z} \text{ for all } y \in \mathbb{Z}^d\}$$

of \mathbb{Z}^d . We refer to the rank of H as the *degeneracy index* of D . We write $m := \text{rank } H$ for the degeneracy index of D and recall that there exist a basis $\{p_1, p_2, \dots, p_d\}$ of \mathbb{Z}^d and positive integers a_1, a_2, \dots, a_m with $a_i \mid a_{i+1}$ such that $\{a_1 p_1, \dots, a_m p_m\}$ is a basis for H (see, for example, [15, Theorem 2.6]). We consider such a basis in what follows.

Lemma 4.4. *Let $D \in M_d(\mathbb{R})$ be an antisymmetric matrix and let \mathcal{A}_D be the associated noncommutative torus, with canonical unitary generators U_1, \dots, U_d . Let $m = \text{rank } H$ be the degeneracy index of D and let $\{p_1, p_2, \dots, p_d\}$ be a basis for \mathbb{Z}^d as above. Consider the compact group*

$$\Lambda := \mathbb{Z}_{a_1} \times \dots \times \mathbb{Z}_{a_m} \times \mathbb{T}^{d-m} \subset \mathbb{T}^d,$$

where \mathbb{Z}_{a_j} is the cyclic group of order a_j viewed as the subgroup of \mathbb{T} generated by a primitive a_j^{th} root of unity. Then there is a continuous action γ' of \mathbb{T}^d on \mathcal{A}_D satisfying $\gamma'_\lambda(U_{p_i}) = \lambda_i U_{p_i}$ for all $\lambda = (\lambda_1, \dots, \lambda_d) \in \mathbb{T}^d$ and such that if we let γ^Λ denote the restriction of γ' to Λ , \mathcal{A}_D^Λ denote the corresponding fixed point algebra, and $Z(\mathcal{A}_D)$ denote the center of \mathcal{A}_D , then

$$C^*(U_b \mid b \in H) = \mathcal{A}_D^\Lambda = Z(\mathcal{A}_D). \quad (4.3)$$

Proof. For each $x = (x_1, x_2, \dots, x_d) \in \mathbb{Z}^d$ we set $U_x := U_1^{x_1} U_2^{x_2} \dots U_d^{x_d}$. Let $B \in \text{GL}_d(\mathbb{Z})$ be the matrix whose i -th column is p_i . Since

$$U_{p_i} U_{p_j} = e^{-2\pi i \langle p_i \mid D p_j \rangle} U_{p_j} U_{p_i} = e^{-2\pi i \langle B e_i \mid D B e_j \rangle} U_{p_j} U_{p_i} = e^{-2\pi i (B^T D B)_{i,j}} U_{p_j} U_{p_i}$$

there is a homomorphism $\pi_B: \mathcal{A}_{B^T D B} \rightarrow \mathcal{A}_D$ that sends the canonical generator $\tilde{U}_i \in \mathcal{A}_{B^T D B}$ to $U_{p_i} \in \mathcal{A}_D$ for $i = 1, \dots, d$. This is in fact an isomorphism with inverse given by the map $\pi_{B^{-1}}: \mathcal{A}_D \rightarrow \mathcal{A}_{B^T D B}$ that sends the generator U_i to $\tilde{U}_{B^{-1} e_i}$, $i = 1, \dots, d$ (see [28, Remark 1.2]). The action $\gamma': \mathbb{T}^d \rightarrow \text{Aut}(\mathcal{A}_D)$ as in the statement of the lemma can now be defined by using the isomorphism π_B to conjugate the canonical gauge action of \mathbb{T}^d on $\mathcal{A}_{B^T D B}$ into an action on \mathcal{A}_D . This proves the first assertion.

In order to prove the first equality in (4.3), observe that $b \in H$ iff $b = x_1 a_1 p_1 + \dots + x_m a_m p_m$ for some $x_1, \dots, x_m \in \mathbb{Z}$. So the product $U_b = U_1^{b_1} \dots U_d^{b_d}$ can be rearranged using the cocycle to yield

$$U_b = \alpha U_{p_1}^{a_1 x_1} \dots U_{p_m}^{a_m x_m}$$

for some scalar $\alpha \in \mathbb{T}$. Hence $C^*(U_b \mid b \in H) = C^*(U_{p_i}^{a_i} \mid i \in \{1, \dots, m\})$.

Next, recall that the action γ' is determined by

$$\gamma'_\lambda(U_{p_1}^{n_1} U_{p_2}^{n_2} \dots U_{p_d}^{n_d}) = (\lambda_1^{n_1} \lambda_2^{n_2} \dots \lambda_d^{n_d}) U_{p_1}^{n_1} U_{p_2}^{n_2} \dots U_{p_d}^{n_d} \quad (\lambda \in \mathbb{T}^d).$$

So let γ^Λ be the restriction of γ' to Λ . If $E^\Lambda: \mathcal{A}_D \rightarrow \mathcal{A}_D^\Lambda$ denotes the conditional expectation obtained by averaging over Λ , then

$$E^\Lambda(U_{p_1}^{n_1} U_{p_2}^{n_2} \dots U_{p_d}^{n_d}) = \begin{cases} U_{p_1}^{n_1} U_{p_2}^{n_2} \dots U_{p_d}^{n_d} & \text{if } \lambda_1^{n_1} \lambda_2^{n_2} \dots \lambda_d^{n_d} = 1 \text{ for all } \lambda \in \Lambda, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly $\lambda_1^{n_1} \lambda_2^{n_2} \dots \lambda_d^{n_d} = 1$ for every $\lambda \in \Lambda$ iff $n_j = 0 \pmod{a_j}$ for $1 \leq j \leq m$ and $n_j = 0$ for $m+1 \leq j \leq d$, that is, iff $U_{p_1}^{n_1} U_{p_2}^{n_2} \dots U_{p_d}^{n_d} = U_{p_1}^{a_1 x_1} \dots U_{p_m}^{a_m x_m}$ for some $x_1, \dots, x_m \in \mathbb{Z}$. This implies that $C^*(U_b \mid b \in H) = \mathcal{A}_D^\Lambda$ because E^Λ is a contraction and products of the form $U_{p_1}^{n_1} U_{p_2}^{n_2} \dots U_{p_d}^{n_d}$ have dense linear span in \mathcal{A}_D .

For the second equality in (4.3), take $b, c \in \mathbb{Z}^d$. Then $U_b U_c = e^{2\pi i \langle b \mid \Theta_d c \rangle} U_c U_b$, and hence $U_b \in Z(\mathcal{A}_D)$ if and only if $b \in H$. This shows that $\mathcal{A}_D^\Lambda \subset Z(\mathcal{A}_D)$ because we have already shown that $C^*(U_b \mid b \in H) = \mathcal{A}_D^\Lambda$. In order to establish the reverse inclusion, we use the spectral

subspaces of γ^Λ . Since Λ is a closed subgroup of \mathbb{T}^d , its dual group $\hat{\Lambda}$ is the quotient of \mathbb{Z}^d by the annihilator of Λ . If we use the duality pairing of \mathbb{T}^d with \mathbb{Z}^d given by $\langle z, a \rangle = z^a = \prod_{j=1}^d z_j^{a_j}$, then $\Lambda^\perp = \prod_{j=1}^m a_j \mathbb{Z} \times \{0\}^{d-m}$ and we have a natural identification $\hat{\Lambda} \cong (\mathbb{Z}/a_1 \mathbb{Z}) \times \cdots \times (\mathbb{Z}/a_m \mathbb{Z}) \times \mathbb{Z}^{d-m}$. Explicitly, denoting by \bar{b} the image of $b \in \mathbb{Z}^d$ in $(\mathbb{Z}/a_1 \mathbb{Z}) \times \cdots \times (\mathbb{Z}/a_m \mathbb{Z}) \times \mathbb{Z}^{d-m}$, the corresponding duality pairing between Λ and $\hat{\Lambda}$ is given by $\langle \lambda, \bar{b} \rangle := \lambda^{\bar{b}} = \lambda_1^{b_1} \lambda_2^{b_2} \cdots \lambda_d^{b_d}$ for $\lambda \in \Lambda$. The corresponding spectral projection $E_{\bar{b}}^\Lambda$ is given by

$$E_{\bar{b}}^\Lambda(x) := \int_{\Lambda} \lambda^{-\bar{b}} \gamma_\lambda^\Lambda(x) d\lambda \quad (x \in \mathcal{A}_D),$$

and its range is the \bar{b} -th spectral subspace

$$E_{\bar{b}}^\Lambda(\mathcal{A}_D) = \overline{\text{span}}\{U_{p_1}^{c_1} \cdots U_{p_m}^{c_m} U_{p_{m+1}}^{b_{m+1}} \cdots U_{p_d}^{b_d} \mid c_i - b_i \in a_i \mathbb{Z} \text{ for all } 1 \leq i \leq m\}.$$

Suppose $x \in Z(\mathcal{A}_D)$. Then $\gamma_\lambda^\Lambda(x) \in Z(\mathcal{A}_D)$ for every λ , and hence also $E_{\bar{b}}^\Lambda(x) \in Z(\mathcal{A}_D)$. Thus

$$E_{\bar{b}}^\Lambda(x) U_{p_i} = e^{2\pi i \langle \sum_{j=1}^d b_j p_j \mid D p_i \rangle} U_{p_i} E_{\bar{b}}^\Lambda(x) = e^{2\pi i \langle \sum_{j=1}^d b_j p_j \mid D p_i \rangle} E_{\bar{b}}^\Lambda(x) U_{p_i}$$

for every $i = 1, 2, \dots, d$. We deduce that $E_{\bar{b}}^\Lambda(x) = 0$ when $\bar{b} \neq 0$. This implies $x = E^\Lambda(x)$ (see, for example, [12, Proposition 17.13]), establishing the inclusion $Z(\mathcal{A}_D) \subset E^\Lambda(\mathcal{A}_D) = \mathcal{A}_D^\Lambda$, and proving the second equality in (4.3). \square

Remark 4.5. Notice that the matrix $D' = B^T D B$ from the proof of Lemma 4.4 has the form

$$D' = \left[\begin{array}{c|c} D'_{m \times m} & D'_{m \times (d-m)} \\ \hline D'_{(d-m) \times m} & D'_{(d-m) \times (d-m)} \end{array} \right],$$

where $D'_{(d-m) \times (d-m)}$ is nondegenerate and the remaining entries of D' are integers. Therefore, \mathcal{A}_D is isomorphic to a d -dimensional noncommutative torus $\mathcal{A}_{D'}$ whose center $Z(\mathcal{A}_{D'})$ is generated by powers of the first m canonical unitary generators of $\mathcal{A}_{D'}$. The last $d - m$ canonical unitaries generate a simple C^* -subalgebra of $\mathcal{A}_{D'}$.

Proposition 4.6. *Let $E^\Lambda: \mathcal{A}_D \rightarrow Z(\mathcal{A}_D)$ be the canonical conditional expectation associated to the action γ^Λ from Lemma 4.4. Then the map $\omega \mapsto \omega \circ E^\Lambda$ is an affine homeomorphism of the state space of $Z(\mathcal{A}_D)$ onto the space of tracial states of \mathcal{A}_D .*

Proof. Clearly $\omega \mapsto \omega \circ E^\Lambda$ is an affine bijection of states of the center to the set of states that factor through E^Λ . The inverse of this map is simply the restriction of states. We first show every tracial state of \mathcal{A}_D factors through the conditional expectation E^Λ . Let τ be a trace of \mathcal{A}_D and let $b \in \mathbb{Z}^d$ be such that $b \notin H$. Let $U_b = U_1^{b_1} U_2^{b_2} \cdots U_d^{b_d}$ and take $c \in \mathbb{Z}^d$ so that $\langle b \mid Dc \rangle \notin \mathbb{Z}$. Then

$$\tau(U_b) = \tau(U_b U_c U_c^*) = \tau(U_c^* U_b U_c) = e^{-2\pi i \langle b \mid Dc \rangle} \tau(U_b).$$

So we must have $\tau(U_b) = 0$ and hence $\tau = (\tau \upharpoonright_{Z(\mathcal{A}_D)}) \circ E^\Lambda$ since $C^*(U_b \mid b \in H) = Z(\mathcal{A}_D)$ by Lemma 4.4.

Next we show that $\omega \circ E^\Lambda$ is indeed a trace of \mathcal{A}_D for every state ω of $Z(\mathcal{A}_D)$. Recall from (2.5) that \mathcal{A}_D is generated by a universal projective unitary representation \bar{v} of \mathbb{Z}^d with cocycle σ_D . Since $\{\bar{v}_b \mid b \in \mathbb{Z}^d\}$ has dense linear span, it suffices to show that the commutator of any two of these elements lies in the kernel of E^Λ . Let $b, c \in \mathbb{Z}^d$. Then

$$E^\Lambda(\bar{v}_b \bar{v}_c - \bar{v}_c \bar{v}_b) = (\sigma_D(b, c) - \sigma_D(c, b)) E^\Lambda(\bar{v}_{b+c})$$

obviously vanishes when $b + c \notin H$ because then $E^\Lambda(\bar{v}_{b+c}) = 0$. Assume now that $b + c \in H$. Then

$$1 = e^{-2\pi i \langle b+c \mid Dc \rangle} = \sigma_D(b+c, c)^2 = \sigma_D(b+c, c) \overline{\sigma_D(c, b+c)}.$$

Hence $\sigma_D(b+c, c) = \sigma_D(c, b+c)$, which implies $\sigma_D(b, c) = \sigma_D(c, b)$. This shows that $E^\Lambda(\bar{v}_b \bar{v}_c - \bar{v}_c \bar{v}_b)$ also vanishes when $b+c \notin H$, and thus $\omega \circ E^\Lambda$ is a trace of \mathcal{A}_D . \square

We can now show that the center of \mathcal{A}_D is isomorphic to a torus of dimension $m = \text{rank } H$. This characterisation can also be derived from the proof of [10, Lemma 2.3].

Proposition 4.7. *Let $D \in M_d(\mathbb{R})$ be an antisymmetric matrix and let \mathcal{A}_D be the associated noncommutative torus, with canonical unitary generators U_1, \dots, U_d . Let $m = \text{rank } H$ and let $\{p_1, \dots, p_d\}$ be a basis for \mathbb{Z}^d as in Lemma 4.4. Then there is an isomorphism $C(\mathbb{T}^m) \cong Z(\mathcal{A}_D)$ that sends z_j to $U_{p_j}^{a_j}$ for $j = 1, \dots, m$, where $z_j: \mathbb{T}^m \rightarrow \mathbb{T} \hookrightarrow \mathbb{C}$ is the projection onto the j -th coordinate.*

Proof. Since $\{U_{p_i}^{a_i} \mid i = 1, 2, \dots, m\}$ is a commuting family of m unitaries generating $Z(\mathcal{A}_D)$, there is a canonical surjective homomorphism $\pi_H: C(\mathbb{T}^m) \rightarrow Z(\mathcal{A}_D)$ such that $\pi_H(z_j) = U_{p_j}^{a_j}$ for $j = 1, \dots, m$.

Now let $a = (a_1, a_2, \dots, a_m) \in \mathbb{Z}^m$ and consider the action of \mathbb{T}^m on $C(\mathbb{T}^m)$ given by the composite of the translation action with the group homomorphism $\nu \mapsto \nu^a = \prod_{j=1}^m \nu_j^{a_j}$. Let γ' be the action of \mathbb{T}^d on \mathcal{A}_D as in Lemma 4.4. Identify \mathbb{T}^m with the closed subgroup $\mathbb{T}^m \times 1_{d-m}$ of \mathbb{T}^d and notice that for all $\nu, \lambda \in \mathbb{T}^m$, one has $z_j(\nu^a \lambda) = \nu_j^{a_j} \lambda_j = \nu_j^{a_j} z_j(\lambda)$ and also $\gamma'_\nu(U_{p_j}^{a_j}) = \nu_j^{a_j} U_{p_j}^{a_j}$. Thus the homomorphism π_H is \mathbb{T}^m -equivariant with respect to the action on $C(\mathbb{T}^m)$ described above and the action on $Z(\mathcal{A}_D)$ obtained by restricting γ' to $Z(\mathcal{A}_D)$ and then to \mathbb{T}^m . By [11, Proposition 2.9], we deduce that π_H is injective, and thus an isomorphism. \square

Remark 4.8. We believe that the description of the center $Z(\mathcal{A}_D)$ in Proposition 4.7 and the characterisation of tracial states of \mathcal{A}_D in Proposition 4.6 are known to experts. Since we were not able to find an explicit source, we have included precise statements and detailed proofs for completeness and ease of reference.

5. MAIN RESULTS

When we combine the results of the preceding two sections we obtain our main theorem, which is a parametrisation of KMS_β states in terms of states on the center of the noncommutative torus \mathcal{A}_{Θ_d} . By Proposition 4.7, this center is isomorphic to $C(\mathbb{T}^m)$, where m is the degeneracy index of Θ_d . Hence the simplex of KMS_β states is affinely weak* homeomorphic to the space of probability measures on the classical torus \mathbb{T}^m .

Theorem 5.1. *Suppose that $r = (r_1, \dots, r_k, 0, \dots, 0) \in \mathbb{R}^n$ is a vector with strictly positive first k coordinates and that*

$$\Theta = \left[\begin{array}{c|c} \Theta_k & \Lambda \\ \hline -\Lambda^T & \Theta_d \end{array} \right]$$

is an antisymmetric $n \times n$ real matrix with diagonal blocks Θ_k and Θ_d of sizes $k \times k$ and $d \times d$, respectively. Let m be the degeneracy index of Θ_d . Then there is an affine weak homeomorphism of the space $M_1(\mathbb{T}^m)$ of probability measures on \mathbb{T}^m onto the space of KMS_β states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha^r)$. Specifically, if $\{p_1, \dots, p_m\}$ is a basis for \mathbb{Z}^d such that $\{a_1 p_1, \dots, a_m p_m\}$ is a basis for $H = \{x \in \mathbb{Z}^d \mid \langle x \mid \Theta_d y \rangle \in \mathbb{Z} \text{ for all } y \in \mathbb{Z}^d\}$ as in Lemma 4.4, then the affine homeomorphism can be chosen so that the extremal KMS_β state $\varphi_{\beta, z}$ associated to the unit point mass at $z \in \mathbb{T}^m$ is given by*

$$\varphi_{\beta, z}(L_p L_x L_y^* L_q^*) = \delta_{p, q} [x - y \in H] \lambda_{x-y} z^c \prod_{j=1}^k \frac{e^{-\beta r_j p_j} (1 - e^{-\beta r_j})}{1 - e^{-\beta r_j + 2\pi i (\Theta(x-y) \mid e_j)}}$$

where $c = (c_1, \dots, c_m)$ is the vector of coefficients of $x-y$ with respect to the basis $\{a_1 p_1, \dots, a_m p_m\}$ of H and $\lambda_{x-y} \in \{-1, 1\}$ is such that $\pi(L_x L_y^) = \lambda_{x-y} U_{a_1 p_1}^{c_1} \dots U_{a_m p_m}^{c_m}$ in \mathcal{A}_{Θ_d} .*

Proof. By Lemma 4.3, the homomorphism $\pi: \mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d}) \rightarrow \mathcal{A}_{\Theta_d}$ that sends the isometry L_{e_j} to the unitary U_j yields a weak* homeomorphism between tracial state spaces via $\tau \mapsto \tau \circ \pi$. By Proposition 4.6, the map that sends a state ω of the center $Z(\mathcal{A}_{\Theta_d})$ to the tracial state $\omega \circ E^\Lambda$ of \mathcal{A}_{Θ_d} is an affine weak* homeomorphism. Combining these maps we conclude that given an isomorphism $\phi: Z(\mathcal{A}_{\Theta_d}) \xrightarrow{\cong} C(\mathbb{T}^m)$, each probability measure μ on \mathbb{T}^m gives rise to a tracial state $\omega_\mu \circ E^\Lambda \circ \pi$ of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$, where

$$\omega_\mu(b) = \int_{\mathbb{T}^m} \phi(b) d\mu \quad (b \in Z(\mathcal{A}_{\Theta_d})).$$

Such an isomorphism $\phi: Z(\mathcal{A}_{\Theta_d}) \xrightarrow{\cong} C(\mathbb{T}^m)$ always exists by Proposition 4.7, and thus it follows from Proposition 4.2 that the space of probability measures on \mathbb{T}^m and the KMS_β simplex of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha^r)$ are affinely weak* homeomorphic.

Suppose that the isomorphism $\phi: Z(\mathcal{A}_{\Theta_d}) \xrightarrow{\cong} C(\mathbb{T}^m)$ in question is the one coming from the basis $\{a_1 p_1, \dots, a_m p_m\}$ for H built in Proposition 4.7. When we let μ be the unit point mass at $z \in \mathbb{T}^m$ and let $\omega_z \circ E^\Lambda \circ \pi$ play the role of τ in Proposition 4.2, we see from (4.2) that the weak* homeomorphism described above gives us precisely a extremal KMS_β state $\varphi_{\beta, z}$ given by the formula in the statement of the theorem. That the scalar $\lambda_{x-y} \in \mathbb{T}$ such that $\pi(L_x L_y^*) = \lambda_{x-y} U_{a_1 p_1}^{c_1} U_{a_2 p_2}^{c_2} \dots U_{a_m p_m}^{c_m}$ in \mathcal{A}_{Θ_d} is either 1 or -1 for $x - y \in H$ follows because $\langle x | \Theta_d y \rangle = \langle x - y | \Theta_d y \rangle \in \mathbb{Z}$, and as in (2.5) we have

$$\lambda_{x-y} = e^{\pi i \langle x | \Theta_d y \rangle} e^{-\pi i \langle (x-y) - c_1 a_1 p_1 | \Theta_d c_1 a_1 p_1 \rangle} \dots e^{-\pi i \langle c_m a_m p_m | \Theta_d c_m a_m p_m \rangle}. \quad \square$$

As consequences of Theorem 5.1 for particular choices of Θ we get the following two corollaries. The first one addresses the question of when these systems have unique equilibrium at each β .

Corollary 5.2. *If $r > 0$ or if Θ_d is nondegenerate, then the system $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha^r)$ has a unique KMS_β state for each $\beta > 0$.*

At the other extreme, the second one gives conditions under which the phase transition is maximal, recovering the phase transition result for the building blocks from [2].

Corollary 5.3. *If $\Theta_d \in M_d(\mathbb{Z})$, then $m = d$ and the simplex of KMS_β states is parametrised by the probability measures on the d -torus.*

6. EQUILIBRIUM AT $\beta = \infty$ AND $\beta = 0$

In this section we study equilibrium at the extremal inverse temperature values ∞ and 0 for the system $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha^r)$. As before, Θ is an antisymmetric $n \times n$ real matrix and $r = (r_1, \dots, r_k, 0_d)$ is a vector in $[0, \infty)^n$ with strictly positive first k coordinates.

6.1. Ground states and KMS_∞ states. A state φ of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ is a *ground state* for α if and only if the function $z \mapsto \varphi(A \alpha_z(B))$ is bounded on the upper half plane for all analytic elements $A, B \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. See, for example, [4, Proposition 5.3.19] and [26, Proposition 8.12.3]. Although initially ground states were also called, indistinctly, ∞ -KMS states, relatively recent results on phase transitions at $\beta = \infty$ related to Bost–Connes systems have required that a distinction be made between generic ground states as defined above and those that can be obtained as limits of KMS_β states for $\beta \rightarrow \infty$. See [7, Definition 3.7] for the definitions and [22, Theorem 7.1] for an example in which the two concepts differ. As a result, it has become customary to reserve the terminology "KMS $_\infty$ states" for the aforementioned limits.

We wish to compute first the ground states and the KMS_∞ states of our systems, and we begin with the following characterisation of ground states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$.

Proposition 6.1. *Let $n = k + d$ with $k, d \in \mathbb{N}$ and let $r = (r_1, \dots, r_k, 0_d)$ with $r_j > 0$ for $j \leq k$. Let α be the dynamics determined by r and suppose that φ is a state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. Then φ is a ground state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$ if and only if for all $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$, one has*

$$\varphi(L_p L_x L_y^* L_q^*) = 0 \text{ unless } p = q = 0. \quad (6.1)$$

Proof. Suppose first that φ is a ground state and let $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$. Then the function

$$z \mapsto \varphi(L_p L_x L_y^* \alpha_z(L_q^*)) = e^{-i\langle q|r \rangle z} \varphi(L_p L_x L_y^* L_q^*)$$

is bounded on the upper half plane, and so either $q = 0$ or $\varphi(L_p L_x L_y^* L_q^*) = 0$. Taking adjoints shows that either $p = 0$ or $\varphi(L_p L_x L_y^* L_q^*) = 0$. This proves that ground states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$ satisfy (6.1).

Suppose now that φ is a state such that (6.1) holds for all $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$. In order to conclude that φ is a ground state, it suffices to show that the function $\varphi(X \sigma_z(Y))$ is bounded on the upper half plane for every $X = L_p L_x L_y^* L_q^*$ and $Y = L_a L_s L_t^* L_b^*$ with $p, q, a, b \in \mathbb{N}^k \times 0_d$ and $x, y, s, t \in 0_k \times \mathbb{N}^d$, because the elements of this type are analytic, α -invariant, and have a dense linear span in $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. Using Definition 2.6(3) we see that $XY = \lambda L_{p-q \vee a} L_x L_y^* L_s L_t^* L_{b-a+q \vee a}^*$ for some $\lambda \in \mathbb{T}$, so (6.1) yields

$$\varphi(X \alpha_z(Y)) = e^{i\langle a-b|r \rangle z} \varphi(XY) = \begin{cases} e^{i\langle q \vee a|r \rangle z} \lambda \varphi(L_x L_y^* L_s L_t^*) & \text{if } q - p = a - b = q \vee a, \\ 0 & \text{otherwise.} \end{cases}$$

Since $\langle q \vee a|r \rangle \geq 0$ the function $z \mapsto \varphi(X \alpha_z(Y))$ is bounded on the upper half plane. This shows that φ is a ground state for $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$ and completes the proof of the proposition. \square

Next we see that ground states are supported on the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$.

Lemma 6.2. *Let Q be the projection from (3.5) and suppose that φ is a ground state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Then*

$$\varphi(L_x L_y^*) = \varphi(Q L_x L_y^* Q), \quad (x, y \in 0_k \times \mathbb{N}^d).$$

Proof. Let φ be a ground state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Using the expansion formula for Q given in (3.7) we have

$$\varphi(Q L_x L_y^* Q) = \varphi(L_x L_y^* Q) = \sum_{J \subseteq \{1, \dots, k\}} (-1)^{|J|} \varphi(L_x L_y^* \prod_{j \in J} (L_{e_j} L_{e_j}^*)). \quad (6.2)$$

Since φ is a ground state, it follows from Proposition 6.1 that $\varphi(L_p L_x L_y^* L_q^*) = 0$ whenever $p, q \in \mathbb{N}^k \times 0_d$ are not both zero. So the terms for $J \neq \emptyset$ in the sum on the right hand side of (6.2) will vanish. Hence $\varphi(Q L_x L_y^* Q) = \varphi(L_x L_y^*)$ as wished. \square

Proposition 6.3. *Let Q be the projection from (3.5). For each state ω of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$, define*

$$T_\infty(\omega)(X) := \omega(QXQ), \quad (X \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)).$$

Then

- (1) T_∞ is an affine weak* homeomorphism of the state space of $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ onto the ground state space of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$;
- (2) T_∞ restricted to tracial states is an affine weak* homeomorphism of the tracial state space of $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ onto the KMS_∞ state space of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$.

Proof. For part (1), take a state ω of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$. Clearly $T_\infty(\omega)$ is a state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. Also, Lemma 3.4(1) implies that for $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$

$$T_\infty(\omega)(L_p L_x L_y^* L_q^*) = \omega(Q L_p L_x L_y^* L_q^* Q)$$

vanishes unless $p = 0 = q$. Hence $T_\infty(\omega)$ satisfies equation (6.1) of Proposition 6.1 and so it is a ground state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. This shows that T_∞ maps the state space of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ into the ground state space of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Clearly T_∞ is a continuous affine map and it is also injective because ω can be recovered from $T_\infty(\omega)$ by restricting it to the corner.

In order to prove that T_∞ is surjective, suppose φ is a ground state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Let ω_φ be the restriction of φ to the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$, that is,

$$\omega_\varphi(X) := \varphi(X) \quad (X \in Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q).$$

It follows from Lemma 6.2 that $\omega_\varphi(Q) = \varphi(Q) = 1$, and so ω_φ is a state of $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$. We claim that $T_\infty(\omega_\varphi) = \varphi$. To see this, it suffices to show that $T_\infty(\omega_\varphi)(X) = \varphi(X)$ for $X = L_p L_x L_y^* L_q^*$, where $p, q \in \mathbb{N}^k \times 0_d$ and $x, y \in 0_k \times \mathbb{N}^d$. Since φ and $T_\infty(\omega_\varphi)$ are ground states, $T_\infty(\omega_\varphi)(X) = \varphi(X) = 0$ when p and q are not both zero by Proposition 6.1. Suppose now $p = q = 0$. Then

$$\varphi(L_x L_y^*) = \varphi(Q L_x L_y^* Q) = \omega_\varphi(Q L_x L_y^* Q) = T_\infty(\omega_\varphi)(L_x L_y^*),$$

where the first equality follows from Lemma 6.2. This shows that $\varphi = T_\infty(\omega_\varphi)$, proving that T_∞ is surjective. Therefore T_∞ is an affine weak* homeomorphism as asserted, finishing the proof of part (1).

For part (2) it suffices to verify that T_∞ maps the tracial state space of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ onto the space of KMS_∞ states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Suppose first ω is a tracial state of the corner $Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ and let $(\beta_j)_{j \in \mathbb{N}}$ be any sequence of positive real numbers such that $\lim_{j \rightarrow \infty} \beta_j = \infty$. By Proposition 3.6 each $T_{\beta_j}(\omega)$ is a KMS_{β_j} state for α . Since $Q L_p = 0$ for $p \in \mathbb{N}^k \times 0_d \setminus \{0\}$ by Lemma 3.4(1), for every $X \in Q\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)Q$ we have

$$T_{\beta_j}(\omega)(X) = \frac{1}{Z(\beta_j)} \sum_{p \in \mathbb{N}^k \times 0_d} e^{-\beta_j \langle p | r \rangle} \omega(Q L_p^* (Q X Q) L_p Q) = \frac{1}{Z(\beta_j)} \omega(X).$$

Since $T_{\beta_j}(\omega)(Q)^{-1} = Z(\beta_j) \rightarrow 1$ by Lemma 3.5, see (3.8), this shows that $T_\infty(\omega)$ is KMS_∞ .

Suppose now that φ is a KMS_∞ state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. Then its restriction ω_φ to the corner is a limit of tracial states and satisfies $\omega_\varphi(Q) = \lim_j Z(\beta_j)^{-1} = 1$, so it is a tracial state itself. Hence $\varphi = T_\infty(\omega_\varphi)$ lies in the range of the restriction of T_∞ to tracial states. This completes the proof of part (2). \square

6.2. Invariant traces and KMS_{0+} states. When $\beta = 0$ the usual KMS_β condition for a C^* -dynamical system (A, α) just says that the state has to be a trace, cf. [4, Definition 5.3.1]. Nevertheless, in many applications and equivalent formulations it is necessary to add the extra assumption of α -invariance. See, for example, [4, Theorem 5.3.22]. In fact, this assumption is often included in the definition, so that the KMS_0 states are the α -invariant traces on A , cf. [26, 8.12.2].

Here we would like to draw a parallel to the distinction of KMS_∞ states among all ground states by defining KMS_{0+} states to be the weak* limits of KMS_β states as $\beta \rightarrow 0^+$. By [4, Proposition 5.3.25] we see that KMS_{0+} states are traces; in fact, they are also KMS_0 states because α -invariance is preserved under weak* limits.

By taking limits in the formula for KMS_β states obtained in Proposition 4.2 we can give a complete description of the space of KMS_{0+} states of our system $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$.

Proposition 6.4. *Identify $C^*(L_x \mid x \in 0_k \times \mathbb{N}^d)$ with $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ canonically and let φ be a state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. Then φ is a KMS_{0+} state of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$ if and only if it is a trace and satisfies*

$$\varphi(L_p L_x L_y^* L_q^*) = \delta_{p,q} \varphi(L_x L_y^*) \quad p, q \in \mathbb{N}^k \times 0_d, x, y \in 0_k \times \mathbb{N}^d.$$

As a consequence, the map that sends a KMS_{0+} state φ of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$ to its restriction to $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ is an affine weak homeomorphism from the space of KMS_{0+} states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$*

onto the space of the tracial states τ of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ such that for all $x, y \in 0 \times \mathbb{N}^d \cong \mathbb{N}^d$

$$\tau(L_x L_y^*) = 0 \quad \text{unless} \quad \langle \Theta(x-y) | e_j \rangle \in \mathbb{Z} \quad \text{for } 1 \leq j \leq n. \quad (6.3)$$

Proof. Clearly if φ is a KMS_{0+} state, then it is a trace and satisfies $\varphi(L_p L_x L_y^* L_q^*) = \delta_{p,q} \varphi(L_x L_y^*)$. In order to prove the converse and the second part in the statement, we begin by considering the individual ratios from (4.1), namely

$$\frac{e^{-\beta r_j p_j} (1 - e^{-\beta r_j})}{1 - e^{-\beta r_j + 2\pi i \langle \Theta(x-y) | e_j \rangle}},$$

for each $x, y \in 0_k \times \mathbb{Z}^d$ and each $j = 1, 2, \dots, k$. As $\beta \rightarrow 0$, the numerator tends to 0 while the denominator tends to $1 - e^{2\pi i \langle \Theta(x-y) | e_j \rangle}$. Thus the j -th ratio tends to zero unless $\langle \Theta(x-y) | e_j \rangle \in \mathbb{Z}$, in which case the limit is clearly 1. Hence

$$\lim_{\beta \rightarrow 0^+} \prod_{j=1}^k \frac{e^{-\beta r_j p_j} (1 - e^{-\beta r_j})}{1 - e^{-\beta r_j + 2\pi i \langle \Theta(x-y) | e_j \rangle}} = \begin{cases} 1 & \text{if } \langle \Theta(x-y) | e_j \rangle \in \mathbb{Z} \quad 1 \leq j \leq k \\ 0 & \text{otherwise.} \end{cases} \quad (6.4)$$

So let φ be a state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})$ and suppose that

$$\varphi(L_p L_x L_y^* L_q^*) = \delta_{p,q} \varphi(L_x L_y^*) = \delta_{p,q},$$

for all $p, q \in \mathbb{N}^k \times 0_d$, and $x, y \in 0_k \times \mathbb{N}^d$. Set $\tau := \varphi \upharpoonright_{\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})}$. Then τ is a trace of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$, and because it is the restriction of a trace of $\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})$, it satisfies $\tau(L_x L_y^*) = 0$ unless $\langle \Theta(x-y) | e_j \rangle = 0$ for $j = 1, \dots, n$ by Proposition 4.6. Thus τ satisfies (6.3). Take a sequence $(\beta_n)_{n \in \mathbb{N}}$ of positive real numbers converging to 0. Taking limits in formula (4.1) for $T_{\beta_n}(\tau \circ \rho_Q^{-1})$ and using (6.4), we see that the weak* limit of $(T_{\beta_n}(\tau \circ \rho_Q^{-1}))_{n \in \mathbb{N}}$ is precisely φ . Hence φ is KMS_{0+} state as wished.

Since a KMS_{0+} state φ of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta}), \alpha)$ satisfies

$$\varphi(L_p L_x L_y^* L_q^*) = \delta_{p,q} \varphi(L_x L_y^*),$$

it is determined by its restriction τ to $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$. Also, we deduce from the above that φ is the weak* limit of $(T_{\beta_n}(\tau \circ \rho_Q^{-1}))_{n \in \mathbb{N}}$, and τ is a tracial state of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ that satisfies (6.3). This gives the second part in the statement and finishes the proof of the proposition. \square

Recall from the beginning of Section 3 that the dynamics determined by a vector r on $\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})$ is a subgroup of the gauge action, given by $\alpha_t = \gamma_{e^{i(r)t}}$ for $t \in \mathbb{R}$. With our convention that the first k coordinates of r are strictly positive and the rest are zero, $e^{i(r)t} = (e^{itr_1}, e^{itr_2}, \dots, e^{itr_n}) \in \mathbb{T}^k \times \{1_d\}$. By continuity, a state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_{\Theta})$ is α^r invariant if and only if it is invariant under the action of the closure of $\{(e^{itr_1}, e^{itr_2}, \dots, e^{itr_k}) \mid t \in \mathbb{R}\}$ in \mathbb{T}^k , viewed as a compact subgroup of $\mathbb{T}^k \times \{1_d\}$.

In order to determine the KMS_0 states, namely the α -invariant traces, we consider the group

$$H_n^r := \{w \in \mathbb{Z}^n \mid \langle w | \Theta v \rangle \in \mathbb{Z} \text{ for all } v \in \mathbb{Z}^n\} \cap \{w \in \mathbb{Z}^n \mid \langle w | r \rangle = 0\}$$

and its annihilator

$$\Lambda_n^r := \{z \in \mathbb{T}^n \mid z^h = \prod_{j=1}^n z_j^{h_j} = 1 \text{ for all } h \in H_n^r\}.$$

Let U_1, \dots, U_n be the canonical unitary generators of \mathcal{A}_{Θ} . The conditional expectation obtained from averaging with respect to the restriction of the gauge action on \mathcal{A}_{Θ} to Λ_n^r satisfies

$$E^{\Lambda_n^r}(U_v) = \begin{cases} U_v & \text{if } v \in H_n^r \\ 0 & \text{otherwise,} \end{cases} \quad (6.5)$$

where $U_v = U_1^{v_1} \dots U_n^{v_n}$ with $v = (v_1, \dots, v_n) \in \mathbb{Z}^n$.

Proposition 6.5. *The map $\omega \mapsto \omega \circ E_n^{\Lambda_r} \circ \pi$ is an affine weak* homeomorphism of the state space of the fixed point algebra of the action of Λ_n^r (which is the α -invariant part of $Z(\mathcal{A}_\Theta)$) and the space of KMS_0 states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$. If H_n^r has nonzero intersection with $\mathbb{Z}^k \times 0_d$, then the space of KMS_{0+} states is properly contained in the space of KMS_0 states.*

Proof. Notice that the range of the conditional expectation $E_n^{\Lambda_r}$ on \mathcal{A}_Θ is simply

$$\overline{\text{span}}\{U_v \mid v \in H_n^r\} = \overline{\text{span}}\{U_v \mid \Theta v \in \mathbb{Z}^n\} \cap \overline{\text{span}}\{U_w \mid \langle w \mid r \rangle = 0\}.$$

In fact, one can easily verify using (6.5) that $E_n^{\Lambda_r}$ is the composition of the conditional expectation onto $Z(\mathcal{A}_\Theta)$ with the conditional expectation onto the fixed point algebra of the dynamics on \mathcal{A}_Θ corresponding to α . So by Lemma 4.3 and Proposition 4.6, $\omega \circ E_n^{\Lambda_r} \circ \pi$ is a tracial state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ for every state ω on the fixed point algebra of Λ_n^r . Since $E_n^{\Lambda_r}(U_v) = 0$ unless $\langle v \mid r \rangle = 0$, it is clear that $\omega \circ E_n^{\Lambda_r} \circ \pi$ is α -invariant. To show that the map $\omega \mapsto \omega \circ E_n^{\Lambda_r} \circ \pi$ is also surjective, let ω be a tracial state of \mathcal{A}_Θ such that the corresponding tracial state $\omega \circ \pi$ of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ is α -invariant and let $v, w \in \mathbb{N}^n$. Then for all $t \in \mathbb{R}$ one has

$$(\omega \circ \pi)(L_v L_w^*) = e^{it\langle v-w \mid r \rangle} (\omega \circ \pi)(L_v L_w^*),$$

which implies $(\omega \circ \pi)(L_v L_w^*) = 0$ if $\langle v-w \mid r \rangle \neq 0$. By Proposition 4.6, we also have $(\omega \circ \pi)(L_v L_w^*) = 0$ if $\Theta(v-w) \notin \mathbb{Z}^n$. Hence $\omega \circ \pi = \omega \circ E_n^{\Lambda_r} \circ \pi$ and we then deduce that the map $\omega \mapsto \omega \circ E_n^{\Lambda_r} \circ \pi$ is an affine weak* homeomorphism from the space of states of the fixed point algebra of Λ_n^r onto the space of α -invariant tracial states of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$.

For the last assertion, suppose the group H_n^r is not contained in $0_k \times \mathbb{Z}^d$. Then we can find $v, w \in \mathbb{N}^n$ such that $v-w \in H_n^r$ and the first k coordinates of $v-w$ are not all zero. Write $v = p + x$ with $p \in \mathbb{N}^k \times 0_d$ and $x \in 0_k \times \mathbb{N}^d$ and $w = q + y$ with $q \in \mathbb{N}^k \times 0_d$ and $y \in 0_k \times \mathbb{N}^d$. It follows that $p - q \neq 0$, and $\pi(L_p L_x L_y^* L_q^*)$ is an element of the center $Z(\mathcal{A}_\Theta)$ that is fixed by α , so that $\pi(L_p L_x L_y^* L_q^*) = (E_n^{\Lambda_r} \circ \pi)(L_p L_x L_y^* L_q^*)$. Let ω be a state of the range of $E_n^{\Lambda_r}$ that does not vanish on $\pi(L_p L_x L_y^* L_q^*)$. Then $\omega \circ E_n^{\Lambda_r} \circ \pi$ is an α -invariant tracial state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ that does not vanish on $L_p L_x L_y^* L_q^*$. This gives a tracial state of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ that is not a KMS_{0+} state because $p \neq q$. \square

Example 6.6. Let θ be an irrational number in $(0, 1/2)$ and consider the matrix

$$\Theta = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \theta \\ 0 & 0 & -\theta & 0 \end{bmatrix}. \quad (6.6)$$

Fix $a > 0$ and let α be the dynamics on $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ defined by $r = (a, a, 0, 0)$. Then $k = d = 2$ and the lower right corner $\Theta_d = \begin{bmatrix} 0 & \theta \\ -\theta & 0 \end{bmatrix}$ has degeneracy index 0, so that $\mathcal{A}_{\Theta_d} \cong A_{2\theta}$ is simple and has a unique tracial state. Hence there is a unique KMS_{0+} state. But the center of \mathcal{A}_Θ is nontrivial and, more importantly, it contains nontrivial α -invariant elements, such as $U_1 U_2^*$. As a consequence, there are more α -invariant traces on $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ than just the unique KMS_{0+} state.

7. AN ALTERNATIVE APPROACH VIA PRODUCT SYSTEMS

In this section we realise $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ as the Nica–Toeplitz algebra of a compactly aligned product system over \mathbb{N}^k in which the underlying coefficient algebra is precisely the Toeplitz noncommutative torus $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$, where Θ_d denotes the $d \times d$ bottom-right corner submatrix of Θ , and σ_{Θ_d} is the corresponding 2-cocycle on \mathbb{N}^d . The motivation is to apply the characterisation of KMS_β states of Nica–Toeplitz algebras from [3] and compare it to our parametrisation.

We begin by defining an action of \mathbb{N}^k by endomorphisms of $\mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$, which will then be used to construct a product system over \mathbb{N}^k with the desired properties. In order to lighten the notation, we let $A := \mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ throughout this section.

Lemma 7.1. *Let $\Theta \in M_n(\mathbb{R})$ be an antisymmetric matrix and let $k, d \in \mathbb{N}$ with $n = k + d$. Let Θ_d be the $d \times d$ bottom-right corner submatrix of Θ . Then for each $p \in \mathbb{N}^k$, there is an automorphism $\rho_p: A \rightarrow A$ that sends an isometry $L_x^{\sigma_{\Theta_d}}$ to $\bar{L}_x^{\sigma_{\Theta_d}} := \sigma_{\Theta}(p, x)^2 L_x^{\sigma_{\Theta_d}}$ for all $x \in \mathbb{N}^d$. Moreover, the map $p \mapsto \rho_p$ is an action of \mathbb{N}^k by automorphisms of A .*

Proof. Fix $p \in \mathbb{N}^k$. As in the statement of the lemma, for $x \in \mathbb{N}^d$ we let

$$\bar{L}_x^{\sigma_{\Theta_d}} = \sigma_{\Theta}(p, x)^2 L_x^{\sigma_{\Theta_d}} = e^{-2\pi i \langle p | \Theta x \rangle} L_x^{\sigma_{\Theta_d}} \in A.$$

Because σ_{Θ} is a bicharacter, the map $x \mapsto \bar{L}_x^{\sigma_{\Theta_d}}$ is an isometric σ_{Θ_d} -representation of \mathbb{N}^d in A . Such a representation is Nica covariant since

$$\bar{L}_x^{\sigma_{\Theta_d}} (\bar{L}_x^{\sigma_{\Theta_d}})^* = L_x^{\sigma_{\Theta_d}} (L_x^{\sigma_{\Theta_d}})^*$$

for all $x \in \mathbb{N}^d$, and so $x \mapsto \bar{L}_x^{\sigma_{\Theta_d}}$ is a covariant isometric σ_{Θ_d} -representation. By universal property, we obtain an endomorphism $\rho_p: A \rightarrow A$ mapping $L_x^{\sigma_{\Theta_d}}$ to $\bar{L}_x^{\sigma_{\Theta_d}}$. Its inverse is the endomorphism induced by the covariant isometric σ_{Θ_d} -representation $x \mapsto \sigma_{\Theta}(x, p)^2 L_x^{\sigma_{\Theta_d}}$, and hence ρ_p is in fact an automorphism. Using again that σ_{Θ} is a bicharacter, we deduce that the map $p \mapsto \rho_p \in \text{Aut}(A)$ is a semigroup action $\rho: \mathbb{N}^k \rightarrow \text{Aut}(A)$ on A . This completes the proof of the lemma. \square

Since any automorphism of A is, in particular, an injective endomorphism with hereditary range, the semigroup homomorphism $\rho: \mathbb{N}^k \rightarrow \text{Aut}(A)$ gives rise to a natural product system $A_{\rho} = (A_{\rho_p})_{p \in \mathbb{N}^k}$ over A (see, for example, [31, Example 3.15]). We recall here the structure and operations of A_{ρ} . The correspondence $A_{\rho_p}: A \rightsquigarrow A$ is simply A as a complex vector space. If we write $a\delta_p$ for the element in A_{ρ_p} that is the image of $a \in A$ under the canonical identification, then the A -valued inner product on A_{ρ_p} is given by $\langle a\delta_p | b\delta_p \rangle = \rho_p^{-1}(a^*b)$. The right action of A on A_{ρ_p} is implemented by ρ_p , that is, $a\delta_p \cdot b = a\rho_p(b)\delta_p$ for all $a, b \in A$, while the left action on A_{ρ_p} is simply multiplication of elements in A , so $b \cdot a\delta_p = (ba)\delta_p$. The multiplication map $\mu_{p,q}: A_{\rho_p} \otimes_A A_{\rho_q} \rightarrow A_{\rho_{p+q}}$ is given on an elementary tensor $a\delta_p \otimes b\delta_q$ by $\mu_{p,q}(a\delta_p \otimes b\delta_q) = a\rho_p(b)\delta_{pq}$. With these operations, $A_{\rho} = (A_{\rho_p})_{p \in \mathbb{N}^k}$ is a product system with coefficient algebra A .

Lemma 7.2. *Let $\rho: \mathbb{N}^k \rightarrow \text{Aut}(A)$ be the semigroup action from Lemma 7.1 and let $A_{\rho} = (A_{\rho_p})_{p \in \mathbb{N}^k}$ be the associated product system over \mathbb{N}^k as explained above. Then A_{ρ} is compactly aligned.*

Proof. In order to see that A_{ρ} is compactly aligned, notice that because ρ_p is an automorphism for all $p \in \mathbb{N}^k$, each A_{ρ_p} is an imprimitivity A -bimodule with left A -valued inner product given by $\langle\langle a\delta_p | b\delta_p \rangle\rangle = ab^*$. Thus for all $p, q \in \mathbb{N}^k$ one has

$$\langle\langle A_{\rho_p} | A_{\rho_p} \rangle\rangle \langle\langle A_{\rho_q} | A_{\rho_q} \rangle\rangle = A = \langle\langle A_{\rho_{p \vee q}} | A_{\rho_{p \vee q}} \rangle\rangle.$$

This implies that A_{ρ} is compactly aligned (see also Definition 3.7 and Remark 3.8 of [31]). \square

Let $A_{\rho}^{\sigma} = (A_{\rho_p}^{\sigma})_{p \in \mathbb{N}^k}$ be the product system obtained from A_{ρ} by twisting the multiplication maps with σ_{Θ} . Specifically, for all $p \in \mathbb{N}^k$, $A_{\rho_p}^{\sigma} = A_{\rho_p}$ as an A -correspondence, and the isomorphism $\mu_{p,q}^{\sigma}: A_{\rho_p}^{\sigma} \otimes_A A_{\rho_q}^{\sigma} \rightarrow A_{\rho_{p+q}}^{\sigma}$ is given by $\sigma_{\Theta}(p, q)\mu_{p,q}$, for $p, q \in \mathbb{N}^k$. Notice that the multiplication maps are indeed associative because μ is and σ_{Θ} is a 2-cocycle, and A_{ρ}^{σ} is compactly aligned because A_{ρ} is so.

Proposition 7.3. *Let $A_\rho^\sigma = (A_{\rho_p}^\sigma)_{p \in \mathbb{N}^k}$ be the product system as above. Let $\iota: A \rightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ be the embedding obtained from the canonical isomorphism $A \cong C^*(L_x: x \in 0_k \times \mathbb{N}^d)$ from Lemma 4.1 and identify \mathbb{N}^k with $\mathbb{N}^k \times 0_d$ canonically. Then the map that sends $a\delta_p \in A_{\rho_p}^\sigma$ to $\iota(a)L_p \in \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ induces an isomorphism $\mathcal{N}\mathcal{T}_{A_\rho^\sigma} \cong \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$.*

Proof. For each $p \in \mathbb{N}^k$, let ψ_p denote the map that sends $a\delta_p \in A_{\rho_p}^\sigma$ to $\iota(a)L_p \in A$. We will prove that $\psi = \{\psi_p\}_{p \in \mathbb{N}^k}$ is a Nica covariant representation of A_ρ^σ . We first show that for all $a \in A$ and $p \in \mathbb{N}^k$, we have

$$L_p \iota(a) L_p^* = \iota(\rho_p(a)) L_p L_p^*.$$

Indeed, it suffices to establish this for $a = L_x^{\sigma_\Theta} (L_y^{\sigma_\Theta})^*$ with $x, y \in \mathbb{N}^d$. We compute

$$L_p \iota(a) L_p^* = L_p L_x L_y^* L_p^* = \sigma_\Theta(p, x)^2 \sigma_\Theta(y, p)^2 L_x L_y^* L_p L_p^* = \iota(\rho_p(a)) L_p L_p^*.$$

Now in order to see that ψ preserves the multiplication in A_ρ^σ , we compute for $a, b \in A$ and $p, q \in \mathbb{N}^d$,

$$\begin{aligned} \psi_p(a\delta_p) \psi_q(b\delta_q) &= \iota(a) L_p \iota(b) L_q = \iota(a) L_p \iota(b) L_p^* L_p L_q \\ &= \iota(a) \iota(\rho_p(b)) L_p (L_p^* L_p) L_q \\ &= \sigma_\Theta(p, q) \iota(a \rho_p(b)) L_{p+q} \\ &= \psi_{p+q}(\mu_{p,q}^\sigma(a\delta_p \otimes b\delta_q)), \end{aligned}$$

proving that $\psi = \{\psi_p\}_{p \in \mathbb{N}^k}$ is multiplicative. Finally,

$$\begin{aligned} \psi_p(a\delta_p)^* \psi_p(b\delta_p) &= L_p^* \iota(a^*b) L_p = L_p^* \iota(\rho_p(\rho_{p^{-1}}(a^*b))) L_p \\ &= L_p^* \iota(\rho_p(\rho_p^{-1}(a^*b))) L_p L_p^* L_p \\ &= L_p^* L_p \iota(\rho_p^{-1}(a^*b)) L_p^* L_p \\ &= \iota(\rho_p^{-1}(a^*b)) = \iota(\langle a\delta_p | b\delta_p \rangle). \end{aligned}$$

This shows that ψ also preserves the inner product of $A_{\rho_p}^\sigma$ for each $p \in \mathbb{N}^k$, and so is a representation of A_ρ^σ . That ψ is Nica covariant follows because $L_p L_p^* L_q L_q^* = L_{p \vee q} L_{p \vee q}^*$ in $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ for all $p, q \in \mathbb{N}^k$. Thus $\psi = \{\psi_p\}_{p \in \mathbb{N}^k}$ induces a homomorphism $\tilde{\psi}: \mathcal{N}\mathcal{T}_{A_\rho^\sigma} \rightarrow \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ by universal property, which is clearly surjective.

In order to produce the inverse of $\tilde{\psi}$, we will apply the universal property of $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$. Let $j \in \{1, \dots, n\}$. If $j \leq k$, we set $\bar{L}_{e_j} := \delta_{e_j} \in \mathcal{N}\mathcal{T}_{A_\rho^\sigma}$, and for $j > k$, we let $\bar{L}_{e_j} := L_{e_{j-k}}^{\sigma_\Theta}$. Notice that we have introduced no new notation for the image of A_ρ^σ in $\mathcal{N}\mathcal{T}_{A_\rho^\sigma}$ under the canonical representation. We will show that the set of isometries $\{\bar{L}_{e_j} \mid 1 \leq j \leq n\}$ satisfies the relations (2.6) from Proposition 2.14. Indeed, if $i, j \leq k$, then the multiplication in A_ρ^σ yields

$$\bar{L}_{e_i} \bar{L}_{e_j} = \sigma_\Theta(e_i, e_j)^2 \bar{L}_{e_j} \bar{L}_{e_i} = e^{-2\pi i \theta_{i,j}} \bar{L}_{e_j} \bar{L}_{e_i}.$$

The same will be true for $i, j > k$ because σ_{Θ_d} is simply the restriction of σ_Θ to $(0_k \times \mathbb{N}^d) \times (0_k \times \mathbb{N}^d)$. Now for $i \leq k < j$ or $j \leq k < i$, the relation $\bar{L}_{e_i} \bar{L}_{e_j} = \sigma_\Theta(e_i, e_j)^2 \bar{L}_{e_j} \bar{L}_{e_i}$ holds because of the right action of A on $A_{\rho_{e_l}}$ for each $1 \leq l \leq k$. One can similarly prove that

$$\bar{L}_{e_i}^* \bar{L}_{e_j} = e^{2\pi i \theta_{i,j}} \bar{L}_{e_j} \bar{L}_{e_i}^*$$

for all $i, j \in \{1, \dots, n\}$. By Proposition 2.14, there is a homomorphism $\varphi: \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta) \rightarrow \mathcal{N}\mathcal{T}_{A_\rho^\sigma}$ mapping L_{e_j} to \bar{L}_{e_j} for $j = 1, \dots, n$. This is precisely the inverse of $\tilde{\psi}$. \square

The dynamics $\bar{\alpha}$ on $\mathcal{N}\mathcal{T}_{A_\rho^\sigma}$ induced by the dynamics α on $\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ via the isomorphism from Proposition 7.3 satisfies

$$\bar{\alpha}_t(a\delta_p) = N(p)^{it} a\delta_p, \quad a \in A, \quad p \in \mathbb{N}^k.$$

Thus $\bar{\alpha}$ has the form specified in equation (1.7) of [3], for the homomorphism $N: \mathbb{N}^k \rightarrow (0, +\infty)$ given by $p \mapsto e^{\langle p|r \rangle}$, where $r = (r_1, \dots, r_k, 0_d)$ with $r_j > 0$ for $j \leq k$ and $\langle p|r \rangle$ is defined by identifying \mathbb{N}^k with $\mathbb{N}^k \times 0_d$. Every KMS_β state of $(\mathcal{N}\mathcal{T}_{A_\rho^\sigma}, \bar{\alpha})$ for $\beta > 0$ is gauge-invariant by Proposition 3.3.

Thus [3, Corollary 3.3] shows that the map $\phi \mapsto \phi|_A$ is a one-to-one correspondence from KMS_β states of $(\mathcal{N}\mathcal{T}_{A_\rho^\sigma}, \bar{\alpha})$ to tracial states of the coefficient algebra A that satisfy the inequality (3.2) of [3]. It follows from [3, Remark 3.2] that a tracial state τ of A satisfies such an inequality if and only if it has the form

$$\tau(a) = \sum_{m \in \mathbb{N}^k} e^{-\beta \langle m|r \rangle} \tau_0(\langle \delta_m | a \delta_m \rangle) \quad (a \in A),$$

for some trace τ_0 of A , which is uniquely determined by τ and satisfies $\tau_0(1) = Z(\beta)^{-1}$. Note that condition (3.3) of [3] is automatically satisfied here. The KMS_β state ϕ of $(\mathcal{N}\mathcal{T}_{A_\rho^\sigma}, \bar{\alpha})$ extending the tracial state τ of A is then given on an element $b \in \mathcal{N}\mathcal{T}_{A_\rho^\sigma}$ by the formula

$$\phi(b) = \sum_{m \in \mathbb{N}^k} e^{-\beta \langle m|r \rangle} \tau_0(\langle \delta_m | b \delta_m \rangle),$$

where the left action of $\mathcal{N}\mathcal{T}_{A_\rho^\sigma}$ on $A_{\rho_m}^\sigma$ is obtained by restricting the left action of $\mathcal{N}\mathcal{T}_{A_\rho^\sigma}$ on the Fock space $\mathcal{F}(A_\rho^\sigma) = \bigoplus_{p \in \mathbb{N}^k} A_{\rho_p}^\sigma$ of A_ρ^σ . Using the terminology of Corollary 6.10(ii) of [3] this implies that for $\beta > 0$ all KMS_β states of our system are of finite type. Combining this with the isomorphism $\mathcal{N}\mathcal{T}_{A_\rho^\sigma} \cong \mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta)$ from Proposition 7.3, we obtain from [3] a one-to-one correspondence between tracial states of $A = \mathcal{T}_r(\mathbb{N}^d, \sigma_{\Theta_d})$ and KMS_β states of $(\mathcal{T}_r(\mathbb{N}^n, \sigma_\Theta), \alpha)$ for which the resulting map from traces to KMS_β states coincides with the map given in Proposition 4.2. We emphasize that this application of the results from [3] to describe the phase transition for $\beta > 0$ depends on the choice of a product system that is adapted to the dynamics in such a way that KMS_β states are gauge-invariant, a choice that is guided by our Proposition 3.3.

REFERENCES

- [1] Sriwulan Adji, Marcelo Laca, May Nilsen, and Iain Raeburn, *Crossed products by semigroups of endomorphisms and the Toeplitz algebras of ordered groups*, Proc. Amer. Math. Soc. **122** (1994), no. 4, 1133–1141, doi: 10.2307/2161182. MR **1215024**
- [2] Zahra Afsar, Astrid an Huef, Iain Raeburn, and Aidan Sims, *Equilibrium states on higher-rank Toeplitz non-commutative solenoids*, Ergodic Theory Dynam. Systems **40** (2020), no. 11, 2881–2912, doi: 10.1017/etds.2019.20. MR4157468
- [3] Zahra Afsar, Nadia S. Larsen, and Sergey Neshveyev, *KMS states on Nica-Toeplitz C^* -algebras*, Comm. Math. Phys. **378** (2020), no. 3, 1875–1929, doi: 10.1007/s00220-020-03711-6. MR **4150892**
- [4] Ola Bratteli and Derek W. Robinson, *Operator algebras and quantum statistical mechanics. 2*, 2nd ed., Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997. Equilibrium states. Models in quantum statistical mechanics. doi: 10.1007/978-3-662-03444-6 MR1441540
- [5] Nathan Brownlowe, Mitchell Hawkins, and Aidan Sims, *The Toeplitz noncommutative solenoid and its Kubo-Martin-Schwinger states*, Ergodic Theory Dynam. Systems **39** (2019), no. 1, 105–131, doi: 10.1017/etds.2017.7.
- [6] Robert C. Busby and Harvey A. Smith, *Representations of Twisted Group Algebras*, Transactions of the American Mathematical Society **149** (1970), no. 2, 503–537.
- [7] Alain Connes and Matilde Marcolli, *Noncommutative geometry, quantum fields and motives*, American Mathematical Society Colloquium Publications, vol. 55, American Mathematical Society, Providence, RI; Hindustan Book Agency, New Delhi, 2008. doi: 10.1090/coll/055 MR2371808
- [8] John Crisp and Marcelo Laca, *Boundary quotients and ideals of Toeplitz C^* -algebras of Artin groups*, J. Funct. Anal. **242** (2007), 127–156, doi: 10.1016/j.jfa.2006.08.001.
- [9] Siegfried Echterhoff, S. Kaliszewski, John Quigg, and Iain Raeburn, *A categorical approach to imprimitivity theorems for C^* -dynamical systems*, Mem. Amer. Math. Soc. **180** (2006), no. 850, viii+169, doi: 10.1090/memo/0850. MR2203930

- [10] George A. Elliott, *On the K -theory of the C^* -algebra generated by a projective representation of a torsion-free discrete abelian group*, Operator algebras and group representations, Vol. I (Neptun, 1980), Monogr. Stud. Math., vol. 17, Pitman, Boston, MA, 1984, pp. 157–184. MR **731772**
- [11] Ruy Exel, *Circle actions on C^* -algebras, partial automorphisms, and a generalized Pimsner–Voiculescu exact sequence*, J. Funct. Anal. **122** (1994), no. 2, 361–401, doi: 10.1006/jfan.1994.1073. MR **1276163**
- [12] ———, *Partial dynamical systems, Fell bundles and applications*, Mathematical Surveys and Monographs, vol. 224, Amer. Math. Soc., Providence, RI, 2017.
- [13] Neal J. Fowler and Iain Raeburn, *Discrete Product Systems and Twisted Crossed Products by Semigroups*, J. Funct. Anal. **155** (1998), no. 1, 171 – 204, doi: 10.1006/jfan.1997.3227.
- [14] Neal J. Fowler, *Discrete product systems of Hilbert bimodules*, Pacific J. Math. **204** (2002), no. 2, 335–375, doi: 10.2140/pjm.2002.204.335. MR1907896
- [15] László Fuchs, *Abelian Groups*, Springer Monographs in Mathematics, Springer International Publishing, 2015. doi: 10.1007/978-3-319-19422-6
- [16] Philip Green, *The local structure of twisted covariance algebras*, Acta Math. **140** (1978), 191–250, doi: 10.1007/BF02392308. MR **493349**
- [17] Astrid an Huef, Marcelo Laca, Iain Raeburn, and Aidan Sims, *KMS states on the C^* -algebras of finite graphs*, J. Math. Anal. Appl. **405** (2013), no. 2, 388–399, doi: 10.1016/j.jmaa.2013.03.055. MR3061018
- [18] ———, *KMS states on the C^* -algebras associated to higher-rank graphs*, J. Funct. Anal. **266** (2014), 265–283.
- [19] Marcelo Laca and Iain Raeburn, *Extending Multipliers from Semigroups*, Proc. Amer. Math. Soc. **123** (1995), 355–362, doi: 10.2307/2160888.
- [20] ———, *Semigroup Crossed Products and the Toeplitz Algebras of Nonabelian Groups*, J. Funct. Anal. **139** (1996), no. 2, 415–440, doi: 10.1006/jfan.1996.0091. MR **1402771**,
- [21] ———, *A semigroup crossed product arising in number theory*, J. London Math. Soc. (2) **59** (1999), no. 1, 330–344, doi: 10.1112/S0024610798006620. MR1688505
- [22] ———, *Phase transition on the Toeplitz algebra of the affine semigroup over the natural numbers*, Adv. Math. **225** (2010), no. 2, 643–688, doi: 10.1016/j.aim.2010.03.007. MR2671177
- [23] Frédéric Latrémolière and Judith Packer, *Noncommutative solenoids*, New York J. Math. **24a** (2018), 155–191.
- [24] Alexandru Nica, *C^* -algebras generated by isometries and Wiener–Hopf operators*, J. Operator Theory **27** (1992), no. 1, 17–52. MR **1241114**
- [25] Chi-Keung Ng, *Discrete coactions on C^* -algebras*, J. Aust. Math. Soc. **60** (1996), 118–127, doi: 10.1017/S1446788700037423.
- [26] Gert K. Pedersen, *C^* -algebras and their automorphism groups*, Pure and Applied Mathematics (Amsterdam), Academic Press, London, 2018. Second edition of [MR0548006]; Edited and with a preface by Søren Eilers and Dorte Olesen. MR3839621
- [27] Dorte Olesen, Gert K. Pedersen, and Masamichi Takesaki, *Ergodic Actions of Compact Abelian groups*, J. Operator Theory **3** (1980), no. 2, 237–269.
- [28] N. Christopher Phillips, *Every simple higher dimensional noncommutative torus is an AT algebra*, arXiv:math/0609783 [math.OA] (2006).
- [29] Marc A. Rieffel, *Morita equivalence for operator algebras*, Proc. Sympos. Pure Math., vol. 38, Amer. Math. Soc., Providence, RI, 1982, pp. 285–298, doi: 10.1090/pspum/038.1.
- [30] ———, *Noncommutative tori—a case study of noncommutative differentiable manifolds*, Geometric and topological invariants of elliptic operators (Brunswick, ME, 1988), Contemp. Math., vol. 105, Amer. Math. Soc., Providence, RI, 1990, pp. 191–211, doi: 10.1090/conm/105. MR **1047281**
- [31] Camila F. Sehnem, *Fell bundles over quasi-lattice ordered groups and C^* -algebras of compactly aligned product systems*, Münster J. of Math. **14** (2021), no. 1, 223–263, doi: 10.17879/59019509488.
- [32] Joseph Slawny, *On factor representations and the C^* -algebra of canonical commutation relations*, Comm. Math. Phys. **24** (1972), no. 2, 151–170.

(Z. Afsar) SCHOOL OF MATHEMATICS AND STATISTICS, SYDNEY UNIVERSITY, NSW 2006, AUSTRALIA
Email address: `zahra.afsar@sydney.edu.au`

(M. Laca) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA, VICTORIA, P.O BOX
1700 STN CSC, BC V8W 2Y2, CANADA
Email address: `laca@uvic.ca`

(J. Ramagge) DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, U.K..
Email address: `jacqui.ramagge@durham.ac.uk`

(C. F. Sehnem) SCHOOL OF MATHEMATICS AND STATISTICS, VICTORIA UNIVERSITY OF WELLINGTON, P.O.
BOX 600, WELLINGTON 6140, NEW ZEALAND.
Email address: `camila.sehnem@vuw.ac.nz`