

AXIOMATIZING AECs AND APPLICATIONS

SAMSON LEUNG

ABSTRACT

For any abstract elementary class (AEC) \mathbf{K} with $\lambda = \text{LS}(\mathbf{K})$, the following holds:

1. K has an axiomatization in $L_{(2^\lambda)^+, \lambda^+}$, allowing game quantification. If \mathbf{K} has arbitrarily large models, the λ -amalgamation property and is categorical both in λ and λ^+ , then it has an axiomatization in L_{λ^+, λ^+} with game quantification. These extend Kueker's [Kue08] result which assumes finite character and $\lambda = \aleph_0$.
2. If K is universal and categorical in λ , then it is axiomatizable in L_{λ^+, λ^+} .
3. Shelah's celebrated presentation theorem asserts that for any AEC \mathbf{K} there is a first-order theory in an expansion of $L(\mathbf{K})$, and a set Γ of 2^λ many T -types such that $K = PC(T, \Gamma, L(\mathbf{K}))$. We provide a better bound on $|\Gamma|$ in terms of $I_2(\lambda, \mathbf{K})$.
4. We present additional applications which extend, simplify and generalize results of Shelah [She87, She01] and Shelah-Vasey [SV18]. Some of our main results generalize to μ -AECs.

TABLE OF CONTENTS

1	Introduction	2
2	Preliminaries	5
3	Encoding an AEC	7
4	A variation on Shelah's presentation theorem	18
5	Generalization to μ -AECs	24

Date: February 6, 2023

AMS 2020 Subject Classification: Primary 03C48. Secondary: 03C45, 03C55. Key words and phrases. Shelah's presentation theorem; infinitary logic; categoricity spectrum; axiomatization; Hanf number.

1 INTRODUCTION

In the proof of Shelah’s presentation theorem [She09, I Lemma 1.9], functions are added to capture isomorphism axioms and Löwenheim-Skolem axiom. [SV21, Theorem 2.1] claimed that any abstract elementary class (AEC) \mathbf{K} can be axiomatized by an $L_{\beth_2(\lambda)+3, \lambda^+}$ sentence where λ is the Löwenheim-Skolem number, and such an axiomatization is in $L(\mathbf{K})$.

Let $\chi := \lambda + I_2(\lambda, \mathbf{K})$, where $I_2(\lambda, \mathbf{K})$ is the number of nonisomorphic pairs (M, N) such that $M, N \in K_\lambda$ and $M \leq_{\mathbf{K}} N$. In Main Theorem 3.7, we will axiomatize an AEC \mathbf{K} by a sentence $\sigma_{\mathbf{K}}$ in L_{χ^+, λ^+} , allowing game quantification. As $\chi \leq 2^\lambda$, we have that $\sigma_{\mathbf{K}}$ is in $L_{(2^\lambda)^+, \lambda^+}$, improving Shelah and Villaveces’ result. Modulo the use of game quantification, our result is optimal for uncountable λ as it is known that there is an AEC that cannot be axiomatized by an $L_{\infty, \lambda}$ sentence [Hen19]. Under extra assumptions, we can axiomatize an AEC by a sentence in L_{λ^+, λ^+} (Proposition 3.10 and Theorem 3.12). By slightly modifying $\sigma_{\mathbf{K}}$, we can encode the \mathbf{K} -substructure relation (Proposition 3.16 and Proposition 3.18) by a formula σ_{\leq} in L_{χ^+, λ^+} . As above, we can improve the results under extra assumptions (Corollary 3.20).

As an application of our axiomatization of AECs, we derive a variation on the presentation theorem, where any AEC is a PC_χ class (Theorem 4.1) and game quantification is not used. Our presentation theorem is stronger than Shelah’s as the bound of $|\Gamma|$ in some cases is smaller than $2^{\text{LS}(\mathbf{K})}$. It also lowers the threshold of the existence theorem from successive categoricity (Theorem 4.8). The axiomatization strategy is also applicable to the μ -AEC analogs, giving a stronger presentation theorem (Theorem 5.6) than [BGL⁺16, Theorem 3.2].

In the following, we provide two tables. The first table summarizes the known results in literature. The definition of $L(\omega)$ can be found in [Kue08, Definition 1.12] and $L_{\chi^+, \lambda^+}(\omega \cdot \omega)$ is defined in Definition 2.6. The second table summarizes the main results in this paper. We write AL = arbitrarily large models, AP = amalgamation property, JEP = joint embedding property, NMM = no maximal models. Monster model means $AP + JEP + NMM$. The entries of the second table link to the related theorems (see Observation 3.15 for cases where we do not assume λ -categoricity).

Known results

Assumptions on \mathbf{K}	K is	References
None	axiomatizable in $L_{(2^{2^{\lambda^+}})_{+++}, \lambda^+}$	[SV21, Theorem 2.1]
None	$PC_{\lambda, 2^\lambda}$	[She87, Lemma 1.8]
None	reducts of a theory in L'_{χ^+, λ^+} where $L' \supseteq L$	[BB16, Theorem 3.2.3]
$\lambda = \aleph_0$, \aleph_0 -AP, stable in \aleph_0 , $I(\aleph_0, \mathbf{K}) \leq \aleph_0$	PC_{\aleph_0}	[SV18, Theorem 4.2]
$\lambda = \aleph_0$	closed under $\equiv_{\infty, \omega_1}$	[Kue08, Theorem 2.5]
$\lambda = \aleph_0$, $\exists \kappa = \kappa^{\aleph_0} (I(\kappa, \mathbf{K}) \leq \kappa)$	axiomatizable in L_{∞, ω_1}	[Kue08, Theorem 2.11]
$\lambda = \aleph_0$, $\exists \kappa = \kappa^{\aleph_0} (I(< \kappa, \mathbf{K}) \leq \kappa)$	axiomatizable in L_{κ^+, ω_1}	[Kue08, Theorem 2.11]
Finitary, $\lambda = \aleph_0$	closed under $\equiv_{\infty, \omega}$	[Kue08, Theorem 3.4]
Finitary	closed under $\equiv_{\infty, \omega}$	[Joh10, Theorem 3.7]
Finitary, $\lambda = \aleph_0$	axiomatizable in $L(\omega)$	[Kue08, Theorem 3.7]
Finitary, $\lambda = \aleph_0$, $\exists \kappa (I(\kappa, \mathbf{K}) \leq \kappa)$	axiomatizable in $L_{\infty, \omega}$	[Kue08, Theorem 3.10]
Finitary, $\exists \kappa = \kappa^{< \lambda} (I(\kappa, \mathbf{K}) \leq \kappa)$	axiomatizable in $L_{\infty, \lambda}$	[Joh10, Theorem 3.10]
Finitary, $\lambda = \aleph_0$, $\exists \kappa (I(< \kappa, \mathbf{K}) \leq \kappa)$	axiomatizable in $L_{\kappa^+, \omega}$	[Kue08, Theorem 3.10]
Finitary, $\exists \kappa = \kappa^{< \lambda} (I(< \kappa, \mathbf{K}) \leq \kappa)$	axiomatizable in $L_{\kappa^+, \lambda}$	[Joh10, Theorem 3.10]
$\lambda = \aleph_0$, $\exists \kappa (I(\kappa, \mathbf{K}) = 1)$	$K_{\geq \kappa}$ is closed under $\equiv_{\infty, \omega}$	[Kue08, Theorem 5.1]
Finitary, $\lambda = \aleph_0$, $\exists \kappa (I(\kappa, \mathbf{K}) = 1)$	$K_{\geq \kappa}$ axiomatizable in $L_{\omega_1, \omega}$	[Kue08, Theorem 5.2]
$\lambda = \aleph_0$, $\exists \kappa = \kappa^{\aleph_0} (I(\kappa, \mathbf{K}) = 1)$	Models in $K_{\geq \kappa}$ are $\equiv_{\infty, \omega_1}$	[Kue08, Theorem 5.3a]
Previous row + monster model	$K_{\geq \kappa}$ axiomatizable in $L_{(2^\omega)^+, \omega_1}$	[Kue08, Theorem 5.3c]
$\lambda > \aleph_0$	K is closed under $\equiv_{\infty, \lambda^+}$	[Kue08, Theorem 7.2]
Monster model, $\lambda > \aleph_0$, $\exists \kappa (I(\kappa, \mathbf{K}) = 1 \wedge \text{cf}(\kappa) > \lambda)$	$K_{\geq \kappa}$ axiomatizable in $L_{2^{\kappa^+}, \kappa^+}$	[Kue08, Theorem 7.4]
Finitary, monster model, $\exists \kappa (I(\kappa, \mathbf{K}) = 1 \wedge \text{cf}(\kappa) > \lambda)$	$K_{\geq \kappa}$ axiomatizable in $L_{\infty, \lambda}$	[Joh10, Theorem 3.11]

New results

Assumptions on \mathbf{K}	Axiomatization in	K is
None	$L_{\chi^+, \lambda^+}(\omega \cdot \omega)$	PC_χ
Universal class, $I(\lambda, \mathbf{K}) \leq \lambda$	L_{λ^+, λ^+}	PC_λ
$AL, \lambda\text{-}AP, I(\lambda, \mathbf{K}) \leq \lambda, I(\lambda^+, \mathbf{K}) = 1$	$L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$	PC_λ
$2^\lambda < 2^{\lambda^+}, AL, I(\lambda, \mathbf{K}) = 1, I(\lambda^+, \mathbf{K}) = 1$	$L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$	PC_λ
$\lambda\text{-}AP, I(\lambda, \mathbf{K}) \leq \lambda, I(\lambda^+, \mathbf{K}) = 1$ stable in λ	$L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$	PC_λ
$2^\lambda < 2^{\lambda^+}, I(\lambda, \mathbf{K}) = 1, I(\lambda^+, \mathbf{K}) = 1,$ stable in λ	$L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$	PC_λ

Note that the last row is a significant improvement of [SV18, Theorem 4.2], using much simpler and general methods while covering the case when λ is uncountable. We highlight the differences between our result and [BB16, Theorem 3.2.3]:

1. They expand the base vocabulary to τ^* by adding new predicates of arity λ , and their theory T^* in the expanded language is more semantic and longer; our axiomatization keeps the original language $L(\mathbf{K})$ and is purely syntactic.
2. Their relational presentation theorem characterizes K as reducts of models of T^* , \mathbf{K} -substructure as reducts of τ^* -substructure; our axiomatization is simply in $L_{\chi^+, \lambda^+}(\omega \cdot \omega)$ and we pin down the formula that determines \mathbf{K} -substructure.
3. Their expanded language τ^* has size χ ; our axiomatization uses the original language so has size $\leq \lambda$ (but both approaches require taking χ -conjunctions). We expand the language to size χ only when we derive a variation on Shelah's presentation theorem.
4. In [BB16, Theorem 3.2.3], they do not require types to be omitted because their theory T^* is in the infinitary logic. We omit types in our variation to Shelah's presentation theorem so as to represent K as a first-order PC class.

Our approach in this paper was inspired by Villaveces' question of the complexity of the example in [Leu23, Proposition 4.1], which has high instability but low complexity $\gamma = 1$.

Also, Grossberg suggested in May 2021 that [SV21] could have a significant improvement. This motivated us to look for a simpler axiomatization of an AEC, without using trees or other combinatorial machinery in [SV21, Theorem 2.4]. At the cost of game quantification, we lower the complexity of junctions in their paper from $\beth_2(\lambda)^{+3}$ to χ^+ .

This paper was written while the author was working on a Ph.D. under the direction of Rami Grossberg at Carnegie Mellon University and we would like to thank Prof. Grossberg for his guidance and assistance in my research in general and in this work in particular.

2 PRELIMINARIES

Let L be a finitary language, $\lambda_1 \geq \lambda_2$ be infinite cardinals. We write L_{λ_1, λ_2} the set of formulas generated by L , allowing $< \lambda_2$ free variables and $< \lambda_2$ quantifiers, in addition to conjunctions and disjunctions of $< \lambda_1$ subformulas. Given an L -structure M , we write $|M|$ the universe of M and $\|M\|$ the cardinality of M .

Definition 2.1. Let $L \subseteq L'$ be two languages (they can be infinitary), T be an L' -theory and Γ be a set of L' -types. Let μ be a regular cardinal. If L, L' are $(< \mu)$ -ary, we define

$$EC^\mu(T, \Gamma) := \{M : M \text{ is an } L'\text{-structure, } M \models T, M \text{ omits } \Gamma\}$$

$$PC^\mu(T, \Gamma) := \{M \upharpoonright L : M \text{ is an } L'\text{-structure, } M \models T, M \text{ omits } \Gamma\}$$

When $\mu = \aleph_0$, we omit the superscript \aleph_0 .

Let λ, χ be infinite cardinals, and assume $|T| \leq \lambda$ and $|\Gamma| \leq \chi$. If $K = EC^\mu(T, \Gamma)$, we call K an $EC_{\lambda, \chi}^\mu$ class. If $K = PC^\mu(T, \Gamma, L)$, we call K a $PC_{\lambda, \chi}^\mu$ class. We omit the superscript \aleph_0 when $\mu = \aleph_0$. We omit λ, χ if the sizes of T and Γ are not specified. PC_λ stands for $PC_{\lambda, \lambda}$.

Definition 2.2. Let L be a finitary language. An abstract elementary class $\mathbf{K} = \langle K, \leq_{\mathbf{K}} \rangle$ in L satisfies the following axioms:

1. K is a class of L -structures and $\leq_{\mathbf{K}}$ is a partial order on $K \times K$.
2. For $M_1, M_2 \in K$, $M_1 \leq_{\mathbf{K}} M_2$ implies $M_1 \subseteq M_2$ (as L -substructures).
3. Isomorphism axioms:

- (a) If $M \in K$, N is an L -structure, $M \cong N$, then $N \in K$.
 - (b) Let $M_1, M_2, N_1, N_2 \in K$. If $f : M_1 \cong M_2$, $g : N_1 \cong N_2$, $g \supseteq f$ and $M_1 \leq_{\mathbf{K}} N_1$, then $M_2 \leq_{\mathbf{K}} N_2$.
4. Coherence: Let $M_1, M_2, M_3 \in K$. If $M_1 \leq_{\mathbf{K}} M_3$, $M_2 \leq_{\mathbf{K}} M_3$ and $M_1 \subseteq M_2$, then $M_1 \leq_{\mathbf{K}} M_2$.
 5. Löwenheim-Skolem axiom: There exists an infinite cardinal $\lambda \geq |L(\mathbf{K})|$ such that: for any $M \in K$, $A \subseteq |M|$, there is some $N \in K$ with $A \subseteq |N|$, $N \leq_{\mathbf{K}} M$ and $\|N\| \leq \lambda + |A|$. We call the minimum such λ the Löwenheim-Skolem number $\text{LS}(\mathbf{K})$.
 6. Chain axioms: Let α be an ordinal and $\langle M_i : i < \alpha \rangle \subseteq K$ such that for $i < j < \alpha$, $M_i \leq_{\mathbf{K}} M_j$.
 - (a) Then $M := \bigcup_{i < \alpha} M_i$ is in K and for all $i < \alpha$, $M_i \leq_{\mathbf{K}} M$.
 - (b) Let $N \in K$. If in addition for all $i < \alpha$, $M_i \leq_{\mathbf{K}} N$, then $M \leq_{\mathbf{K}} N$.

Let $\lambda \geq \text{LS}(\mathbf{K})$ be a cardinal. We define $K_\lambda := \{M \in K : \|M\| = \lambda\}$ and $\mathbf{K}_\lambda := \langle K_\lambda, \leq_{\mathbf{K}} \upharpoonright K_\lambda \times K_\lambda \rangle$. When the context is clear, we omit the subscript of $\leq_{\mathbf{K}}$ and write \leq . We will only consider $\mathbf{K}_{\geq \text{LS}(\mathbf{K})}$ in place of \mathbf{K} , which is still an AEC.

- Definition 2.3.**
1. Let I be an index set. A directed system $\langle M_i : i \in I \rangle \subseteq K$ indexed by I satisfies the following: for any $i, j \in I$, there is $k \in I$ such that $M_i \leq M_k$ and $M_j \leq M_k$.
 2. Let μ be an infinite cardinal. A μ -directed system $\langle M_i : i \in I \rangle \subseteq K$ indexed by I satisfies the following: for any $J \subseteq I$ of size $< \mu$, there is $k \in I$ such that for all $j \in J$, $M_j \leq M_k$, (thus a system is directed iff it is \aleph_0 -directed.)

Fact 2.4. [She09, I Observation 1.16] Let $\langle M_i : i \in I \rangle \subseteq K$ be a directed system. Then

1. $M := \bigcup_{i \in I} M_i \in K$
2. For all $i \in I$, $M_i \leq M$.
3. Let $N \in K$. If in addition for all $i \in I$, $M_i \leq N$, then $M \leq N$.

Fact 2.5. [She09, II Claim 1.8(2)] If $M \leq N$ in K , then there are index sets I_1 and I_2 , directed systems $\langle M_i : i \in I_1 \rangle$ and $\langle N_i : i \in I_2 \rangle$ of union M , N respectively, $I_1 \subseteq I_2$ and $M_i = N_i$ for all $i \in I_1$.

Definition 2.6. Let L be a language, λ, χ be infinite cardinals and δ be an ordinal. $L_{\lambda, \chi}(\delta)$ extends $L_{\lambda, \chi}$ by allowing δ -game quantification: if ϕ is a formula in $L_{\lambda, \chi}(\delta)$ with free variables $(x_\alpha, y_\alpha)_{\alpha < \delta}$ and $l(x_\alpha), l(y_\alpha) < \chi$, then $(\forall x_\alpha \exists y_\alpha)_{\alpha < \delta} \phi$ is a formula in $L_{\lambda, \chi}(\delta)$. An L -structure M satisfies $(\forall x_\alpha \exists y_\alpha)_{\alpha < \delta} \phi$ if Player II has a winning strategy in the following game of δ rounds: in the α -th round, Player I chooses some tuple $a_\alpha \subseteq M$ of length $l(x_\alpha)$ and Player II responds by choosing some tuple $b_\alpha \subseteq M$ of length $l(y_\alpha)$. Player II wins if $M \models \phi[a_\alpha, b_\alpha]_{\alpha < \delta}$ where for $\alpha < \delta$, x_α is substituted by a_α and y_α is substituted by b_α .

If δ above is finite, then $L_{\lambda, \chi}(\delta) = L_{\lambda, \chi}$. The use of game quantifiers in AECs can be found in [Kue08, Theorems 2.9, 3.7] which handle the case $\text{LS}(\mathbf{K}) = \aleph_0$. Our version is consistent with $L(\omega)$ there and is called a *closed game quantifier* in [Kol85, Chapter X.2].

3 ENCODING AN AEC

In this section, we fix an AEC \mathbf{K} in a language L .

Definition 3.1. 1. Let $\lambda \geq \text{LS}(\mathbf{K})$. $I(\lambda, \mathbf{K}) := \{M/\cong : M \in K_\lambda\}$.

2. Let $M_1 \leq N_1$, $M_2 \leq N_2$. We write $(M_1, N_1) \cong (M_2, N_2)$ if there exists $f : N_1 \cong N_2$ such that $f \upharpoonright M_1 : M_1 \cong M_2$.

3. Let $\lambda \geq \text{LS}(\mathbf{K})$. $I_2(\lambda, \mathbf{K}) := \{(M, N)/\cong : M \leq N \text{ in } K_\lambda\}$.

Example 3.2. Depending on λ and \mathbf{K} , $I(\lambda, \mathbf{K})$ and $I_2(\lambda, \mathbf{K})$ may not be the same:

1. If \mathbf{K} is the class of the $L_{\omega_1, \omega}$ theory $\forall x \bigvee_{i < \omega} (x = c_i)$ where c_i are constants, then $I(\aleph_0, \mathbf{K}) = I_2(\aleph_0, \mathbf{K}) = 1$.
2. If \mathbf{K} is the class of the first-order theory of pure equality, then $I(\lambda, \mathbf{K}) = 1$ but $I_2(\lambda, \mathbf{K}) = \lambda$ for any infinite λ .

3. If \mathbf{K} is the class of the first-order theory of dense linear orders without endpoints, then $I(\aleph_0, \mathbf{K}) = 1$ but $I_2(\aleph_0, \mathbf{K}) = 2^{\aleph_0}$. The latter is because one can choose $M = \mathbb{Q}$ while $N_r = \{a + br : a, b \in \mathbb{Q}\}$ where r is an irrational number.

Proposition 3.3. $I(\lambda, \mathbf{K}) \leq I_2(\lambda, \mathbf{K}) \leq 2^\lambda$.

Proof. For any $M \leq N_1$ and $M \leq N_2$ in K , if $N_1 \not\cong N_2$, then $(M, N_1) \not\cong (M, N_2)$ by definition, hence the first inequality. Using the fact that $I(\lambda, \mathbf{K}) \leq 2^\lambda$, we can bound $I_2(\lambda, \mathbf{K}) \leq \lambda^\lambda \cdot I(\lambda, \mathbf{K}) \leq 2^\lambda$. \square

Question 3.4. Assuming stability or categoricity, is it possible to obtain a better bound than Proposition 3.3?

Until the end of this section, we write $\lambda := \text{LS}(\mathbf{K})$.

Observation 3.5. List the representatives of $\{M/\cong : M \in K_\lambda\}$ by $\langle M_i : i < I(\lambda, \mathbf{K}) \rangle$ and those of $\{(M, N)/\cong : M \leq N \text{ in } K_\lambda\}$ by $\langle (M_j, N_j) : j < I_2(\lambda, \mathbf{K}) \rangle$. For $i < I(\lambda, \mathbf{K})$, let $\phi_i(x)$ be an L_{λ^+, λ^+} formula that encodes the isomorphism type of M_i with a fixed enumeration of the universe $|M_i| = \{m_k^i : k < \lambda\}$. For variables $x = \langle x_k : k < \lambda \rangle$,

$$\phi_i(x) := \bigwedge \{ \theta(x_{\alpha_0}, \dots, x_{\alpha_{s-1}}) : M_i \models \theta[m_{\alpha_0}^i, \dots, m_{\alpha_{s-1}}^i], s < \omega, \alpha_0, \dots, \alpha_{s-1} < \lambda, \\ \theta \text{ is an atomic } L\text{-formula or its negation with } s \text{ free variables} \}$$

Namely for any L -structure N and any $a \in |N|$ of length λ , if $N \models \phi_i[a]$ then $a \cong M_i$ (with the fixed enumeration).

Similarly, for $j < I_2(\lambda, \mathbf{K})$, let $\psi_j(x, y)$ be an L_{λ^+, λ^+} formula that encodes the isomorphism type of (M_j, N_j) with fixed enumerations, where $|M_j| = \langle m_k^j : k < \lambda \rangle$, $|N_j| = \langle n_k^j : k < \lambda \rangle$. For variables $x = \langle x_k : k < \lambda \rangle$ and $y = \langle y_k : k < \lambda \rangle$,

$$\psi_j(x; y) := \bigwedge \{ \theta(x_{\alpha_0}, \dots, x_{\alpha_{s-1}}; y_{\beta_0}, \dots, y_{\beta_{t-1}}) : N_j \models \theta[m_{\alpha_0}^j, \dots, m_{\alpha_{s-1}}^j; n_{\beta_0}^j, \dots, n_{\beta_{t-1}}^j], \\ s < \omega, t < \omega, \alpha_0, \dots, \alpha_{s-1}, \beta_0, \dots, \beta_{t-1} < \lambda, \\ \theta \text{ is an atomic } L\text{-formula or its negation with } s + t \text{ free variables} \}$$

Namely for any L -structure N and any $a, b \in |N|$ both of length λ , if $N \models \psi_j[a, b]$ then $a \cong M_j$, $b \cong N_j$ (with the fixed enumerations) and $\text{ran}(a) \subseteq \text{ran}(b)$.

It is also possible to encode the re-enumerations of the isomorphism types in ϕ_i and ψ_j , but we will do that directly in the sentence $\sigma_{\mathbf{K}}$ in Main Theorem 3.7 and σ_{\leq} in Proposition 3.18(1), so as to be more consistent with the format of Theorem 4.1.

Definition 3.6. Let $\alpha, \beta < \lambda^+$, $a = \langle a_i : i < \alpha \rangle$ and $b = \langle b_i : i < \beta \rangle$. $a \subseteq b$ stands for $\text{ran}(a) \subseteq \text{ran}(b)$, which can be expressed by the L_{λ^+, λ^+} formula

$$\bigwedge_{i < \alpha} \bigvee_{j < \beta} a_i = b_j$$

$a \approx b$ stands for $\text{ran}(a) = \text{ran}(b)$, which can be expressed by the L_{λ^+, λ^+} formula

$$\left(\bigwedge_{i < \alpha} \bigvee_{j < \beta} a_i = b_j \right) \wedge \left(\bigwedge_{j < \beta} \bigvee_{i < \alpha} b_j = a_i \right)$$

Main Theorem 3.7. K is axiomatizable by an $L_{(\lambda + I_2(\lambda, \mathbf{K}))^+, \lambda^+}(\omega \cdot \omega)$ sentence $\sigma_{\mathbf{K}}$. In other words, for any L -structure M , $M \in K$ iff $M \models \sigma_{\mathbf{K}}$.

Proof. The following variables $(x_\alpha, y_\alpha)_{\alpha < \omega \cdot \omega}$ are all of length λ .

$$\begin{aligned} \sigma_{\mathbf{K}} := & (\forall x_\alpha \exists y_\alpha)_{\alpha < \omega \cdot \omega} \bigwedge_{\alpha < \omega \cdot \omega} \left[(x_\alpha \subseteq y_\alpha) \wedge \exists z_\alpha \left((y_\alpha \approx z_\alpha) \wedge \bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(z_\alpha) \right) \wedge \right. \\ & \left. \bigwedge_{\beta < \alpha < \omega \cdot \omega} \exists u_{\beta, \alpha} \exists v_{\beta, \alpha} \left((y_\beta \approx u_{\beta, \alpha}) \wedge (y_\alpha \approx v_{\beta, \alpha}) \wedge \bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(u_{\beta, \alpha}, v_{\beta, \alpha}) \right) \right] \end{aligned}$$

In words, $\sigma_{\mathbf{K}}$ stipulates the iterated use of Löwenheim-Skolem and coherence axioms $(\omega \cdot \omega)$ many times.

Suppose $M \in K$, we show that Player II can win the associated game in $\sigma_{\mathbf{K}}$. In the α -th round, Player I provides some x_α of length λ . By Löwenheim-Skolem axiom, pick any $y_\alpha \leq M$ of size λ such that $\text{ran}(x_\alpha) \cup \bigcup_{\beta < \alpha} \text{ran}(y_\beta) \subseteq \text{ran}(y_\alpha)$. By inductive hypothesis, for $\beta < \alpha$, we have $y_\beta \leq M$. By coherence axiom, $y_\beta \leq y_\alpha$ as desired.

Suppose $M \models \sigma_{\mathbf{K}}$. By Fact 2.4(1), it suffices to build a directed system $\langle M_a \in K_\lambda : a \in I \rangle$ of union M . We choose I to be the set of finite tuples a in M . Let a, b be finite tuples in M , we order $a \leq_I b$ iff $\text{ran}(a) \subseteq \text{ran}(b)$. We will inductively build all M_a in ω stages. At stage n we handle finite tuples of length $n + 1$.

- Stage 0: apply $\sigma_{\mathbf{K}}$ to each singleton s in M and substitute $x_0 = s$. We obtain $y_0 = M_s$ which is a \mathbf{K} -structure. Only the 0-th round of the game is used for each singleton.

for L -substructure. The first row represents the game of s while the last row represents the game of t . Each vertical column contains identical \mathbf{K} -structures.

Define $M^* := \bigcup_{k < \omega} N_k$. By requirements (1), (2) and chain axioms, $M^* \geq M_a$ and $M^* \geq M_b$. Also, s has used the first $\omega \cdot (m_s + 1)$ rounds while t has used the first $\omega \cdot (m_t + 1)$ rounds. This finishes the construction of M^* .

- Stage $n + 1$: Now for tuples c of length $n + 2$, we build M_c . Break down c as the union of two tuples of length $\leq n + 1$ (there might be more than one way), say a and b . As above assume M_a is generated by some singleton s and M_b by some singleton t . Then we can find M^* with $M^* \geq M_a$, $M^* \geq M_b$ and M^* is the union of bounded many y_α 's of the game of some singleton s . We cannot immediately define $M_c := M^*$ because M^* depends on the choice of the decomposition a, b . Since there are finitely many possible decompositions of a finite tuple c , we can continue the game of s and extend M^* to M_c which includes all M^* from other decompositions of c (M_c might not be unique but it is a \mathbf{K} -superstructure to all those M_a with $\text{ran}(a) \subseteq \text{ran}(c)$; M_c is also generated by other games of singletons but we just need one representative s for M_c).

After the construction is completed, $\langle M_a \in K_\lambda : a \in I \rangle$ is directed by our inductive step. Their union is M because for each element u in M , $u \in M_{\{u\}}$. \square

Remark 3.8. Our theorem generalizes [Kue08, Theorems 2.9, 3.7] which use the ω -game quantification. Modulo the $(\omega \cdot \omega)$ -game quantification, our result also generalizes [Kue08, Theorems 5.3, 7.4] which assumes a monster model and categoricity in a higher cardinal.

We will encode the \mathbf{K} -substructure relation in Proposition 3.18. Before that we investigate possible improvements of Main Theorem 3.7. The following questions were suggested by Grossberg:

Question 3.9. Is it possible to axiomatize an AEC \mathbf{K} in L_{λ^+, λ^+} instead of $L_{(\lambda + I_2(\lambda, \mathbf{K}))^+, \lambda^+}$ (with or without game quantification), assuming

1. stability?

2. categoricity in λ and λ^+ ?

For (2), Grossberg also suggested that [MAV18] should allow improvements of Main Theorem 3.7. Indeed it is possible when \mathbf{K} is a universal class. A partial converse can be found in [MAV18, Theorem 3.5].

Proposition 3.10. *If \mathbf{K} is a universal class, then it is axiomatizable by an $L_{(\lambda+I(\lambda,\mathbf{K}))^+,\lambda^+}$ sentence $\sigma_{\mathbf{K}}$. In particular if \mathbf{K} is categorical in λ , then it is axiomatizable by an L_{λ^+,λ^+} sentence.*

Proof. Since models are ordered by L -substructures, we can avoid game quantification and replace the ψ_j 's by subset relations when defining $\sigma_{\mathbf{K}}$ in Main Theorem 3.7. Namely,

$$\begin{aligned} \sigma_{\mathbf{K}} := & \forall a \exists m_a \left(a \subseteq m_a \wedge \bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(m_a) \right) \wedge \\ & \forall m_b \forall m_c \left[\left(\bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(m_b) \wedge \bigvee_{j < I(\lambda, \mathbf{K})} \phi_j(m_c) \right) \right. \\ & \left. \rightarrow \exists m_d \left(\bigvee_{k < I(\lambda, \mathbf{K})} \phi_k(m_d) \wedge (m_b \subseteq m_d) \wedge (m_c \subseteq m_d) \right) \right] \end{aligned}$$

□

We also have some approximations to Question 3.9(2), using game quantification. In the following we abbreviate *amalgamation property* as *AP* and *arbitrarily large models* as *AL*.

- Fact 3.11.**
1. If \mathbf{K} has *AL*, λ -*AP*, is categorical in λ^+ , then it is stable in λ . Hence for any $M \in K_\lambda$, there is $N \in K_\lambda$ which is a (λ, ω) -limit model over M .
 2. Let \mathbf{K} have λ -*AP* and $M_1, M_2, M_3 \in K_\lambda$. If $M_1 \leq M_2$ and M_3 is a (λ, ω) -limit model over M_2 , then M_3 is also a (λ, ω) -limit model over M_1 .
 3. Let $M, N, N' \in K_\lambda$. If N, N' are both (λ, ω) -limit models over M , then $N \cong_M N'$.

Theorem 3.12. *If \mathbf{K} has *AL*, λ -*AP* and is categorical in λ and λ^+ , then it is axiomatizable by an $L_{\lambda^+,\lambda^+}(\omega \cdot \omega)$ sentence σ . *AL* can be replaced by stability in λ .*

Proof. By Fact 3.11(1), fix $M, N \in K_\lambda$ such that N is (λ, ω) -limit over M . Let $\phi(x)$ code the isomorphism type of M and $\psi(x, y)$ code the isomorphism type of (M, N) . From the proof of Main Theorem 3.7, it suffices to replace $\bigvee_{i < I(\lambda, \mathbf{K})} \phi_i$ and $\bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j$ there by some λ -junctions. Since \mathbf{K} is λ -categorical, we can replace the first disjunction by ϕ . We also replace the second disjunction by a coherence formula involving ψ . Finally we add a disjunction to $\sigma_{\mathbf{K}}$ specifying models of size λ :

$$\sigma := \exists w (\phi(w) \wedge \forall x (x \subseteq w)) \vee \left\{ \begin{aligned} & (\forall x_\alpha \exists y_\alpha)_{\alpha < \omega \cdot \omega} \bigwedge_{\alpha < \omega \cdot \omega} \left[(x_\alpha \subseteq y_\alpha) \wedge \exists z_\alpha \left((y_\alpha \approx z_\alpha) \wedge \phi(z_\alpha) \right) \wedge \bigwedge_{\beta < \alpha < \omega \cdot \omega} \exists u_{\beta, \alpha} \exists v_{\beta, \alpha} \exists w \exists z \right. \\ & \left. \left((u_{\beta, \alpha} \subseteq v_{\beta, \alpha}) \wedge (y_\beta \approx u_{\beta, \alpha}) \wedge (y_\alpha \approx v_{\beta, \alpha}) \wedge (w \approx z) \wedge \psi(u_{\beta, \alpha}, w) \wedge \psi(v_{\beta, \alpha}, z) \right) \right] \end{aligned} \right\}$$

Suppose M^* is an L -structure and $M^* \models \sigma$. By coherence, the last line of σ implies $y_\beta \leq y_\alpha$. Then either $M^* \in K_\lambda$ or it can build a directed system $\langle M_\alpha \in K_\lambda : \alpha \in I \rangle$ of union M^* as in Main Theorem 3.7. By Fact 2.4(1) $M^* \in K$.

Suppose $M^* \in K_\lambda$, then it satisfies the first disjunct of σ . Otherwise $M^* \in K_{>\lambda}$. We need to verify that for any $M_1, M_2 \in K_\lambda$, if $M_1 \leq M_2 \leq M^*$ then there is $M_3 \leq M^*$ such that $(M_1, M_3) \cong (M_2, M_3) \cong (M, N)$. By *AL* and categoricity in λ^+ , M^* is λ^+ -saturated, so we can build $M_3 \in K_\lambda$ which is (λ, ω) -limit over M_2 .

$$\begin{array}{ccccc} M_3 & \xrightarrow{\cong} & N' & \xrightarrow{\cong} & N \\ (\lambda, \omega) \uparrow & \nearrow & \nwarrow & \uparrow & (\lambda, \omega) \\ M_2 & & & & \\ & \nwarrow & & & \\ & & M_1 & \xrightarrow{\cong} & M \end{array}$$

By Fact 3.11(2), M_3 is (λ, ω) -limit to both M_1 . Since \mathbf{K} is categorical in λ , $M_1 \cong M$ and we can extend this isomorphism to $M_3 \cong N'$ for some $N' \geq M$. Then N' is a (λ, ω) -limit over M . By Fact 3.11(3), $N' \cong_M N$ so $(M_1, M_3) \cong (M, N)$. Similarly $(M_2, M_3) \cong (M, N)$. Therefore, $M^* \models \sigma$ as desired. \square

Using a well-known result of Shelah, we can replace the assumption by *AP* by a set-theoretic one.

Fact 3.13. [She87, Theorem 3.5] Assume $2^\lambda < 2^{\lambda^+}$. If $I(\lambda, \mathbf{K}) = 1$ and $1 \leq I(\lambda^+, \mathbf{K}) < 2^{\lambda^+}$, then \mathbf{K} has λ -AP.

Corollary 3.14. Assume $2^\lambda < 2^{\lambda^+}$. If \mathbf{K} has AL and is categorical in λ and λ^+ , then it is axiomatizable by an $L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$ sentence σ' . AL can be replaced by stability in λ .

Proof. Combine Fact 3.13 and Theorem 3.12. □

In other words, under WGCH Question 3.9 has a positive answer when we assume both (1) and (2) of the hypotheses there and use game quantification.

Observation 3.15. 1. As in Proposition 3.10, we can replace categoricity in λ by $I(\lambda, \mathbf{K}) \leq \lambda$. We keep the original format of the theorem statements to better answer Question 3.9.

2. Let $\kappa \geq \lambda^+$. In Theorem 3.12, if we replace categoricity in λ^+ by κ , and further assume $(< \kappa)$ -AP, then models in K_κ are saturated [Vas17, Corollary 4.11(3)]. The same argument allows us to axiomatize $K_{\geq \kappa}$ by an $L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$ sentence. If κ is regular, then we can replace AL by stability in $[\lambda, \kappa)$.

3. John Baldwin pointed out that [She99] can reduce the successive categoricity assumption to a single categoricity. Indeed, assuming AP and categoricity in a successor above H_2 (the second Hanf number), we have categoricity in H_2 and AL. Thus we can axiomatize $K_{\geq H_2}$ by a sentence in $L_{H_2^+, H_2^+}(\omega \cdot \omega)$.

4. Marcos Mazari-Armida observed that [She01] provides another variation on Corollary 3.14: if we assume categoricity in λ , λ^+ , λ^{++} as well as $2^\lambda < 2^{\lambda^+} < 2^{\lambda^{++}}$, then we have stability in λ and λ -AP. Hence we can axiomatize K by a sentence in $L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$.

We now encode the \mathbf{K} -substructure relation. First we handle the case when the smaller model has size λ . In [Vas18, Section 6], the language is expanded by adding a new predicate for the substructure relation. In [BB16, Theorem 3.2.3], $I_2(\lambda, \mathbf{K})$ many new predicates are added. Here we explicitly define the predicates in $L_{(\lambda + I_2(\lambda, \mathbf{K}))^+, \lambda^+}(\omega \cdot \omega)$ without expanding the language.

Proposition 3.16. *There is an $L_{(\lambda+I_2(\lambda, \mathbf{K}))^+, \lambda^+}(\omega \cdot \omega)$ formula $\sigma_{\leq}(x)$ that encodes the \mathbf{K} -substructure relation: for any $M \in K$, $a \subseteq |M|$ of size λ , $M \models \sigma_{\leq}[a]$ iff $a \leq M$ (the enumeration of a does not matter).*

Proof. Our definition $\sigma_{\leq}(x)$ will be similar to that of $\sigma_{\mathbf{K}}$:

$$\begin{aligned} \sigma_{\leq}(x) := & (\forall x_{\alpha} \exists y_{\alpha})_{\alpha < \omega \cdot \omega} \left(\exists u \exists v (x \approx u) \wedge (y_0 \approx v) \wedge \bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(u, v) \right) \wedge \\ & \bigwedge_{\alpha < \omega \cdot \omega} \left[(x_{\alpha} \subseteq y_{\alpha}) \wedge \exists z_{\alpha} \left((y_{\alpha} \approx z_{\alpha}) \wedge \bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(z_{\alpha}) \right) \wedge \right. \\ & \left. \bigwedge_{\beta < \alpha < \omega \cdot \omega} \exists u_{\beta, \alpha} \exists v_{\beta, \alpha} \left((y_{\beta} \approx u_{\beta, \alpha}) \wedge (y_{\alpha} \approx v_{\beta, \alpha}) \wedge \bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(u_{\beta, \alpha}, v_{\beta, \alpha}) \right) \right] \end{aligned}$$

The enumeration of a does not matter by our definition of \approx . If $a \in K_{\lambda}$ and $a \leq M$, then $M \models \sigma_{\leq}[a]$ by Löwenheim-Skolem and coherence axioms. In particular, given x_0 in $\sigma_{\leq}[a]$, we choose $y_0 \leq M$ that contains both x_0 and a . Then coherence guarantees that $a \leq y_0$. Conversely suppose $a \subseteq |M|$ of size λ and $M \models \sigma_{\leq}[a]$. As in Main Theorem 3.7, we can build a directed system $\langle M_{\alpha} \in K_{\lambda} : \alpha \in I \rangle$ of union M with the additional requirement that for any $\alpha \in I$, $M_{\alpha} \geq a$. By Fact 2.4(1)(2), $M \in K$ and $M_{\alpha} \leq M$ for all $\alpha \in I$. By transitivity of \leq , $a \leq M$ as desired. \square

Remark 3.17. Using the terminology in [SV21, Theorem 2.1], we only used the first two levels of the canonical tree because a K -substructure relation only concerns two levels. The price to pay is game quantification.

Since our infinitary language only allows λ many free variables, it cannot directly encode substructures of size greater than λ . We propose two solutions: the first solution is a substructure relation whose underlying language is the singleton $\{\sigma_{\leq}\}$. The second solution involves relativizing σ_{\leq} to a new predicate.

Proposition 3.18. *Let $M, N \in K$ and σ_{\leq} be defined as in Proposition 3.16.*

1. $M \leq N$ iff $M \subseteq_{\{\sigma_{\leq}\}} N$ (if $a \subseteq |M|$ is of size λ , then $M \models \sigma_{\leq}[a]$ iff $N \models \sigma_{\leq}[a]$).
2. Let R be a new predicate where $N^R = |M|$ closed under permutations. $M \leq N$ iff $(N, R) \models \forall b (\sigma_{\leq}^R(b) \rightarrow \sigma_{\leq}(b))$ where σ_{\leq}^R is the relativized version of σ_{\leq} inside R .

Namely,

$$\begin{aligned} \sigma_{\leq}^R(x) := & (\forall x_\alpha \exists y_\alpha)_{\alpha < \omega \cdot \omega} \left(\bigwedge_{\alpha < \omega \cdot \omega} R(x_\alpha) \right) \rightarrow \left\{ \left(\bigwedge_{\alpha < \omega \cdot \omega} R(y_\alpha) \right) \wedge \right. \\ & \left. \left(\exists u \exists v (x \approx u) \wedge (y_0 \approx v) \wedge \bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(u, v) \right) \wedge \right. \\ & \left. \bigwedge_{\alpha < \omega \cdot \omega} \left[(x_\alpha \subseteq y_\alpha) \wedge \exists z_\alpha \left((y_\alpha \approx z_\alpha) \wedge \bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(z_\alpha) \right) \wedge \right. \right. \\ & \left. \left. \bigwedge_{\beta < \alpha < \omega \cdot \omega} \exists u_{\beta, \alpha} \exists v_{\beta, \alpha} \left((y_\beta \approx u_{\beta, \alpha}) \wedge (y_\alpha \approx v_{\beta, \alpha}) \wedge \bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(u_{\beta, \alpha}, v_{\beta, \alpha}) \right) \right] \right\} \end{aligned}$$

Proof. 1. If $M \leq N$ and let $a \subseteq M$. Using Proposition 3.16, if $M \models \sigma_{\leq}[a]$, then $a \leq M \leq N$ showing $N \models \sigma_{\leq}[a]$. If $N \models \sigma_{\leq}[a]$, then $a \leq N$. By coherence, $a \leq M$ and so $M \models \sigma_{\leq}[a]$. Conversely, build a directed system $\langle M_\alpha \in K_\lambda : \alpha \in I \rangle$ inside M such that for all $\alpha \in I$, $M_\alpha \leq M$. Then $M \models \sigma_{\leq}[M_\alpha]$. Since $M \subseteq_{\{\sigma_{\leq}\}} N$, we have $N \models \sigma_{\leq}[M_\alpha]$ and $M_\alpha \leq N$. The result follows from Fact 2.4(3).

2. If $M \leq N$ and $N \models \sigma_{\leq}^R[b]$ for some $b \subseteq |N|$, we need to show that $N \models \sigma_{\leq}[b]$. By assumption we can build a directed system of union M and have $b \leq M$. By transitivity of \leq , $b \leq N$ and the conclusion follows. Conversely, by Fact 2.4(3), it suffices to build a directed system $\langle M_\alpha \in K_\lambda : \alpha \in I \rangle$ of union M such that for all $\alpha \in I$, $M_\alpha \leq N$. Since $(N, R) \models \forall b (\sigma_{\leq}^R(b) \rightarrow \sigma_{\leq}(b))$, we can require $M_\alpha \leq M$ instead of $M_\alpha \leq N$. Such construction is possible by Löwenheim-Skolem and coherence axioms. □

Using game quantification, we derive a simple proof to [Kue08, Theorem 7.2], which uses back-and-forth arguments. [Vas18, Theorem 6.21] proved similarly by transferring the AEC to its substructure expansion and translating results between $\equiv_{\infty, \lambda^+}$ and λ^+ back-and-forth systems.

Corollary 3.19. *Let M, N be L -structures. If either M or N is in K and $M \preceq_{L_{\infty, \lambda^+(\omega \cdot \omega)}} N$, then $M \leq N$ (and both are in K).*

Proof. Since $M \preceq_{L_{\infty, \lambda^+}(\omega \cdot \omega)} N$, $M \subseteq_{L_{(\lambda + I_2(\lambda, \mathbf{K}))^+, \lambda^+}(\omega \cdot \omega)} N$. In particular $M \models \sigma_{\mathbf{K}}$ iff $N \models \sigma_{\mathbf{K}}$. By Main Theorem 3.7, either M, N is in K implies both are in K . On the other hand, the assumption implies $M \subseteq_{\{\sigma_{\leq}\}} N$. By Proposition 3.18(1), $M \leq N$. \square

One can ask the same Question 3.9 for \mathbf{K} -substructure relation instead of models of K . The following are variations on Proposition 3.16:

Corollary 3.20. *There is a formula $\sigma_{\leq}(x)$ in $L_{\lambda^+, \lambda^+}(\omega \cdot \omega)$ that encodes the \mathbf{K} -substructure relation (for any $M \in K$, $a \subseteq |M|$ of size λ , $M \models \sigma_{\leq}[a]$ iff $a \leq M$), assuming one of the following:*

1. \mathbf{K} is a universal class, in which case $\sigma_{\leq}(x)$ is in L_{λ^+, λ^+} (categoricity is not needed).
2. \mathbf{K} has AL, λ -AP and is categorical in λ, λ^+ and we restrict the use of σ_{\leq} to models of size $\geq \lambda^+$. AL can be replaced by stability in λ .
3. $2^\lambda < 2^{\lambda^+}$, \mathbf{K} has AL and is categorical in λ, λ^+ and we restrict the use of σ_{\leq} to models of size $\geq \lambda^+$. AL can be replaced by stability in λ .

Proof. 1. $\sigma_{\leq}(x)$ requires that x is closed under functions.

2. Combine the proofs of Theorem 3.12 and Proposition 3.16: from Proposition 3.16, it suffices to encode $a \leq b$ where both have length λ (then we can replace $\bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(a, b)$). Fix $M, N \in K_\lambda$ such that N is (λ, ω) -limit over M . Let $\phi(x)$ code the isomorphism type of M and $\psi(x, y)$ code the isomorphism type of (M, N) . From the proof of Theorem 3.12, ψ is the only isomorphism type of pairs inside a λ^+ -saturated model. Thus $a \leq b$ can be encoded as

$$a \subseteq b \wedge \exists z_0 \exists z_1 \exists a' \exists b' ((a \approx a') \wedge (b \approx b') \wedge (z_0 \approx z_1) \wedge \psi(a', z_0) \wedge \psi(b', z_1)).$$

3. Combine (2) and Fact 3.13. \square

With the exact same proof as in Proposition 3.18, we can show:

Corollary 3.21. *Let $M, N \in K$ and σ_{\leq} be defined as in Corollary 3.20.*

1. $M \leq N$ iff $M \subseteq_{\{\sigma_{\leq}\}} N$ (if $a \subseteq |M|$ is of size λ , then $M \models \sigma_{\leq}[a]$ iff $N \models \sigma_{\leq}[a]$).
2. Let R be a new predicate where $N^R = |M|$ closed under permutations. $M \leq N$ iff $(N, R) \models \forall b (\sigma_{\leq}^R(b) \rightarrow \sigma_{\leq}(b))$ where σ_{\leq}^R is the relativized version of σ_{\leq} inside R .

4 A VARIATION ON SHELAH'S PRESENTATION THEOREM

We will give a variation to Shelah's presentation theorem via our encoding of AECs. The presentation theorem statement is adapted from [Bal09, Theorem 4.15] (see also [Gro02, Theorem 3.4]).

Theorem 4.1. *Let \mathbf{K} be an AEC in L and with Löwenheim-Skolem number $\text{LS}(\mathbf{K})$. Define $\chi := \text{LS}(\mathbf{K}) + I_2(\text{LS}(\mathbf{K}), \mathbf{K})$. There exists an expansion $L' \supseteq L$ of size χ , an L' -theory T and a set of L' -types Γ of size χ such that*

1. $K = PC(T, \Gamma, L)$.
2. If $M', N' \in EC(T, \Gamma)$ and $M' \subseteq_{L'} N'$, then $M' \upharpoonright L \leq_{\mathbf{K}} N' \upharpoonright L$.
3. If $M \leq_{\mathbf{K}} N$, there are L' -expansions of M, N to M', N' such that $M' \subseteq_{L'} N'$.

Remark 4.2. When $I_2(\text{LS}(\mathbf{K}), \mathbf{K}) < 2^{\text{LS}(\mathbf{K})}$, our result is stronger than Shelah's presentation theorem as his encoding totally ignores the model theoretic complexity of \mathbf{K} and is using $2^{\text{LS}(\mathbf{K})}$ many types in Γ .

Proof. We will adapt $\sigma_{\mathbf{K}}$ in Main Theorem 3.7 to the first-order language. For a tuple of variables a and $n < \omega$, we write a^n to emphasize a has length n . For $k < n$, we write $a^n(k)$ the k -th coordinate of a^n .

As in the original presentation theorem, expand L to L' which includes $\{f_k^n : k < \text{LS}(\mathbf{K}), n < \omega\}$ where for each $n < \omega$, $k < \text{LS}(\mathbf{K})$, f_k^n is an n -ary function. For $n < \omega$, we will require that $\{f_k^n : k < \text{LS}(\mathbf{K})\}$ maps an n -tuple to a K -structure of size $\text{LS}(\mathbf{K})$ containing that tuple. This will be achieved by

$$\sigma^n := \forall a^n \bigwedge_{l < n} \bigvee_{k < \text{LS}(\mathbf{K})} (f_k^n(a^n) = a^n(l)) \wedge \bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(\{f_k^n(a^n) : k < \text{LS}(\mathbf{K})\})$$

In the above definition, although ϕ_i 's have $\text{LS}(\mathbf{K})$ many free variables, it is just an $\text{LS}(\mathbf{K})$ -conjunction of (negation of) atomic formulas with n free variables (from a^n). So each ϕ_i is inside $L'_{\text{LS}(\mathbf{K})^+, \omega}$.

Also, we want to require that the K -structures generated are directed with respect to the tuple input. However, $\{f_k^n : k < \text{LS}(\mathbf{K}), n < \omega\}$ might not be compatible with the enumerations of *pairs* of models, say $M_j \leq N_j$. Hence we expand L' further to include $\{g_k^{m,l}, h_k^{m,l} : k < \text{LS}(\mathbf{K}), m + l < \omega\}$ where for $m + l < \omega$, $k < \text{LS}(\mathbf{K})$, $g_k^{m,l}$ and $h_k^{m,l}$ are $(m + l)$ -ary functions and correctly enumerate a pair of models. The following will take care of the re-enumerations of $\{f_k^n : k < \text{LS}(\mathbf{K})\}$ for each $n < \omega$.

$$\begin{aligned} \sigma^{m,n,l} := & \forall b^m \forall c^n \forall d^l \left[\text{ran}(b^m) \cup \text{ran}(c^n) \subseteq \text{ran}(d^l) \rightarrow \right. \\ & \left(\{f_k^m(b^m) : k < \text{LS}(\mathbf{K})\} \approx \{g_k^{m,l}(b^m; d^l) : k < \text{LS}(\mathbf{K})\} \wedge \right. \\ & \{f_k^l(d^l) : k < \text{LS}(\mathbf{K})\} \approx \{h_k^{m,l}(b^m; d^l) : k < \text{LS}(\mathbf{K})\} \wedge \\ & \{f_k^n(c^n) : k < \text{LS}(\mathbf{K})\} \approx \{g_k^{n,l}(c^n; d^l) : k < \text{LS}(\mathbf{K})\} \wedge \\ & \left. \{f_k^l(d^l) : k < \text{LS}(\mathbf{K})\} \approx \{h_k^{n,l}(c^n; d^l) : k < \text{LS}(\mathbf{K})\} \wedge \right. \\ & \bigvee_{i < I_2(\text{LS}(\mathbf{K}), \mathbf{K})} \psi_i(\{g_k^{m,l}(b^m; d^l) : k < \text{LS}(\mathbf{K})\}, \{h_k^{m,l}(b^m; d^l) : k < \text{LS}(\mathbf{K})\}) \wedge \\ & \left. \bigvee_{j < I_2(\text{LS}(\mathbf{K}), \mathbf{K})} \psi_j(\{g_k^{n,l}(c^n; d^l) : k < \text{LS}(\mathbf{K})\}, \{h_k^{n,l}(c^n; d^l) : k < \text{LS}(\mathbf{K})\}) \right) \left. \right] \end{aligned}$$

Similar to the case of σ^n , the formulas ϕ_i, ψ_j and the connective \approx are simply $\text{LS}(\mathbf{K})$ -junctions of (negation of) atomic formulas, which are inside $L'_{\text{LS}(\mathbf{K})^+, \omega}$.

To convert $\{\sigma^n : n < \omega\} \cup \{\sigma^{m,n,l} : m, n, l < \omega\}$ into first-order sentences, we use Chang's presentation theorem (see [Gro21, Chapter 1 Theorem 8.16]) which adds χ -many new predicates to L' to represent the χ -conjunctions and disjunctions, and χ -many L' -types to omit. This gives our final T, Γ and L' .

It remains to check the three items in the theorem statement.

1. $K \subseteq PC(T, \Gamma, L)$ by Löwenheim-Skolem and coherence axioms. Let $M \in PC(T, \Gamma, L)$ and $M' \in EC(T, \Gamma)$ such that $M' \upharpoonright L = M$. Then $M' \upharpoonright L$ is the union of a directed system of K -structures of size $\text{LS}(\mathbf{K})$. By Fact 2.4(1) $M \in K$. Hence $PC(T, \Gamma, L) \subseteq K$.

2. By Fact 2.4(3).

3. By Fact 2.5.

□

Another question raised by Grossberg is the following:

Question 4.3. Is it possible to lower the bound of $|\Gamma|$ below $\text{LS}(\mathbf{K}) + I_2(\text{LS}(\mathbf{K}), \mathbf{K})$ in general? What if we also assume tameness or stability?

Remark 4.4. 1. In the above proof, we did not use σ_{\leq} because $\{f_k^n : k < \text{LS}(\mathbf{K}), n < \omega\}$ already plays its role. We could have done the same in Proposition 3.18 but the approach via σ_{\leq} is cleaner and does not add new function symbols.

2. One might want to encode $\bigvee_{i < I_2(\lambda, \mathbf{K})} \psi_i$ etc by omitting types without raising $|T|$ above $\text{LS}(\mathbf{K})$. However, it amounts to list all pairs of isomorphism types that are not any of the ψ_i 's. This will raise $|\Gamma|$ to $2^{\text{LS}(\mathbf{K})}$ which is equivalent to the original presentation theorem.

3. In [BB16, Theorem 3.2.3], new predicates are essentially added for our ϕ_i and ψ_j . Also, the requirement in [BB16, Definition 3.2.1(4)] encodes our $\{\sigma^{m,n,l} : m, n, l < \omega\}$ to build a directed system, but our version is purely syntactic and more direct. We used Chang's representation theorem to bring down the infinitary logic to first-order with omitting types. In [BB16], their theory T^* is still in the infinitary logic and thus does not need to omit types.

We derive the known results in the following corollary. Item (1) appeared for the first time in [Gro21] while (2) and (3) were undoubtedly known to Chang [Cha68].

Corollary 4.5. *Let \mathbf{K} be an AEC and $\chi := \text{LS}(\mathbf{K}) + I_2(\text{LS}(\mathbf{K}), \mathbf{K})$.*

1. *K is a $PC_{\chi}(= PC_{\chi,1})$ class.*

2. *The Hanf number of \mathbf{K} is bounded above by $\beth_{\delta(\chi,1)}$.*

3. If $\chi = \aleph_0$ or χ is a strong limit with $\text{cf}(\chi) = \aleph_0$, then the Hanf number of \mathbf{K} is bounded above by \beth_{χ^+} .

Proof. 1. Combine Theorem 4.1 and Shelah's 1-type coding [She90, VII Lemma 5.1(4)].

2. Combine (1) and [She90, VII Theorem 5.3].

3. Combine (2) and the fact that $\delta(\chi, 1) = \chi^+$ [She90, VII Theorem 5.5(5)].

□

We finish this section with one more application.

Definition 4.6. Let \mathbf{K} be an AEC. $K^{<} := \{ \langle |M|, |N| \rangle : N < M \}$ is a class of structures whose language consists of a single unary predicate.

In 1994, motivated by [She87, Theorem 3.8], Grossberg suggested the following conjecture (see Problem (5) in [She01, Introduction]):

Conjecture 4.7. Let \mathbf{K} be an AEC, $\lambda \geq \text{LS}(\mathbf{K})$. If $I(\lambda, \mathbf{K}) = I(\lambda^+, \mathbf{K}) = 1$, then $K_{\lambda^{++}} \neq \emptyset$.

[She87, Theorem 3.8] has two additional hypotheses:

1. Both K and $K^{<}$ are PC_λ ; and
2. $\delta(\lambda, 1) = \lambda^+$.

Much of [She01] is dedicated to the special cases of Grossberg's conjecture under various strong assumptions (including non-ZFC axioms).

Here we delete hypothesis (1) above and work in ZFC. In addition to hypothesis (2), we assume that $\lambda \geq \text{LS}(\mathbf{K}) + I_2(\text{LS}(\mathbf{K}), \mathbf{K})$.

Theorem 4.8. Let \mathbf{K} be an AEC, $\chi := \text{LS}(\mathbf{K}) + I_2(\text{LS}(\mathbf{K}), \mathbf{K})$, $\lambda \geq \chi$ with $\delta(\lambda, 1) = \lambda^+$. If \mathbf{K} is categorical in λ and λ^+ , then \mathbf{K} has a model of cardinality λ^{++} .

Remark 4.9. Our theorem applies to the case $\text{LS}(\mathbf{K}) = \aleph_1$, $\lambda = \beth_\omega$ while [She87, Theorem 3.8] cannot handle uncountable $L(\mathbf{K})$ or $\text{LS}(\mathbf{K})$.

Proof. We check that hypothesis (1) above is satisfied. Since $\lambda \geq \chi$, it suffices to show that both K and $K^<$ are PC_χ classes. K is a PC_χ class by Corollary 4.5. To show that $K^<$ is also a PC_χ class, we will use the proof of Theorem 4.1, add a new predicate R in L' and encode Proposition 3.18(2) by the new functions $\{f_k^n : k < \text{LS}(\mathbf{K}), n < \omega\}$ (to lighten the notation, we omit the encoding of re-enumerations, but it is the same strategy as in Theorem 4.1). At the end, we will only leave R in the reduct of the language.

The details are as follows: we expand the language L to L' which includes a new predicate R and the functions $\{f_k^n : k < \text{LS}(\mathbf{K}), n < \omega\}$ as in Theorem 4.1. For $n < \omega$, we abbreviate $\{f_k^n : k < \text{LS}(\mathbf{K})\}$ as \bar{f}^n and require that it maps an n -tuple to a model of size $\text{LS}(\mathbf{K})$ containing the tuple. This can be achieved by

$$\sigma^n := \forall a^n \bigwedge_{l < n} \bigvee_{k < \text{LS}(\mathbf{K})} (f_k^n(a^n) = a^n(l)) \wedge \bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(\bar{f}^n(a^n))$$

We also require that given an n -tuple inside R , the model generated is within R :

$$\sigma_R^n := \forall a^n \subseteq R (\bar{f}^n(a^n) \subseteq R)$$

Next, we want to require that the models generated are directed with respect to the tuple input. For $m, n, l < \omega$,

$$\sigma^{m,n,l} := \forall b^m \forall c^n \forall d^l \left[\text{ran}(b^m) \cup \text{ran}(c^n) \subseteq \text{ran}(d^l) \rightarrow \left(\bigvee_{i < I_2(\lambda, \mathbf{K})} \psi_i(\bar{f}^m(b^m), \bar{f}^l(d^l)) \wedge \bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(\bar{f}^n(c^n), \bar{f}^l(d^l)) \right) \right]$$

The final requirement is that R is a proper subset of the model:

$$\sigma_p := \exists x (\neg R(x))$$

Notice that for $m, n, l < \omega$, the sentences $\sigma^n, \sigma_R^n, \sigma^{m,n,l}$ are χ -junctions of (negation of) atomic formulas, so we can use Chang's presentation theorem to convert them to first-order formulas, by adding χ -many new predicates to L' to represent the χ -conjunctions and disjunctions, and χ -many L' -types to omit. This gives our T, Γ and L' .

We check that $K^< = PC(T, \Gamma, \{R\})$. If $\langle |M|, |N| \rangle$ is in $K^<$, then $N < M$ are in K . Expand the language to L' and define $R^M := N$. Inside N , build a directed system of

\mathbf{K} -substructures of size $\text{LS}(\mathbf{K})$, indexed by the finite tuples of N . This determines $\bar{f}^n \upharpoonright R^n$ for $n < \omega$. Now inside M , we extend the directed system to be indexed by the finite tuples of M . This determines \bar{f}^n completely for $n < \omega$. Also, M satisfies σ^n , σ_R^n and $\sigma^{m,n,l}$ for $m, n, l < \omega$. Hence M under the expanded language is in $EC(T, \Gamma)$ and its reduct to $\{R\}$ is in $PC(T, \Gamma, \{R\})$. Conversely, if $M \in PC(T, \Gamma, \{R\})$, expand M to M' such that $M' \in EC(T, \Gamma)$ and define $N' := R^{M'}$. By $\{\sigma^n, \sigma_R^n, \sigma^{m,n,l} : m, n, l < \omega\}$, $N' \upharpoonright L$ is the union of the directed system of \mathbf{K} -structures of size $\text{LS}(\mathbf{K})$. By Fact 2.4(1), $N' \upharpoonright L \in K$. By $\{\sigma^n, \sigma^{m,n,l} : m, n, l < \omega\}$, the directed system can be extended to union $M' \upharpoonright L$. By Fact 2.4(1) again, $M' \upharpoonright L \in K$. By Fact 2.4(2), each \mathbf{K} -structure of the directed system is a \mathbf{K} -substructure $M' \upharpoonright L$. But then the models of the original system that generates $N' \upharpoonright L$ are all \mathbf{K} -substructures of $M' \upharpoonright L$. By Fact 2.4(3), $N' \upharpoonright L \leq M' \upharpoonright L$. By σ_p , $N' \upharpoonright L < M' \upharpoonright L$. In other words, $\langle |M'|, |N'| \rangle = \langle |M|, R^M \rangle = M \in K^<$. \square

As in Section 3, we can add extra assumptions to improve our results:

Corollary 4.10. *1. If \mathbf{K} is a universal class, then $\chi := \text{LS}(\mathbf{K}) + I_2(\text{LS}(\mathbf{K}), \mathbf{K})$ in Theorem 4.1, Corollary 4.5 and Theorem 4.8 can be replaced by $\chi := \text{LS}(\mathbf{K})$.*

2. If \mathbf{K} is categoical in $\text{LS}(\mathbf{K})$ and $\text{LS}(\mathbf{K})^+$, has AL and either

(a) \mathbf{K} has $\text{LS}(\mathbf{K})$ -AP; or

(b) $2^{\text{LS}(\mathbf{K})} < 2^{\text{LS}(\mathbf{K})^+}$

then K and $K^<$ are both $PC_{\text{LS}(\mathbf{K})}$ classes when restricted to models of size $\geq \text{LS}(\mathbf{K})^+$.

In either case, AL can be replaced by stability in $\text{LS}(\mathbf{K})$.

3. In (2), K can be made to a $PC_{\text{LS}(\mathbf{K})}$ class.

Proof sketch. 1. Combine the proof of Theorem 4.1, Corollary 4.5 and Theorem 4.8 with Proposition 3.10. The point is that we do not need to encode \mathbf{K} -substructure relation.

2. In Theorem 3.12 and Corollary 3.14, we used coherence to encode $a \leq b$ in the infinitary language: $a \subseteq b$ and there is c which is $(\text{LS}(\mathbf{K}), \omega)$ -limit over a and b . Thus we can add two more sets of functions $\{d_k^{m,l}, e_k^{m,l} : k < \text{LS}(\mathbf{K}), m + l < \omega\}$ (which

represent c) in addition to the original $\{g_k^{m,l}, h_k^{m,l} : k < \text{LS}(\mathbf{K}), m + l < \omega\}$ (which represent a and b) in Theorem 4.1.

3. It remains to handle the case when the models are of size $\text{LS}(\mathbf{K})$: add a disjunct to the theory T in Theorem 4.1 which stipulates that the model is generated by an element a^1 :

$$\exists a^1 \left(\forall x^1 \bigvee_{k < \text{LS}(\mathbf{K})} (f_k^1(a^1) = x^1) \wedge \phi(\{f_k^1(a^1) : k < \text{LS}(\mathbf{K})\}) \right).$$

□

Remark 4.11. 1. Observation 3.15 also applies to the above corollary. If $\text{LS}(\mathbf{K}) = \aleph_0$ in (3), then we obtain: if \mathbf{K} is stable in \aleph_0 , has \aleph_0 -AP, $I(\aleph_0, \mathbf{K}) \leq \aleph_0$ and $I(\aleph_1, \mathbf{K}) = 1$, then K is PC_{\aleph_0} . This special case (with the extra assumption of categoricity in \aleph_1) provides an alternative proof to [SV18, Theorem 4.2] which uses results from descriptive set theory.

2. We do not know if (3) also applies to $K^{<}$, for a similar reason in Corollary 3.20(2),(3).

5 GENERALIZATION TO μ -AECs

Our strategy of encoding AECs is also applicable to μ -AECs.

Definition 5.1. [BGL⁺16, Definitions 2.1,3.1] Let L be a $(< \mu)$ -ary language. A μ -AEC $\mathbf{K} = \langle K, \leq_{\mathbf{K}} \rangle$ in L satisfies the axioms (1)(2)(3)(4) in Definition 2.2 in addition to

- a. Directed system axioms: if $\langle M_i \in K : i \in I \rangle$ is a μ -directed system, then $M := \bigcup_{i \in I} M_i \in K$ and for all $i \in I$, $M_i \leq_{\mathbf{K}} M$. If in addition $N \in K$ with $M_i \leq_{\mathbf{K}} N$ for all $i \in I$, then $M \leq_{\mathbf{K}} N$.
- b. Löwenheim-Skolem axiom: there exists a cardinal $\lambda = \lambda^{< \mu} \geq |L(\mathbf{K})| + \mu$ such that for any $M \in K$, $A \subseteq |M|$, there is $N \leq_{\mathbf{K}} M$ such that $A \subseteq |N|$ with $\|N\| \leq |A|^{< \mu} + \lambda$. We call the minimum such cardinal the Löwenheim-Skolem number $\text{LS}(\mathbf{K})$.

The analogs of Main Theorem 3.7, Proposition 3.16 and Proposition 3.18 hold in μ -AECs.

Proposition 5.2. *Let \mathbf{K} be a μ -AEC in L and $\lambda := \text{LS}(\mathbf{K})$. K is axiomatizable by an $L_{(\lambda+I_2(\lambda, \mathbf{K}))^+, \lambda^+}$ sentence $\sigma_{\mathbf{K}}$. In other words, for any L -structure M , $M \in K$ iff $M \models \sigma_{\mathbf{K}}$.*

Proof. Similar to the proof in Main Theorem 3.7. The difference is that we allow the iterated use of Löwenheim-Skolem and coherence axioms $(\mu \cdot \mu)$ -many times instead of $(\omega \cdot \omega)$ -many. We give the details below:

As usual, list the isomorphism types $\{M/\cong : M \in K_\lambda\}$ by $\langle M_i : i < I(\lambda, \mathbf{K}) \rangle$ and those of $\{(M, N)/\cong : M \leq N \text{ in } K_\lambda\}$ by $\langle (M_j, N_j) : j < I_2(\lambda, \mathbf{K}) \rangle$. For $i < I(\lambda, \mathbf{K})$, let $\phi_i(x)$ be an L_{λ^+, λ^+} formula that encodes the isomorphism type of M_i with a fixed enumeration of the universe $|M| = \langle m_k^i : k < \lambda \rangle$. For variables $x = \langle x_k : k < \lambda \rangle$,

$$\phi_i(x) := \bigwedge \left\{ \theta(x_{\alpha_0}, \dots, x_{\alpha_\xi}) : M_i \models \theta[m_{\alpha_0}^i, \dots, m_{\alpha_\xi}^i]; \xi < \mu, \alpha_0, \dots, \alpha_\xi < \lambda, \right. \\ \left. \theta \text{ is an atomic } L\text{-formula or its negation with } s \text{ free variables} \right\}$$

Notice that ϕ_i is a conjunction of $\lambda^{<\mu} = \lambda$ many formulas so it is inside L_{λ^+, λ^+} . Similarly for $j < I_2(\lambda, \mathbf{K})$, let $\psi_j(x, y)$ be an L_{λ^+, λ^+} formula that encodes the isomorphism type of (M_j, N_j) with fixed enumerations, where $|M_j| = \{m_k^j : k < \lambda\}$, $|N_j| = \{n_k^j : k < \lambda\}$. For variables $x = \langle x_k : k < \lambda \rangle$ and $y = \langle y_k : k < \lambda \rangle$,

$$\psi_j(x; y) := \bigwedge \left\{ \theta(x_{\alpha_0}, \dots, x_{\alpha_\xi}; y_{\beta_0}, \dots, y_{\beta_{\xi'}}) : N_j \models \theta[m_{\alpha_0}^j, \dots, m_{\alpha_\xi}^j; n_{\beta_0}^j, \dots, n_{\beta_{\xi'}}^j], \right. \\ \left. \xi, \xi' < \mu; \alpha_0, \dots, \alpha_\xi, \beta_0, \dots, \beta_{\xi'} < \lambda, \right. \\ \left. \theta \text{ is an atomic } L\text{-formula or its negation with } \xi + \xi' \text{ free variables} \right\}$$

ψ_j is also a conjunction of $\lambda^{<\mu} = \lambda$ many formulas so it is inside L_{λ^+, λ^+} . The axiomatization $\sigma_{\mathbf{K}}$ consists of two components (the variables all have length λ):

$$\sigma_{\mathbf{K}} := (\forall x_\alpha \exists y_\alpha)_{\alpha < \mu \cdot \mu} \bigwedge_{\alpha < \mu \cdot \mu} \left[(x_\alpha \subseteq y_\alpha) \wedge \exists z_\alpha \left((y_\alpha \approx z_\alpha) \wedge \bigvee_{i < I(\lambda, \mathbf{K})} \phi_i(z_\alpha) \right) \wedge \right. \\ \left. \bigwedge_{\beta < \alpha < \mu \cdot \mu} \exists u_{\beta, \alpha} \exists v_{\beta, \alpha} \left((y_\beta \approx u_{\beta, \alpha}) \wedge (y_\alpha \approx v_{\beta, \alpha}) \wedge \bigvee_{j < I_2(\lambda, \mathbf{K})} \psi_j(u_{\beta, \alpha}, v_{\beta, \alpha}) \right) \right]$$

Suppose $M \in K$, we show that Player II can win the associated game in $\sigma_{\mathbf{K}}$. In the α -th round, Player I provides some x_α of length λ . By Löwenheim-Skolem axiom, pick any

$y_\alpha \leq M$ of size λ such that $\text{ran}(x_\alpha) \cup \bigcup_{\beta < \alpha} \text{ran}(y_\beta) \subseteq \text{ran}(y_\alpha)$. By inductive hypothesis, for $\beta < \alpha$, we have $y_\beta \leq M$. By coherence axiom, $y_\beta \leq y_\alpha$ as desired.

Suppose $M \models \sigma_{\mathbf{K}}$. We will build a μ -directed system $\langle M_a \in K_\lambda : a \in I \rangle$ of union M , with I being the set of tuples of length $(< \mu)$ in M , ordered by inclusion. By directed system axioms, $M \in K$. In the proof of Main Theorem 3.7, we showed that given M_a and M_b generated by the games of the singletons s and t , it is possible to find $M^* \geq M_a, M_b$ by extending those games by ω -many rounds. In the μ -AEC case, without the usual chain axioms we do not know if $M^* \in K$, so we extend those games by μ -many rounds instead to obtain an increasing (but not necessarily continuous) chain $\langle N_k : k < \mu \rangle$ and define $M^* = \bigcup_{k < \mu} N_k \in K$. A similar argument shows: let $\delta < \mu$ and if for $\alpha < \delta$, M_α is generated by the game of some tuple a_α of length $< \mu$, then we can extend the games by μ -many rounds to obtain M^* that extends all M_α . This allows us to get past the limit stages which were absent in the original proof, and continue to build M_a for $l(a) < \mu$. Given a tuple c of length $< \mu$, there are less than μ -many ways to decompose c into a union of a singleton and a tuple of length $< \mu$. Thus we can still combine all copies of M^* from the decompositions of M_c as in the original proof. \square

Proposition 5.3. *Let \mathbf{K} be a μ -AEC in L and $\lambda := \text{LS}(\mathbf{K})$. There is a formula in $L_{(\lambda + I_2(\lambda, \mathbf{K}))^+, \lambda^+, (\mu \cdot \mu)}$ that encodes the \mathbf{K} -substructure relation: for any $M \in K$, $a \subseteq |M|$ of size λ , $M \models \sigma_{\leq}[a]$ iff $a \leq M$ (the enumeration of a does not matter).*

Proof. Define $\sigma_{\leq}(x)$ as in Proposition 3.16 but replace $\omega \cdot \omega$ by $\mu \cdot \mu$. The enumeration of a does not matter by our definition of \approx . If $a \in K_\lambda$ and $a \leq M$, then $M \models \sigma_{\leq}[a]$ by Löwenheim-Skolem and coherence axioms. Conversely suppose $a \subseteq |M|$ of size λ and $M \models \sigma_{\leq}[a]$. As in Proposition 5.2, we can build a μ -directed system $\langle M_\alpha \in K_\lambda : \alpha \in I \rangle$ of union M such that for any $\alpha \in I$, $M_\alpha \geq a$. By directed system axioms, $M \in K$ and $M_\alpha \leq M$ for all $\alpha \in I$. By transitivity of \leq , $a \leq M$ as desired. \square

Proposition 5.4. *Let \mathbf{K} be a μ -AEC in L and $\lambda := \text{LS}(\mathbf{K})$. Let $M, N \in K$.*

1. $M \leq N$ iff $M \subseteq_{\{\sigma_{\leq}\}} N$ (if $a \subseteq |M|$ is of size λ , then $M \models \sigma_{\leq}[a]$ iff $N \models \sigma_{\leq}[a]$).

2. Let R be a new predicate where $N^R = |M|$ closed under permutations. $M \leq N$ iff $(N, R) \models \forall b (\sigma_{\leq}^R(b) \rightarrow \sigma_{\leq}(b))$ where σ_{\leq}^R is the relativized version of σ_{\leq} inside R (replace $(\omega \cdot \omega)$ by $(\mu \cdot \mu)$ in the definition of σ_{\leq}^R in Proposition 3.18).

Proof. Similar to the proof in Proposition 3.18. The difference is that instead of building \aleph_0 -directed systems, we build μ -directed systems. We give details below:

1. If $M \leq N$ and let $a \subseteq M$. If $M \models \sigma_{\leq}[a]$, then $a \leq M \leq N$ showing $N \models \sigma_{\leq}[a]$. If $N \models \sigma_{\leq}[a]$, then $a \leq N$. By coherence, $a \leq M$ and so $M \models \sigma_{\leq}[a]$. Conversely, build a μ -directed system $\langle M_\alpha \in K_\lambda : \alpha \in I \rangle$ inside M such that for all $\alpha \in I$, $M_\alpha \leq M$. Then $M \models \sigma_{\leq}[M_\alpha]$. Since $M \subseteq_{\{\sigma_{\leq}\}} N$, we have $N \models \sigma_{\leq}[M_\alpha]$ and $M_\alpha \leq N$. The result follows from directed system axioms.
2. If $M \leq N$ and $N \models \sigma_{\leq}^R[b]$ for some $b \subseteq |N|$, we need to show that $N \models \sigma_{\leq}[b]$. By assumption we can build a μ -directed system of union M and have $b \leq M$. By transitivity of \leq , $b \leq N$ and the conclusion follows. Conversely, by directed system axioms, it suffices to build a μ -directed system $\langle M_\alpha \in K_\lambda : \alpha \in I \rangle$ of union M such that for all $\alpha \in I$, $M_\alpha \leq N$. Since $(N, R) \models \forall b (\sigma_{\leq}^R(b) \rightarrow \sigma_{\leq}(b))$, we can require $M_\alpha \leq M$ instead of $M_\alpha \leq N$. Such construction is possible by Löwenheim-Skolem and coherence axioms.

□

As an application of Proposition 5.2 and Proposition 5.4, we generalize Corollary 3.19:

Corollary 5.5. *Let \mathbf{K} be a μ -AEC in L , $\lambda := \text{LS}(\mathbf{K})$ and M, N be L -structures. If either M or N is in K and $M \preceq_{L_{\infty, \lambda^+}(\mu \cdot \mu)} N$, then $M \leq N$ (and both are in K).*

Proof. Same proof as in Corollary 3.19: Since $M \preceq_{L_{\infty, \lambda^+}} N$, $M \subseteq_{L_{(\lambda + I_2(\lambda, \mathbf{K}))^+, \lambda^+(\mu \cdot \mu)}} N$. In particular $M \models \sigma_{\mathbf{K}}$ iff $N \models \sigma_{\mathbf{K}}$. By Proposition 5.2, either M, N is in K implies both are in K . On the other hand, the assumption implies $M \subseteq_{\{\sigma_{\leq}\}} N$. By Proposition 5.4(1), $M \leq N$. □

We now state the μ -AEC version of Theorem 4.1, which is a variation to [BGL⁺16, Theorem 3.2].

Theorem 5.6. *Let \mathbf{K} be a μ -AEC in L and with Löwenheim-Skolem number $\text{LS}(\mathbf{K})$. Define $\chi := \text{LS}(\mathbf{K}) + I_2(\text{LS}(\mathbf{K}), \mathbf{K})$. There exists a $(< \mu)$ -ary expansion $L' \supseteq L$ of size χ , an L' -theory T and a set of L' -types Γ of size χ such that*

1. $K = PC^\mu(T, \Gamma, L)$.
2. If $M', N' \in EC(T, \Gamma)$ and $M' \subseteq_{L'} N'$, then $M' \upharpoonright L \leq_{\mathbf{K}} N' \upharpoonright L$.
3. If $M \leq_{\mathbf{K}} N$, there are L' -expansions of M, N to M', N' such that $M' \subseteq_{L'} N'$.

Proof sketch. Repeat the same argument in Theorem 4.1 by replacing ω by μ , in particular:

1. Superscripts of f, g, h will be $\alpha, \beta, \gamma < \mu$ instead of $m, n, l < \omega$.
2. We require that the K -substructures generated by $\{f_k^\alpha : k < \text{LS}(\mathbf{K}), \alpha < \omega\}$ are μ -directed instead of \aleph_0 -directed.
3. The sentences $\{\sigma^\alpha : \alpha < \mu\} \cup \{\sigma^{\alpha, \beta, \gamma} : \alpha, \beta, \gamma < \mu\}$ are in $L'_{\chi^+, \mu}$.
4. For $i < I(\text{LS}(\mathbf{K}), \mathbf{K})$ and $j < I_2(\text{LS}(\mathbf{K}), \mathbf{K})$, the formulas ϕ_i, ψ_j are still $\text{LS}(\mathbf{K})$ -conjunctions because $\text{LS}(\mathbf{K})^{<\mu} = \text{LS}(\mathbf{K})$.
5. Chang's presentation theorem generalizes to μ -AECs and converts a $L'_{\chi^+, \mu}$ theory of size χ into a PC^μ_χ .
6. When checking the items of the theorem statement, notice that by definition of a μ -AEC, directed system axioms (instead of chain axioms) are built-in. Meanwhile, Fact 2.5 generalizes to μ -directed systems.

□

Unlike Corollary 4.5, the above result does not lead to the Hanf number computation because the languages are not finitary while well-ordering is definable. In particular there is no reasonable bound to the Hanf number of L_{\aleph_1, \aleph_1} [Dic75, Chapter 5.1B]. As asked in [BGL⁺16, Remark 3.3]:

Question 5.7. Let $\mu \geq \aleph_1$. Does the Hanf number exist for μ -AECs?

REFERENCES

- [Bal09] John T. Baldwin. *Categoricity*. University lecture series; v. 50. American Mathematical Society, Providence, R.I, 2009.
- [BB16] Will Boney and John Baldwin. Hanf numbers and presentation theorems in AECs. 2016. Preprint: <https://arxiv.org/abs/1511.02935>.
- [BGL⁺16] Will Boney, Rami Grossberg, Michael Lieberman, Jiří Rosický, and Sebastien Vasey. μ -abstract elementary classes and other generalizations. *Journal of Pure and Applied Algebra*, 220(9):3048–3066, 2016.
- [Cha68] Chen Chung Chang. Some remarks on the model theory of infinitary languages. In Jon Barwise, editor, *The Syntax and Semantics of Infinitary Languages*, pages 36–63, Berlin, Heidelberg, 1968. Springer.
- [Dic75] M. A. Dickmann. *Large Infinitary Languages*, volume 83 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 1975.
- [Gro02] Rami Grossberg. Classification theory for abstract elementary classes. *Logic and Algebra*, ed. Yi Zhang, *Contemporary Mathematics*, Vol 302, AMS:165–204, 2002.
- [Gro21] Rami Grossberg. *A Course in Model Theory I: Introduction*. 2021. A book in preparation.
- [Hen19] Simon Henry. An abstract elementary class nonaxiomatizable in $L_{\infty, \kappa}$. *The Journal of Symbolic Logic*, 84(3):1240–1251, 2019.
- [Joh10] Gregory Johnson. Abstract elementary classes with Löwenheim-Skolem number cofinal with ω . *Notre Dame Journal of Formal Logic*, 51(3):361–371, 2010.
- [Kol85] Ph. G. Kolaitis. Game quantification. In J. Barwise and S. Feferman, editors, *Model-Theoretic Logics*, New York, 1985. Springer-Verlag.
- [Kue77] David Kueker. Countable approximations and Löwenheim-Skolem theorems. *Annals of Mathematical Logic*, 11(1):57–103, 1977.
- [Kue08] David Kueker. Abstract elementary classes and infinitary logics. *Annals of Pure and Applied Logic*, 156(2):274–286, 2008.

- [Leu23] Samson Leung. Hanf number of the first stability cardinal in AECs. *Annals of Pure and Applied Logic*, 174(2):103201, 2023.
- [MAV18] Marcos Mazari-Armida and Sebastien Vasey. Universal classes near \aleph_1 . *The Journal of Symbolic Logic*, 83(4):1633–1643, 2018.
- [She87] Saharon Shelah. Classification of nonelementary classes. II. Abstract elementary classes. In *Classification Theory (Chicago, IL, 1985)*, volume 1292 of *Lecture Notes in Math.*, pages 419–497. Springer, Berlin, 1987.
- [She90] Saharon Shelah. *Classification Theory and the Number of Non-isomorphic Models*. Studies in logic and the foundations of mathematics; v. 92. North-Holland, Amsterdam, rev. edition, 1990.
- [She99] Saharon Shelah. Categoricity for abstract classes with amalgamation. *Ann. Pure Appl. Logic*, 98(1-3):261–294, 1999.
- [She01] Saharon Shelah. Categoricity of an abstract elementary class in two successive cardinals. *Israel J. Math.*, 126:29–128, 2001.
- [She09] Saharon Shelah. *Classification Theory for Abstract Elementary Classes*, volume 18 of *Studies in Logic (London)*. College Publications, London, 2009.
- [SV18] Saharon Shelah and Sebastien Vasey. Abstract elementary classes stable in \aleph_0 . *Annals of Pure and Applied Logic*, 169(7):565–587, 2018.
- [SV21] Saharon Shelah and Andrés Villaveces. Infinitary logics and abstract elementary class. *Proceedings of the American Mathematical Society*, pages 1–, 2021.
- [Vas17] Sebastien Vasey. Saturation and solvability in abstract elementary classes with amalgamation. *Archive for Mathematical Logic*, 56(5):671–690, 2017.
- [Vas18] Sebastien Vasey. Math 269X - model theory for abstract elementary classes, spring 2018 lecture notes. <https://svasey.github.io/academic-homepage-may-2020/aec-spring-2018/aec-lectur>
April 2018.

Email: wangchil@andrew.cmu.edu

URL: <http://www.math.cmu.edu/~wangchil/>

Address: Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh PA 15213, USA