

BROWN MEASURE OF THE SUM OF AN ELLIPTIC OPERATOR AND A FREE RANDOM VARIABLE IN A FINITE VON NEUMANN ALGEBRA

PING ZHONG

ABSTRACT. Given a $n \times n$ random matrix X_n with i.i.d. entries of unit variance, the circular law says that the empirical spectral distribution (ESD) of X_n/\sqrt{n} converges to the uniform measure on the unit disk. Let M_n be a deterministic matrix that converges in $*$ -moments to an operator x_0 . It is known from the work by Śniady and Tao–Vu that the ESD of $X_n/\sqrt{n} + M_n$ converges to the Brown measure of $x_0 + c$, where c is Voiculescu’s circular operator. We obtain a formula for the Brown measure of $x_0 + c$ which provides a description of the limit distribution. This answers a question of Biane–Lehner for arbitrary operator x_0 .

The twisted elliptic operators include circular operator and semi-circular operator. Let c_t be a circular operator with variance t and let $g_{t,\gamma}$ be a twisted elliptic operator free from x_0 . The Brown measure of $x_0 + g_{t,\gamma}$ is not calculated directly because of degeneracy. We show that the Brown measure of $x_0 + g_{t,\gamma}$ is the push-forward measure of the Brown measures of $x_0 + c_t$ under a natural map, provided that this map is one-to-one and non-singular. We calculate explicit formula for the case x_0 is selfadjoint. In addition, we prove that the Brown measure of the sum of a Haar unitary operator and a twisted elliptic element is supported in a deformed ring where the inner boundary is a circle and the outer boundary is an ellipse, and its density is constant along a family of ellipses.

These results generalize some known results about free additive Brownian motions where the free random variable x_0 is assumed to be selfadjoint. The approach is based on a Hermitian reduction and subordination functions.

CONTENTS

1. Introduction	2
1.1. Brown measure of free random variables and random matrices	2
1.2. Statements of the results	3
1.3. Discussions on methodologies	6
2. Preliminaries	7
2.1. Free probability and subordination functions	7
2.2. The Brown Measure	8
2.3. Hermitian reduction method for the sum of two free random variables	9
2.4. The elliptic operator and the operator-valued semicircular element	11
3. The Fuglede-Kadison determinant and subordination functions	12
3.1. Free convolution with a semicircular distribution	12
3.2. The operator-valued subordination functions	15
3.3. The coupling Fuglede-Kadison determinants	17
4. The Brown measure of the addition with an elliptic operator	21
4.1. The gradient functions	22
4.2. The addition with a circular operator	25
4.3. A general result on the support of the Brown measure	27

Supported in part by Collaboration Grants for Mathematicians from Simons Foundation.

4.4. Addition with an elliptic operator	29
5. Addition with a selfadjoint operator	32
5.1. Subordination functions	32
5.2. The push-forward map	34
5.3. Brown measure of addition with a selfadjoint operator	37
6. Example of non-selfadjoint operators	41
6.1. The quasi-nilpotent DT operator	41
6.2. The Haar unitary	43
References	49

1. INTRODUCTION

1.1. Brown measure of free random variables and random matrices. Let \mathcal{M} be a von Neumann algebra with a faithful, normal, tracial state ϕ . The Fuglede-Kadison determinant of $x \in \mathcal{M}$ is defined by

$$\Delta(x) = \exp \left(\int_0^\infty \log t d\mu_{|x|}(t) \right),$$

where $\mu_{|x|}$ is the spectral measure of $|x|$ with respect to ϕ . Brown [14] proved that the function $L_x(\lambda) = \log \Delta(x - \lambda \mathbf{1})$ is a subharmonic function whose Riesz measure is the unique, compactly supported probability measure μ_x on \mathbb{C} with the property that

$$\log \Delta(x - \lambda \mathbf{1}) = \int_{\mathbb{C}} \log |z - \lambda| d\mu_x(z), \quad \lambda \in \mathbb{C}.$$

The measure μ_x is called the *Brown measure* of x . In other words, μ_x is the distributional Laplacian of the function $L_x(\lambda)$.

Voiculescu's free probability theory is a suitable framework to describe the limits of the joint distribution of a family of random matrix models [42]. The convergence of *-mixed moments of suitable random matrix models have been studied well. The *-mixed moments of free random variables can be described using either analytic or combinatorial tools. For non-normal random matrix models, very little is known about the limit of the empirical spectral distribution (ESD) of a polynomial of independent random matrices, even for the sum or product of two random matrices. The Brown measure of an operator in \mathcal{M} is an analogue of eigenvalue distribution of a finite dimensional matrix. The Brown measures of the sum or product or a polynomial of free random variables are natural candidates for the limits of the ESD of the sum or product or a polynomial of suitable random matrix models as the size of the matrices tends to infinity.

Let X_n be an $n \times n$ random matrix whose entries are independent identically distributed copies of a complex random variable with zero mean and unit variance. The circular law says that the ESD of X_n/\sqrt{n} converges weakly to the uniform measure on the unit disc which is also the Brown measure of Voiculescu's circular operator. The circular law was established in the 1960s by Ginibre [21] for Gaussian distributed entries and was proved by Tao and Vu [41] under the minimal assumptions after a long list of partial progresses (see [13] and references therein). In fact, Tao and Vu proved results stronger than circular law. In particular, they showed the existence of the limit of the summation $X_n/\sqrt{n} + M_n$ where M_n is a deterministic $n \times n$ matrix satisfying some technical conditions. In [39], Śniady showed that the ESD of $X_n/\sqrt{n} + M_n$ converges to the Brown measure of $x_0 + c_t$ provided that X_n is a Ginibre ensemble and M_n converges in *-moments to x_0 . Hence, by

combining Tao–Vu’s replacement principle [41, Section 2], we conclude that the ESD of $X_n/\sqrt{n} + M_n$ converges to the Brown measure of $x_0 + c_t$ under the minimal requirements on X_n when M_n converges in $*$ -moments to x_0 .

In the above-cited paper, Tao and Vu did not pursue what the limit actually is (see [41, Theorem 1.17]) and they mentioned that the limit distribution ESD of $X_n/\sqrt{n} + M_n$ was established in a work of Krishnapur–Vu for the case where M_n is a diagonal matrix (equivalently, x_0 is a normal operator as the limit) and X_n is a Ginibre ensemble. But that work has not appeared even as a preprint (communicated with Krishnapur). Here we study the Brown measure of the sum of free circular operator and a $*$ -free random variable x_0 with an arbitrary distribution, not necessarily normal, which fills up the gap of the description of the limit ESD of $X_n/\sqrt{n} + M_n$. The case when x_0 is selfadjoint was known in the author’s joint work with Ho [28] using PDE methods. This answers an earlier question of Biane–Lehner [12, Section 5] in general case.

The twisted elliptic operators generalize circular operator, semi-circular operator and elliptic operator. Let c_t be a circular operator with variance t , and let $g_{t,\gamma}$ be a twisted elliptic operator, and let x_0 be an operator $*$ -free from $\{c_t, g_{t,\gamma}\}$. The calculation of the Brown measure of $x_0 + g_{t,\gamma}$ is more evolved than $x_0 + c_t$ because there is possibly degeneracy. We show that the Brown measure of $x_0 + g_{t,\gamma}$ is the push-forward measure of the Brown measures of $x_0 + c_t$ under a natural map, provided that this map is one-to-one and non-singular. We calculate explicit density Brown measure formulas for the case where x_0 is selfadjoint. In addition, we calculate the Brown measure of the sum of a Haar unitary operator and a twisted elliptic element. The Brown measure is supported in a deformed ring where the inner boundary is a circle and the outer boundary is an ellipse, and its density is constant along a family of ellipses. This can be viewed as a deformation of the limit distribution in the single ring theorem [22] in random matrix theory.

The present work extends previous results [12, 28, 31, 32] for the sum of a selfadjoint operator with a circular operator or a (non-twisted) elliptic operator. All these work rely on some PDE methods. We use a completely different approach based on Hermitian reduction and subordination functions. The new method provides a conceptual explanation why subordination functions appeared in Brown measure formulas. The method and subordination results developed in this paper are likely to be useful in the study of non-normal random matrices.

1.2. Statements of the results. Let $c_t \in \mathcal{M}$ be Voiculescu’s circular operator with variance t . Let $g_{t,\gamma}$ be a twisted elliptic operator with parameters $t > 0$ and $\gamma \in \mathbb{C}$ such that $|\gamma| \leq t$ (see Section 2.4 for the definition). Such operator has the same distribution as an operator of the form $e^{i\theta}(s_1 + is_2)$ where $\theta \in [0, 2\pi]$, and s_1, s_2 are semicircular operators that are freely independent in the sense of Voiculescu. The case when $\gamma = t$, the operator $g_{t,\gamma}$ is the semicircular operator g_t with mean zero and variance t , and the case when $\gamma \in \mathbb{R}$ the operator $g_{t,\gamma}$ is an elliptic operator.

Let $x_0 \in \mathcal{M}$ be a random variable that is free from $\{c_t, g_{t,\gamma}\}$. We show that the Brown measure of $x_0 + c_t$ can be calculated directly using subordination functions in free probability under some mild assumptions. The Brown measure of $x_0 + g_{t,\gamma}$ is not calculated directly. Instead, we show that there is a natural push forward map between the Brown measure of $x_0 + c_t$ and the Brown measure of $x_0 + g_{t,\gamma}$ under certain assumptions. We then show that a large family of operators including all selfadjoint operators satisfy the assumption and this allows us to extend results in [28, 31, 32]. Our main result is also applicable to non-selfadjoint operators.

To describe our main results, we need some terminology. Fix $t > 0$ and $x_0 \in \mathcal{M}$. Consider the open set (see Proposition 4.2)

$$\Xi_t = \left\{ \lambda \in \mathbb{C} : \phi \left[((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}))^{-1} \right] > \frac{1}{t} \right\}.$$

Fix $\lambda \in \Xi_t$ and let $w = w(0; \lambda, t)$ be a function of λ taking positive values such that

$$\phi((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w^2 \mathbf{1})^{-1} = \frac{1}{t}.$$

We denote

$$\Phi_{t,\gamma}(\lambda) = \lambda + \gamma \cdot p_\lambda^{(0)}(w), \quad \lambda \in \Xi_t,$$

where

$$p_\lambda^{(0)}(w) = -\phi \left[(x_0 - \lambda \mathbf{1})^* ((x_0 - \lambda \mathbf{1})(x_0 - \lambda \mathbf{1})^* + w^2 \mathbf{1})^{-1} \right].$$

We show that the function w of λ is the imaginary part of the subordination function that appears in free additive convolution of two probability measures on \mathbb{R} (see Proposition 3.5 for precise statement) and this function is a real analytic function of $\lambda \in \Xi_t$. The function $\Phi_{t,\gamma}$ is a real analytic function of λ in the set Ξ_t .

Theorem A (See Theorem 3.11 and Theorem 3.12). *For every $\lambda \in \Xi_t$, we have*

$$(1.1) \quad \Delta(x_0 + c_t - \lambda)^2 = \Delta((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w^2 \mathbf{1}) [\exp(-w^2/t)],$$

and

$$(1.2) \quad \Delta(x_0 + g_{t,\gamma} - z)^2 = \Delta(x_0 + c_t - \lambda \mathbf{1})^2 \exp(H(\lambda)),$$

where $z = \Phi_{t,\gamma}(\lambda)$ and $H(\lambda) = \Re \left(\gamma (p_\lambda^{(0)}(w))^2 \right)$.

We point out that we also have $\Delta(x_0 + c_t - \lambda \mathbf{1}) = \Delta(x_0 - \lambda \mathbf{1})$ for any $\lambda \in \mathbb{C} \setminus \Xi_t$ (see Theorem 3.12) which implies that $\mu_{x_0+c_t}$ and μ_{x_0} coincide in the interior of $\mathbb{C} \setminus \Xi_t$. Moreover, we obtain a general result on the support of the Brown measure (see Theorem 4.9) which allows us to deduce that the interior of $\mathbb{C} \setminus \Xi_t$ is not in the support of μ_{x_0} . Hence, our focus is the Brown measure of $\mu_{x_0+c_t}$ within the set Ξ_t . The above Fuglede-Kadison formulas are fundamental in our study which allows us to calculate the Brown measure formulas. The Brown measure of $x_0 + c_t$ can be described as follows.

Theorem B (See Theorem 4.6 and Theorem 4.10). *The Brown measure of $x_0 + c_t$ is supported in Ξ_t and is absolutely continuous with respect to Lebesgue measure in the open set Ξ_t . The density of the Brown measure at any $\lambda \in \Xi_t$ is strictly positive and can be expressed as*

$$(1.3) \quad \frac{1}{\pi} \left(\frac{1}{t} - \frac{\partial}{\partial \lambda} \left(\phi(x_0^* ((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w^2 \mathbf{1})^{-1}) \right) \right).$$

At a first glance, the implicit density formula (1.3) is not good enough for applications. It turns out that the subordination function $w(0; \lambda, t)$ can be calculated explicitly for a large family of operators that include all selfadjoint operators and many non-selfadjoint operators. This generalizes a result in our earlier work with Ho [32] in which x_0 is selfadjoint. Unlike [32] and its generalizations for semicircular operators and elliptic operators [28, 31], we do not use PDE methods.

Theorem C (See Theorem 4.16). *Assume $\mu_{x_0+c_t}(\partial(\Xi_t)) = 0$. If $\Phi_{t,\gamma}$ is one-to-one and non-singular in the set Ξ_t , then the Brown measure of $x_0+g_{t,\gamma}$ is the push-forward measure of the Brown measure of x_0+c_t by the map $\lambda \mapsto \Phi_{t,\gamma}(\lambda)$. Let $\Gamma \subset \Xi_t$ be a simply connected domain in the set Ξ_t with piecewise smooth boundary such that $\bar{\Gamma} \subset \Xi_t$. Then*

$$(1.4) \quad \mu_{x_0+c_t}(\Gamma) = \mu_{x_0+g_{t,\gamma}}(\Phi_{t,\gamma}(\Gamma)).$$

This result generalizes results from [28, 31, 32] in two directions. Theorem C is applicable for operators x_0 not necessarily selfadjoint. In addition, the twisted elliptic operator include semicircular operators and elliptic operators as special cases. What is more, the proof of Theorem C provides a conceptual explanation about why such push-forward map exists. It is based on the following observation: (i) The gradient of $L_{x_0+c_t}$ at λ is equal to the gradient of $L_{x_0+g_{t,\gamma}}$ at $\Phi_{t,\gamma}(\lambda)$ for any $\lambda \in \Xi_t$; (ii) The map $\Phi_{t,\gamma}$ is the sum of the identity map and a linear combination of the gradient function. See Lemma 4.4 and Theorem 4.16 for detailed descriptions.

We verify that for a large family of operators x_0 , the map $\Phi_{t,\gamma}$ is one-to-one and non-singular in Ξ_t . However, we believe that the condition that $\Phi_{t,\gamma}$ is one-to-one and non-singular in Theorem C is redundant. We will address this question in a future project.

Theorem D (See Theorem 5.14 and Example 5.18). *Let x_0 be a selfadjoint operator that is $*$ -free from $g_{t,\gamma}$. For any $|\gamma| \leq t$ with $\gamma \neq t$, the map $\Phi_{t,\gamma}$ is one-to-one and non-singular in Ξ_t . The Brown measure of $x_0+g_{t,\gamma}$ is the push-forward measure of the Brown measure of x_0+c_t under the map $\Phi_{t,\gamma}$.*

Moreover, the Brown measure $\mu_{x_0+g_{t,\gamma}}$ is concentrated on $\Phi_{t,\gamma}(\Xi_t)$ and the density is given by

$$(1.5) \quad d\mu_{x_0+g_{t,\gamma}}(z) = \frac{1}{2\pi\tau_1} \frac{d\psi_t(a)}{d\delta(a)} dz_1 dz_2, \quad z \in \Phi_{t,\gamma}(\Xi_t)$$

where $z = z_1 + iz_2 = \Phi_{t,\gamma}(a + ib)$, $\tau_1 = t - \Re(\gamma)$, and ψ_t, δ are two increasing homeomorphisms of \mathbb{R} onto \mathbb{R} . In particular, if $\gamma \in \mathbb{R}$ (equivalently, $g_{t,\gamma}$ is an elliptic operator), then $z_1 = \delta(a)$ depending only on a , in which case the Brown measure is constant along vertical lines.

The density formula could be understood as follows. For any selfadjoint operator x_0 , it is known [10] that there is a continuous function v_t such that

$$\Xi_t = \{a + ib \in \mathbb{C} : |b| < v_t(a)\}.$$

See Section 5.1 for a review. It is shown [32] that the density of x_0+c_t is constant along vertical segments in Ξ_t . In this case, for any fixed $a \in \mathbb{R}$ such that the vertical line through a intersects the set Ξ_t , the map $b \mapsto \Phi_{t,\gamma}(a + ib)$ is an affine transformation of b . Hence, the Brown measure of $x_0+g_{t,\gamma}$ is expected to be constant along the trajectory of $\Phi_{t,\gamma}(a + ib)$ as b varies in $(-v_t(a), v_t(a))$ such that $a + ib$ changes within Ξ_t . The formula (1.5) describe precisely this observation. Indeed, as $\Phi_{t,\gamma}$ is one-to-one under the assumption of Theorem D, the set $\Phi_{t,\gamma}(\Xi_t)$ can be parametrized by $a + ib \in \Xi_t$ under the push-forward map $\Phi_{t,\gamma}$. Hence, we can say that the density formula (1.5) depends on only one parameter and is constant in one direction. Theorem D can be viewed as an analogue result for the free additive convolution in a recent work of Hall–Ho [29] concerning free multiplicative Brownian motions. See Remark 5.19 for details.

Finally, we demonstrate the application of Theorem C to some non-selfadjoint operators. We study the quasi-nilpotent DT operator (introduced by Dykema and Haagerup [15]) and Haar unitary operator. For simplification, we only state the result for the case $g_{t,\gamma}$ is a semicircular operator g_t and x_0 is a Haar unitary operator u .

Theorem E (See Theorem 6.10 and Theorem 6.11). *The support of the Brown measure of $u + g_t$ is the deformed single ring where the inner boundary is the circle centered at the origin with radius $\sqrt{(1-t)_+}$ and the outer boundary is the ellipse centered at the origin with semi-axes $\frac{2t+1}{\sqrt{t+1}}$ and $\frac{1}{\sqrt{1+t}}$. The Brown measure is absolutely continuous and its density is strictly positive in the support.*

The Brown measure of $x_0 + g_t$ is the push-forward measure of the Brown measure $x_0 + c_t$ by the map $\Phi_{t,t}$. The map sends the family of circles centered at the origin with radius r to the family of ellipses centered at the origin. Moreover, the Brown measure of $x_0 + g_t$ is constant along the ellipses.

1.3. Discussions on methodologies. Our approach is based on a Hermitian reduction method and subordination functions. The Hermitian reduction method was already used for the calculation of Brown measure of quasi-nilpotent DT operators in Aagaard–Haagerup’s work [1]. In physics literature, to our knowledge, the method was first used in two independent work [19] and [33]. In the work by Jarosz and Nowak [34, 35], the authors used Hermitian reduction approach to study the support of the Brown measure of $x_0 + ig_t$ for selfadjoint x_0 , where the method is not mathematically rigorous as written. The Hermitian reduction also appeared in Voiculescu’s earlier work [43] which serves a motivation to introduce free probability theory with amalgamation. The idea is to study the Brown measure of a non-normal free random variable x by considering the Hermitian matrix

$$X = \begin{bmatrix} 0 & x \\ x^* & 0 \end{bmatrix}$$

and the 2×2 matrix-valued Cauchy transform

$$G_X \left(\begin{bmatrix} i\varepsilon & \lambda \\ \lambda & i\varepsilon \end{bmatrix} \right) = \mathbb{E} \left(\begin{bmatrix} i\varepsilon & \lambda - x \\ (\lambda - x)^* & i\varepsilon \end{bmatrix}^{-1} \right)$$

where \mathbb{E} is the entry-wise conditional expectation $\mathbb{E} : M_2(\mathcal{M}) \rightarrow M_2(\mathbb{C})$. The entries of the matrix-valued Cauchy transform G_X carry important information for the calculation of the Brown measure.

The Hermitian reduction method becomes more powerful by combing with subordination functions and this approach was outlined in Belinschi–Śniady–Speicher’s work [5]. In particular, it is showed [5] that one can iterate certain fixed point equation for subordination functions to approximate boundary values of subordination functions, and then get approximation of Brown measure of an arbitrary polynomial of free random variables by some numerical schemes. For the sums $x_0 + c_t$ or $x_0 + g_{t,\gamma}$, we consider their Hermitian reductions and treat them as the summation of selfadjoint free random variables in the framework of operator-valued free probability. We then use the subordination functions in operator-valued free probability theory to study the Brown measure of $x_0 + c_t$ or $x_0 + g_{t,\gamma}$. It turns out there are nice formulas for subordination functions and this allows us to obtain explicit Brown measure formulas using subordination functions. Our approach is greatly influenced by the work of Aagaard–Haagerup [1], and Haagerup–Schultz [25], and Belinschi–Śniady–Speicher [5]. To our best knowledge, the Hermitian reduction methods have been only used to calculate the explicit Brown measure formula of a single operator [1, 5, 25]. The approach used in this paper demonstrates that they are also applicable to study the explicit formula of the Brown measure of the sum of two free random variables $x_0 + g_{t,\gamma}$. We expect that more applications of these methods are possible.

The paper has six sections. After the Introduction and Preliminaries, in Section 3 we study the sum of a circular operator and a free random variable. We obtain a formula for the Fuglede-Kadison determinant $\Delta(x_0 + g_{t,\gamma} - \lambda \mathbf{1})$ and $\Delta(x_0 + c_t - \lambda \mathbf{1})$ using subordination functions. In Section 4, we study the Brown measures of $x_0 + c_t$ and $x_0 + g_{t,\gamma}$. In Section 5, we calculate the Brown measure of the sum of a twisted elliptic operator and a selfadjoint operator. Finally, we calculate explicitly some non-selfadjoint examples in Section 6.

2. PRELIMINARIES

2.1. Free probability and subordination functions. We recall the definition of freeness with amalgamation over a subalgebra [40, 43]. An *operator-valued W^* -probability space* $(\mathcal{A}, \mathbb{E}, \mathcal{B})$ consists of a von Neumann algebra \mathcal{A} , a unital $*$ -subalgebra $\mathcal{B} \subset \mathcal{A}$, and a *conditional expectation* $\mathbb{E} : \mathcal{A} \rightarrow \mathcal{B}$. Thus, \mathbb{E} is a linear, unital linear positive map satisfying: (1) $\mathbb{E}(b) = b$ for all $b \in \mathcal{B}$, and (2) $\mathbb{E}(b_1 x b_2) = b_1 \mathbb{E}(x) b_2$ for all $x \in \mathcal{A}$, $b_1, b_2 \in \mathcal{B}$. Let $(\mathcal{A}_i)_{i \in I}$ be a family of subalgebras $\mathcal{B} \subset \mathcal{A}_i \subset \mathcal{A}$. We say that $(\mathcal{A}_i)_{i \in I}$ are *free with amalgamation* over \mathcal{B} with respect to the conditional expectation \mathbb{E} (or free with amalgamation in $(\mathcal{A}, \mathbb{E}, \mathcal{B})$) if

$$\mathbb{E}(x_1 x_2 \cdots x_n) = 0$$

for every $n \geq 1$, there are indices $i_1, i_2, \dots, i_n \in I$ such that $i_1 \neq i_2, i_2 \neq i_3, \dots, i_{n-1} \neq i_n$, and for $j = 1, 2, \dots, n$, we have $x_j \in \mathcal{A}_{i_j}$ such that $\mathbb{E}(x_1) = \mathbb{E}(x_2) = \dots = \mathbb{E}(x_n) = 0$.

Let $(\mathcal{A}, \mathbb{E}, \mathcal{B})$ be an operator-valued W^* -probability space. The elements in \mathcal{A} are called (noncommutative) random variables. We call

$$\mathbb{H}^+(\mathcal{B}) = \{b \in \mathcal{B} : \exists \varepsilon > 0, \Im(b) \geq \varepsilon \mathbf{1}\}$$

the Siegel upper half-plane of \mathcal{B} , where we use the notation $\Im(b) = \frac{1}{2i}(b - b^*)$. We set $\mathbb{H}^-(\mathcal{B}) = \{-b : b \in \mathbb{H}^+(\mathcal{B})\}$. The \mathcal{B} -valued Cauchy transform G_X of any selfadjoint operator $X \in \mathcal{A}$ is defined by

$$G_X(b) = \mathbb{E}[(b - X)^{-1}], \quad b \in \mathbb{H}^+(\mathcal{B}).$$

The \mathcal{B} -valued Cauchy transform G_X is a map from $\mathbb{H}^+(\mathcal{B})$ to $\mathbb{H}^-(\mathcal{B})$. The Cauchy transform is one-to-one in $\{b \in \mathbb{H}^+(\mathcal{B}) : \|b^{-1}\| < \varepsilon\}$ for ε sufficiently small, and Voiculescu's amalgamated R -transform is now defined for $X \in \mathcal{A}$ by

$$R_X(b) = G_X^{(-1)}(b) - b^{-1}$$

for b being invertible element of \mathcal{B} suitably close to zero.

Let X, Y be two selfadjoint operators that are free with amalgamation in $(\mathcal{A}, \mathbb{E}, \mathcal{B})$. The R -transform linearizes the free convolution in the sense that if X, Y are free with amalgamation in $(\mathcal{A}, \mathbb{E}, \mathcal{B})$, then

$$R_{X+Y}(b) = R_X(b) + R_Y(b)$$

for b in some suitable domain. There exist two analytic self-maps Ω_1, Ω_2 of the upper half-plane $\mathbb{H}^+(\mathcal{B})$ so that

$$(2.1) \quad (\Omega_1(b) + \Omega_2(b) - b)^{-1} = G_X(\Omega_1(b)) = G_Y(\Omega_2(b)) = G_{X+Y}(b),$$

for all $b \in \mathbb{H}^+(\mathcal{B})$. We refer the reader to [4, 11, 44] for details.

When \mathcal{M} is a von Neumann algebra, $\mathcal{B} = \mathbb{C}\mathbf{1}$ consists of scalar multiples of identity, and ϕ is a normal, faithful tracial state on \mathcal{M} . Then the pair (\mathcal{M}, ϕ) replaces the triple

$(\mathcal{M}, \mathbb{C}\mathbf{1}, \phi)$. We say (\mathcal{M}, ϕ) is a tracial W^* -probability space. For any selfadjoint element $x \in \mathcal{M}$, let $\mu = \mu_a$ be its spectral measure in (\mathcal{M}, ϕ) determined by

$$\phi(f(x)) = \int_{\mathbb{R}} f(u) d\mu_x(u)$$

for all $f \in C(\sigma(x))$. The Cauchy transform of μ (or the Cauchy transform of x) can be written as

$$G_\mu(z) = G_x(z) = \phi((z - x)^{-1}) = \int_{\mathbb{R}} \frac{1}{z - u} d\mu(u), \quad z \in \mathbb{C}^+.$$

We also set $F_\mu(z) = F_x(z) = 1/G_\mu(z)$. The reciprocal Cauchy transform F_μ maps the upper half plane \mathbb{C}^+ into itself. The \mathcal{R} -transform of μ is now an analytic function

$$(2.2) \quad \mathcal{R}_\mu(z) = G_\mu^{(-1)}(z) - \frac{1}{z}$$

where $G_\mu^{(-1)}$ denotes the inverse function to G_μ , that is defined in a truncated Stolz angle $\{z \in \mathbb{C} : \Im z > \beta, |\Re z| < \alpha(\Im z)\}$ for some $\alpha, \beta > 0$.

Suppose that the selfadjoint random variables $x, y \in \mathcal{M}$ are freely independent. Denote by μ_1 the spectral measure of x , and μ_2 the spectral measure of y , and $\mu_1 \boxplus \mu_2$ the *free additive convolution* of μ_1 and μ_2 in the sense that $\mu_1 \boxplus \mu_2 := \mu_{x+y}$. The \mathcal{R} -transform (2.2) also linearizes the free additive convolution [7] such that $\mathcal{R}_{\mu_1 \boxplus \mu_2}(z) = \mathcal{R}_{\mu_1}(z) + \mathcal{R}_{\mu_2}(z)$ in the domain where all the three \mathcal{R} -transforms are defined. In this scalar case, there exists a unique pair of analytic maps $\omega_1, \omega_2 : \mathbb{C}^+ \rightarrow \mathbb{C}^+$ such that

$$F_{\mu_1 \boxplus \mu_2}(z) = F_{\mu_1}(\omega_1(z)) = F_{\mu_2}(\omega_2(z)) = \omega_1(z) + \omega_2(z) - z$$

for all $z \in \mathbb{C}^+$. The above subordination relations can also be written in terms of Cauchy transform. That is, $G_{\mu_1 \boxplus \mu_2}(z) = G_{\mu_1}(\omega_1(z)) = G_{\mu_2}(\omega_2(z))$. The existence of subordination functions leads to many regularity results (see [2, 8] and the survey paper [6, Chapter 6]). The regularity of subordination functions is important in our approach. See Lemma 3.6 for example. For a probability measure μ on \mathbb{R} , denote $H_\mu(z) = F_\mu(z) - z$. In [3], Belinschi–Bercovici showed that ω_1, ω_2 can be obtained from the following fixed point equations

$$\omega_1(z) = z + H_{\mu_2}(z + H_{\mu_1}(\omega_1(z))), \quad \omega_2(z) = z + H_{\mu_1}(z + H_{\mu_2}(\omega_2(z))).$$

Although the subordination functions, in general, cannot be computed explicitly, they play a key role in our study by adopting this fixed point approach. See Subsection 3.1 for details.

2.2. The Brown Measure. The spectral theorem does not apply to non-normal operators. The Brown measure of an operator in \mathcal{M} was introduced by Brown [14] and is a natural replacement of the spectral distribution of a non-normal operator. Given $x \in \mathcal{M}$, the *Fuglede–Kadison determinant* $\Delta(x)$ [20] of x is defined as

$$\Delta(x) = \exp[\phi(\log(|x|))] \in [0, \infty).$$

Define a function L_x on \mathbb{C} by

$$L_x(\lambda) = \log \Delta(x - \lambda \mathbf{1}) = \phi[\log(|a - \lambda|)] \in [-\infty, \infty), \quad \lambda \in \mathbb{C}.$$

This function is subharmonic. The *Brown measure* [14] of x is then defined to be the distributional Laplacian of the subharmonic function $L_x(\lambda)$. That is,

$$(2.3) \quad \mu_x = \frac{1}{2\pi} \nabla^2 L_x(\lambda) = \frac{2}{\pi} \frac{\partial}{\partial \lambda} \frac{\partial}{\partial \bar{\lambda}} \log \Delta(x - \lambda \mathbf{1}).$$

In fact, μ_x is a probability measure supported on a subset of the spectrum of x . When $\mathcal{M} = M_n(\mathbb{C})$ and ϕ is the normalized trace on $M_n(\mathbb{C})$, for $x \in M_n(\mathbb{C})$, we have

$$L_x(\lambda) = \log |\det(x - \lambda I)|^{1/n} = \frac{1}{n} \sum_{i=1}^n \log |\lambda - \lambda_i|,$$

where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of x . Hence the eigenvalue distribution of x can be recovered by taking the distributional Laplacian

$$\frac{1}{n} (\delta_{\lambda_1} + \dots + \delta_{\lambda_n}) = \frac{1}{2\pi} \nabla^2 L_x(\lambda).$$

When $x - \lambda$ is not invertible, it is useful to consider the regularized function

$$L_{x,\varepsilon}(\lambda) = \frac{1}{2} \phi(\log((x - \lambda \mathbf{1})^*(x - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1})), \quad \varepsilon > 0.$$

Then, by the tracial property of ϕ , we have $L_x(\lambda) = \lim_{\varepsilon \rightarrow 0} L_{x,\varepsilon}(\lambda)$, and the Brown measure is calculated as

$$\mu_x = \frac{1}{4\pi} \nabla^2 \left(\lim_{\varepsilon \rightarrow 0} \phi(\log((x - \lambda \mathbf{1})^*(x - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1})) \right).$$

This regularization process makes the calculation of general operator in \mathcal{M} even more challenging. However, for the summation $x_0 + c_t$ and $x_0 + g_{t,\gamma}$, we are able to identify the domain Ξ_t , so that $L_{x_0+c_t}(\lambda)$ is real analytic for any $\lambda \in \Xi_t$, and $L_{x_0+g_{t,\gamma}}(z)$ is real analytic for any $z \in \Phi_{t,\gamma}(\Xi_t)$. Hence, the Brown measures in this paper can be calculated in classic sense. See Lemma 4.4 for details.

For convenience, for $x \in \mathcal{M}$, $\lambda \in \mathbb{C}$ and $\varepsilon \geq 0$, we denote

$$(2.4) \quad S(x, \lambda, \varepsilon) = 2L_{x,\varepsilon}(\lambda) = \phi \left(((x - \lambda \mathbf{1})^*(x - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1}) \right).$$

Then $L_x(\lambda) = \frac{1}{2} S(x, \lambda, 0)$.

2.3. Hermitian reduction method for the sum of two free random variables. Let (\mathcal{M}, ϕ) be a tracial W^* -probability space. We equip the algebra $M_2(\mathcal{M})$, the 2×2 matrices with entries from \mathcal{M} , with the conditional expectation $\mathbb{E} : M_2(\mathcal{M}) \rightarrow M_2(\mathbb{C})$ given by

$$(2.5) \quad \mathbb{E} \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} \phi(a_{11}) & \phi(a_{12}) \\ \phi(a_{21}) & \phi(a_{22}) \end{bmatrix}.$$

Then the triple $(M_2(\mathcal{M}), \mathbb{E}, M_2(\mathbb{C}))$ is a operator-valued W^* -probability space. Given $x \in \mathcal{M}$, let

$$(2.6) \quad X = \begin{bmatrix} 0 & x \\ x^* & 0 \end{bmatrix} \in M_2(\mathcal{M}),$$

which a selfadjoint element in $M_2(\mathcal{M})$. For $\varepsilon > 0$ the element

$$(2.7) \quad \Theta(\lambda, \varepsilon) = \begin{bmatrix} i\varepsilon & \lambda \\ \bar{\lambda} & i\varepsilon \end{bmatrix} \in M_2(\mathbb{C})$$

belongs to the domain of the $M_2(\mathbb{C})$ -valued Cauchy G_X . We now record that

$$(2.8) \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \begin{bmatrix} d(ad - bc)^{-1} & -b(ad - bc)^{-1} \\ -c(ad - bc)^{-1} & a(ad - bc)^{-1} \end{bmatrix}$$

where $a, d \in \mathbb{C}$ and $b, c \in \mathcal{M}$ such that $ad - bc$ is invertible (which is equivalent to $ad - cb$ is invertible). We then have

$$(2.9) \quad \begin{aligned} & (\Theta(\lambda, \varepsilon) - X)^{-1} \\ &= \begin{bmatrix} -i\varepsilon((\lambda - x)(\lambda - x)^* + \varepsilon^2)^{-1} & (\lambda - x)((\lambda - x)^*(\lambda - x) + \varepsilon^2)^{-1} \\ (\lambda - x)^*((\lambda - x)(\lambda - x)^* + \varepsilon^2)^{-1} & -i\varepsilon((\lambda - x)^*(\lambda - x) + \varepsilon^2)^{-1} \end{bmatrix}. \end{aligned}$$

and

$$(2.10) \quad G_X(\Theta(\lambda, \varepsilon)) = \mathbb{E}((\Theta(\lambda, \varepsilon) - X)^{-1}) = \begin{bmatrix} g_{X,11}(\lambda, \varepsilon) & g_{X,12}(\lambda, \varepsilon) \\ g_{X,21}(\lambda, \varepsilon) & g_{X,22}(\lambda, \varepsilon) \end{bmatrix}.$$

where

$$\begin{aligned} g_{X,11}(\lambda, \varepsilon) &= -i\varepsilon\phi\left(\left((\lambda - x)(\lambda - x)^* + \varepsilon^2\right)^{-1}\right) \\ g_{X,12}(\lambda, \varepsilon) &= \phi\left((\lambda - x)\left((\lambda - x)^*(\lambda - x) + \varepsilon^2\right)^{-1}\right) \\ g_{X,21}(\lambda, \varepsilon) &= \phi\left((\lambda - x)^*\left((\lambda - x)(\lambda - x)^* + \varepsilon^2\right)^{-1}\right) \\ g_{X,22}(\lambda, \varepsilon) &= -i\varepsilon\phi\left(\left((\lambda - x)^*(\lambda - x) + \varepsilon^2\right)^{-1}\right). \end{aligned}$$

We note that by the tracial property of ϕ , we have

$$(2.11) \quad g_{X+Y,11}(\lambda, \varepsilon) = g_{X+Y,22}(\lambda, \varepsilon), \quad g_{X,21}(\lambda, \varepsilon) = \overline{g_{X,12}(\lambda, \varepsilon)}$$

We observe that entries of the Cauchy transform (2.10) have symmetry similar to the matrix $\Theta(\lambda, \varepsilon)$. This can be explained as follows. Define the map $J : M_2(\mathbb{C}) \rightarrow M_2(\mathbb{C})$ by

$$b \mapsto J(b) = -b^*.$$

Then we have $G_X(J(b)) = J(G_X(b))$. Notice that $J(\Theta(\lambda, \varepsilon)) = \Theta(\lambda, \varepsilon)$ and hence $G_X(\Theta(\lambda, \varepsilon))$ has symmetric property (2.11).

Equations (2.10) show that two diagonal entries of $G_X(\Theta(\varepsilon, \lambda))$ carry important information to calculate the Brown measure of x . Let x and y be two $*$ -free random variables. We have to understand the $\mathcal{M}_2(\mathbb{C})$ -valued distribution of

$$\begin{bmatrix} 0 & x + y \\ (x + y)^* & 0 \end{bmatrix} = X + Y = \begin{bmatrix} 0 & x \\ x^* & 0 \end{bmatrix} + \begin{bmatrix} 0 & y \\ y^* & 0 \end{bmatrix}$$

in terms of the $\mathcal{M}_2(\mathbb{C})$ -valued distributions of X and of Y . Note that X and Y are free over $\mathcal{M}_2(\mathbb{C})$. The subordination functions in this context are two analytic self-maps Ω_1, Ω_2 of the upper half-plane $\mathbb{H}^+(\mathcal{M}_2(\mathbb{C}))$ so that

$$(2.12) \quad (\Omega_1(b) + \Omega_2(b) - b)^{-1} = G_X(\Omega_1(b)) = G_Y(\Omega_2(b)) = G_{X+Y}(b),$$

for every $b \in \mathbb{H}^+(\mathcal{M}_2(\mathbb{C}))$. We shall be concerned with $b = \Theta(\varepsilon, \lambda)$. Indeed, we have, by (2.10),

$$G_{X+Y}(\Theta(\lambda, \varepsilon)) = \begin{bmatrix} g_{X+Y,11}(\lambda, \varepsilon) & g_{X+Y,12}(\lambda, \varepsilon) \\ g_{X+Y,21}(\lambda, \varepsilon) & g_{X+Y,22}(\lambda, \varepsilon) \end{bmatrix}$$

where

$$\begin{aligned}
g_{X+Y,11}(\lambda, \varepsilon) &= -i\varepsilon\phi\left(\left((\lambda-x-y)(\lambda-x-y)^* + \varepsilon^2\right)^{-1}\right) \\
g_{X+Y,12}(\lambda, \varepsilon) &= \phi\left(\left((\lambda-x-y)((\lambda-x-y)^*(\lambda-x-y) + \varepsilon^2)^{-1}\right)\right) \\
g_{X+Y,21}(\lambda, \varepsilon) &= \phi\left(\left((\lambda-x-y)^*((\lambda-x-y)(\lambda-x-y)^* + \varepsilon^2)^{-1}\right)\right) \\
g_{X+Y,22}(\lambda, \varepsilon) &= -i\varepsilon\phi\left(\left((\lambda-x-y)^*(\lambda-x-y) + \varepsilon^2\right)^{-1}\right).
\end{aligned}$$

The idea of calculating the Brown measure of $x + y$ is to separate the information of X and Y in some tractable way. We achieve this by using subordination functions (2.12).

2.4. The elliptic operator and the operator-valued semicircular element. In free probability, the semicircle law plays a similar role to that of the Gaussian distribution in classical probability. The semicircle law σ_t with variance $t > 0$ is supported in the interval $[-2\sqrt{t}, 2\sqrt{t}]$ with density

$$d\sigma_t(x) = \frac{1}{2\pi t} \sqrt{4t - x^2} \chi_{[-2\sqrt{t}, 2\sqrt{t}]}.$$

In the tracial W^* -probability space (\mathcal{M}, ϕ) , for $t > 0$, Voiculescu's circular operator with variance t , denoted by c_t , is defined as

$$c_t = \frac{1}{\sqrt{2}}(g_t + ig'_t)$$

where $\{g_t, g'_t\}$ is a free semicircular family and each of them has variance t .

Let $t > 0$ and $\gamma \in \mathbb{C}$ such that $|\gamma| \leq t$. The *twisted elliptic operator* operator $g_{t,\gamma}$ can be constructed as follows. Let $\{g_{t_1}, g_{t_2}\}$ be semicircular operators with zero expectation and variance t_1, t_2 respectively such that $\{g_{t_1}, g_{t_2}\}$ are freely independent. For $\theta \in [0, 2\pi]$, consider the operator $y_{t_1, t_2, \theta} = e^{i\theta}(g_{t_1} + ig_{t_2})$, by choosing t_1, t_2 such that $t_1 + t_2 = t$, $t_1 - t_2 = |\gamma|$ and $e^{i2\theta} = \gamma/|\gamma|$, we can check directly that $g_{t,\gamma}$ and $y_{t_1, t_2, \theta}$ have the same $*$ -distribution, whose only nonzero free cumulants are given by

$$\kappa(y, y^*) = \kappa(y^*, y) = t, \quad \kappa(y, y) = \gamma, \quad \kappa(x^*, x^*) = \overline{\gamma},$$

where $y = g_{t,\gamma}$. The operator $g_{t,\gamma}$ include the following operators as special cases: (i) if $\gamma = 0$, y is a circular operator with variance t ; (ii) if $\gamma = t$, y is a semicircular operator g_t with variance t ; (iii) if $\gamma = -t$, then y has the distribution as ig_t ; (iv) if $\gamma \in [-t, t]$, then y has the same distribution as an elliptic operator.

In the operator-valued W^* -probability space $(\mathcal{A}, \mathbb{E}, \mathcal{B})$, following Voiculescu [43] and Speicher [40], we say $Y \in \mathcal{A}$ is \mathcal{B} -Gaussian or an operator-valued semicircular element if and only if the R -transform has a particular simple form

$$(2.13) \quad R_Y(b) = \mathbb{E}_{\mathcal{B}}(YbY).$$

Condition (2.13) says that only \mathcal{B} -cumulants of length two survive. Note that a linear combination of two operator-valued semicircular elements in $(\mathcal{A}, \mathbb{E}, \mathcal{B})$ is again an operator-valued semicircular element.

The following result is well-known. It is a special case of [36, Example 19 in Section 9.4]. One can also deduce it from a general formula about a relation between matrix-valued and scalar-valued free cumulants in [37, Theorem 6.2].

Proposition 2.1. *Let $g_{t,\gamma}$ be a twisted elliptic operator with parameters t, γ in the tracial W^* -probability space (\mathcal{M}, ϕ) and $\lambda \in \mathbb{C}$. Denote*

$$Y = \begin{bmatrix} 0 & g_{t,\gamma} \\ g_{t,\gamma}^* & 0 \end{bmatrix} \in M_2(\mathcal{M}).$$

Then Y is an operator-valued semicircular element in the operator-valued W^ -probability space $(M_2(\mathcal{M}), \mathbb{E}, M_2(\mathbb{C}))$.*

3. THE FUGLEDE-KADISON DETERMINANT AND SUBORDINATION FUNCTIONS

In a tracial W^* -probability space (\mathcal{M}, ϕ) , given $t > 0$ and $\gamma \in \mathbb{C}$ such that $|\gamma| \leq t$, let $y = g_{t,\gamma}$ be a twisted elliptic operator and let x_0 be a random variable that is $*$ -free from y . In this section, we study the Fuglede-Kadison determinant $\Delta(x_0 + y - \lambda \mathbf{1})$ for $\lambda \in \mathbb{C}$. We denote

$$(3.1) \quad X = \begin{bmatrix} 0 & x_0 \\ x_0^* & 0 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & y \\ y & 0 \end{bmatrix}.$$

Note that X and Y are free over $\mathcal{M}_2(\mathbb{C})$. There exist two analytic self-maps Ω_1, Ω_2 of upper half-plane $\mathbb{H}(\mathcal{M}_2(\mathbb{C}))$ of $\mathcal{M}_2(\mathbb{C})$ such that

$$(3.2) \quad (\Omega_1(b) + \Omega_2(b) - b)^{-1} = G_X(\Omega_1(b)) = G_Y(\Omega_2(b)) = G_{X+Y}(b),$$

for all $b \in \mathcal{M}_2(\mathbb{C})$ with $\Im b > 0$. We choose

$$\Theta(\lambda, \varepsilon) = \begin{bmatrix} i\varepsilon & \lambda \\ \bar{\lambda} & i\varepsilon \end{bmatrix}$$

where $\varepsilon > 0$ and $\lambda \in \mathbb{C}$. Our strategy is to find more explicit formulations for subordination functions Ω_1, Ω_2 .

3.1. Free convolution with a semicircular distribution. Given $\lambda \in \mathbb{C}$ such that

$$(3.3) \quad \phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}))^{-1} \right] > \frac{1}{t},$$

let w be a positive function of such λ such that

$$(3.4) \quad \phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w^2 \mathbf{1})^{-1} \right] = \frac{1}{t}.$$

In this section, we show that $w = w(0; \lambda, t)$ is a subordination function and it is a real analytic function of λ as long as (3.3) holds.

Proposition 3.1. *For a random variable $x_0 \in \mathcal{M}$, define*

$$h(\varepsilon, \lambda) = \varepsilon \phi((|\lambda - x_0|^2 + \varepsilon^2)^{-1}), \quad \varepsilon > 0.$$

Then the function

$$(\varepsilon, a, b) \mapsto h(\varepsilon, a + bi)$$

is real analytic on $(0, \infty) \times \mathbb{R} \times \mathbb{R}$.

Proof. Since

$$h(\varepsilon, \lambda) = \frac{1}{2} \int_0^\infty \left(\frac{1}{\varepsilon + iu} + \frac{1}{\varepsilon - iu} \right) d\mu_{|\lambda - x_0|}(u), \quad \varepsilon > 0,$$

the function $\varepsilon \mapsto h(\varepsilon, \lambda)$ has a complex analytic extension

$$\tilde{h} : \{z \in \mathbb{C} | \Im z > 0\} \rightarrow \mathbb{C}$$

given by the same formula. It follows that h is analytic in ε . For $\lambda = a + bi$, it is clear that $(a, b) \mapsto h(\varepsilon, \lambda)$ is real analytic on $\mathbb{R} \times \mathbb{R}$ when $s > 0$. \square

Lemma 3.2. For a probability measure μ on $[0, \infty)$, let

$$h_\mu(s) = \int_0^\infty \frac{s}{s^2 + u^2} d\mu(u)$$

and put

$$(3.5) \quad k(s, \varepsilon) = \frac{s - \varepsilon}{h_\mu(s)}, \quad s > 0, \quad \varepsilon > 0.$$

Then k is an analytic function on $(0, \infty) \times (0, \infty)$. Moreover, for $\varepsilon > 0$, the map $s \mapsto k(s, \varepsilon)$ is a strictly increasing bijection of (ε, ∞) onto $(0, \infty)$, and for $\varepsilon = 0$, the map $s \mapsto k(s, 0)$ is a strictly increasing bijection of $(0, \infty)$ onto $(\lambda_1(\mu)^2, \infty)$, where

$$\lambda_1(\mu)^2 = \left(\int_0^\infty \frac{1}{u^2} d\mu(u) \right)^{-1}.$$

Proof. It is clear that k is analytic. Moreover, for $\varepsilon > 0$,

$$(3.6) \quad k(s, \varepsilon) = \frac{s - \varepsilon}{s} \left(\int_0^\infty \frac{1}{s^2 + u^2} d\mu(u) \right)^{-1},$$

which is a product two increasing and positive functions of s on (ε, ∞) . The monotonicity properties of $s \mapsto k(s, \varepsilon)$ follows for $\varepsilon \geq 0$. \square

Definition 3.3. For $\lambda \in \mathbb{C}$, set $\mu = \mu_{|x_0 - \lambda|}$. Let h_μ, k as in Lemma 3.2. For $\varepsilon, t \in (0, \infty)$, let $w(\varepsilon; \lambda, t)$ denote the unique solution $s \in (\varepsilon, \infty)$ to the equation $k(s, \varepsilon) = t$ following Lemma 3.2 and for $t \in (\lambda_1(\mu)^2, \infty)$, let $w(0; \lambda, t)$ denote the unique solution $s \in (0, \infty)$ to the equation $k(s, 0) = t$.

Remark 3.4. The condition $k(s, \varepsilon) = t$ is equivalent to

$$(3.7) \quad \int_0^\infty \frac{s}{s^2 + u^2} d\mu_{|x_0 - \lambda|}(u) = \frac{s - \varepsilon}{t}.$$

The condition $k(s, 0) = t$ is equivalent to

$$(3.8) \quad \int_0^\infty \frac{1}{s^2 + u^2} d\mu_{|x_0 - \lambda|}(u) = \frac{1}{t}.$$

and it can be rewritten as (3.4).

For $\mu \in \text{Prob}(\mathbb{R}, \mathbb{B})$ let $\tilde{\mu}$ denote the symmetrization of μ . That is, $\tilde{\mu} \in \text{Prob}(\mathbb{R}, \mathbb{B})$ is given by

$$\tilde{\mu}(B) = \frac{1}{2}(\mu(B) + \mu(-B)), \quad (B \in \mathbb{B}).$$

Our next result shows that $w(\varepsilon; \lambda, t)$ is the subordination of a symmetric probability measure with a semicircular distribution.

Proposition 3.5. Let $\mu_1 = \tilde{\mu}_{|x_0 - \lambda|}$ and μ_2 be the semicircular distribution with variance t . Denote $\mu = \mu_1 \boxplus \mu_2$. Let ω_1, ω_2 be subordination functions such that

$$(3.9) \quad F_\mu(z) = F_{\tilde{\mu}_1}(\omega_1(z)) = F_{\mu_2}(\omega_2(z)).$$

Then for any $\varepsilon > 0$, $w(\varepsilon; \lambda, t)$ is the imaginary part of $\omega_1(i\varepsilon)$. That is, $w(\varepsilon; \lambda, t) = \Im \omega_1(i\varepsilon)$.

Proof. We have $\omega_1(z) + \omega_2(z) = z + F_\mu(z)$. Notice that, by symmetry of μ_1, μ_2 and μ , we have

$$(3.10) \quad G_\mu(i\varepsilon) = \int_{\mathbb{R}} \frac{1}{i\varepsilon - u} d\mu(u) = \int_{\mathbb{R}} \frac{-i\varepsilon}{\varepsilon^2 + u^2} d\mu(u) = -ih_\mu(\varepsilon).$$

and similarly

$$(3.11) \quad G_{\mu_1}(i\varepsilon) = -ih_{\mu_1}(\varepsilon), \quad G_{\mu_2}(i\varepsilon) = -ih_{\mu_2}(\varepsilon).$$

Let $H_1(z) = F_{\mu_1}(z) - z$ and $H_2(z) = F_{\mu_2}(z) - z$. Then ω_1 satisfy the following fixed point equation

$$\omega_1(z) = z + H_2(z + H_1(\omega_1(z))).$$

Since μ_2 is the semicircle distribution with variance t , its Cauchy transform satisfies

$$\frac{1}{G_{\mu_2}(z)} + tG_{\mu_2}(z) = z,$$

Hence $H_2(z) = -tG_{\mu_2}(z)$ and the fixed point equation reads

$$\omega_1(z) - z = -tG_{\mu_2}(z + H_1(\omega_1(z))).$$

It is clear that $\omega(i\varepsilon)$ is pure imaginary. Set $W_1(\varepsilon) = \Im\omega_1(i\varepsilon)$. In particular, the fixed point equation implies (by plugging $z = i\varepsilon$)

$$(3.12) \quad iW_1(\varepsilon) - i\varepsilon = -tG_{\mu_2}(i\varepsilon + H_1(iW_1(\varepsilon)))$$

Note that $G_{\mu_2}^{-1}(z) = \frac{1}{z} + tz$. Using the relation $G_{\mu_1}(i\varepsilon) = -ih_{\mu_1}(\varepsilon)$, then a simple algebraic computation simplifies (3.12) as

$$h_{\mu_1}(W_1(\varepsilon)) = \frac{W_1(\varepsilon) - \varepsilon}{t}.$$

Hence, by Definition 3.3, we have $w(\varepsilon; \lambda, t) = W_1(\varepsilon) = \Im\omega(i\varepsilon)$. □

We end this subsection with a regularity result of subordination functions.

Lemma 3.6. *The function $(\varepsilon, t) \mapsto w(\varepsilon; \lambda, t)$ is real analytic in $(0, \infty) \times (0, \infty)$. Moreover,*

$$\lim_{\varepsilon \rightarrow 0^+} w(\varepsilon; \lambda, t) = \begin{cases} w(0; \lambda, t), & \text{for } t \in (\lambda_1(\mu)^2, \infty); \\ 0, & \text{for } t \leq \lambda_1(\mu)^2, \end{cases}$$

where $\mu = \mu_{|x_0 - \lambda|}$ and $\lambda_1(\mu)^2 = \left(\int_0^\infty \frac{1}{u^2} d\mu(u)\right)^{-1}$. In addition, for $t > \lambda_1(\mu)^2$ and $\lambda = a + bi$, the function $(a, b) \mapsto w(0; \lambda, t)$ is real analytic.

Proof. Let $\Omega = \{(s, \varepsilon) \in \mathbb{R}^2 : 0 < \varepsilon < s\}$. Consider the analytic function

$$F(s, \varepsilon) = (k(s, \varepsilon), \varepsilon), \quad (s, \varepsilon) \in \Omega.$$

By Lemma 3.2, F is one-to-one map of Ω onto $(0, \infty) \times (0, \infty)$. Moreover, its inverse function $F^{-1} : (0, \infty) \times (0, \infty) \rightarrow \Omega$ is given by

$$F^{-1}(t, \varepsilon) = (w(\varepsilon; \lambda, t), \varepsilon), \quad t, \varepsilon > 0.$$

We now calculate the Jacobian of F as

$$J(F)(s, \varepsilon) = \frac{\partial}{\partial s} k(s, \varepsilon) > 0$$

by (3.6). Hence, F^{-1} is analytic on $(0, \infty) \times (0, \infty)$. Consequently, $w(\varepsilon; \lambda, t)$ is analytic on $(0, \infty) \times (0, \infty)$.

For $t \in (\lambda_1(\mu)^2, \infty)$ with $\mu = \mu_{|x_0 - \lambda|}$, since

$$\frac{\partial}{\partial s} k(s, 0) = \frac{\partial}{\partial s} \frac{s}{h_\mu(s)} > 0,$$

where h_μ is defined in Lemma 3.2. It follows that F is also analytic in some neighborhood U_0 of $(w(0; \lambda, t), 0)$ and F has an analytic inverse F^{-1} in a neighborhood V_0 of $F(s, 0) = (t, 0)$. Now

$$(3.13) \quad \lim_{\varepsilon \rightarrow 0^+} F^{-1}(t, \varepsilon) = F^{-1}(t, 0) = (w(0; \lambda, t), 0).$$

Hence,

$$\lim_{\varepsilon \rightarrow 0^+} w(\varepsilon; \lambda, t) = w(0; \lambda, t).$$

Moreover, the function $w(0; \lambda, t)$ is a real analytic function of (a, b) where $\lambda = a + bi$.

Note that for fixed $\varepsilon > 0$, the map $t \mapsto w(\varepsilon; \lambda, t)$ is an increasing function of t . Hence, if $t \leq \lambda_1(\mu)^2$, then for any $t' < \lambda_1(\mu)^2$, we have

$$\limsup_{\varepsilon \rightarrow 0^+} w(\varepsilon; \lambda, t) \leq \limsup_{\varepsilon \rightarrow 0^+} w(\varepsilon; \lambda, t') = w(0; \lambda, t').$$

But $t' \mapsto w(0; \lambda, t')$ is a bijection of $(\lambda_1(\mu)^2, \infty)$ onto $(0, \infty)$. It then follows that

$$\lim_{\varepsilon \rightarrow 0^+} w(\varepsilon; \lambda, t) = 0$$

whenever $t \leq \lambda_1(\mu)^2$. \square

3.2. The operator-valued subordination functions. Let $y = g_{t, \gamma}$ be a twisted elliptic operator in (\mathcal{M}, ϕ) and let $x_0 \in \mathcal{M}$ be a random variable that is $*$ -free from y . We recall that

$$G_X(\Theta(\lambda, \varepsilon)) = \begin{bmatrix} g_{X,11}(\lambda, \varepsilon) & g_{X,12}(\lambda, \varepsilon) \\ g_{X,21}(\lambda, \varepsilon) & g_{X,22}(\lambda, \varepsilon) \end{bmatrix},$$

and

$$G_{X+Y}(\Theta(\lambda, \varepsilon)) = \begin{bmatrix} g_{X+Y,11}(\lambda, \varepsilon) & g_{X+Y,12}(\lambda, \varepsilon) \\ g_{X+Y,21}(\lambda, \varepsilon) & g_{X+Y,22}(\lambda, \varepsilon) \end{bmatrix}.$$

The main result in this section is the following.

Theorem 3.7. *Let $y = g_{t, \gamma} \in \mathcal{M}$ and x_0 be a random variable that is free from y . For any $\varepsilon > 0$ and $\lambda \in \mathbb{C}$, using notations in (3.1) and (3.2), let $w(\varepsilon; \lambda, t)$ be the unique solution of w for the equation with parameters ε, t (see Definition 3.3)*

$$\frac{w - \varepsilon}{w} \left(\int_0^\infty \frac{1}{w^2 + u^2} d\mu_{|x_0 - \lambda|}(u) \right)^{-1} = t$$

and

$$(3.14) \quad z = \lambda + \gamma \cdot \phi \left((\lambda - x_0)^* ((\lambda - x_0)(\lambda - x_0)^* + w(\varepsilon; \lambda, t)^2)^{-1} \right)$$

Then $\Omega_1(\Theta(z, \varepsilon)) = \Theta(\lambda, w(\varepsilon; \lambda, t))$. That is

$$(3.15) \quad \Omega_1 \left(\begin{bmatrix} i\varepsilon & z \\ \bar{z} & i\varepsilon \end{bmatrix} \right) = \begin{bmatrix} iw(\varepsilon; \lambda, t) & \lambda \\ \bar{\lambda} & iw(\varepsilon; \lambda, t) \end{bmatrix}.$$

Moreover, the subordination equation $G_{X+Y}(\Theta(z, \varepsilon)) = G_X(\Omega_1(\Theta(z, \varepsilon)))$ is equivalent to

$$(3.16) \quad g_{X+Y,11}(z, \varepsilon) = g_{X,11}(\lambda, w(\varepsilon; \lambda, t)), \quad g_{X+Y,12}(z, \varepsilon) = g_{X,12}(\lambda, w(\varepsilon; \lambda, t)).$$

Proof. The only nonzero free cumulants of $\{y, y^*\}$ are

$$\kappa(y, y^*) = \kappa(y^*, y) = t, \quad \kappa(y, y) = \gamma, \quad \kappa(y^*, y^*) = \bar{\gamma}.$$

The operator $Y = \begin{bmatrix} 0 & y \\ y^* & 0 \end{bmatrix}$ is an operator-valued semicircular element whose R -transform is explicitly given by

$$R_Y(b) = \mathbb{E}(YbY), \quad b \in M_2(\mathbb{C}).$$

Hence, for $b = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$,

$$R_Y \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} a_{22}\kappa(y, y^*) & a_{21}\kappa(y, y) \\ a_{12}\kappa(y^*, y^*) & a_{11}\kappa(y^*, y) \end{bmatrix} = \begin{bmatrix} a_{22}t & a_{21}\gamma \\ a_{12}\bar{\gamma} & a_{11}t \end{bmatrix}.$$

In particular, for any $\varepsilon > 0$ and $z \in \mathbb{C}$, we have

$$R_Y(\Theta(z, \varepsilon)) = R_Y \begin{bmatrix} i\varepsilon & z \\ \bar{z} & i\varepsilon \end{bmatrix} = \begin{bmatrix} i\varepsilon t & \bar{z}\gamma \\ z\bar{\gamma} & i\varepsilon t \end{bmatrix}.$$

Since X, Y are free with amalgamation in the operator-valued W^* -probability space $(M_2(\mathcal{M}), \mathbb{E}, M_2(\mathbb{C}))$, we have

$$R_{X+Y}(b) = R_X(b) + R_Y(b).$$

Hence

$$G_{X+Y}^{(-1)}(b) = G_X^{(-1)}(b) + R_Y(b).$$

By replacing b with $G_{X+Y}(\beta)$, we obtain a formula for the subordination function

$$(3.17) \quad \Omega_1(\beta) = G_X^{(-1)}(G_{X+Y}(\beta)) = \beta - R_Y(G_{X+Y}(\beta))$$

for β in a neighborhood of infinity. Hence, for and $z \in \mathbb{C}$ and ε large, we have

$$(3.18) \quad \Omega_1(\Theta(z, \varepsilon)) = \Theta(z, \varepsilon) - R_Y(G_{X+Y}(\Theta(z, \varepsilon))).$$

We will show that (3.18) holds for any $z = \lambda + \gamma \cdot g_{X,12}(\lambda, w(\varepsilon))$ with $\lambda \in \mathbb{C}$ and $\varepsilon > 0$. Indeed, recall that

$$G_{X+Y}(\Theta(z, \varepsilon)) = \begin{bmatrix} g_{X+Y,11}(z, \varepsilon) & g_{X+Y,12}(z, \varepsilon) \\ g_{X+Y,21}(z, \varepsilon) & g_{X+Y,22}(z, \varepsilon) \end{bmatrix}$$

holds for any $z \in \mathbb{C}$ and $\varepsilon > 0$. Hence,

$$R_Y(G_{X+Y}(\Theta(z, \varepsilon))) = \begin{bmatrix} t \cdot g_{X+Y,22}(z, \varepsilon) & \gamma \cdot g_{X+Y,21}(z, \varepsilon) \\ \bar{\gamma} \cdot g_{X+Y,12}(z, \varepsilon) & t \cdot g_{X+Y,11}(z, \varepsilon) \end{bmatrix},$$

where we reminder the reader that

$$g_{X+Y,11}(z, \varepsilon) = g_{X+Y,22}(z, \varepsilon), \quad g_{X+Y,12}(z, \varepsilon) = \overline{g_{X+Y,21}(z, \varepsilon)}.$$

Therefore, we can rewrite (3.18) as

$$\Omega_1(\Theta(z, \varepsilon)) = \begin{bmatrix} i\varepsilon - t \cdot g_{X+Y,22}(z, \varepsilon) & z - \gamma \cdot g_{X+Y,21}(z, \varepsilon) \\ \bar{z} - \bar{\gamma} \cdot g_{X+Y,12}(z, \varepsilon) & i\varepsilon - t \cdot g_{X+Y,11}(z, \varepsilon) \end{bmatrix}.$$

Denote

$$(3.19) \quad \varepsilon_0 = \varepsilon + it \cdot g_{X+Y,22}(z, \varepsilon),$$

and

$$(3.20) \quad \lambda = z - \gamma \cdot g_{X+Y,21}(z, \varepsilon).$$

Then

$$(3.21) \quad \Omega_1(\Theta(z, \varepsilon)) = \Theta(\lambda, \varepsilon_0) = \begin{bmatrix} i\varepsilon_0 & \lambda \\ \bar{\lambda} & i\varepsilon_0 \end{bmatrix}$$

Hence, the Cauchy transform of X at $\Omega_1(\Theta(z, \varepsilon))$ is given by

$$G_X(\Omega_1(\Theta(z, \varepsilon))) = \mathbb{E}[(\Omega_1(\Theta(z, \varepsilon)) - X)^{-1}] = \begin{bmatrix} g_{X,11}(\lambda, \varepsilon_0) & g_{X,12}(\lambda, \varepsilon_0) \\ g_{X,21}(\lambda, \varepsilon_0) & g_{X,22}(\lambda, \varepsilon_0) \end{bmatrix}.$$

On the other hand, we have

$$G_{X+Y}(\Theta(z, \varepsilon)) = \begin{bmatrix} g_{X+Y,11}(z, \varepsilon) & g_{X+Y,12}(z, \varepsilon) \\ g_{X+Y,21}(z, \varepsilon) & g_{X+Y,22}(z, \varepsilon) \end{bmatrix}.$$

Therefore the subordination relation $G_{X+Y}(\Theta(z, \varepsilon)) = G_X(\Omega_1(\Theta(z, \varepsilon)))$ is equivalent to

$$(3.22) \quad g_{X+Y,11}(z, \varepsilon) = g_{X,11}(\lambda, \varepsilon_0), \quad g_{X+Y,12}(z, \varepsilon) = g_{X,12}(\lambda, \varepsilon_0)$$

where $\varepsilon, \lambda, \varepsilon_0, z$ satisfy the relation (3.19) and (3.20).

We reminder the reader that

$$g_{X,11}(\lambda, \varepsilon_0) = g_{X,22}(\lambda, \varepsilon_0), \quad g_{X,12}(\lambda, \varepsilon_0) = \overline{g_{X,21}(\lambda, \varepsilon_0)}.$$

Now the relation (3.19) can be rewritten as

$$\varepsilon_0 = \varepsilon + itg_{X,22}(\lambda, \varepsilon_0)$$

which is equivalent to

$$\varepsilon_0 = \varepsilon + t \cdot \varepsilon_0 \phi \left(((\lambda - x_0)^*(\lambda - x_0) + \varepsilon_0^2)^{-1} \right).$$

This can be further rewritten as

$$\frac{\varepsilon_0 - \varepsilon}{\varepsilon_0} \left(\int_0^\infty \frac{1}{\varepsilon_0^2 + u^2} d\mu_{|x_0 - \lambda|}(u) \right)^{-1} = t.$$

Hence, $\varepsilon_0 = w(\varepsilon; \lambda, t)$ is the unique solution to the above equation following Definition 3.3.

We argue that (3.18) or equivalently (3.21) holds for any $\varepsilon > 0$ and $\lambda \in \mathbb{C}$. For any $\lambda \in \mathbb{C}$ and $\varepsilon > 0$, we choose $\varepsilon_0 = w(\varepsilon; \lambda, t)$ and, following (3.20) we let

$$z = \lambda + \gamma \cdot g_{X,21}(\lambda, \varepsilon_0) = \lambda + \gamma \cdot \phi \left((\lambda - x_0)^* ((\lambda - x_0)(\lambda - x_0)^* + \varepsilon_0^2)^{-1} \right).$$

Then, the above calculation shows that (3.18) holds if ε is large. Notice that $R_Y(G_{X+Y}(b))$ is an analytic function defined on $\mathbb{H}^+(M_2(\mathbb{C}))$. Hence (3.18) and (3.21) hold for any $\lambda \in \mathbb{C}$, and $\varepsilon > 0$ by analytic continuation provided that they satisfy (3.19) and (3.20). Consequently, (3.22) holds for any $\lambda \in \mathbb{C}$, and $\varepsilon > 0$. This finishes the proof. \square

3.3. The coupling Fuglede-Kadison determinants. To help us remember entries of the Cauchy transform as in (2.10) are derivatives, we introduce the following notation.

Notation 3.8. For any $t > 0$, let c_t be a circular operator with variance t . For $\gamma \in \mathbb{C}$ such that $|\gamma| \leq t$, denote by $y = g_{t,\gamma}$ the twisted elliptic operator with parameters t, γ . For any $x \in \mathcal{M}$, $\lambda \in \mathbb{C}$ and $\varepsilon > 0$, we have

$$\begin{aligned} S(x, \lambda, \varepsilon) &= \log \Delta((x - \lambda \mathbf{1})^*(x - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1}) \\ &= \log \Delta((x - \lambda \mathbf{1})(x - \lambda \mathbf{1})^* + \varepsilon^2 \mathbf{1}), \quad \varepsilon > 0. \end{aligned}$$

It is convenient to introduce the following notations

$$\begin{aligned}
p_\lambda^{c,(t)}(\varepsilon) &= \frac{\partial S}{\partial \lambda}(x_0 + c_t, \lambda, \varepsilon) \\
&= -\phi \left[(x_0 + c_t - \lambda \mathbf{1})^* ((x_0 + c_t - \lambda \mathbf{1})(x_0 + c_t - \lambda \mathbf{1})^* + \varepsilon^2 \mathbf{1})^{-1} \right] \\
p_z^{g,(t,\gamma)}(\varepsilon) &= \frac{\partial S}{\partial z}(x_0 + y, z, \varepsilon) \\
&= -\phi \left[(x_0 + y - z \mathbf{1})^* ((x_0 + y - z \mathbf{1})(x_0 + y - z \mathbf{1})^* + \varepsilon^2 \mathbf{1})^{-1} \right] \\
p_\lambda^{(0)}(\varepsilon) &= \frac{\partial S}{\partial \lambda}(x_0, \lambda, \varepsilon) = -\phi \left[(x_0 - \lambda \mathbf{1})^* ((x_0 - \lambda \mathbf{1})(x_0 - \lambda \mathbf{1})^* + \varepsilon^2 \mathbf{1})^{-1} \right] \\
q_\varepsilon^{c,(t)}(\lambda) &= \frac{1}{2} \frac{\partial S}{\partial \varepsilon}(x_0 + c_t, \lambda, \varepsilon) = \varepsilon \phi \left[((x_0 + c_t - \lambda \mathbf{1})^*(x_0 + c_t - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1})^{-1} \right] \\
q_\varepsilon^{g,(t,\gamma)}(\lambda) &= \frac{1}{2} \frac{\partial S}{\partial \varepsilon}(x_0 + y, \lambda, \varepsilon) = \varepsilon \phi \left[((x_0 + y - \lambda \mathbf{1})^*(x_0 + y - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1})^{-1} \right] \\
q_\varepsilon^{(0)}(\lambda) &= \frac{1}{2} \frac{\partial S}{\partial \varepsilon}(x_0, \lambda, \varepsilon) = \varepsilon \phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1})^{-1} \right]
\end{aligned}$$

We also set

$$p_\lambda^{c,(t)}(\varepsilon) = \frac{\partial S}{\partial \lambda}(x_0 + c_t, \lambda, \varepsilon), \quad p_z^{g,(t,\gamma)}(\varepsilon) = \frac{\partial S}{\partial z}(x_0 + y, z, \varepsilon), \quad p_\lambda^{(0)}(\varepsilon) = \frac{\partial S}{\partial \lambda}(x_0, \lambda, \varepsilon).$$

We note that

$$p_\lambda^{c,(t)}(\varepsilon) = \overline{p_\lambda^{c,(t)}(\varepsilon)}, \quad p_z^{g,(t,\gamma)}(\varepsilon) = \overline{p_z^{g,(t,\gamma)}(\varepsilon)}, \quad p_\lambda^{(0)}(\varepsilon) = \overline{p_\lambda^{(0)}(\varepsilon)}.$$

and the Cauchy transform can be written as

$$(3.23) \quad G_{X+Y}(\Theta(\lambda, \varepsilon)) = \mathbb{E} [(\Theta(\lambda, \varepsilon) - X - Y)^{-1}] = \begin{bmatrix} -iq_\varepsilon^{(t,\gamma)}(\lambda) & p_\lambda^{(t,\gamma)}(\varepsilon) \\ p_\lambda^{(t,\gamma)}(\varepsilon) & -iq_\varepsilon^{(t,\gamma)}(\lambda) \end{bmatrix}.$$

Corollary 3.9. *The subordination relation (3.16) is equivalent to*

$$(3.24) \quad q_{w(\varepsilon)}^{(0)}(\lambda) = q_\varepsilon^{g,(t,\gamma)}(z) = q_\varepsilon^{c,(t)}(\lambda), \quad p_\lambda^{(0)}(w(\varepsilon)) = p_z^{g,(t,\gamma)}(\varepsilon) = p_\lambda^{(0)}(w(\varepsilon)).$$

where $w(\varepsilon) = w(\varepsilon; \lambda, t)$.

Proof. We note that $c_t = g_{t,0}$. Then (3.14) reads $z = \lambda$ if $\gamma = 0$. The result follows from (3.16) and (3.23). \square

The following proof was inspired by the proof of [25, Lemma 4.14].

Lemma 3.10. *Let $y = g_{t,\gamma}$ and x_0 be a random variable free from y . For any $\lambda \in \mathbb{C}$ and $(\varepsilon, t) \in (0, \infty) \times (0, \infty)$, we have the coupling Fugulede-Kadison determinant formula*

$$(3.25) \quad \begin{aligned} \Delta((x_0 + y - z \mathbf{1})^*(x_0 + y - z \mathbf{1}) + \varepsilon^2 \mathbf{1}) &= \Delta((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(\varepsilon)^2 \mathbf{1}) \\ &\times \exp \left[\Re \left(\gamma \cdot (p_\lambda^{(0)}(w(\varepsilon)))^2 \right) - \frac{(w(\varepsilon) - \varepsilon)^2}{t} \right] \end{aligned}$$

where $z = \lambda + \gamma \cdot p_\lambda^{(0)}(w(\varepsilon))$ and $w(\varepsilon) = w(\varepsilon; \lambda, t)$ is defined in Definition 3.3.

Proof. Fix $\lambda \in \mathbb{C}$, then $w(\varepsilon) = w(\varepsilon; \lambda, t)$ and $p_\lambda^{(0)}(w(\varepsilon))$ are then completely determined by ε . We denote

$$(3.26) \quad H_{x_0+y-z}(\varepsilon) = \frac{1}{2} \log \Delta((x_0 + y - z \mathbf{1})^*(x_0 + y - z \mathbf{1}) + \varepsilon^2 \mathbf{1})$$

and

$$H_{x_0-\lambda}(\varepsilon) = \frac{1}{2} \log \Delta((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1}).$$

We have

$$H_{x_0-\lambda}(\varepsilon) = \frac{1}{2} \int_0^\infty \log(u^2 + \varepsilon^2) d\mu_{|x_0-\lambda \mathbf{1}|}(u).$$

Hence

$$(3.27) \quad \lim_{\varepsilon \rightarrow \infty} (H_{x_0-\lambda}(\varepsilon) - \log \varepsilon) = \frac{1}{2} \lim_{\varepsilon \rightarrow \infty} \log \left(1 + \frac{u^2}{\varepsilon^2} \right) d\mu_{|x_0-\lambda \mathbf{1}|}(u) = 0.$$

Observe that $\lim_{\varepsilon \rightarrow \infty} (w(\varepsilon) - \varepsilon) = 0$. Consequently, $\lim_{\varepsilon \rightarrow \infty} p_\lambda^{(0)}(w(\varepsilon)) = 0$. Hence

$$\lim_{\varepsilon \rightarrow \infty} z = \lim_{\varepsilon \rightarrow \infty} (\lambda + t p_\lambda^{(0)}(w(\varepsilon))) = \lambda$$

and by a similar estimation as (3.27)

$$\lim_{\varepsilon \rightarrow \infty} (H_{x_0+y-z}(\varepsilon) - \log \varepsilon) = 0$$

Recall that $z = \lambda + \gamma \cdot p_\lambda^{(0)}(w(\varepsilon))$ and $p_\lambda^{(0)}(w(\varepsilon)) = p_z^{(t,\gamma)}(\varepsilon)$, we have

$$z = \lambda + p_z^{(t,\gamma)}(\varepsilon).$$

Therefore,

$$\begin{aligned} & \frac{d}{d\varepsilon} H_{x_0+y-z}(\varepsilon) \\ &= \frac{1}{2} \frac{d}{d\varepsilon} S(x_0 + y, z, \varepsilon) \\ &= q_\varepsilon^{(t,\gamma)}(z) + \frac{1}{2} \left(\gamma \cdot p_z^{(t,\gamma)}(\varepsilon) \frac{d}{d\varepsilon} p_z^{(t,\gamma)}(\varepsilon) + \bar{\gamma} \cdot p_{\bar{z}}^{(t,\gamma)}(\varepsilon) \frac{d}{d\varepsilon} p_{\bar{z}}^{(t,\gamma)}(\varepsilon) \right) \\ &= q_{w(\varepsilon)}^{(0)}(\lambda) + \frac{1}{2} \left(\gamma \cdot p_\lambda^{(0)}(w(\varepsilon)) \frac{d}{d\varepsilon} p_\lambda^{(0)}(w(\varepsilon)) + \bar{\gamma} \cdot p_{\bar{\lambda}}^{(0)}(w(\varepsilon)) \frac{d}{d\varepsilon} p_{\bar{\lambda}}^{(0)}(w(\varepsilon)) \right) \end{aligned}$$

and

$$\frac{d}{d\varepsilon} H_{x_0-\lambda}(\varepsilon) = q_\varepsilon^{(0)}(\lambda).$$

Recall that (see Definition 3.3 and Remark 3.4)

$$\frac{w(\varepsilon) - \varepsilon}{t} = q_{w(\varepsilon)}^{(0)}(\lambda)$$

We then have

$$\begin{aligned} & \int_{\varepsilon_0}^\varepsilon q_{w(u)}^{(0)}(\lambda) du \\ &= \int_{\varepsilon_0}^\varepsilon q_{w(u)}^{(0)}(\lambda) \frac{d}{du} w(u) du + \int_{\varepsilon_0}^\varepsilon q_{w(u)}^{(0)}(\lambda) \frac{d}{du} (u - w(u)) du \\ &= (H_{x_0-\lambda}(w(\varepsilon)) - H_{x_0-\lambda}(w(\varepsilon_0))) - t \cdot \int_{\varepsilon_0}^\varepsilon q_{w(u)}^{(0)}(\lambda) \left(\frac{d}{du} q_{w(u)}^{(0)}(\lambda) \right) \\ &= (H_{x_0-\lambda}(w(\varepsilon)) - H_{x_0-\lambda}(w(\varepsilon_0))) - \frac{t}{2} \left((q_{w(\varepsilon)}^{(0)}(\lambda))^2 - (q_{w(\varepsilon_0)}^{(0)}(\lambda))^2 \right) \end{aligned}$$

and

$$\begin{aligned} & \gamma \int_{\varepsilon_0}^{\varepsilon} p_{\lambda}^{(0)}(w(u)) \left(\frac{d}{du} p_{\lambda}^{(0)}(w(u)) \right) du + \bar{\gamma} \int_{\varepsilon_0}^{\varepsilon} p_{\bar{\lambda}}^{(0)}(w(u)) \left(\frac{d}{du} p_{\bar{\lambda}}^{(0)}(w(u)) \right) \\ &= \frac{\gamma}{2} \left((p_{\lambda}^{(0)}(w(\varepsilon)))^2 - (p_{\lambda}^{(0)}(w(\varepsilon_0)))^2 \right) + \frac{\bar{\gamma}}{2} \left((p_{\bar{\lambda}}^{(0)}(w(\varepsilon)))^2 - (p_{\bar{\lambda}}^{(0)}(w(\varepsilon_0)))^2 \right) \end{aligned}$$

Hence, there exists a constant C such that

$$\begin{aligned} H_{x_0+y-z}(\varepsilon) &= H_{x_0-\lambda}(w(\varepsilon)) \\ &+ \frac{1}{4} \left(\gamma \cdot (p_{\lambda}^{(0)}(w(\varepsilon)))^2 + \bar{\gamma} \cdot (p_{\bar{\lambda}}^{(0)}(w(\varepsilon)))^2 - 2t \cdot (q_{w(\varepsilon)}^{(0)}(\lambda))^2 \right) + C. \end{aligned}$$

Observe that $\lim_{\varepsilon \rightarrow \infty} (w(\varepsilon) - \varepsilon) = 0$. Consequently, $\lim_{\varepsilon \rightarrow \infty} p_{\lambda}^{(0)}(w(\varepsilon)) = 0$. Hence

$$\lim_{\varepsilon \rightarrow \infty} z = \lim_{\varepsilon \rightarrow \infty} (\lambda + t p_{\lambda}^{(0)}(w(\varepsilon))) = \lambda$$

and by a similar estimation as (3.27)

$$\lim_{\varepsilon \rightarrow \infty} (H_{x_0+y-z}(\varepsilon) - \log \varepsilon) = 0$$

Moreover,

$$\lim_{\varepsilon \rightarrow \infty} q_{w(\varepsilon)}^{(0)}(\lambda) = 0.$$

We conclude that C must be zero. Therefore,

$$\begin{aligned} \exp(2H_{x_0+y-z}(\varepsilon)) &= \exp(2H_{x_0-\lambda}(w(\varepsilon))) \\ &\times \exp \left[\frac{1}{2} \left(\gamma \cdot (p_{\lambda}^{(0)}(w(\varepsilon)))^2 + \bar{\gamma} \cdot (p_{\bar{\lambda}}^{(0)}(w(\varepsilon)))^2 - 2t \cdot (q_{w(\varepsilon)}^{(0)}(\lambda))^2 \right) \right] \end{aligned}$$

Finally, replace $q_{w(\varepsilon)}^{(0)}(\lambda)$ by $\frac{w(\varepsilon)-\varepsilon}{t}$. The result follows by recalling (3.26). \square

Theorem 3.11. *Let $y = g_{t,\gamma}$ and x_0 be a random variable free from x . For $\lambda \in \mathbb{C}$, set $\mu = \mu_{|x_0-\lambda|}$ and let $w(0; \lambda, t)$ be as in Definition 3.3. If $\phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}))^{-1} \right] > \frac{1}{t}$, then $w(0; \lambda, t) \in (0, \infty)$ and*

$$\begin{aligned} & \Delta((x_0+y-z\mathbf{1})^*(x_0+y-z\mathbf{1})) = \Delta((x_0-\lambda\mathbf{1})^*(x_0-\lambda\mathbf{1}) + w(0; \lambda, t)^2\mathbf{1}) \\ (3.28) \quad & \times \exp \left[\frac{1}{2} \cdot \left(\gamma \cdot (p_{\lambda}^{(0)}(w(0; \lambda, t)))^2 + \bar{\gamma} \cdot (p_{\bar{\lambda}}^{(0)}(w(0; \lambda, t)))^2 - \frac{2w(0; \lambda, t)^2}{t} \right) \right], \end{aligned}$$

where $z = \lambda + \gamma \cdot p_{\lambda}^{(0)}(w(0; \lambda, t))$.

Proof. By Lemma 3.6, we know that $w(0; \lambda, t) > 0$. The Fuglede-Kadison determinant is continuous in norm topology in the set of invertible element. The result follows from Lemma 3.10 by letting ε tend to zero. \square

Theorem 3.12. *For $\lambda \in \mathbb{C}$, set $\mu = \mu_{|x_0-\lambda|}$ and let $w(0; \lambda, t)$ be as in Definition 3.3.*

(1) *If $\phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}))^{-1} \right] > \frac{1}{t}$, then*

$$\begin{aligned} & \Delta(x_0 + c_t - \lambda \mathbf{1})^2 = \Delta((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2\mathbf{1}) \\ (3.29) \quad & \times \exp \left(- \frac{(w(0; \lambda, t))^2}{t} \right), \end{aligned}$$

where $w(0; \lambda, t)$ is defined in Definition 3.3, which reads

$$(3.30) \quad \phi \left[\left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1} \right)^{-1} \right] = \frac{1}{t}.$$

(2) If $\phi \left[\left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) \right)^{-1} \right] \leq \frac{1}{t}$, then

$$\Delta(x_0 + c_t - \lambda \mathbf{1}) = \Delta(x_0 - \lambda \mathbf{1}).$$

Proof. The circular operator c_t corresponds to $\gamma = 0$ for a twisted elliptic operator. Hence the first part follows from Theorem 3.11. For the second part, note that

$$\phi \left[\left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) \right)^{-1} \right] \leq \frac{1}{t} \iff \lambda_1(\mu)^2 \geq t$$

and

$$\Delta(x_0 + c_t - \lambda \mathbf{1})^2 = \lim_{\varepsilon \rightarrow 0^+} \Delta \left((x_0 + c_t - \lambda \mathbf{1})^* (x_0 + c_t - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1} \right).$$

By Lemma 3.6, we know if $\phi \left[\left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) \right)^{-1} \right] \leq \frac{1}{t}$, then $\lim_{\varepsilon \rightarrow 0^+} w(\varepsilon; \lambda, t) = 0$. Hence by (3.25), we have

$$\begin{aligned} \Delta(x_0 + c_t - \lambda \mathbf{1})^2 &= \lim_{\varepsilon \rightarrow 0^+} \Delta \left((x_0 + c_t - \lambda \mathbf{1})^* (x_0 + c_t - \lambda \mathbf{1}) + \varepsilon^2 \mathbf{1} \right) \\ &= \lim_{\varepsilon \rightarrow 0^+} \Delta \left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(\varepsilon)^2 \mathbf{1} \right) \times \exp \left[-\frac{(w(\varepsilon) - \varepsilon)^2}{t} \right] \\ &= \Delta(x_0 - \lambda \mathbf{1})^2 \end{aligned}$$

provided that $\phi \left[\left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) \right)^{-1} \right] \leq \frac{1}{t}$. Moreover, the condition (3.30) follows from (3.8) for $\mu = \mu_{|x_0 - \lambda|}$. \square

Corollary 3.13. If $\phi \left[\left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) \right)^{-1} \right] > \frac{1}{t}$, then

$$\Delta(x_0 + g_{t,\gamma} - z \mathbf{1})^2 = \Delta(x_0 + c_t - \lambda \mathbf{1})^2 \exp \left[\Re(\gamma(p_\lambda^{(0)}(w(0; \lambda, t)))^2) \right]$$

where $z = \lambda + \gamma \cdot p_\lambda^{(0)}(w(0; \lambda, t))$.

4. THE BROWN MEASURE OF THE ADDITION WITH AN ELLIPTIC OPERATOR

We study the Brown measure of $x_0 + c_t$ and $x_0 + g_{t,\gamma}$. We show that the Brown measure of $x_0 + c_t$ can be expressed as a formula using subordination function $w(\varepsilon; \lambda, t)$ under some mild assumptions. We then show that there is a natural map $\Phi_{t,\gamma}$ defined as

$$(4.1) \quad \Phi_{t,\gamma}(\lambda) = \lambda + \gamma \cdot p_\lambda^{(0)}(w(0; \lambda, t)), \quad \lambda \in \mathbb{C}$$

where

$$p_\lambda^{(0)}(w(0; \lambda, t)) = -\phi \left[(x_0 - \lambda \mathbf{1})^* \left((x_0 - \lambda \mathbf{1})(x_0 - \lambda \mathbf{1})^* + w(0; \lambda, t)^2 \mathbf{1} \right)^{-1} \right]$$

such that the gradient of $\log \Delta(x_0 + c_t - \lambda \mathbf{1})$ at λ is connected with the gradient of $\log \Delta(x_0 + g_{t,\gamma} - z)$ at $z = \Phi_{t,\gamma}(\lambda)$. If $\Phi_{t,\gamma}$ is one-to-one and non-singular within the support of the Brown measure of $x_0 + c_t$, then the Brown measure of $x_0 + g_{t,\gamma}$ is the push-forward of the Brown measure of $x_0 + c_t$ under the map $\Phi_{t,\gamma}$.

4.1. The gradient functions.

Notation 4.1. We denote

$$(4.2) \quad \Xi_t = \left\{ \lambda : \phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}))^{-1} \right] > \frac{1}{t} \right\}.$$

Following Definition 3.3 and Lemma 3.6, for given $\lambda \in \mathbb{C}$ and $t \in (0, \infty)$, if $\lambda \in \Xi_t$, we let $w(0; \lambda, t) > 0$ be such that

$$\phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1} \right] = \frac{1}{t};$$

and if $\phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}))^{-1} \right] \leq \frac{1}{t}$, set $w(0; \lambda, t) = 0$. Moreover, set

$$(4.3) \quad H(\lambda) = \Re \left(\gamma (p_\lambda^{(0)}(w(0; \lambda, t)))^2 \right)$$

for $\lambda \in \Xi_t$.

Proposition 4.2. The set Ξ_t is open for any $t > 0$.

Proof. For any $\varepsilon \geq 0$, denote the function f_ε of λ by

$$f_\varepsilon(\lambda) = \phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + \varepsilon \mathbf{1})^{-1} \right].$$

For any $\varepsilon > 0$, the function $f_\varepsilon(\lambda)$ is a continuous function of λ . Observe that f_0 is the limit of the increasing sequence of f_ε , hence it is lower semi-continuous. The set Ξ_t can be rewritten as $\Xi_t = \{\lambda : f_0(\lambda) > 1/t\}$ and therefore Ξ_t is open for any $t > 0$. \square

Lemma 4.3. For any $t > 0$. The function $\lambda \mapsto S(x_0 + c_t, \lambda, 0)$ is a real analytic function for $\lambda \in \Xi_t$, and we have

$$p_\lambda^{c_t(t)}(0) = p_\lambda^{(0)}(w(0; \lambda, t)),$$

Proof. By Corollary 3.9, we have

$$p_\lambda^{c_t(t)}(\varepsilon) = p_\lambda^{(0)}(w(\varepsilon; \lambda, t)),$$

Recall that, by Lemma 3.6, for $\lambda \in \Xi_t$, $\lim_{\varepsilon \rightarrow 0} w(\varepsilon; \lambda, t) = w(0; \lambda, t) \in (0, \infty)$. The result then follows by letting ε tend to zero. \square

To move forward, we need the following assumption on the map $\Phi_{t,\gamma}$. We will show that the condition is satisfied by a large family of operators.

Condition A. For any $t > 0$, $\gamma \in \mathbb{C}$ such that $|\lambda| \leq t$, and $\lambda \in \Xi_t$. Let c_t be a circular operator with variance t and let $g_{t,\gamma}$ be an elliptic operator as in Section 2.4. Let x_0 be a random variable that is free from $\{c_t, g_{t,\gamma}\}$. The map $\lambda \mapsto \Phi_{t,\gamma}(\lambda)$ is one-to-one and nonsingular at all $\lambda \in \Xi_t$, where

$$\Phi_{t,\gamma}(\lambda) = \lambda + \gamma \cdot p_\lambda^{(0)}(w(0; \lambda, t)).$$

and $w(0; \lambda, t)$ is determined by

$$\phi((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1} = \frac{1}{t}.$$

For $Q \in \mathcal{M}$, $\lambda \in \mathbb{C}$, set $Q_\lambda = Q - \lambda \mathbf{1}$. We remind the reader that

$$L_{Q,\varepsilon}(\lambda) = \frac{1}{2} S(Q, \lambda, \varepsilon) = \frac{1}{2} \phi(\log((Q - \lambda \mathbf{1})(Q - \lambda \mathbf{1})^* + \varepsilon^2 \mathbf{1}))$$

and $L_Q(\lambda) = L_{Q,0}(\lambda)$. For $\varepsilon \geq 0$, if $(\lambda_1, \lambda_2) \mapsto L_{Q,\varepsilon}(\lambda)$ is differentiable at $\lambda = \lambda_1 + i\lambda_2$, we record that the gradient of $L_{Q,\varepsilon}$ is given by

$$(4.4) \quad \frac{\partial}{\partial \lambda_1} L_{Q,\varepsilon}(\lambda) = -\Re(\phi(Q_\lambda^*(Q_\lambda Q_\lambda^* + \varepsilon^2 \mathbf{1})^{-1})),$$

$$(4.5) \quad \frac{\partial}{\partial \lambda_2} L_{Q,\varepsilon}(\lambda) = \Im(\phi(Q_\lambda^*(Q_\lambda Q_\lambda^* + \varepsilon^2 \mathbf{1})^{-1})),$$

where $\lambda_1 = \Re(\lambda)$, $\lambda_2 = \Im(\lambda)$. That is,

$$\frac{\partial}{\partial \lambda} L_{Q,\varepsilon}(\lambda) = \frac{1}{2} \left[\frac{\partial}{\partial \lambda_1} L_{Q,\varepsilon}(\lambda) - i \frac{\partial}{\partial \lambda_2} L_{Q,\varepsilon}(\lambda) \right].$$

We then denote the derivative with respect to real variables λ_1, λ_2 as

$$\begin{aligned} p_{\lambda_1} &= 2\Re(p_\lambda^{(0)}(w(0; \lambda, t))) = 2 \left(\frac{\partial}{\partial \lambda_1} L_{x_0,\varepsilon}(\lambda) \right) \Big|_{\varepsilon=w(0;\lambda,t)} \\ &= \left(\frac{\partial}{\partial \lambda_1} S(x_0, \lambda, \varepsilon) \right) \Big|_{\varepsilon=w(0;\lambda,t)} \end{aligned}$$

and

$$\begin{aligned} p_{\lambda_2} &= -2\Im(p_\lambda^{(0)}(w(0; \lambda, t))) = -2 \left(\frac{\partial}{\partial \lambda_2} L_{x_0,\varepsilon}(\lambda) \right) \Big|_{\varepsilon=w(0;\lambda,t)} \\ &= - \left(\frac{\partial}{\partial \lambda_2} S(x_0, \lambda, \varepsilon) \right) \Big|_{\varepsilon=w(0;\lambda,t)}. \end{aligned}$$

That is

$$p_\lambda^{(0)}(w(0; \lambda, t)) = \frac{1}{2}(p_{\lambda_1} - ip_{\lambda_2}).$$

For $\gamma = \gamma_1 + i\gamma_2$, then $\Phi_{t,\gamma}$ can be written as

$$\begin{aligned} (4.6) \quad \Phi_{t,\gamma}(\lambda) &= \lambda + \gamma p_\lambda^{(0)}(w(0; \lambda, t)) \\ &= (\lambda_1 + i\lambda_2) + \frac{1}{2}(\gamma_1 + i\gamma_2)(p_{\lambda_1} - ip_{\lambda_2}) \\ &= \left(\lambda_1 + \frac{1}{2}(\gamma_1 p_{\lambda_1} + \gamma_2 p_{\lambda_2}) \right) + i \left(\lambda_2 + \frac{1}{2}(\gamma_2 p_{\lambda_1} - \gamma_1 p_{\lambda_2}) \right). \end{aligned}$$

Hence, the Jacobian of the map $\Phi_{t,\gamma}$ in coordinates (λ_1, λ_2) is given by

$$(4.7) \quad \text{Jacobian}(\Phi_{t,\gamma})(\lambda) = \begin{bmatrix} 1 + \frac{1}{2} \left(\gamma_1 \frac{\partial p_{\lambda_1}}{\partial \lambda_1} + \gamma_2 \frac{\partial p_{\lambda_2}}{\partial \lambda_1} \right) & \frac{1}{2} \left(\gamma_1 \frac{\partial p_{\lambda_1}}{\partial \lambda_2} + \gamma_2 \frac{\partial p_{\lambda_2}}{\partial \lambda_2} \right) \\ \frac{1}{2} \left(\gamma_2 \frac{\partial p_{\lambda_1}}{\partial \lambda_1} - \gamma_1 \frac{\partial p_{\lambda_2}}{\partial \lambda_1} \right) & 1 + \frac{1}{2} \left(\gamma_2 \frac{\partial p_{\lambda_1}}{\partial \lambda_2} - \gamma_1 \frac{\partial p_{\lambda_2}}{\partial \lambda_2} \right) \end{bmatrix}.$$

Lemma 4.4. *Given $t > 0$, $\gamma \in \mathbb{C}$ such that $|\lambda| \leq t$, and $\lambda \in \Xi_t$, assume that $\text{Jacobian}(\Phi_{t,\gamma})(\lambda) \neq 0$, then the function $z \mapsto S(x_0 + g_{t,\gamma}, z, 0)$ is a real analytic function of z in a neighborhood of $z \in \Phi_{t,\gamma}(\Xi_t)$. Moreover, we have*

$$(4.8) \quad p_z^{g,(t,\gamma)}(0) = p_\lambda^{c,(t)}(0), \quad p_{\bar{z}}^{g,(t,\gamma)}(0) = p_{\bar{\lambda}}^{c,(t)}(0),$$

where $z = \Phi_{t,\gamma}(\lambda)$.

In particular, under Condition A, the functions $z \mapsto S(x_0 + g_{t,\gamma}, z, 0)$ is a real analytic function of z for $z \in \Phi_{t,\gamma}(\Xi_t)$, and the identities (4.8) hold for any $\lambda \in \Xi_t$.

Proof. From Corollary 3.13 (by comparing Theorem 3.12 with Theorem 3.11), we see that, for any $\lambda \in \Xi_t$,

(4.9)

$$\Delta((x_0 + y - z)^*(x_0 + y - z)) = \Delta((x_0 + c_t - \lambda \mathbf{1})^*(x_0 + c_t - \lambda \mathbf{1})) \exp(H(\lambda)),$$

where $z = \Phi_{t,\gamma}(\lambda)$ and $H(\lambda)$ was defined as (4.3). That is,

$$S(x_0 + y, z, 0) = S(x_0 + c_t, \lambda, 0) + H(\lambda).$$

Note that both sides of the above equation are real differentiable functions in Ξ_t . Put $p_\lambda = p_\lambda^{(0)} w(0; \lambda, t)$. Take the derivative $\frac{\partial}{\partial \lambda}$, we obtain

$$\begin{aligned} \frac{\partial S(x_0 + y, z, 0)}{\partial \lambda} &= p_z^{g,(t,\gamma)}(0) \left(\frac{\partial \Phi_{t,\gamma}(\lambda)}{\partial \lambda} \right) + p_{\bar{z}}^{g,(t,\gamma)}(0) \left(\frac{\partial \overline{\Phi_{t,\gamma}(\lambda)}}{\partial \lambda} \right) \\ &= p_z^{g,(t,\gamma)}(0) \left(1 + \gamma \frac{\partial p_\lambda}{\partial \lambda} \right) + p_{\bar{z}}^{g,(t,\gamma)}(0) \left(\bar{\gamma} \frac{\partial p_{\bar{\lambda}}}{\partial \lambda} \right), \end{aligned}$$

and $\frac{\partial S(x_0 + c_t, \lambda, 0)}{\partial \lambda} = p_\lambda^{c,(t)}(0)$, and

$$\frac{\partial H(\lambda)}{\partial \lambda} = \gamma \cdot p_\lambda \cdot \frac{\partial p_\lambda}{\partial \lambda} + \bar{\gamma} \cdot p_{\bar{\lambda}} \cdot \frac{\partial p_{\bar{\lambda}}}{\lambda}.$$

Hence,

$$p_z^{g,(t,\gamma)}(0) \left(1 + \gamma \frac{\partial p_\lambda}{\partial \lambda} \right) + p_{\bar{z}}^{g,(t,\gamma)}(0) \left(\bar{\gamma} \frac{\partial p_{\bar{\lambda}}}{\partial \lambda} \right) = p_\lambda^{c,(t)}(0) + \gamma \cdot p_\lambda \cdot \frac{\partial p_\lambda}{\partial \lambda} + \bar{\gamma} \cdot p_{\bar{\lambda}} \cdot \frac{\partial p_{\bar{\lambda}}}{\lambda}.$$

By using $p_\lambda = p_\lambda^{c,(t)}(0)$ in Lemma 4.3, we can then rewrite the above identity as

$$(4.10) \quad \left(p_z^{g,(t,\gamma)}(0) - p_\lambda^{c,(t)}(0) \right) \left(1 + \gamma \frac{\partial p_\lambda}{\partial \lambda} + \bar{\gamma} \frac{\partial p_{\bar{\lambda}}}{\partial \lambda} \right) = 0.$$

We shall show that (4.10) implies that

$$p_z^{g,(t,\gamma)}(0) - p_\lambda^{c,(t)}(0) = 0$$

Indeed, put $p_z^{g,(t,\gamma)}(0) - p_\lambda^{c,(t)}(0) = c + id$, by using (4.6), a simple algebraic computation yields that the identity (4.10) is equivalent to

$$\begin{bmatrix} c & d \end{bmatrix} \cdot M = 0$$

where M is the matrix

$$(4.11) \quad M = \begin{bmatrix} 1 + \frac{1}{2} \left(\gamma_1 \frac{\partial p_{\lambda_1}}{\partial \lambda_1} + \gamma_2 \frac{\partial p_{\lambda_2}}{\partial \lambda_1} \right) & \frac{1}{2} \left(\gamma_1 \frac{\partial p_{\lambda_1}}{\partial \lambda_2} + \gamma_2 \frac{\partial p_{\lambda_2}}{\partial \lambda_2} \right) \\ \frac{1}{2} \left(\gamma_1 \frac{\partial p_{\lambda_2}}{\partial \lambda_1} - \gamma_2 \frac{\partial p_{\lambda_1}}{\partial \lambda_1} \right) & -1 + \frac{1}{2} \left(\gamma_1 \frac{\partial p_{\lambda_2}}{\partial \lambda_2} - \gamma_2 \frac{\partial p_{\lambda_1}}{\partial \lambda_2} \right) \end{bmatrix}.$$

If $\text{Jacobian}(\Phi_{t,\gamma})(\lambda) \neq 0$, then the map $\Phi_{t,\gamma}$ is non-singular in neighborhood of λ . Hence,

$$\det(M) = (-1) \text{Jacobian}(\Phi_{t,\gamma})(\lambda) \neq 0.$$

We then conclude that $c = d = 0$. That is,

$$p_z^{g,(t,\gamma)}(0) - p_\lambda^{c,(t)}(0) = 0.$$

Therefore,

$$p_{\bar{z}}^{g,(t,\gamma)}(0) - p_{\bar{\lambda}}^{c,(t)}(0) = 0$$

as well. Under Condition A, the map $\Phi_{t,\gamma}$ is non-singular at any $\lambda \in \Xi_t$. The the above argument works for any $\lambda \in \Xi_t$. This concludes the statement. \square

Remark 4.5. It is interesting to compare Lemma 4.4 with Lemma 4.3. Choose $z = \lambda + \gamma p_\lambda^{(0)}(w(\varepsilon; \lambda, t))$. If the map $\Phi_{t,\gamma}$ associated with x_0 is singular, it turns out $\lim_{\varepsilon \rightarrow 0^+} p_z^{g,(t,\gamma)}(\varepsilon)$ has a limit that depends on λ . Hence, the condition in Lemma 4.4 is necessary. See Example 5.16.

4.2. The addition with a circular operator. The following result generalizes [32, Theorem 3.10] where it assumed that x_0 is selfadjoint.

Theorem 4.6. *The Brown measure is absolutely continuous with respect to Lebesgue measure in the open set Ξ_t . The density of the Brown measure at $\lambda \in \Xi_t$ is given by*

$$(4.12) \quad \frac{1}{\pi} \left(\frac{1}{t} - \frac{\partial}{\partial \lambda} \left(\phi(x_0^*((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1}) \right) \right)$$

where $w = w(0; \lambda, t)$ is determined by

$$\phi((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1} = \frac{1}{t}.$$

It can also be expressed as

$$(4.13) \quad \frac{1}{\pi} \left(\frac{|\phi((\lambda \mathbf{1} - x_0)(h^{-1})^2)|^2}{\phi((h^{-1})^2)} + \omega(0; \lambda, t)^2 \phi(h^{-1}k^{-1}) \right)$$

where $h = h(\lambda, w(0; \lambda, t))$ and $k = k(\lambda, w(0; \lambda, t))$ for

$$h(\lambda, w) = (\lambda \mathbf{1} - x_0)^*(\lambda \mathbf{1} - x_0) + w^2$$

and

$$k(\lambda, w) = (\lambda \mathbf{1} - x_0)(\lambda \mathbf{1} - x_0)^* + w^2.$$

In particular, the density of the Brown measure of $x_0 + c_t$ is strictly positive in the set Ξ_t .

Proof. For $\lambda \in \Xi_t$, we have $\phi \left[((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}))^{-1} \right] > \frac{1}{t}$. We have

$$\begin{aligned} & \Delta((x_0 + c_t - \lambda \mathbf{1})^*(x_0 + c_t - \lambda \mathbf{1})) \\ &= \Delta((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1}) \cdot \exp \left[-t(q_{w(0; \lambda, t)}^{(0)}(\lambda))^2 \right] \end{aligned}$$

by Theorem 3.12. In addition, $w(0; \lambda, t) \in (0, \infty)$ and $(a, b) \mapsto w(0; \lambda, t)$ for $\lambda = a + bi$ is real analytic by Lemma 3.6. Hence, $\lambda \mapsto \log \Delta(x_0 + c_t - \lambda \mathbf{1})$ is real analytic. We put

$$g(\lambda) = \log \Delta((x_0 + c_t - \lambda \mathbf{1})^*(x_0 + c_t - \lambda \mathbf{1}))$$

The Brown measure can be calculated as

$$d\mu_{x_0+c_t}(\lambda) = \frac{1}{\pi} \frac{\partial^2}{\partial \bar{\lambda} \partial \lambda} g(\lambda).$$

where the Laplacian can be calculated in the usual sense.

Then Lemma 4.3 is equivalent to

$$\frac{\partial}{\partial \lambda} g(\lambda) = \phi \left(h^{-1} \frac{\partial h}{\partial \lambda} \right)$$

where $\frac{\partial h}{\partial \lambda} = (\lambda \mathbf{1} - x_0)^*$. We can continue to take the derivative directly

$$\begin{aligned} \frac{\partial^2}{\partial \bar{\lambda} \partial \lambda} &= \frac{\partial}{\partial \bar{\lambda}} \phi \left(h^{-1} \frac{\partial h}{\partial \lambda} \right) \\ &= \frac{\partial}{\partial \bar{\lambda}} \phi \left((\lambda \mathbf{1} - x_0)^* ((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1} \right) \\ &= \frac{\partial}{\partial \bar{\lambda}} \phi \left(\frac{\bar{\lambda}}{t} - \phi(x_0^* ((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1}) \right) \\ &= \frac{1}{t} - \frac{\partial}{\partial \bar{\lambda}} \left(\phi(x_0^* ((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1}) \right). \end{aligned}$$

Then the first formula (4.12) is established.

We adapt the calculation in [25, Lemma 2.8] to get another form of the density formula. By [27, Lemma 3.2], since $g(\lambda)$ is a real analytic function of λ , we have

$$\begin{aligned} \frac{\partial^2}{\partial \bar{\lambda} \partial \lambda} g(\lambda) &= \frac{\partial}{\partial \bar{\lambda}} \phi \left(h^{-1} \frac{\partial h}{\partial \lambda} \right) \\ (4.14) \quad &= \phi \left(-h^{-1} \left(\frac{\partial h}{\partial \bar{\lambda}} + 2w \frac{\partial w}{\partial \bar{\lambda}} \right) h^{-1} \frac{\partial h}{\partial \lambda} + h^{-1} \frac{\partial^2 h}{\partial \bar{\lambda} \partial \lambda} \right) \\ &= \phi \left(-h^{-1} \frac{\partial h}{\partial \bar{\lambda}} h^{-1} \frac{\partial h}{\partial \lambda} + h^{-1} \frac{\partial^2 h}{\partial \bar{\lambda} \partial \lambda} \right) - 2w \frac{\partial w}{\partial \bar{\lambda}} \phi \left(\frac{\partial h}{\partial \lambda} (h^{-1})^2 \right) \end{aligned}$$

and

$$\frac{\partial h}{\partial \bar{\lambda}} = \frac{\partial h(\lambda, w)}{\partial \bar{\lambda}} = \lambda \mathbf{1} - x_0;$$

and

$$\frac{\partial^2 h}{\partial \bar{\lambda} \partial \lambda} = \frac{\partial}{\partial \bar{\lambda}} (\lambda \mathbf{1} - x_0)^* = 1.$$

We now apply the identity $x(x^*x + \varepsilon \mathbf{1})^{-1} = (xx^* + \varepsilon \mathbf{1})^{-1}x$ to $x = \lambda \mathbf{1} - x_0$ and $\varepsilon = w^2$, we find that

$$\begin{aligned} -\frac{\partial h}{\partial \bar{\lambda}} h^{-1} \frac{\partial h}{\partial \lambda} + \frac{\partial^2 h}{\partial \bar{\lambda} \partial \lambda} &= 1 - x(x^*x + w^2 \mathbf{1})^{-1}x^* \\ &= 1 - (xx^* + w^2 \mathbf{1})^{-1}xx^* \\ &= w^2 (xx^* + w^2 \mathbf{1})^{-1} \\ &= w^2 k^{-1}. \end{aligned}$$

Now, $w(0; \lambda, t)$ is determined by (see Remark 3.4)

$$\phi((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1})^{-1} = \frac{1}{t}$$

which can be rewritten as

$$\phi(h^{-1}) = \frac{1}{t}.$$

Take implicit differentiation $\frac{\partial}{\partial \bar{\lambda}}$ and apply again [27, Lemma 3.2], we then obtain

$$\phi \left(h^{-1} \left(\frac{\partial h}{\partial \bar{\lambda}} + 2w \frac{\partial w}{\partial \bar{\lambda}} \right) h^{-1} \right) = 0,$$

where $\frac{\partial h}{\partial \lambda} = \lambda \mathbf{1} - x_0$. This implies

$$-2w \frac{\partial w}{\partial \lambda} = \frac{\phi \left(\frac{\partial h}{\partial \lambda} (h^{-1})^2 \right)}{\phi((h^{-1})^2)}.$$

By the tracial property, we observe that

$$\phi \left(\frac{\partial h}{\partial \lambda} (h^{-1})^2 \right) = \overline{\phi \left(\frac{\partial h}{\partial \lambda} (h^{-1})^2 \right)}.$$

We therefore can continue to simplify (4.14) as

$$\begin{aligned} \frac{\partial^2}{\partial \lambda \partial \lambda} g(\lambda) &= \phi \left(-h^{-1} \frac{\partial h}{\partial \lambda} h^{-1} \frac{\partial h}{\partial \lambda} + h^{-1} \frac{\partial^2 h}{\partial \lambda \partial \lambda} \right) - 2w \frac{\partial w}{\partial \lambda} \phi \left(\frac{\partial h}{\partial \lambda} (h^{-1})^2 \right) \\ &= w^2 \phi(h^{-1} k^{-1}) + \frac{\left| \phi \left(\frac{\partial h}{\partial \lambda} (h^{-1})^2 \right) \right|^2}{\phi((h^{-1})^2)} \\ &= w^2 \phi(h^{-1/2} k^{-1} h^{-1/2}) + \frac{\left| \phi((\lambda \mathbf{1} - x_0)(h^{-1})^2) \right|^2}{\phi((h^{-1})^2)} > 0 \end{aligned}$$

for any $\lambda \in \Xi_t$. This finishes the proof. \square

4.3. A general result on the support of the Brown measure. Theorem 4.6 does not tell us the Brown measure of $x_0 + c_t$ outside the open set Ξ_t . We will show that the Brown measure of $x_0 + c_t$ is supported in the closure of Ξ_t .

Lemma 4.7. *The Brown measures of x_0 and $x_0 + c_t$ coincide in the complement of the closure $\overline{\Xi_t}$. That is*

$$\mu_{x_0}|_{(\overline{\Xi_t})^c} = \mu_{x_0+c_t}|_{(\overline{\Xi_t})^c}.$$

In particular, $\mu_{x_0+c_t}(\overline{\Xi_t}) = 1$ if and only if $\mu_{x_0}(\overline{\Xi_t}) = 1$.

Proof. For $\lambda \in (\Xi_t)^c$, by Theorem 3.12, we have

$$\log \Delta(x_0 - \lambda \mathbf{1}) = \log \Delta(x_0 + c_t - \lambda \mathbf{1}).$$

Hence,

$$\int_{\mathbb{C}} \log |z - \lambda| d\mu_{x_0}(z) = \int_{\mathbb{C}} \log |z - \lambda| d\mu_{x_0+c_t}(z)$$

for any $\lambda \in (\Xi_t)^c$. Then two Brown measures coincide in the open set $(\overline{\Xi_t})^c$ due to the Unicity Theorem of logarithmic potential (see [38, Theorem 2.1 in Chapter II]). \square

We are grateful to Hari Bercovici for providing us a proof of the following result which is a refinement of an argument in [24, Proposition 4.6].

Lemma 4.8. *Let μ be a finite Borel measure on \mathbb{C} . Define $I : \mathbb{C} \rightarrow (0, \infty]$ by*

$$I(\lambda) = \int_{\mathbb{C}} \frac{1}{|z - \lambda|^2} d\mu(z).$$

Then $I(\lambda)$ is infinite almost everywhere relative to μ .

Proof. If it is not true, by restricting to a bounded subset of \mathbb{C} where $I(\lambda)$ is bounded if necessary, we may assume that $I(\lambda)$ is finite almost everywhere relative to μ and $\text{supp}(\mu)$ is contained in the ball $D(0, r)$ centered at the origin with radius r . We may further assume that there exists $C > 0$ such that

$$I(\lambda) < C, \quad \forall \lambda \in \mathbb{C}.$$

Fix any $\lambda \in \text{supp}(\mu)$, for any $\delta > 0$ there exists $\varepsilon = \varepsilon(\lambda) \in (0, 1)$ such that

$$\int_{|z-\lambda|<\varepsilon} \frac{1}{|z-\lambda|^2} d\mu(z) < \delta$$

by continuity of the integration. Hence

$$\mu(D(\lambda, \varepsilon)) < \delta\varepsilon^2$$

where $D(\lambda, \varepsilon)$ is the disc centered at λ with radius ε . The Vitali covering lemma tells us there is a countable family of discs $(D(\lambda_i, \varepsilon_i))_{i=1}^{\infty}$ that are pairwise disjoint, such that

$$\text{supp}(\mu) \subset \bigcup_{i=1}^{\infty} (5 \cdot D(\lambda_i, \varepsilon_i)).$$

By choosing larger r if needed, we may choose ε_i so that $5 \cdot D(\lambda_i, \varepsilon_i) \subset D(0, r)$. Hence,

$$\mu(\mathbb{C}) \leq 5\delta \sum_{i=1}^{\infty} \varepsilon_i^2 \leq 5\delta r^2.$$

Note that r can be chosen to be independent of δ since μ is compactly supported. This implies that μ is zero measure and we conclude that $I(\lambda)$ is infinite almost everywhere relative to μ . \square

The following result provides a new characterization of the support of the Brown measure of an arbitrary operator. The result also improves [30, Theorem 1.2] where a condition of local boundedness of $(T - \lambda\mathbf{1})^{-1}$ in L^4 -norm is required.

Theorem 4.9. *Let $\lambda \in \mathbb{C}$ and $T \in \mathcal{M}$. If $\tau(|T - \lambda|^{-2}) < \infty$, then*

$$\int_{\mathbb{C}} \frac{1}{|z-\lambda|^2} d\mu_T(z) \leq \tau(|T - \lambda|^{-2}).$$

If $\tau(|T - \lambda|^{-2}) < L$ in some neighborhood of λ_0 for some $L > 0$, then $\lambda_0 \notin \text{supp}(\mu_T)$.

Proof. Observe that for any $t \in (0, 1)$, we have $2 \log t > -\frac{1}{t^2}$. Hence,

$$\begin{aligned} 2 \int_0^1 \log t d\mu_{|T-\lambda|}(t) &> - \int_0^1 \frac{1}{t^2} d\mu_{|T-\lambda|}(t) \\ &> - \int_0^{\infty} \frac{1}{t^2} d\mu_{|T-\lambda|}(t) \\ &= -\tau(|T - \lambda|^{-2}) > -\infty. \end{aligned}$$

By [25, Proposition 2.16], the operator $T - \lambda$ has an inverse $(T - \lambda)^{-1}$ that is possibly unbounded operator affiliated with \mathcal{M} and the Brown measure $\mu_{(T-\lambda)^{-1}}$ of $(T - \lambda)^{-1}$ can be defined. Moreover, $\mu_{(T-\lambda)^{-1}}$ is the push-forward measure of μ_T via the map $z \mapsto (z - \lambda)^{-1}$. Hence, by [25, Theorem 2.19], we obtain

$$(4.15) \quad \int_{\mathbb{C}} \frac{1}{|z-\lambda|^2} d\mu_T(z) = \int_{\mathbb{C}} |z|^2 d\mu_{(T-\lambda)^{-1}}(z) \leq \|(T - \lambda)^{-1}\|_2^2 = \tau(|T - \lambda|^{-2}).$$

If $\tau(|T - \lambda|^{-2}) < L$ in some neighborhood of λ_0 for some $L > 0$. It then follows by Lemma 4.8 that $\mu_T(U) = 0$ thanks to (4.15). Hence, $\lambda_0 \notin \text{supp}(\mu_T)$. \square

Theorem 4.10. *For any x_0 $*$ -free from c_t , the Brown measure $\mu_{x_0+c_t}$ of $x_0 + c_t$ is supported in the closure $\overline{\Xi}_t$.*

Proof. For any $\lambda_0 \in (\Xi_t)^c$, there is some neighborhood U of λ_0 such that $\tau(|x_0 - \lambda|^{-2}) \leq 1/t$ for $\lambda \in U$. By Theorem 4.9, $\mu_{x_0}(U) = 0$. Hence, $\mu_{x_0}((\Xi_t)^c) = 0$. By Lemma 4.7, the Brown measure of $x_0 + c_t$ coincides with the Brown measure of x_0 within the open set $(\Xi_t)^c$. Hence, $\mu_{x_0+c_t}((\Xi_t)^c) = 0$ and thus the Brown measure is supported in the closure $\overline{\Xi_t}$, thanks to Theorem 4.6. \square

Remark 4.11. It is known [32] that all selfadjoint operators x_0 satisfy the condition that $\mu_{x_0+c_t}(\partial(\Xi_t)) = 0$ (see Proposition 5.12 and Section 5.3 for a review). We expect that this condition is satisfied provided that the boundary set $\partial(\Xi_t)$ is smooth enough.

4.4. Addition with an elliptic operator. We now turn to study the Brown measure of $x_0 + g_{t,\gamma}$. We first establish a useful formula for the Brown measure. Let μ_Q be the Brown measure of an operator Q in the tracial W^* -probability space (\mathcal{M}, ϕ) . Denote $Q_\lambda = Q - \lambda$, and recall that

$$L_Q(\lambda) = \log \Delta(Q - \lambda \mathbf{1}), \quad \text{and} \quad L_{Q,\varepsilon}(\lambda) = \log \Delta((Q_\lambda^* Q_\lambda + \varepsilon^2)^{1/2})$$

for $\lambda \in \mathbb{C}$ and $\varepsilon > 0$. The following result can be obtained using the same argument as in [1, Lemma 4.2]. For reader's convenience, we include a proof.

Lemma 4.12. *Let Γ be a simply connected domain in the support of μ_Q with piecewise smooth boundary. If $\mu_Q(\partial\Gamma) = 0$, then,*

$$\mu_Q(\Gamma) = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi} \int_{\partial\Gamma} -\frac{\partial L_{Q,\varepsilon}}{\partial \lambda_2} d\lambda_1 + \frac{\partial L_{Q,\varepsilon}}{\partial \lambda_1} d\lambda_2 = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\pi} \int_{\partial\Gamma} \frac{\partial L_{Q,\varepsilon}}{\partial \vec{\mathbf{n}}} ds$$

where $\lambda_1 = \Re \lambda$, $\lambda_2 = \Im \lambda$; and $\vec{\mathbf{n}}$ is the outer normal vector field of $\partial\Gamma$.

If L_Q is real analytic in a neighborhood of Γ , then

$$\mu_Q(\Gamma) = \frac{1}{2\pi} \int_{\partial\Gamma} -\frac{\partial L_Q}{\partial \lambda_2} d\lambda_1 + \frac{\partial L_Q}{\partial \lambda_1} d\lambda_2 = \frac{1}{2\pi} \int_{\partial\Gamma} \frac{\partial L_Q}{\partial \vec{\mathbf{n}}} ds.$$

Proof. For $\varepsilon > 0$, it is known that $\nabla^2 L_{Q,\varepsilon} \geq 0$ and for each $\varepsilon > 0$, the measure

$$\mu_{Q,\varepsilon} = \frac{1}{2\pi} \nabla^2 L_{Q,\varepsilon}(\lambda) d\lambda_1 d\lambda_2$$

is a probability measure on \mathbb{C} . Moreover, $\lim_{\varepsilon \rightarrow 0^+} \mu_{Q,\varepsilon} = \mu_Q$ in the weak* topology on probability measures on \mathbb{C} . Hence for all $\psi \in C_0(\mathbb{C})$, we have

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{C}} \psi d\mu_{Q,\varepsilon} = \int_{\mathbb{C}} \psi d\mu_Q.$$

Let $(\psi_n)_{n=1}^\infty$ be an increasing sequence of $C_0(\mathbb{C})$ -functions with $0 \leq \psi_n \leq 1$ for all $n \in \mathbb{N}$ that converges to the characteristic function 1_Γ . It follows that

$$\begin{aligned} \mu_Q(\Gamma) &= \lim_{n \rightarrow \infty} \int_{\mathbb{C}} \psi_n d\mu_Q = \lim_{n \rightarrow \infty} \left(\lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{C}} \psi_n d\mu_{Q,\varepsilon} \right) \\ &\leq \lim_{n \rightarrow \infty} \left(\liminf_{\varepsilon \rightarrow 0^+} \int_{\mathbb{C}} 1_\Gamma d\mu_{Q,\varepsilon} \right) = \liminf_{\varepsilon \rightarrow 0^+} \mu_{Q,\varepsilon}(\Gamma). \end{aligned}$$

Similarly, by choosing a decreasing sequence $(\eta_n)_{n=1}^\infty$ of $C_0(\mathbb{C})$ -functions with $0 \leq \eta_n \leq 1$ that converges to $1_{\overline{\Gamma}}$, we deduce that

$$\mu_Q(\overline{\Gamma}) \geq \limsup_{\varepsilon \rightarrow 0^+} \mu_{Q,\varepsilon}(\overline{\Gamma}).$$

Consequently, if $\mu_Q(\partial\Gamma) = 0$, then

$$\limsup_{\varepsilon \rightarrow 0^+} \mu_{Q,\varepsilon}(\overline{\Gamma}) \leq \mu_Q(\Gamma) \leq \liminf_{\varepsilon \rightarrow 0^+} \mu_{Q,\varepsilon}(\Gamma),$$

and therefore $\mu_Q(\Gamma) = \lim_{\varepsilon \rightarrow 0^+} \mu_{Q,\varepsilon}(\Gamma)$.

The probability measure can be rewritten as, by applying Green's theorem,

$$\begin{aligned} \mu_{Q,\varepsilon}(\Gamma) &= \frac{1}{2\pi} \int_{\mathbb{R}} \left(\frac{\partial^2 L_{Q,\varepsilon}(\lambda)}{\partial \lambda_1^2} + \frac{\partial^2 L_{Q,\varepsilon}(\lambda)}{\partial \lambda_2^2} \right) d\lambda_1 d\lambda_2 \\ &= \frac{1}{2\pi} \int_{\partial\Gamma} -\frac{\partial L_{Q,\varepsilon}(\lambda)}{\partial \lambda_2} d\lambda_1 + \frac{\partial L_{Q,\varepsilon}(\lambda)}{\partial \lambda_1} d\lambda_2 \\ &= \frac{1}{2\pi} \int_{\partial\Gamma} \frac{\partial L_{Q,\varepsilon}}{\partial \vec{\mathbf{n}}} ds. \end{aligned}$$

The above application of Green's theorem is valid by replacing $\mu_{Q,\varepsilon}, L_{Q,\varepsilon}$ with μ_Q and L_Q provided that L_Q is real analytic in a neighborhood of Γ . This finishes the proof. \square

Proposition 4.13. *Let $\Gamma \subset \Xi_t$ be a simply connected domain in the set Ξ_t with piecewise smooth boundary such that $\bar{\Gamma} \subset \Xi_t$. Then,*

$$\mu_{x_0+c_t}(\Gamma) = \frac{1}{2\pi} \int_{\partial\Gamma} -\frac{\partial L_{x_0+c_t}}{\partial \lambda_2} d\lambda_1 + \frac{\partial L_{x_0+c_t}}{\partial \lambda_1} d\lambda_2,$$

where $L_{x_0+c_t}(\lambda) = \log \Delta(x_0 + c_t - \lambda \mathbf{1})$.

Proof. Note that for any $\lambda \in \Xi_t$, we have $w(0; \lambda, t) \in (0, \infty)$ by Theorem 3.6 and $L_{x_0+c_t}$ is differentiable with derivative

$$\lim_{\varepsilon \rightarrow 0^+} \frac{\partial}{\partial \lambda_1} L_{x_0+c_t,\varepsilon}(\lambda) = \frac{\partial}{\partial \lambda_1} L_{x_0+c_t}(\lambda) = \Re(p_\lambda^{c,(t)}(0))$$

due to Theorem 3.12 (see also Lemma 4.4). Similarly,

$$\lim_{\varepsilon \rightarrow 0^+} \frac{\partial}{\partial \lambda_2} L_{x_0+c_t,\varepsilon}(\lambda) = \frac{\partial}{\partial \lambda_2} L_{x_0+c_t}(\lambda) = \Im(p_\lambda^{c,(t)}(0))$$

Now, the result follows by taking the limit as ε tends to zero for the formula in Lemma 4.12. \square

Proposition 4.14. *Under Condition A, let $\Gamma \subset \Phi_{t,\gamma}(\Xi_t)$ be a simply connected domain in the set $\Phi_{t,\gamma}(\Xi_t)$ with piecewise smooth boundary such that $\bar{\Gamma} \subset \Phi_{t,\gamma}(\Xi_t)$. Then,*

$$\mu_{x_0+g_{t,\gamma}}(\Gamma) = \frac{1}{2\pi} \int_{\partial\Gamma} -\frac{\partial L_{x_0+g_{t,\gamma}}}{\partial z_2} dz_1 + \frac{\partial L_{x_0+g_{t,\gamma}}}{\partial z_1} dz_2,$$

where $L_{x_0+g_{t,\gamma}}(z) = \log \Delta(x_0 + g_{t,\gamma} - z)$.

Proof. The proof is identical to the proof for Proposition 4.13, thanks to Lemma 4.4. \square

Proposition 4.15. *Under Condition A, let $\Phi_{t,\gamma}^{-1}(z) = \lambda = \lambda_1 + i\lambda_2$ for $z \in \Phi_{t,\gamma}(\Xi_t)$. Then the Brown measure of $x_0 + g_{t,\gamma}$ is absolutely continuous at $\Phi_{t,\gamma}(\Xi_t)$, and if $\gamma \neq 0$, the density is given by*

(4.16)

$$d\mu_{x_0+g_{t,\gamma}} = \frac{1}{2\pi} \left(\frac{\partial^2 L_{x_0+g_{t,\gamma}}}{\partial z_2 \partial z_1} \right) dz_1 dz_2 = \frac{1}{2\pi\gamma} \left[\left(\frac{\partial \lambda_2}{\partial z_2} - \frac{\partial \lambda_1}{\partial z_1} \right) + i \left(\frac{\partial \lambda_1}{\partial z_2} + \frac{\partial \lambda_2}{\partial z_1} \right) \right].$$

where $z_1 = \Im(z), z_2 = \Re(z)$.

Proof. We recall that $z = \Phi_{t,\gamma}(\lambda) = \lambda + \gamma \cdot p_\lambda^{(0)}(w(0; \lambda, t))$, and by Lemma 4.4,

$$p_\lambda^{(0)}(w(0; \lambda, t)) = p_z^{g,(t,\gamma)}(0) = \frac{\partial S(x_0 + g_{t,\gamma}, z, 0)}{\partial z}.$$

Hence

$$\frac{\partial S(x_0 + g_{t,\gamma}, z, 0)}{\partial z} = p_z^{g,(t,\gamma)}(0) = \frac{z - \lambda}{\gamma} = \frac{1}{\gamma}(z - \Phi_{t,\gamma}^{-1}(z)).$$

Therefore,

$$\begin{aligned} \frac{\partial^2 S(x_0 + g_{t,\gamma}, z, 0)}{\partial \bar{z} \partial z} &= \frac{1}{2\gamma} \left(\frac{\partial}{\partial z_1} + i \frac{\partial}{\partial z_2} \right) (z - \lambda_1 - i\lambda_2) \\ &= \frac{1}{2\gamma} \left(\frac{\partial \lambda_2}{\partial z_2} - \frac{\partial \lambda_1}{\partial z_1} \right) + \frac{i}{2\gamma} \left(\frac{\partial \lambda_1}{\partial z_2} + \frac{\partial \lambda_2}{\partial z_1} \right). \end{aligned}$$

The result then follows by recalling that $L_{x_0+g_{t,\gamma}}(\lambda) = \frac{1}{2}S(x_0 + g_{t,\gamma}, \lambda, 0)$ and converting the complex derivative. \square

Theorem 4.16. *Under Condition A, in addition if $\mu_{x_0+c_t}(\partial(\Xi_t)) = 0$, then the Brown measure of $x_0 + g_{t,\gamma}$ is the push-forward measure of the Brown measure of $\mu_{x_0+c_t}$ under the map $\lambda \mapsto \Phi_{t,\gamma}(\lambda)$. Let $\Gamma \subset \Xi_t$ be a simply connected domain in the set Ξ_t with piecewise smooth boundary such that $\bar{\Gamma} \subset \Xi_t$. Then*

$$(4.17) \quad \mu_{x_0+c_t}(\Gamma) = \mu_{x_0+g_{t,\gamma}}(\Phi_{t,\gamma}(\Gamma)).$$

Proof. Under Condition A, the domain $\Phi_{t,\gamma}(\Gamma)$ is a simply connected domain with piecewise smooth boundary. For $\lambda = \lambda_1 + i\lambda_2 \in \Xi_t$ and $z = z_1 + iz_2 \in \Phi_{t,\gamma}(\Xi_t)$, we denote the vector fields

$$P^c(\lambda_1, \lambda_2) = \Re(p_\lambda^{c,(t)}(0)), \quad Q^c(\lambda_1, \lambda_2) = -\Im(p_\lambda^{c,(t)}(0));$$

and

$$P^g(z_1, z_2) = \Re(p_z^{g,(t,\gamma)}(0)), \quad Q^g(z_1, z_2) = -\Im(p_z^{g,(t,\gamma)}(0)).$$

Recall that, for $\lambda \in \Xi_t$,

$$z = \Phi_{t,\gamma}(\lambda) = \lambda + \gamma \cdot p_\lambda^{(0)}(w(0; \lambda, t)),$$

and, by Lemma 4.4, we have

$$p_\lambda^{(0)}(w(0; \lambda, t)) = p_\lambda^{c,(t)}(0) = \frac{\partial L_{x_0+c_t}(\lambda)}{\partial \lambda} = p_z^{g,(t,\gamma)}(0) = \frac{\partial L_{x_0+g_{t,\gamma}}(z)}{\partial z},$$

where $z = \Phi_{t,\gamma}(\lambda)$. We note that

$$p_\lambda^{c,(t)}(0) = \frac{1}{2} \left(\frac{\partial L_{x_0+c_t}(\lambda)}{\partial \lambda_1} - i \frac{\partial L_{x_0+c_t}(\lambda)}{\partial \lambda_2} \right) = P^c(\lambda_1, \lambda_2) - iQ^c(\lambda_1, \lambda_2).$$

Hence, we have

$$P^c(\lambda_1, \lambda_2) = P^g(z_1, z_2) = \frac{\partial L_{x_0+c_t}(\lambda)}{\partial \lambda_1},$$

and

$$Q^c(\lambda_1, \lambda_2) = Q^g(z_1, z_2) = \frac{\partial L_{x_0+c_t}(\lambda)}{\partial \lambda_2}.$$

Let $\gamma = \gamma_1 + i\gamma_2$, then we have

$$(4.18) \quad \begin{aligned} z_1 &= \lambda_1 + \gamma_1 P^c(\lambda_1, \lambda_2) + \gamma_2 Q^c(\lambda_1, \lambda_2); \\ z_2 &= \lambda_2 + \gamma_2 P^c(\lambda_1, \lambda_2) - \gamma_1 Q^c(\lambda_1, \lambda_2). \end{aligned}$$

Denote the differential form α in $\Phi_{t,\gamma}(\Xi_t)$ as

$$(4.19) \quad \alpha = -\frac{\partial L_{x_0+g_{t,\gamma}}}{\partial z_2} dz_1 + \frac{\partial L_{x_0+g_{t,\gamma}}}{\partial z_1} dz_2 = -Q^g dz_1 + P^g dz_2.$$

Under Condition A, we can change variables from z to λ , which really means that we pull back the 1-form α by the map $\Phi_{t,\gamma}$. Letting β be the pulled-back form, and also using formulas (4.18) for z_1, z_2 , we get

$$\begin{aligned} \beta &= -Q^c d(\lambda_1 + \gamma_1 P^c + \gamma_2 Q^c) \\ &\quad + P^c d(\lambda_2 + \gamma_2 P^c - \gamma_1 Q^c). \end{aligned}$$

We can write this as

$$(4.20) \quad \begin{aligned} \beta &= -Q^c d\lambda_1 + P^c d\lambda_2 - \gamma_1 (P^c dQ^c + Q^c dP^c) + \gamma_2 (P^c dP^c - Q^c dQ^c) \\ &= -Q^c d\lambda_1 + P^c d\lambda_2 + d \left[-\gamma_1 P^c Q^c + \frac{1}{2} \gamma_2 ((P^c)^2 - (Q^c)^2) \right]. \end{aligned}$$

Hence, by Proposition 4.14 and the definition of 1-form α, β , we have

$$\begin{aligned} \mu_{x_0+g_{t,\gamma}}(\Phi_{t,\gamma}(\Gamma)) &= \frac{1}{2\pi} \int_{\partial\Phi_{t,\gamma}(\Gamma)} -\frac{\partial L_{x_0+g_{t,\gamma}}}{\partial z_2} dz_1 + \frac{\partial L_{x_0+g_{t,\gamma}}}{\partial z_1} dz_2 \\ &= \frac{1}{2\pi} \int_{\partial\Phi_{t,\gamma}(\Gamma)} \alpha = \frac{1}{2\pi} \int_{\partial\Gamma} \beta \\ &= \frac{1}{2\pi} \int_{\partial\Gamma} -Q^c(\lambda_1, \lambda_2) d\lambda_1 + P^c(\lambda_1, \lambda_2) d\lambda_2 \\ &= \frac{1}{2\pi} \int_{\partial\Gamma} -\frac{\partial L_{x_0+c_t}}{\partial \lambda_2} d\lambda_1 + \frac{\partial L_{x_0+c_t}}{\partial \lambda_1} d\lambda_2 \\ &= \mu_{x_0+c_t}(\Gamma), \end{aligned}$$

where we used (4.20) to deduce the fourth identity. \square

5. ADDITION WITH A SELFADJOINT OPERATOR

The special case when x_0 is selfadjoint has drawn much attention in previous work [28, 31, 32]. In this section, we apply our main result to generalize main results in those work. The generalization can be viewed as the addition analogue of recent work [29] concerning free multiplicative Brownian motions.

5.1. Subordination functions. Let x_0 be selfadjoint and $\mu = \mu_{x_0}$ be its spectral measure. We first study the subordination function as in Definition 3.3 and Proposition 3.5. The set Ξ_t in Notation 4.1 is expressed as

$$(5.1) \quad \Xi_t = \left\{ \lambda = a + bi : \int_{\mathbb{R}} \frac{1}{(u-a)^2 + b^2} d\mu(u) > \frac{1}{t} \right\}.$$

For $\lambda \in \Xi_t$ (which is equivalent to $t > \lambda_1(\mu)^2$), the condition subordination function $w(0; \lambda, t)$ is determined by the condition (3.8), that can be rewritten as

$$(5.2) \quad \int_{\mathbb{R}} \frac{1}{(u-a)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u) = \frac{1}{t}.$$

Following Biane work [10, Section 3] on the spectral measure of $x_0 + g_t$, we set

$$U_t = \left\{ a \in \mathbb{R} : \int_{\mathbb{R}} \frac{1}{(u-a)^2} d\mu(u) > \frac{1}{t} \right\},$$

and define v_t as follows

$$(5.3) \quad v_t(a) = \inf \left\{ y > 0 : \int_{\mathbb{R}} \frac{1}{(a-x)^2 + y^2} d\mu(x) \leq \frac{1}{t} \right\}, \quad a \in \mathbb{R}.$$

We then set

$$\Omega_t = \{a + bi : |b| > v_t(a)\}.$$

It follows that $v_t(a)^2 = b^2 + w(0; \lambda, t)^2$ with $\lambda = a + bi$ if $\lambda \in \Xi_t$; and $v_t(a) = 0$ if $\{a + ib : b \in \mathbb{R}\} \cap \Xi_t = \emptyset$. By regularity of $w(0; \lambda, t)$ in Lemma 3.6, we see that v_t is a continuous function (see [10] for the original definition and [32, Section 2.3] for a review).

Proposition 5.1. *The subordination function $w(0; \lambda, t)$ as in Definition 3.3 can be expressed as*

$$w(0; \lambda, t) = \begin{cases} \sqrt{v_t(a)^2 - b^2}, & \text{for } \lambda = a + bi \in X_t; \\ 0, & \text{for } \lambda = a + bi \notin \Xi_t. \end{cases}$$

In particular, $a \mapsto v_t(a)$ is a continuous function defined on \mathbb{R} , and is real analytic and $v_t(a) > 0$ if $\{a + bi : b \in \mathbb{R}\} \cap \Xi_t \neq \emptyset$.

Moreover, we have

$$\Xi_t \cap \mathbb{R} = U_t$$

and

$$\Xi_t = \{a + ib \in \mathbb{C} : |b| < v_t(a)\}$$

and

$$\overline{\Omega_t} = \{a + bi : |b| \geq v_t(a)\} = (\Xi_t)^c.$$

Definition 5.2. *For $t > 0$ and v_t as defined in (5.3), we set*

$$(5.4) \quad \psi_t(a) = a + t \int_{\mathbb{R}} \frac{a-u}{(a-u)^2 + v_t(a)^2} d\mu(u), \quad a \in \mathbb{R},$$

and

$$(5.5) \quad h_t(a) = t \int_{\mathbb{R}} \frac{a-u}{(a-u)^2 + v_t(a)^2} d\mu(u), \quad a \in \mathbb{R}.$$

The following result play a key role in the following discussion. It is taken from [10, Lemma 5] and [32, Theorem 3.14].

Proposition 5.3. *The function $a \mapsto \psi_t(a)$ is a homeomorphism of \mathbb{R} onto \mathbb{R} . Moreover, if $v_t(a) > 0$ or equivalently $\{a + ib : b \in \mathbb{R}\} \cap \Xi_t \neq \emptyset$, then*

$$0 < \frac{d\psi_t(a)}{da} < 2.$$

Consequently, if $v_t(a) > 0$, then

$$(5.6) \quad -1 < \frac{dh_t(a)}{da} < 1.$$

The function v_t can be used to give the following implicit formula of the density of the spectral measure of $x_0 + g_t$ where g_t is a semicircular with mean zero and variance t .

Proposition 5.4. [10, Corollary 3 and Proposition 3] *The spectral measure of $x_0 + g_t$ is absolutely continuous and its density is given by*

$$p_t(\psi_t(a)) = \frac{v_t(a)}{\pi t}.$$

Moreover, the support of μ is contained in the closure of U_t . Consequently, $\text{supp}(\mu) \cap \Xi_t = \emptyset$.

Finally, we need the following useful result taken from [10].

Lemma 5.5. [10, Lemma 3] *The Cauchy transform of G_μ has a continuous extension to $\overline{\Omega_t}$ which is Lipschitz with Lipschitz constant $\leq 1/t$, and one has*

$$|G_\mu(z)|^2 \leq \int_{\mathbb{R}} \frac{1}{|z-u|^2} d\mu(u) \leq \frac{1}{t}$$

for $z \in \overline{\Omega_t}$.

5.2. The push-forward map. Let x_0 be a selfadjoint operator and $\mu = \mu_{x_0}$ be its spectral measure. We study the map $\Phi_{t,\gamma}$ as defined in Section 4. We note that when $\gamma = t$, the operator $g_{t,\gamma}$ is a semicircular operator g_t with mean zero and variance t and the Brown measure of $x_0 + g_t$ is just the spectral measure of $x_0 + g_t$ that is reviewed in Subsection 5.1. Hence, we will assume that $\gamma \neq t$ throughout this section. Let $\tau = t - \gamma$ and $\tau = \tau_1 + i\tau_2$. Since $|\gamma| \leq t$, we see that $\tau_1 > 0$ if $\gamma \neq t$.

Using notations in Subsection 5.1, we have

$$\Phi_{t,\gamma}(\lambda) = \lambda + \gamma \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(u-a)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u), \quad \lambda \in \mathbb{C},$$

and, if $\lambda \in \Xi_t$ then $\Phi_{t,\gamma}(\lambda)$ can be rewritten as

$$\Phi_{t,\gamma}(\lambda) = \lambda + \gamma \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(u-a)^2 + v_t(a)^2} d\mu(u), \quad \lambda \in \Xi_t.$$

Proposition 5.6. *For any $|\gamma| \leq t$ and $\gamma \neq t$, we have*

$$\Phi_{t,\gamma}(\lambda) = \begin{cases} \psi_t(a) - \frac{\tau}{t} h_t(a) + i \frac{\tau b}{t}, & \text{for } \lambda \in \Xi_t; \\ \lambda + \gamma G_\mu(\lambda) & \text{for } \lambda \notin \Xi_t. \end{cases}$$

Proof. We rewrite $\Phi_{t,\gamma}(\lambda)$ as

$$\Phi_{t,\gamma}(\lambda) = \Phi_{t,t}(\lambda) - \tau \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(u-a)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u);$$

and if $\lambda \in \Xi_t$, we can further rewrite it as

$$\begin{aligned} \Phi_{t,\gamma}(\lambda) &= \Phi_{t,t}(\lambda) - \tau \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(u-a)^2 + v_t(a)^2} d\mu(u) \\ &= \Phi_{t,t}(\lambda) - \frac{\tau}{t} h_t(a) + i \frac{\tau b}{t}, \quad \lambda \in \Xi_t, \end{aligned}$$

where we used (5.2) and the definition of h_t in (5.5). If $\lambda \in \Xi_t$, we have

$$\begin{aligned} \Phi_{t,t}(\lambda) &= \lambda + t \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(u-a)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u) \\ &= \psi_t(a) + i \left(b - t \int_{\mathbb{R}} \frac{b}{(u-a)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u) \right) = \psi_t(a) \end{aligned}$$

where we used (5.2) and the definition of ψ_t in (5.4). This proves the case when $\lambda \in \Xi_t$.

Recall that $(\Xi_t)^c = \overline{\Omega_t}$ and $\text{supp}(\mu) \subset U_t \subset \Xi_t$ (recall Proposition 5.4). When $\lambda \notin \Xi_t$, then $w(0; \lambda, t) = 0$. Hence, when $\lambda = a + bi \notin \Xi_t$,

$$\begin{aligned} \Phi_{t,\gamma}(\lambda) &= \lambda + \gamma \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(u-a)^2 + b^2} d\mu(u) \\ &= \lambda + \gamma \int_{\mathbb{R}} \frac{1}{\lambda - u} d\mu(u) = \lambda + \gamma G_\mu(\lambda). \end{aligned}$$

This finishes the proof. \square

The following result says that $\Phi_{t,\gamma}$ is injective on $(\Xi_t)^c$. It can be viewed as a generalization of [10, Lemma 4] where the latter studies the inverse map of subordination function for the free additive convolution of $x_0 + g_t$ (see also [32, Section 2.4] and [29, Proposition 5.5]). We include a proof for convenience.

Proposition 5.7. *The map $\Phi_{t,\gamma}(\lambda)$ is an injective map when restricted to $(\Xi_t)^c$ and is conformal when λ is in the interior of $(\Xi_t)^c$.*

Proof. If $\lambda \in (\Xi_t)^c$, by Proposition 5.6, we have $\Phi_{t,\gamma}(\lambda) = \lambda + \gamma G_\mu(\lambda)$. Hence, for $\alpha_1, \alpha_2 \in (\Xi_t)^c$, we have

$$\begin{aligned} \Phi_{t,\gamma}(\alpha_1) - \Phi_{t,\gamma}(\alpha_2) &= \alpha_1 - \alpha_2 + \gamma(G_\mu(\alpha_1) - G_\mu(\alpha_2)) \\ &= (\alpha_1 - \alpha_2) \left(1 + \gamma \frac{G_\mu(\alpha_1) - G_\mu(\alpha_2)}{\alpha_1 - \alpha_2} \right). \end{aligned}$$

Then, by Cauchy-Schwartz inequality,

$$\begin{aligned} \left| \frac{G_\mu(\alpha_1) - G_\mu(\alpha_2)}{\alpha_1 - \alpha_2} \right| &= \left| \int_{\mathbb{R}} \frac{1}{(\alpha_1 - u)(\alpha_2 - u)} d\mu(u) \right| \\ &\leq \left(\int_{\mathbb{R}} \frac{d\mu(u)}{|\alpha_1 - u|^2} \int_{\mathbb{R}} \frac{d\mu(u)}{|\alpha_2 - u|^2} \right)^{1/2} \leq \frac{1}{t} \end{aligned}$$

where we used Lemma 5.5 in the final step. It can be shown that we can not have equality in the Cauchy-Schwartz inequality used above. Indeed, if equality holds, then

$$\left| \int_{\mathbb{R}} \frac{1}{(\alpha_1 - u)(\alpha_2 - u)} d\mu(u) \right| = \frac{1}{t}$$

for some $\alpha_1, \alpha_2 \in (\Xi_t)^c = \overline{\Omega_t}$. One can show that μ must be a Dirac measure (see [10, Lemma 4] for details). Therefore,

$$\Phi_{t,\gamma}(\alpha_1) - \Phi_{t,\gamma}(\alpha_2) \neq 0$$

for $\alpha_1 \neq \alpha_2$ and $\alpha_1, \alpha_2 \in (\Xi_t)^c$. \square

Lemma 5.8. *For $|\gamma| \leq t$ and $\gamma \neq t$, let $\tau = t - \gamma$ and $\tau = \tau_1 + i\tau_2$, then we have $0 < \frac{|\tau|^2}{t\tau_1} \leq 2$.*

Proof. The assumption implies directly that $\tau_1 > 0$. Write $\gamma = \gamma_1 + i\gamma_2$, then $\tau_1 = t - \gamma_1$ and $\tau_2 = \gamma_2$. Hence

$$|\tau|^2 - 2t\tau_1 = (t - \gamma_1)^2 + \gamma_2^2 - 2t(t - \gamma_1) = |\gamma|^2 - t^2 \leq 0.$$

The result then follows. \square

Theorem 5.9. *For any $|\gamma| \leq t$ and $\gamma \neq t$, the map $\Phi_{t,\gamma}$ is a smooth, injective map on Ξ_t . Moreover, the determinant of the Jacobian of $\Phi_{t,\gamma}$ is strictly positive at all $\lambda \in \Xi_t$ and can be expressed*

$$(5.7) \quad \det(\text{Jacobian}(\Phi_{t,\gamma}))(\lambda) = \frac{\tau_1}{t} \left[1 + \left(1 - \frac{|\tau|^2}{t\tau_1} \right) \frac{dh_t(a)}{da} \right], \quad \lambda = a + ib,$$

where $\tau = \tau_1 + i\tau_2$ and h_t is defined in 5.5.

Proof. By Proposition 5.6, for $\lambda \in \Xi_t$, we have

$$\Phi_{t,\gamma}(\lambda) = \psi_t(a) - \frac{\tau}{t}h_t(a) + i\frac{\tau b}{t}.$$

Recall $\psi_t(a) = a + h_t(a)$. Hence, for $\Phi_{t,\gamma}(\lambda) = z_1 + iz_2$, we deduce

$$(5.8) \quad z_1 = z_1(a, b) = a + \left(1 - \frac{\tau_1}{t}\right)h_t(a) - \frac{\tau_2}{t}b, \quad z_2 = z_2(a, b) = -\frac{\tau_2}{t}h_t(a) + \frac{\tau_1}{t}b.$$

We then have

$$\text{Jacobian}(\Phi_{t,\gamma}(\lambda)) = \begin{bmatrix} 1 + \left(1 - \frac{\tau_1}{t}\right)h'_t(a) & -\frac{\tau_2}{t} \\ -\frac{\tau_2}{t}h'_t(a) & \frac{\tau_1}{t} \end{bmatrix}.$$

It follows that

$$\det(\text{Jacobian}(\Phi_{t,\gamma}))(\lambda) = \frac{\tau_1}{t} \left[1 + \left(1 - \frac{|\tau|^2}{t\tau_1}\right) \frac{dh_t(a)}{da} \right].$$

By Proposition 5.3, $-1 < h'_t(a) < 1$, hence we have

$$-1 < \left(1 - \frac{|\tau|^2}{t\tau_1}\right) \frac{dh_t(a)}{da} < 1$$

thanks to Lemma 5.8. This implies that $\det(\text{Jacobian}(\Phi_{t,\gamma}))(\lambda)$ is independent of b and $\det(\text{Jacobian}(\Phi_{t,\gamma}))(\lambda) > 0$ for all $\lambda \in \Xi_t$.

We next show that the map is injective in Ξ_t . If $a_1 + ib_1, a_2 + ib_2 \in \Xi_t$ and $\Phi_{t,\gamma}(a_1 + ia_2) = \Phi_{t,\gamma}(b_1 + ib_2)$. Then we have, by using (5.8),

$$\begin{aligned} a_1 - a_2 + \left(1 - \frac{\tau_1}{t}\right)(h_t(a_1) - h_t(a_2)) &= -\frac{\tau_2}{t}(b_2 - b_1) \\ -\frac{\tau_2}{t}(h_t(a_1) - h_t(a_2)) &= \frac{\tau_1}{t}(b_2 - b_1), \end{aligned}$$

which yields (by canceling $b_2 - b_1$),

$$(a_1 - a_2) + \left(1 - \frac{|\tau|^2}{t\tau_1}\right)(h_t(a_1) - h_t(a_2)).$$

By the first part of the proof, we see that the function $a \mapsto \frac{\tau_1}{t} \left[1 + \left(1 - \frac{|\tau|^2}{t\tau_1}\right) \frac{dh_t(a)}{da} \right]$ is strictly positive. Consequently, $a_1 = a_2$. We then deduce that $b_1 = b_2$. The injectivity property is established. \square

Corollary 5.10. *For any $|\gamma| \leq t$ and $\gamma \neq t$, the map $\Phi_{t,\gamma}$ is a homeomorphism of \mathbb{C} to \mathbb{C} .*

Proof. We show that $\Phi_{t,\gamma}(\Xi_t) \cap \Phi_{t,\gamma}((\Xi_t)^c) = \emptyset$. Recall that

$$(\Xi_t)^c = \{a + bi : |b| \geq v_t(a)\} = \overline{\Omega}_t.$$

For $\lambda_0 = a + ib$ so that $b \geq v_t(a)$, if $\Phi_{t,\gamma}(\lambda_0) \in \Phi_{t,\gamma}(\Xi_t)$, consider the vertical half-line $\{\lambda = a + id : d \geq b\}$ starting at λ_0 . Since $\lim_{d \rightarrow \infty} \Phi_{t,\gamma}(a + id) = \infty$, we then see that there is another point $\lambda_1 = a + ib_1$ where $b_1 > b$ so that $\Phi_{t,\gamma}(\lambda_1) \in \partial(\Phi_{t,\gamma}(\Xi_t))$, the boundary of $\Phi_{t,\gamma}(\Xi_t)$. On the other hand $\Phi_{t,\gamma}(a + iv_t(b)) \in \partial(\Phi_{t,\gamma}(\Xi_t))$, and both λ_1 and $a + iv_t(b)$ are in $(\Xi_t)^c$, this contradicts to Proposition 5.7. Therefore,

$$\Phi_{t,\gamma}(\Xi_t) \cap \Phi_{t,\gamma}((\Xi_t)^c) = \emptyset.$$

Since $\Phi_{t,\gamma}$ is a continuous function on \mathbb{C} and it maps a neighborhood of infinity to some neighborhood of infinity, one can then deduce that $\Phi_{t,\gamma}(\mathbb{C}) = \mathbb{C}$ by some standard arguments in topology. We conclude that $\Phi_{t,\gamma}$ is a homeomorphism of \mathbb{C} to \mathbb{C} due to Proposition 5.7 and Theorem 5.9. \square

5.3. Brown measure of addition with a selfadjoint operator. We apply results in Section 4 to give a new proof for the Brown measure of $x_0 + c_t$ and recover a result in our previous work with Ho [32]. We then study the Brown measure of $x_0 + g_{t,\gamma}$ that extends results by Hall and Ho [28, 31] to all twisted elliptic operators.

Theorem 5.11. [32, Theorem 3.10] *For $\lambda = a + bi \in \Xi_t$, then the Brown measure of $x_0 + c_t$ is absolutely continuous at λ and the density at λ is given by*

$$d\mu_{x_0+c_t}(a+ib) = \frac{1}{\pi t} \left(1 - \frac{t}{2} \frac{d}{da} \int_{\mathbb{R}} \frac{u}{(a-u)^2 + v_t(a)^2} d\mu(u) \right) da db,$$

which can also be expressed as

$$(5.9) \quad d\mu_{x_0+c_t}(a+ib) = \frac{1}{2\pi t} \frac{d\psi_t(a)}{da} da db.$$

In particular, the density is constant along the vertical directions.

Proof. One can deduce the density formula directly from the general formula in Theorem 4.6. We note that

$$\begin{aligned} \frac{\partial}{\partial \bar{\lambda}} \left(\phi(x_0^*((x_0 - \lambda \mathbf{1})^*(x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2)^{-1}) \right) \\ = \frac{\partial}{\partial \bar{\lambda}} \left(\int_{\mathbb{R}} \frac{u}{(a-u)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u) \right) \\ = \frac{1}{2} \frac{\partial}{\partial a} \left(\int_{\mathbb{R}} \frac{u}{(a-u)^2 + v_t(a)^2} \mu(u) \right) \end{aligned}$$

where we used the fact that the integration is independent of b . The first formula then follows from (4.12). Using the formula for ψ_t in (5.4), we have

$$\begin{aligned} \frac{d\psi_t(a)}{da} &= \frac{d}{da} \left(a + t \int_{\mathbb{R}} \frac{a-u}{(a-u)^2 + v_t(a)^2} d\mu(u) \right) \\ &= 2 - t \frac{d}{da} \int_{\mathbb{R}} \frac{u}{(a-u)^2 + v_t(a)^2} d\mu(u) \end{aligned}$$

where we used (5.2). This establish the second formula. \square

Proposition 5.12. [32, Lemma 3.11 and Proposition 3.16] *The Brown measure $\mu_{x_0+c_t}$ concentrates in Ξ_t . That is $\mu_{x_0+c_t}(\Xi_t) = 1$. The spectral measure of $x_0 + g_t$ is the push-forward of the Brown measure $\mu_{x_0+c_t}$ under the map $\lambda \mapsto \Phi_{t,t}$, where*

$$\Phi_{t,t} = \lambda + t \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(u-a)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u).$$

Lemma 5.13. *Let $\delta(a) = \delta_{t,\gamma}(a) = a + \left(1 - \frac{|\tau|^2}{t\tau_1}\right) h_t(a)$, then $a \mapsto \delta(a)$ is a homeomorphism of \mathbb{R} onto \mathbb{R} . Then,*

$$0 < \delta'(a) < 2$$

for all $a \in U_t$ and $a \in (\overline{U_t})^c$. Moreover, if $a \in U_t$,

$$\delta'(a) = \frac{t}{\tau_1} \cdot \det(\text{Jacobian}(\Phi_{t,\gamma}))(\lambda).$$

Proof. Using the characterization of Ξ_t and U_t in Proposition 5.1, it follows that the result holds for $a \in U_t$. Indeed, we have

$$\delta'(a) = \frac{t}{\tau_1} \cdot \det(\text{Jacobian}(\Phi_{t,\gamma}))(\lambda)$$

for any $\lambda \in \Xi_t$ such that $\Re(\lambda) = a$. If $a \in (\overline{U_t})^c$, by Proposition 5.4, then $\text{supp}(\mu) \cap (\overline{U_t})^c = \emptyset$. Hence, $v_t(a) = 0$ and

$$h_t(a) = t \int_{\mathbb{R}} \frac{1}{a-u} d\mu(u) = tG_\mu(a).$$

It follows that $|h'_t(a)| \leq 1$ thanks to Lemma 5.5. Note that $a \mapsto h_t(a)$ is strictly convex on any open interval in $(\overline{U_t})^c$. Hence, $h'_t(a)$ can not take local maximum in the open interval. We then conclude that $0 < h'_t(a) < 1$ if $a \in (\overline{U_t})^c$. \square

Theorem 5.14. *Let x_0 be a selfadjoint operator that is free from $g_{t,\gamma}$. For any $|\lambda| \leq t$ with $\lambda \neq t$, the map $\Phi_{t,\gamma}$ satisfy Condition A. The Brown measure of $x_0 + g_{t,\gamma}$ is the push-forward map of the Brown measure of $x_0 + c_t$ under the map $\Phi_{t,\gamma}$.*

Moreover, the Brown measure $\mu_{x_0+g_{t,\gamma}}$ takes the full measure on $\Phi_{t,\gamma}(\Xi_t)$ and the density is given by

$$(5.10) \quad d\mu_{x_0+g_{t,\gamma}}(z) = \frac{1}{2\pi\tau_1} \frac{d\psi_t(a)}{d\delta(a)} dz_1 dz_2, \quad z \in \Phi_{t,\gamma}(\Xi_t)$$

where $z = z_1 + iz_2 = \Phi_{t,\gamma}(a + ib)$ and

$$\delta(a) = a + \left(1 - \frac{|\tau|^2}{t\tau_1}\right) h_t(a).$$

Proof. By Theorem 5.9, we deduce that $\Phi_{t,\gamma}$ satisfies Condition A. Hence, Theorem 4.16 applies in this case. Recall that if $a + ib \in \Xi_t$, then $a \in U_t$. Using the density formula for $x_0 + c_t$ in (5.9) and determinant of the Jacobian of $\Phi_{t,\gamma}$ as in (5.7), it then follows that the density of $x_0 + g_{t,\gamma}$ can be expressed as

$$\frac{1}{2\pi t} \frac{d\psi_t(a)/da}{\det(\text{Jacobian}(\Phi_{t,\gamma})(\lambda))} = \frac{1}{2\pi\tau_1} \frac{d\psi_t(a)/da}{d\delta(a)/da} = \frac{1}{2\pi\tau_1} \frac{d\psi_t(a)}{d\delta(a)}.$$

The result is established. \square

Corollary 5.15. [12, Example 5.3] *The Brown measure of $g_{t,\gamma}$ is the uniform measure in the rotated ellipse with parametrization*

$$e^{i\alpha} \left(\sqrt{t} e^{i\theta} + \frac{|\gamma|}{\sqrt{t}} e^{-i\theta} \right),$$

for $\theta \in [0, 2\pi]$, where α is such that $\gamma = |\gamma| e^{2i\alpha}$.

Proof. Take $x_0 = 0$ and $\mu = \delta_0$. In this case, the formula for the set Ξ_t is simplified as

$$\Xi_t = \{\lambda = a + ib : |\lambda| < \sqrt{t}\},$$

which is also the support of the circular operator c_t . Then the condition determining $w(0; \lambda, t)$ is written as

$$\frac{1}{|\lambda|^2 + w(0; \lambda, t)^2} = \frac{1}{t}, \quad |\lambda| < \sqrt{t}.$$

Hence $w(0; \lambda, t) = \sqrt{t - |\lambda|^2}$. Then the push forward map is

$$\Phi_{t,\gamma}(\lambda) = \lambda + \gamma \frac{\bar{\lambda}}{|\lambda|^2 + w(0; \lambda, t)^2} = \lambda + \frac{\gamma \bar{\lambda}}{t}$$

for $|\lambda| < \sqrt{t}$. Write $\gamma = |\lambda|e^{i\alpha}$. Then for any $0 < r < \sqrt{t}$, the circle centered at origin with radius r is mapped to the ellipse with parametrization

$$re^{i\theta} + \frac{r|\gamma|}{t}e^{i(\alpha-\theta)}.$$

Moreover, for $|\lambda| < \sqrt{t}$ and $\lambda = a + ib$, we have

$$h_t(a) = t \frac{a}{|\lambda|^2 + w(0; \lambda, t)^2} = a.$$

Hence, the determinant of the Jacobian of the push forward map $\Phi_{t,\gamma}$ is the constant

$$\det(\text{Jacobian}(\Phi_{t,\gamma}))(\lambda) = \frac{\tau_1}{t} \left[1 + \left(1 - \frac{|\tau|^2}{t\tau_1} \right) \right], \quad \lambda = a + ib,$$

where $\tau_1 + i\tau_2 = \tau = t - \gamma$. Recall that the density of the Brown measure of c_t is the uniform measure on the circle $\{\lambda : |\lambda| < \sqrt{t}\}$. Therefore, the Brown measure of $g_{t,\gamma}$ is the uniform measure on the ellipse with parametrization $\sqrt{t}e^{i\theta} + |\gamma|e^{i(\alpha-\theta)}/\sqrt{t}$ for $\theta \in [0, 2\pi]$, where α is determined by $\gamma = |\gamma|e^{2i\alpha}$. The result follows. \square

We now discuss some special cases which allow us to recover main results in [28, 31].

Example 5.16 (The semicircular operator). If $\gamma = t$, the operator $g_{t,\gamma}$ is a semicircular operator g_t with mean zero and variance t . In this case $\tau = 0$. Hence, if $\lambda \in \Xi_t$, then

$$\Phi_{t,t}(\lambda) = \psi_t(a) = a + t \int_{\mathbb{R}} \frac{a-u}{(a-u)^2 + v_t(a)^2} d\mu(u), \quad a \in \mathbb{R},$$

and $\delta(a) = a$. More precisely, we have

$$\Phi_{t,t}(a+ib) = \psi_t(a), \quad \text{for } -v_t(a) \leq b \leq v_t(a).$$

In this case, Condition A is not satisfied. However, the spectral measure of $x_0 + g_t$ is the push-forward of the Brown measure of $x_0 + c_t$ under the map $\Phi_{t,t}$ by Proposition 5.12.

Moreover, the statement of Lemma 4.4 does hold in this case as well. Indeed, by Theorem 3.7, for any $\varepsilon > 0$, we have

$$p_z^{g,(t,t)}(\varepsilon) = p_\lambda^{(0)}(w(\varepsilon))$$

where $w(\varepsilon) = w(\varepsilon; \lambda, t)$ and $z = \Phi_{t,\gamma}(\lambda)$. By the Notation 3.8, This can be rewritten as

$$\begin{aligned} & -\phi \left[(x_0 + g_t - z)^* \left((x_0 + g_t - z)^* (x_0 + g_t - z) + \varepsilon^2 \mathbf{1} \right)^{-1} \right] \\ & = -\phi \left[(x_0 - \lambda \mathbf{1})^* \left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(\varepsilon)^2 \mathbf{1} \right)^{-1} \right]. \end{aligned}$$

By Lemma 3.6, for any $\lambda \in \Xi_t$ (equivalently, $t > \lambda_1(\mu_{|x_0-\lambda|})$), we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} p_\lambda^{(0)}(w(\varepsilon)) & = -\phi \left[(x_0 - \lambda \mathbf{1})^* \left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1} \right)^{-1} \right] \\ & = p_\lambda^{(0)}(w(0; \lambda, t)). \end{aligned}$$

Hence, for any $\lambda \in \Xi_t$, by choosing $z = \lambda(t) = \lambda + \gamma p_\lambda^{(0)}(w(\varepsilon))$, we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} p_z^{g,(t,t)}(\varepsilon) & = -\phi \left[(x_0 - \lambda \mathbf{1})^* \left((x_0 - \lambda \mathbf{1})^* (x_0 - \lambda \mathbf{1}) + w(0; \lambda, t)^2 \mathbf{1} \right)^{-1} \right] \\ & = p_\lambda^{(0)}(w(0; \lambda, t)). \end{aligned}$$

We note that $p_\lambda^{(0)}(w(0; \lambda, t))$ can be expressed as

$$\begin{aligned} p_\lambda^{(0)}(w(0; \lambda, t)) &= \int_{\mathbb{R}} \frac{\bar{\lambda} - u}{(a - u)^2 + b^2 + w(0; \lambda, t)^2} d\mu(u) \\ &= \int_{\mathbb{R}} \frac{a - u}{(a - u)^2 + v_t(a)^2} d\mu(u) - \frac{ib}{t}, \end{aligned}$$

where we used (5.2) and the fact that $b^2 + w(0; \lambda, t)^2 = v_t(a)^2$ for $a + ib \in \Xi_t$.

To summarize, for $a \in \mathbb{R}$ fixed so that $v_t(a) > 0$, then: (1) the limit $p_z^{g, (t, t)}(\varepsilon)$ as ε tends to zero have different limit as long as (z, ε) tends to $(\psi_t(a), 0)$ along different paths $(\lambda(t), \varepsilon)$ depending on b ; (2) although the $\lim_{\varepsilon \rightarrow 0^+} \lambda(t) = \Phi_{t, t}(a + ib) = \psi_t(a)$ for any $-v_t(a) \leq b \leq v_t(a)$, the limit of the partial derivative $\lim_{\varepsilon \rightarrow 0^+} p_z^{g, (t, t)}(\varepsilon)$ detects the value b and *remembers* where it came from. Namely, by looking at the limit $\lim_{\varepsilon \rightarrow 0^+} \lambda(t) = \Phi_{t, t}(a + ib)$ we can not identify b since $\Phi_{t, t}$ is not one-to-one, but the limit of the partial derivative $\lim_{\varepsilon \rightarrow 0^+} p_z^{g, (t, t)}(\varepsilon)$ does the work. This phenomena is different from what Lemma 4.4 told us when $\Phi_{t, \gamma}$ is one-to-one.

Example 5.17 (The imaginary multiple of a semicircular operator). If $\gamma = it$, the operator $g_{t, \gamma}$ has the same distribution as ig_t . In this case, $\tau = 2t$, and for $\lambda \in \Xi_t$, we have

$$\begin{aligned} \Phi_{t, it}(\lambda) &= \psi_t(a) - 2h_t(a) + 2ib \\ &= a - h_t(a) + 2ib \\ &= t \int_{\mathbb{R}} \frac{u}{(u - a)^2 + v_t(a)^2} d\mu(u) + 2ib \end{aligned}$$

and

$$\delta_{t, it}(a) = a - h_t(a) = t \int_{\mathbb{R}} \frac{u}{(u - a)^2 + v_t(a)^2} d\mu(u).$$

Example 5.18 (The elliptic operator). If $\gamma = s \in \mathbb{R}$ with $-t < s < t$, the operator $g_{t, \gamma}$ has the same distribution as an elliptic operator. In this case, $\tau = t - s$, and for $\lambda \in \Xi_t$, we have

$$\begin{aligned} \Phi_{t, s}(\lambda) &= \psi_t(a) - \frac{t - s}{t} h_t(a) + i \frac{t - s}{t} b \\ &= a + \frac{s}{t} h_t(a) + i \frac{t - s}{t} b \\ &= \left(a + s \int_{\mathbb{R}} \frac{a - u}{(a - u)^2 + v_t(a)^2} d\mu(u) \right) + i \frac{(t - s)b}{t}, \end{aligned}$$

and

$$\delta_{t, s}(a) = a + s \int_{\mathbb{R}} \frac{a - u}{(a - u)^2 + v_t(a)^2} d\mu(u).$$

We also note that $\Phi_{t, s}(\lambda) = z_1 + iz_2$ where

$$z_1 = \delta_{t, s}(a), \quad z_2 = \frac{(t - s)b}{t}$$

and $\psi_t(a)$ can be written as

$$\psi_t(a) = z_1 + \frac{t - s}{t} h_t(a) = z_1 + (t - s) \int_{\mathbb{R}} \frac{a - u}{(a - u)^2 + v_t(a)^2} d\mu(u).$$

Therefore, the density of the Brown measure at its support $\overline{\Phi_{t,s}(\Xi_t)}$ is given by

$$\begin{aligned} & d\mu_{x_0+g_{t,s}}(z_1 + iz_2) \\ &= \frac{1}{2\pi(t-s)} \left(1 + (t-s) \frac{d}{dz_1} \int_{\mathbb{R}} \frac{a-u}{(a-u)^2 + v_t(a)^2} d\mu(u) \right) dz_1 dz_2, \end{aligned}$$

if $v_t(a) > 0$. Here we remind the reader that the map $a \mapsto z_1 = \delta_{t,s}(a)$ is a homeomorphism from \mathbb{R} to \mathbb{R} . This recovers the main result in [28, 31]

Remark 5.19. [The twisted (ν, δ) -coordinate.] Fix $a \in \mathbb{R}$, as in the proof of Theorem 5.9 (see (5.8)), the map $b \mapsto \Phi_{t,\gamma}(a + bi)$ is an affine transform of b . The density formula of $x_0 + c_t$ in Theorem 5.11 is independent of b . Hence, the density formula of $x_0 + g_{t,\gamma}$ must depend only on one parameter. It is indeed the case as in Theorem 5.14, where the density is expressed in terms of parameter a coming from Ξ_t , the support of $x_0 + c_t$.

We now describe an analogue of the formulation in the recent work [29] where the authors study the free multiplicative Brownian motions as follows. For $z = z_1 + iz_2$, consider twisted (ν, δ) -coordinate determined by

$$a + ib = i\tau\nu + \delta,$$

where $\nu, \delta \in \mathbb{R}$. They can be written as

$$\delta = a + \frac{\tau_2}{\tau_1}b, \quad \nu = \frac{b}{\tau_1}.$$

Using notations in the proof of Theorem 5.9 and the formula of $\Phi_{t,\gamma}(\lambda)$ as in (5.8), for $\lambda = \lambda_1 + i\lambda_2 \in \Xi_t$, we have

$$\begin{aligned} \delta(\Phi_{t,\gamma}(\lambda)) &= z_1 + \frac{\tau_2}{\tau_1}z_2 \\ &= \left[a + \left(1 - \frac{\tau_1}{t} \right) h_t(a) - \frac{\tau_2}{t}b \right] + \frac{\tau_2}{\tau_1} \left(-\frac{\tau_2}{t}h_t(a) + \frac{\tau_1}{t}b \right) \\ &= a + \left(1 - \frac{|\tau|^2}{t\tau_1} \right) h_t(a) = \delta_{t,\gamma}(a), \end{aligned}$$

and

$$\nu(\Phi_{t,\gamma}(\lambda)) = \frac{z_2}{\tau_1} = \frac{1}{t} \left[b - \frac{\tau_2}{\tau_1}h_t(a) \right].$$

We can then say that the Brown measure $\mu_{x_0+g_{t,\gamma}}$ is constant along the ν -direction.

6. EXAMPLE OF NON-SELFADJOINT OPERATORS

In this section, we compute two examples of x_0 for the Brown measure of $x_0 + c_t$ and $x_0 + g_{t,\gamma}$. The first example is quasi-nilpotent DT operator, and the second example is Haar unitary operator.

6.1. The quasi-nilpotent DT operator. The quasi-nilpotent DT-operator T was introduced by Dykema–Haagerup [15]. The operator played a key role in Brown measure of free random variables and invariant subspace problem in type II_1 factors [16, 26]. It can be described as the limit in $*$ -moments of random matrices of the form

$$T^{(n)} = \begin{bmatrix} 0 & t_{1,2} & \cdots & t_{1,n} \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & t_{n-1,n} \\ 0 & \cdots & 0 & 0 \end{bmatrix}$$

where $\{\Re(t_{i,j}), \Im(t_{i,j})\}$ is a set of $n(n-1)$ independent identically distributed Gaussian random variables with mean zero and variance $\frac{1}{2n}$. We refer the reader to [16, 15, 1] for the construction of T in a finite von Neumann algebra.

Proposition 6.1. *Given $\lambda \in \mathbb{C}$ Let $\sigma \in \mathbb{R}$ with $1 + |\lambda|^2\sigma > 0$ and $\mu^2 = -\frac{e^\sigma}{\sigma}(1 + |\lambda|^2\sigma)$. We have*

$$\begin{cases} \phi(((T - \lambda\mathbf{1})^*(T - \lambda\mathbf{1}) + \mu^2)^{-1}) = e^{-\sigma} - 1 \\ \phi((T - \lambda\mathbf{1})((T - \lambda\mathbf{1})^*(T - \lambda\mathbf{1}) + \mu^2)^{-1}) = \lambda\sigma \\ \phi((T - \lambda\mathbf{1})^*((T - \lambda\mathbf{1})^*(T - \lambda\mathbf{1}) + \mu^2)^{-1}) = \bar{\lambda}\sigma. \end{cases}$$

Proof. It can be obtained by taking the expectation of (3.3) in [1]. See [1, Pages 586-588]. \square

Proposition 6.2. [1, Theorem 4.3] *For any $t > 0$, the Brown measure of $T + c_t$ is the uniform measure on the closed disk $\bar{B}(0, \frac{1}{\sqrt{\log(1+1/t)}})$.*

Proof. Choose σ so that $e^{-\sigma} - 1 = \frac{1}{t}$. Then $\sigma = -\log(1 + 1/t)$. Consider the solution of $\varepsilon > 0$ for the equation

$$\phi((T - \lambda\mathbf{1})^*(T - \lambda\mathbf{1}) + \varepsilon^2) = \frac{1}{t},$$

we then deduce that $w(0; \lambda, t)^2 = \mu^2 = -\frac{e^\sigma}{\sigma}(1 + |\lambda|^2\sigma)$ by applying the results in Section 5 for $x_0 = T$. Hence, $w(0; \lambda, t) > 0$ if and only if $(1 + |\lambda|^2\sigma) > 0$. Therefore,

$$\begin{aligned} \Xi_t &= \left\{ \lambda : \phi(((T - \lambda\mathbf{1})^*(T - \lambda\mathbf{1}) + \mu^2)^{-1}) > \frac{1}{t} \right\} \\ &= \{ \lambda : w(0; \lambda, t) > 0 \} = \{ \lambda : 1 + |\lambda|^2\sigma > 0 \} \\ &= \left\{ \lambda : |\lambda| < \frac{1}{\sqrt{\log(1 + 1/t)}} \right\}. \end{aligned}$$

In addition, by the Proof of Theorem 4.6, we have

$$\begin{aligned} &\frac{\partial}{\partial \lambda} \log \Delta((T + c_t - \lambda\mathbf{1})^*(T + c_t - \lambda\mathbf{1})) \\ &= p_\lambda^{(t)}(0) = p_\lambda^{(0)}(w(0; \lambda, t)) \\ &= -\phi((T - \lambda\mathbf{1})^*((T - \lambda\mathbf{1})^*(x_0 - \lambda\mathbf{1}) + w(0; \lambda, t)^2)^{-1}) \\ &= -\phi((T - \lambda\mathbf{1})^*((T - \lambda\mathbf{1})^*(x_0 - \lambda\mathbf{1}) + \mu^2)^{-1}) = -\bar{\lambda}\sigma. \end{aligned}$$

Hence, the density of $T + c_t$ at $\lambda \in \Omega_t = B(0, \frac{1}{\sqrt{\log(1+1/t)}})$ is given by

$$d\mu_{T+c_t} = \frac{1}{\pi} \frac{\partial^2}{\partial \lambda^2} \log \Delta((T + c_t - \lambda\mathbf{1})^*(T + c_t - \lambda\mathbf{1})) = -\frac{\sigma}{\pi} = \frac{\log(1 + 1/t)}{\pi}.$$

It follows that the Brown measure of $T + c_t$ is equal to the uniform measure on the $\bar{B}(0, \frac{1}{\sqrt{\log(1+1/t)}})$. \square

We then calculate the push-forward map from $T + c_t$ to $T + g_{t,\gamma}$. We have, by Notation 4.1, that

$$\Phi_{t,\gamma}(\lambda) = \lambda - \gamma \phi((T - \lambda\mathbf{1})^*((T - \lambda\mathbf{1})^*(x_0 - \lambda\mathbf{1}) + w(0; \lambda, t)^2)^{-1}) = \lambda - \gamma \bar{\lambda}\sigma,$$

where $\sigma = -\log(1 + 1/t)$. Therefore,

$$\Phi_{t,\gamma}(\lambda) = \lambda + \overline{\lambda}\gamma \cdot \log(1 + 1/t).$$

Recall that $|\gamma| \leq t$. In particular, it follows that $\Phi_{t,\gamma}$ is one-to-one in the disk $B(0, \frac{1}{\sqrt{\log(1+1/t)}})$. Hence, Theorem 4.16 applies and it follows that the Brown measure of $T + g_{t,\gamma}$ is supported in the ellipse with parametrization

$$\frac{1}{\sqrt{\log(1 + 1/t)}} e^{i\theta} + |\gamma| \sqrt{\log(1 + 1/t)} e^{i(\alpha - \theta)}, \quad \theta \in [0, 2\pi]$$

where $\alpha = \arg \gamma$. In fact, the Brown measure of $T + g_{t,\gamma}$ is the uniform measure on its support.

6.2. The Haar unitary. The Haar unitary operator u is R -diagonal. Hence, the Brown measure of $u + c_t$ is R -diagonal and the density formula is determined by the S -transform of $(u + c_t)^*(u + c_t)$ [24, 25]. The Brown measure of an R -diagonal operator is the limit of the ESD of certain non-normal random matrix model called single ring theorem named after Feinberg and Zee (see [22] and the earlier physics paper [18, 17]). However, it is not easy to obtain a precise formula for the Brown measure of $u + c_t$ by calculating the S -transform. See [23, Section 3.1] for a relevant calculation on the Cauchy transform of $c + \lambda u$ where c is the standard circular operator with variance one and $\lambda \in \mathbb{C} \setminus \{0\}$. Here we calculate the push-forward map from the Brown measure of $u + c_t$ to $u + g_{t,\gamma}$ and as a by product we are able to calculate the Brown measure of $u + c_t$.

For $t > 0$, following Definition 3.3 and Remark 3.4, we would like to determine the choice of s so that

$$\int_0^\infty \frac{1}{s^2 + x^2} d\mu_{|u - \lambda 1|}(x) = \frac{1}{t}.$$

To this end, we write down the following elementary results for convenience.

Given $\lambda_0 \neq 0$, consider the quadratic equation

$$(6.1) \quad (z - \lambda_0)(1 - z\overline{\lambda_0}) + s^2 z = 0.$$

Rewrite it as

$$\overline{\lambda_0} z^2 - (|\lambda_0|^2 + 1 + s^2)z + \lambda_0 = 0.$$

We set

$$(6.2) \quad \delta = |\lambda_0|^2 + 1 + s^2.$$

Then (6.1) has two solutions

$$(6.3) \quad Z_1 = \frac{\delta - \sqrt{\delta^2 - 4|\lambda_0|^2}}{2\overline{\lambda_0}},$$

$$(6.4) \quad Z_2 = \frac{\delta + \sqrt{\delta^2 - 4|\lambda_0|^2}}{2\overline{\lambda_0}}.$$

Note that

$$Z_1 Z_2 = \lambda_0 / \overline{\lambda_0}, \quad |Z_1 Z_2| = 1,$$

and

$$(6.5) \quad \begin{aligned} \delta^2 - 4|\lambda_0|^2 &\geq (|\lambda_0|^2 + 1)^2 - 4|\lambda_0|^2 = (|\lambda_0|^2 - 1)^2 \geq 0 \\ \delta^2 - 4|\lambda_0|^2 &\geq (|\lambda_0|^2 + 1)^2 - 4|\lambda_0|^2 = (|\lambda_0|^2 - 1)^2 \geq 0 \end{aligned}$$

for any s . Hence, for any $\lambda_0 \neq 0$, we have $|Z_1| \leq 1$ and $|Z_2| \geq 1$. Moreover, when $s > 0$, we have $|Z_1| < 1$ and $|Z_2| > 1$.

Lemma 6.3. *For any λ_0 , we have*

$$(6.6) \quad \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{|e^{i\theta} - \lambda_0|^2 + s^2} d\theta = \frac{1}{\sqrt{\delta^2 - 4|\lambda_0|^2}}$$

where $\delta = |\lambda_0|^2 + 1 + s^2$ is given by (6.2) and the integration is infinity if $\delta^2 = 4|\lambda_0|^2$. In particular, when $s = 0$,

$$(6.7) \quad \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{|e^{i\theta} - \lambda_0|^2} d\theta = \begin{cases} \frac{1}{1-|\lambda_0|^2} & \text{if } 0 \leq |\lambda_0| < 1, \\ \frac{1}{|\lambda_0|^2-1} & \text{if } |\lambda_0| > 1, \\ \infty & \text{if } |\lambda_0| = 1. \end{cases}$$

Proof. We note that

$$\frac{1}{|e^{i\theta} - \lambda_0|^2 + s^2} = \frac{e^{i\theta}}{(e^{i\theta} - \lambda_0)(1 - e^{i\theta}\overline{\lambda_0}) + s^2 e^{i\theta}}.$$

Using the formula (6.3) and (6.4) for roots of the quadratic equation (6.1), we have

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{|e^{i\theta} - \lambda_0|^2 + s^2} d\theta &= \frac{1}{2\pi i} \int_{|z|=1} \frac{1}{(z - \lambda_0)(1 - z\overline{\lambda_0}) + s^2 z} dz \\ &= \frac{1}{2\pi i} \int_{|z|=1} \frac{1}{-\overline{\lambda_0}(z - Z_1)(z - Z_2)} dz \\ &= \frac{-\lambda_0}{|\lambda_0|^2(Z_1 - Z_2)} \\ &= \frac{1}{\sqrt{\delta^2 - 4|\lambda_0|^2}} \end{aligned}$$

provided that $\delta^2 - 4|\lambda_0|^2 \neq 0$. When $s = 0$ and $|\lambda_0| \neq 1$, the above calculation still works.

For the case that $\delta^2 - 4|\lambda_0|^2 = 0$. Recalling (6.5), hence $\delta^2 - 4|\lambda_0|^2 = 0$ if and only if $|\lambda_0| = 1$ and $s = 0$. It is clear the integration is infinity in this case. \square

Lemma 6.4. *For any λ_0 , we have*

$$(6.8) \quad \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{i\theta}}{|e^{i\theta} - \lambda_0|^2 + s^2} d\theta = \begin{cases} \frac{\lambda_0}{\sqrt{\delta^2 - 4|\lambda_0|^2}} \frac{\delta - \sqrt{\delta^2 - 4|\lambda_0|^2}}{2|\lambda_0|^2}, & \text{if } \lambda_0 \neq 0 \\ 0, & \text{if } \lambda_0 = 0. \end{cases}$$

where $\delta = |\lambda_0|^2 + 1 + s^2$ is given by (6.2) and the integration is infinity if the denominator is zero. In particular, when $u = 0$.

$$(6.9) \quad \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{i\theta}}{|e^{i\theta} - \lambda_0|^2} d\theta = \begin{cases} \frac{\lambda_0}{1-|\lambda_0|^2} & \text{if } |\lambda_0| < 1 \\ \frac{\lambda_0}{|\lambda_0|^2(|\lambda_0|^2-1)} & \text{if } |\lambda_0| > 1 \\ \infty & \text{if } |\lambda_0| = 1. \end{cases}$$

Similarly,

$$(6.10) \quad \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{-i\theta}}{|e^{i\theta} - \lambda_0|^2 + s^2} d\theta = \begin{cases} \frac{\overline{\lambda_0}}{\sqrt{\delta^2 - 4|\lambda_0|^2}} \frac{\delta - \sqrt{\delta^2 - 4|\lambda_0|^2}}{2|\lambda_0|^2}, & \text{if } \lambda_0 \neq 0 \\ 0, & \text{if } \lambda_0 = 0. \end{cases}$$

where $\delta = |\lambda_0|^2 + 1 + s^2$ is given by (6.2) and the integration is infinity if the denominator is zero. In particular, when $u = 0$.

$$(6.11) \quad \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{-i\theta}}{|e^{i\theta} - \lambda_0|^2} d\theta = \begin{cases} \frac{\bar{\lambda}_0}{1-|\lambda_0|^2} & \text{if } |\lambda_0| < 1 \\ \frac{\lambda_0}{|\lambda_0|^2(|\lambda_0|^2-1)} & \text{if } |\lambda_0| > 1 \\ \infty & \text{if } |\lambda_0| = 1. \end{cases}$$

Proof. Recall that $|Z_1| < 1$ and $|Z_2| > 1$. Using the formula (6.3) and (6.4) for roots of the quadratic equation (6.1), we have

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{i\theta}}{|e^{i\theta} - \lambda_0|^2 + s^2} d\theta &= \frac{1}{2\pi i} \int_{|z|=1} \frac{z}{(z - \lambda_0)(1 - z\bar{\lambda}_0) + s^2 z} dz \\ &= \frac{1}{2\pi i} \int_{|z|=1} \frac{z}{-\bar{\lambda}_0(z - Z_1)(z - Z_2)} dz \\ &= \frac{-\lambda_0 Z_1}{|\lambda_0|^2(Z_1 - Z_2)} \\ &= \frac{\lambda_0}{\sqrt{\delta^2 - 4|\lambda_0|^2}} \frac{\delta - \sqrt{\delta^2 - 4|\lambda_0|^2}}{2|\lambda_0|^2} \end{aligned}$$

provided that $\delta^2 - 4|\lambda_0|^2 \neq 0$ and $\lambda_0 \neq 0$. If $\lambda_0 = 0$, the integration is clearly equal to zero. Note that $\delta^2 - 4|\lambda_0|^2 = 0$ if and only if $|\lambda_0| = 1$ and $s = 0$ (see (6.5)). The integration is infinity in this case. Hence (6.8) and (6.9) are proved.

Finally, when $s = 0$, we have $\delta^2 - 4|\lambda_0|^2 = (|\lambda_0|^2 - 1)^2$. Plugging this into (6.8), we obtain (6.7). By taking conjugation, we obtain (6.10) and (6.11). \square

We are now ready to describe the push-forward map $\Phi_{t,\gamma}$ between the Brown measures of $u + c_t$ and $u + g_t$.

Proposition 6.5. *Using notations in Notation 4.1 and Lemma 4.4, when $T = u$ is a Haar unitary operator, we have*

$$q_\varepsilon^{(0)}(\lambda) \Big|_{\varepsilon=s} = s\phi[(u - \lambda_0)^*(u - \lambda_0) + s^2]^{-1} = \frac{s}{\sqrt{\delta^2 - 4|\lambda_0|^2}}$$

by (6.6), where $\delta = |\lambda_0|^2 + 1 + s^2$ is given by (6.2). In particular, when $s = 0$,

$$(6.12) \quad \left(q_\varepsilon^{(0)}(\lambda) / \varepsilon \right) \Big|_{\varepsilon=0} = \begin{cases} \frac{1}{1-|\lambda_0|^2} & \text{if } 0 \leq |\lambda_0| < 1, \\ \frac{1}{|\lambda_0|^2-1} & \text{if } |\lambda_0| > 1, \\ \infty & \text{if } |\lambda_0| = 1 \end{cases}$$

by (6.7).

Proposition 6.6. *Using notations in Notation 4.1 and Lemma 4.4, when $T = u$, we have*

$$\begin{aligned} p_\lambda^{(0)}(s) &= \tau \left((\lambda - u)^* [(\lambda - u)^*(\lambda - u) + s^2]^{-1} \right) \\ &= \frac{\bar{\lambda}}{\sqrt{\delta^2 - 4|\lambda|^2}} - \frac{\bar{\lambda}}{\sqrt{\delta^2 - 4|\lambda|^2}} \frac{\delta - \sqrt{\delta^2 - 4|\lambda|^2}}{2|\lambda|^2} \end{aligned}$$

where $\delta = |\lambda|^2 + 1 + s^2$.

Proof. We have

$$\tau\left((\lambda - u)^*[(\lambda - u)^*(\lambda - u) + s^2]^{-1}\right) = \frac{1}{2\pi} \int_0^{2\pi} \frac{\bar{\lambda} - e^{-i\theta}}{|e^{i\theta} - \lambda|^2 + s^2} d\theta.$$

The result then follows from Lemma 6.3 and Lemma 6.4. \square

A direct application of Lemma 6.3 yields that

$$(6.13) \quad \Xi_t = \left\{ \lambda : \phi\left[\left((u - \lambda \mathbf{1})^*(u - \lambda \mathbf{1})\right)^{-1}\right] > \frac{1}{t} \right\} = \{\lambda : \lambda_1 < |\lambda| < \lambda_2\}$$

where

$$(6.14) \quad \lambda_1 = \sqrt{(1-t)_+}, \quad \lambda_2 = \sqrt{1+t}.$$

Proposition 6.7. *The support of the Brown measure of $u + c_t$ is the single ring*

$$\overline{\Xi}_t = \{\lambda : \lambda_1 \leq |\lambda| \leq \lambda_2\}.$$

Moreover, the Brown measure is absolutely continuous and the density of $u + c_t$ is strictly positive on its support.

Proof. It follows from (6.13) and Theorem 4.10. We can also calculate the inner radii r_1 and outer radii r_2 of the support of the Brown measure of $u + c_t$ respectively. By [24], we know that $r_2^2 = \phi((u + c_t)^2) = 1 + t$. Hence, $r_2 = \lambda_2$. By [45, Section 7], we have $r_1 = \sqrt{(1-t)_+} = \lambda_1$. Therefore the support of the Brown measure of $u + c_t$ is the set $\overline{\Xi}_t = \{\lambda : \lambda_1 \leq |\lambda| \leq \lambda_2\}$. \square

We can now describe the subordination function $w(0; \lambda, t)$ in this context. For any $\lambda \in \overline{\Xi}_t$, following Notation 4.1, $w(0; \lambda, t)$ is determined by the condition

$$\phi\left[\left((u - \lambda \mathbf{1})^*(u - \lambda \mathbf{1}) + w(0; \lambda, t)^2\right)^{-1}\right] = \frac{1}{t}.$$

By Proposition 6.5, this condition is equivalent to

$$\delta^2 - 4|\lambda|^2 = (|\lambda|^2 + 1 + w(0; \lambda, t)^2)^2 - 4|\lambda|^2 = t^2.$$

It follows that

$$(6.15) \quad w(0; \lambda, t) = \sqrt{4|\lambda|^2 + t^2} - (|\lambda|^2 + 1), \quad \lambda_1 < |\lambda| < \lambda_2.$$

We now describe the map $\Phi_{t,\gamma}$ more closely. For $\lambda \in \overline{\Xi}_t$, we set $s = |\lambda|^2 - 1$, where $\lambda_1^2 - 1 < s < \lambda_2^2 - 1$. Let $\delta = |\lambda|^2 + 1 + w(0; \lambda, t)^2$. Since $\delta^2 - 4|\lambda|^2 = t^2$, we now have

$$\frac{\delta - \sqrt{\delta^2 - 4|\lambda|^2}}{2|\lambda|^2} = \frac{\delta - t}{2|\lambda|^2} = \frac{\sqrt{4(1+s) + t^2} - t}{2(1+s)}.$$

Therefore, for $\lambda_1^2 - 1 < s < \lambda_2^2 - 1$ and λ such that $|\lambda|^2 = s + 1$, by Proposition 6.6, we have,

$$(6.16) \quad \begin{aligned} \Phi_{t,\gamma}(\lambda) &= \lambda + \gamma p_\lambda^{(0)}(w(0; \lambda, t)) \\ &= \lambda + \gamma \left(\frac{\bar{\lambda}}{\sqrt{\delta^2 - 4|\lambda|^2}} - \frac{\bar{\lambda}}{\sqrt{\delta^2 - 4|\lambda|^2}} \frac{\delta - \sqrt{\delta^2 - 4|\lambda|^2}}{2|\lambda|^2} \right) \end{aligned}$$

where $\delta = |\lambda|^2 + 1 + w(0; \lambda, t)^2$.

We first look at the case when $\gamma = t$. Since $\delta^2 - 4|\lambda|^2 = t^2$, for $\lambda = re^{i\theta}$ where $r = \sqrt{s+1}$, $\Phi_{t,t}$ as in (6.16) can be further simplified as

$$(6.17) \quad \begin{aligned} \Phi_{t,t}(re^{i\theta}) &= re^{i\theta} + tre^{-i\theta} \left(\frac{1}{t} + \frac{1}{2r^2} - \frac{\sqrt{4r^2+t^2}}{2r^2t} \right) \\ &= re^{i\theta} + re^{-i\theta} \left(1 - \frac{\sqrt{4(1+s)+t^2-t}}{2(1+s)} \right) \\ (6.18) \quad &= a(s) \cos(\theta) + ib(s) \sin(\theta) \end{aligned}$$

where, $\lambda_1^2 - 1 < s < \lambda_2^2 - 1$, and

$$(6.19) \quad \begin{cases} a(s) = \left(2\sqrt{1+s} - \frac{\sqrt{4(1+s)+t^2-t}}{2\sqrt{1+s}} \right); \\ b(s) = \frac{\sqrt{4(1+s)+t^2-t}}{2\sqrt{1+s}}. \end{cases}$$

In particular, if $\lambda_1^2 - 1 = s$, we have $1+s = \lambda_1^2 = (1-t)_+$ and $4(1+s) + t^2 = (4(1-t)_+ + t^2)$. Then

$$\begin{cases} a(s) = a^- = \sqrt{(1-t)_+} \\ b(s) = b^- = \sqrt{(1-t)_+} \end{cases} \quad \text{for } s = ((1-t)_+) - 1.$$

Similarly, when $\lambda_2^2 - 1 = s$, we have $s = t$, we have

$$\begin{cases} a(s) = a^+ = \frac{2t+1}{\sqrt{t+1}} \\ b(s) = b^+ = \frac{1}{\sqrt{1+t}} \end{cases} \quad \text{for } s = t.$$

Proposition 6.8. *For $\lambda_1^2 - 1 < s < \lambda_2^2 - 1$, the circle $C_s := \{\lambda : |\lambda|^2 = 1+s\}$ is mapped to the ellipse centered at the origin with semi-axes $a(t)$ and $b(t)$ as in (6.19). Moreover, the function $s \mapsto a(s)$ is a strictly increasing function from $(\lambda_1^2 - 1, \lambda_2^2 - 1)$ onto (a^-, a^+) ; and the the function $s \mapsto b(s)$ is a strictly increasing function from $(\lambda_1^2 - 1, \lambda_2^2 - 1)$ onto (b^-, b^+) .*

Proof. We have

$$a'(s) = \frac{\frac{t^2}{\sqrt{4s+t^2+4}} + 4s - t + 4}{4(s+1)^{3/2}}, \quad \text{and} \quad b'(s) = \frac{t(\sqrt{4s+t^2+4} - t)}{4(s+1)^{3/2}\sqrt{4s+t^2+4}}.$$

Hence $b'(s) > 0$. To show $a'(s) > 0$, we set

$$f(s) = \frac{t^2}{\sqrt{4s+t^2+4}} + 4s - t + 4$$

and calculate its derivatives

$$f'(s) = 4 - \frac{2t^2}{(4s+t^2+4)^{3/2}}$$

and

$$f''(s) = \frac{12t^2}{(4s+t^2+4)^{5/2}} > 0.$$

Hence $f'(s) > f'(\lambda_1^2 - 1) = f'((1-t)_+) > 0$. Consequently, f is increasing. We then check that

$$f((1-t)_+) > 0.$$

It follows that $a'(s) > 0$ for $\lambda_1^2 - 1 < s < \lambda_2^2 - 1$. \square

Theorem 6.9. *The Brown measure of $u + c_t$ is given by*

$$(6.20) \quad \mu_{u+c_t}(\{\lambda : |\lambda| \leq r\}) = r^2 \left(\frac{1}{t} + \frac{1}{2r^2} - \frac{\sqrt{4r^2 + t^2}}{2r^2 t} \right),$$

where $\sqrt{(1-t)_+} \leq r \leq \sqrt{1+t}$.

Proof. It is shown [45, Corollary 4.4] that, for $\sqrt{(1-t)_+} \leq |\lambda| \leq \sqrt{1+t}$,

$$\begin{aligned} p_\lambda^{c_t(t)}(0) &= \frac{\partial S}{\partial \lambda}(u + c_t, \lambda, 0) \\ &= \frac{\partial}{\partial \lambda} \log \Delta((u + c_t - \lambda \mathbf{1})^*(u + c_t - \lambda \mathbf{1})) \\ &= \frac{\mu_{u+c_t}(\{\lambda : |\lambda| \leq r\})}{\lambda}, \end{aligned}$$

which yields $\Phi_{t,t}(\lambda) = \lambda + \frac{t}{\lambda} \mu_{u+c_t}(\{\lambda : |\lambda| \leq r\})$. Hence, the result follows by comparing this general formula with the explicit formula of $\Phi_{t,t}(\lambda)$ as in (6.18). \square

Theorem 6.10. *The support of the Brown measure of $u + g_t$ is the deformed single ring where the inner boundary is the circle centered at the origin with radius $\sqrt{(1-t)_+}$ and the outer boundary is the ellipse centered at the origin with semi-axes $\frac{2t+1}{\sqrt{t+1}}$ and $\frac{1}{\sqrt{1+t}}$. The Brown measure is absolutely continuous and its density is strictly positive in the support.*

Moreover, the Brown measure of $u + g_t$ is the push-forward map of the Brown measure of $u + c_t$ under the map $\Phi_{t,t}$ defined as

$$\Phi_{t,t}(re^{i\theta}) = a(s) \cos(\theta) + ib(s) \sin(\theta), \quad r = \sqrt{s+1},$$

where $((1-t)_+ - 1 < s < t$ and $a(s), b(s)$ are given by (6.19).

Proof. It follows from Proposition 6.8 that the map $\Phi_{t,t}$ is one-to-one and onto from the single ring $\{\lambda : \lambda_1 < |\lambda| < \lambda_2\}$, the support of the Brown measure of $u + c_t$, to the deformed ring. Hence, Theorem 4.16 applies and the push-forward map is given by (6.17) and (6.19). Since the Brown measure of $u + c_t$ is strictly positive on its support, the result then follows. \square

We now fix $\gamma = |\gamma|e^{2i\alpha}$ such that $|\gamma| \leq t$ and $\lambda_1^2 - 1 < s < \lambda_2^2 - 1$ and $r^2 = |\lambda|^2 = s+1$, then similar to the case $\gamma = t$ as in (6.16), we have

$$(6.21) \quad \begin{aligned} \Phi_{t,\gamma}(re^{i\theta}) &= re^{i\theta} + re^{i(2\alpha-\theta)} \frac{|\gamma|}{t} \left(1 - \frac{\sqrt{4(1+s)+t^2}-t}{2(1+s)} \right) \\ &= e^{i\alpha} \left[\sqrt{s+1} e^{i(\theta-\alpha)} + \frac{|\gamma|}{t} e^{i(\alpha-\theta)} \left(\sqrt{s+1} - \frac{\sqrt{4(1+s)+t^2}-t}{2\sqrt{1+s}} \right) \right], \end{aligned}$$

which is an ellipse twisted by angle α with semi-axes

$$(6.22) \quad \begin{cases} f(s) &= \frac{t-|\gamma|}{t} \sqrt{s+1} + \frac{|\gamma|}{t} a(s), \\ g(s) &= \frac{t-|\gamma|}{t} \sqrt{s+1} + \frac{|\gamma|}{t} b(s), \end{cases}$$

where $a(s), b(s)$ are given by (6.19). It follows by Proposition 6.8 that both f and g are increasing function of s in the interval $(\lambda_1^2 - 1, \lambda_2^2 - 1)$.

Then we deduce the following result in the same was as Theorem 6.10. We leave details for interested readers.

Theorem 6.11. *The support of the Brown measure of $u + g_{t,\gamma}$ is the deformed single ring where the inner boundary is the circle centered at the origin with radius $\sqrt{(1-t)_+}$, and the outer boundary is the ellipse twisted by angle α , centered at the origin, with semi-axes $\frac{t+|\gamma|+1}{\sqrt{t+1}}$ and $\frac{t-|\gamma|+1}{\sqrt{t+1}}$.*

Moreover, the Brown measure of $u + g_{t,\gamma}$ is the push-forward map of the Brown measure of $u + c_t$ under the map $\Phi_{t,\gamma}$ defined as

$$\Phi_{t,\gamma}(re^{i\theta}) = e^{i\alpha} (f(s) \cos(\theta - \alpha) + ig(s) \sin(\theta - \alpha)), \quad r = \sqrt{s+1}$$

where $((1-t)_+ - 1 < s < t)$, $f(s), g(s)$ are defined by (6.22).

Remark 6.12. It is interesting to observe that the outer boundary of $u + g_{t,\gamma}$ is the same as the outer boundary of the Brown measure of $c + g_{t,\gamma}$ where c is the standard circular operator with variance one. The outer boundary for $c + g_{t,\gamma}$ can be checked by using Corollary 5.15. This observation is a consequence of a general result in [9].

The Brown measure of $u + g_{t,\gamma}$ is a rotation of the Brown measure of $u + y$ where y is an elliptic operator by the angle α with $e^{2i\alpha} = \gamma/|\gamma|$. This is can be seen directly by the fact that $g_{t,\gamma}$ is a twisted elliptic operator of the form $e^{i\alpha}(g_1 + ig_2)$ where g_1, g_2 are free independent semicircular elements with variances t_1, t_2 respectively so that $t_1 + t_2 = t$ and $t_1 - t_2 = |\gamma|$. Then $u + g_{t,\gamma}$ has the same distribution as $e^{i\alpha}(u + g_1 + ig_2)$.

Acknowledgment. The author would like to express his deep gratitude to his former advisor, Hari Bercovici, for many valuable discussions during the course of the investigation and for careful reading of earlier versions of this paper. Special thanks to Serban Belinschi for his interest in this work and many helpful comments on Hermitian reduction method and subordination functions. The author also wants to thank Brian Hall, Ching-Wei Ho, Dang-Zheng Liu, Weihua Liu, Alexandru Nica, Zhuang Niu, and Zhi Yin for discussions on the topics of Brown measures. He also wants to thank Edward Saff for communications on potential theory and Piotr Śniady for helpful comments on the work [39].

REFERENCES

- [1] AAGAARD, L., AND HAAGERUP, U. Moment formulas for the quasi-nilpotent DT-operator. *Internat. J. Math.* 15, 6 (2004), 581–628.
- [2] BELINSCHI, S. T. The Lebesgue decomposition of the free additive convolution of two probability distributions. *Probab. Theory Related Fields* 142, 1-2 (2008), 125–150.
- [3] BELINSCHI, S. T., AND BERCOVICI, H. A new approach to subordination results in free probability. *J. Anal. Math.* 101 (2007), 357–365.
- [4] BELINSCHI, S. T., MAI, T., AND SPEICHER, R. Analytic subordination theory of operator-valued free additive convolution and the solution of a general random matrix problem. *J. Reine Angew. Math.* 732 (2017), 21–53.
- [5] BELINSCHI, S. T., ŚNIADY, P., AND SPEICHER, R. Eigenvalues of non-Hermitian random matrices and Brown measure of non-normal operators: Hermitian reduction and linearization method. *Linear Algebra and its Applications* 537 (2018), 48 – 83.
- [6] BERCOVICI, H. *Free convolution*, Free probability and operator algebras - Chapter V, *Münster Lectures in Mathematics*. European Mathematical Society (EMS), Zürich, 2016. Lecture notes from the masterclass held in Münster.
- [7] BERCOVICI, H., AND VOICULESCU, D. Free convolution of measures with unbounded support. *Indiana Univ. Math. J.* 42, 3 (1993), 733–773.
- [8] BERCOVICI, H., AND VOICULESCU, D. Regularity questions for free convolution. In *Nonselfadjoint operator algebras, operator theory, and related topics*, vol. 104 of *Oper. Theory Adv. Appl.* Birkhäuser, Basel, 1998, pp. 37–47.
- [9] BERCOVICI, H., AND ZHONG, P. preprint in preparation.

- [10] BIANE, P. On the free convolution with a semi-circular distribution. *Indiana Univ. Math. J.* 46, 3 (1997), 705–718.
- [11] BIANE, P. Processes with free increments. *Math. Z.* 227, 1 (1998), 143–174.
- [12] BIANE, P., AND LEHNER, F. Computation of some examples of Brown’s spectral measure in free probability. *Colloq. Math.* 90, 2 (2001), 181–211.
- [13] BORDENAVE, C., AND CHAFAÏ, D. Around the circular law. *Probab. Surv.* 9 (2012), 1–89.
- [14] BROWN, L. G. Lidskii’s theorem in the type II case. In *Geometric methods in operator algebras (Kyoto, 1983)*, vol. 123 of *Pitman Res. Notes Math. Ser.* Longman Sci. Tech., Harlow, 1986, pp. 1–35.
- [15] DYKEMA, K., AND HAAGERUP, U. DT-operators and decomposability of Voiculescu’s circular operator. *Amer. J. Math.* 126, 1 (2004), 121–189.
- [16] DYKEMA, K., AND HAAGERUP, U. Invariant subspaces of the quasinilpotent DT-operator. *J. Funct. Anal.* 209, 2 (2004), 332–366.
- [17] FEINBERG, J., SCALETTAR, R., AND ZEE, A. “Single ring theorem” and the disk-annulus phase transition. *J. Math. Phys.* 42, 12 (2001), 5718–5740.
- [18] FEINBERG, J., AND ZEE, A. Non-Gaussian non-Hermitian random matrix theory: phase transition and addition formalism. *Nuclear Phys. B* 501, 3 (1997), 643–669.
- [19] FEINBERG, J., AND ZEE, A. Non-Hermitian random matrix theory: method of Hermitian reduction. *Nuclear Phys. B* 504, 3 (1997), 579–608.
- [20] FUGLEDE, B., AND KADISON, R. V. Determinant theory in finite factors. *Ann. of Math. (2)* 55 (1952), 520–530.
- [21] GINIBRE, J. Statistical Ensembles of Complex, Quaternion and Real Matrices. *J. Math. Phys.* 6 (1965), 440–449.
- [22] GUIONNET, A., KRISHNAPUR, M., AND ZEITOUNI, O. The single ring theorem. *Ann. of Math. (2)* 174, 2 (2011), 1189–1217.
- [23] HAAGERUP, U., KEMP, T., AND SPEICHER, R. Resolvents of R-diagonal operators. *Trans. Amer. Math. Soc.* 362, 11 (2010), 6029–6064.
- [24] HAAGERUP, U., AND LARSEN, F. Brown’s spectral distribution measure for R-diagonal elements in finite von Neumann algebras. *J. Funct. Anal.* 176, 2 (2000), 331 – 367.
- [25] HAAGERUP, U., AND SCHULTZ, H. Brown measures of unbounded operators affiliated with a finite von Neumann algebra. *Math. Scand.* 100, 2 (2007), 209–263.
- [26] HAAGERUP, U., AND SCHULTZ, H. Invariant subspaces for operators in a general II_1 -factor. *Publ. Math. Inst. Hautes Études Sci.*, 109 (2009), 19–111.
- [27] HAAGERUP, U., AND THORBJØRNSEN, S. A new application of random matrices: $\text{Ext}(C_{\text{red}}^*(F_2))$ is not a group. *Ann. of Math. (2)* 162, 2 (2005), 711–775.
- [28] HALL, B. C., AND HO, C.-W. The Brown measure of the sum of a self-adjoint element and an imaginary multiple of a semicircular element, 2020.
- [29] HALL, B. C., AND HO, C.-W. The Brown measure of a family of free multiplicative brownian motions, 2021.
- [30] HALL, B. C., AND KEMP, T. Brown measure support and the free multiplicative Brownian motion. *Adv. Math.* 355 (2019), 106771, 36.
- [31] HO, C.-W. The Brown measure of the sum of a self-adjoint element and an elliptic element, 2020.
- [32] HO, C.-W., AND ZHONG, P. Brown measures of free circular and multiplicative brownian motions with self-adjoint and unitary initial conditions. *arXiv:1908.08150*, to appear in *J. Eur. Math. Soc.* (2019).
- [33] JANIK, R. A., NOWAK, M. A., PAPP, G., AND ZAHED, I. Non-Hermitian random matrix models. *Nuclear Phys. B* 501, 3 (1997), 603–642.
- [34] JAROSZ, A., AND NOWAK, M. A. A novel approach to non-hermitian random matrix models. *preprint arXiv:math-ph/0402057*.
- [35] JAROSZ, A., AND NOWAK, M. A. Random Hermitian versus random non-Hermitian operators—unexpected links. *J. Phys. A* 39, 32 (2006), 10107–10122.
- [36] MINGO, J. A., AND SPEICHER, R. *Free probability and random matrices*, vol. 35 of *Fields Institute Monographs*. Springer, New York; Fields Institute for Research in Mathematical Sciences, Toronto, ON, 2017.
- [37] NICA, A., SHLYAKHTENKO, D., AND SPEICHER, R. R-cyclic families of matrices in free probability. *J. Funct. Anal.* 188, 1 (2002), 227–271.
- [38] SAFF, E. B., AND TOTIK, V. *Logarithmic potentials with external fields*, vol. 316 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1997. Appendix B by Thomas Bloom.
- [39] ŚNIADY, P. Random regularization of Brown spectral measure. *J. Funct. Anal.* 193, 2 (2002), 291–313.

- [40] SPEICHER, R. Combinatorial theory of the free product with amalgamation and operator-valued free probability theory. *Mem. Amer. Math. Soc.* 132, 627 (1998), x+88.
- [41] TAO, T., AND VU, V. Random matrices: universality of ESDs and the circular law. *Ann. Probab.* 38, 5 (2010), 2023–2065. With an appendix by Manjunath Krishnapur.
- [42] VOICULESCU, D. Limit laws for random matrices and free products. *Invent. Math.* 104, 1 (1991), 201–220.
- [43] VOICULESCU, D. Operations on certain non-commutative operator-valued random variables. No. 232. 1995, pp. 243–275. Recent advances in operator algebras (Orléans, 1992).
- [44] VOICULESCU, D. The coalgebra of the free difference quotient and free probability. *Int. Math. Res. Not.* 2000, 2 (01 2000), 79–106.
- [45] ZHONG, P. Brown measure of R-diagonal operators, revisited. *preprint*.

PING ZHONG, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF WYOMING, LARAMIE, WY 82070, PZHONG@UWYO.EDU