

PERTURBATION FOR GROUP INVERSE IN A BANACH ALGEBRA

DAYONG LIU, TUGCE PEKACAR CALCI, HANDAN KOSE,
AND HUANYIN CHEN

ABSTRACT. We present new additive results for the group inverse in a Banach algebra under certain perturbations. The upper bound of $\|(a+b)^\# - a^d\|$ is thereby given. These extend the main results in [X. Liu, Y. Qin and H. Wei, Perturbation bound of the group inverse and the generalized Schur complement in Banach algebra, *Abstr. Appl. Anal.*, 2012, 22 pages. <https://doi.org/10.1155/2012/629178>].

1. INTRODUCTION

Let \mathcal{A} be a complex Banach algebra with an identity 1 and $a \in \mathcal{A}$. The Drazin inverse of a is the unique x in \mathcal{A} satisfying

$$ax = xa, \quad xax = x, \quad a^{k+1}x = a^k$$

for some nonnegative integers k . We denote x by a^D . The least such integer k is called the index of a , denoted by $\text{ind}(a) = k$. If $\text{ind}(a) = 1$, we say a is group invertible, that is, $ax = xa, xax = x, axa = a$. The element x will be denoted by $a^\#$. An element x is a generalized Drazin inverse of a in \mathcal{A} if

$$ax = xa, \quad xax = x, \quad a - a^2x \in \mathcal{A}^{\text{qnil}}.$$

The generalized Drazin inverse is unique if it exists. We denote x by a^d . Here, $\mathcal{A}^{\text{qnil}}$ is the sets of all quasinilpotent elements in \mathcal{A} , i.e., $a \in \mathcal{A}^{\text{qnil}} \Leftrightarrow \lim_{n \rightarrow \infty} \|a^n\|^{\frac{1}{n}} = 0$.

2010 *Mathematics Subject Classification.* 15A09, 47L10, 32A65.

Key words and phrases. group inverse; generalized Drazin inverse; additive property; perturbation; Banach algebra.

Corresponding author: Huanyin Chen, huanyinchen@aliyun.com.

The preceding generalized inverses play important roles in matrix and operator theory (see [2, 3, 7, 10, 12, 13]). In [11], Yang and Liu studied the Drazin inverse of the sum of complex matrices P and Q under the condition $QP^2 = 0$ and $QPQ = 0$. The generalized Drazin inverse of $a + b$ in a Banach algebra was studied in [5, Theorem 2.1] under the condition $ba^2 = 0$ and $b^2 = 0$. In [8], Liu et al. investigated the Drazin inverse $(P + Q)^D$ of two complex matrices P and Q under the conditions $P^2Q = PQP$ and $Q^2P = QPQ$. In [13], Zou et al. presented the generalized Drazin inverse of $a + b$ in a Banach algebra under $a^2b = aba$ and $b^2a = bab$.

Liu et al. investigated the perturbation of group inverse under the conditions $\|a^d b a a^d\| < 1, a^\pi b a = 0$ (see [6, Theorem 2.8]). They further studied the group invertibility under the condition $\|a^d b a a^d\| < 1, a^\pi b a = a b a^\pi$ (see [6, Theorem 2.12]). These inspire us to explore the perturbation for group invertibility in a more general setting. The preceding results of X. Liu et al. are thereby generalzied to wider cases.

In Section 2, we investigate the group invertibility of $a + b$ in a Banach algebra under the condition $a^\pi b a^2 = 0, a^\pi b a b = 0$. The upper bound of $\|(a + b)^\# - a^d\|$ is thereby given. In Section 3, we are concerned with the additive property for group inverse under the condition $a^2 b a^\pi = a^\pi a b a, a^\pi b^2 a = b a b a^\pi$ and obtain the bound of $\|(a + b)^\# - a^d\|$.

Throughout the paper, all Banach algebras are complex with an identity. We use \mathcal{A}^d to denote the set of all g-Drazin invertible elements in \mathcal{A} . We use a^π to stand for the spectral idempotent $1 - a a^d$ of an element $a \in \mathcal{A}^d$.

2. ORTHOGONAL CONDITIONS

In this section we establish additive property of group inverse in a Banach algebra under certain orthogonal perturbation conditions. We begin with

Lemma 2.1. *Let \mathcal{A} be a Banach algebra, $a, b \in \mathcal{A}^d$,*

$$x = \begin{pmatrix} a & 0 \\ c & b \end{pmatrix}_p \text{ or } \begin{pmatrix} b & c \\ 0 & a \end{pmatrix}_p.$$

Then

$$x^d = \begin{pmatrix} a^d & 0 \\ z & b^d \end{pmatrix}_p, \text{ or } \begin{pmatrix} b^d & z \\ 0 & a^d \end{pmatrix}_p,$$

where

$$z = (b^d)^2 \left(\sum_{i=0}^{\infty} (b^d)^i c a^i \right) a^\pi + b^\pi \left(\sum_{i=0}^{\infty} b^i c (a^d)^i \right) (a^d)^2 - b^d c a^d.$$

Proof. See [12, Lemma 1.1]. \square

Lemma 2.2. *Let e be an idempotent in a Banach algebra \mathcal{A} , and let $a, ea \in \mathcal{A}^d$. If $ea(1-e) = 0$, then $a(1-e) \in \mathcal{A}^d$, and*

$$(ea)^d = ea^d, (a(1-e))^d = a^d(1-e).$$

Proof. See [12, Lemma 2.2]. \square

We are ready to prove:

Theorem 2.3. *Let \mathcal{A} be a Banach algebra, and let $a, b, a^\pi b \in \mathcal{A}^d$. If*

$$\|a^d b a a^d\| < 1, a^\pi b a^2 = 0, a^\pi b a b = 0,$$

then $a + b$ has group inverse if and only if $a^\pi(a + b)$ has group inverse. In this case,

$$\begin{aligned} & \|(a + b)^\# - a^d\| \\ & \leq \left(\frac{\|a^d\|}{1 - \|a^d b a a^d\|} \right)^2 \|b a^\pi\| \left(\|a^\pi\| + \sum_{n=0}^{\infty} \|a^\pi a\|^{n+1} \|a^\pi b^d\|^{n+1} \right. \\ & \quad \left. + \sum_{n=0}^{\infty} \|a^\pi a\|^{n+2} \|a^\pi b^d\|^{n+2} + \|a^\pi b^d\| (\|a^\pi a\| + \|a^\pi b\|) \right) \\ & \quad + \left(1 + \frac{\|a^d b\|}{1 - \|a^d b a a^d\|} \right) \left(\|a^d\| + \sum_{n=0}^{\infty} \|a^\pi a\|^n \|a^\pi b^d\|^{n+1} \right. \\ & \quad \left. + \sum_{n=0}^{\infty} \|a^\pi a\|^{n+1} \|a^\pi b^d\|^{n+2} \right). \end{aligned}$$

Proof. Let $p = a a^d$. Then we have

$$a = \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix}_p, b = \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix}_p,$$

where $a_1 \in (p\mathcal{A}p)^{-1}$, $a_2 \in ((1-p)\mathcal{A}(1-p))^{\text{qmil}}$.

Furthermore,

$$a^d = \begin{pmatrix} a_1^{-1} & 0 \\ 0 & 0 \end{pmatrix}_p \quad \text{and} \quad a^\pi = \begin{pmatrix} 0 & 0 \\ 0 & a^\pi \end{pmatrix}_p.$$

Since $a^\pi b a^2 = 0$, we see that $a^\pi b a a^d = a^\pi b a^2 (a^d)^2 = 0$. This implies that $b_3 = 0$, and so

$$a + b = \begin{pmatrix} a_1 + b_1 & b_2 \\ 0 & a_2 + b_4 \end{pmatrix}_p.$$

Since $\|a^d b a a^d\| < 1$, we see that $\|a_1^{-1} b_1\| < 1$, and so $a_1 + b_1 = a_1(1 + a_1^{-1} b_1) \in (p\mathcal{A}p)^{-1}$. One easily checks that $a_2 = a - a^2 a^d = a a^\pi \in ((1-p)\mathcal{A}(1-p))^{\text{qnil}}$. On the other hand, we have $b_4 = a^\pi b a^\pi = a^\pi b - a^\pi b a^2 (a^d)^2 = a^\pi b \in \mathcal{A}^d$, and so $b_4 \in ((1-p)\mathcal{A}(1-p))^d$. Moreover, we have

$$\begin{aligned} b_4 a_2^2 &= a^\pi b (1 - a a^d) a^2 a^\pi = a^\pi b a^2 a^\pi = 0, \\ b_4 a_2 b_4 &= a^\pi b (1 - a a^d) a b a^\pi = a^\pi b a b a^\pi = 0. \end{aligned}$$

Noting that $a_2^d = 0$, by virtue of [11, Theorem 2.1], we have

$$\begin{aligned} (a_2 + b_4)^d &= \sum_{n=0}^{\infty} a_2^n (b_4^d)^{n+1} + \sum_{n=0}^{\infty} a_2^n (b_4^d)^{n+2} a_2 \\ &= \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a. \end{aligned}$$

Therefore, by [1, Theorem 2.3],

$$(a + b)^d = \begin{pmatrix} (a_1 + b_1)^{-1} & z \\ 0 & (a_2 + b_4)^d \end{pmatrix}_p,$$

where

$$z = \sum_{n=0}^{\infty} (a_1 + b_1)^{-n-2} b_2 (a_2 + b_4)^n (a_2 + b_4)^\pi - (a_1 + b_1)^{-1} b_2 (a_2 + b_4)^d.$$

We check that

$$a_2 + b_4 = a^\pi a + a^\pi b = a^\pi (a + b).$$

If $a + b \in \mathcal{A}^\#$, we easily see that $a_2 + b_4 \in \mathcal{A}^\#$, and so $a^\pi (a + b) \in \mathcal{A}^\#$. We now assume that $a^\pi (a + b) \in \mathcal{A}^\#$. Then $(a_2 + b_4)^d = (a_2 + b_4)^\#$.

Obviously,

$$\begin{aligned}(a+b)^d(a+b) &= (a+b)(a+b)^d, \\ (a+b)^d &= (a+b)^d(a+b)(a+b)^d.\end{aligned}$$

It is easy to verify that

$$\begin{aligned}& ((a_1 + b_1)z + b_2(a_2 + b_4)^\#)(a_2 + b_4) \\ &= ((a_1 + b_1)^{-1}b_2(a_2 + b_4)^\pi - b_2(a_2 + b_4)^\#)(a_2 + b_4) \\ &\quad + b_2(a_2 + b_4)^\#(a_2 + b_4) \\ &= 0.\end{aligned}$$

Then we have

$$\begin{aligned}& (a+b)(a+b)^d(a+b) \\ &= \begin{pmatrix} a_1 + b_1 & b_2 \\ 0 & a_2 + b_4 \end{pmatrix} \begin{pmatrix} (a_1 + b_1)^{-1} & z \\ 0 & (a_2 + b_4)^d \end{pmatrix} \\ &\quad \begin{pmatrix} a_1 + b_1 & b_2 \\ 0 & a_2 + b_4 \end{pmatrix} \\ &= a + b.\end{aligned}$$

Therefore $(a+b)^\# = (a+b)^d$, i.e., $a+b$ has group inverse. That is,

$$(a_2 + b_4)^\# = \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a.$$

Since $\|a_1^{-1}b_1\| < 1$, we easily check that

$$\begin{aligned}(a_1 + b_1)^{-1} &= (1 + a_1^{-1}b_1)^{-1}a_1^{-1} \\ &= \sum_{n=0}^{\infty} (-1)^n (a_1^{-1}b_1)^n a_1^{-1} \\ &= \sum_{n=0}^{\infty} (-1)^n (a^d b a a^d)^n a^d \\ &= \sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d.\end{aligned}$$

On the other hand,

$$\begin{aligned}
(a_1 + b_1)^{-1} &= a_1^{-1}(1 + b_1 a_1^{-1})^{-1} \\
&= a_1^{-1} \sum_{n=0}^{\infty} (-1)^n (b a_1^{-1})^n \\
&= a^d \sum_{n=0}^{\infty} (-1)^n (b a^d)^n.
\end{aligned}$$

Also we have

$$\begin{aligned}
z &= (a_1 + b_1)^{-2} b_2 (a_2 + b_4)^\pi - (a_1 + b_1)^{-1} b_2 (a_2 + b_2)^\# \\
&= \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 \cdot a a^d b a^\pi \\
&\quad \cdot \left(a^\pi - a^\pi (a + b) \left(\sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a \right) \right) \\
&\quad - \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right) \cdot a a^d b a^\pi \\
&\quad \cdot \left(\sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a \right) \\
&= \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 b a^\pi \left(a^\pi - \sum_{n=0}^{\infty} (a^\pi a)^{n+1} (a^\pi b^d)^{n+1} \right. \\
&\quad \left. - \sum_{n=0}^{\infty} (a^\pi a)^{n+1} (a^\pi b^d)^{n+2} a^\pi a - a^\pi b^d a^\pi (a + b) \right) \\
&\quad - \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^{n+1} \right) \\
&\quad \cdot \left(\sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a \right).
\end{aligned}$$

Hence

$$(a + b)^\# = (a_1 + b_1)^{-1} + z + (a_2 + b_4)^\#.$$

We compute that

$$(a + b)^\# - a^d = \sum_{n=1}^{\infty} (-1)^n (a^d b)^n a^d$$

$$\begin{aligned}
 & + \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 ba^\pi \left(a^\pi - \sum_{n=0}^{\infty} (a^\pi a)^{n+1} (a^\pi b^d)^{n+1} \right. \\
 & \left. - \sum_{n=0}^{\infty} (a^\pi a)^{n+1} (a^\pi b^d)^{n+2} a^\pi a - a^\pi b^d a^\pi (a+b) \right) \\
 & - \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^{n+1} \right) \left(\sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a \right) \\
 & + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a \\
 = & \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 ba^\pi \left(a^\pi - \sum_{n=0}^{\infty} (a^\pi a)^{n+1} (a^\pi b^d)^{n+1} \right. \\
 & \left. - \sum_{n=0}^{\infty} (a^\pi a)^{n+1} (a^\pi b^d)^{n+2} a^\pi a - a^\pi b^d a^\pi (a+b) \right) \\
 & + \sum_{n=0}^{\infty} (-1)^n (a^d b)^n \left[a^d + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+1} + \sum_{n=0}^{\infty} (a^\pi a)^n (a^\pi b^d)^{n+2} a^\pi a \right].
 \end{aligned}$$

Clearly, we have

$$\begin{aligned}
 & \left\| \sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right\| \leq \sum_{n=1}^{\infty} \|a^d b a a^d\|^{n-1} \|a^d\| = \frac{\|a^d\|}{1 - \|a^d b a a^d\|}, \\
 & \left\| \sum_{n=0}^{\infty} (-1)^n (a^d b)^n \right\| \leq 1 + \left\| \sum_{n=0}^{\infty} (-1)^n (a^d b a a^d)^n \right\| \|a^d b\| = 1 + \frac{\|a^d b\|}{1 - \|a^d b a a^d\|}.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 & \|(a+b)^\# - a^d\| \\
 & \leq \left(\frac{\|a^d\|}{1 - \|a^d b a a^d\|} \right)^2 \|ba^\pi\| \left(\|a^\pi\| + \sum_{n=0}^{\infty} \|a^\pi a\|^{n+1} \|a^\pi b^d\|^{n+1} \right. \\
 & \left. + \sum_{n=0}^{\infty} \|a^\pi a\|^{n+2} \|a^\pi b^d\|^{n+2} + \|a^\pi b^d\| (\|a^\pi a\| + \|a^\pi b\|) \right) \\
 & + \left(1 + \frac{\|a^d b\|}{1 - \|a^d b a a^d\|} \right) \left(\|a^d\| + \sum_{n=0}^{\infty} \|a^\pi a\|^n \|a^\pi b^d\|^{n+1} \right)
 \end{aligned}$$

$$+ \sum_{n=0}^{\infty} \|a^\pi a\|^{n+1} \|a^\pi b^d\|^{n+2} \Big). \quad \square$$

Corollary 2.4. *Let \mathcal{A} be a Banach algebra, and let $a, b, a^\pi b \in \mathcal{A}^d$. If*

$$\begin{aligned} a^\pi b a^2 &= 0, a^\pi b a b = 0, \\ \max\{\|a^d b a a^d\|, \|a^\pi a\| \|a^\pi b^d\|\} &< 1, \end{aligned}$$

then $a + b$ has group inverse if and only if $a^\pi(a + b)$ has group inverse. In this case,

$$\begin{aligned} \|(a + b)^\# - a^d\| &\leq \left(\frac{\|a^d\|}{1 - \|a^d b a a^d\|} \right)^2 \|b a^\pi\| \left(\|a^\pi\| + \frac{\|a^\pi a\| \|a^\pi b^d\|}{1 - \|a^\pi a\| \|a^\pi b^d\|} \right. \\ &\quad \left. + \frac{(\|a^\pi a\| \|a^\pi b^d\|)^2}{1 - \|a^\pi a\| \|a^\pi b^d\|} + \|a^\pi b^d\| (\|a^\pi a\| + \|a^\pi b\|) \right) \\ &\quad + \left(1 + \frac{\|a^d b\|}{1 - \|a^d b a a^d\|} \right) \left(\|a^d\| + \frac{\|a^\pi b^d\|}{1 - \|a^\pi a\| \|a^\pi b^d\|} \right. \\ &\quad \left. + \frac{\|a^\pi a\| \|a^\pi b^d\|^2}{1 - \|a^\pi a\| \|a^\pi b^d\|} \right). \end{aligned}$$

Proof. By hypothesis, $\|a^\pi a\| \|a^\pi b^d\| < 1$. Therefore we complete the proof of Theorem 2.3. \square

The following example illustrates that Theorem 2.3 is a nontrivial generalization of [6, Theorem 2.8].

Example 2.5. *Let*

$$a = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, b = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix} \in \mathbb{C}^{3 \times 3}.$$

Clearly, $a^d = b^d = 0$ and $a^\pi = 1$. Therefore we have

$$\|a a^d b a^d\| = 0 < 1, a^\pi b a^2 = 0 \text{ and } a^\pi b a b = 0.$$

But

$$a^\pi b a = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \neq 0.$$

3. COMMUTATIVE CONDITIONS

In this section we shall establish the representation of group inverse of $a + b$ under a kind of commutative perturbation condition which is weaker than $a^\pi b a = a b a^\pi$. The following lemma is crucial.

Lemma 3.1. *Let \mathcal{A} be a Banach algebra, and let $a \in \mathcal{A}^{qnil}$, $b \in \mathcal{A}$. If $a^2 b = a b a$, $b^2 a = b a b$, then $a + b \in \mathcal{A}^d$. In this case,*

$$(a + b)^d = \sum_{n=0}^{\infty} (b^d)^{n+1} (-a)^n + b^\pi a \sum_{n=0}^{\infty} (-1)^n (n+1) (b^d)^{n+2} a^n.$$

Proof. Since $a \in \mathcal{A}^{qnil}$, we have $a^d = 0$. This completes the proof by [13, Theorem 3.3]. \square

We are ready to prove:

Theorem 3.2. *Let \mathcal{A} be a Banach algebra, and let $a, b, a^\pi b \in \mathcal{A}^d$. If*

$$\|a^d b a a^d\| < 1, a^2 b a^\pi = a^\pi a b a, a^\pi b^2 a = b a b a^\pi,$$

then $a + b$ has group inverse if and only if $(a + b)a^\pi$ has group inverse. In this case,

$$\begin{aligned} & \|(a + b)^\# - a^d\| \\ & \leq \frac{\|a^d b\| \|a^d\|}{1 - \|a^d b\|} + \|a^\pi b a a^d\| \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 \\ & \quad + \|(a + b)a^\pi b a a^d\| \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 \sum_{n=0}^{\infty} \|b^d a^\pi\|^{n+1} \|a a^\pi\|^n \\ & \quad + \|b^\pi a a^\pi\| \|(a + b)a^\pi b a a^d\| \sum_{n=0}^{\infty} (n+1) \|b^d a^\pi\|^{n+2} \|a a^\pi\|^n \\ & \quad \cdot \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 + \sum_{n=0}^{\infty} \|b^d a^\pi\|^{n+1} \|a a^\pi\|^n \end{aligned}$$

$$\begin{aligned}
& + \|a^\pi b a a^d\| \frac{\|a^d\|}{1 - \|a^d b\|} \sum_{n=0}^{\infty} \|b^d a^\pi\|^{n+1} \|a a^\pi\|^n \\
& + \|b^\pi a a^\pi\| \sum_{n=0}^{\infty} (n+1) \|b^d a^\pi\|^{n+2} \|a a^\pi\|^n \\
& + \|b^\pi a a^\pi\| \|a^\pi b a a^d\| \frac{\|a^d\|}{1 - \|a^d b\|} \sum_{n=0}^{\infty} (n+1) \|b^d a^\pi\|^{n+2} \|a a^\pi\|^n.
\end{aligned}$$

Proof. Let $p = a a^d$. Then we have

$$a = \begin{pmatrix} a_1 & 0 \\ 0 & a_2 \end{pmatrix}_p, b = \begin{pmatrix} b_1 & b_3 \\ b_4 & b_2 \end{pmatrix}_p,$$

where a_1 is invertible, a_2 is quasinilpotent. Then

$$a^d = \begin{pmatrix} a_1^{-1} & 0 \\ 0 & 0 \end{pmatrix}_p \quad \text{and} \quad a^\pi = \begin{pmatrix} 0 & 0 \\ 0 & a^\pi \end{pmatrix}_p.$$

Since $a^2 b a^\pi = a^\pi a b a$, we check that

$$b_3 = a a^d b a^\pi = (a^d)^2 a^2 b a^\pi = (a^d)^2 a^\pi a b a = 0.$$

Therefore,

$$a + b = \begin{pmatrix} a_1 + b_1 & 0 \\ b_4 & a_2 + b_2 \end{pmatrix}.$$

Obviously, $a^d b = a^d b a a^d = a_1^{-1} b_1$. Since $\|a^d b a a^d\| < 1$, we have $\|a_1^{-1} b_1\| < 1$. Thus $a_1 + b_1 = a_1(1 + a_1^{-1} b_1)$ is invertible in $p\mathcal{A}p$. Moreover, $a_1 + b_1$ is group invertible in $p\mathcal{A}p$. In view of $a^2 b a^\pi = a^\pi a b a$, we get $b_2 = a^\pi b a^\pi = b a^\pi - (a^d)^2 a^2 b a^\pi = b a^\pi - (a^d)^2 a^\pi a b a = b a^\pi$. That is, $a_2 + b_2 = (a + b) a^\pi$. Therefore, by [9, Theorem 2.1] and the invertibility of $a_1 + b_1$, we derive that $a + b$ is group invertible if and only if $(a + b) a^\pi$ is group invertible. In this case,

$$\begin{aligned}
& (a + b)^\# \\
& = \begin{pmatrix} (a_1 + b_1)^{-1} & 0 \\ (a_2 + b_2)^\pi b_4 (a_1 + b_1)^{-2} - (a_2 + b_2)^\# b_4 (a_1 + b_1)^{-1} & (a_2 + b_2)^\# \end{pmatrix}.
\end{aligned}$$

As $a_2 = aa^\pi \in \mathcal{A}^{\text{qnil}}$ and $b_2 = a^\pi ba^\pi$, it follows from $a^2ba^\pi = a^\pi aba$ that

$$aa^\pi \cdot aa^\pi \cdot a^\pi ba^\pi = aa^\pi \cdot a^\pi ba^\pi \cdot aa^\pi.$$

That is, $a_2^2b_2 = a_2b_2a_2$. Likewise, $b_2^2a_2 = b_2a_2b_2$. By hypothesis, $a^\pi b \in \mathcal{A}^d$, by using Cline's formula, $b_2 \in \mathcal{A}^d$. In light of Lemma 3.1, $a_2 + b_2 \in ((1-p)\mathcal{A}(1-p))^d$. Therefore,

$$\begin{aligned} (a_2 + b_2)^\# &= (a_2 + b_2)^d \\ &= \sum_{n=0}^{\infty} (-1)^n (b_2^d)^{n+1} a_2^n + b_2^\pi a_2 \sum_{n=0}^{\infty} (-1)^n (n+1) (b_2^d)^{n+2} a_2^n \\ &= \sum_{n=0}^{\infty} (-1)^n (b^d a^\pi)^{n+1} (aa^\pi)^n \\ &\quad + b^\pi aa^\pi \sum_{n=0}^{\infty} (-1)^n (n+1) (b^d a^\pi)^{n+2} (aa^\pi)^n. \end{aligned}$$

We derive that

$$\begin{aligned} &(a + b)^\# \\ &= (a_1 + b_1)^{-1} + (a_2 + b_2)^\pi b_4 (a_1 + b_1)^{-2} \\ &\quad - (a_2 + b_2)^\# b_4 (a_1 + b_1)^{-1} + (a_2 + b_2)^\# \\ &= \sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d + \left[a^\pi - \left(\sum_{n=0}^{\infty} (-1)^n (b^d a^\pi)^{n+1} (aa^\pi)^n \right. \right. \\ &\quad \left. \left. + b^\pi aa^\pi \sum_{n=0}^{\infty} (-1)^n (n+1) (b^d a^\pi)^{n+2} (aa^\pi)^n \right) (a + b) a^\pi \right] \\ &\quad \cdot a^\pi b a a^d \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 + \left(\sum_{n=0}^{\infty} (-1)^n (b^d a^\pi)^{n+1} (aa^\pi)^n \right. \\ &\quad \left. + b^\pi aa^\pi \sum_{n=0}^{\infty} (-1)^n (n+1) (b^d a^\pi)^{n+2} (aa^\pi)^n \right) \\ &\quad \cdot \left(a^\pi - a^\pi b a a^d \sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d + a^\pi b a a^d \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 \\
&\quad - \sum_{n=0}^{\infty} (-1)^n (b^d a^\pi)^{n+1} (a a^\pi)^n (a+b) a^\pi b a a^d \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 \\
&\quad - b^\pi a a^\pi \sum_{n=0}^{\infty} (-1)^n (n+1) (b^d a^\pi)^{n+2} (a a^\pi)^n (a+b) a^\pi b a a^d \\
&\quad \cdot \left(\sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \right)^2 + \sum_{n=0}^{\infty} (-1)^n (b^d a^\pi)^{n+1} (a a^\pi)^n \\
&\quad - \sum_{n=0}^{\infty} (-1)^n (b^d a^\pi)^{n+1} (a a^\pi)^n a^\pi b a a^d \sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d \\
&\quad + b^\pi a a^\pi \sum_{n=0}^{\infty} (-1)^n (n+1) (b^d a^\pi)^{n+2} (a a^\pi)^n \\
&\quad - b^\pi a a^\pi \sum_{n=0}^{\infty} (-1)^n (n+1) (b^d a^\pi)^{n+2} (a a^\pi)^n a^\pi b a a^d \sum_{n=0}^{\infty} (-1)^n (a^d b)^n a^d.
\end{aligned}$$

From this, we obtain the estimation of norm

$$\begin{aligned}
\|(a+b)^\# - a^d\| &\leq \frac{\|a^d b\| \|a^d\|}{1 - \|a^d b\|} + \|a^\pi b a a^d\| \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 \\
&\quad + \|(a+b) a^\pi b a a^d\| \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 \sum_{n=0}^{\infty} \|b^d a^\pi\|^{n+1} \|a a^\pi\|^n \\
&\quad + \|b^\pi a a^\pi\| \|(a+b) a^\pi b a a^d\| \sum_{n=0}^{\infty} (n+1) \|b^d a^\pi\|^{n+2} \|a a^\pi\|^n \\
&\quad \cdot \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 + \sum_{n=0}^{\infty} \|b^d a^\pi\|^{n+1} \|a a^\pi\|^n \\
&\quad + \|a^\pi b a a^d\| \frac{\|a^d\|}{1 - \|a^d b\|} \sum_{n=0}^{\infty} \|b^d a^\pi\|^{n+1} \|a a^\pi\|^n
\end{aligned}$$

$$\begin{aligned}
 & + \|b^\pi aa^\pi\| \sum_{n=0}^{\infty} (n+1) \|b^d a^\pi\|^{n+2} \|aa^\pi\|^n. \\
 & + \|b^\pi aa^\pi\| \|a^\pi baa^d\| \frac{\|a^d\|}{1 - \|a^d b\|} \sum_{n=0}^{\infty} (n+1) \|b^d a^\pi\|^{n+2} \|aa^\pi\|^n. \quad \square
 \end{aligned}$$

As an immediate consequence of Theorem 3.2, we now derive

Corollary 3.3. *Let \mathcal{A} be a Banach algebra, and let $a, b, a^\pi b \in \mathcal{A}^d$. If*

$$\begin{aligned}
 & a^2 b a^\pi = a^\pi a b a, \quad a^\pi b^2 a = b a b a^\pi, \\
 & \max\{\|a^d b a a^d\|, \|a a^\pi\| \|b^d a^\pi\|\} < 1,
 \end{aligned}$$

then $a + b$ has group inverse if and only if $(a + b)a^\pi$ has group inverse. In this case,

$$\begin{aligned}
 & \|(a + b)^\# - a^d\| \\
 & \leq \frac{\|a^d b\| \|a^d\|}{1 - \|a^d b\|} + \|a^\pi b a a^d\| \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 \\
 & + \|(a + b)a^\pi b a a^d\| \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 \frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|a a^\pi\|} \\
 & + \|b^\pi a a^\pi\| \|(a + b)a^\pi b a a^d\| \left(\frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|a a^\pi\|} \right)^2 \\
 & \cdot \left(\frac{\|a^d\|}{1 - \|a^d b\|} \right)^2 + \frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|a a^\pi\|} \\
 & + \|a^\pi b a a^d\| \frac{\|a^d\|}{1 - \|a^d b\|} \frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|a a^\pi\|} \\
 & + \|b^\pi a a^\pi\| \left(\frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|a a^\pi\|} \right)^2 \\
 & + \|b^\pi a a^\pi\| \|a^\pi b a a^d\| \frac{\|a^d\|}{1 - \|a^d b\|} \left(\frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|a a^\pi\|} \right)^2
 \end{aligned}$$

Proof. The first part follows from Theorem 3.2. For the second part, by $\|aa^\pi\| \|b^d a^\pi\| < 1$, it is easy to compute that

$$\sum_{n=0}^{\infty} \|b^d a^\pi\|^{n+1} \|aa^\pi\|^n = \frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|aa^\pi\|},$$

$$\sum_{n=0}^{\infty} (n+1) \|b^d a^\pi\|^{n+2} \|aa^\pi\|^n = \left(\frac{\|b^d a^\pi\|}{1 - \|b^d a^\pi\| \|aa^\pi\|} \right)^2.$$

So the upper bound of the norm $\|(a+b)^\# - a^d\|$ can be obtained by Theorem 3.2. \square

Corollary 3.4. (see [6, Theorem 2.12]) *Let \mathcal{A} be a Banach algebra, and let $a, b \in \mathcal{A}^d$. If*

$$\|a^d b a a^d\| < 1, a^\pi b a = a b a^\pi,$$

then $a+b$ has group inverse if and only if $a^\pi(a+b)$ has group inverse. In this case,

$$\|(a+b)^\# - a^d\| \leq \frac{\|a^d b\| \|a^d\|}{1 - \|a^d b\|} + \|a^\pi\| \sum_{n=0}^{\infty} \|b^d\|^{n+1} \|a\|^n.$$

Proof. Since $a^\pi b a = a b a^\pi$, we have $aa^d b a^\pi = a^d(a^\pi b a) = 0$. Hence $aa^d b = aa^d b a a^d$. Also we have $a^\pi b a a^d = a b a^\pi a^d = 0$, and so $b a a^d = a a^d b a a^d$. Accordingly, $a^d a b = b a a^d$. Hence $a^\pi b = b a^\pi$. Therefore we check that

$$\begin{aligned} a^2 b a^\pi &= a(a b a^\pi) = a(a^\pi b a) = a^\pi a b a, \\ a^\pi b^2 a &= a^\pi b a^\pi b a = a^\pi b a b a^\pi = a^\pi b a b = b a^\pi a b = b a b a^\pi. \end{aligned}$$

Since $a^\pi b a = a b a^\pi$, we have $a^\pi b a a^d = a b a^\pi a^d = 0$. By virtue of Theorem 3.2, we obtain the upper bound of the norm

$$\|(a+b)^\# - a^d\| \leq \frac{\|a^d b\| \|a^d\|}{1 - \|a^d b\|} + \|a^\pi\| \sum_{n=0}^{\infty} \|b^d\|^{n+1} \|a\|^n. \quad \square$$

We present a numerical example to demonstrate Theorem 3.2 as follow.

Example 3.5. Let $a = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, b = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \in \mathbb{C}^{4 \times 4}$.

Then

$$a^d = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, a^\pi = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

$$b^d = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2} \end{pmatrix}, b^\pi = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

We compute that

$$a^d b a a^d = \text{diag}\left(\frac{1}{2}, 0, 0, 0\right),$$

and so $\|a^d b a a^d\| < 1$. Moreover, we have $a^2 b a^\pi = a^\pi a b a, a^\pi b^2 a = b a b a^\pi$. But we have

$$a b a^\pi = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix}, a^\pi b a = 0,$$

and then $a^\pi a b \neq a^\pi b a$.

In this case,

$$a^d b = \text{diag}\left(\frac{1}{2}, 0, 0, 0\right), b^d a^\pi = \text{diag}\left(0, 0, 0, \frac{1}{2}\right), a^\pi b a a^d = 0,$$

$$a a^\pi = b^\pi a a^\pi = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

By the upper bound in Theorem 3.2, we should have

$$\|(a + b)^\# - a^d\| \leq 2 + 0 + 0 + 0 + 1 + 0 + 1 + 0 = 4.$$

Indeed, since

$$(a + b)^{\#} = \begin{pmatrix} \frac{2}{3} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{4} \\ 0 & 0 & 0 & \frac{1}{2} \end{pmatrix},$$

we compute that

$$\|(a + b)^{\#} - a^d\| = \frac{13}{12} < 4.$$

REFERENCES

- [1] N. Castro González and J. J. Koliha, New additive results for the g-Drazin inverse, *Proc. R. Soc. Edinb.*, **134A**(2004), 1085–1097.
- [2] H. Chen and M. Sheibani, Generalized Drazin inverses in a ring, *Filomat*, **32**(2018), 5289–5295.
- [3] H. Chen and M. Sheibani, g-Drazin inverses for operator matrices, *Oper. Matrices*, **14**(2020), 23–31.
- [4] D. S. Cvetkovic-Ilic, X. Liu and Y. Wei, Some additive results for the generalized Drazin inverse in a Banach algebra, *Electron. J. Linear Algebra*, **22**(2011), 1049–1058.
- [5] X. Liu and Y. Qin, Perturbation of the generalized Drazin inverse in Banach algebra, *Acta Math. Sinica, Chinese Series*, **57**(2014), 35–46.
- [6] X. Liu, Y. Qin and H. Wei, Perturbation bound of the group inverse and the generalized schur complement in Banach algebra, *Abstr. Appl. Anal.*, 2012, 22 pages. <https://doi.org/10.1155/2012/629178>.
- [7] X. Liu, X. Qin and J. Benítez, New additive results for the generalized Drazin inverse in a Banach algebra, *Filomat*, **30**(2016), 2289–2294.
- [8] X. Liu, S. Wu and Y. Yu, On the Drazin inverse of the sum of two matrices, *J. Appl. Math.*, 2011, 13 pages. <https://doi.org/10.1155/2011/831892>.
- [9] N. Mihajlovic and D. S. Djordjevic, On group invertibility in rings, *Filomat*, **33**(2019), 6141–6150.
- [10] D. Mosić, H. Zou and J. Chen, The generalized Drazin inverse of the sum in a Banach algebra, *Ann. Funct. Anal.*, **8**(2017), 90–105.
- [11] H. Yang and X. Liu, The Drazin inverse of the sum of two matrices and its applications, *J. Comput. Appl. Math.*, **235**(2011), 1412–1417.
- [12] D. Zhang and D. Mosić, Explicit formulae for generalized Drazin inverse of block matrices over a Banach algebra, *Filomat*, **32**(2018), 5907–5917.
- [13] H. Zou, D. Mosić and J. Chen, Generalized Drazin invertibility of the product and sum of two elements in a Banach algebra and its applications, *Turkish. J. Math.*, **41**(2017), 548–563.

DAYONG LIU, COLLEGE OF SCIENCE, CENTRAL SOUTH UNIVERSITY OF FORESTRY AND TECHNOLOGY, CHANGSHA, CHINA

Email address: <liudy@csuft.edu.cn>

TUGCE PEKACAR CALCI, DEPARTMENT OF MATHEMATICS, ANKARA UNIVERSITY, ANKARA, TURKEY

Email address: <tcalci@ankara.edu.tr>

HANDAN KOSE, DEPARTMENT OF MATHEMATICS, AHI EVRAN UNIVERSITY, KIRSEHIR, TURKEY

Email address: handan.kose@ahievran.edu.tr

HUANYIN CHEN, DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU, CHINA

Email address: <huanyinchen@aliyun.com>