

DIMENSION INDEPENDENT ATOMIC DECOMPOSITION FOR DYADIC MARTINGALE \mathbb{H}^1

MACIEJ PALUSZYNSKI AND JACEK ZIENKIEWICZ

ABSTRACT. We introduce atoms for dyadic atomic \mathbb{H}^1 for which the equivalence between atomic and maximal function definitions is dimension independent. We give the sharp, up to $\log(d)$ factor, estimates for the $\mathbb{H}^1 \rightarrow L^1$ norm estimates for the special maximal function.

We define a martingale \mathbb{H}^1 space on \mathbb{R}^d

$$(1) \quad \begin{aligned} M^* f &= \sup_n |\mathbb{E}_n f|, \\ \|f\|_{\mathbb{H}^1} &= \|M^* f\|_{L^1}, \end{aligned}$$

where \mathbb{E}_n is the conditional expectation operator associated with the dyadic grid of scale 2^n . There are various equivalent definitions of \mathbb{H}^1 . In particular, it has been proved in [2] that an equivalent norm can be defined by

$$(2) \quad \begin{aligned} S^* f &= \left(\sum_n |\mathbb{E}_n f - \mathbb{E}_{n+1} f|^2 \right)^{\frac{1}{2}}, \\ \|f\|_{\mathbb{H}^1} &\sim \|S^* f\|_{L^1}, \end{aligned}$$

with the equivalence constants independent of d . Similarly as in the Euclidean case, the atomic decompositions of martingale \mathbb{H}^1 have been proved based either on the maximal function or the square function definitions, see [10]. We note that although the atomic norm obtained in [10] (based on the atomic decomposition) is equivalent to the maximal norm for any single d , the equivalence constants depend on d . The aim of this short note is to fine tune the definition of atoms, for which the equivalence between atomic and maximal function norms is uniform in d . By the results of [2], the same decomposition works for the square function (2) norm.

Date: August 25, 2021.

2010 Mathematics Subject Classification. 42B25, 60G46.

Key words and phrases. Hardy space, atomic decomposition, dyadic martingale.

The motivation for our results is their possible applications. We note that the proposed atomic decomposition can be used to obtain dimension explicit estimates for various classical operators acting on martingale \mathbb{H}^1 . In this note we apply Theorem 3 to estimate the $\mathbb{H}^1 \rightarrow L^1$ norm of special radial maximal function modeling classical Hardy-Littlewood maximal operator. Similar argument works for the heat semigroup, see remark at the end of the paper. We are going to address further questions, in particular dimension explicit estimates for classical SIO, in the future, see Remark 6.

The study of dimension dependence of classical estimates is not new, see papers [6], [9], which have motivated this current research.

We define atoms.

Definition 1. *A function a_Q on \mathbb{R}^d is an atom associated with a dyadic cube Q if*

- (a) $\int a_Q = 0$, $\text{supp } a_Q \subset Q$,
- (b) $\|a_Q\|_{L^1} \leq 1$,
- (c) $\|a_Q\|_{L^\infty} \leq \frac{2^{d+1}}{|Q|}$, d - dimension,
- (d) *we have a decomposition*

$$\left\{x : |a_Q(x)| > \frac{1}{|Q|}\right\} \subset \bigcup_s Q_s,$$

where Q_s are essentially disjoint dyadic cubes, $Q_s \subset Q$, satisfying the following two conditions:

- for $Q_s^\#$ being the dyadic parent of Q_s (one scale above)

$$\frac{1}{|Q_s^\#|} \left| \int_{Q_s^\#} a_Q(x) dx \right| \leq \frac{C}{|Q|},$$

where constant C is independent of the dimension d ,

- a_Q is constant on each Q_s .

Lemma 2. *For an atom a_Q we have*

$$\|a_Q\|_{\mathbb{H}^1} \leq C,$$

where the constant C is independent of the dimension d .

Proof. Pick a dyadic cube $\tilde{Q} \subset Q$. We need to compute the average of a_Q over \tilde{Q} . Suppose \tilde{Q} is a cube other than any of the Q_s 's. Let $\{Q_j^\#\}$ be a family of all maximal $Q_s^\# \subset \tilde{Q}$. We denote by $\langle f \rangle_R$ the average of

f over a set R . Then $\langle a_Q \rangle_{\tilde{Q}} = \langle a^\# \rangle_{\tilde{Q}}$, where $a^\#$ is obtained from a_Q by replacing its value on $Q_j^\#$ by the constant $\langle a_Q \rangle_{Q_j^\#}$. By Definition 1 (d) we have $|a^\#| \leq \frac{C}{|Q|}$ (C independent of the dimension). Now suppose \tilde{Q} is one of the Q_s 's. Then a_Q is constant on \tilde{Q} and averaging leaves its value unchanged. Hence $M^*a_Q(x) \leq \frac{C}{|Q|} + |a(x)|$ and the desired L^1 estimate follows by Definition 1 (b). \square

Remark 1. *If we remove condition (d) from Definition 1 and assume the L^∞ estimate (c) on the entire Q , the statement of the above lemma will remain true, but with linear dependence of the implied constant on the dimension d . In order to see this, one has to use $\|M^*\|_{L^p \rightarrow L^p} \leq \frac{C}{p-1}$ for $p = 1 + \frac{1}{d}$.*

We note, that conditions (b) and (c) of the definition of the atom are suitable for application of the near- L^1 estimates. This approach however, do not seem to lead for the optimal bounds, see Remark 4.

Remark 2. *It seems of interest to find the dimensional statement of the result from [7].*

Theorem 3 (Atomic decomposition). *For $f \in \mathbb{H}^1$ there exist a sequence of atoms $\{a_{Q_i}\}$ and a sequence of constants $\{\lambda_i\}$ such that*

$$f = \sum_i \lambda_i a_{Q_i} \quad \text{in } \mathbb{H}^1,$$

and

$$\sum_i |\lambda_i| \leq \|f\|_{\mathbb{H}^1}.$$

Remark 3. *A similar atomic decomposition can be obtained in a similar, direct way using the square function definition of \mathbb{H}^1 . We omit the details.*

Proof. Clearly, we can assume that

$$\int f = 0, \quad \text{supp } f \subset Q, \quad |Q| = 1.$$

We let the family (finite or infinite) $\{Q_{i_1}\}_{i_1}$ consist of the maximal dyadic subcubes of Q , for which the average of f is non-zero. By differentiation, a.e outside of the union of $\{Q_{i_1}\}_{i_1}$ we have $f = 0$. We will define inductively consecutive generations of subcubes. The

first generation is the family $\{Q_{i_1}\}_{i_1}$. We now construct the second generation of subcubes $\{Q_{i_1, i_2}\}_{i_2} \subset Q_{i_1}$. Recall, that we denote by $\langle f \rangle_Q$ the average of f over Q . Let an integer $R(i_1)$ be defined by

$$2^{R(i_1)} \leq |\langle f \rangle_{Q_{i_1}}| < 2^{R(i_1)+1}, \quad i_1 = 1, 2, \dots$$

The cubes Q_{i_1, i_2} are the maximal subcubes of Q_{i_1} for which

$$|\langle f \rangle_{Q_{i_1, i_2}}| \geq 2^{R(i_1)+2}.$$

In other words $\{Q_{i_1, i_2}\}$ are the moments of the first “break” through the level $2^{R(i_1)+2}$. We define integers $\alpha(i_1, i_2)$ and $R(i_1, i_2)$ by

$$\begin{aligned} |Q_{i_1, i_2}| &= 2^{-\alpha(i_1, i_2)}, \\ 2^{R(i_1, i_2)} &\leq |\langle f \rangle_{Q_{i_1, i_2}}| < 2^{R(i_1, i_2)+1}. \end{aligned}$$

We iterate the procedure for each of the cubes Q_{i_1, i_2} , and we obtain a family of cubes

$$\{Q_{i_1, i_2, \dots, i_l}\}_{l=1, 2, \dots}$$

We now write the decomposition of f into “pre-atoms”

$$\begin{aligned} f &= -\mathbf{1}_Q \langle f \rangle_Q + f \cdot \mathbf{1}_{Q \setminus \cup_{i_1} Q_{i_1}} + \sum_{i_1} \mathbf{1}_{Q_{i_1}} \langle f \rangle_{Q_{i_1}} + \\ &+ \sum_{i_1} \left(-\mathbf{1}_{Q_{i_1}} \langle f \rangle_{Q_{i_1}} + f \cdot \mathbf{1}_{Q_{i_1} \setminus \cup_{i_2} Q_{i_1, i_2}} + \sum_{i_2} \mathbf{1}_{Q_{i_1, i_2}} \langle f \rangle_{Q_{i_1, i_2}} \right) + \\ &+ \sum_{i_1, i_2} \left(-\mathbf{1}_{Q_{i_1, i_2}} \langle f \rangle_{Q_{i_1, i_2}} + f \cdot \mathbf{1}_{Q_{i_1, i_2} \setminus \cup_{i_3} Q_{i_1, i_2, i_3}} + \right. \\ &\quad \left. + \sum_{i_3} \mathbf{1}_{Q_{i_1, i_2, i_3}} \langle f \rangle_{Q_{i_1, i_2, i_3}} \right) + \\ &\dots \\ &+ \sum_{i_1, i_2, \dots, i_s} \left(-\mathbf{1}_{Q_{i_1, \dots, i_s}} \langle f \rangle_{Q_{i_1, \dots, i_s}} + f \cdot \mathbf{1}_{Q_{i_1, \dots, i_s} \setminus \cup_{i_{s+1}} Q_{i_1, \dots, i_s, i_{s+1}}} + \right. \\ &\quad \left. + \sum_{i_{s+1}} \mathbf{1}_{Q_{i_1, \dots, i_s, i_{s+1}}} \langle f \rangle_{Q_{i_1, \dots, i_s, i_{s+1}}} \right) + \\ &\dots \end{aligned}$$

We call “pre-atoms” associated with dyadic cubes Q_{i_1, \dots, i_s} the functions $a_{Q_{i_1, \dots, i_s}}$, which are the normalized elements of the above decomposition

$$a_{Q_{i_1, \dots, i_s}} = \frac{\omega_{Q_{i_1, \dots, i_s}}}{\lambda_{Q_{i_1, \dots, i_s}}},$$

where

$$\begin{aligned}\omega_{Q_{i_1, \dots, i_s}} &= -\mathbf{1}_{Q_{i_1, \dots, i_s}} \langle f \rangle_{Q_{i_1, \dots, i_s}} + \\ &\quad + f \cdot \mathbf{1}_{Q_{i_1, \dots, i_s} \setminus \bigcup_{i_{s+1}} Q_{i_1, \dots, i_{s+1}}} + \\ &\quad + \sum_{i_{s+1}} \mathbf{1}_{Q_{i_1, \dots, i_{s+1}}} \langle f \rangle_{Q_{i_1, \dots, i_{s+1}}},\end{aligned}$$

and

$$\begin{aligned}\lambda_{Q_{i_1, \dots, i_s}} &= 2^{R(i_1, \dots, i_s)+1} |Q_{i_1, \dots, i_s}| + \\ &\quad + 2^{R(i_1, \dots, i_s)+2} |Q_{i_1, \dots, i_s}| + \\ &\quad + \sum_{i_{s+1}} 2^{R(i_1, \dots, i_{s+1})+1} |Q_{i_1, \dots, i_{s+1}}|.\end{aligned}$$

We include in the above the first "pre-atom" of the decomposition

$$\omega_Q = f \cdot \mathbf{1}_{Q \setminus \bigcup_{i_1} Q_{i_1}} + \sum_{i_1} \mathbf{1}_{Q_{i_1}} \langle f \rangle_{Q_{i_1}}, \quad a_Q = \frac{\omega_Q}{\lambda_Q},$$

where

$$\lambda_Q = 1 + \sum_{i_1} 2^{R(i_1)} |Q_{i_1}|.$$

We immediately obtain

- $\int a_{Q_{i_1, \dots, i_s}} = 0$, $\text{supp } a_{Q_{i_1, \dots, i_s}} \subset Q_{i_1, \dots, i_s}$,
- the decomposition

$$f = \sum_{s=1}^{\infty} \sum_{i_1, \dots, i_s} \lambda_{Q_{i_1, \dots, i_s}} \cdot a_{Q_{i_1, \dots, i_s}}, \quad \text{in } \mathbb{H}^1.$$

Observe, that by the definition of $R(i_1, \dots, i_s)$ we have

$$2^{R(i_1, \dots, i_s)} \leq |\langle f \rangle_{Q_{i_1, \dots, i_s}}| < 2^{R(i_1, \dots, i_s)+1},$$

and similarly for the cubes $Q_{i_1, \dots, i_{s+1}}$ ($R(i_1, \dots, i_s)$ replaced by $R(i_1, \dots, i_{s+1})$).

On the cube Q_{i_1, \dots, i_s} , outside $\bigcup_{i_{s+1}} Q_{i_1, \dots, i_{s+1}}$ we have

$$|f| \leq 2^{R(i_1, \dots, i_s)+2}$$

(from the definition of $Q_{i_1, \dots, i_{s+1}}$ and differentiation of integrals). Fur-

ther, for $x \in Q_{i_1, \dots, i_s} \setminus \bigcup_{i_{s+1}} Q_{i_1, \dots, i_{s+1}}$

$$|a_{Q_{i_1, \dots, i_s}}(x)| \leq \frac{2^{R(i_1, \dots, i_s)+1} + 2^{R(i_1, \dots, i_s)+2}}{\lambda_{Q_{i_1, \dots, i_s}}} \leq \frac{1}{|Q_{i_1, \dots, i_s}|},$$

while for $x \in Q_{i_1, \dots, i_{s+1}}$

$$|a_{Q_{i_1, \dots, i_s}}(x)| \leq \frac{|\langle f \rangle_{Q_{i_1, \dots, i_{s+1}}} - \langle f \rangle_{Q_{i_1, \dots, i_s}}|}{\lambda_{Q_{i_1, \dots, i_s}}} \leq \frac{1}{|Q_{i_1, \dots, i_s}|}.$$

Thus, we obtain

$$\|a_{Q_{i_1, \dots, i_s}}\|_{L^1} \leq 1.$$

Observe

$$\begin{aligned} \sum_{s=1}^{\infty} \sum_{i_1, \dots, i_s} |\lambda_{Q_{i_1, \dots, i_s}}| &= \sum_{s=1}^{\infty} \sum_{i_1, \dots, i_s} (2^{R(i_1, \dots, i_s)+1} + 2^{R(i_1, \dots, i_s)+2}) |Q_{i_1, \dots, i_s}| + \\ &\quad + \sum_{s=1}^{\infty} \sum_{i_1, \dots, i_{s+1}} 2^{R(i_1, \dots, i_{s+1})+1} |Q_{i_1, \dots, i_{s+1}}| \\ &\leq C \sum_{s=1}^{\infty} \sum_{i_1, \dots, i_s} 2^{R(i_1, \dots, i_s)+1} |Q_{i_1, \dots, i_s}| \\ &= C \sum_{k=1}^{\infty} 2^k \sum_{\substack{s=1, 2, \dots \\ i_1, \dots, i_s \\ R(i_1, \dots, i_s)=k}} |Q_{i_1, \dots, i_s}| \\ &= (**) \end{aligned}$$

where the constant C is absolute. We make the following 2 observations:

- (a) for a fixed $R(i_1, \dots, i_s) = R(j_1, \dots, j_t)$ the cubes Q_{i_1, \dots, i_s} and Q_{j_1, \dots, j_t} are essentially disjoint. This follows, since if they weren't essentially disjoint, one would have to contain the other, which is impossible (unless, of course, they are the same cube).
- (b) we have

$$Q_{i_1, \dots, i_s} \subset \{x : M^* f(x) > 2^{R(i_1, \dots, i_s)}\}$$

so

$$(**) \leq C \sum_{k=1}^{\infty} 2^k |\{x : M^* f(x) > 2^k\}| \leq C \|M^* f\|_{L^1},$$

with the constant C absolute. We will further decompose the $a_{Q_{i_1, \dots, i_s}}$.

Let $Q_{i_1, \dots, i_{s+1}}^{\#}$ be the dyadic, immediate, parent of $Q_{i_1, \dots, i_{s+1}}$. The cubes $Q_{i_1, \dots, i_{s+1}}^{\#}$ need not to be disjoint, so we fix the maximal one. By the construction it is contained in Q_{i_1, \dots, i_s} . We denote by Q_1, \dots, Q_n those cubes among the immediate dyadic descendants of $Q_{i_1, \dots, i_{s+1}}^{\#}$ which belong to the set $\{Q_{i_1, \dots, i_{s+1}}\}_{s+1}$, while we denote by Q_{n+1}, \dots, Q_{2^d} the remaining descendants ($0 < n \leq 2^d$). Since the cube $Q_{i_1, \dots, i_{s+1}}^{\#}$ is not one of the chosen cubes, thus

$$|\langle f \rangle_{Q_{i_1, \dots, i_{s+1}}^{\#}}| < 2^{R(i_1, \dots, i_s)+2}.$$

Similarly,

$$\begin{aligned} |\langle f \rangle_{Q_k}| &< 2^{R(i_1, \dots, i_s)+2} & k = n+1, \dots, 2^d, \\ |\langle f \rangle_{Q_k}| &\geq 2^{R(i_1, \dots, i_s)+2} & k = 1, \dots, n. \end{aligned}$$

We have

$$\begin{aligned} \sum_{k=1}^n |Q_k| \langle f \rangle_{Q_k} &= |Q_{i_1, \dots, i_{s+1}}^\#| \langle f \rangle_{Q_{i_1, \dots, i_{s+1}}^\#} - \sum_{k=n+1}^{2^d} |Q_k| \langle f \rangle_{Q_k} \\ &\leq |Q_{i_1, \dots, i_{s+1}}^\#| \cdot 2^{R(i_1, \dots, i_s)+2} + \sum_{k=n+1}^{2^d} |Q_k| \cdot 2^{R(i_1, \dots, i_s)+2} \\ &\leq |Q_{i_1, \dots, i_{s+1}}^\#| \cdot 2^{R(i_1, \dots, i_s)+3}. \end{aligned}$$

Let us denote

$$C_k = a_{Q_{i_1, \dots, i_s}} \cdot \mathbf{1}_{Q_k}, \quad k = 1, \dots, n,$$

and we obtain

$$\left| \sum_{k=1}^n C_k \right| \leq \frac{2^d \cdot 2^{R(i_1, \dots, i_s)+3}}{\lambda_{Q_{i_1, \dots, i_s}}} \leq \frac{2 \cdot 2^d}{|Q_{i_1, \dots, i_s}|}.$$

Let

$$\tilde{C} = \frac{1}{n} \sum_{k=1}^n C_k,$$

and let us adjust the value of the pre-atom $a_{Q_{i_1, \dots, i_s}}$ on cubes Q_k from C_k to \tilde{C} ($k = 1, \dots, n$). Call the adjusted pre-atom $\tilde{a}_{Q_{i_1, \dots, i_s}}$. Observe that the new functions have the same support, the same mean, and the L^1 norm is still ≤ 1 . Additionally, the new functions satisfy:

- outside $\bigcup_{i_{s+1}} Q_{i_1, \dots, i_{s+1}}$ we have

$$|\tilde{a}_{Q_{i_1, \dots, i_s}}(x)| \leq \frac{1}{|Q_{i_1, \dots, i_s}|},$$

- on $\bigcup_{i_{s+1}} Q_{i_1, \dots, i_{s+1}}$ we have

$$|\tilde{a}_{Q_{i_1, \dots, i_s}}(x)| \leq \frac{2 \cdot 2^d}{|Q_{i_1, \dots, i_s}|},$$

-

$$\left\{ x : |\tilde{a}_{Q_{i_1, \dots, i_s}}(x)| \geq \frac{1}{|Q_{i_1, \dots, i_s}|} \right\} \subset \bigcup_{i_{s+1}} Q_{i_1, \dots, i_{s+1}},$$

while each cube $Q_{i_1, \dots, i_{s+1}}$ is contained in appropriate parent $Q^\# = Q_{i_1, \dots, i_{s+1}}^\#$, for which we have

$$\begin{aligned} \frac{1}{|Q^\#|} \left| \int_{Q^\#} \tilde{a}_{Q_{i_1, \dots, i_s}}(x) dx \right| &= \frac{1}{|Q^\#|} \left| \sum_{k=1}^n \int_{Q_k} \tilde{a}_{Q_{i_1, \dots, i_s}}(x) dx + \right. \\ &\quad \left. + \sum_{k=n+1}^{2^d} \int_{Q_k} \tilde{a}_{Q_{i_1, \dots, i_s}}(x) dx \right| \\ &= \frac{1}{|Q^\#|} \left| \sum_{k=1}^n \tilde{C} + \sum_{k=n+1}^{2^d} \int_{Q_k} \tilde{a}_{Q_{i_1, \dots, i_s}}(x) dx \right| \\ &= \frac{1}{|Q^\#|} \left| \int_{Q^\#} a_{Q_{i_1, \dots, i_s}}(x) dx \right|. \end{aligned}$$

Denote by $b = b_{Q_{i_1, \dots, i_{s+1}}^\#} = a_{Q_{i_1, \dots, i_s}} - \tilde{a}_{Q_{i_1, \dots, i_s}}$.

Observe that $b \neq 0$ only on $S = \bigcup_{k=1}^n Q_k$. By the construction we have $\|b\|_{L^1(S)} \leq 2\|a_{Q_{i_1, \dots, i_s}}\|_{L^1(S)}$. Moreover, since b is constant on each Q_k , $1 \leq k \leq n$ it satisfies $|Q_{i_1, \dots, i_{s+1}}^\#| \|b\|_{L^\infty} \leq 2^d \|b\|_{L^1}$ and hence $\frac{b}{\|b\|_{L^1}}$ is an atom in the sense of Definition 1.

Now we consider one of cubes Q_{n+1}, \dots, Q_{2^d} , say $Q = Q_{n+1}$. Let again $Q^\# = Q_{i_1, \dots, i_{s+1}}^\#$ be maximal contained in Q . We repeat the above procedure to $Q^\#$ and obtain new atom $\tilde{\tilde{a}}_{Q_{i_1, \dots, i_s}}$ obtained by subsequent modification of $\tilde{a}_{Q_{i_1, \dots, i_s}}$ on subcubes of the new cube $Q^\#$ and atom $b_1 = b_{Q_{i_1, \dots, i_{s+1}}^\#}$. Observe that the supports of b, b_1 are disjoint. We continue recurrently packing up the cubes $Q^\#$. As we finish, we have the atom $\tilde{\tilde{\tilde{a}}}$ (multiple tildas) satisfying conditions of Definition 1 (verification of that is immediate once we observe that a has been modified only on cubes Q_s for which we use Definition 1 (d)), and a sequence of correction atoms b_j 's of disjoint supports. We thus have $\sum_j \|b_j\|_{L^1} \leq 2\|a_{Q_{i_1, \dots, i_s}}\|_{L^1}$. The theorem follows. \square

Example. Let φ be a radial kernel, with support in ball of radius $1 + \frac{1}{d}$, with its radial profile constant on ball of radius 1, linear for $1 \leq |x| \leq 1 + \frac{1}{d}$, such that $\int \varphi = 1$. Let φ_t be the L^1 normalized dilation, $\varphi_t(x) = \frac{1}{t^d} \varphi(\frac{x}{t})$. We will prove the following

Theorem 4. For an atom a satisfying axioms of Definition 1, supported on the cube $[0, 1]^d$, we have

$$(3) \quad \left\| \sup_{t \leq \frac{1}{d}} \varphi_t * a \right\|_{L^1} \leq Cd \log(d) \|a\|_{L^1}.$$

with C an absolute constant. As a consequence, the operator norm of the the maximal function

$$(4) \quad Mf(x) = \left| \sup_{t > 0} \varphi_t * f(x) \right|$$

acting from $H_d^1 \rightarrow L^1$, is at most $Cd \log(d)$.

We also prove the lower estimate $C \frac{d}{\log(d)}$, see comments and the end of the proof.

Proof. Let us fix an atom \mathbf{a} supported on $\mathbf{Q} = [0, 1]^d$. We begin with a sequence of lemmas.

Lemma 5. Let Q be a cube with sidelength ρ , $y \in Q$, $t \geq d^{\frac{3}{2}}\rho$ and

$$\mathbf{1}_t = \mathbf{1}_B, \quad \text{where } B = B\left(0, t\left(1 + \frac{2}{d}\right)\right),$$

(ball of center 0 and radius $t(1 + \frac{1}{d})$). We then have

$$|\varphi_t(x - y_c) - \varphi_t(x - y)| \leq C \cdot \frac{d}{t} \cdot \mathbf{1}_t(x - y) \cdot \frac{1}{|B|} \cdot \|y\|,$$

where y_c is the center of Q . Consequently, for an atom a supported on a cube Q , satisfying $\|a\|_{L^\infty} \leq 1$, with Q and t as above, we have

$$|\varphi_t * a(0)| \leq C \cdot \frac{d\sqrt{d}\rho}{t} \cdot \|\mathbf{1}_t\|_{L^1(Q)}.$$

As a corollary to Lemma 5 we immediately have

Corollary 6. Suppose a_1, a_2, \dots are atoms supported on disjoint cubes of sidelengths ρ , all contained in some cube Q . Assume $\|a_i\|_{L^\infty} \leq 1$ and $t \geq d^{\frac{3}{2}}\rho$. Then

$$(5) \quad \left| \varphi_t * \sum_i a_i(0) \right| \leq C \cdot \frac{d^{3/2}\rho}{t} \cdot \|\mathbf{1}_t\|_{L^1(Q)}.$$

We recall that according to Definition 1 (d) for each atom a we have distinguished cubes Q_s such that a is constant (with value no greater than $2^d/|Q_s|$) on each of the Q_s 's. We will call these distinguished cubes black. For a black cube Q the value of a on Q will be denoted α_Q .

Lemma 7. *Let us fix an integer s with $2^{-s} \simeq td^{-\frac{3}{2}}$. Let \mathcal{M} be the family of the maximal $Q^\#$ with sidelength $\leq 2^{-s}$ (cubes $Q^\#$ are the parent cubes of black cubes given by Definition 1 (d) for the fixed atom \mathbf{a}). The atom \mathbf{a} , decomposes as a sum*

$$\mathbf{a} = a_1^s + a_2^s + a_3^s,$$

where

$$a_1^s(x) = \begin{cases} 0 & : \text{on any black cube with sidelength} \geq 2^{-s}, \\ \frac{\mathbf{1}_{Q^\#(x)}}{|Q^\#|} \int_{Q^\#} \mathbf{a} & : x \in Q^\#, Q^\# \in \mathcal{M}, \\ a(x) & : \text{otherwise,} \end{cases}$$

$$a_2^s(x) = \sum_i \mathbf{1}_{Q_i}(x) \mathbf{a}(x),$$

where Q_i 's are all the black cubes with sidelengths $\geq 2^{-s}$, and

$$a_3^s(x) = \begin{cases} \mathbf{a}(x) - \frac{\mathbf{1}_{Q^\#(x)}}{|Q^\#|} \int_{Q^\#} \mathbf{a} & : x \in Q^\#, Q^\# \in \mathcal{M}, \\ 0 & : \text{otherwise.} \end{cases}$$

Clearly we have

$$|\varphi_t * a_1^s| \leq C,$$

and, moreover, for $t \leq \frac{1}{d}$

$$\text{supp}(\varphi_t * a_1^s) \subset (1 + \frac{4}{d})\mathbf{Q}.$$

(For a cube Q and a positive number s sQ means cube with the same center as Q , and sidelength s times the sidelength of Q .)

Corollary 8. *We have*

$$\begin{aligned} |\varphi_t * \mathbf{a}(x)| &\leq C \cdot \mathbf{1}_{(1+\frac{4}{d})\mathbf{Q}}(x) + \sum_{\substack{Q_i \text{ black,} \\ l(Q_i) \geq d^{\frac{5}{2}} 2^{-s}}} |\alpha_{Q_i}| \mathbf{1}_{(1+4/d)Q_i}(x) + \\ &+ \sum_{\substack{Q_i \text{ black,} \\ 2^{-s} \leq l(Q_i) < d^{\frac{5}{2}} 2^{-s}}} |\alpha_{Q_i}| \cdot |\varphi_t * \mathbf{1}_{Q_i}(x) + \varphi_t * a_3^s(x)|, \\ &= I + II + III + IV \end{aligned}$$

where we denote by $l(Q)$ the sidelength of Q .

First two summands give rise to the L^1 control of the maximal function with constants independent of the dimension. For the third summand we have uniform in t estimate

$$III \leq \sum_{Q_i \text{ black,}} |\alpha_{Q_i}| \sup_{d^{-1}l(Q_i) \leq t \leq l(Q_i)d^{\frac{3}{2}}} \varphi_t * \mathbf{1}_{Q_i}(x)$$

and since

$$\int_{d^{-1}l(Q_i) \leq t \leq l(Q_i)d^{\frac{3}{2}}} \|\partial_t \varphi_t\|_{L^1} \leq Cd \log(d)$$

and

$$(6) \quad \begin{aligned} J &= \sup_{d^{-1}l(Q_i) \leq t \leq l(Q_i)d^{\frac{3}{2}}} \varphi_t * \mathbf{1}_{Q_i}(x) \\ &\leq \int_{d^{-1}l(Q_i) \leq t \leq l(Q_i)d^{\frac{3}{2}}} |\partial_t \varphi_t| * \mathbf{1}_{Q_i}(x) dt \end{aligned}$$

we have $\|III\|_{L^1} \leq Cd \log(d) \|a\|_{L^1}$.

The last summand IV will be estimated by Corollary 6.

Lemma 9. *We have*

$$\begin{aligned} a_3^s(x) &= \sum_{n \geq s} (E_{-n} \mathbf{a}(x) - E_{-n-1} \mathbf{a}(x)) \\ &= \sum_{n \geq s} \sum_{Q \in \mathcal{D}_n} \left(\frac{\mathbf{1}_Q}{|Q|} \int_Q \mathbf{a} - \sum_{\substack{Q' \subset Q \\ Q' \in \mathcal{C}(Q)}} \frac{\mathbf{1}_{Q'}}{|Q'|} \int_{Q'} \mathbf{a} \right) \\ &= \sum_{n \geq s} \sum_{\substack{Q \in \mathcal{D}_n \\ \text{all } Q' \in \mathcal{C}(Q), \text{ not black}}} + \text{remainder} \\ &= I + II, \end{aligned}$$

where \mathcal{D}_n denotes the family of dyadic cubes of sidelengths 2^{-n} , and $\mathcal{C}(Q)$ denotes the family of immediate dyadic descendants of a cube Q .

Observe, that for a fixed $n \geq s$, $Q \in \mathcal{D}_n$, Q of type contained in I , the $a_Q = a_3^s \cdot \mathbf{1}_Q$

$$a_Q(x) = \frac{\mathbf{1}_Q(x)}{|Q|} \int_Q \mathbf{a} - \sum_{\substack{Q' \subset Q \\ Q' \in \mathcal{C}(Q)}} \frac{\mathbf{1}_{Q'}(x)}{|Q'|} \int_{Q'} \mathbf{a}$$

satisfy the assumptions of Corollary 6 with $\rho = 2^{-n} = 2^{-s} 2^{-l}$, $2^{-s} d^{\frac{3}{2}} \simeq t$. Hence, by (5) we obtain $|\sum_{Q \in \mathcal{D}_n} \varphi_t * a_Q| \leq C 2^{-l}$, and we can sum up with respect to l . As a result, we obtain a dimension free L^∞ bound. Let $Q \in \mathcal{D}_n$, $n \geq s$ be such that at least one $Q' \in \mathcal{C}(Q)$ is black. We will then say that Q has type 2. We decompose a_Q further into average 0 functions

$$a_Q(x) = b_Q(x) + e_Q(x)$$

where

$$b_Q(x) = \frac{\mathbf{1}_Q(x)}{|Q|} \int_{\cup_{Q':Q' \text{ black}} Q'} a - \sum_{Q' \in \mathcal{C}(Q) \text{ black}} \frac{\mathbf{1}_{Q'}(x)}{|Q'|} \int_{Q'} a$$

and

$$e_Q(x) = \frac{\mathbf{1}_Q(x)}{|Q|} \int_{\cup_{Q':Q' \text{ not black}} Q'} a - \sum_{Q' \in \mathcal{C}(Q) \text{ not black}} \frac{\mathbf{1}_{Q'}(x)}{|Q'|} \int_{Q'} a$$

Observe, that the family $e_Q(x)$ satisfies again the condition of the Corollary 6, so by the preceding case argument we get

$$\left| \sum_{Q \in \mathcal{D}_n, Q \text{ of type 2}} \varphi_t * e_Q \right| \leq C \cdot 2^{-l}$$

and we again sum up to obtain a dimension free L^∞ bound.

We are left with the estimate for

$$J = \sum_{n \geq s} \left| \sum_{Q \in \mathcal{D}_n, Q \text{ of type 2}} \varphi_t * b_Q(x) \right|$$

where we have an additional relation $2^{-s} d^{\frac{3}{2}} \simeq t$. We have

$$J \leq \sum_{Q \text{ of type 2}} \left| \sup_{t \geq d^{\frac{3}{2}} l(Q)} \varphi_t * b_Q(x) \right|$$

and the right hand side does not depend on t . Observe, that by standard cancellation argument

$$(7) \quad \left\| \sup_{t \geq d^{\frac{3}{2}} l(Q)} \varphi_t * b_Q \right\|_{L^1} \leq C \text{diam}(Q) \|b_Q\|_{L^1} \sup_{t \geq d^{\frac{3}{2}} l(Q)} \frac{d \cdot \mathbf{1}_{B(0, t(1+2/d))}(x)}{t^{d+1} |B(0, 1 + 2/d)|}$$

We have, for $|x| \geq (1 + 1/d) d^{\frac{3}{2}} l(Q)$

$$\sup_{t \geq d^{\frac{3}{2}} l(Q)} \frac{d \cdot \mathbf{1}_{B(0, t(1+2/d))}(x)}{t^{d+1} |B(0, 1 + 2/d)|} = \frac{d(1 + 2/d)^d}{|B(0, 1 + 2/d)| |x|^{d+1}}$$

and integrating in polar coordinates, the expression (7) has L^1 norm bounded by Cd . Since the case $|x| \leq (1 + \frac{1}{d}) d^{\frac{3}{2}} l(Q)$ is immediate, the main estimate (3) follows.

The estimates of the maximal function over the intervals $\frac{1}{d} \leq t \leq d^{\frac{3}{2}}$ and $t \geq d^{\frac{3}{2}}$ follows similarly to (6), (7). We leave the details for the reader. Theorem 4 follows. \square

We now briefly sketch the argument leading to the maximal function estimates from below. We recall that $B, |B|$ denote the unit ball in \mathbb{R}^d and its Lebesgue measure.

First observe, that for $A = 2^{2\lceil \log(d) \rceil} \approx d^2$, the function

$$(8) \quad h(x) = 2^{-d} \mathbf{1}_{[-1,1]^d}(x) - (2A)^{-d} \mathbf{1}_{[-A,A]^d}(x) = h_1(x) - h_2(x)$$

defined on \mathbb{R}^d has \mathbb{H}^1 norm of order $\log(d)$. This can be easily checked by the formula

$$(9) \quad h(x) = \sum_{s=0}^{2\lceil \log(d) \rceil - 1} 2^{-ds} \mathbf{1}_{[-2^s, 2^s]^d}(x) - 2^{-d(s+1)} \mathbf{1}_{[-2^{s+1}, 2^{s+1}]^d}(x) = \sum_{s=0}^{2\lceil \log(d) \rceil - 1} h^s(x)$$

It can be easily checked, that the expectation of each h^s over a grid of the dyadic cubes of sidelength 2^l , $l \geq s+1$ vanish, and that the expectation over a dyadic cubes of sidelength 2^l , $l \leq s$ leaves h^s unchanged. Consequently h^s has its \mathbb{H}^1 norm equal to 2.

Then we consider linearized maximal operator $Th(x) = \varphi_{t(x)} * h(x)$, where we will assume $t(x) = |x| + 4 \leq 3d$ for $|x| \leq 2d$. Then observe that $Th_2(x) = (2A)^{-d}$ for $|x| \leq Cd$ and this function restricted to the ball of radius $2d$ is of the L^1 norm of order $O(1)$ (and even smaller).

Now observe, that the L^1 norm $Th_1(x)$ restricted to the ring $d \leq |x| \leq 2d$ is at least cd , where $c > 0$ is dimension free. The crucial observation is that if $t(x) = |x| + 4$, the ball $B(t)$ covers all points in the support of h_1 lying below (that is in the direction of x) the hyperplane passing through 0 and perpendicular to x . Since $\varphi(x) \geq \frac{c_0}{|B|} \mathbf{1}_B$, where c_0 is a dimension free constant, as a result we have $Th_1(x) \geq \frac{c_0}{2|B|} (|x| + 4)^{-d}$. and the statement follows by integration in polar coordinates.

Remark 4. *The above argument applied to the classical heat maximal function leads to the corresponding lower and upper bounds: $\frac{Cd^{\frac{1}{2}}}{\log(d)}$, $Cd^{\frac{1}{2}} \log(d)$. We note, that “near L^1 ” approach based on Rota’s theorem seems to give upper estimate equal to Cd .*

Remark 5. *The following, easy to prove, inequality is very useful in obtaining the estimates from below*

$$(10) \quad Ma(x) \geq Mr(a)(x)$$

where M denotes the maximal function with respect to a radial kernel, and $r(a)$ is the radialisation of a function a . Using (10) one can obtain lower bound Cd for an example considered in Theorem 4.

Remark 6. *The following lemma can be used to obtain an L^2 version of our argument.*

Lemma. *Let*

$$T(f) = \sum_{Q^\#} \sum_{Q_c \sim Q^\#} \frac{\mathbf{1}_{Q^\#}}{|Q^\#|} \int_{Q_c} f,$$

where Q_c are the black cubes, whose immediate dyadic parent is $Q^\#$.

Then

$$\int_Q |T(f)|^2 \leq A \int_Q |f_c|,$$

where f_c is the restriction of f to the black cubes, and A is a universal constant.

Proof. We have

$$\int_Q |T(f)|^2 = \int_Q \sum_{Q_1^\#, Q_2^\# \subset Q} \sum_{\substack{Q_c^1 \sim Q_1^\# \\ Q_c^2 \sim Q_2^\#}} \frac{\mathbf{1}_{Q_1^\#} \mathbf{1}_{Q_2^\#}}{|Q_1^\#| |Q_2^\#|} \int_{Q_c^1} f \int_{Q_c^2} \bar{f} = (*).$$

Clearly, only intersecting $Q_1^\#, Q_2^\#$ need to be considered, and there are two possibilities: $Q_1^\# \subseteq Q_2^\#$ or $Q_2^\# \subseteq Q_1^\#$. We proceed with the first case, the second is similar.

$$\begin{aligned} (*) &= \sum_{Q_1^\# \subseteq Q_2^\#} \sum_{\substack{Q_c^1 \sim Q_1^\# \\ Q_c^2 \sim Q_2^\#}} \frac{1}{|Q_2^\#|} \int_{Q_c^1} f \int_{Q_c^2} \bar{f} \\ &= \sum_{Q_2^\# \subset Q} \frac{1}{|Q_2^\#|} \int_{\cup_{Q_c \sim Q_2^\#} Q_c \text{-black}} f \sum_{Q_1^\# \subseteq Q_2^\#} \int_{\cup_{Q_c \sim Q_1^\#} Q_c \text{-black}} \bar{f} \end{aligned}$$

All sums and unions are over essentially disjoint cubes. In particular

$$\sum_{Q_1^\# \subseteq Q_2^\#} \int_{\cup_{Q_c \sim Q_1^\#} Q_c \text{-black}} \bar{f} = \int_{\cup_{Q_c \subset Q_2^\#} Q_c \text{-black}} \bar{f}.$$

Since

$$\left| \int_{Q_2^\#} \bar{f} \right| \leq C |Q_2^\#|, \quad C \text{ - universal,}$$

by the definition of the ‘‘parent’’ cubes, and

$$\left| \int_{\substack{Q_2^\# \setminus \cup_{Q_c} \\ Q_c \text{-black}}} \bar{f} \right| \leq C |Q_2^\#|, \quad C \text{ - universal,}$$

by the definition of the black cubes, we also have

$$\left| \int_{\cup_{Q_c \subset Q_2^\#} Q_c \text{-black}} \bar{f} \right| \leq C |Q_2^\#|, \quad C \text{ - universal.}$$

This concludes the proof:

$$\begin{aligned}
(*) &= |(*)| \\
&\leq \sum_{Q_2^\# \subset Q} \frac{1}{|Q_2^\#|} \left| \int_{\cup_{Q_c \sim Q_2^\#} Q_c - \text{black}} f \right| C |Q_2^\#| \\
&\leq C \sum_{Q_2^\# \subset Q} \int_{\cup_{Q_c \sim Q_2^\#} Q_c - \text{black}} |f| \\
&= C \int_Q |f_c|.
\end{aligned}$$

□

REFERENCES

- [1] Coifman, R. R., *A real variable characterization of H^p* , Studia Math., **51** (1974), pp. 269–274.
- [2] Davis, B., *On the integrability of the martingale square function*, Israel J. Math., **8** (1970), pp. 187–190.
- [3] Herz, C. S., *H^p spaces of martingales*, $0 < p \leq 1$, Zeit f. War., **28** (1974), pp. 189–205.
- [4] Herz, C. S., *Bounded mean oscillation and regulated martingales*, Trans. Amer. Math. Soc. **193** (1974), pp. 199–215.
- [5] Latter, R. H., *A decomposition of $H^p(\mathbb{R}^n)$ in terms of atoms*, Studia Math., **62** (1978), pp. 92–101.
- [6] Naor, A. and Tao, T., *Random martingales and localization of maximal inequalities*, J. Func. Anal., **259** (2010), pp. 731–779.
- [7] Paluszynski, M. and Zienkiewicz, J., *A Remark On Atomic Decompositions of Martingale Hardy's Spaces*, J. Geom. Anal., **31** (2021), pp. 8866–8878.
- [8] Stein, E. M., *Harmonic Analysis: Real-variable Methods, Orthogonality and Oscillatory Integrals*, Princeton Univ. Press, Princeton N. J., 1993.
- [9] Stein, E. M. and Strömberg, J.-O., *Behavior of maximal functions in \mathbb{R}^n for large n* , Ark. Mat. **21** (1983), pp. 259–269.
- [10] Weisz, F., *Martingale Hardy Spaces and their Applications in Fourier Analysis*, (Lect. Notes Math., vol. 1568) Berlin, Heidelberg, New York, Springer 1994.