

ASPECTS OF COULOMB GASES

DJALIL CHAFAÏ

ABSTRACT. Coulomb gases are special probability distributions, related to potential theory, that appear at many places in pure and applied mathematics and physics. In these short expository notes, we focus on some models, ideas, and structures. We present briefly selected mathematical aspects, mostly related to exact solvability and first and second order global asymptotics. A particular attention is devoted to two-dimensional exactly solvable models of random matrix theory such as the Ginibre model. Thematically, these notes lie between probability theory, mathematical analysis, and statistical physics, and aim to be very accessible. They form a contribution to a volume of the *Panoramas et Synthèses* series around the workshop *États de la recherche en mécanique statistique*, organized by Société Mathématique de France, held at Institut Henri Poincaré, Paris, in the fall of 2018 (<https://statmech2018.sciencesconf.org/>).

CONTENTS

| | |
|---|----|
| Notation | 2 |
| 1. Coulomb electrostatics and equilibrium measures | 2 |
| 2. Coulomb gases | 6 |
| 2.1. One-dimensional log-gases as Coulomb gases | 6 |
| 2.2. Beta-Ginibre gas | 7 |
| 2.3. From multivariate statistics to atomic physics | 7 |
| 2.4. The Wigner jellium and electrons in metals | 7 |
| 2.5. Random polynomials | 8 |
| 3. First order global asymptotics and large deviations | 8 |
| 3.1. The large deviations principle | 8 |
| 3.2. First order global asymptotics | 9 |
| 3.3. Weakly confining versus strongly confining potential | 9 |
| 3.4. Concentration of measure | 9 |
| 4. Edge behavior | 10 |
| 5. Global fluctuations and Gaussian free field | 10 |
| 6. Aspects of general exact solvability | 11 |
| 7. Exactly solvable two-dimensional gases from random matrix theory | 13 |
| 7.1. Ginibre model | 13 |
| 7.2. More determinantal models | 20 |
| 8. Comments and open problems | 21 |
| Acknowledgments | 22 |
| References | 22 |

There are several introductory texts around Coulomb gases. We refer for instance to [ER05, DG09, Dei99, AGZ10, ABDF11, For10] for the relation to random matrices, to [Ser15] for the relation to analysis and Ginzburg–Landau vortices, to [Bou15, GZ19b] and references therein for a relation to geometry, to [But17] and references therein for a relation to random polynomials, to [Rou15] for a relation to Fock–Hartree quantum theory and Bose–Einstein condensates, to [Ser18b] and [Lew21] for an overview from a mathematical analysis/physics perspective, and to [LACTMS19] for a statistical physics point of view.

Date: Summer 2021, compiled August 26, 2021.

2000 Mathematics Subject Classification. Primary: 82B21; Secondary: 82D05; 81V45; 60F05.

Key words and phrases. Asymptotic analysis; Exactly solvable model; High dimensional phenomenon; Mean-field particle system; Random matrix; Spectral analysis; Determinantal point process; Coulomb gas; Potential theory; Wigner jellium; Langevin dynamics; Large deviations principle; Boltzmann–Gibbs measure.

NOTATION

The Euclidean norm of $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ is $|x| = \sqrt{x_1^2 + \dots + x_d^2}$. It is the modulus if $d = 2$ with the identification $\mathbb{C} = \mathbb{R}^2$. We set $i = (0, 1) \in \mathbb{C}$. The real and imaginary parts of $z \in \mathbb{C}$ are denoted $\Re z$ and $\Im z$. The Lebesgue measure is denoted dx . Let (E, τ) be a topological space with Borel σ -field $\mathcal{B}(E)$. We denote by $\mathcal{C}_b(E, \mathbb{R})$ the set of bounded continuous functions $E \rightarrow \mathbb{R}$, and by $\mathcal{M}_1(E)$ the set of probability measures on $(E, \mathcal{B}(E))$. If μ_1, μ_2, \dots, μ are in $\mathcal{M}_1(\mathbb{R}^d)$, then $\lim_{n \rightarrow \infty} \mu_n = \mu$ weakly, denoted

$$\mu_n \xrightarrow[n \rightarrow \infty]{\mathcal{C}_b} \mu, \text{ when for all } f \in \mathcal{C}_b(E, \mathbb{R}) \text{ we have } \lim_{n \rightarrow \infty} \int f d\mu_n = \int f d\mu.$$

This defines a sequential topology on $\mathcal{M}_1(E)$, giving a Borel σ -field $\mathcal{B}(\mathcal{M}_1(E))$. A random probability measure on E is a random variable taking values in $\mathcal{M}_1(E)$. By $X \sim \mu$ we mean that the random variable X has law μ . We denote by $\stackrel{d}{=}$ and \xrightarrow{d} the equality and the convergence in law respectively.

1. COULOMB ELECTROSTATICS AND EQUILIBRIUM MEASURES

The Coulomb kernel is identical to the Newton kernel. Mathematically, potential theory deals with the analysis of the Laplacian and its Green function and the behavior of harmonic functions. In some sense, it emerges naturally from the gravitation theory of Johannes Kepler and Isaac Newton, as well as from the modeling of electrostatics, namely the study of the distribution of static electric charges on conductors and their interactions. From this last point of view, it takes its historical roots in the works of Charles-Augustin de Coulomb, Joseph-Louis Lagrange, and Carl Friedrich Gauss. Some mathematical parts of potential theory were developed later on by – among others – Johann Peter Gustav Lejeune Dirichlet, Victor Gustave Robin, Henri Poincaré, David Hilbert, Charles-Jean de La Vallée Poussin [de 37], Marcel Brelot [Bre67], Otto Frostman [Fro35], Oliver Dimon Kellogg [Kel67], Gustave Choquet [Cho54], and their followers. Deep links with Markov processes and probability theory were explored in particular by Joseph Leo Doob in [Doo01], Gilbert Hunt, Claude Dellacherie and Paul-André Meyer [DM78], and their followers. Nowadays the basic objects and structures of potential theory appear at many places in mathematics and physics, providing in general a very useful electrostatic modeling or interpretation.

Cet aspect probabiliste, que Brelot regrettait tant d'être arrivé trop tard pour maîtriser, est sans doute le plus bel exemple d'interaction féconde entre deux théories : La théorie du potentiel, née dans le ciel (Kepler 1618, Newton 1665) et la théorie des probabilités née d'un coup de dés (Pascal 1654), donc presque simultanément, devaient après trois siècles et des petits pas l'une vers l'autre (Wiener 1923, puis P. Levy, Doob), prendre avec G. Hunt (1957) pleinement conscience que leurs parties les plus vivaces ne sont que deux faces complémentaires d'un même bel objet, et qu'on ne peut bien comprendre l'une sans connaître l'autre (le traité Dellacherie-P.A. Meyer veut en donner la preuve).

Gustave Choquet, *La vie et l'oeuvre de Marcel Brelot (1903 – 1987)* [Cho90].

Let $d \geq 1$. The *Coulomb kernel* g in \mathbb{R}^d is given by $g(0) = +\infty$, and, for all $x \in \mathbb{R}^d \setminus \{0\}$,

$$g(x) = \begin{cases} \log \frac{1}{|x|} & \text{if } d = 2, \\ \frac{1}{(d-2)|x|^{d-2}} & \text{if not.} \end{cases} \quad (1.1)$$

We say that $(x, y) \mapsto G(x, y) = g(x - y)$ is the Green function of the Laplace operator $\Delta = \partial_1^2 + \dots + \partial_d^2$, and g is the fundamental solution of the Poisson equation, see for instance [LL01, Theorem 6.20]. Indeed, denoting δ_0 the Dirac mass at the origin, we have, in the sense of Schwartz distributions,

$$-\Delta g = c_d \delta_0 \quad \text{and} \quad c_d = d\omega_d = \frac{2\pi^{d/2}}{\Gamma(d/2)} \quad (1.2)$$

where ω_d is the volume of the unit ball (its surface is $d\omega_d$). The case $d = 3$ is physical for electrostatics modeling in the ambient space. The case $d = 2$ also appears at many places in mathematical physics. The case $d = 1$ serves historically as a toy model, less singular but exactly solvable.

For simplicity, we suppose from now on that $d \geq 2$.

| | |
|------|--|
| ... | ... |
| 1660 | Newton, ... |
| 1770 | Coulomb, ... |
| 1800 | Gauss, ... |
| 1870 | Boltzmann, Gibbs, ... |
| 1900 | Thomson, de la Vallée Poussin, ... |
| 1920 | Fock, Hartree, ... |
| 1930 | Wigner, Wishart, ... |
| 1940 | Doob, Onsager, ... |
| 1950 | Choquet, Hunt, Wigner, ... |
| 1960 | Dyson, Ginibre, Mehta, Selberg, ... |
| 1970 | Kosterlitz, Landkof, Pastur, Thouless, ... |
| 1980 | Deift, Laughlin, Lebowitz, Saff, Voiculescu, ... |
| 1990 | Ben Arous, Edelman, Guionnet, Johansson, ... |
| 2000 | Forrester, Erdős, Lewin, Serfaty, Yau, ... |
| ... | ... |

TABLE 1.1. The arrow of time and some of the main actors mentioned in the text or in the references. As mentioned in [Cha15], the *Stigler law of eponymy* states that “*No scientific discovery is named after its original discoverer.*”, attributed by Stephen Stigler to Robert K. Merton. This is also known as the *Arnold principle* by some people.




FIGURE 1.1. Coulomb kernel in dimension 1 (solid line) 2 (dotted line) and 3 (dashed line).

For all $\mu \in \mathcal{M}_1(\mathbb{R}^d)$ such that $\log(1 + |\cdot|)\mathbf{1}_{d=2} \in L^1(\mu)$, the *Coulomb energy* of μ is

$$\mathcal{E}(\mu) = \frac{1}{2} \iint g(x-y) d\mu(x) d\mu(y) \in (-\infty, +\infty]. \quad (1.3)$$

Note that if $d = 2$ then $\mathcal{E}(\mu) = +\infty$ if μ has a Dirac mass. If μ models the distribution of unit charges (say electrons) in \mathbb{R}^d then $\mathcal{E}(\mu)$ is the electrostatic self-interaction energy of the configuration μ .

We say that a Borel set $B \in \mathcal{B}(\mathbb{R}^d)$ is of *positive capacity* when $\text{supp}(\mu) \subset B$ and $\mathcal{E}(\mu) < \infty$ for some $\mu \in \mathcal{M}_1(\mathbb{R}^d)$, and is of *zero capacity* when it does not carry a probability measure μ with $\mathcal{E}(\mu) < \infty$.

For all $\mu \in \mathcal{M}_1(\mathbb{R}^d)$ with $\log(1 + |\cdot|)\mathbf{1}_{d=2} \in L^1(\mu)$, the *Coulomb potential* of μ at $x \in \mathbb{R}^d$ is defined by

$$U_\mu(x) = \int g(x-y) d\mu(y) = (g * \mu)(x).$$

We have $U_\mu(x) \in (-\infty, +\infty]$, and $U_\mu(x) = +\infty$ if μ has a Dirac mass at point x . We also have

$$\mathcal{E}(\mu) = \frac{1}{2} \int U_\mu(x) d\mu(x). \quad (1.4)$$

Since $g \in L^1_{\text{loc}}(\mathbb{R}^2, dx)$, the Fubini–Tonelli theorem gives $U_\mu \in L^1_{\text{loc}}(\mathbb{R}^2, dx)$, hence $U_\mu < +\infty$ almost everywhere. Moreover $U_\mu = g * \mu$, and, in the sense of Schwartz distributions, we get from (1.2) that

$$\Delta U_\mu = -c_d \mu. \quad (1.5)$$

In particular, this gives the formula

$$\mathcal{E}(\mu) = \frac{1}{2} \int U_\mu d\mu = -\frac{1}{2c_d} \int U_\mu \Delta U_\mu dx. \quad (1.6)$$

When $d \geq 3$, the functional \mathcal{E} does not take negative values on probability measures because $g \geq 0$. However, when $d = 2$, the functional \mathcal{E} may take negative values on compactly supported probability measures, due to the change of sign of g when $d = 2$ inside and outside the unit ball. For instance if μ_r is the uniform law on the circle $\{x \in \mathbb{C} : |x| = r\}$ of radius $r > 0$ then, for all $x \in \mathbb{C}^2$,

$$U_{\mu_r}(x) = -\log(r) \mathbf{1}_{|x| \leq r} - \log|x| \mathbf{1}_{|x| > r}, \quad \text{and} \quad \mathcal{E}(\mu_r) = -\frac{\log(r)}{2}, \quad (1.7)$$

which is negative if $r > 1$. See for instance [ST97, (0.5.5) and (I.1.6)] for these computations. Similarly, if μ_R is the uniform law on the disc $\{x \in \mathbb{C} : |x| \leq R\}$ of radius $R > 0$ then we find that for all $x \in \mathbb{C}^2$,

$$U_{\mu_R}(x) = -\frac{1}{2} \left(\frac{|x|^2}{R^2} - 1 + 2 \log R \right) \mathbf{1}_{|x| \leq R} - \log|x| \mathbf{1}_{|x| > R}, \quad \text{and} \quad \mathcal{E}_{\mu_R} = \frac{1}{4} - \log(R). \quad (1.8)$$

The functionals U and \mathcal{E} extend to signed measures. If $\eta = \mu - \nu$ where μ and ν are two compactly supported probability measures on \mathbb{R}^d , then U_η vanishes at infinity and an integration by parts gives

$$\mathcal{E}(\eta) = \frac{1}{2} \int U_\eta d\eta = -\frac{1}{2c_d} \int U_\eta \Delta U_\eta dx = \frac{1}{2c_d} \int |\nabla U_\eta|^2 dx, \quad (1.9)$$

see [Ser15]. This shows that \mathcal{E} does not take negative values on signed measures with total mass zero.

The right hand side of (1.9) is the “carré du champ” in potential theory [Rot76, Hir78] while $-\nabla U_\mu$ is the electric field – “champ électrique” in French – generated by the configuration of charges μ .

Let us introduce now $V : \mathbb{R}^d \rightarrow (-\infty, +\infty]$ such that (we say then that V is an *admissible potential*):

- the function V is lower semi-continuous;
- the set $\{x \in \mathbb{R}^d : V(x) < +\infty\}$ has positive capacity;
- the function V is not beaten by the Coulomb kernel at infinity, namely

$$\liminf_{|x| \rightarrow \infty} (V(x) - \log|x| \mathbf{1}_{d=2}) > -\infty. \quad (1.10)$$

The *electrostatic energy* with *external potential* V is defined from $\mathcal{M}_1(\mathbb{R}^d)$ to $(-\infty, +\infty]$ by

$$\mathcal{E}_V(\mu) = \frac{1}{2} \iint (g(x-y) + V(x) + V(y)) \mu(dx) \mu(dy). \quad (1.11)$$

This makes sense since the function under the double integral is bounded below on $\mathbb{R}^d \times \mathbb{R}^d$ thanks to (1.10). Finally, for all $\mu \in \mathcal{M}_1(\mathbb{R}^d)$, if both $\log(1 + |\cdot|) \mathbf{1}_{d=2}$ and V are in $L^1(\mu)$, then

$$\mathcal{E}_V(\mu) = \mathcal{E}(\mu) + \int V(x) d\mu(x). \quad (1.12)$$

The external potential plays typically the role of a confinement.

The convexity of the quadratic form \mathcal{E}_V is related to a Bochner positivity of the kernel g , see [Lan72, HP00, CGZ14, BHS19]. Indeed for all $\lambda \in (0, 1)$ and $\mu, \nu \in \mathcal{M}_1(\mathbb{R}^d)$ with $\mathcal{E}(\mu) < +\infty$ and $\mathcal{E}(\nu) < +\infty$ and $V \in L^1(\mu) \cap L^1(\nu)$,

$$\frac{\lambda \mathcal{E}_V(\mu) + (1-\lambda) \mathcal{E}_V(\nu) - \mathcal{E}_V(\lambda\mu + (1-\lambda)\nu)}{\lambda(1-\lambda)} = \mathcal{E}(\mu - \nu) = \frac{1}{2c_d} \int |\nabla U_{\mu-\nu}|^2 dx \geq 0.$$

We are now ready for the general concept of equilibrium measure and its properties. The following couple of theorems is a classic in potential theory. For a proof, we refer for instance to the books [Lan72, Hel14, ST97, Ser15] and to the articles [BAG97, CGZ14, Ser18b].

Theorem 1.1 (Equilibrium measure). *The following properties hold true:*

| Dimension d | Potential V | Equilibrium measure μ_V |
|---------------|--|--|
| ≥ 1 | $\infty \mathbf{1}_{ \cdot > r}$ | Uniform on sphere $\{x \in \mathbb{R}^d : x = r\}$ |
| ≥ 1 | finite and \mathcal{C}^2 | With density $\frac{\Delta V}{c_d}$ on the interior of its support (absolutely continuous part of μ_*) |
| ≥ 1 | $\frac{1}{2} \cdot ^2$ | Uniform on unit ball with density $\frac{\mathbf{1}_{ \cdot \leq 1}}{\omega_d}$ |
| (Ginibre) 2 | $\frac{1}{2} \cdot ^2$ | Uniform on unit disc with density $\frac{\mathbf{1}_{ \cdot \leq 1}}{\pi}$ |
| (Spherical) 2 | $\frac{1}{2} \log(1 + \cdot ^2)$ | Heavy-tailed with density $\frac{1}{\pi(1+ \cdot ^2)^2}$ |
| (CUE) 2 | $\infty \mathbf{1}_{([a,b] \times \{0\})^c}$ | Arcsine on $[a, b] \times \{0\}$, density $s \mapsto \frac{\mathbf{1}_{s \in [a,b]}}{\pi \sqrt{(s-a)(b-s)}}$ |
| (GUE) 2 | $\frac{ \cdot ^2}{2} \mathbf{1}_{\mathbb{R} \times \{0\}} + \infty \mathbf{1}_{(\mathbb{R} \times \{0\})^c}$ | Semicircle on $[-2, 2] \times \{0\}$, density $s \mapsto \frac{\sqrt{4-s^2}}{2\pi} \mathbf{1}_{s \in [-2,2]}$ |

TABLE 1.2. Basic examples of equilibrium measures. Some other examples are given in Section 7.2. The last four examples appear as limiting spectral distributions of random matrices. The last two examples are singular in the sense that the potential is infinite outside the real line. They appear as one-dimensional log-gases from random matrices. In this case, the equilibrium measure cannot be deduced as a specialization of the second example, and its computation is a bit more subtle, see for instance [ST97].

- (1) \mathcal{E}_V is strictly convex on its domain, is lower semi-continuous, with compact level sets;
- (2) $\inf_{\mathcal{M}_1(\mathbb{R}^d)} \mathcal{E}_V < +\infty$;
- (3) there exists a unique $\mu_V \in \mathcal{M}_1(\mathbb{R}^d)$, called the equilibrium measure, such that

$$\mathcal{E}_V(\mu_V) = \inf_{\mu \in \mathcal{M}_1(\mathbb{R}^d)} \mathcal{E}_V(\mu) \quad \text{in other words} \quad \mu_V = \arg \min_{\mathcal{M}_1(\mathbb{R}^d)} \mathcal{E}_V.$$

Some examples of equilibrium measures are gathered in Table 1.2.

Theorem 1.2 (Properties of the equilibrium measure). *The following properties hold true:*

- (1) the equilibrium measure μ is compactly supported if

$$\lim_{|x| \rightarrow \infty} (V(x) - \log |x| \mathbf{1}_{d=2}) = +\infty; \quad (1.13)$$

- (2) the equilibrium measure μ_V has finite Coulomb energy $\mathcal{E}(\mu_V) \in \mathbb{R}$;
- (3) we have $\text{supp}(\mu_V) \subset \{x \in \mathbb{R}^d : V(x) \leq R\}$ for some constant $R < \infty$;
- (4) the following Euler–Lagrange equations hold:

- $U_{\mu_V}(x) + V(x) \leq c_V$ for all $x \in \text{supp}(\mu_V)$,
 - $U_{\mu_V}(x) + V(x) \geq c_V$ for all $x \in \mathbb{R}^d$ except on a set of zero capacity,
- where c_V is a quantity called the modified Robin constant defined by

$$c_V = \mathcal{E}(\mu_V) - \int V d\mu_V.$$

In particular, for all $x \in \text{supp}(\mu_V)$ except on a set of zero capacity, we have

$$U_{\mu_V}(x) + V(x) = c.$$

In particular, we have the equality in the sense of distributions

$$\mu_V = \frac{\Delta V}{c_d},$$

and the interior of $\text{supp}(\mu_V)$ does not intersect $\{\Delta V < 0\}$.

Remark 1.3 (Logarithmic kernels and Riesz kernels).

- The logarithmic kernel in dimension d is given by

$$-\log|x|, \quad x \in \mathbb{R}^d, x \neq 0;$$

- The Riesz kernel k_s in \mathbb{R}^d with parameter $s > 0$ is given by

$$\frac{1}{s|x|^s}, \quad x \in \mathbb{R}^d, x \neq 0.$$

- The Coulomb kernel in dimension $d \neq 2$ matches the Riesz kernel with $s = d - 2$;
- The logarithmic kernel for all $d \geq 1$ can be seen as the Riesz kernel with $s = 0$. Indeed, it suffices to remove the singularity in the sense that for all $x \in \mathbb{R}^d$ with $x \neq 0$,

$$\lim_{s \rightarrow 0} \frac{|x|^{-s} - 1}{s} = \partial_{s=0} |x|^{-s} = -\log|x|.$$

In particular the Coulomb kernel in dimension $d = 2$ is the Riesz kernel with $s \rightarrow 0$.

- For all $\alpha \in (0, d)$, the Riesz kernel with $s = d - \alpha$ is the fundamental solution of the fractional Laplace operator $\Delta_\alpha = \Delta^{\frac{\alpha}{2}}$, a Fourier multiplier, non-local operator if $\alpha \neq 2$, see [CGZ14, RS16].

We refer for instance to [BHS19] for more analytic properties of these kernels and various applications.

2. COULOMB GASES

Let $d \geq 2$, $n \geq 1$, $\beta > 0$, and let g and V as before. Suppose moreover that V is such that

$$\int_{\mathbb{R}^d} e^{-n\beta(V(x) - \log(1+|x|)\mathbf{1}_{d=2})} dx < \infty. \quad (2.1)$$

By using the fact that $g \geq 0$ when $d \geq 3$ and $|x - y| \leq (1 + |x|)(1 + |y|)$ when $d = 2$, we get then

$$Z_n = \int_{(\mathbb{R}^d)^n} e^{-\beta E_n(x_1, \dots, x_n)} dx_1 \cdots dx_n < \infty$$

where

$$E_n(x_1, \dots, x_n) = n \sum_{i=1}^n V(x_i) + \frac{1}{2} \sum_{i \neq j} g(x_i - x_j).$$

The Coulomb gas P_n is the Boltzmann–Gibbs probability measure on $(\mathbb{R}^d)^n$ given by

$$dP_n(x_1, \dots, x_n) = \frac{e^{-\beta E_n(x_1, \dots, x_n)}}{Z_n} dx_1 \cdots dx_n. \quad (2.2)$$

It models a “gas of electrons” in \mathbb{R}^d of charge $1/n$, at positions x_1, \dots, x_n , inverse temperature βn^2 , energy $(1/n^2)E_n(x_1, \dots, x_n)$, subject to Coulomb pair interaction and external field of potential V , namely

$$\beta E_n(x_1, \dots, x_n) = \beta n^2 \left(\frac{1}{n} \sum_{i=1}^n V(x_i) + \frac{1}{n^2} \sum_{i < j} g(x_i - x_j) \right). \quad (2.3)$$

Beware that we should interpret P_n as a way to model a random static configuration of charged particles. We deal here with electrostatics rather than with electrodynamics. The charged particles do not move and there is no magnetic field. We have only an electric field.

In view of Remark 1.3, we could also define log-gases and Riesz gases. We do not follow this idea in these notes, for simplicity and because the Coulomb case is by far the most important in applications.

2.1. One-dimensional log-gases as Coulomb gases. The formula (2.2) makes sense provided that $Z_n > 0$. Actually the integration in (2.1) should be interpreted as with respect to the trace of the Lebesgue measure or Hausdorff measure on $\{V < +\infty\} \subset \mathbb{R}^d$. Similarly the integration in (2.2) should be interpreted as with respect to the trace of the Lebesgue measure or Hausdorff measure on $\{V < +\infty\}^n \subset (\mathbb{R}^d)^n$. This allows to incorporate in the Coulomb gas model (2.2) the one-dimensional log-gases of random matrix theory, by taking $d = 2$ and $V = +\infty$ on S^c where S is a one-dimensional subset of \mathbb{R}^2 , typically $S = \{x \in \mathbb{R}^2 : x_2 = 0\}$ or $S = \{x \in \mathbb{R}^2 : |x| = 1\}$. This includes all beta Hermite/Laguerre/Jacobi

ensembles, Gaussian Unitary/Orthogonal/Symplectic Ensembles, etc. For instance, the famous Gaussian Unitary Ensemble (GUE) corresponds to take $d = 2$ and

$$x \in \mathbb{R}^2 \mapsto V(x) = \begin{cases} \frac{|x|^2}{2} & \text{if } x \in S = \mathbb{R} \times \{0\} \\ +\infty & \text{if not} \end{cases}.$$

For simplicity, we do not study further the one-dimensional log-gases, in particular the ones coming from random matrix theory. We refer to the books [Dei99, Meh04, ER05, DG09, For10, AGZ10, PS11]. Actually, most of the models that we consider in the sequel are fully dimensional in the sense that V is finite everywhere. The simplest models that we focus on are two-dimensional: beta-Ginibre gases.

2.2. Beta-Ginibre gas. The case $d = 2$ is known as the *two-dimensional one-component plasma*. We call it the *beta gas* for short. Its density with respect to the Lebesgue measure on $(\mathbb{R}^2)^n = \mathbb{C}^n = \mathbb{R}^{2n}$ is

$$(z_1, \dots, z_n) \in \mathbb{C}^n \mapsto \frac{e^{-n\beta \sum_{j=1}^n V(z_j)}}{Z_n} \prod_{i < j} |z_i - z_j|^\beta. \quad (2.4)$$

The quadratic potential case $V = \frac{1}{2} |\cdot|^2$ is sometimes referred to as the *beta-Ginibre gas*. In the special case $\beta = 2$ and $V = \frac{1}{2} |\cdot|^2$, that we call the *Ginibre gas*, the density of P_n can be written as

$$(z_1, \dots, z_n) \in \mathbb{C}^n \mapsto n^n \varphi_n(\sqrt{n}z_1, \dots, \sqrt{n}z_n) \quad \text{with} \quad \varphi_n(z_1, \dots, z_n) = \frac{e^{-\sum_{j=1}^n |z_j|^2}}{\pi^n \prod_{k=1}^n k!} \prod_{i < j} |z_i - z_j|^2. \quad (2.5)$$

The beta gas (2.4) with $V = \frac{1}{2} |\cdot|^2$ and $\beta \in \{2, 4, 6, \dots\}$ matches the squared modulus of the Laughlin wave function of the fractional quantum hall effect [Lau87, Gir05]. The Ginibre gas (2.5) matches the density of the eigenvalues of Gaussian random matrices [Gin65] (see Section 7.1 for more details), the distribution of vortices in the Ginzburg–Landau modeling of superconductivity [Ser15], and rotating trapped fermions in two dimensions [LACTMS19]. The beta gas (2.4) with $\beta = 2$ such as the Ginibre gas (2.5) has a determinantal structure which provides exact solvability (see Section 7.1 for more details).

2.3. From multivariate statistics to atomic physics. Historically, Coulomb gases emerged in mathematical statistics in the years 1920/30 in the study of the spectral decomposition of empirical covariance matrices of Gaussian samples. We speak nowadays about Laguerre ensembles and Wishart random matrices. In the 1950s, Eugene P. Wigner discovered by accident this model when reading a statistics textbook, and this led him to use random matrices for the modeling of energy levels of heavy nuclei in atomic physics, see for instance [Cha14, BW11]. His work generated an enormous trend of activity in statistical physics in the 1960s, with the works of Gaudin, Mehta, Dyson, Ginibre, Marchenko, Pastur, among others. The term *Coulomb gas* is already in the abstract of the first seminal article of Dyson [Dys62b] and of Ginibre [Gin65]. The terms *Fermi gas* and *one-component plasma* are also used.

2.4. The Wigner jellium and electrons in metals. It turns out that Coulomb gases are related to another famous model of mathematical physics also due to Wigner. Let $S \subset \mathbb{R}^d$ be compact and let μ be a positive measure on \mathbb{R}^d with $\mu(\mathbb{R}^d) = \alpha > 0$. The Coulomb gas with potential

$$V = \begin{cases} -\frac{1}{n} U_\mu & \text{on } S, \\ +\infty & \text{outside } S \end{cases}$$

is known as a *Wigner jellium* with background μ , and is said to be *charge neutral* when $n = \alpha$. The background μ models a positive charge α smeared out on $\text{supp}(\mu)$. This model, or more precisely its thermodynamic limit as $|S| \rightarrow +\infty$, was derived by Wigner in 1938 as an approximation of the Hartree–Fock quantum model in order to model electrons in metals [Wig38], see also the 1904 pre-quantum work by Thomson [Tho04] on electrons and the structure of atoms. Conversely, a Coulomb gas with smooth potential V can be seen as a jellium with background μ of density $\frac{\Delta V}{c_d}$. From this point of view, by looking at (1.8), the complex Ginibre ensemble can be seen as a Jellium with full space Lebesgue background. The measure μ is positive when V is sub-harmonic (meaning that $\Delta V \geq 0$). If V is not sub-harmonic then μ is no-longer positive but we may interpret it as an opposite charge on the subset $\{\Delta V < 0\}$. We refer for instance to [CGZJ20a, CGZJ20b] for a bibliography and a discussion. The term *jellium* was apparently coined by Conyers Herring, the smeared charge being viewed as a positive “jelly”, see [Hug06].

2.5. Random polynomials. The Coulomb or log gases emerging from random matrix theory describe the law of the eigenvalues of a random matrix, the roots of the characteristic polynomial. This random polynomial has random dependent coefficients. We could study the distribution of the roots of random polynomials with random independent coefficients. Actually this question emerged from various fields of research including algebraic and geometric analysis and number theory, for instance with the works of Littlewood and Offord in the 1920s, independently of the works of the statisticians on the spectral analysis of empirical covariance matrices. The simplest model that we could imagine is a random polynomial with independent and identically distributed coefficients. This model is known as Kac polynomials, and the distribution of the roots was computed in the Gaussian case by John Hammersley in 1956. There are several other natural models of random polynomials and plenty of works on such models. The gases emerging from these models are two-dimensional but differ from Coulomb gases due to the presence in the energy of an additional non-quadratic term with respect to the empirical measure. For more details, we refer for instance to [BBL96, BBLu92, KZ13, But17] and references therein. When the degree tends to infinity, such models give rise to random analytic functions, see for instance [KZ14, HKPV09, But17] and references therein.

3. FIRST ORDER GLOBAL ASYMPTOTICS AND LARGE DEVIATIONS

Let P_n be the Coulomb gas as in (2.2). If x_1, \dots, x_n are pairwise distinct elements of \mathbb{R}^d , which holds almost everywhere with respect to P_n in $(\mathbb{R}^d)^n$, we get from (2.3) that

$$E_n(x_1, \dots, x_n) = n^2 \mathcal{E}_V^\#(\mu_{x_1, \dots, x_n})$$

where

$$\mathcal{E}_V^\#(\mu) = \int V d\mu + \frac{1}{2} \iint \mathbf{1}_{u \neq v} g(u-v) d\mu(u) d\mu(v) \quad \text{and} \quad \mu_{x_1, \dots, x_n} = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}.$$

The probability measure P_n is *exchangeable* in the sense that it is invariant by permutation of the n particles. The system is *mean-field* in the sense that each particle interacts with all the other particles via their empirical measure. The density of P_n at (x_1, \dots, x_n) is a function of μ_{x_1, \dots, x_n} and rewrites

$$\frac{\exp\left(-\beta n^2 \mathcal{E}_V^\#(\mu_{x_1, \dots, x_n})\right)}{Z_n}. \quad (3.1)$$

In terms of asymptotic analysis, we expect that $\mathcal{E}_V^\# \approx \mathcal{E}_V$ as $n \rightarrow \infty$, and the Laplace method suggests that under P_n , the empirical measure μ_{x_1, \dots, x_n} concentrates as $n \rightarrow \infty$ around the minimizers of \mathcal{E}_V . Since there is a unique minimizer known as the equilibrium measure μ_V , we expect that the empirical measure μ_{x_1, \dots, x_n} under P_n converges towards μ_V as $n \rightarrow \infty$. More precisely, for all n , let us define

$$X_n = (X_{n,1}, \dots, X_{n,n}) \sim P_n \quad \text{and} \quad \mu_n = \frac{1}{n} \sum_{k=1}^n \delta_{X_{n,k}}.$$

3.1. The large deviations principle. For all Borel subsets $A \subset \mathcal{M}_1(\mathbb{R}^d)$, $\mathbb{P}(\mu_n \in A) = P_n(\mu_{x_1, \dots, x_n} \in A)$. The following theorem translates mathematically the intuition above based on the Laplace principle: $\mathbb{P}(\mu_n \in A) \approx_{n \rightarrow \infty} e^{-\beta n^2 \inf_A (\mathcal{E}_V - \mathcal{E}_V(\mu_V))}$. The difficulty lies in the singularity of the Coulomb interaction.

Theorem 3.1 (Large deviations principle). *We have*

$$\lim_{n \rightarrow \infty} \frac{\log Z_n}{\beta n^2} = -\mathcal{E}_V(\mu_V).$$

Moreover the sequence $(\mu_n)_n$ satisfies to a large deviations principle of speed n^2 and good rate function $\mathcal{E}_V - \mathcal{E}_V(\mu_V)$, in other words for all Borel subset of $A \subset \mathcal{M}_1(\mathbb{R}^d)$, we have

$$\mathcal{E}_V(\mu_V) - \inf_{\text{int}(A)} \mathcal{E}_V \leq \underline{\lim}_{n \rightarrow \infty} \frac{\log \mathbb{P}(\mu_n \in A)}{\beta n^2} \leq \overline{\lim}_{n \rightarrow \infty} \frac{\log \mathbb{P}(\mu_n \in A)}{\beta n^2} \leq \mathcal{E}_V(\mu_V) - \inf_{\text{clo}(A)} \mathcal{E}_V$$

where $\text{int}(A)$ and $\text{clo}(A)$ are the interior and closure of A respectively.

About the proof. The first proof of such a result dates back to [BAG97] and concerns one-dimensional log-gases. It is inspired by the work of Voiculescu on a Boltzmann point of view over free entropy and random matrices. Later contributions include [PH98, Har12, CGZ14]. The approach developed in [DLR20, Ber18b, Gar19] is very efficient. \square

Theorem 3.1 remains valid when $\beta = \beta_n$ provided that

$$\lim_{n \rightarrow \infty} n\beta_n = +\infty.$$

This can be called the “low temperature regime”. In the “high temperature regime” $\beta = \beta_n$ with

$$\lim_{n \rightarrow \infty} n\beta_n = \kappa \in (0, +\infty),$$

then Theorem 3.1 remains valid provided that we replace \mathcal{E}_V by the new functional

$$\mathcal{E} + \frac{1}{\kappa} \text{Entropy}(\cdot \mid \nu_{V,\kappa}) = \mathcal{E}_V + \frac{1}{\kappa} \text{Entropy}(\cdot \mid dx) + c_{V,\kappa}$$

where $\nu_{V,\kappa}$ has density proportional to $e^{-\kappa V}$, and where Entropy is the Kullback–Leibler divergence or relative entropy. Note that $-\text{Entropy}(\cdot \mid dx)$ is by definition the Boltzmann–Shannon entropy. We should also replace μ_V in Theorem 3.1 by the minimizer of this new functional. This is also known as the *crossover regime*, interpolating between μ_V and $\nu_{V,\kappa}$. Formally, if we turn off the interaction by taking $g = 0$ and if we take $\beta_n = \kappa/n$ then P_n is the product probability measure $\nu_{V,\kappa}^{\otimes n}$ and the large deviations principle becomes the classical Sanov theorem associated to the law of large numbers for independent random variables. The crossover regime is considered for instance in [CLMP92, BG99, Gar19, AB19, AS19].

3.2. First order global asymptotics. The (weak) convergence in $\mathcal{M}_1(\mathbb{R}^d)$ is metrized by the bounded-Lipschitz distance defined by

$$d_{\text{BL}}(\mu, \nu) = \sup \left\{ \int f d(\mu - \nu) : \|f\|_{\infty} \leq 1, \|f\|_{\text{Lip}} \leq 1 \right\}$$

where the supremum runs over all measurable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and where

$$\|f\|_{\infty} = \sup_x |f(x)| \quad \text{and} \quad \|f\|_{\text{Lip}} = \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|}.$$

Now for all $r \geq 0$, by Theorem 3.1 with $A = A_r = \{\mu \in \mathcal{M}_1(\mathbb{R}^d) : d_{\text{BL}}(\mu, \mu_V) \geq r\}$, for n large enough,

$$e^{-c_r \beta n^2} \leq \mathbb{P}(d_{\text{BL}}(\mu_n, \mu_V) \geq r) \leq e^{-C_r \beta n^2}, \quad (3.2)$$

where $c_r, C_r > 0$ are constants depending on A_r and \mathcal{E}_V but not on n . In particular, for all $\varepsilon > 0$,

$$\sum_n \mathbb{P}(d_{\text{BL}}(\mu_n, \mu_V) > \varepsilon) < \infty.$$

By the Borel–Cantelli lemma, it follows that regardless of the way we choose a common probability space to define the sequence of random vectors $(X_n)_n$, we have, almost surely,

$$\lim_{n \rightarrow \infty} d_{\text{BL}}(\mu_n, \mu_V) = 0. \quad (3.3)$$

This is a sort of law of large numbers for our system of exchangeable particles. They are not independent due to the Coulomb interaction, and the information about the interaction remains in μ_V . We refer to [Ser15, Ber18a, Gar19] for the relation to the notion of “Gamma convergence”.

3.3. Weakly confining versus strongly confining potential. We could say that V is *weakly confining* when (1.10) holds, and that V is *strongly confining* when (1.13) holds. The integrability condition (2.1) may hold for weakly confining potentials. An example of a two dimensional Coulomb gas with a weakly confining potential is given by the Forrester–Krishnapur spherical ensemble considered in the sequel, for which the equilibrium measure is not compactly supported and is heavy-tailed.

3.4. Concentration of measure. The proof of Theorem 3.1 can be adapted in order to provide quantitative (meaning non-asymptotic) estimates for deviation probabilities. Namely, for all $r \geq 0$,

$$\mathbb{P}(d_{\text{BL}}(\mu_n, \mu_V) \geq r) = \frac{1}{Z_n} \int_{d_{\text{BL}}(\mu_n, \mu_V) \geq r} e^{-\beta n^2 \mathcal{E}_V^{\neq}(\mu_{x_1, \dots, x_n})} dx_1 \cdots dx_n.$$

Now if we could approximate $\mathcal{E}_V^{\neq}(\mu_{x_1, \dots, x_n})$ with $\mathcal{E}_V(\mu_{x_1, \dots, x_n})$ and use an inequality of the form

$$d_{\text{BL}}(\mu, \mu_V) \leq c(\mathcal{E}_V(\mu) - \mathcal{E}_V(\mu_V)),$$

and use a bound of the form

$$\log Z_n \geq n^2 \mathcal{E}_V(\mu_V) + n(\beta \mathcal{E}(\mu_V) + c_V), \quad (3.4)$$

then we would obtain a concentration of measure inequality of the form

$$\mathbb{P}(\mathrm{d}(\mu_n, \mu_V) \geq r) \leq e^{-cn^2r^2 + o(n^2)}.$$

for all n and all $r \geq r_n$ for some threshold r_n . Actually the quantity $\mathcal{E}_V(\mu_{x_1, \dots, x_n})$ is infinite due to the atomic nature of μ_{x_1, \dots, x_n} and the method requires then a regularization procedure. The details are in [CHM18]. The method, inspired by [MMS14], is related to [RS16]. See also [GZ19a, MS19, Ber19a, PG20] for other variations on this topic. Moreover we could replace the bounded-Lipschitz distance by a Kantorovich–Wasserstein distance, provided a growth assumption on V .

Such concentration inequalities around the equilibrium measure provide typically an upper bound on the speed of the almost sure convergence. More precisely if r_n is such that $\sum_n e^{-cn^2r_n^2 + o(n^2)} < \infty$, then by the Borel–Cantelli lemma, we get that almost surely, for n large enough,

$$\mathrm{d}_{\mathrm{BL}}(\mu_n, \mu_V) \leq r_n.$$

On the other hand, in the case of one-dimensional log-gases with strongly convex potential V such as the Gaussian unitary ensemble, another approach is possible for concentration of measure, related to logarithmic Sobolev inequalities, see for instance [CL20] and references therein.

4. EDGE BEHAVIOR

We suppose in this section that V is strongly confining, in particular the equilibrium measure μ_V is compactly supported (Theorem 1.2). The convergence (3.3) holds in a weak sense, which does not imply the convergence of the support. The most general result about the convergence of the support is probably [Ame21], and appears as a refinement of [CHM18, Theorem 1.12]. When V is rotationally invariant, this provides constants $c, r_*, p > 0$ such that for all n and $r \geq r_*$,

$$\mathbb{P}\left(\max_{1 \leq k \leq n} |X_{n,k}| \geq r\right) \leq e^{-cnr^p}.$$

The fluctuation at the edge is a difficult subject which is well understood for one-dimensional log-gases, for which it gives rise to Tracy–Widom laws. For strongly confined rotationally invariant determinantal two-dimensional Coulomb gases, it gives rise to Gumbel laws. An explicit analysis of the Ginibre Coulomb gas is presented in the sequel (Theorem 7.14), see also [CP14, JQ17, CGZJ20b, Seo20, Ame21, BGZ18, GZ18, BGZNW21, CGZJ20a] for more results in the same spirit.

5. GLOBAL FLUCTUATIONS AND GAUSSIAN FREE FIELD

Formally, from (1.9) we could write

$$\mathcal{E}(\mu) = \frac{1}{2} \langle -c_d^{-1} \Delta U_\mu, U_\mu \rangle + \langle -c_d^{-1} \Delta V, U_\mu \rangle,$$

and thus

$$\beta n^2 \mathcal{E}(\mu) = \frac{1}{2} \langle -\beta c_d^{-1} \Delta U_{n\mu}, U_{n\mu} \rangle + \langle -n\beta c_d^{-1} \Delta V, U_{n\mu} \rangle.$$

In view of the Coulomb gas formula (3.1), this suggests to interpret as $n \rightarrow \infty$ the random function $U_{n\mu_{x_1, \dots, x_n}}$ under P_n as a Gaussian with covariance operator $K = c_d(-\beta\Delta)^{-1}$. Actually such an object is known as a *Gaussian Free Field* (GFF). Next, again from (1.9), this suggests to interpret formally as $n \rightarrow \infty$ the random measure $n\mu_{x_1, \dots, x_n} = -c_d^{-1} \Delta U_{\mu_{x_1, \dots, x_n}} = A U_{\mu_{x_1, \dots, x_n}}$ under P_n as a Gaussian random measure with covariance operator $A^2 K = (-c_d^{-1} \Delta)^2 (c_d(-\beta\Delta)^{-1}) = -(\beta c_d)^{-1} \Delta$. This argument would involve in principle a change of variable and a Jacobian, that we do not consider here. This leads naturally to conjecture that for a smooth enough test function $f : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$n \left(\int f d\mu_n - \mathbb{E} \int f d\mu_n \right) = \sum_{k=1}^n f(X_{n,k}) - \mathbb{E}(f(X_{n,k})) \xrightarrow[n \rightarrow \infty]{\mathrm{d}} \mathcal{N}\left(0, \frac{1}{\beta c_d} \int |\nabla f|^2 dx\right). \quad (5.1)$$

This can be seen as the “central limit theorem” statement associated to the “law of large numbers” statement (3.3). The limiting variance could be perturbed by edge effects depending on the relative position of the support and regularity of f and μ_V . We could have also an additional bias correction.

The Coulomb interaction together with the confinement produces a rigidity of the global configuration and reduces the variance of linear statistics. Indeed (5.1) comes with an n scaling that differs from the usual \sqrt{n} scaling for independent random variables (no interaction).

The covariance of the limiting Gaussian in (5.1) is easily guessed from the Hessian at the minimizer of the rate function in the large deviations principle of Theorem 3.1. This CLT–LDP link is well known. The GFF is an example of a log-correlated Gaussian field [DRSV17], a fashionable subject.

A statement similar to (5.1) is proved rigorously in [RV07] for the complex Ginibre ensemble by using its exact solvability (determinantal structure). See also [AHM15, AHM11]. Extensions to non-exactly solvable two-dimensional Coulomb gases are considered in [BBNY19, LS18, Ser20a, LZ20].

For one-dimensional log-gases emerging from random matrix theory, central limit theorems such as (5.1) were established using the Laplace transform and “loop equations” in [Joh98]. See also [PS11, BGG17, BLS18, HL21] and references therein for extensions and generalizations.

6. ASPECTS OF GENERAL EXACT SOLVABILITY

Theorem 6.1 is taken from [Cha19a] and [CL20] (see also [CFS21]).

Theorem 6.1 (Exact distributions for special linear statistics of general gases). *Let $X = (X_1, \dots, X_n)$ be a random vector of $(\mathbb{R}^d)^n$, $n, d \geq 1$, with density proportional to*

$$e^{-\sum_i V(x_i)} \prod_{i < j} W(x_i - x_j)$$

where $V : \mathbb{R}^d \rightarrow [0, +\infty]$ and $W : \mathbb{R}^d \rightarrow [0, +\infty]$ are measurable.

- If V and W are homogeneous in the sense that for some $a, b \geq 0$, and for all $\lambda \geq 0$ and $x \in \mathbb{R}^d$,

$$V(\lambda x) = \lambda^a V(x) \quad \text{and} \quad W(\lambda x) = \lambda^b W(x),$$

then

$$V(X_1) + \dots + V(X_n) \sim \text{Gamma}\left(\frac{nd}{a} + \frac{n(n-1)b}{2a}, 1\right).$$

- If $V = \gamma|\cdot|^2$ for some $\gamma > 0$ then

$$X_1 + \dots + X_n \sim \mathcal{N}\left(0, \frac{n}{2\gamma} I_d\right),$$

and moreover the orthogonal projection π on the subspace $\{(z, \dots, z) : z \in \mathbb{R}^d\}$ of $(\mathbb{R}^d)^n$ satisfies $\pi(X) = \frac{X_1 + \dots + X_n}{n}(1, \dots, 1)$, and furthermore $\pi(X)$ and $\pi^\perp(X) = X - \pi(X)$ are independent.

Proof of Theorem 6.1. Recall that linear change of variable is valid for integrals of measurable functions. *First formula.* For all $\theta > 0$, we have, with the substitution $x_i = \left(\frac{1}{1+\theta}\right)^{1/a} y_i$,

$$\int_{(\mathbb{R}^d)^n} e^{-\theta \sum_i V(x_i)} e^{-\sum_i V(x_i)} \prod_{i < j} W(x_i - x_j) dx = \left(\frac{1}{1+\theta}\right)^{\frac{nd}{a} + \frac{(n^2-n)b}{2a}} \int_{(\mathbb{R}^d)^n} e^{-\sum_i V(y_i)} \prod_{i < j} W(y_i - y_j) dy.$$

We recognize the Laplace transform of $\text{Gamma}\left(\frac{nd}{a} + \beta \frac{n(n-1)b}{2a}, 1\right)$, namely

$$\int_0^\infty e^{-\theta x} x^{\alpha-1} e^{-\lambda x} dx = \int_0^\infty x^{\alpha-1} e^{-(\lambda+\theta)x} dx = \left(\frac{\lambda}{\lambda+\theta}\right)^\alpha \frac{\Gamma(\alpha)}{\lambda^\alpha},$$

therefore $\sum_i V(X_i) \sim \text{Gamma}\left(\frac{nd}{a} + \frac{n(n-1)b}{2a}, 1\right)$.

Second formula. For all $\theta \in \mathbb{R}^d$, we have, with the substitution $y_i = x_i + \frac{1}{2\gamma}\theta$ (a translation or shift),

$$\int_{(\mathbb{R}^d)^n} e^{-\theta \cdot \sum_i x_i} e^{-\sum_i V(x_i)} \prod_{i < j} W(x_i - x_j) dx = e^{\frac{n}{2\gamma} \theta \cdot \frac{\theta}{2}} \int_{(\mathbb{R}^d)^n} e^{-\sum_i V(y_i)} \prod_{i < j} W(y_i - y_j) dy,$$

and we recognize the Laplace transform of the Gaussian law $\mathcal{N}\left(0, \frac{n}{2\gamma} I_d\right)$. Finally the properties related to $\pi(X)$ follow from the quadratic nature of V and the shift invariance of W , and correspond to a factorization of the law of X , namely, denoting $\pi^\perp(x) = x - \pi(x)$ and using $|x|^2 = |\pi(x)|^2 + |\pi^\perp(x)|^2$ (Pythagoras theorem) and $x_i - x_j = \pi^\perp(x)_i - \pi^\perp(x)_j$ (from the definition of π), we get

$$e^{-\sum_i V(x_i)} \prod_{i < j} W(x_i - x_j) = e^{-\gamma|\pi(x)|^2} \times e^{-\gamma|\pi^\perp(x)|^2} \prod_{i < j} W(\pi^\perp(x)_i - \pi^\perp(x)_j).$$

This provides the independence of $\pi(X)$ and $\pi^\perp(X)$ as well as the fact that $\pi(X) \sim \mathcal{N}\left(0, \frac{1}{2\gamma} I_d\right)$. \square

Corollary 6.2 (Exact laws for beta-Ginibre gases). *Let us consider $X_n = (X_{n,1}, \dots, X_{n,n}) \sim P_n$ where P_n is as in (2.4) with $\beta > 0$ and $V = \frac{1}{2}|\cdot|^2$. In other words, the density of P_n in \mathbb{C}^n is given by*

$$(z_1, \dots, z_n) \in \mathbb{C}^n \mapsto \frac{e^{-n\frac{\beta}{2}(|z_1|^2 + \dots + |z_n|^2)}}{Z_n} \prod_{i < j} |z_i - z_j|^\beta.$$

Then

$$X_{n,1} + \dots + X_{n,n} \sim \mathcal{N}\left(0, \frac{I_2}{\beta}\right) \quad \text{and} \quad |X_n|^2 = |X_{n,1}|^2 + \dots + |X_{n,n}|^2 \sim \text{Gamma}\left(n + \beta \frac{n(n-1)}{4}, \beta \frac{n}{2}\right),$$

and in particular

$$\mathbb{E}(|X_{n,1} + \dots + X_{n,n}|^2) = \frac{2}{\beta} \quad \text{and} \quad \mathbb{E}(|X_n|^2) = \mathbb{E}(|X_{n,1}|^2 + \dots + |X_{n,n}|^2) = \frac{2}{\beta} + \frac{n-1}{2}.$$

When $\beta = 2$ we recover the Ginibre gas (2.5). Beyond this case, and up to our knowledge, it seems that there is no useful matrix model with independent entries for which the spectrum follows this β gas.

With $\beta = \frac{1}{n}$, we get as $n \rightarrow \infty$ that the variance of the Gauss – Ginibre crossover is $2 + \frac{1}{2} = \frac{5}{2}$.

Proof of Corollary 6.2. It suffices to use Theorem 6.1 with $d = 2$, $V = n\frac{\beta}{2}|\cdot|^2$, $W = |\cdot|^\beta$, for which $a = 2$ and $b = \beta$, and the scaling property $\sigma Z \sim \text{Gamma}(\alpha, \frac{\lambda}{\sigma})$ when $Z \sim \text{Gamma}(\alpha, \lambda)$, for any $\sigma > 0$.

Note that in the determinantal case $\beta = 2$, Theorem 7.13 gives that $n|X|^2 \sim \text{Gamma}(1 + 2 + \dots + n, 1)$ since it has the law of a sum of n independent random variables of law $\text{Gamma}(1, 1), \dots, \text{Gamma}(n, 1)$. \square

Remark 6.3 (Real case). *For all $\beta > 0$, $n \geq 2$, let P_n be the law on \mathbb{R}^n with density*

$$(x_1, \dots, x_n) \in \mathbb{R}^n \mapsto \frac{e^{-n\frac{\beta}{4}(x_1^2 + \dots + x_n^2)}}{Z_n} \prod_{i < j} |x_i - x_j|^\beta.$$

The normalization Z_n can be explicitly computed via a Mehta – Selberg integral [FW08]. It is a quadratically confined one-dimensional log-gas known as the real beta Hermite gas. The case $\beta = 2$ corresponds to GUE. If $X_n = (X_{n,1}, \dots, X_{n,n}) \sim P_n$, then the proof of Theorem 6.2 provides

$$X_{n,1} + \dots + X_{n,n} = \mathcal{N}\left(0, \frac{2}{\beta}\right) \quad \text{and} \quad X_{n,1}^2 + \dots + X_{n,n}^2 \sim \text{Gamma}\left(\frac{n}{2} + \frac{\beta n(n-1)}{4}, \frac{\beta n}{4}\right),$$

and in particular,

$$\mathbb{E}((X_{n,1} + \dots + X_{n,n})^2) = \frac{2}{\beta} \quad \text{and} \quad \mathbb{E}(X_{n,1}^2 + \dots + X_{n,n}^2) = \frac{2}{\beta} + n - 1.$$

Alternatively, these formulas can also be derived by using the tridiagonal random matrix model of Dumitriu and Edelman [DE02] valid for all real beta Hermite gases, see for instance [CL20].

Remark 6.4 (Langevin dynamics). *The Boltzmann – Gibbs measure P_n defined in (2.2) is the invariant law of the Kolmogorov diffusion process $(X_t)_{t \geq 0}$ solution of the stochastic differential equation*

$$dX_t = \sqrt{2\frac{\alpha}{\beta}} dB_t - \alpha \nabla E_n(X_t) dt \tag{6.1}$$

where $(B_t)_{t \geq 0}$ is a standard Brownian motion on $(\mathbb{R}^d)^n$, and where $\alpha > 0$ is an arbitrary parameter which corresponds to a deterministic time change. The infinitesimal generator of the associated Markov semi-group is the second order linear differential operator without constant term

$$L = \alpha \left(\frac{1}{\beta} \Delta - \nabla E_n \cdot \nabla \right). \tag{6.2}$$

The most standard parametrizations are $\alpha = 1$, which allows to interpret $1/\beta$ as the temperature of the Brownian part, and $\alpha = \beta$. See for instance [Cha15, Roy07, BGL14]. Since E_n is a two-body interaction energy, the operator L can be seen as a mean-field particle approximation of a McKean – Vlasov dynamics, see for instance [BCF18, Ser20b]. The singularity of g makes non-obvious the well-posedness or absence of explosion of (6.1), and we refer to [RS93, AGZ10, EY17] for one-dimensional log-gases, and to [BCF18] for the (two-dimensional) beta-Ginibre gas. See also [BD20] and [AB19] for more recent results on such dynamics. Historically (6.1) emerges as the description of the dynamics of the spectrum of Hermitian Ornstein – Uhlenbeck processes, and is nowadays called a Dyson process, named after [Dys62a]. We say

that (6.1) is a gradient dynamics because the drift is the gradient of a function. From the point of view of statistical physics a stochastic differential equation such as (6.1) is also known as an overdamped Langevin dynamics, which is a degenerate version of the true (kinetic under-damped) Langevin dynamics, see for instance [CF19]. Langevin dynamics can be used for the numerical simulation of P_n , see for instance [CF19, CFS21] and references therein. Dynamics such as (6.1) can be used as an interpolation device between X_0 and $X_\infty \sim P_n$, possibly by using conservation laws related to eigenfunctions. In this spirit, and following [BCF18, CL20], we could prove Corollary 6.2 by using the fact that $\sum_i x_i$ and $\sum_i |x_i|^2$ are essentially eigenfunctions of (6.2), producing Ornstein–Uhlenbeck and Cox–Ingersoll–Ross processes for which the targeted Gaussian and Gamma laws are invariant.

7. EXACTLY SOLVABLE TWO-DIMENSIONAL GASES FROM RANDOM MATRIX THEORY

The spectrum of several random matrix models are gases in dimension $d \in \{1, 2\}$ with $W = g$. The cases $\beta \in \{1, 2, 4\}$ play often a special role related to algebra. We refer to [Meh04, ER05, For10] for more details on the zoology of random matrices. It is natural to ask if there exists a random matrix model with independent entries for which the spectrum is distributed according to the beta gas (2.4). Up to our knowledge, the answer is negative for (2.4) in general but positive for the Ginibre gas (2.5).

7.1. Ginibre model. A (complex) Ginibre random matrix \mathbf{M} is an $n \times n$ complex matrix such that

$$\left\{ \Re \mathbf{M}_{i,j}, \Im \mathbf{M}_{i,j} : 1 \leq i, j \leq n \right\} \quad (7.1)$$

are independent and Gaussian random variables of law $\mathcal{N}(0, \frac{1}{2})$. In other words the complex random variables $\{\mathbf{M}_{i,j} : 1 \leq i, j \leq n\}$ are independent and Gaussian of law $\mathcal{N}(0, \frac{1}{2}I_2)$. Note that $\mathbb{E}(|\mathbf{M}_{i,j}|^2) = 1$.

Let $(\lambda_1, \dots, \lambda_n)$ be the eigenvalues of \mathbf{M} seen as an exchangeable random vector of \mathbb{C}^n . This means that we randomize the numbering of the eigenvalues with an independent uniform random permutation of $\{1, \dots, n\}$. Equivalently, this corresponds to consider the random multi-set encoding the spectrum, keeping by this way the possible multiplicities but discarding the numbering of the eigenvalues.

Theorem 7.1 (From the Ginibre random matrix to the Ginibre gas). *The exchangeable random vector*

$$\left(\frac{\lambda_1}{\sqrt{n}}, \dots, \frac{\lambda_n}{\sqrt{n}} \right)$$

is distributed according to the Ginibre gas (2.5). In other words $(\lambda_1, \dots, \lambda_n)$ has density φ_n as in (2.5).

Idea of the proof. The set of $n \times n$ complex matrices with multiple eigenvalues has zero Lebesgue measure. Since the law of \mathbf{M} is absolutely continuous, it follows that almost surely \mathbf{M} is diagonalizable with distinct eigenvalues. The density is proportional to $(M^* = \overline{M}^\top$ is the conjugate-transpose of M)

$$M \mapsto e^{-\sum_{i,j=1}^n |M_{i,j}|^2} = e^{-\text{Trace}(MM^*)}.$$

In order to compute the law of the spectrum of \mathbf{M} , an idea is to use for instance the Schur unitary decomposition as a change of variable. Namely, if M is diagonalizable, then the Schur decomposition is the matrix factorization $M = U(D + N)U^*$ where U is unitary, D is diagonal, and N is upper triangular with null diagonal (nilpotent). The matrix D carries the eigenvalues of M . We have the decoupling

$$\text{Trace}(MM^*) = \text{Trace}(DD^*) + \text{Trace}(NN^*).$$

This allows to integrate out (N, U) in the density and to get that the law of the eigenvalues of \mathbf{M} is given by (2.5). The term $\prod_{i < j} |x_i - x_j|^2$ is the modulus of the determinant of the Jacobian of the change of variable. We obtain that for every symmetric bounded (or positive) measurable function $F : \mathbb{C}^n \rightarrow \mathbb{R}$,

$$\mathbb{E}[F(\lambda_1, \dots, \lambda_n)] = \int_{\mathbb{C}^n} F(z_1, \dots, z_n) \varphi_n(z_1, \dots, z_n) dz_1 \cdots dz_n$$

where $dz_1 \cdots dz_n$ stands for the Lebesgue measure on $\mathbb{C}^n = \mathbb{R}^{2n}$. The result goes back to [Gin65]. The scheme of proof that we follow here can be found in [KS11], see also [Meh04, For10, Ch. 15]. \square

Remark 7.2 (Immediate properties of Ginibre random matrices).

- Since the law of \mathbf{M} is absolutely continuous, almost surely $\mathbf{M}\mathbf{M}^* \neq \mathbf{M}^*\mathbf{M}$ (non-normality);
- By the law of large numbers, almost surely, as $n \rightarrow \infty$, $\frac{1}{\sqrt{n}}\mathbf{M}$ has orthonormal rows/columns;
- The law of \mathbf{M} is bi-unitary invariant: if U and V are unitary then UMV and \mathbf{M} have same law;

- The Hermitian random matrices $\frac{1}{\sqrt{2}}(\mathbf{M} + \mathbf{M}^*)$ and $\frac{1}{\sqrt{2i}}(\mathbf{M} - \mathbf{M}^*)$, the matrix real and imaginary parts of \mathbf{M} , are independent and belong to the Gaussian Unitary Ensemble (GUE): their density is proportional to $H \mapsto e^{-\frac{1}{2}\text{Trace}(H^2)}$; Conversely, if \mathbf{H}_1 and \mathbf{H}_2 are independent copies of the Gaussian Unitary Ensemble then the random matrices $\frac{1}{\sqrt{2}}(\mathbf{H}_1 + i\mathbf{H}_2)$ and \mathbf{M} have same law.

The exact solvability of the Ginibre gas (2.5) is largely due to a determinantal structure studied below, itself related to the fact that $\beta = 2$ and $W = g$. More precisely, first of all, from (2.5) we have

$$\varphi_n(z_1, \dots, z_n) = \frac{\prod_{k=1}^n \gamma(z_k)}{\prod_{k=1}^n k!} \prod_{i < j} |z_i - z_j|^2 \quad (7.2)$$

where γ is the density of $\mathcal{N}(0, \frac{1}{2}I_2)$ given for all $z \in \mathbb{C}$ by

$$\gamma(z) = \frac{e^{-|z|^2}}{\pi}.$$

Theorem 7.3 (Determinantal structure and marginals). *For all $n \geq 1$ and $(z_1, \dots, z_n) \in \mathbb{C}^n$,*

$$\varphi_n(z_1, \dots, z_n) = \frac{1}{n!} \det [K_n(z_i, z_j)]_{1 \leq i, j \leq n}$$

where the kernel K_n is given for all $z, w \in \mathbb{C}$ by

$$K_n(z, w) = \sqrt{\gamma(z)\gamma(w)} \sum_{\ell=0}^{n-1} \frac{(z\bar{w})^\ell}{\ell!}.$$

More generally, for all $1 \leq k \leq n$, the marginal density

$$(z_1, \dots, z_k) \in \mathbb{C}^k \mapsto \varphi_{n,k}(z_1, \dots, z_k) = \int_{\mathbb{C}^{n-k}} \varphi_n(z_1, \dots, z_n) dz_{k+1} \cdots dz_n$$

satisfies, for all $(z_1, \dots, z_k) \in \mathbb{C}^k$,

$$\varphi_{n,k}(z_1, \dots, z_k) = \frac{(n-k)!}{n!} \det [K_n(z_i, z_j)]_{1 \leq i, j \leq k}.$$

In particular for $k = n$ we get $\varphi_{n,n} = \varphi_n$, while for $k = 1$ we get, for all $z \in \mathbb{C}$,

$$\varphi_{n,1}(z) = \frac{\gamma(z)}{n} \sum_{\ell=0}^{n-1} \frac{|z|^{2\ell}}{\ell!}.$$

We say that the spectrum of \mathbf{M} is a Gaussian determinantal point process, see [HKPV09, Ch. 4].

The “ k -point correlation” is $R_{n,k}(z_1, \dots, z_k) = \frac{n!}{(n-k)!} \varphi_{n,k}(z_1, \dots, z_k) = \det [K_n(z_i, z_j)]_{1 \leq i, j \leq k}$.

Idea of proof. Following for instance [Meh04, Sec. 5.2 and Ch. 15], we get, starting with (7.2),

$$\begin{aligned} \varphi_n(z_1, \dots, z_n) &= \frac{\prod_{k=1}^n \gamma(z_k)}{\prod_{k=1}^n k!} \prod_{1 \leq i < j \leq n} (z_i - z_j) \prod_{1 \leq i < j \leq n} \overline{(z_i - z_j)} \\ &= \frac{\prod_{k=1}^n \gamma(z_k)}{n!} \det \left[\frac{z_j^{i-1}}{\sqrt{(i-1)!}} \right]_{1 \leq i, j \leq n} \det \left[\frac{\overline{z_j}^{i-1}}{\sqrt{(i-1)!}} \right]_{1 \leq i, j \leq n} \\ &= \frac{1}{n!} \det [K_n(z_i, z_j)]_{1 \leq i, j \leq n}. \end{aligned}$$

On the other hand, the orthogonality of $\{ \frac{z^\ell}{\sqrt{\ell!}} : 0 \leq \ell \leq n-1 \}$ in $L^2(\mathbb{C}, \gamma)$ gives the identities

$$\int_{\mathbb{C}} K_n(x, x) dx = n \quad \text{and} \quad \int_{\mathbb{C}} K_n(x, y) K_n(y, z) dy = K_n(x, z), \quad x, z \in \mathbb{C}.$$

Finally the formula for $\varphi_{n,k}$ follows by expanding the determinant in φ_n and using these identities. \square

Theorem 7.4 (Mean circular Law). *Let $\lambda_1, \dots, \lambda_n$ be as in Theorem 7.1 and let us define*

$$\mu_n = \frac{1}{n} \sum_{k=1}^n \delta_{\frac{\lambda_k}{\sqrt{n}}}.$$

Let μ_∞ be the uniform distribution on the unit disc $\{z \in \mathbb{C} : |z| \leq 1\}$ with density $z \in \mathbb{C} \mapsto \frac{\mathbf{1}_{|z| \leq 1}}{\pi}$. Then

$$\mathbb{E}\mu_n \xrightarrow[n \rightarrow \infty]{\mathcal{C}_b} \mu_\infty.$$

Proof. Let $\varphi_{n,1}$ be as in Theorem 7.3. For all $f \in \mathcal{C}_b(\mathbb{C}, \mathbb{R})$, we have, using Theorem 7.1,

$$\mathbb{E} \int f d\mu_n = \frac{1}{n} \sum_{k=1}^n \int_{\mathbb{C}^n} f\left(\frac{z_k}{\sqrt{n}}\right) \varphi_n(z_1, \dots, z_n) dz_1 \cdots dz_n = n \int_{\mathbb{C}} f(z) \varphi_{n,1}(\sqrt{n}z) dz.$$

Thus $\mathbb{E}\mu_n$ has density $n\varphi_{n,1}(\sqrt{n}\bullet)$. By Theorem 7.3 and Lemma 7.5, if $K \subset \{z \in \mathbb{C} : |z| \neq 1\}$ is compact,

$$\lim_{n \rightarrow \infty} \sup_{z \in K} \left| n\varphi_{n,1}(\sqrt{n}z) - \frac{\mathbf{1}_{|z| \leq 1}}{\pi} \right| = \frac{1}{\pi} \lim_{n \rightarrow \infty} \sup_{z \in K} \left| e^{-n|z|^2} e_n(n|z|^2) - \mathbf{1}_{|z| \leq 1} \right| = 0.$$

It follows then by dominated convergence that $\mathbb{E}\mu_n \xrightarrow[n \rightarrow \infty]{\mathcal{C}_b} \mu_\infty$. \square

Lemma 7.5 (Exponential series). *For every $n \geq 1$ and $z \in \mathbb{C}$,*

$$|e_n(nz) - e^{nz} \mathbf{1}_{|z| \leq 1}| \leq r_n(z)$$

where $e_n(z) = \sum_{\ell=0}^{n-1} \frac{z^\ell}{\ell!}$ is the truncated exponential series and

$$r_n(z) = \frac{e^n}{\sqrt{2\pi n}} |z|^n \left(\frac{n+1}{n(1-|z|)+1} \mathbf{1}_{|z| \leq 1} + \frac{n}{n(|z|-1)+1} \mathbf{1}_{|z| > 1} \right).$$

Proof of Lemma 7.5. As in Mehta [Meh04, Ch. 15], for every $n \geq 1$, $z \in \mathbb{C}$, if $|z| \leq n$ then

$$\left| e^z - e_n(z) \right| = \left| \sum_{\ell=n}^{\infty} \frac{z^\ell}{\ell!} \right| \leq \frac{|z|^n}{n!} \sum_{\ell=0}^{\infty} \frac{|z|^\ell}{(n+1)^\ell} = \frac{|z|^n}{n!} \frac{n+1}{n+1-|z|},$$

while if $|z| > n$ then

$$|e_n(z)| \leq \sum_{\ell=0}^{n-1} \frac{|z|^\ell}{\ell!} \leq \frac{|z|^{n-1}}{(n-1)!} \sum_{\ell=0}^{n-1} \frac{(n-1)^\ell}{|z|^\ell} \leq \frac{|z|^{n-1}}{(n-1)!} \frac{|z|}{|z| - n + 1}.$$

Therefore, for every $n \geq 1$ and $z \in \mathbb{C}$,

$$|e_n(nz) - e^{nz} \mathbf{1}_{|z| \leq 1}| \leq \frac{n^n}{n!} \left(|z|^n \frac{n+1}{n+1-|nz|} \mathbf{1}_{|z| \leq 1} + |z|^{n-1} \frac{|nz|}{|nz| - n + 1} \mathbf{1}_{|z| > 1} \right).$$

It remains to use the Stirling bound $\sqrt{2\pi n} n^n \leq n! e^n$ to get the first result. \square

Remark 7.6 (Probabilistic view). *There is a probabilistic interpretation of Lemma 7.5. For all $z \in \mathbb{C}$,*

$$\lim_{n \rightarrow \infty} n\varphi_{n,1}(\sqrt{n}z) = \frac{1}{\pi} \left(\mathbf{1}_{|z| < 1} + \frac{1}{2} \mathbf{1}_{|z|=1} \right).$$

Namely, by rotational invariance, it suffices to consider the case $z = r > 0$. Next, if Y_1, \dots, Y_n are independent and identically distributed random variables following the Poisson law of mean r^2 , then

$$e^{-nr^2} e_n(nr^2) = \mathbb{P}(Y_1 + \dots + Y_n < n) = \mathbb{P}\left(\frac{Y_1 + \dots + Y_n}{n} < 1\right).$$

Now $\lim_{n \rightarrow \infty} \frac{Y_1 + \dots + Y_n}{n} = r^2$ almost surely by the strong law of large numbers, and thus the probability in the right-hand side above tends as $n \rightarrow \infty$ to 0 if $r > 1$ and to 1 if $r < 1$. In other words, for all $r \neq 1$,

$$\lim_{n \rightarrow \infty} e^{-nr^2} e_n(nr^2) = \mathbf{1}_{r < 1}.$$

It remains to note that for $r = 1$ by the central limit theorem we get

$$\mathbb{P}\left(\frac{Y_1 + \dots + Y_n}{n} < 1\right) = \mathbb{P}\left(\frac{Y_1 + \dots + Y_n - n}{\sqrt{n}} < 0\right) \xrightarrow[n \rightarrow \infty]{} \frac{1}{2}.$$

Remark 7.7 (Incomplete gamma function). *It is well known that the Gamma and the Poisson laws are connected. Namely, if $X \sim \text{Gamma}(n, \lambda)$ with $n \geq 1$ and $\lambda > 0$ and $Y \sim \text{Poisson}(r)$ with $r > 0$ then*

$$\mathbb{P}(X \geq \lambda r) = \frac{1}{(n-1)!} \int_r^\infty x^{n-1} e^{-x} dx = e^{-r} \sum_{\ell=0}^{n-1} \frac{r^\ell}{\ell!} = \mathbb{P}(Y \geq n).$$

Also we could use Gamma random variables instead of Poisson random variables in Remark 7.6. Note also that the integral in the middle of the formula above is the incomplete Gamma function $\Gamma(n, r)$. This allows to benefit from the asymptotic analysis of this special function, see [KS11] and references therein.

Theorem 7.8 (Strong circular law). *With the notations of Theorem 7.4, almost surely,*

$$\mu_n \xrightarrow[n \rightarrow \infty]{\mathcal{C}_b} \mu_\infty.$$

Note that this convergence holds regardless of the way we define the random matrices on the same probability space when n varies. This is an instance of the concept of *complete convergence*, see [Yuk98].

Idea of the proof. The argument, due to Jack Silverstein, is in [Hwa86]. It is similar to the quick proof of the strong law of large numbers for independent random variables with bounded fourth moment. It suffices to establish the result for an arbitrary compactly supported $f \in \mathcal{C}_b(\mathbb{C}, \mathbb{R})$. Let us define

$$S_n = \int_{\mathbb{C}} f d\mu_n \quad \text{and} \quad S_\infty = \frac{1}{\pi} \int_{|z| \leq 1} f(z) dz.$$

Suppose for now that we have

$$\mathbb{E}[(S_n - \mathbb{E}S_n)^4] = \mathcal{O}\left(\frac{1}{n^2}\right). \quad (7.3)$$

By monotone convergence or by the Fubini–Tonelli theorem,

$$\mathbb{E} \sum_{n=1}^{\infty} (S_n - \mathbb{E}S_n)^4 = \sum_{n=1}^{\infty} \mathbb{E}[(S_n - \mathbb{E}S_n)^4] < \infty$$

and thus $\sum_{n=1}^{\infty} (S_n - \mathbb{E}S_n)^4 < \infty$ almost surely, which implies $\lim_{n \rightarrow \infty} S_n - \mathbb{E}S_n = 0$ almost surely. Since $\lim_{n \rightarrow \infty} \mathbb{E}S_n = S_\infty$ by Theorem 7.4, we get that almost surely

$$\lim_{n \rightarrow \infty} S_n = S_\infty.$$

Finally, one can swap the universal quantifiers on ω and f thanks to the separability of the set of compactly supported continuous bounded functions $\mathbb{C} \rightarrow \mathbb{R}$ equipped with the supremum norm. To establish the fourth moment bound (7.3), we set

$$S_n - \mathbb{E}S_n = \frac{1}{n} \sum_{k=1}^n Z_k \quad \text{with} \quad Z_k = f\left(\frac{\lambda_k}{\sqrt{n}}\right) - \mathbb{E}f\left(\frac{\lambda_k}{\sqrt{n}}\right).$$

Next, we obtain, with $\sum_{k_1, \dots}$ running over distinct indices in $1, \dots, n$,

$$\begin{aligned} \mathbb{E}[(S_n - \mathbb{E}S_n)^4] &= \frac{1}{n^4} \sum_{k_1} \mathbb{E}[Z_{k_1}^4] \\ &\quad + \frac{4}{n^4} \sum_{k_1, k_2} \mathbb{E}[Z_{k_1} Z_{k_2}^3] \\ &\quad + \frac{3}{n^4} \sum_{k_1, k_2} \mathbb{E}[Z_{k_1}^2 Z_{k_2}^2] \\ &\quad + \frac{6}{n^4} \sum_{k_1, k_2, k_3} \mathbb{E}[Z_{k_1} Z_{k_2} Z_{k_3}^2] \\ &\quad + \frac{1}{n^4} \sum_{k_1, k_2, k_3, k_4} \mathbb{E}[Z_{k_1} Z_{k_2} Z_{k_3} Z_{k_4}]. \end{aligned}$$

The first three terms of the right are $\mathcal{O}(n^{-2})$ since $\max_{1 \leq k \leq n} |Z_k| \leq \|f\|_\infty$. The expressions of $\varphi_{n,3}$ and $\varphi_{n,4}$ from Theorem 7.3 allow to show that the remaining two terms are also $\mathcal{O}(n^{-2})$, see [Hwa86]. \square

The following theorem includes Theorem 7.4, which corresponds to the case $k = 1$.

Theorem 7.9 (Chaoticity). *Let μ_∞ be the uniform distribution on the unit disc $\{z \in \mathbb{C} : |z| \leq 1\}$. For all $1 \leq k \leq n$, denoting by $P_{n,k}$ the k -dimensional marginal distribution of the Ginibre gas (2.5), we have*

$$P_{n,k} \xrightarrow[n \rightarrow \infty]{C_b} \mu_\infty^{\otimes k}.$$

Idea of the proof. The measures $P_{n,k}$ and μ_∞ have densities $\varphi_{n,k}$ and $z \in \mathbb{C} \mapsto \varphi_\infty(z) = \pi^{-1} \mathbf{1}_{|z| \leq 1}$.

The case $k = 1$ is nothing else but Theorem 7.4, namely $P_{n,1} \xrightarrow[n \rightarrow \infty]{C_b} \mu_\infty$. This comes via dominated convergence from the fact that $\lim_{n \rightarrow \infty} \varphi_{n,k} = \varphi_\infty$ uniformly on compact subsets of $\{z \in \mathbb{C} : |z| \neq 1\}$.

Let us consider now the case $k = 2$. Here again, by dominated convergence, it suffices to show that

$$\lim_{n \rightarrow \infty} \varphi^{n,2} = \varphi_\infty^{\otimes 2}$$

uniformly on compact subsets of $\{(z_1, z_2) \in \mathbb{C}^2 : |z_1| \neq 1, |z_2| \neq 1, z_1 \neq z_2\}$.

By Theorem 7.3, for all $z_1, z_2 \in \mathbb{C}$,

$$\begin{aligned} \varphi^{n,2}(z_1, z_2) &= \frac{n}{n-1} \frac{e^{-n(|z_1|^2 + |z_2|^2)}}{\pi^2} (e_n(n|z_1|^2)e_n(n|z_2|^2) - |e_n(nz_1\bar{z}_2)|^2) \\ &= \frac{n}{n-1} \varphi^{n,1}(z_1)\varphi^{n,1}(z_2) - \frac{n}{n-1} \frac{e^{-n(|z_1|^2 + |z_2|^2)}}{\pi^2} |e_n(nz_1\bar{z}_2)|^2 \end{aligned} \quad (7.4)$$

where e_n is as in Lemma 7.5. It follows that for any $n \geq 2$ and $z_1, z_2 \in \mathbb{C}$,

$$\begin{aligned} \Delta_n(z_1, z_2) &= \varphi^{n,2}(z_1, z_2) - \varphi^{n,1}(z_1)\varphi^{n,1}(z_2) \\ &= \frac{1}{n-1} \varphi^{n,1}(z_1)\varphi^{n,1}(z_2) - \frac{n}{n-1} \frac{e^{-n(|z_1|^2 + |z_2|^2)}}{\pi^2} |e_n(nz_1\bar{z}_2)|^2. \end{aligned} \quad (7.5)$$

In particular, using $\varphi^{n,2} \geq 0$ for the lower bound,

$$-\varphi^{n,1}(z_1)\varphi^{n,1}(z_2) \leq \Delta_n(z_1, z_2) \leq \frac{1}{n-1} \varphi^{n,1}(z_1)\varphi^{n,1}(z_2).$$

From this and Lemma 7.5 we first deduce that for any compact subset K of $\{z \in \mathbb{C} : |z| > 1\}$

$$\lim_{n \rightarrow \infty} \sup_{\substack{z_1 \in \mathbb{C} \\ z_2 \in K}} |\Delta_n(z_1, z_2)| = \lim_{n \rightarrow \infty} \sup_{\substack{z_1 \in K \\ z_2 \in \mathbb{C}}} |\Delta_n(z_1, z_2)| = 0.$$

It remains to show that $\Delta_n(z_1, z_2) \rightarrow 0$ as $n \rightarrow \infty$ when z_1 and z_2 are in compact subsets of $\{(z_1, z_2) \in \mathbb{C}^2 : |z_1| < 1, |z_2| < 1\}$. In this case $|z_1\bar{z}_2| \leq 1$, and Lemma 7.5 gives

$$|e_n(nz_1\bar{z}_2)|^2 \leq 2e^{2n\Re(z_1\bar{z}_2)} + 2r_n^2(z_1\bar{z}_2).$$

Next, using the elementary identity $2\Re(z_1\bar{z}_2) = |z_1|^2 + |z_2|^2 - |z_1 - z_2|^2$, we get

$$e^{-n(|z_1|^2 + |z_2|^2)} |e_n(nz_1\bar{z}_2)|^2 \leq 2e^{-n|z_1 - z_2|^2} + 2e^{-n(|z_1|^2 + |z_2|^2)} r_n^2(z_1\bar{z}_2). \quad (7.6)$$

Since $|z_1\bar{z}_2| \leq 1$, the formula for r_n in Lemma 7.5 gives

$$e^{-n(|z_1|^2 + |z_2|^2)} r_n^2(z_1\bar{z}_2) \leq e^{-n(|z_1|^2 + |z_2|^2 - 2 - \log|z_1|^2 - \log|z_2|^2)} \frac{(n+1)^2}{2\pi n}.$$

Using (7.5), (7.6) and the bounds $\varphi^{n,1} \leq \pi^{-1}$ and $u - 1 - \log u > 0$ for $0 < u < 1$, it follows that $\Delta_n(z_1, z_2)$ tends to 0 as $n \rightarrow \infty$ uniformly in z_1, z_2 on compact subsets of

$$\{(z_1, z_2) \in \mathbb{C}^2 : |z_1| < 1, |z_2| < 1, z_1 \neq z_2\}.$$

This finishes the proof of the case $k = 2$. The case $k \geq 3$ follows from the case $k = 2$ by Lemma 7.11. \square

Remark 7.10 (Impossibility of global uniform convergence of densities). *The convergence of $\varphi_{n,1}$ cannot hold uniformly on arbitrary compact sets of \mathbb{C} since the point-wise limit is not continuous on the unit circle. Similarly, the convergence of $\varphi_{n,2}$ cannot hold on $\{(z, z) : z \in \mathbb{C}, |z| < 1\}$ since $\varphi^{n,2}(z, z) = 0$ for any $n \geq 2$ and $z \in \mathbb{C}$ while $\varphi_\infty(z)\varphi_\infty(z) = \pi^{-2} \neq 0$ when $|z| < 1$.*

Lemma 7.11 (Chaoticity). *Let E be a Polish space. For all $n \geq 1$, let $P_n \in \mathcal{M}_1(E^n)$ be exchangeable, and for all $1 \leq k \leq n$, let $P_{n,k} \in \mathcal{M}_1(E^k)$ be its k -dimensional marginal distribution. For all $n \geq 1$, let us pick a random vector $X_n = (X_{n,1}, \dots, X_{n,n}) \sim P_n$ and let us define the random empirical measure*

$$\mu_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_{n,i}}.$$

For all $\mu \in \mathcal{M}_1(E)$, the following properties are equivalent:

- (1) $\mu_n \xrightarrow[n \rightarrow \infty]{C_b} \delta_\mu$ in $\mathcal{M}_1(\mathcal{M}_1(E))$ (here we see μ_n a random variable taking values in $\mathcal{M}_1(E)$);
- (2) $P_{n,k} \xrightarrow[n \rightarrow \infty]{C_b} \mu^{\otimes k}$ for any fixed $k \geq 1$ (note that $P_{n,k}$ has a meaning as soon as $n \geq k$);
- (3) $P_{n,2} \xrightarrow[n \rightarrow \infty]{C_b} \mu^{\otimes 2}$;

where these weak convergences are with respect to continuous and bounded test functions.

Proof. Folkloric in the domain of mean field particle systems. We refer to [BCF18] and references therein. \square

Theorem 7.12 (Central limit phenomenon). *Let μ_n be as in Theorem 7.4. Then, for all measurable $f : \mathbb{C} \rightarrow \mathbb{R}$ which are C^1 in a neighborhood of the unit disc $D = \{z \in \mathbb{C} : |z| \leq 1\}$ of the complex plane,*

$$n \left[\int f d\mu_n - \mathbb{E} \int f d\mu_n \right] = \sum_{k=1}^n \left[f\left(\frac{\lambda_k}{\sqrt{n}}\right) - \mathbb{E} f\left(\frac{\lambda_k}{\sqrt{n}}\right) \right] \xrightarrow[n \rightarrow \infty]{d} \mathcal{N}\left(\frac{1}{4\pi} \|f\|_{\mathbb{H}^1(D)}^2 + \frac{1}{2} \|f\|_{\mathbb{H}^{1/2}(\partial D)}^2\right)$$

where

$$\|f\|_{\mathbb{H}^1(D)}^2 = \int_D |\nabla f|^2 dz \quad \text{and} \quad \|f\|_{\mathbb{H}^{1/2}(\partial D)}^2 = \sum_{k \in \mathbb{Z}} |k| |\hat{f}(k)|^2$$

where $\hat{f}(k)$ is the k -th Fourier coefficient of f on $\partial D = \{z \in \mathbb{C} : |z| = 1\}$, namely

$$\hat{f}(k) = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) e^{-ik\theta} d\theta.$$

Note that $\|f\|_{\mathbb{H}^{1/2}(\partial D)} = 0$ if f is analytic on a neighborhood of ∂D .

About the proof. It is known [CL95] that the cumulants of linear statistics of determinantal processes have a nice form that can be used to prove a central limit theorem. This idea is followed in [RV07] in order to produce the result, via combinatorial identities, and via reduction to polynomial test functions. The method is used for more general two-dimensional determinantal gases in [AHM11, AHM15]. \square

Theorem 7.13 (Distribution of the moduli in the Ginibre model). *Let $(\lambda_1, \dots, \lambda_n)$ be the exchangeable random vector considered in Theorem 7.1. Then the following equality in distribution holds*

$$(|\lambda_1|, \dots, |\lambda_n|) \stackrel{d}{=} (Z_{\sigma(1)}, \dots, Z_{\sigma(n)})$$

where Z_1, \dots, Z_n are independent non-negative random variables with¹²³

$$Z_k^2 \sim \text{Gamma}(k, 1), \quad 1 \leq k \leq n,$$

and where σ is a uniform random permutation of $\{1, \dots, n\}$ independent of Z_1, \dots, Z_n . Equivalently, for all symmetric bounded measurable $F : \mathbb{R}^n \rightarrow \mathbb{R}$, we have $\mathbb{E}(F(|\lambda_1|, \dots, |\lambda_n|)) = \mathbb{E}(F(Z_1, \dots, Z_n))$.

This is an equality between two exchangeable laws on \mathbb{R}^n , in other words an equality in law between two configurations of unlabeled random points in \mathbb{R} (multi-sets). Note in particular that for all $1 \leq k \leq n$, taking $F(x_1, \dots, x_n) = \sum_{\substack{i_1, \dots, i_k \\ \text{distinct}}} f(x_{i_1}) \cdots f(x_{i_k})$ gives equality of k -point correlation functions.

¹The law $\text{Gamma}(a, \lambda)$ has density $x \in \mathbb{R} \mapsto \frac{\lambda^a}{\Gamma(a)} x^{a-1} e^{-\lambda x} \mathbf{1}_{x \geq 0}$, and $\text{Gamma}(a, \lambda) * \text{Gamma}(b, \lambda) = \text{Gamma}(a+b, \lambda)$.

²Note that $(\sqrt{2}Z_k)^2 \sim \text{Gamma}(k, \frac{1}{2}) = \text{Exponential}(\frac{1}{2})^{*k} = \chi^2(2k)$ since $\chi^2(n) = \text{Gamma}(\frac{n}{2}, \frac{1}{2})$ for all $n \geq 1$.

³For $n = 1$ we recover the Box–Muller formula $|X|^2 \sim \chi^2(2) = \text{Gamma}(1, \frac{1}{2}) = \text{Exponential}(\frac{1}{2})$ with $X \sim \mathcal{N}(0, I_2)$.

Proof. From Theorem 7.1, the exchangeable random vector $(\lambda_1, \dots, \lambda_n)$ has density φ_n . It follows that the density of the exchangeable random vector $(|\lambda_1|, \dots, |\lambda_n|)$ is obtained from φ_n by integrating the phases in polar coordinates. In polar coordinates $x_k = r_k e^{i\theta_k}$, the density φ_n writes

$$(r_1, \dots, r_n, \theta_1, \dots, \theta_n) \mapsto \frac{e^{-\sum_{j=1}^n r_j^2}}{\pi^n \prod_{k=1}^n k!} \prod_{j < k} |r_j e^{i\theta_j} - r_k e^{i\theta_k}|^2.$$

Now we have, denoting Σ_n the symmetric group of permutations of $\{1, \dots, n\}$,

$$\begin{aligned} \prod_{j < k} |r_j e^{i\theta_j} - r_k e^{i\theta_k}|^2 &= \prod_{j < k} (r_j e^{i\theta_j} - r_k e^{i\theta_k}) \prod_{j < k} \overline{(r_j e^{i\theta_j} - r_k e^{i\theta_k})} \\ &= \det \left[r_j^{k-1} e^{i(k-1)\theta_j} \right]_{1 \leq j, k \leq n} \det \left[r_j^{k-1} e^{-i(k-1)\theta_j} \right]_{1 \leq j, k \leq n} \\ &= \left(\sum_{\sigma \in \Sigma_n} (-1)^{\text{sign}(\sigma)} \prod_{j=1}^n r_j^{\sigma(j)-1} e^{i(\sigma(j)-1)\theta_j} \right) \left(\sum_{\sigma' \in \Sigma_n} (-1)^{\text{sign}(\sigma')} \prod_{j=1}^n r_j^{\sigma'(j)-1} e^{-i(\sigma'(j)-1)\theta_j} \right) \\ &= \sum_{\sigma, \sigma' \in \Sigma_n} (-1)^{\text{sign}(\sigma) + \text{sign}(\sigma')} \prod_{j=1}^n r_j^{\sigma(j) + \sigma'(j) - 2} e^{i((\sigma(j) - \sigma'(j))\theta_j)}. \end{aligned}$$

If we integrate the phases, we note that only the terms with $\sigma = \sigma'$ contribute to the result, namely

$$\int_0^{2\pi} \dots \int_0^{2\pi} \prod_{j < k} |r_j e^{i\theta_j} - r_k e^{i\theta_k}|^2 d\theta_1 \dots d\theta_n = (2\pi)^n \sum_{\sigma \in \Sigma_n} \prod_{j=1}^n r_j^{2(\sigma(j)-1)} = (2\pi)^n \text{per} \left[r_j^{2k} \right]_{1 \leq j, k \leq n}$$

where “per” stands for “permanent”. Therefore, the (exchangeable) density of the moduli is given by

$$\int_0^{2\pi} \dots \int_0^{2\pi} \frac{e^{-\sum_{j=1}^n r_j^2}}{\pi^n \prod_{k=1}^n k!} \prod_{j < k} |r_j e^{i\theta_j} - r_k e^{i\theta_k}|^2 d\theta_1 \dots d\theta_n = \text{perm} \left[\frac{2}{k!} r_j^{2k} e^{-r_j^2} \right]_{1 \leq j, k \leq n}.$$

But if $f_1, \dots, f_n : \mathbb{R} \rightarrow \mathbb{R}$ are probability density functions then $(x_1, \dots, x_n) \mapsto \text{perm}[f_j(x_k)]_{1 \leq j, k \leq n}$ is the density of the random vector $(X_{\sigma(1)}, \dots, X_{\sigma(n)})$ where X_1, \dots, X_n are independent real random variables with densities f_1, \dots, f_n and where σ is a uniform random permutation of $\{1, \dots, n\}$ independent of X_1, \dots, X_n . Also the desired result follows from the formula above and the fact that for all $1 \leq k \leq n$, a non-negative random variable Z_k has density $r \mapsto \frac{2}{k!} r^{2k} e^{-r^2} \mathbf{1}_{r \geq 0}$ if and only if $Z_k^2 \sim \text{Gamma}(k, 1)$.

This proof, essentially due to Kostlan [Kos92], see also [Rid03], relies on the determinantal nature of φ_n in (2.5), and remains usable for general determinantal processes, see for instance [HKPV09]. \square

Theorem 7.14 (Spectral radius). *With the notation of Theorem 7.1, almost surely*

$$\rho_n = \max_{1 \leq k \leq n} \frac{|\lambda_k|}{\sqrt{n}} \xrightarrow[n \rightarrow \infty]{} 1.$$

Moreover, denoting $\kappa_n = \log \frac{n}{2\pi} - 2 \log(\log(n))$,

$$\sqrt{4n\kappa_n} \left(\rho_n - 1 - \sqrt{\frac{\kappa_n}{4n}} \right) \xrightarrow[n \rightarrow \infty]{d} \text{Gumbel}^4.$$

Note the the second statement (Gumbel fluctuation) implies that $\rho_n \xrightarrow[n \rightarrow \infty]{} 1$ in probability.

Regarding the convergence, see [BCGZ20] for a random analytic function point of view, related to the central limit theorem. Regarding the fluctuation, see [Joh07, Ben10] for an interpolation with the Tracy–Widom fluctuation at the edge of GUE.

Idea of proof. By Theorem 7.13 Since $Z_k^2 \stackrel{d}{=} E_1 + \dots + E_k$ where E_1, \dots, E_k are independent and identically distributed exponential random variables of unit mean, we get, for every $r > 0$,

$$\mathbb{P}(\rho_n \leq \sqrt{nr}) = \prod_{1 \leq k \leq n} \mathbb{P} \left(\frac{E_1 + \dots + E_k}{n} \leq r^2 \right).$$

By the law of large numbers, this tends as $n \rightarrow \infty$ to 0 or 1 depending on the position of r with respect to 1. Moreover the central limit theorem suggests that ρ_n behaves as $n \rightarrow \infty$ as the maximum of

⁴If $X \sim \text{Exp}(1)$ then $-\log(X)$ has Gumbel law and cumulative distribution function $\mathbb{P}(-\log(X) \leq x) = e^{-e^{-x}}$.

independent and identically distributed Gaussian random variables, a situation for which it is known that the fluctuation follows the Gumbel law. The full proof is in [Rid03] and involves crucially a quantitative central limit theorem and the Borel–Cantelli lemma. The approach is robust and remains valid beyond the Ginibre gas, for determinantal gases, see for instance [CP14, JQ17, GZ18] and references therein. \square

Remark 7.15 (Real or quaternionic Ginibre model). *How about an analogue of Theorem 7.1 when the entries of \mathbf{M} are real Gaussian or real quaternionic Gaussian instead of complex Gaussian? Some answers are already in [Gin65]. In these cases, the density of the eigenvalues can be computed but it is not the beta gas (2.4) with $\beta \in \{1, 4\}$. This is in contrast with the $G(O/U/S)E$ triplet of the Hermitian random matrix Dysonian universe [Dys62c]. See for instance [Ede97, Dub18b] and references therein.*

Remark 7.16 (Large deviations). *The large deviations principle for the beta Ginibre gas (2.5) was established in [HP00, PH98], using a method inspired from [BAG08], itself inspired from [Voi93, Voi94]. It does not rely on the determinantal structure, and allows to extend Theorem 7.8 to all $\beta > 0$.*

7.2. More determinantal models. It is well known that the ratio of two independent real standard Gaussian random variables follows a Cauchy distribution. The following theorem can be seen as a matrix version of this phenomenon.

Theorem 7.17 (Forrester–Krishnapur spherical ensemble). *Let \mathbf{M}_1 and \mathbf{M}_2 be independent copies of the Ginibre random matrix defined in (7.1). Then as an exchangeable random vector of \mathbb{C}^n , the eigenvalues of $\mathbf{M}_1\mathbf{M}_2^{-1}$ have density*

$$(z_1, \dots, z_n) \in \mathbb{C}^n \mapsto \frac{1}{Z_n} \frac{\prod_{j < k} |z_j - z_k|^2}{\prod_{j=1}^n (1 + |z_j|^2)^{n+1}}.$$

This corresponds to the beta gas (2.4) with $V = \frac{1}{2} \frac{n+1}{n} \log(1 + |\cdot|^2)$ and $\beta = 2$. Moreover its push-forward on the Riemann sphere using inverse stereographic projection is the uniform law on the sphere.

See [Kri09] and [HKPV09, FK09, For10] for a proof. The set of singular $n \times n$ complex matrices is a hyper-surface of zero Lebesgue measure in \mathbb{C}^{n^2} and therefore, almost surely, the Ginibre random matrix \mathbf{M} in (7.1) is invertible (its law is absolutely continuous with respect to the Lebesgue measure on \mathbb{C}^{n^2}).

From Theorem 1.1, the equilibrium measure of the gas is heavy tailed with density

$$z \in \mathbb{C} \mapsto \frac{1}{\pi(1 + |z|^2)^2}.$$

A large deviations principle for the empirical measure associated to the Coulomb gas is proved in [Har12] in relation with the sphere. The convergence of the empirical measure is also considered in [Bor11]. The fluctuation at the edge is discussed in [CP14] and studied in [JQ17] by using the idea of Kostlan behind Theorem 7.13 thanks to the determinantal structure. A beta version of the model is considered in [CMMOC18] and studied using transportation of measure.

Theorem 7.18 (Życzkowski–Sommers ensemble). *Let $\mathbf{U} = (\mathbf{U}_{j,k})_{1 \leq j, k \leq m}$ be a random $m \times m$ unitary matrix following the (Haar) uniform law on this compact group of matrices. Then, for all $1 \leq n < m$, as an exchangeable random vector of \mathbb{C}^n , the eigenvalues of the truncation $(\mathbf{U}_{j,k})_{1 \leq j, k \leq n}$ have density*

$$(z_1, \dots, z_n) \in \mathbb{C}^n \mapsto \frac{\prod_{j=1}^n (1 - |z_j|^2)^{m-n-1}}{Z_n} \prod_{1 \leq j < k \leq n} |z_j - z_k|^2.$$

This corresponds to the beta gas (2.4) with $V = V_{n,m} = \frac{m-n-1}{n} \log \frac{1}{1-|z|^2}$ and $\beta = 2$.

See [ZS00] for a proof, and [FK09] for the special case $m \geq 2n$ and a link with the pseudo-sphere and Schur transformation. Following [PR05] and references therein, if $\lim_{m,n \rightarrow \infty} \frac{n}{m} = \alpha \in (0, 1)$ then the empirical measure converges towards the heavy tailed probability measure with density

$$z \in \mathbb{C} \mapsto \frac{(1 - \alpha)}{\pi\alpha(1 - |z|^2)^2} \mathbf{1}_{|z| \leq \sqrt{\alpha}}.$$

In a sense this law interpolates between the uniform law on the unit disc ($\alpha \rightarrow 0$ after scaling by $\sqrt{\alpha}$) and the uniform law on the unit circle ($\alpha \rightarrow 1$). A large deviations principle is obtained in [PR05], concentration inequalities are derived in [MS19], while the fluctuation at the edge is studied in [JQ17].

Theorem 7.19 (Product of Ginibre random matrices). *Let $m \geq 1$ and let $\mathbf{M}_1, \dots, \mathbf{M}_m$ be independent and identically distributed copies of the $n \times n$ in (7.1). Then, as an exchangeable random vector of \mathbb{C}^n , the eigenvalues of the scaled product $n^{-\frac{m}{2}} \mathbf{M}_1 \cdots \mathbf{M}_m$ have density*

$$\frac{\prod_{j=1}^n w_m(\sqrt{n}|z_j|)}{Z_n} \prod_{j < k} |z_j - z_k|^2$$

where w_k is the Meijer G-function given by the recursive formula

$$w_1(z) = e^{-|z|^2} \quad \text{and} \quad w_k(z) = 2\pi \int_0^\infty w_{k-1}\left(\frac{z}{r}\right) \frac{e^{-r^2}}{r} dr.$$

This corresponds to the beta gas (2.4) with $V = V_{n,m} = -\frac{1}{n} \log w_m(\sqrt{n}\bullet)$ and $\beta = 2$.

See [AB12] for a proof. Following [GT11, Bor11], it converges to the equilibrium measure with density

$$z \in \mathbb{C} \mapsto \frac{|z|^{\frac{2}{m}-2}}{m\pi} \mathbf{1}_{|z| \leq 1}.$$

We recover the uniform law on the unit disc when $m = 1$. The edge fluctuation is considered in [JQ17].

Remark 7.20 (Determinantal gases and random normal matrices). *An $n \times n$ complex matrix is normal when $MM^* = M^*M$. The random matrices in theorems 7.1, 7.17, 7.18, 7.19 are not normal. Let us comment now on models of normal random matrices. Let \mathcal{N}_n be the hyper-surface of \mathbb{C}^{n^2} of all $n \times n$ normal matrices. Let $V : \mathbb{C} \rightarrow \mathbb{R}$ be \mathcal{C}^2 and such that $V(z) \geq c \log(1 + |z|^2)$ for some constant $c > 0$. Following [CZ98, EF05], let us consider the probability measure on \mathcal{N}_n with density proportional to $M \mapsto e^{-n \text{Trace}(V(M))}$ with respect to the Hausdorff measure on \mathcal{N}_n . This produces random normal matrices, and their eigenvalues, seen as an exchangeable random vector, have density given by the gas (2.4) with $\beta = 2$. This random (normal) matrix model is referred to as the random normal matrix model. The fluctuation of the empirical measure is studied in [AHM15, AHM11], while the fluctuation at the edge is studied in [CP14, JQ17, GZ18].*

The power of a Ginibre matrix has also a nice determinantal structure, see [Dub18a].

8. COMMENTS AND OPEN PROBLEMS

We have skipped several important old and new results on Coulomb gases. The main themes are local versus global, first versus second order, macroscopics versus microscopics, non-universal versus universal.

Universality. The first order global convergence $\lim_{n \rightarrow \infty} \mu_n = \mu_V$, that we call macroscopics, is not universal in the sense that the limit μ_V still depends on V . The second order convergence provided by the central limit theorem (5.1) is universal in the sense that the limit should not depend on V . Similarly, for a two-dimensional Coulomb gas with radial confining potential V , the limit of the edge depends on V but its fluctuation does not and is universal. Universality emerges often in a second order asymptotic analysis, as for classical limit theorems of probability theory.

Microscopics. A second order analysis corresponds to the asymptotic analysis of $n(\mu_n - \nu_V)$ as $n \rightarrow \infty$. This can be seen as a microscopic analysis while the convergence $\mu_n \rightarrow \mu_V$ is a macroscopic analysis. This corresponds to a second order Taylor formula for the quadratic form \mathcal{E}_V , in other words in a special factorization, leading to a new object called the *renormalized energy*. This was the subject of an series of works by Étienne Sandier and Sylvia Serfaty, and by Sylvia Serfaty and other co-authors. See for instance [Ser18b, Ser18a, LS17, Ser15] and references therein. An outcome of this refined analysis is a second order asymptotics for the free energy. More precisely, recall that the Boltzmann–Shannon entropy of the Boltzmann–Gibbs measure P_n in (2.2) is defined by

$$\mathcal{S}(P_n) = - \int_{(\mathbb{R}^d)^n} f_n(x_1, \dots, x_n) \log f_n(x_1, \dots, x_n) dx_1 \cdots dx_n$$

where f_n is the density of P_n . Its Helmholtz free energy is given by

$$\int E_n dP_n - \frac{\mathcal{S}(P_n)}{\beta} = - \frac{\log Z_n}{\beta},$$

see [Cha15]. Now following [LS17], if μ_V has density f_V with a finite Boltzmann–Shannon entropy

$$\mathcal{S}(\mu_V) = - \int_{\mathbb{R}^d} f_V \log f_V dx,$$

then we have an asymptotic expansion of the free energy as $n \rightarrow \infty$ as

$$-\frac{\log Z_n}{\beta} = \begin{cases} \frac{n^2}{2} \mathcal{E}_V(\mu_V) - \frac{n \log n}{4} + n(c_\beta + c'_\beta \mathcal{S}(\mu_V)) + no_n(1) & \text{if } d = 2 \\ \frac{n^2}{2} \mathcal{E}_V(\mu_V) + n(c_{\beta,d,V} + c'_{\beta} \mathcal{S}(\mu_V)) + no_n(1) & \text{if } d \neq 2 \end{cases},$$

where $c_\beta, c'_\beta, c_{\beta,d,V}$ are constants which can be made explicit.

Edge. The most elementary open question related to Coulomb gases is perhaps the law of fluctuation at the edge, even in the case of rotationally invariant confining potential for arbitrary values of d and β . The Gumbel fluctuation is known for instance to be universal for a class of two dimensional ($d = 2$) determinantal ($\beta = 2$) Coulomb gases with radial confining potential, see [CP14]. The same question for arbitrary β is open, and the same question for arbitrary dimension $d \geq 3$ and $\beta > 0$ is also open.

Crystallization. A conjecture related to Coulomb gases is the emergence of rigid structures at low temperatures. This is known as crystallization and was proved in special cases, for instance for one-dimensional Coulomb gases. See for instance [Ser15, BL15, Ser18b, Ser18a, PS20] and references therein.

More. Among all the important results on Coulomb gases that we have not yet mentioned, we may cite the approximate transport maps for universality considered in [FG16, BFG15], the rigidity analysis for hierarchical Coulomb gases considered in [Cha19b], the local density for two-dimensional Coulomb gases considered in [BBNY17], the Dobrushin–Lanford–Ruelle equations considered in [DHLM21], the Coulomb gas properties on the sphere considered in [BH19], the local laws and rigidity considered in [AS21], the quasi-Monte-Carlo method on the sphere considered in [Ber19b], and the Berezinskii–Kosterlitz–Thouless transition [KP17, GS20].

ACKNOWLEDGMENTS

We would like to thank David García-Zelada and Kilian Rashel for their helpful remarks.

REFERENCES

- [AB12] G. AKEMANN & Z. BURDA – “Universal microscopic correlation functions for products of independent Ginibre matrices”, *J. Phys. A* **45** (2012), no. 46, p. 465201, 18.
- [AB19] G. AKEMANN & S.-S. BYUN – “The high temperature crossover for general 2D Coulomb gases”, *J. Stat. Phys.* **175** (2019), no. 6, p. 1043–1065 (English).
- [ABDF11] G. AKEMANN, J. BAIK & P. DI FRANCESCO (éds.) – *The Oxford handbook of random matrix theory*, Oxford University Press, Oxford, 2011.
- [AGZ10] G. W. ANDERSON, A. GUIONNET & O. ZEITOUNI – *An introduction to random matrices*, Cambridge Studies in Advanced Mathematics, vol. 118, Cambridge University Press, Cambridge, 2010.
- [AHM11] Y. AMEUR, H. K. HEDENMALM & N. MAKAROV – “Fluctuations of eigenvalues of random normal matrices”, *Duke Math. J.* **159** (2011), no. 1, p. 31–81.
- [AHM15] Y. AMEUR, H. HEDENMALM & N. MAKAROV – “Random normal matrices and Ward identities”, *Ann. Probab.* **43** (2015), no. 3, p. 1157–1201.
- [Ame21] Y. AMEUR – “A localization theorem for the planar Coulomb gas in an external field”, *Electronic Journal of Probability* **26** (2021), no. none, p. 1 – 21.
- [AS19] S. ARMSTRONG & S. SERFATY – “Thermal approximation of the equilibrium measure and obstacle problem”, preprint arXiv:1912.13018, 2019.
- [AS21] S. ARMSTRONG & S. SERFATY – “Local laws and rigidity for Coulomb gases at any temperature”, *Ann. Probab.* **49** (2021), no. 1, p. 46–121 (English).
- [BAG97] G. BEN AROUS & A. GUIONNET – “Large deviations for Wigner’s law and Voiculescu’s non-commutative entropy”, *Probab. Theory Related Fields* **108** (1997), no. 4, p. 517–542.
- [BAG08] G. BEN AROUS & A. GUIONNET – “The spectrum of heavy tailed random matrices”, *Comm. Math. Phys.* **278** (2008), no. 3, p. 715–751.
- [BBL96] E. BOGOMOLNY, O. BOHIGAS & P. LEBOEUF – “Quantum chaotic dynamics and random polynomials”, *J. Statist. Phys.* **85** (1996), no. 5-6, p. 639–679.
- [BBLu92] E. BOGOMOLNY, O. BOHIGAS & P. LEBEUF – “Distribution of roots of random polynomials”, *Phys. Rev. Lett.* **68** (1992), no. 18, p. 2726–2729.
- [BBNY17] R. BAUERSCHMIDT, P. BOURGADE, M. NIKULA & H.-T. YAU – “Local density for two-dimensional one-component plasma”, *Comm. Math. Phys.* **356** (2017), no. 1, p. 189–230.
- [BBNY19] ———, “The two-dimensional Coulomb plasma: quasi-free approximation and central limit theorem”, *Adv. Theor. Math. Phys.* **23** (2019), no. 4, p. 841–1002.
- [BCF18] F. BOLLEY, D. CHAFAÏ & J. FONTBONA – “Dynamics of a planar Coulomb gas”, *Ann. Appl. Probab.* **28** (2018), no. 5, p. 3152–3183.

- [BCGZ20] C. BORDENAVE, D. CHAFAÏ & D. GARCÍA-ZELADA – “Convergence of the spectral radius of a random matrix through its characteristic polynomial”, preprint arXiv:2012.05602 to appear in *Probability Theory and Related Fields*, 2020.
- [BD20] P. BOURGADE & G. DUBACH – “The distribution of overlaps between eigenvectors of Ginibre matrices”, *Probab. Theory Relat. Fields* **177** (2020), no. 1-2, p. 397–464 (English).
- [Ben10] M. BENDER – “Edge scaling limits for a family of non-Hermitian random matrix ensembles”, *Probab. Theory Related Fields* **147** (2010), no. 1-2, p. 241–271.
- [Ber18a] R. BERMAN – “The Coulomb gas, potential theory and phase transitions”, preprint arXiv:1811.10249, 2018.
- [Ber18b] R. J. BERMAN – “On large deviations for Gibbs measures, mean energy and gamma-convergence”, *Constr. Approx.* **48** (2018), no. 1, p. 3–30.
- [Ber19a] R. BERMAN – “Sharp deviation inequalities for the 2D Coulomb gas and Quantum hall states, I”, preprint arXiv:1906.08529v1, 2019.
- [Ber19b] R. J. BERMAN – “The spherical ensemble and quasi-Monte-Carlo designs”, preprint arXiv:1906.08533v1, 2019.
- [BFG15] F. BEKERMANN, A. FIGALLI & A. GUIONNET – “Transport maps for β -matrix models and universality”, *Comm. Math. Phys.* **338** (2015), no. 2, p. 589–619.
- [BG99] T. BODINEAU & A. GUIONNET – “About the stationary states of vortex systems”, *Ann. Inst. H. Poincaré Probab. Statist.* **35** (1999), no. 2, p. 205–237.
- [BGG17] A. BORODIN, V. GORIN & A. GUIONNET – “Gaussian asymptotics of discrete β -ensembles”, *Publ. Math. Inst. Hautes Études Sci.* **125** (2017), p. 1–78.
- [BGL14] D. BAKRY, I. GENTIL & M. LEDOUX – *Analysis and geometry of Markov diffusion operators*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 348, Springer, Cham, 2014.
- [BGZ18] R. BUTEZ & D. GARCIA-ZELADA – “Extremal particles of two-dimensional Coulomb gases and random polynomials on a positive background”, preprint arXiv:1811.12225 to appear in *Annals of Applied Probability*, 2018.
- [BGZNW21] R. BUTEZ, D. GARCÍA-ZELADA, A. NISHRY & A. WENNMAN – “Universality for outliers in weakly confined coulomb-type systems”, preprint arXiv:2104.03959v1, 2021.
- [BH19] C. BELTRÁN & A. HARDY – “Energy of the Coulomb gas on the sphere at low temperature”, *Arch. Ration. Mech. Anal.* **231** (2019), no. 3, p. 2007–2017.
- [BHS19] S. V. BORODACHOV, D. P. HARDIN & E. B. SAFF – *Discrete energy on rectifiable sets*, New York, NY: Springer, 2019 (English).
- [BL15] X. BLANC & M. LEWIN – “The crystallization conjecture: a review”, *EMS Surv. Math. Sci.* **2** (2015), no. 2, p. 225–306.
- [BLS18] F. BEKERMANN, T. LEBLÉ & S. SERFATY – “CLT for fluctuations of β -ensembles with general potential”, *Electron. J. Probab.* **23** (2018), p. Paper no. 115, 31.
- [Bor11] C. BORDENAVE – “On the spectrum of sum and product of non-Hermitian random matrices”, *Electronic Communications in Probability* **16** (2011), p. 104–113.
- [Bou15] S. BOUCKSOM – “Limite thermodynamique et théorie du potentiel”, *Gaz. Math.* (2015), no. 146, p. 16–26.
- [Bre67] M. BRELOT – *Lectures on potential theory*, Notes by K. N. Gowrisankaran and M. K. Venkatesha Murthy. Second edition, revised and enlarged with the help of S. Ramaswamy. Tata Institute of Fundamental Research Lectures on Mathematics, No. 19, Tata Institute of Fundamental Research, Bombay, 1967.
- [But17] R. BUTEZ – “Polynômes aléatoires, gaz de Coulomb, et matrices aléatoires (Random polynomials, Coulomb gas and random matrices)”, Thèse, Paris-Dauphine / PSL, 2017.
- [BW11] O. BOHIGAS & H. A. WEIDENMÜLLER – “History—an overview”, in *The Oxford handbook of random matrix theory*, Oxford Univ. Press, Oxford, 2011, p. 15–39.
- [CF19] D. CHAFAÏ & G. FERRÉ – “Simulating Coulomb and Log-Gases with Hybrid Monte Carlo Algorithms”, *J. Stat. Phys.* **174** (2019), no. 3, p. 692–714.
- [CFS21] D. CHAFAÏ, G. FERRÉ & G. STOLTZ – “Coulomb gases under constraint: some theoretical and numerical results”, *SIAM J. Math. Anal.* **53** (2021), no. 1, p. 181–220 (English).
- [CGZ14] D. CHAFAÏ, N. GOZLAN & P.-A. ZITT – “First-order global asymptotics for confined particles with singular pair repulsion”, *Ann. Appl. Probab.* **24** (2014), no. 6, p. 2371–2413.
- [CGZJ20a] D. CHAFAÏ, D. GARCÍA-ZELADA & P. JUNG – “At the edge of a one-dimensional jellium”, preprint 2012.04633 to appear in *Bernoulli*, 2020.
- [CGZJ20b] D. CHAFAÏ, D. GARCÍA-ZELADA & P. JUNG – “Macroscopic and edge behavior of a planar jellium”, *Journal of Mathematical Physics* **61** (2020), no. 3, p. 033304.
- [Cha14] D. CHAFAÏ – “Wigner about level spacing and Wishart”, blogpost <http://djalil.chafai.net/blog/2014/09/26/wigner-about-level-spacing-and-wishart/>, 2014.
- [Cha15] D. CHAFAÏ – “From Boltzmann to random matrices and beyond”, *Ann. Fac. Sci. Toulouse Math. (6)* **24** (2015), no. 4, p. 641–689.
- [Cha19a] D. CHAFAÏ – “An unexpected distribution”, blogpost <https://djalil.chafai.net/blog/2019/12/15/an-unexpected-distribution/>, 2019.
- [Cha19b] S. CHATTERJEE – “Rigidity of the three-dimensional hierarchical Coulomb gas”, *Probab. Theory Related Fields* **175** (2019), no. 3-4, p. 1123–1176.
- [CHM18] D. CHAFAÏ, A. HARDY & M. MAÏDA – “Concentration for Coulomb gases and Coulomb transport inequalities”, *J. Funct. Anal.* **275** (2018), no. 6, p. 1447–1483.
- [Cho54] G. CHOQUET – “Theory of capacities”, *Ann. Inst. Fourier, Grenoble* **5** (1953–1954), p. 131–295 (1955).

- [Cho90] ———, “La vie et l’oeuvre de Marcel Brelot (1903-1987)”, *Cahiers du séminaire d’histoire des mathématiques* **11** (1990), p. 1–31 (fre).
- [CL95] O. COSTIN & J. L. LEBOWITZ – “Gaussian fluctuation in random matrices”, *Phys. Rev. Lett.* **75** (1995), no. 1, p. 69–72.
- [CL20] D. CHAFAÏ & J. LEHEC – “On Poincaré and logarithmic Sobolev inequalities for a class of singular Gibbs measures”, in *Geometric aspects of functional analysis. Vol. I*, Lecture Notes in Math., vol. 2256, Springer, Cham, [2020] ©2020, p. 219–246.
- [CLMP92] E. CAGLIOTI, P.-L. LIONS, C. MARCHIORO & M. PULVIRENTI – “A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description”, *Comm. Math. Phys.* **143** (1992), no. 3, p. 501–525.
- [CMMOC18] T. CARROLL, J. MARZO, X. MASSANEDA & J. ORTEGA-CERDÀ – “Equidistribution and β -ensembles”, *Ann. Fac. Sci. Toulouse Math. (6)* **27** (2018), no. 2, p. 377–387.
- [CP14] D. CHAFAÏ & S. PÉCHÉ – “A note on the second order universality at the edge of Coulomb gases on the plane”, *J. Stat. Phys.* **156** (2014), no. 2, p. 368–383.
- [CZ98] L.-L. CHAU & O. ZABORONSKY – “On the structure of correlation functions in the normal matrix model”, *Comm. Math. Phys.* **196** (1998), no. 1, p. 203–247.
- [de 37] C.-J. DE LA VALLÉE POUSSIN – “Les nouvelles méthodes de la théorie du potentiel et le problème généralisé de Dirichlet.”, Paris: Hermann & Cie. 47 p. (1937)., 1937.
- [DE02] I. DUMITRIU & A. EDELMAN – “Matrix models for beta ensembles”, *J. Math. Phys.* **43** (2002), no. 11, p. 5830–5847.
- [Dei99] P. A. DEIFT – *Orthogonal polynomials and random matrices: a Riemann-Hilbert approach*, Courant Lecture Notes in Mathematics, vol. 3, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [DG09] P. DEIFT & D. GIOEV – *Random matrix theory: invariant ensembles and universality*, Courant Lecture Notes in Mathematics, vol. 18, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2009.
- [DHLM21] D. DEREUDRE, A. HARDY, T. LEBLÉ & M. MAÏDA – “DLR equations and rigidity for the sine-beta process”, *Comm. Pure Appl. Math.* **74** (2021), no. 1, p. 172–222.
- [DLR20] P. DUPUIS, V. LASCHOS & K. RAMANAN – “Large deviations for configurations generated by Gibbs distributions with energy functionals consisting of singular interaction and weakly confining potentials”, *Electron. J. Probab.* **25** (2020), p. 41 (English), Id/No 46.
- [DM78] C. DELLACHERIE & P.-A. MEYER – *Probabilities and potential*, North-Holland Mathematics Studies, vol. 29, North-Holland Publishing Co., Amsterdam, 1978.
- [Doo01] J. L. DOOB – *Classical potential theory and its probabilistic counterpart*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1984 edition.
- [DRSV17] B. DUPLANTIER, R. RHODES, S. SHEFFIELD & V. VARGAS – “Log-correlated Gaussian fields: an overview”, in *Geometry, analysis and probability. In Honor of Jean-Michel Bismut. Selected papers based on the presentations at the conference ‘Control, index, traces and determinants – the journey of a probabilist’, Orsay, France, May 27–31, 2013*, Basel: Birkhäuser/Springer, 2017, p. 191–216 (English).
- [Dub18a] G. DUBACH – “Powers of Ginibre eigenvalues”, *Electron. J. Probab.* **23** (2018), p. Paper No. 111, 31.
- [Dub18b] ———, “Symmetries of the Quaternionic Ginibre Ensemble”, preprint arXiv:1811.03724, 2018.
- [Dys62a] F. J. DYSON – “A Brownian-motion model for the eigenvalues of a random matrix”, *J. Mathematical Phys.* **3** (1962), p. 1191–1198.
- [Dys62b] ———, “Statistical theory of the energy levels of complex systems. I”, *J. Mathematical Phys.* **3** (1962), p. 140–156.
- [Dys62c] ———, “The threefold way. Algebraic structure of symmetry groups and ensembles in quantum mechanics”, *J. Mathematical Phys.* **3** (1962), p. 1199–1215.
- [Ede97] A. EDELMAN – “The probability that a random real Gaussian matrix has k real eigenvalues, related distributions, and the circular law”, *J. Multivariate Anal.* **60** (1997), no. 2, p. 203–232.
- [EF05] P. ELBAU & G. FELDER – “Density of eigenvalues of random normal matrices”, *Comm. Math. Phys.* **259** (2005), no. 2, p. 433–450.
- [ER05] A. EDELMAN & N. R. RAO – “Random matrix theory”, *Acta Numer.* **14** (2005), p. 233–297.
- [EY17] L. ERDŐS & H.-T. YAU – *A dynamical approach to random matrix theory*, Courant Lecture Notes in Mathematics, vol. 28, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2017.
- [FG16] A. FIGALI & A. GUIONNET – “Universality in several-matrix models via approximate transport maps”, *Acta Math.* **217** (2016), no. 1, p. 81–176.
- [FK09] P. J. FORRESTER & M. KRISHNAPUR – “Derivation of an eigenvalue probability density function relating to the poincaré disk”, *J. of Physics A: Math. and Theor.* **42** (2009), no. 38, p. 385204.
- [For10] P. J. FORRESTER – *Log-gases and random matrices*, London Mathematical Society Monographs Series, vol. 34, Princeton University Press, Princeton, NJ, 2010.
- [Fro35] O. FROSTMAN – “Potentiel d’Équilibre et Capacité des Ensembles”, Thèse, Faculté des sciences de Lund, 1935.
- [FW08] P. J. FORRESTER & S. O. WARNAAR – “The importance of the Selberg integral”, *Bull. Am. Math. Soc., New Ser.* **45** (2008), no. 4, p. 489–534 (English).

- [Gar19] D. GARCÍA-ZELADA – “A large deviation principle for empirical measures on Polish spaces: application to singular Gibbs measures on manifolds”, *Ann. Inst. Henri Poincaré, Probab. Stat.* **55** (2019), no. 3, p. 1377–1401 (English).
- [Gin65] J. GINIBRE – “Statistical ensembles of complex, quaternion, and real matrices”, *J. Mathematical Phys.* **6** (1965), p. 440–449.
- [Gir05] S. M. GIRVIN – “Introduction to the fractional quantum Hall effect”, in *The quantum Hall effect*, Prog. Math. Phys., vol. 45, Birkhäuser, Basel, 2005, p. 133–162.
- [GS20] C. GARBAN & A. SEPÚLVEDA – “Statistical reconstruction of the Gaussian free field and KT transition”, preprint arXiv:2002.12284, 2020.
- [GT11] F. GÖTZE & A. TIKHOMIROV – “On the asymptotic spectrum of products of independent random matrices”, preprint arXiv:1012.2710v3, 2011.
- [GZ18] D. GARCÍA-ZELADA – “Edge fluctuations for random normal matrix ensembles”, preprint arXiv:1812.11170v3, 2018.
- [GZ19a] D. GARCÍA-ZELADA – “Concentration for Coulomb gases on compact manifolds”, *Electron. Commun. Probab.* **24** (2019), p. Paper No. 12, 18.
- [GZ19b] D. GARCÍA-ZELADA – “Aspects géométriques et probabilistes des gaz de Coulomb (Geometric and probabilistic aspects of Coulomb gases)”, Thèse, Paris-Dauphine / PSL, 2019.
- [Har12] A. HARDY – “A note on large deviations for 2D Coulomb gas with weakly confining potential”, *Electron. Commun. Probab.* **17** (2012), p. no. 19, 12.
- [Hel14] L. L. HELMS – *Potential theory*, second éd., Universitext, Springer, London, 2014.
- [Hir78] F. HIRSCH – “Opérateurs carré du champ (d’après J. P. Roth)”, in *Séminaire Bourbaki, 29e année (1976/77)*, Lecture Notes in Math., vol. 677, Springer, Berlin, 1978, p. Exp. No. 501, pp. 167–182.
- [HKPV09] J. B. HOUGH, M. KRISHNAPUR, Y. PERES & B. VIRÁG – *Zeros of Gaussian analytic functions and determinantal point processes*, University Lecture Series, vol. 51, AMS, Providence, RI, 2009.
- [HL21] A. HARDY & G. LAMBERT – “CLT for circular beta-ensembles at high temperature”, *J. Funct. Anal.* **280** (2021), no. 7, p. Paper No. 108869, 40.
- [HP00] F. HIAI & D. PETZ – *The semicircle law, free random variables and entropy*, Mathematical Surveys and Monographs, vol. 77, American Mathematical Society, Providence, RI, 2000.
- [Hug06] R. HUGHES – “Theoretical practice: the Bohm-Pines quartet”, *Perspectives on science* **14** (2006), no. 4, p. 457–524.
- [Hwa86] C.-R. HWANG – “A brief survey on the spectral radius and the spectral distribution of large random matrices with i.i.d. entries”, in *Random matrices and their applications (Brunswick, Maine, 1984)*, Contemp. Math., vol. 50, Amer. Math. Soc., Providence, RI, 1986, p. 145–152.
- [Joh98] K. JOHANSSON – “On fluctuations of eigenvalues of random Hermitian matrices”, *Duke Math. J.* **91** (1998), no. 1, p. 151–204.
- [Joh07] ———, “From Gumbel to Tracy-Widom”, *Probab. Theory Related Fields* **138** (2007), no. 1-2, p. 75–112.
- [JQ17] T. JIANG & Y. QI – “Spectral radii of large non-Hermitian random matrices”, *J. Theoret. Probab.* **30** (2017), no. 1, p. 326–364.
- [Kel67] O. D. KELLOGG – “Foundations of potential theory.”, Berlin-Heidelberg-New York: Springer-Verlag 1967. IX, 384 p. with 30 fig. (1967)., 1967.
- [Kos92] E. KOSTLAN – “On the spectra of Gaussian matrices”, *Linear Algebra Appl.* **162/164** (1992), p. 385–388, Directions in matrix theory (Auburn, AL, 1990).
- [KP17] V. KHARASH & R. PELED – “The Fröhlich-Spencer proof of the Berezinskii-Kosterlitz-Thouless transition”, preprint arXiv:1711.04720, 2017.
- [Kri09] M. KRISHNAPUR – “From random matrices to random analytic functions”, *Ann. Probab.* **37** (2009), no. 1, p. 314–346.
- [KS11] B. A. KHORUZHENKO & H.-J. SOMMERS – “Non-Hermitian ensembles”, in *The Oxford handbook of random matrix theory*, Oxford Univ. Press, Oxford, 2011, p. 376–397.
- [KZ13] Z. KABLUCHKO & D. ZAPOROZHETS – “Roots of random polynomials whose coefficients have logarithmic tails”, *Ann. Probab.* **41** (2013), no. 5, p. 3542–3581.
- [KZ14] ———, “Asymptotic distribution of complex zeros of random analytic functions”, *Ann. Probab.* **42** (2014), no. 4, p. 1374–1395.
- [LACTMS19] B. LACROIX-A-CHEZ-TOINE, S. N. MAJUMDAR & G. SCHEHR – “Rotating trapped fermions in two dimensions and the complex ginibre ensemble: Exact results for the entanglement entropy and number variance”, *Phys. Rev. A* **99** (2019), p. 021602.
- [Lan72] N. S. LANDKOF – *Foundations of modern potential theory*, Springer-Verlag, New York-Heidelberg, 1972, Translated from the Russian by A. P. Doohovskoy, Die Grundlehren der mathematischen Wissenschaften, Band 180.
- [Lau87] R. B. LAUGHLIN – “Elementary theory: the incompressible quantum fluid”, in *The Quantum Hall Effect* (R. E. Prange & S. M. Girvin, éd.), Springer US, New York, NY, 1987, p. 233–301.
- [Lew21] M. LEWIN – “Riesz and Coulomb gases: what’s known and unknown”, Survey talk at “Séminaire Matrices Et Graphes Aléatoires” (MEGA), Institut Henri Poincaré (IHP), <https://www.ceremade.dauphine.fr/dokuwiki/mega:seminaire>, 2021.
- [LL01] E. H. LIEB & M. LOSS – *Analysis*, second éd., Graduate Studies in Mathematics, vol. 14, American Mathematical Society, Providence, RI, 2001.

- [LS17] T. LEBLÉ & S. SERFATY – “Large deviation principle for empirical fields of log and Riesz gases”, *Invent. Math.* **210** (2017), no. 3, p. 645–757.
- [LS18] ———, “Fluctuations of two dimensional Coulomb gases”, *Geom. Funct. Anal.* **28** (2018), no. 2, p. 443–508.
- [LZ20] T. LEBLÉ & O. ZEITOUNI – “A local CLT for linear statistics of 2D Coulomb gases”, arXiv:2005.12163, 2020.
- [Meh04] M. L. MEHTA – *Random matrices*, third éd., Pure and Applied Mathematics, vol. 142, Acad. Press, 2004.
- [MMS14] M. MAÏDA & E. MAUREL-SEGALA – “Free transport-entropy inequalities for non-convex potentials and application to concentration for random matrices”, *Probab. Theory Related Fields* **159** (2014), no. 1-2, p. 329–356.
- [MS19] E. MECKES & K. STEWART – “On the eigenvalues of truncations of random unitary matrices”, *Electron. Commun. Probab.* **24** (2019), p. 12 (English), Id/No 57.
- [PG20] D. PADILLA-GARZA – “Concentration inequality around the thermal equilibrium measure of Coulomb gases”, preprint arXiv:2010.00194, 2020.
- [PH98] D. PETZ & F. HIAI – “Logarithmic energy as an entropy functional”, in *Advances in differential equations and mathematical physics (Atlanta, GA, 1997)*, Contemp. Math., vol. 217, Amer. Math. Soc., Providence, RI, 1998, p. 205–221.
- [PR05] D. PETZ & J. RÉFFY – “Large deviation for the empirical eigenvalue density of truncated Haar unitary matrices”, *Probab. Theory Related Fields* **133** (2005), no. 2, p. 175–189.
- [PS11] L. PASTUR & M. SHCHERBINA – *Eigenvalue distribution of large random matrices*, Mathematical Surveys and Monographs, vol. 171, American Mathematical Society, Providence, RI, 2011.
- [PS20] M. PETRACHE & S. SERFATY – “Crystallization for Coulomb and Riesz interactions as a consequence of the Cohn-Kumar conjecture”, *Proc. Am. Math. Soc.* **148** (2020), no. 7, p. 3047–3057 (English).
- [Rid03] B. RIDER – “A limit theorem at the edge of a non-Hermitian random matrix ensemble”, *J. Phys. A* **36** (2003), no. 12, p. 3401–3409, Random matrix theory.
- [Rot76] J.-P. ROTH – “Opérateurs dissipatifs et semi-groupes dans les espaces de fonctions continues”, *Ann. Inst. Fourier (Grenoble)* **26** (1976), no. 4, p. ix, 1–97.
- [Rou15] N. ROUGERIE – “De Finetti theorems, mean-field limits and Bose–Einstein condensation”, Cours Peccot lecture notes arXiv:1506.05263v1, 2015.
- [Roy07] G. ROYER – *An initiation to logarithmic Sobolev inequalities*, SMF/AMS Texts and Monographs, vol. 14, American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2007, Translated from the 1999 French original by Donald Babbitt.
- [RS93] L. C. G. ROGERS & Z. SHI – “Interacting Brownian particles and the Wigner law”, *Probab. Theory Related Fields* **95** (1993), no. 4, p. 555–570.
- [RS16] N. ROUGERIE & S. SERFATY – “Higher-dimensional Coulomb gases and renormalized energy functionals”, *Comm. Pure Appl. Math.* **69** (2016), no. 3, p. 519–605.
- [RV07] B. RIDER & B. VIRÁG – “The noise in the circular law and the Gaussian free field”, *Int. Math. Res. Not. IMRN* (2007), no. 2, p. Art. ID rnm006, 33.
- [Seo20] S.-M. SEO – “Edge Scaling Limit of the Spectral Radius for Random Normal Matrix Ensembles at Hard Edge”, *J. Stat. Phys.* **181** (2020), no. 5, p. 1473–1489.
- [Ser15] S. SERFATY – *Coulomb gases and Ginzburg-Landau vortices*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2015.
- [Ser18a] S. SERFATY – “Systèmes de points en interaction coulombienne”, *Gaz. Math.* (2018), no. 157, p. 29–37.
- [Ser18b] S. SERFATY – “Systems of points with Coulomb interactions”, *Eur. Math. Soc. Newsl.* (2018), no. 110, p. 16–21.
- [Ser20a] ———, “Gaussian fluctuations and free energy expansion for 2D and 3D Coulomb gases at any temperature”, preprint arXiv:2003.11704, 2020.
- [Ser20b] ———, “Mean field limit for Coulomb-type flows”, *Duke Math. J.* **169** (2020), no. 15, p. 2887–2935, With an appendix by Mitia Duerinckx and Serfaty.
- [ST97] E. B. SAFF & V. TOTIK – *Logarithmic potentials with external fields*, Fundamental Principles of Mathematical Sciences, vol. 316, Springer-Verlag, 1997, Appendix B by Thomas Bloom.
- [Tho04] J. J. THOMSON – “XXIV. On the structure of the atom: an investigation of the stability and periods of oscillation of a number of corpuscles arranged at equal intervals around the circumference of a circle; with application of the results to the theory of atomic structure”, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **7** (1904), no. 39, p. 237–265.
- [Voi93] D. VOICULESCU – “The analogues of entropy and of Fisher’s information measure in free probability theory. I”, *Comm. Math. Phys.* **155** (1993), no. 1, p. 71–92.
- [Voi94] ———, “The analogues of entropy and of Fisher’s information measure in free probability theory. II”, *Invent. Math.* **118** (1994), no. 3, p. 411–440.
- [Wig38] E. WIGNER – “Effects of the electron interaction on the energy levels of electrons in metals”, *Trans. Faraday Soc.* **34** (1938), p. 678–685.
- [Yuk98] J. E. YUKICH – *Probability theory of classical Euclidean optimization problems*, Lecture Notes in Mathematics, vol. 1675, Springer-Verlag, Berlin, 1998.
- [ZS00] K. ŻYCZKOWSKI & H.-J. SOMMERS – “Truncations of random unitary matrices”, *J. Phys. A* **33** (2000), no. 10, p. 2045–2057.