

ON BORSUK–ULAM THEOREMS AND CONVEX SETS

M. C. CRABB

ABSTRACT. The Intermediate Value Theorem is used to give an elementary proof of a Borsuk-Ulam theorem of Adams, Bush and Frick [1] that, if $f : S^1 \rightarrow \mathbb{R}^{2k+1}$ is a continuous function on the unit circle S^1 in \mathbb{C} such that $f(-z) = -f(z)$ for all $z \in S^1$, then there is a finite subset X of S^1 of diameter at most $\pi - \pi/(2k+1)$ (in the standard metric in which the circle has circumference of length 2π) such the convex hull of $f(X)$ contains $0 \in \mathbb{R}^{2k+1}$.

1. INTRODUCTION

We shall use the Intermediate Value Theorem to give an elementary proof of the following Borsuk-Ulam theorem of Adams, Bush and Frick in which $k \geq 1$ is a natural number, $\zeta = e^{2\pi i/(2k+1)} \in \mathbb{C}$ and the metric d on the unit circle, $S(\mathbb{C})$, in \mathbb{C} is given by $d(z, e^{i\theta}z) = |\theta|$ if $|\theta| \leq \pi$.

Theorem 1.1. ([1, Theorems 1 and 5]). *Let $f : S^1 = S(\mathbb{C}) \rightarrow \mathbb{R}^{2k+1}$ be a continuous map such that $f(-z) = -f(z)$ for all $z \in S(\mathbb{C})$. Then there exist $e_i \in \{\pm 1\}$, $i = 0, \dots, 2k$, and $z \in S(\mathbb{C})$ such that 0 lies in the convex hull of the image $f(X)$ of the finite set $X = \{e_i \zeta^i z \mid i = 0, \dots, 2k\}$, which is a subset of $S(\mathbb{C})$ with diameter at most $\pi - \pi/(2k+1)$.*

Moreover, there is an example of such a map f with the property that any finite subset $X \subseteq S(\mathbb{C})$ such that 0 lies in the convex hull of $f(X)$ has diameter greater than or equal to $\pi - \pi/(2k+1)$.

Cohomological methods, using little more than knowledge of the \mathbb{F}_2 -cohomology of a real projective space, will also, in Section 3, establish a Borsuk–Ulam theorem for higher dimensional spheres strengthening [1, Theorem 3]. The unit sphere $S(\mathbb{R}^n)$ in \mathbb{R}^n is equipped with the standard metric d : $\cos(d(u, v)) = \langle u, v \rangle$, $0 \leq d(u, v) \leq \pi$.

Theorem 1.2. *Let $m, n \geq 1$ be positive integers such that $m \leq 2^r \leq n < 2^{r+1}$, where $r \geq 0$ is a non-negative integer. Suppose that $f : S^{n-1} = S(\mathbb{R}^n) \rightarrow \mathbb{R}^{m+n-1}$ is a continuous map such that $f(-v) = -f(v)$ for all $v \in S(\mathbb{R}^n)$. Then there exists a finite subset $X \subseteq S(\mathbb{R}^n)$ with cardinality at most $m+n$ and diameter at most $\pi - \arccos(1/n)$ such that 0 lies in the convex hull of $f(X)$ in \mathbb{R}^{m+n-1} .*

Moreover, there is an example of such a map f with the property that any finite subset $X \subseteq S(\mathbb{R}^n)$ such that 0 lies in the convex hull of $f(X)$ has diameter greater than or equal to $\pi - \arccos(1/n)$.

Consider, in general, a continuous map $f : S(\mathbb{R}^n) \rightarrow \mathbb{R}^{m+n-1}$, where $m \geq 0$ and $n > 1$, such that $f(-v) = -f(v)$ for all $v \in S(\mathbb{R}^n)$. The classical Borsuk-Ulam

Date: August 2022.

2020 Mathematics Subject Classification. Primary 05E45, 52A20, 55M25, 55R25; Secondary 54E35, 55R40.

Key words and phrases. Intermediate Value theorem, Borsuk–Ulam theorem, Euler class.

theorem asserts, in one form, that, if $m = 0$, there is a point $x \in S(\mathbb{R}^n)$ such that $f(x) = 0$. If $n = 2$, the proof is an elementary exercise using the Intermediate Value Theorem, while for general n a proof (one of many) can be given using the \mathbb{F}_2 -cohomology of real projective space. If $m > 0$, the theorem clearly fails; for the inclusion $S(\mathbb{R}^n) \subseteq \mathbb{R}^n \hookrightarrow \mathbb{R}^{m+n-1}$ has no zero. But the Adams-Bush-Frick theorems show that there is a finite subset $X \subseteq S(\mathbb{R}^n)$ with the property that zero is a convex linear combination of the values $f(x)$, $x \in X$, and, to make the assertion non-trivial (because $\frac{1}{2}f(x) + \frac{1}{2}f(-x) = 0$ for any x), satisfying the condition that X does not contain any pair of antipodal points. And this condition is refined, for given m and n , by bounding the diameter of X . A precise statement is given in Corollary 2.4.

For the wider context of these results the reader is referred to [1, 2].

Acknowledgment. I am grateful to the authors of [1], H. Adams, J. Bush and F. Frick, for their helpful comments on the 2019 version of this note.

2. THE INTERMEDIATE VALUE THEOREM

Lemma 2.1. *Let $f : S(\mathbb{C}) \rightarrow \mathbb{R}^{2k+1}$ be a continuous map such that $f(-z) = -f(z)$. Suppose that w_0, \dots, w_{2k} are any $2k+1$ points in $S(\mathbb{C})$. Then there exist $e_i \in \{\pm 1\}$, $\lambda_i \geq 0$, for $i = 0, \dots, 2k$, with $\sum \lambda_i = 1$, and $z \in S(\mathbb{C})$, such that $\sum_{i=0}^{2k} \lambda_i f(e_i z w_i) = 0$.*

If $f(w_0), \dots, f(w_{2k})$ lie in a $2k$ -dimensional subspace of \mathbb{R}^{2k+1} , we can require that $z = 1$.

Proof. Consider the determinant map

$$\varphi : S(\mathbb{C}) \rightarrow \Lambda^{2k+1} \mathbb{R}^{2k+1}, z \mapsto f(z w_0) \wedge \cdots \wedge f(z w_{2k}).$$

Then $\varphi(-z) = (-1)^{2k+1} \varphi(z) = -\varphi(z)$. So, by the Intermediate Value Theorem, φ has a zero.

If $\varphi(z) = 0$, the vectors $f(z w_i)$ in \mathbb{R}^{2k+1} are linearly dependent and there exist $\mu_i \in \mathbb{R}$, not all zero such that $\sum_i \mu_i f(w_i) = 0$. We may assume, by scaling, that $\sum |\mu_i| = 1$. Choose $\lambda_i \geq 0$ and $e_i = \pm 1$ so that $\lambda_i e_i = \mu_i$. \square

Lemma 2.2. *For $z \in S(\mathbb{C})$ and $e_i = \pm 1$, $i = 0, \dots, 2k$, the distance $d(e_i z \zeta^i, e_j z \zeta^j)$ is less than or equal to $\pi - \pi/(2k+1)$.*

Proof. Indeed, the $2(2k+1)$ points $\pm z \zeta^i$ on the unit circle lie at the vertices of a regular polygon. \square

These two lemmas already prove the first part of Theorem 1.1.

Example 2.3. ([1, Theorem 5]). Let P be the k -dimensional complex vector space of complex polynomials $p(z)$ of degree $\leq 2k-1$ such that $p(-z) = -p(z)$. Let $g : S(\mathbb{C}) \rightarrow P^*$ be the evaluation map to the dual $P^* = \text{Hom}_{\mathbb{C}}(P, \mathbb{C})$. Suppose that w_0, \dots, w_{2k} are points of $S(\mathbb{C})$ such that 0 lies in the convex hull of the $g(w_i)$. Then $d(w_i, w_j) \geq \pi - \pi/(2k+1)$ for some i, j .

For the sake of completeness we include a concise version of the proof in [1].

Proof. Assume that the $2k+1$ points w_i^2 are distinct. By relabelling we may arrange that $w_1 = e^{i\theta} w_0$, where θ is the minimum of the distances $d(w_i, w_j)$, $i \neq j$.

Suppose that $\sum \lambda_i g(w_i) = 0$, where the $\lambda_i \in \mathbb{R}$ are not all equal to zero. This means that $\sum \lambda_i p(w_i) = 0$ for each $p \in P$, and hence, since $\lambda_i \in \mathbb{R}$ and $w_i^{-1} =$

$\overline{w_i}$, that $\sum \lambda_i p(w_i^{-1}) = 0$ too (because $\sum \lambda_i p(\overline{w_i})$ can be written as the complex conjugate of $\sum \lambda_i \overline{p(w_i)}$ with $\overline{p} \in P$). For $r \neq s$, $0 \leq r, s \leq 2k$, we may write

$$z^{-2k+1} \prod_{j \neq r, j \neq s} (z^2 - w_j^2) = p_+(z) + p_-(z^{-1}),$$

for unique polynomials $p_+, p_- \in P$. Then we find, because $\sum \lambda_i p_+(w_i) = 0$ and $\sum \lambda_i p_-(w_i^{-1}) = 0$, that

$$\sum_i \lambda_i w_i^{-2k+1} \prod_{j \neq r, j \neq s} (w_i^2 - w_j^2) = \sum_i \lambda_i (p_+(w_i) + p_-(w_i^{-1})) = 0,$$

that is,

$$\lambda_r \prod_{j \neq r} (w_r w_j^{-1} - w_r^{-1} w_j) = \lambda_s \prod_{j \neq s} (w_s w_j^{-1} - w_s^{-1} w_j).$$

It follows that, for some non-zero $c \in \mathbb{R}$,

$$\lambda_i \delta_i = c, \text{ where } \delta_i = \prod_{j \neq i} (w_i w_j^{-1} - w_i^{-1} w_j),$$

for all i . (Notice that δ_i , being the product of the $2k$ purely imaginary numbers $w_i \overline{w_j} - \overline{w_i} w_j$, is real.) In particular, all the λ_i are non-zero.

Given that 0 lies in the convex hull of the points $g(w_i)$, we can now assume further that all λ_i are non-negative, and so, because they are non-zero, strictly positive. Then the $\delta_i \in \mathbb{R}$ all have the same sign. We show that there is some i such that $w_i = -e^{it\theta} w_0$ for $0 < t < 1$.

Indeed, write $\psi(t) = \prod_{1 < j \leq 2k} (e^{it\theta} w_0 w_j^{-1} - e^{-it\theta} w_0^{-1} w_j) \in i\mathbb{R}$, for $0 \leq t \leq 1$. Then $\delta_0 = (w_0 w_1^{-1} - w_0^{-1} w_1) \psi(0)$ and $\delta_1 = (w_1 w_0^{-1} - w_1^{-1} w_0) \psi(1)$. So, by the Intermediate Value theorem again, $\psi(t) = 0$ for some t , and then $w_i^2 = (e^{it\theta} w_0)^2$ for some i , $1 < i \leq 2k$. But $w_i \neq e^{it\theta} w_0$, by the minimality of θ . So $w_i = -e^{it\theta} w_0$. Now $d(w_i, w_0) = \pi - t\theta$ and $d(w_i, w_1) = \pi - (1-t)\theta$. But clearly $\theta \leq 2\pi/(2k+1)$ and either $t \geq 1/2$ or $1-t \geq 1/2$. \square

Lemmas 2.1 and 2.2, together with Example 2.3, establish Theorem 1.1, using the fact¹ that, if Y is a finite subset of a real vector space such that the convex hull of Y contains 0 but no proper subset of Y has this property, then the set Y is affinely independent.

Corollary 2.4. *Let $m \geq 0$ and $n > 1$ be integers. Then there is a non-negative real number $\delta < 1$ with the property that, for any continuous map $f : S(\mathbb{R}^n) \rightarrow \mathbb{R}^{m+n-1}$ such that $f(-v) = -f(v)$ for all $v \in S(\mathbb{R}^n)$, there is a finite subset X of $S(\mathbb{R}^n)$ of cardinality at most $m+n$ and diameter at most $\pi - \arccos(\delta) < \pi$ such that 0 lies in the convex hull of $f(X)$.*

The classical Borsuk-Ulam theorem deals with the case $m = 0$: we may take $\delta = 0$ so that X consists of a single point.

Proof. By restricting f to $S(\mathbb{R}^2) \subseteq S(\mathbb{R}^n)$ and including \mathbb{R}^{m+n-1} in \mathbb{R}^{2k+1} for the smallest $k \geq 1$ such that $m+n-1 \leq 2k+1$, we see from Theorem 1.1 that the assertion is true with $\delta = \cos(\pi/(2k+1))$. \square

¹We have $\lambda_y > 0$, $y \in Y$, such that $\sum \lambda_y y = 0$. Suppose that $\mu_y \in \mathbb{R}$, $y \in Y$, satisfy $\sum \mu_y y = 0$ and $\sum \mu_y = 0$. Then, for any $t \in \mathbb{R}$ such that $\lambda_y \geq |t\mu_y|$ for all y , $\sum (\lambda_y + t\mu_y)y = 0$, $\sum (\lambda_y - t\mu_y)y = 0$, $\lambda_y \pm t\mu_y \geq 0$, and so $\lambda_y > \pm t\mu_y$, that is, $\lambda_y > |t\mu_y|$, for all y . Hence $\mu_y = 0$ for all y .

There is an easy extension of Lemma 2.1 to higher dimensions.

Lemma 2.5. *For integers $n, k \geq 1$, suppose that $f : S(\mathbb{C}^n) \rightarrow \mathbb{R}^{2k+1}$ is a continuous map such that $f(-v) = -f(v)$ for all $v \in S(\mathbb{C}^n)$. Then, for any $2k + 1$ vectors w_0, \dots, w_{2k} in $S(\mathbb{C}^n)$, there exist $e_i \in \{\pm 1\}$, $\lambda_i \geq 0$, for $i = 0, \dots, 2k$, with $\sum \lambda_i = 1$, and $z \in S(\mathbb{C})$ such that $\sum_{i=0}^{2k} \lambda_i f(e_i z w_i) = 0$.*

Proof. This can be established, using the same arguments as in the proof of Lemma 2.1, by looking at the function $\varphi : S(\mathbb{C}) \rightarrow \Lambda^{2k+1} \mathbb{R}^{2k+1}$ defined by $\varphi(z) = f(zw_0) \wedge \dots \wedge f(zw_{2k})$. \square

Example 2.6. For $n > 1$, write $\mathbf{e}_1, \dots, \mathbf{e}_n$ for the standard orthonormal \mathbb{C} -basis of \mathbb{C}^n , let $\eta = e^{2\pi i/(2l+1)}$ where $l \geq 1$ is a positive integer, and fix an integer r in the range $1 \leq r \leq n$.

If k satisfies $2k + 1 \leq (2l + 1)^r \binom{n}{r}$, we can choose distinct vectors w_0, \dots, w_{2k} from the set

$$\{(\sum_{s=1}^r \eta^{a_s} \mathbf{e}_{i_s})/\sqrt{r} \mid 1 \leq i_1 < i_2 < \dots < i_r \leq n, a_s = 0, \dots, 2l\} \subseteq S(\mathbb{C}^n).$$

Then $|\langle w_i, w_j \rangle| \leq \delta = 1 - (1 - \cos(\pi/(2l + 1)))/r$ for $i \neq j$. Hence $d(e_i z w_i, e_j z w_j) \leq \pi - \arccos(\delta)$.

Application of Lemma 2.5 to the case $r = 1$, for which $\arccos(\delta) = \pi/(2l + 1)$, gives a result which is close to [1, Theorem 2], but slightly weaker. For general r , if l is large, $\arccos(\delta)$ is close to $(\pi/(2l + 1))/\sqrt{r}$. In particular, the case $r = n$ gives a much stronger result than that provided by [1, Theorem 2] when k is sufficiently large. (A similar observation is made in [2, Remark 4.2].)

3. COHOMOLOGICAL METHODS

In this section we shall use cohomological methods in higher dimensions. We begin by recalling some standard facts about projective bundles.

Consider a real vector bundle $E \rightarrow B$ (or simply ' E ') of dimension n over a finite complex B . The *mod 2 Euler class* of E will be written as

$$e(E) \in H^n(B; \mathbb{F}_2)$$

in cohomology with \mathbb{F}_2 -coefficients. (Thus, $e(E)$ is the top Stiefel-Whitney class $w_n(E)$.) The crucial property for our applications is that, if $e(E) \neq 0$, then every section $s : B \rightarrow E$ has a zero, that is, there is some point $b \in B$ such that $s(b) \in E_b$, the vector space fibre of E at b , is equal to $0 \in E_b$. For calculations we shall use the *multiplicativity* of the Euler class: $e(E \oplus E') = e(E) \cdot e(E')$ for two vector bundles E and E' over B .

The *projective bundle* $P(E) \rightarrow B$ of E is the quotient of the sphere bundle $S(E)$ of E by the involution -1 : $P(E) = S(E)/\{\pm 1\}$; elements of $P(E)$ can be written as $[u]$, where $u \in S(E)$. The *Hopf line bundle* over $P(E)$ is denoted by H ; its fibre at $[u]$ is the line $\mathbb{R}u$ generated by u .

The cohomology of the space $P(E)$, as an algebra over $H^*(B; \mathbb{F}_2)$, is a quotient of the polynomial ring $H^*(B; \mathbb{F}_2)[T]$ in an indeterminate T :

$$H^*(P(E); \mathbb{F}_2) = H^*(B; \mathbb{F}_2)[T]/(T^n + \dots + w_i(E)T^{n-i} + \dots + w_n(E)),$$

where $w_i(E) \in H^i(B; \mathbb{F}_2)$ is the i th *Stiefel-Whitney class* of E , and the Euler class $e(H)$ corresponds to T .

The multiplicativity of the Euler class translates into the multiplicativity of the *total Stiefel-Whitney class* $w(E) = 1 + w_1(E) + \dots + w_i(E) + \dots + w_n(E)$: $w(E \oplus E') = w(E) \cdot w(E')$.

Working now towards a proof of Theorem 1.2 we consider, for $n > 1$, an inscribed regular n -simplex in the sphere $S(\mathbb{R}^n)$ with vertices v_0, \dots, v_n . Write $n = 2^r + s$, where $r \geq 1$ and $0 \leq s < 2^r$. An involution τ in the orthogonal group $O(\mathbb{R}^n)$ is specified by: $\tau(v_i) = v_{i+s}$ for $i < s$ and $\tau(v_i) = v_i$ for $i = 2s, \dots, n$. Let V_+ and V_- be the ± 1 -eigenspaces of τ : $V_{\pm} = \{u \in \mathbb{R}^n \mid \tau u = \pm u\}$. Thus V_+ has dimension $n - s = 2^r$ with a basis $v_i + v_{i+s}$ ($0 \leq i < s$), v_i ($2s < i \leq n$), and V_- has dimension s with a basis $v_i - v_{i+s}$ ($0 \leq i < s$). The special orthogonal group $SO(V_+)$, which is isomorphic to $SO(\mathbb{R}^{2^r})$, is included in $SO(\mathbb{R}^n)$ by the map $g \mapsto (g, 1) : V_+ \oplus V_- = \mathbb{R}^n \rightarrow V_+ \oplus V_- = \mathbb{R}^n$ as the subgroup $G \leq SO(\mathbb{R}^n)$ consisting of those elements which fix the s vectors $v_i - v_{i+s}$, $0 \leq i < s$.

Theorem 3.1. (Compare [1, Theorem 3]). *For an integer $n > 1$, let v_0, \dots, v_n be the vertices of an inscribed regular n -simplex in $S(\mathbb{R}^n)$. Write $n = 2^r + s$, where $r \geq 1$ and $0 \leq s < 2^r$, and let $G \leq SO(\mathbb{R}^n)$ be the subgroup $\{g \in SO(\mathbb{R}^n) \mid g(v_i - v_{i+s}) = v_i - v_{i+s} \text{ for } 0 \leq i < s\}$.*

Let $f : S(\mathbb{R}^n) \rightarrow \mathbb{R}^{n+2^r-1}$ be a continuous map such that $f(-v) = -f(v)$ for all $v \in S(\mathbb{R}^n)$. Then there exist $e_i \in \{\pm 1\}$, $i = 0, \dots, n$, and an element $g \in G$ such that 0 lies in the convex hull of the image $f(X)$ of the finite subset $X = \{e_i g v_i \mid i = 0, \dots, n\} \subseteq S(\mathbb{R}^n)$, which has diameter at most $\pi - \arccos(1/n)$.

Moreover, the example of the inclusion $f : S(\mathbb{R}^n) \subseteq \mathbb{R}^n \hookrightarrow \mathbb{R}^{n+2^r-1}$ has the property that any finite subset $X \subseteq S(\mathbb{R}^n)$ such that 0 lies in the convex hull of $f(X)$ has diameter greater than or equal to $\pi - \arccos(1/n)$.

Proof. We continue to use the notation introduced before the statement of the theorem.

Consider the map $\sigma : SO(V_+) \times S(\mathbb{R}^{n+1}) \rightarrow \mathbb{R}^{n+2^r-1}$

$$(g, (\mu_0, \dots, \mu_n)) \mapsto \sum_{i=0}^n \mu_i f(g v_i), \quad g \in SO(V_+), \quad \sum_{i=0}^n \mu_i^2 = 1.$$

We shall show that σ has a zero.

Clearly $\sigma(g, -\mu) = -\sigma(g, \mu)$. Now notice that $-g \in SO(V_+)$ maps to $(-g, 1) = -(g, 1)\tau \in SO(\mathbb{R}^n)$. So $\sigma(-g, \mu) = -\sigma(g, T\mu)$, where T is the linear involution of \mathbb{R}^{n+1} satisfying, for $\mu = (\mu_0, \dots, \mu_n) \in \mathbb{R}^{n+1}$, $(T\mu)_i = \mu_{i+s}$ for $i < s$ and $(T\mu)_i = \mu_i$ for $i = 2s, \dots, n$.

Let L be the real line bundle over the projective special orthogonal group $PSO(V_+)$ associated with the double cover $SO(V_+) \rightarrow PSO(V_+) = SO(V_+)/\{\pm 1\}$ or, explicitly, the quotient of $SO(V_+) \times \mathbb{R}$ by the involution $(g, t) \mapsto (-g, -t)$. And let E be the $(n+1)$ -dimensional real vector bundle over $PSO(V_+)$ associated to the involution $-T$ of \mathbb{R}^{n+1} , that is, E is the quotient of $SO(V_+) \times \mathbb{R}^{n+1}$ by the involution $(g, \mu) \mapsto (-g, -T\mu)$. The -1 -eigenspace of $-T$ consists of those $\mu \in \mathbb{R}^{n+1}$ such that $\mu_i = \mu_{i+s}$ for $0 \leq i < s$ and has dimension $n+1-s = 2^r+1$, and the $+1$ -eigenspace has the complementary dimension s . The bundle E is thus isomorphic to the direct sum $(L \otimes \mathbb{R}^{2^r+1}) \oplus \mathbb{R}^s$ of 2^r+1 copies of L and the trivial bundle with fibre \mathbb{R}^s .

The map σ determines a section s of $H \otimes \mathbb{R}^{n+2^r-1}$ (the tensor product of the pullback of the Hopf line bundle H over $P(\mathbb{R}^{n+1})$ with the trivial bundle with fibre

\mathbb{R}^{n+2^r-1}):

$$s([[g, \pm\mu]]) = [g, \mu] \otimes \sigma(g, \mu), \quad \text{for } g \in \text{SO}(V_+), \mu \in S(\mathbb{R}^{n+1}),$$

where the element $[[g, \mu]] = [[g, -\mu]] \in P(E)$ is determined by the class $[g, \mu] \in S(E)$ of $(g, \mu) \in \text{SO}(V_+) \times S(\mathbb{R}^{n+1})$. We shall show that s has a zero, and hence that σ has a zero, by proving that the mod 2 Euler class

$$e(H \otimes \mathbb{R}^{n+2^r-1}) = e(H)^{n+2^r-1} \in H^{n+2^r-1}(P(E); \mathbb{F}_2)$$

is non-zero.

Let us write $x = e(L)$ and $y = e(H)$. Now we recall from [3, Lemma (8.1)] that $x^{2^r-1} \neq 0$ and $x^{2^r} = 0$. (In order to make this account as self-contained as possible, an elementary proof of this fact is included in an Appendix as Proposition A.1.) So the total Stiefel-Whitney class $w(E)$ of E is determined, using multiplicativity, by $w(L) = 1 + x$ as $w(E) = w(L)^{2^r+1} = (1+x)^{2^r+1} = 1+x$ (because $(1+x)^{2^r} = 1+x^{2^r} = 1$). Hence

$$H^*(P(E); \mathbb{F}_2) = H^*(\text{PSO}(V_+); \mathbb{F}_2)[y]/(y^{n+1} + xy^n),$$

(meaning that the quotient of the polynomial ring $H^*(\text{PSO}(V_+); \mathbb{F}_2)[T]$ by the principal ideal generated by $T^{n+1} + xT^n$ is isomorphic to $H^*(P(E); \mathbb{F}_2)$ by the homomorphism mapping T to y). Using the relation $y^{n+1} = xy^n$, we conclude that $y^{n+2^r-1} = x^{2^r-1}y^n$ is non-zero.

Hence σ has a zero, that is, there exist $g \in \text{SO}(V_+)$ and μ_0, \dots, μ_n , not all zero, such that $\sum_i \mu_i f(gv_i) = 0$. The existence of the required $e_i \in \{\pm 1\}$ follows as in the proof of Lemma 2.1.

The $n+1$ points v_i satisfy $\sum_i v_i = 0$ and $\langle v_i, v_j \rangle = c$, say, for $i \neq j$. Since $\sum_{i=0}^n \langle v_0, v_i \rangle = 0$, we see that $1 + nc = 0$, so that $c = -1/n$. Hence $d(v_i, v_j) = \pi - \arccos(1/n)$ and $d(v_i, -v_j) = \arccos(1/n)$ if $i \neq j$.

So $d(e_i gv_i, e_j gv_j) = d(e_i v_i, e_j v_j) \leq \pi - \arccos(1/n)$ for $i \neq j$.

The final assertion follows, as in the proof of Theorem 1.1, from Lemma 3.2 below. \square

Lemma 3.2. *Suppose that w_0, \dots, w_n are points of $S(\mathbb{R}^n)$, such that $\sum \lambda_i w_i = 0$, where $\lambda_i \geq 0$ and $\sum \lambda_i = 1$. Then $d(w_i, w_j) \geq \pi - \arccos(1/n)$ for some i, j .*

Proof. Put $d = \min\{\langle w_i, w_j \rangle \mid i \neq j\}$. Then

$$0 = \sum_i \sum_j \lambda_j \langle w_i, w_j \rangle = \sum_i \lambda_i \langle w_i, w_i \rangle + \sum_i \sum_{j \neq i} \lambda_j \langle w_i, w_j \rangle \geq 1 + nd.$$

So $d \leq -1/n$. If $\langle w_i, w_j \rangle \leq -1/n$, then $d(w_i, w_j) \geq \pi - \arccos(1/n)$. \square

The result in Theorem 3.1 is optimal when $n = 2^r$ is a power of 2 and so $n + 2^r - 1 = 2n - 1$, as the following construction demonstrates.

Proposition 3.3. *For any even integer $n > 1$, there is a map $f : S(\mathbb{R}^n) \rightarrow \mathbb{R}^{2n}$ such that for every finite subset X of $S(\mathbb{R}^n)$ with diameter $\leq \pi - \arccos(1/n)$ the zero vector $0 \in \mathbb{R}^{2n}$ does not lie in the convex hull of $f(X)$.*

Proof. Consider a map $f : S(\mathbb{R}^n) \rightarrow \mathbb{R}^{2n} = \mathbb{R}^n \oplus \mathbb{R}^n$ of the form $f(v) = (v, \varphi(v)v)$, where $\varphi : S(\mathbb{R}^n) \rightarrow \mathbb{R}$ is a continuous function such that $\varphi(-v) = \varphi(v)$.

Suppose that $X \subseteq S(\mathbb{R}^n)$ is a finite subset of diameter at most $\pi - \arccos(1/n)$ such that $0 = (0, 0)$ lies in the convex hull of $f(X)$.

Considering the first component, we see that there are points w_0, \dots, w_m of X for some m with $1 \leq m \leq n$ and $\lambda_0, \dots, \lambda_m \geq 0$ with $\sum \lambda_i = 1$, such that $\sum \lambda_i w_i = 0$ and $\langle w_i, w_j \rangle \geq -1/n$ for all i, j . (Indeed, let $Y \subseteq X$ be a minimal subset of X such that 0 lies in the convex hull of Y . As we have already recalled in the proof of Theorem 1.1, the set Y must be affinely independent and so has cardinality, $m + 1$ say, less than or equal to $n + 1$.)

Then, for each i ,

$$-\lambda_i = \sum_{j \neq i} \lambda_j \langle w_i, w_j \rangle \geq -(1 - \lambda_i)/n,$$

and so $\lambda_i \leq 1/(n + 1)$. Hence $m = n$ and $\lambda_i = 1/(n + 1)$ for all i . And then $\sum_{j \neq i} \langle w_i, w_j \rangle = -1$. We conclude that $\langle w_i, w_j \rangle = -1/n$ for $i \neq j$. Hence w_0, \dots, w_n are the vertices of an inscribed regular n -simplex in $S(\mathbb{R}^n)$.

If $v \in S(\mathbb{R}^n)$ is a vector distinct from each of w_0, \dots, w_n , then $\langle v, w_i \rangle < -1/n$ for some i . (Indeed, if $v = \sum_{j \neq i} t_j w_j$, where $t_j \geq 0$, then $\langle v, w_i \rangle < -1/n$. For $1 = \|v\|^2 = (\sum_{j \neq i} t_j)^2 - ((n + 1)/n) \sum_{j \neq j', j \neq i \neq j'} t_j t_{j'}$. So $\sum_{j \neq i} t_j > 1$. Hence $\langle v, w_i \rangle = -(\sum_{j \neq i} t_j)/n < -1/n$.) Thus X is precisely the set $\{w_0, \dots, w_n\}$.

Since the w_i are the vertices of a regular n -simplex, if $\sum_i \mu_i w_i = 0$, $\mu_i \in \mathbb{R}$, then the coefficients μ_i must be equal. So, looking now at the second component of f , we conclude that the real numbers $\varphi(w_i)$, $i = 0, \dots, n$, are equal. (For $\sum_i \lambda_i f(w_i) = 0$ means that $\sum_i \lambda_i w_i = 0$ and $\sum_i \lambda_i \varphi(w_i) w_i = 0$, and we have shown that $\lambda_i = 1/(n + 1)$.)

To construct a map f such that 0 does not lie in the convex hull of $f(X)$ for any finite set with diameter at most $\pi - \arccos(1/n)$, we write down a function φ which is not constant on the vertices w_0, \dots, w_n of any regular n -simplex. For example, choose a non-zero vector $u \in \mathbb{R}^n$ and take $\varphi(v) = \langle u, v \rangle^2$. Then $0 = \sum \langle u, w_i \rangle = \sum_{i=0}^n \epsilon_i \sqrt{\varphi(w_i)}$, where $\epsilon_i = \pm 1$. Since $n + 1$ is odd, if $\varphi(w_i) = c^2$ for all i , then $c = 0$, which is impossible because $u \neq 0$. \square

Proof of Theorem 1.2. This follows at once from Theorem 3.1 by including \mathbb{R}^{m+n-1} in \mathbb{R}^{n+2^r-1} . \square

Methods similar to those used to derive Theorem 3.1 give the following generalization of Lemma 2.5.

Lemma 3.4. *For some integers $r \geq 0$ and $n, k \geq 1$, suppose that $f : S(\mathbb{R}^{2^r n}) \rightarrow \mathbb{R}^{2^r k + 2^r - 1}$ is a continuous map such that $f(-v) = -f(v)$ for all $v \in S(\mathbb{R}^{2^r n})$. We include the special orthogonal group $\text{SO}(\mathbb{R}^{2^r})$ in $\text{SO}(\mathbb{R}^{2^r n})$ by the diagonal homomorphism $\text{SO}(\mathbb{R}^{2^r}) \rightarrow \text{SO}(\mathbb{R}^{2^r}) \times \dots \times \text{SO}(\mathbb{R}^{2^r}) \subseteq \text{SO}(\mathbb{R}^{2^r n})$. Then, for any $2^r k + 1$ vectors $w_0, \dots, w_{2^r k}$ in $S(\mathbb{R}^{2^r n})$, there exist $e_i \in \{\pm 1\}$, $\lambda_i \geq 0$, for $i = 0, \dots, 2^r k$, with $\sum \lambda_i = 1$, and $g \in \text{SO}(\mathbb{R}^{2^r})$, such that $\sum_{i=0}^{2^r k} \lambda_i f(e_i g w_i) = 0$.*

If, for some $\delta < 1$, the vectors w_i satisfy $|\langle w_i, w_j \rangle| \leq \delta$ for all $i \neq j$, then $d(e_i g w_i, e_j g w_j) \leq \pi - \arccos(\delta) < \pi$ for $i \neq j$.

Proof. We assume that $r \geq 1$. The simpler case $r = 0$ is discussed in Remark 3.5. Consider the map $\sigma : \text{SO}(\mathbb{R}^{2^r}) \times S(\mathbb{R}^{2^r k + 1}) \rightarrow \mathbb{R}^{2^r k + 2^r - 1}$

$$(g, (\mu_0, \dots, \mu_{2^r k})) \mapsto \sum_{i=0}^{2^r k} \mu_i f(g w_i), \quad g \in \text{SO}(\mathbb{R}^{2^r}), \quad \sum_{i=0}^{2^r k} \mu_i^2 = 1.$$

Since $\sigma(-g, \mu) = -\sigma(g, \mu) = \sigma(g, -\mu)$, the map σ determines a section s of the vector bundle $\mathbb{R}^{2^r k + 2^r - 1} \otimes (L \otimes H)$ over the space $(\mathrm{SO}(\mathbb{R}^{2^r})/\{\pm 1\}) \times (S(\mathbb{R}^{2^r k + 1})/\{\pm 1\})$, where L and H are the line bundles associated with the double covers $\mathrm{SO}(\mathbb{R}^{2^r}) \rightarrow \mathrm{SO}(\mathbb{R}^{2^r})/\{\pm 1\}$ and $S(\mathbb{R}^{2^r k + 1}) \rightarrow S(\mathbb{R}^{2^r k + 1})/\{\pm 1\}$ of $\mathrm{PSO}(\mathbb{R}^{2^r})$ and the real projective space $P(\mathbb{R}^{2^r k + 1})$, respectively.

The \mathbb{F}_2 -Euler class of $\mathbb{R}^{2^r k + 2^r - 1} \otimes (L \otimes H)$ is equal to

$$e(L \otimes H)^{2^r k + 2^r - 1} = (x + y)^{2^r k + 2^r - 1} \in H^{2^r k + 2^r - 1}(\mathrm{PSO}(\mathbb{R}^{2^r}) \times P(\mathbb{R}^{2^r k + 1}); \mathbb{F}_2),$$

where $x = e(L)$ and $y = e(H)$. Since $x^{2^r - 1} \neq 0$ and $x^{2^r} = 0$ (by Proposition A.1 again), $y^{2^r k} \neq 0$ and $y^{2^r k + 1} = 0$,

$$(x + y)^{2^r k + 2^r - 1} = (x + y)^{2^r - 1} (x^{2^r} + y^{2^r})^k = (x + y)^{2^r - 1} y^{2^r k} = x^{2^r - 1} y^{2^r k}$$

is non-zero. Hence the section s has a zero, and, therefore, so too has σ . The proof is completed as before. \square

Remark 3.5. When $r = 0$, so that $g = 1$ (because $\mathrm{SO}(\mathbb{R}^1)$ is trivial), the result specializes to the main theorem of [2, Theorem 1]. In that case, the map σ determines a section s of $\mathbb{R}^k \otimes H$ over $P(\mathbb{R}^{k+1})$, and s has a zero just because $y^k = e(H)^k$ is non-zero in $H^k(P(\mathbb{R}^{k+1}); \mathbb{F}_2)$.

The case $r = 1$ is precisely Lemma 2.5 with the identifications $\mathbb{R}^{2n} = \mathbb{C}^n$ and $\mathrm{SO}(\mathbb{R}^2) = \mathrm{U}(\mathbb{C}) = S(\mathbb{C})$.

When $r = 2$, we can write $\mathbb{R}^{4n} = \mathbb{H}^n$ and use quaternion multiplication to show that the element $g \in \mathrm{SO}(\mathbb{R}^4)$ may be chosen to lie in the subgroup $\mathrm{Sp}(\mathbb{H}) = S(\mathbb{H})$ of unit quaternions. (And, when $r = 3$, we could use Cayley multiplication.)

Corollary 3.6. *Let $m, n \geq 1$ be integers with $m \leq n$. Then there are $m + n$ vectors $w_0, \dots, w_{m+n-1} \in S(\mathbb{R}^n)$ with the property that, for any continuous map $f : S(\mathbb{R}^n) \rightarrow \mathbb{R}^{m+n-1}$ with $f(-v) = -f(v)$, there exist $e_0, \dots, e_{m+n-1} \in \{\pm 1\}$ such that 0 lies in the convex hull of the image $f(X)$ of the finite set $X = \{e_i w_i \mid i = 0, \dots, m+n-1\}$ with cardinality $m+n$ and diameter at most $\pi - \arccos(1/\lfloor n/m \rfloor)$.*

Notice that the vectors w_0, \dots, w_{m+n-1} are independent of f ; compare Theorem 1.2. The construction we use can be seen as a special case of [4, Lemma 16]. Compare, also, Example 2.6.

Proof. Write $n = qm + l$, where $q = \lfloor n/m \rfloor \geq 1$ and $0 \leq l < m$. We can assume that $q > 1$. Let u_0, \dots, u_q be the vertices of an inscribed regular q -simplex in $S(\mathbb{R}^q)$ and let $\mathbf{e}_1, \dots, \mathbf{e}_m$ be an orthonormal basis of \mathbb{R}^m . We give $\mathbb{R}^q \otimes \mathbb{R}^m$ the standard inner product: $\langle u \otimes v, u' \otimes v' \rangle = \langle u, u' \rangle \cdot \langle v, v' \rangle$. In other words, $\mathbb{R}^q \otimes \mathbb{R}^m$ is the orthogonal direct sum $\bigoplus_{j=1}^m \mathbb{R}^q \otimes \mathbb{R} \mathbf{e}_j$. Thus $\langle u_i \otimes \mathbf{e}_j, u_{i'} \otimes \mathbf{e}_{j'} \rangle$ is equal to 1 if $i = i', j = j'$, equal to $-1/q$ if $i \neq i', j = j'$, and is zero otherwise.

Let $w_0, \dots, w_{k-l} \in \mathbb{R}^{qm} \subseteq \mathbb{R}^{qm} \oplus \mathbb{R}^l = \mathbb{R}^n$, where $k = n + m - 1$, correspond to the $(q+1)m = n + m - l = k - l + 1$ vectors $u_i \otimes \mathbf{e}_j$, in some order, under a chosen isometric isomorphism $\mathbb{R}^{qm} \cong \mathbb{R}^q \otimes \mathbb{R}^m$ and let $w_{k-l+1}, \dots, w_{k-l+l}$ be an orthonormal basis of $\mathbb{R}^l \subseteq \mathbb{R}^{qm} \oplus \mathbb{R}^l = \mathbb{R}^n$. The result follows from Lemma 3.4 with $r = 0$ and $\delta = 1/q$. \square

4. CONCLUSION

We finish with a reformulation of the main results.

Definition 4.1. Given integers $m \geq 1$ and $n > 1$, set $k = m + n - 1$. Let $\delta(m, n)$ be the smallest real number $\delta \in (0, 1]$ with the property that, for any continuous map $f : S(\mathbb{R}^n) \rightarrow \mathbb{R}^k$ satisfying $f(-v) = -f(v)$ for all $v \in S(\mathbb{R}^n)$, there is a finite subset $X \subseteq S(\mathbb{R}^n)$ of cardinality less than or equal to $m + n$ and diameter at most $\pi - \arccos(\delta)$ such that 0 lies in the convex hull of $f(X)$.

The existence of $\delta(m, n)$ follows from a compactness argument. For a given f , the subspace

$$\{(\lambda_0, \dots, \lambda_k, w_0, \dots, w_k) \in [0, 1]^{k+1} \times S(\mathbb{R}^n)^{k+1} \mid \sum_{i=0}^k \lambda_i = 1, \sum_{i=0}^k \lambda_i f(w_i) = 0\}$$

of $[0, 1]^{k+1} \times S(\mathbb{R}^n)^{k+1}$ is compact and non-empty. So the subset $\Delta(f) \subseteq [0, 1]$ consisting of those δ for which there is a subset X of diameter $\leq \pi - \arccos(\delta)$ such that 0 is in the convex hull of $f(X)$ is closed. Hence $\bigcap_f \Delta(f)$ is closed (and contains 1).

The example of the inclusion of $S(\mathbb{R}^n)$ in $\mathbb{R}^n \subseteq \mathbb{R}^k$ shows, by Lemma 3.2, that $\delta(m, n) \geq 1/n$. Corollary 2.4 asserts that $\delta(m, n) < 1$.

Theorem 4.2. *The function $\delta(m, n)$ defined for $m \geq 1$ and $n > 1$ has the following properties.*

- (i). For all m and n , $1/n \leq \delta(m, n) < 1$.
- (ii). For all $k \geq 1$, $\delta(2k-1, 2) = \delta(2k, 2) = \cos(\pi/(2k+1))$ (Theorem 1.1, Example 2.3).
- (iii). If there is an integer r such that $1 \leq m \leq 2^r \leq n$, then $\delta(m, n) = 1/n$ (Theorem 1.2).
- (iv). For all $k \geq 1$, $\delta(2k+1, 2k) > 1/(2k)$ (Proposition 3.3).
- (v). Fix $m \geq 1$. Then $\delta(m, n) \rightarrow 0$ as $n \rightarrow \infty$, and so $\pi - \arccos(\delta(m, n)) \rightarrow \pi/2$. (Compare [2, Corollary 3.1].)
- (vi). Fix $n > 1$. Then $\delta(m, n) \rightarrow 1$ as $m \rightarrow \infty$, and so $\pi - \arccos(\delta(m, n)) \rightarrow \pi$.

Proof. Part (v) follows at once from (iii). Only (vi) still requires proof. If $m \leq m'$, then clearly $\delta(m, n) \leq \delta(m', n)$.

Given n and a positive integer $l \geq 1$, set $m + n - 1 = k = \binom{n+2l}{n-1}$. We shall establish the easy estimate $\delta(m, n) \geq 1/k^{1/(2l+1)}$. Since k , the dimension of the space of homogeneous polynomials of degree $2l+1$ in n variables, is clearly bounded by $2l+1 \leq k \leq (2l+1)^{n-1}$, we see that, as $l \rightarrow \infty$ (with n fixed), $m \rightarrow \infty$ and $k^{1/(2l+1)} \rightarrow 1$.

Consider the map $f : S(\mathbb{R}^n) \rightarrow (\mathbb{R}^n)^{\otimes(2l+1)}$ given by the $(2l+1)$ -th power $f(v) = v \otimes \cdots \otimes v$ and mapping into the k -dimensional vector subspace fixed by the permutation action of \mathfrak{S}_{2l+1} . Give $(\mathbb{R}^n)^{\otimes(2l+1)}$ the standard inner product:

$$\langle u_1 \otimes \cdots \otimes u_{2l+1}, v_1 \otimes \cdots \otimes v_{2l+1} \rangle = \prod_i \langle u_i, v_i \rangle,$$

so that each vector $f(v)$ lies in the unit sphere.

By Lemma 3.2, if w_0, \dots, w_k are points of $S(\mathbb{R}^n)$ such that 0 lies in the convex hull of the unit vectors $f(w_0), \dots, f(w_k)$, then $\langle f(w_i), f(w_j) \rangle = \langle w_i, w_j \rangle^{2l+1} \leq -1/k$ for some $i \neq j$. So $\langle w_i, w_j \rangle \leq -1/k^{1/(2l+1)}$. \square

Remark 4.3. The example used to prove part (vi) of Theorem 4.2 shows more precisely that, for any real number $\alpha < 1/(n-1)$,

$$m^\alpha(1 - \delta(m, n)) \rightarrow 0 \text{ as } m \rightarrow \infty.$$

Proof. The construction shows that $\delta((2l+1)^{n-1}, n) \geq 1/(2l+1)^{(n-1)/(2l+1)}$. Write $x = 1 - \delta((2l+1)^{n-1}, n)$. Then

$$(2l+1)^{n-1} \geq (1-x)^{-(2l+1)} = \sum_{s \geq 0} \binom{2l+s}{s} x^s \geq \binom{2l+r}{r} x^r \geq (2l+1)^r x^r / r!$$

for any $r \geq 1$. Thus $x \leq (r!)^{1/r} (2l+1)^{(n-1)/r-1}$.

Given α , choose an integer r so large that $\epsilon = 1/(n-1) - \alpha - 1/r$ is greater than 0. Then, with $m = (2l+1)^{n-1}$, we have $m^\alpha(1 - \delta(m, n)) \leq (r!)^{1/r} / m^\epsilon$. \square

Remark 4.4. In the opposite direction, an example shows that, for any real number $\alpha > 2/(n-1)$,

$$m^\alpha(1 - \delta(m, n)) \rightarrow \infty \text{ as } m \rightarrow \infty.$$

Proof. We shall use Lemma 3.4 with $r = 0$.

Let $V = \{(t_1, \dots, t_n) \in \mathbb{R}^n \mid \sum_{i=1}^n t_i = 1\}$ and let $\Delta \subseteq V$ be the simplex $\{(t_i) \in V \mid t_i \geq 0\}$. The map $\pi : V \rightarrow S(\mathbb{R}^n)$, $v \mapsto v/\|v\|$ gives a diffeomorphism from V to an open subspace of the sphere mapping geodesics (straight line segments) to geodesics (segments of great circles). Hence, there is a constant $c_n > 0$ such that $d(\pi(u), \pi(v)) \geq c_n \|u - v\|$ for all u, v in the compact subspace Δ .

Fix an integer $l \geq 1$ and let $W = \{(t_i) \in \Delta \mid lt_i \in \mathbb{Z} \text{ for all } i\}$. For $u, v \in W$, $u \neq v$, we see that $\|u - v\|^2 \geq 2/l^2$. List the vectors $\pi(v)$, $v \in W$, as w_0, \dots, w_k , where $k+1 = \binom{l+n-1}{n-1} \geq l^{n-1}/(n-1)!$. So, for $i \neq j$, $d(w_i, w_j) \geq c_n \sqrt{2}/l$.

This shows that $\delta(k+1-n, n) \leq \cos(c_n \sqrt{2}/l) = 1 - c_n^2/l^2 + \dots$. Finally, observe that $m = k+1-n \geq \frac{1}{2} l^{n-1}/(n-1)!$ if l is sufficiently large. \square

APPENDIX A. ON THE COHOMOLOGY OF $\text{PO}(\mathbb{R}^{2^r})$

In this appendix we give an elementary proof of the result from [3, Lemma (8.1)] that we used in the proofs of Theorem 3.1 and Lemma 3.4.

Proposition A.1. *Write $V = \mathbb{R}^{2^r n}$, where $r \geq 1$ and $n \geq 1$ is odd. Let L be the real line bundle over the projective orthogonal group $\text{PO}(V)$ associated with the double cover $\text{O}(V) \rightarrow \text{PO}(V)$. Then $e(L) \in H^1(\text{PO}(V); \mathbb{F}_2)$ satisfies $e(L)^{2^r-1} \neq 0$, $e(L)^{2^r} = 0$. Moreover, the restriction of $e(L)^{2^r-1}$ to the projective special orthogonal group $\text{PSO}(V) \leq \text{PO}(V)$ is non-zero.*

Proof. Notice, first, that the vector bundle $L \otimes V$ is trivial: it has a natural trivialization $L \otimes V \rightarrow V$ mapping $g \otimes v$, where g is an element of $\text{O}(V)$, regarded as a generator of the fibre of L at $[g] \in \text{PO}(V)$, and $v \in V$, to gv . Hence $1 = w(L \otimes V) = (1 + e(L))^{2^r n} = (1 + e(L)^{2^r})^n$ and, since n is odd, $e(L)^{2^r} = 0$.

The diagonal map $\text{O}(\mathbb{R}^{2^r}) \rightarrow (\text{O}(\mathbb{R}^{2^r}))^n \leq \text{O}(\mathbb{R}^{2^r n})$ induces an inclusion of $\text{PO}(\mathbb{R}^{2^r})$ in $\text{PO}(V)$. To prove that $e(L)^{2^r-1}$ is non-zero it is, therefore, enough to deal with the case $n = 1$, and for the remainder of the discussion we take $V = \mathbb{R}^{2^r}$.

Given a Euclidean vector bundle E over a base B , we write $\text{O}(\mathbb{R}^i, E)$ for the Stiefel bundle over B with fibre at $b \in B$ the Stiefel manifold $\text{O}(\mathbb{R}^i, E_b)$ of isometric linear maps from \mathbb{R}^i to the fibre E_b of E at $b \in B$.

Let H be the Hopf bundle over the real projective space $P(V)$ of V . Now there is a natural projection

$$\pi : O(V, H \otimes V) \rightarrow PO(V)$$

taking an isometric linear map $v \mapsto u \otimes g(v) : V \rightarrow \mathbb{R}u \otimes V$, where $g \in O(V)$, in the fibre over $[u] \in P(V)$ to $[g] \in PO(V)$. The pullback π^*L is identified with the lift of H (because $(-u) \otimes g(v) = u \otimes (-g)v$).

In terms of $x = e(H) \in H^1(P(V); \mathbb{F}_2)$, the total Stiefel-Whitney class of $H \otimes V$ is equal to $(1+x)^{2^r} = 1$, so that $w_j(H \otimes V) = 0$ for $j \geq 1$.

For each i , $0 \leq i < 2^r$, the fibration

$$p_i : O(\mathbb{R}^{i+1}, H \otimes V) \rightarrow O(\mathbb{R}^i, H \otimes V)$$

(restricting from $\mathbb{R}^{i+1} = \mathbb{R}^i \oplus \mathbb{R}$ to \mathbb{R}^i and interpreted when $i = 0$ as the projection $p_0 : S(H \otimes V) \rightarrow P(V)$) is the sphere bundle $S(\zeta_i)$ of the complementary $(2^r - i)$ -dimensional vector bundle ζ_i over $O(\mathbb{R}^i, H \otimes V)$. Thus $\zeta_i \oplus \mathbb{R}^i$ is the pullback of $H \otimes V$, and $e(\zeta_i) = w_{2^r-i}(\zeta_i)$ is the lift of $w_{2^r-i}(H \otimes V)$, which is zero. Hence, from the Gysin sequence of the sphere bundle, the induced homomorphism

$$p_i^* : H^*(O(\mathbb{R}^i, H \otimes V); \mathbb{F}_2) \rightarrow H^*(O(\mathbb{R}^{i+1}, H \otimes V); \mathbb{F}_2)$$

is injective. In the top dimension $i = 2^r - 1$, when ζ_i is a line bundle with $w_1 = 0$, p_i is a trivial bundle with fibre S^0 and p_i^* is an injection on each component.

Hence, $(\pi^*e(L))^{2^r-1}$, which is the image of $e(H)^{2^r-1} = x^{2^r-1} \in H^*(P(V); \mathbb{F}_2)$ under the composition

$$p_0 \circ \cdots \circ p_{2^r-1} : O(V, H \otimes V) = O(\mathbb{R}^{2^r}, H \otimes V) \rightarrow P(V) = O(\mathbb{R}^0, H \otimes V),$$

is non-zero on each of the two components. \square

Remark A.2. Simpler proofs can be given in the low dimensional cases $V = \mathbb{R}^{2^r}$, $r = 1, 2, 3$. In those dimensions there is a bilinear map $\cdot : V \times V \rightarrow V$ satisfying $\|u \cdot v\| = \|u\| \cdot \|v\|$ for $u, v \in V$, given by complex, quaternionic or Cayley multiplication. This determines a map $S(V) \rightarrow O(V)$ taking $u \in S(V)$ to the orthogonal transformation $v \mapsto u \cdot v$, which induces an embedding of $P(V)$ into $PO(V)$. The line bundle L restricts to the Hopf bundle over $P(V)$, and so we see that $e(L)^{2^r-1} \neq 0$.

Remark A.3. A result of Gitler and Handel [5, Theorem 1.6] on projective Stiefel manifolds can be established by the same method. For $V = \mathbb{R}^{2^r n}$, with n odd, consider, for $1 \leq l \leq n$, the projective Stiefel manifold $PO(\mathbb{R}^{2^r l}, V) = O(\mathbb{R}^{2^r l}, V)/\{\pm 1\}$ and the associated real line bundle L . Then $e(L)^{2^r m-1}$ is non-zero if $\binom{n}{j}$ is even for $n - m < j < l$, and $e(L)^{2^r m} = 0$ if $n - m < l$ and $\binom{n}{m}$ is odd.

For $e(\zeta_i)$ is the lift of $w_{2^r n-i}(H \otimes V) = \binom{2^r n}{2^r n-i} x^{2^r n-i}$, which is 0 if i is not divisible by 2^r and equal to $\binom{n}{n-j} x^{2^r(n-j)}$ if $i = 2^r j$; and the bundle $L \otimes V$ has a canonical trivial summand $\mathbb{R}^{2^r l}$, so that $w_i(L \otimes V) = 0$ for $i > 2^r(n-l)$.

REFERENCES

- [1] H. Adams, J. Bush and F. Frick, *Metric thickenings, Borsuk–Ulam theorems, and orbitopes*. *Mathematika* **66** (2020), 79–102.
- [2] H. Adams, J. Bush and F. Frick, *The topology of projective codes and the distribution of zeros of odd maps*. arXiv math.GT 2106.14677 (2021).
- [3] P. F. Baum and W. Browder, *The cohomology of quotients of classical groups*. *Topology* **3** (1965), 305–336.

- [4] B. Bukh and C. Cox, *Nearly orthogonal vectors and small antipodal spherical codes*. Israel J. Math. **238** (2020), 359–388.
- [5] S. Gitler and D. Handel, *The projective Stiefel manifolds–I*. Topology **7** (1968), 39–45.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ABERDEEN, ABERDEEN AB24 3UE, UK
Email address: `m.crabb@abdn.ac.uk`