

**FELL BUNDLES OVER A COUNTABLE DISCRETE GROUP
AND STRONG MORITA EQUIVALENCE FOR INCLUSIONS OF
 C^* -ALGEBRAS**

KAZUNORI KODAKA

ABSTRACT. We consider two saturated Fell bundles over a countable discrete group, whose unit fibers are σ -unital C^* -algebras. Then by taking the reduced cross-sectional C^* -algebras, we get two inclusions of C^* -algebras. We suppose that they are strongly Morita equivalent as inclusions of C^* -algebras. Also, we suppose that one of the inclusions of C^* -algebras is irreducible, that is, the relative commutant of one of the unit fiber algebras, which is a σ -unital C^* -algebra, in the multiplier C^* -algebra of the reduced cross-sectional C^* -algebra is trivial. We show that the two saturated Fell bundles are then equivalent up to some automorphism of the group.

1. INTRODUCTION

Let G be a countable discrete group and let $\mathcal{A} = \{A_t\}_{t \in G}$ be a Fell bundle over G . Let $C_r^*(\mathcal{A})$ be the reduced cross-sectional C^* -algebra of \mathcal{A} and $A_e = A$, a C^* -algebra, where e is the unit element in G . Then we obtain an inclusion of C^* -algebras $A \subset C_r^*(\mathcal{A})$ and we call it the inclusion of C^* -algebras induced by \mathcal{A} . By Abadie and Ferraro [1, Sections 3, 4], it is easy to show that if Fell bundles $\mathcal{A} = \{A_t\}_{t \in G}$ and $\mathcal{B} = \{B_t\}_{t \in G}$ over G are equivalent with respect to an equivalence bundle $\mathcal{X} = \{X_t\}_{t \in G}$ over G such that

$$\overline{\mathcal{A}\langle X_t, X_s \rangle} = A_{ts^{-1}} \quad , \quad \overline{\langle X_t, X_s \rangle_{\mathcal{B}}} = B_{t^{-1}s}$$

for any $t, s \in G$, then the inclusions of C^* -algebras induced by \mathcal{A} and \mathcal{B} are strongly Morita equivalent, where $\overline{\mathcal{A}\langle X_t, X_s \rangle}$ means the closure of linear span of the set

$$\{\mathcal{A}\langle x, y \rangle \mid x \in X_t, y \in X_s\}$$

and $\overline{\langle X_t, X_s \rangle_{\mathcal{B}}}$ means the closure of the same set as above.

In this paper, we shall show the inverse direction as follows: Let $\mathcal{A} = \{A_t\}_{t \in G}$ and $\mathcal{B} = \{B_t\}_{t \in G}$ be saturated Fell bundles over G . We suppose that $A_e = A$ and $B_e = B$ are σ -unital C^* -algebras and that $A' \cap C_r^*(\mathcal{A}) = \mathbf{C}1$. If the inclusions of C^* -algebras induced by \mathcal{A} and \mathcal{B} are strongly Morita equivalent, then there is an automorphism f of G such that \mathcal{A} and \mathcal{B}^f are equivalent with respect to an equivalence bundle $\mathcal{X} = \{X_t\}_{t \in G}$ such that

$$\overline{\mathcal{A}\langle X_t, X_s \rangle} = A_{ts^{-1}} \quad , \quad \overline{\langle X_t, X_s \rangle_{\mathcal{B}}} = B_{t^{-1}s}$$

for any $t, s \in G$, where \mathcal{B}^f is a Fell bundle over G induced by $\mathcal{B} = \{B_t\}_{t \in G}$ and f , that is, $\mathcal{B}^f = \{B_{f(t)}\}_{t \in G}$.

We prove this result in the following way: Let \mathbf{K} be the C^* -algebra of all compact operators on a countably infinite dimensional Hilbert space. Let \mathcal{A}^S and \mathcal{B}^S be the Fell bundles over G induced by \mathcal{A}, \mathbf{K} and \mathcal{B}, \mathbf{K} , respectively. Then since A and B are σ -unital, \mathcal{A}^S and \mathcal{B}^S satisfy the assumptions of Exel [3, Theorem 7.3]. Since

2010 *Mathematics Subject Classification*. Primary 46L05.

Key words and phrases. cross-sectional C^* -algebras, equivalence bundles, Fell bundles, inclusions of C^* -algebras, strong Morita equivalence.

\mathcal{A}^S and \mathcal{B}^S are saturated, by [3, Theorem 7.3] there are twisted actions (α, w_α) and (β, w_β) of G on the C^* -algebras $A \otimes \mathbf{K}$ and $B \otimes \mathbf{K}$, which are the unit fibre algebras of \mathcal{A}^S and \mathcal{B}^S , such that \mathcal{A}^S and \mathcal{B}^S are isomorphic to the semidirect product bundles of $A \otimes \mathbf{K}, G$ and $B \otimes \mathbf{K}, G$ constructed by (α, w_α) and (β, w_β) as Fell bundles over G , respectively. Since the inclusions of C^* -algebras induced by \mathcal{A} and \mathcal{B} are strongly Morita equivalent, so are the inclusions of C^* -algebras induced by \mathcal{A}^S and \mathcal{B}^S . Hence the inclusions of C^* -algebras induced by (α, w_α) and (β, w_β) are strongly Morita equivalent. Then since the inclusions are irreducible, by [6, Theorem 5.5] there is an automorphism f of G such that (α, w_α) and (β^f, w_β^f) are strongly Morita equivalent, where (β^f, w_β^f) is the twisted action of G on $B \otimes \mathbf{K}$ induced by (β, w_β) and f , that is,

$$\beta_t^f = \beta_{f(t)} \quad , \quad w_\beta^f(t, s) = w_\beta(f(t), f(s))$$

for any $t, s \in G$. Using this, we can prove the result.

2. PRELIMINARIES

Let A be a C^* -algebra and we denote by $M(A)$ the multiplier C^* -algebra of A . Let α be an automorphism of A . Then there is a unique strictly continuous automorphism of $M(A)$ extending α by Jensen and Thomsen [4, Corollary 1.1.15]. We denote it by $\underline{\alpha}$.

Let G be a countable discrete group and e the unit element in G . Let $\mathcal{A} = \{A_t\}_{t \in G}$ be a Fell bundle over G and let $A_e = A$, a C^* -algebra. Also, let $\mathcal{B} = \{B_t\}_{t \in G}$ be a Fell bundle over G and let $B_e = B$, a C^* -algebra. Following Abadie and Ferraro [1, Definitions 2.1 and 2.2], we give the definition of an equivalence bundle:

Definition 2.1. (1) A *right Hilbert \mathcal{B} -bundle* is a complex Banach bundle over G , $\mathcal{X} = \{X_t\}_{t \in G}$ with continuous maps

$$\mathcal{X} \times \mathcal{B} \rightarrow \mathcal{X}, (x, b) \mapsto xb \quad \text{and} \quad \langle -, - \rangle_{\mathcal{B}} : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{B}, (x, y) \mapsto \langle x, y \rangle_{\mathcal{B}}$$

such that:

- (1R) $X_r B_s \subset X_{rs}$ and $\langle X_r, X_s \rangle_{\mathcal{B}} \subset B_{r^{-1}s}$ for all $r, s \in G$.
- (2R) $X_r \times B_s \rightarrow X_{rs}, (x, b) \mapsto xb$ is bilinear for all $r, s \in G$.
- (3R) $X_s \rightarrow B_{r^{-1}s}, y \mapsto \langle x, y \rangle_{\mathcal{B}}$ is linear for all $x \in X_r$ and $s \in G$.
- (4R) $\langle x, yb \rangle_{\mathcal{B}} = \langle x, y \rangle_{\mathcal{B}} b$ and $\langle x, y \rangle_{\mathcal{B}}^* = \langle y, x \rangle_{\mathcal{B}}$ for all $x, y \in \mathcal{X}$ and $b \in \mathcal{B}$.
- (5R) $\langle x, x \rangle_{\mathcal{B}} \geq 0$ for all $x \in \mathcal{X}$ and $\langle x, x \rangle_{\mathcal{B}} = 0$ implies $x = 0$. Beside, each fiber X_t is complete with respect to the norm $x \mapsto \|\langle x, x \rangle_{\mathcal{B}}\|^{1/2}$.
- (6R) For all $x \in \mathcal{X}, \|x\|^2 = \|\langle x, x \rangle_{\mathcal{B}}\|$.
- (7R) $\overline{\{\langle X_s, X_s \rangle_{\mathcal{B}} \mid s \in G\}} = B_e$.

(2) A *left Hilbert \mathcal{A} -bundle* is a complex Banach bundle over G , $\mathcal{X} = \{X_t\}_{t \in G}$ with continuous maps

$$\mathcal{A} \times \mathcal{X} \rightarrow \mathcal{X}, (a, x) \mapsto ax \quad \text{and} \quad \langle -, - \rangle_{\mathcal{A}} : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{A}, (x, y) \mapsto \langle x, y \rangle_{\mathcal{A}}$$

such that:

- (1L) $A_r X_s \subset X_{rs}$ and $\langle X_r, X_s \rangle_{\mathcal{A}} \subset A_{r^{-1}s}$ for all $r, s \in G$.
- (2L) $A_r \times X_s \rightarrow X_{rs}, (a, x) \mapsto ax$ is bilinear for all $r, s \in G$.
- (3L) $X_s \rightarrow A_{r^{-1}s}, y \mapsto \langle a, y \rangle_{\mathcal{A}}$ is linear for all $x \in X_r$ and $s \in G$.
- (4L) $\langle ax, y \rangle_{\mathcal{A}} = a \langle x, y \rangle_{\mathcal{A}}$ and $\langle x, y \rangle_{\mathcal{A}}^* = \langle y, x \rangle_{\mathcal{A}}$ for all $x, y \in \mathcal{X}$ and $a \in \mathcal{A}$.
- (5L) $\langle x, x \rangle_{\mathcal{A}} \geq 0$ for all $x \in \mathcal{X}$ and $\langle x, x \rangle_{\mathcal{A}} = 0$ implies $x = 0$. Beside, each fiber X_t is complete with respect to the norm $x \mapsto \|\langle x, x \rangle_{\mathcal{A}}\|^{1/2}$.
- (6L) For all $x \in \mathcal{X}, \|x\|^2 = \|\langle x, x \rangle_{\mathcal{A}}\|$.
- (7L) $\overline{\{\langle X_s, X_s \rangle_{\mathcal{A}} \mid s \in G\}} = A_e$.

(3) We say that \mathcal{X} is an $\mathcal{A} - \mathcal{B}$ -equivalence bundle if \mathcal{X} is both a left Hilbert \mathcal{A} -bundle, a right Hilbert \mathcal{B} -bundle and ${}_A\langle x, y \rangle_z = x\langle y, z \rangle_B$ for all $x, y, z \in \mathcal{X}$. Besides, we say \mathcal{A} is equivalent to \mathcal{B} if there exists an $\mathcal{A} - \mathcal{B}$ -equivalence bundle.

Definition 2.2. Let $\mathcal{A} = \{A_t\}_{t \in G}$ be a Fell bundle over G . We say that $\mathcal{A} = \{A_t\}_{t \in G}$ is saturated if $\overline{A_t A_t^*} = A_e$ for any $t \in G$.

Let $\mathcal{A} = \{A_t\}_{t \in G}$ be a saturated Fell bundle over G and let $A_e = A$, a C^* -algebra. We suppose that A is a σ -unital C^* -algebra.

Let $C_r^*(\mathcal{A})$ be the reduced cross-sectional C^* -algebra of \mathcal{A} . Then for any $t \in G$, A_t is regarded as a closed subspace of $C_r^*(\mathcal{A})$ since G is discrete.

Let \mathbf{K} be the C^* -algebra of all compact operators on a countably infinite dimensional Hilbert space. Let $\{e_{ij}\}_{i, j \in \mathbf{N}}$ be a system of matrix units of \mathbf{K} .

Let $A_t \otimes \mathbf{K}$ be the closure of linear span of the subset

$$\{x \otimes k \in C_r^*(\mathcal{A}) \otimes \mathbf{K} \mid x \in A_t, k \in \mathbf{K}\}.$$

Let $\mathcal{A}^S = \{A_t \otimes \mathbf{K}\}_{t \in G}$. Then \mathcal{A}^S is a saturated Fell bundle over G and $A \otimes \mathbf{K}$ is its unit fibre algebra. Clearly $C_r^*(\mathcal{A}^S) = C_r^*(\mathcal{A}) \otimes \mathbf{K}$. Since \mathcal{A} is saturated, we can regard A_t as an $A - A$ -equivalence bimodule for any $t \in G$ and by the definition of the product in $C_r^*(\mathcal{A}^S)$ we can regard $A_t \otimes \mathbf{K}$ as the tensor product of the $A - A$ -equivalence bimodule A_t and the trivial $\mathbf{K} - \mathbf{K}$ -equivalence bimodule \mathbf{K} , which is $A \otimes \mathbf{K} - A \otimes \mathbf{K}$ -equivalence bimodule for any $t \in G$. Thus \mathcal{A}^S is a saturated Fell bundle over G . Since $A \otimes \mathbf{K}$ is σ -unital, by [3, Theorem 7.3], there is a twisted action (α, w_α) of G on $A \otimes \mathbf{K}$ such that \mathcal{A}^S is isomorphic to the semidirect product bundle over G induced by (α, w_α) as Fell bundles over G .

Lemma 2.1. *With the above notation, if $A' \cap M(C_r^*(\mathcal{A})) = \mathbf{C}1$, then $(A \otimes \mathbf{K})' \cap M(A \rtimes_{\alpha, w_\alpha, r} G) = \mathbf{C}1$.*

Proof. Since $A' \cap M(C_r^*(\mathcal{A})) = \mathbf{C}1$, by [6, Lemma 3.1] $(A \otimes \mathbf{K})' \cap M(C_r^*(\mathcal{A}) \otimes \mathbf{K}) = \mathbf{C}1$. Since \mathcal{A}^S is isomorphic to the semidirect product bundle over G of $A \otimes \mathbf{K}$ induced by (α, w_α) as Fell bundles over G , the inclusions $A \otimes \mathbf{K} \subset C_r^*(\mathcal{A}) \otimes \mathbf{K}$ and $A \otimes \mathbf{K} \subset (A \otimes \mathbf{K}) \rtimes_{\alpha, w_\alpha, r} G$ are isomorphic as inclusions of C^* -algebras. Thus $(A \otimes \mathbf{K})' \cap M((A \otimes \mathbf{K}) \rtimes_{\alpha, w_\alpha, r} G) = \mathbf{C}1$. \square

Let (α, w_α) and (β, w_β) be twisted actions of a countable discrete group G on C^* -algebras A and B , respectively. We suppose that (α, w_α) and (β, w_β) are strongly Morita equivalent with respect to a twisted action λ of G on an $A - B$ -equivalence bimodule X , that is, λ is a map from G to $\text{Aut}(X)$ satisfying the following:

- (1) $\alpha_t({}_A\langle x, y \rangle) = {}_A\langle \lambda_t(x), \lambda_t(y) \rangle$,
- (2) $\beta_t(\langle x, y \rangle_B) = \langle \lambda_t(x), \lambda_t(y) \rangle_B$,
- (3) $(\lambda_t \circ \lambda_s)(x) = w_\alpha(t, s)\lambda_{ts}(x)w_\beta(t, s)^*$

for any $t, s \in G$, $x, y \in X$, where we regard X as a Hilbert $M(A) - M(B)$ -bimodule as in [6, Preliminaries].

Let u and v be unitary representations of G to $M(A \rtimes_{\alpha, w_\alpha, r} G)$ and $M(B \rtimes_{\beta, w_\beta, r} G)$ implementing α and β , respectively, that is, $\alpha_t = \text{Ad}(u_t)$ and $\beta_t = \text{Ad}(v_t)$ for any $t \in G$. Let $\mathcal{A} = \{Au_t\}_{t \in G}$ and $\mathcal{B} = \{Bv_t\}_{t \in G}$ be the semidirect product Fell bundles over G induced by (α, w_α) and (β, w_β) , respectively.

For any $t \in G$, let $X_t = Xv_t$ as Banach spaces. We regard $\mathcal{X} = \{X_t\}_{t \in G}$ as an $\mathcal{A} - \mathcal{B}$ -equivalence bundle in the following (See [3] and [5]): For any $au_t \in Au_t$,

$bv_t \in Bv_t, xv_s \in Xv_s, yv_t \in Xv_t,$

$$(au_t)(xv_s) = a\lambda_t(x)w_\beta(t, s)v_{ts},$$

$$(xv_s)(bv_t) = x\beta_s(b)w_\beta(s, t)v_{st},$$

$${}_A\langle xv_s, yv_t \rangle = {}_A\langle x, (\lambda_s \circ \lambda_t^{-1})(y) \rangle w_\alpha(t, s^{-1})\alpha_{ts^{-1}}(w_\alpha(s, s^{-1}))^* u_{ts^{-1}},$$

$$\langle xv_s, yv_t \rangle_B = \beta_s^{-1}(\langle x, y \rangle_B)w_\beta(s^{-1}, s)^* w_\beta(s^{-1}, t)v_{s^{-1}t}.$$

Lemma 2.2. *With the above notation, $\mathcal{X} = \{X_t\}_{t \in G}$ is an \mathcal{A} – \mathcal{B} -equivalence bundle over G such that*

$$\overline{{}_A\langle X_t, X_s \rangle} = A_{ts^{-1}}, \quad \overline{\langle X_t, X_s \rangle_B} = B_{t^{-1}s}$$

for any $t, s \in G$.

Proof. By the definition of $\mathcal{X} = \{X_t\}_{t \in G}$, it is clear that \mathcal{X} has Conditions (1R)-(4R) and (1L)-(4L) in Definition 2.1. For any $xv_t \in Xv_t$,

$$\|\langle xv_t, xv_t \rangle_B\| = \|\beta_t^{-1}(\langle x, x \rangle_B)\| = \|\langle x, x \rangle_B\| = \|x\|^2 = \|xv_t\|^2,$$

$$\|{}_A\langle xv_t, xv_t \rangle\| = \|{}_A\langle x, x \rangle\| = \|xv_t\|^2.$$

Hence we see that $\{X_t\}_{t \in G}$ has Conditions (5R), (6R) and (5L), (6L) in Definition 2.1. Furthermore, since X is an A – B -equivalence bimodule, we can see that \mathcal{X} satisfies that

$$\overline{{}_A\langle X_t, X_s \rangle} = A_{ts^{-1}}, \quad \overline{\langle X_t, X_s \rangle_B} = B_{t^{-1}s}$$

for any $t, s \in G$. □

3. STRONG MORITA EQUIVALENCE

Let $\mathcal{A} = \{A_t\}_{t \in G}$ and $\mathcal{B} = \{B_t\}_{t \in G}$ be saturated Fell bundles over G . Let $A_e = A$ and $B_e = B$ be C^* -algebras. Let \mathcal{A}^S and \mathcal{B}^S be the saturated Fell bundles over G induced by \mathcal{A}, \mathbf{K} and \mathcal{B}, \mathbf{K} , respectively. Let (α, w_α) and (β, w_β) be the twisted actions of G on $A \otimes \mathbf{K}$ and $B \otimes \mathbf{K}$ such that \mathcal{A}^S and \mathcal{B}^S are isomorphic to the semidirect product bundles of $A \otimes \mathbf{K}$ and $B \otimes \mathbf{K}$ induced by (α, w_α) and (β, w_β) , which are defined in Section 2, respectively. We suppose that $A \subset C_r^*(\mathcal{A})$ and $B \subset C_r^*(\mathcal{B})$ are strongly Morita equivalent and that $A' \cap M(C_r^*(\mathcal{A})) = \mathbf{C}1$.

Lemma 3.1. *With the above notation and assumptions, there is an automorphism f of G such that (α, w_α) is strongly Morita equivalent to (β, w_β^f) , (β^f, w_β^f) is the twisted action of G on $B \otimes \mathbf{K}$ induced by (β, w_β) and f , which is defined by*

$$\beta_t^f = \beta_{f(t)}, \quad w_\beta^f(t, s) = w_\beta(f(t), f(s))$$

for any $t, s \in G$.

Proof. Since $A' \cap M(C_r^*(\mathcal{A})) = \mathbf{C}1$, by Lemma 2.1

$$(A \otimes \mathbf{K})' \cap M((A \otimes \mathbf{K}) \rtimes_{\alpha, w_\alpha, r} G) = \mathbf{C}1.$$

Hence by [6, Theorem 5.5] we obtain the conclusion. □

Let $\mathcal{B}^{S, f} = \{B_{f(t)} \otimes \mathbf{K}\}_{t \in G}$ be the semidirect product bundle of $B \otimes \mathbf{K}$ induced by (β^f, w_β^f) . Then by Lemma 2.2 and Lemma 3.1, there is an \mathcal{A}^S – $\mathcal{B}^{S, f}$ -equivalence bundle $\mathcal{Y} = \{Y_t\}_{t \in G}$ such that

$$\overline{{}_A\langle Y_t, Y_s \rangle} = A_{ts^{-1}} \otimes \mathbf{K} \quad , \quad \overline{\langle Y_t, Y_s \rangle_{\mathcal{B}^{S, f}}} = B_{f(t^{-1}s)} \otimes \mathbf{K}$$

for any $t, s \in G$.

Let $\mathcal{B}^f = \{B_{f(t)}\}_{t \in G}$ be the Fell bundle over G induced by $\mathcal{B} = \{B_t\}_{t \in G}$ and the automorphism f of G .

Lemma 3.2. *With the above notation, \mathcal{A} and \mathcal{B}^f are equivalent with respect to an $\mathcal{A} - \mathcal{B}^f$ -equivalence bundle $\mathcal{X} = \{X_t\}_{t \in G}$ such that*

$$\overline{\mathcal{A}\langle X_t, X_s \rangle} = A_{ts^{-1}} \quad , \quad \overline{\langle X_t, X_s \rangle_{\mathcal{B}^f}} = B_{f(t^{-1}s)}$$

Proof. Let $X_t = (1 \otimes e_{11})Y_t(1 \otimes e_{11})$ for any $t \in G$. Let $t, s \in G$ and let $a_t \in A_t$, $b_t \in B_t$, $x_s \in Y_s$ and $x_t, y_t \in Y_t$. Then

$$\begin{aligned} a_t(1 \otimes e_{11})x_s(1 \otimes e_{11}) &= (1 \otimes e_{11})(a_t \otimes e_{11})x_s(1 \otimes e_{11}) \in X_{ts}, \\ (1 \otimes e_{11})x_s(1 \otimes e_{11})b_t &= (1 \otimes e_{11})x_s(b_t \otimes e_{11})(1 \otimes e_{11}) \in X_{st}, \end{aligned}$$

$$\begin{aligned} &{}_{\mathcal{A}^s}\langle (1 \otimes e_{11})x_s(1 \otimes e_{11}), (1 \otimes e_{11})y_t(1 \otimes e_{11}) \rangle \\ &= (1 \otimes e_{11})_{\mathcal{A}^s}\langle x_s(1 \otimes e_{11}), y_t(1 \otimes e_{11}) \rangle(1 \otimes e_{11}) \in (1 \otimes e_{11})(A_{st^{-1}} \otimes \mathbf{K})(1 \otimes e_{11}), \end{aligned}$$

where we identify $(1 \otimes e_{11})(A_{st^{-1}} \otimes \mathbf{K})(1 \otimes e_{11})$ with $A_{st^{-1}}$. Also,

$$\begin{aligned} &\langle (1 \otimes e_{11})x_s(1 \otimes e_{11}), (1 \otimes e_{11})y_t(1 \otimes e_{11}) \rangle_{\mathcal{B}^{s,f}} \\ &= (1 \otimes e_{11})\langle (1 \otimes e_{11})x_s, (1 \otimes e_{11})y_t \rangle_{\mathcal{B}^{s,f}}(1 \otimes e_{11}) \in (1 \otimes e_{11})(B_{f(s^{-1}t)} \otimes \mathbf{K})(1 \otimes e_{11}), \end{aligned}$$

where we identify $(1 \otimes e_{11})(B_{f(s^{-1}t)} \otimes \mathbf{K})(1 \otimes e_{11})$ with $B_{f(s^{-1}t)}$. Hence \mathcal{X} has Conditions (1R)-(4R) and (1L)-(4L) in Definition 2.1. Furthermore,

$$\begin{aligned} &{}_{\mathcal{A}^s}\langle (1 \otimes e_{11})x_t(1 \otimes e_{11}), (1 \otimes e_{11})x_t(1 \otimes e_{11}) \rangle \\ &= (1 \otimes e_{11})_{\mathcal{A}^s}\langle x_t(1 \otimes e_{11}), x_t(1 \otimes e_{11}) \rangle(1 \otimes e_{11}), \\ &\langle (1 \otimes e_{11})x_t(1 \otimes e_{11}), (1 \otimes e_{11})x_t(1 \otimes e_{11}) \rangle_{\mathcal{B}^{s,f}} \\ &= (1 \otimes e_{11})\langle (1 \otimes e_{11})x_t, (1 \otimes e_{11})x_t \rangle_{\mathcal{B}^{s,f}}(1 \otimes e_{11}). \end{aligned}$$

These equations implies that \mathcal{X} has Conditions (5R) and (5L) in Definition 2.1. It is clear that \mathcal{X} has Conditions (6R) and (6L) in Definition 2.1. Moreover, let $c, d \in B \otimes \mathbf{K}$. Then

$$\begin{aligned} &{}_{\mathcal{A}^s}\langle (1 \otimes e_{11})x_sc(1 \otimes e_{11}), (1 \otimes e_{11})y_td(1 \otimes e_{11}) \rangle \\ &= (1 \otimes e_{11})_{\mathcal{A}^s}\langle x_sc(1 \otimes e_{11})d^*, y_t \rangle(1 \otimes e_{11}). \end{aligned}$$

Since $1 \otimes e_{11}$ is full in $B \otimes \mathbf{K}$, that is, $\overline{(B \otimes \mathbf{K})(1 \otimes e_{11})(B \otimes \mathbf{K})} = B \otimes \mathbf{K}$ and $\overline{Y_t(B \otimes \mathbf{K})} = Y_t$ by [2, Proposition 1.7(i)],

$$\overline{{}_{\mathcal{A}^s}\langle (1 \otimes e_{11})x_s(1 \otimes e_{11}), (1 \otimes e_{11})y_t(1 \otimes e_{11}) \rangle} = A_{st^{-1}} \otimes e_{11}.$$

In the same way, we can see that

$$\overline{\langle (1 \otimes e_{11})x_s(1 \otimes e_{11}), (1 \otimes e_{11})y_t(1 \otimes e_{11}) \rangle_{\mathcal{B}^{s,f}}} = B_{f(s^{-1}t)} \otimes e_{11}.$$

Hence

$$\overline{\mathcal{A}\langle X_t, X_s \rangle} = A_{ts^{-1}} \quad , \quad \overline{\langle X_t, X_s \rangle_{\mathcal{B}^f}} = B_{f(t^{-1}s)}$$

for any $t, s \in G$. Since for any $x, y, z \in \mathcal{Y}$, ${}_{\mathcal{A}^s}\langle x, y \rangle z = x \langle y, z \rangle_{\mathcal{B}^{s,f}}$, we can see that

$$\begin{aligned} &{}_{\mathcal{A}}\langle (1 \otimes e_{11})x(1 \otimes e_{11}), (1 \otimes e_{11})y(1 \otimes e_{11}) \rangle(1 \otimes e_{11})z(1 \otimes e_{11}) \\ &= (1 \otimes e_{11})x(1 \otimes e_{11})\langle (1 \otimes e_{11})y(1 \otimes e_{11}), (1 \otimes e_{11})z(1 \otimes e_{11}) \rangle_{\mathcal{B}^f} \end{aligned}$$

for any $x, y, z \in \mathcal{Y}$. It follows that \mathcal{X} is an $\mathcal{A} - \mathcal{B}^f$ -equivalence bundle. Therefore, we obtain the conclusion. \square

Theorem 3.3. *Let G be a countable discrete group and let $\mathcal{A} = \{A_t\}_{t \in G}$ and $\mathcal{B} = \{B_t\}_{t \in G}$ be saturated Fell bundles over G . We suppose that $A_e = A$ and $B_e = B$ are σ -unital C^* -algebras and that $A' \cap C_r^*(A) = \mathbf{C}1$, where e is the unit element in G and $C_r^*(A)$ is the reduced cross-sectional C^* -algebra of \mathcal{A} . If the inclusions of C^* -algebras induced by \mathcal{A} and \mathcal{B} are strongly Morita equivalent, then*

there is an automorphism f of G such that \mathcal{A} and \mathcal{B}^f are equivalent with respect to an $\mathcal{A} - \mathcal{B}^f$ -equivalence bundle $\mathcal{X} = \{X_t\}_{t \in G}$ over G such that

$$\overline{\mathcal{A}\langle X_t, X_s \rangle} = A_{ts^{-1}} \quad , \quad \overline{\langle X_t, X_s \rangle_{\mathcal{B}^f}} = B_{f(t^{-1}s)}$$

for any $t, s \in G$, where \mathcal{B}^f is a Fell bundle over G induced by $\mathcal{B} = \{B_t\}_{t \in G}$ and f , that is, $\mathcal{B}^f = \{B_{f(t)}\}_{t \in G}$.

Proof. This is immediate by Lemmas 3.1, 3.2 and the discussions before Lemma 3.2. \square

REFERENCES

- [1] F. Abadie and D. Ferraro, *Equivalence of Fell bundles over groups*, J. Operator Theory **81** (2019), 273–319.
- [2] L. G. Brown, J. Mingo and N-T. Shen, *Quasi-multipliers and embeddings of Hilbert C^* -bimodules*, Can. J. Math., **46** (1994), 1150–1174.
- [3] R. Exel, *Twisted partial actions: A classification of regular C^* -algebraic bundles*, Proc. London. Math. Soc., **74** (1997), 417–443.
- [4] K. K. Jensen and K. Thomsen, *Elements of KK -theory*, Birkhäuser, 1991.
- [5] T. Kajiwara and Y. Watatani, *Crossed products of Hilbert C^* -bimodules by countable discrete groups*, Proc. Amer. Math. Soc., **126** (1998), 841–851.
- [6] K. Kodaka, *Strong Morita equivalence for inclusions of C^* -algebras induced by twisted actions of countable discrete group*, preprint, arXiv:1910.03774, Math. Scand. to appear.
- [7] K. Kodaka and T. Teruya, *The strong Morita equivalence for inclusions of C^* -algebras and conditional expectations for equivalence bimodules*, J. Aust. Math. Soc., **105** (2018), 103–144.

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, RYUKYU UNIVERSITY, NISHIHARA-CHO, OKINAWA, 903-0213, JAPAN

E-mail address: kodaka@math.u-ryukyu.ac.jp