

Inverse boundary value problems for polyharmonic operators with non-smooth coefficients

R.M. Brown*

Department of Mathematics
University of Kentucky
Lexington, KY 40506 0027

L.D. Gauthier

Department of Mathematics
University of Kentucky
Lexington, KY 40506 0027

Abstract

We consider inverse boundary value problems for polyharmonic operators and in particular, the problem of recovering the coefficients of terms up to order one. The main interest of our result is that it further relaxes the regularity required to establish uniqueness. The proof relies on an averaging technique introduced by Haberman and Tataru for the study of an inverse boundary value problem for a second order operator.

1 Introduction

The goal of this paper is to study an inverse boundary value problem for a family of operators whose principal part is the polyharmonic operator $(-\Delta)^m$ for $m \geq 2$. We fix a bounded domain $\Omega \subset \mathbb{R}^d$, $d \geq 3$, and assume the boundary of Ω is at least Lipschitz. We consider an operator L given by

$$Lu = (-\Delta)^m u + Q \cdot Du + qu \quad (1.1)$$

where we use $D = -i\nabla$ and we assume that the coefficients Q and q lie in Sobolev spaces of negative order. To give an example of the problems we study, we consider the Navier boundary value problem for a polyharmonic operator with lower order terms

$$\begin{cases} Lu = 0, & \text{in } \Omega \\ (-\Delta)^j u = \phi_j, & \text{on } \partial\Omega, j = 0, \dots, m-1. \end{cases} \quad (1.2)$$

For this solution u , we find $\frac{\partial}{\partial\nu}(-\Delta)^j u$, $j = 0, \dots, m-1$ where $\partial/\partial\nu$ is the derivative in the direction of outer unit normal. The problem we study is whether the coefficients q and Q are uniquely determined by the Cauchy data at the boundary

$$\left(u, \frac{\partial u}{\partial\nu}, -\Delta u, \dots, (-\Delta)^{m-1} u, \frac{\partial}{\partial\nu}(-\Delta)^{m-1} u\right)$$

*R. Brown is partially supported by a grant from the Simons Foundation (#422756).

for all solutions. Our main result gives regularity conditions on the coefficients that guarantee a positive answer to this question. The primary interest of our work is that we lower the smoothness required to establish this result by up to one derivative as compared to earlier work of Assylbekov and Iyer [1].

We briefly review some of the history of inverse boundary value problems. An early study of inverse boundary value problems for second order operators was carried out by Calderón [13] who considered a linearized problem and introduced the idea of looking at what we now call complex geometrical optics (CGO) solutions. Sylvester and Uhlmann [31] obtained the first uniqueness result for the full nonlinear problem. They studied the inverse conductivity problem where the goal is to show that a scalar function γ (the conductivity) is uniquely determined from the Cauchy data for solutions of the operator $\operatorname{div} \gamma \nabla$. The study of inverse boundary value problems for the biharmonic operator was begun by Krupchyk, Lassas and Uhlmann [22] who establish that first order terms are uniquely determined by the Cauchy data. The stability of the solution of this problem was considered by Choudhury and Krishnan [15]. Krupchyk, Lassas and Uhlmann [21] study inverse boundary value problems for the biharmonic operators with partial data on the boundary. Khrishnan and Ghosh [17] extend this work to consider inverse boundary value problems for the polyharmonic operators with partial data. Bhattacharyya and Ghosh [6, 7] consider second order perturbations of biharmonic and polyharmonic. The most recent paper [7] establishes the uniqueness of a second order term from boundary data. Work of Yan [34] establishes uniqueness for first order perturbations of a biharmonic operator on transversally isotropic manifolds. She extends this work to give a constructive recovery procedure for continuous potentials [35]. Serov [30] establishes a Borg-Levinson theorem which shows that certain coefficients for biharmonic operator are determined by boundary spectral information. His proof proceeds by establishing a connection between the boundary spectral information and a Dirichlet to Neumann map.

Our interest here is to study the regularity hypotheses that are needed to show that the coefficients are uniquely determined by the Cauchy data at the boundary. We review some of the history related to smoothness hypotheses needed to obtain uniqueness in inverse boundary value problems for second order equations. An important technique in the study of inverse boundary value problems for the conductivity equation $\operatorname{div} \gamma \nabla$ is a change of dependent variable which transforms the conductivity equation into a Schrödinger equation. A calculation shows that if $\operatorname{div} \gamma \nabla u = 0$, then $v = \sqrt{\gamma} u$ will be a solution of $(-\Delta + q)v = 0$ with $q = \Delta \sqrt{\gamma} / \sqrt{\gamma}$. Thus results for the conductivity equation with γ having s derivatives is implied by a result for Schrödinger operators with potentials that have $s - 2$ derivatives, at least for $s \geq 1$. It will be helpful to keep this correspondence in mind in the discussion below. For the conductivity equation, work of Brown [11] and Brown and Torres [12] establish the uniqueness of the coefficient γ in the conductivity operator when γ has roughly $3/2$ derivatives. Päivärinta, Panchenko and Uhlmann [26] construct CGO solutions for C^1 -conductivities but are not able to use these solutions to establish a uniqueness result. An important breakthrough was made by B. Haberman and D. Tataru [18] who establish uniqueness for C^1 -conductivities. Haberman and Tataru's innovation is to develop a family of spaces and show that in an average sense the behavior of the potentials is better than one might expect from looking at the pointwise behavior. This averaging gives better estimates than obtained, for example, by Päivärinta, Panchenko and Uhlmann

and leads to uniqueness for C^1 -conductivities. Further extensions include work of Caro and Rogers [14] who establish uniqueness for Lipschitz conductivities and an additional work of Haberman [19] that considers conductivities with one derivative in L^d (where d is the dimension) at least for $d = 3, 4$. Recent extensions of these ideas may be found in articles by Ham, Kwon and Lee [20] and Ponce-Vanegas [27, 28]. It is a conjecture of Uhlmann that the least regular Sobolev space for which we can prove uniqueness is $W^{1,d}$. However, there are few tools for treating conductivities which have less than one derivative when $d \geq 3$ nor is there an obvious path to constructing examples. If $d = 2$, Astala and Päiväranta [4] establish uniqueness for conductivities that are only bounded and measurable.

The key innovation of our research is to adapt the arguments of Haberman and Tataru to polyharmonic operators. We will consider operators whose principal part is the polyharmonic operator and include lower order terms involving at most one derivative as in (1.1). For these operators we are able to establish uniqueness for potentials q which lie in Sobolev spaces with smoothness index $-s > -m/2 - 1$. We believe that the cutoff in the smoothness index should be $-m$ for operators where the principal part is $(-\Delta)^m$. This is consistent with the result of Haberman and Tataru when $m = 1$. Our result allows us to get arbitrarily close to this cut off when $m = 2$. However, when $m \geq 3$ we believe that there is a large gap between our results and the optimal result.

The investigation of uniqueness in inverse problems for polyharmonic operators with coefficients of negative order was begun by Assylbekov [2, 3] and Assylbekov and Iyer [1]. In the work of Assylbekov and Iyer, the authors establish uniqueness when q lies in the Sobolev space $W^{-s,p}$ with $s < m/2$ and $p \geq 2d/m$ and Q lies in a Sobolev space $W^{-s+1,p}$ with $s < m/2$ and $p > 2d/m$, at least in the case $m < d$. We refer the interested reader to the work of Assylbekov and Iyer for the complete statement. The result of Assylbekov and Iyer is in the spirit of the work of the first author and Torres [12] who studied the inverse conductivity problem for conductivities with $3/2$ derivatives. A related result is due to Krupchyk and Uhlmann who consider polyharmonic operators with potentials in L^p spaces [23]. They are able to establish uniqueness when $q \in L^p$ for $p > d/2m$. Since L^p embeds into $W^{-s,q}$ for $1/p = 1/q + s/d$ our result below in Theorem 1.3 will give uniqueness for potentials in L^p for $p > d/m$ which is a weaker result for L^p spaces than Krupchyk and Uhlmann give. This is one of several indications that our result, while an advance over earlier work, is not the last word on this subject. It also remains an open problem to extend the techniques we discuss for nonsmooth first order perturbations to second order perturbations as studied in the works of Bhattacharyya and Ghosh mentioned earlier. We refer readers to the articles cited for additional work on these problems.

The main result of this work is the following result. The reader will need to refer to later sections of the paper for definitions of some of the notations occurring in this result.

Theorem 1.3. *Fix $m \geq 2$ and $d \geq 3$ and let $\Omega \subset \mathbb{R}^d$ be a bounded domain with smooth boundary. Let $L_j = (-\Delta)^m + Q^j \cdot D + q^j$ with $q^j \in \tilde{W}^{-s,p}(\Omega)$ and each component Q^j lies in $\tilde{W}^{-s+1,p}(\Omega)$ with $s < m/2 + 1$, $1/p + (s - m)/d < 0$, and $p \geq 2$. If each of the operators L_1 and L_2 have the same linear space of Cauchy data for (1.2), then $Q^1 = Q^2$ and $q^1 = q^2$.*

The remainder of the paper is devoted to the proof of this theorem as well as a more general statement where the hypothesis involving the Cauchy data is replaced by an assumption stated using quadratic forms. The paper is organized as follows. Section 2 gives notation

and definitions. Section 3 describes the spaces X^λ which were introduced by Haberman and Tataru [18] and are modeled on spaces used by Bourgain [9, 10] for the study of dispersive wave equations. We consider the operator obtained by conjugating the operator in (1.1) by exponentially growing harmonic functions and study its action on the X^λ spaces. Section 4 gives an extension of the averaging method of Haberman and Tataru which we use to average in X^λ spaces. In section 5, we describe the Cauchy data we use to establish uniqueness and relate the Cauchy data to a Dirichlet to Neumann map and to bilinear forms that we use in our uniqueness theorem. In section 6, we construct CGO solutions. We begin with an analog to the Päivärinta, Panchenko and Uhlmann result for second order equations [26, Theorem 1.1]. We then show how the methods of Haberman and Tataru give solutions which are better behaved. We use these solutions to give our uniqueness result. The hypothesis of this uniqueness result is stated in terms of bilinear forms. We remark that the smoothness of the domain Ω enters only in our definition of the Cauchy data and the Dirichlet to Neumann map. Our work with bilinear forms only requires the domain to be Lipschitz. Finally, an appendix collects a few facts about Sobolev spaces that are needed to work with our operators which have coefficients from Sobolev spaces of negative order.

The authors would like to thank K. Krupchyk for suggesting this problem and the referees whose careful reading of our manuscript led to several suggestions for improvement.

2 Notation

We will let Ω be a bounded, Lipschitz domain by which we mean a bounded connected open set whose boundary is locally the graph of a Lipschitz function (see, for example, Verchota [32] for a definition). In a few places, especially section 5, we will require the boundary to be smooth. We define the Fourier Transform on functions by

$$\hat{f}(\xi) = \int_{\mathbb{R}^d} e^{-ix \cdot \xi} f(x) dx.$$

We will let $\mathcal{S}(\mathbb{R}^d)$ be the Schwartz space on \mathbb{R}^d and $\mathcal{S}'(\mathbb{R}^d)$, the dual space of tempered distributions. We use $\langle \cdot, \cdot \rangle$ for the bilinear pairing of \mathcal{S}' and \mathcal{S} and the pairing of duality for various Sobolev spaces. We will use the notation $\langle x \rangle = (1 + |x|^2)^{1/2}$ and thus, we can write $((1 - \Delta)^{s/2} u)^\wedge(\xi) = \hat{u}(\xi) \langle \xi \rangle^s$.

Our hypotheses on the smoothness of the coefficients will be given in terms of Sobolev spaces. For $s \in \mathbb{R}$ and $1 < p < \infty$ we let $W^{s,p}(\mathbb{R}^d)$ be the Sobolev space of order s with the norm $\|u\|_{W^{s,p}(\mathbb{R}^d)} = \|((1 - \Delta)^{s/2} u)\|_{L^p(\mathbb{R}^d)}$. We also let $W^{s,p}(\Omega)$ denote the Sobolev space on Ω obtained by by restricting distributions in $W^{s,p}(\mathbb{R}^d)$ to Ω . We let $W_0^{s,p}(\Omega)$ denote the closure of $C_0^\infty(\Omega)$ in the norm of $W^{s,p}(\Omega)$. We define

$$\tilde{W}^{s,p}(\Omega) = \{u \in W^{s,p}(\mathbb{R}^d) : \text{supp}(u) \subset \bar{\Omega}\}.$$

Note that this notation is misleading since $\tilde{W}^{s,p}(\Omega)$ may not be a space of distributions in Ω . It is useful to note that for Lipschitz domains (and some more general domains) that $C_0^\infty(\Omega)$ is dense in the space $\tilde{W}^{s,p}(\Omega)$. See McLean [24, Theorem 3.29] for the case $p = 2$ and a proof that may be generalized to $p \in (1, \infty)$. The distinction between $W_0^{s,p}(\Omega)$ and

$\tilde{W}^{s,p}(\Omega)$ is subtle. For many values of s and p , the spaces agree. However, in certain cases the space $W_0^{s,p}(\Omega)$ will be a larger space. The norm in $\tilde{W}^{s,p}(\Omega)$ is the norm of a distribution which is zero outside $\bar{\Omega}$. The norm of an element u in $W_0^{s,p}(\Omega)$ is the infimum of the norm in \mathbb{R}^d over all possible extensions u to \mathbb{R}^d and may be smaller. The last function space we will need is the Hölder space $C^\theta(\mathbb{R}^d)$. For $\theta \in (0, 1]$, this space is the collection of functions on \mathbb{R}^d for which the norm

$$\|f\|_{C^\theta(\mathbb{R}^d)} = \sup_{x \in \mathbb{R}^d} |f(x)| + \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|^\theta}$$

is finite.

If the domain Ω is Lipschitz, then we have a characterization of the dual space of $W^{s,p}(\Omega)$,

$$W^{s,p}(\Omega)^* = \tilde{W}^{-s,p'}(\Omega), \quad 1 < p < \infty, s \geq 0. \quad (2.1)$$

The case $p = 2$ may be found in the monograph of McLean [24, Theorem 3.30] and his argument extends to $p \in (1, \infty)$. The spaces $\tilde{W}^{s,p}(\Omega)$ will be useful for the study of solutions with nonzero boundary data since for certain s and p we will be able to define the product of a function in a Sobolev space of positive order with the coefficients taken from Sobolev spaces of negative order. From the Sobolev embedding theorem and the product rule, we have that the bilinear map $(u, v) \rightarrow vD^\alpha u$ maps $W^{m,2}(\Omega) \times W^{m,2}(\Omega) \rightarrow W^{m-|\alpha|,t}(\Omega)$ provided $|\alpha| \leq m$ and where t is given by

$$\begin{cases} \frac{1}{t} = 1 - \frac{m}{d}, & \frac{m}{d} < \frac{1}{2} \\ \frac{1}{t} = \frac{1}{2}, & \frac{m}{d} > \frac{1}{2} \\ \frac{1}{t} > \frac{1}{2}, & \frac{m}{d} = \frac{1}{2}. \end{cases} \quad (2.2)$$

This is a special case of a more general result in the monograph of Runst and Sickel [29, §4.4.4, Theorems 1,2]. Thus, if $f \in \tilde{W}^{|\alpha|-m,t'}(\Omega)$ and $u \in W^{m,2}(\Omega)$, then we may define $fD^\alpha u \in \tilde{W}^{-m,2}(\Omega)$. If $v \in W^{m,2}(\Omega)$, then we define $\langle fD^\alpha u, v \rangle$ as equal to $\langle f, vD^\alpha u \rangle$ and we have the estimate

$$|\langle fD^\alpha u, v \rangle| \lesssim \|f\|_{\tilde{W}^{|\alpha|-m,t'}(\Omega)} \|u\|_{W^{m,2}(\Omega)} \|v\|_{W^{m,2}(\Omega)}. \quad (2.3)$$

We now describe a standard notion of weak solution that we will use to give a careful definition of solutions to the equation $Lu = 0$. We let $B_0 : W^{m,2}(\Omega) \times W^{m,2}(\Omega) \rightarrow \mathbb{C}$ be a form given by

$$B_0(u, v) = \int_{\Omega} \sum_{|\alpha|=|\beta|=m} a_{\alpha,\beta} D^\alpha u D^\beta v \, dx$$

with the property that

$$B_0(u, v) = \int_{\Omega} [(-\Delta)^m u] v \, dx, \quad u \in C^\infty(\Omega), v \in C_0^\infty(\Omega). \quad (2.4)$$

There are two common choices for the form B_0 . The first is useful for the Navier boundary value problem

$$B_0(u, v) = \begin{cases} \int_{\Omega} (-\Delta)^{m/2} u (-\Delta)^{m/2} v \, dx & \text{if } m \text{ is even} \\ \int_{\Omega} \nabla (-\Delta)^{(m-1)/2} u \cdot \nabla (-\Delta)^{(m-1)/2} v \, dx & \text{if } m \text{ is odd} \end{cases} \quad (2.5)$$

For some purposes, the form

$$B_0(u, v) = \int_{\Omega} \sum_{|\alpha|=m} \frac{m!}{\alpha!} D^\alpha u D^\alpha v \, dx. \quad (2.6)$$

is more convenient because it is easier to see that this form is coercive. We note that in our uniqueness theorem based on forms, Theorem 6.6, will hold for any choice of B_0 satisfying (2.4).

To define weak solutions to the equation $Lu = 0$ we require that

$$Q \in \tilde{W}^{1-m, t'}(\Omega) \quad \text{and} \quad q \in \tilde{W}^{-m, t'}(\Omega), \quad \text{with } t \text{ as in (2.2)}. \quad (2.7)$$

We say u is a *weak solution* of

$$Lu = (-\Delta)^m u + Q \cdot Du + qu = 0$$

if $u \in W^{m, 2}(\Omega)$ and

$$B_0(u, v) + \langle Q \cdot Du, v \rangle + \langle qu, v \rangle = 0, \quad v \in W_0^{m, 2}(\Omega).$$

The estimate (2.3) and the assumption (2.7) guarantee that the bilinear form for the equation is continuous on $W^{m, 2}(\Omega) \times W^{m, 2}(\Omega)$. We note that the condition (2.7) is close to the weakest condition that allows us to discuss weak solutions of the equation $Lu = 0$; we will need to impose additional regularity on Q and q in order to study the inverse problem that is the focus of this paper.

We will adopt the convention of using $a \lesssim b$ to mean that there is a constant C so that $a \leq Cb$. The constants C are allowed to depend on the L^p -index, the order of a Sobolev space, the dimension, the domain and the coefficients Q and q appearing in the operator. However, it is an important point that C will always be independent of h for h small as this is needed to evaluate limits as h tends to zero. We use $a \approx b$ to mean $a \lesssim b$ and $b \lesssim a$.

3 CGO solutions and X^λ spaces

In this section, we begin the construction of CGO solutions of (1.1). We let \mathcal{V} denote the collection of vectors

$$\mathcal{V} = \{\zeta \in \mathbb{C}^d : \zeta = \mu_1 + i\mu_2, \mu_j \in \mathbb{R}^d, \mu_i \cdot \mu_j = \delta_{ij}\}. \quad (3.1)$$

For $\zeta \in \mathcal{V}$ and $h \neq 0$, the function $e^{x \cdot \zeta / h}$ will be harmonic. For $0 < h \leq h_0 < 1$ we will construct solutions of $Lu = 0$ of the form

$$u(x) = e^{x \cdot \zeta / h} (a(x) + \psi(x)) \text{ in } \Omega \quad (3.2)$$

where the amplitude a will be smooth. If we substitute (3.2) into the equation $Lu = 0$ and multiply on the left by $h^{2m} e^{-x \cdot \zeta / h}$, we obtain an equation for ψ

$$L_\zeta \psi = -L_\zeta a. \quad (3.3)$$

We introduce the operator $P_\zeta(hD)$ defined by

$$P_\zeta(hD) = P(hD) = e^{-x \cdot \zeta/h} (-h^2 \Delta) e^{x \cdot \zeta/h},$$

where we will view this as a semi-classical operator with symbol $P_\zeta(\xi) = P(\xi) = |\xi|^2 - 2i\zeta \cdot \xi$. With this notation we may write

$$L_\zeta = P(hD)^m + h^{2m} Q \cdot \left(\frac{\zeta}{ih} + D \right) + h^{2m} q. \quad (3.4)$$

We will give estimates for solutions of the equation $L_\zeta u = f$, at least for h small. Since the symbol of the operator $P(hD)$ vanishes on a set of codimension 2, the operator is not elliptic and finding solutions is an interesting problem. To study the equation (3.3), it is useful to use function spaces that work well with $P(hD)$. Similar spaces were used by Bourgain [9] in his study of evolution equations. The spaces described below were introduced in the study of the inverse conductivity problem by Haberman and Tataru [18]. For $\lambda \in \mathbb{R}$, $\zeta \in \mathcal{V}$, and $h > 0$, we introduce a space $X_{h\zeta}^\lambda$ by

$$X_{h\zeta}^\lambda = X^\lambda = \{u \in \mathcal{S}'(\mathbb{R}^d) : \hat{u} \text{ is a function and } \|u\|_{X^\lambda} < \infty\}$$

where the norm in X^λ is given by

$$\|u\|_{X^\lambda}^2 = \int_{\mathbb{R}^d} |\hat{u}(\xi)|^2 (h + |P_\zeta(h\xi)|)^{2\lambda} d\xi.$$

We note that we will frequently omit the dependence of the operator $P_\zeta(hD)$ on ζ and the dependence of the space X^λ on h and ζ when these can be determined from the context. The dual space of X^λ is $(X^\lambda)^* = X^{-\lambda}$. Since the weight $(h + |P(h\xi)|)^\lambda$ is bounded above and below on compact sets, it is clear that the Schwartz space is dense in X^λ .

Our next goal is to construct an operator that is a right inverse to $P(hD)$ on Ω . While we are interested in constructing solutions in Ω , our right inverse will act on the spaces X^λ which consist of distributions on \mathbb{R}^d . To construct our operator, we choose $\phi \in C_0^\infty(\mathbb{R}^d)$ with $\phi = 1$ in a neighborhood of $\bar{\Omega}$. Let $J_\phi = \phi |P(hD)|^{-1/2}$ and $I_\phi = \phi P(hD)^{-1} \phi$. We then have $I_\phi = J_\phi \frac{|P(hD)|}{P(hD)} J_\phi^*$. What is left to show is that I_ϕ is a bounded operator between suitable spaces and a right inverse.

We begin by recalling an estimate from Haberman and Tataru which they use to give an estimate for u in L_{loc}^2 in terms of $|P(hD)|^{1/2} u$.

Lemma 3.5 ([18, Lemma 2.1]). *Let v and w be nonnegative weights on \mathbb{R}^d . If $\phi \in \mathcal{S}(\mathbb{R}^d)$, then*

$$\|\phi * f\|_{L_w^2} \lesssim_\phi \min \left(\sup_\xi \int J(\xi, \eta) d\eta, \sup_\eta \int J(\xi, \eta) d\xi \right)^{1/2} \|f\|_{L_v^2},$$

where

$$J(\xi, \eta) = |\phi(\xi - \eta)| \frac{w(\xi)}{v(\eta)}.$$

Lemma 3.6. *Let $h > 0$ and $0 \leq s \leq 1$. We have*

$$\int_{B(x,r)} \frac{|\xi|^s}{|P(h\xi)|} d\xi \lesssim \frac{r^{d+1}}{h^{1+s}}, \quad x \in \mathbb{R}^d, \quad r > 0. \quad (3.7)$$

If $\phi \in \mathcal{S}(\mathbb{R}^d)$, then

$$\int_{B(x,r)} \frac{|\phi(x-\xi)||\xi|^s}{|P(h\xi)|} d\xi \lesssim \frac{1}{h^{1+s}}. \quad (3.8)$$

Proof. We begin with the proof of (3.7). By applying the change of variables, $\xi \rightarrow \xi/h$ we may reduce to the case where $h = 1$. To show (3.7) with $h = 1$, we will consider three cases: a) $|x| < 8$, $r \leq 1$, b) $|x| \geq 8$, $r \leq 1$, c) $r > 1$. We let $\Sigma = \{\xi : P(\xi) = 0\}$ and since $P(\xi) = |\xi + \text{Im } \zeta \cdot \xi|^2 - 1 - 2i \text{Re } \zeta \cdot \xi$, we have that Σ is a sphere of radius 1 and codimension two centered at $-\text{Im } \zeta$. Furthermore, one can check that $\nabla P(\xi) \neq 0$ on Σ and thus it follows that $P(\xi) \approx \text{dist}(\xi, \Sigma)$ on $B(0, 9)$. To establish (3.7) in case a) we need to estimate integrals of $|\xi|^s/|P(\xi)| \sim 1/\text{dist}(\xi, \Sigma)$. We can obtain the necessary estimate by using a change of variables to straighten out a portion of the sphere to reduce to integrating the distance to a plane of codimension two in \mathbb{R}^d .

To establish (3.7) in case b), we use that $|\xi|^s/|P(\xi)| \approx |\xi|^{s-2} \lesssim 1$ for $|\xi| \geq 7$. From this, we have

$$\int_{B(x,r)} \frac{|\xi|^s}{|P(\xi)|} d\xi \lesssim r^d \leq r^{d-1}$$

where the last inequality uses that $r \leq 1$. Finally, in case c) we use that $|P(\xi)| \approx |\xi|^2$ for $|\xi|$ large to obtain

$$\int_{B(x,r)} \frac{|\xi|^s}{|P(\xi)|} d\xi \leq \int_{\{|\xi| \leq 8\}} \frac{|\xi|^s}{|P(\xi)|} d\xi + \int_{\{|\xi| > 8\} \cap B(x,r)} |\xi|^{s-2} d\xi.$$

We use case a) to bound the first term and obtain that the previous display is at most $1 + r^{d-2+s} \lesssim r^{d-1}$. The last inequality follows since $r \geq 1$ and $s \leq 1$. We note that if $d = 2$ and $s = 0$, the intermediate estimate involves a logarithm, but the conclusion still holds.

We turn to the proof of (3.8) and give the proof for $h > 0$. We write $\mathbb{R}^d = \cup_{k \geq 0} A_k$ where $A_0 = B(x, 1)$ and for $k \geq 1$ $A_k = B(x, 2^k) \setminus B(x, 2^{k-1})$. If $\phi \in \mathcal{S}(\mathbb{R}^d)$, then it is rapidly decreasing and, in particular, we have $|\phi(x-\xi)| \lesssim 2^{-kd}$ if $\xi \in A_k$. We break the integral in (3.8) into integrals over the sets A_k , and use (3.7) to obtain

$$\int_{\mathbb{R}^d} \frac{|\phi(x-\xi)||\xi|^s}{|P(\xi)|} d\xi \lesssim \frac{1}{h^{1+s}} \sum_{k=1}^{\infty} 2^{k(d-1)} \cdot 2^{-kd} \lesssim \frac{1}{h^{1+s}}.$$

□

Lemma 3.9. *Let $\phi \in \mathcal{S}(\mathbb{R}^d)$. For any $\eta \in \mathbb{R}^d$,*

$$\int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} (|P(h\eta)| - |P(h\xi)|) d\xi \lesssim 1$$

Proof. Using the triangle inequality and the definition of $P(h\xi)$, we have

$$||P(h\eta)| - |P(h\xi)|| \lesssim h|\xi - \eta|(h|\xi + \eta| + 1). \quad (3.10)$$

Using (3.10), we obtain the following inequality.

$$\begin{aligned} \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} (||P(h\eta)| - |P(h\xi)||) d\xi &\lesssim \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} h|\xi - \eta| d\xi \\ &+ \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} h^2|\xi - \eta|^2 d\xi \\ &+ \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} h^2|\xi - \eta||\xi| d\xi. \end{aligned}$$

Using Lemma 3.6 and recalling that $0 < h \leq 1$ completes the proof. \square

We now have enough tools to show that I_ϕ is a bounded right inverse. We begin with an estimate for J_ϕ which will be the main step in establishing estimates for I_ϕ .

Theorem 3.11. *The operator $J_\phi : X^\lambda \rightarrow X^{\lambda+1/2}$ is a bounded operator for all $\lambda \in \mathbb{R}$.*

Proof. We have that

$$(J_\phi u)^\wedge = \hat{\phi} * (\hat{u}/|P(h\xi)|^{1/2}).$$

Since $\hat{\phi}$ is again a Schwartz function, we replace $\hat{\phi}$ by ϕ , and we will show that for a Schwartz function, ϕ , we have

$$\sup_\eta \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{(h + |P(h\xi)|)^{2\lambda+1}}{(h + |P(h\eta)|)^{2\lambda}} d\xi \lesssim 1, \quad 0 \leq h \leq 1. \quad (3.12)$$

Then an application of Lemma 3.5 will give the result of the Theorem.

To establish (3.12), we begin by adding and subtracting $(h + |P(h\eta)|)^{2\lambda+1}$ and then adding and subtracting $|P(h\xi)|$, we can show the following inequality. For $\eta \in \mathbb{R}^d$,

$$\begin{aligned} &\int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{(h + |P(h\xi)|)^{2\lambda+1}}{(h + |P(h\eta)|)^{2\lambda}} d\xi \\ &\leq \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} h d\xi + \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} (||P(h\eta)| - |P(h\xi)||) d\xi + \int_{\mathbb{R}^d} |\phi(\xi - \eta)| d\xi \\ &\quad + \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{|(h + |P(h\xi)|)^{2\lambda+1} - (h + |P(h\eta)|)^{2\lambda+1}|}{(h + |P(h\eta)|)^{2\lambda}} d\xi \\ &= I + II + III + IV. \end{aligned}$$

Using (3.8) of Lemma 3.6 we see that I is bounded by a constant which is independent of η and h , Lemma 3.9 gives the same estimate for II , and the estimate for III is easy. This leaves us to bound IV .

Now, we turn to an application of the mean value theorem. We set $f(t) = t^{2\lambda+1}$ for $t \geq 0$. For $\eta, \xi \in \mathbb{R}^d$, let $a = \min(h + |P(h\xi)|, h + |P(h\eta)|)$ and $b = ||P(h\xi)| - |P(h\eta)||$. Then, we get that $a + b = \max(h + |P(h\xi)|, h + |P(h\eta)|)$. The mean value theorem states that there exists θ between a and $a + b$ such that

$$f(a + b) - f(a) = (2\lambda + 1)\theta^{2\lambda}b.$$

We will go through the proof when $\lambda \geq 0$ as the proof for $\lambda < 0$ is similar. Let $A_0 = B(\eta, 1)$ and $A_k = B(\eta, 2^k) \setminus B(\eta, 2^{k-1})$ for $k \geq 1$. Using (3.10) and then the mean value theorem, we obtain the upper bound

$$\begin{aligned} IV &= \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{|(h + |P(h\xi)|)^{2\lambda+1} - (h + |P(h\eta)|)^{2\lambda+1}|}{(h + |P(h\eta)|)^{2\lambda}} d\xi \\ &\leq \sum_{k=0}^{\infty} \int_{A_k} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{\theta^{2\lambda} h |\xi - \eta| (h|\xi + \eta| + 1)}{(h + |P(h\eta)|)^{2\lambda}} d\xi. \end{aligned}$$

We use $|\theta| \leq h + |P(h\eta)| + |P(h\xi)|$ and that $\lambda \geq 0$ to obtain an upper bound for $\theta^{2\lambda}$. We now consider four cases. Since ϕ is a Schwartz function for $N = 2 + d + 4\lambda$, we have $|\phi(\xi)| \lesssim \langle \xi \rangle^{-N}$.

Case 1: $|h\eta| < 8$ and $2^k \leq 32/h$. Then, in A_k , we have $|\xi + \eta| \leq 48/h$, $|P(h\xi)| \lesssim |P(h\eta)| + h2^k$, and $|\theta|^{2\lambda} \lesssim (h2^k + |P(h\eta)|)^{2\lambda}$. This gives us

$$\begin{aligned} \int_{A_k} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{\theta^{2\lambda} h |\xi - \eta| (h|\xi + \eta| + 1)}{(h + |P(h\eta)|)^{2\lambda}} d\xi &\lesssim \int_{A_k} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{(h2^k + |P(h\eta)|)^{2\lambda} h 2^k}{(h + |P(h\eta)|)^{2\lambda}} d\xi \\ &\lesssim 2^k 2^{-Nk} \int_{A_k} \frac{h}{|P(h\xi)|} \left(\frac{h2^k + |P(h\eta)|}{h + |P(h\eta)|} \right)^{2\lambda} d\xi \\ &\lesssim 2^k 2^{-Nk} \int_{A_k} \frac{h}{|P(h\xi)|} (2^k)^{2\lambda} d\xi \\ &\lesssim 2^{k(-N+2\lambda+d)}. \end{aligned}$$

Case 2: $|h\eta| < 8$ and $2^k \geq 32/h$. Then, in A_k , we have $2^{k-1} \geq 16/h$, $16/h \leq |\xi - \eta|$, $|\xi| \geq 8/h$, and $h|\eta + \xi| \leq h2^k + 8$. Using similar techniques to Case 1, we obtain

$$\int_{A_k} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{\theta^{2\lambda} h |\xi - \eta| (h|\xi + \eta| + 1)}{(h + |P(h\eta)|)^{2\lambda}} d\xi \lesssim 2^{k(-N+d+1+4\lambda)}.$$

Case 3: $|h\eta| \geq 8$ and $2^k \leq |\eta|/8$. Then, $A_k \subset \{\xi : 7|\eta|/8 \leq |\xi| \leq 9|\eta|/8\}$. Therefore, in A_k , we have $|\eta| \approx |\xi|$. Therefore we have $1 \leq h|\xi|/8$, $1 + h|\xi + \eta| \lesssim h|\xi|$, and $|P(h\eta)| \approx |h\eta|^2 \approx |h\xi|^2 \approx |P(h\xi)|$. We then obtain

$$\int_{A_k} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{\theta^{2\lambda} h |\xi - \eta| (h|\xi + \eta| + 1)}{(h + |P(h\eta)|)^{2\lambda}} d\xi \lesssim 2^{k(-N+1+d)}.$$

Case 4: $|h\eta| \geq 8$ and $2^k \geq |\eta|/8$. Then, in A_k , we have $|\xi| \lesssim 2^k$, $|P(h\xi)| \lesssim h2^k + h^2 2^{2k}$, $|P(h\eta)| \lesssim h2^k + h^2 2^{2k}$, and $|\eta + \xi| \lesssim 2^k$. We then obtain

$$\int_{A_k} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{\theta^{2\lambda} h |\xi - \eta| (h|\xi + \eta| + 1)}{(h + |P(h\eta)|)^{2\lambda}} d\xi \lesssim 2^{k(-N+1+4\lambda+d)}.$$

Our choice of N gives

$$\sum_{k=0}^{\infty} \int_{A_k} \frac{|\phi(\xi - \eta)|}{|P(h\xi)|} \frac{\theta^{2\lambda} h |\xi - \eta| (h|\xi + \eta| + 1)}{(h + |P(h\eta)|)^{2\lambda}} d\xi \lesssim \sum_{k=0}^{\infty} 2^{-k}.$$

This gives us that

$$\sup_{\eta} \int_{\mathbb{R}^d} \frac{|\phi(\xi - \eta)| (h + |P(h\xi)|)^{2\lambda+1}}{|P(h\xi)| (h + |P(h\eta)|)^{2\lambda}} d\xi < \infty$$

which completes our proof of (3.12) when $\lambda \geq 0$. We leave it to the reader to adapt this argument to the case $\lambda \leq 0$. \square

Corollary 3.13. *The map $I_\phi : X^\lambda \rightarrow X^{\lambda+1}$ is a bounded operator for all $\lambda \in \mathbb{R}$.*

Proof. Recall that $I_\phi = J_\phi \frac{|P(hD)|}{P(hD)} J_\phi^*$. From Theorem 3.11 we have and $J_\phi : X^{\lambda+1/2} \rightarrow X^{\lambda+1}$. By passing to the adjoint, the boundedness of $J_\phi : X^{-\lambda-1/2} \rightarrow X^{-\lambda}$ implies that $J_\phi^* : X^\lambda \rightarrow X^{\lambda+1/2}$. Since $\frac{|P(hD)|}{P(hD)}$ has a symbol of modulus one, it is an isometry on X^λ for all λ . It follows that $I_\phi : X^\lambda \rightarrow X^{\lambda+1}$. \square

The next Lemma will imply that I_ϕ is a right inverse to $P(hD)$ on Ω .

Lemma 3.14. *We have $P(hD)I_\phi u = u$ in Ω .*

Proof. First, note that $D\phi = 0$ in Ω and $\phi^2 = \phi = 1$ in Ω . Thus, we see that

$$[hD, \phi]u = hD(\phi u) - \phi hDu = uhD\phi + \phi hDu - \phi hDu = uhD\phi = 0, \quad \text{in } \Omega$$

Since $[hD, \phi] = 0$ in Ω , we have

$$P(hD)I_\phi u = P(hD)\phi P(hD)^{-1}\phi u = \phi P(hD)P(hD)^{-1}\phi u = \phi^2 u = u, \quad \text{in } \Omega$$

as we wanted. \square

We now construct solutions to the equation $L_\zeta \psi = f$ where the operator L_ζ is introduced in (3.4). We consider the integral equation

$$\psi + h^{2m} I_\phi^m (Q \cdot (\frac{\zeta}{ih} + D)\phi + q\psi) = I_\phi^m f \quad (3.15)$$

and will show that the map

$$\psi \rightarrow h^{2m} I_\phi^m (Q \cdot (\frac{\zeta}{ih} + D)\psi) + h^{2m} I_\phi^m (q\psi) \quad (3.16)$$

is a contraction on $X^{m/2}$ for suitable conditions on q , Q , and h . Recalling Lemma 3.14, we can show that a solution of our integral equation is a distribution solution of the equation $L_\zeta \psi = f$. Since (3.20) below implies that ψ is in $W^{m,2}(\mathbb{R}^d)$, also obtain that ψ is a weak solution. To prove we have a contraction map, we need estimates for the product $q\psi$ when q is a distribution of negative order and ψ lies in X^λ for some $\lambda > 0$.

We begin with a simple, but useful estimate that will allow us to relate smoothness in the spaces X^λ with smoothness in L^2 -Sobolev spaces. Suppose $0 < h \leq 1$ and $0 \leq s \leq 2\lambda$, then we have

$$\sup_{\xi \in \mathbb{R}^d} \frac{\langle \xi \rangle^s}{(h + |P(h\xi)|)^\lambda} \lesssim h^{-\lambda-s}. \quad (3.17)$$

To establish (3.17), we consider the case $|h\xi| \geq 8$ and $|h\xi| < 8$ separately. If $|h\xi| \geq 8$, we have $|P(h\xi)| \approx h^2|\xi|^2$ and thus $\langle \xi \rangle^s / (h + |P(h\xi)|)^\lambda \lesssim \langle \xi \rangle^{s-2\lambda} h^{-2\lambda} \lesssim h^{-s}$. Since $0 < h \leq 1$, this is better than the estimate (3.17). In the case that $|h\xi| < 8$, we that $\langle \xi \rangle^s \lesssim h^{-s}$ and that $(h + |p(h\xi)|)^{-\lambda} \lesssim h^{-\lambda}$. Multiplying these two inequalities leads to (3.17). Our next step uses the estimate (3.17) to establish estimates relating L^2 -Sobolev spaces and X^λ spaces.

Proposition 3.18. *If $|\alpha| \leq 2(\lambda_2 - \lambda_1)$, then*

$$\|D^\alpha u\|_{X^{\lambda_1}} \lesssim h^{-|\alpha|+\lambda_1-\lambda_2} \|u\|_{X^{\lambda_2}}. \quad (3.19)$$

and if $0 \leq s \leq 2\lambda$, then

$$\|u\|_{W^{s,2}(\mathbb{R}^d)} \lesssim h^{-s-\lambda} \|u\|_{X^\lambda} \quad (3.20)$$

$$\|u\|_{X^{-\lambda}} \lesssim h^{-s-\lambda} \|u\|_{W^{-s,2}(\mathbb{R}^d)}. \quad (3.21)$$

Proof. We begin with the proof of (3.19). We have

$$\|D^\alpha u\|_{X^{\lambda_1}}^2 \leq \sup_{\xi \in \mathbb{R}^d} |\xi|^{2|\alpha|} (h + |P(h\xi)|)^{2(\lambda_1-\lambda_2)} \|u\|_{X^{\lambda_2}}^2 \lesssim h^{2(-|\alpha|+\lambda_1-\lambda_2)} \|u\|_{X^{\lambda_2}}^2$$

where the second inequality uses (3.17).

The proof of (3.20) is similar. Since the embedding of $W^{-s,2}(\mathbb{R}^d)$ into $X^{-\lambda}$ is the adjoint of the embedding $X^\lambda \subset W^{s,2}$, the estimate (3.21) follows from (3.20). \square

Proposition 3.22. *Suppose $f \in L^\infty(\mathbb{R}^d)$, ζ , ζ_1 and ζ_2 are in \mathcal{V} (see (3.1)) and $0 < h \leq 1$. Then when $0 \leq |\alpha| + |\beta| \leq 2\lambda$ we have the trilinear estimate*

$$|\langle D^\alpha f D^\beta u, v \rangle| \lesssim h^{-2\lambda-|\alpha|-|\beta|} \|f\|_\infty \|u\|_{X_{h\zeta_1}^\lambda} \|v\|_{X_{h\zeta_2}^\lambda}. \quad (3.23)$$

Let $Tu = (D^\alpha f)(D + \frac{\zeta}{ih})^\beta u$ with $f \in L^\infty$ and assume $|\alpha| + |\beta| \leq 2\lambda$, then

$$\|Tu\|_{X_{h\zeta_2}^{-\lambda}} \lesssim \|f\|_\infty h^{-2\lambda-|\alpha|-|\beta|} \|u\|_{X_{h\zeta_1}^\lambda}. \quad (3.24)$$

If in addition, we have $f \in C^\theta(\mathbb{R}^d)$ and $|\alpha| \geq 1$, then we have

$$\|Tu\|_{X_{h\zeta_2}^{-\lambda}} \lesssim \|f\|_{C^\theta(\mathbb{R}^d)} h^{-2\lambda-|\alpha|-|\beta|+\theta} \|u\|_{X_{h\zeta_1}^\lambda}. \quad (3.25)$$

Proof. We first consider the case where u and v are in the Schwartz class. Using the definition of $D^\alpha f D^\beta u$ and the product rule we obtain

$$\langle D^\alpha f D^\beta u, v \rangle = (-1)^{|\alpha|} \int_{\mathbb{R}^d} f \sum_{\alpha_1+\alpha_2=\alpha} \frac{\alpha!}{\alpha_1!\alpha_2!} D^{\beta+\alpha_1} u D^{\alpha_2} v dx.$$

Using the Cauchy-Schwartz inequality and the estimate (3.19) with $\lambda_1 = 0$ and $\lambda_2 = \lambda$ quickly gives (3.23). The result for general u and v follows because the Schwartz class is dense in the space X^λ .

To establish (3.24), we write

$$\langle Tu, v \rangle = (-1)^{|\alpha|} \int_{\mathbb{R}^d} \sum_{\beta_1 + \beta_2 = \beta} \frac{\beta!}{\beta_1! \beta_2!} f D^\alpha (D^{\beta_1} u (\zeta / (ih))^{\beta_2} v) dx. \quad (3.26)$$

The estimate (3.24) follows from (3.23).

In the case when f is Hölder continuous, we write $f = f_h + f^h$ where f_h is a mollification of f and the decomposition satisfies $\|D^\alpha f_h\|_\infty \lesssim h^{\theta - |\alpha|}$ for $|\alpha| \geq 1$ and $\|f^h\|_\infty \lesssim h^\theta$. We use this decomposition of f in (3.26) and consider the two terms. For the first term, we use that $|\alpha| \geq 1$, integrate by parts and use (3.19) to obtain

$$\left| \int_{\mathbb{R}^d} \sum_{\beta_1 + \beta_2 = \beta} \frac{\beta!}{\beta_1! \beta_2!} f_h D^\alpha (D^{\beta_1} u (\zeta / (ih))^{\beta_2} v) dx \right| \lesssim \|f\|_{C^\theta(\mathbb{R}^d)} h^{-2\lambda - |\alpha| - |\beta| + \theta} \|u\|_{X^\lambda} \|v\|_{X^\lambda}.$$

Since $\|f^h\|_\infty \lesssim \|f\|_{C^\theta(\mathbb{R}^d)} h^\theta$, we may use (3.24) and obtain the same estimate for the term involving f^h ,

$$\left| \int_{\mathbb{R}^d} \sum_{\beta_1 + \beta_2 = \beta} \frac{\beta!}{\beta_1! \beta_2!} f^h D^\alpha (D^{\beta_1} u (\zeta / (ih))^{\beta_2} v) dx \right| \lesssim \|f\|_{C^\theta(\mathbb{R}^d)} h^{-2\lambda - |\alpha| - |\beta| + \theta} \|u\|_{X^\lambda} \|v\|_{X^\lambda}.$$

□

Our next proposition shows that multiplying by a smooth function preserves the class X^λ .

Proposition 3.27. *Let $v \in C^\infty(\mathbb{R}^d)$ and assume that v and all derivatives of v are bounded on \mathbb{R}^d . Then for $\lambda \in \mathbb{R}$, we have*

$$\|uv\|_{X^\lambda} \lesssim \|u\|_{X^\lambda}. \quad (3.28)$$

The implied constant in (3.28) depends on λ and $\|D^\alpha v\|_\infty$ for $|\alpha| \leq c(|\lambda|)$.

Proof. We begin by giving a proof by induction for (3.28) when $\lambda \geq 0$ is an integer. Since $X^0 = L^2$, the base case $\lambda = 0$ is elementary. Next, we assume (3.28) holds for $\lambda \geq 0$ and establish the inequality for $\lambda + 1$. Using Plancherel's theorem we have $\|uv\|_{X^{\lambda+1}} \leq h\|uv\|_{X^\lambda} + \|P(hD)(uv)\|_{X^\lambda}$. Then using the product rule, we have

$$\begin{aligned} \|uv\|_{X^{\lambda+1}} &\lesssim h\|uv\|_{X^\lambda} + \|vP(hD)u\|_{X^\lambda} + h\|uDv\|_{X^\lambda} + h^2\|u\Delta v\|_{X^\lambda} + h^2\|Du \cdot Dv\|_{X^\lambda} \\ &\lesssim (h + h^2)\|u\|_{X^\lambda} + \|P(hD)u\|_{X^\lambda} + h^2\|Du\|_{X^\lambda} \\ &\lesssim \|u\|_{X^{\lambda+1}}. \end{aligned}$$

The second inequality uses our induction hypothesis, the third inequality follows from (3.19) and uses that $0 < h \leq 1$.

Next, we may use complex interpolation to establish that the map $u \rightarrow uv$ is bounded on X^λ for all $\lambda \geq 0$. Finally, the adjoint of this map will also be bounded on $X^{-\lambda}$ which gives (3.28) for $\lambda < 0$. □

We now give conditions which imply that (3.16) is a contraction map. Proposition A.3 implies that the conditions on q and Q in our next theorem follow from the hypotheses in Theorem 1.3.

Theorem 3.29. *Suppose that*

$$Q_j = \sum_{|\beta| \leq m-1} D^\beta Q_{j\beta}, \quad q = \sum_{|\beta| \leq m} D^\beta q_\beta \quad (3.30)$$

with $Q_{j\beta}, q_\beta$ in $C^\theta(\mathbb{R}^d)$ for some $\theta \in (0, 1]$ and each function is supported in $\bar{\Omega}$. Then there is a value h_0 depending on θ and the Hölder norms of the functions $Q_{j\beta}, q_\beta$ so that if $0 < h \leq h_0$, we may find a function $\psi \in X^{m/2}$ which satisfies $L_\zeta \psi = f$ in Ω . The solution will satisfy the estimate

$$\|\psi\|_{X^{m/2}} \lesssim \|f\|_{X^{-m/2}}.$$

Proof. If Q and q are as in (3.30), then according to Proposition 3.22 the map (3.16) will be a contraction on $X^{m/2}$ at least for h sufficiently small. Thus, we may solve the integral equation (3.15) for ψ and the solution will satisfy the estimate $\|\psi\|_{X^{m/2}} \lesssim \|I_\phi^m f\|_{X^{m/2}}$. Then m applications of Corollary 3.13 imply the estimate of the Theorem. To show that ψ is a weak solution of the differential equation $L_\zeta \psi = f$ we need to show that it is a solution in the sense of distributions, which follows from Lemma 3.14 and that ψ is in $W^{m,2}(\Omega)$. We use (3.20) to show that $X^{m/2} \subset W^{m,2}(\mathbb{R}^d)$. \square

4 Average Estimate

The estimate (3.21) of Proposition 3.18 gives an estimate for the norm of a distribution in $X^{-\lambda}$ in terms of the $W^{-s,2}$ norm. However, as observed by Haberman and Tataru [18], there is an improvement available if we look at averages of $X_{\tau\zeta}^\lambda$ norms over certain families of vectors ζ and for τ in an interval $[h, 2h]$. The improvement is that the averages grow less rapidly in h . This improved behavior is used in section 6 to recover q and Q by studying limits as h tends to zero of expressions defined using our solutions. In this section, we describe the averages we are interested in and give an extension of Haberman and Tataru's estimate.

Fix $\xi_0 \in \mathbb{R}^d \setminus \{0\}$ and let $\mu_1, \mu_2 \in \mathbb{R}^d$ such that $\xi_0 \cdot \mu_1 = \xi_0 \cdot \mu_2 = 0$ and $\mu_i \cdot \mu_j = \delta_{ij}$. We set

$$\begin{aligned} \mu_1(\theta) &= \operatorname{Re}(e^{i\theta}(\mu_1 + i\mu_2)) = \mu_1 \cos(\theta) - \mu_2 \sin(\theta) \\ \mu_2(\theta) &= \operatorname{Im}(e^{i\theta}(\mu_1 + i\mu_2)) = \mu_1 \sin(\theta) + \mu_2 \cos(\theta) \end{aligned}$$

and then for τ small we put

$$\begin{aligned} \zeta^1(\tau, \theta) &= \mu_1(\theta) + i\sqrt{1 - \frac{\tau^2 |\xi_0|^2}{4}} \mu_2(\theta) - i\frac{\tau \xi_0}{2}, \\ \zeta^2(\tau, \theta) &= -\mu_1(\theta) - i\sqrt{1 - \frac{\tau^2 |\xi_0|^2}{4}} \mu_2(\theta) - i\frac{\tau \xi_0}{2}. \end{aligned} \quad (4.1)$$

Throughout this section, we will let $P(\tau\xi) = |\tau\xi|^2 - 2i\zeta^k(\tau, \theta) \cdot \tau\xi$ for $k = 1$ or 2 . We will choose $h > 0$ such that $h \leq 1/(4|\xi_0|)$ and we will have that $h \leq \tau \leq 2h$. For $\xi \in \mathbb{R}^d$, we

define ξ^\perp to be the projection of ξ onto the plane spanned by $\{\mu_1, \mu_2\}$. We will now prove an estimate for the average of the norms

$$\|f\|_{X_{\tau\zeta^k(\tau,\theta)}^{-\lambda}}, \quad h \leq \tau \leq 2h, \quad \theta \in [0, 2\pi] \quad (4.2)$$

We begin with a technical lemma which gives an estimate for the Jacobian of a change of variables that will be used in studying the average of the norms in (4.2).

Lemma 4.3. *Let $\xi, \xi_0 \in \mathbb{R}^d \setminus \{0\}$, and $0 < \tau \leq \min(1, 1/(2|\xi_0|))$ and let $P(\tau\xi)$ be as defined after (4.1). We define a change of variables $z(\tau, \theta) = (z_1(\tau, \theta), z_2(\tau, \theta))$ by*

$$z_1(\tau, \theta) = \frac{\operatorname{Re} P(\tau\xi)}{\tau^2} \quad \text{and} \quad z_2(\tau, \theta) = \frac{\operatorname{Im} P(\tau\xi)}{\tau^2}. \quad (4.4)$$

Then the Jacobian J for this change of variables has the lower bound

$$\frac{2|\xi^\perp|^2}{\tau^3} \leq J.$$

Proof. We give the proof for ζ^1 . The case of ζ^2 is similar and we omit the details. A calculation shows that

$$\begin{aligned} J &= \left| \det \begin{pmatrix} \frac{-2\mu_2(\theta) \cdot \xi}{\tau^2} \left(\sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}} + \frac{\tau^2|\xi_0|^2}{4\sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}}} \right) & \frac{2\mu_1(\theta) \cdot \xi}{\tau} \sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}} \\ 2\mu_1(\theta) \cdot \xi/\tau^2 & 2\mu_2(\theta) \cdot \xi/\tau \end{pmatrix} \right| \\ &= \left| -\frac{|\xi_0|^2}{\sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}}} \frac{|\mu_2(\theta) \cdot \xi|^2}{\tau} - 4\frac{\sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}}}{\tau^3} |\mu_2(\theta) \cdot \xi|^2 - 4\frac{\sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}}}{\tau^3} |\mu_1(\theta) \cdot \xi|^2 \right| \\ &= \frac{4|\xi^\perp|^2}{\tau^3} \left(|\xi_0|^2 \tau^2 \frac{1}{4\sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}}} \frac{|\mu_2(\theta) \cdot \xi|^2}{|\xi^\perp|^2} + \sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}} \right). \end{aligned}$$

The condition that $\tau \leq 1/(2|\xi_0|)$, gives at least the following bound

$$\frac{1}{2} \leq \sqrt{1 - \frac{\tau^2|\xi_0|^2}{4}} \leq 1.$$

Therefore we can conclude

$$2\frac{|\xi^\perp|^2}{\tau^3} \leq J. \quad \square$$

Proposition 4.5. *Let $P(\tau\xi)$ depend τ and θ as defined in the beginning of this section. If $0 < \epsilon \leq 1$ and $h\langle \xi_0 \rangle^2 \lesssim 1$, we have*

$$\frac{1}{h} \int_0^{2\pi} \int_h^{2h} \frac{1}{(\tau + |P(\tau\xi)|)^{2-2\epsilon}} d\tau d\theta \lesssim \frac{1}{\tau^{4-4\epsilon} \langle \xi \rangle^{4-4\epsilon}}.$$

Proof. We consider three cases depending on ξ . Case 1: $|\xi| \leq \max(16|\xi_0|, 1)$ Case 2a: $|\xi| > \max(16|\xi_0|, 1)$ and $|\xi|^2 \leq 16|\xi^\perp|/h$, Case 2b: $|\xi| > \max(16|\xi_0|, 1)$ and $|\xi|^2 > 16|\xi^\perp|/h$.

In Case 1, we use that $(\tau + |P(\tau\xi)|)^{2\epsilon-2} \leq h^{2\epsilon-2}$ and $\langle \xi \rangle \lesssim \langle \xi_0 \rangle$ to obtain that

$$\frac{1}{h} \int_0^{2\pi} \int_h^{2h} \frac{1}{(\tau + |P(\tau\xi)|)^{2-2\epsilon}} d\tau d\theta \lesssim \frac{1}{h^{2-2\epsilon}} \lesssim \frac{\langle \xi_0 \rangle^{4-4\epsilon} h^{2-2\epsilon}}{h^{4-4\epsilon} \langle \xi \rangle^{4-4\epsilon}}$$

which gives the desired result if we require that $h^{2-2\epsilon} \langle \xi_0 \rangle^{4-4\epsilon} \lesssim 1$.

In Case 2a) we will use the change of variable from Lemma 4.3. The conditions $|\xi|^2 \leq 16|\xi^\perp|/h$ and $|\xi| \geq 16|\xi_0|$ imply that $|z| = |z_1 + iz_2| \leq 21|\xi^\perp|/h$. Using this observation and the estimate for the Jacobian from Lemma 4.3 gives

$$\begin{aligned} \frac{1}{h} \int_0^{2\pi} \int_h^{2h} \frac{1}{(\tau + |P(\tau\xi)|)^{2-2\epsilon}} d\tau d\theta &\lesssim \frac{h^2}{|\xi^\perp|^2} \int_{|\tau| \leq 21|\xi^\perp|/h} \frac{1}{(h^2|z|)^{2-2\epsilon}} dz \\ &\lesssim \frac{1}{h^{4-4\epsilon}} \left(\frac{h}{|\xi^\perp|} \right)^{2-2\epsilon} \lesssim \frac{1}{h^{4-4\epsilon} \langle \xi \rangle^{4-4\epsilon}}. \end{aligned}$$

The last step uses that $|\xi| \geq 1$ and $|\xi|^2 \leq 16|\xi^\perp|/h$.

Finally, in Case 2b) our conditions on ξ imply that $|P(\tau\xi)| \geq h^2 \langle \xi \rangle^2$ which quickly gives the conclusion of the Proposition. \square

We will use Proposition 4.5 to establish an estimate relating the average of the $X^{-\lambda}$ -norm of a distribution to the norm in an L^2 -Sobolev space.

Theorem 4.6. *For any $0 < \epsilon < 1$ and $2 - 2\epsilon \leq s \leq 2\lambda$ we have*

$$\frac{1}{h} \int_0^{2\pi} \int_h^{2h} \|f\|_{X_{\zeta(\tau, \theta)}^{-\lambda}}^2 d\tau d\theta \lesssim h^{-2(s+\lambda-1+\epsilon)} \|f\|_{W^{-s, 2}(\mathbb{R}^d)}^2.$$

Proof. We have

$$\begin{aligned} \frac{1}{h} \int_0^{2\pi} \int_h^{2h} \|f\|_{X_{\zeta(\tau, \theta)}^{-\lambda}}^2 d\tau d\theta &= \int_{\mathbb{R}^d} |\hat{f}(\xi)|^2 \frac{1}{h} \int_0^{2\pi} \int_h^{2h} \int_{\mathbb{R}^d} \frac{1}{(\tau + |P_\zeta(\tau\xi)|)^{2\lambda}} |\hat{f}(\xi)|^2 d\tau d\theta d\xi \\ &\lesssim \sup_\xi \frac{\langle \xi \rangle^{2s-4+4\epsilon}}{(h + |P_\zeta(h\xi)|)^{2\lambda-2+2\epsilon}} \\ &\quad \times \int_{\mathbb{R}^d} |\hat{f}(\xi)|^2 \langle \xi \rangle^{-2s+4-4\epsilon} \frac{1}{h} \int_0^{2\pi} \int_h^{2h} \frac{1}{(s + |P_\zeta(\tau\xi)|)^{2-2\epsilon}} d\tau d\theta d\xi \\ &\lesssim h^{-2(s+\lambda-1+\epsilon)} \|f\|_{W^{-s, 2}(\mathbb{R}^d)}^2. \end{aligned}$$

The last inequality depends on Proposition 4.5 and (3.17). The conditions on s are needed to apply (3.17). \square

5 Cauchy data, bilinear forms and Dirichlet to Neumann maps

In this section, we establish the existence of the Cauchy data for a weak solution and discuss a Dirichlet to Neumann map. Our solution to the inverse boundary value problem in section 6 establishes that the map from the coefficients Q and q to a bilinear form is injective. In this section, we establish that the bilinear form can be determined from Cauchy data or a Dirichlet to Neumann map. Thus, our main theorem which gives uniqueness of coefficients in terms of an hypothesis on bilinear forms will imply the solution of traditional formulations of the inverse boundary value problem with hypotheses on Cauchy data or Dirichlet to Neumann maps.

We define boundary operators $\delta_{2j}u = (-\Delta)^j u|_{\partial\Omega}$ and $\delta_{2j+1} = \frac{\partial}{\partial\nu}(-\Delta)^j u$, both for $j = 0, \dots, m-1$. By applying Green's identity to B_0 for $u, v \in C^\infty(\bar{\Omega})$ we obtain

$$B_0(u, v) - \int_{\Omega} [(-\Delta)^m u]v = \sum_{j=0}^{m-1} \int_{\partial\Omega} (-1)^j \delta_j v \delta_{2m-1-j} u \, d\sigma \quad (5.1)$$

If u is a solution of an equation with principal part $(-\Delta)^m$, the vector of distributions on the boundary $(\delta_0 u, \dots, \delta_{2m-1} u)$ are the *Cauchy data* of the solution u . (Or at least one possible choice for the Cauchy data.) Here and throughout this section, we are using the form B_0 given in (2.5). The next Proposition shows that we can define Cauchy data for a weak solution of $(-\Delta)^m u = F$.

Proposition 5.2. *We assume $\partial\Omega$ is smooth. Suppose $u \in W^{m,2}(\Omega)$ and is a weak solution of $(-\Delta)^m u = F$ in Ω with $F \in \tilde{W}^{-m,2}(\Omega)$. Then we may define the Cauchy data for u , $\delta_j u \in W^{m-\frac{1}{2}-j,2}(\partial\Omega)$ for $j = 0, \dots, 2m-1$.*

Proof. For $j = 0, \dots, m-1$ the trace theorem gives us $\delta_j u \in W^{m-\frac{1}{2}-j,2}(\partial\Omega)$. Next, for $\phi_j \in W^{m-\frac{1}{2}-j,2}(\partial\Omega)$ for $j = 0, \dots, m-1$ we can solve the Dirichlet problem

$$\begin{cases} (-\Delta)^m v = 0 & \text{in } \Omega \\ \delta_j v = \phi_j & \text{on } \partial\Omega. \end{cases}$$

See, for example, the monograph of Gazzola *et. al.* [16, Theorem 2.14]. We use the identity below to define $\delta_j u$ for $j = m, \dots, 2m-1$:

$$B_0(u, v) - \langle F, v \rangle = \sum_{j=0}^{m-1} (-1)^j \langle \delta_{2m-1-j} u, \delta_j v \rangle_{\partial\Omega}.$$

Here we are using $\langle \cdot, \cdot \rangle_{\partial\Omega}$ to denote the pairing of duality for Sobolev spaces on the boundary. Note that the lefthand side of this identity is defined due to our choice of the form B_0 and our assumption that F lies in the dual space to $W^{m,2}(\Omega)$, $\tilde{W}^{-m,2}(\Omega)$. \square

Remark. 1. We note that the identity (5.1) allows us to show that for a smooth solution, the weak definition of the Cauchy data in Proposition 5.2 agrees with the classical definition of the operators $\delta_j u$.

2. Our discussion of the Cauchy data requires us to assume that the boundary is smooth. For example, even to define Sobolev spaces of order k on the boundary, we would normally need the boundary to be C^k . In addition, the existence result for the Dirichlet problem for the polyharmonic operator that we quote also assumes the boundary is smooth.

3. It is possible that the work of Verchota on boundary value problems for polyharmonic operators on Lipschitz domains [33] can be used to provide a formulation of the Cauchy data for solutions on Lipschitz domains. However, this would take us far afield from the themes we are studying here and we have not pursued this direction.

For a weak solution of $Lu = 0$ with L as defined in (1.1) and coefficients satisfying (2.7), we have that qu and $Q \cdot Du$ lie in $\tilde{W}^{-m,2}(\Omega)$ (see (2.3)), thus Proposition 5.2 implies the existence of the Cauchy data for u .

We now consider two operators $L_j = (-\Delta)^m u + Q^j \cdot Du + q^j u$ for $j = 1, 2$ with Q^j and q^j satisfying the conditions (2.7) and let $B_j(u, v) = B_0(u, v) + \langle Q^j \cdot Du, v \rangle + \langle q^j u, v \rangle$ be the form used to define weak solutions of the operator L_j . We say $B_1 = B_2$ if for each $u_1 \in W^{m,2}(\Omega)$ a solution of $L_1 u_1 = 0$, there is a solution u_2 of $L_2 u_2 = 0$ with $u_1 - u_2 \in W_0^{m,2}(\Omega)$ and $B_1(u_1, v) = B_2(u_2, v)$ for all $v \in W^{m,2}(\Omega)$. We also assume that we have the same result with the roles of L_1 and L_2 reversed.

Now, suppose that $B_1 = B_2$ and u_1 is a solution to $L_1 u_1 = 0$ and v_2 a solution of $L_2^t v_2 = 0$. If there is a solution of $L_2 u_2 = 0$ with $u_1 - u_2 \in W_0^{m,2}(\Omega)$ then

$$B_1(u_1, v_2) = B_2(u_2, v_2)$$

and since v_2 is a solution of $L_2^t v_2 = 0$, we have

$$B_2(u_1 - u_2, v_2) = 0.$$

Thus, we can conclude

$$B_1(u_1, v_2) = B_2(u_1, v_2). \quad (5.3)$$

Next we say that two operators L_1 and L_2 have the same Cauchy data if whenever $u_1 \in W^{m,2}(\Omega)$ is a weak solution of $L_1 u_1 = 0$, there exists $u_2 \in W^{m,2}(\Omega)$ so that $L_2 u_2 = 0$ and $\delta_j u_1 = \delta_j u_2$, $j = 0, \dots, 2m - 1$. We observe that the condition $\delta_j u_1 = \delta_j u_2$, $j = 0, \dots, m - 1$ implies $u_1 - u_2 \in W_0^{m,2}(\Omega)$ and then the definition of the Cauchy data using the form implies

$$B_0(u_1, v) + \langle (Q^1 \cdot D + q^1)u_1, v \rangle = B_0(u_2, v) + \langle (Q^2 \cdot D + q^2)u_2, v \rangle$$

and thus we have that if the Cauchy data for the two operators agree, then the forms are equal. This argument may be reversed so that we have that the equality of the forms is equivalent to the equality of the Cauchy data.

Finally, in the case that we have unique solvability of a boundary value problem, we may view the linear space of Cauchy data as the graph of a Dirichlet to Neumann type map. For example, if we have unique solvability for the Navier boundary value problem (1.2) with data $(\phi_0, \phi_2, \dots, \phi_{2m-2}) \in \oplus_{j=0}^{m-1} W^{m-1/2-2j,2}(\partial\Omega)$, then we may define $\Lambda : \oplus_{j=0}^{m-1} W^{m-1/2-2j,2}(\partial\Omega) \rightarrow \oplus_{j=0}^{m-1} W^{m-3/2-2j,2}(\partial\Omega)$ by

$$\Lambda(\phi_0, \dots, \phi_{2m-2}) = \left(\frac{\partial}{\partial \nu} u, \frac{\partial}{\partial \nu} (-\Delta u), \dots, \frac{\partial}{\partial \nu} (-\Delta)^{m-1} u \right)$$

where u is the solution to (1.2) with data $(\phi_0, \dots, \phi_{2m-2})$. It is clear that under the assumption of the unique solvability of the Navier boundary value problem, the equality of the Dirichlet to Neumann maps for two different operators implies the two operators have the same Cauchy data. One may also define Dirichlet to Neumann maps for other boundary value problems such as the Dirichlet problem. We leave it to the interested reader to write down the details connecting these maps to Cauchy data and then the bilinear form.

We will want to work on a simply connected domain in order to recover Q . To do so, let B be a ball such that $\Omega \subset\subset B$. Let $L_j = (-\Delta)^m + Q^j \cdot D + q^j$ in Ω and an operator \tilde{L}_j in B . We define the forms of the operators as before and we have $\tilde{B}_j = B_j + B'$ where B' is the form B_0 , with domain $W^{m,2}(B \setminus \Omega) \times W^{m,2}(B \setminus \Omega)$.

Lemma 5.4. *If $B_1 = B_2$ then $\tilde{B}_1 = \tilde{B}_2$.*

Proof. Let $\tilde{u}_1 \in W^{m,2}(B)$ such that $L_1 \tilde{u}_1 = 0$ and let $u_1 = \tilde{u}_1|_{\Omega}$. Since $B_1 = B_2$ there exists $u_2 \in W^{m,2}(\Omega)$ such that $u_1 - u_2 \in W_0^{m,2}(\Omega)$ and $L_2 u_2 = 0$. Then, for any $v \in W_0^{m,2}(B)$

$$\tilde{B}_1(\tilde{u}_1, v) = B_1(u_1, v) + B'(\tilde{u}_1, v) = B_2(u_2, v) + B'(\tilde{u}_1, v) = \tilde{B}_2(\tilde{u}_2, v)$$

where

$$\tilde{u}_2 = \begin{cases} \tilde{u}_1 & \text{in } B \setminus \bar{\Omega} \\ u_2 & \text{in } \bar{\Omega}. \end{cases}$$

It follows that if $\tilde{L}_1 \tilde{u}_1 = 0$ then $\tilde{L}_2 \tilde{u}_2 = 0$. We have shown that if $\tilde{u}_1 \in W^{m,2}(B)$ is a solution to $\tilde{L}_1 \tilde{u}_1 = 0$ then there exists $\tilde{u}_2 \in W^{m,2}(B)$ with $\tilde{L}_2 \tilde{u}_2 = 0$ and $\tilde{u}_1 - \tilde{u}_2 \in W_0^{m,2}(B)$ since $\tilde{u}_1 - \tilde{u}_2$ is compactly supported in B . Reversing the roles of 1 and 2, it follows that $\tilde{B}_1 = \tilde{B}_2$. \square

6 Main Result

In this section, we give a construction of CGO solutions. We begin with a simple result that may be viewed as an analog of a theorem of Päivärinta *et. al.* [26, Theorem 1.1] in the scale of the X^λ spaces. Our next step is to give an improvement of this result that relies on the averaging estimate from Section 4. We then prove our main result on the uniqueness of the coefficients.

Proposition 6.1. *Let $\zeta \in \mathcal{V}$, suppose $q \in \tilde{W}^{-s,p}(\Omega)$ and $Q \in \tilde{W}^{1-s,p}(\Omega)$ with s and p satisfying $1/p + (s - m)/d < 0$ and $s < m$. Suppose $a \in C^\infty(\mathbb{R}^d)$ and $P(hD)^m a = 0$. For h small, we may find a solution ψ so that $u = e^{x \cdot \zeta/h}(a + \psi)$ is a CGO solution in Ω and with $X^\lambda = X_{h\zeta}^\lambda$ we have $\psi \in X^{m/2}$ and the estimate*

$$\|\psi\|_{X^{m/2}} \lesssim h^{2m} (\|Q \cdot Da\|_{X^{-m/2}} + h^{-1} \|Qa\|_{X^{-m/2}} + \|aq\|_{X^{-m/2}}). \quad (6.2)$$

If in addition we have $p \geq 2$, then $\|\psi\|_{X^{m/2}} \lesssim h^{3m/2-s}$.

Proof. By Sobolev embedding [5, Theorem 6.5.1], we have $\tilde{W}^{-s,p}(\Omega) \subset \tilde{W}^{-s_1,p_1}(\Omega)$ provided $s_1 > s$, $p, p_1 \in (1, \infty)$ and

$$\frac{1}{p} + \frac{s}{d} \leq \frac{1}{p_1} + \frac{s_1}{d}.$$

Thus we may choose $p_1 \in (1, \infty)$ and $r \in (0, 1)$ so that $q \in \tilde{W}^{r-m, p_1}(\Omega)$ and $Q \in \tilde{W}^{1+r-m, p_1}(\Omega)$ and $1/p_1 - r/d < 0$. By Proposition A.3 the coefficients q and Q have a representation as in (3.30), the hypothesis of Theorem 3.29. This follows because Morrey's Lemma implies functions in $W^{r, p_1}(\mathbb{R}^d)$ are Hölder continuous. Thus we may use Theorem 3.29 to solve $L_\zeta \psi = -L_\zeta a$ and the solution will satisfy $\|\psi\|_{X^{m/2}} \lesssim \|L_\zeta a\|_{X^{-m/2}}$. The first estimate (6.2) follows from this. To obtain the estimate in terms of h , we use (3.21) and Proposition 3.27 to estimate the $X^{-m/2}$ norm in terms of the Sobolev norm. \square

The next Lemma is rather technical but important. It provides a sequence of CGO solutions that are instrumental in the proof of the main theorem.

Lemma 6.3. *Suppose $1/p + (s-m)/d < 0$, $p \geq 2$, and $s < \frac{m}{2} + 1$. Assume we have potentials q^k, Q^k for $k = 1, 2$ which satisfy $q^k \in \tilde{W}^{-s, p}(\Omega)$ and $Q^k \in \tilde{W}^{-s+1, p}(\Omega)$. Assume $a^k \in C^\infty(\mathbb{R}^d)$ and $P(hD)^m a^k = 0$. Given $\xi_0 \in \mathbb{R}^d \setminus \{0\}$ and unit vectors μ_1 and μ_2 so that $\{\xi_0, \mu_1, \mu_2\}$ are mutually orthogonal, there exist sequences $\{h_j\}, \{\theta_j\}$ with $\lim_{j \rightarrow \infty} h_j = 0$ and $\lim_{j \rightarrow \infty} \theta_j = \theta_0$ with the following properties. If $\zeta_j^k = \zeta^k(h_j, \theta_j)$ is as in (4.1) and $X_{k,j}^\lambda = X_{h_j \zeta_j^k}^\lambda$, then we may find sequences $\{\psi_j^k\} \subset X_{k,j}^{m/2}$, $k = 1, 2$ so that $e^{x \cdot \zeta_j^k / h_j} (a^k + \psi_j^k)$ is a CGO solution to $L_k u = 0$ and*

$$\|q^\ell\|_{X_{k,j}^{-m/2}} + h_j^{-1} \|Q^\ell\|_{X_{k,j}^{-m/2}} + h_j^{-3/2} \|Q^\ell\|_{X_{k,j}^{(1-m)/2}} \lesssim h_j^{-m/2-s+1-\epsilon}, \quad k, \ell = 1, 2 \quad (6.4)$$

$$\|\psi_j^k\|_{X_{k,j}^{m/2}} \lesssim h_j^{\frac{3m}{2}-s+1-\epsilon}. \quad (6.5)$$

Proof. We may use Proposition 6.1 to find solutions ψ^k for each h and θ which lie in the space $X_k^{m/2}$ where we use $X_k^\lambda = X_{h\zeta^k(h, \theta)}^\lambda$. These solutions ψ^k satisfy the estimate (6.2). Using Proposition 3.27, we can bound the right hand side of (6.2) by $h^{2m} (\|q^k\|_{X_k^{-m/2}} + h^{-1} \|Q^k\|_{X_k^{-m/2}})$. If we sum the estimates of Theorem 4.6 for $k = 1, 2$, may find sequences $\{h_j\}$ and $\{\theta_j\}$ so that for $k, \ell = 1, 2$ we have the estimates in the $X_{k,j}^{-m/2}$ norm in (6.4). The estimate for the remaining term then follows from (3.19). After perhaps passing to a sub-sequence, we have that the sequence θ_j will be convergent and we let θ_0 be the limit of this sequence. The estimate (6.5) follows from (6.4) and (6.2). \square

We are now ready to give the proof of our result establishing that the bilinear form for the operator L uniquely determines the coefficients q and Q .

Theorem 6.6. *Suppose that we have two operators L_k , $k = 1, 2$ as in (1.1) and we have $s < m/2 + 1$ and $p \geq 2$ which satisfy $1/p + (s-m)/d < 0$, $p \geq 2$. Suppose the coefficients of the operators L_k satisfy $Q^k \in \tilde{W}^{-s+1, p}(\Omega)$ and $q^k \in \tilde{W}^{-s, p}(\Omega)$. If the bilinear forms for the operators satisfy $B_1 = B_2$, then $Q^1 = Q^2$ and $q^1 = q^2$.*

Proof. We begin by applying Lemma 5.4 to assume Ω is a ball. Since the Sobolev spaces $\tilde{W}^{-s, 2}(\Omega)$ are increasing as $-s$ decreases, we may assume that $s > m/2$ and choose ϵ so that $s = \frac{m}{2} + 1 - \frac{3}{2}\epsilon$ for $0 < \epsilon < \frac{1}{3}$. Let $\xi_0 \in \mathbb{R}^d$ and find unit vectors $\mu_1, \mu_2 \in \mathbb{R}^d$ such that $\{\xi_0, \mu_1, \mu_2\}$ are mutually orthogonal. Given amplitudes a^k which satisfy $P(hD)^m a^k = 0$. We apply Lemma 6.3 to the operators L_1 and L_2^t to find a sequence of solutions $\{\psi_j^k\}$, a sequence

of spaces $\{X_{k,j}^{m/2}\}$ and numerical sequences $\{h_j\}$ and $\{\theta_j\}$ so that $u_k = e^{\zeta_j^k \cdot x/h_j} (a^k + \psi_j^k)$ are CGO solutions of $L_1 u_1 = 0$ and $L_2^t u_2 = 0$ which satisfy the estimates (6.4) and (6.5). It is clear from the definitions of ζ^k in (3.1) that $\zeta_j^1 + \zeta_j^2 = -ih_j \xi_0$. Since we have that the forms for L_1 and L_2 are equal (see (5.3)), we have

$$\begin{aligned}
0 &= ih_j [B_1(u_1, u_2) - B_2(u_1, u_2)] \\
&= \zeta_j^1 \cdot \langle (Q^1 - Q^2)a^1, a^2 e^{-ix \cdot \xi_0} \rangle + \zeta_j^1 \cdot \langle (Q^1 - Q^2)\psi_j^1, a^2 e^{-ix \cdot \xi_0} \rangle + \zeta_j^1 \cdot \langle (Q^1 - Q^2)a^1, e^{-ix \cdot \xi_0} \psi_j^2 \rangle \\
&\quad + \zeta_j^1 \cdot \langle (Q^1 - Q^2)\psi_j^1, e^{-ix \cdot \xi_0} \psi_j^2 \rangle + ih_j \langle (Q^1 - Q^2) \cdot Da^1, a^2 e^{-ix \cdot \xi_0} \rangle \\
&\quad + ih_j \langle (Q^1 - Q^2) \cdot Da^1, \psi_j^2 e^{-ix \cdot \xi_0} \rangle + ih_j \langle (Q^1 - Q^2) \cdot D\psi_j^1, a^2 e^{-ix \cdot \xi_0} \rangle \\
&\quad + ih_j \langle (Q^1 - Q^2) \cdot D\psi_j^1, e^{-ix \cdot \xi_0} \psi_j^2 \rangle + ih_j \langle (Q^1 - Q^2)(a^1 + \psi_j^1), e^{-ix \cdot \xi_0} (a^2 + \psi_j^2) \rangle \\
&= I + II + III + IV + V + VI + VII + VIII + IX.
\end{aligned}$$

Since the only dependence on h in I is through the formula for ζ_j^1 (see (3.1)), we have

$$\lim_{j \rightarrow \infty} I = \lim_{j \rightarrow \infty} \zeta_j^1 \cdot \langle (Q^1 - Q^2)a^1, a^2 e^{-ix \cdot \xi_0} \rangle = -ie^{i\theta_0} \langle (\mu_1 + i\mu_2) \cdot (Q^1 - Q^2), a^1 a^2 e^{-ix \cdot \xi_0} \rangle$$

For II , we use the duality of $X_{1,j}^{m/2}$ and $X_{1,j}^{-m/2}$, the estimates (3.28), (6.4) and (6.5) to obtain

$$|II| = |\zeta_j^1 \cdot \langle (Q^1 - Q^2)\psi_j^1, a^2 e^{-ix \cdot \xi_0} \rangle| \lesssim \|(Q^1 - Q^2)\|_{X_{1,j}^{-m/2}} \|\psi_j^1\|_{X_{1,j}^{m/2}} \lesssim h_j^{m-2s+2-2\epsilon} = h_j^\epsilon.$$

where the last equality is our definition of ϵ . For III , we use the same argument with $X_{1,j}^\lambda$ replaced by $X_{2,j}^\lambda$

$$|III| = |\zeta_j^1 \cdot \langle (Q^1 - Q^2)a^1, e^{-ix \cdot \xi_0} \psi_j^2 \rangle| \lesssim \|(Q^1 - Q^2)\|_{X_{2,j}^{-m/2}} \|\psi_j^2\|_{X_{2,j}^{m/2}} \lesssim h_j^{m-2s+2-2\epsilon} = h_j^\epsilon.$$

For IV , we write Q^k in the form (3.30), use the estimate (3.23) and the estimates (6.5) for ψ_j^k to conclude

$$|IV| = |\zeta_j^1 \cdot \langle (Q^1 - Q^2)\psi_j^1, e^{-ix \cdot \xi_0} \psi_j^2 \rangle| \lesssim h_j^{1-2m} \|\psi_j^1\|_{X_{1,j}^{m/2}} \|\psi_j^2\|_{X_{2,j}^{m/2}} \lesssim h_j^{m+1-2s+2-2\epsilon} \lesssim h_j^{1+\epsilon}.$$

Since the amplitudes are smooth and there is no dependence on ψ_j^j , we have $|V| = |ih_j \langle (Q^1 - Q^2) \cdot Da^1, a^2 e^{-ix \cdot \xi_0} \rangle| \lesssim h_j$. For VI , we use duality in $X_{1,j}^{(m-1)/2}$, (3.19) and (6.5) to obtain

$$\begin{aligned}
|VI| &= |ih_j \langle (Q^1 - Q^2) \cdot Da^1, \psi_j^2 e^{-ix \cdot \xi_0} \rangle| \\
&\lesssim h_j \|(Q^1 - Q^2)\|_{X_{1,j}^{(1-m)/2}} \|D\psi_1\|_{X_{1,j}^{(m-1)/2}} \lesssim h_j^{m+1-2s+2-2\epsilon} = h_j^{1+\epsilon}
\end{aligned}$$

by (6.5) and (3.19) We may estimate VII using the same argument we used for III giving

$$|VII| = |ih_j \langle (Q^1 - Q^2) \cdot D\psi_j^1, a^2 e^{-ix \cdot \xi_0} \rangle| \lesssim h_j.$$

For $VIII$, we use the representation as in (3.30) and (3.23) to obtain

$$|VIII| = |ih_j \langle (Q^1 - Q^2) \cdot D\psi_j^1, e^{-ix \cdot \xi_0} \psi_j^2 \rangle| \lesssim h_j h_j^{-2m} \|\psi_j^1\|_{X_{1,j}^{m/2}} \|\psi_j^2\|_{X_{2,j}^{m/2}} \lesssim h_j h_j^{m-2s+2-\epsilon} = h_j^{1+\epsilon}.$$

where we have also used (6.5). For IX , we expand the product $(a^1 + \psi_j^1)(a^2 + \psi_j^2)$ and use similar arguments to obtain

$$\begin{aligned}
|IX| &= |ih_j \langle (q^1 - q^2)(a^1 + \psi_j^1), e^{-ix \cdot \xi_0} (a^2 + \psi_j^2) \rangle| \\
&\lesssim h_j (|\hat{q}^1(\xi_0) - \hat{q}^2(\xi_0)| + \|q^1 - q^2\|_{X_{2,j}^{-m/2}} \|\psi_j^2\|_{X_{2,j}^{m/2}} + \|q^1 - q^2\|_{X_{1,j}^{-m/2}} \|\psi_1\|_{X_{1,j}^{m/2}} \\
&\quad + h^{-2m} \|\psi_1\|_{X_{1,j}^{m/2}} \|\psi_{1,j}\|_{X_{m/2}}) \\
&\lesssim h_j (1 + h_j^{m-2s+2-2\epsilon}) = h_j + h_j^{1+\epsilon}.
\end{aligned}$$

Thus we can conclude

$$0 = \lim_{j \rightarrow \infty} ie^{-i\theta_0} h_j (B(u_1, u_2) - B_2(u_1, u_2)) = \langle (\mu_1 + i\mu_2) \cdot (Q^1 - Q^2), a^1 a^2 e^{-ix \cdot \xi_0} \rangle \quad (6.7)$$

and if we set $a^1 = a^2 = 1$, then we conclude

$$(\mu_1 + i\mu_2) \cdot (\hat{Q}^1(\xi_0) - \hat{Q}^2(\xi_0)) = 0.$$

If we repeat this argument with $-\mu_2$ replacing μ_2 we obtain

$$(\mu_1 - i\mu_2) \cdot (\hat{Q}^1(\xi_0) - \hat{Q}^2(\xi_0)) = 0.$$

Adding the last two displays we have $\mu \cdot (\hat{Q}^1(\xi_0) - \hat{Q}^2(\xi_0)) = 0$ if μ is perpendicular to ξ_0 . Choosing μ parallel to $e_k \xi_{0,j} - e_j \xi_{0,k}$ we conclude that

$$\frac{\partial}{\partial x_k} (Q_j^1 - Q_j^2) - \frac{\partial}{\partial x_j} (Q_k^1 - Q_k^2) = 0.$$

This gives us that $\text{curl}(Q^1 - Q^2) = 0$. Since we assume Ω is a ball, we may use Proposition A.4 to find g which is supported in $\bar{\Omega}$ and so that $Q^1 - Q^2 = Dg$. Now, we use (6.7) with $a^1 = 1$ and $a^2 = (\mu_1 - i\mu_2) \cdot x/2$. This choice for a^2 gives that $(\mu_1 + i\mu_2) \cdot Da^2 = -i$. Using that $(Q^1 - Q^2) = Dg$, we send $h_j \rightarrow 0$ and integrate by parts to obtain that

$$0 = (\mu_1 + i\mu_2) \cdot \langle Dg, e^{-ix \cdot \xi_0} a_2 \rangle = i \langle g, e^{-ix \cdot \xi_0} \rangle$$

where we used that $(\mu_1 + i\mu_2) \cdot \xi_0 = 0$. This gives us that $g = 0$ and we have that $Q^1 = Q^2$ as desired. Now that we have $Q^1 = Q^2$, we turn to the proof that $q^1 = q^2$. We use our CGO solutions with $a^1 = a^2 = 1$ to obtain

$$0 = B_1(u_1, u_2) - B_2(u_1, u_2) = \langle (q^1 - q^2)(1 + \psi_1), e^{-ix \cdot \xi_0} (1 + \psi_2) \rangle$$

Arguing as above (see the argument for the terms $I-IV$), we have

$$\lim_{j \rightarrow \infty} \langle (q^1 - q^2)(1 + \psi_1^j), (1 + \psi_2^j) e^{-ix \cdot \xi_0} \rangle = \hat{q}^1(\xi_0) - \hat{q}^2(\xi_0).$$

Since $\hat{q}^1 - \hat{q}^2 = 0$, we have $q^1 = q^2$. □

Proof of Main Theorem, Theorem 1.3. In section 5, we show that if the Dirichlet to Neumann to maps for two operators are equal, then the two forms are equal. Thus this result follows immediately from Theorem 6.6. □

A Sobolev spaces

We establish several representation theorems for elements of Sobolev spaces that are used in our arguments. Our results will depend on techniques developed by D. Mitrea, M. Mitrea, and S. Monniaux [25] who give a version of the Poincaré Lemma in their study of the Poisson problem for the exterior derivative. They construct a homotopy for the de Rham complex in a Lipschitz domain that is star-shaped with respect to each point in a small ball. The homotopy operators have the useful property that they preserve the class of forms that are supported in the domain. We briefly summarize what we need from their work and refer the reader to their article for precise definitions and complete statements of their results. They work with differential forms and initially assume the coefficients lie in $C_0^\infty(\Omega)$. Thus, let Λ denote the exterior algebra on \mathbb{R}^d and let Λ^j be the elements of degree j . We will use $C_0^\infty(\Omega; \Lambda^j)$ to denote the differential forms of degree j on Ω with coefficients in $C_0^\infty(\Omega)$. In [25, Theorem 4.1], Mitrea *et. al.* find a family of operators $J_j : C_0^\infty(\Omega; \Lambda^j) \rightarrow C_0^\infty(\Omega; \Lambda^{j-1})$, for $j = 1, \dots, d$ with the property that if $\omega \in C_0^\infty(\Omega; \Lambda^j)$

$$\omega = \begin{cases} J_1(d\omega), & j = 0 \\ J_{j+1}(d\omega) + dJ_j(\omega), & j = 1, \dots, d-1 \\ dJ_d(\omega) + \langle \omega, \theta V_d \rangle \theta V_d, & j = d \end{cases} \quad (\text{A.1})$$

Here, we are temporarily using d to denote both the dimension and the exterior derivative on \mathbb{R}^d . In the third line, $\theta \in C_0^\infty(\Omega)$ is determined in the construction of the operator J_d , $V_d = dx_1 \wedge \dots \wedge dx_d$ is the standard volume form on \mathbb{R}^d and the pairing of duality $\langle \cdot, \cdot \rangle$ is defined in section 2 of Mitrea *et. al.* [25]. In addition, the operators J_j map

$$J_j : \tilde{W}^{s,p}(\Omega; \Lambda^j) \rightarrow \tilde{W}^{s+1,p}(\Omega; \Lambda^{j-1}). \quad (\text{A.2})$$

To see this, consider J_j acting on a larger domain, and use the estimate of Theorem 3.8 in Mitrea *et. al.* and the property that J_j preserves the property of being supported in Ω .

Proposition A.3. *Assume that Ω is a Lipschitz domain. Suppose that $f \in \tilde{W}^{-s,p}(\Omega)$ with $s = k - r$ where $0 \leq r < 1$ and $k \geq 1$ is an integer. Then we may write*

$$f = \sum_{|\alpha|=k} D^\alpha F_\alpha + F_0$$

with $F_\alpha \in \tilde{W}^{r,p}(\Omega)$ and $F_0 \in C_c^\infty(\Omega)$

Proof. Using a partition of unity, we may reduce to the case where the domain Ω is star-shaped with respect to each point in a ball. We fix $f \in \tilde{W}^{s,p}(\Omega)$ and apply the case $j = d$ of (A.1) to the form $\omega = fV_d$. If we suppress the differentials, this can be rewritten as

$$f = D \cdot F + \langle u, \theta \rangle \theta$$

with $F = (F_1, \dots, F_d)$ and $F_j \in \tilde{W}^{s+1,p}(\Omega)$. If $k = 1$, we are done and for $k \geq 2$, we repeat this argument to represent each F_j as a smooth term and derivatives of functions in $\tilde{W}^{s+2,p}(\Omega)$. After k steps, we obtain the desired representation. \square

We remark that the specific case of Mitrea’s results used in the proof of Proposition A.3 may be found in earlier work of Bogovskii [8].

Proposition A.4. *Suppose that Ω is a ball in \mathbb{R}^d and $Q = (Q_1, \dots, Q_d)$ lies $\tilde{W}^{s,p}(\mathbb{R}^d)$ with $\text{curl } Q = 0$, then we may find $g \in \tilde{W}^{s+1,p}(\Omega)$ so that $Q = Dg$.*

Proof. We form a differential form $\omega = \sum_{k=1}^d Q_k dx_k$. Our hypothesis $\text{curl } Q = 0$ gives immediately that $d\omega = 0$ and then the Proposition follows from the case $j = 1$ of (A.1). \square

References

- [1] Yernat Assylbekov and Karthik Iyer. Determining rough first order perturbations of the polyharmonic operator. *Inverse Probl. Imaging*, 13(5):1045–1066, 2019.
- [2] Yernat M. Assylbekov. Inverse problems for the perturbed polyharmonic operator with coefficients in Sobolev spaces with non-positive order. *Inverse Problems*, 32(10):105009, 22, 2016.
- [3] Yernat M. Assylbekov. Corrigendum: Inverse problems for the perturbed polyharmonic operator with coefficients in Sobolev spaces with non-positive order (2016 Inverse Problems 32 105009) [MR3627033]. *Inverse Problems*, 33(9):099501, 2, 2017.
- [4] K. Astala and L. Päivärinta. Calderón’s inverse conductivity problem in the plane. *Ann. of Math. (2)*, 163(1):265–299, 2006.
- [5] J. Bergh and J. Löfström. *Interpolation spaces*. Springer-Verlag, Berlin, New York, 1976.
- [6] Sombuddha Bhattacharyya and Tuhin Ghosh. Inverse boundary value problem of determining up to a second order tensor appear in the lower order perturbation of a polyharmonic operator. *J. Fourier Anal. Appl.*, 25(3):661–683, 2019.
- [7] Sombuddha Bhattacharyya and Tuhin Ghosh. An inverse problem on determining second order symmetric tensor for perturbed biharmonic operator. *Math. Annalen*, 2021. Published online, 16 October 2021.
- [8] M. E. Bogovskii. Solutions of some problems of vector analysis, associated with the operators div and grad . In *Theory of cubature formulas and the application of functional analysis to problems of mathematical physics*, volume 1980 of *Trudy Sem. S. L. Soboleva*, No. 1, pages 5–40, 149. Akad. Nauk SSSR Sibirsk. Otdel. Inst. Mat., Novosibirsk, 1980.
- [9] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations. *Geom. Funct. Anal.*, 3(2):107–156, 1993.
- [10] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. II. The KdV-equation. *Geom. Funct. Anal.*, 3(3):209–262, 1993.

- [11] R.M. Brown. Global uniqueness in the impedance imaging problem for less regular conductivities. *SIAM J. Math. Anal.*, 27:1049–1056, 1996.
- [12] R.M. Brown and R.H. Torres. Uniqueness in the inverse conductivity problem for conductivities with $3/2$ derivatives in L^p , $p > 2n$. *J. Fourier Anal. Appl.*, 9(6):563–574, 2003.
- [13] A. P. Calderón. On an inverse boundary value problem. In *Seminar on Numerical Analysis and its Applications to Continuum Physics*, pages 65–73, Rio de Janeiro, 1980. Soc. Brasileira de Matemática.
- [14] Pedro Caro and Keith M. Rogers. Global uniqueness for the Calderón problem with Lipschitz conductivities. *Forum Math. Pi*, 4:e2, 28, 2016.
- [15] Anupam Pal Choudhury and Venkateswaran P. Krishnan. Stability estimates for the inverse boundary value problem for the biharmonic operator with bounded potentials. *J. Math. Anal. Appl.*, 431(1):300–316, 2015.
- [16] Filippo Gazzola, Hans-Christoph Grunau, and Guido Sweers. *Polyharmonic boundary value problems*, volume 1991 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2010. Positivity preserving and nonlinear higher order elliptic equations in bounded domains.
- [17] Tuhin Ghosh and Venkateswaran P. Krishnan. Determination of lower order perturbations of the polyharmonic operator from partial boundary data. *Appl. Anal.*, 95(11):2444–2463, 2016.
- [18] B. Haberman and D. Tataru. Uniqueness in Calderón’s problem with Lipschitz conductivities. *Duke Math. J.*, 162(3):496–516, 2013.
- [19] Boaz Haberman. Uniqueness in Calderón’s problem for conductivities with unbounded gradient. *Comm. Math. Phys.*, 340(2):639–659, 2015.
- [20] Seheon Ham, Yehyun Kwon, and Sanghyuk Lee. Uniqueness in the Calderón problem and bilinear restriction estimates. *J. Funct. Anal.*, 281(8):Paper No. 109119, 58, 2021.
- [21] Katsiaryna Krupchyk, Matti Lassas, and Gunther Uhlmann. Determining a first order perturbation of the biharmonic operator by partial boundary measurements. *J. Funct. Anal.*, 262(4):1781–1801, 2012.
- [22] Katsiaryna Krupchyk, Matti Lassas, and Gunther Uhlmann. Inverse boundary value problems for the perturbed polyharmonic operator. *Trans. Amer. Math. Soc.*, 366(1):95–112, 2014.
- [23] Katsiaryna Krupchyk and Gunther Uhlmann. Inverse boundary problems for polyharmonic operators with unbounded potentials. *J. Spectr. Theory*, 6(1):145–183, 2016.
- [24] William McLean. *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, Cambridge, 2000.

- [25] D. Mitrea, M. Mitrea, and S. Monniaux. The Poisson problem for the exterior derivative operator with Dirichlet boundary condition in nonsmooth domains. *Commun. Pure Appl. Anal.*, 7(6):1295–1333, 2008.
- [26] Lassi Päivärinta, Alexander Panchenko, and Gunther Uhlmann. Complex geometrical optics solutions for Lipschitz conductivities. *Rev. Mat. Iberoamericana*, 19(1):57–72, 2003.
- [27] Felipe Ponce-Vanegas. The bilinear strategy for Calderón’s problem, 2019. arXiv:1908.04050 [math.AP].
- [28] Felipe Ponce-Vanegas. Reconstruction of the derivative of the conductivity at the boundary. *Inverse Probl. Imaging*, 14(4):701–718, 2020.
- [29] T. Runst and W. Sickel. *Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations*. Walter de Gruyter & Co., Berlin, 1996.
- [30] V. S. Serov. Borg-Levinson theorem for perturbations of the bi-harmonic operator. *Inverse Problems*, 32(4):045002, 19, 2016.
- [31] J. Sylvester and G. Uhlmann. A global uniqueness theorem for an inverse boundary value problem. *Annals of Math.*, 125:153–169, 1987.
- [32] G.C. Verchota. Layer potentials and regularity for the Dirichlet problem for Laplace’s equation on Lipschitz domains. *J. Funct. Anal.*, 59:572–611, 1984.
- [33] Gregory Verchota. The Dirichlet problem for the polyharmonic equation in Lipschitz domains. *Indiana Univ. Math. J.*, 39(3):671–702, 1990.
- [34] Lili Yan. Inverse boundary problems for biharmonic operators in transversally anisotropic geometries. arXiv:2012.14273 [math.AP].
- [35] Lili Yan. Reconstructing a potential perturbation of the biharmonic operator on transversally anisotropic manifolds. arXiv:2109.07712.

posted to arXiv on 26 August 2021,
revised 7 December 2021