PARABOLICITY OF ZERO-TWIST TIGHT FLUTE SURFACES AND UNIFORMIZATION OF THE LOCH NESS MONSTER

JOHN A. ARREDONDO, ISRAEL MORALES, CAMILO RAMÍREZ MALUENDAS

ABSTRACT. We study the zero-twist flute surface and we associate to each one of them a sequence of positive real numbers $\mathbf{x}=(x_n)_{n\in\mathbb{N}_0}$, with a torsion-free Fuchsian group $\Gamma_{\mathbf{x}}$ such that the convex core of $\mathbb{H}^2/\Gamma_{\mathbf{x}}$ is isometric to a zero-twist tight flute surface $S_{\mathbf{x}}$. Moreover, we prove that the Fuchsian group $\Gamma_{\mathbf{x}}$ is of the first kind if and only if the series $\sum x_n$ diverges. As consequence of the recent work of Basmajian, Hakobian and Šarić, we obtain that the zero-twist flute surface $S_{\mathbf{x}}$ is of parabolic type if and only $\sum x_n$ diverges. In addition, we present an uncountable family of hyperbolic surfaces homeomorphic to the Loch Ness Monster. More precisely, we associate to each sequence $\mathbf{y}=(y_n)_{n\in\mathbb{Z}}$, where $y_n=(a_n,b_n,c_n,d_n,e_n)\in\mathbb{R}^5$ and $a_n\leq b_n\leq c_n\leq d_n\leq e_n\leq a_{n+1}$, a Fuchsian group $G_{\mathbf{y}}$ such that $\mathbb{H}^2/G_{\mathbf{y}}$ is homeomorphic to the Loch Ness Monster.

INTRODUCTION

A fundamental open question in the classification of Riemann surfaces, known as the *type problem* is:

whether a Riemann surface S supports a Green's function? In the context of simply connected Riemann surfaces we have that the Poincaré disc Δ admits Green's functions, while the Riemann sphere $\widehat{\mathbb{C}}$ and the complex plane \mathbb{C} do not. From this, recall that a Riemann surface S is **elliptic** if and only if S is compact (equivalently, closed). On the other hand, an open Riemann surface S is said to be **parabolic**, if S does not carry a negative non-constant subharmonic function. All open Riemann surface which are not parabolic will be called **hyperbolic** (see [12, p.164], [19, Section 2]). It is well known that the type problem becomes equivalent to answering the question

which open Riemann surfaces are of parabolic type?

The interest in solving this problem has captured the attention of the mathematical community for almost one hundred years. Some of the first partial results to this question, in modern language can be read in the theorems of

1

²⁰⁰⁰ Mathematics Subject Classification. 20H10, 57N05, 57N16, 30F20, 30F25, 30F45, 57K20.

Key words and phrases. Parabolic type, Fenchel-Nielsen parameters, zero-twist tight flute surface, Loch Ness monster.

section 6 in [2, p. 204]. After these results, numerous known characterizations of parabolic surfaces, that are equivalent to the type problem have been produced from potential theory, theory of functions, dynamics and geometry of surfaces, among others. This is a short list of known characterizations of Riemann surfaces of parabolic type: if the Riemann surface S is the quotient of the Poincaré unit disk Δ by a Fuchsian group Γ , then S is parabolic if and only if

- The series $\sum_{\gamma \in \Gamma} \left(\frac{1 ||\gamma(\mathbf{0})||}{1 + ||\gamma(\mathbf{0})||} \right)$ diverges [18, Theorem 5.2.1],
- The geodesic flow on the unit tangent bundle of *S* is ergodic [18, Theorem 8.3.4];
- The Mostow rigidity holds for Γ [1], [7], [24];
- The group Γ has the Bowen's property [7], [13].

Parabolicity of zero-twist tight flute surfaces. Open Riemann surfaces are classified, up to homeomorphisms, by their genus $g(S) \in \mathbb{N}_0 \cup \{\infty\}$, and a pair of nested topological spaces $\operatorname{Ends}_{\infty}(S) \subseteq \operatorname{Ends}(S)$ homeomorphic to a pair of nested closed subsets of the Cantor space. In this context, the flute surfaces are the unique infinite-type Riemann surfaces, up to homeomorphism, of genus zero and with space of ends homeomorphic to the ordinal number $\omega + 1$ (§1.1). The classof flute surfaces is one of the simplest families where we can ask about the parabolicity problem. However, even in this class the problem of parabolicity is widely open (see [9], [17]). A better manage family in the class of Riemann flute surfaces are the so-called tight flute surfaces. Such surfaces are hyperbolic surfaces obtained by starting with a geodesic pair of pants P_0 with two punctures and one boundary geodesic and consecutively gluing geodesic pair of pants P_n $(n \ge 1)$ with one cusp and two boundary geodesics in an infinite chain (§1.5). A tight flute surface is determined by its Fenchel-Nielsen parameters (§1.2) $(\{l_n, t_n\})_{n \in \mathbb{N}}$ where l_n and t_n are the length and twist parameter of the boundary closed geodesic α_n of the surface obtained after gluing n pair of pants. This surface is denoted by $S = S(\{l_n, t_n\})_{n \in \mathbb{N}}$. If $t_n = 0$ for all $n \in \mathbb{N}$, then we say that $S = S(\{l_n, 0\})_{n \in \mathbb{N}}$ is a zero-twist flute surface.

In this article, we aim to present a new way of describing zero-twist flute surfaces; this description differs from the Fenchel-Nielsen parameters. Our main contribution is as follows:

Theorem 0.1. For each zero-twist tight flute surface $S = S(\{l_n, 0\})_{n \in \mathbb{N}_0}$, there exists a unique sequence $\mathbf{x} = (x_n)_{n \in \mathbb{N}_0}$ of positive real numbers, which

defines the Fuchsian group $\Gamma_{\mathbf{x}}$ generated by

(1)
$$g_n(z) := \frac{\left(1 + \frac{2s_{n-1}}{s_n - s_{n-1}}\right)z - 2s_{n-1}\left(1 + \frac{s_{n-1}}{s_n - s_{n-1}}\right)}{-\frac{2}{s_n - s_{n-1}}z + \left(1 + \frac{2s_{n-1}}{s_n - s_{n-1}}\right)} \in PSL(2, \mathbb{R}),$$

with $n \in \mathbb{N}_0$ and $s_n = \sum_{i=0}^n x_i$, such that the convex core of the surface $\mathbb{H}^2/\Gamma_{\mathbf{x}}$ is isometric to S. Moreover, Γ_x is of the first kind if and only if the series $\sum x_n$ diverges. In this case, $\mathbb{H}^2/\Gamma_{\mathbf{x}}$ coincides with S and it is geodesically complete.

Our construction above is inspired on A. Basmajian work in [8], where the author present a realization theorem showing that $(\{l_n, t_n\})_{n \in \mathbb{N}}$ are the Fenchel-Nielsen coordinates of a tight flute surface if and only if $r_n > \ln 2$ for all n odd, and $r_n < \ln 2$ for all n even, where $r_n := \sum_{i=1}^n (-1)^{i+1} l_i$, see [8, Theorem 2]. Observe that these conditions does not depend on twist parameters. In contrast with our construction of a zero-twist tight flute surface in terms of the sequence $\mathbf{x} = (x_n)_{n \in \mathbb{N}_0} \in \mathbb{R}_+^{\mathbb{N}_0}$, we have that it does not depend of any additional condition on the sequence. Additionally, we can recover the Fenchel-Nielsen parameters of $S = S(\{l_n, 0\})_{n \in \mathbb{N}_0}$ using

(2)
$$l_n = \ln\left(\frac{s_{n-1} + s_n + 2\sqrt{s_{n-1}s_n}}{s_{n-1} + s_n - 2\sqrt{s_{n-1}s_n}}\right).$$

Let $\mathbb{R}^{\mathbb{N}_0}_+$ be the space of all sequences of positive real numbers indexed by the set \mathbb{N}_0 endowed with the compact-open topology, and denote by \mathcal{F} the set of all zero-twist flute surfaces, that is, $\mathcal{F} := \{S = S(\{l_n, 0\})_{n \in \mathbb{N}_0}\}.$ Accordingly to Theorem 0.1, there is a bijective map $\sigma: \mathbb{R}^{\mathbb{N}_0}_+ \to \mathcal{F}$, which associates to each sequence in $\mathbb{R}^{\mathbb{N}_0}_+$ a zero-twist flute surface. We denote by $S_{\mathbf{x}}$ the image of $\mathbf{x} \in \mathbb{R}^{\mathbb{N}_0}_+$ under σ .

Recently, A. Basmajian, H. Hakobyan and D. Šarić in [9, Theorems 1.1 and 1.2] give sufficient conditions on Fenchel-Nielsen parameters of a surface to determine parabolicity. In the particular case of zero-twist flute surfaces, they were able to characterize parabolicty.

Theorem 0.2 ([9, Theorem 1.5]). A zero-twist tight flute surface $S(\{l_n, 0\})_{n \in \mathbb{N}}$ is of parabolic type if and only if one of the following holds:

- (1) The surface $S(\{l_n, 0\})_{n \in \mathbb{N}}$ is complete. (2) The series $\sum_{n=1}^{\infty} e^{-\frac{l_n}{2}} = \infty$.

Using equation (2) and Theorem 0.1, we have that items (1) and (2) in Theorem 0.2 above are equivalent. Thus we obtain the following:

Corollary 0.1. A zero-twist flute surface S_x is of parabolic type if and only if $\sum x_n$ diverges.

Theorem 0.1 also gives an explicit set of generators for $\Gamma_{\mathbf{x}}$. We observe that if $\mathbf{x} \in \{1, 2\}^{\mathbb{N}_0}$ then $\Gamma_{\mathbf{x}}$ is a subgroup of PSL(2, \mathbb{Z}). This proves

Corollary 0.2. The set of all zero-twist tight flute surfaces uniformized by subgroups of $PSL(2, \mathbb{Z})$ can be identified to a homeomorphic copy of the Cantor set.

An uncountable family of hyperbolic Loch Ness monsters. Recall that the Loch Ness monster surface is the unique surface (up to homeomorphism) which has infinite genus and only one end (§1.1). In [6] the authors introduced a Fuchsian group uniformizing the Loch Ness monster. We generalize their construction and find an uncountable family of hyperbolic surfaces such that each one of these surface is homeomorphic to the Loch Ness Monster.

Denote by N the set of all sequences $\mathbf{y} := (y_n)_{n \in \mathbb{Z}}$ of elements $y_n = (a_n, b_n, c_n, d_n, e_n) \in \mathbb{R}^5$ satisfying

(3)
$$a_n < b_n < c_n < d_n < e_n \text{ and } e_n \le a_{n+1}$$
.

Let $\mathbf{y} := (y_n)_{n \in \mathbb{Z}} \in \mathcal{N}$. For each $n \in \mathbb{Z}$, let f_n and g_n be elements of $\mathrm{PSL}(2,\mathbb{R})$ mapping σ_n onto $\tilde{\sigma}_n$ and ρ_n onto $\tilde{\rho}_n$, respectively, where σ_n , ρ_n , $\tilde{\sigma}_n$ and $\tilde{\rho}_n$ are the half-circles in the hyperbolic plane \mathbb{H}^2 depicted in Figure 1.




FIGURE 1. Half-circles σ_n , ρ_n , $\tilde{\sigma}_n$ and $\tilde{\rho}_n$.

Define

(4)
$$G_{\mathbf{v}} := \langle f_n, g_n : n \in \mathbb{Z} \rangle \leq \mathrm{PSL}(2, \mathbb{R}).$$

The result we establish is the following:

Theorem 0.3. For each $\mathbf{y} \in \mathcal{N}$, the group $G_{\mathbf{y}}$ is Fuchsian and such that $S_{\mathbf{y}} := \mathbb{H}^2/G_{\mathbf{y}}$ is topologically equivalent to the Loch Ness Monster. Moreover, $G_{\mathbf{y}}$ is of the first kind if and only if $e_n = a_{n+1}$ for all $n \in \mathbb{Z}$, $\lim_{n \to \infty} e_n = \infty$ and $\lim_{n \to -\infty} a_n = -\infty$.

Organization of the paper. In Section 1 we compile some classic results and concepts of the theory of surfaces necessary for the development of this work. Section 2 is dedicated to the proof of Theorem 0.1. In Section 3 we give the proof of Theorem 0.3.

1. Preliminaries

1.1. **Topological surfaces.** Topological orientable surfaces are classified, up to homeomorphisms, by their genus $g(S) \in \mathbb{N}_0 \cup \{\infty\}$, and a pair of nested topological spaces $\operatorname{Ends}_{\infty}(S) \subseteq \operatorname{Ends}(S)$ homeomorphic to a pair of nested closed subsets of the Cantor set. The spaces $\operatorname{Ends}(S)$ and $\operatorname{Ends}_{\infty}(S)$ are called the *end space* and the *non-planar ends space* of S (or, ends accumulated by genus), respectively. Moreover, any pair of nested closed subsets of the Cantor set can be realized as the space of ends of a connected, orientable topological surface. We call to the isolated elements of the end space *punctures* of the surface. For more details, we refer the reader to [20].

Theorem 1.1 (Classification of topological surfaces, [15, §7], [20, Theorem 1]). Two orientable surfaces S_1 and S_2 having the same genus are topological equivalent if and only if there exists a homeomorphism $f : \operatorname{Ends}(S_1) \to \operatorname{Ends}(S_2)$ such that $f(\operatorname{Ends}_{\infty}(S_1)) = \operatorname{Ends}_{\infty}(S_2)$.

A topological surface is of *infinite-type* if it has fundamental group infinitely generated. The simplest infinite-type surface that we can find are the flutes and the Loche Ness monster.

Definition 1.1 ([8, p. 423]). The *flute surface* is the unique topological surface, up to homeomorphism, of genus zero and with ends space homeomorphic to the ordinal number $\omega + 1$, see Figure 2.




FIGURE 2. A flute surface.

Definition 1.2. The **Loch Ness monster** is the unique infinite-type surface, up to homeomorphism, of infinite genus with exactly one end, see Figure 3.




FIGURE 3. The Loch Ness monster.

In section 3 we use the following result to prove the first part of Theorem 0.3.

Lemma 1.1 ([22, §5.1., p. 320]). The surface S has exactly one end if and only if for all compact subset $K \subset S$ there is a compact subset $K' \subset S$ such that $K \subset K'$ and $S \setminus K'$ is connected.

We say that a simple close curved on a surface *S* is *essential* if it is not isotopic to the boundary loop of a disk or a punctured disk. Recall that a *topological pair of pants* is a topological surface homeomorphic to a three punctured sphere.

Definition 1.3. We say that a collection of pairwise disjoint essential curves in a surface S is a **pair of pants decomposition** of S if it decomposes the surface into a disjoint union of pair of pants.

Observe that any topological surface admits a pair of pants decomposition [4].

1.2. **Fenchel-Nielsen parameters of a hyperbolic surfaces.** For the results presented in this section we refer the reader to [11, Chapter 10] for the definition of the Fenchel-Nielsen parameters in the context of finite-type surfaces, and to [3] for surfaces of infinite-type.

A geodesic pair of pants is a hyperbolic surface P of finite hyperbolic area such that its interior is homeomorphic to a pair of pants and with at least one boundary a closed geodesic, see Figure 4. A tight pair of pants is a pair of pants that has exactly one puncture, see Figure 4-b. Similarly, a collection of pairwise disjoint of essential geodesic curves in a surface S is geometric pair of pants decomposition of S if it decomposes the surface into geodesic pair of pants. In [3, Theorem 4.5] the authors give sufficient conditions

under which a topological and geodesic decomposition of a surface in pair of pants is related .




FIGURE 4. Three possible kinds of hyperbolic pants with geodesic boundary. In particular, b) represents a tight pair of pants.

1.2.1. Length and twist parameters. Let P be a geodesic pair of pants and fix α a boundary curve of P. We choose a marked point x in α in the following way: Let β be either a boundary component (different from α) or a puncture of P. Let γ be the orthogeodesics between α and β . Then the marked point x in α is defined as the intersection point of α with γ . In Figure 5 is shown such marked point x in α which is obtained from drawing the orthogeodesic γ connecting the boundary component α with β . The hyperbolic length of the geodesic α is denoted by $l(\alpha)$.




FIGURE 5. The point $x \in \alpha$ is the intersection of α with the orthogeodesic γ .

Let P and P' be two geodesic pair of pants with α and α' boundary geodesic curve of P and P', respectively. Let γ and γ' be the geodesic in P and P', respectively, that define the marked points x in α and x' in α' as above. Suppose that α and α' have the same hyperbolic length, that is, $l(\alpha) = l(\alpha')$. Then we identify α and α' , by gluing with an isometry T, to obtain a surface S from P and P'. This isometric identification is determined by the pair $\{l(\alpha), t(\alpha)\}$, such that the *length parameter* $l(\alpha)$ is the hyperbolic length of α ,

and the *twist parameter* $t(\alpha) \in \left(-\frac{1}{2}, \frac{1}{2}\right]$ corresponds to the relative position of the marked points x in α and x' in α' , which is given as follows:

- * If x = T(x'), then $t(\alpha) = 0$.
- * If $x \neq T(x')$, then $|t(\alpha)| \leq \frac{1}{2}$ is equal to the hyperbolic length of the shorter arc σ having endpoints x and T(x') contained in $\alpha \setminus \{x, T(x')\}$, divided by $|t(\alpha)|$. If $|t(\alpha)| = \frac{1}{2}$, then $|t(\alpha)| = \frac{1}{2}$. If $|t(\alpha)| < \frac{1}{2}$ then we orient α with the orientation induced from P. If σ is the arc from x to T(x') then $|t(\alpha)| = |t(\alpha)|$; otherwise $|t(\alpha)| = -|t(\alpha)|$. See Figure 6. a).




FIGURE 6

Remark 1.1. If the twist parameter $t(\alpha) = 0$, then $\gamma \cup \gamma'$ is a geodesic in S that is orthogonal to the closed geodesic α , see Figure 6. b).

Definition 1.4. Let S be a surface obtained by gluing countably many geodesic pair of pants in the way described above. These gluing have associated a geometric pair of pants decomposition $\{\alpha_i : i \in \mathbb{N}\}\$ of S. Let $l_i(\alpha) = l_i$ and $t_i(\alpha) = t_i$ be the length and twist parameter respectively of α_i . The collection of pairs

$$(\{l_i, t_i\})_{i \in \mathbb{N}},$$

is called the **Fenchel-Nielsen parameters** of S. As the hyperbolic metric on S is uniquely determined by its Fenchel-Nielsen parameters, then

$$S := S(\{l_i, t_i\})_{i \in \mathbb{N}}.$$

The surface S might not be complete in the induced hyperbolic metric. V. Álvarez and J. M. Rodríguez in [4] showed that the boundary of the metric completion of S consists of simple closed geodesics and bi-infinite simple geodesics. Moreover, they proved that by attaching funnels to the closed

geodesics and attaching geodesic half-planes to the bi-infinite geodesics of the boundary of the metric completion of S, we obtain a surface \hat{S} homeomorphic to S with a geodesically complete hyperbolic metric such that the inclusion $i:S\hookrightarrow \hat{S}$ is an isometric embedding. Conversely, any geodesically complete surface is obtained by attaching funnels and half-planes to the convex core of the surface, see also [10].

We are in a position to define the so-called *tight flute surfaces*.

Definition 1.5 ([8, p. 423]). A tight flute surface S is a surface obtained by starting with a geodesic pair of pants P_0 with two punctures and then consecutively gluing tight pairs of pants P_n , $n \ge 1$.

If S is a tight flute surface, we denote by α_n the closed geodesic of the boundary of the surface obtained after gluing n geodesic pairs of pants and let l_n and t_n be the length and twist parameters associated to α_n . In terms of Fenchel-Nielsen parameters this surface is denoted by $S := S(\{l_n, t_n\})_{n \in \mathbb{N}}$.

Definition 1.6. If all twist parameters of a tight flute surface are zero, that is $S := S(\{l_n, 0\})_{n \in \mathbb{N}}$, then we called it a **zero-twist tight flute surface**.

2. Proof of Theorem 0.1

We begin by associating to each zero-twist flute surface S a sequence \mathbf{x} of positive real numbers and a Fuchsian group $\Gamma_{\mathbf{x}}$ such that the convex core of $\mathbb{H}^2/\Gamma_{\mathbf{x}}$ is isometric to S. We do this by constructing explicitly a hyperbolic ideal polygon \mathcal{P} by cutting S along an infinite collection of bi-infinite geodesics which serves as fundamental domain for $\Gamma_{\mathbf{x}}$.

Let $S = S(\{l_n, 0\})_{n \in \mathbb{N}_0}$ be a zero-twist flute surface (see Figure 7). For each $n \in \mathbb{N}$, let α_n be the closed geodesic in S, which comes from gluing the geodesic pair of pants P_{n-1} and P_n . Let $\mathbf{0}$ and s_0 be the punctures of P_0 and for $n \geq 1$, let s_n be the unique puncture of P_n . Now we describe how to obtain P by removing a collection $\{\gamma_n\}_{n \in \mathbb{N}_0}$ of geodesics in S having endpoints on the punctures of S. Let be the geodesics $\gamma_0 \subset P_0$ connecting the two punctures of P_0 , and γ_n connecting the punctures s_{n-1} and s_n , the last one is orthogonal to the close geodesic α_{n-1} , see Remark 1.1. Finally, we draw on S the geodesic ray β orthogonal to each α_n and having one endpoint in $\mathbf{0}$ (see Figure 7). Note that S admits a reflection τ whose connected components of fixed points are the geodesic arcs γ_n and β .

Now, we cut S along each γ_n . Then S turns into an ideal hyperbolic polygon \mathcal{P} with infinitely many sides. More precisely, the edges and vertices of \mathcal{P} are given as follows:

* For each $n \ge 0$, let γ_n^+ and γ_n^- be the sides of \mathcal{P} coming from cutting S along γ_n .




FIGURE 7. A zero-twist tight flute surface.

* For each $n \ge 0$, let s_n^+ and s_n^- be the ideal vertices of \mathcal{P} coming from the puncture s_n .

Remark 2.1. For each $n \ge 0$, the sides γ_n^+ and γ_n^- of the ideal hyperbolic polygon \mathcal{P} are identified by an element g_n of $PSL(2,\mathbb{R})$. If we identify the sides γ_n^+ and γ_n^- using g_n , then we recover the zero-twist tight surface $S(\{l_n,0\})_{n\in\mathbb{N}_0}$.

The ideal hyperbolic polygon \mathcal{P} can be thought in the hyperbolic plane \mathbb{H}^2 satisfying the following properties (see Figure 8):

- * Its vertices are on the real axis. Up to take a real translation, we can assume that the vertex $\mathbf{0}$ of \mathcal{P} is equal to the complex number zero and that the geodesic ray β coincides with the imaginary axis. The collection of vertices $(s_n^+)_{n\in\mathbb{N}_0}$ defines a strictly increasing sequence of positive real numbers. The collection of vertices $(s_n^-)_{n\in\mathbb{N}_0}$ defines a strictly decreasing sequence of negative real numbers. Given that \mathcal{P} has a reflexive symmetry fixing the ray β , then s_n^+ and s_n^- are symmetric with respect to the imaginary axis, it means, $s_n^- = -s_n^+$, for all $n \ge 0$.
- * For each $n \in \mathbb{N}_0$, the edges γ_n^+ and γ_n^- of \mathcal{P} are half-circles having the same radius and endpoints s_{n-1}^+ and s_n^+ ; and s_{n-1}^- and s_n^- , respectively. Thus, the intersection of any two of those edges is either empty or they meet at the same point in the real line. Moreover, if we choose γ one of these half-circles, then the other half-circles are in the exterior of γ .




FIGURE 8. Realization of the ideal hyperbolic polygon \mathcal{P} in \mathbb{H}^2 .

To the zero-twist flute surface S we associated the sequence $\mathbf{x} = (x_n)_{n \in \mathbb{N}_0}$ of positive real numbers that satisfies

(6)
$$s_n^+ = \sum_{i=0}^n x_i.$$

Thus, to the sequence x we associated the group Γ_x given by

(7)
$$\Gamma_{\mathbf{x}} = \langle g_n : n \in \mathbb{N}_0 \rangle.$$

Now we prove that $\Gamma_{\mathbf{x}}$ satisfies the properties described in Theorem 0.1. For each $n \in \mathbb{N}_0$, let $s_n := s_n^+$. On the hyperbolic plane \mathbb{H}^2 , we draw the half-circles: γ_0^+ and γ_0^- having endpoints $\mathbf{0}$ and s_0 , and $\mathbf{0}$ and $-s_0$, respectively; For each $n \ge 1$, γ_n^+ and γ_n^- having endpoints s_{n-1} and s_n , and $-s_{n-1}$ and $-s_n$, respectively. See Figure 8. We remark that γ_n^+ and γ_n^- are symmetric with respect to the imaginary axis. Then the element g_n of PSL(2, \mathbb{R}) is given by

(8)
$$g_n(z) := \frac{\left(1 + \frac{2s_{n-1}}{s_n - s_{n-1}}\right)z - 2s_{n-1}\left(1 + \frac{s_{n-1}}{s_n - s_{n-1}}\right)}{-\frac{2}{s_n - s_{n-1}}z + \left(1 + \frac{2s_{n-1}}{s_n - s_{n-1}}\right)},$$

which sends γ_n^+ onto γ_n^- , for each $n \ge 0$. Observe that g_0 is parabolic and, for each $n \ge 1$, g_n is hyperbolic with trace equal to

$$\left| 2\left(1 + \frac{2s_{n-1}}{s_n - s_{n-1}}\right) \right| > 2.$$

As $\Gamma_{\mathbf{x}}$ is obtained by side-pairing the sides of the convex ideal polygon \mathcal{P} , then it is a non elementary group and its set of generators is composed by non elliptic elements. Thus, $\Gamma_{\mathbf{x}}$ is a Fuchsian group (see [12, Theorem 8.3.1, p. 198]), with fundamental domain \mathcal{P} for its action on \mathbb{H}^2 . In what follows we describe \mathcal{P} more precisely.

Recall that if γ is a half-circle in \mathbb{H}^2 having as center the point $z \in \mathbb{R}$ and radius r > 0, then $\mathbb{H}^2 \setminus \gamma$ has two connected components. The connected component of $\mathbb{H}^2 \setminus \gamma$ equal to the set $\{w \in \mathbb{H}^2 : |z - w| > r\}$ is called the *exterior of* γ and it is denoted by $\operatorname{Ext}(\gamma)$. The complement of the closure in \mathbb{H}^2 of $\operatorname{Ext}(\gamma)$ is called the *interior of* γ and is denoted by $\operatorname{Int}(\gamma)$, see Figure 9.




FIGURE 9. Exterior and interior of a half-circle γ in \mathbb{H}^2 .

The limit $\lim_{n\to\infty} s_n$ in $\mathbb{R} \cup \{\infty\}$ is well defined because the sequence $(s_n)_{n\in\mathbb{N}_0}$ is the partial sums of \mathbf{x} . Let a be such limit. If a is infinity, then \mathcal{P} is equal to the closure (in \mathbb{H}^2) of

$$D:=\bigcap_{n\in\mathbb{N}_0}\left(\operatorname{Ext}(\gamma_n^+)\cap\operatorname{Ext}(\gamma_n^-)\right)\subset\mathbb{H}^2.$$

Otherwise, if a is a positive real number, then we denote by γ the half-circle in \mathbb{H}^2 having a and -a as ends points. In this case \mathcal{P} is given by

$$\overline{D} \cap \operatorname{Int}(\gamma) \subset \mathbb{H}^2$$
.

Notice that \overline{D} is a fundamental domain for $\Gamma_{\mathbf{x}}$ and $\overline{D}/\Gamma_{\mathbf{x}}$ is a complete hyperbolic surface. Therefore, if a is infinity then $\mathcal{P}/\Gamma_{\mathbf{x}} = \overline{D}/\Gamma_{\mathbf{x}}$ is isometric to S and $\Gamma_{\mathbf{x}}$ is a Fuchsian group of the first kind. Otherwise, if a is finite then $\mathcal{P}/\Gamma_{\mathbf{x}}$ is isometric to S and it is not a complete hyperbolic surface because the projection points of γ are limit points which are not in the surface. However, $\mathcal{P}/\Gamma_{\mathbf{x}}$ is the convex core of $\overline{D}/\Gamma_{\mathbf{x}}$. Here finished the proof of Theorem 0.1.

3. Proof of Theorem 0.3

From the introduction we recall that \mathcal{N} denotes the set of all sequences $\mathbf{y} := (y_n)_{n \in \mathbb{Z}}$ of elements $y_n = (a_n, b_n, c_n, d_n, e_n) \in \mathbb{R}^5$ satisfying

(9)
$$a_n < b_n < c_n < d_n < e_n \text{ and } e_n \le a_{n+1}$$

For each $n \in \mathbb{Z}$, let f_n and g_n be elements of PSL $(2, \mathbb{R})$ mapping σ_n onto $\tilde{\sigma}_n$ and ρ_n onto $\tilde{\rho}_n$, respectively, where σ_n , ρ_n , $\tilde{\sigma}_n$ and $\tilde{\rho}_n$ are the half-circles in the hyperbolic plane \mathbb{H}^2 depicted in Figure 1. Let

(10)
$$G_{\mathbf{v}} := \langle f_n, g_n : n \in \mathbb{Z} \rangle \leq \mathrm{PSL}(2, \mathbb{R}).$$

To start with the proof of Theorem 0.3, the following lemma is required.

Lemma 3.1. Let σ and $\tilde{\sigma}$ denote the half-circles having endpoints a and b; c and d respectively, where a < b < c < d (see Figure 10). Then the element f of $PSL(2,\mathbb{R})$ which sends the half-circle σ onto the half-circle $\tilde{\sigma}$ is hyperbolic.

Proof. By hypothesis, the half-circles σ and $\tilde{\sigma}$ are disjoint, and each one of the half-circles is contained in the exterior of the other half-circle. Explic-




FIGURE 10. Half-circles σ and $\tilde{\sigma}$.

itly, they have centers at the real numbers $O=a+\frac{b-a}{2}$ and $\tilde{O}=c+\frac{d-c}{2}$ respectively, and radius $r=\frac{b-a}{2}$ and $\tilde{r}=\frac{d-c}{2}$, respectively. Then the element f of $PSL(2,\mathbb{R})$ is given by

(11)
$$f(z) = \frac{-r\tilde{r}}{z - O} + \tilde{O}.$$

Let us note that the fixed points of f are

(12)
$$z = \frac{-(r+\tilde{r}) \pm \sqrt{\delta}}{2},$$

where $\delta = (O + \tilde{O})^2 - 4(O\tilde{O} + r\tilde{r}) = (O - \tilde{O})^2 - 4r\tilde{r}$. Since $0 < r + \tilde{r} < |\tilde{O} - O|$, then

$$0 \leq (r-\tilde{r})^2 = (r+\tilde{r})^2 - 4r\tilde{r} < (\tilde{O}-O)^2 - 4r\tilde{r} = \delta.$$

It implies that $\sqrt{\delta}$ is a real number. Therefore, f is hyperbolic.

The group G_y is Fuchsian. By Lemma 3.1, G_y is generated by hyperbolic elements. So, by Theorem 8.2.1 in [12], G_y is a Fuchsian group. By construction, the collection of half-circles $C = \{\sigma_n, \tilde{\sigma}_n, \rho_n, \tilde{\rho}_n : n \in \mathbb{Z}\}$ are pairwise disjoint, and each one of the half-circles in C is contained in the

exterior of each one of the other half-circles in C. From Theorem 3.3.5 in [14], a fundamental region $D(G_v)$ for the Fuchsian group G_v is given by

(13)
$$D(G_{\mathbf{y}}) = \bigcap_{n \in \mathbb{Z}} (\operatorname{Ext}(\sigma_n) \cap \operatorname{Ext}(\tilde{\sigma}_n) \cap \operatorname{Ext}(\rho_n) \cap \operatorname{Ext}(\tilde{\rho}_n)) \subset \mathbb{H}^2.$$

Given that the intersection of any two different elements belonged to C is either empty or at infinity, the last means, they meet at the same point in the real line, then G_y acts freely and properly discontinuously on whole \mathbb{H}^2 , see [16, p. 29]. Hence, the quotient space $S := \mathbb{H}^2/G_y$ is a complete Riemann surface.

In order to prove that *S* is topologically equivalent to the Loch Ness monster, we must verify that *S* has only one end and infinite genus. The proof uses the same ideas appearing in [5].

The surface S has only one end. By Lemma 1.1 is enough to prove that for any compact subset K of S, there exists a compact subset K' of S such that $K \subset K'$ and the space $S \setminus K'$ is connected.

Let K be a compact subset of S. Observe that $\tilde{K} := \pi^{-1}(K)$ is a compact subset of $\overline{D(G_y)}$ where $\pi:\overline{D(G_y)}\to S=\overline{D(G_y)}/G_y$ is the quotient map. Thus, \tilde{K} is a compact subset of \mathbb{H}^2 . Therefore, there exist closed intervals I_x and I_y in the real and imaginary axes, respectively, such that, $\pi_x(\tilde{K})\subset I_x$ and $\pi_y(\tilde{K})\subset I_y$, where $\pi_x:\mathbb{H}^2\to\mathbb{R}$ is the standard projection map on the real axis and $\pi_y:\mathbb{H}^2\to(0,+\infty)$ is the projection map on the imaginary axis. By construction of the fundamental domain for G_y we have that $\tilde{K}':=\overline{D(\Gamma)}\cap I_x\times I_y$ is a compact subset of $\overline{D(G_y)}$ such that $\tilde{K}\subset \tilde{K}'$. Thus, $K':=\pi(\tilde{K}')$ is a compact subset of S which contains S. Finally, we can verify that $S\setminus K'$ is connected.

The surface S has infinite genus. For each $n \in \mathbb{Z}$, we define the strip

$$\tilde{S}_n := D(G_{\mathbf{v}}) \cap \{z \in \mathbb{H}^2 : a_n < \operatorname{Re}(z) < e_n\} \subset \mathbb{H}^2.$$

Observe that $\tilde{S}_n/\langle f_n, g_n \rangle$ defines, under the projection map π , a subsurface S_n of S topologically equivalent to a torus minus a disk. By construction, any for two different integers $m \neq n$, the subsurfaces S_n and S_m are disjoint because the strips $\tilde{S}_n \cap \tilde{S}_m = \emptyset$. This shows that S has infinite genus.

The group G_y is of the first kind if and only if $e_n = a_{n+1}$ for all $n \in \mathbb{Z}$, $\lim_{n \to \infty} e_n = \infty$ and $\lim_{n \to -\infty} a_n = -\infty$. Suppose that G_y if of the first kind but that the conclusion is false. In any case it is possible to find an interval $I \subset \mathbb{R}$ not contained in the limit set of G_y which leads to a contradiction. Indeed, if $e_n \neq a_{n+1}$ for some $n \in \mathbb{Z}$ then $I = (e_n, a_{n+1})$ is such interval.

If $\lim_{n\to\infty} e_n = r < \infty$ then $I = (a, \infty)$. Finally, if $\lim_{n\to-\infty} a_n = R > -\infty$, then $I = (-\infty, R)$.

Now, suppose that $e_n=a_{n+1}$ for all $n\in\mathbb{Z}$, $\lim_{n\to\infty}e_n=\infty$ and $\lim_{n\to-\infty}a_n=-\infty$ but that $G_{\mathbf{y}}$ is not of the first kind. Let x be a point in $\partial\mathbb{H}^2$ such that it is not in the limit set $\Lambda(G_{\mathbf{y}})$ of $G_{\mathbf{y}}$. Then there exists an open interval $L\subset\partial\mathbb{H}^2$ with $x\in L$ and $\Lambda(G_{\mathbf{y}})\cap L=\emptyset$. Let C be the half-circle in \mathbb{H}^2 with endpoints the endpoints of L. Given that $\mathbb{H}^2=\bigcup_{g\in G_{\mathbf{y}}}g(\overline{D(G_{\mathbf{y}})})$ then there exists $g\in G_{\mathbf{y}}$ such that $C\cap g(D(G_{\mathbf{y}}))\neq\emptyset$. Therefore, $g^{-1}(C)\cap D(G_{\mathbf{y}})\neq\emptyset$. By hypothesis we have that $g^{-1}(L)\cap\Lambda(G_{\mathbf{y}})\neq\emptyset$. Given that the limit set is $G_{\mathbf{y}}$ -invariant, then we obtain that $L\cap\Lambda(G_{\mathbf{y}})\neq\emptyset$, it is a contradiction because we suppose that $\Lambda(G_{\mathbf{y}})\cap L=\emptyset$.

From the point of view of parabolicity the following questions appears naturally: Which points y of N define Fuchsian groups uniformizing Loch Ness monsters of parabolic type?

ACKNOWLEDGEMENTS

We thank José Hernández Santiago, Rubén Hidalgo and Jesús Muciño for his effort to review our paper and for the provided us with great suggestions and comments.

I. Morales was partially supported by "Proyecto CONACyT Ciencia de Frontera 217392 cerrando brechas y extendiendo puentes en Geometría y Topología".

Camilo Ramírez Maluendas expresses his gratitude to Universidad Nacional de Colombia sede Manizales, and to the research program "Reconstrucción del tejido social en zonas posconflicto en Colombia" SIGP code: 57579 with the project entitled: "Competencias empresariales y de innovación para el desarrollo económico y la inclusión productiva de las regiones afectadas por el conflicto colombiano" SIGP code 58907. Contract number: FP44842-213-2018. He dedicates this work to his beautiful family, Marbella and Emilio, in gratitude for their love and support.

REFERENCES

- [1] Stephen Agard, A geometric proof of Mostow's rigidity theorem for groups of divergence type, Acta Mathematica **151** (1983), 231–252.
- [2] Lars Valerian Ahlfors and Leo Sario, *Riemann surfaces*, Princeton university press, 2015.
- [3] Daniele Alessandrini, Lixin Liu, Athanase Papadopoulos, Weixu Su, and Zongliang Sun, On Fenchel-Nielsen coordinates on Teichmüller spaces of surfaces of infinite type, Ann. Acad. Sci. Fenn. Math. **36** (2011), no. 2, 621-659.

- [4] Venancio Álvarez and José M. Rodríguez, *Structure theorems for Riemann and topological surfaces*, Journal of the London Mathematical Society **69** (2004), no. 1, 153-168.
- [5] John A. Arredondo and Camilo Ramírez Maluendas, *On the infinite Loch Ness monster*, Comment. Math. Univ. Carolin. **58** (2017), no. 4, 465-479.
- [6] ______, On Infinitely generated Fuchsian groups of the Loch Ness monster, the Cantor tree and the Blooming Cantor tree, Complex Manifolds 7 (2020), no. 1, 73–92.
- [7] Kari Astala and Michel Zinsmeister, *Mostow rigidity and Fuchsian groups*, C. R. Acad. Sci. Paris Ser. I Math. **311** (1990), no. 1, 301-306.
- [8] Ara Basmajian, *Hyperbolic Structures for Surfaces of Infinite Type*, Transactions of the American Mathematical Society **336** (1993), no. 1, 421-444.
- [9] Ara Basmajian, Hrant Hakobyan, and Dragomir Šarić, *The type problem for Riemann surfaces via Fenchel-Nielsen parameters*, arXiv preprint arXiv:2011.03166 (2020).
- [10] Ara Basmajian, *Geodesically complete hyperbolic structures*, Mathematical Proceedings of the Cambridge Philosophical Society, 2019, pp. 219-242.
- [11] Benson Farb and Dan Margalit, A primer on mapping class groups (pms-49), Princeton university press, 2011.
- [12] Alan F. Beardon, *The geometry of discrete groups*, Vol. 91, Springer Science & Business Media, 2012.
- [13] Christopher J. Bishop, *Divergence Groups Have the Bowen Property*, Annals of Mathematics **154** (2001), no. 1, 205–217.
- [14] Svetlana Katok, *Fuchsian groups*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1992.
- [15] Béla Kerékjártó, *Vorlesungen über Topologie I*, Mathematics: Theory & Applications, Springer, Berlín, 1923.
- [16] Bernard Maskit, *Kleinian groups*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 287, Springer-Verlag, Berlin, 1988.
- [17] Katsuhiko Matsuzaki and José M. Rodríguez, *Planar Riemann surfaces with uniformly distributed cusps: parabolicity and hyperbolicity*, Mathematische Nachrichten **290** (2017), no. 7, 1097–1112.
- [18] Peter J. Nicholls and and others, *The ergodic theory of discrete groups*, Vol. 143, Cambridge University Press, 1989.
- [19] Makoto Ohtsuka, *Dirichlet problems on Riemann surfaces and conformal mappings*, Nagoya Mathematical Journal **3** (1951), 91–137.
- [20] Ian Richards, *On the classification of noncompact surfaces*, Trans. Amer. Math. Soc. **106** (1963), 259-269.
- [21] Leo Sario and Mitsuru Nakai, *Classification theory of Riemann surfaces*, Vol. 164, Springer Science & Business Media, 2012.
- [22] Ernst Specker, *Die erste Cohomologiegruppe von Überlagerungen und Homotopie-eigenschaften dreidimensionaler Mannigfaltigkeiten*, Commentarii Mathematici Helvetici **23** (1949), no. 1, 303–333.
- [23] M. Tsuji, Potential theory in modern function theory, Chelsea Publishing Co., New York, 1975.
- [24] Pekka Tukia, *Differentiability and rigidity of Möbius groups*, Inventiones mathematicae **82** (1985), no. 3, 557–578.

JOHN A. ARREDONDO,

DEPARTAMENTO DE MATEMÁTICAS, FUNDACIÓN UNIVERSITARIA KONRAD LORENZ, BOGOTÁ, COLOMBIA.

Email address: alexander.arredondo@konradlorenz.edu.co

ISRAEL MORALES,

INSTITUTO DE MATEMÁTICAS UNAM UNIDAD OAXACA, OAXACA DE JUÁREZ, OAXACA. MÉXICO.

Email address: fast.imj@gmail.com

CAMILO RAMÍREZ MALUENDAS,

Departamento de Matemáticas y Estadística, Universidad Nacional de Colombia, Sede Manizales, Manizales 170004, Colombia.

Email address: camramirezma@unal.edu.co