

Infinite convolutions of probability measures on Polish semigroups

Kouji YANO⁽¹⁾⁽²⁾

August 31, 2021

Abstract

This expository paper is intended for a short self-contained introduction to the theory of infinite convolutions of probability measures on Polish semigroups. We give the proofs of the Rees decomposition theorem of completely simple semigroups, the Ellis–Żelazko theorem, the convolution factorization theorem of convolution idempotents, and the convolution factorization theorem of cluster points of infinite convolutions.

Keywords and phrases: Polish semigroup; Rees decomposition; Ellis–Żelazko theorem; convolution idempotent; infinite convolution

AMS 2010 subject classifications: 60B15 (60F05; 60G50)

1 Introduction

As a natural generalization of random walks on an integer lattice, the theory of infinite convolutions of probability measures on topological semigroups has been extensively studied and widely applied to various problems. For this theory, there are celebrated textbooks Rosenblatt [41], Mukherjea–Tserpes [32] and Högnäs–Mukherjea [16], which include a lot of applications of the theory; see also Mukherjea’s lecture notes [29] for applications to random matrices, and Ito–Sera–Yano [17] for applications to the problem of resolution of σ -fields.

The aim of this paper is to help the reader to gain the basic knowledge of this theory conveniently. We mainly follow [16] and we make some modifications on the proofs. For a potential application, we develop the theory for topological semigroups with a Polish topology, while the textbooks [41, 32, 16] deal with those with a locally compact Hausdorff second countable topology.

The goal of this paper is the convolution factorization theorem of cluster points of infinite convolutions, which will be stated as Theorem 4.9. The key to the proof is the convolution factorization theorem of convolution idempotents, which will be stated as

⁽¹⁾Graduate School of Science, Kyoto University.

⁽²⁾The research of Kouji Yano was supported by JSPS KAKENHI grant no.’s JP19H01791 and JP19K21834 and by JSPS Open Partnership Joint Research Projects grant no. JPJSBP120209921. This research was supported by RIMS and by ISM.

Theorem 4.6, and the study of probability measures with convolution invariance, which will be stated as Proposition 4.5. Theorems 4.6 and 4.9 are based on the product decomposition theorem for completely simple semigroups, which will be called the *Rees decomposition* and stated as Theorem 2.8. To show that the algebraic decomposition is compatible with a Polish topology, we need Ellis–Żelazko theorem, which will be stated as Theorem 3.2.

The Ellis theorem [11](1957) asserts that an algebraic group where the product mapping is separately continuous is a topological group, where the topology is locally compact Hausdorff second countable. It was extended by Żelazko [51](1960) for a completely metrizable topology.

The study of infinite convolutions on compact groups dates back to Kawada–Itô [19](1940), which was generalized by Urbanik [49](1957), Kloss [22](1959), and Stromberg [45](1960) and for locally compact groups by Tortrat [48](1964) and Csiszár [7](1966). The convolution invariance Proposition 4.5 is due to Mukherjea [26](1972), which originates from the Choquet–Deny equation [2](1960); for later studies, see [50, 38, 9, 8, 37, 23, 47]. Theorem 4.6 for convolution idempotents is due to Mukherjea–Tserpes [31] (1971); for earlier studies, see Collins [6](1962), Pym [36](1962), Heble–Rosenblatt [14](1963), Schwarz [44](1964), Choy [3](1970), Duncan [10](1970), and Sun–Tserpes [46](1970); see also [12]. Theorem 4.9 for cluster points of infinite convolutions is due to Rosenblatt [39](1960) in the compact case and to Mukherjea [28](1979) in the locally compact case; for studies earlier than [28], see Glicksberg [13](1959), Collins [5](1962), Schwarz [43](1964), Rosenblatt [40](1965), Lin [24](1966), Mukherjea [27](1977), and Mukherjea–Sun [30](1978); for related papers, see [33, 42, 25, 1].

This paper is organized as follows. In Section 2 we review the theory of algebraic semigroups. In Section 3 we study the theory of Polish semigroups, where the Ellis–Żelazko theorem is proved and utilized. Section 4 is devoted to the convolution factorization theorems of convolution idempotents and of cluster points of infinite convolutions.

Acknowledgements

The author would like to express our gratitude to Takao Hirayama for having a hard time to learn the theory together as beginners. He also thanks Yu Ito and Toru Sera for fruitful discussions.

2 Algebraic semigroup

We say that a non-empty set S is a *semigroup* if it is endowed with multiplication

$$S \times S \ni (a, b) \mapsto ab \in S \tag{2.1}$$

which is associative, i.e.,

$$(ab)c = a(bc), \quad a, b, c \in S. \quad (2.2)$$

For two subsets A and B of S , we denote their product by

$$AB = \{ab : a \in A, b \in B\}. \quad (2.3)$$

We write $A^1 = A$ and $A^n = A^{n-1}A$ for $n \geq 2$. We sometimes identify an element $a \in S$ with the singleton $\{a\}$; for instance, $aS = \{a\}S = \{ab : b \in S\}$. An element $e \in S$ is called *identity* if

$$xe = ex = x, \quad x \in S. \quad (2.4)$$

It is obvious that identity is unique if it exists. For a semigroup S with identity e , we say that $y \in S$ is called the *inverse* of $x \in S$ if $xy = yx = e$. It is obvious that the inverse of an element $x \in S$ is unique if it exists. A *group* is a semigroup S with identity such that all elements have their inverses.

2.1 Left and right simplicity

Let S be a semigroup. A non-empty subset I is called a *left ideal* [*right ideal*] (of S) if $SI \subset I$ [$IS \subset I$]. If S contains no proper left ideal [right ideal], then it is called *left simple* [*right simple*]. A non-empty subset I is called a *ideal* if it is both a left and a right ideal, i.e., $SI \cup IS \subset I$. If S contains no proper ideal, then it is called *simple*. Note that both left and right simple implies simple, but the converse is not necessarily true.

Lemma 2.1. *For a subsemigroup S of a semigroup S_0 , the following are equivalent:*

- (i) S is a minimal left ideal of S_0 .
- (ii) $S = S_0a$ for all $a \in S$.

Proof. Suppose S is a minimal left ideal. Since S_0a for $a \in S$ is a left ideal of S_0 contained in S , we have $S = S_0a$ by minimality.

Suppose $S = S_0a$ for all $a \in S$. Let I be a left ideal of S_0 such that $I \subset S$. For any $a \in I$, we have $S = S_0a \subset S_0I \subset I$, which shows that S is a minimal left ideal of S_0 . \square

Lemma 2.2. *For a subsemigroup S of a semigroup S_0 , the following are equivalent:*

- (i) S is a minimal ideal of S_0 .
- (ii) $S = S_0aS_0$ for all $a \in S$.

The proof of Lemma 2.2 is almost the same as that of Lemma 2.1, and so we omit it.

Lemma 2.3. *For a semigroup S , the following are equivalent:*

- (i) S is left simple.
- (ii) For any semigroup S_0 of which S is a left ideal, S is a minimal left ideal of S_0 .
- (iii) S is a minimal left ideal of S itself (if and only if $S = Sa$ for all $a \in S$ by Lemma 2.1).
- (iv) There exists a semigroup S_0 such that S is a minimal left ideal of S_0 .
- (v) For any $a, b \in S$, the equation $xa = b$ has at least one solution $x \in S$.

Proof. [(i) \Rightarrow (ii)] Suppose that S is a left ideal of a semigroup S_0 and let I be a left ideal of S_0 such that $I \subset S$. Then $SI \subset S_0I \subset I$, and so I is a left ideal of S . Since S is left simple, we have $I = S$, which shows that S is a minimal left ideal of S_0 .

[(ii) \Rightarrow (iii) \Rightarrow (iv)] These are obvious.

[(iv) \Rightarrow (i)] Suppose that S is a minimal left ideal of S_0 and let I be a left ideal of S . Since $S_0SI \subset SI \subset I \subset S$, we see that SI is a left ideal of S_0 with $SI \subset S$. Hence $SI = S$ by minimality. Since $I \subset S = SI \subset I$, we have $I = S$, which implies that S is left simple.

[(iii) \Rightarrow (v)] This is obvious by $S \subset Sa$.

[(v) \Rightarrow (iii)] Let $a \in S$. Then we have $S \subset Sa$ by (v). Since S is a semigroup, we have $Sa \subset S$. Hence we have $S = Sa$. \square

For the simplicity, we have the following.

Lemma 2.4. *For a semigroup S , the following are equivalent:*

- (i) S is simple.
- (ii) For any semigroup S_0 of which S is an ideal, S is a minimal ideal of S_0 .
- (iii) S is a minimal ideal of S itself (if and only if $S = SaS$ for all $a \in S$ by Lemma 2.2).
- (iv) There exists a semigroup S_0 such that S is a minimal ideal of S_0 .
- (v) For any $a, b \in S$, the equation $xay = b$ has at least one solution $(x, y) \in S \times S$.

The proof of Lemma 2.4 is almost the same as that of Lemma 2.3, and so we omit it.

Proposition 2.5. *A semigroup S which is both left and right simple is a group.*

Proof. Let $a \in S$. By Lemma 2.3, we have $ea = a$ for some $e \in S$. For any $x \in S$, we have $x = ay$ for some $y \in S$, and so we have $ex = eay = ay = x$. Similarly, there exists $e' \in S$ such that $xe' = x$ for all $x \in S$. Then we obtain $e' = ee' = e$, and thus e is identity of S .

Let $x \in S$. By Lemma 2.3, we have $xy = e$ and $y'x = e$ for some $y, y' \in S$. Since $y' = y'e = y'xy = ey = y$, we see that y is the inverse of x . \square

2.2 Left and right groups

Let S be a semigroup. An element $e \in S$ is called an *idempotent* if $e^2 = e$. We denote the set of all idempotents of S by

$$E(S) = \{e \in S : e^2 = e\}. \quad (2.5)$$

Note that, if e is an idempotent, then any element of Se is invariant under right multiplication by e , i.e., $x \in Se$ implies $xe = x$. A semigroup S is called *left group* [*right group*] if S is left simple [*right simple*] and contains at least one idempotent.

A semigroup S is called *left cancellative* [*right cancellative*] if, for any $a, x, y \in S$ with $ax = ay$ [$xa = ya$], we have $x = y$. An element $e \in S$ is called a *left identity* [*right identity*] if $ex = x$ [$xe = x$] for all $x \in S$.

Lemma 2.6. *Let S be a semigroup. If S is either right cancellative or left simple, then any idempotent of S is a right identity.*

Proof. Suppose S is right cancellative and let $e \in E(S)$. Then $xee = xe$ implies $xe = x$.

Suppose S is left simple and let $e \in E(S)$. By Lemma 2.3, we have $S = Se$, which yields that $xe = x$ for all $x \in S$. \square

Proposition 2.7. *For a semigroup S , the following are equivalent:*

- (i) S is a left group.
- (ii) S is left simple and right cancellative.
- (iii) For any $a, b \in S$, the equation $xa = b$ has a unique solution $x \in S$.

Proof. [(i) \Rightarrow (ii)] Let $e \in E(S)$ be fixed. By Lemma 2.6, we see that e is a right identity.

Suppose $xa = ya$. By Lemma 2.3, we have $ba = e$ for some $b \in S$. We then have $abab = aeb = ab$, so that $ab \in E(S)$ and ab is a right identity. We then obtain $x = xab = yab = y$.

[(ii) \Rightarrow (iii)] Existence follows from left simplicity and Lemma 2.3. Uniqueness follows from right cancellativity.

[(iii) \Rightarrow (i)] By (iii), we have $S = Sa$ for all $a \in S$, which shows by Lemma 2.3 that S is left simple.

Let $a \in S$ and take $e \in S$ such that $ea = a$ by (iii). Then we have $e^2a = ea = a$, which leads to $e^2 = e$ by right cancellativity. \square

2.3 Rees decomposition

Let S be a semigroup. An idempotent $e \in E(S)$ is called *primitive* if

$$ex = xe = x \in E(S) \text{ implies } x = e. \quad (2.6)$$

We say that S is *completely simple* if S is simple and contains a primitive idempotent.

Theorem 2.8 (Rees decomposition). *Let S be a completely simple semigroup and let e be a primitive idempotent of S . Set*

$$L := E(Se), \quad G := eSe, \quad R := E(eS). \quad (2.7)$$

Then the following assertions hold:

- (i) $LG = Se$ is a left group and $GR = eS$ is a right group.
- (ii) $RL \subset G$ and $eL = Re = \{e\}$.
- (iii) $G = Se \cap eS$ is a group where e is its identity.
- (iv) $S = LGR$ (This factorization will be called the Rees decomposition of S at e , and G will be called the group factor at e).
- (v) The product mapping

$$\psi : L \times G \times R \ni (x, g, y) \mapsto (xgy) \in LGR \quad (2.8)$$

is bijective with its inverse given as

$$\psi^{-1} : LGR \ni z \mapsto (ze(eze)^{-1}, eze, (eze)^{-1}ez) \in L \times G \times R. \quad (2.9)$$

Proof. (i) It is obvious that Se is a left ideal of S . Let I be a left ideal of S such that $I \subset Se$. Let $a \in I$. Note that $ae = a$ since $a \in Se$. By simplicity of S and Lemma 2.4, we have $uav = e$ for some $u, v \in S$. Set $r = eu$ and $s = eve$. We then have

$$ras = eu(ae)ve = euave = e, \quad er = r, \quad es = se = s. \quad (2.10)$$

If we set $t = sra$, then $et = te = t$ and

$$t^2 = s(ras)ra = sera = sra = t, \quad (2.11)$$

which yields $t = e$ by primitivity. Since $e = t = sra \in srI \subset I$, we have $Se \subset SI \subset I$, which shows $I = Se$ and that Se is a minimal left ideal of S . By Lemma 2.3, we see that Se is left simple. Since Se contains an idempotent e , we see that Se is a left group. By a similar argument we see that eS is a right group.

Let us show $LG = Se$. It is obvious that $LG \subset Se$. Let $a \in Se$. Set $g := ea \in eSe = G$ and set $b = ag^{-1} \in Se$. Since $g^{-1} = g^{-1}e$, we have

$$b^2 = ag^{-1}ag^{-1} = ag^{-1}(ea)g^{-1} = ag^{-1} = b. \quad (2.12)$$

Hence we have $b \in E(Se) = L$ and $a = ae = ag^{-1}g = bg \in LG$. We now have $LG = Se$. We also have $GR = eS$ similarly.

(ii) $RL \subset (eS)(Se) \subset eSe = G$.

Let $x \in L = E(Se)$. Since $(ex)^2 = e(ex)x = exx = ex$ and $e(ex) = (ex)e = ex$, we have $ex = e$ by primitivity. We thus see that $eL = \{e\}$. We have $Re = \{e\}$ similarly.

(iii) It is obvious that $G = eSe = eS \cap Se$, since $x \in eS \cap Se$ implies $x = ex = xe = exe$. It is also obvious that e is identity of G . Let $g \in G$. Since $G \subset eSe$, we have $g = ea$ for some $a \in Se$. By the left simplicity of Se and by Lemma 2.3, we have $ba = e$ for some $b \in Se$. Since $(ab)^2 = a(ba)b = aeb = ab$, we see by Lemma 2.6 that ab is right identity. Hence $ab = abe = e$, which shows that b is the inverse of a .

(iv) $LGR = LGGR = SeeS = SeS = S$ by Lemma 2.4.

(v) Let $z = xgy$ with $(x, g, y) \in L \times G \times R$. Since $x = xx = xex$ and since $exgye \in eSe = G$, we have

$$x = xe = x(exgye)(exgye)^{-1} = ze(eze)^{-1}. \quad (2.13)$$

We have $y = (eze)^{-1}ez$ similarly. Since $ex = ye = e$ by (ii), we obtain

$$g = ege = (ex)g(ye) = eze. \quad (2.14)$$

The proof is now complete. \square

Corollary 2.9. *Under the same assumptions and notation as Theorem 2.8, it holds that $\{Sy = LGy : y \in R\}$ is the family of all minimal left ideals of S .*

Proof. Any minimal left ideal of S is of the form Sz for some $z \in S$. We represent $z = xgy$ and then we obtain $Sz = LG(Rx)gy = LGy$, since $RL \subset G$.

Conversely, for any $z \in LGy$, we have $z = xgy$ for some $(x, g) \in L \times G$, so that we have $LGyz = LG(yx)gy = LGy$, which shows by Lemma 2.1 that LGy is a minimal left ideal. \square

Corollary 2.10. *Under the same assumptions and notation as Theorem 2.8, the following assertions hold:*

(i) *For $z = xgy$ with $(x, g, y) \in L \times G \times R$, z is idempotent if and only if $g = (yx)^{-1}$.*

(ii) All idempotents of S are primitive.

(iii) Let e' be another idempotent of S and represent it as $e' = a(ba)^{-1}b$ for $(a, b) \in L \times R$. Let $S = L'G'R'$ denote the Rees decomposition of S at e' . Then

$$L'G' = LGb, \quad G' = aGb, \quad G'R' = aGR. \quad (2.15)$$

Proof. (i) Suppose $z^2 = z$. Then $xgyxgy = xgy$. Since $eL = Re = \{e\}$, we have $gyxg = g$, which shows $g = (yx)^{-1}$.

Conversely, suppose $g = (yx)^{-1}$. Then $z^2 = x(gyxg)y = xgy = z$.

(ii) Let $e_1, e_2 \in S$ be two idempotents of S and represent them as $e_i = a_i(b_i a_i)^{-1}b_i$ for $(a_i, b_i) \in L \times R$, $i = 1, 2$. Suppose $e_1 e_2 = e_2 e_1 = e_2$. Then $a_1((b_1 a_1)^{-1}(b_1 a_2)(b_2 a_2)^{-1})b_2 = a_2((b_2 a_2)^{-1}(b_2 a_1)(b_1 a_1)^{-1})b_1 = a_2(b_2 a_2)^{-1}b_2$, which shows $a_1 = a_2$ and $b_1 = b_2$ by the injectivity of the product mapping ψ . Hence we have $e_1 = e_2$, which shows that e_1 is a primitive idempotent.

(iii) We have $L'G' = Se' = LG(Ra)(ba)^{-1}b = LGb$ and $G'R' = aGR$ similarly. We also have $G' = e'Se' = a(ba)^{-1}(bL)G(Ra)(ba)^{-1}b = aGb$. \square

Corollary 2.11. *A left group S is completely simple. The Rees decomposition of S at $e \in E(S)$ is given as $S = LG$ with $R = \{e\}$.*

Proof. Suppose $ex = xe = x \in E(S)$. By Lemma 2.3, we have $yx = e$ for some $y \in S$. Hence $x = ex = yxx = yx = e$, which shows that e is an primitive idempotent. Hence S is completely simple. Let $S = LGR$ denote the Rees decomposition of S at e . Since $S = Se$ by Lemma 2.3 and since $Re = \{e\}$, we obtain $S = Se = LGR = LG$. \square

For later use we prove the following proposition.

Proposition 2.12. *Suppose that a semigroup S contains a minimal left ideal A and a minimal right ideal B as well. Then BA is a group and its identity is a primitive idempotent of S . If, in addition, S is simple, then S is completely simple.*

Proof. Since $(BA)(BA) = (BAB)A \subset BA$, we see that BA is a subsemigroup of S . To prove right simplicity of BA , let I be a right ideal of BA . Since IB is a right ideal of S and $IB \subset BAB \subset B$, we see that $IB = B$ by minimality. Hence $BA = IBA \subset I$, which shows right simplicity of BA . By a similar argument we obtain left simplicity of BA . We thus conclude by Proposition 2.5 that BA is a group.

Let e be the identity of BA and suppose $ex = xe = x \in E(S)$. Then $x = xx = exxe \in (BAS)(SBA) \subset BA$. Since BA is a group and since $x^2 = x$, we have $x = xx^{-1} = e$, which shows that e is a primitive idempotent of S . \square

2.4 Kernel

A minimal ideal of a semigroup S will be called a *kernel* of S .

Theorem 2.13. *Let S be a semigroup. Then the following assertions hold:*

- (i) *If S contains a minimal left ideal, then S contains a unique kernel K , and $SzS = K$ for all $z \in K$.*
- (ii) *If S contains a minimal left ideal and a minimal right ideal as well, then the unique kernel of S is completely simple.*
- (iii) *If S contains a completely simple kernel K , then it is the unique kernel of S . Let $K = LGR$ denote the Rees decomposition at e . Then $Sz = Kz = LGz$ for all $z \in K$.*

Proof. (i) Let \mathcal{A} denote the family of all minimal left ideals of S and suppose \mathcal{A} is not empty. We shall prove that $K := \bigcup \mathcal{A}$ is a unique kernel of S .

Let $z \in K$ and take $A \in \mathcal{A}$ such that $z \in A$. Then $Sz = A$ by Lemma 2.1. For $x \in S$, we see that $Ax \in \mathcal{A}$; in fact, for any left ideal I of S such that $I \subset Ax$, we see that $J = \{a \in A : ax \in I\} \subset A$ is a left ideal of S , so that $J = A$ by minimality and thus $I = Ax$. Hence $SzS = AS = \bigcup_{x \in S} Ax \subset \bigcup \mathcal{A} = K$, which shows by Lemma 2.2 that K is a kernel of S .

Let K' be another kernel of S . Since $K \cap K'$ contains KK' which is not empty, we see that $K \cap K'$ is an ideal contained both in K and in K' . Thus $K \cap K' = K = K'$ by minimality.

(ii) By (i) and Lemma 2.2, we see that the unique kernel K of S is both a minimal left ideal of K and a minimal right ideal of K . By Proposition 2.12, we see that K is completely simple.

(iii) Suppose K is a completely simple kernel of S with a primitive idempotent e . By Theorem 2.8, K contains a left group Ke . By Lemma 2.3, we see that Ke is a minimal left ideal of S . Hence by (i) the kernel of S is unique.

For $z \in K$, we represent $z = xgy \in LGR$. Then by (i) Sz is a minimal left ideal of K containing y . By Corollary 2.9, we see that $Sz = LGy = LGz = Kz$. \square

3 Topological semigroup

A semigroup S is called *topological* if S is endowed with a topology such that the product mapping $S \times S \ni (x, y) \mapsto xy \in S$ is jointly continuous. A semigroup S is called *Polish* if S is a topological semigroup with respect to a Polish topology, i.e. a separable and completely metrizable topology.

It is well-known (see, e.g. [20, Theorem 1.5.3]) that locally compact Polish is equivalent to locally compact Hausdorff with a countable base. It is elementary that compact Polish is equivalent to compact metrizable.

For $a \in S$ and $A \subset S$, we write

$$a^{-1}A = \{x \in S : ax \in A\}, \quad Aa^{-1} = \{x \in S : xa \in A\}. \quad (3.1)$$

If S contains identity e and $a \in S$ has its inverse $a^{-1} \in S$, then $(a^{-1})A = a^{-1}A$; in fact,

$$(a^{-1})A = \{a^{-1}x \in S : x \in A\} = \{y \in S : ay \in A\} = a^{-1}A. \quad (3.2)$$

Lemma 3.1. *Let S be a Polish semigroup. Then the following assertions hold:*

(i) *For $a \in S$ and for a closed [open, Borel] subset A , both $a^{-1}A$ and Aa^{-1} are also closed [open, Borel].*

(ii) *If A is a subsemigroup of S , then so is its closure \overline{A} .*

(iii) *Let A be a closed subsemigroup of S . Then $E(A)$, eA , Ae and eAe are closed for all $e \in E(A)$.*

(iv) *For two compact subsets K and K' , the product KK' is also compact.*

Proof. (i) If we write $\psi_a : S \rightarrow S$ for the translation $\psi_a(x) = ax$, then $a^{-1}A = \psi_a^{-1}(A)$. Since ψ_a is continuous, we obtain the desired results.

(ii) Let $a, b \in \overline{A}$ and take $\{a_n\}, \{b_n\} \subset A$ such that $a_n \rightarrow a$ and $b_n \rightarrow b$. Then we have $ab = \lim a_n b_n \in \overline{A}$.

(iii) Let $\{e_n\} \subset E(A)$ such that $e_n \rightarrow e \in S$. Since A is closed, we have $e \in A$. Since $e_n^2 = e_n$ for all n , we have $e^2 = e$, which shows $e \in E(A)$.

Let $\{x_n\} \subset eA$ such that $x_n \rightarrow x \in S$. Since $eA \subset A$ and since A is closed, we have $x \in A$. Then $ex = \lim ex_n = \lim x_n = x$, which shows $x = ex \in eA$.

(iv) Let $\psi : S \times S \rightarrow S$ denote the jointly continuous product mapping: $\psi(x, y) = xy$. Since $KK' = \psi(K \times K')$ and $K \times K'$ is compact, we see that KK' is compact. \square

3.1 Topological group

A group S is called *topological* if G is a topological semigroup and the inverse mapping $G \ni g \mapsto g^{-1} \in G$ is continuous.

Theorem 3.2 (Ellis [11] and Żelazko [51]). *If a group G is a topological semigroup with respect to a completely metrizable topology, then it is a topological group.*

Proof. We borrow the proof from Pfister [35]. Let e denote the identity of G and let d be a complete metric of G .

Let U_0 be a open neighborhood of e . By the joint continuity of the product mapping, we can construct a sequence $\{U_n\}_{n=1}^\infty$ of open balls of e such that the radius of U_n decreases to 0 and $\overline{U_n} \overline{U_n} \subset U_{n-1}$ for $n = 1, 2, \dots$, where $\overline{U_n}$ stands for the closure of U_n .

Let $\{x_n\}_{n=1}^\infty$ be a subsequence of an arbitrary sequence of G which converges to e . It then suffices to construct a subsequence $\{n(k)\}_{k=1}^\infty$ of $\{1, 2, \dots\}$ such that $x_{n(k)}^{-1} \rightarrow e$. We write $y_k := x_{n(1)} \cdots x_{n(k)}$.

Set $n(0) = 0$ and $y_0 = x_0 = e$. If we have $n(0), n(1), \dots, n(k-1)$, then we can take $n(k) > n(k-1)$ such that $x_{n(k)} \in U_k$ and $d(y_k, y_{k-1}) < 2^{-k}$, since $y_{k-1}x_n \rightarrow y_{k-1}$ as $n \rightarrow \infty$. By completeness of d , we see that y_k converges to a limit $y \in G$. Let n be fixed for a while. Since yU_{n+1} is a neighborhood of y , we see that $y_{k-1} \in yU_{n+1}$ for large k . For $j > k$, we have $U_{j-1}U_j \subset U_jU_j \subset U_{j-1}$, and hence

$$y_k^{-1}y_j = x_{n(k+1)} \cdots x_{n(j-1)}x_{n(j)} \in U_{k+1} \cdots U_j \subset U_k, \quad (3.3)$$

which implies $y_k^{-1}y \in \overline{U_k} \subset U_{k-1}$. We now obtain

$$x_{n(k)}^{-1} = (y_{k-1}^{-1}y_k)^{-1} = y_k^{-1}y_{k-1} \in y_k^{-1}yU_{n+1} \subset U_{k-1}U_{n+1} \subset U_{n+1}U_{n+1} \subset U_n \quad (3.4)$$

for large k . Thus we obtain $x_{n(k)}^{-1} \rightarrow e$. \square

Corollary 3.3. *Suppose that a Polish semigroup S contains a completely simple kernel K . Let $K = LGR$ denote the Rees decomposition of K at $e \in E(K)$. Then it holds that L, G, R and K are closed subsets, and that the product mapping*

$$\psi : L \times G \times R \ni (x, g, y) \mapsto xgy \in LGR \quad (3.5)$$

is homeomorphic.

Proof. By Corollary 2.13, we have $Ke = Se$, $eK = eS$ and $eKe = eSe$. By Lemma 3.1, we see that $L = E(Ke)$, $G = eKe$ and $R = E(eK)$ are all closed. By Theorem 3.2, we see that G is a Polish group. We now see that the inverse

$$\psi^{-1} : LGR \ni z \mapsto (ze(eze)^{-1}, eze, (eze)^{-1}ez) \in L \times G \times R \quad (3.6)$$

is continuous. Consequently, we see that K is closed. \square

3.2 Compact semigroup

Theorem 3.4. *A compact Polish semigroup S contains a compact completely simple kernel.*

Proof. Let \mathcal{I} denote the family of all closed left ideals of S . The family \mathcal{I} contains S and is endowed with a partial order by the usual inclusion. For any linearly ordered subfamily \mathcal{J} of \mathcal{I} has a lower bound in \mathcal{I} ; in fact, the intersection $\bigcap \mathcal{J}$ is not empty by compactness of S and is a closed left ideal of S such that $\bigcap \mathcal{J} \subset J$ for all $J \in \mathcal{J}$. Hence, by Zorn's lemma, we see that \mathcal{I} contains a minimal element, say A .

Let us prove that A is a minimal left ideal of S . Let I be a left ideal of S such that $I \subset A$. For $a \in I$, we have $Sa \in \mathcal{I}$ and $Sa \subset SI \subset I \subset A$, which yields $Sa = I = A$ by the minimality of A in \mathcal{I} . This shows that A is a minimal left ideal of S .

Similarly we see that S contains a minimal right ideal. By Theorem 2.13, we see that S contains a completely simple kernel K . By Corollary 3.3, we see that K is a closed subset of S , and hence K is compact. \square

Proposition 3.5. *Let S be a Polish semigroup and let $a \in S$. Suppose that any subsequence of $\{a^n\}_{n=1}^\infty$ has a convergent further subsequence. Then the set C of all cluster points of $\{a^n\}_{n=1}^\infty$ is a compact abelian group. If we denote the identity of C by e , then $C = \{e, ae, a^2e, \dots\}$.*

Proof. Let C denote the set of all cluster points of $\{a^n\}_{n=1}^\infty$. By the assumption, we see that C is a compact abelian semigroup. By Theorem 3.4, we see that C contains a compact completely simple kernel K . Since the Rees decomposition of K is $LGR = GRL = G$ by commutativity, we see that K is a compact abelian group. Let e denote the identity of K . Then, for any $x \in C$, we can find a subsequence $\{n(k)\}$ of $\{1, 2, \dots\}$ such that $x = e \lim_{k \rightarrow \infty} a^{n(k)} \in \overline{eC} \subset \overline{KC} \subset K \subset C$, which shows $K = eC = C$. It is now easy to see that $C = \{e, ae, a^2e, \dots\}$. \square

Remark 3.6. In the settings of Proposition 3.5, suppose that the sequence $\{a^n\}_{n=1}^\infty$ has multiple points. Let p and q be the smallest positive integers such that $a^{q+p} = a^q$. Then we have $\{a^n : n = 1, 2, \dots\} = \{a, a^2, \dots, a^{q+p-1}\}$ and

$$K = \{a^q, a^{q+1}, \dots, a^{q+p-1}\} = \{e, ae, \dots, a^{p-1}e\} \quad (3.7)$$

with $e = a^{rp}$, where r is the unique integer such that $q \leq rp \leq q + p - 1$.

4 Convolutions of probability measures on Polish semigroups

4.1 Convolutions

Let S be a Polish semigroup. Let $\mathcal{B}(S)$ denote the family of all Borel sets of S and $\mathcal{P}(S)$ the family of all probability measures on $(S, \mathcal{B}(S))$.

For $\mu, \nu \in \mathcal{P}(S)$, we define the *convolution* $\mu * \nu \in \mathcal{P}(S)$ of μ and ν by

$$\mu * \nu(B) = \iint 1_B(xy) \mu(dx) \nu(dy), \quad B \in \mathcal{B}(S). \quad (4.1)$$

Since $1_B(xy) = 1_{By^{-1}}(x) = 1_{x^{-1}B}(y)$, we have

$$\mu * \nu(B) = \int \mu(By^{-1})\nu(dy) = \int \nu(x^{-1}B)\mu(dx), \quad B \in \mathcal{B}(S). \quad (4.2)$$

For $a \in S$, we write δ_a for the Dirac mass at a : $\delta_a(B) = 1_B(a)$. It is obvious that

$$\mu * \delta_x(B) = \mu(Bx^{-1}), \quad \delta_x * \mu(B) = \mu(x^{-1}B), \quad B \in \mathcal{B}(S), \quad (4.3)$$

which will be called *translations* of μ .

For $\mu \in \mathcal{P}(S)$, we denote its *topological support* by

$$\mathcal{S}(\mu) = \{x \in S : \mu(U) > 0 \text{ for all open neighborhood } U \text{ of } x\}. \quad (4.4)$$

It is obvious that $\mathcal{S}(\mu)$ is closed and $\mu(\mathcal{S}(\mu)^c) = 0$.

Lemma 4.1. *For $\mu, \nu \in \mathcal{P}(S)$, it holds that*

$$\mathcal{S}(\mu * \nu) = \overline{\mathcal{S}(\mu)\mathcal{S}(\nu)}. \quad (4.5)$$

Proof. Let $a \in \mathcal{S}(\mu)$ and $b \in \mathcal{S}(\nu)$. For any open neighborhood U of ab , the joint continuity of the product mapping allows us to take open neighborhoods U_1 of a and U_2 of b such that $U_1U_2 \subset U$, so that

$$\mu * \nu(U) \geq \iint 1_{U_1U_2}(xy)\mu(dx)\nu(dy) \geq \mu(U_1)\nu(U_2) > 0, \quad (4.6)$$

which yields $ab \in \mathcal{S}(\mu * \nu)$ and hence $\overline{\mathcal{S}(\mu)\mathcal{S}(\nu)} \subset \mathcal{S}(\mu * \nu)$.

Let $a \in \overline{\mathcal{S}(\mu)\mathcal{S}(\nu)}^c$. Then we can take an open neighborhood U of a such that $U \subset \{\mathcal{S}(\mu)\mathcal{S}(\nu)\}^c$, so that

$$\mu * \nu(U) \leq \iint 1_{\{\mathcal{S}(\mu)\mathcal{S}(\nu)\}^c}(xy)\mu(dx)\nu(dy) \leq \mu(\mathcal{S}(\mu)^c) + \nu(\mathcal{S}(\nu)^c) = 0, \quad (4.7)$$

which shows $a \in \mathcal{S}(\mu * \nu)^c$ and hence $\mathcal{S}(\mu * \nu) \subset \overline{\mathcal{S}(\mu)\mathcal{S}(\nu)}$. \square

Proposition 4.2. *Let S be a completely simple Polish semigroup. Let $S = LGR$ denote the Rees decomposition at $e \in E(S)$. For the inverse of the product mapping $\psi : L \times G \times R \rightarrow LGR$, we denote*

$$(z^L, z^G, z^R) := \psi^{-1}(z) = (ze(eze)^{-1}, eze, (eze)^{-1}ez) \in L \times G \times R, \quad z \in LGR. \quad (4.8)$$

For $\mu \in \mathcal{P}(S)$, we define

$$\mu^L(B) = \mu(z : z^L \in B), \quad \mu^G(B) = \mu(z : z^G \in B), \quad \mu^R(B) = \mu(z : z^R \in B) \quad (4.9)$$

for $B \in \mathcal{B}(S)$. Then, for $\mu, \nu \in \mathcal{P}(S)$, it holds that

$$(\mu * \nu)^L = \mu^L, \quad (\mu * \nu)^R = \nu^R. \quad (4.10)$$

Proof. This is obvious by noting that $(z_1 z_2)^L = z_1^L$ and $(z_1 z_2)^R = z_2^R$. \square

We equip $\mathcal{P}(S)$ with the topology of weak convergence: $\mu_n \rightarrow \mu$ if and only if $\int f d\mu_n \rightarrow \int f d\mu$ for all $f \in C_b(S)$, the class of all bounded continuous functions on S . It is well-known (see, e.g. [34, Theorems 6.2 and 6.5 of Chapter 2]) that $\mathcal{P}(S)$ is a Polish space.

Proposition 4.3. *Let S be a Polish semigroup. Then the convolution mapping $\mathcal{P}(S) \times \mathcal{P}(S) \ni (\mu, \nu) \mapsto \mu * \nu \in \mathcal{P}(S)$ is jointly continuous. Consequently, $\mathcal{P}(S)$ is a Polish semigroup.*

Proof. Note that, if we take independent random variables X and Y taking values in S such that $X \stackrel{d}{=} \mu$ and $Y \stackrel{d}{=} \nu$, then $\mu * \nu$ coincides with the law of the product XY . The desired result now follows from the Skorokhod coupling theorem (see, e.g. [18, Theorem 4.30]), which asserts that $\mu_n \rightarrow \mu$ implies that we can take random variables $\{X_n\}, X$ taking values in S such that $X_n \stackrel{d}{=} \mu_n$, $X \stackrel{d}{=} \mu$ and $X_n \rightarrow X$ a.s. \square

4.2 Translation invariance

Let S be a Polish semigroup. A probability measure $\mu \in \mathcal{P}(S)$ is called ℓ^* -invariant [r^* -invariant] if $\delta_x * \mu = \mu$ [$\mu * \delta_x = \mu$] for all $x \in S$.

Theorem 4.4. *Let S be a Polish semigroup and let $\mu \in \mathcal{P}(S)$. Suppose that μ is both ℓ^* -invariant and r^* -invariant. Then $\mathcal{S}(\mu)$ is a compact Polish group, and μ coincides with the normalized unimodular Haar measure on $\mathcal{S}(\mu)$ (see e.g. [4, Chapter 9] for the Haar measure).*

Proof. Note that

$$\mathcal{S}(\mu) = \mathcal{S}(\delta_x * \mu) = \overline{x\mathcal{S}(\mu)}, \quad x \in S, \quad (4.11)$$

which implies that $\mathcal{S}(\mu)$ is a left ideal of S . Similarly $\mathcal{S}(\mu)$ is a right ideal of S , and hence $\mathcal{S}(\mu)$ is an ideal of S .

Let us prove that, for any $x \in \mathcal{S}(\mu)$, the subsemigroup xS is left-cancellative. Let $y, a, b \in S$ be such that $(xy)(xa) = (xy)(xb)$. Since $\mathcal{S}(\mu) = \mathcal{S}(\mu * \delta_{xyx}) = \overline{\mathcal{S}(\mu)xyx}$, we can take $\{z_n\} \subset \mathcal{S}(\mu)$ such that $z_n xyx \rightarrow x$, and hence

$$xa = \lim z_n xyxa = \lim z_n xyxb = xb, \quad (4.12)$$

which shows that xS is left-cancellative. Similarly Sx is right-cancellative.

Let $a, b \in \mathcal{S}(\mu)$ be fixed. We shall prove that the subsemigroup $D := a\mathcal{S}(\mu)b$ contains an idempotent. Note that

$$\mu(D) = (\delta_a * \mu * \delta_b)(D) = \mu(a^{-1}Db^{-1}) \geq \mu(\mathcal{S}(\mu)) = \mu(S) = 1, \quad (4.13)$$

which shows $\mu(D) = 1$. For $x \in D$, we have

$$\mu(D) \leq \mu(x^{-1}(xD)) = (\mu * \delta_x)(xD) = \mu(xD) \leq \mu(D), \quad (4.14)$$

which shows $\mu(xD) = \mu(D) = 1$. We define two mappings $\theta, \beta : S \times S \rightarrow S \times S$ by

$$\theta(x, y) = (x, xy), \quad \beta(x, y) = (y, x). \quad (4.15)$$

Since $(x, y) \in \theta(D \times D)$ if and only if $x \in D$ and $y \in xD$, we have

$$(\mu \otimes \mu)(\beta \circ \theta(D \times D)) = (\mu \otimes \mu)(\theta(D \times D)) = \int_D \mu(xD)\mu(dx) = \mu(D)^2 = 1. \quad (4.16)$$

This shows that $\beta \circ \theta(D \times D) \cap \theta(D \times D)$ is not empty, so that $(vw, v) = (x, xy)$ for some $v, w, x, y \in D$. We now have $x(yw) = vwyw = x(yw)^2$, which implies $yw = (yw)^2$ by left-cancellativity of D .

Let $e := yw \in E(D) = E(a\mathcal{S}(\mu)b)$. By the left- and right-cancellativity of $a\mathcal{S}(\mu)b$ and by Lemma 2.6, we see that e is identity of $a\mathcal{S}(\mu)b$. By Lemma 3.1, we see that

$$\mathcal{S}(\mu) = \overline{ea\mathcal{S}(\mu)b} = e \overline{(a\mathcal{S}(\mu)b)} = e\mathcal{S}(\mu) \subset a\mathcal{S}(\mu)b\mathcal{S}(\mu) \subset a\mathcal{S}(\mu) \subset \mathcal{S}(\mu), \quad (4.17)$$

which shows $a\mathcal{S}(\mu) = \mathcal{S}(\mu)$. Similarly we have $\mathcal{S}(\mu)b = \mathcal{S}(\mu)$. By Lemma 2.3, Proposition 2.5 and Theorem 3.2, we see that $\mathcal{S}(\mu)$ is a Polish group.

By the ℓ^* -invariance, we have $\mu * \mu = \mu$. We now apply [34, Theorem 3.1 of Chapter 3] to obtain the desired result. \square

4.3 Convolution invariance

Proposition 4.5 (Mukherjea [26]). *Let S be a Polish semigroup and let $\mu, \nu \in \mathcal{P}(S)$. Suppose*

$$\nu = \mu * \nu = \nu * \mu. \quad (4.18)$$

Then, for any $x \in \mathcal{S}(\mu)$ and any $a \in \mathcal{S}(\nu)$, it holds that

$$\nu * \delta_{xa} = \nu * \delta_a, \quad \delta_{ax} * \nu = \delta_a * \nu. \quad (4.19)$$

Proof. Let $a \in \mathcal{S}(\nu)$, $f \in C_b(S)$ and $\varepsilon > 0$ be fixed for a while, and set

$$g(x) = \max \left\{ \int f d(\nu * \delta_x) - \int f d(\nu * \delta_a) - \varepsilon, 0 \right\}, \quad x \in S. \quad (4.20)$$

It is obvious that $g \in C_b(S)$, g is non-negative and $g(a) = 0$. By $\nu = \nu * \mu$, we have

$$\int f d(\nu * \delta_x) - \int f d(\nu * \delta_a) - \varepsilon \quad (4.21)$$

$$= \int \left\{ \int f d(\nu * \delta_{yx}) - \int f d(\nu * \delta_a) - \varepsilon \right\} \mu(dy) \leq \int g(yx)\mu(dy), \quad (4.22)$$

so that we have

$$g(x) \leq \int g(yx)\mu(dy), \quad x \in S. \quad (4.23)$$

In addition, by $\nu = \mu * \nu$, we have

$$\int \left\{ g(x) - \int g(yx)\mu(dy) \right\} \nu(dx) = \int g d\nu - \int g d(\mu * \nu) = 0, \quad (4.24)$$

which shows that the equality in (4.23) holds for ν -a.e. $x \in S$. Since g is continuous, we see that the equality in (4.23) holds for all $x \in \mathcal{S}(\nu)$. Since $a \in \mathcal{S}(\nu)$ and $g(a) = 0$, we see, again by continuity of g , that

$$g(ya) = 0, \quad y \in \mathcal{S}(\mu). \quad (4.25)$$

Since $\varepsilon > 0$ is arbitrary, we obtain

$$\int f d(\nu * \delta_{ya}) \leq \int f d(\nu * \delta_a), \quad a \in \mathcal{S}(\nu), \quad y \in \mathcal{S}(\mu). \quad (4.26)$$

Since $\nu = \nu * \mu$, we have $\int \{ \int f d(\nu * \delta_{ya}) - \int f d(\nu * \delta_a) \} \mu(dy) = 0$, which implies

$$\int f d(\nu * \delta_{ya}) = \int f d(\nu * \delta_a), \quad a \in \mathcal{S}(\nu), \quad y \in \mathcal{S}(\mu), \quad f \in C_b(S). \quad (4.27)$$

Since $f \in C_b(S)$ is arbitrary, we obtain $\nu * \delta_{ya} = \nu * \delta_a$ for all $a \in \mathcal{S}(\nu)$ and $y \in \mathcal{S}(\mu)$. We obtain $\delta_{ay} * \nu = \delta_a * \nu$ similarly. \square

4.4 Convolution idempotent

We denote the n -fold convolution by μ^n , i.e. $\mu^1 = \mu$ and $\mu^n = \mu^{n-1} * \mu$ for $n = 2, 3, \dots$

Theorem 4.6 (Mukherjea–Tserpes [31]). *Let S be a Polish semigroup and let $\mu \in \mathcal{P}(S)$. Suppose that $\mu^2 = \mu$. Then $\mathcal{S}(\mu)$ is completely simple and its group factor is compact. Let $\mathcal{S}(\mu) = LGR$ denote the Rees decomposition at $e \in E(\mathcal{S}(\mu))$. Then μ admits the convolution factorization*

$$\mu = \mu^L * \omega_G * \mu^R, \quad (4.28)$$

where μ^L and μ^R have been introduced in (4.9) and ω_G stands for the normalized unimodular Haar measure on the compact Polish group G .

Remark 4.7. The convolution factorization (4.28) is equivalent to the following assertion: If we let Z be a random variable whose law is μ , then

$$Z^L, Z^G \text{ and } Z^R \text{ are independent and the law of } Z^G \text{ is } \omega_G. \quad (4.29)$$

Here $(Z^L, Z^G, Z^R) = \psi^{-1}(Z)$ with $\psi : L \times G \times R \rightarrow LGR$ denoting the product mapping; see Proposition 4.2.

Proof of Theorem 4.6. Since $\mathcal{S}(\mu) = \overline{\mathcal{S}(\mu)\mathcal{S}(\mu)}$, we see that $\mathcal{S}(\mu)$ is a closed subsemigroup of S . By Proposition 4.5, we see that, for any $a \in \mathcal{S}(\mu)$,

$$\mu * \delta_{xa} = \mu * \delta_a, \quad \delta_{ax} * \mu = \delta_a * \mu, \quad x \in \mathcal{S}(\mu). \quad (4.30)$$

Then, for $a \in \mathcal{S}(\mu)$, we have

$$\mu * \delta_{ay} = \mu * \delta_a \ (y \in \mathcal{S}(\mu * \delta_a)), \quad \delta_{za} * \mu = \delta_a * \mu \ (z \in \mathcal{S}(\delta_a * \mu)) \quad (4.31)$$

In fact, for $y \in \mathcal{S}(\mu * \delta_a) = \overline{\mathcal{S}(\mu)a}$, we may take $\{x_n\} \subset \mathcal{S}(\mu)$ such that $x_n a \rightarrow y$, so that $\mu * \delta_a = \mu * \delta_{ax_n a} \rightarrow \mu * \delta_{ay}$.

Let $a \in \mathcal{S}(\mu)$ be fixed and set $\nu = \overline{\delta_a * \mu * \delta_a}$. Then $\mathcal{S}(\nu) = \overline{a\mathcal{S}(\mu)a}$ is a closed subsemigroup of S . For any $y \in \mathcal{S}(\nu) = \overline{a\mathcal{S}(\mu)a}$, we may take $\{x_n\} \subset \mathcal{S}(\mu)$ such that $ax_n a \rightarrow y$, so that, using (4.30), we have

$$\nu = \delta_a * \mu * \delta_a = \delta_{ax_n a^2} * \mu * \delta_a = \delta_{ax_n a} * \nu \rightarrow \delta_y * \nu, \quad (4.32)$$

which shows that $\nu|_{\mathcal{S}(\nu)}$ is ℓ^* -invariant. We see similarly that $\nu|_{\mathcal{S}(\nu)}$ is r^* -invariant. We may now apply Theorem 4.4 to see that $\mathcal{S}(\nu) = \overline{a\mathcal{S}(\mu)a}$ is a compact Polish group. Its identity is an idempotent of $\mathcal{S}(\mu)$.

Let $e \in E(\mathcal{S}(\mu))$. By the above argument with $a = e$, we see that $G := e\mathcal{S}(\mu)e$ is a compact Polish group (note that $e\mathcal{S}(\mu)e$ is closed by Lemma 3.1). Set $A := \mathcal{S}(\mu)e$. For $y \in A$, using (4.31), we have

$$\overline{Ay} = \overline{\mathcal{S}(\mu)ey} = \mathcal{S}(\mu * \delta_{ey}) = \mathcal{S}(\mu * \delta_e) = \mathcal{S}(\mu)e = A. \quad (4.33)$$

Since $Ay \cap e\mathcal{S}(\mu)e$ is a left ideal of the group $e\mathcal{S}(\mu)e$, we see that $Ay \cap e\mathcal{S}(\mu)e = e\mathcal{S}(\mu)e$, i.e. $e\mathcal{S}(\mu)e \subset Ay$, which shows $e \in Ay$. Hence

$$A = Ae \subset AAy \subset Ay \subset \overline{Ay} = A, \quad (4.34)$$

which yields $Ay = A$ for all $y \in A$. By Lemma 2.3, we see that A is a left group. We see similarly that $B := e\mathcal{S}(\mu)$ is a right group. By Theorem 2.13, we see that $\mathcal{S}(\mu)$ contains a completely simple kernel K , which is closed by Corollary 3.3.

By (4.30), we have

$$\mu * \delta_e * \mu = \int (\mu * \delta_e * \delta_a) \mu(da) = \int (\mu * \delta_a) \mu(da) = \mu * \mu = \mu. \quad (4.35)$$

By Lemma 2.4, we have $K = \mathcal{S}(\mu)e\mathcal{S}(\mu)$, and hence we obtain

$$K = \overline{K} = \overline{\mathcal{S}(\mu)e\mathcal{S}(\mu)} = \mathcal{S}(\mu * \delta_e * \mu) = \mathcal{S}(\mu), \quad (4.36)$$

which shows that $\mathcal{S}(\mu)$ is completely simple.

By (4.31), we see that $\mu * \delta_e$ is r^* -invariant on $A = \mathcal{S}(\mu)e = LG$, so that $\mu * \delta_e = \mu * \delta_e * \omega_G$. Hence, for any $B \in \mathcal{B}(\mathcal{S}(\mu))$,

$$\mu(B) = (\mu * \delta_e * \mu)(B) = (\mu * \delta_e * \omega_G * \mu)(B) \quad (4.37)$$

$$= \int \mu(dz_1) \int \mu(dz_2) \int \omega_G(dg) 1_B(z_1 e g z_2) \quad (4.38)$$

$$= \int \mu(dz_1) \int \mu(dz_2) \int \omega_G(dg) 1_B(z_1^L g z_2^R) = (\mu^L * \omega_G * \mu^R)(B), \quad (4.39)$$

which completes the proof. \square

The following proposition is a converse to Theorem 4.6.

Proposition 4.8. *Let S be a Polish semigroup and let $\mu_1, \mu_2 \in \mathcal{P}(S)$. Let G be a compact Polish subgroup of S and suppose that $\mathcal{S}(\mu_2 * \mu_1) \subset G$. Then $\mu := \mu_1 * \omega_G * \mu_2$ satisfies $\mu^2 = \mu$.*

Proof. For any $B \in \mathcal{B}(S)$, we have

$$\mu^2(B) = (\mu_1 * \omega_G * \mu_2 * \mu_1 * \omega_G * \mu_2)(B) \quad (4.40)$$

$$= \int \mu_1(dz_1) \int \omega_G(dg_1) \int (\mu_2 * \mu_1)(dg_2) \int \omega_G(dg_3) \int \mu_2(dz_2) 1_B(z_1 g_1 g_2 g_3 z_2) \quad (4.41)$$

$$= \int \mu_1(dz_1) \int \omega_G(dg_1) \int \mu_2(dz_2) 1_B(z_1 g_1 z_2) = (\mu_1 * \omega_G * \mu_2)(B) = \mu(B), \quad (4.42)$$

which completes the proof. \square

4.5 Infinite convolutions

Theorem 4.9 (Rosenblatt [39] and Mukherjea [28]). *Let S_0 be a Polish semigroup and let $\mu \in \mathcal{P}(S_0)$. Suppose that the sequence $\{\mu^n\}_{n=1}^\infty$ is tight. Let S denote the closure of the semigroup generated by $\mathcal{S}(\mu)$, i.e.*

$$S := \overline{\bigcup_{n=1}^{\infty} \mathcal{S}(\mu)^n}. \quad (4.43)$$

Then the following assertions hold:

(i) *There exists $\nu \in \mathcal{P}(S)$ such that $\nu^2 = \nu$, $\mu * \nu = \nu * \mu = \nu$ and*

$$\mu_n := \frac{1}{n} \sum_{k=1}^n \mu^k \xrightarrow[n \rightarrow \infty]{} \nu. \quad (4.44)$$

(ii) The family \mathcal{K} of cluster points of $\{\mu^n : n = 1, 2, \dots\}$ is a compact abelian group such that

$$\mathcal{S}(\nu) = \overline{\bigcup_{\lambda \in \mathcal{K}} \mathcal{S}(\lambda)}. \quad (4.45)$$

(iii) Let η denote the identity of \mathcal{K} . Then $\mathcal{S}(\eta)$ is a completely simple semigroup. Let $\mathcal{S}(\eta) = LHR$ denote the Rees decomposition at $e \in E(\mathcal{S}(\eta))$. Then H is a compact group and η admits the convolution factorization

$$\eta = \eta_L * \omega_H * \eta_R. \quad (4.46)$$

(iv) $\mathcal{S}(\nu)$ is a completely simple kernel of S containing the idempotent e . The Rees decomposition of $\mathcal{S}(\nu)$ at e is of the form $\mathcal{S}(\nu) = LGR$, where G is a compact group containing H , and ν admits the convolution factorization

$$\nu = \eta_L * \omega_G * \eta_R. \quad (4.47)$$

(v) For $g \in G$, we write $\omega_{gH} := \delta_g * \omega_H$. It holds that H is a closed normal subgroup of G and that there exists a Polish group isomorphism $F : \mathcal{K} \rightarrow G/H$ such that

$$\lambda = \eta^L * \omega_{F(\lambda)} * \eta^R, \quad (4.48)$$

Consequently, there exists $\gamma \in G$ such that $\mu^k * \eta$ admits the convolution factorization

$$\mu^k * \eta = \eta^L * \omega_{\gamma^k H} * \eta^R, \quad k = 1, 2, \dots, \quad (4.49)$$

and furthermore, \mathcal{K} and G/H may be represented as

$$\mathcal{K} = \overline{\{\eta, \mu * \eta, \mu^2 * \eta, \dots\}}, \quad G/H = \overline{\{H, \gamma H, \gamma^2 H, \dots\}}. \quad (4.50)$$

Remark 4.10. If the order of the group \mathcal{K} or G/H is finite, say p , then

$$\mathcal{K} = \{\eta, \mu * \eta, \dots, \mu^{p-1} * \eta\}, \quad G/H = \{H, \gamma H, \dots, \gamma^{p-1} H\} \quad (4.51)$$

with $\gamma^p \in H$. It is now obvious that $\lim_{n \rightarrow \infty} \mu^n$ converges if and only if $p = 1$.

Proof of Theorem 4.9. (i) Let $\|\cdot\|$ denote the total variation norm. For $j = 1, 2, \dots$, we have

$$\|\mu_n - \mu^j * \mu_n\| \leq \frac{1}{n} \left\| \sum_{k=1}^n \mu^k - \sum_{k=1}^n \mu^{k+j} \right\| = \frac{1}{n} \left\| \sum_{k=1}^j \mu^k - \sum_{k=n+1}^{n+j} \mu^k \right\| \leq \frac{2j}{n} \xrightarrow{n \rightarrow \infty} 0. \quad (4.52)$$

Since $\{\mu^n\}$ is tight, we see that $\{\mu_n\}$ is also tight. Let ν_1, ν_2 be cluster points of $\{\mu_n\}$. For $i = 1, 2$, we see by (4.52) that $\mu^j * \nu_i = \nu_i * \mu^j = \nu_i$ for $j = 1, 2, \dots$, so that $\mu_n * \nu_i = \nu_i * \mu_n = \nu_i$ for $n = 1, 2, \dots$, which implies $\nu_1 = \nu_1 * \nu_2 = \nu_2 * \nu_1 = \nu_2$. Hence we see that $\{\mu_n\}$ converges to some $\nu \in \mathcal{P}(S_0)$ and we have $\nu^2 = \nu$ and $\mu * \nu = \nu * \mu = \nu$. We

may apply Theorem 4.6 to see that $\mathcal{S}(\nu)$ is a completely simple semigroup and its group factor is compact.

(ii) Let us prove that $\mathcal{S}(\nu)$ and $\mathcal{S}(\mathcal{K}) := \overline{\bigcup_{\lambda \in \mathcal{K}} \mathcal{S}(\lambda)}$ are ideals of S . Let $a \in S$, $x \in \mathcal{S}(\nu)$ and $y \in \mathcal{S}(\mathcal{K})$. Then we may take $\{a_n\} \subset \mathcal{S}(\mu)^{m(n)} \subset \mathcal{S}(\mu^{m(n)})$ and $\{y_n\} \subset \mathcal{S}(\lambda_n)$ such that $a_n \rightarrow a$ and $y_n \rightarrow y$. Since

$$a_n x \in \mathcal{S}(\mu^{m(n)})\mathcal{S}(\nu) \subset \mathcal{S}(\mu^{m(n)} * \nu) = \mathcal{S}(\nu), \quad (4.53)$$

$$a_n y_n \in \mathcal{S}(\mu^{m(n)})\mathcal{S}(\lambda_n) \subset \mathcal{S}(\mu^{m(n)} * \lambda_n) \subset \mathcal{S}(\mathcal{K}), \quad (4.54)$$

we obtain $ax = \lim a_n x \in \mathcal{S}(\nu)$ and $ay = \lim a_n y_n \in \mathcal{S}(\mathcal{K})$, which shows that $\mathcal{S}(\nu)$ and $\mathcal{S}(\mathcal{K})$ are both left ideals of S . Similarly we see that they are also right ideals of S .

Let U be an open subset containing $\mathcal{S}(\nu)$. We shall prove that $\mu^n(U) \rightarrow 1$. Let $\varepsilon > 0$. By tightness, we may take a compact subset K_1 such that $\inf_n \mu^n(K_1) > 1 - \varepsilon$. We may take a compact subset $K_2 \subset \mathcal{S}(\nu)$ such that $\nu(K_2) > 1 - \varepsilon$. Since $K_1 K_2 \subset \mathcal{S}(\nu) \subset U$, we have $K_1 \times K_2 \subset \tilde{U} := \{(x, y) \in S_0 \times S_0 : xy \in U\}$. By the Wallace theorem (see, e.g., [21, Theorem 12 of Chapter 5]), we may take open subsets V_1 and V_2 such that $K_1 \subset V_1$, $K_2 \subset V_2$ and $V_1 \times V_2 \subset \tilde{U}$, which implies $V_1 V_2 \subset U$. Since $\mu_n \rightarrow \nu$, we have $\liminf_n \mu_n(V_2) \geq \nu(V_2) \geq \nu(K_2) > 1 - \varepsilon$. We may then take some n_0 such that $\mu^{n_0}(V) > 1 - \varepsilon$. We now have

$$\mu^{n+n_0}(U) = \iint 1_U(xy) \mu^n(dx) \mu^{n_0}(dy) \geq \mu^n(V_1) \mu^{n_0}(V_2) > (1 - \varepsilon)^2, \quad (4.55)$$

which leads to $\mu^n(U) \rightarrow 1$.

By the tightness assumption, we may apply Proposition 3.5 to see that \mathcal{K} is a compact abelian group. Let $\lambda \in \mathcal{K}$ and let $x \in S(\lambda)$. Suppose that $x \notin \mathcal{S}(\nu)$. We could then take disjoint open sets U and V such that $\mathcal{S}(\nu) \subset U$ and $x \in V$. If we let $\delta := \lambda(V)/2 > 0$, then $\mu^n(V) > \delta$ for infinitely many n , and then $\liminf_n \mu^n(U) \leq \liminf_n \mu^n(V^c) \leq 1 - \delta$, which would contradict $\mu^n(U) \rightarrow 1$. Hence we obtain $\mathcal{S}(\mathcal{K}) \subset \mathcal{S}(\nu)$. Since $\mathcal{S}(\nu)$ is a minimal ideal of S by Lemma 2.4 and since $\mathcal{S}(\mathcal{K})$ is an ideal of S , we see that $\mathcal{S}(\mathcal{K}) = \mathcal{S}(\nu)$.

(iii) By Theorem 4.6, we see that $\mathcal{S}(\eta)$ is a completely simple semigroup. Let $\mathcal{S}(\eta) = LHR$ denote the Rees decomposition at $e \in E(\mathcal{S}(\eta))$ (hence $RL \subset H$). Then the group factor H is compact and η admits the convolution factorization (4.46).

(iv) We have already seen in (i) that $\mathcal{S}(\nu)$ is a completely simple kernel of S . Since $\mathcal{S}(\eta) \subset \mathcal{S}(\mathcal{K}) = \mathcal{S}(\nu)$, we have $e \in E(\mathcal{S}(\nu))$. Let $\mathcal{S}(\nu) = L'GR'$ denote the Rees decomposition at e . As a consequence of Theorem 4.6, we see that ν admits the convolution factorization $\nu = \eta^{L'} * \omega_G * \eta^{R'}$. Since $\mathcal{S}(\eta) \subset \mathcal{S}(\nu)$ and $L = E(\mathcal{S}(\eta)e)$ etc., we see that $L \subset L'$, $H \subset G$ and $R \subset R'$.

Let us prove that $L' = L$ and $R' = R$. Let $z = xgy \in L'GR'$. Since $\mathcal{S}(\nu) = \mathcal{S}(\mathcal{K})$, we may take $z_n \in \mathcal{S}(\lambda_n)$ such that $z_n \rightarrow z$. Since \mathcal{K} is abelian, we have $\lambda_n * \lambda_n^{-1} = \lambda_n^{-1} * \lambda_n = \eta$, and by Proposition 4.2 we have $\lambda_n^{L'} = \eta^{L'} = \eta^L$ and $\lambda_n^{R'} = \eta^{R'} = \eta^R$. Hence we obtain $x_n := z_n^{L'} \in \mathcal{S}(\lambda_n^{L'}) = \mathcal{S}(\eta^L) = L$ and $y_n := z_n^{R'} \in \mathcal{S}(\lambda_n^{R'}) = \mathcal{S}(\eta^R) = R$, and thus $x = \lim x_n \in L$ and $y = \lim y_n \in R$, which shows $L' = L$ and $R' = R$.

(v) Let $\lambda \in \mathcal{K}$. For $z = xgy \in \mathcal{S}(\lambda) \subset \mathcal{S}(\nu) = LGR$, since $RL \subset H$, we have

$$xgy \in LgHR \subset LHRxgyLHR \subset \mathcal{S}(\eta)\mathcal{S}(\lambda)\mathcal{S}(\eta) \subset \mathcal{S}(\eta * \lambda * \eta) = \mathcal{S}(\lambda). \quad (4.56)$$

Hence we have $\mathcal{S}(\lambda) = LG_\lambda R$ for $G_\lambda := \bigcup\{gH : z = xgy \in \mathcal{S}(\lambda)\} \subset G$, and we also have $G_\lambda = \bigcup\{Hg : z = xgy \in \mathcal{S}(\lambda)\}$ similarly. Note that $G_\lambda H = HG_\lambda = G_\lambda$. Take $g_\lambda \in G$ such that $Hg_\lambda^{-1} \subset G_{\lambda^{-1}}$. Then we obtain

$$LHg_\lambda^{-1}G_\lambda R \subset LG_{\lambda^{-1}}RLG_\lambda R \subset \mathcal{S}(\lambda^{-1})\mathcal{S}(\lambda) \subset \mathcal{S}(\lambda^{-1} * \lambda) \subset \mathcal{S}(\eta) = LHR, \quad (4.57)$$

which yields that $Hg_\lambda^{-1}G_\lambda \subset H$ and hence $G_\lambda = g_\lambda H$. Similarly, we obtain $G_\lambda = Hg_\lambda$.

For any $h \in H$ and $g \in G \subset \mathcal{S}(\nu) = \mathcal{S}(\mathcal{K})$, we may take $z_n = x_n g_n y_n \in \mathcal{S}(\lambda_n)$ such that $z_n \rightarrow g$ and consequently $g_n \rightarrow g$. In a similar way to (4.57), we have

$$g_n h g_n^{-1} \in (g_n H)(H g_n^{-1}) = G_{\lambda_n} G_{\lambda_n^{-1}} \subset \mathcal{S}(\eta) = LHR, \quad (4.58)$$

which shows $g_n h g_n^{-1} \in eLHRe = H$. Letting $n \rightarrow \infty$, we obtain $ghg^{-1} \in H$, which shows that H is a normal subgroup of G . Since G and H are both compact, we see by [15, Theorem 5.22] that the quotient group $G/H = \{gH : g \in G\}$ is also compact. Let $\pi : G \rightarrow G/H$ denote the natural projection.

Since

$$\mathcal{S}(\eta^R * \lambda * \eta^L) = \overline{\mathcal{S}(\eta^R)\mathcal{S}(\lambda)\mathcal{S}(\eta^L)} = \overline{RLG_\lambda RL} \subset \overline{Hg_\lambda H H} = g_\lambda H, \quad (4.59)$$

we obtain the convolution factorization

$$\lambda = \eta\lambda\eta = \eta^L * \omega_H * (\eta^R * \lambda * \eta^L) * \omega_H * \eta^R = \eta^L * \omega_{g_\lambda H} * \eta^R. \quad (4.60)$$

We now define the mapping $F : \mathcal{K} \rightarrow G/H$ by $F(\lambda) := g_\lambda H$. For $\lambda_1, \lambda_2 \in \mathcal{K}$, then

$$\lambda_1 * \lambda_2 = \eta^L * \omega_{g_{\lambda_1} H} * (\eta^R * \eta^L) * \omega_{g_{\lambda_2} H} * \eta^R = \eta^L * \omega_{(g_{\lambda_1} g_{\lambda_2} H)} * \eta^R, \quad (4.61)$$

since $RL \subset H$, which shows that F is a group homomorphism. Injectivity of F is obvious by (4.60). Let $g \in G$. As we have seen it above, we may take $z_n = x_n g_n y_n \in \mathcal{S}(\lambda_n)$ such that $g_n \rightarrow g$ and $g_n H = g_{\lambda_n} H$. Then, by (4.60), we have

$$\lambda_n = \eta^L * \omega_{g_{\lambda_n} H} * \eta^R \rightarrow \eta^L * \omega_{gH} * \eta^R =: \lambda. \quad (4.62)$$

This shows that $\lambda \in \mathcal{K}$ and $F(\lambda) = gH$, which yields surjectivity of F . Suppose $\mathcal{K} \ni \lambda_n \rightarrow \lambda \in \mathcal{K}$. By (4.60), we have

$$\omega_{F(\lambda_n)} = \delta_e * \lambda_n * \delta_e \rightarrow \delta_e * \lambda * \delta_e = \omega_{F(\lambda)} \quad \text{in } \mathcal{P}(G), \quad (4.63)$$

which shows by the continuity of the natural projection π that

$$\delta_{F(\lambda_n)} = \omega_{F(\lambda_n)} \circ \pi^{-1} \rightarrow \omega_{F(\lambda)} \circ \pi^{-1} = \delta_{F(\lambda)} \quad \text{in } \mathcal{P}(G/H), \quad (4.64)$$

which implies $F(\lambda_n) \rightarrow F(\lambda)$ and we have seen continuity of F . Since \mathcal{K} is compact and G/H is Hausdorff, we see by [21, Theorem 9 of Chapter 5] that F is homeomorphic. Since $F(\mu * \eta) \in G/H$, we may take $\gamma \in G$ such that $F(\mu * \eta) = \gamma H$, and then we obtain (4.49) since $(\mu * \eta)^k = \mu^k = \mu^k * \eta$ and F is a group homomorphism.

Finally, let us prove the representations (4.50). Since any $\lambda \in \mathcal{K}$ can be represented as $\lambda = \lambda * \eta = \lim \mu^{n(k)} * \eta$, we see that $\mathcal{K} = \overline{\{\eta, \mu * \eta, \mu^2 * \eta, \dots\}}$. Since for any $g \in G$ we have $F(\lambda) = gH$ for some $\lambda = \lim \mu^{n(k)} * \eta \in \mathcal{K}$, we obtain $gH = F(\lambda) = \lim F(\mu^{n(k)} * \eta) = \lim \gamma^{n(k)} H$ in G/H , which yields $G/H = \overline{\{H, \gamma H, \gamma^2 H, \dots\}}$. \square

References

- [1] G. Budzban and A. Mukherjea. Subsemigroups of completely simple semigroups and weak convergence of convolution products of probability measures. *Semigroup Forum*, 68(3):400–410, 2004.
- [2] G. Choquet and J. Deny. Sur l'équation de convolution $\mu = \mu * \sigma$. *C. R. Acad. Sci. Paris*, 250:799–801, 1960.
- [3] S. T. L. Choy. Idempotent measures on compact semigroups. *Proc. London Math. Soc. (3)*, 20:717–733, 1970.
- [4] D. L. Cohn. *Measure theory*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser/Springer, New York, second edition, 2013.
- [5] H. S. Collins. Convergence of convolution iterates of measures. *Duke Math. J.*, 29:259–264, 1962.
- [6] H. S. Collins. Idempotent measures on compact semigroups. *Proc. Amer. Math. Soc.*, 13:442–446, 1962.
- [7] I. Csiszár. On infinite products of random elements and infinite convolutions of probability distributions on locally compact groups. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 5:279–295, 1966.
- [8] P. L. Davies and D. N. Shanbhag. A generalization of a theorem of Deny with applications in characterization theory. *Quart. J. Math. Oxford Ser. (2)*, 38(149):13–34, 1987.
- [9] Y. Derriennic. Sur le théorème de point fixe de Brunel et le théorème de Choquet-Deny. *Ann. Sci. Univ. Clermont-Ferrand II Probab. Appl.*, (4):107–111, 1985.
- [10] J. Duncan. Primitive idempotent measures on compact semigroups. *Proc. Edinburgh Math. Soc. (2)*, 17:95–103, 1970.
- [11] R. Ellis. A note on the continuity of the inverse. *Proc. Amer. Math. Soc.*, 8:372–373, 1957.

- [12] H. Furstenberg and Y. Katznelson. Idempotents in compact semigroups and Ramsey theory. *Israel J. Math.*, 68(3):257–270, 1989.
- [13] I. Glicksberg. Convolution semigroups of measures. *Pacific J. Math.*, 9:51–67, 1959.
- [14] M. Heble and M. Rosenblatt. Idempotent measures on a compact topological semigroup. *Proc. Amer. Math. Soc.*, 14:177–184, 1963.
- [15] E. Hewitt and K. A. Ross. *Abstract harmonic analysis. Vol. I*, volume 115 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin-New York, second edition, 1979. Structure of topological groups, integration theory, group representations.
- [16] G. Högnäs and A. Mukherjea. *Probability measures on semigroups*. Probability and its Applications (New York). Springer, New York, second edition, 2011. Convolution products, random walks, and random matrices.
- [17] Y. Ito, T. Sera, and K. Yano. Resolution of sigma-fields for multiparticle finite-state action evolutions with infinite past. Preprint, arXiv:2008.12407.
- [18] O. Kallenberg. *Foundations of modern probability*. Probability and its Applications (New York). Springer-Verlag, New York, second edition, 2002.
- [19] Y. Kawada and K. Itô. On the probability distribution on a compact group. I. *Proc. Phys.-Math. Soc. Japan (3)*, 22:977–998, 1940.
- [20] A. S. Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [21] J. L. Kelley. *General topology*. Graduate Texts in Mathematics, No. 27. Springer-Verlag, New York-Berlin, 1975. Reprint of the 1955 edition [Van Nostrand, Toronto, Ont.].
- [22] B. M. Kloss. Probability distributions on bicomact topological groups. *Theor. Probability Appl.*, 4:237–270, 1959.
- [23] K.-S. Lau and W. B. Zeng. The convolution equation of Choquet and Deny on semigroups. *Studia Math.*, 97(2):115–135, 1990.
- [24] Y.-F. Lin. Not necessarily abelian convolution semigroups of probability measures. *Math. Z.*, 91:300–307, 1966.
- [25] C.-C. Lo and A. Mukherjea. Convergence in distribution of products of $d \times d$ random matrices. *J. Math. Anal. Appl.*, 162(1):71–91, 1991.
- [26] A. Mukherjea. On the convolution equation $P = PQ$ of Choquet and Deny for probability measures on semigroups. *Proc. Amer. Math. Soc.*, 32:457–463, 1972.
- [27] A. Mukherjea. Limit theorems for convolution iterates of a probability measure on completely simple or compact semigroups. *Trans. Amer. Math. Soc.*, 225:355–370, 1977.

- [28] A. Mukherjea. Limit theorems: stochastic matrices, ergodic Markov chains, and measures on semigroups. In *Probabilistic analysis and related topics, Vol. 2*, pages 143–203. Academic Press, New York-London, 1979.
- [29] A. Mukherjea. *Topics in products of random matrices*, volume 87 of *Tata Institute of Fundamental Research Lectures on Mathematics*. Published by Narosa Publishing House, New Delhi; for the Tata Institute of Fundamental Research, Mumbai, 2000.
- [30] A. Mukherjea and T. C. Sun. Convergence of products of independent random variables with values in a discrete semigroup. *Z. Wahrsch. Verw. Gebiete*, 46(2):227–236, 1978/79.
- [31] A. Mukherjea and N. A. Tserpes. Idempotent measures on locally compact semigroups. *Proc. Amer. Math. Soc.*, 29:143–150, 1971.
- [32] A. Mukherjea and N. A. Tserpes. *Measures on topological semigroups: convolution products and random walks*. Lecture Notes in Mathematics, Vol. 547. Springer-Verlag, Berlin-New York, 1976.
- [33] A. Nakassis. Limit behavior of the convolution iterates of a probability measure on a semigroup of matrices. *J. Math. Anal. Appl.*, 70(2):337–347, 1979.
- [34] K. R. Parthasarathy. *Probability measures on metric spaces*. AMS Chelsea Publishing, Providence, RI, 2005. Reprint of the 1967 original.
- [35] H. Pfister. Continuity of the inverse. *Proc. Amer. Math. Soc.*, 95(2):312–314, 1985.
- [36] J. S. Pym. Idempotent measures on semigroups. *Pacific J. Math.*, 12:685–698, 1962.
- [37] C. R. Rao and D. N. Shanbhag. Further extensions of the Choquet-Deny and Deny theorems with applications in characterization theory. *Quart. J. Math. Oxford Ser. (2)*, 40(159):333–350, 1989.
- [38] A. Raugi. Un théorème de Choquet-Deny pour les semi-groupes abéliens. In *Théorie du potentiel (Orsay, 1983)*, volume 1096 of *Lecture Notes in Math.*, pages 502–520. Springer, Berlin, 1984.
- [39] M. Rosenblatt. Limits of convolution sequences of measures on a compact topological semigroup. *J. Math. Mech.*, 9:293–305, 1960.
- [40] M. Rosenblatt. Products of independent identically distributed stochastic matrices. *J. Math. Anal. Appl.*, 11:1–10, 1965.
- [41] M. Rosenblatt. *Markov processes. Structure and asymptotic behavior*. Die Grundlehren der mathematischen Wissenschaften, Band 184. Springer-Verlag, New York-Heidelberg, 1971.
- [42] M. Rosenblatt. Convolution sequences of measures on the semigroup of stochastic matrices. In *Random matrices and their applications (Brunswick, Maine, 1984)*, volume 50 of *Contemp. Math.*, pages 215–220. Amer. Math. Soc., Providence, RI, 1986.

- [43] Š. Schwarz. Convolution semigroup of measures on compact noncommutative semigroups. *Czechoslovak Math. J.*, 14(89):95–115, 1964.
- [44] Š. Schwarz. Product decomposition of idempotent measures on compact semigroups. *Czechoslovak Math. J.*, 14(89):121–124, 1964.
- [45] K. Stromberg. Probabilities on a compact group. *Trans. Amer. Math. Soc.*, 94:295–309, 1960.
- [46] T.-C. Sun and N. A. Tserpes. Idempotent measures on locally compact semigroups. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 15:273–278, 1970.
- [47] G. J. Székely and W. B. Zeng. The Choquet-Deny convolution equation $\mu = \mu * \sigma$ for probability measures on abelian semigroups. *J. Theoret. Probab.*, 3(2):361–365, 1990.
- [48] A. Tortrat. Lois tendues, convergence en probabilité et équation $P * P' = P$. *C. R. Acad. Sci. Paris*, 258:3813–3816, 1964.
- [49] K. Urbanik. On the limiting probability distribution on a compact topological group. *Fund. Math.*, 44:253–261, 1957.
- [50] J. Woś. The convolution equation of Choquet and Deny for probability measures on discrete semigroups. *Colloq. Math.*, 47(1):143–148, 1982.
- [51] W. Żelazko. A theorem on B_0 division algebras. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.*, 8:373–375, 1960.