

CONCENTRATION INEQUALITIES FOR SOME NEGATIVELY DEPENDENT BINARY RANDOM VARIABLES

RADOSŁAW ADAMCZAK AND BARTŁOMIEJ POLACZYK

ABSTRACT. We investigate concentration properties of functions of random vectors with values in the discrete cube, satisfying the stochastic covering property (SCP) or the strong Rayleigh property (SRP).

Our result for SCP measures include subgaussian inequalities of bounded-difference type extending classical results by Pemantle and Peres and their counterparts for matrix-valued setting strengthening recent estimates by Aoun, Banna and Youssef. Under a stronger assumption of the SRP we obtain Bernstein-type inequalities for matrix-valued functions, generalizing recent bounds for linear combinations of positive definite matrices due to Kyng and Song.

We also treat in detail the special case of independent Bernoulli random variables conditioned on their sum for which we obtain strengthened estimates, deriving in particular modified log-Sobolev inequalities, Talagrand's convex distance inequality and, as corollaries, concentration results for convex functions and polynomials, as well as improved estimates for matrix-valued functions. These results generalize inequalities for the uniform measure on slices of the discrete cube, studied extensively by many authors. Our approach is based on recent results by Hermon and Salez and a general framework involving modified log-Sobolev inequalities on the discrete cube, which is of independent interest.

Keywords: concentration of measure, strong Rayleigh measure, stochastic covering property

AMS Classification: 60E15, 60B20, 60J28

1. INTRODUCTION

Investigating families of binary random variables with negatively dependent coordinates is an important problem from the point of view of computer science and combinatorics, which in the recent years has attracted considerable attention, see, e.g., [43, 49, 14, 44, 34, 26, 10, 6, 8, 32]. A wide and important class of such variables is constituted by those satisfying the *strong Rayleigh property* (abbrev. SRP) introduced by Borcea et al. [14]. More precisely, a probability measure π on the hypercube $\mathcal{B}_n := \{0, 1\}^n$ satisfies the SRP if its generating polynomial

$$\mathbb{C}^n \ni z \mapsto \sum_{x \in \mathcal{B}_n} \pi(x) \prod_{i=1}^n z_i^{x_i}$$

has no roots z whose all coordinates lie in the (strict) upper half-plane. The examples of such measures are, e.g., the law of independent Bernoulli random variables conditioned on their sum, determinantal measures, uniform measure on the bases of balanced matroids, laws of point processes or measures obtained by running exclusion dynamics on the cube, cf. Pemantle and Peres [44].

The main purpose of this article is to deepen the understanding of the concentration of measure phenomenon in the context of strong Rayleigh distributions and related classes of probability measures on the discrete cube. In some of our considerations we will exploit only a more general notion of the *stochastic covering property* (abbrev. SCP, cf. Definition 2.1) introduced by Pemantle and Peres [44], since this condition already turns out to provide a useful framework for proving concentration results [44, 29, 8, 34, 32, 2]. On the other hand for some more specialized inequalities we will restrict our attention to independent Bernoulli variables conditioned on their sum taking some fixed value. Distributions of this type generalize the uniform measure on slices of the discrete cube, related to the Bernoulli–Laplace model, which has been extensively studied, e.g., by Lee–Yau, Bobkov–Tetali, Gao–Quastel [37, 13, 25] and more recently by Samson [48] and Sambale–Sinulis [46]. The non-uniform distribution given by conditioned Bernoulli variables has found applications, e.g.,

Date: Last changes: August 31, 2021.

Research partially supported by the National Science Centre, Poland, via the Sonata Bis grant no. 2015/18/E/ST1/00214 (RA) and the Preludium grant no. 2020/37/N/ST1/02667 (BP).

in survey sampling being a model of sampling without replacement from a finite population, with prescribed inclusion probabilities, which maximizes the entropy (often referred to as *conditional Poisson sampling*). We refer to [20, 19, 21, 52, 10] for properties and applications of this family of distributions.

1.1. State of the art. The landmark paper that initiated the study of concentration phenomenon implied by the SCP is due to Pemantle and Peres [44] who, using the martingale method, proved a sub-Gaussian concentration bound for measures satisfying the SCP and functions that are Lipschitz with respect to the Hamming distance $d_H(x, y) = \sum_i \mathbf{1}_{x_i \neq y_i}$. Recently, Hermon and Salez [29], building on the works [40, 31], retrieved this estimate by proving that the SCP implies the modified log-Sobolev inequality.

These findings in terms of concentration of measure can be summarized as follows (we use the notation $\pi(f) := \int f d\pi$ for a probability measure π on \mathcal{B}_n and $f: \mathcal{B}_n \rightarrow \mathbb{R}$).

Theorem 1.1 (Pemantle–Peres [44]). *For a probability measure π on \mathcal{B}_n satisfying the SCP and any $f: \mathcal{B}_n \rightarrow \mathbb{R}$ such that*

$$|f(x) - f(y)| \leq d_H(x, y) \quad \forall x, y \in \mathcal{B}_n$$

the following estimate holds for all $t > 0$

$$(1.1) \quad \pi(f > \pi(f) + t) \leq \exp(-t^2/8n).$$

If π is k -homogeneous (i.e., it is supported on the set of binary vectors with exactly k coefficients equal to one), then n in the above expression can be replaced with k .

Recently, a sub-exponential version of Theorem 1.1 for matrix-valued functions has been shown by Aoun et al. [8], who develop a general framework for deducing concentration bounds for matrix-valued functions from the Poincaré inequality. A Bernstein-type bound for measures with the SRP, which in certain situations may give stronger concentration, has been also developed by Kyng and Song [34] for functions of the form $f(x) = \sum_{i=1}^n x_i C_i$, where C_i are nonnegative definite matrices (see Theorem 2.8 and Remark 2.10 below).

While concentration estimates and functional inequalities for general SCP measures are relatively recent, investigation of uniform measures on slices of the discrete cube in this context has much longer history. Such measures are of interest in relation to the Bernoulli–Laplace models of statistical physics and with uniform sampling without replacement. In particular Lee and Yau studied the Poincaré and log-Sobolev inequalities for such measures, whereas Bobkov–Tetali [13] and independently Gao–Quastel [25] investigated modified log-Sobolev inequalities relevant for concentration estimates. Strong concentration results for this case can be also obtained by projection from Talagrand’s convex distance inequality for uniform measure on the symmetric group [51]. Samson [48] complemented this approach by proving corresponding transportation inequalities. Very recently Sambale and Sinulis [46], investigating general multislices, recovered convex distance inequalities by means of functional inequalities and also obtained concentration for polynomials. One should stress that concentration results for slices of the cube provided by the above references are substantially stronger than those coming from more general inequalities for SCP or SRP measures.

The uniform measure on slices of the cube can be seen as a special case of the distribution of independent Bernoulli random variables conditioned on their sum, when all the variables have the same probability of success. Such general distributions are known to be strong Rayleigh. To our best knowledge there has not been much work concerning refined concentration inequalities for general measures of this type. The only exception we are aware of is a recent article [10] by Bertail and Cléménçon in which the authors, motivated by applications to survey sampling, obtain precise Bernstein-type inequalities for linear functionals.

1.2. Our contribution. We develop two types of results.

Firstly, for general measures satisfying the SCP we extend the martingale argument from Pemantle and Peres [44] and generalize Theorem 1.1 to Lipschitz functions with respect to more general weighted Hamming distances $d_\alpha(x, y) = \sum \alpha_i \mathbf{1}_{x_i \neq y_i}$ obtaining a bounded-difference type inequality. We use the approach developed for the scalar case together with matrix bounded-difference inequality due to Tropp [54] to get an analogous concentration for matrix-valued functions, strengthening the results of Aoun et al. [8], in particular obtaining a subgaussian inequality in place of a subexponential one. Under a stronger assumption of the SRP we are also able to strengthen the Bernstein-type inequality of Kyng and Song and extend it from linear combinations with coefficients in nonnegative definite matrices to general functions satisfying a matrix bounded-difference type assumptions.

Secondly, building up on the work of Hermon and Salez [29], we develop an abstract condition which implies Talagrand’s convex distance inequality, matrix-Bernstein inequality and higher order concentration for tetrahedral polynomials. Finally, we prove that this condition is satisfied for the distribution of Bernoulli random variables conditioned on their sum being equal to some constant, yielding all the aforementioned results in this case.

1.3. Organization of the article. In Section 2 we present our results concerning concentration for general measures satisfying the SCP/SRP. In Section 3 we specialize our analysis to Bernoulli random variables conditioned on their sum being equal to some constant. Then, in Section 4 we formulate an abstract framework that allows to deduce the results of Section 3. Finally, all the proofs are presented in Sections 5, 6 and 7.

2. CONCENTRATION UNDER THE SCP AND SRP

In this section we present our concentration results for general measures satisfying the SCP or SRP. Let us start with introducing some notation. For $x = (x_1, \dots, x_n) \in \mathcal{B}_n := \{0, 1\}^n$ and any $S \subset \{1, 2, \dots, n\} =: [n]$ we use the shorthand notation $x_S = (x_i)_{i \in S}$. For any $r \in [n]$ we denote $x_{>r} = (x_i)_{i > r}$ (and analogously with relations other than $>$)¹. We also write x^i for the vector obtained from $x \in \mathcal{B}_n$ by flipping its i -th coordinate and x^{ij} for the vector obtained by swapping the i -th and j -th coordinate, i.e., $x^i = x \pm e_i$ and if $x_i \neq x_j$ then $x^{ij} = x \pm e_i \mp e_j$ for $i, j \in [n]$, where $e_i \in \mathcal{B}_n$ is the vector with one on the i -th and zeros on the remaining coordinates; whereas $x^{ij} = x$ if $x_i = x_j$. We remark that the notation x^{ij} should not be confused with $(x^i)^j$. The law of a random variable X is denoted by $\mathcal{L}(X)$, whereas $\mathcal{L}(X|A)$ stands for the conditional law of X given an event A (with an analogous convention for conditioning with respect to σ -fields or other random variables).

Below, we recall the definition of the SCP.

Definition 2.1 (Stochastic covering property). For $x, y \in \mathcal{B}_n$, we say that x covers y , denoted $x \triangleright y$, if $x = y$ or $x = y + e_i$ for some $i \in [n]$.

A random variable X taking values in \mathcal{B}_n satisfies the SCP if for any $S \subset [n]$ and any $x, y \in \mathcal{B}_n$ satisfying $x_S \triangleright y_S$ there exists a coupling (U, V) between the conditional distributions $\mathcal{L}(X_{S^c} | X_S = y_S)$ and $\mathcal{L}(X_{S^c} | X_S = x_S)$ such that $U \triangleright V$. A measure π satisfies the SCP if X with law π does so.

Remark 2.2. As indicated in the introduction, the SCP is implied by the SRP, cf. [44]. The opposite however is not true, as is demonstrated, e.g., by Cryan et al. in [22, Appendix A], where the authors show that yet another possible generalization of the SRP, the strong log-concavity, is incomparable with the SCP. In particular, they construct a distribution that is supported on the bases of a matroid, and that satisfies the SCP and violates the log-concavity (and whence the SRP as well).

For a finite sequence x , we denote by x^\downarrow the non-increasing rearrangement of the elements of x and for $\alpha \in [0, \infty)^n =: \mathbb{R}_+^n$ and $x, y \in \mathcal{B}_n$ we define the α -weighted Hamming distance $d_\alpha(x, y) = \sum_i \alpha_i \mathbf{1}_{x_i \neq y_i}$. Finally, for $p \in [1, \infty]$, $|\cdot|_p$ is the ℓ_p norm on \mathbb{R}^n and $|\cdot| := |\cdot|_2$ denotes the Euclidean norm.

The first main result of this paper is the following generalization of Theorem 1.1.

Theorem 2.3. *For a probability measure π on \mathcal{B}_n satisfying the SCP, any $f: \mathcal{B}_n \rightarrow \mathbb{R}$ and $\alpha \in \mathbb{R}_+^n$ such that*

$$|f(x) - f(y)| \leq d_\alpha(x, y) \quad \forall x, y \in \mathcal{B}_n$$

the following estimate holds for all $t > 0$:

$$\pi(f > \pi(f) + t) \leq \exp(-t^2/8|\alpha|^2).$$

If π is k -homogeneous then $8|\alpha|^2$ in the above estimate can be replaced with $16 \sum_{i=1}^k (\alpha_i^\downarrow)^2$.

Remark 2.4. Theorem 2.3 implies Theorem 1.1 (up to an absolute constant in the exponent) by taking $\alpha = (1, 1, \dots, 1)$. Moreover, by considering functions of the form $f(x) = \sum_i c_i x_i$ with $|c|^2 \ll n|c|_\infty^2$ in the non-homogeneous or $\sum_{i=1}^k (c_i^\downarrow)^2 \ll k|c|_\infty^2$ in the k -homogeneous case, one can see that Theorem 2.3 can give substantially better concentration estimates than Theorem 1.1.

¹We adopt the convention that if $x \in \mathcal{B}_n$ then $x_{>n} = \emptyset$ and as a consequence, e.g., $\mathbb{P}(\cdot | X_{>n} = \emptyset) = \mathbb{P}(\cdot)$.

We now formulate the matrix analogue of Theorem 2.3. To this end, let us denote the space of d -dimensional Hermitian matrices by \mathcal{H}_d , the identity matrix in \mathcal{H}_d by I_d , the maximal eigenvalue of $H \in \mathcal{H}_d$ by $\lambda_{\max}(H)$ and the operator norm of H by $\|H\|$.

Theorem 2.5. *For a probability measure π on \mathcal{B}_n satisfying the SCP, any $f: \mathcal{B}_n \rightarrow \mathcal{H}_d$ and $\alpha \in \mathbb{R}_+^n$ such that*

$$(2.1) \quad \|f(x) - f(y)\| \leq d_\alpha(x, y) \quad \forall x, y \in \mathcal{B}_n$$

the following estimate holds for all $t > 0$

$$\pi(\lambda_{\max}(f - \pi(f)) > t) \leq d \exp(-t^2/32|\alpha|^2).$$

If π is k -homogeneous then $32|\alpha|^2$ in the above estimate can be replaced with $64 \sum_{i=1}^k (\alpha_i^\downarrow)^2$.

Remark 2.6. Recently, Aoun et al. [8] showed that for any k -homogeneous probability measure π on \mathcal{B}_n satisfying the SCP and any $f: \mathcal{B}_n \rightarrow \mathcal{H}_d$ such that

$$\|f(x) - f(y)\| \leq d_H(x, y) \quad \forall x, y \in \mathcal{B}_n$$

the following estimate applies

$$(2.2) \quad \pi(\lambda_{\max}(f - \pi(f)) > t) \leq d \exp\left(-\frac{t^2}{8k + 2t\sqrt{2k}}\right).$$

The exponent in (2.2) is proportional to $-t/2\sqrt{2k}$ for t big enough and whence for such t Theorem 2.5 applied with $\alpha = (1, \dots, 1)$ strengthens on (2.2) (and on an analogous result from [32]) as it yields a sub-Gaussian estimate.

Remark 2.7. Using semigroup techniques together with matrix concentration results implied by the Poincaré inequality due to Aoun et al. [8], we are also able to derive a sub-exponential concentration inequality for general measures satisfying the SCP under weaker assumptions on f than those of Theorem 2.5, cf. Remark 3.7.

When comparing the inequality of Theorem 2.5 or the results from [8] with results for matrix-valued functions of independent random variables, one can ask if it is possible to weaken the assumptions on the function f and instead of the Lipschitz constant with respect to d_α use some weaker parameter, involving bounds on the increments of the function in terms of the positive semidefinite order. In many situations one encounters functions for which $(f(x) - f(x^i))^2 \preceq C_i^2$ where C_i are some positive semidefinite matrices and \preceq stands for the positive semidefinite order (note that considering arbitrary matrices C_i is a generalization of the condition (2.1), which corresponds to the special case $C_i^2 = \alpha_i^2 I_d$). The simplest, yet important situation of this type is given by $f(x) = \sum_{i=1}^n x_i C_i$. Inequalities for such functions together with algorithmic applications were considered by Kyng and Song in [34]. It turns out that their approach can be adapted to the setting of general functions, yielding the following theorem.

Theorem 2.8. *Let π be a k -homogeneous probability measure \mathcal{B}_n satisfying the strong Rayleigh property and $f: \mathcal{B}_n \rightarrow \mathcal{H}_d$ be such that there exists a sequence $C_1, \dots, C_n \in \mathcal{H}_d$ satisfying*

$$(2.3) \quad (f(x) - f(x^i))^2 \preceq C_i^2 \quad \forall x \in \mathcal{B}_n, i \in [n].$$

Then for any $t > 0$,

$$(2.4) \quad \pi(\lambda_{\max}(f - \pi(f)) > t) \leq d \exp\left(-\frac{t^2}{8\|\pi(\tilde{f})\| \log(ek) + \frac{4}{3}Kt}\right),$$

where $\tilde{f}(x) = \sum_{i=1}^n x_i C_i^2$ and $K = \max_{i \leq n} \|C_i\|$.

Remark 2.9. In fact, the only place in the proof of Theorem 2.8 where we use the SRP in its full strength is to get that $\mathbb{P}(X_i = 1 | X_{i_1} = 1, \dots, X_{i_k} = 1) \leq \mathbb{P}(X_i = 1)$ for $X \sim \pi$ and any $i, k \in [n]$ and $i_1, \dots, i_k \subset [n] \setminus \{i\}$. Therefore, in Theorem 2.8 it suffices to assume that π satisfies the SCP and negative association, which is implied by the SRP, cf. [44].

Remark 2.10. It is natural to expect that $\log(ek)$ in (2.4) is just an artefact of the proof. Removing it even just for functions of the form $f(x) = \sum_{i=1}^n x_i C_i$ for positive semidefinite matrices C_i would lead to improvement of certain algorithmic constructions related to graph sparsifiers obtained via random spanning trees, cf. [34].

Let us also point out that even though Theorem 2.8 applies to general functions, when specialized to the function f as discussed above, it improves slightly on the results from [34],

which instead of $\|\pi(\tilde{f})\|$ use a larger quantity $K\|\pi(\hat{f})\|$ with $\hat{f}(x) = \sum_{i=1}^n x_i C_i$ (recall that C_i 's are nonnegative definite). One should stress however that we rely on the approach worked out in [34].

3. CONCENTRATION FOR CONDITIONAL BERNOULLIS

In this section, we present our concentration results concerning Bernoulli random variables conditioned on their sum being constant. These include Talagrand's convex distance inequality, matrix-Bernstein inequality and concentration for polynomials.

We start with introducing the notation. For a sequence $p = (p_1, \dots, p_n) \in (0, 1)^n$, let $B = (B_1, \dots, B_n)$ be a sequence of independent Bernoulli random variables with probabilities of success p_i , i.e., $\mathbb{P}(B_i = 1) = 1 - \mathbb{P}(B_i = 0) = p_i$ for $i \in [n]$. Finally, set $X = (X_1, \dots, X_n) \sim \mathcal{L}(B \mid \sum_i B_i = k)$ for some $k \in \{0, \dots, n\}$ and denote the distribution of X by $\pi(p, k)$.

Our first contribution is a counterpart of the celebrated convex distance inequality, introduced for the first time by Talagrand [50] for product measures on the cube.

Theorem 3.1. *If $\pi \sim \pi(p, k)$ for some $p \in (0, 1)^n$ and $k \in \{0, \dots, n\}$, then for any $A \subset \mathcal{B}_n$,*

$$\pi(A)\pi(d_T^2(\cdot, A)/84) \leq 1,$$

where

$$d_T(x, A) = \sup_{\alpha: |\alpha| \leq 1} d_\alpha(x, A) \quad \text{for } x \in \mathcal{B}_n, A \subset \mathcal{B}_n.$$

Let $\mathbb{M}_\pi f$ denote any median of f with respect to the measure π . A classical consequence of Theorem 3.1 is the following fact regarding the concentration around the median of convex functions [17]. Let us recall the classical observation that subgaussian concentration around median and mean for convex Lipschitz functions are equivalent up to the change of constants by a universal factor.

Corollary 3.2. *If $\pi \sim \pi(p, k)$ for some $p \in (0, 1)^n$ and $k \in \{0, \dots, n\}$, then for any convex function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ that is L -Lipschitz with respect to the standard Euclidean distance on \mathbb{R}^n and any $t > 0$,*

$$\pi(|f - \mathbb{M}_\pi f| > t) \leq 4 \exp(-t^2/84L^2).$$

Remark 3.3. If one is interested just in the lower tail of a convex function, then one can in fact replace the Lipschitz constant L by $\pi(|\nabla f|)$ or even certain quantiles of $|\nabla f|$. We do not pursue this direction here and refer the reader to [3].

Remark 3.4. If $f: \mathcal{B}_n \rightarrow \mathbb{R}$ is d_α 1-Lipschitz, then it can be extended to a function on \mathbb{R}^n which is $|\alpha|$ -Lipschitz with respect to the standard Euclidean distance. Therefore, Corollary 3.2 counterparts Theorem 2.3 in the sense that it yields the same concentration profile while allowing for a weaker Lipschitz condition on f at the cost of assuming convexity.

Our next result concerns concentration for matrix-valued functions under weaker assumptions than those in Theorem 2.5.

Theorem 3.5. *Let $\pi \sim \pi(p, k)$ for some $p \in (0, 1)^n$ and $k \in \{0, \dots, n\}$. Assume that $f: \mathcal{B}_n \rightarrow \mathcal{H}_d$ is such that there is a sequence of positive semidefinite matrices C_1, \dots, C_n satisfying*

$$(3.1) \quad (f(x) - f(x^i))^2 \preceq C_i^2, \quad \forall x \in \mathcal{B}_n, i \in [n],$$

where \preceq denotes the partial ordering of the set of positive semidefinite matrices. Define the variance proxy

$$\sigma^2 = 16 \sup \left\{ \left\| \sum_{i \in \mathcal{I}} C_i^2 \right\| : |\mathcal{I}| = k, \mathcal{I} \subset [n] \right\}.$$

Then for any $t > 0$,

$$\pi(\lambda_{\max}(f - \mu(f)) > t) \leq d \exp(-t^2/(\sigma^2 + \sigma t)).$$

Remark 3.6. Condition (3.1) implies that f is 1-Lipschitz with respect to the distance d_α with $\alpha_i = \|C_i\|$. On the other hand, for many choices of matrices C_1, \dots, C_n it happens that $\sigma^2 \ll \sum_{i=1}^k (\|C_i\|^2)^\downarrow$ as $n, k \rightarrow \infty$. Therefore, while yielding only sub-exponential concentration as opposed to the sub-Gaussian one given by Theorem 2.5, Theorem 3.5 may improve significantly on Theorem 2.5 through better parameters in the exponent.

Remark 3.7. By an adaptation of the proof of Theorem 3.5, one can obtain a similar result for general k -homogeneous measures satisfying the SCP condition with the variance proxy parameter

$$\sigma^2 = 8 \sup \left\{ \left\| \sum_{i \in \mathcal{I}} C_i^2 \right\| + k \max_{i \notin \mathcal{I}} \|C_i^2\| : |\mathcal{I}| \leq k, \mathcal{I} \subset [n] \right\}.$$

Finally, let us turn to the higher order concentration. By the Fourier–Walsh expansion (see e.g., [41]), every function $f: \mathcal{B}_n \rightarrow \mathbb{R}$ can be written in a unique way as a tetrahedral polynomial, i.e., a polynomial which is affine with respect to every variable (in particular the degree of the polynomial is at most n). Therefore in what follows we restrict our attention to this representation. In particular, when we speak about the gradient $\nabla f = (\partial_1 f, \dots, \partial_n f)$ or higher order derivatives $\nabla^k f$, we always think of the usual derivatives of the polynomial function on \mathbb{R}^n given by the tetrahedral representation of f (sometimes referred to as the harmonic extension of f). We remark that the directional derivatives $\partial_i f$ coincide on \mathcal{B}_n with the discrete derivatives of f given by $D_i f(x) = f(\max(x, x^i)) - f(\min(x, x^i))$, where the maximum and minimum are taken coordinatewise.

In order to formulate concentration of measure estimates for tetrahedral polynomials, we need to introduce a family of injective tensor product norms on d -index matrices (d -tensors). Let us recall the notation introduced by Latała in [35].

Let $|I|$ be the cardinality of a set I and for $\mathbf{i} = (i_1, \dots, i_d) \in [n]^d$ let $|\mathbf{i}| = \max_{j \leq d} i_j$ and $|\mathbf{i}_I| = \max_{j \in I} i_j$. Denote by P_d the set of partitions of $[d]$ into nonempty, pairwise disjoint sets. For a partition $\mathcal{I} = \{I_1, \dots, I_k\} \in P_d$, and a d -indexed matrix $A = (a_{\mathbf{i}})_{\mathbf{i} \in [n]^d}$, define

$$\|A\|_{\mathcal{I}} = \sup \left\{ \sum_{\mathbf{i} \in [n]^d} a_{\mathbf{i}} \prod_{l=1}^k x_{\mathbf{i}_{I_l}}^{(l)} : |(x_{\mathbf{i}_{I_l}}^{(l)})| \leq 1, 1 \leq l \leq k \right\},$$

where $|(x_{\mathbf{i}_{I_l}}^{(l)})| = \sqrt{\sum_{|\mathbf{i}_{I_l}| \leq n} x_{\mathbf{i}_{I_l}}^2}$. Therefore, for example,

$$\begin{aligned} \|(a_{ij})_{i,j \leq n}\|_{\{1,2\}} &= \sup \left\{ \sum_{i,j \leq n} a_{ij} x_{ij} : \sum_{i,j \leq n} x_{ij}^2 \leq 1 \right\} = \sqrt{\sum_{i,j \leq n} a_{ij}^2} = \|(a_{ij})_{i,j \leq n}\|_{HS}, \\ \|(a_{ij})_{i,j \leq n}\|_{\{1\}\{2\}} &= \sup \left\{ \sum_{i,j \leq n} a_{ij} x_i y_j : \sum_{i \leq n} x_i^2 \leq 1, \sum_{j \leq n} y_j^2 \leq 1 \right\} = \|(a_{ij})_{i,j \leq n}\|, \\ \|(a_{ijk})_{i,j,k \leq n}\|_{\{1,2\}\{3\}} &= \sup \left\{ \sum_{i,j,k \leq n} a_{ijk} x_{ij} y_k : \sum_{i,j \leq n} x_{ij}^2 \leq 1, \sum_{k \leq n} y_k^2 \leq 1 \right\}, \end{aligned}$$

where $\|\cdot\|_{HS}$ and $\|\cdot\|$ denote the Hilbert–Schmidt and the operator norm respectively.

Theorem 3.8. *If $\pi \sim \pi(p, k)$ for some $p \in (0, 1)^n$ and $k \in \{0, \dots, n\}$, then for any tetrahedral polynomial $f: \mathcal{B}_n \rightarrow \mathbb{R}$ of degree d ,*

$$\pi(|f - \pi(f)| \geq t) \leq 2 \exp \left(-\frac{1}{C_d} \min_{1 \leq r \leq d} \min_{\mathcal{J} \in P_r} \left(\frac{t}{\|\pi(\nabla^r f)\|_{\mathcal{J}}} \right)^{2/|\mathcal{J}|} \right),$$

where C_d is a constant depending only on the degree d of f .

Inequalities of this type for polynomials of arbitrary degree were introduced for the first time by Latała [35] for tetrahedral polynomials in i.i.d. standard Gaussian variables. Subsequently they were extended to general polynomials in independent subgaussian random variables and to certain dependent situations related to Glauber dynamics (see [4, 1, 47, 46, 2]). We remark that in the independent, subgaussian case and $d = 2$ they reduce to the well known Hanson–Wright inequality for quadratic forms, which has proved useful in non-asymptotic analysis of random matrices and in asymptotic geometric analysis (see, e.g., [55, Chapter 6]). It is worth mentioning that in the Gaussian case they may be reversed up to the value of the absolute constants, thus Theorem 3.8 shows that the measures $\pi(p, k)$ exhibit Gaussian type concentration for polynomials. While calculating the norms $\|\cdot\|_{\mathcal{J}}$ is usually difficult, estimating them is sometimes possible, leading to applications involving subgraph counts (in the Erdős–Rényi case or for some models of random graphs with dependencies [4, 47, 46]) or to statistical applications, e.g., in testing Ising models [23] and signal processing [56].

4. ABSTRACT FORMULATIONS

In this section we recall some notions from the theory of Markov semigroups and formulate the abstract counterparts of the results of Section 3 and of Theorem 2.3. We believe that

the results presented in this section might be of separate interest as they provide a general framework for proving concentration on the hypercube. We stress that most of the proof techniques that we exploit were known previously – our main contribution is the abstract formulation of these results by means of the novel stability condition (cf. Definition 4.3).

Throughout this section we will rely on the usual notions from the theory of Markov processes and Dirichlet forms specialized to finite state space. We will briefly recall them and refer to [39, 9, 13] for details.

4.1. Modified log-Sobolev inequalities. Let L be the generator of a jump Markov process on some finite probability space (M, π) . In what follows we will sometimes treat L as a linear operator on \mathbb{R}^M and sometimes identify it with the corresponding matrix, indexed by the elements of M .

Assume that L satisfies the detailed-balance condition

$$(4.1) \quad \forall x, y \in M \quad \pi(x)L(x, y) = \pi(y)L(y, x),$$

which implies that π is a stationary measure for the Markov process and L is self-adjoint on $L^2(\pi)$. In this article we consider only Markov processes satisfying the above condition, which may not be stated explicitly in all the results.

For a given L , we define $\Delta(L) := \max_x -L(x, x) = \max_x \sum_{y: y \neq x} L(x, y)$ and write $\mathcal{E}(f, g) = -\pi(fLg)$ for the Dirichlet form associated with L . In particular $\mathcal{E}(f, g) = \pi(\Gamma(f, g))$, where $\Gamma: \mathbb{R}^M \times \mathbb{R}^M \rightarrow \mathbb{R}^M$ given by

$$(4.2) \quad \Gamma(f, g)(x) = \frac{1}{2} \sum_{y \in M} (f(x) - f(y))(g(x) - g(y))L(x, y)$$

is the corresponding carré-du-champ operator. We use shorthand notation $\Gamma(f, f) =: \Gamma(f)$ and observe that by the detailed-balance condition (4.1) we have $\pi(\Gamma(f)) = \pi(\Gamma_+(f))$, where

$$(4.3) \quad \Gamma_+(f)(x) = \sum_{y \in M} (f(x) - f(y))_+^2 L(x, y).$$

Finally, we denote by $\rho(L)$ the best (the greatest) constant such that the following modified log-Sobolev inequality is satisfied

$$(4.4) \quad \rho(L) \text{Ent}_\pi(f) \leq \mathcal{E}(f, \log f)$$

for all non-constant functions $f: M \rightarrow [0, \infty)$, where $\text{Ent}_\pi(f) = \pi(f \log f) - \pi(f) \log \pi(f)$ is the entropy functional (we adopt the convention $0 \log 0 = 0$). We remark that $\rho(L)$ is positive iff L is irreducible on the support of π (see the discussion in [13] and [38, Chapter 12]). In what follows we will restrict our attention to this situation, without mentioning this assumption explicitly in each statement.

A classical observation, often referred to as Herbst's argument (cf. the monographs [36] by Ledoux and [17] by Boucheron et al.), says that for any $f: M \rightarrow \mathbb{R}$,

$$(4.5) \quad \pi(f > \pi(f) + t) \leq \exp(-t^2 \rho(L) / 4 \|\Gamma_+(f)\|_\infty),$$

where $\|\cdot\|_\infty$ stands for the norm in $L^\infty(\pi)$.

4.2. Flip-swap random walks. After Hermon and Salez [29], we say that a kernel L generates a flip-swap random walk if $L(x, y) > 0$ implies that $x = y^i$ for some $i \in [n]$ (i.e., x and y differ by a flip) or $x = y^{ij}$ for some $i \neq j, i, j \in [n]$ (i.e., x and y differ by a swap). The main contribution of [29] can be stated in the following way.

Theorem 4.1 (Hermon–Salez [29]). *For any measure π on \mathcal{B}_n satisfying the SCP, there exists a kernel L generating a reversible flip-swap random walk with stationary measure π such that $\rho(L) \geq 1$ and $\Delta(L) \leq n$. If π is also k -homogeneous, then $\Delta(L) \leq 2k$ as well.*

Theorem 4.1, by means of Herbst's argument (4.5), implies (up to an absolute constant in the exponent) the estimate from Theorem 1.1 after observing that for a flip-swap random walk and any $f: \mathcal{B}_n \rightarrow \mathbb{R}$ that is 1-Lipschitz with respect to the Hamming distance d_H

$$(4.6) \quad \|\Gamma_+(f)\|_\infty \leq \Delta(L) \cdot \max_{x, y \in \mathcal{B}_n} \{ (f(y) - f(x))_+^2 : L(x, y) > 0 \} \leq 4\Delta(L).$$

Finally, Theorem 4.1 is constructive – in Section 7 we revisit the inductive construction of L from [29] in the context of conditional Bernoulli distribution.

Remark 4.2. There are many examples of flip-swap random walks on the hypercube in the literature, including, e.g., the Bernoulli–Laplace model, Glauber dynamics or base exchange random walk on matroids, cf. e.g., [13, 28, 47, 22]. We note that the results of this section apply to any flip-swap random walk as long as we have control of its stability (cf. Definition 4.3) constant.

It turns out that for the proofs of all the statements of Section 3 it suffices to demonstrate that the following condition is true for some reversible generator L with stationary measure $\pi(p, k)$ for which the modified log-Sobolev inequality (4.4) is known.

Definition 4.3 (Stability condition). Let L be a generator of a flip-swap random walk on \mathcal{B}_n with invariant probability distribution π . We say that the pair (L, π) meets the stability condition with constant $R \geq 0$ (i.e., is R -stable) if it satisfies the modified log-Sobolev inequality (4.4) and

$$(4.7) \quad \max_{x \in \text{supp } \pi; i \in [n]} \sum_{y: y_i \neq x_i} L(x, y) \leq R\rho(L).$$

If it is clear from the context which measure π is associated with L , we will often omit it in the discussion and simply say that L is R -stable.

Remark 4.4. If π is not concentrated on a single point, then a random walk on \mathcal{B}_n with a generator L that satisfies the modified log-Sobolev inequality (4.4) may be at best 0.25-stable (i.e., $R \geq 0.25$). Indeed, in this case there exists i such that $\pi(\{x_i = 1\}), \pi(\{x_i = 0\}) > 0$. If L satisfies the modified log-Sobolev inequality, then it also satisfies the Poincaré inequality $\frac{1}{2}\rho(L) \text{Var}_\pi(f) \leq \mathcal{E}(f, f)$, see e.g., [2, Proposition B.5]. Therefore, by the stability condition (4.7) applied to the function $f(x) = \mathbf{1}_{\{x_i=1\}}$ and reversibility of L we get that

$$\begin{aligned} R\rho(L)\pi(\{x_i = 1\}) &\geq \sum_{x: x_i=1} \sum_{y: y_i=0} L(x, y)\pi(x) \\ &= \sum_{x, y} (x_i - y_i)_+^2 L(x, y)\pi(x) \\ &= \mathcal{E}(f, f) \\ &\geq \frac{1}{2}\rho(L) \text{Var}_\pi(f) = \frac{1}{2}\rho(L)\pi(\{x_i = 1\})\pi(\{x_i = 0\}), \end{aligned}$$

which gives $R \geq 0.5 \cdot \pi(\{x_i = 0\})$. Similarly, by considering $f(x) = \mathbf{1}_{\{x_i=0\}}$ we get that $R \geq 0.5 \cdot \pi(\{x_i = 1\})$ as well, yielding $R \geq 0.25$.

This bound is optimal, as can be seen for π being the uniform measure on \mathcal{B}_n and $L(x, y) = 1$ if there exists i such that $y = x^i$, $L(x, y) = -n$ if $y = x$ and $L(x, y) = 0$ otherwise (this corresponds to the special case of Glauber dynamics, in which at rate n , a random coordinate is flipped). In this case $\rho(L) = 4$ (see [13, Example 3.7], note a different normalization of both the Dirichlet form and the constant in the modified log-Sobolev inequality), whereas for all $x \in \mathcal{B}_n$

$$\max_i \sum_{y: y_i \neq x_i} L(x, y) = L(x, x^i) = 1 = 0.25 \cdot \rho(L).$$

Let us illustrate the notion of R -stability with another classical example.

Example 4.5 (Bernoulli–Laplace model). Let π be the uniform measure on the slice of \mathcal{B}_n consisting of elements with exactly k ones and let L be given by $Lf(x) = \frac{1}{n} \sum_{i < j} (f(x^{ij}) - f(x))$ (thus the corresponding Markov process at rate $(n-1)/2$ swaps a uniformly chosen pair of coordinates). In the matrix form this corresponds to $L(x, y) = \frac{1}{n}$ if $x \neq y$ and $y = x^{ij}$, $L(x, x) = -k(n-k)/n$ and $L(x, y) = 0$ otherwise. It has been proved in [25] and independently in [13] that $\rho_0(L) \geq 1/2$. At the same time $\sum_{y: y_i \neq x_i} L(x, y)$ equals to $(n-k)/n$ if $x_i = 1$ and to k/n otherwise. This shows that L is 2-stable, independently of n and k . As mentioned in the introduction, the uniform measure on the slice of the discrete cube can be interpreted as the distribution of i.i.d. Bernoulli variables conditioned on their sum being equal to k . In Proposition 7.3 we generalize the above observation on stability and show that if μ is the law of general independent Bernoulli variables conditioned on their sum being equal to a fixed constant, there exists a 2-stable generator of a random walk reversible with respect to μ .

Remark 4.6. Observe that the notion of stability is invariant under scaling of L (change of time), i.e., if L is R -stable then so is aL for any $a > 0$. This leads to a tensorization

property for measures admitting an R -stable generator. More precisely, let π_1, \dots, π_m be measures on $\mathcal{B}_{n_1}, \dots, \mathcal{B}_{n_m}$, for which there exist reversible flip-swap random walks with R -stable generators L_1, \dots, L_m . By changing time, we can assume without loss of generality that $\rho(L_i) = \rho$ for all $i \leq m$. Let $n = n_1 + \dots + n_m$ and consider the product measure $\pi = \pi_1 \otimes \dots \otimes \pi_m$ on \mathcal{B}_n together with the generator $L = L_1 + \dots + L_m$, where we think of L_i as acting only on the i -th block of coordinates on $\mathcal{B}_n = \mathcal{B}_{n_1} \times \dots \times \mathcal{B}_{n_m}$, i.e., we identify L_i with its tensor product with identity on $\otimes_{j \neq i} \mathbb{R}^{\mathcal{B}_{n_j}}$. In the matrix form we have the representation

$$L(x, y) = \sum_{i=1}^m L_i(P_i x, P_i y) \prod_{j \neq i} \mathbf{1}_{\{P_j x = P_j y\}},$$

where $P_j: \mathcal{B}_n \rightarrow \mathcal{B}_j$ is the projection onto the j -th factor in the product $\mathcal{B}_n = \mathcal{B}_{n_1} \times \dots \times \mathcal{B}_{n_m}$. Thanks to the well known tensorization property of the entropy (see, e.g., [7, Chapter 3]) we have $\rho(L) = \rho$. Moreover, for $i \in (n_1 + \dots + n_{j-1}, n_1 + \dots + n_j]$,

$$\sum_{y \in \mathcal{B}_n: y_i \neq x_i} L(x, y) = \sum_{y \in \mathcal{B}_{n_j}: y_l \neq (P_j x)_l} L_j((P_j x), y) \leq R\rho,$$

where $l = i - (n_1 + \dots + n_{j-1})$. Thus L is indeed R -stable.

This observation allows in particular to extend all the theorems of Section 3 to product of measures $\pi(n, k)$ allowing for more general conditioning of Bernoulli variables.

4.3. Abstract formulations. Finally, let us present the counterparts of the results of Section 2 and of Theorem 2.3 from Section 3 in the abstract language of the stability condition (4.7). We stress here that it is the sole property needed for their proofs, which are deferred to Section 7.

We start with a bounded-difference type inequality for real valued functions.

Proposition 4.7. *If a flip-swap random walk on \mathcal{B}_n with stationary distribution π and generator L satisfies the stability condition (4.7), then for any $f: \mathcal{B}_n \rightarrow \mathbb{R}$ and $\alpha \in \mathbb{R}_+^n$ such that*

$$|f(x) - f(y)| \leq d_\alpha(x, y) \quad \forall x, y \in \mathcal{B}_n$$

the following estimate holds for all $t > 0$

$$\pi(f > \pi(f) + t) \leq \exp\left(-\frac{t^2}{8R|\alpha|^2}\right).$$

In the above estimate one can also replace $8|\alpha|^2$ with $16 \sum_{i=1}^n [\Delta(L)/R\rho(L)] (\alpha_i^\downarrow)^2$.

Remark 4.8. Using the definitions of R -stability and of $\Delta(L)$ one can see that $\Delta(L)/R\rho(L) \leq n$ and if π is k -homogeneous, then $\Delta(L)/R\rho(L) \leq k$.

Let us now pass to the matrix-valued case.

Proposition 4.9. *Let a flip-swap random walk on \mathcal{B}_n with stationary distribution π and generator L satisfy the stability condition (4.7). Assume also that $f: \mathcal{B}_n \rightarrow \mathcal{H}_d$ is such that there is a sequence of positive semidefinite matrices C_1, \dots, C_n satisfying*

$$(4.8) \quad (f(x) - f(x^i))^2 \preceq C_i^2 \quad \forall x \in \mathcal{B}_n, i \in [n],$$

where \preceq denotes the positive semidefinite order on the set of symmetric matrices. Set the variance proxy

$$\sigma^2 = 8R \cdot \sup \left\{ \left\| \sum_{i \in \mathcal{I}} C_i^2 \right\| : |\mathcal{I}| = \lceil \Delta(L)/R\rho(L) \rceil, \mathcal{I} \subset [n] \right\}.$$

Then for any $t > 0$,

$$\pi(\lambda_{\max}(f - \pi(f)) > t) \leq d \exp\left(-t^2/(\sigma^2 + \sigma t)\right).$$

Our next proposition is the convex distance inequality under R -stability.

Proposition 4.10. *If a flip-swap random walk on \mathcal{B}_n with some stationary distribution π and a generator L satisfies the stability condition (4.7), then for any set $A \subset \mathcal{B}_n$*

$$\pi(A) \pi\left(\exp\left(\frac{1}{40R+4} \cdot d_T^2(\cdot, A)\right)\right) \leq 1.$$

Finally, let us state the concentration result for polynomials in an abstract version.

Proposition 4.11. *If a flip-swap random walk on \mathcal{B}_n with some stationary distribution π and a generator L satisfies the stability condition (4.7), then for any tetrahedral polynomial $f: \mathcal{B}_n \rightarrow \mathbb{R}$ of degree d*

$$\pi(|f - \pi(f)| \geq t) \leq 2 \exp\left(-\frac{1}{C_d} \min_{1 \leq r \leq d} \min_{\mathcal{J} \in P_r} \left(\frac{t}{R^{r/2} \|\pi(\nabla^r f)\|_{\mathcal{J}}}\right)^{2/|\mathcal{J}|}\right),$$

where C_d is a constant depending only on the degree d of f .

Remark 4.12. Although Proposition 4.7 gives a worse constant in the exponent than Theorem 2.3 even in the case of conditional Bernoulli distributions $\pi(p, k)$, we state it here as in principle it does not assume that π satisfies the SCP and thus potentially can be applied in other settings.

Remark 4.13. The above propositions can be transferred to more general random walks that change at each step at most a fixed number of coordinates N (with $N = 2$ in case of flip-swap random walks). We do not pursue this direction though and do not write all the theorems in full generality for the sake of readability.

Remark 4.14. Recently, Cryan et al. [22] have shown a version of Theorem 4.1 for k -homogeneous strongly log-concave measures. Strong log-concavity is yet another possible generalization of the SRP, which is in general incomparable with the SCP [22]. It is known, cf. Brändén and Huh [18], that any k -homogeneous strongly log-concave measure is supported on the set of bases of some matroid of rank k . Using this fact, and extending the previous results for uniform measures on the bases of matroids by Anari et al. [6] and Kaufman and Oppenheim [33], Cryan et al. [22] explicitly construct a base-exchange random walk, which has any given strongly log-concave measure as a stationary distribution, and verify it satisfies the modified log-Sobolev inequality (4.4).

Since the base-exchange random walk proposed therein is a particular instance of a flip-swap random walk, a natural question is whether it satisfies the stability condition (4.7), which would allow to deduce concentration results presented in this section. Unfortunately, the answer seems to be negative in full generality as can be seen already in the case of independent Bernoulli random variables $B = (B_1, \dots, B_n)$ with different probabilities of success $\mathbb{P}(B_i = 1) = p_i$ conditioned on their sum being k , i.e., for the distribution $\pi(p, k) \sim \mathcal{L}(B | \sum_i B_i = k)$. If one chooses $p_1 \rightarrow 1^-$ and $p_j = c$ for $j > 1$ and some $c \in (0, 1)$, then it is straightforward to verify that the base-exchange random walk of [22] is at best k -stable. Therefore, applying propositions of Section 4.3 to the base-exchange random walk gives much worse concentration constants than those of Section 3. On the other hand, the flip-swap random walk proposed by Hermon and Salez [29] with stationary measure $\pi(p, k)$, which we use to prove the results of Section 3, turns out to be 2-stable.

In view of the above, it is an interesting problem to analyze what other known kernels satisfy the stability condition (4.7) with good (dimension-independent) constant and to look for some other criteria that would allow to deduce this condition.

5. PROOFS OF THE RESULTS OF SECTION 2

In this section we provide proofs of Theorems 2.3, 2.5 and 2.8. All these results follow by modifications of the martingale argument due to Pemantle and Peres [44].

Let $X \sim \pi$ be a random variable with values in \mathcal{B}_n satisfying the SCP and denote $\text{supp } X = \{i \in \{1, \dots, n\} : X_i = 1\}$. In the non-homogeneous case define a filtration $\mathcal{F} = (\mathcal{F}_l)_{l=0}^n$ by letting simply $\mathcal{F}_0 = \{\emptyset, \Omega\}$ and $\mathcal{F}_l = \sigma(X_1, \dots, X_l)$ for $l = 1, \dots, n$. In the k -homogeneous case introduce a family of random variables Y_1, \dots, Y_k given by the conditions

$$(5.1) \quad \begin{aligned} \mathcal{L}(Y_1 | X) &= \text{Unif}(\text{supp } X \setminus \{1, \dots, k\}) \quad \text{and} \\ \mathcal{L}(Y_l | X, Y_1, \dots, Y_{l-1}) &= \text{Unif}(\text{supp } X \setminus \{1, \dots, k, Y_1, \dots, Y_{l-1}\}), \quad \text{for } l = 2, \dots, k, \end{aligned}$$

where $\text{Unif}(A)$ stands for the uniform distribution on the set A , and for notational simplicity we set $\text{Unif}(\emptyset)$ to be the Dirac mass at 0 and $X_0 \equiv 1$ (i.e., we add to X an additional coordinate providing no information and if the above sampling scheme yields all elements from $\text{supp } X$ before sampling some Y_l , we set Y_l to zero for all $i \geq l$). Finally, define a filtration $\mathcal{G} = (\mathcal{G}_l)_{l=0}^{2k}$ setting $\mathcal{G}_0 = \{\emptyset, \Omega\}$ and $\mathcal{G}_l = \sigma(X_1, \dots, X_l)$ for $l \in [k]$, $\mathcal{G}_{k+r} = \sigma(X_1, \dots, X_k, Y_1, \dots, Y_r)$ for $r \in [k]$.

In other words in the first k -steps the subsequent values of X at the first k coordinates are revealed, while in the last k steps one reveals in a uniformly random order the remaining coordinates at which X takes the value 1. Note that if α is nonincreasing (which we may

assume without loss of generality) and f is 1-Lipschitz with respect to d_α then the first part of this sampling scheme promotes the coordinates which may have the greatest impact on the value of $f(X)$. The construction can be thought of as a modification of the sampling scheme proposed by Pemantle and Peres in which one immediately starts revealing in a random order the coordinates at which X takes the value 1.

The proof of Theorems 2.3 and 2.5 will be based on the following two lemmas.

Lemma 5.1. *Let $\alpha \in \mathbb{R}_+^n$ be nonincreasing and let $f: \mathcal{B}_n \rightarrow \mathcal{H}_d$ be 1-Lipschitz with respect to the distance d_α . Assume that X is a \mathcal{B}_n -valued random vector satisfying the SCP. Let $M_l = \mathbb{E}[f(X) | \mathcal{F}_l] - \mathbb{E}[f(X) | \mathcal{F}_{l-1}]$ for $l \in [n]$. Then for every $l \in [n]$,*

$$(5.2) \quad M_l^2 \preceq 4\alpha_l^2 I_d.$$

Lemma 5.2. *In the setting of Lemma 5.1 assume additionally that X is k -homogeneous. For $l \in [2k]$ define $N_l = \mathbb{E}[f(X) | \mathcal{G}_l] - \mathbb{E}[f(X) | \mathcal{G}_{l-1}]$. Then for $l \in [k]$,*

$$(5.3) \quad N_l^2 \preceq 4\alpha_l^2 I_d,$$

while for $l = k+1, \dots, 2k$,

$$(5.4) \quad N_l^2 \preceq 4\alpha_k^2 I_d.$$

We postpone for now the proof of the above lemmas and firstly show how they imply Theorems 2.3 and 2.5. To this end let us recall the matrix version of the Azuma-Hoeffding inequality due to Tropp [54, Theorem 7.1], which asserts that if D_l , $l = 1, \dots, n$ are \mathcal{H}_d -valued martingale differences and $D_l^2 \preceq C_l^2$ for some deterministic matrices $C_l \in \mathcal{H}_d$, then for all $t \geq 0$,

$$\mathbb{P}\left(\lambda_{\max}\left(\sum_{l=1}^n D_l\right) \geq t\right) \leq de^{-t^2/8\sigma^2},$$

where $\sigma^2 = \|\sum_{l=1}^n C_l^2\|$. Note also that for $d = 1$ the classical Azuma-Hoeffding inequality (see, e.g., [24, Theorem 5.8]) allows to replace the constant $1/8$ by $1/2$.

Proof of Theorems 2.3 and 2.5. Since the SCP is invariant under permutations of coordinates of X , we may and do assume that $\alpha = \alpha^\downarrow$. By Lemma 5.1 the martingale differences M_l satisfy $M_l^2 \preceq C_l^2 := 4\alpha_l^2 I_d$. Clearly

$$(5.5) \quad \left\| \sum_{l=1}^n C_l^2 \right\| = 4|\alpha|^2.$$

If X is k -homogeneous, then by Lemma 5.2, $N_l^2 \preceq \tilde{C}_l^2 := 4\alpha_{\min(l,k)}^2 I_d$. In this case

$$(5.6) \quad \left\| \sum_{l=1}^{2k} \tilde{C}_l^2 \right\| = 4\left[\left(\sum_{l=1}^k \alpha_l^2\right) + k\alpha_k^2\right] \leq 8\sum_{l=1}^k \alpha_l^2.$$

We have $f(X) = \sum_{l=1}^n M_l$, whereas in the k -homogeneous case $f(X) = \sum_{l=1}^{2k} N_l$ (observe that after $2k$ -steps of the sampling procedure all the nonzero coordinates of X are revealed and so X is \mathcal{G}_{2k} -measurable). Thus the conclusion of Theorem 2.3 follows by applying estimates (5.5) and (5.6) for $d = 1$ together with the classical Azuma-Hoeffding inequality. Similarly, Theorem 2.5 follows from the matrix version of the Azuma-Hoeffding inequality. \square

It remains to prove Lemmas 5.1 and 5.2.

Proof of Lemma 5.1. Let $A_l^x = \{X_1 = x_1, \dots, X_l = x_l\}$ for $x = (x_1, \dots, x_n) \in \mathcal{B}_n$ and $l = 0, \dots, n$. Then, for $l = 1, \dots, n$ and any $x \in \mathcal{B}_n$ such that $\mathbb{P}(A_l^x) > 0$,

$$\begin{aligned} \mathbb{E}[f(X) | A_l^x] - \mathbb{E}[f(X) | A_{l-1}^x] &= \mathbb{E}[f(X) | A_{l-1}^x, X_l = x_l] - \mathbb{E}[f(X) | A_{l-1}^x] \\ &= \mathbb{P}(X_l \neq x_l | A_{l-1}^x) (\mathbb{E}[f(X) | A_{l-1}^x, X_l = x_l] - \mathbb{E}[f(X) | A_{l-1}^x, X_l \neq x_l]). \end{aligned}$$

If $\mathbb{P}(X_l \neq x_l | A_{l-1}^x) \neq 0$, then by the SCP there exist a coupling (\hat{X}, \hat{Y}) between the distributions $\mathcal{L}(X | A_{l-1}^x, X_l = x_l)$ and $\mathcal{L}(X | A_{l-1}^x, X_l \neq x_l)$ that is supported on the set $\{(y, z) \in \mathcal{B}_n^2 : d_H((y_i)_{i>l}, (z_i)_{i>l}) \leq 1\}$. Using this coupling, the Lipschitz property of f , Jensen's inequality and the fact that $\alpha_i \leq \alpha_l$ for any $i > l$, we get that

$$\begin{aligned} \|\mathbb{E}[f(X) | A_l^x] - \mathbb{E}[f(X) | A_{l-1}^x]\| &\leq \mathbb{P}(X_l \neq x_l | A_{l-1}^x) \mathbb{E}\|f((x_i)_{i \leq l}, \hat{X}_{i>l}) - f((x_i)_{i \leq l}, 1 - x_l, \hat{Y}_{i>l})\| \\ &\leq \mathbb{P}(X_l \neq x_l | A_{l-1}^x) \cdot 2\alpha_l \leq 2\alpha_l, \end{aligned}$$

which is equivalent to (5.2). \square

Proof of Lemma 5.2. Note that for $l \leq k$, we have $\mathcal{G}_l = \mathcal{F}_l$. As a consequence $N_l = M_l$, where M_l are martingale increments defined in Lemma 5.1, which implies (5.3).

Consider now $l > k$ of the form $l = k + r$ and for $x = (x_1, \dots, x_k) \in \mathcal{B}_k$ and $v = (v_1, \dots, v_k) \in (\{0\} \cup \{k+1, \dots, n\})^k$ set $A_l^{x,v} = \{X_1 = x_1, \dots, X_k = x_k, Y_1 = v_1, \dots, Y_r = v_r\}$. Then \mathcal{F}_l is generated by the sets $A_l^{x,v}$. By the definition of the variables Y_r , we have $\{Y_r = i\} \subseteq \{X_i = 1\}$ and so for any x, v such that $\mathbb{P}(A_l^{x,v}) > 0$,

$$(5.7) \quad \mathbb{E}[f(X) | A_l^{x,v}] = \frac{\mathbb{E}[f(X) \mathbf{1}_{A_{l-1}^{x,v}} \mathbf{1}_{\{X_{v_r}=1\}} \mathbf{1}_{\{Y_r=v_r\}}]}{\mathbb{P}(A_{l-1}^{x,v}, X_{v_r} = 1, Y_r = v_r)}.$$

For $s \in [r]$ let $m_s = |\{i \in [k]: x_i = 1\}| + |\{j \in [s-1]: v_j \neq 0\}|$ be the number of ones sampled by the time $k+s-1$. It follows from (5.1) that if $m_s < k$ then $\mathbb{P}(A_{k+s}^{x,v}) > 0$ implies that $v_s \neq 0$ and $\mathbb{P}(Y_s = v_s | X, Y_1, \dots, Y_{s-1}) = \frac{1}{k-m_s}$ on $A_{k+s-1}^{x,v} \cap \{X_{v_s} = 1\}$, whereas if $m_s = k$, then $\mathbb{P}(A_{k+s}^{x,v}) > 0$ implies that $v_s = 0$ and $\mathbb{P}(Y_s = v_s | X, Y_1, \dots, Y_{s-1}) = 1$ on $A_{k+s-1}^{x,v} \cap \{X_{v_s} = 1\} = A_{k+s-1}^{x,v}$. Going back to (5.7) and using this observation for $s = r, \dots, 1$, we obtain that

$$\mathbb{E}[f(X) | A_l^{x,v}] = \mathbb{E}[f(X) | B_l^{x,v}],$$

where $B_l^{x,v} = \{X_1 = x_1, \dots, X_k = x_k, X_{v_1} = \dots = X_{v_{l-k}} = 1\}$. We thus obtain

$$\begin{aligned} & \mathbb{E}[f(X) | A_l^{x,v}] - \mathbb{E}[f(X) | A_{l-1}^{x,v}] \\ &= \mathbb{P}(X_{v_r} \neq 1 | B_{l-1}^{x,v}) (\mathbb{E}[f(X) | B_{l-1}^{x,v}, X_{v_r} = 1] - \mathbb{E}[f(X) | B_{l-1}^{x,v}, X_{v_r} \neq 1]). \end{aligned}$$

Note that the right-hand side may be non-zero only if $v_r \neq 0$. In this case using the inequality $\alpha_{v_s} \leq \alpha_k$ for $s \in [k]$ we can conclude as in the proof of Lemma 5.1. \square

Let us now pass to the proof of Theorem 2.8.

Proof of Theorem 2.8. We will rely on the martingale used in the article by Pemantle and Peres. Let X be a random vector with law π and define the random variables Y_l for $l \leq n$ as

$$(5.8) \quad \begin{aligned} \mathcal{L}(Y_1 | X) &= \text{Unif}(\text{supp } X) \quad \text{and} \\ \mathcal{L}(Y_l | X, Y_1, \dots, Y_{l-1}) &= \text{Unif}(\text{supp } X \setminus \{Y_1, \dots, Y_{l-1}\}), \quad \text{for } l = 2, \dots, k, \end{aligned}$$

i.e., Y_l 's reveal in a uniformly random order the elements of $\text{supp } X$. Let $\mathcal{H}_0 = \{\emptyset, \Omega\}$ and $\mathcal{H}_l = \sigma(Y_1, \dots, Y_l)$ for $l = 1, \dots, k$. Then $f(X) - \mathbb{E}f(X) = \sum_{l=1}^k \mathbb{E}[f(X) | \mathcal{H}_l] - \mathbb{E}[f(X) | \mathcal{H}_{l-1}] =: \sum_{l=1}^k D_l$. We will use the matrix version of Freedman's inequality due to Tropp [53], which asserts (in a version specialized for our application) that if $\|D_l\| \leq a$ a.s. for all l , and $\|\sum_{l=1}^k \mathbb{E}[D_l^2 | \mathcal{H}_{l-1}]\| \leq \sigma^2$ a.s., then for any $t \geq 0$,

$$(5.9) \quad \mathbb{P}(\|f(X) - \mathbb{E}f(X)\| \geq t) \leq 2d \exp\left(-\frac{t^2}{2\sigma^2 + 2at/3}\right).$$

Consider thus a sequence of pairwise distinct $v_1, \dots, v_k \in [n]$ and denote $A_l^v = \{Y_1 = v_1, \dots, Y_l = v_l\}$. Similarly as in the proof of Lemma 5.2, if $\mathbb{P}(A_l^v) > 0$, then we have

$$\mathbb{E}[f(X) | A_l^v] = \mathbb{E}[f(X) | B_l^v],$$

where $B_l^v = \{X_{v_1} = \dots = X_{v_l} = 1\}$. Therefore we have

$$(5.10) \quad D_l \mathbf{1}_{A_l^v} = \mathbb{P}(X_{v_l} = 0 | B_{l-1}^v) (\mathbb{E}[f(X) | B_{l-1}^v, X_{v_l} = 1] - \mathbb{E}[f(X) | B_{l-1}^v, X_{v_l} = 0]) \mathbf{1}_{A_l^v}.$$

Since the SRP implies the SCP, there exists a coupling (\tilde{Z}, \hat{Z}) between the distributions $\mathcal{L}(X | B_l^v)$ and $\mathcal{L}(X | B_{l-1}^v, X_{v_l} = 0)$ such that \tilde{Z} and \hat{Z} differ just at coordinate v_l and one additional coordinate (at which by k -homogeneity \tilde{Z} necessarily takes the value one). Let \tilde{Y}_l be this coordinate. We have

$$(5.11) \quad \mathbb{E}[f(X) | B_{l-1}^v, X_{v_l} = 1] - \mathbb{E}[f(X) | B_{l-1}^v, X_{v_l} = 0] = \mathbb{E}[f(\tilde{Z}) - f(\hat{Z})],$$

Since $\hat{Z}^{\tilde{Y}_l} = \tilde{Z}^{v_l}$, we have

$$(5.12) \quad \left(\mathbb{E}[f(X) | B_{l-1}^v, X_{v_l} = 1] - \mathbb{E}[f(X) | B_{l-1}^v, X_{v_l} = 0] \right)^2 = \left(\mathbb{E}[f(\tilde{Z}) - f(\hat{Z})] \right)^2 \\ \preceq \mathbb{E} \left[(f(\tilde{Z}) - f(\hat{Z}))^2 \right] = \mathbb{E} \left[(f(\tilde{Z}) - f(\tilde{Z}^{v_l}) + f(\hat{Z}^{\tilde{Y}_l}) - f(\hat{Z}))^2 \right] \\ \preceq 2 \mathbb{E} \left[(f(\tilde{Z}) - f(\tilde{Z}^{v_l}))^2 \right] + 2 \mathbb{E} \left[(f(\hat{Z}^{\tilde{Y}_l}) - f(\hat{Z}))^2 \right] \preceq 2C_{v_l}^2 + 2\mathbb{E}C_{\tilde{Y}_l}^2,$$

where in the first and second inequality we used the operator convexity of the function $x \mapsto x^2$ (see [11, Example V.1.3]), and in the last inequality the assumption (2.3).

In particular, using (5.10), we obtain $\|D_l^2\| \leq 4 \max_i \|C_i^2\|$, so $\|D_l\| \leq 2K$. Moreover as on A_l^v we have $Y_l = v_l$, by (5.10) and (5.12) we get that

$$D_l^2 \mathbf{1}_{A_l^v} \preceq 2(C_{Y_l}^2 + \mathbb{E}C_{\tilde{Y}_l}^2) \mathbb{P}(X_{v_l} = 0 | B_{l-1}^v)^2 \mathbf{1}_{A_l^v}.$$

Let us now slightly change our notation and think of \tilde{Y}_l as of random variable defined on the same probability space as X , with conditional distribution with respect to the σ -field \mathcal{H}_l given on each of its atoms A_l^v by the above construction, using the corresponding coupling (which depends on v_1, \dots, v_{l-1}). Then the above inequality can be written as

$$(5.13) \quad D_l^2 \preceq 2 \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} (C_{\tilde{Y}_l}^2 + \mathbb{E}[C_{\tilde{Y}_l}^2 | A_l^v]) \mathbb{P}(X_{v_l} = 0 | B_{l-1}^v)^2 \mathbf{1}_{A_l^v}.$$

Let us now go back to the equations (5.10) and (5.11) and let us apply them in the special case of the function $\tilde{f}(x) = \sum_{i=1}^n x_i C_i^2$, denoting the corresponding martingale increment by \tilde{D}_l . We obtain that

$$\tilde{D}_l \mathbf{1}_{A_l^v} = \mathbb{P}(X_{v_l} = 0 | B_{l-1}^v) (C_{Y_l}^2 - \mathbb{E}[C_{Y_l}^2 | A_l^v]) \mathbf{1}_{A_l^v}.$$

Thus we get that

$$0 = \mathbb{E}[\tilde{D}_l | A_{l-1}^v] = \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} \mathbb{E} \left[\mathbb{P}(X_{v_l} = 0 | B_{l-1}^v) \mathbf{1}_{A_l^v} (C_{Y_l}^2 - \mathbb{E}[C_{Y_l}^2 | A_l^v]) \mid A_{l-1}^v \right],$$

i.e.,

$$\sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} \mathbb{E} \left[\mathbb{P}(X_{v_l} = 0 | B_{l-1}^v) \mathbf{1}_{A_l^v} C_{Y_l}^2 \mid A_{l-1}^v \right] \\ = \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} \mathbb{E} \left[\mathbb{P}(X_{v_l} = 0 | B_{l-1}^v) \mathbf{1}_{A_l^v} \mathbb{E}[C_{Y_l}^2 | A_l^v] \mid A_{l-1}^v \right],$$

which combined with the estimate (5.13) on D_l^2 (replacing $\mathbb{P}(X_{v_l} = 0 | B_{l-1}^v)^2$ by $\mathbb{P}(X_{v_l} = 0 | B_{l-1}^v)$) gives

$$\mathbb{E}[D_l^2 | A_{l-1}^v] \preceq 2 \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} \mathbb{E} \left[(C_{Y_l}^2 + \mathbb{E}[C_{Y_l}^2 | A_l^v]) \mathbb{P}(X_{v_l} = 0 | B_{l-1}^v) \mathbf{1}_{A_l^v} \mid A_{l-1}^v \right] \\ = 4 \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} \mathbb{E} \left[C_{Y_l}^2 \mathbb{P}(X_{v_l} = 0 | B_{l-1}^v) \mathbf{1}_{A_l^v} \mid A_{l-1}^v \right] \\ \preceq 4 \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} C_{v_l}^2 \mathbb{P}(A_l^v | A_{l-1}^v) \\ = 4 \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} C_{v_l}^2 \frac{1}{k-l+1} \mathbb{P}(X_{v_l} = 1 | B_{l-1}^v) \\ \preceq 4 \sum_{v_l \in [n] \setminus \{v_1, \dots, v_{l-1}\}} C_{v_l}^2 \frac{1}{k-l+1} \mathbb{P}(X_{v_l} = 1),$$

where in the last inequality we used [34, Lemma 1.10], which asserts that $\mathbb{P}(X_{v_l} = 1) \geq \mathbb{P}(X_{v_l} = 1 | B_{l-1}^v)$ (we remark that this is the only place in the proof in which we use the full strength of the strong Rayleigh property).

Extending the summation to $[n]$, we thus obtain

$$\mathbb{E}[D_l^2 | \mathcal{H}_{l-1}] \preceq 4 \sum_{v=1}^n C_v^2 \mathbb{P}(X_v = 1) \frac{1}{k-l+1},$$

whence

$$\sum_{l=1}^k \mathbb{E}[D_l^2 | \mathcal{H}_{l-1}] \preceq 4 \sum_{v=1}^n C_v^2 \mathbb{P}(X_v = 1) \log(ek) \preceq 4 \log(ek) \cdot \mathbb{E} \left[\sum_{v=1}^n X_v C_v^2 \right].$$

Combining this with the already obtained bound on $\|D_l\|$ allows us to apply (5.9) with $a = 2K$ and $\sigma^2 = 4\|\mathbb{E}\sum_{v=1}^n X_v C_v^2\| \log(ek)$, which ends the proof of the theorem. \square

6. PROOFS OF THE RESULTS OF SECTION 4

6.1. Propositions 4.7 and 4.9. The main idea behind the proof of Proposition 4.7 is to find an estimate on $\|\Gamma_+(f)\|_\infty$ in terms of α , refining (4.6), and then to use the Herbst argument. We will need the following lemma which we state in the matrix setting as it will be useful for the proof of Proposition 4.9 as well.

Lemma 6.1. *Let $t = (t_1, \dots, t_n)$ be a sequence of nonnegative numbers and let $D_1, \dots, D_n \in \mathcal{H}_d$ be positive semidefinite matrices. Then for any $T_1 \geq |t|_1$ and $T_\infty \geq |t|_\infty$*

$$(6.1) \quad \left\| \sum_{i=1}^n t_i D_i \right\| \leq T_\infty \cdot \sup \left\{ \left\| \sum_{i \in \mathcal{I}} D_i \right\| : \mathcal{I} \subset [n], |\mathcal{I}| \leq \lceil T_1/T_\infty \rceil \right\}.$$

Proof. By homogeneity we may assume without loss of generality that $T_\infty = 1$. We may also assume that T_1 is a positive integer. Let

$$\mathcal{X} = \left\{ x \in [0, 1]^n : \sum_{i=1}^n x_i \leq T_1 \right\}, \quad \mathcal{Y} = \left\{ y \in \{0, 1\}^n : \sum_{i=1}^n y_i \leq T_1 \right\}.$$

Since the right-hand side of (6.1) equals to $\max\{\|\sum_{i=1}^n y_i D_i\| : y \in \mathcal{Y}\}$, whereas the left-hand side is a convex function of t , the lemma will follow once we prove that $\mathcal{X} \subset \text{conv } \mathcal{Y}$. To this end, by the Krein–Milman theorem, it is enough to show that \mathcal{Y} is the set of all extreme points of the closed convex set \mathcal{X} . Consider any $x \in \mathcal{X} \setminus \mathcal{Y}$. Let $i_0 \in [n]$ be such that $x_{i_0} \in (0, 1)$. If $\sum_i x_i < T_1$ then for ε sufficiently close to zero, $x + \varepsilon e_{i_0}, x - \varepsilon e_{i_0} \in \mathcal{X}$ and so $x = \frac{1}{2}(x + \varepsilon e_{i_0}) + \frac{1}{2}(x - \varepsilon e_{i_0})$ is not an extreme point of \mathcal{X} . If $\sum_i x_i = T_1$, then since T_1 is an integer, there exists $i_1 \neq i_0$ such that $x_{i_1} \in (0, 1)$. Then $x = \frac{1}{2}u + \frac{1}{2}v$, where $u = x + \varepsilon e_{i_0} - \varepsilon e_{i_1}, v = x - \varepsilon e_{i_0} + \varepsilon e_{i_1}$. For ε close to zero $u, v \in \mathcal{X}$, thus again, x is not an extreme point. \square

Proof of Proposition 4.7. We recall that for $x \in \mathcal{B}_n$ and $i, j \in [n]$, x^i and x^{ij} denote the vectors obtained from x by flipping the i -th and swapping the i -th and j -th coordinates respectively. For any $x \in \mathcal{B}_n$, using the definition (4.3) of Γ_+ , Lipschitz property of f and inequality $(a+b)^2 \leq 2(a^2 + b^2)$ we get

$$(6.2) \quad \begin{aligned} \Gamma_+(f)(x) &= \sum_{i=1}^n (f(x) - f(x^i))_+^2 L(x, x^i) + \frac{1}{2} \sum_{i,j=1}^n (f(x) - f(x^{ij}))_+^2 L(x, x^{ij}) \\ &\leq \sum_{i=1}^n \alpha_i^2 L(x, x^i) + \frac{1}{2} \sum_{i,j=1}^n (\alpha_i + \alpha_j)^2 L(x, x^{ij}) \mathbf{1}_{\{x \neq x^{ij}\}} \\ &\leq \sum_{i=1}^n \alpha_i^2 L(x, x^i) + 2 \sum_{i=1}^n \alpha_i^2 \sum_{j=1}^n L(x, x^{ij}) \mathbf{1}_{\{x \neq x^{ij}\}} \leq 2 \sum_{i=1}^n \alpha_i^2 \sum_{y: y_i \neq x_i} L(x, y). \end{aligned}$$

Therefore, by the stability condition (4.7) we estimate $\|\Gamma_+(f)\|_\infty \leq 2R\rho(L)|\alpha|^2$. Herbst's argument (4.5) allows to conclude the first part.

The second part of the proposition follows by observing that for a flip-swap random walk

$$\sum_{i=1}^n \sum_{y: y_i \neq x_i} L(x, y) \leq 2 \cdot \Delta(L)$$

so by (6.2), Lemma 6.1 applied in the scalar setting $d = 1$ with $t_i = 2 \sum_{y: y_i \neq x_i} L(x, y)$, $D_i = \alpha_i^2$, $T_1 = 4\Delta(L)$ and $T_\infty = 4R\rho(L)$ we can estimate

$$\|\Gamma_+(f)\|_\infty \leq 4R\rho(L) \sum_{i=1}^{\lceil \Delta(L)/R\rho(L) \rceil} (\alpha_i^\downarrow)^2$$

and conclude again in virtue of Herbst's argument (4.5). \square

The proof of Proposition 4.9 follows along similar lines to the proof of Proposition 4.7, the difference being that in the end, instead of Herbst's argument, we apply the concentration result of Aoun et al. [8], which asserts that if L satisfies the matrix Poincaré inequality with constant $C_P > 0$

$$(6.3) \quad \text{Var}(f) \preceq -C_P \pi(fLf) \quad \forall f: \mathcal{B}_n \rightarrow \mathcal{H}_d,$$

(where L acts on the matrix-valued function f elementwise and fLf is the matrix product), then it satisfies the exponential concentration bound of the form

$$(6.4) \quad \pi(\lambda_{\max}(f - \pi(f)) > t) \leq d \exp\left(\frac{-t^2}{2C_P v_f + t\sqrt{2C_P v_f}}\right),$$

where $v_f = \sup_x \|\Gamma(f)(x)\|$ (where Γ is defined via (4.2), again with matrix multiplication, and $\|\cdot\|$ stands for the operator norm). Note that for $d = 1$, (6.3) is just the usual scalar Poincaré inequality.

Proof of Proposition 4.9. For any $x \in \mathcal{B}_n$ and $i, j \in [n]$, using operator convexity of the function $x \mapsto x^2$ (see [11, Example V.1.3]) we get that

$$(6.5) \quad (f(x) - f(x^{ij}))^2 = [(f(x) - f(x^i)) + (f(x^i) - f(x^{ij}))]^2 \\ \preceq 2(f(x) - f(x^i))^2 + 2(f(x^i) - f(x^{ij}))^2.$$

Therefore, by the definition (4.2) of Γ , by the assumed Lipschitz property (4.8) of f and by (6.5), for any $x \in \mathcal{B}_n$,

$$(6.6) \quad \Gamma(f)(x) = \frac{1}{2} \sum_{i=1}^n (f(x) - f(x^i))^2 L(x, x^i) + \frac{1}{4} \sum_{i,j=1}^n (f(x) - f(x^{ij}))^2 L(x, x^{ij}) \\ \preceq \frac{1}{2} \sum_{i=1}^n C_i^2 L(x, x^i) + \frac{1}{2} \sum_{i,j=1}^n (C_i^2 + C_j^2) L(x, x^{ij}) \mathbf{1}_{\{x \neq x^{ij}\}} \\ \preceq \sum_{i=1}^n C_i^2 \cdot \left[\sum_{y: y_i \neq x_i} L(x, y) \right].$$

As both hand sides of (6.6) are positive semidefinite, their norms compare as well. Therefore, as in the proof of Proposition 4.7, by Lemma 6.1 with $t_i = \sum_{y: y_i \neq x_i} L(x, y)$, $T_1 = 2\Delta(L)$, $T_\infty = 2R\rho(L)$ and $D_i = C_i^2$

$$\sup_{x \in \mathcal{B}_n} \|\Gamma(f)(x)\| \leq 2R\rho(L) \cdot \sup \left\{ \left\| \sum_{i \in \mathcal{I}} C_i^2 \right\| : \mathcal{I} \subset [n], |\mathcal{I}| \leq \lceil \Delta(L)/R\rho(L) \rceil \right\}.$$

Since L satisfies the (scalar) modified log-Sobolev inequality (4.4), then it satisfies the (scalar) Poincaré inequality with constant $C_P = 2/\rho(L)$ (see, e.g., [13, p. 292], noting slightly different definitions of constants in functional inequalities used therein) and whence by [30, Proposition 2.2] or [27, Theorem 1.1] it satisfies the matrix Poincaré inequality (6.3) with the same constant, which yields the conclusion in virtue of (6.4). \square

6.2. Proposition 4.10. The proof of Proposition 4.10 is based on the idea introduced by Boucheron et al. [16] and then developed by Paulin [42]. We follow the exposition from the works of Sambale and Sinulis [45, 46]. We start with the following lemmas.

Lemma 6.2. *For any flip-swap random walk with generator L satisfying the stability condition (4.7) and for any $A \subset \mathcal{B}_n$*

$$(6.7) \quad \Gamma_+(d_T^2(\cdot, A))(x) \leq 8R\rho(L) \cdot d_T^2(x, A).$$

Moreover, for any $x, y \in \mathcal{B}_n$ and any set $A \subset \mathcal{B}_n$,

$$(6.8) \quad d_T^2(x, A) - d_T^2(y, A) \leq d_H(x, y).$$

Proof. For $x \in \mathcal{B}_n$, $\alpha \in \mathbb{R}^n$ and a probability measure μ on \mathcal{B}_n , let $h_x(\mu, \alpha) = \sum_i \alpha_i \mu(z: z_i \neq x_i)$. By Sion's minmax theorem, cf. [17, p. 227],

$$(6.9) \quad d_T(x, A) = \inf_{\mu \in \mathcal{M}(A)} \sup_{\alpha \in B_2^n} h_x(\mu, \alpha),$$

where $\mathcal{M}(A)$ is the set of probability measures on A and $B_2^n = \{x \in \mathbb{R}^n : |x| \leq 1\}$ is the unit ball in \mathbb{R}^n . Let $\alpha^* \in \mathbb{R}_+^n \cap B_2^n$, $\mu^* \in \mathcal{M}(A)$ be such that $d_T(x, A) = h_x(\mu^*, \alpha^*)$ and set $\nu_y = \operatorname{argmin}_{\nu \in \mathcal{M}(A)} h_y(\nu, \alpha^*)$. Then

$$\begin{aligned} \Gamma_+(d_T(\cdot, A))(x) &= \sum_y [h_x(\mu^*, \alpha^*) - \inf_{\nu \in \mathcal{M}(A)} \sup_{\alpha \in B_2^n} h_y(\nu, \alpha)]_+^2 L(x, y) \\ &\leq \sum_y [h_x(\mu^*, \alpha^*) - h_y(\nu_y, \alpha^*)]_+^2 L(x, y) \\ &\leq \sum_y [h_x(\nu_y, \alpha^*) - h_y(\nu_y, \alpha^*)]_+^2 L(x, y) \\ &= \sum_y \left[\sum_i \alpha_i^* (\nu_y(z: z_i \neq x_i) - \nu_y(z: z_i \neq y_i)) \right]_+^2 L(x, y) \\ &\leq \sum_y \left[\sum_i \alpha_i^* \mathbf{1}_{\{x_i \neq y_i\}} \right]_+^2 L(x, y) \\ &\leq 2 \sum_i (\alpha_i^*)^2 \sum_{y: y_i \neq x_i} L(x, y) \leq 2R\rho(L), \end{aligned}$$

where the penultimate inequality follows since L is a flip-swap random walk and therefore $L(x, y) > 0$ implies that $d_H(x, y) \leq 2$ and so at most two elements of the sum $\sum_i \alpha_i^* \mathbf{1}_{\{x_i \neq y_i\}}$ are non-zero, whence we may apply the inequality $(a+b)^2 \leq 2(a^2+b^2)$. The last inequality is a consequence of the condition $\alpha^* \in B_2^n$ and the stability condition (4.7). We conclude (6.7) using the definition of Γ_+ and estimating $(a-b)_+^2(a+b)_+^2 \leq 4a^2(a-b)_+^2$.

To show the second part, note that (6.9) together with the Cauchy–Schwarz inequality imply that

$$d_T^2(x, A) = \inf_{\mu \in \mathcal{M}(A)} \sum_i (\mu(z: z_i \neq x_i))^2 = \sum_i (\mu_x^*(z: z_i \neq x_i))^2$$

for some $\mu_x^* \in \mathcal{M}(A)$. Therefore, for any $x, y \in \mathcal{B}_n$

$$d_T^2(x, A) - d_T^2(y, A) \leq \sum_i \left[(\mu_x^*(z: z_i \neq x_i))^2 - (\mu_x^*(z: z_i \neq y_i))^2 \right] \leq \sum_i \mathbf{1}_{\{x_i \neq y_i\}},$$

as desired. \square

Using the inequality $1 - e^{-z} \leq z$ we observe that for any $f: \mathcal{B}_n \rightarrow \mathbb{R}$,

$$\mathcal{E}(e^f, f) = \sum_x \pi(x) e^{f(x)} \left[\sum_y (f(x) - f(y))_+ (1 - e^{f(y)-f(x)}) L(x, y) \right] \leq \pi(e^f \Gamma_+(f)).$$

Therefore, the modified log-Sobolev inequality (4.4) implies the following inequality stated in Bobkov and Götze [12]:

$$(6.10) \quad \rho(L) \operatorname{Ent}_\pi(e^f) \leq \pi(e^f \tilde{\Gamma}(f)^2)$$

with operator $\tilde{\Gamma}(f) = \sqrt{\Gamma_+(f)}$ (note that in [12] $\tilde{\Gamma}$ is denoted by Γ , we use $\tilde{\Gamma}$ to avoid a conflict of notation). As a consequence, the hypothesis of [12, Theorem 2.1] (formula (1.1)) therein holds under the assumption of the modified log-Sobolev inequality (4.4) (with $c = 2/\rho(L)$). As a result, the following lemma follows directly by the derivation of [12, equation (2.4)] with a slight adjustment of constants (see also [5]).

Lemma 6.3. *If a measure π on \mathcal{B}_n satisfies the modified log-Sobolev inequality (4.4) and $f: \mathcal{B}_n \rightarrow [0, \infty)$ is such that $\Gamma_+(f) \leq Cf$ for some constant $C > 0$, then for all $t > C/\rho(L)$,*

$$(6.11) \quad \pi(\exp(f/t)) \leq \exp\left(\frac{\pi(f)}{t - C/\rho(L)}\right).$$

We are finally in position to prove Proposition 4.10.

Proof of Proposition 4.10. To lighten notation, denote $f(x) = d_T^2(x, A)$ for $x \in \mathcal{B}_n$ and some fixed set $A \subset \mathcal{B}_n$. Denote also $h(z) = (e^z - 1)/z$ for $z \in [0, \infty)$ and $Df_y(x) = f(x) - f(y)$ for $x, y \in \mathcal{B}_n$, and note that h is an increasing function. Starting with the modified log-Sobolev

inequality (4.4) we have for all $\lambda > 0$,

$$\begin{aligned}
 \text{Ent}_\pi(e^{-\lambda f}) &\leq \lambda/\rho(L) \cdot \mathcal{E}(e^{-\lambda f}, -f) \\
 \text{(by reversibility of } L) &= \lambda/\rho(L) \sum_{x,y} (Df_y(x))_+ (e^{-\lambda f(y)} - e^{-\lambda f(x)}) L(x,y) \pi(x) \\
 &= \lambda^2/\rho(L) \sum_x \pi(x) e^{-\lambda f(x)} \left[\sum_y (Df_y(x))_+^2 h(\lambda Df_y(x)) L(x,y) \right] \\
 \text{(by (6.8))} &\leq \lambda^2 h(2\lambda)/\rho(L) \cdot \pi(e^{-\lambda f} \Gamma_+(f)) \\
 \text{(by (6.7))} &\leq 8R\lambda^2 h(2\lambda) \cdot \pi(e^{-\lambda f} f) \\
 &\leq 8R\lambda^2 h(2\lambda) \cdot \pi(e^{-\lambda f}) \pi(f),
 \end{aligned}$$

where the last inequality follows by convexity of the function $t \mapsto t \log t$. Therefore, using the entropy method (cf., e.g., [17, Chapter 6]) and monotonicity of h , we have for every $\lambda > 0$,

$$\begin{aligned}
 \pi(\exp(\lambda(\pi(f) - f))) &= \exp\left(\lambda \int_0^\lambda \frac{d}{ds} \left[\frac{1}{s} \log \pi(e^{-sf}) \right] ds\right) \\
 &= \exp\left(\lambda \int_0^\lambda \frac{\text{Ent}_\pi(e^{-sf})}{s^2 \pi(e^{-sf})} ds\right) \\
 &\leq \exp\left(\lambda \cdot 8R\pi(f) \int_0^\lambda h(2s) ds\right) \leq \exp(4R\lambda(e^{2\lambda} - 1)\pi(f)).
 \end{aligned}$$

By Chebyshev's exponential inequality

$$\pi(A) = \pi(\pi(f) - f \geq \pi(f)) \leq \exp\left(\lambda(4R(e^{2\lambda} - 1) - 1)\pi(f)\right).$$

Taking $\lambda = \frac{1}{2} \log(1 + \frac{1}{8R})$ and estimating $\log(1+x) \geq x/(x+1)$ for $x \geq 1$ gives

$$(6.12) \quad \pi(A) \leq \exp\left(-\frac{1}{4} \log\left(1 + \frac{1}{8R}\right) \pi(f)\right) \leq \exp\left(-\frac{\pi(f)}{32R+4}\right).$$

We conclude by dividing (6.12) by its right hand side and using Lemma 6.3 with $t = 4 + 40R$ and $C = 8R\rho(L)$ (in virtue of Lemma 6.2). \square

6.3. Proposition 4.11. Before we move to the proof of Proposition 4.11, let us comment a bit on a background result. Using the equivalence between the modified log-Sobolev inequality (4.4) and the family of Beckner inequalities together with the approach developed by Boucheron et al. [15], it was shown in [2, Proposition 3.1] that the following moment estimate is implied by the modified log-Sobolev inequality. Below we will denote by $\|\cdot\|_p$ the norm in $L^p(\pi)$.

Proposition 6.4. *If a probability measure π on \mathcal{B}_n satisfies the modified log-Sobolev inequality (4.4), then for any $p \geq 2$,*

$$(6.13) \quad \|(f - \pi(f))_+\|_p \leq C \sqrt{p/\rho(L)} \|\sqrt{\Gamma_+(f)}\|_p,$$

where $C = \sqrt{3\sqrt{e}/(\sqrt{e}-1)}$.

A general method of deriving estimates for polynomials from moment inequalities of the form (6.13) has been presented in [4] in the continuous case, and in [1, 2] in the context of Glauber dynamics. To obtain results for flip-swap random walks we will adapt a version of this method introduced recently by Sambale and Sinulis [46] for multislices.

Proof of Proposition 4.11. Below we write C to denote universal constants and C_a to denote constants depending only on the parameter a . In both cases the constants may change values between occurrences. Let $f: \mathcal{B}_n \rightarrow \mathbb{R}$ be a tetrahedral polynomial. By ∂_i we denote the partial derivative with respect to the i -th coordinate. If $x, y \in \mathcal{B}_n$ differ at the i -th coordinate only, then by the fact that f is linear in each coordinate

$$|f(x) - f(y)| = |\partial_i f(x)|.$$

Similarly, if x and y differ only by a swap of the i -th and j -th coordinate, we have

$$\begin{aligned}
 |f(x) - f(y)| &= |\partial_i f(x)(y_i - x_i) + \partial_j f(x)(y_j - x_j) + \partial_i \partial_j f(x)(y_i - x_i)(y_j - x_j)| \\
 &\leq |\partial_i f(x)| + |\partial_j f(x)| + |\partial_i \partial_j f(x)|.
 \end{aligned}$$

Thus

$$\begin{aligned} \Gamma(f)(x) &= \frac{1}{2} \sum_{i=1}^n (f(x) - f(x^i))^2 L(x, x^i) + \frac{1}{2} \sum_{1 \leq i < j \leq n} (f(x) - f(x^{ij}))^2 L(x, x^{ij}) \\ &\leq \frac{1}{2} \sum_{i=1}^n |\partial_i f(x)|^2 L(x, x^i) + \frac{3}{2} \sum_{\substack{1 \leq i < j \leq n \\ x^{ij} \neq x}} (|\partial_i f(x)|^2 + |\partial_j f(x)|^2 + |\partial_i \partial_j f(x)|^2) L(x, x^{ij}) \\ &\leq R\rho(L) \left(3.5 \sum_{i=1}^n |\partial_i f(x)|^2 + 0.75 \sum_{i,j=1}^n |\partial_i \partial_j f(x)|^2 \right), \end{aligned}$$

where in the last inequality we used the stability condition (4.7). Note that since f is tetrahedral, $\partial_i \partial_i f(x) = 0$ for all i .

Combining the above equality with Proposition 6.4 we obtain that for every tetrahedral polynomial $f: \mathcal{B}_n \rightarrow \mathbb{R}$

$$(6.14) \quad \|f - \pi(f)\|_p \leq C\sqrt{p}\sqrt{R} \left(\|\nabla f\|_p + \|\nabla^2 f\|_{HS} \right),$$

where C is a universal constant.

In the subsequent part of the proof we are going to need some auxiliary notation. For d -tensors $A = (a_i)_{i \in [n]^d}$, $B = (b_i)_{i \in [n]^d}$ define

$$\langle A, B \rangle = \sum_{i \in [n]^d} a_i b_i.$$

Let us now consider a family of stochastically independent random tensors $\{G^I: I \subseteq \mathbb{N}, |I| \in \{1, 2\}\}$, given by $G^{\{m\}} = (g_i^{\{m\}})_{i \in [m]}$, $G^{\{l,k\}} = (g_{i,j}^{\{l,k\}})_{i,j \in [n]}$, with coefficients being i.i.d. standard Gaussian variables. Denote by $P_{d, \leq 2}$ the family of all partitions of the set $[d]$ into non-empty subsets of cardinality at most 2. Finally, for any positive integers d and l and $\mathcal{J} = \{J_1, \dots, J_l\} \in P_{d, \leq 2}$ define the random d -tensor $G_{\mathcal{J}} = (\prod_{j=1}^l g_{i_j}^{J_j})_{i \in [n]^d}$. For instance $G_{\{\{1,3\}, \{2\}\}} = (g_{i_1 i_3}^{\{1,3\}} g_{i_2}^{\{2\}})_{i_1, i_2, i_3 \in [n]}$.

Using the fact that the p -th moment of a mean zero Gaussian variable with variance σ^2 is for $p \geq 2$ comparable to $\sqrt{p}\sigma$ up to universal constants, we can rewrite (6.14) as

$$(6.15) \quad \|f(X) - \mathbb{E} f(X)\|_p \leq C\sqrt{R} \left(\|\nabla f(X), G^{\{1\}}\|_p + \|\nabla^2 f(X), G^{\{1,2\}}\|_p \right),$$

where X is a random vector with law π , independent of the family $\{G^I\}$.

The inequality (6.15) constitutes a basis for the induction argument leading to the following inequality valid for any $f: \mathcal{B}_n \rightarrow \mathbb{R}$, $d \geq 1$ and $p \geq 2$,

$$(6.16) \quad \begin{aligned} \|f(X) - \mathbb{E} f(X)\|_p &\leq C_d \left(\sum_{l=d}^{2d} \sum_{\mathcal{J} \in P_{l, \leq 2}} R^{|\mathcal{J}|/2} \|\nabla^l f(X), G_{\mathcal{J}}\|_p \right. \\ &\quad \left. + \sum_{l=1}^{2d-2} \sum_{\mathcal{J} \in P_{l, \leq 2}} R^{|\mathcal{J}|/2} \|\mathbb{E}_X \nabla^l f(X), G_{\mathcal{J}}\|_p \right). \end{aligned}$$

Before we prove the above estimate, let us show how it implies the statement of the proposition. If f is a tetrahedral polynomial of degree d , then $\nabla^l f = 0$ for $l > d$, moreover $\nabla^d f$ is constant and so $\nabla^d f(X) = \mathbb{E} \nabla^d f(X)$. Thus (6.16) reduces to

$$\|f(X) - \mathbb{E} f(X)\|_p \leq C_d \sum_{l=1}^d \sum_{\mathcal{J} \in P_{l, \leq 2}} R^{|\mathcal{J}|/2} \|\mathbb{E}_X \nabla^l f(X), G_{\mathcal{J}}\|_p.$$

We can now use moment estimates for tetrahedral homogeneous polynomials in i.i.d. standard Gaussian variables due to Latała [35], which assert that for any l -tensor $A = (a_i)_{i \in [n]^l}$ and $p \geq 2$,

$$\|\langle A, G_{\{\{1\}, \dots, \{l\}\}} \rangle\|_p \leq C_l \sum_{\mathcal{J} \in P_l} p^{|\mathcal{J}|/2} \|A\|_{\mathcal{J}}.$$

Applying this inequality to $\langle \mathbb{E}_X \nabla^l f(X), G_{\mathcal{J}} \rangle$ (we treat here $\mathbb{E}_X \nabla^l f(X)$ as a $|\mathcal{J}|$ -tensor by merging the indices according to the partition \mathcal{J}), we obtain

$$\|f(X) - \mathbb{E} f(X)\|_p \leq C_d \sum_{l=1}^d \sum_{\mathcal{J} \in P_{l, \leq 2}} R^{|\mathcal{J}|/2} \sum_{\mathcal{I} \in P_l: \mathcal{I} \succ \mathcal{J}} p^{|\mathcal{I}|/2} \|\mathbb{E} \nabla^l f(X)\|_{\mathcal{I}},$$

where $\mathcal{I} \succ \mathcal{J}$ if every element of \mathcal{I} is a union of certain elements of \mathcal{J} . Rearranging the terms and taking into account that in a non-trivial case R is bounded away from zero by an absolute constant (see Remark 4.4), which gives $R^{|\mathcal{J}|/2} \leq C_d R^{l/2}$ for $\mathcal{J} \in P_{l, \leq 2}$, we get

$$\|f(X) - \mathbb{E} f(X)\|_p \leq C_d \sum_{l=1}^d \sum_{\mathcal{I} \in P_l} R^{l/2} p^{|\mathcal{I}|/2} \|\mathbb{E} \nabla^l f(X)\|_{\mathcal{I}}$$

for $p \geq 2$. This implies the tail inequality of the proposition in the standard way by the use of Chebyshev's inequality $\mathbb{P}(|f(X) - \mathbb{E} f(X)| \geq e \|f(X) - \mathbb{E} f(X)\|_p) \leq e^{-p}$ followed by an appropriate change of variables and adjustment of constants. We leave the details to the reader and turn to the proof of (6.16).

We will proceed by induction on d . For $d = 1$, using the definitions of $G_{\{1\}}$ and $G_{\{\{1,2\}\}}$ one can easily see that (6.16) reads as

$$\begin{aligned} \|f(X) - \mathbb{E} f(X)\|_p \leq & C \left(\sqrt{R} \|\langle \nabla f(X), G_{\{1\}} \rangle\|_p + \sqrt{R} \|\langle \nabla^2 f(X), G_{\{\{1,2\}\}} \rangle\|_p \right. \\ & \left. + R \|\langle \nabla^2 f(X), G_{\{\{1\}, \{2\}\}} \rangle\|_p \right), \end{aligned}$$

which is clearly weaker than (6.15). Let us thus assume that the inequality holds for all positive integers smaller than d . Applying the inequality with $d-1$ and combining it with the triangle inequality in L_p we get (recall that the value of C_d may change between occurrences)

$$\begin{aligned} (6.17) \quad \|f(X) - \mathbb{E} f(X)\|_p \leq & C_d \left(\sum_{l=d-1}^{2d-2} \sum_{\mathcal{J} \in P_{l, \leq 2}} R^{|\mathcal{J}|/2} \|\langle \nabla^l f(X), G_{\mathcal{J}} \rangle\|_p \right. \\ & \left. + \sum_{l=1}^{2d-4} \sum_{\mathcal{J} \in P_{l, \leq 2}} R^{|\mathcal{J}|/2} \|\langle \mathbb{E}_X \nabla^l f(X), G_{\mathcal{J}} \rangle\|_p \right) \\ \leq & C_d \left(\sum_{l=d-1}^{2d-2} \sum_{\mathcal{J} \in P_{l, \leq 2}} R^{|\mathcal{J}|/2} \|\langle \nabla^l f(X), G_{\mathcal{J}} \rangle - \langle \mathbb{E}_X \nabla^l f(X), G_{\mathcal{J}} \rangle\|_p \right. \\ & \left. + \sum_{l=1}^{2d-2} \sum_{\mathcal{J} \in P_{l, \leq 2}} 2R^{|\mathcal{J}|/2} \|\langle \mathbb{E}_X \nabla^l f(X), G_{\mathcal{J}} \rangle\|_p \right). \end{aligned}$$

An application of inequality (6.15) conditionally on $G_{\mathcal{J}}$ to the functions $g_{l, \mathcal{J}}(x) = \langle \nabla^l f(x), G_{\mathcal{J}} \rangle$ for $l = d-1, \dots, 2d-2$ and $\mathcal{J} \in P_{l, \leq 2}$ (note that $g_{l, \mathcal{J}}$'s are tetrahedral polynomials), followed by the Fubini theorem, gives

$$\begin{aligned} \|\langle \nabla^l f(X), G_{\mathcal{J}} \rangle - \langle \mathbb{E}_X \nabla^l f(X), G_{\mathcal{J}} \rangle\|_p \leq & C \sqrt{R} \left(\|\langle \nabla^{l+1} f(X), G_{\mathcal{J} \cup \{\{l+1\}\}} \rangle\|_p \right. \\ & \left. + \|\langle \nabla^{l+2} f(X), G_{\mathcal{J} \cup \{\{l+1, l+2\}\}} \rangle\|_p \right), \end{aligned}$$

which combined with (6.17) concludes the induction step, thus proving (6.16). \square

7. PROOFS OF THE RESULTS OF SECTION 3

By virtue of the abstract results of Section 4, all the results of Section 3 follow from the observation that there exist a flip-swap random walk on \mathcal{B}_n with stationary measure $\pi = \pi(p, k)$ that satisfy the stability condition (4.7) with constant $R = 2$ for all $p \in (0, 1)^n$ and $k = 0, \dots, n$ (cf. Proposition 7.3). The rest of this section is devoted to proving this fact.

We start with the following lemma, the proof of which is postponed until the end of this section.

Lemma 7.1. *For every $n \in \mathbb{N}$, $p \in (0, 1)^n$ and $k \in [n]$, there exist a coupling (Z, Z') of measures $\pi(p, k)$ and $\pi(p, k-1)$ such that for all $x \in \text{supp } \pi(p, k-1)$, and $r \in [n]$ such that $x_r = 0$,*

$$(7.1) \quad \mathbb{P}(Z = x + e_r \mid Z' = x) = \mathbb{E} \left[\frac{\mathbf{1}_{\{Z_r=1\}}}{\sum_{l=1}^n \mathbf{1}_{\{Z_l=1\}} \mathbf{1}_{\{x_l=0\}}} \right]$$

and for all $x \in \text{supp } \pi(p, k)$, and $r \in [n]$ such that $x_r = 1$,

$$(7.2) \quad \mathbb{P}(Z' = x - e_r \mid Z = x) = \mathbb{E} \left[\frac{\mathbf{1}_{\{Z'_r=0\}}}{\sum_{l=1}^n \mathbf{1}_{\{Z'_l=0\}} \mathbf{1}_{\{x_l=1\}}} \right].$$

Our approach to defining an R -stable generator L_π will be based on the inductive construction of Hermon and Salez [29] in the k -homogeneous case, which we will now recall. The construction works for any k -homogeneous probability measure π on \mathcal{B}_n , satisfying the SCP and produces a generator of a π -reversible flip-swap random walk Q^* such that $\rho(Q^*) \geq 1$ and $\Delta(Q^*) \leq 2k$.

To simplify the notation, we are going to treat vectors $x_{\neq l}$ for $x \in \mathcal{B}_n$ and $l \in [n]$ sometimes as elements of $\{0, 1\}^{[n] \setminus \{l\}}$ (this is how they were defined at the beginning of Section 2) and sometimes as elements of \mathcal{B}_{n-1} (with the natural identification, i.e., preserving the order of coordinates). The exact meaning will be clear from the context. The same convention will apply to random vectors, e.g., to $X_{\neq l}$.

In the case $n = 1$, we let Q be the trivial generator on a one-point space. Clearly then $\rho(Q) = \infty$ and $\Delta(Q) = 0$.

For $n > 1$, $l \in [n]$ and $x, y \in \text{supp } \pi$, $x \neq y$, we set

$$(7.3) \quad Q^{(l)}(x, y) = \begin{cases} \mathbb{P}(U = y_{\neq l} \mid V = x_{\neq l}) \mathbb{P}(X_l \neq x_l) & \text{if } x_l \neq y_l, \\ Q_{x_l}^{(l)}(x_{\neq l}, y_{\neq l}) & \text{else,} \end{cases}$$

where X is a random vector with law π and (U, V) is any coupling between $\mathcal{L}(X_{\neq l} \mid X_l = y_l)$ and $\mathcal{L}(X_{\neq l} \mid X_l = x_l)$ given by the SCP² and $Q_{x_l}^{(l)}$ is any flip-swap generator on \mathcal{B}_{n-1} with stationary distribution $\mathcal{L}(X_{\neq l} \mid X_l = x_l)$ such that $\rho(Q_{x_l}^{(l)}) \geq 1$ and $\Delta(Q_{x_l}^{(l)}) \leq 2(k - x_l)$, the existence of which is provided by the induction scheme. We define the diagonal elements of $Q^{(l)}$ so that the row sums vanish. Finally put

$$(7.4) \quad Q^* = \frac{1}{n} \sum_{l=1}^n Q^{(l)}.$$

Then by (the proof of) [29, Theorem 2], we have $\rho(Q^*) \geq 1$, $\Delta(Q^*) \leq 2k$.

Now we are in position to construct the generator L_π . Let $X \sim \pi = \pi(p, k)$ for some $p \in (0, 1)^n$ and $k \in \{0, \dots, n\}$. Observe that for any $y_l \in \{0, 1\}$, we have $\mathcal{L}(X_{\neq l} \mid X_l = y_l) = \pi(p_{\neq l}, k - y_l)$, in particular in the above recursive construction we can restrict our attention to the class of conditional Bernoulli distributions and use as $Q_{x_l}^{(l)}$ the generators defined for such measures in dimension $n - 1$. Moreover, for (U, V) we can take the coupling (Z, Z') (if $y_l = 0$) or (Z', Z) (if $y_l = 1$) given by Lemma 7.1 applied in dimension $n - 1$ with $p_{\neq l}$ instead of p (note that since the right-hand side of (7.1) summed over r such that $x_r = 0$ gives one, we indeed have $Z \triangleright Z'$, which makes this coupling a legitimate choice in the Hermon–Salez construction). Let us define L_π as the outcome of the Hermon–Salez construction with the above choices of $Q_{x_l}^{(l)}$ and (U, V) . Thus, formally for $n = 1$ we let L_π be the trivial generator and for $n > 1$ and $l \in [n]$ we set

$$(7.5) \quad L_\pi = \frac{1}{n} \sum_{l=1}^n L^{(l)}$$

with

$$(7.6) \quad L^{(l)}(x, y) = \begin{cases} \mathbb{P}(U = y_{\neq l} \mid V = x_{\neq l}) \mathbb{P}(X_l \neq x_l) & \text{if } x_l \neq y_l, \\ L_{\pi_l}(x_{\neq l}, y_{\neq l}) & \text{else,} \end{cases}$$

for $x \neq y$, where (U, V) is the coupling of $\pi(p_{\neq l}, k - y_l)$ and $\pi(p_{\neq l}, k - x_l)$ given by Lemma 7.1, and $\pi_l = \pi(p_{\neq l}, k - y_l)$ (again the diagonal elements are adjusted so that the row sums vanish).

Then, the results by Hermon and Salez, specialized to L_π give

Proposition 7.2. *The generator L_π constructed according to (7.5) generates a reversible flip-swap random walk with stationary measure π such that $\rho(L_\pi) \geq 1$ and $\Delta(L_\pi) \leq 2k$.*

Our main result concerning conditional Bernoulli distributions, underlying all the results from Section 3 is

²We note a small typo in the cited preprint version of [29] – for Q to be self-adjoint we need to have $\pi(x)$ in the denominator of each expression in [29, equation (90)]. This change is also consistent with the subsequent part of the proof in [29].

Proposition 7.3. *The generator L_π constructed according to (7.5) with stationary measure π satisfies the stability condition (4.7) with $R = 2$.*

Proof of Proposition 7.3. We proceed by induction in the dimension n .

For $n = 1$ the only possibilities are $k = 0$ and $k = 1$ and in both cases the left-hand side of (4.7) vanishes. Thus the stability condition (4.7) is satisfied with any nonnegative R .

Assume the induction hypothesis holds for $n - 1$ and fix $x \in \text{supp } \pi$ and $i \in [n]$. We may and do assume that $k \in \{1, \dots, n - 1\}$ as otherwise L_π trivializes.

Since $\rho(L_\pi) \geq 1$, it is enough to show that

$$(7.7) \quad \max_{x \in \text{supp } \pi; i \in [n]} \sum_{y: y_i \neq x_i} L_\pi(x, y) \leq 2.$$

As in the definition of L_π we will denote by X a random variable with distribution π .

If $x_i = 0$, then by (7.5)

$$(7.8) \quad \begin{aligned} \sum_{y: y_i \neq x_i} L_\pi(x, y) &= \sum_{j: x_j = 1} \frac{1}{n} \sum_{l=1}^n L^{(l)}(x, x^{ij}) \\ &= \sum_{l \in [n] \setminus \{i, j\}} \frac{1}{n} \sum_{j: x_j = 1} L^{(l)}(x, x^{ij}) + \frac{1}{n} \sum_{j: x_j = 1} L^{(i)}(x, x^{ij}) + \frac{1}{n} \sum_{j: x_j = 1} L^{(j)}(x, x^{ij}), \end{aligned}$$

where we recall that $x^{ij} = x + e_i - e_j$. We estimate each term on the right hand side separately.

For $l \in [n]$ let ζ_l be the unique increasing bijection between $[n] \setminus \{l\}$ and $[n - 1]$. If $l \neq i, j$, then for $y = x^{ij}$ we have $y_l = x_l$ and so, by (7.6), $L^{(l)}(x, y) = L_{\pi_l}(x_{\neq l}, y_{\neq l})$, where $\pi_l = \pi(p_{\neq l}, k - x_l)$. Thus, denoting $r_l = \zeta_l(i)$, we get

$$(7.9) \quad \begin{aligned} \sum_{l \in [n] \setminus \{i, j\}} \frac{1}{n} \sum_{j: x_j = 1} L^{(l)}(x, x^{ij}) &= \frac{1}{n} \sum_{l \in [n] \setminus \{i\}} \sum_{j \neq l: x_j = 1} L_{\pi_l}(x_{\neq l}, (x^{ij})_{\neq l}) \\ &= \frac{1}{n} \sum_{l \in [n] \setminus \{i\}} \sum_{y \in \mathcal{B}_{n-1}: y_{r_l} \neq (x_{\neq l})_{r_l}} L_{\pi_l}(x_{\neq l}, y) \leq \frac{n-1}{n} \cdot 2, \end{aligned}$$

where the last inequality follows from the induction assumption applied to π_l .

The second term of (7.8) is estimated again using the definition (7.6). Indeed, if $x_j = 1$, then for $y = x^{ij}$ we have $x_i \neq y_i$. Thus, recalling that (U, V) is a coupling between the laws $\mathcal{L}(X_{\neq i} | X_i = 1)$ and $\mathcal{L}(X_{\neq i} | X_i = 0)$ such that $V \triangleright U$, we obtain

$$(7.10) \quad \begin{aligned} \frac{1}{n} \sum_{j: x_j = 1} L^{(i)}(x, x^{ij}) &= \frac{1}{n} \sum_{j: x_j = 1} \mathbb{P}(U = (x^{ij})_{\neq i} | V = x_{\neq i}) \mathbb{P}(X_i \neq x_i) \\ &= \frac{1}{n} \mathbb{P}(X_i = 1) \leq \frac{1}{n}. \end{aligned}$$

Let us pass to the last term of (7.8). We stress that this is the crucial part of the proof, the only one in which we use the specific form of the coupling (U, V) used in the construction of L_π .

To estimate this last term we use (7.1) from Lemma 7.1 combined with the fact that if $x_i = 0$ and $x_j = 1$, then for $y = x^{ij}$, $y_j = 0 \neq x_j$ and so (U, V) from (7.6) is the coupling between the laws $\pi(p_{\neq j}, k)$ and $\pi(p_{\neq j}, k - 1)$ given by Lemma 7.1 (in dimension $n - 1$). For $j \in [n]$ consider a \mathcal{B}_{n-1} -valued random vector $Z^{(j)} \sim \mathcal{L}(X_{\neq j} | X_j = 0) = \pi(p_{\neq j}, k)$. Note also that since X, x have the same number of ones, we have

$$(7.11) \quad \sum_{l=1}^n \mathbf{1}_{\{X_l=0\}} \mathbf{1}_{\{x_l=1\}} = \sum_{l=1}^n \mathbf{1}_{\{X_l=1\}} \mathbf{1}_{\{x_l=0\}}.$$

Putting all the above observations together, we obtain

$$\begin{aligned}
(7.12) \quad \frac{1}{n} \sum_{j: x_j=1} L^{(j)}(x, x^{ij}) &\stackrel{\text{Lemma 7.1, (7.6)}}{=} \frac{1}{n} \sum_{j: x_j=1} \mathbb{E} \left[\frac{\mathbf{1}_{\{Z_i^{(j)}=1\}}}{\sum_{l \neq j} \mathbf{1}_{\{Z_l^{(j)}=1\}} \mathbf{1}_{\{x_l=0\}}} \right] \mathbb{P}(X_j = 0) \\
&= \frac{1}{n} \sum_{j: x_j=1} \mathbb{E} \left[\mathbf{1}_{\{X_i=1\}} \frac{\mathbf{1}_{\{X_j=0\}}}{\sum_{l \neq j} \mathbf{1}_{\{X_l=1\}} \mathbf{1}_{\{x_l=0\}}} \right] \\
&\stackrel{x_j=1}{=} \frac{1}{n} \sum_{j: x_j=1} \mathbb{E} \left[\mathbf{1}_{\{X_i=1\}} \frac{\mathbf{1}_{\{X_j=0\}}}{\sum_l \mathbf{1}_{\{X_l=1\}} \mathbf{1}_{\{x_l=0\}}} \right] \\
&\stackrel{(7.11)}{=} \frac{1}{n} \sum_{j: x_j=1} \mathbb{E} \left[\mathbf{1}_{\{X_i=1\}} \frac{\mathbf{1}_{\{X_j=0\}}}{\sum_l \mathbf{1}_{\{X_l=0\}} \mathbf{1}_{\{x_l=1\}}} \right] \\
&= \frac{1}{n} \mathbb{P}(X_i = 1) \leq \frac{1}{n}.
\end{aligned}$$

Combining the estimates (7.9), (7.10) and (7.12) with (7.8) yields (7.7) and thus the stability condition (4.7) with $R = 2$ in the case $x_i = 0$. The case $x_i = 1$ is analogous, the main difference being that in (7.12) we use (7.2) in place of (7.1) from Lemma 7.1.

Together the two cases give the induction step and conclude the proof of the proposition. \square

Let us conclude this section with the proof of Lemma 7.1.

Proof of Lemma 7.1. For $x \in \mathcal{B}_n$, let $\kappa(x) = \sum_i x_i$ and let B be a vector of independent Bernoulli random variables with probabilities of success given by $p = (p_1, \dots, p_n)$. Consider three \mathcal{B}_n -valued random variables: $\widehat{Z} \sim \mathcal{L}(B \mid \kappa(B) = k)$, $Z' \sim \mathcal{L}(B \mid \kappa(B) = k - 1)$ and Z such that for all $x, y \in \mathcal{B}_n$,

$$(7.13) \quad \mathbb{P}(Z = y \mid Z' = x) = h(y, x),$$

where

$$h(y, x) = \mathbb{E} \left[\frac{\mathbf{1}_{\{\widehat{Z}_r=1\}}}{\sum_l \mathbf{1}_{\{\widehat{Z}_l=1\}} \mathbf{1}_{\{x_l=0\}}} \right]$$

if $y = x + e_r$ for some $r \in [n]$ and $\kappa(x) = k - 1$, and $h(y, x) = 0$ otherwise. Note that for $x \in \mathcal{B}_n$ such that $\kappa(x) = k - 1$, $\sum_l \mathbf{1}_{\{\widehat{Z}_l=1\}} \mathbf{1}_{\{x_l=0\}} > 0$ with probability one, so $h(y, x)$ is well defined. Moreover, for such x

$$\sum_{y \in \mathcal{B}_n} h(y, x) = \sum_{r: x_r=0} h(x + e_r, x) = 1,$$

which guarantees the existence of the couple (Z, Z') satisfying (7.13). Thus to prove (7.1) it is enough to show that $Z \sim \widehat{Z}$, i.e., that $\sum_{x \in \mathcal{B}_n} h(y, x) \mathbb{P}(Z' = x) = \mathbb{P}(\widehat{Z} = y)$ for any $y \in \mathcal{B}_n$ such that $\kappa(y) = k$.

Observe that for any $r \in [n]$ such that $x_r = 0$ and $\kappa(x) = k - 1$

$$(7.14) \quad \frac{\mathbb{P}(Z' = x)}{\mathbb{P}(\widehat{Z} = x + e_r)} = \frac{\mathbb{P}(B = x)}{\mathbb{P}(B = x + e_r)} \frac{\mathbb{P}(\kappa(B) = k)}{\mathbb{P}(\kappa(B) = k - 1)} = \frac{1 - p_r}{p_r} \frac{\mathbb{P}(\kappa(B) = k)}{\mathbb{P}(\kappa(B) = k - 1)}.$$

Moreover, for any $f: \mathcal{B}_n \rightarrow \mathbb{R}$ and $r \in [n]$,

$$(7.15) \quad \mathbb{E} [f(B) \mathbf{1}_{\{B_r=1\}}] \frac{1 - p_r}{p_r} = \mathbb{E} [f(B + e_r) \mathbf{1}_{\{B_r=0\}}].$$

We use (7.14) and (7.15) to get that for any such y and any $r \in [n]$ such that $y_r = 1$ and $\kappa(y) = k$,

$$\begin{aligned}
 (7.16) \quad & \frac{h(y, y - e_r) \mathbb{P}(Z' = y - e_r)}{\mathbb{P}(\widehat{Z} = y)} \stackrel{(7.14)}{=} h(y, y - e_r) \frac{1 - p_r}{p_r} \frac{\mathbb{P}(\kappa(B) = k)}{\mathbb{P}(\kappa(B) = k - 1)} \\
 & = \mathbb{E} \left[\frac{\mathbf{1}_{\{B_r=1\}} \mathbf{1}_{\{\kappa(B)=k\}}}{\mathbf{1}_{\{B_r=1\}} + \sum_{l \neq r} \mathbf{1}_{\{B_l=1\}} \mathbf{1}_{\{y_l=0\}}} \right] \frac{(1 - p_r)/p_r}{\mathbb{P}(\kappa(B) = k - 1)} \\
 & \stackrel{(7.15)}{=} \mathbb{E} \left[\frac{\mathbf{1}_{\{B_r=0\}} \mathbf{1}_{\{\kappa(B)=k-1\}}}{\mathbf{1}_{\{B_r=0\}} + \sum_{l \neq r} \mathbf{1}_{\{B_l=1\}} \mathbf{1}_{\{y_l=0\}}} \right] \frac{1}{\mathbb{P}(\kappa(B) = k - 1)} \\
 & = \mathbb{E} \left[\frac{\mathbf{1}_{\{B_r=0\}} \mathbf{1}_{\{\kappa(B)=k-1\}}}{\mathbf{1}_{\{B_r=0\}} + \sum_{l \neq r} \mathbf{1}_{\{B_l=0\}} \mathbf{1}_{\{y_l=1\}}} \right] \frac{1}{\mathbb{P}(\kappa(B) = k - 1)} \\
 & = \mathbb{E} \left[\frac{\mathbf{1}_{\{Z'_r=0\}}}{\sum_l \mathbf{1}_{\{Z'_l=0\}} \mathbf{1}_{\{y_l=1\}}} \right]
 \end{aligned}$$

where the penultimate step comes from the fact that for any u, v such that $\kappa(u) = \kappa(v)$ one has $\sum \mathbf{1}_{\{u=0\}} \mathbf{1}_{\{v=1\}} = \sum \mathbf{1}_{\{u=1\}} \mathbf{1}_{\{v=0\}}$ applied to $u = \xi_r(B)$, $v = \xi_r(y)$, where ξ_r is the projection from \mathcal{B}_n to \mathcal{B}_{n-1} obtained by skipping the r -th coordinate (note that if $B_r = 0$ and $\kappa(B) = k - 1$ then $\kappa(u) = \kappa(v) = k - 1$). Therefore by (7.16)

$$\frac{\sum_x h(y, x) \mathbb{P}(Z' = x)}{\mathbb{P}(\widehat{Z} = y)} = \frac{\sum_{r: y_r=1} h(y, y - e_r) \mathbb{P}(Z' = y - e_r)}{\mathbb{P}(\widehat{Z} = y)} = 1,$$

which completes the proof of (7.1). The equality (7.2) follows again by (7.16). \square

REFERENCES

1. Radosław Adamczak, Michał Kotowski, Bartłomiej Polaczyk, and Michał Strzelecki, *A note on concentration for polynomials in the Ising model*, Electron. J. Probab. **24** (2019), Paper No. 42, 22. MR 3949267 6, 17
2. Radosław Adamczak, Bartłomiej Polaczyk, and Michał Strzelecki, *Modified log-Sobolev inequalities, Beckner inequalities and moment estimates*, arXiv preprint arXiv:2007.10209 (2020). 1, 6, 8, 17
3. Radosław Adamczak and Michał Strzelecki, *On the convex Poincaré inequality and weak transportation inequalities*, Bernoulli **25** (2019), no. 1, 341–374. MR 3892322 5
4. Radosław Adamczak and Paweł Wolff, *Concentration inequalities for non-Lipschitz functions with bounded derivatives of higher order*, Probab. Theory Related Fields **162** (2015), no. 3-4, 531–586. MR 3383337 6, 17
5. Shigeki Aida, Takao Masuda, and Ichirō Shigekawa, *Logarithmic Sobolev inequalities and exponential integrability*, J. Funct. Anal. **126** (1994), no. 1, 83–101. MR 1305064 16
6. Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant, *Log-concave polynomials II: High-dimensional walks and an FPRAS for counting bases of a matroid*, STOC'19—Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing, ACM, New York, 2019, pp. 1–12. MR 4003314 1, 10
7. Cécile Ané, Sébastien Blachère, Djilil Chafaï, Pierre Fougères, Ivan Gentil, Florent Malrieu, Cyril Roberto, and Grégory Scheffer, *Sur les inégalités de Sobolev logarithmiques*, Panoramas et Synthèses [Panoramas and Synthèses], vol. 10, Société Mathématique de France, Paris, 2000, With a preface by Dominique Bakry and Michel Ledoux. MR 1845806 9
8. Richard Aoun, Marwa Banna, and Pierre Youssef, *Matrix Poincaré inequalities and concentration*, Adv. Math. **371** (2020), 107251, 33. MR 4108222 1, 2, 4, 15
9. Dominique Bakry, Michel Ledoux, and Laurent Saloff-Coste, *Markov semigroups at Saint-Flour*, Probability at Saint-Flour, Springer, Heidelberg, 2012. MR 3075390 7
10. Patrice Bertail and Stephan Cléménçon, *Bernstein-type exponential inequalities in survey sampling: conditional Poisson sampling schemes*, Bernoulli **25** (2019), no. 4B, 3527–3554. MR 4010964 1, 2
11. Rajendra Bhatia, *Matrix analysis*, Graduate Texts in Mathematics, vol. 169, Springer-Verlag, New York, 1997. MR 1477662 13, 15
12. S. G. Bobkov and F. Götze, *Exponential integrability and transportation cost related to logarithmic Sobolev inequalities*, J. Funct. Anal. **163** (1999), no. 1, 1–28. MR 1682772 16
13. Sergey G. Bobkov and Prasad Tetali, *Modified logarithmic Sobolev inequalities in discrete settings*, J. Theoret. Probab. **19** (2006), no. 2, 289–336. MR 2283379 1, 2, 7, 8, 15
14. Julius Borcea, Petter Brändén, and Thomas M. Liggett, *Negative dependence and the geometry of polynomials*, J. Amer. Math. Soc. **22** (2009), no. 2, 521–567. MR 2476782 1
15. Stéphane Boucheron, Olivier Bousquet, Gábor Lugosi, and Pascal Massart, *Moment inequalities for functions of independent random variables*, Ann. Probab. **33** (2005), no. 2, 514–560. MR 2123200 17
16. Stéphane Boucheron, Gábor Lugosi, and Pacal Massart, *On concentration of self-bounding functions*, Electron. J. Probab. **14** (2009), no. 64, 1884–1899. MR 2540852 15
17. Stéphane Boucheron, Gábor Lugosi, and Pascal Massart, *Concentration inequalities. A nonasymptotic theory of independence*, Oxford University Press, Oxford, 2013. MR 3185193 5, 7, 15, 17
18. Petter Brändén and June Huh, *Lorentzian polynomials*, Ann. of Math. (2) **192** (2020), no. 3, 821–891. MR 4172622 10

19. Sean X. Chen, *General properties and estimation of conditional Bernoulli models*, J. Multivariate Anal. **74** (2000), no. 1, 69–87. MR 1790614 2
20. Sean X. Chen and Jun S. Liu, *Statistical applications of the Poisson-binomial and conditional Bernoulli distributions*, Statist. Sinica **7** (1997), no. 4, 875–892. MR 1488647 2
21. Xiang-Hui Chen, Arthur P. Dempster, and Jun S. Liu, *Weighted finite population sampling to maximize entropy*, Biometrika **81** (1994), no. 3, 457–469. MR 1311090 2
22. Mary Cryan, Heng Guo, and Giorgos Mousa, *Modified log-Sobolev inequalities for strongly log-concave distributions*, Ann. Probab. **49** (2021), no. 1, 506–525. MR 4203344 3, 8, 10
23. Yuval Dagan, Constantinos Daskalakis, Nishanth Dikkala, and Anthimos Vardis Kandiros, *Learning Ising models from one or multiple samples*, Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing (New York, NY, USA), STOC 2021, Association for Computing Machinery, 2021, p. 161–168. 6
24. Devdatt P. Dubhashi and Alessandro Panconesi, *Concentration of measure for the analysis of randomized algorithms*, Cambridge University Press, Cambridge, 2009. MR 2547432 11
25. Fuqing Gao and Jeremy Quastel, *Exponential decay of entropy in the random transposition and Bernoulli-Laplace models*, Ann. Appl. Probab. **13** (2003), no. 4, 1591–1600. MR 2023890 1, 2, 8
26. Kevin Garbe and Jan Vondrak, *Concentration of Lipschitz functions of negatively dependent variables*, arXiv preprint arXiv:1804.10084 (2018). 1
27. Ankit Garg, Tarun Kathuria, and Nikhil Srivastava, *Scalar Poincaré implies matrix Poincaré*, Electron. Commun. Probab. **26** (2021), Paper No. 20, 4. MR 4245747 15
28. Sharad Goel, *Modified logarithmic Sobolev inequalities for some models of random walk*, Stochastic Process. Appl. **114** (2004), no. 1, 51–79. MR 2094147 8
29. Jonathan Hermon and Justin Salez, *Modified log-Sobolev inequalities for strong-Rayleigh measures*, arXiv preprint arXiv:1902.02775 (2019). 1, 2, 3, 7, 10, 20
30. De Huang and Joel A. Tropp, *Nonlinear matrix concentration via semigroup methods*, Electron. J. Probab. **26** (2021), Paper No. 8, 31. MR 4216521 15
31. Mark Jerrum, Jung-Bae Son, Prasad Tetali, and Eric Vigoda, *Elementary bounds on Poincaré and log-Sobolev constants for decomposable Markov chains*, Ann. Appl. Probab. **14** (2004), no. 4, 1741–1765. MR 2099650 2
32. Tarun Kathuria, *A Matrix Bernstein Inequality for Strong Rayleigh Distributions*, arXiv preprint arXiv:2011.13340 (2020). 1, 4
33. Tali Kaufman and Izhhar Oppenheim, *High order random walks: beyond spectral gap*, Approximation, randomization, and combinatorial optimization. Algorithms and techniques, LIPIcs. Leibniz Int. Proc. Inform., vol. 116, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2018, pp. Art. No. 47, 17. MR 3857285 10
34. Rasmus Kyng and Zhao Song, *A matrix Chernoff bound for strongly Rayleigh distributions and spectral sparsifiers from a few random spanning trees*, 59th Annual IEEE Symposium on Foundations of Computer Science—FOCS 2018, IEEE Computer Soc., Los Alamitos, CA, 2018, pp. 373–384. MR 3899605 1, 2, 4, 5, 13
35. Rafał Łatała, *Estimates of moments and tails of Gaussian chaoses*, Ann. Probab. **34** (2006), no. 6, 2315–2331. MR 2294983 6, 18
36. Michel Ledoux, *The concentration of measure phenomenon*, Mathematical Surveys and Monographs, vol. 89, American Mathematical Society, Providence, RI, 2001. MR 1849347 7
37. Tzong-Yow Lee and Horng-Tzer Yau, *Logarithmic Sobolev inequality for some models of random walks*, Ann. Probab. **26** (1998), no. 4, 1855–1873. MR 1675008 1
38. David A. Levin and Yuval Peres, *Markov chains and mixing times*, American Mathematical Society, Providence, RI, 2017, Second edition of [MR2466937], With contributions by Elizabeth L. Wilmer, With a chapter on “Coupling from the past” by James G. Propp and David B. Wilson. MR 3726904 7
39. Thomas M. Liggett, *Continuous time Markov processes*, Graduate Studies in Mathematics, vol. 113, American Mathematical Society, Providence, RI, 2010, An introduction. MR 2574430 7
40. Sheng Lin Lu and Horng-Tzer Yau, *Spectral gap and logarithmic Sobolev inequality for Kawasaki and Glauber dynamics*, Comm. Math. Phys. **156** (1993), no. 2, 399–433. MR 1233852 2
41. Ryan O’Donnell, *Analysis of Boolean functions*, Cambridge University Press, New York, 2014. MR 3443800 6
42. Daniel Paulin, *The convex distance inequality for dependent random variables, with applications to the stochastic travelling salesman and other problems*, Electron. J. Probab. **19** (2014), no. 68, 34. MR 3248197 15
43. Robin Pemantle, *Towards a theory of negative dependence*, vol. 41, 2000, Probabilistic techniques in equilibrium and nonequilibrium statistical physics, pp. 1371–1390. MR 1757964 1
44. Robin Pemantle and Yuval Peres, *Concentration of Lipschitz functionals of determinantal and other strong Rayleigh measures*, Combin. Probab. Comput. **23** (2014), no. 1, 140–160. MR 3197973 1, 2, 3, 4, 10
45. Holger Sambale and Arthur Sinulis, *Modified log-Sobolev inequalities and two-level concentration*, arXiv preprint arXiv:1905.06137 (2019). 15
46. ———, *Concentration inequalities on the multislice and for sampling without replacement*, arXiv preprint arXiv:2010.16289 (2020). 1, 2, 6, 15, 17
47. ———, *Logarithmic Sobolev inequalities for finite spin systems and applications*, Bernoulli **26** (2020), no. 3, 1863–1890. MR 4091094 6, 8
48. Paul-Marie Samson, *Transport-entropy inequalities on locally acting groups of permutations*, Electron. J. Probab. **22** (2017), Paper No. 62, 33. MR 3690287 1, 2
49. Qi-Man Shao, *A comparison theorem on moment inequalities between negatively associated and independent random variables*, J. Theoret. Probab. **13** (2000), no. 2, 343–356. MR 1777538 1

50. Michel Talagrand, *An isoperimetric theorem on the cube and the Kintchine-Kahane inequalities*, Proc. Amer. Math. Soc. **104** (1988), no. 3, 905–909. MR 964871 5
51. ———, *Concentration of measure and isoperimetric inequalities in product spaces*, Inst. Hautes Études Sci. Publ. Math. (1995), no. 81, 73–205. MR 1361756 2
52. Yves Tillé, *Sampling algorithms*, Springer Series in Statistics, Springer, New York, 2006. MR 2225036 2
53. Joel A. Tropp, *Freedman’s inequality for matrix martingales*, Electron. Commun. Probab. **16** (2011), 262–270. MR 2802042 12
54. ———, *User-friendly tail bounds for sums of random matrices*, Found. Comput. Math. **12** (2012), no. 4, 389–434. MR 2946459 2, 11
55. Roman Vershynin, *High-dimensional probability*, Cambridge Series in Statistical and Probabilistic Mathematics, vol. 47, Cambridge University Press, Cambridge, 2018, An introduction with applications in data science, With a foreword by Sara van de Geer. MR 3837109 6
56. Nicolas Verzelen and Elisabeth Gassiat, *Adaptive estimation of high-dimensional signal-to-noise ratios*, Bernoulli **24** (2018), no. 4B, 3683–3710. MR 3788186 6

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, POLAND
Email address: R.Adamczak, B.Polaczyk @mimuw.edu.pl