

# Extremal Polygonal Cacti for General Sombor Index\*

Jiachang Ye, Jianguo Qian<sup>†</sup>

School of Mathematical Sciences, Xiamen University,

Xiamen, 361005, P.R. China

## Abstract

The Sombor index of a graph  $G$  was recently introduced by Gutman from the geometric point of view, defined as  $SO(G) = \sum_{uv \in E(G)} \sqrt{d(u)^2 + d(v)^2}$ , where  $d(u)$  is the degree of a vertex  $u$ . For two real numbers  $\alpha$  and  $\beta$ , the  $\alpha$ -Sombor index and general Sombor index of  $G$  are two generalized forms of the Sombor index defined as  $SO_\alpha(G) = \sum_{uv \in E(G)} (d(u)^\alpha + d(v)^\alpha)^{1/\alpha}$  and  $SO_\alpha(G; \beta) = \sum_{uv \in E(G)} (d(u)^\alpha + d(v)^\alpha)^\beta$ , respectively. A  $k$ -polygonal cactus is a connected graph in which every block is a cycle of length  $k$ . In this paper, we establish a lower bound on  $\alpha$ -Sombor index for  $k$ -polygonal cacti and show that the bound is attained only by chemical  $k$ -polygonal cacti. The extremal  $k$ -polygonal cacti for  $SO_\alpha(G; \beta)$  with some particular  $\alpha$  and  $\beta$  are also considered.

**Keywords:** general Sombor index, polygonal cactus, extremal problem

**Mathematics Subject Classification:** 05C07, 05C09, 05C90

## 1 Introduction

We consider only connected simple graphs. For a graph  $G$ , we denote by  $V(G)$  and  $E(G)$  the vertex set and edge set of  $G$ , respectively. For a vertex  $v \in V(G)$ , we denote by  $d_G(v)$ , or  $d(v)$  if no confusion can occur, the degree of  $v$ . A vertex  $v$  is called a *cut vertex* of  $G$  if  $G - v$  is not connected.

In mathematical chemistry, particularly in QSPR/QSAR investigation, a large number of topological indices were introduced in an attempt to characterize the physical-chemical

---

\*This paper was supported by the National Natural Science Foundation of China [No. 11971406].

<sup>†</sup>Corresponding author. E-mail: jgqian@xmu.edu (J.G. Qian)

properties of molecules. Among these indices, the vertex-degree-based indices play important roles [4, 7, 8]. Probably the most studied are, for examples, the Randić connectivity index  $R(G)$  [18], the first and second Zagreb indices  $M_1(G)$  and  $M_2(G)$  [9], which were introduced for the total  $\pi$ -energy of alternant hydrocarbons.

A vertex-degree-based index of a graph  $G$  can be generally represented as the sum of a real function  $f(d(u), d(v))$  associated with the edges of  $G$  [10], i.e.,

$$I_f(G) = \sum_{uv \in E(G)} f(d(u), d(v)),$$

where  $f(s, t) = f(t, s)$ . In the literature,  $I_f(G)$  is also called the *connectivity function* [22] or *bond incident degree index* [1, 21, 23].

Recently, Gutman [10] introduced an idea to view an edge  $e = uv$  as a geometric point, namely the degree-point, that is, to view the ordered pair  $(d(u), d(v))$  as the coordinate of  $e$ . Therefore, it is interesting to consider the function  $f(s, t)$  from the geometric point of view. A natural considering is to define  $f(s, t)$  as a geometric distance from the degree-point  $(s, t)$  to the origin. In this sense, the first Zagreb index, i.e.,  $M_1(G) = \sum_{uv \in E(G)} (d(u) + d(v)) = \sum_{uv \in E(G)} (|d(u)| + |d(v)|)$ , is exactly the index defined on the Manhattan distance. Along this direction, a more natural considering would be to define  $f(s, t)$  as the Euclidean distance, i.e.,  $f(s, t) = \sqrt{s^2 + t^2}$ . Indeed, based on this idea, Gutman [10] introduced the *Sombor index* defined by

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d(u)^2 + d(v)^2}$$

and further determined the extremal trees for the index. In [3], Das et al. established some bounds on the Sombor index and some relations between Sombor index and the Zagreb indices and, in [19], Redžpović studied chemical applicability of the Sombor index. Further, Cruz et al. [2] characterized the extremal chemical graphs and hexagonal systems for the Sombor index.

More recently, for positive real number  $\alpha$ , Réti et al. [20] defined the  $\alpha$ -*Sombor index* as

$$SO_\alpha(G) = \sum_{uv \in E(G)} (d(u)^\alpha + d(v)^\alpha)^{1/\alpha},$$

which could be viewed as the one based on Minkowski distance. In the same paper, they also characterized the extremal graphs with few cycles for  $\alpha$ -Sombor index.

In this paper we consider a more generalized form of Sombor index defined as

$$SO_\alpha(G; \beta) = \sum_{uv \in E(G)} (d(u)^\alpha + d(v)^\alpha)^\beta,$$

where  $\alpha, \beta$  are real numbers. We note that this form is a natural generalization of the Sombor index, which was also introduced elsewhere, e.g., the first  $(\alpha, \beta) - KA$  index in [12] and the general Sombor index in [11]. In addition to the first Zagreb, Sombor and the  $\alpha$ -Sombor index listed above, the general Sombor index also includes many other known indices, e.g., the modified first Zagreb index  $(\alpha = -3, \beta = 1)$  [17], forgotten index  $(\alpha = 2, \beta = 1)$  [6], inverse degree index  $(\alpha = -2, \beta = 1)$  [5], modified Sombor index  $(\alpha = 2, \beta = -1/2)$  [13], first Banhatti-Sombor index  $(\alpha = -2, \beta = 1/2)$  [15] and general sum-connectivity index  $(\alpha = 1, \beta \in \mathbb{R})$  [24].

A *block* in a graph is a cut edge or a maximal 2-connected component. A *cactus* is a connected graph in which every block is a cut edge or a cycle. Equivalently, a cactus has no edge lies in more than one cycle. In the following, we call a  $k$ -cycle (a cycle of length  $k$ ) a  $k$ -*polygon*. If each block of a cactus  $G$  is a  $k$ -polygon, then  $G$  is called a  $k$ -*polygonal cactus* or *polygonal cactus* with no confusion.

In this paper, we consider the extremal  $k$ -polygonal cacti for  $SO_\alpha(G; \beta)$ . In the following section we establish a lower bound on  $\alpha$ -Sombor index for  $k$ -polygonal cacti and show that the bound is attained only by chemical  $k$ -polygonal cacti. In the third section we characterize the extremal polygonal cactus with maximum  $SO_\alpha(G; \beta)$  for (i)  $\alpha > 1$  and  $\beta \geq 1$ ; and (ii)  $1/2 \leq \alpha < 1$  and  $\beta = 2$ , respectively. In the fourth section, we characterize the extremal polygonal cacti with minimum  $SO_\alpha(G; \beta)$  for  $\alpha > 1$  and  $\beta \geq 1$ .

## 2 Polygonal cacti with minimum $\alpha$ -Sombor index

For convenience, in what follows we denote  $r_\alpha(s, t; \beta) = (s^\alpha + t^\alpha)^\beta$ ,  $r_\alpha(s, t) = (s^\alpha + t^\alpha)^{1/\alpha}$  and  $r(s, t) = \sqrt{s^2 + t^2}$ , where  $s > 0, t > 0, \alpha, \beta \in \mathbb{R}$  and  $\alpha \neq 0$ . For integers  $n$  with  $n \geq 1$  and  $k$  with  $k \geq 3$ , we denote by  $\mathcal{G}_{n,k}$  the class of  $k$ -polygonal cacti with  $n$  polygons.

In this section we consider the  $\alpha$ -Sombor index  $SO_\alpha(G)$ , i.e.,  $SO_\alpha(G; 1/\alpha)$ . For  $G \in \mathcal{G}_{n,k}$ , it is clear that  $|V(G)| = nk - n + 1, |E(G)| = nk$  and every vertex of  $G$  has even degree no more than  $2n$ . Further, it is clear that  $v$  is a cut vertex of  $G$  if and only if  $v$  has degree

no less than 4, i.e.,  $d_G(v) \geq 4$ . A polygon is called a *pendent polygon* if it contains exactly one cut-vertex of  $G$ . For  $2 \leq s \leq t \leq 2n$ , we denote by  $n_{s,t} = n_{s,t}(G)$  the number of edges in  $G$  that join two vertices of degrees  $s$  and  $t$ . Let  $X = \{(s, t) : s, t \in \{2, 4, \dots, 2n\}, s \leq t\}$  and  $Y = X \setminus \{(2, 2), (2, 4), (4, 4)\}$ .

**Definition 2.1.** [16] Let  $\pi = (w_1, w_2, \dots, w_n)$  and  $\pi' = (w'_1, w'_2, \dots, w'_n)$  be two non-increasing sequences of nonnegative real numbers. We write  $\pi \triangleleft \pi'$  if and only if  $\pi \neq \pi'$ ,  $\sum_{i=1}^n w_i = \sum_{i=1}^n w'_i$ , and  $\sum_{i=1}^j w_i \leq \sum_{i=1}^j w'_i$  for all  $j = 1, 2, \dots, n$ .

A function  $\zeta(x)$  defined on a convex set  $X$  is called *strictly convex* if

$$\zeta(\mu x_1 + (1 - \mu)x_2) < \mu\zeta(x_1) + (1 - \mu)\zeta(x_2) \quad (1)$$

for any  $0 < \mu < 1$  and  $x_1, x_2 \in X$  with  $x_1 \neq x_2$ .

**Lemma 2.1.** [16] Let  $\pi = (w_1, w_2, \dots, w_n)$  and  $\pi' = (w'_1, w'_2, \dots, w'_n)$  be two non-increasing sequences of nonnegative real numbers. If  $\pi \triangleleft \pi'$ , then for any strictly convex function  $\zeta(x)$ , we have  $\sum_{i=1}^n \zeta(w_i) < \sum_{i=1}^n \zeta(w'_i)$ .

**Lemma 2.2.** Let  $\alpha > 1$  and  $n \geq 3$ . Then

- (i).  $r_\alpha(2n, 2) - r_\alpha(2n - 2, 4) > 0$ ;
- (ii).  $r_\alpha(6, 2) + r_\alpha(2, 2) - 2r_\alpha(4, 2) > 0$ .

*Proof.* Since  $\alpha > 1$  and  $n \geq 3$ , then by Lemma 2.1, we have  $(2n)^\alpha + 2^\alpha > (2n - 2)^\alpha + 4^\alpha$ . Hence (i) holds clearly. Let  $g(x) = r_\alpha(x, 2) = (x^\alpha + 2^\alpha)^{1/\alpha}$ , where  $x > 0$  and  $\alpha > 1$ . Since  $g''(x) = \frac{(\alpha-1)2^\alpha x^{\alpha-2}}{(x^\alpha + 2^\alpha)^{2-1/\alpha}} > 0$ , then  $g(x)$  is strictly convex. Then by Lemma 2.1,  $g(6) + g(2) > 2g(4)$ . Hence (ii) also holds.  $\square$

**Lemma 2.3.** Let  $\alpha \neq 0$  and  $G \in \mathcal{G}_{n,k}$ , where  $n \geq 1$  and  $k \geq 3$ . Then

$$\begin{aligned} SO_\alpha(G) &= (4n - 4)(2^\alpha + 4^\alpha)^{1/\alpha} + 2(nk - 4n + 4)2^{1/\alpha} \\ &\quad + (6 \times 2^{1/\alpha} - 2(2^\alpha + 4^\alpha)^{1/\alpha}) n_{4,4} + \sum_{(s,t) \in Y} \eta(s, t; \alpha) n_{s,t}, \end{aligned}$$

where  $\eta(s, t; \alpha) = (s^\alpha + t^\alpha)^{1/\alpha} - 2 \left( \frac{1}{s} + \frac{1}{t} \right) 2^{1/\alpha}$ .

*Proof.* By the definition of  $n_{s,t}$ , it is not difficult to see that

$$\begin{cases} nk - n + 1 &= \sum_{(s,t) \in X} \left( \frac{1}{s} + \frac{1}{t} \right) n_{s,t}, \\ nk &= \sum_{(s,t) \in X} n_{s,t} \end{cases} \quad (2)$$

because each vertex of  $G$  contributes 1 to each of the two sides in the first equation and each edge of  $G$  contributes 1 to each of the two sides in the second equation. Write (2) as

$$\begin{cases} 4n_{2,2} + 3n_{2,4} &= 4(nk - n + 1) - 2n_{4,4} - 4 \sum_{(s,t) \in Y} \left(\frac{1}{s} + \frac{1}{t}\right) n_{s,t}, \\ n_{2,2} + n_{2,4} &= nk - n_{4,4} - \sum_{(s,t) \in Y} \left(\frac{1}{s} + \frac{1}{t}\right) n_{s,t}. \end{cases} \quad (3)$$

Therefore,

$$\begin{cases} n_{2,4} &= 4n - 4 - 2n_{4,4}, \\ n_{2,2} &= nk - 4n + 4 + n_{4,4} - \sum_{(s,t) \in Y} \left(\frac{1}{s} + \frac{1}{t}\right) n_{s,t}. \end{cases} \quad (4)$$

Consequently, by (4) we have

$$\begin{aligned} SO_\alpha(G) &= (4^\alpha + 4^\alpha)^{1/\alpha} n_{4,4} + (2^\alpha + 4^\alpha)^{1/\alpha} n_{2,4} + (2^\alpha + 2^\alpha)^{1/\alpha} n_{2,2} \\ &\quad + \sum_{(s,t) \in Y} (s^\alpha + t^\alpha)^{1/\alpha} n_{s,t} \\ &= (4n - 4)(2^\alpha + 4^\alpha)^{1/\alpha} + 2(nk - 4n + 4)2^{1/\alpha} \\ &\quad + (6 \times 2^{1/\alpha} - 2(2^\alpha + 4^\alpha)^{1/\alpha}) n_{4,4} \\ &\quad + \sum_{(s,t) \in Y} \left( (s^\alpha + t^\alpha)^{1/\alpha} - 2 \left( \frac{1}{s} + \frac{1}{t} \right) 2^{1/\alpha} \right) n_{s,t}. \end{aligned}$$

□

For  $\alpha \neq 0$  and positive integer  $p$ , let  $\delta_{\alpha,p}(s, t) = ((s+p)^\alpha + t^\alpha)^{1/\alpha} - (s^\alpha + t^\alpha)^{1/\alpha}$ , where  $s, t > 0$ .

**Lemma 2.4.** *If  $s, t > 0$  and  $p$  is an arbitrary positive integer, then*

- (i).  $r_\alpha(s, t; \beta)$  strictly increases in  $s$  for fixed  $t$ , and in  $t$  for fixed  $s$  when  $\alpha, \beta > 0$ ;
- (ii).  $\delta_{\alpha,p}(s, t) > 0$  and  $\delta_{\alpha,p}(s, t)$  strictly decreases in  $t$  for fixed  $s$  when  $\alpha > 1$ ;
- (iii).  $\delta_{\alpha,p}(s, t)$  strictly increases in  $s$  for fixed  $t$  when  $\alpha > 1$ .

*Proof.* (i) follows directly since  $\alpha, \beta > 0$ .

Since  $-1 < 1/\alpha - 1 < 0$  when  $\alpha > 1$ ,

$$\frac{\partial \delta_{\alpha,1}(s, t)}{\partial t} = t^{\alpha-1} \left( ((s+1)^\alpha + t^\alpha)^{1/\alpha-1} - (s^\alpha + t^\alpha)^{1/\alpha-1} \right) < 0.$$

We also note that  $\delta_{\alpha,1}(s, t) > 0$ , hence (ii) follows as  $\delta_{\alpha,p}(s, t) = \sum_{i=0}^{p-1} \delta_{\alpha,1}(s+i, t)$ .

Finally, since  $1 - 1/\alpha > 0$  when  $\alpha > 1$ ,

$$\frac{\partial \delta_{\alpha,1}(s, t)}{\partial s} = \frac{((s+1)^\alpha s^\alpha + (s+1)^\alpha t^\alpha)^{1-1/\alpha} - ((s+1)^\alpha s^\alpha + s^\alpha t^\alpha)^{1-1/\alpha}}{(((s+1)^\alpha + t^\alpha)(s^\alpha + t^\alpha))^{1-1/\alpha}} > 0.$$

Hence (iii) follows as  $\delta_{\alpha,p}(s,t) = \sum_{i=0}^{p-1} \delta_{\alpha,1}(s+i,t)$ .  $\square$

The *distance*  $d_G(u,v)$  between two vertices  $u$  and  $v$  of a connected graph  $G$  is defined as usual as the length of a shortest path that connects  $u$  and  $v$ . In general, for two subgraphs  $G_1$  and  $G_2$  of  $G$ , we define the distance between  $G_1$  and  $G_2$  by  $d_G(G_1, G_2) = \min\{d_G(u,v) : u \in V(G_1), v \in V(G_2)\}$ . For  $n \geq 2$ , a *star-like cactus*  $S_{n,k}$  is defined intuitively as a  $k$ -polygonal cactus such that all polygons have a vertex in common. It is clear that  $S_{n,k}$  is unique and contains exactly one vertex of degree  $2n$  while all other vertices have degree two.

**Lemma 2.5.** *Let  $\alpha > 1$  and  $G \in \mathcal{G}_{n,k}$ , where  $n \geq 3$  and  $k \geq 3$ . If  $G$  contains a vertex of degree at least 6, then  $SO_\alpha(G)$  is not minimum in  $\mathcal{G}_{n,k}$ .*

*Proof.* Suppose to the contrary that  $G$  contains  $q$  ( $q \geq 1$ ) vertices of degree at least 6 and  $SO_\alpha(G)$  is minimum in  $\mathcal{G}_{n,k}$ .

**Case 1.**  $q \geq 2$ .

Let  $u_1$  and  $w$  be two vertices of degree at least 6 such that  $d_G(u_1, w)$  is maximum and let  $P$  be a shortest path connecting  $u_1$  and  $w$ . Since  $d_G(u_1) \geq 6$ ,  $u_1$  is contained in at least three polygons, exactly one of which, say  $C$ , has at least two common vertices with  $P$ . Let  $C_1 = u_1 u_2 \cdots u_k u_1$  and  $C_2 = u_1 z_2 \cdots z_k u_1$  be two polygons other than  $C$  that contain  $u_1$  as a common vertex. Since  $d_G(u_1, w)$  is maximum, we have  $d_G(v) \leq 4$  for every  $v \in \{u_2, u_k, z_2, z_k\}$ . Let  $C_3 = v_1 v_2 \cdots v_k v_1$  be a pendent polygon that lies in the same component with  $w$  in  $G - u_1$  and the distance  $d_G(u_1, C_3)$  is as large as possible, where  $d_G(v_1) = 2a \geq 4$  and  $d_G(v_2) = d_G(v_3) = 2$ .

Without loss of generality, assume  $d_G(u_1) = 2b \geq d_G(w) \geq 6$ . Let  $G' = G - u_1 u_2 - u_1 u_k + v_2 u_2 + v_2 u_k$ . Then by Lemma 2.4, we have

$$\begin{aligned}
SO_\alpha(G) - SO_\alpha(G') &> (r_\alpha(2b, d(u_2)) - r_\alpha(4, d(u_2))) + (r_\alpha(2b, d(u_k)) - r_\alpha(4, d(u_k))) \\
&+ (r_\alpha(2, 2) - r_\alpha(4, 2)) + (r_\alpha(2, 2a) - r_\alpha(4, 2a)) \\
&+ (r_\alpha(2b, d(z_2)) - r_\alpha(2b - 2, d(z_2))) \\
&+ (r_\alpha(2b, d(z_k)) - r_\alpha(2b - 2, d(z_k))) \\
&> (r_\alpha(6, 4) - r_\alpha(4, 4)) + (r_\alpha(6, 4) - r_\alpha(4, 4)) \\
&+ (r_\alpha(2, 2) - r_\alpha(4, 2)) + (r_\alpha(2, 4) - r_\alpha(4, 4)) \\
&+ (r_\alpha(6, 4) - r_\alpha(4, 4)) + (r_\alpha(6, 4) - r_\alpha(4, 4)) \\
&= 8(3^\alpha + 2^\alpha)^{1/\alpha} - 18 \times 2^{1/\alpha} \\
&> 8\sqrt{6} \times 2^{1/\alpha} - 18 \times 2^{1/\alpha} \\
&> 0,
\end{aligned}$$

which contradicts the minimality of  $G$ .

**Case 2.**  $q = 1$ .

If  $G \not\cong S_{n,k}$ , then the discussion for this case is similar to that for Case 1 by choosing  $u_1$  to be the vertex with degree at least 6 and  $C_3 = v_1v_2 \cdots v_kv_1$  to be a pendent polygon such that  $d_G(u_1, C_3)$  is maximum. Otherwise,  $G \cong S_{n,k}$ . Let  $u_1$  be the vertex with degree  $2n \geq 6$ ,  $C_1 = u_1u_2 \cdots u_ku_1$  and  $C_2 = u_1z_2 \cdots z_ku_1$  be two pendent polygons. Let  $G' = G - u_1u_2 - u_1u_k + z_2u_2 + z_ku_k$ . Then by Lemma 2.4 and Lemma 2.2, we have

$$\begin{aligned}
SO_\alpha(G) - SO_\alpha(G') &= (2n \times r_\alpha(2n, 2) + (nk - 2n) \times r_\alpha(2, 2)) \\
&- ((2n - 3) \times r_\alpha(2n - 2, 2) + r_\alpha(2n - 2, 4)) \\
&+ 3r_\alpha(4, 2) + (nk - 2n - 1)r_\alpha(2, 2)) \\
&= 2n \times r_\alpha(2n, 2) + r_\alpha(2, 2) - (2n - 3) \times r_\alpha(2n - 2, 2) \\
&- r_\alpha(2n - 2, 4) - 3r_\alpha(4, 2) \\
&> 3r_\alpha(2n, 2) + r_\alpha(2, 2) - r_\alpha(2n - 2, 4) - 3r_\alpha(4, 2) \\
&> 2r_\alpha(2n, 2) + r_\alpha(2, 2) - 3r_\alpha(4, 2) \\
&> r_\alpha(6, 2) + r_\alpha(2, 2) - 2r_\alpha(4, 2) \\
&> 0,
\end{aligned}$$

which contradicts the minimality of  $G$ .

□

Recall that a graph is called a chemical graph if it has no vertex of degree more than 4. For  $G \in \mathcal{G}_{n,k}$ , we call  $G$  a *chemical  $(n, k)$ -cactus*, or *chemical cactus* for short, if  $G$  has no vertex of degree greater than 4. It is clear that every cut vertex in a chemical cactus has degree 4, which connects exactly two polygons. The following corollary follows directly from Lemma 2.3 and Lemma 2.5, which shows that the minimum value of  $SO_\alpha(G)$  among all cacti in  $\mathcal{G}_{n,k}$  is attained only by chemical cacti.

**Corollary 2.1.** *For  $\alpha > 1$ ,  $n \geq 3$ ,  $k \geq 3$  and  $G \in \mathcal{G}_{n,k}$ , if  $G$  attains the minimum value of  $SO_\alpha(G)$ , then  $G$  is a chemical cactus and*

$$SO_\alpha(G) = (4n - 4)(2^\alpha + 4^\alpha)^{1/\alpha} + 2(nk - 4n + 4)2^{1/\alpha} + (6 \times 2^{1/\alpha} - 2(2^\alpha + 4^\alpha)^{1/\alpha}) n_{4,4}(G).$$

In the following we will determine the minimum value of  $SO_\alpha(G)$  among all chemical cacti. By Corollary 2.1, this is equivalent to determine the maximum value of  $n_{4,4}(G)$  as  $6 \times 2^{1/\alpha} - 2(2^\alpha + 4^\alpha)^{1/\alpha} < 6 \times 2^{1/\alpha} - 2(2 \times 3^\alpha)^{1/\alpha} = 0$  by Lemma 2.1. For a chemical cactus  $H$ , we call a polygon  $C$  in  $H$  a *saturated polygon* if every vertex on  $C$  is a cut vertex, i.e., a vertex of degree 4. Further, we call a chemical cactus  $H$  *nice-saturated* if the following two conditions hold:

- 1).  $H$  has as many as possible saturated polygons;
- 2). the cut vertices on each polygon of  $H$  are successively arranged.

For a chemical cactus  $H$ , let  $T(H)$  be the tree whose vertices are the polygons in  $H$  and two vertices are adjacent provided their corresponding polygons has a common vertex. It is clear that  $T(H)$  is a tree with maximum vertex degree no more than  $k$ . Let  $p$  be the number of the vertices of degree  $k$  in  $T(H)$ , and let  $d_1, d_2, \dots, d_s$  be the degrees of all the vertices in  $H$  that are neither of degree 1 nor of degree  $k$ , i.e.,  $1 < d_i < k$  for each  $i \in \{1, 2, \dots, s\}$ . Since  $T(H)$  is a tree, we have

$$kp + d_1 + d_2 + \dots + d_s + (n - p - s) = 2n - 2 \quad (5)$$

and every saturated polygon in  $H$  corresponds to a vertex of degree  $k$  in  $T(H)$ . Further,  $T(H)$  has as many as possible vertices of degree  $k$  if and only if  $d_1 + d_2 + \dots + d_s - s < k - 1$ .

This implies that

$$\frac{n-2}{k-1} - 1 < p = \frac{n-2 - (d_1 + d_2 + \dots + d_s - s)}{k-1} \leq \frac{n-2}{k-1}. \quad (6)$$



$$+ (6 \times 2^{1/\alpha} - 2(2^\alpha + 4^\alpha)^{1/\alpha}) \left( n - 2 + \left\lfloor \frac{n-2}{k-1} \right\rfloor \right),$$

and the equality holds if and only if  $G$  is a nice-saturated chemical cactus.

*Proof.* By (7), (5) and (6), if  $G$  is minimum, then

$$n_{4,4}(G) = kp + (d_1 - 1) + (d_2 - 1) + \cdots + (d_s - 1) = n - 2 + p = n - 2 + \left\lfloor \frac{n-2}{k-1} \right\rfloor.$$

Hence, the theorem follows directly from Corollary 2.1 and Lemma 2.6.  $\square$

### 3 Polygonal cactus with maximum general Sombor index

In this section we will characterize the polygonal cactus with maximum general Sombor index for the two cases  $\alpha \geq 1, \beta > 1$ ; and  $\alpha = 2, 1/2 \leq \beta < 1$ , respectively.

**Lemma 3.1.** *Let  $\triangle ABM$  be a triangle in Euclidean space and  $O$  the midpoint of the triangle side  $AB$ . Then  $|MA|^{2\beta} + |MB|^{2\beta} > 2|MO|^{2\beta}$  for any real number  $\beta \geq \frac{1}{2}$ , where  $|MA|$  is the length of the side  $MA$ .*

*Proof.* Let  $|MA| = a$ ,  $|MB| = b$  and  $|MO| = d$ . When  $a = b$ , the lemma follows directly. Without loss of generality, we now assume  $a > b > 0$ . By the triangle inequality,  $d < \frac{a+b}{2} < a$  and so  $(a, b) \triangleright \left( \frac{a+b}{2}, \frac{a+b}{2} \right)$ . Hence, by Lemma 2.1,  $a^{2\beta} + b^{2\beta} \geq 2 \left( \frac{a+b}{2} \right)^{2\beta} > 2d^{2\beta}$  when  $\beta \geq \frac{1}{2}$ .  $\square$

**Lemma 3.2.** *Let  $s > 2$  and  $t > 2$ . Then*

- (i).  $r_\alpha(s+2, 2; \beta) - r_\alpha(s-2, 2; \beta) > 0$  for any  $\alpha > 0$  and  $\beta > 0$ ;
- (ii).  $r_\alpha(s+2, t; \beta) + r_\alpha(s-2, t; \beta) \geq 2r_\alpha(s, t; \beta)$  for any  $\alpha \geq 1$  and  $\beta > 1$ ;
- (iii).  $r_\alpha(s+2, t-2; \beta) + r_\alpha(s-2, t+2; \beta) \geq 2r_\alpha(s, t; \beta)$  for any  $\alpha \geq 1$  and  $\beta > 1$ .

*Proof.* (i) follows directly.

For (ii), by Lemma 2.1 and the monotonicity of  $r_\alpha(s, t; \beta)$ , we have

$$\begin{aligned} r_\alpha(s+2, t; \beta) + r_\alpha(s-2, t; \beta) &= ((s+2)^\alpha + t^\alpha)^\beta + ((s-2)^\alpha + t^\alpha)^\beta \\ &\geq 2 \left( \frac{(s+2)^\alpha + (s-2)^\alpha}{2} + t^\alpha \right)^\beta \\ &\geq 2(s^\alpha + t^\alpha)^\beta \\ &= 2r_\alpha(s, t; \beta). \end{aligned}$$

Hence, (ii) holds.

The discussion for (iii) is analogous to that for (ii).  $\square$

**Theorem 3.1.** *Let  $n \geq 3, k \geq 3$  and  $G \in \mathcal{G}_{n,k}$ . If  $\alpha \geq 1$  and  $\beta > 1$ ; or  $\alpha = 2$  and  $\frac{1}{2} \leq \beta < 1$ , then*

$$SO_\alpha(G; \beta) \leq 2n((2n)^\alpha + 2^\alpha)^\beta + n(k-2)(2^{\alpha+1})^\beta$$

and the equality holds if and only if  $G \cong S_{n,k}$ .

*Proof.* We first assume that  $\alpha \geq 1$  and  $\beta > 1$ .

Let  $G$  be such that  $SO_\alpha(G; \beta)$  is as large as possible. Further, let  $C_1 = z_1 z_2 \cdots z_k z_1$  and  $C_2 = v_1 v_2 \cdots v_k v_1$  be two pendent polygons such that  $d_G(C_1, C_2)$  is as large as possible, where  $z_1$  and  $v_1$  are the cut-vertices of  $C_1$  and  $C_2$ , respectively.

If  $G \cong S_{n,k}$ , then the theorem follows directly. We now assume  $G \not\cong S_{n,k}$ . Then,  $z_1 \neq v_1$ . Let  $G_1 = G - v_1 v_2 - v_1 v_k + z_1 v_2 + z_1 v_k$  and  $G_2 = G - z_1 z_2 - z_1 z_k + v_1 z_2 + v_1 z_k$ . We consider the following two cases:

**Case 1.**  $z_1$  and  $v_1$  are adjacent in  $G$ .

In this case, we have  $SO_\alpha(G_1; \beta) - SO_\alpha(G; \beta) =$

$$\begin{aligned} & \sum_{v \in N_G(v_1) \setminus \{z_1\}} (r_\alpha(d_G(v_1) - 2, d_G(v); \beta) - r_\alpha(d_G(v_1), d_G(v); \beta)) \\ & + \sum_{z \in N_G(z_1) \setminus \{v_1\}} (r_\alpha(d_G(z_1) + 2, d_G(z); \beta) - r_\alpha(d_G(z_1), d_G(z); \beta)) \\ & + 2r_\alpha(d_G(z_1) + 2, 2; \beta) - 2r_\alpha(d_G(v_1) - 2, 2; \beta) \\ & + r_\alpha(d_G(v_1) - 2, d_G(z_1) + 2; \beta) - r_\alpha(d_G(v_1), d_G(z_1); \beta), \text{ and} \end{aligned}$$

$SO_\alpha(G_2; \beta) - SO_\alpha(G; \beta) =$

$$\begin{aligned} & \sum_{v \in N_G(v_1) \setminus \{z_1\}} (r_\alpha(d_G(v_1) + 2, d_G(v); \beta) - r_\alpha(d_G(v_1), d_G(v); \beta)) \\ & + \sum_{z \in N_G(z_1) \setminus \{v_1\}} (r_\alpha(d_G(z_1) - 2, d_G(z); \beta) - r_\alpha(d_G(z_1), d_G(z); \beta)) \\ & + 2r_\alpha(d_G(v_1) + 2, 2; \beta) - 2r_\alpha(d_G(z_1) - 2, 2; \beta) \\ & + r_\alpha(d_G(v_1) + 2, d_G(z_1) - 2; \beta) - r_\alpha(d_G(v_1), d_G(z_1); \beta). \end{aligned}$$

Recall that  $z_1$  and  $v_1$  are the cut-vertices of  $C_1$  and  $C_2$ , respectively. Therefore,  $d_G(z_1) \geq 4$  and  $d_G(v_1) \geq 4$ . Combining with Lemma 3.2, we have

$$\begin{aligned} r_\alpha(d_G(v_1) + 2, d_G(v); \beta) + r_\alpha(d_G(v_1) - 2, d_G(v); \beta) &> 2r_\alpha(d_G(v_1), d_G(v); \beta), \\ r_\alpha(d_G(z_1) + 2, d_G(z); \beta) + r_\alpha(d_G(z_1) - 2, d_G(z); \beta) &> 2r_\alpha(d_G(z_1), d_G(z); \beta), \\ r_\alpha(d_G(v_1) + 2, d_G(z_1) - 2; \beta) + r_\alpha(d_G(v_1) - 2, d_G(z_1) + 2; \beta) &> 2r_\alpha(d_G(v_1), d_G(z_1); \beta), \\ r_\alpha(d_G(z_1) + 2, 2; \beta) - r_\alpha(d_G(z_1) - 2, 2; \beta) &> 0, \quad \text{and} \\ r_\alpha(d_G(v_1) + 2, 2; \beta) - r_\alpha(d_G(v_1) - 2, 2; \beta) &> 0. \end{aligned}$$

This means that  $SO_\alpha(G_1; \beta) > SO_\alpha(G; \beta)$  or  $SO_\alpha(G_2; \beta) > SO_\alpha(G; \beta)$ , a contradiction.

**Case 2.**  $z_1$  and  $v_1$  are not adjacent in  $G$ .

In this case, we have

$$\begin{aligned} SO_\alpha(G_1; \beta) - SO_\alpha(G; \beta) &= \sum_{v \in N_G(v_1)} (r_\alpha(d_G(v_1) - 2, d_G(v); \beta) - r_\alpha(d_G(v_1), d_G(v); \beta)) \\ &\quad + \sum_{z \in N_G(z_1)} (r_\alpha(d_G(z_1) + 2, d_G(z); \beta) - r_\alpha(d_G(z_1), d_G(z); \beta)) \\ &\quad + 2r_\alpha(d_G(z_1) + 2, 2; \beta) - 2r_\alpha(d_G(v_1) - 2, 2; \beta), \quad \text{and} \\ SO_\alpha(G_2; \beta) - SO_\alpha(G; \beta) &= \sum_{v \in N_G(v_1)} (r_\alpha(d_G(v_1) + 2, d_G(v); \beta) - r_\alpha(d_G(v_1), d_G(v); \beta)) \\ &\quad + \sum_{z \in N_G(z_1)} (r_\alpha(d_G(z_1) - 2, d_G(z); \beta) - r_\alpha(d_G(z_1), d_G(z); \beta)) \\ &\quad + 2r_\alpha(d_G(v_1) + 2, 2; \beta) - 2r_\alpha(d_G(z_1) - 2, 2; \beta). \end{aligned}$$

Recall that  $d_G(z_1) \geq 4$  and  $d_G(v_1) \geq 4$ . Similar to Case 1, by Lemma 3.2, we have

$$\begin{aligned} r_\alpha(d_G(v_1) + 2, d_G(v); \beta) + r_\alpha(d_G(v_1) - 2, d_G(v); \beta) &> 2r_\alpha(d_G(v_1), d_G(v); \beta), \\ r_\alpha(d_G(z_1) + 2, d_G(z); \beta) + r_\alpha(d_G(z_1) - 2, d_G(z); \beta) &> 2r_\alpha(d_G(z_1), d_G(z); \beta), \\ r_\alpha(d_G(z_1) + 2, 2; \beta) - r_\alpha(d_G(z_1) - 2, 2; \beta) &> 0 \quad \text{and} \\ r_\alpha(d_G(v_1) + 2, 2; \beta) - r_\alpha(d_G(v_1) - 2, 2; \beta) &> 0, \end{aligned}$$

which means that  $SO_\alpha(G_1; \beta) > SO_\alpha(G; \beta)$  or  $SO_\alpha(G_2; \beta) > SO_\alpha(G; \beta)$ , a contradiction.

Therefore,  $S_{n,k}$  is the unique maximal polygonal cactus. Further, we have

$$SO_\alpha(S_{n,k}; \beta) = 2nr_\alpha(2n, 2; \beta) + n(k-2)r_\alpha(2, 2; \beta) = 2n((2n)^\alpha + 2^\alpha)^\beta + n(k-2)(2^{\alpha+1})^\beta.$$

The discussion for the case that  $\alpha = 2$  and  $\frac{1}{2} \leq \beta < 1$  is analogous by Lemma 3.2 (i) and Lemma 3.1.  $\square$

## 4 Polygonal cacti with minimum general Sombor index

A symmetric function  $\varphi(s, t)$  defined on positive real numbers is called *escalating* [22] if

$$\varphi(s_1, s_2) + \varphi(t_1, t_2) \geq \varphi(s_2, t_1) + \varphi(s_1, t_2) \quad (8)$$

for any  $s_1 \geq t_1 > 0$  and  $s_2 \geq t_2 > 0$ , and the inequality holds if  $s_1 > t_1 > 0$  and  $s_2 > t_2 > 0$ . Further, an escalating function  $\varphi(s, t)$  is called *special escalating* [23] if

$$4\varphi(2l, 2) - \varphi(2l - 2, 4) - \varphi(2l - 2, 2) - \varphi(4, 2) - \varphi(4, 4) \geq 0 \quad (9)$$

for  $l \geq 3$  and

$$\varphi(s_1, s_2) - \varphi(t_1, t_2) \geq 0 \quad (10)$$

for any  $s_1 \geq t_1 \geq 2$  and  $s_2 \geq t_2 \geq 2$ .

**Lemma 4.1.** *If  $s, t > 0$ , then  $r_\alpha(s, t; \beta) = (s^\alpha + t^\alpha)^\beta$  is special escalating for  $\alpha \geq 1$  and  $\beta > 1$ .*

*Proof.* Set  $\varphi(s, t) = (s^\alpha + t^\alpha)^\beta$ . Since  $\alpha \geq 1$  and  $\beta > 1$ , we have  $(s_1^\alpha + s_2^\alpha, t_1^\alpha + t_2^\alpha) \triangleright (s_2^\alpha + t_1^\alpha, s_1^\alpha + t_2^\alpha)$  when  $s_1 > t_1 > 0$  and  $s_2 > t_2 > 0$ . Then by Lemma 2.1, the inequality in (8) strictly holds. Further, it is clear that the equality in (8) holds when  $s_1 = t_1 > 0$  or  $s_2 = t_2 > 0$ . This means that  $(s^\alpha + t^\alpha)^\beta$  is escalating.

In addition, by Lemma 2.1 and the monotonicity of  $(s^\alpha + t^\alpha)^\beta$ , if  $l \geq 3$  and  $\alpha \geq 1$  then  $(2l)^\alpha + 2^\alpha \geq (2l - 2)^\alpha + 4^\alpha > (2l - 2)^\alpha + 2^\alpha$ ,  $(2l)^\alpha + 2^\alpha > 4^\alpha + 2^\alpha$  and  $(2l)^\alpha + 2^\alpha \geq 6^\alpha + 2^\alpha \geq 4^\alpha + 4^\alpha$ . Hence, (9) follows directly as  $\beta > 1$ .

Finally, it is easy to see that (10) holds when  $\alpha \geq 1$  and  $\beta > 1$  by the monotonicity of  $(s^\alpha + t^\alpha)^\beta$ . Therefore,  $(s^\alpha + t^\alpha)^\beta$  is special escalating.  $\square$

A  $k$ -polygonal cactus  $G$  is called a *cactus chain* if each polygon in  $G$  has at most two cut-vertices and each cut-vertex is the common vertex of exactly two polygons. It is clear

that each cactus chain has exactly  $n - 2$  non-pendent polygons and two pendent polygons for  $n \geq 2$ . We denote by  $\mathcal{A}_{n,k}$  the class consisting of those cactus chains such that each pair of cut-vertices that lies in the same polygon of  $G$  are adjacent. In contrast, we denote by  $\mathcal{B}_{n,k}$  the class consisting of those cactus chains such that each pair of cut-vertices that lies in the same polygon of  $G$  are not adjacent. It can be seen that  $\mathcal{A}_{n,k}$  is unique for  $k \geq 3$  and  $\mathcal{B}_{n,3} = \emptyset$ .

**Theorem 4.1.** [23] *Let  $f(s, t)$  be a special escalating function and  $G$  be a cactus of  $\mathcal{G}_{n,k}$ , where  $n \geq 3$  and  $k \geq 3$ .*

(i). *If  $k = 3$ , then*

$$I_f(G) \geq 2f(2, 2) + 2nf(4, 2) + (n - 2)f(4, 4)$$

*with equality holding if and only if  $G \in \mathcal{A}_{n,3}$ .*

(ii). *If  $k \geq 4$ , then*

$$I_f(G) \geq (kn - 4n + 4)f(2, 2) + (4n - 4)f(4, 2),$$

*where the equality holds if  $G \in \mathcal{B}_{n,k}$ . Furthermore, if  $k \in \{4, 5\}$ , then the equality holds if and only if  $G \in \mathcal{B}_{n,k}$ .*

**Corollary 4.1.** *Let  $n \geq 3, k \geq 3, \alpha \geq 1, \beta > 1$  and  $G \in \mathcal{G}_{n,k}$ .*

(i). *If  $k = 3$ , then  $SO_\alpha(G; \beta) \geq 2(2^{\alpha+1})^\beta + 2n(4^\alpha + 2^\alpha)^\beta + (n - 2)(2 \cdot 4^\alpha)^\beta$ , where the equality holds if and only if  $G \in \mathcal{A}_{n,3}$ .*

(ii). *If  $k \geq 4$ , then  $SO_\alpha(G; \beta) \geq (kn - 4n + 4)(2^{\alpha+1})^\beta + (4n - 4)(4^\alpha + 2^\alpha)^\beta$ , where the equality holds if  $G \in \mathcal{B}_{n,k}$ . Furthermore, if  $k \in \{4, 5\}$ , then the equality holds if and only if  $G \in \mathcal{B}_{n,k}$ .*

*Proof.* In Theorem 4.1, set  $f(s, t) = r_\alpha(s, t; \beta)$ . Then the corollary follows immediately by Lemma 4.1 and a simple calculation.  $\square$

## 5 Acknowledgement

This work was supported by the National Natural Science Foundation of China [Grant number: 11971406].

## References

- [1] A. Ali, D. Dimitrov, On the extremal graphs with respect to bond incident degree indices, *Discrete Appl. Math.* 238 (2018) 32-40.
- [2] R. Cruz, I. Gutman, J. Rada, Sombor index of chemical graphs, *Appl. Math. Comput.* 399 (2021) 126018.
- [3] K.C. Das, A.S. Çevik, I.N. Cangul, Y. Shang, On Sombor index, *Symmetry* 13, 140 (2021), <https://doi.org/10.3390/sym13010140>.
- [4] E. Deutsch, S. Klavžar,  $M$ -polynomial revisited: Bethe cacti and an extension of Gutman's approach, *J. Appl. Math. Comput.* 60 (2019) 253-264.
- [5] S. Fajtlowicz, On conjectures of Graffiti-II, *Congr. Numer.* 60 (1987) 187-197.
- [6] B. Furtula, I. Gutman, A forgotten topological index, *J. Math. Chem.* 53 (2015) 1184-1190.
- [7] I. Gutman, Degree-based topological indices, *Croat. Chem. Acta* 86 (2013) 351-361.
- [8] I. Gutman, J. Tošović, Testing the quality of molecular structure descriptors. Vertex-degree-based topological indices, *J. Serb. Chem. Soc.* 78 (2013) 805-810.
- [9] I. Gutman, N. Trinajstić, Graph theory and molecular orbitals. Total  $\pi$ -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* 17 (1972) 535-538.
- [10] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices, *MATCH Commun. Math. Comput. Chem.* 86 (2021) 11-16.
- [11] J.C. Hernández, J.M. Rodríguez, O. Rosario, J.M. Sigarreta, Optimal inequalities and extremal problems on the general Sombor index, paper submitted arXiv:2108.05224.
- [12] V.R. Kulli, The  $(a, b)$ - $KA$  indices of polycyclic aromatic hydrocarbons and benzenoid systems, *International Journal of Mathematics Trends and Technology* 65 (2019) 115-120.

- [13] V.R. Kulli, I. Gutman, Computation of Sombor indices of certain networks, *SSRG Int. J. Appl. Chem.* 8 (2021) 1-5.
- [14] X. Li, J. Zheng, A unified approach to the extremal trees for different indices, *MATCH Commun. Math. Comput. Chem.* 54 (2005) 195-208.
- [15] Z. Lin, T. Zhou, V.R. Kullic, L. Miao, On the first Banhatti-Sombor index, preprint arXiv:2104.03615.
- [16] A.W. Marshall , I. Olkin, *Inequalities: Theory of Majorization and Its Applications*, Academic Press, New York, 1979.
- [17] S. Nikolić, G. Kovačević, A. Miličević, N. Trinajstić, The Zagreb indices 30 years after, *Croat. Chem. Acta* 76 (2003) 113-124.
- [18] M. Randić, On characterization of molecular branching, *J. Amer. Chem. Soc.* 97 (1975) 6609-6615.
- [19] I. Redžpović, Chemical applicability of Sombor indices, *J. Serb. Chem. Soc.* 86 (2021) 445-457.
- [20] T. Réti, T. Došlić, A. Ali, On the Sombor index of graphs, *Contrib. Math.* 3 (2021) 11-18.
- [21] D. Vukičević, J. Durdević, Bond additive modeling 10. Upper and lower bounds of bond incident degree indices of catacondensed uoranthenes, *Chem. Phys. Lett.* 515 (2011) 186-189.
- [22] H. Wang, Functions on adjacent vertex degrees of trees with given degree sequence, *Cent. Eur. J. Math.* 12 (2014) 1656-1663.
- [23] J. Ye, M. Liu, Y. Yao, K.C. Das, Extremal polygonal cacti for bond incident degree indices, *Discrete Appl. Math.* 257 (2019) 289-298.
- [24] B. Zhou, N. Trinajstić, On general sum-connectivity index, *J. Math. Chem.* 47 (2010) 210-218.