

**q -ANALOGUES OF π -RELATED FORMULAE FROM
JACKSON'S ${}_8\phi_7$ -SERIES VIA INVERSION APPROACH**

XIAOJING CHEN AND WENCHANG CHU

ABSTRACT. By making use of the multiplicate form of the extended Carlitz inverse series relations, we establish two general ‘dual’ theorems of Jackson’s summation formula for well–poised ${}_8\phi_7$ -series. Their duplicate forms under the partition pattern $n = \lfloor \frac{n}{2} \rfloor + \lfloor \frac{n+1}{2} \rfloor$ are explored and yield numerous q -series identities whose limiting cases as $q \rightarrow 1$ result in classical π -related Ramanujan–like series of convergence rate “ $\frac{1}{16}$ ” including one for $1/\pi^2$ discovered by Guillera (2003). The triplicate dual formulae under the partition pattern $n = \lfloor \frac{n}{3} \rfloor + \lfloor \frac{n+1}{3} \rfloor + \lfloor \frac{n+2}{3} \rfloor$ are examined via the “reverse bisection method”, which leads us to twenty new q -series identities together with their classical counterparts of convergence rate “ $\frac{1}{27}$ ” when $q \rightarrow 1$.

1. INTRODUCTION AND MOTIVATION

Let \mathbb{N} and \mathbb{N}_0 be the sets of natural numbers and nonnegative integers, respectively. For an indeterminate x , the shifted factorial is defined by

$$(x)_0 \equiv 1 \quad \text{and} \quad (x)_n = x(x+1)\cdots(x+n-1) \quad \text{for } n \in \mathbb{N}$$

with the following shortened multiparameter notation

$$\left[\begin{matrix} \alpha, \beta, \dots, \gamma \\ A, B, \dots, C \end{matrix} \right]_n = \frac{(\alpha)_n (\beta)_n \cdots (\gamma)_n}{(A)_n (B)_n \cdots (C)_n}.$$

Analogously, the shifted factorials with the base q are given by $(x; q)_0 \equiv 1$ and

$$(x; q)_n = (1-x)(1-qx)\cdots(1-q^{n-1}x) \quad \text{for } n \in \mathbb{N}.$$

The product and quotient of the q -shifted factorials will be abbreviated respectively to

$$\begin{aligned} (\alpha, \beta, \dots, \gamma; q)_n &= (\alpha; q)_n (\beta; q)_n \cdots (\gamma; q)_n, \\ \left[\begin{matrix} \alpha, \beta, \dots, \gamma \\ A, B, \dots, C \end{matrix} \middle| q \right]_n &= \frac{(\alpha; q)_n (\beta; q)_n \cdots (\gamma; q)_n}{(A; q)_n (B; q)_n \cdots (C; q)_n}. \end{aligned}$$

When $|q| < 1$, the infinite product $(x; q)_\infty$ is well defined, from which we introduce the q -gamma function [25, §1.10]

$$\Gamma_q(x) = (1-q)^{1-x} \frac{(q; q)_\infty}{(q^x; q)_\infty} \quad \text{and} \quad \lim_{q \rightarrow 1^-} \Gamma_q(x) = \Gamma(x).$$

Then for real numbers $\{a_k, c_k\}$, it is easy to check the following limiting relation

$$\lim_{q \rightarrow 1} \left[\begin{matrix} q^{a_1}, q^{a_2}, \dots, q^{a_n} \\ q^{c_1}, q^{c_2}, \dots, q^{c_n} \end{matrix} \middle| q \right]_\infty = \prod_{k=1}^n \frac{\Gamma(c_k)}{\Gamma(a_k)} \quad \text{provided that} \quad \sum_{k=1}^n a_k = \sum_{k=1}^n c_k.$$

2020 *Mathematics Subject Classification*. Primary 33D15, Secondary 33C20, 65B10.

Key words and phrases. Basic hypergeometric series; Bisection series; Jackson’s formula for well–poised ${}_8\phi_7$ -series; Duplicate inversions; Triplicate inversions; Ramanujan–like π -related series; Guillera’s infinite series for $1/\pi^2$.

Email addresses: chu.wenchang@unisalento.it and upcxjchen@outlook.com.

One century ago, Ramanujan [38] discovered (without proofs) 17 π -related infinite series, that were demonstrated rigorously by Borwein brothers [4] during the 1980s. Three typical formulae are reproduced as follows:

$$\begin{aligned}\frac{4}{\pi} &= \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ 1, 1, 1 \end{matrix} \right]_n \frac{1+6n}{4^n}. \\ \frac{8}{\pi} &= \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \\ 1, 1, 1 \end{matrix} \right]_n \frac{3+20n}{(-4)^n}. \\ \frac{16}{\pi} &= \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ 1, 1, 1 \end{matrix} \right]_n \frac{5+42n}{64^n}.\end{aligned}$$

By making use of computer algebra, Guillera [30–32] found several remarkable identities including the following one

$$\frac{32}{\pi^2} = \sum_{n=0}^{\infty} \frac{1}{16^n} \left[\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \\ 1, 1, 1, 1, 1 \end{matrix} \right]_n \{3 + 34n + 120n^2\}. \quad (1)$$

Recently, there has been a growing interest in searching for q -analogues of Ramanujan-like series (cf. [9,10,33–36]). By means of the modified Abel lemma on summation by parts, the second author [19] (see also [10]) established the following q -analogue of (1):

$$\begin{aligned}\left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{5}{2}} \\ q, q, q, q \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2} \frac{(q^{\frac{1}{2}}; q)_n^3 (q^{\frac{3}{2}}; q)_{2n} (1 - q^{3n + \frac{3}{2}})}{(q; q)_n^5 (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n+1}^3 (1 - q^{3/2})} \\ &\times \left\{ q^n + \frac{(1 - q^n)(1 - q^{3n + \frac{1}{2}})(1 + q^{n + \frac{1}{2}})^3}{(1 - q^{2n + \frac{1}{2}})(1 - q^{3n + \frac{3}{2}})} \right\}.\end{aligned} \quad (2)$$

In 1973, Gould and Hsu [29] found an useful pair of inverse series relations. Carlitz [5] obtained their q -analogues in the same year. These inversions together with their multiplicate forms (cf. [7, 8, 16]) have been shown powerful in proving identities of hypergeometric series (cf. [11, 12, 14, 26]) and q -series (cf. [15, 16, 21, 27]). By applying the multiplicate form of the extended Carlitz inverse series relations (cf. [11, 15]) to Jackson's well-poised ${}_8\phi_7$ -series, we shall provide a new proof (see Example 1) for identity (2). Numerous further q -series identities will be proved likewise. These identities are significant not only for their existence, but also for the novelty of their limiting series as $q \rightarrow 1$, namely, the classical Ramanujan-like series. Nine remarkable ones are highlighted in advance as

follows:

$$\text{Example 40: } \frac{\pi\Gamma^2(\frac{1}{3})}{6\Gamma^2(\frac{5}{6})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 1, \frac{1}{2}, \frac{2}{3} \\ \frac{4}{3}, \frac{4}{3}, \frac{4}{3} \end{matrix} \right]_n \{3 + 7n\}.$$

$$\text{Example 41: } \frac{48\pi\Gamma^2(\frac{2}{3})}{\Gamma^2(\frac{1}{6})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 1, \frac{1}{2}, \frac{1}{3} \\ \frac{5}{3}, \frac{5}{3}, \frac{5}{3} \end{matrix} \right]_n \{9 + 28n + 21n^2\}.$$

$$\text{Example 42: } \frac{9\sqrt{3}}{2^{\frac{4}{3}}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, \frac{2}{3}, \frac{1}{6} \\ 1, 1, 1 \end{matrix} \right]_n \{2 + 21n\}.$$

$$\text{Example 43: } \frac{27\sqrt{3}}{2^{\frac{5}{3}}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, \frac{2}{3}, \frac{5}{6} \\ 1, 1, 1 \end{matrix} \right]_n \{5 + 42n\}.$$

$$\text{Example 44: } \frac{81\sqrt{3}}{28 \cdot 2^{2/3}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{2}{3}, \frac{7}{3}, -\frac{1}{6} \\ 1, 1, 2 \end{matrix} \right]_n.$$

$$\text{Example 51: } \frac{3\sqrt{3}}{\pi\sqrt[3]{4}} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, -\frac{1}{3}, -\frac{1}{6} \\ 1, 1, 1 \end{matrix} \right]_n \{1 - 63n^2\}.$$

$$\begin{aligned} \text{Example 57: } \frac{32}{\pi^2} &= \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^6 (\frac{1}{2})_{2n}^3}{(2n)!^3 (3n)!^2} \times \frac{(1+2n)(1+4n)^2(3+8n)}{(1+3n)^2} \\ &\times \left\{ 1 + \frac{32n(1+3n)^2(1+8n)}{(1+2n)(1+4n)^2(3+8n)} + \frac{(1+4n)(3+4n)(5+8n)}{16(2+3n)^2(3+8n)} \right\}. \end{aligned}$$

$$\begin{aligned} \text{Example 59: } \frac{32}{\pi^2} &= \sum_{n=0}^{\infty} (3+8n) \frac{(\frac{1}{2})_n^3 (\frac{3}{2})_{2n}^3 (\frac{3}{2})_{3n}}{n!(2n)!(3n+1)!^3} \\ &\times \left\{ 1 + \frac{16n(1+3n)^3(1+10n)}{(1+4n)^3(1+6n)(3+8n)} + \frac{3(1+2n)^3(7+10n)}{16(2+3n)^3(3+8n)} \right\}. \end{aligned}$$

$$\begin{aligned} \text{Example 60: } \frac{9\pi^2}{16} &= \sum_{n=0}^{\infty} \frac{(2n)!(4n+1)!}{(6n+1)!(1+2n)(1+3n)} \\ &\times \left\{ 1 + \frac{9(1+2n)(1+3n)(1+5n)}{2(1+4n)^2(1+6n)} + \frac{3(1+n)(1+2n)^2(4+5n)}{2(2+3n)(3+4n)(5+6n)^2} \right\}. \end{aligned}$$

The rest of the paper will be organized as follows. By applying the multiplicate form of the extended Carlitz inverse series relations to Jackson’s q -analogue of Dougall’s well–poised ${}_7F_6$ -sum, we shall derive, in the next section, two general dual formulae (Theorems 3 and 4). Then under the partition pattern $n = \lfloor \frac{n}{2} \rfloor + \lfloor \frac{n+1}{2} \rfloor$, we shall deduce, in Section 3, from the duplicate inversions a number of q -series identities whose limiting cases as $q \rightarrow 1$ result in classical π -related Ramanujan–like series of convergence rate “ $\frac{1}{16}$ ”. The triplicate inversions under the partition pattern $n = \lfloor \frac{n}{3} \rfloor + \lfloor \frac{n+1}{3} \rfloor + \lfloor \frac{n+2}{3} \rfloor$ will be utilized in Section 4 to establish further formulae for q -series, where the “reverse bisection method” is employed to reduce the resulting series dramatically to simpler but more elegant ones.

2. INVERSION APPROACH TO JACKSON'S WELL-POISED ${}_8\phi_7$ -SERIES

In 1973, Gould and Hsu [29] discovered a useful pair of inverse series relations, which can equivalently be reproduced below. Let $\{a_i, b_i\}$ be two complex sequences such that the φ -polynomials defined by

$$\varphi(x; 0) \equiv 1 \quad \text{and} \quad \varphi(x; n) = \prod_{k=0}^{n-1} (a_k + xb_k) \quad \text{for } n \in \mathbb{N} \quad (3)$$

differ from zero for $x, n \in \mathbb{N}_0$. Then there hold the inverse series relations

$$f(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \varphi(k; n) g(k), \quad (4)$$

$$g(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \frac{a_k + kb_k}{\varphi(n; k+1)} f(k). \quad (5)$$

Define the Gaussian binomial coefficient by

$$\begin{bmatrix} m \\ n \end{bmatrix} = \frac{(q; q)_m}{(q; q)_n (q; q)_{m-n}} = \frac{(q^{m-n+1}; q)_n}{(q; q)_n}, \quad \text{where } m, n \in \mathbb{N}.$$

Carlitz [5] found the following q -analogues

$$f(n) = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} \varphi(q^{-k}; n) g(k), \quad (6)$$

$$g(n) = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\binom{n-k}{2}} \frac{a_k + q^{-k} b_k}{\varphi(q^{-n}; k+1)} f(k); \quad (7)$$

which were utilized to derive q -series identities [13, 15, 27]. There is an extended form (cf. [11, 15]):

$$f(n) = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} \varphi(q^k \sigma; n) \varphi(q^{-k}; n) \frac{1 - q^{2k} \sigma}{(q^n \sigma; q)_{k+1}} g(k), \quad (8)$$

$$g(n) = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\binom{n-k}{2}} \frac{(a_k + q^k \sigma b_k)(a_k + q^{-k} b_k)}{\varphi(q^n \sigma; k+1) \varphi(q^{-n}; k+1)} (q^k \sigma; q)_n f(k). \quad (9)$$

In order to facilitate the subsequent application, we can reformulate the above inversions as

$$F(n) = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} \frac{\phi(q^k; n)}{\psi(q^n; k+1)} \frac{\psi(q^k; k+1)}{\psi(q^k; k)} G(k), \quad (10)$$

$$G(n) = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\binom{n-k}{2}} \frac{\psi(q^k; n)}{\phi(q^n; k+1)} \frac{\phi(q^k; k+1)}{\phi(q^k; k)} F(k); \quad (11)$$

where the polynomial sequences $\{\phi(x; n), \psi(x; n)\}$ are defined respectively by

$$\phi(x; n) = \varphi(x\sigma; n) \varphi(x^{-1}; n) \quad \text{and} \quad \psi(x; n) = (x\sigma; q)_n. \quad (12)$$

For the fundamental identity of Dougall [24] about the very well-poised terminating ${}_7F_6$ -series, Jackson [37] (cf. Bailey [2, §8.3]) discovered, one century ago, the following q -analogue:

$$\begin{aligned} \Omega_n(a; b, c, d) &:= \sum_{k=0}^n \frac{1 - q^{2k} a}{1 - a} \left[\begin{matrix} a, b, c, d, e, q^{-n} \\ q, qa/b, qa/c, qa/d, qa/e, q^{n+1} a \end{matrix} \middle| q \right]_k q^k \\ &= \left[\begin{matrix} qa, qa/bc, qa/bd, qa/cd \\ qa/b, qa/c, qa/d, qa/bcd \end{matrix} \middle| q \right]_n, \quad \text{where } q^{n+1} a^2 = bcde. \end{aligned} \quad (13)$$

For a real number x , denote by $[x]$ the greatest integer $\leq x$. Suppose that for all $n \in \mathbb{N}_0$, there exist eight integers $\{\varepsilon_b, \varepsilon_c, \varepsilon_d, \varepsilon_e, \lambda_b, \lambda_c, \lambda_d, \lambda_e\} \subset \mathbb{N}_0$ with $\boxed{\Lambda := \lambda_b + \lambda_c + \lambda_d + \lambda_e}$ such that

$$n = \left\lfloor \frac{\varepsilon_b + n\lambda_b}{\Lambda} \right\rfloor + \left\lfloor \frac{\varepsilon_c + n\lambda_c}{\Lambda} \right\rfloor + \left\lfloor \frac{\varepsilon_d + n\lambda_d}{\Lambda} \right\rfloor + \left\lfloor \frac{\varepsilon_e + n\lambda_e}{\Lambda} \right\rfloor. \quad (14)$$

For the sake of brevity, denote this partition by

$$n = \langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_d + \langle n \rangle_e,$$

where

$$\begin{aligned} \langle n \rangle_b &:= \left\lfloor \frac{\varepsilon_b + n\lambda_b}{\Lambda} \right\rfloor, \\ \langle n \rangle_c &:= \left\lfloor \frac{\varepsilon_c + n\lambda_c}{\Lambda} \right\rfloor, \\ \langle n \rangle_d &:= \left\lfloor \frac{\varepsilon_d + n\lambda_d}{\Lambda} \right\rfloor, \\ \langle n \rangle_e &:= \left\lfloor \frac{\varepsilon_e + n\lambda_e}{\Lambda} \right\rfloor. \end{aligned}$$

Then Jackson's formula (13) can be reformulated as

$$\begin{aligned} \Omega_n(a; bq^{\langle n \rangle_b}, cq^{\langle n \rangle_c}, dq^{\langle n \rangle_d}) &= \frac{(qa; q)_n (bc/a; q)_{\langle n \rangle_b + \langle n \rangle_c} (qa/bc; q)_{\langle n \rangle_d + \langle n \rangle_e}}{(b/a; q)_{\langle n \rangle_b} (c/a; q)_{\langle n \rangle_c} (d/a; q)_{\langle n \rangle_d} (qa/bcd; q)_{\langle n \rangle_e}} \\ &\times \frac{(bd/a; q)_{\langle n \rangle_b + \langle n \rangle_d} (qa/bd; q)_{\langle n \rangle_c + \langle n \rangle_e}}{(qa/b; q)_{\langle n \rangle_c + \langle n \rangle_d + \langle n \rangle_e} (qa/c; q)_{\langle n \rangle_b + \langle n \rangle_d + \langle n \rangle_e}} \\ &\times \frac{(cd/a; q)_{\langle n \rangle_c + \langle n \rangle_d} (qa/cd; q)_{\langle n \rangle_b + \langle n \rangle_e}}{(qa/d; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_e} (bcd/a; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_d}}. \end{aligned}$$

In view of the relation (which is also true when “ b ” is replaced by one of “ c, d, e ”)

$$\frac{(bq^{\langle n \rangle_b}; q)_k}{(q^{1-\langle n \rangle_b} a/b; q)_k} = q^{k\langle n \rangle_b} \frac{(b; q)_k (q^k b; q)_{\langle n \rangle_b} (q^{-k} b/a; q)_{\langle n \rangle_b}}{(qa/b; q)_k (b; q)_{\langle n \rangle_b} (b/a; q)_{\langle n \rangle_b}},$$

it is not difficult to check that Jackson's formula is equivalent to the following q -binomial sum

$$\begin{aligned} &\sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} \frac{1 - q^{2k} a}{(q^n a; q)_{k+1}} \begin{bmatrix} a, b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{bmatrix}_k q^{\binom{k+1}{2}} \\ &\quad \times (q^k b; q)_{\langle n \rangle_b} (q^k c; q)_{\langle n \rangle_c} (q^k d; q)_{\langle n \rangle_d} (q^{1+k} a^2/bcd; q)_{\langle n \rangle_e} \\ &\quad \times (q^{-k} b/a; q)_{\langle n \rangle_b} (q^{-k} c/a; q)_{\langle n \rangle_c} (q^{-k} d/a; q)_{\langle n \rangle_d} (q^{1-k} a/bcd; q)_{\langle n \rangle_e} \\ &= (a; q)_n (b; q)_{\langle n \rangle_b} (c; q)_{\langle n \rangle_c} (d; q)_{\langle n \rangle_d} (qa^2/bcd; q)_{\langle n \rangle_e} \\ &\quad \times \frac{(bc/a; q)_{\langle n \rangle_b + \langle n \rangle_c} (bd/a; q)_{\langle n \rangle_b + \langle n \rangle_d} (cd/a; q)_{\langle n \rangle_c + \langle n \rangle_d}}{(qa/b; q)_{\langle n \rangle_c + \langle n \rangle_d + \langle n \rangle_e} (qa/c; q)_{\langle n \rangle_b + \langle n \rangle_d + \langle n \rangle_e}} \\ &\quad \times \frac{(qa/bc; q)_{\langle n \rangle_d + \langle n \rangle_e} (qa/bd; q)_{\langle n \rangle_c + \langle n \rangle_e} (qa/cd; q)_{\langle n \rangle_b + \langle n \rangle_e}}{(qa/d; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_e} (bcd/a; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_d}}. \end{aligned}$$

Observing that this equation matches exactly to (10) specified by “ $\sigma = a$ ” and

$$\begin{aligned} \phi(x; n) &= (bx; q)_{\langle n \rangle_b} (cx; q)_{\langle n \rangle_c} (dx; q)_{\langle n \rangle_d} (qa^2 x/bcd; q)_{\langle n \rangle_e} \\ &\quad \times (b/ax; q)_{\langle n \rangle_b} (c/ax; q)_{\langle n \rangle_c} (d/ax; q)_{\langle n \rangle_d} (qa/bcdx; q)_{\langle n \rangle_e}, \\ \psi(x; n) &= (ax; q)_n; \end{aligned}$$

as well as

$$\begin{aligned}
F(n) &= (a; q)_n (b; q)_{\langle n \rangle_b} (c; q)_{\langle n \rangle_c} (d; q)_{\langle n \rangle_d} (qa^2/bcd; q)_{\langle n \rangle_e} \\
&\times \frac{(bc/a; q)_{\langle n \rangle_b + \langle n \rangle_c} (bd/a; q)_{\langle n \rangle_b + \langle n \rangle_d} (cd/a; q)_{\langle n \rangle_c + \langle n \rangle_d}}{(qa/b; q)_{\langle n \rangle_c + \langle n \rangle_d + \langle n \rangle_e} (qa/c; q)_{\langle n \rangle_b + \langle n \rangle_d + \langle n \rangle_e}} \\
&\times \frac{(qa/bc; q)_{\langle n \rangle_d + \langle n \rangle_e} (qa/bd; q)_{\langle n \rangle_c + \langle n \rangle_e} (qa/cd; q)_{\langle n \rangle_b + \langle n \rangle_e}}{(qa/d; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_e} (bcd/a; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_d}}, \\
G(n) &= \left[\begin{matrix} a, b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{matrix} \middle| q \right]_n q^{\binom{n+1}{2}};
\end{aligned}$$

we may state the dual relation corresponding to (11) as

$$G(n) = \sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\binom{n-k}{2}} \frac{\psi(q^k; n)}{\phi(q^n; k+1)} \frac{\phi(q^k; k+1)}{\phi(q^k; k)} F(k).$$

This can explicitly be highlighted, as the following lemma.

Lemma 1. *The following identity holds*

$$\left[\begin{matrix} b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{matrix} \middle| q \right]_n = \sum_{k=0}^n q^{k-n} \frac{(q^n a; q)_k (q^{-n}; q)_k}{\phi(q^n; k+1)} \frac{\phi(q^k; k+1)}{\phi(q^k; k)} H(k),$$

where $H(n)$ is the quotient of shifted factorials given by

$$\begin{aligned}
H(n) &= \frac{(b; q)_{\langle n \rangle_b} (c; q)_{\langle n \rangle_c} (d; q)_{\langle n \rangle_d} (qa^2/bcd; q)_{\langle n \rangle_e}}{(q; q)_n (bcd/a; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_d}} \\
&\times \frac{(bc/a; q)_{\langle n \rangle_b + \langle n \rangle_c} (qa/bc; q)_{\langle n \rangle_d + \langle n \rangle_e}}{(qa/b; q)_{\langle n \rangle_c + \langle n \rangle_d + \langle n \rangle_e}} \\
&\times \frac{(bd/a; q)_{\langle n \rangle_b + \langle n \rangle_d} (qa/bd; q)_{\langle n \rangle_c + \langle n \rangle_e}}{(qa/c; q)_{\langle n \rangle_b + \langle n \rangle_d + \langle n \rangle_e}} \\
&\times \frac{(cd/a; q)_{\langle n \rangle_c + \langle n \rangle_d} (qa/cd; q)_{\langle n \rangle_b + \langle n \rangle_e}}{(qa/d; q)_{\langle n \rangle_b + \langle n \rangle_c + \langle n \rangle_e}}.
\end{aligned}$$

Recall that for all $k \in \mathbb{N}_0$, we have

$$\langle k+1 \rangle_b + \langle k+1 \rangle_c + \langle k+1 \rangle_d + \langle k+1 \rangle_e = k+1$$

and

$$\begin{aligned}
\phi(q^n; k+1) &= (q^n b; q)_{\langle k+1 \rangle_b} (q^n c; q)_{\langle k+1 \rangle_c} (q^n d; q)_{\langle k+1 \rangle_d} (q^{1+n} a^2/bcd; q)_{\langle k+1 \rangle_e} \\
&\times (q^{-n} b/a; q)_{\langle k+1 \rangle_b} (q^{-n} c/a; q)_{\langle k+1 \rangle_c} (q^{-n} d/a; q)_{\langle k+1 \rangle_d} (q^{1-n} a/bcd; q)_{\langle k+1 \rangle_e}.
\end{aligned}$$

When $n \rightarrow \infty$, it is not hard to check the limiting relation

$$\begin{aligned}
q^{k-n} \frac{(q^n a; q)_k (q^{-n}; q)_k}{\phi(q^n; k+1)} &\Rightarrow -q^{\binom{k+1}{2} - \binom{(k+1)b}{2} - \binom{(k+1)c}{2} - \binom{(k+1)d}{2} - \binom{(k+1)e}{2}} \\
&\times \left(\frac{a}{b}\right)^{\langle k+1 \rangle_b} \left(\frac{a}{c}\right)^{\langle k+1 \rangle_c} \left(\frac{a}{d}\right)^{\langle k+1 \rangle_d} \left(\frac{bcd}{qa}\right)^{\langle k+1 \rangle_e}.
\end{aligned}$$

Now let $n \rightarrow \infty$ in Lemma 1 and then determine the limit of the right member through the Weierstrass M -test on uniformly convergent series (cf. Stromberg [39, §3.106]). After some routine simplification, the resulting limiting relation may be stated as in the following proposition.

Proposition 2. *The following identity holds*

$$\left[\begin{array}{c} b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{array} \middle| q \right]_{\infty} = \sum_{k=0}^{\infty} \Theta(k+1) \left\{ \frac{\phi(q^k; k+1)}{-\phi(q^k; k)} \right\} H(k),$$

where the monomial $\Theta(m)$ is defined by

$$\begin{aligned} \Theta(m) &= q^{\binom{m}{2} - \binom{m}{2}_b - \binom{m}{2}_c - \binom{m}{2}_d - \binom{m}{2}_e} \\ &\times \left(\frac{a}{b}\right)^{\langle m \rangle_b} \left(\frac{a}{c}\right)^{\langle m \rangle_c} \left(\frac{a}{d}\right)^{\langle m \rangle_d} \left(\frac{bcd}{qa}\right)^{\langle m \rangle_e}. \end{aligned}$$

To facilitate understanding, we explicitly display the expression from this proposition:

$$\begin{aligned} \left[\begin{array}{c} b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{array} \middle| q \right]_{\infty} &= \sum_{k=0}^{\infty} \left\{ \frac{\phi(q^k; k+1)}{-\phi(q^k; k)} \right\} \\ &\times q^{\binom{k+1}{2} - \binom{k+1}{2}_b - \binom{k+1}{2}_c - \binom{k+1}{2}_d - \binom{k+1}{2}_e} \\ &\times \left(\frac{a}{b}\right)^{\langle k+1 \rangle_b} \left(\frac{a}{c}\right)^{\langle k+1 \rangle_c} \left(\frac{a}{d}\right)^{\langle k+1 \rangle_d} \left(\frac{bcd}{qa}\right)^{\langle k+1 \rangle_e} \\ &\times \frac{(b; q)_{\langle k \rangle_b} (c; q)_{\langle k \rangle_c} (d; q)_{\langle k \rangle_d} (qa^2/bcd; q)_{\langle k \rangle_e}}{(q; q)_k (bcd/a; q)_{\langle k \rangle_b + \langle k \rangle_c + \langle k \rangle_d}} \\ &\times \frac{(bc/a; q)_{\langle k \rangle_b + \langle k \rangle_c} (qa/bc; q)_{\langle k \rangle_d + \langle k \rangle_e}}{(qa/b; q)_{\langle k \rangle_c + \langle k \rangle_d + \langle k \rangle_e}} \\ &\times \frac{(bd/a; q)_{\langle k \rangle_b + \langle k \rangle_d} (qa/bd; q)_{\langle k \rangle_c + \langle k \rangle_e}}{(qa/c; q)_{\langle k \rangle_b + \langle k \rangle_d + \langle k \rangle_e}} \\ &\times \frac{(cd/a; q)_{\langle k \rangle_c + \langle k \rangle_d} (qa/cd; q)_{\langle k \rangle_b + \langle k \rangle_e}}{(qa/d; q)_{\langle k \rangle_b + \langle k \rangle_c + \langle k \rangle_e}}. \end{aligned}$$

Classifying the summation indices $\boxed{k \rightarrow k + n\Lambda}$ with respect to their residues modulo Λ , we can split the sum in Proposition 2 into Λ partial sums:

$$\begin{aligned} &\left[\begin{array}{c} b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{array} \middle| q \right]_{\infty} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\Lambda-1} \Theta(1+k+n\Lambda) \left\{ \frac{\phi(q^{k+n\Lambda}; 1+k+n\Lambda)}{-\phi(q^{k+n\Lambda}; k+n\Lambda)} \right\} H(k+n\Lambda) \\ &= \sum_{n=0}^{\infty} H(n\Lambda) \Theta(n\Lambda) \sum_{k=0}^{\Lambda-1} \frac{\Theta(1+k+n\Lambda)}{\Theta(n\Lambda)} \left\{ \frac{\phi(q^{k+n\Lambda}; 1+k+n\Lambda)}{-\phi(q^{k+n\Lambda}; k+n\Lambda)} \right\} \frac{H(k+n\Lambda)}{H(n\Lambda)}. \end{aligned}$$

In general, by singling out the initial δ -terms, where $\boxed{0 \leq \delta < \Lambda}$, we have two further expressions, that will efficiently be utilized to derive concrete q -series identities in the next two sections.

Theorem 3 (Nonterminating series identity).

$$\begin{aligned} \left[\begin{array}{c} b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{array} \middle| q \right]_{\infty} &= \sum_{k=0}^{\delta-1} \Theta(k+1) \left\{ \frac{\phi(q^k; k+1)}{-\phi(q^k; k)} \right\} H(k) \\ &+ \sum_{n=0}^{\infty} H(n\Lambda) \Theta(n\Lambda) \sum_{i=\delta}^{\delta-1+\Lambda} \frac{\Theta(1+i+n\Lambda)}{\Theta(n\Lambda)} \left\{ \frac{\phi(q^{i+n\Lambda}; 1+i+n\Lambda)}{-\phi(q^{i+n\Lambda}; i+n\Lambda)} \right\} \frac{H(i+n\Lambda)}{H(n\Lambda)}. \end{aligned}$$

Theorem 4 (Nonterminating series identity).

$$\begin{aligned} & \left[\begin{array}{c} b, c, d, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{array} \middle| q \right]_{\infty} = \sum_{k=0}^{\delta-1} \Theta(k+1) \left\{ \frac{\phi(q^k; k+1)}{-\phi(q^k; k)} \right\} H(k) \\ & + \sum_{n=1}^{\infty} H(n\Lambda) \Theta(n\Lambda) \sum_{j=\delta-\Lambda}^{\delta-1} \frac{\Theta(1+j+n\Lambda)}{\Theta(n\Lambda)} \left\{ \frac{\phi(q^{j+n\Lambda}; 1+j+n\Lambda)}{-\phi(q^{j+n\Lambda}; j+n\Lambda)} \right\} \frac{H(j+n\Lambda)}{H(n\Lambda)}. \end{aligned}$$

3. SERIES FROM DUPLICATE INVERSIONS

For all $n \in \mathbb{N}_0$, it is well known that $\boxed{n = \lfloor \frac{n}{2} \rfloor + \lfloor \frac{1+n}{2} \rfloor}$. Then it is not difficult to check that Jackson's formula (13) is equivalent to the following one

$$\Omega_n(a; q^{\lfloor \frac{n}{2} \rfloor} b, c, q^{\lfloor \frac{1+n}{2} \rfloor} d) = \left[\begin{array}{c} qa, bd/a \\ qa/c, bcd/a \end{array} \middle| q \right]_n \left[\begin{array}{c} qa/cd, bc/a \\ qa/d, b/a \end{array} \middle| q \right]_{\lfloor \frac{n}{2} \rfloor} \left[\begin{array}{c} qa/bc, cd/a \\ qa/b, d/a \end{array} \middle| q \right]_{\lfloor \frac{1+n}{2} \rfloor}$$

with its parameters subject to $\boxed{qa^2 = bcde}$. In this case, the partition pattern defined by (14) can be explicitly determined as

$$\boxed{\Lambda = 2: \quad \begin{array}{cccc} \varepsilon_b = 0, & \varepsilon_c = 0, & \varepsilon_d = 1, & \varepsilon_e = 0; \\ \lambda_b = 1, & \lambda_c = 0, & \lambda_d = 1, & \lambda_e = 0. \end{array}} \quad (15)$$

§3.1. Letting $\boxed{\delta = 0}$ in Theorem 3, we derive, under the above partition pattern, the following quite general formula involving four free parameters $\{a, b, c, d\}$.

Theorem 5 (Nonterminating series identity).

$$\begin{aligned} & \left[\begin{array}{c} b, c, qd, qa^2/bcd \\ qa/b, qa/c, a/d, bcd/a \end{array} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} W_n(a, b, c, d) \left(\frac{q^n a^2}{bd} \right)^n \\ & \quad \times \frac{(bd/a; q)_{2n} (b, qd, q^2 a/bc, bc/a, qa/cd, cd/a; q)_n}{(q, qa/c, bcd/a; q)_{2n} (qa/b, qa/d; q)_n}, \end{aligned}$$

where the weight function is given by

$$\begin{aligned} W_n(a, b, c, d) &= \frac{(1 - qa/bc)(1 - q^n cd/a)(1 - q^{2n} bd/a)(1 - q^{1+3n} b)}{(1 - a/d)(1 - q^{1+2n})(1 - q^{1+2n} a/c)(1 - q^{2n} bcd/a)} \\ & \times \left\{ q^n \frac{a}{d} + \frac{(1 - q^n a/d)(1 - q^{1+2n})(1 - q^{1+2n} a/c)(1 - q^{2n} bcd/a)(1 - q^{3n} d)}{(1 - q^{1+n} a/bc)(1 - q^n d)(1 - q^n cd/a)(1 - q^{2n} bd/a)(1 - q^{1+3n} b)} \right\}. \end{aligned}$$

Letting $\boxed{a = b = c = d = q^{\frac{1}{2}}}$ in this theorem and then simplifying slightly the resulting expression, we get immediately the following infinite series identity, that was derived by the second author [19] by making use of the modified Abel lemma on summation by parts.

Example 1 ($\boxed{a = b = c = d = q^{\frac{1}{2}}}$ in Theorem 5).

$$\begin{aligned} & \left[\begin{array}{c} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{5}{2}} \\ q, q, q, q \end{array} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2} \frac{(q^{\frac{1}{2}}; q)_n^3 (q^{\frac{3}{2}}; q)_{2n} (1 - q^{3n + \frac{3}{2}})}{(q; q)_n^5 (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n+1}^3 (1 - q^{3/2})} \\ & \quad \times \left\{ q^n + \frac{(1 - q^n)(1 - q^{3n + \frac{1}{2}})(1 + q^{n + \frac{1}{2}})^3}{(1 - q^{2n + \frac{1}{2}})(1 - q^{3n + \frac{3}{2}})} \right\}. \end{aligned}$$

When $q \rightarrow 1$, we recover the following beautiful series discovered by Guillera [30, 31]:

$$\frac{32}{\pi^2} = \sum_{n=0}^{\infty} \frac{1}{16^n} \left[\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \\ 1, 1, 1, 1, 1 \end{matrix} \right]_n \{3 + 34n + 120n^2\}.$$

Further q -analogues of Ramanujan-like π -related series can be derived similarly, where the parameter settings for $\{a, b, c, d\}$ are highlighted in the headers for all the examples.

Example 2 ($a = q^{\frac{3}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{3}{2}}, d = q^{\frac{1}{2}}$ in Theorem 5).

$$\left[\begin{matrix} q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}} \\ q, q, q^2, q^2 \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2+2n} \frac{(q^{\frac{1}{2}}; q)_n^3 (q^{-\frac{1}{2}}; q)_{2n} (1 + q^{1/2})}{(q^2; q^2)_n^3 (q; q)_n (q^2; q)_n (-q^{\frac{1}{2}}; q)_n^3} \\ \times \frac{1 - q^{3n+\frac{1}{2}}}{1 - q^{1/2}} \left\{ 1 + \frac{q^{1+n} (1 - q^{2n-\frac{1}{2}}) (1 - q^{3n+\frac{3}{2}})}{(1 - q^{1+n}) (1 - q^{3n+\frac{1}{2}}) (1 + q^{n+\frac{1}{2}})^3} \right\}.$$

When $q \rightarrow 1$, we recover the following known series (cf. [22]):

$$\frac{128}{\pi^2} = \sum_{n=0}^{\infty} \frac{1}{16^n} \left[\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, -\frac{1}{4} \\ 1, 1, 1, 2, 2 \end{matrix} \right]_n \{13 + 118n + 120n^2\}.$$

Example 3 ($a = q^{\frac{3}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{2}}, d = q^{\frac{3}{2}}$ in Theorem 5).

$$\left[\begin{matrix} q^{\frac{3}{2}}, q^{\frac{5}{2}}, q^{\frac{5}{2}}, q^{\frac{5}{2}} \\ q^2, q^2, q^2, q^3 \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2+n} \frac{(q^{\frac{1}{2}}; q)_n^2 (q^{-\frac{1}{2}}; q)_n (q^{\frac{3}{2}}; q)_n (q^{\frac{5}{2}}; q)_n^2 (q^{\frac{3}{2}}; q)_{2n}}{(q^2; q^2)_{2n}^2 (q^3; q)_{2n} (q; q)_n (q^2; q)_n} \\ \times \frac{1 - q^{3n+\frac{3}{2}}}{1 - q^{3/2}} \left\{ q^n + \frac{(1 - q^n) (1 - q^{2n+1})^2 (1 - q^{2n+2})}{(1 - q^{n+\frac{1}{2}}) (1 - q^{n+\frac{3}{2}})^2 (1 - q^{2n+\frac{1}{2}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{256}{3\pi^2} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, -\frac{1}{2}, \frac{3}{2}, \frac{1}{4}, \frac{3}{4} \\ 1, 1, 1, 2, 2 \end{matrix} \right]_n \frac{80n^3 + 148n^2 + 80n + 9}{16^n}.$$

Example 4 ($a = q^{\frac{3}{2}}, b = q^{\frac{3}{2}}, c = q^{\frac{1}{2}}, d = q^{\frac{3}{2}}$ in Theorem 5).

$$\left[\begin{matrix} q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{5}{2}}, q^{\frac{7}{2}} \\ q, q^2, q^3, q^3 \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2} \frac{(q^{\frac{1}{2}}; q)_n^2 (q^{\frac{5}{2}}; q)_n (q^{\frac{5}{2}}; q)_{2n} (1 - q^{3n+\frac{5}{2}})}{(q; q)_n^2 (q^2; q^2)_n (q^4; q^2)_n^2 (-q^{\frac{3}{2}}; q)_n^3 (1 - q^{5/2})} \\ \times \left\{ q^n + \frac{(1 - q^n) (1 - q^{2n+1}) (1 - q^{2n+2})^2 (1 - q^{3n+\frac{3}{2}})}{(1 - q^{n+\frac{1}{2}})^2 (1 - q^{n+\frac{3}{2}}) (1 - q^{2n+\frac{3}{2}}) (1 - q^{3n+\frac{5}{2}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{512}{\pi^2} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{4}, \frac{5}{4} \\ 1, 1, 1, 2, 2 \end{matrix} \right]_n \frac{240n^3 + 532n^2 + 336n + 45}{16^n}.$$

Example 5 ($a = q^{-\frac{1}{2}}, b = q^{-\frac{5}{6}}, c = q^{\frac{1}{6}}, d = q^{\frac{1}{6}}$ in Theorem 5).

$$\left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{7}{6}}, q^{\frac{7}{6}}, q^{\frac{13}{6}} \\ q^2, q^{\frac{1}{3}}, q^{\frac{4}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2+\frac{2}{3}n} \frac{(q^{\frac{1}{6}}, q^{-\frac{1}{6}}, q^{-\frac{5}{6}}, q^{\frac{7}{6}}, q^{\frac{11}{6}}, q^{\frac{13}{6}}; q)_n (q^{\frac{5}{6}}; q)_{2n}}{(q^{\frac{1}{3}}; q)_n (q^{\frac{4}{3}}; q)_n (q; q)_{2n} (q^2; q)_{2n} (q^{\frac{4}{3}}; q)_{2n}} \\ \times \frac{1 - q^{3n+\frac{1}{6}}}{1 - q^{\frac{1}{6}}} \left\{ 1 - \frac{(1 - q^{\frac{2}{3}-n}) (1 - q^{2n+\frac{1}{3}}) (1 - q^{2n}) (1 - q^{2n+1})}{(1 - q^{n+\frac{1}{6}}) (1 - q^{n+\frac{5}{6}}) (1 - q^{n+\frac{7}{6}}) (1 - q^{2n-\frac{1}{6}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{180\Gamma(\frac{2}{3})^3}{\pi^2} = \sum_{n=0}^{\infty} \left[\frac{1}{6}, -\frac{1}{6}, \frac{5}{6}, -\frac{5}{6}, \frac{7}{6}, -\frac{1}{12}, \frac{5}{12}, \frac{19}{18} \right]_n \frac{35 + 228n - 540n^2 - 2160n^3}{16^n}.$$

Example 6 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{6}}, c = q^{-\frac{5}{6}}, d = q^{\frac{7}{6}}$ in Theorem 5).

$$\begin{aligned} \left[q^{\frac{3}{2}}, q^{\frac{1}{6}}, q^{\frac{13}{6}}, q^{\frac{19}{6}} \mid q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{2}{3}n} \frac{(q^{\frac{1}{6}}, q^{\frac{5}{6}}, q^{\frac{7}{6}}, q^{-\frac{7}{6}}, q^{\frac{13}{6}}, q^{\frac{19}{6}}; q)_n (q^{\frac{11}{6}}; q)_{2n}}{(q^{\frac{1}{3}}; q)_n (q^{\frac{4}{3}}; q)_n (q; q)_{2n} (q^2; q)_{2n} (q^{\frac{10}{3}}; q)_{2n}} \\ &\times \frac{1 - q^{3n + \frac{7}{6}}}{1 - q^{\frac{7}{6}}} \left\{ 1 - \frac{(1 - q^{\frac{2}{3} - n})(1 - q^{2n + \frac{7}{3}})(1 - q^{2n})(1 - q^{2n+1})}{(1 - q^{n - \frac{1}{6}})(1 - q^{n + \frac{7}{6}})(1 - q^{n + \frac{13}{6}})(1 - q^{2n + \frac{5}{6}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{960\Gamma(\frac{2}{3})^3}{7\pi^2} = \sum_{n=0}^{\infty} \left[\frac{1}{6}, -\frac{1}{6}, \frac{7}{6}, -\frac{7}{6}, \frac{13}{6}, \frac{5}{12}, \frac{11}{12}, \frac{25}{18} \right]_n \frac{65 + 372n - 756n^2 - 2160n^3}{16^n}.$$

Example 7 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{6}}, c = q^{\frac{7}{6}}, d = q^{\frac{7}{6}}$ in Theorem 5).

$$\begin{aligned} \left[q^{-\frac{1}{2}}, q^{\frac{1}{6}}, q^{\frac{7}{6}}, q^{\frac{13}{6}} \mid q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 - \frac{n}{3}} \frac{(q^{\frac{1}{6}}, q^{\frac{1}{6}}, q^{\frac{5}{6}}, q^{-\frac{5}{6}}, q^{\frac{7}{6}}, q^{\frac{11}{6}}; q)_n (q^{\frac{5}{6}}; q)_{2n}}{(q^{-\frac{2}{3}}; q)_n (q^{\frac{4}{3}}; q)_n (q; q)_{2n} (q^2; q)_{2n} (q^{\frac{5}{3}}; q)_{2n}} \\ &\times \frac{1 - q^{3n + \frac{7}{6}}}{1 - q^{\frac{7}{6}}} \left\{ 1 + q^{n - \frac{2}{3}} \frac{(1 - q^{n + \frac{1}{6}})(1 - q^{n + \frac{7}{6}})(1 - q^{n + \frac{11}{6}})(1 - q^{2n + \frac{5}{6}})}{(1 - q^{n - \frac{2}{3}})(1 - q^{2n + \frac{1}{3}})(1 - q^{2n+1})(1 - q^{2n+2})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{7776\Gamma(\frac{2}{3})^3}{7\pi^2} = \sum_{n=0}^{\infty} \left[\frac{1}{6}, \frac{5}{6}, \frac{5}{6}, -\frac{5}{6}, \frac{7}{6}, \frac{11}{6}, \frac{5}{12}, \frac{11}{12}, \frac{25}{18} \right]_n \frac{360n^2 + 546n + 191}{16^n}.$$

Example 8 ($a = q^{\frac{1}{2}}, b = q^{\frac{5}{6}}, c = q^{-\frac{1}{6}}, d = q^{-\frac{1}{6}}$ in Theorem 5).

$$\begin{aligned} \left[q^{\frac{3}{2}}, q^{\frac{5}{6}}, q^{\frac{11}{6}}, q^{\frac{17}{6}} \mid q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{4}{3}n} \frac{(q^{\frac{1}{6}}; q)_n^2 (q^{\frac{5}{6}}; q)_n^2 (q^{\frac{11}{6}}; q)_n^2 (q^{\frac{7}{6}}; q)_{2n} (1 - q^{3n + \frac{11}{6}})}{(q^{\frac{2}{3}}; q)_n (q^{\frac{5}{3}}; q)_n (q; q)_{2n} (q^2; q)_{2n} (q^{\frac{8}{3}}; q)_{2n} (1 - q^{\frac{11}{6}})} \\ &\times \left\{ 1 - \frac{(1 - q^{-n - \frac{2}{3}})(1 - q^{2n + \frac{5}{3}})(1 - q^{2n})(1 - q^{2n+1})(1 - q^{3n - \frac{1}{6}})}{(1 - q^{n - \frac{5}{6}})(1 - q^{n - \frac{1}{6}})(1 - q^{n + \frac{5}{6}})(1 - q^{2n + \frac{1}{6}})(1 - q^{3n + \frac{11}{6}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{32\Gamma(\frac{1}{3})^3}{5\pi^2} = \sum_{n=0}^{\infty} \left[\frac{1}{6}, -\frac{1}{6}, \frac{5}{6}, -\frac{5}{6}, \frac{11}{6}, \frac{1}{12}, \frac{7}{12}, \frac{23}{18} \right]_n \frac{2160n^3 + 1188n^2 - 84n + 11}{16^n}.$$

Example 9 ($a = q^{\frac{3}{2}}, b = c = d = q$ in Theorem 5).

$$\begin{aligned} \left[q, q, q, q \mid q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + n} \frac{(q^{\frac{1}{2}}; q)_n^2 (q^2; q)_n (q^{\frac{3}{2}}; q)_{2n} (1 - q^{3n+2})}{(q^3; q^2)_n (q^{\frac{5}{2}}; q)_{2n} (-q; q)_n (1 - q^{\frac{3}{2}})} \\ &\times \left\{ q^{n + \frac{1}{2}} + \frac{(1 + q^{n + \frac{1}{2}})(1 - q^{2n + \frac{3}{2}})^2 (1 - q^{3n+1})}{(1 - q^{n+1})(1 - q^{2n + \frac{1}{2}})(1 - q^{3n+2})} \right\}. \end{aligned}$$

When $q \rightarrow 1$, we recover the following known series (cf. [22]):

$$\frac{9\pi^2}{8} = \sum_{n=0}^{\infty} \left[\begin{matrix} 1, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4} \\ \frac{3}{2}, \frac{5}{4}, \frac{5}{4}, \frac{7}{4}, \frac{7}{4} \end{matrix} \middle| q \right]_n \frac{60n^3 + 111n^2 + 64n + 11}{16^n}.$$

Example 10 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{3}}, c = q^{\frac{1}{3}}, d = q^{-\frac{2}{3}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^2, q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{-\frac{2}{3}} \\ q^{-\frac{1}{2}}, q^{\frac{1}{6}}, q^{\frac{7}{6}}, q^{\frac{7}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{4}{3}n} \frac{(q^{\frac{1}{3}}, q^{-\frac{2}{3}}, q^{\frac{5}{6}}, q^{-\frac{5}{6}}, q^{\frac{11}{6}}; q)_n (q^{-\frac{5}{6}}; q)_{2n} (1 - q^{3n - \frac{2}{3}})}{(q^{\frac{7}{6}}; q)_n (q^{-\frac{1}{2}}; q)_{2n} (q; q)_{2n} (q^{\frac{7}{6}}; q)_{2n} (1 - q^{n + \frac{1}{6}})} \\ &\times \left\{ 1 + q^{n + \frac{7}{6}} \frac{(1 - q^{n - \frac{2}{3}})(1 - q^{n - \frac{5}{6}})(1 - q^{n + \frac{5}{6}})(1 - q^{2n - \frac{5}{6}})(1 - q^{3n + \frac{4}{3}})}{(1 - q^{n + \frac{7}{6}})(1 - q^{2n - \frac{1}{2}})(1 - q^{2n + 1})(1 - q^{2n + \frac{7}{6}})(1 - q^{3n - \frac{2}{3}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{98\pi^2}{3\Gamma(\frac{2}{3})^3} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{3}, -\frac{2}{3}, \frac{5}{6}, -\frac{5}{6}, \frac{11}{6}, \frac{10}{9}, \frac{1}{12}, -\frac{5}{12} \\ 1, \frac{3}{2}, \frac{1}{4}, \frac{3}{4}, \frac{13}{6}, \frac{1}{9}, \frac{13}{12}, \frac{19}{12} \end{matrix} \middle| q \right]_n \frac{118 + 45n - 1098n^2 - 1080n^3}{16^n}.$$

Example 11 ($a = q^{\frac{3}{2}}, b = q^{\frac{1}{3}}, c = q^{\frac{1}{3}}, d = q^{\frac{4}{3}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^2, q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{7}{3}} \\ q^{\frac{1}{2}}, q^{\frac{1}{6}}, q^{\frac{13}{6}}, q^{\frac{13}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{4}{3}n} \frac{(q^{\frac{1}{3}}, q^{\frac{4}{3}}, q^{\frac{5}{6}}, q^{-\frac{5}{6}}, q^{\frac{11}{6}}; q)_n (q^{\frac{1}{6}}; q)_{2n}}{(q^{\frac{13}{6}}; q)_n (q^{\frac{1}{2}}; q)_{2n} (q; q)_{2n} (q^{\frac{13}{6}}; q)_{2n}} \\ &\times \frac{1 - q^{3n + \frac{4}{3}}}{1 - q^{\frac{4}{3}}} \left\{ 1 + q^{n + \frac{1}{6}} \frac{(1 - q^{n + \frac{4}{3}})(1 - q^{n + \frac{11}{6}})(1 - q^{2n + \frac{1}{6}})}{(1 - q^{2n + \frac{1}{2}})(1 - q^{2n + 1})(1 - q^{2n + \frac{13}{6}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{637\pi^2}{16\Gamma(\frac{2}{3})^3} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{3}, \frac{4}{3}, \frac{5}{6}, -\frac{5}{6}, \frac{11}{6}, \frac{13}{9}, \frac{1}{12}, \frac{7}{12} \\ 1, \frac{3}{2}, \frac{3}{4}, \frac{5}{4}, \frac{13}{6}, \frac{4}{9}, \frac{19}{12}, \frac{25}{12} \end{matrix} \middle| q \right]_n \frac{1080n^3 + 2286n^2 + 1395n + 161}{16^n}.$$

Example 12 ($a = q^{\frac{1}{2}}, b = q^{\frac{2}{3}}, c = q^{-\frac{1}{3}}, d = q^{-\frac{1}{3}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^2, q^{-\frac{1}{3}}, q^{-\frac{1}{3}}, q^{\frac{2}{3}} \\ q^{-\frac{1}{2}}, q^{-\frac{1}{6}}, q^{\frac{5}{6}}, q^{\frac{11}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{2}{3}n} \frac{(q^{\frac{2}{3}}, q^{-\frac{1}{3}}, q^{\frac{7}{6}}, q^{-\frac{7}{6}}, q^{\frac{13}{6}}; q)_n (q^{-\frac{1}{6}}; q)_{2n} (1 - q^{3n - \frac{1}{3}})}{(q^{\frac{5}{6}}; q)_n (q^{-\frac{1}{2}}; q)_{2n} (q; q)_{2n} (q^{\frac{11}{6}}; q)_{2n} (1 - q^{n - \frac{1}{6}})} \\ &\times \left\{ 1 + q^{n + \frac{5}{6}} \frac{(1 - q^{n - \frac{1}{3}})(1 - q^{n - \frac{7}{6}})(1 - q^{n + \frac{7}{6}})(1 - q^{2n - \frac{1}{6}})(1 - q^{3n + \frac{5}{3}})}{(1 - q^{n + \frac{5}{6}})(1 - q^{2n - \frac{1}{2}})(1 - q^{2n + 1})(1 - q^{2n + \frac{11}{6}})(1 - q^{3n - \frac{1}{3}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{275\pi^2}{\Gamma(\frac{1}{3})^3} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{2}{3}, -\frac{1}{3}, \frac{7}{6}, -\frac{7}{6}, \frac{13}{6}, \frac{11}{9}, -\frac{1}{12}, \frac{5}{12} \\ 1, \frac{3}{2}, \frac{1}{4}, \frac{3}{4}, \frac{11}{6}, \frac{2}{9}, \frac{17}{12}, \frac{23}{12} \end{matrix} \middle| q \right]_n \frac{125 - 351n - 1602n^2 - 1080n^3}{16^n}.$$

Example 13 ($a = q^{\frac{3}{2}}, b = q^{\frac{2}{3}}, c = q^{-\frac{1}{3}}, d = q^{\frac{5}{3}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^2, q^{-\frac{1}{3}}, q^{\frac{2}{3}}, q^{\frac{8}{3}} \\ q^{\frac{1}{2}}, q^{-\frac{1}{6}}, q^{\frac{11}{6}}, q^{\frac{17}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{2}{3}n} \frac{(q^{\frac{2}{3}}, q^{\frac{5}{3}}, q^{\frac{7}{6}}, q^{-\frac{7}{6}}, q^{\frac{13}{6}}; q)_n (q^{\frac{5}{6}}; q)_{2n}}{(q^{\frac{11}{6}}; q)_n (q^{\frac{1}{2}}; q)_{2n} (q; q)_{2n} (q^{\frac{17}{6}}; q)_{2n}} \\ &\times \frac{1 - q^{3n + \frac{5}{3}}}{1 - q^{\frac{5}{3}}} \left\{ 1 + q^{n - \frac{1}{6}} \frac{(1 - q^{n + \frac{5}{3}})(1 - q^{n + \frac{13}{6}})(1 - q^{2n + \frac{5}{6}})}{(1 - q^{2n + \frac{1}{2}})(1 - q^{2n + 1})(1 - q^{2n + \frac{17}{6}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{2805\pi^2}{4\Gamma(\frac{1}{3})^3} = \sum_{n=0}^{\infty} \left[\frac{2}{3}, \frac{5}{3}, \frac{7}{6}, -\frac{7}{6}, \frac{13}{6}, \frac{14}{9}, \frac{5}{12}, \frac{11}{12} \right]_n \frac{1080n^3 + 2790n^2 + 2151n + 478}{16^n}.$$

Example 14 ($a = q^{\frac{3}{2}}, b = c = q, d = q^{\frac{3}{4}}$ in Theorem 5).

$$\left[\begin{matrix} q, q \\ q^{\frac{1}{2}}, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2 + \frac{5}{4}n} \frac{(q^{\frac{1}{4}}; q)_n (q^{\frac{3}{4}}; q)_n (q^{\frac{1}{4}}; q)_{2n} (1 - q^{3n + \frac{3}{4}})}{(q^{\frac{3}{2}}; q)_{2n} (q^{\frac{5}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q^{n + \frac{1}{2}})} \\ \times \left\{ 1 + q^{n + \frac{3}{4}} \frac{(1 - q^{n + \frac{1}{4}})(1 - q^{n + \frac{1}{2}})(1 - q^{2n + \frac{1}{4}})(1 - q^{3n + 2})}{(1 - q^{2n + 1})(1 - q^{2n + \frac{5}{4}})(1 - q^{2n + \frac{3}{2}})(1 - q^{3n + \frac{3}{4}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$15\pi = \sum_{n=0}^{\infty} \left[\frac{1}{2}, \frac{3}{4}, \frac{1}{8}, \frac{5}{8} \right]_n \frac{120n^2 + 151n + 47}{16^n}.$$

Example 15 ($a = q^{\frac{5}{2}}, b = q, c = q^2, d = q^{\frac{3}{4}}$ in Theorem 5).

$$\left[\begin{matrix} q, q^2, q^{\frac{9}{4}} \\ q^{\frac{3}{2}}, q^{\frac{5}{2}}, q^{\frac{3}{4}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2 + \frac{13}{4}n} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{4}}; q)_n (q^{\frac{3}{4}}; q)_{2n}^2 (q^{-\frac{3}{4}}; q)_{2n} (1 - q^{3n + \frac{3}{4}})}{(q^{\frac{5}{2}}, q^{\frac{7}{4}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (q^{\frac{5}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q^{\frac{3}{4}})} \\ \times \left\{ 1 + q^{n + \frac{7}{4}} \frac{(1 - q^{n + \frac{1}{4}})(1 - q^{n + \frac{1}{2}})(1 - q^{n + \frac{3}{4}})(1 - q^{2n - \frac{3}{4}})(1 - q^{3n + 2})}{(1 - q^{n + \frac{7}{4}})(1 - q^{2n + 1})(1 - q^{2n + \frac{5}{4}})(1 - q^{2n + \frac{3}{2}})(1 - q^{3n + \frac{3}{4}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{63\pi}{2} = \sum_{n=0}^{\infty} \left[\frac{1}{2}, \frac{3}{4}, \frac{1}{8}, -\frac{3}{8} \right]_n \frac{120n^2 + 235n + 99}{16^n}.$$

Example 16 ($a = q^{\frac{3}{2}}, b = q, c = q^{\frac{5}{6}}, d = q$ in Theorem 5).

$$\left[\begin{matrix} q, q^2, q^{\frac{5}{6}}, q^{\frac{7}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{4}{3}}, q^{\frac{5}{3}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2 + n} \frac{(q; q)_n (q^{\frac{1}{3}}; q)_n^2 (q^{\frac{2}{3}}; q)_n^2 (q^{\frac{1}{2}}; q)_{2n} (1 - q^{3n + 1})}{(q^{\frac{1}{2}}; q)_{2n}^2 (q^{\frac{3}{2}}; q)_n (q^{\frac{4}{3}}; q)_{2n} (q^{\frac{5}{3}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q)} \\ \times \left\{ 1 + q^{n + \frac{1}{2}} \frac{(1 - q^{n + \frac{1}{3}})(1 - q^{n + \frac{2}{3}})(1 - q^{n + 1})(1 - q^{2n + \frac{1}{2}})(1 - q^{3n + 2})}{(1 - q^{n + \frac{1}{2}})(1 - q^{2n + 1})(1 - q^{2n + \frac{4}{3}})(1 - q^{2n + \frac{5}{3}})(1 - q^{3n + 1})} \right\}.$$

When $q \rightarrow 1$, we recover another known series (cf. [22]):

$$\frac{20\pi}{9\sqrt{3}} = \sum_{n=0}^{\infty} \left[1, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{3}{4}, \frac{8}{5} \right]_n \frac{12n^2 + 15n + 4}{16^n}.$$

Example 17 ($a = q^{\frac{3}{2}}, b = c = q, d = q^{\frac{5}{6}}$ in Theorem 5).

$$\left[\begin{matrix} q, q, q^{\frac{5}{6}}, q^{\frac{7}{6}} \\ q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{1}{3}}, q^{\frac{2}{3}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2 + \frac{7}{6}n} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{3}}; q)_n (q^{\frac{2}{3}}; q)_n (q^{\frac{5}{6}}; q)_n (1 - q^{3n + \frac{5}{6}})}{(q^{\frac{3}{2}}, q^{\frac{2}{3}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q^{2n + \frac{1}{3}})} \\ \times \left\{ 1 + q^{n + \frac{2}{3}} \frac{(1 - q^{n + \frac{1}{3}})(1 - q^{n + \frac{1}{2}})(1 - q^{n + \frac{5}{6}})(1 - q^{2n + \frac{1}{3}})(1 - q^{3n + 2})}{(1 - q^{n + \frac{2}{3}})(1 - q^{2n + 1})(1 - q^{2n + \frac{4}{3}})(1 - q^{2n + \frac{5}{6}})(1 - q^{3n + \frac{5}{6}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{36\pi}{5\sqrt{3}} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{6}, \frac{5}{6}, \frac{14}{9} \\ \frac{3}{2}, \frac{5}{3}, \frac{5}{4}, \frac{7}{4}, \frac{7}{6}, \frac{5}{9} \end{matrix} \right]_n \frac{60n^2 + 64n + 13}{16^n}.$$

Example 18 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{2}}, d = q^{\frac{1}{4}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{3}{2}} \\ q, q \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{n}{4}} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{4}}; q)_n (q^{\frac{3}{4}}; q)_n (q^{\frac{1}{4}}; q)_{2n} (1 - q^{3n + \frac{1}{4}})}{(q; q)_n^3 (q^{\frac{3}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}^2 (1 - q^{\frac{1}{2}})} \\ &\times \left\{ 1 + \frac{q^{n + \frac{1}{4}} (1 - q^{n + \frac{1}{4}}) (1 - q^{n + \frac{1}{2}}) (1 - q^{2n + \frac{1}{4}}) (1 - q^{3n + \frac{3}{4}})}{(1 - q^{2n + \frac{3}{4}}) (1 - q^{2n + 1})^2 (1 - q^{3n + \frac{1}{4}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following remarkable series:

$$\frac{48}{\pi} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{4}, \frac{3}{4}, \frac{1}{8}, \frac{5}{8} \\ 1, 1, 1, \frac{7}{8}, \frac{11}{8} \end{matrix} \right]_n \frac{480n^2 + 212n + 15}{16^n}.$$

Example 19 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{2}}, d = q^{\frac{3}{4}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{4}}, q^{\frac{7}{4}} \\ q, q, q^{-\frac{1}{4}}, q^{\frac{5}{4}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 - \frac{n}{4}} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{4}}; q)_n (q^{\frac{3}{4}}; q)_n^2 (q^{\frac{3}{4}}; q)_{2n}}{(q; q)_n^3 (q^{-\frac{1}{4}}; q)_n (q^{\frac{5}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}^2} \\ &\times \frac{1 - q^{3n + \frac{3}{4}}}{1 - q^{\frac{3}{4}}} \left\{ 1 + q^{n - \frac{1}{4}} \frac{(1 - q^{n + \frac{3}{4}})^2 (1 - q^{n + \frac{1}{2}}) (1 - q^{2n + \frac{3}{4}}) (1 - q^{3n + \frac{3}{4}})}{(1 - q^{n - \frac{1}{4}}) (1 - q^{2n + 1})^2 (1 - q^{2n + \frac{5}{4}}) (1 - q^{3n + \frac{3}{4}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{80}{3\pi} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{4}, \frac{3}{4}, \frac{3}{8}, \frac{7}{8} \\ 1, 1, 1, \frac{9}{8}, \frac{13}{8} \end{matrix} \right]_n \frac{640n^3 + 560n^2 + 112n + 7}{16^n}.$$

Example 20 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{2}}, d = q^{\frac{1}{3}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{2}{3}}, q^{\frac{4}{3}} \\ q, q, q^{\frac{1}{6}}, q^{\frac{5}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{n}{6}} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{3}}; q)_n^2 (q^{\frac{2}{3}}; q)_n (q^{\frac{1}{3}}; q)_{2n}}{(q; q)_n^3 (q^{\frac{1}{6}}; q)_n (q^{\frac{5}{6}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}^2} \\ &\times \frac{1 - q^{3n + \frac{1}{3}}}{1 - q^{\frac{1}{3}}} \left\{ 1 + \frac{q^{n + \frac{1}{6}} (1 - q^{n + \frac{1}{3}})^2 (1 - q^{n + \frac{1}{2}}) (1 - q^{2n + \frac{1}{3}}) (1 - q^{3n + \frac{2}{3}})}{(1 - q^{n + \frac{1}{6}}) (1 - q^{2n + \frac{5}{6}}) (1 - q^{2n + 1})^2 (1 - q^{3n + \frac{1}{3}})} \right\}. \end{aligned}$$

When $q \rightarrow 1$, we recover the following known series (cf. [22]):

$$\frac{15\sqrt{3}}{\pi} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{3}, \frac{1}{3}, \frac{2}{3}, \frac{2}{3} \\ 1, 1, 1, \frac{11}{12}, \frac{17}{12} \end{matrix} \right]_n \frac{135n^2 + 75n + 8}{16^n}.$$

Example 21 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{6}}, d = q^{\frac{1}{2}}$ in Theorem 5).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{3}{2}}, q^{\frac{5}{2}}, q^{\frac{7}{6}}, q^{\frac{11}{6}} \\ q, q^2, q^{\frac{5}{3}}, q^{\frac{7}{3}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2} \frac{(q^{\frac{1}{2}}, q^{\frac{1}{6}}, q^{\frac{5}{6}}, q^{\frac{7}{6}}, q^{\frac{11}{6}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (1 - q^{3n + \frac{3}{2}})}{(q; q)_n^3 (q^{\frac{5}{3}}; q)_{2n} (q^{\frac{7}{3}}; q)_{2n} (-q; q^{\frac{1}{2}})_{2n} (1 - q^{\frac{3}{2}})} \\ &\times \left\{ q^n + \frac{(1 - q^n) (1 - q^{2n + \frac{2}{3}}) (1 - q^{2n + 1}) (1 - q^{2n + \frac{4}{3}}) (1 - q^{3n + \frac{1}{2}})}{(1 - q^{n + \frac{1}{6}}) (1 - q^{n + \frac{1}{2}}) (1 - q^{n + \frac{5}{6}}) (1 - q^{2n + \frac{1}{2}}) (1 - q^{3n + \frac{3}{2}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{256}{3\pi\sqrt{3}} = \sum_{n=0}^{\infty} \left[\frac{1}{2}, \frac{1}{4}, \frac{3}{4}, \frac{1}{6}, \frac{5}{6} \right]_n \frac{720n^3 + 804n^2 + 236n + 15}{16^n}.$$

Example 22 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{2}}, d = q^{\frac{5}{6}}$ in Theorem 5).

$$\begin{aligned} & \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{6}}, q^{\frac{11}{6}} \\ q, q, q^{-\frac{1}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2 - \frac{n}{3}} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{6}}; q)_n (q^{\frac{5}{6}}; q)_n^2 (q^{\frac{5}{6}}; q)_{2n}}{(q; q)_n^3 (q^{-\frac{1}{3}}; q)_n (q^{\frac{4}{3}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}^2} \\ & \times \frac{1 - q^{3n + \frac{5}{6}}}{1 - q^{\frac{5}{6}}} \left\{ 1 + q^{n - \frac{1}{3}} \frac{(1 - q^{n + \frac{1}{2}})(1 - q^{n + \frac{5}{6}})^2 (1 - q^{2n + \frac{5}{6}})(1 - q^{3n + \frac{3}{2}})}{(1 - q^{n - \frac{1}{3}})(1 - q^{2n + 1})^2 (1 - q^{2n + \frac{4}{3}})(1 - q^{3n + \frac{5}{6}})} \right\}. \end{aligned}$$

When $q \rightarrow 1$, we recover the following known series (cf. [22]):

$$\frac{384}{\pi\sqrt{3}} = \sum_{n=0}^{\infty} \left[\frac{1}{2}, \frac{5}{6}, \frac{5}{6}, \frac{5}{12}, \frac{11}{12} \right]_n \frac{1080n^2 + 798n + 55}{16^n}.$$

§3.2. Analogously, letting $\boxed{\delta = 1}$ in Theorem 4, we obtain, under the partition pattern (15), another summation formula.

Theorem 6 (Nonterminating series identity).

$$\begin{aligned} & \left[\begin{matrix} b, c, qd, qa^2/bcd \\ qa/b, qa/c, a/d, bcd/a \end{matrix} \middle| q \right]_{\infty} = 1 + \sum_{n=1}^{\infty} W_n(a, b, c, d) \left(\frac{q^n a^2}{bd} \right)^n \\ & \times \frac{(bd/a; q)_{2n} (b, d, qa/bc, bc/a, qa/cd, cd/a; q)_n}{(q, qa/c, bcd/a; q)_{2n} (qa/b, a/d; q)_n}, \end{aligned}$$

where the weight function is given by

$$\begin{aligned} W_n(a, b, c, d) &= \frac{1 - q^{3n}d}{1 - d} \\ & \times \left\{ 1 - \frac{(1 - q^{2n})(1 - q^{-n}b/a)(1 - q^{2n}a/c)(1 - q^{2n-1}bcd/a)(1 - q^{3n-2}b)}{(1 - q^n a/cd)(1 - q^{n-1}b)(1 - q^{n-1}bc/a)(1 - q^{2n-1}bd/a)(1 - q^{3n}d)} \right\}. \end{aligned}$$

By specifying the parameters $\{a, b, c, d\}$ concretely, we can deduce from Theorem 6 a number of q -series identities that are q -analogues of classical series counterparts with the convergence rate " $\frac{1}{16}$ ". Twelve formulae are recorded below as examples.

Example 23 ($a = b = c = d = q^{\frac{1}{2}}$ in Theorem 6).

$$\begin{aligned} & \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{3}{2}} \\ q, q, q, q \end{matrix} \middle| q \right]_{\infty} = \sum_{n=1}^{\infty} q^{n^2} \frac{(q^{\frac{1}{2}}; q)_n^3 (q^{\frac{1}{2}}; q)_{2n}}{(q; q)_n^5 (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}^3} \frac{(1 - q^n)(1 - q^{3n + \frac{1}{2}})}{(1 - q^{1/2})(1 - q^{3/2})} \\ & \times \left\{ 1 - \frac{(1 - q^{-n})(1 - q^{2n})^3 (1 - q^{3n - \frac{3}{2}})}{(1 - q^{2n - \frac{1}{2}})(1 - q^{n - \frac{1}{2}})^3 (1 - q^{3n + \frac{1}{2}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{2}{\pi^2} = \sum_{n=1}^{\infty} \left[\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{4}, -\frac{1}{4} \right]_n \frac{n(1 - 6n)(1 + 4n - 4n^2 + 80n^3)}{16^n}.$$

Example 24 $\left(a = q^{\frac{1}{2}}, b = q^{\frac{5}{6}}, c = q^{\frac{5}{6}}, d = q^{\frac{5}{6}} \right)$ in Theorem 6).

$$\left[q^{\frac{1}{2}}, q^{\frac{5}{6}}, q^{\frac{5}{6}}, q^{\frac{5}{6}} \mid q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2 - \frac{2}{3}n} \frac{(q^{-\frac{1}{6}}; q)_n^2 (q^{\frac{5}{6}}; q)_n^2 (q^{\frac{7}{6}}; q)_n^2 (q^{\frac{7}{6}}; q)_{2n}}{(q^{-\frac{1}{3}}; q)_n (q^{\frac{2}{3}}; q)_n (q; q)_{2n} (q^2; q)_{2n} (q^{\frac{2}{3}}; q)_{2n}} \\ \times \frac{1 - q^{3n + \frac{5}{6}}}{1 - q^{-\frac{1}{2}}} \left\{ 1 - \frac{(1 - q^{\frac{1}{3} - n})(1 - q^{2n - \frac{1}{3}})(1 - q^{2n})(1 - q^{2n+1})(1 - q^{3n - \frac{7}{6}})}{(1 - q^{n - \frac{7}{6}})(1 - q^{n - \frac{1}{6}})(1 - q^{n + \frac{1}{6}})(1 - q^{2n + \frac{1}{6}})(1 - q^{3n + \frac{5}{6}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{63\Gamma(\frac{1}{3})^3}{10\pi^2} = \sum_{n=0}^{\infty} \left[\frac{1}{6}, -\frac{1}{6}, \frac{5}{6}, \frac{7}{6}, -\frac{7}{6}, \frac{1}{12}, \frac{7}{12}, \frac{17}{18} \right]_n \frac{432n^3 - 108n^2 - 132n + 7}{16^n}.$$

Example 25 $\left(a = q^{\frac{5}{2}}, b = q, c = q^2, d = q \right)$ in Theorem 6).

$$\left[q, q, q^2, q^2 \mid q \right]_{\infty} = 1 + q^{\frac{1}{2}} + \sum_{n=1}^{\infty} q^{n^2 + 3n} \frac{(q^{\frac{1}{2}}; q)_n^2 (q; q)_n (q^{-\frac{1}{2}}; q)_{2n}}{(q^{\frac{5}{2}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}} \\ \times \frac{1 - q^{3n+1}}{1 - q^{n+\frac{1}{2}}} \left\{ 1 - \frac{(1 + q^n)(1 - q^{-n - \frac{3}{2}})(1 - q^{2n + \frac{1}{2}})^2(1 - q^{3n-1})}{(1 - q^{n - \frac{1}{2}})^2(1 - q^{2n - \frac{3}{2}})(1 - q^{3n+1})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{3\pi^2}{32} = 1 + \sum_{n=1}^{\infty} \left[1, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{4}, -\frac{3}{4} \right]_n \frac{3 + 3n - 22n^2 - 40n^3}{16^n}.$$

Example 26 $\left(a = q^{\frac{7}{2}}, b = q, c = q^2, d = q \right)$ in Theorem 6).

$$\left[q, q^2, q^2, q^4 \mid q \right]_{\infty} = 1 + \sum_{n=1}^{\infty} q^{n^2 + 5n} \frac{(q^{-\frac{1}{2}}; q)_n^2 (q; q)_n (q^{-\frac{3}{2}}; q)_{2n} (1 - q^{3n+1})}{(q^{\frac{7}{2}}; q)_n (q^{\frac{1}{2}}; q)_{2n} (q^{\frac{5}{2}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q)} \\ \times \frac{(1 - q^{\frac{3}{2}})(1 - q^{n+\frac{1}{2}})}{(1 - q^{\frac{1}{2}})(1 - q^{n+\frac{3}{2}})} \left\{ 1 - \frac{(1 + q^n)(1 - q^{-n - \frac{5}{2}})(1 - q^{2n - \frac{1}{2}})(1 - q^{2n + \frac{3}{2}})(1 - q^{3n-1})}{(1 - q^{n - \frac{3}{2}})(1 - q^{n + \frac{1}{2}})(1 - q^{2n - \frac{5}{2}})(1 - q^{3n+1})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{15\pi^2}{256} = \frac{1}{3} + \sum_{n=1}^{\infty} \left[1, -\frac{1}{2}, -\frac{3}{2}, -\frac{3}{4}, -\frac{5}{4} \right]_n \frac{8n^3 + 2n^2 - 3n + 1}{16^n}.$$

Example 27 $\left(a = q^{\frac{5}{2}}, b = q^2, c = q^2, d = q^{\frac{3}{4}} \right)$ in Theorem 6).

$$\left[q, q^2 \mid q \right]_{\infty} = \sum_{n=0}^{\infty} q^{n^2 + \frac{9}{4}n} \frac{(q^{-\frac{1}{2}}; q)_n (q^{\frac{1}{4}}; q)_n (q^{\frac{3}{4}}; q)_n (q^{\frac{1}{4}}; q)_{2n} (1 - q^{n+1})}{(q^{\frac{5}{2}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (q^{\frac{9}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q^{n+\frac{3}{4}})} \\ \times \frac{1 - q^{3n + \frac{3}{4}}}{1 - q^{\frac{5}{4}}} \left\{ 1 + \frac{q^{-n - \frac{1}{2}}(1 - q^{2n})(1 - q^{3n})(1 - q^{2n + \frac{1}{2}})(1 - q^{2n + \frac{5}{4}})}{(1 - q^{n - \frac{1}{4}})(1 - q^{n+1})(1 - q^{2n - \frac{3}{4}})(1 - q^{3n + \frac{3}{4}})} \right\}.$$

This provides a q -analogue for the following classical series:

$$\frac{5\pi}{16} = \sum_{n=0}^{\infty} \left[-\frac{1}{2}, -\frac{1}{4}, \frac{1}{8}, -\frac{3}{8} \right]_n \frac{40n^2 - 7n + 1}{16^n}.$$

Example 28 ($a = q^{\frac{5}{2}}, b = q^2, c = q^2, d = q^{\frac{1}{4}}$ in Theorem 6).

$$\begin{aligned} \left[\begin{matrix} q, q^2 \\ q^{\frac{3}{2}}, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2 + \frac{11}{4}n} \frac{(q^{-\frac{1}{2}}; q)_n (q^{\frac{1}{4}}; q)_n (q^{-\frac{1}{4}}; q)_n (q^{-\frac{1}{4}}; q)_{2n} (1 - q^{3n + \frac{1}{4}})}{(q^{\frac{1}{2}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (q^{\frac{7}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q^{\frac{1}{4}})} \\ &\times \frac{1 - q^{n+1}}{1 - q^{n + \frac{5}{4}}} \left\{ 1 + \frac{q^{-n - \frac{1}{2}} (1 - q^{2n}) (1 - q^{3n}) (1 - q^{2n + \frac{1}{2}}) (1 - q^{2n + \frac{3}{4}})}{(1 - q^{n + \frac{1}{4}}) (1 - q^{n+1}) (1 - q^{2n - \frac{5}{4}}) (1 - q^{3n + \frac{1}{4}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{25\pi}{16} = \sum_{n=0}^{\infty} \left[\begin{matrix} -\frac{1}{2}, \frac{1}{4}, -\frac{1}{8}, -\frac{5}{8} \\ \frac{1}{2}, \frac{9}{4}, \frac{7}{8}, \frac{11}{8} \end{matrix} \right]_n \frac{120n^2 + 77n + 5}{16^n}.$$

Example 29 ($a = q^{\frac{3}{2}}, b = q, c = q, d = q^{\frac{1}{4}}$ in Theorem 6).

$$\begin{aligned} \left[\begin{matrix} q, q, q^{\frac{7}{4}} \\ q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{4}} \end{matrix} \middle| q \right]_{\infty} &= 1 + \sum_{n=1}^{\infty} q^{n^2 + \frac{7}{4}n} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{4}}; q)_n (q^{-\frac{1}{4}}; q)_n (q^{-\frac{1}{4}}; q)_{2n}}{(q^{\frac{3}{2}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (q^{\frac{3}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}} \\ &\times \frac{1 - q^{3n + \frac{1}{4}}}{1 - q^{\frac{1}{4}}} \left\{ 1 - \frac{(1 + q^n) (1 - q^{-n - \frac{1}{2}}) (1 - q^{2n - \frac{1}{4}}) (1 - q^{2n + \frac{1}{2}}) (1 - q^{3n - 1})}{(1 - q^{n - \frac{1}{2}}) (1 - q^{n + \frac{1}{4}}) (1 - q^{2n - \frac{5}{4}}) (1 - q^{3n + \frac{1}{4}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{5\pi}{9} = \frac{5}{3} + \sum_{n=1}^{\infty} \left[\begin{matrix} -\frac{1}{2}, \frac{1}{4}, -\frac{1}{8}, -\frac{5}{8} \\ \frac{3}{2}, \frac{5}{4}, \frac{3}{8}, \frac{7}{8} \end{matrix} \right]_n \frac{7 - 6n - 80n^2}{16^n}.$$

Example 30 ($a = q^{\frac{5}{2}}, b = q, c = q^2, d = q^{\frac{1}{4}}$ in Theorem 6).

$$\begin{aligned} \left[\begin{matrix} q, q^2, q^{\frac{5}{4}}, q^{\frac{11}{4}} \\ q^{\frac{3}{2}}, q^{\frac{5}{2}}, q^{\frac{3}{4}}, q^{\frac{9}{4}} \end{matrix} \middle| q \right]_{\infty} &= 1 + \sum_{n=1}^{\infty} q^{n^2 + \frac{15}{4}n} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{4}}; q)_n (q^{-\frac{1}{4}}; q)_n (q^{\frac{5}{4}}; q)_n (q^{-\frac{5}{4}}; q)_{2n}}{(q^{\frac{5}{2}}; q)_n (q^{\frac{9}{4}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (q^{\frac{3}{4}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}} \\ &\times \frac{1 - q^{3n + \frac{1}{4}}}{1 - q^{\frac{1}{4}}} \left\{ 1 - \frac{(1 + q^n) (1 - q^{-n - \frac{3}{2}}) (1 - q^{2n - \frac{1}{4}}) (1 - q^{2n + \frac{1}{2}}) (1 - q^{3n - 1})}{(1 - q^{n - \frac{1}{2}}) (1 - q^{n + \frac{1}{4}}) (1 - q^{2n - \frac{9}{4}}) (1 - q^{3n + \frac{1}{4}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{15\pi}{14} = 3 + \sum_{n=1}^{\infty} \left[\begin{matrix} -\frac{1}{2}, \frac{1}{4}, -\frac{5}{8}, -\frac{9}{8} \\ \frac{5}{2}, \frac{9}{4}, \frac{3}{8}, \frac{7}{8} \end{matrix} \right]_n \frac{19 - 62n - 80n^2}{16^n}.$$

Example 31 ($a = q^{\frac{1}{2}}, b = q, c = q^{\frac{1}{6}}, d = q$ in Theorem 6).

$$\begin{aligned} \left[\begin{matrix} q, q^2, q^{\frac{1}{6}}, q^{-\frac{1}{6}} \\ q^{-\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{4}{3}}, q^{\frac{5}{3}} \end{matrix} \middle| q \right]_{\infty} &= 1 + \sum_{n=1}^{\infty} q^{n^2 - n} \frac{(q; q)_n (q^{\frac{1}{3}}; q)_n^2 (q^{\frac{2}{3}}; q)_n^2 (q^{\frac{2}{3}}; q)_{2n}}{(q^{\frac{1}{2}}; q)_n^2 (q^{-\frac{1}{2}}; q)_n (q^{\frac{4}{3}}; q)_{2n} (q^{\frac{5}{3}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}} \\ &\times \frac{1 - q^{3n+1}}{1 - q} \left\{ 1 - \frac{(1 + q^n) (1 - q^{\frac{1}{2} - n}) (1 - q^{2n + \frac{1}{3}}) (1 - q^{2n + \frac{2}{3}}) (1 - q^{3n - 1})}{(1 - q^{n - \frac{2}{3}}) (1 - q^{n - \frac{1}{3}}) (1 - q^{2n + \frac{1}{2}}) (1 - q^{3n + 1})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{4\pi}{81\sqrt{3}} = \frac{2}{3} + \sum_{n=1}^{\infty} \left[\begin{matrix} 1, \frac{2}{3}, -\frac{2}{3}, \frac{1}{4}, \frac{3}{4} \\ \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{5}{6}, \frac{7}{6} \end{matrix} \right]_n \frac{2 + 7n - 20n^2}{16^n}.$$

Example 32 ($a = q^{\frac{1}{2}}, b = q, c = q^{\frac{1}{6}}, d = q^2$) in Theorem 6).

$$\begin{aligned} \left[\begin{matrix} q, q^3, q^{\frac{1}{6}}, q^{-\frac{7}{6}} \\ q^{-\frac{3}{2}}, q^{\frac{1}{2}}, q^{\frac{4}{3}}, q^{\frac{5}{3}} \end{matrix} \middle| q \right]_{\infty} &= 1 + \sum_{n=1}^{\infty} q^{n^2-2n} \frac{(q^2; q)_n (q^{\frac{1}{3}}; q)_n (q^{\frac{2}{3}}; q)_n (q^{-\frac{2}{3}}; q)_n (q^{\frac{5}{3}}; q)_n (q^{\frac{5}{3}}; q)_{2n}}{(q^{\frac{1}{2}}; q)_{2n}^2 (q^{-\frac{3}{2}}; q)_n (q^{\frac{4}{3}}; q)_{2n} (q^{\frac{8}{3}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}} \\ &\times \frac{1 - q^{3n+2}}{1 - q^2} \left\{ 1 - \frac{(1 + q^n)(1 - q^{\frac{1}{2}-n})(1 - q^{2n+\frac{1}{3}})(1 - q^{2n+\frac{5}{3}})(1 - q^{3n-1})}{(1 - q^{n-\frac{5}{3}})(1 - q^{n-\frac{1}{3}})(1 - q^{2n+\frac{3}{2}})(1 - q^{3n+2})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{350\pi}{243\sqrt{3}} = 30 + \sum_{n=1}^{\infty} \left[\begin{matrix} 2, \frac{5}{3}, -\frac{5}{3}, \frac{3}{4}, \frac{5}{4} \\ \frac{1}{2}, \frac{1}{2}, -\frac{3}{2}, \frac{7}{6}, \frac{11}{6} \end{matrix} \right]_n \frac{40 - n - 60n^2}{16^n}.$$

Example 33 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{6}}, d = q^{\frac{1}{2}}$) in Theorem 6).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{5}{6}}, q^{\frac{7}{6}} \\ q, q, q^{\frac{2}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2} \frac{(q^{\frac{1}{2}}; q)_n (q^{\frac{1}{6}}; q)_n^2 (q^{\frac{5}{6}}; q)_n^2 (q^{\frac{1}{2}}; q)_{2n} (1 - q^{3n+\frac{1}{2}})}{(q; q)_n^3 (q^{\frac{2}{3}}; q)_{2n} (q^{\frac{4}{3}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n} (1 - q^{\frac{1}{6}})} \\ &\times \frac{1 - q^n}{1 - q^{\frac{1}{2}}} \left\{ 1 - \frac{(1 - q^{-n})(1 - q^{2n})(1 - q^{2n-\frac{1}{3}})(1 - q^{2n+\frac{1}{3}})(1 - q^{3n-\frac{3}{2}})}{(1 - q^{n-\frac{1}{2}})(1 - q^{n-\frac{1}{6}})(1 - q^{n-\frac{5}{6}})(1 - q^{2n-\frac{1}{2}})(1 - q^{3n+\frac{1}{2}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{10\sqrt{3}}{9\pi} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{1}{2}, \frac{1}{4}, -\frac{1}{4}, \frac{5}{6}, -\frac{5}{6} \\ 1, 1, 1, \frac{1}{3}, \frac{2}{3} \end{matrix} \right]_n \frac{n(120n^2 - 26n + 5)}{16^n}.$$

Example 34 ($a = q^{\frac{1}{2}}, b = q^{\frac{1}{2}}, c = q^{\frac{1}{2}}, d = q^{-\frac{1}{3}}$) in Theorem 6).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{2}{3}}, q^{\frac{4}{3}} \\ q, q, q^{\frac{1}{6}}, q^{\frac{5}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{n^2+\frac{5}{6}n} \frac{(q^{\frac{1}{2}}; q)_n (q^{-\frac{1}{3}}; q)_n^2 (q^{\frac{4}{3}}; q)_n (q^{-\frac{1}{3}}; q)_{2n}}{(q; q)_n^3 (q^{\frac{5}{6}}; q)_n (q^{\frac{1}{6}}; q)_{2n} (-q^{\frac{1}{2}}; q^{\frac{1}{2}})_{2n}^2} \\ &\times \frac{1 - q^{3n-\frac{1}{3}}}{1 - q^{-\frac{1}{3}}} \left\{ 1 - \frac{(1 - q^{-n})(1 - q^{2n})^2 (1 - q^{2n-\frac{5}{6}})(1 - q^{3n-\frac{3}{2}})}{(1 - q^{n-\frac{1}{2}})^2 (1 - q^{n+\frac{1}{3}})(1 - q^{2n-\frac{4}{3}})(1 - q^{3n-\frac{1}{3}})} \right\}. \end{aligned}$$

This provides a q -analogue for the following classical series:

$$\frac{6\sqrt{3}}{\pi} = \sum_{n=0}^{\infty} \left[\begin{matrix} -\frac{1}{2}, \frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}, -\frac{2}{3} \\ 1, 1, 1, \frac{1}{12}, \frac{7}{12} \end{matrix} \right]_n \frac{135n^3 - 48n^2 - 7n + 2}{16^n}.$$

4. SERIES FROM TRIPPLICATE INVERSIONS

For all $n \in \mathbb{N}_0$, it is well known that $n = \lfloor \frac{n}{3} \rfloor + \lfloor \frac{1+n}{3} \rfloor + \lfloor \frac{2+n}{3} \rfloor$. Then it is not difficult to check that Jackson's formula (13) is equivalent to the following one

$$\begin{aligned} \Omega_n(a; q^{\lfloor \frac{n}{3} \rfloor} b, q^{\lfloor \frac{1+n}{3} \rfloor} c, q^{\lfloor \frac{2+n}{3} \rfloor} d) &= \left[\begin{matrix} qa/cd \\ b/a \end{matrix} \middle| q \right]_{\lfloor \frac{n}{3} \rfloor} \left[\begin{matrix} qa/bd \\ c/a \end{matrix} \middle| q \right]_{\lfloor \frac{1+n}{3} \rfloor} \left[\begin{matrix} qa/bc \\ d/a \end{matrix} \middle| q \right]_{\lfloor \frac{2+n}{3} \rfloor} \\ &\times \left[\begin{matrix} qa \\ bcd/a \end{matrix} \middle| q \right]_n \left[\begin{matrix} bc/a \\ qa/d \end{matrix} \middle| q \right]_{\lfloor \frac{2n}{3} \rfloor} \left[\begin{matrix} bd/a \\ qa/c \end{matrix} \middle| q \right]_{\lfloor \frac{1+2n}{3} \rfloor} \left[\begin{matrix} cd/a \\ qa/b \end{matrix} \middle| q \right]_{\lfloor \frac{2+2n}{3} \rfloor} \end{aligned}$$

with its parameters subject to $\boxed{qa^2 = bcde}$. The partition pattern corresponding to (14) becomes

$$\Lambda = 3 : \quad \begin{array}{cccc} \varepsilon_b = 0, & \varepsilon_c = 1, & \varepsilon_d = 2, & \varepsilon_e = 0; \\ \lambda_b = 1, & \lambda_c = 1, & \lambda_d = 1, & \lambda_e = 0. \end{array} \quad (16)$$

§4.1. Specifying with $\boxed{\delta = 0}$ and the above partition pattern in Theorem 3, we can express the corresponding formula as in the following theorem.

Theorem 7 (Nonterminating series identity).

$$\begin{aligned} \left[\begin{array}{c} b, qc, qd, qa^2/bcd \\ qa/b, qa/c, qa/d, bcd/a \end{array} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} W_n(a, b, c, d) \left(\frac{q^{3n} a^3}{bcd} \right)^n \\ &\times \frac{(b, c, qd, qa/bc, qa/bd, qa/cd; q)_n (bc/a, bd/a, cd/a; q)_{2n}}{(qa/b, qa/c, qa/d; q)_{2n} (q, bcd/a; q)_{3n}}, \end{aligned}$$

where the weight function is given by

$$\begin{aligned} W_n(a, b, c, d) &= \frac{(1 - q^{2n} bd/a)(1 - q^{1+n} a/(bc))(1 - q^{2n} cd/a)(1 - q^{1+4n} c)}{(1 - q^{1+3n})(1 - c)(1 - q^{1+2n} a/b)(1 - q^{3n} bcd/a)} \\ &\times \left\{ q^{2n} \frac{a}{d} + \frac{(1 - q^{2n} a/d)(1 - q^{3n} bcd/a)(1 - q^{4n} d)(1 - q^{1+2n} a/b)(1 - q^{1+3n})}{(1 - q^n d)(1 - q^{2n} bd/a)(1 - q^{2n} cd/a)(1 - q^{1+n} a/(bc))(1 - q^{1+4n} c)} \right. \\ &\left. + \frac{q^{1+4n} a^2 (1 - q^n c)(1 - q^{2n} bc/a)(1 - q^{1+n} a/(bd))(1 - q^{1+2n} cd/a)(1 - q^{2+4n} b)}{cd(1 - q^{1+2n} a/c)(1 - q^{1+2n} a/d)(1 - q^{1+3n} bcd/a)(1 - q^{2+3n})(1 - q^{1+4n} c)} \right\}. \end{aligned}$$

According to this theorem, by assigning concrete values for the quadruplet $\{a, b, c, d\}$, we can derive numerous q -series identities after some routine simplifications. Even though the q -series so obtained are quite complicated in general, they can dramatically be reduced, in some cases, to simple ones by means of the ‘‘reverse bisection method’’. We take the formulae from Examples 40 and 42 to illustrate how this approach works.

When $\boxed{a = q^{\frac{4}{3}}, b = c = q, d = q^{\frac{5}{6}}}$, the corresponding formula in Theorem 7 reads as

$$\left[\begin{array}{c} q, q, q^{\frac{5}{6}}, q^{\frac{5}{6}} \\ q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{4}{3}}, q^{\frac{4}{3}} \end{array} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{3n^2 + \frac{7}{6}n} \frac{(q, q, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{3}}, q^{\frac{5}{6}}; q)_n (q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{2}{3}}; q)_{2n}}{(q^{\frac{3}{2}}, q^{\frac{4}{3}}, q^{\frac{4}{3}}; q)_{2n} (q, q^{\frac{3}{2}}; q)_{3n}} W_n(q^{\frac{4}{3}}, q, q, q^{\frac{5}{6}}),$$

where W_n is a rational function of q^n given explicitly by

$$\begin{aligned} W_n(q^{\frac{4}{3}}, q, q, q^{\frac{5}{6}}) &= \frac{(1 - q^{n+\frac{1}{3}})(1 - q^{n+\frac{5}{6}})(1 - q^{2n+\frac{1}{2}})^2(1 - q^{4n+2})}{(1 - q^{2n+\frac{4}{3}})(1 - q^{3n+1})(1 - q^{3n+\frac{3}{2}})} \\ &\times \left\{ q^{2n+\frac{1}{2}} + \frac{(1 - q^{2n+\frac{4}{3}})(1 - q^{3n+1})(1 - q^{3n+\frac{3}{2}})(1 - q^{4n+\frac{5}{6}})}{(1 - q^{n+\frac{1}{3}})(1 - q^{n+\frac{5}{6}})(1 - q^{2n+\frac{1}{2}})(1 - q^{4n+2})} \right. \\ &\left. + \frac{q^{4n+\frac{11}{6}}(1 - q^{n+\frac{1}{2}})(1 - q^{n+1})(1 - q^{2n+\frac{2}{3}})(1 - q^{4n+3})}{(1 - q^{2n+\frac{4}{3}})(1 - q^{3n+2})(1 - q^{3n+\frac{5}{2}})(1 - q^{4n+2})} \right\}. \end{aligned}$$

By factorization with *Mathematica* commands, we can check that

$$P(q^n) := \frac{(1 - q^{2n+\frac{4}{3}})^2(1 - q^{3n+1})(1 - q^{3n+2})(1 - q^{3n+\frac{3}{2}})(1 - q^{3n+\frac{5}{2}})}{(1 - q^{n+\frac{1}{3}})(1 - q^{2n+\frac{1}{2}})} W_n(q^{\frac{4}{3}}, q, q, q^{\frac{5}{6}})$$

is a polynomial of degree 19 in q^n . Then the preceding formula in question can be rewritten, by substitution, as

$$\left[\begin{matrix} q, q, q^{\frac{5}{6}}, q^{\frac{5}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{1}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{3n^2 + \frac{7}{6}n} \frac{(q^{\frac{1}{2}}, q^{\frac{2}{3}}; q)_{2n} (q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{5}{6}}; q^{\frac{1}{2}})_{2n}}{(1-q)(q^{\frac{4}{3}}, q^{\frac{4}{3}}; q)_{2n+1} (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{6n+3}} P(q^n). \quad (17)$$

Supposing that $Q(y)$ is a polynomial (to be determined), the series on the right suggests us to introduce the hypergeometric term

$$T_n := q^{\frac{3}{4}n^2 + \frac{7}{12}n} \frac{(q^{\frac{1}{2}}, q^{\frac{2}{3}}; q)_n (q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{5}{6}}; q^{\frac{1}{2}})_n}{(1-q)(q^{\frac{4}{3}}, q^{\frac{4}{3}}; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n}} Q(q^{\frac{n}{2}}).$$

By computing the difference of two consecutive terms

$$\begin{aligned} T_{2n} \pm T_{2n+1} &= q^{3n^2 + \frac{7}{6}n} \frac{(q^{\frac{1}{2}}, q^{\frac{2}{3}}; q)_{2n} (q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{5}{6}}; q^{\frac{1}{2}})_{2n}}{(1-q)(q^{\frac{4}{3}}, q^{\frac{4}{3}}; q)_{2n+1} (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{6n+3}} \\ &\times \left\{ Q(q^n) (1 - q^{\frac{4}{3}+2n})^2 (1 - q^{\frac{3}{2}+3n}) (1 - q^{2+3n}) (1 - q^{\frac{5}{2}+3n}) \right. \\ &\left. \pm q^{3n + \frac{4}{3}} Q(q^{n+\frac{1}{2}}) (1 - q^{\frac{1}{2}+n})^2 (1 - q^{\frac{5}{6}+n}) (1 - q^{\frac{1}{2}+2n}) (1 - q^{\frac{2}{3}+2n}) \right\}, \end{aligned}$$

we can formally consider the series in (17) as the following ‘‘bisection series’’

$$\left[\begin{matrix} q, q, q^{\frac{5}{6}}, q^{\frac{5}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{1}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} (\pm 1)^n T_n = \sum_{n=0}^{\infty} \{ T_{2n} \pm T_{2n+1} \}, \quad (18)$$

where the signs ‘ \pm ’ indicate that the series can be positive or alternating. By equating the summand of (17) with the expression of ‘‘ $T_{2n} \pm T_{2n+1}$ ’’ and then canceling the common factors, we get a polynomial equation

$$\begin{aligned} P(y) &= Q(y) (1 - q^{\frac{4}{3}} y^2)^2 (1 - q^{\frac{3}{2}} y^3) (1 - q^2 y^3) (1 - q^{\frac{5}{2}} y^3) \\ &\pm q^{\frac{4}{3}} y^3 Q(q^{\frac{1}{2}} y) (1 - q^{\frac{1}{2}} y)^2 (1 - q^{\frac{5}{6}} y) (1 - q^{\frac{1}{2}} y^2) (1 - q^{\frac{2}{3}} y^2), \end{aligned} \quad (19)$$

where q^n is replaced by y for brevity. Now that $P(y)$ is a polynomial of degree 19 in y , we may further take for granted that $Q(y)$ is a polynomial of degree 6 so that both sides of the above equation have the same degree. Writing explicitly

$$Q(y) = a_0 + a_1 y + a_2 y^2 + a_3 y^3 + a_4 y^4 + a_5 y^5 + a_6 y^6$$

and then comparing the coefficients of y^k for $0 \leq k \leq 19$ across equation (19), we get a system of 20 linear equations in seven variables $\{a_0, a_1, a_2, a_3, a_4, a_5, a_6\}$. The system corresponding to the positive sign ‘ $+$ ’ is not compatible. Resolving, instead, the system with the minus sign ‘ $-$ ’ of equations by *Mathematica*, we get the following solution:

$$\left\{ a_0 \rightarrow 1, a_1 \rightarrow q^{\frac{1}{3}}, a_2 \rightarrow q^{\frac{1}{2}} + q^{\frac{2}{3}}, a_3 \rightarrow 0, a_4 \rightarrow -q - q^{\frac{5}{6}}, a_5 \rightarrow -q^{\frac{7}{6}}, a_6 \rightarrow -q^{\frac{3}{2}} \right\}.$$

Therefore, we have determined the polynomial

$$\begin{aligned} Q(y) &= 1 + q^{\frac{1}{3}} y + q^{\frac{1}{2}} y^2 + q^{\frac{2}{3}} y^2 - q^{\frac{5}{6}} y^4 - q y^4 - q^{\frac{7}{6}} y^5 - q^{\frac{3}{2}} y^6 \\ &= q^{\frac{1}{2}} y^2 (1 - q^{\frac{1}{3}} y^2) (1 + q^{\frac{1}{6}}) + (1 + q^{\frac{1}{3}} y) (1 - q^{\frac{7}{6}} y^5). \end{aligned}$$

Finally, substituting $Q(y)$ and T_n into (18) and simplifying slightly the expression, we arrive at

$$\begin{aligned} \left[\begin{matrix} q, q, q^{\frac{5}{6}}, q^{\frac{5}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{1}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q^{\frac{5}{6}}; q)_n (q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{3}}; q^{\frac{1}{2}})_n}{(q^{\frac{4}{3}}, q^{\frac{4}{3}}; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n}} q^{\frac{n(9n+7)}{12}} \\ &\times \frac{1 - q^{\frac{2}{3}}}{1 - q} \left\{ q^{n+\frac{1}{2}} \frac{(1 + q^{\frac{1}{6}})}{(1 + q^{\frac{2}{3}+\frac{1}{3}})} + \frac{1 - q^{\frac{5}{2}n+\frac{7}{6}}}{1 - q^{n+\frac{1}{3}}} \right\}, \end{aligned}$$

which is exactly the identity displayed in Example 40. \square

The same “reverse bisection method” can be applied to derive the formula in Example 42. In fact, by specifying $\boxed{a = b = c = q^{\frac{1}{3}}, d = q^{\frac{1}{6}}}$ in Theorem 7, we can explicitly write down the formula

$$\left[\begin{matrix} q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{6}}, q^{\frac{5}{6}} \\ q, q, q^{\frac{1}{2}}, q^{\frac{7}{6}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{3n^2 + \frac{n}{6}} \frac{(q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{2}{3}}, q^{\frac{1}{6}}, q^{\frac{5}{6}}, q^{\frac{5}{6}}; q)_n (q^{\frac{1}{3}}, q^{\frac{1}{6}}, q^{\frac{1}{6}}; q)_{2n}}{(q, q, q^{\frac{7}{6}}; q)_{2n} (q, q^{\frac{1}{2}}; q)_{3n}} \mathcal{W}_n(q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{6}}),$$

where \mathcal{W}_n is a rational function of q^n defined by

$$\begin{aligned} \mathcal{W}_n(q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{6}}) &= \frac{(1 - q^{n + \frac{2}{3}})(1 - q^{n + \frac{1}{6}})(1 - q^{2n + \frac{1}{6}})^2(1 - q^{4n + \frac{4}{3}})}{(1 - q^{2n+1})(1 - q^{3n+1})(1 - q^{3n + \frac{1}{2}})} \\ &\times \left\{ q^{2n + \frac{1}{6}} + \frac{(1 - q^{2n+1})(1 - q^{3n+1})(1 - q^{3n + \frac{1}{2}})(1 - q^{4n + \frac{4}{3}})}{(1 - q^{n + \frac{2}{3}})(1 - q^{n + \frac{1}{6}})(1 - q^{2n + \frac{1}{6}})(1 - q^{4n + \frac{4}{3}})} \right. \\ &\left. + \frac{q^{4n + \frac{7}{6}}(1 - q^{n + \frac{1}{3}})(1 - q^{n + \frac{5}{6}})(1 - q^{2n + \frac{1}{3}})(1 - q^{4n + \frac{7}{3}})}{(1 - q^{2n+1})(1 - q^{3n+2})(1 - q^{3n + \frac{3}{2}})(1 - q^{4n + \frac{4}{3}})} \right\}. \end{aligned}$$

Analogously, it is can be verified without difficulty that

$$\mathcal{P}(q^n) := \frac{(1 - q^{2n+1})^2(1 - q^{3n+1})(1 - q^{3n+2})(1 - q^{3n + \frac{1}{2}})(1 - q^{3n + \frac{3}{2}})}{(1 - q^{n + \frac{1}{6}})(1 - q^{2n + \frac{1}{6}})} \mathcal{W}_n(q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{6}})$$

is a polynomial of degree 19 in q^n . Replacing \mathcal{W}_n by \mathcal{P} , we get the following formula

$$\left[\begin{matrix} q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{5}{6}} \\ q, q, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{3n^2 + \frac{n}{6}} \frac{(q^{\frac{1}{3}}, q^{\frac{1}{6}}; q)_{2n} (q^{\frac{1}{3}}; q^{\frac{1}{2}})_{2n}^2 (q^{\frac{1}{6}}; q^{\frac{1}{2}})_{2n+1}}{(q, q; q)_{2n+1} (q; q^{\frac{1}{2}})_{6n+3}} \mathcal{P}(q^n). \quad (20)$$

According to the factorial structure of the above summand, we define the hypergeometric term

$$\mathcal{T}_n := q^{\frac{3}{4}n^2 + \frac{n}{12}} \frac{(q^{\frac{1}{3}}, q^{\frac{1}{6}}; q)_n (q^{\frac{1}{3}}; q^{\frac{1}{2}})_n^2 (q^{\frac{1}{6}}; q^{\frac{1}{2}})_{n+1}}{(q, q; q)_n (q; q^{\frac{1}{2}})_{3n}} \mathcal{Q}(q^{\frac{n}{2}}),$$

where $\mathcal{Q}(y)$ is a polynomial to be determined. Taking into account that

$$\begin{aligned} \mathcal{T}_{2n} - \mathcal{T}_{2n+1} &= q^{3n^2 + \frac{n}{6}} \frac{(q^{\frac{1}{3}}, q^{\frac{1}{6}}; q)_{2n} (q^{\frac{1}{3}}; q^{\frac{1}{2}})_{2n}^2 (q^{\frac{1}{6}}; q^{\frac{1}{2}})_{2n+1}}{(q, q; q)_{2n+1} (q; q^{\frac{1}{2}})_{6n+3}} \\ &\times \left\{ \mathcal{Q}(q^n)(1 - q^{1+2n})^2(1 - q^{1+3n})(1 - q^{2+3n})(1 - q^{\frac{3}{2}+3n}) \right. \\ &\left. - q^{3n + \frac{5}{6}} \mathcal{Q}(q^{n + \frac{1}{2}})(1 - q^{\frac{1}{3}+n})^2(1 - q^{\frac{2}{3}+n})(1 - q^{\frac{1}{3}+2n})(1 - q^{\frac{1}{6}+2n}) \right\}, \end{aligned}$$

we may match the series in (20) with the following “bisection series”

$$\left[\begin{matrix} q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{5}{6}} \\ q, q, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} (-1)^n \mathcal{T}_n = \sum_{n=0}^{\infty} \left\{ \mathcal{T}_{2n} - \mathcal{T}_{2n+1} \right\}. \quad (21)$$

By equating the summand of (20) with the expression of “ $\mathcal{T}_{2n} - \mathcal{T}_{2n+1}$ ” and then canceling the common factors, we find the polynomial equation below

$$\begin{aligned} \mathcal{P}(y) &= \mathcal{Q}(y)(1 - qy^2)^2(1 - qy^3)(1 - q^2y^3)(1 - q^{\frac{3}{2}}y^3) \\ &\quad - q^{\frac{5}{6}}y^3 \mathcal{Q}(q^{\frac{1}{2}}y)(1 - q^{\frac{1}{3}}y)^2(1 - q^{\frac{2}{3}}y)(1 - q^{\frac{1}{3}}y^2)(1 - q^{\frac{1}{6}}y^2). \end{aligned} \quad (22)$$

Assuming further that $\mathcal{Q}(y)$ is of degree 6 and then resolving the linear system of equations for the coefficients of $\mathcal{Q}(y)$, we can explicitly determine, as done for $\mathcal{Q}(y)$ in the precedent example, the following polynomial expression

$$\begin{aligned} \mathcal{Q}(y) &= 1 + q^{\frac{1}{6}}y + q^{\frac{1}{6}}y^2 + q^{\frac{1}{3}}y^2 - q^{\frac{1}{6}}y^4 - q^{\frac{1}{3}}y^4 - q^{\frac{1}{3}}y^5 - q^{\frac{1}{2}}y^6 \\ &= q^{\frac{1}{6}}y^2(1 - y^2)(1 + q^{\frac{1}{6}}) + (1 + q^{\frac{1}{6}}y)(1 - q^{\frac{1}{3}}y^5). \end{aligned}$$

By making substitutions of $\mathcal{Q}(y)$ and \mathcal{J}_n into (21), we find, after some simplifications, the reduced formula as displayed in Example 42:

$$\begin{aligned} \left[\begin{matrix} q^{\frac{4}{3}}, q^{\frac{4}{3}}, q^{\frac{5}{6}} \\ q, q, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{3}}, q^{\frac{1}{6}}; q)_n (q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{2}{3}}; q^{\frac{1}{2}})_n}{(q, q; q)_n (q; q^{\frac{1}{2}})_{3n}} q^{\frac{n(9n+1)}{12}} \\ &\times \left\{ q^{n+\frac{1}{6}} \frac{1-q^n}{1-q^{\frac{1}{3}}} + \frac{(1+q^{\frac{n}{2}+\frac{1}{6}})(1-q^{\frac{5n}{2}+\frac{1}{3}})}{(1+q^{\frac{1}{6}})(1-q^{\frac{1}{3}})} \right\}. \quad \square \end{aligned}$$

For any given series, it is trivial to write down its bisection series counterpart. However, reversing this process is quite laborious due to the complexity to figure out exactly the parameter patterns of the target series. Notwithstanding the difficulty, we do succeed, with a help of *Mathematica*, in determining 20 reduced series. The classical series corresponding to the limits as $q \rightarrow 1$ with the convergence rate “ $\frac{-1}{27}$ ” have not been examined sufficiently, in spite of the fact they are π -related series investigated initially by Ramanujan [38] one century ago and explored extensively up to now, for example, in [1, 3, 6, 17–20, 22, 23, 28, 30–33, 40].

Example 35 ($a = q^{\frac{3}{2}}, b = c = d = q$) in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q, q, q, q \\ q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}; q^{\frac{1}{2}})_n^3 (q^{\frac{1}{2}}; q)_n}{(q^{\frac{3}{2}}; q)_n (q; q^{\frac{1}{2}})_{3n}} q^{\frac{3n(n+1)}{4}} \\ &\times (1+q^{n+\frac{1}{2}}) \left\{ 1 + \frac{q^{n+\frac{1}{2}}(1-q^{\frac{n}{2}+\frac{1}{2}})}{(1+q^{n+\frac{1}{2}})(1-q^{\frac{3}{2}n+1})} \right\}. \end{aligned}$$

When $q \rightarrow 1$, we recover the following known series (cf. [19, 22]):

$$\frac{\pi^2}{2} = \sum_{n=0}^{\infty} \left(\frac{-1}{27} \right)^n \left[\begin{matrix} 1, 1 \\ \frac{4}{3}, \frac{5}{3} \end{matrix} \right]_n \frac{5+7n}{1+2n}.$$

Example 36 ($a = q^{\frac{1}{2}}, b = c = d = q^{\frac{1}{3}}$) in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q, q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{1}{3}} \\ q^{\frac{1}{2}}, q^{\frac{1}{6}}, q^{\frac{1}{6}}, q^{\frac{1}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{3}}; q^{\frac{1}{3}})_n^3 (q^{\frac{1}{6}}; q)_n}{(q^{\frac{7}{6}}; q)_n (q^{\frac{1}{2}}; q^{\frac{1}{2}})_{3n}} q^{\frac{n(3n+1)}{4}} \\ &\times (1+q^{n+\frac{1}{6}}) \left\{ 1 + \frac{q^{n+\frac{1}{6}}(1-q^{\frac{n}{2}+\frac{1}{3}})}{(1+q^{n+\frac{1}{6}})(1-q^{\frac{3}{2}n+\frac{1}{2}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{2\pi^2}{\Gamma^3(\frac{2}{3})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27} \right)^n \left[\begin{matrix} \frac{2}{3}, \frac{2}{3}, \frac{1}{6} \\ 1, \frac{4}{3}, \frac{7}{6} \end{matrix} \right]_n \{8+21n\}.$$

Example 37 ($a = q^{\frac{5}{2}}, b = q^{\frac{2}{3}}, c = d = q^{\frac{5}{3}}$) in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q^2, q^{\frac{2}{3}}, q^{\frac{5}{3}}, q^{\frac{5}{3}} \\ q^{\frac{5}{2}}, q^{\frac{5}{6}}, q^{\frac{5}{6}}, q^{\frac{17}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{2}{3}}; q^{\frac{1}{2}})_n (q^{\frac{7}{6}}; q^{\frac{1}{2}})_n^2 (q^{-\frac{1}{6}}; q)_n^2}{(q^{\frac{5}{6}}; q)_n (q^{\frac{17}{6}}; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n}} q^{\frac{n(3n+7)}{4}} \\ &\times \frac{1-q^{\frac{1}{6}}}{1-q} \left\{ q^{n+\frac{5}{6}} + \frac{(1-q^{\frac{3}{2}n+2})}{(1-q^{n+\frac{5}{6}})} (1+q^{\frac{n}{2}+\frac{1}{6}})(1+q^{n+\frac{1}{3}}) \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{825\pi^2}{8\Gamma^3(\frac{1}{3})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{7}{3}, \frac{7}{3}, -\frac{1}{6}, -\frac{1}{6} \\ 1, \frac{5}{3}, \frac{11}{6}, \frac{17}{6} \end{matrix} \right]_n \{53 + 42n\}.$$

Example 38 ($a = q^{\frac{2}{3}}, b = q^{\frac{1}{6}}, c = d = q$) in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q, q, q^{\frac{1}{6}}, q^{\frac{1}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{-\frac{1}{3}}, q^{\frac{2}{3}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q, q^{-\frac{1}{3}}; q^{\frac{1}{2}})_n (q^{\frac{1}{2}}; q)_n (q^{\frac{4}{3}}; q)_n}{(q^{-\frac{1}{3}}; q)_n (q^{\frac{2}{3}}; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n} (1 + q^{\frac{1}{2}})} \\ &\quad \times q^{\frac{n(9n-1)}{12}} \left\{ q^{n+\frac{1}{2}} + \frac{(1 - q^{2n+1})(1 - q^{\frac{3}{2}n+\frac{1}{6}})}{(1 - q^{\frac{n}{2}+\frac{1}{2}})(1 - q^{n-\frac{1}{3}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{3\pi\Gamma^2(\frac{2}{3})}{\Gamma^2(\frac{1}{6})} = \sum_{n=1}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 1, \frac{1}{2}, -\frac{2}{3} \\ \frac{2}{3}, \frac{2}{3}, \frac{5}{3} \end{matrix} \right]_n n\{13 + 21n\}.$$

Example 39 ($a = q^{\frac{1}{3}}, b = q^{-\frac{1}{6}}, c = d = q$) in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q, q, q^{-\frac{1}{6}}, q^{-\frac{1}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{1}{3}}, q^{-\frac{2}{3}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q, q^{-\frac{2}{3}}; q^{\frac{1}{2}})_n (q^{\frac{1}{2}}; q)_n (q^{\frac{5}{3}}; q)_n}{(q^{-\frac{2}{3}}; q)_n (q^{\frac{1}{3}}; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n} (1 + q^{\frac{1}{2}})} \\ &\quad \times q^{\frac{n(9n-5)}{12}} \left\{ q^{n+\frac{1}{2}} + \frac{(1 - q^{2n+1})(1 - q^{\frac{3}{2}n-\frac{1}{6}})}{(1 - q^{\frac{n}{2}+\frac{1}{2}})(1 - q^{n-\frac{2}{3}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{\pi\Gamma^2(\frac{1}{3})}{12\Gamma^2(\frac{5}{6})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 1, \frac{1}{2}, -\frac{4}{3} \\ \frac{1}{3}, \frac{1}{3}, \frac{4}{3} \end{matrix} \right]_n \{21n^2 + 8n - 3\}.$$

Example 40 ($a = q^{\frac{4}{3}}, b = c = q, d = q^{\frac{5}{6}}$) in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q, q, q^{\frac{5}{6}}, q^{\frac{5}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{1}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q^{\frac{5}{3}}; q)_n (q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{3}}; q^{\frac{1}{2}})_n}{(q^{\frac{1}{3}}, q^{\frac{4}{3}}; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n}} q^{\frac{n(9n+7)}{12}} \\ &\quad \times \frac{1 - q^{\frac{2}{3}}}{1 - q} \left\{ q^{n+\frac{1}{2}} \frac{(1 + q^{\frac{1}{6}})}{(1 + q^{\frac{n}{2}+\frac{1}{3}})} + \frac{1 - q^{\frac{5}{2}n+\frac{7}{6}}}{1 - q^{n+\frac{1}{3}}} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{\pi\Gamma^2(\frac{1}{3})}{6\Gamma^2(\frac{5}{6})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 1, \frac{1}{2}, \frac{2}{3} \\ \frac{4}{3}, \frac{4}{3}, \frac{4}{3} \end{matrix} \right]_n \{3 + 7n\}.$$

Example 41 ($a = q^{\frac{5}{3}}, b = q^{\frac{7}{6}}, c = d = q$) in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q, q, q^{\frac{7}{6}}, q^{\frac{7}{6}} \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{2}{3}}, q^{\frac{5}{3}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q, q^{\frac{2}{3}}; q^{\frac{1}{2}})_n (q^{\frac{1}{2}}; q)_n (q^{\frac{1}{3}}; q)_n}{(q^{\frac{2}{3}}; q)_n (q^{\frac{5}{3}}; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n} (1 + q^{\frac{1}{2}})} \\ &\quad \times q^{\frac{n(9n+11)}{12}} \left\{ q^{n+\frac{1}{2}} + \frac{(1 - q^{2n+1})(1 - q^{\frac{3}{2}n+\frac{7}{6}})}{(1 - q^{\frac{n}{2}+\frac{1}{2}})(1 - q^{n+\frac{2}{3}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{48\pi\Gamma^2(\frac{2}{3})}{\Gamma^2(\frac{1}{6})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 1, \frac{1}{2}, \frac{1}{3} \\ \frac{5}{3}, \frac{5}{3}, \frac{5}{3} \end{matrix} \right]_n \{9 + 28n + 21n^2\}.$$

Example 42 ($a = b = c = q^{\frac{1}{3}}, d = q^{\frac{1}{6}}$ in Theorem 7).

$$\left[\begin{matrix} q^{\frac{4}{3}}, q^{\frac{4}{3}}, q^{\frac{5}{6}} \\ q, q, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{3}}, q^{\frac{1}{6}}; q)_n (q^{\frac{1}{3}}, q^{\frac{1}{3}}, q^{\frac{2}{3}}; q^{\frac{1}{2}})_n}{(q, q; q)_n (q; q^{\frac{1}{2}})_{3n}} q^{\frac{n(9n+1)}{12}} \\ \times \left\{ q^{n+\frac{1}{6}} \frac{1-q^n}{1-q^{\frac{1}{3}}} + \frac{(1+q^{\frac{n+\frac{1}{6}}}{2})(1-q^{\frac{5n+\frac{1}{3}}}{2})}{(1+q^{\frac{1}{6}})(1-q^{\frac{1}{3}})} \right\}.$$

When $q \rightarrow 1$, we recover the following elegant series (cf. [19, 20, 22]):

$$\frac{9\sqrt{3}}{2^{\frac{4}{3}}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, \frac{2}{3}, \frac{1}{6} \\ 1, 1, 1 \end{matrix} \right]_n \{2 + 21n\}.$$

Example 43 ($a = b = c = q^{\frac{2}{3}}, d = q^{\frac{5}{6}}$ in Theorem 7).

$$\left[\begin{matrix} q^{\frac{2}{3}}, q^{\frac{5}{3}}, q^{\frac{1}{6}} \\ q, q, q^{\frac{1}{2}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{6}}, q^{\frac{1}{6}}, q^{\frac{1}{3}}; q^{\frac{1}{2}})_n (q^{\frac{5}{3}}; q)_n (q^{\frac{5}{6}}; q)_n}{(q; q)_n^2 (q^{\frac{1}{2}}; q^{\frac{1}{2}})_{3n} (1+q^{\frac{n+\frac{1}{3}}{2}}+q^{n+\frac{2}{3}})} \\ \times q^{\frac{9n^2+5n+4}{12}} \left\{ \frac{1-q^{n-\frac{1}{6}}}{1-q^{\frac{n+\frac{1}{3}}{2}}} + \frac{q^{-\frac{n}{2}-\frac{1}{3}}(1-q^{2n+\frac{2}{3}})^2}{(1-q^{n+\frac{2}{3}})(1-q^{\frac{3n+\frac{1}{2}}{2}})} \right\}.$$

When $q \rightarrow 1$, we recover another elegant series (cf. [19, 20, 22]):

$$\frac{27\sqrt{3}}{2^{\frac{5}{3}}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, \frac{2}{3}, \frac{5}{6} \\ 1, 1, 1 \end{matrix} \right]_n \{5 + 42n\}.$$

Example 44 ($a = b = q^{\frac{5}{3}}, c = q^{\frac{2}{3}}, d = q^{\frac{5}{6}}$ in Theorem 7).

$$\left[\begin{matrix} q^{\frac{5}{3}}, q^{\frac{5}{3}}, q^{\frac{7}{6}}, q^{\frac{7}{6}} \\ q^2, q^2, q^{\frac{3}{2}}, q^{\frac{1}{6}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{2}{3}}, q^{\frac{5}{6}}, q^{\frac{7}{6}}; q^{\frac{1}{2}})_n (q^{-\frac{1}{6}}; q)_n (q^{\frac{2}{3}}; q)_n}{(q; q)_n (q^2; q)_n (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n} (1+q^{\frac{1}{3}})} q^{\frac{n(9n+11)}{12}} \\ \times \left\{ (1+q^{\frac{n+\frac{1}{6}}{2}})(1+q^{\frac{n+\frac{1}{3}}{2}})(1+q^{\frac{n+\frac{1}{2}}{2}})(1+q^{n+\frac{1}{3}}) - q^{\frac{n+1}{2}}(1+q^{\frac{3n+\frac{1}{3}}{2}}) \right\}.$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{81\sqrt{3}}{28 \cdot 2^{2/3}\pi} = \sum_{n=0}^{\infty} \left[\begin{matrix} \frac{2}{3}, \frac{7}{3}, -\frac{1}{6} \\ 1, 1, 2 \end{matrix} \right]_n \left(\frac{-1}{27}\right)^n.$$

Example 45 ($a = q^{\frac{4}{3}}, b = q^{\frac{1}{3}}, c = q^{\frac{4}{3}}, d = q^{\frac{7}{6}}$ in Theorem 7).

$$\left[\begin{matrix} q^{\frac{1}{3}}, q^{\frac{4}{3}}, q^{\frac{7}{6}}, q^{\frac{11}{6}} \\ q, q^2, q^{\frac{3}{2}}, q^{\frac{1}{6}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{2}{3}}, q^{\frac{4}{3}}, q^{-\frac{1}{6}}; q^{\frac{1}{2}})_n (q^{\frac{1}{3}}; q)_n (q^{\frac{7}{6}}; q)_n}{(q; q)_n^2 (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n}} \\ \times q^{\frac{n(9n+7)}{12}} \left\{ q^{n+\frac{1}{6}} + \frac{(1-q^{\frac{3n+\frac{5}{6}}{2}})1-q^{2n+\frac{4}{3}}}{(1-q^{\frac{n+\frac{5}{6}}{2}})(1-q^{n+1})} \right\}.$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{81\sqrt{3}}{2^{\frac{1}{3}}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, -\frac{1}{3}, \frac{7}{6} \\ 1, 1, 2 \end{matrix} \right]_n \{35 + 90n + 63n^2\}.$$

§4.2. The $\boxed{\delta = 1}$ case of Theorem 4 under the partition pattern (16) results in the formula below.

Theorem 8 (Nonterminating series identity).

$$\left[\begin{matrix} b, c, qd, qa^2/bcd \\ qa/b, qa/c, a/d, bcd/a \end{matrix} \middle| q \right]_{\infty} = 1 + \sum_{n=1}^{\infty} W_n(a, b, c, d) \left(\frac{q^{3n-2}a^3}{bcd} \right)^n \\ \times \frac{(b, c, d, qa/bc, qa/bd, a/cd; q)_n (bc/a, bd/a, cd/a; q)_{2n}}{(a/b, a/c, a/d; q)_{2n} (q, bcd/a; q)_{3n}}$$

where the weight function is given by

$$W_n(a, b, c, d) = \frac{(1-q^{3n})(1-b/a)(1-c/a)(1-q^{3n-1}bcd/a)(1-q^{4n-2}b)}{(1-cd/a)(1-1/d)(1-q^{n-1}b)(1-q^{2n-1}bc/a)(1-q^{2n-1}bd/a)} \\ \times \left\{ 1 - \frac{(1-q^{n-1}b)(1-q^{2n-1}bc/a)(1-q^{2n-1}bd/a)(1-q^n a/(cd))(1-q^{4n}d)}{(1-q^{3n})(1-q^{2n}b/a)(1-q^{2n}a/c)(1-q^{3n-1}bcd/a)(1-q^{4n-2}b)} \right. \\ \left. - \frac{(1-q^{3n-1})(1-q^{1-2n}c/a)(1-q^{2n-1}a/d)(1-q^{3n-2}bcd/a)(1-q^{4n-3}c)}{(1-q^{n-1}c)(1-q^{2n-2}bc/a)(1-q^n a/(bd))(1-q^{2n-1}cd/a)(1-q^{4n-2}b)} \right\}.$$

Following the same “reverse bisection approach” illustrated for Examples 40 and 42, we can derive from Theorem 8 further 9 identities.

Example 46 ($\boxed{a = q^{\frac{3}{2}}, b = c = d = q}$ in Theorem 8).

$$\left[\begin{matrix} q, q, q, q^2 \\ q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} = 1 + \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{3}{2}}; q^{\frac{1}{2}})_n (q^{\frac{1}{2}}; q)_n q^{\frac{(n+2)(3n+1)}{4}}}{(q^{\frac{5}{2}}; q)_n (q^2; q^{\frac{1}{2}})_{3n}} \\ \times \frac{(1-q^{\frac{1}{2}})(1-q^{2n+2})}{(1+q^{\frac{1}{2}})(1-q^{\frac{3}{2}})^2} \left\{ 1 + \frac{q^{n+\frac{3}{2}}(1-q^{\frac{n}{2}+\frac{1}{2}})(1-q^{n+\frac{1}{2}})}{(1-q^{\frac{3}{2}n+2})(1-q^{2n+2})} \right\}.$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$9\pi^2 = 89 + \sum_{n=1}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{2}, 1, 3 \\ \frac{5}{2}, \frac{5}{3}, \frac{7}{3} \end{matrix} \right]_n \{17 + 14n\}.$$

Example 47 ($\boxed{a = q^{\frac{1}{2}}, b = c = q^{\frac{2}{3}}, d = q^{-\frac{1}{3}}}$ in Theorem 8).

$$\left[\begin{matrix} q, q^{\frac{2}{3}}, q^{\frac{2}{3}}, q^{\frac{2}{3}} \\ q^{\frac{1}{2}}, q^{\frac{5}{6}}, q^{\frac{5}{6}}, q^{\frac{5}{6}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{6}}; q^{\frac{1}{2}})_n^2 (q^{\frac{2}{3}}; q^{\frac{1}{2}})_n (q^{-\frac{1}{6}}; q)_n}{(q^{\frac{5}{6}}; q)_n (q^{\frac{1}{2}}; q^{\frac{1}{2}})_{3n}} \\ \times q^{\frac{3n(n+1)}{4}} \left\{ 1 + \frac{(1-q^{\frac{3}{2}n})(1+q^{\frac{1}{6}-n})}{1-q^{\frac{n}{2}+\frac{1}{6}}} \right\}.$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{2\pi^2}{\Gamma^3(\frac{1}{3})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, \frac{1}{3} \\ 1, \frac{2}{3} \end{matrix} \right]_n \frac{1 + 21n}{1 - 6n}.$$

Example 48 ($a = q^{\frac{3}{2}}, b = q^{\frac{4}{3}}, c = d = q^{\frac{1}{3}}$ in Theorem 8).

$$\begin{aligned} \left[\begin{matrix} q^2, q^{\frac{1}{3}}, q^{\frac{4}{3}}, q^{\frac{4}{3}} \\ q^{\frac{1}{2}}, q^{\frac{1}{6}}, q^{\frac{7}{6}}, q^{\frac{13}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{3}}, q^{\frac{4}{3}}, q^{\frac{5}{6}}; q^{\frac{1}{2}})_n (q^{-\frac{5}{6}}; q)_n^2 q^{\frac{3n^2+3n+2}{4}}}{(q^{\frac{7}{6}}; q)_n (q^{\frac{13}{6}}; q)_n (q^{\frac{1}{2}}; q^{\frac{1}{2}})_{3n}} \\ &\times \frac{1 + q^{\frac{n}{2} - \frac{2}{3}}}{1 - q^{\frac{5}{6}}} \left\{ q^{n+\frac{1}{6}} + \frac{(1 - q^{2n+\frac{5}{6}})(1 + q^{n-\frac{1}{3}} + q^{\frac{n}{2}-\frac{1}{6}})}{(1 - q^{\frac{n}{2}+\frac{5}{6}})(1 + q^{\frac{n}{2}-\frac{2}{3}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{175\pi^2}{36\Gamma^3(\frac{2}{3})} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{5}{3}, \frac{5}{3}, -\frac{5}{6}, -\frac{5}{6} \\ 1, \frac{1}{3}, \frac{7}{6}, \frac{13}{6} \end{matrix} \right]_n \{25 + 42n\}.$$

Example 49 ($a = q^{\frac{4}{3}}, b = q^{\frac{5}{6}}, c = d = q$ in Theorem 8).

$$\begin{aligned} \left[\begin{matrix} q, q^2, q^{\frac{5}{6}}, q^{\frac{5}{6}} \\ q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{1}{3}}, q^{\frac{4}{3}} \end{matrix} \middle| q \right]_{\infty} &= 1 + \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{5}{6}}; q^{\frac{1}{2}})_n (q^{\frac{1}{2}}, q^{\frac{2}{3}}; q)_n q^{\frac{9n^2+19n+4}{12}}}{(q^{\frac{4}{3}}; q)_n^2 (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n}} \\ &\times \frac{(1 - q^{\frac{1}{2}})(1 + q^{\frac{n}{2}+\frac{1}{3}})}{(1 + q^{\frac{1}{2}})(1 - q^{2+\frac{3n}{2}})} \left\{ q^{\frac{n+1}{2}} + \frac{(1 - q^{2n+\frac{11}{6}})(1 + q^{n+\frac{3}{2}})}{(1 - q^{n+\frac{4}{3}})(1 + q^{n+1} + q^{\frac{n+1}{2}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{4\pi\Gamma^2(\frac{1}{3})}{\Gamma^2(\frac{5}{6})} = 71 + \sum_{n=1}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 3, \frac{1}{2}, \frac{2}{3} \\ \frac{4}{3}, \frac{7}{3}, \frac{7}{3} \end{matrix} \right]_n \{23 + 21n\}.$$

Example 50 ($a = q^{\frac{2}{3}}, b = q^{\frac{1}{6}}, c = d = q$ in Theorem 8).

$$\begin{aligned} \left[\begin{matrix} q, q^2, q^{\frac{1}{6}}, q^{\frac{1}{6}} \\ q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{-\frac{1}{3}}, q^{\frac{2}{3}} \end{matrix} \middle| q \right]_{\infty} &= 1 - \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{-\frac{1}{3}}; q^{\frac{1}{2}})_n (q^{\frac{1}{2}}, q^{\frac{7}{3}}; q)_n q^{\frac{n(9n+11)}{12}}}{(q^{\frac{2}{3}}; q)_n^2 (q^{\frac{3}{2}}; q^{\frac{1}{2}})_{3n}} \\ &\times \frac{(1 - q^{\frac{1}{2}})(1 + q^{\frac{2}{3}})(1 - q^{\frac{2}{3}})}{(1 + q^{\frac{1}{2}})(1 - q^{\frac{1}{3}})(1 - q^{2+\frac{3n}{2}})} \left\{ q^{\frac{n+1}{2}} + \frac{(1 - q^{2n+\frac{7}{6}})(1 + q^{n+\frac{3}{2}})}{(1 - q^{n+\frac{2}{3}})(1 + q^{n+1} + q^{\frac{n+1}{2}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{9\pi\Gamma^2(\frac{2}{3})}{\Gamma^2(\frac{1}{6})} = 1 + \sum_{n=1}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} 3, \frac{1}{2}, -\frac{2}{3} \\ \frac{2}{3}, \frac{5}{3}, \frac{5}{3} \end{matrix} \right]_n \{13 + 21n\}.$$

Example 51 ($a = b = c = q^{\frac{2}{3}}, d = q^{-\frac{1}{6}}$ in Theorem 8).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{2}{3}}, q^{\frac{2}{3}}, q^{\frac{7}{6}} \\ q, q, q^{\frac{1}{2}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{6}}; q^{\frac{1}{2}})_n^2 (q^{\frac{5}{6}}; q^{\frac{1}{2}})_n (q^{-\frac{1}{3}}, q^{\frac{5}{6}}; q)_n}{(q; q)_n^2 (q^{\frac{1}{2}}; q^{\frac{1}{2}})_{3n}} \\ &\times q^{\frac{n(9n+11)}{12}} \left\{ 1 + q^{\frac{1}{6}-n} \frac{(1 - q^{\frac{3}{2}n})(1 - q^{2n-\frac{1}{6}})}{(1 - q^{\frac{n}{2}+\frac{1}{3}})(1 - q^{n-\frac{1}{6}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{3\sqrt{3}}{\pi^{\frac{3}{4}}} = \sum_{n=0}^{\infty} \left(\frac{-1}{27}\right)^n \left[\begin{matrix} \frac{1}{3}, -\frac{1}{3}, -\frac{1}{6} \\ 1, 1, 1 \end{matrix} \right]_n \{1 - 63n^2\}.$$

Example 52 ($a = q^{\frac{1}{3}}, b = q^{\frac{1}{6}}, c = d = q^{\frac{1}{3}}$ in Theorem 8).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{4}{3}}, q^{\frac{4}{3}}, q^{\frac{11}{6}} \\ q, q, q^{\frac{5}{2}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{3}}, q^{\frac{4}{3}}, q^{\frac{1}{6}}; q^{\frac{1}{2}})_n (q^{\frac{4}{3}}, q^{\frac{1}{6}}; q)_n q^{\frac{n(9n+13)}{12}}}{(q; q)_n (q; q)_n (q^2; q^{\frac{1}{2}})_{3n}} \\ &\times \frac{1 - q^{\frac{3n}{2}+1}}{1 - q} \left\{ q^{\frac{n}{2}+\frac{1}{3}} + \frac{(1 - q^{2n+\frac{7}{6}})(1 + q^{n+\frac{7}{6}})}{(1 - q^{n+1})(1 + q^{\frac{n}{2}+\frac{1}{3}} + q^{n+\frac{2}{3}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{729\sqrt{3}}{20\sqrt[3]{2}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27} \right)^n \left[\begin{matrix} \frac{1}{3}, \frac{8}{3}, \frac{1}{6} \\ 1, 2, 2 \end{matrix} \right]_n \{16 + 21n\}.$$

Example 53 ($a = q^{\frac{2}{3}}, b = q^{\frac{5}{6}}, c = d = q^{\frac{2}{3}}$ in Theorem 8).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{5}{3}}, q^{\frac{5}{3}}, q^{\frac{11}{6}}, q^{\frac{13}{6}} \\ q, q^3, q^{\frac{5}{2}}, q^{\frac{5}{6}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{1}{6}}, q^{\frac{1}{3}}, q^{\frac{5}{6}}; q^{\frac{1}{2}})_n (q^{\frac{5}{3}}, q^{\frac{11}{6}}; q)_n}{(q; q)_n (q; q)_n (q^{\frac{5}{2}}; q^{\frac{1}{2}})_{3n}} \\ &\times q^{\frac{(n+2)(9n+5)}{12}} \left\{ 1 + \frac{(1 - q^{2n+\frac{8}{3}})(1 - q^{\frac{3}{2}n+\frac{7}{6}})}{q^{n+\frac{5}{6}}(1 - q^{\frac{n}{2}+\frac{7}{6}})(1 - q^{n+1})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{2187\sqrt{3}}{10\sqrt[3]{4}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27} \right)^n \left[\begin{matrix} \frac{1}{3}, \frac{2}{3}, \frac{11}{6} \\ 1, 2, 2 \end{matrix} \right]_n \{77 + 144n + 63n^2\}.$$

Example 54 ($a = q^{\frac{4}{3}}, b = q^{\frac{7}{6}}, c = q^{\frac{1}{3}}, d = q^{\frac{4}{3}}$ in Theorem 8).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{7}{3}}, q^{\frac{7}{3}}, q^{\frac{11}{6}} \\ q^2, q^2, q^{\frac{5}{2}} \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} (-1)^n \frac{(q^{\frac{2}{3}}, q^{-\frac{1}{6}}, q^{\frac{11}{6}}; q^{\frac{1}{2}})_n (q^{\frac{7}{3}}, q^{\frac{1}{6}}; q)_n q^{\frac{n(9n+19)}{12}}}{(q; q)_n (q^2; q)_n (q^{\frac{5}{2}}; q^{\frac{1}{2}})_{3n}} \\ &\times \left\{ \frac{q^{n+\frac{4}{3}}(1 - q^n)}{(1 - q^2)(1 + q^{\frac{n}{2}+\frac{2}{3}})} + \frac{(1 - q^{\frac{3}{2}n+\frac{11}{6}})(1 - q^{2n+\frac{13}{6}})}{(1 - q^2)(1 - q^{n+2})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{6561\sqrt{3}}{40\sqrt[3]{2}\pi} = \sum_{n=0}^{\infty} \left(\frac{-1}{27} \right)^n \left[\begin{matrix} \frac{4}{3}, \frac{11}{3}, \frac{1}{6} \\ 1, 2, 3 \end{matrix} \right]_n \frac{143 + 285n + 126n^2}{(1 - 3n)(2 + 3n)}.$$

From the above examples, one may get an impression that all the series obtained through triplicate inversions might be reducible, via the ‘‘reverse bisection method’’, to simpler ones. However, this is not true in general. Five counterexamples below demonstrate this fact.

Example 55 ($a = b = c = q^{\frac{1}{2}}, d = q^{\frac{1}{4}}$ in Theorem 7).

$$\begin{aligned} \left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{5}{2}} \\ q, q^2 \end{matrix} \middle| q \right]_{\infty} &= \sum_{n=0}^{\infty} q^{3n^2+\frac{n}{4}} \frac{(q^{\frac{1}{2}}; q)_n^2 (q^{\frac{3}{2}}; q)_n (q^{\frac{3}{4}}; q)_n^2 (q^{\frac{1}{4}}; q)_{n+1} (q^{\frac{1}{2}}; q)_{2n} (q^{\frac{1}{4}}; q)_{2n+1}}{(q; q)_{2n} (q^2; q)_{2n} (q; q)_{3n+1} (q^{\frac{3}{4}}; q)_{3n+1}} \\ &\times \frac{1 - q^{4n+\frac{3}{2}}}{1 - q^{\frac{3}{2}}} \left\{ q^{2n+\frac{1}{4}} + \frac{(1 - q^{2n+1})(1 - q^{3n+1})(1 - q^{3n+\frac{3}{4}})(1 - q^{4n+\frac{1}{4}})}{(1 - q^{n+\frac{1}{4}})(1 - q^{n+\frac{1}{2}})(1 - q^{2n+\frac{1}{4}})(1 - q^{4n+\frac{3}{2}})} \right. \\ &\quad \left. + \frac{q^{4n+\frac{5}{4}}(1 - q^{n+\frac{1}{2}})(1 - q^{n+\frac{3}{4}})(1 - q^{2n+\frac{1}{2}})(1 - q^{4n+\frac{5}{2}})}{(1 - q^{2n+1})(1 - q^{3n+\frac{7}{4}})(1 - q^{3n+2})(1 - q^{4n+\frac{3}{2}})} \right\}. \end{aligned}$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{16}{\pi} = \sum_{n=0}^{\infty} \frac{1}{729^n} \left[\frac{1}{2}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{5}{8}, \frac{9}{8}, \frac{11}{8} \right]_n \\ \times \left\{ 1 + \frac{12(3n+1)(16n+1)}{(8n+1)(8n+3)} + \frac{(4n+1)(4n+3)(8n+5)}{4(3n+2)(8n+3)(12n+7)} \right\}.$$

Example 56 ($a = q^{\frac{3}{2}}$, $b = q^{\frac{5}{6}}$, $c = d = q$) in Theorem 8).

$$\left[q, q^2, q^{\frac{5}{6}}, q^{\frac{7}{6}} \mid q \right]_{\infty} = 1 + \sum_{n=1}^{\infty} q^{\frac{9n^2+11n}{3}} \frac{(q^{-\frac{1}{2}}; q)_n (q; q)_n^2 (q^{\frac{2}{3}}; q)_n^2 (q^{-\frac{1}{6}}; q)_n (q^{-\frac{2}{3}}; q)_{2n}^2}{(q^{\frac{1}{2}}; q)_{2n} (q^{\frac{2}{3}}; q)_{2n} (q; q)_{3n} (q^{\frac{1}{3}}; q)_{3n}} \\ \times \frac{(1-q^{3n})(1-q^{\frac{7}{6}-4n})}{(1-q)(1+q^{\frac{1}{3}})(1-q^{-\frac{1}{6}})} \left\{ 1 - \frac{(1-q^{\frac{1}{2}-2n})(1-q^{3n-1})(1-q^{3n-\frac{2}{3}})(1-q^{4n-2})}{(1-q^n)(1-q^{n-\frac{1}{3}})(1-q^{2n-\frac{5}{3}})(1-q^{4n-\frac{7}{6}})} \right. \\ \left. + \frac{q^{2n+\frac{2}{3}}(1-q^{n-\frac{1}{2}})(1-q^{n-\frac{1}{6}})(1-q^{2n-\frac{2}{3}})^2(1-q^{4n+1})}{(1-q^{3n})(1-q^{2n+\frac{1}{2}})(1-q^{2n+\frac{2}{3}})(1-q^{3n+\frac{1}{3}})(1-q^{4n-\frac{7}{6}})} \right\}.$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{2\pi}{3\sqrt{3}} = 1 + \sum_{n=1}^{\infty} \frac{1}{729^n} \left[1, \frac{1}{2}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{1}{6}, \frac{1}{6} \right]_n \\ \times \left\{ 1 + \frac{3n(3n+1)(9n+1)(24n-7)}{2(2n-1)(3n-1)^2(6n-1)} + \frac{27(3n+1)(4n-1)(9n-2)(9n+1)}{(3n-1)^2(6n-5)(6n-1)} \right\}.$$

Example 57 ($a = b = c = d = q^{\frac{1}{2}}$) in Theorem 7).

$$\left[q^{\frac{1}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}}, q^{\frac{3}{2}} \mid q \right]_{\infty} = \sum_{n=0}^{\infty} q^{3n^2} \frac{(q^{\frac{1}{2}}; q)_n^6 (q^{\frac{3}{2}}; q)_{2n}^3 (1-q^{n+\frac{1}{2}})^2 (1-q^{4n+\frac{3}{2}})}{(q; q)_{2n}^3 (q^2; q)_{3n}^2 (1-q^{2n+1})(1-q^{2n+\frac{1}{2}})} \\ \times \left\{ q^{2n} + \frac{(1+q^{n+\frac{1}{2}})(1-q^{2n})(1-q^{3n+1})^2 (1-q^{4n+\frac{1}{2}})}{(1-q^{n+\frac{1}{2}})(1-q^{2n+\frac{1}{2}})^2 (1-q^{4n+\frac{3}{2}})} \right. \\ \left. + \frac{q^{4n+1}(1-q^{2n+\frac{1}{2}})(1-q^{2n+\frac{3}{2}})(1-q^{4n+\frac{5}{2}})}{(1+q^{n+\frac{1}{2}})^2 (1-q^{3n+2})^2 (1-q^{4n+\frac{3}{2}})} \right\}.$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{32}{\pi^2} = \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^6 (\frac{1}{2})_{2n}^3}{(2n)!^3 (3n)!^2} \times \frac{(1+2n)(1+4n)^2 (3+8n)}{(1+3n)^2} \\ \times \left\{ 1 + \frac{32n(1+3n)^2 (1+8n)}{(1+2n)(1+4n)^2 (3+8n)} + \frac{(1+4n)(3+4n)(5+8n)}{16(2+3n)^2 (3+8n)} \right\}.$$

Example 58 ($a = b = c = d = q^{\frac{1}{2}}$) in Theorem 8).

$$\left[q^{\frac{1}{2}}, q^{\frac{5}{2}}, q^{\frac{5}{2}}, q^{\frac{5}{2}} \mid q \right]_{\infty} = \sum_{n=0}^{\infty} q^{(n+1)(3n+1)} \frac{(q^{\frac{1}{2}}; q)_n^6 (q^{\frac{5}{2}}; q)_{2n}^3 (1-q^{n+\frac{1}{2}})^4 (1-q^{4n+\frac{5}{2}})}{(q^2; q)_{2n}^3 (q^3; q)_{3n}^2 (1-q)^3 (1-q^{2n+\frac{3}{2}})^2} \\ \times \left\{ 1 + \frac{q^{-1-2n}(1-q^{2n+1})^2 (1-q^{3n+2})^2 (1-q^{4n+\frac{3}{2}})}{(1-q^{n+\frac{1}{2}})^2 (1-q^{2n+\frac{1}{2}})(1-q^{2n+\frac{3}{2}})(1-q^{4n+\frac{5}{2}})} \right. \\ \left. + \frac{q^{2+2n}(1-q^{n+\frac{1}{2}})^2 (1-q^{2n+\frac{3}{2}})^2 (1-q^{4n+\frac{5}{2}})}{(1-q^{2n+2})^2 (1-q^{3n+3})^2 (1-q^{4n+\frac{5}{2}})} \right\}.$$

Its limiting case as $q \rightarrow 1$ results in the following classical series:

$$\frac{512}{\pi^2} = \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^6 (\frac{1}{2})_{2n}^3}{(2n)!^3 (3n)!^2} \times \frac{(1+2n)(1+4n)^3(3+4n)(5+8n)}{(1+3n)^2(2+3n)^2} \\ \times \left\{ 1 + \frac{16(2+3n)^2(3+8n)}{(1+4n)(3+4n)(5+8n)} + \frac{(1+2n)^2(3+4n)^2(9+8n)}{576(1+n)^4(5+8n)} \right\}.$$

These last two q -series have the same structure of factorial quotients, and differ to each other only by extra factors of rational functions in q^n . When $q \rightarrow 1$, their limiting series with the convergence rate “ $\frac{1}{729}$ ” are also significant, since they have not been recorded in the literature, even though there have been extensive research activities around infinite series for $1/\pi^2$, mainly made by Guillaera [30–32].

§4.3. Theorems 3 and 4 are so general that can be specified concretely to produce numerous summation formulae. For instance, it is well known that $n = \lfloor \frac{1+n}{3} \rfloor + \lfloor \frac{1+2n}{3} \rfloor$ holds for all $n \in \mathbb{N}_0$. Then it is not difficult to check that Jackson’s formula (13) is equivalent to the following one

$$\Omega_n(a; q^{\lfloor \frac{1+n}{3} \rfloor} b, c, q^{\lfloor \frac{1+2n}{3} \rfloor} d) = \left[\begin{matrix} qa, bd/a \\ qa/c, bcd/a \end{matrix} \middle| q \right]_n \left[\begin{matrix} qa/cd, bc/a \\ qa/d, b/a \end{matrix} \middle| q \right]_{\lfloor \frac{1+n}{3} \rfloor} \left[\begin{matrix} qa/bc, cd/a \\ qa/b, d/a \end{matrix} \middle| q \right]_{\lfloor \frac{1+2n}{3} \rfloor}$$

with its parameters subject to $qa^2 = bcde$. The corresponding partition pattern is determined by

$$\Lambda = 3: \quad \begin{matrix} \varepsilon_b = 1, & \varepsilon_c = 0, & \varepsilon_d = 1, & \varepsilon_e = 0; \\ \lambda_b = 1, & \lambda_c = 0, & \lambda_d = 2, & \lambda_e = 0. \end{matrix}$$

For Theorems 3 and 4, there exist six different summation formulae in accordance with $\delta = 0, 1, 2$. We limit to highlight only the $\delta = 0$ case of Theorem 3 together with two infinite series identities as examples. When $q \rightarrow 1$, the corresponding limiting series have the convergence rate “ $\frac{16}{729}$ ”.

Theorem 9 (Nonterminating series identity).

$$\left[\begin{matrix} b, c, qd, qa^2/bcd \\ qa/b, qa/c, a/d, bcd/a \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} W_n(a, b, c, d) \left(\frac{q^{2n} a^3}{bd^2} \right)^n \\ \times \frac{(qa/cd, b, bc/a; q)_n (qa/bc, qd, cd/a; q)_{2n} (bd/a; q)_{3n}}{(qa/d; q)_n (qa/b; q)_{2n} (q, qa/c, bcd/a; q)_{3n}},$$

where the weight function is given by

$$W_n(a, b, c, d) = \frac{(1 - q^{2n+1} a/bc)(1 - q^{2n} cd/a)(1 - q^{3n} bd/a)(1 - q^{4n+1} b)}{(1 - q^{3n+1})(1 - q^{3n+1} a/c)(1 - a/d)(1 - q^{3n} bcd/a)} \\ \times \left\{ q^n \frac{a}{d} + \frac{(1 - q^n a/d)(1 - q^{3n+1})(1 - q^{3n+1} a/c)(1 - q^{3n} bcd/a)(1 - q^{5n} d)}{(1 - q^{2n+1} a/bc)(1 - q^{2n} cd/a)(1 - q^{2n} d)(1 - q^{3n} bd/a)(1 - q^{4n+1} b)} \right. \\ \left. + \frac{q^{3n+1} a^2 (1 - q^n b)(1 - q^n bc/a)(1 - q^{n+1} a/cd)(1 - q^{3n+1} bd/a)(1 - q^{5n+3} d)}{bd(1 - q^{2n+1} a/b)(1 - q^{3n+2})(1 - q^{3n+2} a/c)(1 - q^{3n+1} bcd/a)(1 - q^{4n+1} b)} \right\}.$$

Example 59 ($a = b = c = d = q^{\frac{1}{2}}$ in Theorem 9).

$$\left[\begin{matrix} q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{5}{2}} \\ q, q, q, q \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{2n^2} \frac{(q^{\frac{1}{2}}; q)_{2n+1}^3 (q^{\frac{1}{2}}; q)_{2n+1}^3 (q^{\frac{3}{2}}; q)_{3n} (1 - q^{4n+\frac{3}{2}})}{(q; q)_n (q; q)_{2n} (q; q)_{3n+1} (1 - q^{3/2})} \\ \times \left\{ q^n + \frac{(1 - q^n)(1 - q^{3n+1})^3 (1 - q^{5n+\frac{1}{2}})}{(1 - q^{3n+\frac{1}{2}})(1 - q^{2n+\frac{1}{2}})^3 (1 - q^{4n+\frac{3}{2}})} \right. \\ \left. + q^{3n+1} \frac{(1 - q^{n+\frac{1}{2}})^3 (1 - q^{3n+\frac{3}{2}})(1 - q^{5n+\frac{7}{2}})}{(1 - q^{3n+2})^3 (1 - q^{2n+1})(1 - q^{4n+\frac{3}{2}})} \right\}.$$

This provides a q -analogue for the following series:

$$\frac{32}{\pi^2} = \sum_{n=0}^{\infty} (3+8n) \frac{(\frac{1}{2})_n^3 (\frac{3}{2})_n^3 (\frac{3}{2})_{3n}}{n!(2n)!(3n+1)!^3} \times \left\{ 1 + \frac{16n(1+3n)^3(1+10n)}{(1+4n)^3(1+6n)(3+8n)} + \frac{3(1+2n)^3(7+10n)}{16(2+3n)^3(3+8n)} \right\}.$$

Example 60 ($a = q^{\frac{3}{2}}, b = c = d = q$ in Theorem 9).

$$\left[\begin{matrix} q, q, q, q \\ q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{1}{2}}, q^{\frac{3}{2}} \end{matrix} \middle| q \right]_{\infty} = \sum_{n=0}^{\infty} q^{2n^2 + \frac{3n}{2}} \frac{(q, q^{\frac{1}{2}}, q^{\frac{1}{2}}; q)_n (q, q^{\frac{3}{2}}, q^{\frac{3}{2}}; q)_{2n} (1 - q^{2n+1})(1 - q^{4n+2})}{(q^{\frac{3}{2}}; q)_n (q^{\frac{3}{2}}; q)_{2n} (q, q^{\frac{3}{2}}; q)_{3n} (1 - q^{3n+1})(1 - q^{3n+\frac{3}{2}})^2} \times \left\{ q^{n+\frac{1}{2}} + \frac{(1 - q^{n+\frac{1}{2}})(1 - q^{3n+1})(1 - q^{3n+\frac{3}{2}})^2(1 - q^{5n+1})}{(1 - q^{2n+1})(1 - q^{2n+\frac{1}{2}})^2(1 - q^{3n+\frac{1}{2}})(1 - q^{4n+2})} + q^{3n+2} \frac{(1 - q^{n+\frac{1}{2}})^2(1 - q^{n+1})(1 - q^{3n+\frac{3}{2}})(1 - q^{5n+4})}{(1 - q^{3n+2})(1 - q^{2n+\frac{3}{2}})(1 - q^{3n+\frac{5}{2}})^2(1 - q^{4n+2})} \right\}.$$

This provides a q -analogue for another series:

$$\frac{9\pi^2}{16} = \sum_{n=0}^{\infty} \frac{(2n)!(4n+1)!}{(6n+1)!(1+2n)(1+3n)} \times \left\{ 1 + \frac{9(1+2n)(1+3n)(1+5n)}{2(1+4n)^2(1+6n)} + \frac{3(1+n)(1+2n)^2(4+5n)}{2(2+3n)(3+4n)(5+6n)^2} \right\}.$$

REFERENCES

- [1] V. Adamchik and S. Wagon, π : A 2000-year search changes direction, *Mathematica in Education and Research* 5:1 (1996), 11–19.
- [2] W. N. Bailey, *Generalized Hypergeometric Series*, Cambridge University Press, Cambridge, 1935.
- [3] N. D. Baruah, B. C. Berndt and H. H. Chan, *Ramanujan’s series for $1/\pi$: a survey*, *Amer. Math. Monthly* 116 (2009), 567–587.
- [4] J. M. Borwein and P. B. Borwein, *Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity*, Wiley, New York, 1987.
- [5] L. Carlitz, *Some inverse series relations*, *Duke Math. J.* 40 (1973), 893–901.
- [6] H. H. Chan and S. Cooper, *Rational analogues of Ramanujan’s series for $1/\pi$* , *Math. Proc. Cambridge Philos. Soc.* 153:2 (2012), 361–383.
- [7] X. J. Chen and W. Chu, *The Pfaff–Saalschütz theorem and terminating ${}_3F_2(4/3)$ -series*, *J. Math. Anal. Appl.* 380:2 (2011), 440–454.
- [8] X. J. Chen and W. Chu, *Multiply inverse series and identities of terminating hypergeometric series*, *Util. Math.* 90 (2013), 115–133.
- [9] X. J. Chen and W. Chu, *q -Analogues of Guillera’s two series for $\pi^{\pm 2}$ with convergence rate $\frac{27}{64}$* , *Int. J. Number Theory* 17:1 (2021), 71–90.
- [10] X. J. Chen and W. Chu, *Hidden q -analogues of Ramanujan-like π -series*, *Ramanujan J.* 54:3 (2021), 625–648.
- [11] W. Chu, *Inversion techniques and combinatorial identities*, *Boll. Unione Mat. Ital.* B-7 (1993, Serie VII), 737–760.
- [12] W. Chu, *Inversion techniques and combinatorial identities: A quick introduction to the hypergeometric evaluations*, *Math. Appl.* 283 (1994), 31–57.
- [13] W. Chu, *Inversion techniques and combinatorial identities: Strange evaluations of basic hypergeometric series*, *Compos. Math.* 91 (1994), 121–144.
- [14] W. Chu, *Inversion techniques and combinatorial identities: A unified treatment for the ${}_7F_6$ -series identities*, *Collect. Math.* 45:1 (1994), 13–43.
- [15] W. Chu, *Inversion techniques and combinatorial identities: Jackson’s q -analogue of the Dougall–Dixon theorem and the dual formulae*, *Compos. Math.* 95 (1995), 43–68.
- [16] W. Chu, *Inversion techniques and combinatorial identities: Balanced hypergeometric series*, *Rocky Mountain J. Math.* 32:2 (2002), 561–587.
- [17] W. Chu, *π -formulae implied by Dougall’s summation theorem for ${}_5F_4$ -series*, *Ramanujan J.* 26:2 (2011), 251–255.
- [18] W. Chu, *Dougall’s bilateral ${}_2H_2$ -series and Ramanujan-like π -formulae*, *Math. Comp.* 80:276 (2011), 2223–2251.

- [19] W. Chu, *q-series reciprocities and further π -formulae*, Kodai Math. J. 41:3 (2018), 512–530.
- [20] W. Chu, *Infinite series identities derived from the very well-poised Ω -sum*, Ramanujan J. 55:1 (2021), 239–270.
- [21] W. Chu and C. Y. Wang, *Bilateral inversions and terminating basic hypergeometric series identities*, Discrete Math. 309:12 (2009), 3888–3904 .
- [22] W. Chu and W. Zhang, *Accelerating Dougall's ${}_5F_4$ -sum and infinite series involving π* , Math. Comp. 83:285 (2014), 475–512.
- [23] L. Dembélé, A. Panchishkin, J. Voight, and W. Zudilin, *Special hypergeometric motives and their L-functions: Asai recognition*, Exp. Math. (2020); Available at <https://doi.org/10.1080/10586458.2020.1737990>.
- [24] J. Dougall, *On Vandermonde's theorem, and some more general expansions*, Proc. Edinb. Math. Soc. 25 (1907), 114–132.
- [25] G. Gasper and M. Rahman, *Basic Hypergeometric Series* (2nd ed.), Cambridge University Press, Cambridge, 2004.
- [26] I. Gessel and D. Stanton, *Strange evaluations of hypergeometric series*, SIAM J. Math. Anal. 13:2 (1982), 295–308.
- [27] I. Gessel and D. Stanton, *Applications of q-Lagrange inversion to basic hypergeometric series*, Trans. Amer. Math. Soc. 277:1 (1983), 173–201.
- [28] J. W. L. Glaisher, *On series for $1/\pi$ and $1/\pi^2$* , Quart. J. Math. 37 (1905), 173–198.
- [29] H. W. Gould and L. C. Hsu, *Some new inverse series relations*, Duke Math. J. 40 (1973), 885–891.
- [30] J. Guillera, *About a new kind of Ramanujan-type series*, Exp. Math. 12:4 (2003), 507–510.
- [31] J. Guillera, *Generators of some Ramanujan formulas*, Ramanujan J. 11:1 (2006), 41–48.
- [32] J. Guillera, *Hypergeometric identities for 10 extended Ramanujan-type series*, Ramanujan J. 15:2 (2008), 219–234.
- [33] J. Guillera, *Proofs of some Ramanujan series for $1/\pi$ using a program due to Zeilberger*, J. Difference Equ. Appl. 24:10 (2018), 1643–1648.
- [34] V. J. W. Guo, *q-Analogues of three Ramanujan-type formulas for $1/\pi$* , Ramanujan J. 52:1 (2020), 123–132.
- [35] V. J. W. Guo and J. C. Liu, *q-Analogues of two Ramanujan-type formulas for $1/\pi$* , J. Difference Equ. Appl. 24:8 (2018), 1368–1373.
- [36] V. J. W. Guo and W. Zudilin, *Ramanujan-type formulae for $1/\pi$: q-analogues*, Integral Transforms Spec. Funct. 29:7 (2018), 505–513.
- [37] F. H. Jackson, *Summation of q-hypergeometric series*, Messenger Math. 50 (1921), 101–112.
- [38] S. Ramanujan, *Modular equations and approximations to π* , Quart. J. Math. (Oxford) 45 (1914), 350–372.
- [39] K. R. Stromberg, *An Introduction to Classical Real Analysis*, Wadsworth, INC. Belmont, California, 1981.
- [40] W. Zudilin, *Ramanujan-type formulae for $1/\pi$: A second wind?*, Modular forms and string duality, 179–188; Amer. Math. Soc., Providence, RI, 2008.

SCHOOL OF STATISTICS
 QUFU NORMAL UNIVERSITY
 QUFU (SHANDONG), P. R. CHINA

Email address: upcxjchen@outlook.com

DEPARTMENT OF MATHEMATICS AND PHYSICS
 UNIVERSITY OF SALENTO (P. O. BOX 193)
 73100 LECCE, ITALY

Email address: chu.wenchang@unisalento.it