

LATTICE PATHS WITH INFINITELY MANY DOWN STEPS – THE NEGATIVE BOUNDARY MODEL

HELMUT PRODINGER

ABSTRACT. We consider a variation of Dyck paths, where additionally to steps $(1, 1)$ and $(1, -1)$ down-steps $(1, -j)$, for $j \geq 2$ are allowed. We give credits to Emeric Deutsch for that. The enumeration of such objects living in a strip is performed. Methods are the kernel method and techniques from linear algebra.

1. INTRODUCTION

Emeric Deutsch [1] had the idea to consider a variation of ordinary Dyck paths, by augmenting the usual up-steps and down-steps by one unit each, by down-steps of size $3, 5, 7, \dots$. This leads to ternary equations, as can be seen for instance from [3].

The present author started to investigate a related but simpler model of down-steps $1, 2, 3, 4, \dots$ and investigated it (named Deutsch paths in honour of Emeric Deutsch) in a series of papers, [2, 4, 5].

This paper is an further member of this series: The condition that (as with Dyck paths) the paths cannot enter negative territory, is relaxed, by introducing a negative boundary $-t$. Here are two recent publications about such a negative boundary: [8] and [7].

Instead of allowing negative altitudes, we think about the whole system shifted up by t units, and start at the point $(0, t)$ instead. This is much better for the generating functions that we are going to investigate. Eventually, the results can be re-interpreted as results about enumerations with respect to a negative boundary.

The setting with flexible initial level t and final level j allows us to consider the Deutsch paths also from left to right (they are not symmetric!), without any new computations.

The next sections achieves this, using the celebrated kernel-method, one of the tools that is dear to our heart [6].

In the following section, an additional upper bound is introduced, so that the Deutsch paths live now in a strip. The way to attack this is linear algebra. Once everything has been computed, one can relax the conditions and let lower/upper boundary go to $\mp\infty$.

2. GENERATING FUNCTIONS AND THE KERNEL METHOD

As discussed, we consider Deutsch paths starting at $(0, t)$ and ending at (n, j) , for $n, t, j \geq 0$. First we consider univariate generating functions $f_j(z)$, where z^n stays for

Date: August 31, 2021.

Key words and phrases. Lattice paths, boundaries, strip, kernel-method, Cramer's rule.

n steps done, and j is the final destination. The recursion is immediate:

$$f_j(z) = \llbracket t = j \rrbracket + z f_{j-1}(z) + z \sum_{k>j} f_k(z),$$

where $f_{-1}(z) = 0$. Next, we consider

$$F(z, u) := \sum_{j \geq 0} f_j(z) u^j,$$

and get

$$\begin{aligned} F(z, u) &= u^t + zuF(z, u) + z \sum_{j \geq 0} u^j \sum_{k>j} f_k(z) \\ &= u^t + zuF(z, u) + z \sum_{k>0} f_k(z) \sum_{0 \leq j < k} u^j \\ &= u^t + zuF(z, u) + z \sum_{k \geq 0} f_k(z) \frac{1 - u^k}{1 - u} \\ &= u^t + zuF(z, u) + \frac{z}{1 - u} [F(z, 1) - F(z, u)] \\ &= \frac{u^t(1 - u) + zF(z, 1)}{z - zu + zu^2 + 1 - u}. \end{aligned}$$

Since the critical value is around $u = 1$, we write the denominator as

$$z(u - 1)^2 + (u - 1)(z - 1) + z = z(u - 1 - r_1)(u - 1 - r_2),$$

with

$$r_1 = \frac{1 - z + \sqrt{1 - 2z - 3z^2}}{2z}, \quad r_2 = \frac{1 - z - \sqrt{1 - 2z - 3z^2}}{2z}.$$

The factor $(u - 1 - r_2)$ is bad, so the numerator must vanish for $[u^t(1 - u) + zF(z, 1)]|_{u=1+r_2}$, therefore

$$zF(z, 1) = (1 + r_2)^t r_2.$$

Furthermore

$$F(z, u) = \frac{\frac{u^t(1-u) + zF(z,1)}{u-r_2}}{z(u-r_1)}.$$

The expressions become prettier using the substitution $z = \frac{v}{1+v+v^2}$; then

$$r_1 = \frac{1}{v}, \quad r_2 = v.$$

It can be proved by induction (or computer algebra) that

$$\frac{u^t(1 - u) + v(1 + v)^t}{u - 1 - v} = -v \sum_{k=0}^{t-1} (1 + v)^{t-1-k} - u^t.$$

Furthermore

$$\frac{1}{z(u - 1 - r_1)} = -\frac{1}{z(1 + r_1)(1 - \frac{u}{1+r_1})},$$

and so

$$f_j(z) = [u^j]F(z, u) = [u^j] \left[v \sum_{k=0}^{t-1} (1+v)^{t-1-k} u^k + u^t \right] \sum_{\ell \geq 0} \frac{u^\ell}{z(1+r_1)^{\ell+1}}.$$

Of interest are two special cases: The case that was studied before [2] is $t = 0$:

$$f_j = \frac{(1+v+v^2)v^j}{(1+v)^{j+1}}.$$

The other special case is $j = 0$ for general t , as it may be interpreted as Deutsch paths read from right to left, starting at level 0 and ending at level $t \geq 1$ (for $t = 0$, the previous formula applies):

$$\begin{aligned} f_0(z) &= [u^0] \left[v \sum_{k=0}^{t-1} (1+v)^{t-1-k} u^k + u^t \right] \sum_{\ell \geq 0} \frac{u^\ell}{z(1+r_1)^{\ell+1}} \\ &= v(1+v)^{t-1} \frac{1}{z(1+r_1)} = v(1+v+v^2)(1+v)^{t-2}. \end{aligned}$$

The next section will present a simplification of the expression for $f_j(z)$, which could be obtained directly by distinguishing cases and summing some geometric series.

3. REFINED ANALYSIS: LOWER AND UPPER BOUNDARY

Now we consider Deutsch paths bounded from below by zero and bounded from above by $m-1$; they start at level t and end at level j after n steps. For that, we use generating functions $\varphi_j(z)$ (the quantity t is a silent parameter here). The recursions that are straight-forward are best organized in a matrix, as the following example shows.

$$\begin{pmatrix} 1 & -z & -z & -z & -z & -z & -z & -z \\ -z & 1 & -z & -z & -z & -z & -z & -z \\ 0 & -z & 1 & -z & -z & -z & -z & -z \\ 0 & 0 & -z & 1 & -z & -z & -z & -z \\ 0 & 0 & 0 & -z & 1 & -z & -z & -z \\ 0 & 0 & 0 & 0 & -z & 1 & -z & -z \\ 0 & 0 & 0 & 0 & 0 & -z & 1 & -z \\ 0 & 0 & 0 & 0 & 0 & 0 & -z & 1 \end{pmatrix} \begin{pmatrix} \varphi_0 \\ \varphi_1 \\ \varphi_2 \\ \varphi_3 \\ \varphi_4 \\ \varphi_5 \\ \varphi_6 \\ \varphi_7 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \Bigg\} t.$$

The goal is now to solve this system. For that the substitution $z = \frac{v}{1+v+v^2}$ is used throughout. The method is to use Cramer's rule, which means that the right-hand side has to replace various columns of the matrix, and determinants have to be computed. At the end, one has to divide by the determinant of the system.


Let D_m be the determinant of the matrix with m rows and columns. The recursion

$$(1+v+v^2)^2 m_{n+2} - (1+v+v^2)(1+v)^2 D_{m+1} + v(1+v)^2 D_m = 0$$

appeared already in [2] and is not difficult to derive and to solve:

$$D_m = \frac{(1+v)^{m-1}}{(1+v+v^2)^m} \frac{1-v^{m+2}}{1-v}.$$

To solve the system with Cramer's rule, we must compute a determinant of the following type,



where the various rows are replaced by the right-hand side. While it is not impossible to solve this recursion by hand, it is very easy to make mistakes, so it is best to employ a computer. Let $D(m; t, j)$ the determinant according to the drawing.

It is not unexpected that the results are different for $j < t$ resp. $j \geq t$. Here is what we found:

$$D(m; t, j) = \frac{(1+v)^{t-j-3+m}(1-v^{j+1})v(1-v^{m-t})}{(1-v)^2(1+v+v^2)^{m-1}}, \quad \text{for } j < t,$$

$$D(m; t, j) = \frac{v^{j-t}(1-v^{t+2})(1-v^{1-j+m})}{(1-v)^2(1+v+v^2)^{m-1}(1+v)^{j-t+3-m}}, \quad \text{for } j \geq t.$$

To solve the system, we have to divide by the determinant D_m , with the result

$$\varphi_j = \frac{D(m; t, j)}{D_m} = \frac{(1+v)^{t-j-2}(1-v^{j+1})v(1-v^{m-t})(1+v+v^2)}{(1-v)(1-v^{m+2})}, \quad \text{for } j < t,$$

$$\varphi_j = \frac{D(m; t, j)}{D_m} = \frac{v^{j-t}(1-v^{t+2})(1-v^{1-j+m})(1+v+v^2)}{(1-v)(1+v)^{j-t+2}(1-v^{m+2})}, \quad \text{for } j \geq t.$$

We found all this using Computer algebra. Some critical minds may argue that this is only experimental. One way of rectifying this would be to show that indeed the functions φ_j solve the system, which consists of summing various geometric series; again, a computer could be helpful for such an enterprise.

Of interest are also the limits for $m \rightarrow \infty$, i.e., no upper boundary:

$$\varphi_j = \lim_{m \rightarrow \infty} \frac{D(m; t, j)}{D_m} = \frac{(1+v)^{t-j-2}(1-v^{j+1})v(1+v+v^2)}{(1-v)}, \quad \text{for } j < t,$$

$$\varphi_j = \frac{v^{j-t}(1-v^{t+2})(1+v+v^2)}{(1-v)(1+v)^{j-t+2}}, \quad \text{for } j \geq t.$$

The special case $t = 0$ appeared already in the previous section:

$$\varphi_j = \frac{v^j(1+v+v^2)}{(1+v)^{j+1}}.$$

Likewise, for $t \geq 1$,

$$\varphi_0 = v(1+v+v^2)(1+v)^{t-2}.$$

In particular, the formulæ show that the expression from the previous section can be simplified in general, which could have been seen directly, of course.

Theorem 1. *The generating function of Deutsch path with lower boundary 0 and upper boundary $m - 1$, starting at $(0, t)$ and ending at (n, j) is given by*

$$\frac{(1+v)^{t-j-2}(1-v^{j+1})v(1-v^{m-t})(1+v+v^2)}{(1-v)(1-v^{m+2})}, \quad \text{for } j < t,$$

$$\frac{v^{j-t}(1-v^{t+2})(1-v^{1-j+m})(1+v+v^2)}{(1-v)(1+v)^{j-t+2}(1-v^{m+2})}, \quad \text{for } j \geq t,$$

with the substitution $z = \frac{v}{1+v+v^2}$.

By shifting everything down, we can interpret the results as Deutsch walks between boundaries $-t$ and $m - 1 - t$, starting at the origin $(0, 0)$ and ending at $(n, j - t)$.

Theorem 2. *The generating function of Deutsch path with lower boundary $-t$ and upper boundary h , starting at $(0, 0)$ and ending at (n, i) with $-t \leq i \leq h$ is given by*

$$\frac{(1+v)^{i-2}(1-v^{i+t+1})v(1-v^{h+1})(1+v+v^2)}{(1-v)(1-v^{h+t+3})}, \quad \text{for } i < 0,$$

$$\frac{v^i(1-v^{t+2})(1-v^{2-i+h})(1+v+v^2)}{(1-v)(1+v)^{i+2}(1-v^{h+t+3})}, \quad \text{for } i \geq 0.$$

It is possible to consider the limits $t \rightarrow \infty$ and/or $h \rightarrow \infty$ resulting in simplified formulae.

4. CONCLUSION

Various parameters could be worked out starting from the present findings. Currently, nothing to that effect has been done.

REFERENCES

- [1] Emeric Deutsch. Problem 10751. *American Mathematical Monthly*, 107 (2000); solution *American Mathematical Monthly*, 108 (2001).
- [2] Helmut Prodinger. Deutsch paths and their enumeration, *Open J. Discret. Appl. Math.* 2021, 4(1), 12–18.
- [3] Helmut Prodinger. Generating functions for a lattice path model introduced by Deutsch, *Special Matrices* 2021, 9: 217–225.
- [4] Helmut Prodinger. Enumeration of Deutsch paths by the adding-a-new-slice method and applications, *Matimyas Matematika, Journal of the Mathematical Society of the Philippines*, 44 (2021), 15–22.
- [5] Helmut Prodinger. Non-decreasing Deutsch paths, *The Fibonacci Quarterly* 59 (2021), 232–236.
- [6] Helmut Prodinger. *The Kernel Method: A Collection of Examples*, Séminaire Lotharingien de Combinatoire, B50f (2004), 19 pp.
- [7] Helmut Prodinger. *On k -Dyck paths with a negative boundary*, preprint.
- [8] Sarah J. Selkirk. *MSc-thesis: On a generalisation of k -Dyck paths*. Stellenbosch University. 2019.
- [9] Neil J. A. Sloane and The OEIS Foundation Inc. The on-line encyclopedia of integer sequences, 2021.

HELMUT PRODINGER, MATHEMATICS DEPARTMENT, STELLENBOSCH UNIVERSITY, 7602 STELLENBOSCH, SOUTH AFRICA.

Email address: hprodinger@sun.ac.za