

THE SPECTRUM OF PERIOD-DOUBLING HAMILTONIAN

QINGHUI LIU, YANHUI QU, AND XIAO YAO

ABSTRACT. In this paper, we show the following: the Hausdorff dimension of the spectrum of period-doubling Hamiltonian is bigger than $\log \alpha / \log 4$, where α is the Golden number; there exists a dense uncountable subset of the spectrum such that for each energy in this set, the related trace orbit is unbounded, which is in contrast with a recent result of Carvalho (Nonlinearity 33, 2020); we give a complete characterization for the structure of gaps and the gap labelling of the spectrum. All of these results are consequences of an intrinsic coding of the spectrum we construct in this paper.

1. INTRODUCTION

1.1. Background and motivation.

Since the pioneer works [17, 24], the discrete Schrödinger operators with substitutional potentials have been extensively studied. In these two works, a crucial concept—trace map, were introduced, which became a standard and powerful tool for studying the spectral properties of this class of operators. Using trace map technique, Bovier and Ghez [7] showed that if the potential is generated by primitive substitution, then, under certain technical condition, the spectrum of the related operator is of zero Lebesgue measure. Liu et. al. [22] removed this technical condition. This is the most general result in this direction. Lenz [19] also proved the same result by using cocycle dynamical technique, which is quite different from the previous two papers.

Three operators are heavily studied: the Fibonacci, Thue-Morse and period-doubling Hamiltonians. The potentials of these Hamiltonians are generated by Fibonacci, Thue-Morse and period-doubling substitutions, respectively. We note that Fibonacci substitution is invertible, while the Thue-Morse and period-doubling substitutions are not. The differences in substitution rules for these models have great impact on the corresponding dynamical systems induced by the trace polynomials. For the Fibonacci case, the induced dynamical system is a diffeomorphism on a cubic surface, which has a close relation with Hénon map on \mathbb{C}^2 . There has been much progress for understanding the dynamics of Hénon map in the past three decades. As a resonance, many powerful tools have been developed in the study of the spectrum of the Fibonacci Hamiltonian, see [8, 13, 14, 15, 16]. While, for the Thue-Morse and period-doubling models, the corresponding two dimensional dynamical systems are completely different: they are not diffeomorphisms at all, see for example [6, 21] and the present work. This makes them extremely hard to study, both from real and complex analytic point of views. To the best of our knowledge, there is no even a basic picture for the dynamics of this kind of maps in the community of complex dynamical system of higher dimension.

Fibonacci Hamiltonian was introduced in [17, 24], which soon became the most popular model for quasicrystal. Let $\{\hat{h}_n : n \geq 1\}$ be the trace polynomials related to Fibonacci Hamiltonian, Casdagli [10] defined the pseudospectrum of the operator:

$$(1) \quad \hat{B}_\infty := \left\{ E \in \mathbb{R} : \{\hat{h}_n(E) : n \geq 1\} \text{ is bounded} \right\}$$

and showed that \hat{B}_∞ is of zero Lebesgue measure. Sütő [27] showed that \hat{B}_∞ coincides with the spectrum $\hat{\sigma}_\lambda$ of Fibonacci Hamiltonian. Dynamically, \hat{B}_∞ can be redefined as follows(see for example [10]). Let

$$\hat{f}(x, y, z) := (y, z, yz - x)$$

be the *trace map* of Fibonacci Hamiltonian, then $\hat{f} : S_\lambda \rightarrow S_\lambda$ is a diffeomorphism, where $\lambda > 0$ is the coupling constant and

$$S_\lambda := \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 - xyz = 4 + \lambda^2\}.$$

Moreover, for any $n \geq 0$ and $E \in \mathbb{R}$, one has

$$\hat{f}(\hat{h}_n(E), \hat{h}_{n+1}(E), \hat{h}_{n+2}(E)) = (\hat{h}_{n+1}(E), \hat{h}_{n+2}(E), \hat{h}_{n+3}(E)).$$

For any E , write the initial condition as $\hat{\ell}(E) := (2, E, E - \lambda)$. Then

$$(2) \quad \hat{B}_\infty = \left\{ E \in \mathbb{R} : \{\hat{f}^n(\hat{\ell}(E)) : n \geq 1\} \text{ is bounded} \right\}.$$

These two papers suggest a way of characterizing the spectrum via boundedness of trace orbits, compare (1). Along this direction, Damanik [11] showed that if the potential is generated by an invertible primitive substitution over two letters, then the spectrum of the related operator still coincides with \hat{B}_∞ . In [10], Casdagli essentially coded the spectrum by a kind of subshift of finite type when the coupling $\lambda > 16$. Raymond [26] developed this approach to give coding for Sturmian Hamiltonian, which has Sturmian sequence as its potential. We note that Sturmian sequence is not necessarily generated by single substitution. Through this coding, he could compute the integrated density of states(IDS) of all the energies in an explicit way when $\lambda > 4$. His approach also gave a proof for gap opening and gap labelling of Sturmian Hamiltonian. The Hausdorff dimension of $\hat{\sigma}_\lambda$ has been well understood, see [26, 18, 8, 12, 13, 14, 16]. In particular the following property is shown in [12]:

$$\lim_{\lambda \rightarrow \infty} (\dim_H \hat{\sigma}_\lambda) \log \lambda = \log(1 + \sqrt{2}).$$

This implies that $\dim_H \hat{\sigma}_\lambda \rightarrow 0$ with the speed $1/\log \lambda$ when $\lambda \rightarrow \infty$. In the remarkable paper [16], almost all interesting spectral properties of Fibonacci Hamiltonian are established for all the coupling $\lambda > 0$.

Next we discuss Thue-Morse Hamiltonian(TMh). This operator has potential generated by Thue-Morse substitution, which is primitive but non invertible, as we have mentioned above. TMh was studied in many works at 1980's via trace maps, see for example [1, 2, 3, 4, 23]. In particular, the gap labelling and gap opening properties of the operator were studied in [2, 4, 23]. However, more detailed structure of the spectrum (such as a coding) is not known compare with the Fibonacci case. This is reflected by the fact that

the computation of the dimension of the spectrum is very difficult. Recently it was shown in [20] that for any $\lambda > 0$, the spectrum $\tilde{\sigma}_\lambda$ of TMH satisfies

$$\dim_H \tilde{\sigma}_\lambda \geq \frac{\log 2}{140 \log 2.1} = 0.00667 \dots$$

Also, a numerical estimation for the box dimension of the spectrum was provided in [25]. Indicated by the results of [10, 27], one may expect that the spectrum also coincides with \tilde{B}_∞ , where

$$(3) \quad \tilde{B}_\infty := \left\{ E \in \mathbb{R} : \{\tilde{h}_n(E) : n \geq 1\} \text{ is bounded} \right\},$$

where $\{\tilde{h}_n : n \geq 0\}$ are the trace polynomials related to TMH. This problem was studied in [21]. There, a Thue-Morse trace map $\tilde{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is defined as

$$\tilde{f}(x, y) := (x^2(y - 2) + 2, x^2y^2(y - 2) + 2).$$

Note that \tilde{f} is not a diffeomorphism. Indeed it is not even surjective and is generically 4-to-1. For any $E \in \mathbb{R}$ and any $n \geq 1$,

$$\tilde{f}(\tilde{h}_n(E), \tilde{h}_{n+1}(E)) = (\tilde{h}_{n+2}(E), \tilde{h}_{n+3}(E)).$$

Write the initial condition as $\tilde{\ell}(E) := (E^2 - \lambda^2 - 2, (E^2 - \lambda^2)^2 - 4E^2 + 2)$, then

$$\tilde{B}_\infty = \{E \in \mathbb{R} : \{\tilde{f}^n(\tilde{\ell}(E)) : n \geq 1\} \text{ is bounded}\}.$$

In [21], it was shown that $\tilde{\mathcal{E}}_\infty := \tilde{\sigma}_\lambda \setminus \tilde{B}_\infty$ is dense in $\tilde{\sigma}_\lambda$ and uncountable. That means there are many energies in the spectrum with unbounded trace orbit. We remark that, by a bit extra work, one indeed can show that $\tilde{B}_\infty \subset \tilde{\sigma}_\lambda$, see for example Theorem 1.3.

Now we come to period-doubling Hamiltonian(PDH), which is the object of this paper. The potential of PDH is generated by periodic-doubling substitution. Luck [23] first studied the trace maps of PDH and presented several conjectures about the labelling and opening of the gaps. Bellissard, Bovier and Ghez [6] confirmed these conjectures. Moreover, they showed that the spectrum is of zero Lebesgue measure. Like the Thue-Morse case, further information on the structure of the spectrum is missing. The first motivation of this paper is the following: try to find out some coding map similar to that of Fibonacci Hamiltonian, then use this coding to study the dimension, the IDS and gap labeling of the spectrum in a concrete manner, as Raymond did for Sturmian Hamiltonian([26]).

Most recently, Carvalho [9] showed that for PDH, the spectrum also coincides with the set of energies related to bounded trace orbit. However, our experience in [21] seems suggest that the opposite result should hold, just like the Thue-Morse case. Our second motivation is to understand the problem of existence of unbounded trace orbit completely.

1.2. Main results.

Now we set up the setting and state the main results of the paper.

1.2.1. Basic definitions.

Let η be the *period-doubling substitution*: $\eta(a) = ab$, $\eta(b) = aa$. It is seen that $\eta^{2^n}(a)$ is both a prefix and suffix of $\eta^{2^{(n+1)}}(a)$. Define a two-sided sequence ξ as

$$\xi := \lim_{n \rightarrow \infty} \eta^{2^n}(a) | \eta^{2^n}(a) = \cdots \xi(-2)\xi(-1) | \xi(0)\xi(1) \cdots .$$

Define the *period-doubling potential* $V_\xi = (V_\xi(n))_{n \in \mathbb{Z}}$ by, $V_\xi(n) = 1$ if $\xi(n) = a$ and $V_\xi(n) = -1$ if $\xi(n) = b$ for $n \in \mathbb{Z}$. Take $\lambda \in \mathbb{R}$ and $\lambda \neq 0$. Let $H_{\lambda V_\xi}$ be the discrete Schrödinger operator acting on $\ell^2(\mathbb{Z})$ with potential λV_ξ , i.e., for any $n \in \mathbb{Z}$,

$$(H_{\lambda V_\xi} \psi)_n = \psi_{n+1} + \psi_{n-1} + \lambda V_\xi(n) \psi_n, \quad \forall \psi \in \ell^2(\mathbb{Z}).$$

We call $H_{\lambda V_\xi}$ the *period-doubling Hamiltonian* (PDH). Denote by $\sigma(H_{\lambda V_\xi})$ the spectrum of $H_{\lambda V_\xi}$. From now on, we only consider PDH, so we will write

$$(4) \quad H_\lambda := H_{\lambda V_\xi} \quad \text{and} \quad \sigma_\lambda := \sigma(H_{\lambda V_\xi}).$$

It is a general fact that $\sigma_{-\lambda} = -\sigma_\lambda$. So in the following, without loss of generality, we always assume $\lambda > 0$.

Given $E \in \mathbb{R}$, define an anti-homomorphism $\tau : \text{FG}(a, b) \rightarrow \text{SL}(2, \mathbb{R})$ as

$$\tau(a) := \begin{bmatrix} E - \lambda & -1 \\ 1 & 0 \end{bmatrix} \quad \text{and} \quad \tau(b) := \begin{bmatrix} E + \lambda & -1 \\ 1 & 0 \end{bmatrix}$$

and $\tau(a_1 \cdots a_n) := \tau(a_n) \cdots \tau(a_1)$. Define

$$(5) \quad h_n(E) := \text{tr}(\tau(\eta^n(a))).$$

h_n is called the n -th *trace polynomial* related to PDH. It is seen that $\deg(h_n) = 2^n$. For each $E \in \mathbb{R}$, we define

$$(6) \quad \mathcal{O}(E) := \{h_n(E) : n \geq 0\}$$

and call $\mathcal{O}(E)$ the *trace orbit* of E . Define

$$(7) \quad B_\infty := \{E \in \mathbb{R} : \mathcal{O}(E) \text{ is bounded}\}.$$

We say $E \in \sigma_\lambda$ is of ∞ -*type* if $\mathcal{O}(E)$ is unbounded. Define

$$(8) \quad \mathcal{E}_\infty := \{E \in \sigma_\lambda : E \text{ is of } \infty\text{-type}\}.$$

1.2.2. Lower bound for the Hausdorff dimension of the spectrum.

At first we have the following estimate for the dimension of the spectrum:

Theorem 1.1. *Let $\alpha = (1 + \sqrt{5})/2$. Then for any $\lambda > 0$,*

$$\dim_H \sigma_\lambda \geq \frac{\log \alpha}{\log 4} = 0.34712 \cdots$$

Remark 1.2. 1) In [25], a numerical estimation for $\dim_H \sigma_\lambda$ is given. The numerical lower bound of $\dim_H \sigma_\lambda$ is around 0.75.

2) Indeed, $\log \alpha$ is the entropy of certain subshift of finite type and $\log 4$ can be viewed as an upper bound of the Lyapunov exponents, see Remark 5.9 for explanation. So the lower bound is kind of Young's dimension formula.

3) As we have mentioned, in [20] we obtained a lower bound $0.00667 \dots$ for the Hausdorff dimension of the spectrum of TMH. Let us make a comparison on the related methods. In [20], we showed that, after a suitable renormalization, the trace polynomials $\{\tilde{h}_n : n \geq 1\}$ are exponentially close to the model family $\{2 \cos 2^n E : n \geq 1\}$. Using this closeness, we can construct a nested covering structure such that the ratios of lengths between the son intervals and the father intervals are bounded from below. By this, we can show that the dimension of the limit set has a universal lower bound (which is $0.00667 \dots$, very small and far from optimal). Also by the construction, the limit set is a subset of the spectrum, so we obtain the dimension lower bound for the spectrum. It is desirable to adapt the method to the PDH case. It is true that one can still find a model family $\{2 \cos 2^n \sqrt{E} : n \geq 1\}$ such that after a suitable renormalization, the trace polynomials $\{h_n : n \geq 1\}$ of PDH are close to them. However it is much harder to show the exponential convergence. It is even harder to construct a good Cantor subset of the spectrum if it was possible. Here we take a totally different approach. Due to the explicit coding of the spectrum (see Theorem 1.5), we can construct a nested covering structure with nice separation property. This structure determines a limit set, which is a subset of the spectrum. Every energy in this subset has a bounded trace orbit and the bound is 2. This bound together with the recurrence relation of the trace polynomials imply that the lengths of the bands in level n are bigger than $C4^{-n}$. Now it is not hard to show that the dimension of the limit set has a good lower bound.

4) We also note that it is doable to apply the method in the present paper to the TMH to give a much better lower bound for the Hausdorff dimension of the spectrum.

1.2.3. Existence of unbounded trace orbits.

Theorem 1.3. $Z \subset B_\infty \subset \sigma_\lambda$ and $\mathcal{E}_\infty = \sigma_\lambda \setminus B_\infty$. Both B_∞ and \mathcal{E}_∞ are dense in σ_λ and uncountable.

Remark 1.4. 1) As we have mentioned, Carvalho [9] claimed that \mathcal{E}_∞ is empty. Here we show the contrary. We will see soon that ∞ -type energies are closely related to the gap edges of the spectrum.

2) In [21], for TMH, we showed that $\tilde{\mathcal{E}}_\infty$ is dense in $\tilde{\sigma}_\lambda$ and uncountable. But we did not show that \tilde{B}_∞ is a subset of $\tilde{\sigma}_\lambda$. Here, we show that $\{B_\infty, \mathcal{E}_\infty\}$ form a partition of σ_λ .

3) Similar to the TMH case, we will construct a period-doubling trace map $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ (see (72)) such that for any $E \in \mathbb{R}$ and any $n \geq 0$,

$$f(h_n(E), h_{n+1}(E)) = (h_{n+2}(E), h_{n+3}(E)).$$

If we write the initial condition as $\ell(E) := (E - \lambda, E^2 - \lambda^2 - 2)$, then

$$B_\infty = \{E \in \mathbb{R} : \{f^n(\ell(E)) : n \geq 1\} \text{ is bounded}\}.$$

We will see that the dynamical properties of f play an essential role for showing the injectivity of the coding map.

Both results are consequences of an intrinsic coding of the spectrum. So we turn to this coding map.

1.2.4. *Coding of the spectrum.*

We will code the spectrum σ_λ via a subshift of finite type. At first we define such a subshift. We will explain the idea behind these definitions in Sec. 1.3.3.

Define an *alphabet* \mathcal{A} as

$$(9) \quad \mathcal{A} := \{0_e, 0_o, 1_e, 1_o, 2_e, 2_o, 3_{el}, 3_{er}, 3_{ol}, 3_{or}\}.$$

Define the *admissible rules* as

$$(10) \quad \begin{aligned} 0_e &\rightarrow \{3_{ol}, 1_o, 3_{or}\}; & 0_o &\rightarrow \{3_{el}, 1_e, 3_{er}\}; & 1_e &\rightarrow \{3_{ol}, 2_o\}; & 1_o &\rightarrow \{2_e, 3_{er}\}; \\ 2_e &\rightarrow \{1_o, 3_{or}, 2_o\}; & 2_o &\rightarrow \{2_e, 3_{el}, 1_e\}; & 3_{el}, 3_{er} &\rightarrow 0_o; & 3_{ol}, 3_{or} &\rightarrow 0_e. \end{aligned}$$

See Figure 1 for the related directed graph \mathbb{G} (we will explain the extra information such as the colors and the labels of the edges, the dashed red square etc. later). Define the *adjacency matrix* $A = [a_{\alpha\beta}]$ of \mathbb{G} as

$$\begin{cases} a_{\alpha\beta} = 1, & \text{if } \alpha \rightarrow \beta; \\ a_{\alpha\beta} = 0, & \text{otherwise.} \end{cases}$$

It is seen that A is irreducible but not aperiodic, indeed it has period 2. Consequently, \mathbb{G} is connected. Let Ω_A be the related subshift of finite type:

$$\Omega_A := \{\omega \in \mathcal{A}^{\mathbb{N}} : a_{\omega_j \omega_{j+1}} = 1, j \geq 0\}.$$




FIGURE 1. The directed graph \mathbb{G} related to the subshift

Define a metric d on Ω_A as $d(\omega, \hat{\omega}) := 2^{-|\omega \wedge \hat{\omega}|}$. It is standard to check that d is indeed a metric and (Ω_A, d) is compact. Define the symbolic space

$$(11) \quad \Omega_\infty := \{\omega \in \Omega_A : \omega_0 \in \{3_{el}, 0_e\}\}.$$

Ω_∞ is a union of two cylinders: $\Omega_\infty = [3_{el}] \cup [0_e]$, so it is compact.

Later we will define a total order \preceq on Ω_∞ , which is essentially induced by the ordinary order \leq on \mathbb{R} . (We will explain this in Sec. 1.3.5 and Sec. 4).

Theorem 1.5. *There exists an order-preserving homeomorphism $\pi : (\Omega_\infty, \preceq) \rightarrow (\sigma_\lambda, \leq)$, where \leq is the standard order on \mathbb{R} .*

We call π the *coding map* of σ_λ .

Remark 1.6. In some sense, we construct a “Markov partition” of the spectrum σ_λ through the coding given in Theorem 1.5. The idea of constructing combinatoric models (Markov partition, Young tower, Yoccoz puzzle, inducing scheme, etc.) play a central role for understanding the hyperbolicity of the dynamical system, which can date back to Bowen. For the spectrum of discrete Schrödinger operator, the idea of constructing certain Markov partition can date back to Casdagli for Fibonacci Hamiltonian ([10]) and Raymond for Sturmian Hamiltonian ([26]). Some recent progress have been made by Cantat ([8]) and Damanik, Gorodetski, Yessen ([13, 14, 15, 16]), which heavily rely on the dynamics of \hat{f} on the surface S_λ .

To some extent, all the constructions of the combinatoric models relies on the bounded distortion (complex bound) arguments. For the PDH model, the existence of the energies with unbounded trace orbits is the main obstruction for the bounded distortion phenomenon, which does not happen in the Fibonacci model. This makes the construction of combinatoric model extremely hard, we need be very careful in discussion of various combinatorics of the zeros, bands, ends of the bands and related issues.

Remark 1.7. Let us make a comparison with [26] on the methods of constructing codings. For both classes of Hamiltonians, let σ_n be the n -th periodic approximation of the spectrum $\sigma(H_\lambda)$, one can show that $\{\sigma_n \cup \sigma_{n+1} : n \geq 1\}$ is a nested sequence of coverings of the spectrum. By choosing an optimal covering in each level, it is not hard to construct a symbolic space Ω and a surjective coding $\pi : \Omega \rightarrow \sigma(H_\lambda)$. Usually it is much harder to get a bijective coding. To achieve this, one need some disjoint condition for bands in σ_n and σ_{n+1} . That is why in [26], the technical assumption $\lambda > 4$ is needed. We remark that when $\lambda \downarrow 0$, the overlapping of σ_n and σ_{n+1} is unavoidable. This presents an obstruction for a bijective coding. In our case, many bands in the n -th optimal covering do overlap with each other (at least for small λ). One can see Figure 3 (b) for some intuition of this situation. A big challenge is to show that as n grows, this kind of overlapping will disappear. By a detailed analysis on the band configurations of $\sigma_n \cup \sigma_{n+1} \cup \sigma_{n+2}$, finally we succeed to show that all the overlapping disappear and gaps of $\sigma(H_\lambda)$ show up. Surprisingly, we find that the edges of this kind of gaps are related to unbounded trace orbits. Indeed, the bijectivity of π follows from the study of the unbounded trace orbits.

1.2.5. Gaps, zeros of trace polynomials and ∞ -type energies.

With this coding map in hand, we can give a complete description for the gaps of the spectrum. We will see that the gap edges, zeros of trace polynomials and ∞ -type energies are related in a delicate but beautiful way.

Define the set of zeros of trace polynomials as

$$(12) \quad \mathcal{Z}^o := \bigcup_{n \text{ odd}} \{h_n = 0\}; \quad \mathcal{Z}^e := \bigcup_{n \text{ even}} \{h_n = 0\}; \quad \mathcal{Z} := \mathcal{Z}^o \cup \mathcal{Z}^e.$$

We will see that, if an energy is a gap edge, then its coding is eventually 2-periodic. Figure 1 suggests that there are 7 types of eventually 2-periodic codings:

$$(13) \quad \begin{cases} \mathcal{E}_l^o & := \{\omega \in \Omega_\infty : \omega = w(2_o1_e)^\infty \text{ for some } w \in \Omega_*\} \\ \mathcal{E}_r^o & := \{\omega \in \Omega_\infty : \omega = w(0_o3_{el})^\infty \text{ for some } w \in \Omega_*\} \\ \mathcal{E}_l^e & := \{\omega \in \Omega_\infty : \omega = w(0_e3_{or})^\infty \text{ for some } w \in \Omega_*\} \\ \mathcal{E}_r^e & := \{\omega \in \Omega_\infty : \omega = w(2_e1_o)^\infty \text{ for some } w \in \Omega_*\} \\ \tilde{\mathcal{E}}_l & := \{\omega \in \Omega_\infty : \omega = w(0_o3_{er})^\infty \text{ for some } w \in \Omega_*\} \\ \tilde{\mathcal{E}}_r & := \{\omega \in \Omega_\infty : \omega = w(0_e3_{ol})^\infty \text{ for some } w \in \Omega_*\} \\ \mathcal{F} & := \{\omega \in \Omega_\infty : \omega = w(2_e2_o)^\infty \text{ for some } w \in \Omega_*\} \end{cases}$$

where Ω_* is the set of finite admissible words. Define two special codings as follows:

$$(14) \quad \omega_* := 3_{el}(0_o3_{el})^\infty; \quad \omega^* := (0_e3_{or})^\infty.$$

Denote the set of gaps of σ_λ by \mathcal{G} . We have the following complete description on the structure of \mathcal{G} .

Theorem 1.8. *With the notations above, we have:*

i) $\min \sigma_\lambda = \pi(\omega_*)$ and $\max \sigma_\lambda = \pi(\omega^*)$.

ii) (Type-I gaps): *There exists a bijection $\iota^o : \pi(\mathcal{E}_l^o) \rightarrow \pi(\mathcal{E}_r^o \setminus \{\omega_*\})$ such that $E < \iota^o(E)$ for any $E \in \pi(\mathcal{E}_l^o)$ and*

$$\mathcal{G}_I^o := \{(E, \iota^o(E)) : E \in \pi(\mathcal{E}_l^o)\} \subset \mathcal{G}.$$

There exists a bijection $\iota^e : \pi(\mathcal{E}_r^e) \rightarrow \pi(\mathcal{E}_l^e \setminus \{\omega^\})$ such that $\iota^e(E) < E$ for any $E \in \pi(\mathcal{E}_r^e)$ and*

$$\mathcal{G}_I^e := \{(\iota^e(E), E) : E \in \pi(\mathcal{E}_r^e)\} \subset \mathcal{G}.$$

iii) (Type-II gaps): *There exists a bijection $\iota : \pi(\tilde{\mathcal{E}}_l) \rightarrow \pi(\tilde{\mathcal{E}}_r)$ such that $E < \iota(E)$ for any $E \in \pi(\tilde{\mathcal{E}}_l)$ and*

$$\mathcal{G}_{II} := \{(E, \iota(E)) : E \in \pi(\tilde{\mathcal{E}}_l)\} \subset \mathcal{G}.$$

iv) $\mathcal{G} = \mathcal{G}_I \cup \mathcal{G}_{II}$, where $\mathcal{G}_I := \mathcal{G}_I^o \cup \mathcal{G}_I^e$.

v) (Coding of zeros): $\pi(\mathcal{E}_l^o) = \mathcal{Z}^o$ and $\pi(\mathcal{E}_r^e) = \mathcal{Z}^e$.

vi) (∞ -type energies): *For any $\mathcal{E} \in \{\mathcal{E}_l^e, \mathcal{E}_r^o, \tilde{\mathcal{E}}_l, \tilde{\mathcal{E}}_r\}$, we have*

$$\pi(\mathcal{E}) \subset \mathcal{E}_\infty \text{ and } \pi(\mathcal{E}) \text{ is dense in } \sigma_\lambda.$$

We say that the energy $E \in \sigma_\lambda$ is of *type ∞_I (type ∞_{II})* if $E \in \pi(\mathcal{E}_l^e \cup \mathcal{E}_r^o)$ ($\pi(\tilde{\mathcal{E}}_l \cup \tilde{\mathcal{E}}_r)$).

Remark 1.9. 1) For a gap in \mathcal{G}_I , one of its end point is a zero of trace polynomials, another endpoint if of type ∞_I . For a gap in \mathcal{G}_{II} , both endpoints of it are of type ∞_{II} . So the codings of all the gap edges are eventually 2-periodic. However, by ii)-iv), $E \in \pi(\mathcal{F})$ is not an edge of any gap. See Theorem 1.11 and Remark 1.12 for the special role played by \mathcal{F} .

2) $\mathcal{G}_I^e, \mathcal{G}_I^o, \mathcal{G}_{II}$ are corresponding to the green, blue and pink edges of \mathbb{G} , respectively, see Figure 1. We also remark that \mathcal{G}_I are exact the gaps in [6] Theorem 4 which open linearly, while \mathcal{G}_{II} are exact the gaps which open exponentially.

1.2.6. Integrated density of states and Gap labelling of the spectrum.

Now we compute the IDS for every energy in the spectrum. The labels on the edges of the graph \mathbb{G} play an essential role in this computation.

Define the infinite binary tree Σ_∞ and the evaluation map $\varepsilon : \Sigma_\infty \rightarrow [0, 1]$ as

$$(15) \quad \Sigma_\infty := \{0, 1\}^\infty; \quad \varepsilon(\sigma) = \sum_{n \geq 1} \frac{\sigma_n}{2^n}.$$

Define a map $\Pi : \Omega_\infty \rightarrow \Sigma_\infty$ as follows: given $\omega \in \Omega_\infty$, follow the infinite path $\omega = \omega_0 \omega_1 \omega_2 \cdots$ in \mathbb{G} , record the label w_i on the edge $\omega_{i-1} \omega_i$ and concatenate them to get

$$(16) \quad \Pi(\omega) := i(\omega) w_1 w_2 w_3 \cdots \in \Sigma_\infty,$$

where $i(\omega) = \emptyset$ if $\omega_0 = 0_e$ and $i(\omega) = 0$ if $\omega_0 = 3_{el}$. Here we use the convention that $\emptyset w = w \emptyset = w$ for a finite word w .

Denote by $N(E)$ the IDS of $E \in \sigma_\lambda$. We have the following:

Theorem 1.10. $\Pi : \Omega_\infty \rightarrow \Sigma_\infty$ is surjective and for any $\omega \in \Omega_\infty$,

$$N(\pi(\omega)) = \varepsilon(\Pi(\omega)).$$

In other words, we have the following commutative diagram:

$$\begin{array}{ccc} \Omega_\infty & \xrightarrow{\pi} & \sigma_\lambda \\ \downarrow \Pi & & \downarrow N \\ \Sigma_\infty & \xrightarrow{\varepsilon} & [0, 1] \end{array}$$

Later, we will show that Π is not injective. Indeed, we have $\Pi(\tilde{\mathcal{E}}_l) = \Pi(\tilde{\mathcal{E}}_r)$, and $\Pi : \Omega_\infty \setminus \tilde{\mathcal{E}}_r \rightarrow \Sigma_\infty$ is bijective, see Proposition 4.5 for detail. We also point out that Π play an essential role in the study of the symbolic space Ω_∞ .

It was conjectured in [23] and proved in [6] that all the possible gaps of σ_λ are open and labeled by dyadic numbers or dyadic numbers divided by 3 in $(0, 1)$. However, to the best of our knowledge, the complete characterization of the labels of these gaps seems unknown in the literature (compare [6] Theorem 4 (iv) and [5] Sec. 5.4). With the coding map in hand, we can determine the exact set of numbers which label the gaps.

It is known that N is constant on a gap G of the spectrum and we denote this number by $N(G)$.

Theorem 1.11. We have the gap labelling:

$$\{N(G) : G \in \mathcal{G}_I\} = \mathcal{D} \cap (0, 1) \quad \text{and} \quad \{N(G) : G \in \mathcal{G}_{II}\} = \mathcal{E},$$

where \mathcal{D} is the set of positive dyadic numbers and

$$(17) \quad \mathcal{E} = \varepsilon(\Pi(\tilde{\mathcal{E}}_1)) = [(\mathcal{D}/3 \setminus \mathcal{D}) \cap (0, 1)] \setminus \varepsilon(\Pi(\mathcal{F})).$$

Remark 1.12. Assume $s \in (0, 1)$ is a dyadic number divided by 3. (17) tells us that s is a label of some gap of σ_λ if and only if $s \notin \varepsilon(\Pi(\mathcal{F}))$.

1.3. Sketch of the main ideas.

Very roughly, our idea is the following: using periodic approximations to construct a family of nested optimal coverings; find out all the possible band configurations between consecutive levels; define the types of bands and figure out the evolution laws of types; finally construct the symbolic space. We explain more details in the following.

1.3.1. Nested structure.

The family of coverings is a nested structure which we define now.

For any $n \geq 0$, let $\mathcal{I}_n = \{I_{n,1}, \dots, I_{n,k_n}\}$ be a family of compact, non degenerate intervals. $\mathcal{I} := \{\mathcal{I}_n : n \geq 0\}$ is called a *nested structure(NS)*, if

- i) \mathcal{I} is *optimal*: for any $n \geq 0$, $i \neq j$, $I_{n,i} \not\subset I_{n,j}$;
- ii) \mathcal{I} is *nested*: for any $I \in \mathcal{I}_{n+1}$, there is a unique $J \in \mathcal{I}_n$ such that $I \subset J$;
- iii) \mathcal{I} is *minimal*: for any $J \in \mathcal{I}_n$, there exists $I \in \mathcal{I}_{n+1}$ such that $I \subset J$.

Assume $\mathcal{I} := \{\mathcal{I}_n : n \geq 0\}$ is a NS. By ii), $\{\bigcup_{i=1}^{k_n} I_{n,i} : n \geq 0\}$ is a decreasing sequence of nonempty compact sets. Define

$$A(\mathcal{I}) := \bigcap_{n \geq 0} \bigcup_{i=1}^{k_n} I_{n,i}.$$

We call $A(\mathcal{I})$ the *limit set* of \mathcal{I} .

Assume $\mathcal{I} = \{\mathcal{I}_n : n \geq 0\}$ is a NS. \mathcal{I} is called a *separating nested structure(SNS)*, if moreover,

- iv) \mathcal{I} is *separating*: for each n , \mathcal{I}_n is a disjoint family.

We will construct a NS such that its limit set is σ_λ . Then we choose a sub-NS from it such that this sub-NS is a SNS and for which we can estimate the dimension.

1.3.2. NS via Periodic approximations.

For all $n \geq 0$, define the n -th *periodic approximation* of σ_λ as

$$(18) \quad \sigma_{\lambda,n} := \{E \in \mathbb{R} : |h_n(E)| \leq 2\}.$$

By Floquet theory, $\sigma_{\lambda,n}$ is made of 2^n non-overlapping intervals. Following the convention, we call these intervals bands. We denote this family of bands by \mathcal{B}_n and code it by $\Sigma_n = \{0, 1\}^n$ as

$$(19) \quad \mathcal{B}_n := \{B_\sigma : \sigma \in \Sigma_n\}.$$

See Sec. 2.1.2 for detail. It is well-known that $\sigma_{\lambda,n} \rightarrow \sigma_\lambda$ in Hausdorff distance. In general, \mathcal{B}_n is not a covering of σ_λ . However we will show that $\{\mathcal{B}_n \cup \mathcal{B}_{n+1} : n \geq 0\}$ is a family of nested coverings. We will see that $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ is always not optimal. To get an optimal covering, we simply delete those $B \in \mathcal{B}_{n+1}$ which is contained in some $\hat{B} \in \mathcal{B}_n$. Denote by \mathcal{B}_n the resulting optimal covering. Formally

$$(20) \quad \mathcal{B}_n := \mathcal{B}_n \cup \{B \in \mathcal{B}_{n+1} : B \not\subset \hat{B} \text{ for any } \hat{B} \in \mathcal{B}_n\}.$$

Now $\mathcal{B} := \{\mathcal{B}_n : n \geq 0\}$ become a NS and its limit set coincides with σ_λ , see Corollary 3.10. In principle, one can construct a symbolic space and the related coding map of the spectrum by following the sequence $\{\mathcal{B}_n : n \geq 0\}$. However, we warn that when $\lambda > 0$ is small, \mathcal{B} is never a SNS. As a consequence, it is very difficult to show the injectivity of the coding map.

To decide which bands in \mathcal{B}_{n+1} should be deleted, we need to study the configurations of bands in $\mathcal{B}_n \cup \mathcal{B}_{n+1}$. This is the content of Lemma 2.3—the first key lemma of the paper.

1.3.3. Types, graph-directed structure and coding.

The next step is to study how the bands in \mathcal{B}_{n+1} are situated in their father bands in \mathcal{B}_n . The crucial concept here is the *type* of the band.

By performing a renormalization process, we observe that every band in \mathcal{B}_n has certain type. The type of $B \in \mathcal{B}_n$ determines its sons $\hat{B} \in \mathcal{B}_{n+1}$ and their types, see Figure 2 for some intuition. By detailed analysis, we obtain an alphabet \mathcal{A} for the types and a directed graph \mathbb{G} for the evolution of types.

Recall that \mathcal{B}_n is the union of \mathcal{B}_n and a subset of \mathcal{B}_{n+1} . Roughly speaking, there are four types for $B \in \mathcal{B}_n$: 0, 1, 2, 3. For $k = 0, 1, 2$, B has type k if $B \in \mathcal{B}_n$ and $\#\partial B \cap \mathcal{R}_{n-1} = k$ where \mathcal{R}_{n-1} is the set of zeros of trace polynomials h_0, \dots, h_{n-1} (see (28)). B has type 3 if $B \notin \mathcal{B}_n$. Since we aim to give an order-preserving coding for the spectrum, the order of the son bands become important. With this in mind, we observe that the bands with same types but with different parities of levels have different evolution laws: they are reflectional symmetric. This observation suggests that we need to distinguish $0_o, 0_e$ etc. This explains the definition (9) except for type 3, for which we need to distinguish left and right further. The evolution laws (10) of types are hidden in Lemma 3.1—the second key lemma of this paper. Now we can build a related directed graph \mathbb{G} .

Once \mathbb{G} is constructed, we can first code the covering \mathcal{B}_n by Ω_n (the set of admissible words of level n , see (42)) as $\mathcal{B}_n := \{I_w : w \in \Omega_n\}$ and then code the spectrum $\sigma_\lambda = A(\mathcal{B})$ by the symbolic space Ω_∞ as $\pi(\omega) := \bigcap_n I_{\omega|_n}$. As we have mentioned, it is easy to show that π is surjective and continuous. However to show that π is injective, we need further information on ∞ -type energies, see Sec. 1.3.5.

1.3.4. Sub-SNS and the lower bound of the dimension.

Now consider the sub-alphabet $\tilde{\mathcal{A}} = \{1_e, 1_o, 2_e, 2_o\}$ and the sub directed graph $\tilde{\mathbb{G}}$ (enclosed by the red square in Figure 1) with restricted admissible rules:

$$1_e \rightarrow 2_o; \quad 1_o \rightarrow 2_e; \quad 2_e \rightarrow 1_o, 2_o; \quad 2_o \rightarrow 1_e, 2_e.$$

Consider the sub-NS $\tilde{\mathcal{B}} := \{\tilde{\mathcal{B}}_n : n \geq 1\}$ defined by

$$\tilde{\mathcal{B}}_n := \{I_{0_e 1_{\sigma} w} : w = w_1 \cdots w_n, w_i \in \tilde{A}\}.$$

Obviously its limit set $A(\tilde{\mathcal{B}})$ is a subset of the spectrum. Indeed, each energy in $A(\tilde{\mathcal{B}})$ has a bounded trace orbit of bound 2. In particular, $A(\tilde{\mathcal{B}}) \subset B_\infty$. Dynamically, if $E \in A(\tilde{\mathcal{B}})$, then for any $n \geq 0$,

$$f^n(\ell(E)) \in [-2, 2] \times [-2, 2].$$

In some sense, $A(\tilde{\mathcal{B}})$ is the portion of σ_λ on which the trace polynomials behave in a uniformly hyperbolic way. We can show that $\tilde{\mathcal{B}}$ is a SNS and there is a uniform estimation for the length of the bands in $\tilde{\mathcal{B}}_n$, see Proposition 5.8. Now by a general result for SNS (see Proposition 5.7), we obtain a lower bound for the Hausdorff dimension of the spectrum.

1.3.5. Order, Gaps and ∞ -type energies.

Now we explain the orders on Ω_∞ . There are two orders on Ω_∞ , both are induced from orders on \mathcal{A} . So we start from the orders on \mathcal{A} .

We have two goals for designing an order on \mathcal{A} : 1) to remember the order of the son-bands inside a father-band; 2) to remember whether two neighbour son-bands overlap. As we have mentioned several times, if λ is small, the overlapping of some son-bands are unavoidable. To obtain a nice coding, it is essential to remember whether two bands overlap. For this purpose, we define the following two ‘‘orders’’ to compare two bands.

Assume $I = [a, b], J = [c, d]$. We define

$$(21) \quad \begin{cases} I \prec J, & \text{if } a < c; b < d, \\ I < J, & \text{if } b < c. \end{cases}$$

$I < J$ means I and J are disjoint and J is on the right of I . While $I \prec J$ means that $I \not\subset J, J \not\subset I$ and J is on the right of I , however I and J may overlap. It is obvious that $I < J$ implies $I \prec J$. Lemmas 3.6-3.8 summarize the order configurations of son bands for all types of bands. These motivate the definitions of two orders \preceq and \leq on \mathcal{A} (see (43) and (44)).

Now we can extend these orders to Ω_∞ in an obvious way to get the two orders \preceq and \leq on Ω_∞ . We will show that \preceq is a total order on Ω_∞ , which is the order appearing in Theorem 1.5. \leq is stronger than \preceq in the following sense: if $\omega < \omega'$, then $\pi(\omega) < \pi(\omega')$; however, if $\omega \prec \omega'$ but $\omega \not< \omega'$, we only have $\pi(\omega) \leq \pi(\omega')$. It is easy to show the former statement, which follows from the disjointness of the bands. It is quite subtle to show the latter statement because the condition means that the bands containing $\pi(\omega)$ and $\pi(\omega')$ overlap with each other. We obtain the inequality by a detailed analysis on the combinatorics of \mathcal{Z} . Note that our ultimate goal is to show the strict inequality, which is even harder. We need to study the dynamics of the trace map of f .

Next we study the gap of σ_λ . Since (Ω_∞, \preceq) is totally ordered, the notion of gap for Ω_∞ is available. We will at first study the gap structure on Ω_∞ (see Theorem 4.3), and then descend them to σ_λ by the coding map π .

Now we discuss ∞ -type energy. Recall that $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ is a covering of the spectrum, this implies that for any $E \in \sigma_\lambda$ and any $n \geq 0$, $|h_n(E)| \leq 2$ or $|h_{n+1}(E)| \leq 2$. Inspired by the Thue-Morse case [21], to produce a unbounded trace orbit, we can ask for the following mechanism: there exists N such that

$$(22) \quad |h_{N+2k}(E)| \leq 2; \quad |h_{N+2k+1}(E)| > 2, \quad \text{for any } k.$$

Dynamically, this means that for n large enough (if N is even),

$$f^n(\ell(E)) \in [-2, 2] \times [-2, 2]^c.$$

If E has coding ω , (22) implies that necessarily ω is eventually $0_*3_*0_*3_* \dots$. With this in mind, we finally succeed to show that all the energies with coding eventually $(0_a3_b)^\infty$ are of ∞ -type.

As one will see in Sec. 6, it is relatively easy to show that E is of ∞ -type if the coding of E is eventually $(0_o3_{el})^\infty$ or $(0_e3_{or})^\infty$; however it is much harder to show that E is of ∞ -type if the coding of E is eventually $(0_o3_{er})^\infty$ or $(0_e3_{ol})^\infty$. For the latter case, we need to study the local dynamics of f around the fixed point $(-1, -1)$. The injectivity of π is also a consequence of this study.

1.3.6. Labelled directed graph and integrated density of states.

At last, we explain the labels on the edges of \mathbb{G} . Assume $B_\sigma \in \mathcal{B}_n$ has type α , $B_\tau \in \mathcal{B}_{n+1}$ has type β and $B_\tau \subset B_\sigma$. Then we will show that $\alpha \rightarrow \beta$ and σ is a prefix of τ . Write $\tau = \sigma\sigma'$, then the label of the edge $\alpha\beta$ is α' . Now by Lemmas 3.5-3.8. we can label all the edges in \mathbb{G} .

Next we compute the IDS for $E \in \mathcal{Z}$. To do this we need to study the structure of \mathcal{Z} , see Proposition 2.6. Once this is computed, since \mathcal{Z} is dense in σ_λ , by the continuity of N , we can compute the IDS for all $E \in \sigma_\lambda$.

Now we come to the last point-gap labelling. By Theorem 1.8, we know the coding of the gap edges. To compute the IDS of gaps, we just follow the path

$$E \rightarrow \pi^{-1}(E) \rightarrow \Pi \circ \pi^{-1}(E) \rightarrow \varepsilon \circ \Pi \circ \pi^{-1}(E).$$

Final remark about the proof: As we will see in all the properties which need to distinguish the parities of the level n that the statements for the odd case are completely symmetric with that for the even case, so are the proofs. So in the rest of the paper, usually we only give the proofs for the odd case and leave the routine checking for the even case to the reader.

1.4. Structure of the paper.

The rest of this paper is organized as follows. In Sec. 2, we study the band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ and give a characterization for \mathcal{B}_n , we also study the properties of \mathcal{Z} . In Sec. 3, we introduce the concept of type and derive the evolution law and the order of types and the labels of the directed edges. In Sec. 4, we study the symbolic space Ω_∞ . In Sec. 5, we prove a weak version of Theorem 1.5, and apply it to prove Theorem 1.1. In Sec. 6, by a detailed study of ∞ -type energies, we prove Theorems 1.3, 1.5 and 1.8. In

Sec. 7, we prove Theorems 1.10 and 1.11. In Sec. 8, we prove the technical lemmas and the related results.

2. OPTIMAL COVERING \mathcal{B}_n AND THE PROPERTY OF \mathcal{Z}

At first, we study the band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ and give a characterization for \mathcal{B}_n . Then we study the property of \mathcal{Z} .

2.1. The optimal covering \mathcal{B}_n .

2.1.1. Coverings of σ_λ via periodic approximations.

It is known that ([6, 23]) the sequence of trace polynomials of PDH $\{h_n : n \geq 0\}$ defined by (5) satisfy the following initial conditions and recurrence relation:

$$(23) \quad \begin{cases} h_0(E) = E - \lambda, & h_1(E) = E^2 - \lambda^2 - 2; \\ h_{n+1}(E) = h_n(E) (h_{n-1}^2(E) - 2) - 2, & (\forall n \geq 1). \end{cases}$$

We have defined $\sigma_{\lambda,n}$ in (18) and \mathcal{B}_n in (19). By Floquet theory (see for example [28] chapter 4), $\sigma_{\lambda,n}$ is the spectrum of $H_{V^{(n)}}$, where $V^{(n)}$ is 2^n -periodic and

$$V^{(n)}(j) = \lambda V_\xi(j), \quad j = 0, \dots, 2^n - 1.$$

$\sigma_{\lambda,n}$ is called the n -th *periodic approximation* of σ_λ . We know that \mathcal{B}_n is a non-overlapping family. Indeed, we can say more:

Proposition 2.1. *For any $n \geq 1$, all the possible gaps of $\sigma_{\lambda,n}$ are open, i.e. \mathcal{B}_n is a disjoint family.*

We will prove this proposition in Sec. 8.1.3. In general, \mathcal{B}_n is not a covering of σ_λ . However, we have

Lemma 2.2. *i) For any $n \geq 0$,*

$$\sigma_{\lambda,n+2} \subset \sigma_{\lambda,n} \cup \sigma_{\lambda,n+1}.$$

Thus $\{\sigma_{\lambda,n} \cup \sigma_{\lambda,n+1} : n \geq 0\}$ is a decreasing sequence. Moreover,

$$(24) \quad \sigma_\lambda = \bigcap_{n \geq 0} (\sigma_{\lambda,n} \cup \sigma_{\lambda,n+1}).$$

ii) We have

$$(25) \quad \max_{B \in \mathcal{B}_n} |B| \rightarrow 0, \quad n \rightarrow \infty.$$

See [20] Lemma A.2. for a proof (there all the results are proven for TMH, but the proof for PDH is the same).

As a consequence, $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ is a covering of σ_λ .

2.1.2. The coding of \mathcal{B}_n .

We mentioned that $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ is not an optimal covering of σ_λ . To construct an optimal covering, we need to throw some bands in \mathcal{B}_{n+1} . To proceed, we give a coding for \mathcal{B}_n .

Assume $n \geq 1$. Let $\Sigma_n := \{0, 1\}^n$ and $\mathbb{M}_n := \{0, 1, \dots, 2^n - 1\}$. By convention, $\Sigma_0 := \{\emptyset\}$ and $\mathbb{M}_0 = \{0\}$. We endow Σ_n with the lexicographical order and denote by \leq . If $\sigma \neq 1^n$ ($\sigma \neq 0^n$), denote the *successor* (*predecessor*) of σ by σ^+ (σ^-). For $\sigma = \sigma_1 \cdots \sigma_n \in \Sigma_n$ and $k \leq n$, we define the k -th prefix of σ as

$$\sigma|_k := \sigma_1 \cdots \sigma_k.$$

Assume $n, k \geq 0$ and $\sigma \in \Sigma_n, \tau \in \Sigma_k$. Then $\sigma \wedge \tau$ is the maximal common prefix of σ and τ ; $\sigma\tau \in \Sigma_{n+k}$ is the concatenation of σ and τ . If σ is a prefix of τ , we write $\sigma \triangleleft \tau$. We denote the length of σ by $|\sigma|$. By convention, $|\emptyset| = 0$.

When $n = 0$, define $\varpi_0 : \Sigma_0 \rightarrow \mathbb{M}_0$ as $\varpi_0(\emptyset) = 0$. When $n \geq 1$, define $\varpi_n : \Sigma_n \rightarrow \mathbb{M}_n$ as

$$(26) \quad \varpi_n(\sigma) := \sum_{j=1}^n \sigma_j 2^{n-j} = 2^n \sum_{j=1}^n \frac{\sigma_j}{2^j}, \quad \text{where } \sigma = \sigma_1 \cdots \sigma_n.$$

It is direct to check that $\varpi_n : \Sigma_n \rightarrow \mathbb{M}_n$ is bijective and order-preserving.

Now we list the 2^n bands in \mathcal{B}_n from left to right with numbering $0, \dots, 2^n - 1$. Then we denote the j -th bands by $B_{\varpi_n^{-1}(j)}$. In this way, we give a coding for \mathcal{B}_n .

For any $n \geq 0$, combining with Proposition 2.1, we have

$$\mathcal{B}_n = \{B_\sigma : \sigma \in \Sigma_n\} \quad \text{and} \quad B_\sigma < B_{\sigma'} \text{ if } \sigma < \sigma'.$$

Now we have

$$(27) \quad \sigma_{\lambda, n} = \bigcup_{B \in \mathcal{B}_n} B = \bigcup_{\sigma \in \Sigma_n} B_\sigma.$$

We say that all the bands in \mathcal{B}_n are of *level- n* .

For any $\sigma \in \Sigma_n$, write

$$[a_\sigma, b_\sigma] := B_\sigma.$$

For any $n \geq 0$, define

$$(28) \quad \mathcal{Z}_n := \{E \in \mathbb{R} : h_n(E) = 0\} \quad \text{and} \quad \mathcal{R}_n := \bigcup_{j=0}^n \mathcal{Z}_j.$$

By Floquet theory, in each band B_σ , there is exactly one zero of h_n , which we denote by z_σ . So we have

$$(29) \quad a_\sigma < z_\sigma < b_\sigma; \quad \mathcal{Z}_n = \{z_\sigma : \sigma \in \Sigma_n\}; \quad z_\sigma < z_\tau \text{ if } \sigma < \tau.$$

As an example, we compute \mathcal{B}_0 and \mathcal{B}_1 . By (23), we have

$$(30) \quad \begin{cases} B_\emptyset = [\lambda - 2, \lambda + 2], & B_0 = [-\sqrt{\lambda^2 + 4}, -\lambda], & B_1 = [\lambda, \sqrt{\lambda^2 + 4}] \\ \mathcal{B}_0 = \{B_\emptyset\}, & \mathcal{B}_1 = \{B_0, B_1\}. \end{cases}$$

2.1.3. The band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ and characterization of \mathcal{B}_n .

Recall that we have defined two “orders” for bands in (21). The band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1}$ is summarized in the following technical lemma:

Lemma 2.3. Fix $n \geq 0$ and $\sigma \in \Sigma_n$.

(i) Assume n is odd. Then $b_{\sigma 0} = z_\sigma, B_{\sigma 0} \subset B_\sigma$ and $a_{\sigma 1} \notin \mathcal{R}_n$. Moreover,

$$\begin{cases} B_{\sigma 0} = [a_\sigma, z_\sigma], & \text{if } a_\sigma \in \mathcal{R}_{n-1} \\ B_{\sigma 0} \subset (a_\sigma, z_\sigma], & \text{if } a_\sigma \notin \mathcal{R}_{n-1} \end{cases}; \quad \begin{cases} B_{\sigma 1} \subset (z_\sigma, b_\sigma], & \text{if } b_\sigma \in \mathcal{R}_{n-1} \\ [z_\sigma, b_\sigma] \prec B_{\sigma 1}, & \text{if } b_\sigma \notin \mathcal{R}_{n-1} \end{cases}$$

(ii) Assume n is even. Then $a_{\sigma 1} = z_\sigma, B_{\sigma 1} \subset B_\sigma$ and $b_{\sigma 0} \notin \mathcal{R}_n$. Moreover,

$$\begin{cases} B_{\sigma 1} = [z_\sigma, b_\sigma], & \text{if } b_\sigma \in \mathcal{R}_{n-1} \\ B_{\sigma 1} \subset [z_\sigma, b_\sigma), & \text{if } b_\sigma \notin \mathcal{R}_{n-1} \end{cases}; \quad \begin{cases} B_{\sigma 0} \subset [a_\sigma, z_\sigma), & \text{if } a_\sigma \in \mathcal{R}_{n-1} \\ B_{\sigma 0} \prec [a_\sigma, z_\sigma], & \text{if } a_\sigma \notin \mathcal{R}_{n-1} \end{cases}$$

iii) $B_{\sigma 0} < B_{\sigma 1}$. If $\sigma > 0^n$, then $B_{\sigma^-} < B_{\sigma 0}$; if $\sigma < 1^n$, then $B_{\sigma 1} < B_{\sigma^+}$.

Here we use the convention that $\mathcal{R}_{-1} = \emptyset$.

One can see Figure 2 for some concrete examples.




FIGURE 2. Band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1} \cup \mathcal{B}_{n+2}$

We will prove this lemma in Sec. 8.2. For the moment, we will use it to characterize the bands in \mathcal{B}_n .

Recall that \mathcal{B}_n is defined by (20). By Proposition 2.1 and Lemma 2.2 i), \mathcal{B}_n is an optimal covering of σ_λ . By the definition, we always have $\mathcal{B}_n \subset \mathcal{B}_n$. Now we give a characterization for a band $B \in \mathcal{B}_{n+1}$ in \mathcal{B}_n .

Proposition 2.4. *Assume $\sigma \in \Sigma_{n+1}$. Then $B_\sigma \in \mathcal{B}_n$ if and only if $B_\sigma \not\subset B_{\sigma|_n}$.*

Proof. Write $\tau := \sigma|_n$. If $B_\sigma \subset B_\tau \in \mathcal{B}_n$, by (20), $B_\sigma \notin \mathcal{B}_n$.

Now assume $B_\sigma \not\subset B_\tau$. By Lemma 2.3 iii),

$$B_{\tau^-} < B_\sigma, \text{ if } \tau > 0^n; \quad B_\sigma < B_{\tau^+}, \text{ if } \tau < 1^n.$$

By the coding and the order of the bands in \mathcal{B}_n , we conclude that $B_\sigma \not\subset B$ for any $B \in \mathcal{B}_n$. So by (20), $B_\sigma \in \mathcal{B}_n$. \square

As an example, by (30), we have

$$(31) \quad B_1 \subset B_\emptyset \quad \text{and} \quad B_0 \prec B_\emptyset.$$

Consequently we have (see Figure 2 (a))

$$\mathcal{B}_0 = \{B_0, B_\emptyset\}.$$

By direct computation, we have (see also Figure 2 (a))

$$\mathcal{B}_1 = \{B_0, B_1, B_{01}, B_{11}\}.$$

2.2. The property of \mathcal{Z} .

At first, we have the following simple but useful observation:

Lemma 2.5. *i) For $n \geq 0$, if $E \in \mathcal{Z}_n$, then*

$$(32) \quad h_{n+1}(E) = -2 \quad \text{and} \quad h_k(E) = 2 \text{ for any } k \geq n+2.$$

Consequently, $\mathcal{Z}_n \cap \mathcal{Z}_m = \emptyset$ if $n \neq m$.

ii) For $n \geq 1$ and $\sigma \in \Sigma_n$,

$$\text{int}(B_\sigma) \cap \mathcal{R}_{n-1} = \emptyset \quad \text{and} \quad \text{int}(B_\sigma) \cap \mathcal{R}_n = \{z_\sigma\}.$$

iii) $\mathcal{Z} \subset B_\infty \cap \sigma_\lambda$ and \mathcal{Z} is dense in σ_λ .

Proof. i) By (23), one check directly that the (32) holds. WLOG assume $n < m$. By (32), $h_{n+1}(\mathcal{Z}_n) = \{-2\}$ and $h_k(\mathcal{Z}_n) = \{2\}$ for $k \geq n+2$. Hence $\mathcal{Z}_n \cap \mathcal{Z}_m = \emptyset$.

ii) If $E \in \mathcal{R}_{n-1}$, then by i), $h_n(E) = \pm 2$. By Floquet theory, $h_n(\text{int}(B_\sigma)) = (-2, 2)$. So $\text{int}(B_\sigma) \cap \mathcal{R}_{n-1} = \emptyset$. Still by Floquet theory, $\text{int}(B_\sigma) \cap \mathcal{Z}_n = \{z_\sigma\}$. Since $\mathcal{R}_n = \mathcal{R}_{n-1} \cup \mathcal{Z}_n$, we have $\text{int}(B_\sigma) \cap \mathcal{R}_n = \{z_\sigma\}$.

iii) By i), $\mathcal{Z} \subset B_\infty$. By i) and (24), $\mathcal{Z} \subset \sigma_\lambda$. Fix any $E \in \sigma_\lambda$. For any $n \geq 0$, by (24), there exists $\sigma^{(n)} \in \Sigma_n \cup \Sigma_{n+1}$ such that $E \in B_{\sigma^{(n)}}$. Since $z_{\sigma^{(n)}} \in B_{\sigma^{(n)}}$, by (25), we have $|z_{\sigma^{(n)}} - E| \rightarrow 0$. So \mathcal{Z} is dense in σ_λ . \square

2.2.1. *The order-preserving coding of \mathcal{Z} .*

Recall that \mathcal{Z} is defined by (12). Define

$$(33) \quad \Sigma_* := \bigcup_{n=0}^{\infty} \Sigma_n.$$

By (29) and Lemma 2.5 i), the map $\sigma \rightarrow z_\sigma$ is a bijection between Σ_* and \mathcal{Z} . Thus we obtain a coding of \mathcal{Z} as

$$\mathcal{Z} = \{z_\sigma : \sigma \in \Sigma_*\}.$$

In the following, we define an order on Σ_* such that this coding preserves the orders.

At first we define an order \preceq on $\{0, \emptyset, 1\}$ as

$$0 \prec \emptyset \prec 1.$$

For each $n \geq 1$, let \preceq be the lexicographical order on $\{0, \emptyset, 1\}^n$.

Now we define an order \preceq on Σ_* as follows. Given $\sigma \in \Sigma_n$ and $\tau \in \Sigma_m$. Assume $n \leq m$. Define $\tilde{\sigma} := \sigma \emptyset^{m-n}$. Then define

$$(34) \quad \begin{cases} \sigma \prec \tau & \text{if } \tilde{\sigma} \prec \tau, \\ \sigma = \tau & \text{if } \tilde{\sigma} = \tau, \\ \tau \prec \sigma & \text{if } \tau \prec \tilde{\sigma}. \end{cases}$$

One check directly that \preceq is a total order on Σ_* .

Proposition 2.6. *Assume $\sigma, \tau \in \Sigma_*$. Then*

$$z_\sigma < z_\tau \quad \text{if and only if} \quad \sigma \prec \tau.$$

It has the following consequence:

Corollary 2.7. *If $n \geq 1$ and $\sigma \in \Sigma_n$, then for any $E \in \text{int}(B_\sigma)$,*

$$\prod_{j=0}^{n-1} h_j(E) \begin{cases} < 0, & \text{if } \sigma_n = 0 \\ > 0, & \text{if } \sigma_n = 1. \end{cases}$$

We will prove Proposition 2.6 and Corollary 2.7 in Sec. 8.1.1.

2.2.2. *\mathcal{Z} and band edges.*

Recall that $\mathcal{R}_{-1} = \emptyset$.

Lemma 2.8. *Assume $n, t \geq 0$ and $\sigma \in \Sigma_n$. Then*

$$i) \quad b_{\sigma 01^t} < a_{\sigma 10^t} \text{ and } z_\sigma = \begin{cases} b_{\sigma 01^t}, & \text{if } n \text{ is odd;} \\ a_{\sigma 10^t}, & \text{if } n \text{ is even.} \end{cases}$$

ii) *If $a_\sigma \in \mathcal{R}_{n-1}$, then $a_{\sigma 0^t} = a_\sigma$; if $b_\sigma \in \mathcal{R}_{n-1}$, then $b_{\sigma 1^t} = b_\sigma$.*

iii) *$a_\sigma \in \mathcal{R}_{n-1}$ if and only if $a_{\sigma 0} \in \mathcal{R}_n$. $b_\sigma \in \mathcal{R}_{n-1}$ if and only if $b_{\sigma 1} \in \mathcal{R}_n$.*

iv) *If $\sigma < 1^n$, then $\#(\{b_\sigma, a_{\sigma^+}\} \cap \mathcal{R}_{n-1}) = 1$.*

We will prove this lemma in Sec. 8.1.3.

2.3. Two useful basic facts.

We record two basic facts, which are frequently used in the whole paper.

The following lemma is fundamental for us.

Lemma 2.9. *For any $n \geq 0$ and $E \in \mathbb{R}$,*

$$(35) \quad h_{n+1}(E) - (h_n^2(E) - 2) = (-1)^n 2\lambda \prod_{c \in \mathcal{R}_n} (E - c) = (-1)^n 2\lambda \prod_{j=0}^n h_j(E).$$

Proof. We prove it by induction. For $n = 0$, we have

$$h_1(E) - (h_0^2(E) - 2) = 2\lambda(E - \lambda) = 2\lambda h_0(E).$$

Then (35) holds for $n = 0$.

Assume (35) holds for $n = k \geq 0$. Now consider $n = k + 1$. By (23) and induction hypothesis, we have

$$\begin{aligned} & h_{k+1}(E) - (h_k^2(E) - 2) \\ &= h_k(E)(h_{k-1}^2(E) - 2) - 2 - (h_k^2(E) - 2) \\ &= -h_k(E)(h_k(E) - (h_{k-1}^2(E) - 2)) \\ &= (-1)^k 2\lambda \prod_{j=0}^k h_j(E) = (-1)^k 2\lambda \prod_{c \in \mathcal{R}_k} (E - c). \end{aligned}$$

Hence, the result holds for $n = k + 1$. By induction, the result follows. \square

The following is standard Floquet theory:

Proposition 2.10. *i) For each $\sigma \in \Sigma_n$, $h_n : B_\sigma \rightarrow [-2, 2]$ is a homeomorphism. If $\sigma_n = 0$ ($\sigma_n = 1$), then $h'_n(E) < 0$ ($h'_n(E) > 0$) for $E \in \text{int}(B_\sigma)$. Consequently*

$$h_n(a_\sigma) = 2, \quad h_n(b_\sigma) = -2 \quad (h_n(a_\sigma) = -2, \quad h_n(b_\sigma) = 2).$$

ii) If $\sigma \neq 1^n$, then the gap $(b_\sigma, a_{\sigma+})$ is open if and only if $h'_n(b_\sigma) \neq 0$ or $h'_n(a_{\sigma+}) \neq 0$.

iii) E is an endpoint of some band of level n iff $h_n(E) = \pm 2$. If $h_n(E) = -2$ and $h'_n(E) < 0$ ($h'_n(E) > 0$), then $E = b_\sigma$ (a_σ) for some $\sigma \in \Sigma_n$. If $h_n(E) = 2$ and $h'_n(E) < 0$ ($h'_n(E) > 0$), then $E = a_\sigma$ (b_σ) for some $\sigma \in \Sigma_n$.

3. TYPES, EVOLUTION LAW, ORDER AND LABELS

In this section, at first we study the band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1} \cup \mathcal{B}_{n+2}$. Then we introduce the concept of type, which is fundamental for this paper. Next we derive the evolution law and the order of types and the labels of the directed edges. At last, we show that $\{\mathcal{B}_n : n \geq 0\}$ is a NS with limit set σ_λ .

3.1. band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1} \cup \mathcal{B}_{n+2}$.

Now we present the second technical lemma of our paper, which summarizes all the possible band configurations of $\mathcal{B}_n \cup \mathcal{B}_{n+1} \cup \mathcal{B}_{n+2}$:

Lemma 3.1. *Fix $n \geq 0$ and $\sigma \in \Sigma_n$.*

i) Assume n is odd. Then $a_{\sigma 10}, b_{\sigma 10} \notin \mathcal{R}_{n+1}$ and

$$B_{\sigma 00} \subset B_\sigma, B_{\sigma 10} \subset \text{int}(B_\sigma); \quad B_{\sigma 01} \subset B_{\sigma 0}; \quad B_{\sigma 0} < B_{\sigma 10} \prec B_{\sigma 1}.$$

Moreover we have

$$\begin{cases} B_{\sigma 00} \subset B_{\sigma 0}, & \text{if } a_\sigma \in \mathcal{R}_{n-1} \\ B_{\sigma 00} \prec B_{\sigma 0}, & \text{if } a_\sigma \notin \mathcal{R}_{n-1} \end{cases} \quad \begin{cases} B_{\sigma 11} \subset B_{\sigma 1} \subset B_\sigma, & \text{if } b_\sigma \in \mathcal{R}_{n-1} \\ B_\sigma \prec B_{\sigma 11}, & \text{if } b_\sigma \notin \mathcal{R}_{n-1} \end{cases}$$

If $a_\sigma \notin \mathcal{R}_{n-1}$, then $a_{\sigma 00}, b_{\sigma 00} \notin \mathcal{R}_{n+1}$ and $B_{\sigma 00} \subset \text{int}(B_\sigma)$.

ii) Assume n is even. Then $a_{\sigma 01}, b_{\sigma 01} \notin \mathcal{R}_{n+1}$ and

$$B_{\sigma 11} \subset B_\sigma, B_{\sigma 01} \subset \text{int}(B_\sigma); \quad B_{\sigma 10} \subset B_{\sigma 1}; \quad B_{\sigma 0} \prec B_{\sigma 01} < B_{\sigma 1}.$$

Moreover we have

$$\begin{cases} B_{\sigma 11} \subset B_{\sigma 1}, & \text{if } b_\sigma \in \mathcal{R}_{n-1} \\ B_{\sigma 1} \prec B_{\sigma 11}, & \text{if } b_\sigma \notin \mathcal{R}_{n-1} \end{cases} \quad \begin{cases} B_{\sigma 00} \subset B_{\sigma 0} \subset B_\sigma, & \text{if } a_\sigma \in \mathcal{R}_{n-1} \\ B_{\sigma 00} \prec B_\sigma, & \text{if } a_\sigma \notin \mathcal{R}_{n-1} \end{cases}$$

If $b_\sigma \notin \mathcal{R}_{n-1}$, then $a_{\sigma 11}, b_{\sigma 11} \notin \mathcal{R}_{n+1}$ and $B_{\sigma 11} \subset \text{int}(B_\sigma)$.

Combine Lemmas 2.3 and 3.1, we have

Corollary 3.2. *i) Assume $n \geq 1$ and $\sigma \in \Sigma_n$. Then*

$$B_\sigma \subset B_{\sigma|_{n-1}} \Leftrightarrow B_\sigma \cap \mathcal{R}_{n-1} \neq \emptyset \Leftrightarrow \partial B_\sigma \cap \mathcal{R}_{n-1} \neq \emptyset.$$

ii) Assume $n \geq 1$ and $\sigma \in \Sigma_n$. Then

$$B_\sigma \subset B_{\sigma|_{n-1}} \text{ or } B_\sigma \subset B_{\sigma|_{n-2}}.$$

(For $n = 1$, $B_{\sigma|_0} := B_\emptyset$ and $B_{\sigma|_{-1}} := \mathbb{R}$).

iii) Assume $n \geq 0$ and $\sigma \in \Sigma_n, \tau \in \Sigma_n \cup \Sigma_{n+1} \cup \Sigma_{n+2}$. If $B_\tau \subset B_\sigma$, then $\sigma \triangleleft \tau$.

We will prove Lemma 3.1 and Corollary 3.2 in Sec. 8.3. Now we apply them to derive the concept of type and figure out the evolution law.

3.2. The types of the bands.

Lemmas 2.3 and 3.1 suggest that for a band B_σ of level n , once we know that whether $a_\sigma, b_\sigma \in \mathcal{R}_{n-1}$, the local configurations of B_σ and $B_{\sigma\tau}$ with $|\tau| \leq 2$ are completely determined. This observation motivates the following definition for types of bands.

Definition 3.3. *Assume $n \geq 0$, $0 \leq \kappa \leq 2$ and $B \in \mathcal{B}_n$.*

At first we assume $B \in \mathcal{B}_n$.

We say B has type $\kappa_o(\kappa_e)$, if n is odd (even), and $\#\partial B \cap \mathcal{R}_{n-1} = \kappa$.

(Recall that $\mathcal{R}_{-1} = \emptyset$.)

Next we assume $B \notin \mathcal{B}_n$. Then $B \in \mathcal{B}_{n+1}$. There exists a unique $\sigma \in \Sigma_{n+1}$ such that $B = B_\sigma$. Since $B \in \mathcal{B}_n$, by Proposition 2.4, $B \not\subset B_{\sigma|n}$. By Corollary 3.2 ii), $B \subset B_{\sigma|n-1}$.

If $B_{\sigma|n} \subset B_{\sigma|n-1}$.

We say B has type $3_{ol}(3_{el})$ if n is odd (even) and $B \prec B_{\sigma|n}$.

We say B has type $3_{or}(3_{er})$ if n is odd (even) and $B_{\sigma|n} \prec B$.

If $B_{\sigma|n} \not\subset B_{\sigma|n-1}$.

We say B has type $3_{ol}(3_{el})$ if n is odd (even) and $B_{\sigma|n} \prec B_{\sigma|n-1}$.

We say B has type $3_{or}(3_{er})$ if n is odd (even) and $B_{\sigma|n-1} \prec B_{\sigma|n}$.

By the definition above, each band in \mathcal{B}_n has one and only one type.

Example 3.4. Recall that $\mathcal{B}_0 = \{B_0, B_\emptyset\}$. By the definition, B_\emptyset has type 0_e , B_0 has type 3_{el} , see Figure 2 (a). $\mathcal{B}_1 = \{B_0, B_1, B_{01}, B_{11}\}$. B_0 has type 0_o , B_1 has type 1_o , B_{01} have type 3_{ol} and B_{11} have type 3_{or} , see Figure 2 (a).

3.3. The evolution law and the order of the types, the label of the directed edges.

At first, we study type 3_* bands.

Lemma 3.5. Assume $B_\sigma \in \mathcal{B}_n$ has type 3_{ol} or 3_{or} (3_{el} or 3_{er}). Then B_σ contains exact one band in \mathcal{B}_{n+1} : B_σ itself with type 0_e (0_o).

Proof. We only show the odd case. Assume $B_\sigma \in \mathcal{B}_n$ has type 3_{ol} or 3_{or} , then n is odd and $\sigma \in \Sigma_{n+1}$. By Proposition 2.4, $B_\sigma \not\subset B_{\sigma|n}$. By Corollary 3.2 i), $B_\sigma \cap \mathcal{R}_n = \emptyset$. Thus $B_\sigma \in \mathcal{B}_{n+1} \subset \mathcal{B}_{n+1}$ has type 0_e . By the definition of \mathcal{B}_{n+1} , B_σ is the only band in \mathcal{B}_{n+1} and contained in B_σ . \square

Before continuing, we do the following observation. Take $B_\sigma \in \mathcal{B}_n$ and $B_\tau \in \mathcal{B}_{n+1}$. Assume $B_\tau \subset B_\sigma$. Then $\sigma \in \Sigma_n$, $\tau \in \Sigma_{n+1}$ or Σ_{n+2} . By Corollary 3.2 iii), $\sigma \triangleleft \tau$. So

$$(36) \quad \tau \in \{\sigma 0, \sigma 1, \sigma 00, \sigma 01, \sigma 10, \sigma 11\}.$$

Next, we study type 0_* bands.

Lemma 3.6. Assume $B_\sigma \in \mathcal{B}_n$.

i) If B_σ has type 0_o , then B_σ contains exact three bands in \mathcal{B}_{n+1} with the order:

$$B_{\sigma 00} \prec B_{\sigma 0} < B_{\sigma 10},$$

and with the types $3_{el}, 1_e, 3_{er}$, respectively. Moreover,

$$b_{\sigma 0} = z_\sigma; \quad B_{\sigma 00}, B_{\sigma 10} \subset \text{int}(B_\sigma).$$

ii) If B_σ has type 0_e , then B_σ contains exact three bands in \mathcal{B}_{n+1} with the order:

$$B_{\sigma 01} < B_{\sigma 1} \prec B_{\sigma 11},$$

and with the type $3_{ol}, 1_o, 3_{or}$, respectively. Moreover,

$$a_{\sigma 1} = z_{\sigma}; \quad B_{\sigma 01}, B_{\sigma 11} \subset \text{int}(B_{\sigma}).$$

Proof. We only show i), so we assume n is odd. By the assumption, $\sigma \in \Sigma_n$ and $a_{\sigma}, b_{\sigma} \notin \mathcal{R}_{n-1}$. Assume $B_{\tau} \in \mathcal{B}_{n+1}$ and $B_{\tau} \subset B_{\sigma}$, then τ satisfies (36).

By Lemma 2.3 i) and Lemma 2.8 iii),

$$B_{\sigma 0} \subset B_{\sigma}; \quad B_{\sigma} \prec B_{\sigma 1}; \quad a_{\sigma 0} \notin \mathcal{R}_n; \quad b_{\sigma 0} = z_{\sigma} \in \mathcal{R}_n.$$

So $B_{\sigma 0} \in \mathcal{B}_{n+1}$ has type 1_e and $B_{\sigma 1} \not\subset B_{\sigma}$.

By Lemma 3.1 i),

$$\begin{cases} B_{\sigma 00}, B_{\sigma 10} \subset \text{int}(B_{\sigma}); & B_{\sigma 00} \prec B_{\sigma 0} < B_{\sigma 10} \prec B_{\sigma 1}, \\ B_{\sigma 01} \subset B_{\sigma 0}; & B_{\sigma} \prec B_{\sigma 11}. \end{cases}$$

So $B_{\sigma 11} \not\subset B_{\sigma}$ and $B_{\sigma 01} \notin \mathcal{B}_{n+1}$; $B_{\sigma 00} \not\subset B_{\sigma 0}, B_{\sigma 10} \not\subset B_{\sigma 1}$. By Proposition 2.4, $B_{\sigma 00}, B_{\sigma 10} \in \mathcal{B}_{n+1}$. Since $B_{\sigma 0} \subset B_{\sigma}$ and $B_{\sigma 00} \prec B_{\sigma 0}$, by the definition, $B_{\sigma 00}$ has type 3_{el} . Since $B_{\sigma 1} \not\subset B_{\sigma}$ and $B_{\sigma} \prec B_{\sigma 1}$, by the definition, $B_{\sigma 10}$ has type 3_{er} . \square

As two examples, B_{\emptyset} has type 0_e and B_0 has type 0_o , see Figure 2 (a) and (b) for the order configurations of their son intervals.

Now, we study type 1_* bands.

Lemma 3.7. *Assume $B_{\sigma} \in \mathcal{B}_n$.*

i) *If B_{σ} has type 1_o , then B_{σ} contains exact two bands in \mathcal{B}_{n+1} with the order:*

$$B_{\sigma 0} < B_{\sigma 10},$$

and with the type $2_e, 3_{er}$, respectively. Moreover $B_{\sigma 0} = [a_{\sigma}, z_{\sigma}]$ and $B_{\sigma 10} \subset \text{int}(B_{\sigma})$.

ii) *If B_{σ} has type 1_e , then B_{σ} contains exact two bands in \mathcal{B}_{n+1} with the order:*

$$B_{\sigma 01} < B_{\sigma 1},$$

and with the type $3_{ol}, 2_o$, respectively. Moreover $B_{\sigma 1} = [z_{\sigma}, b_{\sigma}]$ and $B_{\sigma 01} \subset \text{int}(B_{\sigma})$.

Proof. We only show i), so we assume n is odd. By the assumption, $\sigma \in \Sigma_n$ and $\#\{a_{\sigma}, b_{\sigma}\} \cap \mathcal{R}_{n-1} = 1$. Assume $B_{\tau} \in \mathcal{B}_{n+1}$ and $B_{\tau} \subset B_{\sigma}$, then τ satisfies (36).

At first, we claim that $a_{\sigma} \in \mathcal{R}_{n-1}$ and $b_{\sigma} \notin \mathcal{R}_{n-1}$. Indeed, since $\partial B_{\sigma} \cap \mathcal{R}_{n-1} \neq \emptyset$, by Corollary 3.2 i), $B_{\sigma} \subset B_{\sigma|_{n-1}}$. Write $\hat{\sigma} = \sigma|_{n-1}$, notice that $n-1$ is even. By applying Lemma 2.3 ii) to $\hat{\sigma}$, we have $b_{\hat{\sigma}0} \notin \mathcal{R}_{n-1}$ and $a_{\hat{\sigma}1} = z_{\hat{\sigma}} \in \mathcal{R}_{n-1}$. Since $\#\{a_{\sigma}, b_{\sigma}\} \cap \mathcal{R}_{n-1} = 1$, we have $a_{\sigma} \in \mathcal{R}_{n-1}$ and $b_{\sigma} \notin \mathcal{R}_{n-1}$ in either case of $\sigma = \hat{\sigma}0$ or $\sigma = \hat{\sigma}1$. So the claim holds.

By Lemma 2.3 i), $B_{\sigma 0} = [a_{\sigma}, z_{\sigma}] \subset B_{\sigma}$ and $B_{\sigma} \prec B_{\sigma 1}$. Since $a_{\sigma} \in \mathcal{R}_{n-1}$ and $z_{\sigma} \in \mathcal{R}_n$, $B_{\sigma 0} \in \mathcal{B}_{n+1}$ has type 2_e . $B_{\sigma 1} \not\subset B_{\sigma}$.

By Lemma 3.1 i),

$$B_{\sigma 00}, B_{\sigma 01} \subset B_{\sigma 0}; \quad B_{\sigma 10} \subset \text{int}(B_{\sigma}), \quad B_{\sigma 10} \prec B_{\sigma 1}; \quad B_{\sigma} \prec B_{\sigma 11}, \quad B_{\sigma 0} < B_{\sigma 10}.$$

So $B_{\sigma 11} \not\subset B_\sigma$ and $B_{\sigma 00}, B_{\sigma 01} \notin \mathcal{B}_{n+1}$. Since $B_{\sigma 10} \not\subset B_{\sigma 1}$, by Proposition 2.4, $B_{\sigma 10} \in \mathcal{B}_{n+1}$. Since $B_\sigma \prec B_{\sigma 1}$, $B_{\sigma 10}$ has type 3_{er} . So the result holds. \square

As two examples, B_{00} has type 1_e and B_1 has type 1_o , see Figure 2 (c) and (d) for the order configurations of their son intervals.

At last, we study type 2_* bands.

Lemma 3.8. *Assume $B_\sigma \in \mathcal{B}_n$.*

i) If B_σ has type 2_o , then B_σ contains exact three bands in \mathcal{B}_{n+1} with the order:

$$B_{\sigma 0} < B_{\sigma 10} \prec B_{\sigma 1},$$

and with the type $2_e, 3_{el}, 1_e$, respectively.

ii) If B_σ has type 2_e , then B_σ contains exact three bands in \mathcal{B}_{n+1} with the order:

$$B_{\sigma 0} \prec B_{\sigma 01} < B_{\sigma 1},$$

and with the type $1_o, 3_{or}, 2_o$, respectively.

Proof. We only show i), so we assume n is odd. By the assumption, $\sigma \in \Sigma_n$ and $a_\sigma, b_\sigma \in \mathcal{R}_{n-1}$. By Lemma 2.8 ii), $b_{\sigma 1} = b_\sigma$. Assume $B_\tau \in \mathcal{B}_{n+1}$ and $B_\tau \subset B_\sigma$, then τ satisfies (36).

By Lemma 2.3 i), $B_{\sigma 0} = [a_\sigma, z_\sigma], B_{\sigma 1} \subset B_\sigma$, $a_{\sigma 1} \notin \mathcal{R}_n$ and $b_{\sigma 1} = b_\sigma \in \mathcal{R}_n$. So $B_{\sigma 1} \in \mathcal{B}_{n+1}$ has type 1_e . Since $a_\sigma \in \mathcal{R}_{n-1}$ and $z_\sigma \in \mathcal{R}_n$, $B_{\sigma 0} \in \mathcal{B}_{n+1}$ has type 2_e .

By Lemma 3.1 i),

$$B_{\sigma 00}, B_{\sigma 01} \subset B_{\sigma 0}; \quad B_{\sigma 10} \subset B_{\sigma 1}; \quad B_{\sigma 11} \subset B_{\sigma 1}, \quad B_{\sigma 0} < B_{\sigma 10} \prec B_{\sigma 1}.$$

So $B_{\sigma 00}, B_{\sigma 01}, B_{\sigma 11} \notin \mathcal{B}_{n+1}$. Since $B_{\sigma 10} \not\subset B_{\sigma 1}$, by Proposition 2.4, $B_{\sigma 10} \in \mathcal{B}_{n+1}$. Since $B_{\sigma 1} \subset B_\sigma$ and $B_{\sigma 10} \prec B_{\sigma 1}$, by the definition, $B_{\sigma 10}$ has type 3_{el} . \square

As two examples, B_{10} has type 2_e and B_{101} has type 2_o , see Figure 2 (e) and (f) for the order configurations of their son intervals.

We summarize Lemmas 3.5-3.8 in a symbolic way as follows:

$$(37) \quad 3_{ol}, 3_{or} \xrightarrow{\emptyset} 0_e; \quad 3_{el}, 3_{er} \xrightarrow{\emptyset} 0_o$$

$$(38) \quad \begin{cases} 0_o \xrightarrow{00} 3_{el}; & 0_o \xrightarrow{0} 1_e; & 0_o \xrightarrow{10} 3_{er}; & 3_{el} \prec 1_e < 3_{er}; \\ 0_e \xrightarrow{01} 3_{ol}; & 0_e \xrightarrow{1} 1_o; & 0_e \xrightarrow{11} 3_{or}; & 3_{ol} < 1_o \prec 3_{or}. \end{cases}$$

$$(39) \quad \begin{cases} 1_o \xrightarrow{0} 2_e; & 1_o \xrightarrow{10} 3_{er}; & 2_e < 3_{er}; \\ 1_e \xrightarrow{01} 3_{ol}; & 1_e \xrightarrow{1} 2_o; & 3_{ol} < 2_o. \end{cases}$$

$$(40) \quad \begin{cases} 2_o \xrightarrow{0} 2_e; & 2_o \xrightarrow{10} 3_{el}; & 2_o \xrightarrow{1} 1_e; & 2_e < 3_{el} \prec 1_e; \\ 2_e \xrightarrow{0} 1_o; & 2_e \xrightarrow{01} 3_{or}; & 2_e \xrightarrow{1} 2_o; & 1_o \prec 3_{or} < 2_o. \end{cases}$$

Here $\alpha \xrightarrow{\tau} \beta$ means type α can evolve to type β and the directed edge $\alpha\beta$ is labeled by τ .

Finally, consider $\mathcal{B}_0 = \{B_0, B_\emptyset\}$. By (31), $B_0 \prec B_\emptyset$. By Example 3.4, B_0 has type 3_{el} and B_\emptyset has type 0_e . So we complete the order of the types by

$$(41) \quad 3_{el} \prec 0_e.$$

3.4. $\{\mathcal{B}_n : n \geq 0\}$ as a NS.

At first we have

Proposition 3.9. *For each $\tilde{B} \in \mathcal{B}_{n+1}$, there is a unique band $B \in \mathcal{B}_n$ such that $\tilde{B} \subset B$.*

Proof. Let $\sigma \in \Sigma_{n+1} \cup \Sigma_{n+2}$ be the unique word such that $\tilde{B} = B_\sigma$. If $B \in \mathcal{B}_n$ is such that $\tilde{B} \subset B$, then by Corollary 3.2 iii), the only two possibilities are $B = B_{\sigma|_n}$ or $B_{\sigma|_{n+1}}$.

If \tilde{B} has type 3_* , then $\sigma \in \Sigma_{n+2}$. By Proposition 2.4, $B_\sigma \not\subset B_{\sigma|_{n+1}}$. By Corollary 3.2 ii), $B_\sigma \subset B_{\sigma|_n} \in \mathcal{B}_n \subset \mathcal{B}_n$.

If \tilde{B} has type 0_* , then $\sigma \in \Sigma_{n+1}$ and $\partial B_\sigma \cap \mathcal{R}_n = \emptyset$. By Corollary 3.2 i), $B_\sigma \not\subset B_{\sigma|_n}$. By Proposition 2.4, $B_\sigma \in \mathcal{B}_n$ and $\tilde{B} = B_\sigma$.

If \tilde{B} has type 1_* or 2_* , then $\sigma \in \Sigma_{n+1}$ and $\partial B_\sigma \cap \mathcal{R}_n \neq \emptyset$. By Corollary 3.2 i), $B_\sigma \subset B_{\sigma|_n} \in \mathcal{B}_n$. By the definition of \mathcal{B}_n , $B_\sigma \notin \mathcal{B}_n$.

In all the cases, there is a unique $B \in \mathcal{B}_n$ such that $\tilde{B} \subset B$. □

See Sec. 1.3.1 for the definition of NS. Now we can show that

Corollary 3.10. $\mathcal{B} := \{\mathcal{B}_n : n \geq 0\}$ is a NS and $A(\mathcal{B}) = \sigma_\lambda$.

Proof. By Proposition 2.1, \mathcal{B}_n is a disjoint family. Combine with the definition (20) of \mathcal{B}_n , \mathcal{B} is optimal. By Proposition 3.9, \mathcal{B} is nested. By Lemmas 3.5-3.8, each band in \mathcal{B}_n contains at least one band in \mathcal{B}_{n+1} , so \mathcal{B} is minimal. By the definition of NS (see Sec. 1.3.1), \mathcal{B} is a NS. By (27) and the definition of \mathcal{B}_n ,

$$\bigcup_{B \in \mathcal{B}_n} B = \sigma_{\lambda, n} \cup \sigma_{\lambda, n+1}.$$

By (24), σ_λ is the limit set of \mathcal{B} . □

As a final complement, we show the following

Lemma 3.11. *If $B \in \mathcal{B}_n$ has type $0_e(0_o)$, then h_n is increasing (decreasing) on B .*

Proof. We only show the case that B has type 0_e . The proof of the other case is the same.

If $n = 0$, then $B = B_\emptyset$ and $h_0(E) = E - \lambda$, so the result holds.

Now assume $n \geq 1$ and n is even. Assume $B = B_\sigma$ for some $\sigma \in \Sigma_n$. We claim that $\sigma_n = 1$. If otherwise $\sigma = \tau 0$ for some $\tau \in \Sigma_{n-1}$. Then by Lemma 2.3 i), we have $b_\sigma = z_\tau \in \mathcal{R}_{n-1}$, which contradicts with the fact that B has type 0_e . Now by Proposition 2.10 i), h_n is increasing on B . □

4. SYMBOLIC SPACE Ω_∞

In this section, we define two orders on the symbolic space Ω_∞ and characterize the gaps of it by using the map Π .

We have defined the symbolic space Ω_∞ in Sec. 1.2.4. Through Sec. 3, we understand that \mathcal{A} , defined by (9), is just the set of all the possible types; (10) is just a summary of the evolution laws appeared in (37)-(40). These information tell us how to construct the directed graph \mathbb{G} . The labels of edges of \mathbb{G} can also be read off from (37)-(40). Now the order relations appeared in (38)-(41) suggest the following definitions of orders on Ω_∞ .

4.1. Two orders on Ω_∞ .

Define the set of *admissible words* as

$$(42) \quad \Omega_n := \{w \in \mathcal{A}^{n+1} : w_0 \in \{3_{el}, 0_e\}, a_{w_j w_{j+1}} = 1\}, \quad n \geq 0 \quad \text{and} \quad \Omega_* := \bigcup_{n \geq 0} \Omega_n.$$

For $\omega = \omega_0 \omega_1 \cdots \in \Omega_* \cup \Omega_\infty$, write $\omega|_n := \omega_0 \cdots \omega_n$. For $\omega, \omega' \in \Omega_* \cup \Omega_\infty$, write $\omega \wedge \omega'$ for their maximal common prefix. For $w \in \Omega_n$, define the cylinder as

$$[w] := \{\omega \in \Omega_\infty : \omega|_n = w\}.$$

4.1.1. Two orders on \mathcal{A} .

Define \prec to be the smallest partial order on \mathcal{A} such that the following holds:

$$(43) \quad 3_{ol} \prec 1_o \prec 3_{or} \prec 2_o; \quad 2_e \prec 3_{el} \prec 1_e \prec 3_{er}; \quad 3_{el} \prec 0_e.$$

Define \leq to be the smallest partial order on \mathcal{A} such that the following holds:

$$(44) \quad 3_{ol} < 1_o < 2_o; \quad 3_{ol} < 3_{or} < 2_o; \quad 2_e < 3_{el} < 3_{er}; \quad 2_e < 1_e < 3_{er}.$$

(Here we view partial order as a subset of $\mathcal{A} \times \mathcal{A}$.)

By (43) and (44), it is direct to check that if $\alpha < \beta$, then $\alpha \prec \beta$.

Remark 4.1. The idea for defining (43) and (44) is as follows. At first, we impose

$$3_{ol} < 1_o \prec 3_{or} < 2_o; \quad 2_e < 3_{el} \prec 1_e < 3_{er}; \quad 3_{el} \prec 0_e,$$

which is a summary of the order relations appeared in (38)-(41). Since $<$ is stronger than \prec , by transitivity, we get (43) and (44).

4.1.2. Induced orders on Ω_∞ .

Define a relation \preceq (\leq) on Ω_∞ as follows. For any $\omega \in \Omega_\infty$, let $\omega \preceq \omega$ ($\omega \leq \omega$). Given $\omega, \hat{\omega} \in \Omega_\infty$ and $\omega \neq \hat{\omega}$. Assume n is such that $\omega|_{n-1} = \hat{\omega}|_{n-1}$ and $\omega_n \neq \hat{\omega}_n$. Let $\omega \prec \hat{\omega}$ ($\omega < \hat{\omega}$) if $\omega_n \prec \hat{\omega}_n$ ($\omega_n < \hat{\omega}_n$).

Lemma 4.2. *Both \preceq and \leq are partial orders on Ω_∞ . Moreover, \preceq is a total order. If $\omega < \hat{\omega}$, then $\omega \prec \hat{\omega}$.*

Proof. It is routine to check that both \preceq and \leq are partial orders.

Now we show that \preceq is a total order. Assume $\omega, \hat{\omega} \in \Omega_\infty$ and $\omega \neq \hat{\omega}$. Let $n \geq 0$ be the minimal integer such that $\omega_n \neq \hat{\omega}_n$. If $n = 0$, then $\omega_0 \neq \hat{\omega}_0$. By (11), $\{\omega_0, \hat{\omega}_0\} = \{3_{el}, 0_e\}$. By (43), we can compare ω_0 and $\hat{\omega}_0$. Then by the definition, we can compare ω and $\hat{\omega}$. If $n \geq 1$, then $\omega \wedge \hat{\omega} = \omega|_{n-1} = \hat{\omega}|_{n-1}$ and by (10), $\omega_{n-1} = \hat{\omega}_{n-1} \in \{0_e, 0_o, 1_e, 1_o, 2_e, 2_o\}$. Now by (10) and (43), we can compare ω_n and $\hat{\omega}_n$. So, we can compare ω and $\hat{\omega}$.

Since in \mathcal{A} , $\alpha < \beta$ implies $\alpha \prec \beta$, by the definitions, the last statement holds. \square

Take $\omega, \hat{\omega} \in \Omega_\infty$. Assume $\omega \prec \hat{\omega}$. Define the *open interval* $(\omega, \hat{\omega})$ as

$$(\omega, \hat{\omega}) := \{\tilde{\omega} \in \Omega_\infty : \omega \prec \tilde{\omega} \prec \hat{\omega}\}.$$

We call $(\omega, \hat{\omega})$ a *gap* of Ω_∞ if $(\omega, \hat{\omega}) = \emptyset$. We also say that ω ($\hat{\omega}$) is the *left (right) edge* of the gap.

4.2. Characterization of the gaps of Ω_∞ .

We denote the set of the gaps of Ω_∞ by \mathcal{G} . The following result is a symbolic version of Theorem 1.8:

Theorem 4.3. *For Ω_∞ , we have*

i) $\omega_* \preceq \omega \preceq \omega^*$ for any $\omega \in \Omega_\infty$.

ii) (Type-I gaps): *There exists a bijection $\ell^o : \mathcal{E}_l^o \rightarrow \mathcal{E}_r^o \setminus \{\omega_*\}$ such that $\omega < \ell^o(\omega)$ for any $\omega \in \mathcal{E}_l^o$ and*

$$(45) \quad \mathcal{G}_I^o := \{(\omega, \ell^o(\omega)) : \omega \in \mathcal{E}_l^o\} \subset \mathcal{G}.$$

There exists a bijection $\ell^e : \mathcal{E}_r^e \rightarrow \mathcal{E}_l^e \setminus \{\omega^\}$ such that $\ell^e(\omega) < \omega$ for any $\omega \in \mathcal{E}_r^e$ and*

$$(46) \quad \mathcal{G}_I^e := \{(\ell^e(\omega), \omega) : \omega \in \mathcal{E}_r^e\} \subset \mathcal{G}.$$

iii) (Type-II gaps): *There exists a bijection $\ell : \tilde{\mathcal{E}}_l \rightarrow \tilde{\mathcal{E}}_r$ such that $\omega \prec \ell(\omega)$ for any $\omega \in \tilde{\mathcal{E}}_l$ and*

$$(47) \quad \mathcal{G}_{II} := \{(\omega, \ell(\omega)) : \omega \in \tilde{\mathcal{E}}_l\} \subset \mathcal{G}.$$

i) $\mathcal{G} = \mathcal{G}_I \cup \mathcal{G}_{II}$, where $\mathcal{G}_I := \mathcal{G}_I^o \cup \mathcal{G}_I^e$.

We will prove Theorem 4.3 by using the map Π defined by (16). But at first, we need to know how to define ℓ, ℓ^o and ℓ^e .

4.2.1. Definitions and bijectivities of ℓ, ℓ^o, ℓ^e .

At first, we define $\ell : \tilde{\mathcal{E}}_l \rightarrow \tilde{\mathcal{E}}_r$. Assume $\omega \in \tilde{\mathcal{E}}_l$. Then there exists n such that $\omega = \omega|_n 3_{er} (0_o 3_{er})^\infty$ with $\omega_n \neq 0_o$; or $\omega = \omega|_n (0_o 3_{er})^\infty$ with $\omega_n \neq 3_{er}$. By tracing the graph \mathbb{G} backward, in the former case, $\omega_n = 1_o$ and hence $n \geq 1, \omega_{n-1} = 0_e$ or 2_e ; in the latter




FIGURE 3. Two types of gaps

case, $\omega_n = 3_{el}$ and hence $n \geq 2, \omega_{n-1} = 0_o$ or 2_o , or $n = 0$. Now define

$$(48) \quad \ell(\omega) := \begin{cases} \omega|_{n-1} 3_{or} (0_e 3_{ol})^\infty, & \text{if } \omega = \omega|_{n-1} 1_o 3_{er} (0_o 3_{er})^\infty, n \geq 1 \\ \omega|_{n-1} 1_e 3_{ol} (0_e 3_{ol})^\infty, & \text{if } \omega = \omega|_{n-1} 3_{el} (0_o 3_{er})^\infty, n \geq 2 \\ (0_e 3_{ol})^\infty, & \text{if } \omega = 3_{el} (0_o 3_{er})^\infty. \end{cases}$$

See Figure 3 (b) for an illustration of the definition of ℓ and for some intuition of a gap of ∞_{II} -type.

Next, we define $\ell^o : \mathcal{E}_l^o \rightarrow \mathcal{E}_r^o \setminus \{\omega_*\}$. Assume $\omega \in \mathcal{E}_l^o$. Then there exists n such that either $\omega = \omega|_n (2_o 1_e)^\infty$ with $\omega_n \neq 1_e$; or $\omega = \omega|_n 1_e (2_o 1_e)^\infty$ with $\omega_n \neq 2_o$. By tracing the graph \mathbb{G} backward, in the former case, $\omega_n = 2_e$ and hence $n \geq 2, \omega_{n-1} = 1_o$ or 2_o ; in the latter case, $\omega_n = 0_o$ and hence $n \geq 1$. Now define

$$(49) \quad \ell^o(\omega) := \begin{cases} \omega|_{n-2} 1_o 3_{er} (0_o 3_{el})^\infty, & \text{if } \omega = \omega|_{n-2} 1_o 2_e (2_o 1_e)^\infty, n \geq 2 \\ \omega|_{n-2} 2_o 3_{el} (0_o 3_{el})^\infty, & \text{if } \omega = \omega|_{n-2} 2_o 2_e (2_o 1_e)^\infty, n \geq 2 \\ \omega|_{n-1} 0_o 3_{er} (0_o 3_{el})^\infty, & \text{if } \omega = \omega|_{n-1} 0_o 1_e (2_o 1_e)^\infty, n \geq 1 \end{cases}$$

See Figure 3 (a) for an illustration of the definition of ℓ^o and for some intuition of a gap of ∞_I -type.

Now, we define $\ell^e : \mathcal{E}_r^e \rightarrow \mathcal{E}_l^e \setminus \{\omega_*\}$. Assume $\omega \in \mathcal{E}_r^e$. Then there exists n such that either $\omega = \omega|_n (2_e 1_o)^\infty$ with $\omega_n \neq 1_o$; or $\omega = \omega|_n 1_o (2_e 1_o)^\infty$ with $\omega_n \neq 2_e$. By tracing the graph \mathbb{G} backward, in the former case, $\omega_n = 2_o$ and hence $n \geq 3, \omega_{n-1} = 1_e$ or 2_e ; in the latter case, $\omega_n = 0_e$ and hence $n \geq 0$. Now define

$$(50) \quad \ell^e(\omega) := \begin{cases} \omega|_{n-2} 1_e 3_{ol} (0_e 3_{or})^\infty, & \text{if } \omega = \omega|_{n-2} 1_e 2_o (2_e 1_o)^\infty, n \geq 3 \\ \omega|_{n-2} 2_e 3_{or} (0_e 3_{or})^\infty, & \text{if } \omega = \omega|_{n-2} 2_e 2_o (2_e 1_o)^\infty, n \geq 3 \\ \omega|_{n-1} 0_e 3_{ol} (0_e 3_{or})^\infty, & \text{if } \omega = \omega|_{n-1} 0_e 1_o (2_e 1_o)^\infty, n \geq 0 \end{cases}$$

Lemma 4.4. *i) $\ell : \tilde{\mathcal{E}}_l \rightarrow \tilde{\mathcal{E}}_r$ is a bijection and $\omega \prec \ell(\omega)$ for any $\omega \in \tilde{\mathcal{E}}_l$.*

ii) $\ell^o : \mathcal{E}_l^o \rightarrow \mathcal{E}_r^o \setminus \{\omega_\}$ is a bijection and $\omega < \ell^o(\omega)$ for any $\omega \in \mathcal{E}_l^o$.*

iii) $\ell^e : \mathcal{E}_r^e \rightarrow \mathcal{E}_l^e \setminus \{\omega^*\}$ is a bijection and $\ell^e(\omega) < \omega$ for any $\omega \in \mathcal{E}_r^e$.

Proof. i) By (48) and (43), we have $\omega \prec \ell(\omega)$.

To show the bijection of ℓ , we simply reverse the definition of ℓ and construct another map $\eta : \tilde{\mathcal{E}}_r \rightarrow \tilde{\mathcal{E}}_l$, then check η is the inverse of ℓ . Assume $\omega \in \tilde{\mathcal{E}}_r$. Then either there exists $n \geq 1$ such that $\omega = \omega|_n(0_e 3_{ol})^\infty$ with $\omega_n \neq 3_{ol}$; or there exists $n \geq 0$ such that $\omega = \omega|_n 3_{ol}(0_e 3_{ol})^\infty$ with $\omega_n \neq 0_e$; or $\omega = (0_e 3_{ol})^\infty$. By tracing the graph \mathbb{G} backward, in the first case, $\omega_n = 3_{or}$ and hence $\omega_{n-1} = 0_e$ or 2_e . In the second case, $\omega_n = 1_e$ and hence $n \geq 2, \omega_{n-1} = 0_o$ or 2_o , Now define

$$\eta(\omega) := \begin{cases} \omega|_{n-1} 1_o 3_{er} (0_o 3_{er})^\infty, & \text{if } \omega = \omega|_{n-1} 3_{or} (0_e 3_{ol})^\infty, n \geq 1 \\ \omega|_{n-1} 3_{el} (0_o 3_{er})^\infty, & \text{if } \omega = \omega|_{n-1} 1_e 3_{ol} (0_e 3_{ol})^\infty, n \geq 2 \\ 3_{el} (0_o 3_{er})^\infty, & \text{if } \omega = (0_e 3_{ol})^\infty. \end{cases}$$

One check directly that $\eta \circ \ell = \text{Id}_{\tilde{\mathcal{E}}_l}$ and $\ell \circ \eta = \text{Id}_{\tilde{\mathcal{E}}_r}$. So ℓ is a bijection.

ii) and iii) can be proven similarly, we omit the proof. \square

4.2.2. The property of the map $\Pi : \Omega_\infty \rightarrow \Sigma_\infty$.

Recall that Π is defined by (16). We treat the map Π more precisely as follows.

Define the *label-assigning map* $\mathcal{L} : \{\alpha\beta : \alpha \rightarrow \beta\} \rightarrow \{\emptyset, 0, 1, 00, 01, 10, 11\}$ as follows: $\mathcal{L}(\alpha\beta)$ is the label of the edge $\alpha\beta$.

By (38)-(41), the following holds:

$$(51) \quad \alpha \rightarrow \beta, \beta' \text{ and } \beta < \beta' \Rightarrow \mathcal{L}(\alpha\beta) = 0*; \quad \mathcal{L}(\alpha\beta') = 1*.$$

Define a map $\Pi_* : \Omega_* \rightarrow \Sigma_*$ as

$$(52) \quad \Pi_*(w) := i(w)\mathcal{L}(w_0 w_1)\mathcal{L}(w_1 w_2) \cdots \mathcal{L}(w_{n-1} w_n), \quad w \in \Omega_n,$$

where $i(w) = \emptyset$ if $w_0 = 0_e$ and $i(w) = 0$ if $w_0 = 3_{el}$.

By (37)-(40) and (16), it is direct to check that

$$(53) \quad \Pi_*(\omega|_n) \triangleleft \Pi(\omega); \quad |\Pi_*(\omega|_n)| = n \text{ or } n+1; \quad \Pi(\omega) = \lim_{n \rightarrow \infty} \Pi_*(\omega|_n).$$

Define

$$(54) \quad \Sigma_*^o := \bigcup_{n \geq 0} \Sigma_{2n+1}; \quad \Sigma_*^e := \bigcup_{n \geq 0} \Sigma_{2n}; \quad \Sigma_\infty^{(2)} := \{\sigma(01)^\infty : \sigma \in \Sigma_*\}.$$

We endow Σ_∞ with the lexicographical order \leq .

Proposition 4.5. i) For any $\omega \in \tilde{\mathcal{E}}_l$, we have $\Pi(\omega) = \Pi(\ell(\omega))$. Consequently

$$(55) \quad \Pi(\tilde{\mathcal{E}}_l) = \Pi(\tilde{\mathcal{E}}_r) \subset \Sigma_\infty^{(2)}.$$

ii) Assume $\sigma \in \Sigma_\infty$. If $\#\Pi^{-1}(\{\sigma\}) \geq 2$, then there exists $\omega \in \tilde{\mathcal{E}}_l$ such that

$$\Pi^{-1}(\{\sigma\}) = \{\omega, \ell(\omega)\}.$$

Consequently, $\#\Pi^{-1}(\{\sigma\}) \geq 2$ if and only if $\sigma \in \Pi(\tilde{\mathcal{E}}_l)$.

iii) Π is surjective and order-preserving.

$$(56) \quad \Pi(\omega_*) = 0^\infty; \quad \Pi(\omega^*) = 1^\infty.$$

Moreover, $\Pi : \Omega_\infty \setminus \tilde{\mathcal{E}}_r \rightarrow \Sigma_\infty$ is bijective.

iv) If $\omega \in \mathcal{E}_l^o$, then there exists a unique $\sigma \in \Sigma_*^o$ such that

$$(57) \quad \Pi(\omega) = \sigma 01^\infty; \quad \Pi(\ell^o(\omega)) = \sigma 10^\infty.$$

If $\omega \in \mathcal{E}_r^e$, then there exists a unique $\sigma \in \Sigma_*^e$ such that

$$(58) \quad \Pi(\omega) = \sigma 10^\infty; \quad \Pi(\ell^e(\omega)) = \sigma 01^\infty.$$

The following restrictions of Π are bijections:

$$(59) \quad \begin{cases} \Pi_{ol} : \mathcal{E}_l^o \rightarrow \{\sigma 01^\infty : \sigma \in \Sigma_*^o\}; & \Pi_{or} : \mathcal{E}_r^o \setminus \{\omega_*\} \rightarrow \{\sigma 10^\infty : \sigma \in \Sigma_*^o\} \\ \Pi_{er} : \mathcal{E}_r^e \rightarrow \{\sigma 10^\infty : \sigma \in \Sigma_*^e\}; & \Pi_{el} : \mathcal{E}_l^e \setminus \{\omega^*\} \rightarrow \{\sigma 01^\infty : \sigma \in \Sigma_*^e\} \end{cases}$$

v) $\Pi : \tilde{\mathcal{E}}_l \cup \mathcal{F} \rightarrow \Sigma_\infty^{(2)}$ is bijective. Consequently

$$(60) \quad \tilde{\mathcal{E}}_l \cup \tilde{\mathcal{E}}_r \cup \mathcal{F} = \Pi^{-1}(\Sigma_\infty^{(2)}).$$

Proof. i) Assume $\omega \in \tilde{\mathcal{E}}_l$. By (48), there are three cases.

If $\omega = \omega|_{n-1} 1_o 3_{er} (0_o 3_{er})^\infty$, then $\omega_{n-1} = 0_e$ or 2_e and $\ell(\omega) = \omega|_{n-1} 3_{or} (0_e 3_{ol})^\infty$. By tracing on the graph \mathbb{G} , for $m \geq n$,

$$(61) \quad \begin{cases} \Pi_*(\omega|_m) = \Pi_*(\omega|_{n-1}) \delta (10)^{[(m+1-n)/2]}, \\ \Pi_*(\ell(\omega)|_m) = \Pi_*(\omega|_{n-1}) \delta 1 (01)^{[(m-n)/2]}. \end{cases}$$

Here $\delta = 1$ if $\omega_{n-1} = 0_e$ and $\delta = 0$ if $\omega_{n-1} = 2_e$.

If $\omega = \omega|_{n-1} 3_{el} (0_o 3_{er})^\infty$, then $\omega_{n-1} = 0_o$ or 2_o and $\ell(\omega) = \omega|_{n-1} 1_e 3_{ol} (0_e 3_{ol})^\infty$. By tracing on the graph \mathbb{G} , for $m \geq n$,

$$(62) \quad \begin{cases} \Pi_*(\omega|_m) = \Pi_*(\omega|_{n-1}) \delta 0 (10)^{[(m-n)/2]}, \\ \Pi_*(\ell(\omega)|_m) = \Pi_*(\omega|_{n-1}) \delta (01)^{[(m+1-n)/2]}. \end{cases}$$

Here $\delta = 0$ if $\omega_{n-1} = 0_o$ and $\delta = 1$ if $\omega_{n-1} = 2_o$.

If $\omega = 3_{el} (0_o 3_{er})^\infty$, then $\ell(\omega) = (0_e 3_{ol})^\infty$. Notice that, in this case $i(\omega) = 0$ and $i(\ell(\omega)) = \emptyset$. By tracing on the graph \mathbb{G} , for $m \geq 0$,

$$(63) \quad \begin{cases} \Pi_*(\omega|_m) = 0 (10)^{[m/2]}, \\ \Pi_*(\ell(\omega)|_m) = (01)^{[(m+1)/2]}. \end{cases}$$

In all cases, by (53) we have

$$\Pi(\omega) = \lim_{m \rightarrow \infty} \Pi_*(\omega|_m) = \lim_{m \rightarrow \infty} \Pi_*(\ell(\omega)|_m) = \Pi(\ell(\omega)).$$

Combine with Lemma 4.4 i), (55) holds.

ii) Assume $\#\Pi^{-1}(\{\sigma\}) \geq 2$. Take any $\omega, \hat{\omega} \in \Pi^{-1}(\{\sigma\})$. WLOG, we assume $\omega \prec \hat{\omega}$. Let $w := \omega \wedge \hat{\omega}$, and assume $w \in \Omega_{n-1}$. Then $\omega_n \prec \hat{\omega}_n$. By (51), $\omega_n \not\prec \hat{\omega}_n$. So either $n \geq 1$ and

$$(w_{n-1}, \omega_n, \hat{\omega}_n) \in \{(0_o, 3_{el}, 1_e), (2_o, 3_{el}, 1_e), (0_e, 1_o, 3_{or}), (2_e, 1_o, 3_{or})\};$$

or $n = 0$, $w = \emptyset$ and $\omega_0 = 3_{el}, \hat{\omega}_0 = 0_e$.

At first assume $n \geq 1$ and $(w_{n-1}, \omega_n, \hat{\omega}_n) = (0_o, 3_{el}, 1_e)$. Then

$$\Pi_*(\omega|_n) = \Pi_*(w3_{el}) = \tau 00; \quad \Pi_*(\hat{\omega}|_n) = \Pi_*(w1_e) = \tau 0, \quad \text{where } \tau = \Pi_*(w).$$

Since $\Pi(\omega) = \sigma = \Pi(\hat{\omega})$, we have $\mathcal{L}(\hat{\omega}_n \hat{\omega}_{n+1}) = \mathcal{L}(1_e \hat{\omega}_{n+1}) = 0*$. By checking \mathbb{G} , the only possibility is $\hat{\omega}_{n+1} = 3_{ol}$. So $\mathcal{L}(\hat{\omega}_n \hat{\omega}_{n+1}) = 01$ and $\Pi_*(\hat{\omega}|_{n+1}) = \tau 001$. Again by $\Pi(\omega) = \Pi(\hat{\omega})$, we have

$$\mathcal{L}(\omega_n \omega_{n+1}) \mathcal{L}(\omega_{n+1} \omega_{n+2}) = \mathcal{L}(3_{el} 0_o) \mathcal{L}(0_o \omega_{n+2}) = \mathcal{L}(0_o \omega_{n+2}) = 1 * .$$

By checking \mathbb{G} , the only possibility is $\omega_{n+1} = 3_{er}$. As a consequence $\Pi_*(\omega|_{n+2}) = \tau 0010$. This process can continue infinitely. By induction, we get

$$\omega = w 3_{el} (0_o 3_{er})^\infty \in \tilde{\mathcal{E}}_l; \quad \hat{\omega} = w 1_e 3_{ol} (0_e 3_{ol})^\infty \in \tilde{\mathcal{E}}_r.$$

By (48), $\hat{\omega} = \ell(\omega)$.

For the other four cases, the same proof shows that

$$\omega \in \tilde{\mathcal{E}}_l; \quad \hat{\omega} \in \tilde{\mathcal{E}}_r \quad \text{and} \quad \hat{\omega} = \ell(\omega).$$

This implies that

$$\Pi^{-1}(\{\sigma\}) \subset \tilde{\mathcal{E}}_l \cup \tilde{\mathcal{E}}_r, \quad \#\tilde{\mathcal{E}}_l \cap \Pi^{-1}(\{\sigma\}), \#\tilde{\mathcal{E}}_r \cap \Pi^{-1}(\{\sigma\}) \leq 1.$$

So the first statement follows.

If $\#\Pi^{-1}(\{\sigma\}) \geq 2$, then $\Pi^{-1}(\{\sigma\}) = \{\omega, \ell(\omega)\}$ for some $\omega \in \tilde{\mathcal{E}}_l$, hence $\sigma \in \Pi(\tilde{\mathcal{E}}_l)$. If $\sigma \in \Pi(\tilde{\mathcal{E}}_l)$, then by i), $\sigma \in \Pi(\tilde{\mathcal{E}}_r)$, so $\#\Pi^{-1}(\{\sigma\}) \geq 2$. Thus the second statement holds.

iii) We will prove the Surjectivity of Π in Sec. 5.1.3 by using the coding of \mathcal{B}_n .

Now we show that Π preserves the order. Given $\omega \prec \hat{\omega}$. Then there exists $n \geq 0$ such that $\omega|_{n-1} = \hat{\omega}|_{n-1} =: w$ and $\omega|_n \prec \hat{\omega}|_n$. If $\omega|_n < \hat{\omega}|_n$, then by (51),

$$\mathcal{L}(w_{n-1} \omega_n) = 0*; \quad \mathcal{L}(w_{n-1} \hat{\omega}_n) = 1 * .$$

Hence $\Pi(\omega) < \Pi(\hat{\omega})$.

Now assume $\omega|_n \not\prec \hat{\omega}|_n$. Then

$$(\omega_n, \hat{\omega}_n) \in \{(3_{el}, 1_e), (1_o, 3_{or}), (3_{el}, 0_e)\}.$$

At first assume $(\omega_n, \hat{\omega}_n) = (3_{el}, 1_e)$. So either $\omega = \omega|_{n-1} 3_{el} (0_o 3_{er})^\infty := \tilde{\omega}$ or there exists $m \geq 0$ such that $\omega = \omega|_{n-1} 3_{el} (0_o 3_{er})^m 0_o \alpha \dots$ with $\alpha = 1_e$ or 3_{el} . Since

$$\mathcal{L}(0_o 1_e) = 0; \quad \mathcal{L}(0_o 3_{el}) = 00; \quad \mathcal{L}(0_o 3_{er}) = 10,$$

We conclude that $\Pi(\omega) \leq \Pi(\tilde{\omega})$.

The same argument shows that $\Pi(\ell(\tilde{\omega})) \leq \Pi(\hat{\omega})$. So by i) we have

$$\Pi(\omega) \leq \Pi(\tilde{\omega}) = \Pi(\ell(\tilde{\omega})) \leq \Pi(\hat{\omega}).$$

For the other two cases, the same argument shows that $\Pi(\omega) \leq \Pi(\hat{\omega})$. Hence Π is order preserving.

By (14) and the definition of Π , one get (56) by tracing the graph \mathbb{G} .

By ii) $\Pi^{-1}(\Pi(\tilde{\mathcal{E}}_l)) = \tilde{\mathcal{E}}_l \cup \tilde{\mathcal{E}}_r$ and $\Pi : \tilde{\mathcal{E}}_l \rightarrow \Pi(\tilde{\mathcal{E}}_l)$ is bijective. Since Π is surjective, the restriction $\Pi : \Omega_\infty \setminus (\tilde{\mathcal{E}}_l \cup \tilde{\mathcal{E}}_r) \rightarrow \Sigma_\infty \setminus \Pi(\tilde{\mathcal{E}}_l)$ is also surjective. Again by ii), this restriction is injective. So it is bijective. Combine these two bijections, we conclude that $\Pi : \Omega_\infty \setminus \tilde{\mathcal{E}}_r \rightarrow \Sigma_\infty$ is a bijection.

iv) (57) and (58) follow from (49) and (50) by using the definition of Π and tracing on \mathbb{G} . Now we show that all the restriction maps in (59) are bijections. By iii), they are all injections. Now we show that they are surjections. We take the first one as an example. Since Π is surjective, for each $\sigma \in \Sigma_*^o$, there exists some $\omega \in \Omega_\infty$ such that $\Pi(\omega) = \sigma 0 1^\infty$. We only need to show that $\omega \in \mathcal{E}_l^o$. We delete all the edges of \mathbb{G} with label containing 0. Then only two connected subgraphs are left:

$$0_e \xrightarrow{11} 3_{or} \xrightarrow{\emptyset} 0_e; \quad 1_e \xrightarrow{1} 2_o \xrightarrow{1} 1_e.$$

Each of these subgraphs can generates 1^∞ and 1^∞ can only be generated by these two graphs. Thus $\omega \in \mathcal{E}_l^o$ or \mathcal{E}_l^e . However, by (56) and (58), if $\omega \in \mathcal{E}_l^e$, then $\Pi(\omega) = 1^\infty$ or $\sigma 0 1^\infty$ with $\sigma \in \Sigma_*^e$. So we conclude that $\omega \in \mathcal{E}_l^o$. That is, Π_{ol} is surjective. The other three cases can be proven by exactly the same way.

v) By (13) and tracing on \mathbb{G} , it is seen that $\Pi(\mathcal{F}) \subset \Sigma_\infty^{(2)}$. Combine with (55), we have $\Pi(\tilde{\mathcal{E}}_l \cup \mathcal{F}) \subset \Sigma_\infty^{(2)}$. By iii), The restriction of Π on $\tilde{\mathcal{E}}_l \cup \mathcal{F}$ is injective. Thus we only need to show that $\Sigma_\infty^{(2)} \subset \Pi(\tilde{\mathcal{E}}_l \cup \mathcal{F})$.

Fix any $\sigma \in \Sigma_\infty^{(2)}$. Since Π is surjective, there exists $\omega \in \Omega_\infty$ such that $\Pi(\omega) = \sigma$.

At first, we claim that 1_e does not appear in ω infinitely often (i.o.). Indeed if otherwise, by tracing backward on \mathbb{G} , one of the following words must appear in ω i.o.:

$$1_o 3_{er} 0_o 1_e; \quad 0_o 3_{er} 0_o 1_e; \quad 2_o 3_{el} 0_o 1_e; \quad 0_o 3_{el} 0_o 1_e; \quad 2_e 2_o 1_e; \quad 1_e 2_o 1_e.$$

By applying \mathcal{L} to these words, we see that for the first four cases, 00 will appear and for the last two cases, 11 will appear. So 00 or 11 will appears in $\Pi(\omega)$ i.o., a contradiction.

The same proof shows that 1_o also only appear in ω finite times. Now we go back to the graph \mathbb{G} and delete 1_o and 1_e and all the edges connected to 1_o and 1_e . We also delete $0_e 3_{or}$ and $0_o 3_{el}$ since they have labels 11 and 00 , respectively. Now the remaining part of the graph has only three connected components:

$$0_e \xrightarrow{01} 3_{ol} \xrightarrow{\emptyset} 0_e; \quad 0_o \xrightarrow{10} 3_{er} \xrightarrow{\emptyset} 0_o; \quad 2_o \xrightarrow{0} 2_e \xrightarrow{1} 2_o,$$

and $(01)^\infty$ can be realized by them and only by them. So ω can only be eventually $(ab)^\infty$, with

$$ab \in \{0_o 3_{er}, 0_e 3_{ol}, 2_e 2_o\}.$$

That is, $\omega \in \tilde{\mathcal{E}}_l \cup \tilde{\mathcal{E}}_r \cup \mathcal{F}$. Now combine with i), $\Pi(\omega) \in \Pi(\tilde{\mathcal{E}}_l \cup \mathcal{F})$. Hence, $\Sigma_\infty^{(2)} \subset \Pi(\tilde{\mathcal{E}}_l \cup \mathcal{F})$. Then the restriction of Π on $\tilde{\mathcal{E}}_l \cup \mathcal{F}$ is bijective.

Now by i), (60) follows. □

Remark 4.6. Recall that we have defined an order \preceq on Σ_* by (34). If $(\omega, \hat{\omega}) \in \mathcal{G}_{II}$, by (61)-(63), we conclude that for any $m, m' > |\omega \wedge \hat{\omega}|$,

$$\Pi_*(\omega|_m) \prec \Pi_*(\hat{\omega}|_{m'}).$$

4.2.3. Proof of Theorem 4.3.

Before giving the formal proof, let us explain the mechanism to form a gap in Ω_∞ . At first, if $(\sigma, \hat{\sigma})$ is gap of Σ_∞ , then it can be lifted to a gap of Ω_∞ via Π . It is seen that the gap of Σ_∞ has the form $(\tau 01^\infty, \tau 10^\infty)$. Secondly, if $\Pi^{-1}(\sigma)$ has more than one elements, then by Proposition 4.5 ii), it has exactly two elements, and the elements form a gap. These are the only two ways to form a gap.

Proof of Theorem 4.3. i) Fix any $\omega \neq \omega_*$. By (56), $\Pi(\omega_*) = 0^\infty \notin \Sigma_\infty^{(2)}$. By Proposition 4.5 ii), $\Pi(\omega) \neq 0^\infty$. We claim that $\omega_* \prec \omega$. Indeed, if otherwise, $\omega \preceq \omega_*$, then $\Pi(\omega) \leq \Pi(\omega_*) = 0^\infty$. Since 0^∞ is the minimum of Σ_∞ , we have $\Pi(\omega) = 0^\infty$, a contradiction. So $\omega_* \prec \omega$.

The same argument shows that if $\omega \neq \omega^*$, then $\omega \prec \omega^*$. So the result follows.

ii) Combine with Lemma 4.4, we only need to show (45) and (46). Fix $\omega \in \mathcal{E}_l^\circ$. By (57), there exists $\sigma \in \Sigma_*$ such that $\Pi(\omega) = \sigma 01^\infty =: \tau$ and $\Pi(\ell^\circ(\omega)) = \sigma 10^\infty =: \hat{\tau}$. It is well-known that $(\tau, \hat{\tau})$ is a gap of Σ_∞ . We claim that $(\omega, \ell^\circ(\omega))$ is a gap of Ω_∞ . Indeed if otherwise, there exists $\tilde{\omega}$ such that $\omega \prec \tilde{\omega} \prec \ell^\circ(\omega)$, so

$$\tau \leq \Pi(\tilde{\omega}) \leq \hat{\tau}.$$

So either $\Pi(\tilde{\omega}) = \tau$ or $\hat{\tau}$. In the former case, $\Pi^{-1}(\{\tau\}) \supset \{\omega, \tilde{\omega}\}$. So by Proposition 4.5 ii), $\tau \in \Sigma_\infty^{(2)}$, which is a contradiction. In the latter case, we get a contradiction by the same reasoning. So $(\omega, \ell^\circ(\omega))$ is a gap of Ω_∞ and (45) holds.

The same proof shows that (46) holds.

iii) Combine with Lemma 4.4, we only need to show (47). Take $\omega \in \tilde{\mathcal{E}}_l$, we claim that $(\omega, \ell(\omega))$ is a gap. Indeed if otherwise, there exists $\tilde{\omega}$ such that $\omega \prec \tilde{\omega} \prec \ell(\omega)$, so

$$\Pi(\omega) \leq \Pi(\tilde{\omega}) \leq \Pi(\ell(\omega)).$$

By Proposition 4.5 i), $\Pi(\omega) = \Pi(\ell(\omega)) =: \sigma$. So $\Pi^{-1}(\{\sigma\}) \supset \{\omega, \tilde{\omega}, \ell(\omega)\}$, which contradicts with Proposition 4.5 ii). So (47) holds.

iv) Now assume $(\omega, \hat{\omega}) \in \mathcal{G}$. Then either $\Pi(\omega) = \Pi(\hat{\omega}) =: \sigma$, or $\Pi(\omega) < \Pi(\hat{\omega})$.

In the former case, by Proposition 4.5 ii), $\omega \in \tilde{\mathcal{E}}_l$ and $\hat{\omega} = \ell(\omega)$. So $(\omega, \hat{\omega}) \in \mathcal{G}_{II}$.

In the latter case, we claim that $(\Pi(\omega), \Pi(\hat{\omega}))$ is a gap of Σ_∞ . Indeed, if otherwise, there exists $\sigma \in \Sigma_\infty$ such that $\Pi(\omega) < \sigma < \Pi(\hat{\omega})$. Assume $\Pi(\tilde{\omega}) = \sigma$, then $\tilde{\omega} \neq \omega, \hat{\omega}$. Since Π preserve the order, we must have $\omega \prec \tilde{\omega} \prec \hat{\omega}$, a contradiction.

So there exists $\sigma \in \Sigma_*$ such that $\Pi(\omega) = \sigma 01^\infty$ and $\Pi(\hat{\omega}) = \sigma 10^\infty$. By Proposition 4.5 iv) and ii), either $\omega \in \mathcal{E}_l^\circ, \hat{\omega} = \ell^\circ(\omega)$, or $\hat{\omega} \in \mathcal{E}_r^e, \omega = \ell^e(\hat{\omega})$. So $(\omega, \hat{\omega}) \in \mathcal{G}_I$.

As a result, $\mathcal{G} \subset \mathcal{G}_I \cup \mathcal{G}_{II}$. Combine with ii) and iii), we get the equality. \square

We end with the following lemma, which will be used to show that π is order-preserving.

Lemma 4.7. *Assume $\omega \prec \hat{\omega}$ but $\omega \not\prec \hat{\omega}$. Then there exists $(\tau, \hat{\tau}) \in \mathcal{G}_{II}$ such that $\omega \preceq \tau$ and $\hat{\tau} \preceq \hat{\omega}$. Moreover if $\omega \neq \tau(\hat{\tau} \neq \hat{\omega})$, then $\omega < \tau(\hat{\tau} < \hat{\omega})$.*

Proof. Assume $n \geq 0$ is such that $\omega|_{n-1} = \hat{\omega}|_{n-1} =: w$ and $\omega|_n \prec \hat{\omega}|_n$. Since $\omega \not\prec \hat{\omega}$,

$$(\omega_n, \hat{\omega}_n) \in \{(3_{el}, 1_e), (1_o, 3_{or}), (3_{el}, 0_e)\}.$$

At first assume $(\omega_n, \hat{\omega}_n) = (3_{el}, 1_e)$. So either $\omega = \omega|_{n-1}3_{el}(0_o3_{er})^\infty =: \tau \in \tilde{\mathcal{E}}_l$ or there exists $m \geq 0$ such that $\omega = \omega|_{n-1}3_{el}(0_o3_{er})^m 0_o \alpha \cdots$ with $\alpha = 1_e$ or 3_{el} . In the latter case, by (44), $\omega < \tau$. The same argument shows that $\ell(\tau) \preceq \hat{\omega}$ and if $\hat{\omega} \neq \ell(\tau)$, then $\ell(\tau) < \hat{\omega}$.

For the case that $(\omega_n, \hat{\omega}_n) = (1_o, 3_{or})$ or $(3_{el}, 0_e)$, the proof is the same. \square

5. CODING AND THE HAUSDORFF DIMENSION OF THE SPECTRUM

In this section, we prove a weak version of Theorem 1.5, which is enough for estimating the dimension of the spectrum. Then we prove Theorem 1.1.

5.1. Coding of the spectrum.

5.1.1. Coding of \mathcal{B}_n .

We start with coding \mathcal{B}_n by Ω_n . We summarize Lemmas 3.5-3.8 in the following Corollary, the proof of which is by direct checking.

Corollary 5.1. *Assume $B \in \mathcal{B}_n$ has type α . Then B contains a band $\tilde{B} \in \mathcal{B}_{n+1}$ of type β if and only if $\alpha \rightarrow \beta$. Moreover, if $\alpha \rightarrow \beta$, then B contains exact one band $\tilde{B} \in \mathcal{B}_{n+1}$ of type β .*

Now we can code \mathcal{B}_n by Ω_n inductively as follows.

Recall that $\mathcal{B}_0 = \{B_0, B_\emptyset\}$; B_0 has type 3_{el} and B_\emptyset has type 0_e and by (31), $B_0 \prec B_\emptyset$. We code \mathcal{B}_0 by $\Omega_0 = \{3_{el}, 0_e\}$ as

$$I_{3_{el}} := B_0; \quad I_{0_e} := B_\emptyset.$$

Assume we have coded \mathcal{B}_n by Ω_n :

$$\mathcal{B}_n = \{I_w : w \in \Omega_n\}, \quad I_w \text{ has type } w_n \quad \text{and} \quad I_v \neq I_w \text{ if } v \neq w.$$

Take any $u = u_0 \cdots u_{n+1} \in \Omega_{n+1}$, then $u|_n \in \Omega_n$ and $u_n \rightarrow u_{n+1}$. by Corollary 5.1, there exists a unique $\hat{B} \in \mathcal{B}_{n+1}$ with type u_{n+1} such that $\hat{B} \subset I_{u|_n}$. Define $I_u := \hat{B}$. Then

$$\{I_w : w \in \Omega_{n+1}\} \subset \mathcal{B}_{n+1}.$$

Now take any $\tilde{B} \in \mathcal{B}_{n+1}$, assume it has type β . By Proposition 3.9, there exists a unique $B \in \mathcal{B}_n$ such that $\tilde{B} \subset B$. By induction hypothesis, there is a unique $w \in \Omega_n$ such that $B = I_w$. Again by Corollary 5.1, $w_n \rightarrow \beta$ and $\tilde{B} = I_{w\beta}$ and no other $u \in \Sigma_{n+1}$ satisfies $\tilde{B} = I_u$. Thus

$$\mathcal{B}_{n+1} = \{I_w : w \in \Omega_{n+1}\}, \quad I_w \text{ has type } w_{n+1} \quad \text{and} \quad I_v \neq I_w \text{ if } v \neq w.$$

By induction, we finish the coding process.

Combine with Lemmas 3.5-3.8, the above coding process also implies that for $w \in \Omega_*$,

$$(64) \quad I_w = B_{\Pi_*(w)}.$$

By Corollary 5.1, we also have the following: Assume $v \in \Omega_n$ and $w \in \Omega_{n-1}$, then

$$(65) \quad I_v \subset I_w \Leftrightarrow w \triangleleft v.$$

With this coding, we can explain two orders (43) and (44) as follows:

Corollary 5.2. *Assume $w \in \Omega_n$ and $w_n \rightarrow \alpha, \beta$. Then*

$$I_{w\alpha} \prec I_{w\beta} \Leftrightarrow \alpha \prec \beta; \quad I_{w\alpha} < I_{w\beta} \Leftrightarrow \alpha < \beta.$$

The proof is again by applying Lemmas 3.6-3.8 and checking directly.

5.1.2. Coding of σ_λ .

For any $\omega \in \Omega_\infty$, we have $\omega|_n \in \Omega_n$. Hence $I_{\omega|_n} \in \mathcal{B}_n$. By the construction, we have $I_{\omega|_{n+1}} \subset I_{\omega|_n}$. By (25), $|I_{\omega|_n}| \rightarrow 0$. Hence $\bigcap_{n \geq 0} I_{\omega|_n}$ is a singleton. By Corollary 3.10, this point is in σ_λ .

We define the coding map $\pi : \Omega_\infty \rightarrow \sigma_\lambda$ as

$$\pi(\omega) := \bigcap_{n \geq 0} I_{\omega|_n}.$$

We have the following weak version of Theorem 1.5:

Proposition 5.3. $\pi : (\Omega_\infty, \preceq) \rightarrow (\sigma_\lambda, \leq)$ is continuous, surjective and preserves the orders.

To prove it, we need two lemmas.

Lemma 5.4. *Given $\omega, \hat{\omega} \in \Omega_\infty$. If $\omega < \hat{\omega}$, then $\pi(\omega) < \pi(\hat{\omega})$.*

Proof. Assume n is such that $\omega|_{n-1} = \hat{\omega}|_{n-1}$ and $\omega_n < \hat{\omega}_n$. By Corollary 5.2, $I_{\omega|_n} < I_{\hat{\omega}|_n}$. Since $\pi(\omega) \in I_{\omega|_n}$ and $\pi(\hat{\omega}) \in I_{\hat{\omega}|_n}$, we conclude that $\pi(\omega) < \pi(\hat{\omega})$. \square

Lemma 5.5. *If $(\omega, \hat{\omega}) \in \mathcal{G}_{II}$, then $\pi(\omega) \leq \pi(\hat{\omega})$.*

Proof. Write

$$\sigma^{(n)} = \Pi_*(\omega|_n); \quad \hat{\sigma}^{(n)} = \Pi_*(\hat{\omega}|_n).$$

By Remark 4.6, if $n > |\omega \wedge \hat{\omega}|$, then $\sigma^{(n)} \prec \hat{\sigma}^{(n)}$. By Proposition 2.6, $z_{\sigma^{(n)}} < z_{\hat{\sigma}^{(n)}}$.

On the other hand, by (64),

$$\pi(\omega) \in I_{\omega|_n} = B_{\sigma^{(n)}}; \quad \pi(\hat{\omega}) \in I_{\hat{\omega}|_n} = B_{\hat{\sigma}^{(n)}}.$$

By (25), the length of the bands $B_{\sigma^{(n)}}$ and $B_{\hat{\sigma}^{(n)}}$ tends to zero as n tends to infinity. Since $z_{\sigma^{(n)}} \in B_{\sigma^{(n)}}$ and $z_{\hat{\sigma}^{(n)}} \in B_{\hat{\sigma}^{(n)}}$, we have

$$\pi(\omega) = \lim_n z_{\sigma^{(n)}} \leq \lim_n z_{\hat{\sigma}^{(n)}} = \pi(\hat{\omega})$$

So the result follows. \square

Remark 5.6. In Lemma 6.8, we will show that the strict inequality holds. Here we just mention that the proof is highly nontrivial. We need to study the ∞_{II} energies to finally reach the strict inequality.

Proof of Proposition 5.3. By Corollary 3.10, π is surjective.

Assume $|\omega \wedge \hat{\omega}| = n$, then $d(\omega, \hat{\omega}) = 2^{-n}$. We have

$$|\pi(\omega) - \pi(\hat{\omega})| \leq |I_{\omega \wedge \hat{\omega}}|.$$

Combine with (25), π is continuous.

Assume $\omega \prec \hat{\omega}$. If $\omega < \hat{\omega}$, by Lemma 5.4, $\pi(\omega) < \pi(\hat{\omega})$. If $\omega \not< \hat{\omega}$, by Lemma 4.7, there exists $(\tau, \hat{\tau}) \in \mathcal{G}_{II}$ such that $\omega \preceq \tau$ and $\hat{\tau} \preceq \hat{\omega}$, moreover if $\omega \neq \tau$ ($\hat{\tau} \neq \hat{\omega}$), then $\omega < \tau$ ($\hat{\tau} < \hat{\omega}$). By Lemmas 5.4 and 5.5,

$$\pi(\omega) \leq \pi(\tau) \leq \pi(\hat{\tau}) \leq \pi(\hat{\omega}).$$

So π is order-preserving. □

5.1.3. Surjectivity of Π .

As an application of the coding of \mathcal{B}_n , we can show that Π is surjective.

Given $\sigma \in \Sigma_\infty$. For each $n \geq 0$, we always have $B_{\sigma|_n} \in \mathcal{B}_n$. By the coding of \mathcal{B}_n , there exists a unique $w^{(n)} \in \Omega_n$ such that $I_{w^{(n)}} = B_{\sigma|_n}$.

Claim 1: For each $n \geq 2$, $w^{(n-1)} \triangleleft w^{(n)}$ or $w^{(n-2)} \triangleleft w^{(n)}$.

\triangleleft If $B_{\sigma|_n} \notin \mathcal{B}_{n-1}$, then $B_{\sigma|_n} \subset \tilde{B}$ for some $\tilde{B} \in \mathcal{B}_{n-1}$. by Corollary 3.2 iii), $\tilde{B} = B_{\sigma|_{n-1}}$. That is, $I_{w^{(n)}} \subset I_{w^{(n-1)}}$. By (65), $w^{(n-1)} \triangleleft w^{(n)}$.

If $B_{\sigma|_n} \in \mathcal{B}_{n-1}$, then $B_{\sigma|_n} \not\subset B_{\sigma|_{n-1}}$. By Corollary 3.2 ii), $B_{\sigma|_n} \subset B_{\sigma|_{n-2}}$. Assume $\hat{w}^{(n-1)} \in \Omega_{n-1}$ is such that $B_{\sigma|_n} = I_{\hat{w}^{(n-1)}}$. By (65), we have $w^{(n-2)} \triangleleft \hat{w}^{(n-1)}$. We also have $I_{w^{(n)}} = I_{\hat{w}^{(n-1)}}$, again by (65), we have $\hat{w}^{(n-1)} \triangleleft w^{(n)}$. So, $w^{(n-2)} \triangleleft w^{(n)}$. ▷

By the claim, we conclude that

$$(66) \quad \text{for any } n > k \geq 0, \text{ either } w^{(k)} \triangleleft w^{(n)} \text{ or } w^{(k+1)} \triangleleft w^{(n)}.$$

Define inductively an integer sequence $(m_k)_{k \geq 1}$ and a decreasing sequence $(\Xi_k)_{k \geq 1}$ with $\Xi_k \subset \mathbb{N}$ and $\#\Xi_k = \infty$ as follows. Write $\Xi_0 := \mathbb{N}$. By (66), at least one of $\{n \in \Xi_0 : w^{(0)} \triangleleft w^{(n)}\}$ and $\{n \in \Xi_0 : w^{(1)} \triangleleft w^{(n)}\}$ is infinite. If $\{n \in \Xi_0 : w^{(0)} \triangleleft w^{(n)}\}$ is infinite, define $m_1 := 0$. If otherwise, then $\{n \in \Xi_0 : w^{(1)} \triangleleft w^{(n)}\}$ is infinite, define $m_1 := 1$. Define $\Xi_1 := \{n \in \Xi_0 : w^{(m_1)} \triangleleft w^{(n)}\}$. Then $\#\Xi_1 = \infty$.

Assume m_k and Ξ_k has been defined with the desired property. By (66), at least one of $\{n \in \Xi_k : w^{(m_k+1)} \triangleleft w^{(n)}\}$ and $\{n \in \Xi_k : w^{(m_k+2)} \triangleleft w^{(n)}\}$ is infinite. If $\{n \in \Xi_k : w^{(m_k+1)} \triangleleft w^{(n)}\}$ is infinite, define $m_{k+1} := m_k + 1$. If otherwise, then $\{n \in \Xi_k : w^{(m_k+2)} \triangleleft w^{(n)}\}$ is infinite, define $m_{k+1} := m_k + 2$. Define $\Xi_{k+1} := \{n \in \Xi_k : w^{(m_{k+1})} \triangleleft w^{(n)}\}$. Hence $\Xi_{k+1} \subset \Xi_k$ and is infinite.

By induction, we finish the definition.

Now we claim that for any k , $w^{(m_k)} \triangleleft w^{(m_{k+1})}$. Indeed, by the defining process, we have $w^{(m_k)} \triangleleft w^{(n)}$ for any $n \in \Xi_k$. Choose some $n \in \Xi_{k+1}$, then $n \in \Xi_k$ since $\Xi_{k+1} \subset \Xi_k$. Hence we have $w^{(m_k)}, w^{(m_{k+1})} \triangleleft w^{(n)}$. Since $|w^{(m_k)}| < |w^{(m_{k+1})}|$, we conclude that $w^{(m_k)} \triangleleft w^{(m_{k+1})}$.

Let $\omega = \lim_{k \rightarrow \infty} w^{(m_k)}$. Then we have $\Pi(\omega) = \sigma$. \square

It is an interesting question to find a purely combinatorial proof for the surjectivity of Π .

5.2. Lower bound for the Hausdorff dimension of σ_λ .

In this subsection, at first we present a sufficient condition for estimating from below the Hausdorff dimension of the limit set of a SNS. Then we apply it to a sub-SNS of $\{\mathcal{B}_n : n \geq 0\}$ to obtain a lower bound for the Hausdorff dimension of σ_λ .

5.2.1. Dimension estimation for limit set of SNS.

We need the following general result on the dimension of limit set of SNS, which is also useful in its own right.

Proposition 5.7. *Let $\mathcal{I} = \{\mathcal{I}_n : n \geq 0\}$ be a SNS. Assume \mathcal{I} satisfies:*

i) *There exist $\lambda \in (0, 1)$ and $C > 0$ such that*

$$|I| \geq C\lambda^n, \quad \forall n \geq 0, \forall I \in \mathcal{I}_n.$$

ii) *There exists $C' > 0$ such that for any $n, k \geq 0$, and $I, I' \in \mathcal{I}_n$,*

$$\frac{\#\{J \in \mathcal{I}_{n+k} : J \subset I\}}{\#\{J \in \mathcal{I}_{n+k} : J \subset I'\}} \leq C'.$$

Then the limit set satisfies

$$\dim_H A(\mathcal{I}) \geq \liminf_{n \rightarrow \infty} \frac{\log \#\mathcal{I}_n}{-n \log \lambda}.$$

Proof. For any $n \geq 0$, let

$$\kappa_n = \#\mathcal{I}_n, \quad \alpha = \liminf_{n \rightarrow \infty} \sqrt[n]{\kappa_n}, \quad t = -\log \alpha / \log \lambda.$$

If $\alpha = 1$, we are done. Suppose $\alpha > 1$. Take any $0 < \delta < t$. There exists $0 < \varepsilon < \alpha$ such that, for any large enough n ,

$$t - \delta < \frac{-n \log(\alpha - \varepsilon)}{(n+1) \log \lambda + \log C}.$$

Note that, for any large enough n , $\kappa_n > (\alpha - \varepsilon)^n$.

For any $n \geq 0$, define a probability measure μ_n on \mathbb{R} as follows: μ_n is supported on $\bigcup_{B \in \mathcal{I}_n} B$. For any $B \in \mathcal{I}_n$,

$$\mu_n(B) = \kappa_n^{-1},$$

and μ_n is uniform on the interval B .

Take any $k, n > 0$. Let $B \in \mathcal{I}_n$. Suppose there are m intervals in \mathcal{I}_{n+k} that is contained in B . Then $\mu_{n+k}(B) = m/\kappa_{n+k}$. By ii),

$$m/C' \leq \frac{\kappa_{n+k}}{\kappa_n} \leq C'm.$$

So we have

$$(67) \quad \kappa_n^{-1}/C' \leq \mu_{n+k}(B) \leq C'\kappa_n^{-1}.$$

Take a weak-star limit of $\{\mu_n\}_{n \geq 1}$, say μ . Since \mathcal{I} is a SNS, μ is a probability measure supported on $A(\mathcal{I})$. Moreover, by (67), for any $n > 0$ and $B \in \mathcal{I}_n$,

$$\kappa_n^{-1}/C' \leq \mu(B) \leq C'\kappa_n^{-1}.$$

For any open interval U with $|U|$ small enough, there exists $k \in \mathbb{N}$ such that

$$C\lambda^{k+1} < |U| \leq C\lambda^k.$$

At most two intervals in \mathcal{I}_k intersect U . Then

$$\mu(U) \leq 2C'\kappa_k^{-1} \leq 2C'(\alpha - \varepsilon)^{-k} \leq 2C'|U|^{\frac{-k \log(\alpha - \varepsilon)}{(k+1) \log \lambda + \log C}} \leq 2C'|U|^{t-\delta}.$$

By mass distribution principle,

$$\dim_H A(\mathcal{I}) \geq t - \delta.$$

Since $\delta > 0$ can be arbitrarily small, the result follows. \square

5.2.2. Lower bound for $\dim_H \sigma_\lambda$.

Now we construct a sub-NS of $\{\mathcal{B}_n : n \geq 0\}$ such that it is a SNS and satisfies the conditions in Proposition 5.7.

Define a sub-alphabet $\tilde{A} = \{1_e, 1_o, 2_e, 2_o\}$ of \mathcal{A} and a sub directed graph $\tilde{\mathbb{G}}$ with restricted admissible relation:

$$1_e \rightarrow 2_o; \quad 1_o \rightarrow 2_e; \quad 2_e \rightarrow 1_o, 2_o; \quad 2_o \rightarrow 1_e, 2_e.$$

Consider the sub-NS $\tilde{\mathcal{B}} = \{\tilde{\mathcal{B}}_n : n \geq 1\}$ defined by

$$\tilde{\mathcal{B}}_n := \{I_{0_e 1_o w} \in \mathcal{B}_{n+1} : w = w_1 \cdots w_n, w_i \in \tilde{A}\}.$$

Proposition 5.8. *$\tilde{\mathcal{B}}$ is a SNS and its limit set $A(\tilde{\mathcal{B}}) \subset \sigma_\lambda \cap B_\infty$. Moreover There exists constant $C > 0$ such that for all $n, k \geq 0$ and $I, I' \in \tilde{\mathcal{B}}_n$,*

$$|I| \geq C 4^{-n} \quad \text{and} \quad \frac{\#\{J \in \tilde{\mathcal{B}}_{n+k} : J \subset I\}}{\#\{J \in \tilde{\mathcal{B}}_{n+k} : J \subset I'\}} \leq 2.$$

Proof. One check directly that $\tilde{\mathcal{B}}$ is a sub-NS of \mathcal{B} , so $A(\tilde{\mathcal{B}}) \subset A(\mathcal{B}) = \sigma_\lambda$. Moreover, since $2_e < 1_e$ and $1_o < 2_o$, combine with Corollary 5.2, one conclude that $\tilde{\mathcal{B}}$ is a SNS.

Take any $n > 0$ and any $I \in \tilde{\mathcal{B}}_n$. We claim that for any $1 \leq k \leq n+1$ and any $x \in I$,

$$|h_k(x)| \leq 2.$$

Indeed, assume $I = I_{0_e 1_o w}$ with $w \in \tilde{\mathcal{A}}^n$, then by (64) and tracing on \mathbb{G} , for any $k = 0, \dots, n$ we have

$$I^{(k)} := I_{0_e 1_o w_1 \dots w_k} = B_{1\sigma_1 \dots \sigma_k}, \text{ where } \sigma_i \in \{0, 1\}.$$

Thus $I^{(k)} \in \mathcal{B}_{k+1}$ and $I = I^{(n)} \subset I^{(n-1)} \subset \dots \subset I^{(0)} = B_1$. Since $|h_k| \leq 2$ on any $B \in \mathcal{B}_k$, the claim follows.

The claim implies immediately that $A(\tilde{\mathcal{B}}) \subset B_\infty$.

Recall that

$$h_{k+1}(x) = h_k(x)(h_{k-1}^2(x) - 2) - 2,$$

we have

$$h'_{k+1}(x) = (h_{k-1}^2(x) - 2)h'_k(x) + 2h_{k-1}(x)h_k(x)h'_{k-1}(x).$$

Then, for $x \in I$ and $2 \leq k \leq n$,

$$|h'_{k+1}(x)| \leq 2|h'_k(x)| + 8|h'_{k-1}(x)|.$$

This implies there exists $c > 0$ depending only on B_1 such that, for $x \in I$ and $1 \leq k \leq n+1$,

$$|h'_k(x)| \leq c4^k.$$

By Floquet theory, h_{n+1} is monotone on $I \in \mathcal{B}_{n+1}$ and $h_{n+1}(I) = [-2, 2]$. Writing $I = [x_0, x_1]$, we have

$$4 = \int_{x_0}^{x_1} |h'_{n+1}(x)| dx \leq c(x_1 - x_0)4^{n+1}.$$

Thus $|I| = x_1 - x_0 \geq c^{-1}4^{-n}$.

It is seen that $\tilde{\mathcal{B}}$ is corresponding to a sub-shift of finite type with alphabet $\{1_o, 2_o, 1_e, 2_e\}$ and incidence matrix

$$\tilde{A} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix}.$$

By this, one can show easily that

$$\frac{\#\{J \in \tilde{\mathcal{B}}_{n+k} : J \subset I\}}{\#\{J \in \tilde{\mathcal{B}}_{n+k} : J \subset I'\}} \leq \frac{F_k}{F_{k-1}},$$

where $\{F_k : k \geq 0\}$ is the Fibonacci sequence defined by $F_0 = 1, F_1 = 2$ and $F_{n+1} = F_n + F_{n-1}$. Then the result follows easily. \square

Proof of Theorem 1.1. By the definition of $\tilde{\mathcal{B}}$, it is seen that $\#\tilde{\mathcal{B}}_n = F_n$. It is well known that $F_n \sim \alpha^n$. By Propositions 5.8 and 5.7,

$$(68) \quad \dim_H \sigma_\lambda \geq \dim_H A(\tilde{\mathcal{B}}) \geq \liminf_{n \rightarrow \infty} \frac{\log \#\tilde{\mathcal{B}}_n}{n \log 4} = \frac{\log \alpha}{\log 4}.$$

\square

Remark 5.9. 1) Indeed, we can start from any band B with type in $\tilde{\mathcal{A}}$ and do the same construction by following the sub graph $\tilde{\mathcal{G}}$. Then we construct a Cantor subset of $\sigma_\lambda \cap B$ with the same dimension estimation. This means that the “local” dimension of σ_λ is also uniformly bounded from below.

2) If we view the SNS $\tilde{\mathcal{B}}$ as a dynamical system coded by the subshift $\Omega_{\tilde{\mathcal{A}}}$, then $\log \alpha$ is just the topological entropy of the system and $\log 4$ can be viewed as an upper bound of the lyapunov exponents.

6. ∞ -TYPE ENERGIES, GAPS OF THE SPECTRUM AND ZEROS OF TRACE POLYNOMIALS

In this section, at first, we establish the existence of ∞ -type energies; then we prove Theorems 1.3, 1.5 and 1.8.

6.1. ∞_I -type energies.

We will show that $E \in \pi(\mathcal{E}_l^e \cup \mathcal{E}_r^o)$ is an ∞ -type energy. Recall that in Sec. 1.2.5 we call it ∞_I -type energy. Indeed, we can be more precise:

Proposition 6.1. *i) $\pi(\mathcal{E}_l^e)$ is dense in σ_λ . If $E \in \pi(\mathcal{E}_l^e)$, then*

$$\lim_{n \rightarrow \infty} |h_{2n+1}(E)| \rightarrow \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} |h_{2n}(E)| \rightarrow \sqrt{2}.$$

ii) $\pi(\mathcal{E}_r^o)$ is dense in σ_λ . If $E \in \pi(\mathcal{E}_r^o)$, then

$$\lim_{n \rightarrow \infty} |h_{2n}(E)| \rightarrow \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} |h_{2n+1}(E)| \rightarrow \sqrt{2}.$$

We only prove i), since the proof of ii) is the same.

Lemma 6.2. *Suppose $\sigma \in \Sigma_n$ and B_σ has type 0_e . Then*

$$(69) \quad b_{\sigma 1} < b_{\sigma 111} < b_{\sigma 11} < b_\sigma,$$

and

$$(70) \quad h_{n+1}(E) \geq 2 \quad \text{for} \quad E \in [b_{\sigma 1}, b_\sigma],$$

$$(71) \quad h_n(E) \geq \sqrt{2} \quad \text{for} \quad E \in [a_{\sigma 11}, b_\sigma].$$

Proof. Since B_σ has type 0_e , n is even and $b_\sigma \notin \mathcal{R}_{n-1}$. By Lemma 2.8 i), $a_{\sigma 1} = z_\sigma$. By Lemma 2.8 iii), $b_{\sigma 1^k} \notin \mathcal{R}_{n+k-1}$ for any $k \geq 1$. Applying Lemma 3.1 i) to $B_{\sigma 1}$, we have $B_{\sigma 1} \prec B_{\sigma 111}$. So $b_{\sigma 1} < b_{\sigma 111}$. Applying Lemma 2.3 ii) to $B_{\sigma 11}$, we have $B_{\sigma 111} \subset \text{int}(B_{\sigma 11})$. So $b_{\sigma 111} < b_{\sigma 11}$. By Lemma 3.1 ii), $a_{\sigma 11} \notin \mathcal{R}_{n+1}$ and $B_{\sigma 1} \prec B_{\sigma 11} \subset \text{int}(B_\sigma)$. So $z_\sigma = a_{\sigma 1} < a_{\sigma 11}$ and $b_{\sigma 11} < b_\sigma$. Thus (69) follows.

If $\sigma = 1^n$, then $b_{\sigma 1} = b_{1^{n+1}}$ is the maximal root of $h_{n+1}(E) = 2$. Hence $h_{n+1}(E) \geq 2$ for any $E \in [b_{\sigma 1}, \infty)$. Now assume $\sigma < 1^n$. Since $b_\sigma \notin \mathcal{R}_{n-1}$, by Lemma 2.8 iv), $a_{\sigma^+} \in \mathcal{R}_{n-1}$. By Lemma 2.8 ii), $a_{\sigma^+0} = a_{\sigma^+}$. By Proposition 2.10 i), h_{n+1} is increasing on $B_{\sigma 1}$ and decreasing on B_{σ^+0} , hence,

$$h_{n+1}(b_{\sigma 1}) = 2; \quad h_{n+1}(a_{\sigma^+0}) = 2; \quad h_n([b_{\sigma 1}, a_{\sigma^+0}]) \subset [2, \infty).$$

Since $b_{\sigma 1} < b_\sigma < a_{\sigma^+} = a_{\sigma^+0}$, (70) holds.

By Proposition 2.10 i), we have $h_{n+2}(a_{\sigma 11}) = -2$. So

$$-2 = h_{n+2}(a_{\sigma 11}) = h_{n+1}(a_{\sigma 11})(h_n(a_{\sigma 11})^2 - 2) - 2.$$

Since $a_{\sigma 11} \notin \mathcal{R}_{n+1}$, $h_{n+1}(a_{\sigma 11}) \neq 0$. So we must have $h_n(a_{\sigma 11}) = \pm\sqrt{2}$. By Lemma 3.11, h_n is increasing on B_σ . Since $z_\sigma < a_{\sigma 11} < b_{\sigma 11} < b_\sigma$, we must have $h_n(a_{\sigma 11}) = \sqrt{2}$. Consequently (71) holds. \square

Proof of Proposition 6.1. We only show i). At first we show that $\pi(\mathcal{E}_l^e)$ is dense in σ_λ . Since $\{I_w : w \in \Omega_n\}$ is a covering of the spectrum and $\max\{|I_w| : w \in \Omega_n\} \rightarrow 0$ as $n \rightarrow \infty$, we only need to show that $I_w \cap \pi(\mathcal{E}_l^e) \neq \emptyset$ for any $w \in \Omega_*$. Since \mathbb{G} is connected, for any $\alpha \in \mathcal{A}$, there exists an admissible path $u_\alpha := \alpha\alpha_1 \cdots \alpha_{k-1}0_e$. Assume $w \in \Omega_m$, define $v := w|_{m-1}u_{w_m}$, then I_v has type 0_e , $I_v \subset I_w$ and $\pi(v3_{or}(0_e3_{or})^\infty) \in I_v \cap \pi(\mathcal{E}_l^e)$. Hence $I_w \cap \pi(\mathcal{E}_l^e) \neq \emptyset$.

Assume $E \in \pi(\mathcal{E}_l^e)$, then there exist some even n and $w \in \Omega_n$ with $w_n = 0_e$ such that $E = \pi(w(3_{or}0_e)^\infty)$. write $\sigma = \Pi_*(w)$, then $\sigma \in \Sigma_n$. For any $k \geq 0$, by (52) and tracing on \mathbb{G} , we have $\Pi_*(w(3_{or}0_e)^k) = \sigma 1^{2k}$. By (64),

$$B_{\sigma 1^{2k}} = I_{w(3_{or}0_e)^k}.$$

Write $\sigma^{(k)} := \sigma 1^{2k}$. Then $B_{\sigma^{(k)}}$ has type 0_e . By the definition of π , we have $E \in B_{\sigma^{(k)}}$. We claim that $E \geq b_{\sigma 1^{2k+1}}$. Indeed, by applying (69) to every $\sigma^{(k)}$, we have

$$b_{\sigma 1} < b_{\sigma 1^3} < \cdots < b_{\sigma 1^{2k+1}} < \cdots < b_{\sigma 1^{2k}} < \cdots < b_{\sigma 1^2} < b_\sigma.$$

Since $E \in B_{\sigma^{(k)}}$ and the length of $B_{\sigma^{(k)}}$ tend to 0, we conclude that

$$E = \inf_{k \geq 1} b_{\sigma 1^{2k}} \geq \sup_{k \geq 1} b_{\sigma 1^{2k+1}}.$$

So $E \in [b_{\sigma^{(k)1}}, b_{\sigma^{(k)}}]$. By (70),

$$h_{n+2k+1}(E) \geq 2.$$

Since $E \in B_{\sigma^{(k+1)}} = [a_{\sigma^{(k+1)11}}, b_{\sigma^{(k+1)11}}] \subset B_{\sigma^{(k)}} = [a_{\sigma^{(k)}}, b_{\sigma^{(k)}}]$, by (71),

$$h_{n+2k}(E) \geq \sqrt{2}.$$

By Lemma 3.6 ii), $E \in B_{\sigma 1^4} \subset \text{int}(B_{\sigma 1^2})$. By Lemma 2.5 ii), $E \notin \mathcal{R}_{n+1}$.

Now for any $N \geq 1$, since $E \in B_{\sigma 1^{2N}}$, we have $|h_{n+2N}(E)| \leq 2$. Then by Lemma 2.9,

$$\begin{aligned} |h_{n+2N+1}(E)| &\geq 2\lambda \prod_{j=0}^{n+2N} |h_j(E)| - |h_{n+2N}^2(E) - 2| \\ &\geq 2^{3N/2+1}\lambda \prod_{j=0}^n |h_j(E)| - 6. \end{aligned}$$

Hence $|h_{n+2N+1}(E)| \rightarrow \infty$. Now by (23),

$$|h_{n+2N}^2(E) - 2| = \frac{|h_{n+2N+2}(E) + 2|}{|h_{n+2N+1}(E)|} \leq \frac{4}{|h_{n+2N+1}(E)|} \rightarrow 0.$$

So $|h_{n+2N}(E)| \rightarrow \sqrt{2}$. \square

6.2. ∞_{II} -type energies.

We will show that $E \in \pi(\tilde{\mathcal{E}}_l \cup \tilde{\mathcal{E}}_r)$ is also an ∞ -type energy. Recall that in Sec. 1.2.5 we call it ∞_{II} -type energy.

Proposition 6.3. *i) Both $\pi(\tilde{\mathcal{E}}_l)$ and $\pi(\tilde{\mathcal{E}}_r)$ are dense in σ_λ .*

ii) Assume $(\omega, \hat{\omega}) \in \mathcal{G}_{II}$, then

$$\lim_{n \rightarrow \infty} |h_{2n}(\pi(\omega))| = \lim_{n \rightarrow \infty} |h_{2n+1}(\pi(\hat{\omega}))| = \infty.$$

Although this proposition looks similar with Proposition 6.1, the proof of which is much more involved. We need to study a dynamical system induced by the recurrence relation of trace polynomials, which is inspired by our previous work [21].




FIGURE 4. Dynamics of f

Define the *period-doubling trace map* $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ as

$$(72) \quad f(x, y) = (y(x^2 - 2) - 2, [y(x^2 - 2) - 2](y^2 - 2) - 2).$$

By (23), for any $n \geq 0$ and $E \in \mathbb{R}$, we have

$$(h_{n+2}(E), h_{n+3}(E)) = f(h_n(E), h_{n+1}(E)).$$

We have

$$Df(x, y) = \begin{bmatrix} 2xy & x^2 - 2 \\ 2xy(y^2 - 2) & (x^2 - 2)(3y^2 - 2) - 4y \end{bmatrix}.$$

So we have

$$\det Df(x, y) = 4xy^2(y(x^2 - 2) - 2).$$

Define a simply connected domain U (enclosed by the red curve in Figure 4 (a)) as

$$U := \{(x, y) : x, y < 0, y(x^2 - 2) - 2 < 0\}.$$

Define a compact set D (enclosed by the blue curve in Figure 4 (a)) as

$$D := \{(x, y) : -\sqrt{2} \leq x, y \leq 0; y \leq x^2 - 2\}.$$

Notice that $(-1, -1) \in \partial D$. Write

$$A := (-\sqrt{2}, -\sqrt{2}), \quad B := (-\sqrt{2}, 0), \quad C := (-\sqrt{2 - \sqrt{2}}, -\sqrt{2}), \quad F := (-1, -1).$$

Then A, B, C are the vertices of the “triangle” D . Define

$$\begin{cases} \Gamma := \{(-\sqrt{2}, y) : -\sqrt{2} \leq y \leq 0\}, \\ \Upsilon := \{(x, -\sqrt{2}) : -\sqrt{2} \leq x \leq -\sqrt{2 - \sqrt{2}}\}, \\ \Lambda := \{(x, x^2 - 2) : -\sqrt{2} \leq x \leq -\sqrt{2 - \sqrt{2}}\}. \end{cases}$$

Then $B, C, F \in \Lambda$ and $\partial D = \Gamma \cup \Upsilon \cup \Lambda$, see Figure 4 (b).

6.2.1. Local dynamical property of f .

We summarize the needed properties of f in the following proposition.

Proposition 6.4. *i) $f : U \rightarrow f(U)$ is a diffeomorphism and $D \subset f(U)$. f has a unique fixed point F in U .*

ii) Let $g : f(U) \rightarrow U$ be the inverse of f , then g has a unique fixed point F in $f(U)$. Moreover, $g(D) \subset D \cap U$.

iii) $g(\Lambda) \subset \Lambda$ and $g^n(\Lambda) \downarrow \{F\}$.

iv) Write $g = (g_1, g_2)$. For any $(x, y) \in D$ and $i = 1, 2$,

$$(73) \quad \frac{\partial g_i}{\partial x}(x, y), \quad \frac{\partial g_i}{\partial y}(x, y) > 0.$$

Proof. i) Write $(x_1, y_1) := f(x, y)$, we have

$$y_1 = x_1 \frac{(x_1 + 2)^2}{(x^2 - 2)^2} - 2x_1 - 2.$$

Since $x_1 < 0$ for $(x, y) \in U$, we have $y_1 < -2x_1 - 2$. Moreover, for $-2 \leq x < 0$, $y_1 < x_1^3/4 + x_1^2 - x_1 - 2$. Now by computing the image of the fiber $U_a := U \cap \{y = a\}$ under f for all $a < 0$, one can verify that

$$f(U) = \left\{ (x, y) : x < 0; y < -2x - 2; y < \frac{x^3}{4} + x^2 - x - 2 \text{ if } -2 \leq x < 0 \right\}.$$

One check directly that $D \subset f(U)$. Define a function g as

$$g(x, y) := \left(-\sqrt{2 - \frac{2+x}{\sqrt{2 + \frac{2+y}{x}}}}, -\sqrt{2 + \frac{2+y}{x}} \right).$$

One check directly that g can be defined on $f(U)$ and is smooth, and

$$g \circ f|_U = \text{Id}_U; \quad f \circ g|_{f(U)} = \text{Id}_{f(U)}.$$

So $f : U \rightarrow f(U)$ is a diffeomorphism and g is the inverse of f .

By direct computation, the fixed points of f are

$$(-1, -1), \quad (2, 2), \quad (-\alpha, \alpha - 1), \quad (\alpha^{-1}, -(1 + \alpha^{-1})),$$

where $\alpha = (\sqrt{5} + 1)/2$. So F is the only fixed point of f in U .

ii) The first assertion is a restatement of i). Fix any $(x, y) \in D$. By direct computation,

$$-\sqrt{2} < -1.3 \leq g_1(x, y) \leq -1.2 < -\sqrt{2 - \sqrt{2}}; \quad -\sqrt{2} < -1.3 \leq g_2(x, y) \leq -0.7.$$

Moreover,

$$g_2(x, y) - g_1^2(x, y) + 2 = \frac{y - x^2 + 2}{xg_2(x, y)} \leq 0.$$

By the definitions of D and U , we conclude that $g(D) \subset D \cap U$.

iii) Define $\phi : \mathbb{R} \rightarrow \mathbb{R}$ as $\phi(x) = x^2 - 2$. Write $a := -\sqrt{2}, b := -\sqrt{2 - \sqrt{2}}$. Then

$$\Lambda = \{(x, \phi(x)) : x \in [a, b]\}.$$

By the definition of f , for any $x \in \mathbb{R}$,

$$f(x, \phi(x)) = (\phi^2(x), \phi^3(x)),$$

where ϕ^n is the n -time iteration of ϕ . By solving $\phi^2(x) = a$ and b , there exist a_1, b_1 such that

$$a < a_1 = -1.11 \cdots < -1 < b_1 < b = -0.76 \cdots; \quad \phi^2(a_1) = a; \quad \phi^2(b_1) = b.$$

Since $[\phi^2(x)]' = 4x(x^2 - 2)$, for any $x \in [a_1, b_1]$,

$$(74) \quad [\phi^2(x)]' \geq 3.$$

Define $\Lambda_1 := \{(x, \phi(x)) : x \in [a_1, b_1]\}$, then $\Lambda_1 \subset \Lambda$ and $f(\Lambda_1) = \Lambda$. Since $\Lambda \subset D \subset f(U)$ and $g : f(U) \rightarrow U$ is a diffeomorphism, we conclude that

$$g(\Lambda) = \Lambda_1 \subset \Lambda.$$

Notice that $\phi^2(-1) = -1$. By (74) and contraction principle, for any $n \geq 2$ one can find a_n, b_n such that $a_n > a_1, b_n < b_1$ and $a_n \uparrow -1, b_n \downarrow -1$ and $\phi^{2n}(a_n) = a, \phi^{2n}(b_n) = b$. By repeating the above proof, we get

$$g^n(\Lambda) = \Lambda_n := \{(x, \phi(x)) : x \in [a_n, b_n]\} \downarrow \{F\}.$$

iv) By ii), we have $-1.3 \leq g_1(x, y), g_2(x, y) \leq -0.7$. Thus

$$\begin{aligned} \frac{\partial g_1}{\partial x} &= \frac{1}{2g_1g_2} + \frac{(2 - g_1^2)(2 - g_2^2)}{4xg_1g_2^2} > 0, & \frac{\partial g_1}{\partial y} &= \frac{2 - g_1^2}{4xg_1g_2^2} > 0; \\ \frac{\partial g_2}{\partial x} &= \frac{2 - g_2^2}{2xg_2} > 0, & \frac{\partial g_2}{\partial y} &= \frac{1}{2xg_2} > 0. \end{aligned}$$

So, (73) holds. □

Define a partial order on \mathbb{R}^2 as follows. Assume $p = (x, y)$ and $q = (\tilde{x}, \tilde{y})$, we say $p \leq q$ if $x \leq \tilde{x}, y \leq \tilde{y}$. Assume \mathcal{C} is a continuous curve in \mathbb{R}^2 with a parametrization $\gamma : [a, b] \rightarrow \mathcal{C}$ such that $\gamma(t) \leq \gamma(s)$ whenever $t < s$, then we call \mathcal{C} an *increasing curve* in \mathbb{R}^2 .

Lemma 6.5. *If $p, q \in D$ and $p \leq q$, then $g(p) \leq g(q)$.*

Proof. Notice that, if $p, q \in D$ and $p \leq q$, then for any $\hat{p} \in \mathbb{R}^2$ with $p \leq \hat{p} \leq q$, we have $\hat{p} \in D$. By (73), $g_1(x, \cdot)$, $g_1(\cdot, y)$, $g_2(x, \cdot)$, $g_2(\cdot, y)$ are all strictly increasing. Now assume $p = (x, y)$, $q = (\tilde{x}, \tilde{y})$ and $p \leq q$, then

$$\begin{aligned} g_1(p) &= g_1(x, y) \leq g_1(\tilde{x}, y) \leq g_1(\tilde{x}, \tilde{y}) = g_1(q), \\ g_2(p) &= g_2(x, y) \leq g_2(\tilde{x}, y) \leq g_2(\tilde{x}, \tilde{y}) = g_2(q). \end{aligned}$$

Hence $g(p) \leq g(q)$. □

Now we have the following consequence.

Proposition 6.6. *Assume $p \in D$. Then $f^n(p) \in D$ for any $n \in \mathbb{N}$ if and only if $p = F$.*

Proof. Since $f(F) = F$ and $F \in D$, we have $f^n(F) \in D$ for any n . So the if part holds.

To prove the only if part, we need the following claim:

Claim: We have

$$(75) \quad g^n(D) \downarrow \{(-1, -1)\}.$$

◁ since $g(F) = F$, we have $F \in g^n(D)$ for any $n \geq 0$. By Proposition 6.4 ii), $g(D) \subset D$. So $\{g^n(D) : n \geq 0\}$ is a decreasing sequence of compact sets. To prove (75), we only need to show that the diameter of $g^n(D)$ tends to 0.

Recall that, A, B, C are the vertices of the “triangle” D . Define

$$A_n := g^n(A), \quad B_n := g^n(B), \quad C_n := g^n(C).$$

Since $B, C \in \Lambda$, By Proposition 6.4 iii), $B_n, C_n \rightarrow F$. We also have

$$A \leq A_1 = \left(-\sqrt{2 - \frac{2 - \sqrt{2}}{\sqrt{3 - \sqrt{2}}}}, -\sqrt{3 - \sqrt{2}} \right), \quad A, A_1 \in D.$$

We have $A_n \in D$ since $g(D) \subset D$. By Lemma 6.5,

$$A \leq A_1 \leq A_2 \leq \dots$$

Hence $A_\infty := \lim_n A_n$ exists and $A_\infty \in D$, since g is continuous on D , $g(A_\infty) = A_\infty$. By Proposition 6.4 ii), g has a unique fixed point F in D , so $A_\infty = F$.

Recall that $\Gamma \cup \Upsilon \cup \Lambda$ is the boundary of D . By Proposition 6.4 i) and ii), $g : f(U) \rightarrow U$ is a diffeomorphism and $D \subset f(U)$. Since $g^n(D) \subset D$, we conclude that $g^n(\Gamma) \cup g^n(\Upsilon) \cup g^n(\Lambda)$ is the boundary of $g^n(D)$. Notice that Γ is the vertical interval AB which is an increasing interval. By Lemma 6.5, $g^n(\Gamma)$ is also increasing and has endpoints A_n, B_n . Similarly, $g^n(\Upsilon)$ is increasing and has endpoints A_n, C_n . By Proposition 6.4 iii), $g^n(\Lambda) \subset \Lambda$ is the segment of Λ with endpoints B_n, C_n . So we conclude that

$$g^n(D) \subset [x_{A_n}, x_{C_n}] \times [y_{A_n}, y_{B_n}].$$

Since $A_n, B_n, C_n \rightarrow F$, we conclude that $\text{diam}(g^n(D)) \rightarrow 0$. ▷

By the claim, we can write $D \setminus \{F\}$ as the following disjoint union:

$$D \setminus \{F\} = \bigcup_{n \geq 1} (g^{n-1}(D) \setminus g^n(D)).$$

Now if $p \in D \setminus \{F\}$, then there exists $m \geq 1$ such that $p \in g^{m-1}(D) \setminus g^m(D)$. So we have

$$f^m(p) \in f(f^{m-1}(g^{m-1}(D) \setminus g^m(D))) = f(D \setminus g(D)).$$

By Proposition 6.4 ii), $g(D) \subset D \cap U$. Notice that $B, C \notin U$ and $D \setminus \{B, C\} \subset U$, so $g(D) \subset D \setminus \{B, C\} \subset U$. Since f is injective on U , we conclude that

$$f((D \setminus \{B, C\}) \setminus g(D)) = f(D \setminus \{B, C\}) \setminus D.$$

Hence $f((D \setminus \{B, C\}) \setminus g(D)) \cap D = \emptyset$. On the other hand, by direct computation,

$$f(B) = (-2, 2), f(C) = (0, -2) \notin D.$$

So we have

$$D \cap f(D \setminus g(D)) = D \cap (f((D \setminus \{B, C\}) \setminus g(D)) \cup \{f(B), f(C)\}) = \emptyset.$$

That is, $f^m(p) \notin D$. So the only if part follows. \square

6.2.2. Proof of Proposition 6.3.

Assume $(\omega, \hat{\omega}) \in \mathcal{G}_{II}$. Write $E_1 = \pi(\omega)$ and $E_2 = \pi(\hat{\omega})$. By Lemma 5.5, $E_1 \leq E_2$.

Lemma 6.7. *If $E_1 = E_2 =: E$, then there exists $m \in \mathbb{N}$ such that for any $k \geq m$,*

$$(h_k(E), h_{k+1}(E)) \in \text{int}(D).$$

If $E_1 < E_2$, then as $k \rightarrow \infty$,

$$|h_{2k}(E_1)|, |h_{2k+1}(E_2)| \rightarrow \infty.$$

Proof. By (48), ω takes one of the following two forms:

$$w1_o3_{er}(0_o3_{er})^\infty; \quad w3_{el}(0_o3_{er})^\infty.$$

We prove the lemma for $\omega = w1_o3_{er}(0_o3_{er})^\infty$, and leave the other case to the reader.

Assume $w \in \Omega_n$. In this case, n is even, $w_n = 0_e$ or 2_e and $\hat{\omega} = \ell(\omega) = w3_{or}(0_e3_{ol})^\infty$. For $k \geq 1$, by (61),

$$\begin{cases} \Pi_*(\omega|_{n+2k+1}) = \Pi_*(w)\delta(10)^k =: \sigma^{(k)}, \\ \Pi_*(\hat{\omega}|_{n+2k+2}) = \Pi_*(w)\delta 1(01)^k =: \hat{\sigma}^{(k)}. \end{cases}$$

Here $\delta = 1$ if $w_n = 0_e$ and $\delta = 0$ if $w_n = 2_e$. Notice that $\hat{\sigma}^{(k)} = \sigma^{(k)}1$. By (64),

$$I_{\omega|_{n+2k+1}} = I_{w1_o(3_{er}0_o)^k} = B_{\sigma^{(k)}}, \quad I_{\hat{\omega}|_{n+2k+2}} = I_{w3_{or}0_e(3_{ol}0_e)^k} = B_{\hat{\sigma}^{(k)}}.$$

We have $E_1 \in B_{\sigma^{(k)}}$ and $E_2 \in B_{\hat{\sigma}^{(k)}}$. Notice that $B_{\sigma^{(k)}}$ has type 0_o and $B_{\hat{\sigma}^{(k)}}$ has type 0_e . By applying Lemma 3.6 i) to $B_{\sigma^{(k)}}$, we have

$$(76) \quad z_{\sigma^{(k)}} = b_{\sigma^{(k)}0} < a_{\sigma^{(k+1)}} < E_1 < b_{\sigma^{(k+1)}} < b_{\sigma^{(k)}}.$$

In particular, $E_1 \in \text{int}(B_{\sigma^{(k)}})$. By Proposition 2.10 i), $h_{n+2(k+1)+1}$ is decreasing on $B_{\sigma^{(k+1)}}$, so $h_{n+2(k+1)+1}(b_{\sigma^{(k+1)}}) = -2$. Hence by (23),

$$-2 = h_{n+2(k+1)}(b_{\sigma^{(k+1)}})(h_{n+2k+1}^2(b_{\sigma^{(k+1)}}) - 2) - 2.$$

Since $B_{\sigma^{(k+1)}}$ has type 0_o , $b_{\sigma^{(k+1)}} \notin \mathcal{R}_{n+2(k+1)}$. Thus $h_{n+2k+1}(b_{\sigma^{(k+1)}}) = \pm\sqrt{2}$. Since h_{n+2k+1} is decreasing on $B_{\sigma^{(k)}}$, by (76),

$$(77) \quad -\sqrt{2} = h_{n+2k+1}(b_{\sigma^{(k+1)}}) < h_{n+2k+1}(E_1) < h_{n+2k+1}(z_{\sigma^{(k)}}) = 0.$$

On the other hand, since $E_1 \in \text{int}(B_{\sigma^{(k)}})$ with $|\sigma^{(k)}| = n + 2k + 1$ and $\sigma^{(k)}$ ends with 0, by Lemma 2.9 and Corollary 2.7,

$$(78) \quad h_{n+2k+1}(E_1) - (h_{n+2k}^2(E_1) - 2) = (-1)^{n+2k} 2\lambda \prod_{j=0}^{n+2k} h_j(E_1) < 0.$$

By a symmetric argument, one can show that

$$(79) \quad -\sqrt{2} < h_{n+2k+2}(E_2) < 0 \quad \text{and} \quad h_{n+2k+2}(E_2) - (h_{n+2k+1}^2(E_2) - 2) < 0.$$

Now assume $E_1 = E_2 =: E$. By (77), (78) and (79), for any $k \geq n + 3 =: m$,

$$-\sqrt{2} < h_k(E) < 0; \quad h_{k+1}(E) < h_k^2(E) - 2.$$

By the definition of D , we have $(h_k(E), h_{k+1}(E)) \in \text{int}(D)$.

Next assume $E_1 < E_2$. By (76), we have $z_{\sigma^{(k)}} < E_1$. By a symmetric argument, we have $E_2 < z_{\hat{\sigma}^{(k)}}$. So

$$z_{\sigma^{(k)}} < E_1 < E_2 < z_{\hat{\sigma}^{(k)}}.$$

Since the length of $B_{\sigma^{(k)}}$ and $B_{\hat{\sigma}^{(k)}}$ tend to 0, There exists $K \geq n + 3$ such that for any $k \geq K$, $B_{\sigma^{(k)}} < B_{\hat{\sigma}^{(k)}}$. Recall that $\hat{\sigma}^{(k)} = \sigma^{(k)}1$. So $b_{\sigma^{(k)}} < a_{\hat{\sigma}^{(k)}} = a_{\sigma^{(k)}1}$. By Proposition 2.10 i), h_{n+2k+2} is decreasing on $B_{\sigma^{(k)}0}$ and increasing on $B_{\sigma^{(k)}1}$. By Floquet theory, for any $E \in (b_{\sigma^{(k)}0}, a_{\sigma^{(k)}1})$, we have $h_{n+2k+2}(E) < -2$. By (76), $b_{\sigma^{(k)}0} < E_1 < b_{\sigma^{(k)}} < a_{\sigma^{(k)}1}$, so

$$(80) \quad h_{n+2k+2}(E_1) < -2.$$

We claim that

$$(81) \quad h_{n+2k+1}(E_1) \leq -1.$$

If otherwise, by (77), we have $-1 < h_{n+2k+1}(E_1) < 0$. Then

$$h_{n+2k+3}(E_1) = h_{n+2k+2}(E_1)(h_{n+2k+1}^2(E_1) - 2) - 2 \geq 0,$$

which contradicts with (77).

By (78), $E_1 \notin \mathcal{Z}$. Since $E_1 \in B_{\sigma^{(k)}}$, we have $|h_{n+2k+1}(E_1)| \leq 2$. Now for any $N > K$, by Lemma 2.9 and (80), (81),

$$\begin{aligned} |h_{n+2N+2}(E_1)| &\geq 2\lambda \prod_{j=0}^{n+2N+1} |h_j(E_1)| - |h_{n+2N+1}^2(E_1) - 2| \\ &\geq \lambda 2^{(N-K)/2} \prod_{j=0}^{n+2K} |h_j(E_1)| - 2. \end{aligned}$$

Hence $|h_{n+2N+2}(E_1)| \rightarrow \infty$ as $N \rightarrow \infty$.

By exactly the same proof, we can show that $|h_{n+2N+1}(E_2)| \rightarrow \infty$ as $N \rightarrow \infty$. So the proof is finished. \square

As an application, we can improve Lemma 5.5 as follows:

Lemma 6.8. *If $(\omega, \hat{\omega}) \in \mathcal{G}_{II}$, then $\pi(\omega) < \pi(\hat{\omega})$.*

Proof. We prove it by contradiction. Assume $\pi(\omega) = \pi(\hat{\omega}) =: E$. By Lemma 6.7, there exists $m \in \mathbb{N}$ such that for any $k \geq m$, $(h_k(E), h_{k+1}(E)) \in \text{int}(D)$. Recall that $f(h_k(E), h_{k+1}(E)) = (h_{k+2}(E), h_{k+3}(E))$, thus for any $k \in \mathbb{N}$, we have

$$f^k(h_m(E), h_{m+1}(E)) = (h_{m+2k}(E), h_{m+2k+1}(E)) \in \text{int}(D).$$

By Proposition 6.6, $(h_m(E), h_{m+1}(E)) = F$. However, $F \in \partial D$, a contradiction. So $\pi(\omega) < \pi(\hat{\omega})$. \square

Proof of Proposition 6.3 i) The proof is the same as Proposition 6.1.

ii) It is a direct consequence of Lemma 6.8 and Lemma 6.7. \square

6.3. Proof of Theorems 1.3, 1.5 and 1.8.

Proof of Theorem 1.3. At first we show $B_\infty \subset \sigma_\lambda$. Fix $E \in B_\infty$, we claim that for any $n \geq 0$, $|h_n(E)| \leq 2$ or $|h_{n+1}(E)| \leq 2$. If otherwise, there exists $n_0 \geq 0$ and $\delta > 0$ such that $|h_{n_0}(E)|, |h_{n_0+1}(E)| \geq 2 + \delta$. By (23),

$$|h_{n_0+2}(E)| \geq |h_{n_0+1}(E)|(h_{n_0}^2(E) - 2) - 2 \geq 2 + 10\delta.$$

Using (23) again, we have $|h_{n_0+3}(E)| \geq 2 + 10\delta$. Now by induction, it is seen that

$$|h_{n_0+k}(E)| \geq 2 + 10^{\lfloor k/2 \rfloor} \delta.$$

Thus $|h_n(E)| \rightarrow \infty$, which contradicts with $E \in B_\infty$. So the claim holds. Now by (24), we conclude that $E \in \sigma_\lambda$. Thus $B_\infty \subset \sigma_\lambda$.

By Lemma 2.5 iii), $\mathcal{Z} \subset B_\infty$ and B_∞ is dense in σ_λ . By Proposition 5.8 and (68),

$$\dim_H B_\infty \geq \dim_H A(\tilde{\mathcal{B}}) \geq \log \alpha / \log 4 > 0,$$

so B_∞ is uncountable.

By the definition of \mathcal{E}_∞ , $\mathcal{E}_\infty = \sigma_\lambda \setminus B_\infty$ always holds. Now we show that \mathcal{E}_∞ is dense in σ_λ and uncountable.

By Proposition 6.1, $\pi(\mathcal{E}_i^e) \subset \mathcal{E}_\infty$ and is dense in σ_λ . So \mathcal{E}_∞ is dense in σ_λ .

Next we show that \mathcal{E}_∞ is uncountable. We prove it by contradiction. Otherwise, \mathcal{E}_∞ is countable. Let $\mathcal{E}_\infty = \{\gamma_i\}_{i=1}^\infty$. Since σ_λ has no isolated point, we conclude that $\{\gamma_i\}$ is nowhere dense in σ_λ for each i . Denote

$$T := \sigma_\lambda \setminus \mathcal{E}_\infty = \{E \in \sigma_\lambda : \mathcal{O}(E) \text{ is bounded}\}.$$

We can write T as a countable union of closed subsets,

$$T = \bigcup_{k=1}^{\infty} A_k,$$

where

$$A_k = \bigcap_{n=0}^{\infty} \{E \in \sigma_\lambda : |h_n(E)| \leq k\}.$$

Since \mathcal{E}_∞ is dense in σ_λ and $\mathcal{E}_\infty \cap A_k = \emptyset$, A_k is nowhere dense for each $k \in \mathbb{N}$. Thus

$$\sigma_\lambda = \bigcup_{i=1}^{\infty} \{\gamma_i\} \cup \bigcup_{k=1}^{\infty} A_k,$$

i.e., σ_λ is a countable union of nowhere dense closed sets. This contradicts with Baire's category theorem. \square

We remark that, once we know that \mathcal{E}_∞ is dense in σ_λ , the rest proof of the theorem is purely topological.

Proof of Theorem 1.5. By Proposition 5.3, we only need to show that π is a homeomorphism. Since Ω_∞ is compact and σ_λ is Hausdorff, it suffice to show that π is injective.

Assume $\omega \neq \hat{\omega}$. WLOG, we assume $\omega \prec \hat{\omega}$. If $\omega < \hat{\omega}$, then by Lemma 5.4, $\pi(\omega) < \pi(\hat{\omega})$. If $\omega \not< \hat{\omega}$, by Lemma 4.7, there exists $(\tau, \hat{\tau}) \in \mathcal{G}_{II}$ such that $\omega \preceq \tau$ and $\hat{\tau} \preceq \hat{\omega}$. By Proposition 5.3 and Lemma 6.8,

$$\pi(\omega) \leq \pi(\tau) < \pi(\hat{\tau}) \leq \pi(\hat{\omega}).$$

Hence π is injective. \square

Proof of Theorem 1.8. i)-iv) By Theorem 1.5, (a, b) is a gap of σ_λ if and only if $(\pi^{-1}(a), \pi^{-1}(b))$ is a gap of Ω_∞ . So i)-iv) follows directly from i)-iv) of Theorem 4.3 and Theorem 1.5.

v) At first, we prove $\pi(\mathcal{E}_l^o) = \mathcal{Z}^o$. Recall that $\mathcal{Z}^o = \{z_\sigma : \sigma \in \Sigma_*^o\}$. Fix any $\sigma \in \Sigma_*^o$, by (59), there exists a unique $\omega_\sigma \in \mathcal{E}_l^o$ such that $\Pi(\omega) = \sigma 01^\infty$. Moreover

$$\{\omega_\sigma : \sigma \in \Sigma_*^o\} = \mathcal{E}_l^o.$$

We claim that $\pi(\omega_\sigma) = z_\sigma$. Once we show the claim, we immediately get $\pi(\mathcal{E}_l^o) = \mathcal{Z}^o$. Now we show the claim. At first, by Lemma 2.8 i), we have $z_\sigma = b_{\sigma 01^k}$ for any $k \geq 0$. On the other hand, by the definition of Π , there exists $N \in \mathbb{N}$ such that when $m \geq N$, $\Pi_*(\omega_\sigma|_m) = \sigma 01^m$. Note the $\{I_{\omega_\sigma|_m} : m \geq 1\}$ is a decreasing set sequence, so we have

$$\pi(\omega_\sigma) = \bigcap_{m \geq 0} I_{\omega_\sigma|_m} = \bigcap_{m \geq N} I_{\omega_\sigma|_m} = \bigcap_{m \geq N} B_{\sigma 01^m}.$$

Consequently $z_\sigma \in \bigcap_{m \geq N} B_{\sigma 01^m}$. Thus $\pi(\omega_\sigma) = z_\sigma$.

The proof of $\pi(\mathcal{E}_r^e) = \mathcal{Z}^e$ is the same.

vi) It is a direct consequence of Propositions 6.1 and 6.3. \square

7. IDS AND GAP LABELLING

In this section, we prove Theorems 1.10 and 1.11.

7.1. Computation of IDS.

7.1.1. *The IDS for \mathcal{Z} .*

Recall that $\mathcal{Z} = \{z_\sigma : \sigma \in \Sigma_*\}$; ϖ_n is defined by (26); an order \preceq on Σ_* is defined by (34). We have

Lemma 7.1. *For $\sigma \in \Sigma_n$, we have*

$$(82) \quad N(z_\sigma) = \sum_{i=1}^n \frac{\sigma_i}{2^i} + \frac{1}{2^{n+1}}.$$

Consequently,

$$(83) \quad N(\mathcal{Z}) = \{N(z_\sigma) : \sigma \in \Sigma_*\} = \left\{ \frac{j}{2^n} : n \geq 1; 1 \leq j < 2^n \right\} = \mathcal{D} \cap (0, 1).$$

Proof. By the property of IDS and Proposition 2.6, we have

$$\begin{aligned} N(z_\sigma) &= \lim_{m \rightarrow \infty} \frac{\#\{z \in \mathcal{Z}_m : z \leq z_\sigma\}}{2^m} = \lim_{m \rightarrow \infty} \frac{\#\{\alpha \in \Sigma_m : \alpha \preceq \sigma\}}{2^m} \\ &= \lim_{m \rightarrow \infty} \frac{\#\{\alpha \in \Sigma_m : \alpha \preceq \sigma 01^{m-n-1}\}}{2^m} = \lim_{m \rightarrow \infty} \frac{\varpi_m(\sigma 01^{m-n-1}) + 1}{2^m} \\ &= \lim_{m \rightarrow \infty} \left(\sum_{j=1}^n \frac{\sigma_j}{2^j} + \sum_{j=n+2}^m 2^{-j} + 2^{-m} \right) = \sum_{j=1}^n \frac{\sigma_j}{2^j} + \frac{1}{2^{n+1}}. \end{aligned}$$

Hence we have

$$\{N(z_\sigma) : \sigma \in \Sigma_0\} = \{1/2\}; \quad \{N(z_\sigma) : \sigma \in \Sigma_1\} = \{1/4, 3/4\}; \quad \dots$$

Then (83) holds. □

7.1.2. *The IDS for σ_λ .*

Now we compute the IDS for any energy in the spectrum.

Proof of Theorem 1.10. By Proposition 4.5 iii), Π is surjective.

Now fix any $\omega \in \Omega_\infty$. Write $E = \pi(\omega)$ and $\sigma = \Pi(\omega)$. By (53), $\Pi_*(\omega|_n) \triangleleft \sigma$ for any n and $k_n := |\Pi_*(\omega|_n)| = n$ or $n + 1$. So

$$E = \pi(\omega) = \bigcap_{n \geq 0} I_{\omega|_n} = \bigcap_{n \geq 0} B_{\sigma|_{k_n}}.$$

By (25), $|B_{\sigma|_{k_n}}| \rightarrow 0$ as $n \rightarrow \infty$. Since $z_{\sigma|_{k_n}} \in B_{\sigma|_{k_n}}$, we have $z_{\sigma|_{k_n}} \rightarrow E$. Since N is continuous, by (82), we have

$$N(E) = \lim_{n \rightarrow \infty} N(z_{\sigma|_{k_n}}) = \lim_{n \rightarrow \infty} \sum_{i=1}^{k_n} \frac{\sigma_i}{2^i} + \frac{1}{2^{k_n+1}} = \sum_{i=1}^{\infty} \frac{\sigma_i}{2^i} = \varepsilon(\sigma).$$

That is, $N(\pi(\omega)) = \varepsilon(\Pi(\omega))$. □

7.2. Gap labelling of σ_λ .

Proof of Theorem 1.11. By Theorem 1.8 ii), v) and (83), we have

$$\{N(G) : G \in \mathcal{G}_I\} = \{N(\pi(\omega)) : \omega \in \mathcal{E}_l^o \cup \mathcal{E}_r^e\} = \{N(z) : z \in \mathcal{Z}\} = \mathcal{D} \cap (0, 1).$$

It is well known that the restriction of ε on $\Sigma_\infty^{(2)}$ is injective and

$$(84) \quad (\mathcal{D}/3 \setminus \mathcal{D}) \cap (0, 1) = \varepsilon(\Sigma_\infty^{(2)}).$$

By Theorem 1.8 iii), Theorem 1.10, Proposition 4.5 v), injectivity of ε on $\Sigma_\infty^{(2)}$ and (84), we have

$$\begin{aligned} \{N(G) : G \in \mathcal{G}_{II}\} &= \{N(\pi(\omega)) : \omega \in \tilde{\mathcal{E}}_l\} = \{\varepsilon(\Pi(\omega)) : \omega \in \tilde{\mathcal{E}}_l\} = \varepsilon(\Pi(\tilde{\mathcal{E}}_l)) \\ &= \varepsilon(\Sigma_\infty^{(2)} \setminus \Pi(\mathcal{F})) = \varepsilon(\Sigma_\infty^{(2)}) \setminus \varepsilon(\Pi(\mathcal{F})) \\ &= [(\mathcal{D}/3 \setminus \mathcal{D}) \cap (0, 1)] \setminus \varepsilon(\Pi(\mathcal{F})). \end{aligned}$$

So the result follows. \square

8. PROOF OF TECHNICAL LEMMAS AND RELATED RESULTS

In this section, we give the proofs of the technical lemmas and related results in Sec. 2 and Sec. 3.

8.1. The properties of \mathcal{Z} .

8.1.1. The order of \mathcal{Z} .

The following lemma tells us how the zeros in \mathcal{R}_n are ordered.

Lemma 8.1. $\#\mathcal{R}_n = 2^{n+1} - 1$. Moreover \mathcal{Z}_n interlaces \mathcal{R}_{n-1} . That is, if we list the elements of \mathcal{R}_{n-1} as

$$r_1^{(n-1)} < r_2^{(n-1)} < \dots < r_{2^{n-1}}^{(n-1)},$$

then \mathcal{R}_n is ordered as

$$(85) \quad z_{\varpi_n^{-1}(0)} < r_1^{(n-1)} < z_{\varpi_n^{-1}(1)} < r_2^{(n-1)} < \dots < z_{\varpi_n^{-1}(2^{n-2})} < r_{2^{n-1}}^{(n-1)} < z_{\varpi_n^{-1}(2^{n-1})}.$$

In particular, for any $\sigma \in \Sigma_n$ we have

$$(86) \quad B_\sigma \subset [r_{\varpi_n(\sigma)}^{(n-1)}, r_{\varpi_n(\sigma)+1}^{(n-1)}].$$

Here we use the convention: $r_0^{(n-1)} = -\infty$ and $r_{2^n}^{(n-1)} = \infty$.

Proof. Notice that $\#\mathcal{Z}_n = 2^n$. By Lemma 2.5 i), $\mathcal{Z}_n \cap \mathcal{Z}_m = \emptyset$ for $n \neq m$. Hence,

$$\#\mathcal{R}_n = \sum_{j=0}^n \#\mathcal{Z}_j = 2^{n+1} - 1.$$

We prove (85) by induction. By (23),

$$z_\emptyset = \lambda; \quad z_0 = -\sqrt{2 + \lambda^2}; \quad z_1 = \sqrt{2 + \lambda^2}.$$

So $\mathcal{R}_0 = \mathcal{Z}_0 = \{z_\emptyset\}$ and $\mathcal{Z}_1 = \{z_0, z_1\}$. Since $z_0 < z_\emptyset < z_1$, (85) holds for $n = 1$.

Now assume the result holds for $n < k$. We consider \mathcal{R}_k . By Lemma 2.5 i) and induction hypothesis,

$$(87) \quad \begin{cases} \{r_2^{(k-1)}, r_4^{(k-1)}, \dots, r_{2^{k-2}}^{(k-1)}\} = \mathcal{R}_{k-2} \subset \{h_k = 2\}; \\ \{r_1^{(k-1)}, r_3^{(k-1)}, \dots, r_{2^{k-1}}^{(k-1)}\} = \mathcal{Z}_{k-1} \subset \{h_k = -2\}. \end{cases}$$

Thus in each interval $(r_i^{(k-1)}, r_{i+1}^{(k-1)})$ there is an element of \mathcal{Z}_k . Moreover since $h_k(x) \rightarrow +\infty$ when $|x| \rightarrow \infty$, there is one element of \mathcal{Z}_k in $(r_0^{(k-1)}, r_1^{(k-1)})$ and one element of \mathcal{Z}_k in $(r_{2^{k-1}}^{(k-1)}, r_{2^k}^{(k-1)})$. Since $\#\mathcal{Z}_k = 2^k$, we get all elements of \mathcal{Z}_k . Thus the result holds for $n = k$.

By induction, (85) holds.

By (85), $z_\sigma \in [r_{\varpi_n(\sigma)}^{(n-1)}, r_{\varpi_n(\sigma)+1}^{(n-1)}]$. If $\sigma \neq 0^n$ or 1^n , by (87), $h_n(r_{\varpi_n(\sigma)}^{(n-1)}) = \pm 2$ and $r_{\varpi_n(\sigma)+1}^{(n-1)} = \mp 2$. If $\sigma = 0^n$, $h_n(r_0^{(n-1)}) = \infty$ and $h_n(r_1^{(n-1)}) = -2$. If $\sigma = 1^n$, $h_n(r_{2^n}^{(n-1)}) = \infty$ and $h_n(r_{2^n-1}^{(n-1)}) = -2$. By Proposition 2.10 i), (86) holds. \square

Corollary 8.2. *For any $n, t \geq 0$ and $\sigma \in \Sigma_n$, we have*

$$(88) \quad z_{\sigma 0 1^t} < z_\sigma < z_{\sigma 1 0^t}.$$

Proof. Fix any $\sigma \in \Sigma_n$.

Claim: $z_{\sigma 0} < z_\sigma < z_{\sigma 1}$ are three consecutive numbers in \mathcal{R}_{n+1} .

\triangleleft By (85), z_σ is the $2\varpi_n(\sigma) + 1$ -th number in \mathcal{R}_n . So

$$(89) \quad z_\sigma = r_{2\varpi_n(\sigma)+1}^{(n)} = r_{\varpi_{n+1}(\sigma 0)+1}^{(n)} = r_{\varpi_{n+1}(\sigma 1)}^{(n)}.$$

By applying (85) for \mathcal{R}_{n+1} , we conclude that

$$z_{\sigma 0} < r_{\varpi_{n+1}(\sigma 1)}^{(n)} = z_\sigma < z_{\sigma 1}$$

are three consecutive numbers in \mathcal{R}_{n+1} . \triangleright

Applying the claim to $\sigma 0$, we know that $z_{\sigma 0} < z_{\sigma 0 1}$ are two consecutive numbers in \mathcal{R}_{n+2} . Since $z_\sigma \in \mathcal{R}_{n+2}$ and $z_{\sigma 0} < z_\sigma$, we have $z_{\sigma 0} < z_{\sigma 0 1} < z_\sigma$. A symmetric argument shows that $z_\sigma < z_{\sigma 1 0}$. We can continue this process. By an inductive argument, we get (88). \square

Proof of Proposition 2.6. Since the map $\sigma \rightarrow z_\sigma$ is a bijection between Σ_* and \mathcal{Z} , we only need to show that if $\sigma \prec \tau$, then $z_\sigma < z_\tau$.

Assume $\sigma \prec \tau$ and $|\sigma| = m, |\tau| = n$. Write $\theta = \sigma \wedge \tau$.

If $|\theta| < m, n$, then $\sigma = \theta 0^* \leq \theta 0 1^{m-1-|\theta|} \in \Sigma_m$ and $\tau = \theta 1^* \geq \theta 1 0^{n-1-|\theta|} \in \Sigma_n$. By (29) and Corollary 8.2,

$$z_\sigma \leq z_{\theta 0 1^{m-1-|\theta|}} < z_\theta < z_{\theta 1 0^{n-1-|\theta|}} \leq z_\tau.$$

If $|\theta| = m$, then $\sigma \triangleleft \tau$. Since $\sigma \prec \tau$, we have $\tau = \sigma 1^* \geq \sigma 1 0^{n-1-m}$. Again by (29) and Corollary 8.2,

$$z_\sigma < z_{\sigma 1 0^{n-1-m}} \leq z_\tau.$$

If $|\theta| = n$, the same proof shows that $z_\sigma < z_\tau$. \square

Proof of Corollary 2.7. By (86),

$$\text{int}(B_\sigma) \subset (r_{\varpi_n(\sigma)}^{(n-1)}, r_{\varpi_n(\sigma)+1}^{(n-1)}) =: J_\sigma.$$

So we only need to show the following claim.

Claim: If $\sigma_n = 0 (= 1)$, then $\prod_{j=0}^{n-1} h_j(E) < 0 (> 0)$ for any $E \in J_\sigma$.

\triangleleft We show it by induction. If $n = 1$, then $J_0 = (-\infty, \lambda)$ and $J_1 = (\lambda, \infty)$. Since $h_0(E) = E - \lambda$, $h_0(E) < 0$ for $E \in J_0$ and $h_0(E) > 0$ for $E \in J_1$.

Assume the result holds for $n = k$. Now fix any $\sigma \in \Sigma_k$. By (85), for any j we have

$$r_j^{(k-1)} = r_{2^j}^{(k)}.$$

Together with (89), we have

$$\begin{aligned} J_{\sigma 0} &= (r_{\varpi_{k+1}(\sigma 0)}^{(k)}, r_{\varpi_{k+1}(\sigma 0)+1}^{(k)}) = (r_{\varpi_k(\sigma)}^{(k-1)}, z_\sigma), \\ J_{\sigma 1} &= (r_{\varpi_{k+1}(\sigma 1)}^{(k)}, r_{\varpi_{k+1}(\sigma 1)+1}^{(k)}) = (z_\sigma, r_{\varpi_k(\sigma)+1}^{(k-1)}). \end{aligned}$$

Take $E \in \text{int}(J_{\sigma 0})$. If $\sigma_k = 0$, then by the induction hypothesis, $\prod_{j=0}^{k-1} h_j(E) < 0$. By Proposition 2.10 i), h_k is decreasing in a neighbourhood of z_σ . Since $z_{\sigma^-} < r_{\varpi_k(\sigma)}^{(k-1)} < z_\sigma$, we conclude that $h_k(E) > 0$. If $\sigma_k = 1$, then by the induction hypothesis, $\prod_{j=0}^{k-1} h_j(E) > 0$. By a symmetric argument, $h_k(E) < 0$. So we always have $\prod_{j=0}^k h_j(E) < 0$.

Take $E \in \text{int}(J_{\sigma 1})$. By the same argument, we can show that $\prod_{j=0}^k h_j(E) > 0$.

By induction, the claim holds. \triangleright \square

8.1.2. Local behaviors of trace polynomials at zeros.

Next we study the monotone properties of the trace polynomials around a zero.

Lemma 8.3. *Assume $n \geq 0$ and $\sigma \in \Sigma_n$. Then*

$$(90) \quad \text{sgn}(h'_{n+1}(z_\sigma)) = (-1)^n; \quad \text{sgn}(h'_{n+k}(z_\sigma)) = (-1)^{n+1}, \quad (k \geq 2).$$

Proof. If $n = 0$, then $z_\sigma = z_\emptyset = \lambda$ and $h'_{n+1}(z_\sigma) = h'_1(\lambda) = 2\lambda > 0$.

For $n \geq 1$, by the recurrence relation (23), we have,

$$(91) \quad h'_{n+1}(E) = h'_n(E)(h_{n-1}^2(E) - 2) + 2h_n(E)h_{n-1}(E)h'_{n-1}(E).$$

Take $E = z_\sigma$. Since $h_n(z_\sigma) = 0$, we get

$$(92) \quad h'_{n+1}(z_\sigma) = h'_n(z_\sigma)(h_{n-1}^2(z_\sigma) - 2).$$

By Lemma 2.9, we have

$$(93) \quad -(h_{n-1}^2(z_\sigma) - 2) = h_n(z_\sigma) - (h_{n-1}^2(z_\sigma) - 2) = (-1)^{n-1}2\lambda \prod_{c \in \mathcal{R}_{n-1}} (z_\sigma - c).$$

By Proposition 2.10 i),

$$\text{sgn}(h'_n(z_\sigma)) = (-1)^{\sigma_n+1}.$$

By (85) and (26),

$$\operatorname{sgn} \left(\prod_{c \in \mathcal{R}_{n-1}} (z_\sigma - c) \right) = (-1)^{\#\mathcal{R}_{n-1} - \varpi_n(\sigma)} = (-1)^{\sigma_{n+1}}.$$

Combining (92) and (93), the sign of $h'_{n+1}(z_\sigma)$ is $(-1)^n$. So the first equation of (90) holds.

By Lemma 2.5 i), $h_{n+1}(z_\sigma) = -2, h_{n+k}(z_\sigma) = 2$ for $k \geq 2$. Then by (91), for any $n \geq 0$,

$$\begin{cases} h'_{n+2}(z_\sigma) = -2h'_{n+1}(z_\sigma); \\ h'_{n+3}(z_\sigma) = 2h'_{n+2}(z_\sigma) - 8h'_{n+1}(z_\sigma); \\ h'_{n+k}(z_\sigma) = 2h'_{n+k-1}(z_\sigma) + 8h'_{n+k-2}(z_\sigma), \quad (k \geq 4). \end{cases}$$

From this, we conclude that, there exists an increasing positive integer sequence $\{\tau_k : k \geq 2\}$ such that $\tau_2 = 2, \tau_3 = 12, \tau_{k+2} = 2\tau_{k+1} + 8\tau_k, (k \geq 2)$ and

$$h'_{n+k}(z_\sigma) = -\tau_k h'_{n+1}(z_\sigma), \quad k \geq 2.$$

So the second equation of (90) holds. \square

8.1.3. The relation between \mathcal{Z} and band edges.

Proof of Lemma 2.8. i) By Lemma 2.5 i),

$$(94) \quad h_{n+1}(z_\sigma) = -2; \quad h_{n+1+t}(z_\sigma) = 2, \quad \forall t \geq 1.$$

By Proposition 2.10 iii), z_σ must be an endpoint of some band in \mathcal{B}_{n+1+t} . By Corollary 8.2, $z_{\sigma 01^t} < z_\sigma < z_{\sigma 10^t}$. Since $(\sigma 01^t)^+ = \sigma 10^t$, $B_{\sigma 01^t}$ and $B_{\sigma 10^t}$ are two consecutive bands in \mathcal{B}_{n+1+t} . Hence z_σ can only be $b_{\sigma 01^t}$ or $a_{\sigma 10^t}$.

If n is odd, by Lemma 8.3,

$$h'_{n+1}(z_\sigma) < 0 \quad \text{and} \quad h'_{n+1+t}(z_\sigma) > 0 (t \geq 1).$$

By (94) and Proposition 2.10 iii), $z_\sigma = b_{\sigma 01^t}$.

If n is even, by Lemma 8.3,

$$h'_{n+1}(z_\sigma) > 0 \quad \text{and} \quad h'_{n+1+t}(z_\sigma) < 0 (t \geq 1).$$

By (94) and Proposition 2.10 iii), $z_\sigma = a_{\sigma 10^t}$.

In both cases, by Proposition 2.10 ii), $b_{\sigma 01^t} < a_{(\sigma 01^t)^+} = a_{\sigma 10^t}$.

ii) If $a_\sigma \in \mathcal{R}_{n-1}$, then by i), $\sigma = \hat{\sigma} 10^s$ for some $\hat{\sigma} \in \Sigma_*$, $s \geq 0$ and $z_{\hat{\sigma}} = a_{\hat{\sigma} 10^t}$ for any $t \geq 0$. This implies $a_\sigma = a_{\sigma 0^t}$ for any $t > 0$.

If $b_\sigma \in \mathcal{R}_{n-1}$, then by i), $\sigma = \hat{\sigma} 01^s$ for some $\hat{\sigma} \in \Sigma_*$, $s \geq 0$ and $z_{\hat{\sigma}} = b_{\hat{\sigma} 01^t}$ for any $t \geq 0$. This implies $b_\sigma = b_{\sigma 1^t}$ for any $t > 0$.

iii) If $a_\sigma \in \mathcal{R}_{n-1}$, then by ii), $a_{\sigma 0} = a_\sigma \in \mathcal{R}_{n-1} \subset \mathcal{R}_n$. If $a_{\sigma 0} \in \mathcal{R}_n$, then by i), $\sigma 0 = \hat{\sigma} 10^s$ for some $\hat{\sigma} \in \Sigma_*$, $s \geq 0$ and $z_{\hat{\sigma}} = a_{\hat{\sigma} 10^t}$ for any $t \geq 0$. In this case, we must have $s \geq 1$. So we have $a_\sigma = a_{\hat{\sigma} 10^{s-1}} = z_{\hat{\sigma}} \in \mathcal{R}_{n-1}$. Thus the first assertion holds. The same proof shows that the second assertion holds.

iv) By Lemma 8.1, $\#(z_\sigma, z_{\sigma+}) \cap \mathcal{R}_{n-1} = 1$. By Lemma 2.5 ii),

$$((z_\sigma, b_\sigma) \cup (a_{\sigma+}, z_{\sigma+})) \cap \mathcal{R}_{n-1} = \emptyset.$$

By Floquet theory and Lemma 2.5 i),

$$h_n((b_\sigma, a_{\sigma+})) \subset (-\infty, -2) \cup (2, \infty) \quad \text{and} \quad h_n(\mathcal{R}_{n-1}) \subset \{\pm 2\}.$$

So $(b_\sigma, a_{\sigma+}) \cap \mathcal{R}_{n-1} = \emptyset$. Combine these facts, the result follows. \square

Proof of Proposition 2.1. The possible gap of $\sigma_{\lambda, n}$ has the form: $(b_\sigma, a_{\sigma+})$ with $\sigma \neq 1^n$. By Lemma 2.8 iv), $\{b_\sigma, a_{\sigma+}\} \cap \mathcal{R}_{n-1} = \{z_*\}$. By Lemma 8.3, $h'_n(z_*) \neq 0$. By Proposition 2.10 ii), the gap (b_w, a_{w+}) is open. \square

8.2. Proof of Lemma 2.3.

At first we need the following lemma:

Lemma 8.4. *Assume $n \geq 1$ and $\sigma \in \Sigma_n$.*

i) *If n is odd, then*

$$\begin{cases} h_{n+1}(a_\sigma) > 2, & \text{if } a_\sigma \notin \mathcal{R}_{n-1}; \\ h_{n+1}(b_\sigma) < 2, & \text{if } b_\sigma \notin \mathcal{R}_{n-1}. \end{cases}$$

ii) *If n is even, then*

$$\begin{cases} h_{n+1}(b_\sigma) > 2, & \text{if } b_\sigma \notin \mathcal{R}_{n-1}; \\ h_{n+1}(a_\sigma) < 2, & \text{if } a_\sigma \notin \mathcal{R}_{n-1}. \end{cases}$$

Proof. By (85),

$$\#(\mathcal{R}_n \cap (-\infty, z_\sigma)) = 2\varpi_n(\sigma).$$

i) At first assume $a_\sigma \notin \mathcal{R}_{n-1}$. Since $h_n(a_\sigma) = \pm 2$, and $\mathcal{R}_n = \mathcal{R}_{n-1} \cup \mathcal{Z}_n$, $a_\sigma \notin \mathcal{R}_n$. By Lemma 2.9, we have

$$h_{n+1}(a_\sigma) = h_n^2(a_\sigma) - 2 + (-1)^n 2\lambda \prod_{c \in \mathcal{R}_n} (a_\sigma - c) = 2 - 2\lambda \prod_{c \in \mathcal{R}_n} (a_\sigma - c).$$

By Lemma 2.5 ii), $(a_\sigma, z_\sigma) \cap \mathcal{R}_n = \emptyset$. By Lemma 8.1, we have

$$\begin{aligned} \#(\mathcal{R}_n \cap (a_\sigma, +\infty)) &= 2^{n+1} - 1 - \#(\mathcal{R}_n \cap (-\infty, a_\sigma]) \\ &= 2^{n+1} - 1 - \#(\mathcal{R}_n \cap (-\infty, z_\sigma)) = 2^{n+1} - 1 - 2\varpi_n(\sigma). \end{aligned}$$

Thus $\prod_{c \in \mathcal{R}_n} (a_\sigma - c) < 0$, and consequently $h_{n+1}(a_\sigma) > 2$.

Now assume $b_\sigma \notin \mathcal{R}_{n-1}$. Then by the same argument, $b_\sigma \notin \mathcal{R}_n$. Still by Lemma 2.9,

$$h_{n+1}(b_\sigma) = h_n^2(b_\sigma) - 2 + (-1)^n 2\lambda \prod_{c \in \mathcal{R}_n} (b_\sigma - c) = 2 - 2\lambda \prod_{c \in \mathcal{R}_n} (b_\sigma - c).$$

By Lemma 2.5 ii), $[z_\sigma, b_\sigma] \cap \mathcal{R}_n = [z_\sigma, b_\sigma] \cap \mathcal{R}_n = \{z_\sigma\}$. By Lemma 8.1, we have

$$\begin{aligned} \#(\mathcal{R}_n \cap (b_\sigma, +\infty)) &= 2^{n+1} - 1 - \#(\mathcal{R}_n \cap (-\infty, b_\sigma]) \\ &= 2^{n+1} - 2 - \#(\mathcal{R}_n \cap (-\infty, z_\sigma)) = 2^{n+1} - 2 - 2\varpi_n(\sigma). \end{aligned}$$

Thus $\prod_{c \in \mathcal{R}_n} (b_\sigma - c) > 0$, and consequently $h_{n+1}(b_\sigma) < 2$.

ii) The proof is the same as i). □

Proof of Lemma 2.3. i) By Lemma 2.8 i), $b_{\sigma 0} = z_{\sigma} \in \mathcal{R}_n$. By Lemma 2.8 iv),

$$a_{\sigma 1} = a_{(\sigma 0)^+} \notin \mathcal{R}_n.$$

If $a_{\sigma} \in \mathcal{R}_{n-1}$, by Lemma 2.8 ii), $a_{\sigma} = a_{\sigma 0}$. Hence $B_{\sigma 0} = [a_{\sigma 0}, b_{\sigma 0}] = [a_{\sigma}, z_{\sigma}]$.

If $a_{\sigma} \notin \mathcal{R}_{n-1}$, by Lemma 8.4 i), $h_{n+1}(a_{\sigma}) > 2$. By Proposition 2.10 i), $h_{n+1}(B_{\sigma 0}) = [-2, 2]$. Since $a_{\sigma} < z_{\sigma} = b_{\sigma 0}$, we conclude that $a_{\sigma} < a_{\sigma 0}$. Hence $B_{\sigma 0} \subset (a_{\sigma}, z_{\sigma}]$.

In both cases, we have $B_{\sigma 0} \subset B_{\sigma}$.

By Proposition 2.1, $b_{\sigma 0} < a_{(\sigma 0)^+} = a_{\sigma 1}$. So $z_{\sigma} = b_{\sigma 0} < a_{\sigma 1}$.

If $b_{\sigma} \in \mathcal{R}_{n-1}$, by Lemma 2.8 ii), $b_{\sigma} = b_{\sigma 1}$. So we have $B_{\sigma 1} \subset (z_{\sigma}, b_{\sigma}]$.

If $b_{\sigma} \notin \mathcal{R}_{n-1}$, By Lemma 8.4 i), $h_{n+1}(b_{\sigma}) < 2$. If $\sigma = 1^n$, then $b_{\sigma 1} = b_{1^{n+1}}$ is the largest root of $h_{n+1}(x) = 2$. Since $h_{n+1}(b_{\sigma}) < 2$ and $h_{n+1}(x) \rightarrow \infty$ when $x \rightarrow \infty$, we conclude that $b_{\sigma} < b_{\sigma 1}$. Now assume $\sigma < 1^n$. By Lemma 2.8 iv), $a_{\sigma^+} \in \mathcal{R}_{n-1}$. By Lemma 2.8 ii), $a_{\sigma^+} = a_{\sigma+0}$. By Proposition 2.1,

$$b_{\sigma} < a_{\sigma^+}; \quad b_{\sigma 1} < a_{\sigma+0} = a_{\sigma^+}.$$

By Proposition 2.10 i), h_{n+1} is increasing on $B_{\sigma 1}$ and decreasing on $B_{\sigma+0}$, so

$$h_{n+1}([b_{\sigma 1}, a_{\sigma^+}]) = h_{n+1}([b_{\sigma 1}, a_{\sigma+0}]) \subset [2, \infty).$$

Since $h_{n+1}(b_{\sigma}) < 2$, we conclude that

$$(95) \quad b_{\sigma} < b_{\sigma 1} < a_{\sigma^+}.$$

Together with $z_{\sigma} < a_{\sigma 1}$, we always have $[z_{\sigma}, b_{\sigma}] \prec B_{\sigma 1}$.

ii) The proof is the same as i). In particular, we have that if $\sigma > 0^n$, then

$$(96) \quad b_{\sigma^-} < a_{\sigma 0} < a_{\sigma}.$$

iii) By Proposition 2.1, $B_{\sigma 0} < B_{\sigma 1}$. If n is odd and $\sigma > 0^n$, then $B_{\sigma^-} < B_{\sigma}$ and $B_{\sigma 0} \subset B_{\sigma}$. Hence $B_{\sigma^-} < B_{\sigma 0}$. If n is even and $\sigma > 0^n$, then by (96), $B_{\sigma^-} < B_{\sigma 0}$. If n is odd and $\sigma < 1^n$, then by (95), $B_{\sigma 1} < B_{\sigma^+}$. If n is even and $\sigma < 1^n$, then $B_{\sigma} < B_{\sigma^+}$ and $B_{\sigma 1} \subset B_{\sigma}$. Hence $B_{\sigma 1} < B_{\sigma^+}$. □

8.3. Proof of Lemma 3.1 and Corollary 3.2.

Proof of Lemma 3.1. i) Applying Lemma 2.3 i) to σ and ii) to $\hat{\sigma} = \sigma 0$, we have

$$a_{\sigma 1} \notin \mathcal{R}_n, \quad B_{\sigma 0} \subset B_{\sigma} \quad \text{and} \quad B_{\sigma 0 1} \subset B_{\sigma 0}.$$

Applying Lemma 2.8 iii) and Lemma 2.3 ii), iii) to $\hat{\sigma} = \sigma 1$, we have

$$(97) \quad a_{\sigma 10}, b_{\sigma 10} \notin \mathcal{R}_{n+1}, \quad B_{\sigma 0} = B_{(\sigma 1)^-} < B_{\sigma 10} \prec B_{\sigma 1}.$$

If $a_{\sigma} \in \mathcal{R}_{n-1}$, by Lemma 2.8 ii), $a_{\sigma 0} = a_{\sigma} \in \mathcal{R}_{n-1} \subset \mathcal{R}_n$. Applying Lemma 2.3 ii) to $\hat{\sigma} = \sigma 0$, we have

$$(98) \quad B_{\sigma 0 0} \subset B_{\sigma 0} \subset B_{\sigma}.$$

If $a_\sigma \notin \mathcal{R}_{n-1}$, by Lemma 2.8 iii), $a_{\sigma 0} \notin \mathcal{R}_n$ and hence $a_{\sigma 00} \notin \mathcal{R}_{n+1}$. Applying Lemma 2.3 ii) to $\hat{\sigma} = \sigma 0$, we have

$$b_{\sigma 00} \notin \mathcal{R}_{n+1}, \quad B_{\sigma 00} \prec B_{\sigma 0}.$$

Now we show $a_\sigma < a_{\sigma 00}$. By Lemma 8.4 i), $h_{n+1}(a_\sigma) > 2$. Hence

$$h_{n+2}(a_\sigma) = h_{n+1}(a_\sigma)(h_n^2(a_\sigma) - 2) - 2 > 2.$$

Since $a_{\sigma 01} \in B_{\sigma 01} \subset B_{\sigma 0} \subset B_\sigma$, we have $a_\sigma \leq a_{\sigma 01}$. By Proposition 2.10 i),

$$h_{n+2}(a_{\sigma 01}) = -2.$$

Since $a_{\sigma 00}$ is the maximal solution of $h_{n+2}(E) = 2$ which are less than $a_{\sigma 01}$, we conclude that $a_\sigma < a_{\sigma 00}$. Since $b_{\sigma 00} < b_{\sigma 01}$ and by Lemma 2.8 i), $b_{\sigma 01} = b_{\sigma 0} = z_\sigma$, we have

$$(99) \quad B_{\sigma 00} = [a_{\sigma 00}, b_{\sigma 00}] \subset (a_\sigma, z_\sigma) \subset \text{int}(B_\sigma).$$

Applying Lemma 2.3 ii) to $\hat{\sigma} = \sigma 1$, we always have

$$B_{\sigma 11} \subset B_{\sigma 1}.$$

If $b_\sigma \in \mathcal{R}_{n-1}$, by Lemma 2.3 i), $B_{\sigma 1} \subset B_\sigma$. Hence

$$B_{\sigma 11} \subset B_{\sigma 1} \subset B_\sigma.$$

By (97), $B_{\sigma 0} < B_{\sigma 10} \prec B_{\sigma 1}$. Since $B_{\sigma 0}, B_{\sigma 1} \subset B_\sigma$, we have

$$(100) \quad B_{\sigma 10} = [a_{\sigma 10}, b_{\sigma 10}] \subset (a_{\sigma 0}, b_{\sigma 1}) \subset \text{int}(B_\sigma).$$

If $b_\sigma \notin \mathcal{R}_{n-1}$, we claim that

$$b_{\sigma 10} < b_\sigma < b_{\sigma 11}.$$

In fact, by Lemma 2.8 iv), either $\sigma = 1^n$, or $a_{\sigma+} \in \mathcal{R}_{n-1}$. By Lemma 8.4 i), $h_{n+1}(b_\sigma) < 2$. Hence

$$h_{n+2}(b_\sigma) = h_{n+1}(b_\sigma)(h_n^2(b_\sigma) - 2) - 2 < 2.$$

If $\sigma = 1^n$, then since $h_{n+2}(E) \rightarrow \infty$ as $E \rightarrow \infty$, there is a root of $h_{n+2}(E) = 2$ in (b_σ, ∞) . Since $b_{\sigma 11} = b_{1^{n+2}}$ is the largest root of $h_{n+2}(E) = 2$, we conclude that $b_\sigma < b_{\sigma 11}$. Now assume $\sigma \neq 1^n$, then $a_{\sigma+} \in \mathcal{R}_{n-1}$. By Lemma 2.8 ii), $a_{\sigma+0^t} = a_{\sigma+}$. By Proposition 2.10 i), h_{n+2} is increasing on $B_{\sigma 11}$ and decreasing on $B_{(\sigma 11)^+} = B_{\sigma+00}$. Hence by Floquet theory,

$$h_{n+2}([b_{\sigma 11}, a_{\sigma+}]) = h_{n+2}([b_{\sigma 11}, a_{\sigma+00}]) \subset [2, \infty).$$

Since $b_\sigma < a_{\sigma+}$ and $h_{n+2}(b_\sigma) < 2$, we conclude that $b_\sigma < b_{\sigma 11} < a_{\sigma+}$. As a result, we always have $b_\sigma < b_{\sigma 11}$.

On the other hand, we have

$$-2 = h_{n+2}(b_{\sigma 10}) = h_{n+1}(b_{\sigma 10})(h_n^2(b_{\sigma 10}) - 2) - 2.$$

So $h_{n+1}(b_{\sigma 10}) = 0$ or $|h_n(b_{\sigma 10})| = \sqrt{2}$. By (97), $b_{\sigma 10} \notin \mathcal{R}_{n+1}$. So $|h_n(b_{\sigma 10})| = \sqrt{2}$. If $\sigma = 1^n$, then since $h_n(E) \geq 2$ for any $E \geq b_\sigma$, we have $b_{\sigma 10} < b_\sigma$. Now assume $\sigma \neq 1^n$. By Floquet theory, $h_n([b_\sigma, a_{\sigma+}]) \cap (-2, 2) = \emptyset$. Since $b_{\sigma 10} < b_{\sigma 11} < a_{\sigma+00} = a_{\sigma+}$ and $h_n(b_{\sigma 10}) \in (-2, 2)$, we conclude that $b_{\sigma 10} < b_\sigma$. As a result, we always have $b_{\sigma 10} < b_\sigma$. Hence the claim holds.

By (97), $B_{\sigma 0} < B_{\sigma 10}$, so we have $a_\sigma < z_\sigma = b_{\sigma 0} < a_{\sigma 10} < a_{\sigma 11}$. Together with the claim above, we have

$$B_\sigma \prec B_{\sigma 11}.$$

By $B_{\sigma 0} < B_{\sigma 10}$ and the claim above, we also conclude that

$$(101) \quad B_{\sigma 10} = [a_{\sigma 10}, b_{\sigma 10}] \subset (b_{\sigma 0}, b_\sigma) = (z_\sigma, b_\sigma) \subset \text{int}(B_\sigma).$$

By (98) and (99), $B_{\sigma 00} \subset B_\sigma$ always holds.

By (100) and (101), $B_{\sigma 10} \subset \text{int}(B_\sigma)$ always holds.

ii) The proof is the same as i). □

Proof of Corollary 3.2. i) By Lemma 2.5 ii), $\text{int}(B_\sigma) \cap \mathcal{R}_{n-1} = \emptyset$. So $B_\sigma \cap \mathcal{R}_{n-1} \neq \emptyset$ if and only if $\partial B_\sigma \cap \mathcal{R}_{n-1} \neq \emptyset$. Write $\hat{\sigma} := \sigma|_{n-1}$.

At first assume $\partial B_\sigma \cap \mathcal{R}_{n-1} \neq \emptyset$. If $a_\sigma \in \mathcal{R}_{n-1}$, then by Lemma 2.8 i), there exist $m < n$ even and $\tau \in \Sigma_m$ such that $\sigma = \tau 10^{n-m-1}$ and $z_\tau = a_{\tau 10^{n-m-1}} = a_\sigma$. If $m = n-1$, applying Lemma 2.3 ii) to τ , we have $B_\sigma = B_{\tau 1} \subset B_\tau = B_{\hat{\sigma}}$. If $m \leq n-2$, then $a_{\hat{\sigma}} = z_\tau \in \mathcal{R}_{n-2}$. By Lemma 2.3, we always have $B_\sigma = B_{\hat{\sigma} 0} \subset B_{\hat{\sigma}}$. If $b_\sigma \in \mathcal{R}_{n-1}$, the same proof shows that $B_\sigma \subset B_{\hat{\sigma}}$.

Next assume $B_\sigma \subset B_{\hat{\sigma}}$. If n is even, by applying Lemma 2.3 i) to $\hat{\sigma}$, either $\sigma = \hat{\sigma} 0$, hence $b_\sigma = z_{\hat{\sigma}} \in \mathcal{R}_{n-1}$; or $\sigma = \hat{\sigma} 1$, hence $b_{\hat{\sigma}} \in \mathcal{R}_{n-2}$. In the latter case, by Lemma 2.8 ii), $b_\sigma = b_{\hat{\sigma} 1} = b_{\hat{\sigma}} \in \mathcal{R}_{n-1}$. Thus we always have $\partial B_\sigma \cap \mathcal{R}_{n-1} \neq \emptyset$. If n is odd, the same proof shows that $\partial B_\sigma \cap \mathcal{R}_{n-1} \neq \emptyset$.

ii) Write $\hat{\sigma} := \sigma|_{n-1}$ and $\tilde{\sigma} := \sigma|_{n-2}$. Assume $B_\sigma \not\subset B_{\hat{\sigma}}$. If $n = 1$, by the convention, $B_{\hat{\sigma}} = B_\emptyset$ and $B_{\tilde{\sigma}} = \mathbb{R}$. So $B_\sigma \subset B_{\tilde{\sigma}}$. Now assume $n \geq 2$. At first, assume n is odd. By applying Lemma 2.3 ii) to $\hat{\sigma}$, the only possibility is that $a_{\hat{\sigma}} \notin \mathcal{R}_{n-2}$ and $\sigma = \hat{\sigma} 0$. So $\sigma = \tilde{\sigma} \sigma_{n-1} 0$. Now by applying Lemma 3.1 i) to $\tilde{\sigma}$, we conclude that $B_\sigma = B_{\tilde{\sigma} \sigma_{n-1} 0} \subset B_{\tilde{\sigma}}$. If n is even, the same proof show that $B_\sigma \subset B_{\tilde{\sigma}}$. So ii) follows.

iii) At first assume $\tau \in \Sigma_n$. By Proposition 2.1, we must have $\sigma = \tau$.

Next assume $\tau \in \Sigma_{n+1}$. By Lemma 2.3 iii), if $\sigma \neq \tau|_n$, then $B_\tau \cap B_\sigma = \emptyset$. So we must have $\sigma = \tau|_n$.

Now assume $\tau \in \Sigma_{n+2}$. By ii), $B_\tau \subset B_{\tau|_{n+1}}$ or $B_\tau \subset B_{\tau|_n}$. If $B_\tau \subset B_{\tau|_{n+1}}$, again by Lemma 2.3 iii), if $\sigma \neq \tau|_n$, then $B_{\tau|_{n+1}} \cap B_\sigma = \emptyset$. Consequently $B_\tau \cap B_\sigma = \emptyset$. So we must have $\sigma = \tau|_n$. If $B_\tau \subset B_{\tau|_n}$, by Proposition 2.1, we must have $\sigma = \tau|_n$.

As a result, $\sigma = \tau|_n \triangleleft \tau$. □

Acknowledgement. Liu was supported by the National Natural Science Foundation of China, No. 11871098. Qu was supported by the National Natural Science Foundation of China, No. 11790273 and No. 11871098. Yao was supported by the National Natural Science Foundation of China, No.11901311 and key technologies research and development program, 2020YFA0713300.

APPENDIX

For the reader's convenience, we include two tables of indexes in this appendix. One is for the notations used in this paper, another is for the various orders used in this paper.

Table I: Index of notations

ξ, V_ξ	P-D sequence and potential, see Sec. 1.2.1
$H_\lambda, \sigma_\lambda$	PDH, the spectrum, see (4)
$h_n(E)$	trace polynomial, see (5)
$\mathcal{O}(E)$	trace orbit of E , see (6)
\mathcal{E}_∞	the set of ∞ -type energies, see (8)
$\mathcal{A}, \mathbb{G}, A = [a_{\alpha\beta}]$	alphabet, graph, adjacency matrix, see Sect. 1.2.4
Ω_A, Ω_∞	symbolic spaces, see Sect. 1.2.4
Ω_n, Ω_*	the set of admissible words, see (42)
$\mathcal{Z}^o, \mathcal{Z}^e, \mathcal{Z}, \mathcal{Z}_n, \mathcal{R}_n$	zeros of $\{h_n : n \geq 0\}$, see (12), (28)
$\mathcal{E}_l^o, \mathcal{E}_r^o, \mathcal{E}_l^e, \mathcal{E}_r^e, \tilde{\mathcal{E}}_l, \tilde{\mathcal{E}}_r, \mathcal{F}$	eventually 2-periodic codings, see (13)
ω_*, ω^*	minimum and maximum of Ω_∞ , see (14)
$\mathcal{G}, \mathcal{G}_I, \mathcal{G}_{II}, \mathcal{G}_I^o, \mathcal{G}_I^e$	the set of gaps of σ_λ , see Theorem 1.8
$\mathcal{G}, \mathcal{G}_I, \mathcal{G}_{II}, \mathcal{G}_I^o, \mathcal{G}_I^e$	the set of gaps of Ω_∞ , see Theorem 4.3
$\Sigma_n, \Sigma_*, \Sigma_*^o, \Sigma_*^e, \Sigma_\infty, \Sigma_\infty^{(2)}$	binary trees, see (15), (33), (54), Sec. 2.1.2
$\mathcal{I}, A(\mathcal{I})$	NS(SNS), limit set, see Sec. 1.3.1
$\sigma_{\lambda,n}$	n -th approximation of σ_λ , see (18),(27)
\mathcal{B}_n	family of bands of level n , see (19)
\mathcal{B}_n	optimal covering of level n , see (20)
$\tilde{\mathcal{A}}, \tilde{\mathbb{G}}, \tilde{\mathcal{B}}$	sub-alphabet, graph, NS, see Sec. 1.3.4 and 5.2.2
\mathbb{M}_n, ϖ_n	see Sec. 2.1.2, (26)
$\sigma^+, \sigma^-, \sigma _k, \sigma \wedge \tau, \sigma \triangleleft \tau, \sigma $	see Sec. 2.1.2
$B_\sigma, a_\sigma, b_\sigma, z_\sigma$	band, left and right endpoint, zero, see Sec. 2.1.2
I_w	rename of band, see Sec. 5.1.1
$\pi : (\Omega_\infty, \preceq) \rightarrow (\sigma_\lambda, \leq)$	coding map, see Sec. 1.2.4 and 5.1.2
$\Pi : (\Omega_\infty, \preceq) \rightarrow (\Sigma_\infty, \leq)$	another coding map, see Sec. 1.2.6 and 4.2.2
$\varepsilon : \Sigma_\infty \rightarrow [0, 1]$	evaluation map, see Sec. 1.2.6
$N : \sigma_\lambda \rightarrow [0, 1], N(E), N(G)$	IDS, see Sec. 1.2.6
\mathcal{L}, Π_*	see Sec. 4.2.2
l, l^o, l^e	see Sec. 4.2

In this paper, two types of orders are defined in various spaces: one is \leq , another is \preceq . When \leq is used, it means standard or strong, depending on the context. When \preceq is used, it means non standard or weak.

Table II: Index for various orders

$I < J, I \prec J$	two orders for bands, see (21)
$\sigma \leq \tau$	lexicographical order on Σ_n and Σ_∞ , see Sec. 2.1.2, 4.2.2
$\sigma \preceq \tau$	total order on Σ_* , see (34)
$\alpha < \beta, \alpha \prec \beta$	two partial orders on \mathcal{A} , see (43), (44)
$\omega \leq \omega', \omega \preceq \omega'$	two partial orders on Ω_∞ , see Sec. 4.1.2
$(x, y) \leq (x', y')$	partial order on \mathbb{R}^2 , see Sec. 6.2.1

REFERENCES

- [1] Axel, F., Allouche, J.P., Kleman, M., Mendes-France, M., Peyriere, J., Vibrational modes in a one-dimensional “quasi alloy”, the Morse case. *J. de Phys. C3*, 47,(1986), 181-187.
- [2] Axel, F., Peyrière, J., Extended states in a chain with controlled disorder. *C. R. Acad. Sci. Paris Ser. II Mec. Phys. Chim. Sci. Univers Sci. Terre* 306, 179-182 (1988)
- [3] Axel, F., Peyrière, J., Spectrum and extended states in a harmonic chain with controlled disorder: Effects of the Thue-Morse symmetry. *J. Statist. Phys.* 57, 1013-1047 (1989)
- [4] Bellissard, J., Spectral properties of Schrödinger operator with a Thue-Morse potential. In: *Number Theory and Physics* (Les Houches, 1989), Springer Proc. Phys. 47, pp. 140-150. Springer, Berlin (1990)
- [5] Bellissard, J., Gap labelling theorems for Schrödinger operators. *From number theory to physics* (Les Houches, 1989), 538-630, Springer, Berlin, 1992.
- [6] Bellissard J., Bovier A., Ghez J. M., Spectral properties of a tight binding Hamiltonian with period doubling potential, *Commun. Math. Phys.*, 135(1991), 379–399.
- [7] Bovier A., Ghez J. M., Spectral properties of one-dimensional Schrödinger operators with potentials generated by substitutions, *Commun. Math. Phys.* 158(1993), 45-66.
- [8] Cantat, S., Bers and Hénon, Painlevé and Schrödinger. *Duke Math. J.* 149 (2009), no. 3, 411-460.
- [9] Carvalho T. O., Forward boundedness of the trace map for the period doubling substitution, *non-linearity*, 33(2020), 4390-4406.
- [10] Casdagli, M., Symbolic dynamics for the renormalization map of a quasiperiodic Schrödinger equation, *Comm. Math. Phys.* 107 (1986), no. 2, 295-318.
- [11] Damanik, D., Substitution Hamiltonians with bounded trace map orbits. *J. Math. Anal. Appl.* 249 (2000), no. 2, 393-411.
- [12] Damanik, D., Embree, M., Gorodetski, A., Tcheremchantsev, S.: The fractal dimension of the spectrum of the Fibonacci Hamiltonian. *Commun. Math. Phys.* 280, 499-516 (2008)
- [13] Damanik, D., Gorodetski, A.: Hyperbolicity of the trace map for the weakly coupled Fibonacci Hamiltonian. *Nonlinearity* 22, 123-143 (2009)
- [14] Damanik, D., Gorodetski, A.: Spectral and quantum dynamical properties of the weakly coupled Fibonacci Hamiltonian. *Commun. Math. Phys.* 305, 221-277 (2011)
- [15] Damanik, D., Gorodetski, A.; The density of states measure of the weakly coupled Fibonacci Hamiltonian. *Geom. Funct. Anal.* 22 (2012), no. 4, 976-989.
- [16] Damanik, D.; Gorodetski, A.; Yessen, W., The Fibonacci Hamiltonian. *Invent. Math.* 206 (2016), no. 3, 629-692.
- [17] Kohmoto, M., Kadanoff, L., Tang, C., Localization problem in one dimension: mapping and escape, *Phys. Rev. Lett.* 50 (1983), no. 23, 1870-1872.
- [18] Jitomirskaya, S., Last, Y.: Power-law subordinacy and singular spectra. II. Line operators. *Commun. Math. Phys.* 211, 643-658 (2000)
- [19] Lenz D., Singular spectrum of Lebesgue measure zero for one-dimensional quasicrystals, *Commun. Math. Phys.* 227(2002), 119–130.
- [20] Liu, Q.-H., Qu, Y.-H., On the Hausdorff dimension of the spectrum of Thue-Morse Hamiltonian, *Comm. Math. Phys.* 338 (2015), no. 2, 867-891.

- [21] Liu Q.-H., Qu Y.- H., Yao X., Unbounded trace orbits of Thue-Morse Hamiltonian, *J. Stat. Phys.*, 166(2017), 1509-1557.
- [22] Liu Q.- H., Tan B., Wen Z.- X., Wu J., Measure zero spectrum of a class of Schrödinger operators, *J. Stat. Phys.* 106(2002), 681-691.
- [23] Luck, J. M., Cantor spectra and scaling of gap widths in deterministic aperiodic systems, *Phys. Rev. B* 39, 5834-5849 (1989)
- [24] Ostlund, S., Pandit, R., Rand, D., Schellnhuber, H., Siggia, D., One-dimensional Schrödinger equation with an almost periodic potential, *Phys. Rev. Lett.* 50 (1983), no. 23, 1873-1876.
- [25] Puelz, C.; Embree, M.; Fillman, J., Spectral approximation for quasiperiodic Jacobi operators. *Integral Equations Operator Theory*, 82 (2015), no. 4, 533-554.
- [26] Raymond L., A constructive gap labelling for the discrete Schrödinger operator on a quasiperiodic chain, *Preprint*, 1997.
- [27] Sütö A., The spectrum of a quasiperiodic Schrödinger operator, *Commun. Math. Phys.* 111(1987), 409-415.
- [28] Toda M., Theory of Nonlinear Lattices, 2nd enlarged edn, Solid-State Sciences 20, Springer- Verlag, 1989, Chap. 4.

(Q.H. LIU) DEPARTMENT OF COMPUTER SCIENCE, BEIJING INSTITUTE OF TECHNOLOGY, BEIJING 100081, P. R. CHINA.

Email address: qhliu@bit.edu.cn

(Y.H. QU) DEPARTMENT OF MATHEMATICAL SCIENCE, TSINGHUA UNIVERSITY, BEIJING 100084, P. R. CHINA.

Email address: yhqu@tsinghua.edu.cn

(X. Yao) SCHOOL OF MATHEMATICAL SCIENCES AND LPMC, NANKAI UNIVERSITY, TIANJIN, 300071, P. R. CHINA.

Email address: yaoxiao@nankai.edu.cn